

Guiding coral reef futures in the Anthropocene

Norstrom, Albert V.; Nystrom, Magnus; Jouffray, Jean-Baptiste; Folke, Carl; Graham, Nicholas A.J.; Moberg, Frederik; Olsson, Per; Williams, Gareth

Frontiers in Ecology and the Environment

DOI: 10.1002/fee.1427

Published: 01/11/2016

Peer reviewed version

Cyswllt i'r cyhoeddiad / Link to publication

Dyfyniad o'r fersiwn a gyhoeddwyd / Citation for published version (APA): Norstrom, A. V., Nystrom, M., Jouffray, J.-B., Folke, C., Graham, N. A. J., Moberg, F., Olsson, P., & Williams, G. (2016). Guiding coral reef futures in the Anthropocene. *Frontiers in Ecology and the Environment*. https://doi.org/10.1002/fee.1427

Hawliau Cyffredinol / General rights Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

· Users may download and print one copy of any publication from the public portal for the purpose of private study or research.

- You may not further distribute the material or use it for any profit-making activity or commercial gain
 You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

1	Guiding coral reef futures in the Anthropocene
2	
3	
4	Albert V Norström ^{1*} , Magnus Nyström ¹ , Jean-Baptiste Jouffray ^{1,2} , Carl Folke ^{1,2,3} ,
5	Nicholas A J Graham ^{4,5} , Fredrik Moberg ¹ , Per Olsson ¹ ., Gareth J Williams ^{6,7}
6	
7	
8	¹⁾ Stockholm Resilience Centre, Stockholm University, Sweden
9	*(albert.norstrom@su.se)
10	²⁾ Global Economic Dynamics and the Biosphere Family Erling-Persson Academy
11	Programme, Royal Swedish Academy of Sciences, Sweden
12	³⁾ The Beijer Institute, Royal Swedish Academy of Sciences, Sweden
13	⁴⁾ Lancaster Environment Centre, Lancaster University, Lancaster, LA1 4YQ, UK
14	⁵⁾ ARC Centre of Excellence for Coral Reef Studies, James Cook University,
15	Townsville, QLD 4811, Australia
16	⁶⁾ Center for Marine Biodiversity & Conservation, Scripps Institution of
17	Oceanography, La Jolla, CA, USA
18	⁷⁾ School of Ocean Sciences, Bangor University, Anglesey, LL59 5AB, UK
19	
20	

23 Abstract

25	Human changes to the Earth now rival the great forces of nature, and have shepherded
26	us into a new planetary era – the Anthropocene. Changes include profound, and often
27	surprising, alterations to coral reef ecosystems and the services they provide human
28	societies. Ensuring their future in the Anthropocene will require that key drivers of
29	coral reef change – fishing, water quality and anthropogenic climate change – stay
30	within acceptable levels, or "safe operating spaces". The capacity to remain within
31	these safe operating spaces hinges on understanding the local, but also the
32	increasingly global and cross-scale, socio-economic causes of these human drivers of
33	change. Consequently, even successful local and regional management efforts will
34	fail if current decision making and institution-building around coral reef systems
35	remains fragmented, poorly coordinated, and unable to keep pace with the escalating
36	speed of social, technological and ecological change in the Anthropocene.
37	
38	In a nutshell
39	
40	• Key drivers of coral reef change should be kept at a "safe" distance from
41	dangerous levels or potential thresholds.
42	
43	• Fishable biomass should stay within or above 500-250 kg ha ⁻¹ , and
44	chlorophyll between 0.45-0.55 μ g L ⁻¹ .
45	
46	• CO ₂ concentrations should remain within or below 340-480 ppm and 480-750
47	ppm to avoid mass bleaching events and ocean acidification respectively.
48	
49	• The capacity to stay within the safe operating spaces is challenged by socio-
50	economic factors, including globalized drivers of change such as trade, human
51	migration and land-use change.
52	
53	• Adaptive and multi-level governance that involves state and non-state actors is
54	necessary keep pace with the escalating speed of change in the Anthropocene.

55 **Coral reefs in the Anthropocene**

56

57 There is growing scientific recognition that we live in the Anthropocene, an era where 58 humans have become a dominant force of planetary change (Steffen et al. 2011). 59 Changes include profound alterations of the Earth's marine and terrestrial ecosystems 60 and the services they provide to globally interconnected societies and economies 61 (Carpenter et al. 2009). Human migration, international trade, transnational land 62 acquisitions, spread of invasive species and technology diffusion occur at 63 unprecedented scales, underpinned by a global infrastructure that facilitates 64 movement of people, goods, services, diseases and information (Reid et al. 2010). 65 Actions taken in seemingly independent places increasingly affect the interlinked 66 global social-ecological system in unexpected ways, with surprising mixes of 67 immediate consequences as well as cascading and distant effects (Liu et al. 2013). 68 69 Coral reefs are informative examples of the key social-ecological challenges and 70 interactions playing out in the Anthropocene. They are economic and social assets 71 that have exhibited stability on centennial to millennial scales, but have experienced

an unprecedented decline over the last 50 years (Hughes *et al.* 2010). Changes to reefs

73 in the Anthropocene are multifaceted and complex (Figure 1). Impacts of overfishing

and coastal pollution, which can be managed successfully at local scales, are

increasingly compounded by the more recent, superimposed impacts of global
warming and ocean acidification. These anthropogenic drivers of change are mediated

by underlying traits in the social sphere such as economic systems, demography,

cultural dimensions and societal norms. Many coral reefs have already shown signs of
 transgressing thresholds and have undergone regime shifts to alternate degraded states

80 (Norström *et al.* 2009). In many cases this is resulting in a reduction of ecosystem

81 services, such as tourism and fisheries that provide income and food security (Moberg

- and Folke 1999). On the other end of the spectrum, a few reefs are maintained in a
- 83 semi-pristine state due to their remoteness from direct human impact (Graham and
- 84 McClanahan 2013; Williams et al. 2015). An increasingly common scenario, however,

85 is that reefs change in composition to novel coral-dominated ecosystems while still

86 maintaining key functions and ecosystem services at relatively desirable levels

87 (Graham *et al.* 2014).

89 The interlinked social, economic and ecological challenges of the Anthropocene 90 call for broader transdisciplinary coral reef science that is complemented by 91 management and governance strategies that facilitate the stewardship of coral reefs. 92 Ecosystem stewardship has emerged as a powerful sustainability framework with a 93 central goal to sustain ecosystem capacity to provide services that support human 94 well-being under conditions of uncertainty and change (Chapin et al. 2010). Here we 95 draw on several areas of emerging transdisciplinary social-ecological research to 96 highlight three broad challenges that need to be addressed in the efforts towards 97 sustainable stewardship of coral reefs. We start by describing safe operating spaces 98 for the key drivers of change that must not be transgressed for coral reefs to continue 99 to develop and exist. We then explore some of the critical cross-scale social-100 ecological interactions that will increasingly challenge the capacity to remain within 101 these safe operating spaces, and propose ways to study these social-ecological 102 interconnections. Finally, we outline the governance and institutional factors that need 103 to be in place for navigating coral reefs towards a sustainable future. 104

104

88

105

106 Safe operating spaces for global coral reef change

107

108 Avoiding thresholds that trigger regime shifts is becoming a focal point of resilience-109 based management of coral reefs. However, despite recent advances in predicting 110 thresholds (Mumby et al. 2007; Graham et al. 2015) their global generalizability is 111 confounded by a strong dependence on the historical, geographic and environmental 112 context of the system. Furthermore, the ecosystem consequences of crossing 113 thresholds may lag by decades (or even centuries) and may not be obvious over 114 human time scales (Hughes et al. 2013). In the face of this uncertainty a 115 complementary approach has been to establish safe operating spaces for ecosystems 116 (Scheffer et al. 2015). This concept is different from identifying specific thresholds. 117 Safe operating spaces are set to maintain safe levels of human drivers to avoid the 118 long-term degradation of ecosystems, and societies that depend on them. The concept 119 neither assumes, nor rules out, the existence of thresholds and is applicable in 120 situations with different types of system responses to increased levels of different

121 drivers (Rockström et al. 2009; Hughes et al. 2013) (Figure 2). We set safe operating 122 spaces and zones of uncertainty for the key drivers of change on coral reefs; i) fishing 123 *ii) water quality, and iii) anthropogenic climate change* (i.e. sea surface temperature, 124 aragonite saturation levels, ocean acidification). The safe operating space (green 125 zones in Figure 3) indicates the values of the drivers set at a "safe" distance from 126 potentially dangerous levels or threshold points (where they exist). Defining the safe 127 operating spaces is challenging and involves uncertainty due to interactions among 128 drivers (WebPanel 1), variable responses within and among taxa, geographic variation, 129 data limitation and the scope for acclimation or adaptation of reef-organisms to 130 change (Mumby and Van Woesik 2014; Barkley et al. 2015). Consequently, a zone of 131 uncertainty is associated with each of the drivers (yellow zones in Figure 3). Moving 132 towards the "high risk" (red) zones represents an increasing probability of crossing a 133 critical threshold or accelerated decline (Steffen et al. 2015). The values we provide 134 should be regarded as guidelines that will become more accurate with increasing 135 studies and knowledge.

136

137 Fishing

138

139 Historical overfishing precedes all other pervasive human drivers of change on coral 140 reefs (Jackson et al. 2001). As predatory and herbivorous fish are removed from reef 141 ecosystems, the risk of crossing thresholds and undergoing regime shifts to 142 undesirable reef configurations increases. In order to set a safe operating range for 143 fishing, we draw on recent regional (McClanahan et al. 2011, 2015; Karr et al. 2015) 144 and global (MacNeil et al. 2015) assessments of the threshold and non-linear 145 dynamics associated with fishable biomass - an easily measured proxy of fishing 146 pressure - on reefs. Threshold points in the trend or variance associated with a range 147 of ecosystem processes (e.g. herbivory, predation), state variables (e.g. the ratio of 148 coral to macroalgae cover), fish community life history traits and functional 149 groupings were associated with fishable biomass levels between 25-50% of unfished 150 biomass (calculated from recovery trajectories in marine reserves, and unfished 151 reference sites in each region). The results of these studies suggest that maintaining 152 reefs in a desirable regime (i.e. low macroalgal cover, high coral cover, high fish 153 diversity) requires fishable biomass to be kept above 500 kg ha⁻¹, with a zone of

154 uncertainty between 500-250 kg ha⁻¹ (Figure 3).

155

156 Water quality

157

158 In many parts of the world, water quality (e.g. nutrient loads, pollutants, sediments) in 159 coastal areas is changing in response to rapid urbanization, increasing fertilizer use 160 and land use change. Poor water quality can disrupt coral reproduction and 161 recruitment, smother adult corals and favor algal proliferation (Fabricius 2005). A 162 representative proxy for overall water quality status, which is highly correlated to 163 nutrient status and phytoplankton biomass, is chlorophyll concentration (De'ath and 164 Fabricius 2010). Chlorophyll concentration on reefs is naturally variable (Gove et al. 165 2016) and across uninhabited Pacific coral reefs the abundance of reef-building corals increases as chlorophyll concentration rises from 0.05-0.20 μ g L⁻¹ (Williams et al. 166 167 2015). However, a large-scale assessment of the relationship between chlorophyll and 168 reef condition across the whole of the Great Barrier Reef in Australia, found critical levels of 0.45 μ g L⁻¹ chlorophyll beyond which macroalgal cover increased and hard 169 170 coral richness declined (De'ath and Fabricius 2010). Earlier, smaller-scale, studies 171 from Barbados and Hawaii also showed measurable negative changes at chlorophyll 172 annual means above 0.5 μ g L⁻¹ (Bell 1992). We therefore suggest a safe-operating space value of chlorophyll concentration below 0.45 μ g L⁻¹, and a zone of uncertainty 173 174 between 0.45-0.55 µg L⁻¹, for continental and archipelago reef systems (Figure 3).

175

176 Anthropogenic climate change

177

178 Human-induced increases in atmospheric CO₂ concentrations ([CO₂]_{atm}) have driven 179 rapid rises in sea surface temperatures (SST) and ongoing ocean acidification (OA). 180 The vulnerability of reef-building corals to the unprecedented rates of change in SST 181 has been well documented; when temperatures exceed summer maxima by 1°-2°C for 182 3-4 weeks coral bleaching and mortality occurs. It is the increased intensity and 183 frequency of episodes of ocean warming and associated mass bleaching events (i.e. 184 the significant bleaching of multiple coral species at a regional scale) that is 185 compromising the long-term integrity of coral reefs. If mass bleaching events become 186 annual or biennial events corals may experience chronic decline as a result of reduced

187 growth, calcification, fecundity and greater incidences of disease (Hoegh-Guldberg et 188 al. 2007). Models suggest that avoiding chronic mass bleaching events (i.e. annual or 189 biennial) for the majority of the world's coral reefs requires keeping [CO2]_{atm} levels 190 below 480 ppm (Donner et al. 2005; Hoegh-Guldberg et al. 2007), or even below 450 191 ppm (van Hooidonk *et al.* 2013). However, substantially lower levels of [CO₂]_{atm} have 192 been suggested based on conservative backcasting exercises that associate the advent 193 of highly destructive mass bleaching (e.g. the 1997/1998 mass bleaching event which 194 killed approximately 16% of coral communities globally), with [CO₂]_{atm} values of 340 195 ppm (Veron et al. 2009). We therefore suggest that the safe operating space to avoid 196 chronic mass bleaching ends at 340 ppm, with the zone of uncertainty ranging 197 between 340-480 ppm (Figure 3). With a current global value of 400 ppm it means

that reefs have already entered the zone of uncertainty.

199

200 Absorption of CO₂ by the ocean is reducing water pH and the saturation levels of 201 aragonite (Ω_{arag}), the principle crystalline form of calcium carbonate deposited in 202 coral skeletons. Coral reefs are generally found in regions with Ω_{arag} values greater 203 than 3.3, and this observation underlies projections of global coral reef decline as 204 $[CO_2]_{atm}$ approaches 480 ppm and Ω_{arag} drops below 3.3 (Hoegh-Guldberg 2010). 205 Models parameterized by field observations of coral community calcification as a 206 response to Ω_{arag} , SST and live coral cover values, predict that by the time $[CO_2]_{atm}$ 207 will reach 560 ppm almost all coral reefs will cease to grow and start to dissolve 208 (Silverman et al. 2009). However, internal pH up-regulation at the point of 209 calcification has been shown to reduce the vulnerability of corals to ocean 210 acidification, and varies among species (McCulloch et al. 2012). Studies from 211 naturally low-pH coral communities suggest that adaptation to low pH can occur over 212 long time scales (Barkley et al. 2015), but that many ecological properties might be 213 irreversibly damaged as pH drops below 7.8 at [CO2]atm 750 ppm (Fabricius et al. 214 2011). Consequently, we set a safe upper boundary associated with ocean 215 acidification at 480ppm, and a broad zone of uncertainty between 480-750 ppm 216 (Figure 3). 217 218

219 Coral reef social-ecological dynamics in the Anthropocene

221 The capacity to keep human drivers of change within safe operating spaces is 222 challenged by a broad range of socio-economic interactions and feedbacks between 223 reef systems and the human societies that depend on their goods and services (Panel 224 1). However, social-ecological dynamics in the Anthropocene are seldom just local or 225 place-specific, but rather influenced by multiple global drivers with complex 226 connections to other places that are now more prevalent, and occur more quickly, than 227 ever before (Liu et al. 2013). We highlight three transboundary interactions - trade, 228 human migration and foreign investments in land and large-scale land acquisitions 229 (land grabbing) - that will increasingly define coral reef social-ecological dynamics 230 (Figure 5).

231

220

232 Regional and global analyses suggest that access to external markets can affect 233 coral reef fish resources (Cinner et al. 2013). Aside from local consumptive markets, 234 the global aquarium trade targets over 1800 species of reef fishes and removes up to 235 30 million fish per year (Rhyne *et al.* 2012), while the live reef fish trade (LRFT) 236 involves the exploitation of coral reef fishes from across the Indo-Pacific to satiate 237 consumer demand in luxury seafood restaurants (Johnston and Yeeting 2006). 238 Similarly, many invertebrate reef fisheries are extensively embedded in global trade 239 networks composed by actors operating at different levels, including local fishers, 240 middlemen and consumers in areas far from the reefs themselves. A consequence of 241 this increased market connectivity and nestedness is that many local invertebrate and 242 reef fish stocks are sequentially depleted as a result of the rapid emergence of 243 specialized export markets and quick spatial shifts in exploitation (Scales *et al.* 2007; 244 Eriksson et al. 2015).

245

246 Human migration, in particular to coastal regions, is currently at unprecedented 247 levels (Ozden et al. 2011) and is forecast to increase as a response to the social-248 ecological changes associated with the Anthropocene. Consequently, local social-249 ecological dynamics will increasingly be sculpted by the complex flows of people 250 across and within administrative boundaries. Fishers associated with coral reefs are 251 already highly mobile in many regions and known to move to areas where the fish are 252 more easily caught (Pollnac et al. 2010). Coastal areas are often the targets for 253 internal migration in many countries, particularly as urban centers and industries

promising employment are commonly located at the coast. While mobility can be a
key strategy for coastal communities to cope with global change, it can also
exacerbate reef resource degradation through the concentration of fishing effort,
introduction of new technology and fishing gear, and the deterioration of traditional
rules and practices (Cassels *et al.* 2005).

259

260 A third important cluster of drivers are foreign investments in land and large-261 scale land acquisitions – commonly referred to as land grabbing - that are increasingly 262 driving land use change (Meyfroidt et al. 2013). Land use change is a substantial 263 threat to coral reefs, by directly affecting sediment, pollution and fresh water 264 discharge into coastal zones. Past examples show how large-scale land clearing driven 265 by intensive banana production, and exasperated by tourism development, has 266 depleted coral communities in certain Caribbean reefs (Cramer et al. 2012). More 267 recent modeling efforts are suggesting that human deforestation, primarily driven by 268 demand for agricultural land, mineral exploration and mining, will outweigh climate 269 change as the principal contributor to increased sedimentation of near-shore marine 270 environments in Madagascar (Maina et al. 2013). Similarly, the run-off from export 271 agriculture such as squash in Tonga and oil palm in Papua New Guinea is emerging as 272 a key driver of change in Pacific Island reefs (Hunt 2003).

273

274 Capturing and studying the growing importance of these complex social-275 ecological interconnections on coral reef systems is a key research challenge. 276 Research on land systems change has made progress, from which coral reef social-277 ecological systems research could learn. For example, cross-country statistical 278 analyses have shown that recent tropical deforestation is associated with international 279 trade of agricultural products and remote urban demand, rather than with rural 280 population growth (DeFries et al. 2010). This resonates with coral reef systems, 281 where access to markets (e.g. for exports or satisfying urban demand) is often a better 282 predictor of overall reef fish biomass than other local socio-economic and natural 283 drivers (Cinner et al. 2013). Land systems change research has also explored 284 "displacement" and "cascade effects" - the unintended negative consequences of 285 forest recovery beyond the borders of reforesting countries. For example, recent forest 286 transitions and forest protection policies in both developed and developing countries 287 have outsourced forest exploitation abroad via increased imports of wood and

288 agricultural products (Meyfroidt et al. 2013). Such approaches merge detailed 289 economic (forest product prices, imports and exports of wood products) and 290 environmental (land cover change) data. Similar analyses could be used to investigate 291 whether the positive relationship between socio-economic development and reef 292 condition in some parts of the world is due to displacement of domestic environment 293 impacts through trade, or because of other, local factors such as low dependence on 294 fishing and reduced use of potentially damaging gear (Cinner et al. 2009a). Similarly, 295 while Marine Protected Areas (MPAs) can displace fishing effort at a local scale, the 296 potential leakage of fishing effort across regions and national borders is a key 297 research gap - especially in light of current trends of establishing large mega-reserves 298 in many regions (Graham and McClanahan 2013). The approaches to analyze cross-299 scale linkages in coral reef social-ecological systems will be determined by the 300 specific context, research question and data available. Learning from other disciplines 301 and adapting existing methods and frameworks will speed these advances.

302

303

304 Multi-scale challenges require multi-level governance

305

306 Conventional approaches to deal with the decline of coral reefs, such as MPAs can 307 offer local socioeconomic and ecological benefits but are usually narrow in scope, 308 small-scale and often suffer from weak compliance and enforcement (Pollnac et al. 309 2010). Coral reef management is slowly shifting towards more systemic management 310 strategies that are collaborative (involving both state and non-state actors) and 311 adaptive. There is also increasing focus on ecosystem processes that underpin 312 resilience and actions that target social-ecological interactions across the wider 313 seascape (Panel 1). Advancing social-ecological and adaptive comanagement 314 approaches requires acknowledging the broader governance and institutional (norms 315 and rules) contexts that enable their successful implementation. For example, while 316 monitoring and experimentation are central tenets of adaptively managing coral reefs, 317 they have typically been carried out by scientists. Involving local resource users in the 318 monitoring process enhances incentives to learn about local ecosystem dynamics and 319 facilitates collective action in line with the management objectives (Christie et al. 320 2009; Montambault et al. 2015). Initial support by local communities and government

- bodies is crucial (Olsson *et al.* 2004), and hinges on the management plans building
- 322 on existing rules and institutions, such as traditional tenure and community
- 323 committees. Research has also highlighted the role of key individuals that build
- 324 visions, foster trust and develop partnerships between stakeholders (e.g., community
- 325 groups, religious leaders, government authorities, NGOs and researchers) and
- 326 facilitate the participatory and inclusive process that sets and adapts the management
- 327 strategies to local contexts (Schultz *et al.* 2015).
- 328

329 Local management efforts alone will not be able to keep pace with the escalating 330 speed of technological and ecological change in the Anthropocene. The sustainability 331 challenges of an increasingly interconnected world call for developing governance 332 systems that foster international and cross-sectorial cooperation. An international 333 binding treaty to alleviate coral reef degradation has not materialized, despite a 334 number of favorable factors, such as the presence of supporting business interests, 335 public appeal and the relatively small number of nations involved (Dimitrov 2002). 336 However, the socio-economic and environmental issues facing marine ecosystems are 337 finally receiving a focus equal to their terrestrial counterparts. For example, Goal 14 338 of the newly adopted United Nations Sustainable Development Goals encompasses 339 ten targets for sustainable development in the oceans, while one of Convention of 340 Biological Diversity's Aichi Targets explicitly calls to minimize anthropogenic 341 pressures on coral reefs and maintain their integrity and functioning. This momentum 342 could provide a window of opportunity for organizations such as the International 343 Coral Reef Initiative (ICRI) and the International Society for Reef Studies (ISRS) to 344 more ambitiously engage with high-level policy processes across different domains, 345 such as climate change and trade, and bring issues of coral reef sustainability on the 346 negotiating tables. Crucially, it will require strategic collaborations with emerging 347 regional management initiatives such as the Micronesia Challenge, the Caribbean 348 Challenge Initiative, Western Indian Ocean Coastal Challenge and Coral Triangle 349 Initiative. These serve as practical operating platforms convening political leaders, 350 non-governmental organizations, coastal communities and scientists with the aim of 351 sustainably managing marine and coastal resources (Rosen and Olsson 2013; Johnson 352 et al. 2014). Such multi-level governance systems involving state and non-state actors 353 have emerged in response to other complex transnational and regional collective 354 action problems like ocean acidification (Galaz et al. 2012) and fisheries

- 355 overexploitation (Österblom and Sumaila 2011) when enforceable global agreements
- are missing or have failed. Importantly, it has been shown that they foster learning
- between several types of key individuals and organizations, nurture trust and can
- 358 facilitate collective action toward common goals.
- 359

360 **Conclusions**

361

362 Ensuring sustainable coral reef futures in the Anthropocene will require human 363 drivers of change to stay within safe levels, far from dangerous thresholds. Local and 364 regional actions can enhance resilience and limit the longer-term damage from 365 climate-related effects by keeping fishing and water quality targets within their safe 366 operating spaces. It is critical that such management targets are applied within a 367 broader adaptive management context, which allows for learning and experimentation, 368 and tolerates variability within the safe operating spaces. Management strategies that 369 reduce the short-term variance near the boundary levels run the risk of narrowing the 370 safe operating space, with potentially catastrophic consequences (Carpenter et al. 371 2015). Understanding the social dynamics underlying these drivers of change 372 becomes crucial. New research is required to better capture how social-ecological 373 dynamics are affected by interactions between regions, and across large distances. We 374 reinforce the urgency for coral reef science to deeply engage with emerging regional 375 management initiatives (such as the Micronesia Challenge and Coral Triangle 376 Initiative) and the international policy arena (such as the United Nations Framework 377 Convention on Climate Change) to work for sharp reductions of greenhouse gas 378 emissions and the implementation of the Sustainable Development Goals. With the 379 second global mass bleaching event currently underway, it is clearly urgent to up 380 efforts to help steer reefs toward a more sustainable future 381 382 383

384

385 **Panel 1. Social-ecological research on coral reefs**

387 Coral reef social-ecological systems (SES) research has grown exponentially over the 388 past 25 years (Figure 4), with a strong emphasis at the local or regional scale. One 389 sub-set of coral SES research has focused on ecosystem services and human 390 wellbeing in tropical coastal communities that exhibit livelihood strategies that are 391 strongly tied to coral reefs. Ecosystem services associated with coral reefs extend 392 beyond food production and encompass a broad bundle of provisioning, regulating 393 and cultural services that varies across regions and contexts (Moberg and Folke 1999). 394 Novel insights are uncovering how different social, institutional and knowledge 395 mechanisms determine access to these different ecosystem services, and how 396 preferences for ecosystem services are linked to inherent psychological values held by 397 different kinds of people (Hicks and Cinner 2014; Hicks et al. 2015). Another sub-set 398 of this research has highlighted how the combination of weak or missing institutions, 399 a lack of individual and institutional leadership, few alternative livelihoods and 400 inadequate financial capacity can trap a coral reef SES in undesirable and 401 unsustainable pathways (Cinner 2011; Sale et al. 2014). Finally, a third broad 402 category of research is using different diagnostic SES frameworks to understand how 403 the ecological performance of fisheries and marine reserves is related to different 404 socioeconomic variables of associated coastal communities (Pollnac et al. 2010). 405

406 This body of research is also beginning to underlie novel approaches to 407 management that specifically include the local human communities dependent on 408 coral reefs. For example, different fisheries management tools (such as gear-based 409 management and size-selectivity) can help to maintain key ecosystem functions and 410 significant yields of provisioning and other services (Johnson 2010). The emergence 411 of property rights systems for coral reef fisheries, such as Kenya's recent Beach 412 Management Unit legislation, allows local communities to deal with transgressions 413 committed by outside poachers or globalized "roving-bandit" type exploitation 414 (Cinner et al. 2009b). Combining local knowledge with contemporary science is 415 developing 'hybrid' co-management systems that are having tangible conservation 416 benefits (Aswani et al. 2012). Finally, there are increased calls for adaptive 417 management efforts that emphasize collaborative "management experiments" and the 418 importance of learning from these experiments. For example, viewing the 419 implementation of MPAs as a hypothesis driven process that is monitored would

420	enable managers to	learn	what	works and	better	deal	with	the	uncertain	futures	of

- 421 coral reefs.
- 422
- 423

424 **References**

426	Aswani S, Christie P, Muthiga NA, et al. 2012. The way forward with ecosystem-
427	based management in tropical contexts: reconciling with existing management
428	systems. Mar Policy 36: 1–10.

- Barkley HC, Cohen AL, Golbuu Y, *et al.* 2015. Changes in coral reef communities
 across a natural gradient in seawater pH. *Sci Adv* 1: e1500328.
- Bell P. 1992. Eutrophication and coral reefs: examples in the Great Barrier Reef
 lagoon. *Water Res* 26: 553–68.
- 433 Carpenter SR, Mooney HA, Agard J, *et al.* 2009. Science for managing ecosystem
 434 services: Beyond the Millennium Ecosystem Assessment. *Proc Natl Acad Sci*435 106: 1305-12.
- 436 Carpenter SR, Brock WA, Folke C, *et al.* 2015. Allowing variance may enlarge the
 437 safe operating space for exploited ecosystems. *Proc Natl Acad Sci*: 201511804.
- 438 Cassels S, Curran SR, and Kramer R. 2005. Do migrants degrade coastal
- environments? Migration, natural resource extraction and poverty in North
 Sulawesi, Indonesia. *Hum Ecol* 33: 329–63.
- 441 Chapin FS, Carpenter SR, Kofinas GP, *et al.* 2010. Ecosystem stewardship:
 442 sustainability strategies for a rapidly changing planet. *Trends Ecol Evol* 25: 241–
 443 9.
- 444 Christie P, Pollnac RB, Fluharty DL, *et al.* 2009. Tropical Marine EBM Feasibility: A
 445 Synthesis of Case Studies and Comparative Analyses. *Coast Manag* 37: 374–85.
- 446 Cinner JE. 2011. Social-ecological traps in reef fisheries. *Glob Environ Chang* 21:
 447 835–9.
- 448 Cinner JE, Graham NAJ, Huchery C, and MacNeil MA. 2013. Global effects of local

449 450	human population density and distance to markets on the condition of coral reef fisheries. <i>Conserv Biol</i> 27 : 453–8.
451 452	Cinner JE, McClanahan TR, Daw TM, <i>et al.</i> 2009a. Linking social and ecological systems to sustain coral reef fisheries. <i>Curr Biol</i> 19 : 206–12.
453 454 455	Cinner JE, Wamukota A, Randriamahazo H, and Rabearisoa A. 2009b. Toward institutions for community-based management of inshore marine resources in the Western Indian Ocean. <i>Mar Policy</i> 33 : 489–96.
456 457 458	 Cramer KL, Jackson JBC, Angioletti C V, <i>et al.</i> 2012. Anthropogenic mortality on coral reefs in Caribbean Panama predates coral disease and bleaching. <i>Ecol Lett</i> 15: 561–7.
459 460	De'ath G and Fabricius K. 2010. Water quality as a regional driver of coral biodiversity and macroalgae on the Great Barrier Reef. <i>Ecol Appl</i> 20 : 840–50.
461 462 463	 DeFries RS, Rudel T, Uriarte M, and Hansen M. 2010. Deforestation driven by urban population growth and agricultural trade in the twenty-first century. <i>Nat Geosci</i> 3: 178–81.
464 465	Dimitrov RS. 2002. Confronting nonregimes: science and international coral reef policy. <i>J Environ Dev</i> 11 : 53–78.
466 467 468	Donner SD, Skirving WJ, Little CM, <i>et al.</i> 2005. Global assessment of coral bleaching and required rates of adaptation under climate change. <i>Glob Chang Biol</i> 11 : 2251–65.
469 470	Eriksson H, Österblom H, Crona B, <i>et al.</i> 2015. Contagious exploitation of marine resources. <i>Front Ecol Environ</i> 13 : 435–40.
471 472	Fabricius K. 2005. Effects of terrestrial runoff on the ecology of corals and coral reefs: review and synthesis. <i>Mar Pollut Bull</i> 50 : 125–46.
473 474 475	 Fabricius KE, Langdon C, Uthicke S, <i>et al.</i> 2011. Losers and winners in coral reefs acclimatized to elevated carbon dioxide concentrations. <i>Nat Clim Chang</i> 1: 165–9.
476 477 478	Galaz V, Crona B, Österblom H, <i>et al.</i> 2012. Polycentric systems and interacting planetary boundaries — Emerging governance of climate change–ocean acidification–marine biodiversity. <i>Ecol Econ</i> 81 : 21–32.

- Gove JM, McManus MA, Neuheimer AB, *et al.* 2016 Near-island biological hotspots
 in barren ocean basins. *Nat Commun* 7: 10581.
- 481 Graham NA, Cinner JE, Norström A V, and Nyström M. 2014. Coral reefs as novel
 482 ecosystems: embracing new futures. *Curr Opin Environ Sustain* 7: 9–14.
- 483 Graham NAJ, Jennings S, Macneil MA, *et al.* 2015. Predicting climate-driven regime
 484 shifts versus rebound potential in coral reefs. *Nature* 518: 94–7.
- 485 Graham NAJ and McClanahan TR. 2013. The last call for marine wilderness?
 486 *Bioscience* 63: 397–402.
- Hicks CC and Cinner JE. 2014. Social, institutional, and knowledge mechanisms
 mediate diverse ecosystem service benefits from coral reefs. *Proc Natl Acad Sci*111: 17791–6.
- 490 Hicks CC, Cinner JE, Stoeckl N, and McClanahan TR. 2015. Linking ecosystem
 491 services and human-values theory. *Conserv Biol* 29: 1471-80.
- Hoegh-Guldberg O. 2010. Coral reef ecosystems and anthropogenic climate change. *Reg Environ Chang* 11: 215–27.
- Hoegh-Guldberg O, Mumby PJ, Hooten AJ, *et al.* 2007. Coral reefs under rapid
 climate change and ocean acidification. *Science* 318: 1737–42.
- Hooidonk R van, Maynard J, and Planes S. 2013. Temporary refugia for coral reefs in
 a warming world. *Nat Clim Chang* 3: 508–11.
- Hughes TP, Graham NAJ, Jackson JBC, *et al.* 2010. Rising to the challenge of
 sustaining coral reef resilience. *Trends Ecol Evol* 25: 633–42.
- Hughes TP, Carpenter S, Rockström J, *et al.* 2013. Multiscale regime shifts and
 planetary boundaries. *Trends Ecol Evol* 28: 389-95
- Hunt C. 2003. Economic globalisation impacts on Pacific marine resources. *Mar Policy* 27: 79–85.
- Jackson JBC, Kirby MX, Berger WH, *et al.* 2001. Historical overfishing and the
 recent collapse of coastal ecosystems. *Science* 293: 629–38.
- Johnson AE. 2010. Reducing bycatch in coral reef trap fisheries: Escape gaps as a
 step towards sustainability. *Mar Ecol Prog Ser* 415: 201–9.

508 509 510	 Johnson DE, Martinez C, Vestergaard O, <i>et al.</i> 2014. Building the regional perspective: Platforms for success. <i>Aquat Conserv Mar Freshw Ecosyst</i> 24: 75–93.
511 512	Johnston B and Yeeting B. 2006. Economics and marketing of the live reef fish trade in Asia–Pacific.
513 514	Karr KA, Fujita R, Halpern BS, <i>et al.</i> 2015. Thresholds in Caribbean coral reefs: implications for ecosystem-based fishery management. <i>J Appl Ecol</i> 52 : 402-12.
515 516	Liu J, Hull V, Batistella M, <i>et al.</i> 2013. Framing Sustainability in a Telecoupled World. <i>Ecol Soc</i> 18 .
517 518	MacNeil MA, Graham N a. J, Cinner JE, <i>et al.</i> 2015. Recovery potential of the world's coral reef fishes. <i>Nature</i> 520 : 341–4.
519 520	Maina J, Moel H de, Zinke J, <i>et al.</i> 2013. Human deforestation outweighs future climate change impacts of sedimentation on coral reefs. <i>Nat Commun</i> 4 : 1986.
521 522 523	McClanahan TR, Graham NAJ, MacNeil MA, <i>et al.</i> 2011. Critical thresholds and tangible targets for ecosystem-based management of coral reef fisheries. <i>Proc Natl Acad Sci U S A</i> 108 : 17230–3.
524 525 526	McClanahan TR, Graham NAJ, MacNeil MA, and Cinner JE. 2015. Biomass-based targets and the management of multispecies coral reef fisheries. <i>Conserv Biol</i> 29 : 409–17.
527 528 529	McCulloch M, Falter J, Trotter J, and Montagna P. 2012. Coral resilience to ocean acidification and global warming through pH up-regulation. <i>Nat Clim Chang</i> 2 : 623–7.
530 531 532	Meyfroidt P, Lambin EF, Erb K-H, and Hertel TW. 2013. Globalization of land use: distant drivers of land change and geographic displacement of land use. <i>Curr</i> <i>Opin Environ Sustain</i> 5 : 438–44.
533 534	Moberg F and Folke C. 1999. Ecological goods and services of coral reef ecosystems. <i>Ecol Econ</i> 29 : 215–33.
535 536 537	Montambault JR, Wongbusarakum S, Leberer T, <i>et al.</i> 2015. Use of monitoring data to support conservation management and policy decisions in Micronesia. <i>Conserv Biol</i> 29 : 1279–89.

- Mumby PJ, Hastings A, and Edwards HJ. 2007. Thresholds and the resilience of
 Caribbean coral reefs. *Nature* 450: 98–101.
- Mumby PJ and Woesik R Van. 2014. Consequences of ecological, evolutionary and
 biogeochemical uncertainty for coral reef responses to climatic stress. *Curr Biol*24: R413–23.
- Norström A, Nyström M, Lokrantz J, and Folke C. 2009. Alternative states on coral
 reefs: beyond coral–macroalgal phase shifts. *Mar Ecol Prog Ser* 376: 295–306.
- 545 Olsson P, Folke C, and Berkes F. 2004. Adaptive comanagement for building
 546 resilience in social-ecological systems. *Environ Manage* 34: 75–90.
- 547 Österblom H and Sumaila UR. 2011. Toothfish crises, actor diversity and the
 548 emergence of compliance mechanisms in the Southern Ocean. *Glob Environ*549 *Chang* 21: 972–82.
- Ozden C, Parsons CR, Schiff M, and Walmsley TL. 2011. Where on Earth is
 everybody? The evolution of global bilateral migration 1960-2000. *World Bank Econ Rev* 25: 12–56.
- Pollnac R, Christie P, Cinner JE, *et al.* 2010. Marine reserves as linked socialecological systems. *Proc Natl Acad Sci U S A* 107: 18262–5.
- Reid W V., Chen D, Goldfarb L, *et al.* 2010. Earth system science for global
 sustainability: grand challenges. *Science* 330: 916–7.
- Rhyne AL, Tlusty MF, Schofield PJ, *et al.* 2012. Revealing the appetite of the marine
 aquarium fish trade: the volume and biodiversity of fish imported into the United
 States. *PLoS One* 7: e35808.
- 560 Rockström J, Steffen W, Noone K, *et al.* 2009. Planetary boundaries: exploring the
 561 safe operating space for humanity. *Ecol Soc* 114: art32.
- Sale PF, Agardy T, Ainsworth CH, *et al.* 2014. Transforming management of tropical
 coastal seas to cope with challenges of the 21st century. *Mar Pollut Bull* 85: 8–
 23.
- Scales H, Balmford A, and Manica A. 2007. Impacts of the live reef fish trade on
 populations of coral reef fish off northern Borneo. *Proc Biol Sci* 274: 989–94.

- 567 Scheffer BM, Barrett S, Folke C, *et al.* 2015. Creating a safe operating space for
 568 iconic ecosystems. *Science* 347: 1317–9.
- 569 Silverman J, Lazar B, Cao L, *et al.* 2009. Coral reefs may start dissolving when
 570 atmospheric CO 2 doubles. *Geophys Res Lett* 36: L05606.
- 571 Steffen W, Persson Å, Deutsch L, *et al.* 2011. The Anthropocene: from global change
 572 to planetary stewardship. *Ambio* 40: 739–61.
- 573 Steffen W, Richardson K, Rockstrom J, *et al.* 2015. Planetary boundaries: guiding
 574 human development on a changing planet. *Science* 347: 1259855.
- Veron JEN, Hoegh-Guldberg O, Lenton TM, *et al.* 2009. The coral reef crisis: the
 critical importance of<350 ppm CO2. *Mar Pollut Bull* 58: 1428–36.
- 577 Williams GJ, Gove JM, Eynaud Y, *et al.* 2015. Local human impacts decouple natural
- 578 biophysical relationships on Pacific coral reefs. *Ecography* **38**: 751–61.

581 **Figure captions**

582

583 Figure 1. Examples of multifaceted changes occurring on reefs in the Anthropocene. 584 (a) Many coral reefs have become degraded as a consequence of overfishing 585 (macroalgal dominated, *bottom panel*)(courtesy of J Lokrantz), decreased water 586 quality (corallimorph dominated, *left panel*) and climate change (bleached, *right* 587 panel)(courtesy of A Masslenikov), (b) Some reefs are maintained in a semi-pristine 588 state due to their remoteness from direct human impacts (courtesy of BJ Zgliczynski) 589 (c) Other reefs are changing in composition to novel coral-dominated ecosystems 590 (courtesy of M Vermeij)

591 Figure 2. Three potential ways a coral reef may respond to increased driver levels are 592 illustrated, and all three are congruent with the safe operating space concept. 593 Increased levels of certain drivers (e.g. overfishing) may trigger threshold responses (I 594 and II). In other cases the response may be a smoother acceleration towards a 595 deleterious state (III). The safe operating space (green zones) indicates the range of 596 driver values that are at a "safe" distance from potentially dangerous levels or 597 threshold points. The zone of uncertainty associated with each of the boundaries 598 (yellow zones) encapsulates the gaps in scientific knowledge and uncertainty due to 599 driver interaction, scope for adaptation and geographic variation. As driver values 600 move towards the "high risk" end of the zone of uncertainty, there is an increasing 601 probability of declining ecosystem state. Modified from Rockström et al. 2009 and 602 Hughes et al. 2013

Figure 3. The safe operating spaces, zones of uncertainty and zones of high risk of

the key drivers of change on coral reefs; i) fishing ii) water quality, and iii)

anthropogenic climate change (i.e. sea surface temperature and ocean acidification).

606 Figure 4 (to be embedded in Panel 1). The dramatic increase of coral reef social-

607 ecological research. An ISI Web of Knowledge literature survey showed that the

- number of papers containing the keywords "coral reef" together with either "social-
- 609 ecological", "socio-ecological", "social-environmental" or "socio-environmental" has
- 610 increased exponentially between 1990 (n = 1) and 2014 (n = 106).

611 Figure 5. Three global interactions that shape local social-ecological dynamics of 612 coral reefs: 1) Human migration to coastal areas can result in deterioration of 613 traditional rules and practices, enhance pollution and increase pressures on reef fish 614 stocks. Graph shows net global migration to coastal areas between 1970-2010, and 615 specifically in the regions housing the majority of the worlds coral reefs; 2) Land 616 grabbing is increasingly driving land use change, which is a threat to coral reefs by 617 directly affecting water quality (e.g. nutrient loads, pollutants, sediments). Graph 618 shows cumulative number of concluded land grab deals between 2000-2014 on a 619 global scale, and in countries that have coral reefs; 3) International trade of coral reef 620 products is driven by intensifying foreign consumer demand and better access to 621 markets. Graph shows US imports of chilled reef fish (groupers and snappers) and 622 live coral colonies between 1990-2014. Data sources and methods are explained in 623 WebPanel 2.

624

625