

**Long-term effects of altered pH and temperature on the feeding energetics of the Antarctic sea urchin, *Sterechinus neumayeri***

Morley, Simon, A.; Suckling, Coleen; Clark, Melody, S.; Cross, Emma, L. ; Peck, Lloyd, S.

Biodiversity

DOI:

[10.1080/14888386.2016.1174956](https://doi.org/10.1080/14888386.2016.1174956)

Published: 22/04/2016

Peer reviewed version

[Cyswllt i'r cyhoeddiad / Link to publication](#)

Dyfyniad o'r fersiwn a gyhoeddwyd / Citation for published version (APA):

Morley, S. A., Suckling, C., Clark, M. S., Cross, E. L., & Peck, L. S. (2016). Long-term effects of altered pH and temperature on the feeding energetics of the Antarctic sea urchin, *Sterechinus neumayeri*. *Biodiversity*, 17(1-2), 34-45. <https://doi.org/10.1080/14888386.2016.1174956>

Hawliau Cyffredinol / General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

1 **Long term effects of altered pH and temperature on the feeding**
2 **energetics of the Antarctic sea urchin, *Sterechinus neumayeri*.**

3 Simon A. Morley*^a, Coleen C. Suckling^{a,b}, Melody S. Clark^a, Emma L. Cross^{a,c}
4 and Lloyd S. Peck^a

5 ^aBritish Antarctic Survey, Natural Environment Research Council, High Cross, Madingley Road,
6 Cambridge, Cambridgeshire, CB30ET. UK.

7 ^bCurrent address: School of Ocean Sciences, Bangor University, Anglesey, Wales, LL595AB, UK

8 ^cDepartment of Earth Sciences, University of Cambridge, Cambridge, UK;

9 *Corresponding author: smor@bas.ac.uk, Tel: 01223 221554, Fax: 01223 221259

10 Running Title: Long term OA energy budget

11

12 **Summary**

13 This study investigated the effects of long-term incubation to near-future combined warming
14 (+2 °C) and ocean acidification (-0.3 and -0.5 pH units) stressors, relative to current
15 conditions (-0.3 °C and pH 8.0), on the energetics of food processing in the Antarctic sea
16 urchin, *Sterechinus neumayeri*. After an extended incubation of 40 months, energy absorbed,
17 energy lost through respiration and lost as waste were monitored through two feeding cycles.
18 Growth parameters (mass of somatic and gonad tissues and the CHN content of gonad) were
19 also measured. There were no significant effects of combined ocean acidification (OA) and
20 temperature stressors on the growth of somatic or reproductive tissue. Despite more food
21 being consumed in the low temperature control, once food processing and maintenance costs
22 were subtracted, there were no significant effects of treatment on the scope for growth. The
23 biggest significant differences were between food consumed during the two feeding cycles.
24 More food was consumed by the low temperature (0°C) control animals, indicating a
25 potential effect of the changed conditions on digestive efficiency. Also in November, more
26 food was consumed, with a higher absorption efficiency which resulted in a higher scope for
27 growth in November than September, which may reflect increased energetic needs associated
28 with a switch to summer physiology. The effect of endogenous seasonal cycles and
29 environmental variability on organism capacity is discussed.

30

31 **Key-words** carbonate saturation; climate change; echinoderm; energetics; ocean
32 acidification; physiology; resilience

33

34 **Introduction**

35 To predict future patterns of biodiversity it is essential to understand the mechanisms that will
36 determine organism vulnerability. Of the physical factors affecting ectotherms, temperature is
37 one of the most extensively studied and global patterns of thermal tolerance have improved
38 our understanding of how environment correlates with physiological capacities (Gaston *et al.*
39 2009; Sunday, Bates & Dulvy 2011; Peck *et al.* 2014). Warming oceans increase the body
40 temperature of marine ectotherms, which alters the rates of all organism biochemical
41 reactions (Hochachka & Somero 2002). The vulnerability of organisms to warming therefore
42 depends on the characteristics of their thermal tolerance windows and both their
43 physiological plasticity and adaptive capacity to alter these windows (Angilletta 2009;
44 Somero 2012). Whilst the distributions of many marine species are shifting in response to the
45 rate of environmental warming (Appelhans *et al.* 2014), the effects of temperature do not
46 work in isolation. Within the marine environment the interacting effects of increasing
47 temperature and ocean acidification are predicted to be two of the key factors driving range
48 shifts (Pörtner 2012).

49

50 Ocean acidification is likely to have wide ranging effects on marine invertebrates,
51 particularly those with calcified skeletons. The absorption of anthropogenic carbon dioxide
52 into shallow seas is leading to a reduction in carbonate mineral saturation states, particularly
53 aragonite (McNeil & Matear 2008; Fabry *et al.* 2009). This could either result in altered
54 skeletal structure (Bray, Pancucci-Papadopoulou & Hall-Spencer 2014), potentially altering
55 predator prey interactions (Watson *et al.* 2012), or, if skeletal structure is maintained, the
56 costs of producing skeleton may increase (Wood, Spicer & Widdicombe 2008). Ocean
57 acidification may also alter the balance of metabolic costs, as extra energy is required to

58 maintain the homeostasis of inner body fluids against hypercapnia (internal CO₂) and acidosis
59 (reduction of internal pH; Wood 1993; Pörtner, Bock & Reipschlag 2000; Melzner *et al.*
60 2009; Spicer *et al.* 2011). However, more subtle changes have been identified, which would
61 not necessarily be predicted by the effects of calcium ion concentration on skeletal structure,
62 such as the ability to detect prey, aerobic scope and behaviour (Munday, Crawley & Nilsson
63 2009; Munday *et al.* 2009; Dixson, Munday & Jones 2010).

64

65 The shallow seas around the Antarctic Peninsula have one of the least variable thermal
66 regimes on the planet, with a 3-4 °C annual sea surface temperature range (Peck, Convey &
67 Barnes 2006). Consequently, many Antarctic marine species are stenothermal, with generally
68 poor capacities to cope with elevated temperatures (Pörtner, Peck & Somero 2007).

69 Acclimation is known to take longer in Antarctic marine invertebrates (Morley *et al.* 2011;
70 Peck *et al.* 2014) and their slow generation times and lower fecundity are expected to reduce
71 the capacity for adaptive change (Somero 2010; Peck 2011; Peck *et al.* 2014). Carbon dioxide
72 is more soluble in cold waters (Guinotte & Fabry 2008) and so high latitude oceans are also
73 expected to be amongst the first to become under-saturated with respect to calcite and
74 aragonite (McNeil & Matear 2008). The effects of temperature and ocean acidification are
75 therefore expected to have greater effects on Polar shallow water communities than at lower
76 latitudes (Hofmann *et al.* 2010). Recent studies have, however, shown that aragonite
77 saturation state varies markedly, between 0.8 and 3.9 off the Western Antarctic Peninsula
78 (WAP; Bjork *et al.* 2014; Collard *et al.* 2015). This high natural variability may result in
79 species from the WAP having the physiological capacity to cope with variation in carbonate
80 saturation state.

81

82 The Antarctic sea urchin, *Sterechinus neumayeri*, is an important component of shallow water
83 ecosystems throughout the Southern Ocean (Fabry *et al.* 2009). *S. neumayeri* are omnivorous,
84 benthic pioneer species, occurring in high densities in recent iceberg scours, where a large
85 portion of their diet comes from scavenging on dead organisms. Any major effect of future
86 conditions on this keystone species could lead to dramatic shifts in Antarctic benthic food
87 webs. Due to the high Magnesium calcite composition of echinoid skeletons they are a
88 taxonomic group which was predicted to be particularly susceptible to the effects of ocean
89 acidification (Sewell & Hofmann 2011), although recent studies have shown that some
90 echinoids are quite resilient (Wittmann & Pörtner 2013; Collard *et al.* 2015; Suckling *et al.*
91 2015). Studies are therefore required to determine the capacity of *S. neumayeri* to future
92 temperature and ocean acidification allowing predictions to estimate their future role as a
93 key-stone species in shallow Southern Ocean. In our previous investigations of the same *S.*
94 *neumayeri* used in this current study, whilst reproduction and larval development were
95 partially acclimated, adult somatic, skeletal growth and reproduction were fully acclimated to
96 altered conditions after 8 months (Suckling *et al.* 2015). In our previous study, *S. neumayeri*
97 were fed *ad libitum* and food consumption was not recorded (Suckling *et al.* 2015). The aim
98 of the current project was therefore to determine if there were subtle changes in the energetics
99 of the same adult Antarctic urchins, *Sterechinus neumayeri*, after a further 16 months
100 incubation (40 months in total) to a combination of elevated temperature and $p\text{CO}_2$
101 treatments, which would not have been detected using techniques in our previous study
102 (Suckling *et al.* 2015). Specifically, food consumption and the energetic costs of maintenance
103 and food processing were investigated to examine if acclimation to predicted future
104 conditions resulted in any changes in the energy budget, which could influence the scope for
105 growth and long-term resilience to altered environmental conditions.

106

107 **Material and methods**

108 **Animal collection and incubation**

109

110 Adult *Sterechinus neumayeri* were collected by SCUBA divers in the austral summer of
111 2008-2009 from 5-10 m depth at South Cove, Ryder Bay, Antarctic Peninsula (67°34' S,
112 68°08' W). Environmental conditions in Ryder Bay at 5-10m depth consist of seawater
113 temperatures that range from -1.8 to +2.0 °C, however, temperatures rarely exceed +0.5 °C
114 and salinity remains between 32.5-34.5 (Venables, Clarke & Meredith 2013). The animals
115 were transported to the UK and held in the British Antarctic Survey 0 °C re-circulating
116 aquarium in Cambridge for approximately 2 months before being introduced to the re-
117 circulating CO₂ microcosm system (adapted from Widdicombe and Needham (2007) and
118 fully detailed in Suckling *et al.* (2015)). Seawater was transported to Cambridge from the
119 North Sea which had an aragonite saturation state (0.75) slightly lower than the 0.8 to >3
120 range, but a pH of 8.0 which is within the range of typical values (7.6 to 8.3), for the Western
121 Antarctic Peninsula (Collard *et al.* 2015; Hauri *et al.* 2015). The treatments used in this study
122 were based on the IPCC 'business-as-usual' scenario with the forecasted reduction of 0.3 to
123 0.5 pH units in oceanic surface waters by the year 2100 (Barbarino & Lourenco 2009) and a
124 predicted rise in surface sea temperature of 2.0 °C. The four treatment combinations were: 1)
125 Low temperature control, present day temperature (-0.3 °C) and pH (pH 8.0); 2) High
126 temperature control, elevated temperature (2 °C) and current pH (pH 8.0); 3) -0.3 pH,
127 elevated temperatures (2 °C) and moderate acidification (pH 7.8) and 4) -0.5 pH, elevated
128 temperature (2 °C) and high acidification (pH 7.5). Urchins were incubated in microcosms
129 under the 4 treatment conditions for 40 months (beginning June 2009).

130

131 In the two microcosms with reduced pH treatments (200 L), UV disinfection and 50 μm
132 filtered seawater was delivered to 80L closed cylindrical mixing tanks. CO_2 gas (British
133 Oxygen Company) was introduced via a ceramic diffuser using an Aquamedic pH controlled
134 computer and electrode system and mixed with seawater by an Aquamedic ocean runner
135 power head 2000. Treated seawater was gravity fed to each experimental tank at a rate of
136 $0.56 \pm 0.03 \text{ L min}^{-1}$. The pH control mesocosm had a similar header tank, without a pH
137 computer controller, but with an Aquamedic ocean runner power head 2000. The low
138 temperature control animals were kept in a recirculating aquarium facility with identical pre-
139 treatment of water.

140

141 Seawater pH was initially at control levels in all tanks, with the urchins acclimated to these
142 tank conditions for 14 days prior to starting the incubations. The pH of the sea water in
143 selected tanks was then gradually decreased in equal twice daily increments over a period of
144 3 days until the desired pH target was achieved.

145

146 **Water chemistry**

147

148 Temperature was recorded daily for all treatments ($^{\circ}\text{C}$; Digital Testo 106) and the room
149 temperature adjusted as required. Once weekly temperature, salinity (Tropical Marine Centre
150 V2 Handheld refractometer), pH_{NIST} (temperature compensated; HANNA bench top meter
151 pH/ORP 115 v pH21-01) and TCO_2 (mmol L^{-1} ; Ciba Corning TCO_2 Analyzer 965, Olympic
152 Analytical, UK) were measured and recorded. The TCO_2 analyzer was calibrated with 2 g L^{-1}
153 CO_2 standard prior to measurements. Aquamedic pH probes were calibrated twice weekly
154 with NIST certified pH buffer solutions and CO_2 gas flow into the header tank was adjusted

155 accordingly. Seawater samples were also analysed for phosphate and silicate levels according
156 to Nickell *et al.* (2003).

157

158 Seawater quality in randomly selected individual urchin containers was assessed every 2-3
159 days using Nutrafin Aquarium test kits. Ammonia, nitrite and nitrates were maintained well
160 below 0.4, 0.2 and 5 mg L respectively by a combination of biological filtration, protein
161 skimming and partial seawater exchanges (approximately 5-15% every 2-3 days) to prevent
162 toxicity from metabolic by-products. A 12:12h light dark cycle was maintained throughout.

163

164 **Physiological Measurements**

165 The urchins used in the current study were reared in the same incubation system for a further
166 16 months (in addition to the previous 24 months; Suckling *et al.* 2015) before being used for
167 trials to measure the energetics of feeding and growth in September and November. This
168 coincided with the summer period when energy is partitioned towards maturing gonads in the
169 wild (Brockington & Peck 2001). For each feeding trial, nine or ten *S. neumayeri* were
170 chosen randomly from each treatment. Within each microcosm, specimens were separated by
171 placing them in individually labelled 300 cm³ containers. Each container had a coarse mesh
172 lid that allowed free exchange of water within each microcosm, but retained the urchin and
173 any food or faeces. To measure individual energy budgets, energy absorbed from food was
174 calculated from the quantity of food consumed and the organic mass of faeces produced. The
175 energy lost through maintenance and food processing was calculated from measurements of
176 oxygen consumption and ammonia and urea (nitrogenous waste) production, both before and
177 six days after feeding.

178

179 For the 40 month incubation period, *S. neumayeri* were fed every two weeks (Suckling *et al.*
180 2015) but trials showed that faecal production, and elevated waste production, continued for
181 longer than two weeks, up to 18 days (pers obs). To ensure a full food processing cycle was
182 measured during the experimental period (September to November), from August, *S.*
183 *neumayeri* were therefore fed every 3 weeks. *S. neumayeri* were fed individually with an
184 excess of fish fillets, *Polachius virens*, (0.48 ± 0.03 g wet mass, 4% of mean wet body mass)
185 and allowed to feed for 48 hours before uneaten food was collected and weighed. A high
186 protein diet is representative of the broad diet in the field whilst importantly providing an
187 easily quantifiable ration. The water uptake, and concomitant increase in weight of uneaten
188 food, was measured through trials in the same microcosms. Faeces were collected every 2
189 days and dried until faecal production had stopped. Total faecal dry mass (dried at 60 °C to
190 constant mass) and ash free dry mass (AFDM), calculated by subtraction following ignition
191 for 24 hours at 475 °C, were then determined.

192

193 To measure routine and feeding respiratory costs, oxygen consumption was measured before
194 and 6 days post-feeding (defined as pre and post feeding) using closed cell respirometry (
195 following, Obermüller *et al.* (2010). The night before experiments *S. neumayeri* were
196 transferred from their individual containers into respirometers with mesh lids. Before they
197 were closed, respirometers were flushed with seawater from the experimental system,
198 ensuring that any faeces were removed. Respirometers were matched to the size of *S.*
199 *neumayeri* so that a 10-20% reduction in oxygen was recorded in 3-5 hours, and experiments
200 were stopped before oxygen concentration fell below 80% of saturation values. Oxygen
201 concentration was measured using a Fibox-3 fibre optic oxygen sensor using an individually
202 calibrated oxygen sensitive foil glued into each respirometer (Morley *et al.* 2007). Two or

203 three blanks were run simultaneously to measure background changes in oxygen
204 concentration. The volume of each urchin was measured using Archimedes principle and the
205 volume of water in each respirometer calculated by subtraction.

206

207 At the end of respirometry measurement, the energy lost through nitrogenous waste
208 production was estimated by measuring ammonia and urea in the water in each respirometer.
209 Ammonia concentration in the chamber water was measured with a Turner Designs TD-700
210 fluorometer, fitted with a near UV mercury vapour lamp and a 310-390 nm excitation filter,
211 following the ortho-phthaldialdehyde (OPA) method of (Holmes *et al.* 1999). Samples were
212 analysed in triplicate and calibration was by standard dilution (four concentrations in
213 triplicate). The remaining seawater was frozen at -80 °C and urea concentration was
214 measured with a Lachat Quikchem 8500 flow injection auto-analyser at the Scottish
215 Association for Marine Science using the Lachat Method 10-206-00-1-A for determination of
216 urea in waters by flow injection analysis colorimetry. However, urea concentration in
217 samples was not significantly different from background levels (blank) in 12 of the 16
218 treatment-month-feeding combinations (Z-tests) confirming that *S. neumayeri* is largely
219 ammonotelic. Urea production was, therefore, excluded from further analysis.

220

221 The results from these measurements of oxygen consumption and nitrogen production were
222 used to calculate the atomic O:N ratio. O:N ratios vary from around 3 for protein only
223 catabolism to over 100 for diets dominated by lipids and carbohydrates (Mayzaud & Conover
224 1988). The change in O:N ratio before and after feeding therefore indicates how metabolic
225 substrate use varied through the period of feeding.

226

227 **Calculation of energy budget**

228

229 Energy available for growth was assessed by converting the physiological measurements into
230 energy equivalents, expressed in J individual⁻¹ h⁻¹. The energy budget modified from Winberg
231 (1960) partitioned the energy consumed from food (C) into: respiratory costs (R), waste
232 production (U) as ammonia or faeces (F) and the scope for growth (SfG):

233

234 $C = R + U + F + SfG$

235

236 The energy of the consumed food (C) was calculated using the supplier's (Waitrose)
237 nutritional information. Each 100 g (wet mass) of food contained 340 kJ of energy which was
238 largely in the form of protein (19.3 g of protein, 0.3 g of fat and 0 g of carbohydrate).

239

240 The time course of SDA has been calculated for several Antarctic marine invertebrates (2 to
241 13 days; Peck 1998; Robertson *et al.* 2001; Peck *et al.* 2008) but not *S. neumayeri*, so data
242 from another marine invertebrate, which also has a largely protein based metabolism, *Nacella*
243 *concinna*, was used (Fraser, Clarke & Peck 2002). The peak of SDA of *N. concinna* at 0°C
244 occurred between days 5 to 7 and so the oxygen consumption on day 6 was calculated to be
245 1.6 times the average daily elevation in oxygen consumption through the duration of the SDA
246 (Peck & Veal 2001). Therefore, to estimate the respiratory cost of processing food through
247 the whole *S. neumayeri* SDA, the value for the peak SDA, measured at 6 days post feeding

248 was divided by 1.6 to estimate the average daily increase in standard metabolic rate and
249 nitrogen waste production.

250

251 As the food was largely protein and nitrogenous waste production of *S. neumayeri* is
252 predominantly in the form of ammonia (Brockington & Peck 2001), a literature value of
253 $0.484 \text{ J } \mu\text{molO}_2^{-1}$ was used to convert oxygen consumption into an energy cost (Elliott &
254 Davison 1975). The energy loss through ammonia (U) were also calculated using literature
255 energy conversion factors of $0.348 \text{ J } \mu\text{mol}^{-1}$ (Elliott & Davison 1975).

256

257 Absorbed energy (A) was calculated from the proportion of the consumed AFDM (M_C) that
258 was retained and not egested as faecal AFDM (M_F):

259

$$260 \quad A = ((M_C - M_F) / M_C) * C$$

261

262 The scope for growth was calculated as:

$$263 \quad \text{SfG} = A - (R + U)$$

264

265 **Growth**

266 At the end of experiments in both September and November, *S. neumayeri* volume was
267 measured, urchins were then dissected and wet mass, dry mass and AFDM of gonad (G) and

268 the rest (S; mainly skeleton) of each animal were measured. Measurement of dry and ash
269 mass followed the same protocol as described above for faeces. From these masses Gonad
270 Somatic Index (%) was calculated as:

271

$$272 \text{ GSI} = \text{G}/(\text{G}+\text{S}) \times 100$$

273

274 Prior to drying, a small piece of gonad was weighed, dried and the total carbon, hydrogen and
275 nitrogen contents were measured in a CHN analyser Model CE 440 (Exeter Analytical, Inc.,
276 Massachusetts, USA). Each run was calibrated with acetanilide standards. From the CHN
277 data C:N and C:H ratios were calculated.

278

279 **Statistics**

280 Data were tested for normality with Anderson-Darling tests. Non-normal data were box cox
281 transformed to achieve normality before the fixed effect of treatment and the random effects
282 (to account for repeated measures) of both feeding and month were tested with ANOVA.
283 When a factor had a significant effect, *post hoc* Tukey tests were used. When a factor was
284 still non-normally distributed, even after transformation, differences were analysed using
285 non-parametric Kruskal Wallis tests.

286

287 **Results**

288 **Water chemistry**

289 In each system, once treatment conditions had been reached, water chemistry in the urchin
290 tanks was very stable through the 40 month duration of experiments (Table 1).

291

292 **Energetics**

293 *S. neumayeri* consumed more than twice as much food (105% more) in November than
294 September (ANOVA: $F_{(1,64)} = 35.7$, $P < 0.01$) and in both months more food was consumed
295 in the low temperature control than pH treatments (20-30% more consumed; ANOVA, $F_{(3,64)}$
296 $= 6.6$, $P < 0.01$; Tukey tests, $pH -0.3$, $T = 3.7$ and $pH, -0.5 T = 4.1$, $P < 0.01$; Table 2). The
297 absorption efficiency of organic matter from food was also lower in September than
298 November ($F_{(1,64)} = 40.8$, $P > 0.01$; Table 2). More energy was therefore absorbed (A) from
299 food in November than September 2012 ($F_{(1,64)} = 35.0$, $P < 0.01$; Fig. 1) and low temperature
300 control individuals absorbed significantly more energy than both pH treatments ($F_{(3,64)} = 5.8$,
301 $P < 0.01$; $pH -0.3$, $T = 3.5$ and $pH -0.5$, $T = 3.7$, $P < 0.01$).

302

303 There was no significant difference in oxygen consumption, between months ($F_{(1,139)} = 1.2$, P
304 $= 0.27$; Fig. 2) but there was a significant difference between treatments ($F_{(3,139)} = 3.8$, $P =$
305 0.01). The interaction between month and treatment was just non-significant ($F_{(3,139)} = 2.6$, P
306 $= 0.06$), so overall, lower oxygen consumption was observed in the low temperature control
307 compared to the high temperature control ($T = 2.6$, $P < 0.05$) and the -0.3 pH treatment ($T =$
308 3.0 , $P < 0.05$). Metabolic rate increased post feeding ($F_{(1,139)} = 6.3$, $P = 0.01$), resulting in an
309 increase in energy costs as food was processed and assimilated.

310

311 Ammonia excretion increased post feeding (Kruskal Wallis test: $H = 12.7$, $P < 0.01$; Fig. 3)
312 but there was no significant difference in the magnitude of this increase between months ($H =$
313 0.1 , $P = 0.8$; Fig. 1c) or treatments ($H = 2.2$, $P = 0.54$). The O:N ratio was generally between
314 2 and 4, indicating that the metabolic substrate was almost exclusively protein (Fig. 4). There
315 was no effect of treatment ($H = 6.7$, $P = 0.08$) or month ($H = 1.0$, $P = 0.31$). There was also
316 no significant difference in the change in O:N ratio post feeding between months ($H = 0.0$, P
317 $= 0.94$; Fig. 4) or between treatments ($H = 2.4$, $P = 0.49$).

318

319 Whilst the scope for growth (SfG) was significantly lower in September than November ($H =$
320 15.5 , $P < 0.01$; Fig. 4) there was no significant difference between treatments ($H = 5.3$, $P =$
321 0.15), although the general trend mirrored that of energy gain from food.

322

323 **Composition**

324 There was no significant difference in the organic mass (AFDM) of test (month, $F_{(1,69)} = 1.9$,
325 $P = 0.18$; treatment $F_{(3,69)} = 2.2$, $P = 0.10$), gonad (month, $F_{(1,69)} = 0.2$, $P = 0.70$; treatment
326 $F_{(3,69)} = 1.2$, $P = 0.32$), gonad somatic index (month, $F_{(1,69)} = 1.0$, $P = 0.33$; treatment $F_{(3,69)} =$
327 0.7 , $P = 0.57$) or gonad C:N ratio between months or between treatments (Table 3). There
328 was a small, but significant difference in gonad C:H ratio between treatments ($H = 14.4$, $P <$
329 0.01) but not between months ($H = 0.35$, $P = 0.55$). Gonads in the low temperature control
330 had the lowest C:H ratio compared to higher temperatures.

331

332 **Discussion**

333 **Growth and energetics**

334 This study describes the longest incubation to date of an Antarctic marine invertebrate to the
335 combined stressors of temperature and ocean acidification and significantly extends the
336 published time series (Suckling *et al.* 2015). After forty months exposure, there was little effect
337 of the treatment conditions on adult *Sterechinus neumayeri* somatic and reproductive tissue
338 mass, elemental composition or scope for growth. However there was a significant effect on
339 oxygen consumption and energetics, with lower metabolic rates and energy absorption in the
340 individuals subjected to elevated temperature. *S. neumayeri* held at +2°C had an elevated
341 metabolic rate, as expected, due to the rate increasing effect that temperature has on
342 biochemical reactions (Clarke 1983; Hochachka & Somero 2002). Indeed the data are very
343 similar to our previous study where metabolic rates of *S. neumayeri* was initially elevated in
344 response to incubation at +2°C with combined OA stressors, but any difference became non-
345 significant after 8 months of incubation (Suckling *et al.* 2015). Average metabolic rates after
346 40 months at 2 °C were between 2 and 3 $\mu\text{mol O}_2 \text{h}^{-1} \text{g AFDM}^{-1}$ which is slightly above summer
347 values measured in the wild (Brockington & Peck 2001), where temperatures are above zero,
348 but rarely reach 2°C (Venables, Clarke & Meredith 2013). What is surprising, however, is that
349 the animals at high temperature consumed less food and also absorbed less energy (Fig. 1, Table
350 2). With their elevated metabolic rates, compared to the animals kept at 0°C, they would be
351 expected to consume more food to fuel their elevated metabolism, which was clearly not the
352 case for *S. neumayeri*. The effects of temperature on feeding rate and energy absorption vary
353 between urchin species (Hill & Lawrence 2006; Zhao *et al.* 2015). For example,
354 *Strongylocentrotus intermedius* consumed less food at higher temperatures which led to a
355 reduction in gonad production (Zhao *et al.* 2015). Under increased warming, the metabolic
356 rates of many ectotherms are expected to increase at greater rates than consumption which

357 could in turn lead to a reduction in ingestion efficiency, ultimately resulting in energy deficits
358 (Lemoine & Burkepile 2012). How an increase in temperature will effect energy budgets will
359 depend on the thermal reaction norms of biochemical pathways and the proximity of the
360 elevated temperature to the upper boundary of their thermal window (Angilletta 2009). After
361 40 months in this study, there was, however, no significant difference in animal size,
362 reproductive allocation, or skeletal mass between the different treatments and all individuals
363 were still burning protein as their main food source (Fig. 4). *S.neumayeri*, of the size used here
364 (with test diameters above 20mm), grow very slowly, are difficult to age (Brey *et al.* 1995;
365 Brockington & Peck 2001) and therefore any difference in growth rate may be difficult to
366 detect. It has been estimated that in *S. neumayeri* only 5% of food is allocated to growth, with
367 the remaining 95% going towards reproduction (Brey *et al.* 1995), thus any reduction in
368 nutrition would be expected to affect reproduction first. However, more subtle effect of
369 temperature may lead to differences in energy allocation, some of which may have been missed
370 in the current study.

371

372

373 Two recent studies on echinoderms, albeit on larvae, demonstrated the potential effects of
374 altered pH on the digestive system; with smaller stomachs and reduced feeding performance in
375 the sand dollar *Dendraster excentricus* (Chan, Gruenbaum & O'Donnell 2011) and larger
376 stomachs and increased energetic requirements in the urchin *Stronglycentrotus droebachiensis*
377 (Dorey *et al.* 2013). The importance of feeding and food processing has also been demonstrated
378 in adult urchins. Individuals that were feeding were able to partially compensate extracellular
379 pH while individuals with empty digestive systems were suffering severe metabolic acidosis
380 (Stumpp *et al.* 2012). With reported effects of ocean acidification on energy allocation (Pan,

381 Applebaum & Manahan 2015) and feeding behaviour (Barry *et al.* 2014) an increasing number
382 of studies are reporting an interaction between OA stressors and nutritional status (Sandjensen
383 & Pedersen 1994; Melzner *et al.* 2011; Pan, Applebaum & Manahan 2015). Hence there is *a*
384 *priori* evidence that altered environmental conditions, especially low pH, can affect the
385 energetics of food processing. Which mechanism is most likely to underlie the physiological
386 effects of treatment, particularly the effect of temperature, is impossible to determine without
387 further study.

388

389 Bigger differences were found in this study between the two sample months, September and
390 November, than between treatments. In November, consumption of food and absorption of
391 energy were higher, leading to a higher SfG in all treatments. November is the start of the
392 austral summer, the time of peak spawning of *S. neumayeri* on the WAP (Pörtner, Bock &
393 Reipschlagler 2000) and when spawning in the laboratory was most successful (Suckling *et al.*
394 2015). The presence of seasonal cycle, in spite of *S. neumayeri* being kept in constant
395 temperature and photoperiod conditions for more than 40 months shows that these endogenous
396 rhythms are deeply entrained within this species.

397

398 **Implications for the benthic ecosystem**

399

400 This long term study has shown that the Antarctic sea urchin, *Sterechinus neumayeri*, is
401 relatively robust to the effects of near future ocean acidification. The results of the current study
402 show that temperature had a greater effect on the acclimated physiology of *S. neumayeri* than
403 low seawater pH, although there was an indication of an interactive effect, as is being found in

404 an increasing number of studies of marine ectotherms (Schram *et al.* 2014; Feidantsis *et al.*
405 2015). Recent studies have found that some echinoid taxa have a relatively high capacity to
406 buffer the pH of internal fluids against OA stressors (Sandjensen & Pedersen 1994; Stumpp *et*
407 *al.* 2012; Collard *et al.* 2015). This appears to be in part due to their ability to accumulate
408 bicarbonate in the coelomic fluid to reduce the impact of acidosis (Stumpp *et al.* 2012). *S.*
409 *droebachiensis* studied by Stumpp *et al.* (2012) live in a region that has high seasonal variation
410 in seawater pCO₂ and organism physiological plasticity and resilience are expected to correlate
411 with experienced environmental variation (Gaston *et al.* 2009). The Western Antarctic
412 Peninsula has a stable thermal environment (Venables, Clarke & Meredith 2013) but large
413 variations in pH have been recorded in shallow coastal waters, between pH 7.6 and 8.3 (Bjork
414 *et al.* 2014; Collard *et al.* 2015) which may be correlated with the capacity of *S. neumayeri* to
415 cope with changes in ocean acidification whilst being more sensitive to small changes in
416 temperature.

417

418 The focus of recent laboratory studies towards longer term ocean acidification incubations,
419 particularly for cold water species that have incubated adults for a full reproductive cycle and
420 across multiple generations is providing us with a clearer picture of the capacity of echinoderms
421 to cope with predicted future environmental conditions (Stumpp *et al.* 2012; Dupont *et al.*
422 2013; Suckling *et al.* 2015). As more detailed environmental manipulations are conducted, it
423 is becoming apparent that the subtlety of response is increasingly complex (Munday *et al.* 2009;
424 Kroeker, Micheli & Gambi 2013; Heuer & Grosell 2014). The differences in food consumption
425 and energetics of food processing found in *S. neumayeri*, in the current study, require further
426 studies that combine different ration sizes along with multiple environmental stressors, in order
427 to disentangle the mismatch between food consumption and the energetics of food processing.

428 However, studies to date show that *S. neumayeri* is robust to the impact of near future ocean
429 acidification and may actually benefit from a small rise in environmental temperature (Table
430 4). As *S. neumayeri* are an abundant, keystone, Southern Ocean species, at depths shallower
431 than 20 m, any change in food consumption or conversion efficiency of energy into body
432 tissues could cause a major shift in energy flow through the shallow water ecosystem.

433

434 **Acknowledgements**

435 This study was funded by the UK Natural Environment Research Council core funding to the
436 British Antarctic Survey's Adaptations and Physiology work package. Thanks to Paul
437 Geissler for completing the CHN analysis and Tim Brand at SAMS for analysing the urea
438 samples. Also thanks to Andy Clarke for advice on conducting the ammonia assay. We also
439 thank Joelle Richard, Laura Weir and Guy Hillyard for assisting in animal husbandry and
440 carbonate chemistry. The full dataset can be found at <http://doi.org/r34>.

441

442

443 **References**

- 444 Angilletta, M.J. (2009) *Thermal adaptation: a theoretical and empirical synthesis*. Oxford University
 445 Press.
- 446 Appelhans, Y.S., Thomsen, J., Opitz, S., Pansch, C., Melzner, F. & Wahl, M. (2014) Juvenile sea stars
 447 exposed to acidification decrease feeding and growth with no acclimation potential. *Marine*
 448 *Ecology Progress Series*, **509**, 227-239.
- 449 Barbarino, E. & Lourenco, S.O. (2009) Comparison of CHN analysis and Hach acid digestion to
 450 quantify total nitrogen in marine organisms. *Limnology and Oceanography-Methods*, **7**, 751-
 451 760.
- 452 Barry, J.P., Lovera, C., Buck, K.R., Peltzer, E.T., Taylor, J.R., Walz, P., Whaling, P.J. & Brewer, P.G.
 453 (2014) Use of a Free Ocean CO₂ Enrichment (FOCE) System to Evaluate the Effects of Ocean
 454 Acidification on the Foraging Behavior of a Deep-Sea Urchin. *Environmental Science &*
 455 *Technology*, **48**, 9890-9897.
- 456 Bjork, M.M., Fransson, A., Torstensson, A. & Chierici, M. (2014) Ocean acidification state in western
 457 Antarctic surface waters: controls and interannual variability. *Biogeosciences*, **11**, 57-73.
- 458 Bray, L., Pancucci-Papadopoulou, M.A. & Hall-Spencer, J.M. (2014) Sea urchin response to rising
 459 pCO₂ shows ocean acidification may fundamentally alter the chemistry of marine
 460 skeletons. *Mediterranean Marine Science*, **15**, 510-519.
- 461 Brey, T., Pearse, J., Basch, L., McClintock, J. & Slattery, M. (1995) Growth and production of
 462 *Sterechinus neumayeri* (Echinoidea: Echinodermata) in McMurdo sound, Antarctica. *Mar*
 463 *Biol*, **124**, 279-292.
- 464 Brockington, S. & Peck, L.S. (2001) Seasonality of respiration and ammonium excretion in the
 465 Antarctic echinoid *Sterechinus neumayeri*. *Marine Ecology Progress Series*, **219**, 159-168.
- 466 Chan, K.Y.K., Gruenbaum, D. & O'Donnell, M.J. (2011) Effects of ocean-acidification-induced
 467 morphological changes on larval swimming and feeding. *Journal of Experimental Biology*,
 468 **214**, 3857-3867.
- 469 Clarke, A. (1983) Life in cold water: the physiological ecology of polar marine ectotherms.
 470 *Oceanography and Marine Biology Annual Review*, **21**, 341-453.
- 471 Collard, M., De Ridder, C., David, B., Dehairs, F. & Dubois, P. (2015) Could the acid-base status of
 472 Antarctic sea urchins indicate a better-than-expected resilience to near-future ocean
 473 acidification? *Global Change Biology*, **21**, 605-617.
- 474 Dickson, A.G. & Millero, F.J. (1987) A comparison of the equilibrium-constants for the dissociation of
 475 carbonic-acid in seawater media. *Deep-Sea Research*, **34**, 1733-1743.
- 476 Dixon, D.L., Munday, P.L. & Jones, G.P. (2010) Ocean acidification disrupts the innate ability of fish
 477 to detect predator olfactory cues. *Ecol Lett*, **13**, 68-75.
- 478 Dorey, N., Lancon, P., Thorndyke, M. & Dupont, S. (2013) Assessing physiological tipping point of sea
 479 urchin larvae exposed to a broad range of pH. *Global Change Biology*, **19**, 3355-3367.
- 480 Dupont, S., Dorey, N., Stumpp, M., Melzner, F. & Thorndyke, M. (2013) Long-term and trans-life-
 481 cycle effects of exposure to ocean acidification in the green sea urchin *Strongylocentrotus*
 482 *droebachiensis*. *Mar Biol*, **160**, 1835-1843.
- 483 Elliott, J.M. & Davison, W. (1975) Energy equivalents of oxygen-consumption in animal energetics.
 484 *Oecologia*, **19**, 195-201.
- 485 Fabry, V.J., McClintock, J.B., Mathis, J.T. & Grebmeier, J.M. (2009) Ocean Acidification at High
 486 Latitudes: The Bellweather. *Oceanography*, **22**, 160-171.
- 487 Feidantsis, K., Poertner, H.-O., Antonopoulou, E. & Michaelidis, B. (2015) Synergistic effects of acute
 488 warming and low pH on cellular stress responses of the gilthead seabream *Sparus aurata*.
 489 *Journal of Comparative Physiology B-Biochemical Systemic and Environmental Physiology*,
 490 **185**, 185-205.

491 Fraser, K.P.P., Clarke, A. & Peck, L.S. (2002) Feast of famine in Antarctica: seasonal physiology in the
492 limpet *Nacella concinna*. *Marine Ecology Progress Series*, **242**, 169-177.

493 Gaston, K.J., Chown, S.L., Calosi, P., Bernardo, J., Bilton, D.T., Clarke, A., Clusella-Trullas, S.,
494 Ghalambor, C.K., Konarzewski, M., Peck, L.S., Porter, W.P., Portner, H.O., Rezende, E.L.,
495 Schulte, P.M., Spicer, J.I., Stillman, J.H., Terblanche, J.S. & van Kleunen, M. (2009)
496 Macrophysiology: A Conceptual Reunification. *American Naturalist*, **174**, 595-612.

497 Guinotte, J.M. & Fabry, V.J. (2008) Ocean acidification and its potential effects on marine
498 ecosystems. *Year in Ecology and Conservation Biology 2008*, pp. 320-342.

499 Hauri, C., Doney, S.C., Takahashi, T., Erickson, M., Jiang, G. & Ducklow, H.W. (2015) Two decades of
500 inorganic carbon dynamics along the Western Antarctic Peninsula. *Biogeosciences*
501 *Discussions*, **12**, 6929-6969.

502 Heuer, R.M. & Grosell, M. (2014) Physiological impacts of elevated carbon dioxide and ocean
503 acidification on fish. *American Journal of Physiology - Regulatory, Integrative and*
504 *Comparative Physiology*, **307**, 1061-1074.

505 Hill, S.K. & Lawrence, J.M. (2006) Interactive effects of temperature and nutritional condition on the
506 energy budgets of the sea urchins *Arbacia punctulata* and *Lytechinus variegatus*
507 (Echinodermata : Echinoidea). *Journal of the Marine Biological Association of the United*
508 *Kingdom*, **86**, 783-790.

509 Hochachka, P.W. & Somero, G.N. (2002) *Biochemical Adaptation: mechanisms and processes in*
510 *physiological evolution*. Oxford University Press, New York.

511 Hofmann, G.E., Barry, J.P., Edmunds, P.J., Gates, R.D., Hutchins, D.A., Klinger, T. & Sewell, M.A.
512 (2010) The Effect of Ocean Acidification on Calcifying Organisms in Marine Ecosystems: An
513 Organism-to-Ecosystem Perspective. *Annual Review of Ecology, Evolution, and Systematics*,
514 *Vol 41* (eds D.J. Futuyma, H.B. Shafer & D. Simberloff), pp. 127-147.

515 Holmes, R.M., Aminot, A., Kerouel, R., Hooker, B.A. & Peterson, B.J. (1999) A simple and precise
516 method for measuring ammonium in marine and freshwater ecosystems. *Canadian Journal*
517 *of Fisheries and Aquatic Sciences*, **56**, 1801-1808.

518 Kroeker, K.J., Micheli, F. & Gambi, M.C. (2013) Ocean acidification causes ecosystem shifts via
519 altered competitive interactions. *Nature Climate Change*, **3**, 156-159.

520 Lemoine, N.P. & Burkepile, D.E. (2012) Temperature-induced mismatches between consumption and
521 metabolism reduce consumer fitness. *Ecology*, **93**, 2483-2489.

522 Lewis, E. & Wallace, D.W.R. (1988) Carbon dioxide information analysis center, Oak ridge National
523 Laboratory. US Department of energy, Tennessee.

524 Mayzaud, P. & Conover, R.J. (1988) O:N atomic ratio as a tool to describe zooplankton metabolism.
525 *Marine Ecology Progress Series*, **45**, 289-302.

526 McNeil, B.I. & Matear, R.J. (2008) Southern Ocean acidification: A tipping point at 450-ppm
527 atmospheric CO₂. *Proc Natl Acad Sci U S A*, **105**, 18860-18864.

528 Mehrbach, C., Culbertson, C.H., Hawley, J.E. & Pytkowicz, R.M. (1973) Measurement of apparent
529 dissociation constants of carbonic acid in seawater at atmospheric pressure. *Limnology and*
530 *Oceanography*, **18**, 897-907.

531 Melzner, F., Gutowska, M.A., Langenbuch, M., Dupont, S., Lucassen, M., Thorndyke, M.C., Bleich, M.
532 & Portner, H.O. (2009) Physiological basis for high CO₂ tolerance in marine ectothermic
533 animals: pre-adaptation through lifestyle and ontogeny? *Biogeosciences*, **6**, 2313-2331.

534 Melzner, F., Stange, P., Trübenbach, K., Thomsen, J., Casties, I., Panknin, U., Gorb, S.N. & Gutowska,
535 M.A. (2011) Food Supply and Seawater pCO₂ Impact Calcification and Internal Shell
536 Dissolution in the Blue Mussel *Mytilus edulis*. *PLoS One*, **6**.

537 Morley, S.A., Lemmon, V., Obermueller, B.E., Spicer, J.I., Clark, M.S. & Peck, L.S. (2011) Duration
538 tenacity: A method for assessing acclimatory capacity of the Antarctic limpet, *Nacella*
539 *concinna*. *Journal of Experimental Marine Biology and Ecology*, **399**, 39-42.

540 Morley, S.A., Peck, L.S., Miller, A.J. & Poertner, H.O. (2007) Hypoxia tolerance associated with
541 activity reduction is a key adaptation for *Laternula elliptica* seasonal energetics. *Oecologia*,
542 **153**, 29-36.

543 Munday, P.L., Crawley, N.E. & Nilsson, G.E. (2009) Interacting effects of elevated temperature and
544 ocean acidification on the aerobic performance of coral reef fishes. *Marine Ecology Progress
545 Series*, **388**, 235-242.

546 Munday, P.L., Dixon, D.L., Donelson, J.M., Jones, G.P., Pratchett, M.S., Devitsina, G.V. & Doving, K.B.
547 (2009) Ocean acidification impairs olfactory discrimination and homing ability of a marine
548 fish. *Proc Natl Acad Sci U S A*, **106**, 1848-1852.

549 Nickell, L.A., Black, K.D., Hughes, D.J., Overnell, J., Brand, T., Nickell, T.D., Breuer, E. & Harvey, S.M.
550 (2003) Bioturbation, sediment fluxes and benthic community structure around a salmon
551 cage farm in Loch Creran, Scotland. *Journal of Experimental Marine Biology and Ecology*,
552 **285**, 221-233.

553 Obermüller, B.E., Morley, S.A., Barnes, D.K.A. & Peck, L.S. (2010) Seasonal physiology and ecology of
554 Antarctic marine benthic predators and scavengers. *Marine Ecology Progress Series*, **415**,
555 109-126.

556 Pan, T.C.F., Applebaum, S.L. & Manahan, D.T. (2015) Experimental ocean acidification alters the
557 allocation of metabolic energy. *Proc Natl Acad Sci U S A*, **112**, 4696-4701.

558 Peck, L.S. (1998) Feeding, metabolism and metabolic scope in Antarctic marine ectotherms. *Cold
559 Ocean Physiology* (eds H.O. Pörtner & R.C. Playle), pp. 365-390. Cambridge University Press,
560 Cambridge.

561 Peck, L.S. (2011) Organisms and responses to environmental change. *Mar Genomics*, **4**, 237-243.

562 Peck, L.S., Convey, P. & Barnes, D.K.A. (2006) Environmental constraints on life histories in Antarctic
563 ecosystems: tempos, timings and predictability. *Biological Reviews*, **81**, 75-109.

564 Peck, L.S., Morley, S.A., Richard, J. & Clark, M.S. (2014) Acclimation and thermal tolerance in
565 Antarctic marine ectotherms. *Journal of Experimental Biology*, **217**, 16-22.

566 Peck, L.S. & Veal, R. (2001) Feeding, metabolism and growth in the Antarctic limpet, *Nacella
567 concinna* (Strebel 1908). *Mar Biol*, **138**, 553-560.

568 Peck, L.S., Webb, K.E., Miller, A., Clark, M.S. & Hill, T. (2008) Temperature limits to activity, feeding
569 and metabolism in the Antarctic starfish *Odontaster validus*. *Marine Ecology-Progress Series*,
570 **358**, 181-189.

571 Pörtner, H.O. (2012) Integrating climate-related stressor effects on marine organisms: unifying
572 principles linking molecule to ecosystem-level changes. *Marine Ecology Progress Series*, **470**,
573 273-290.

574 Pörtner, H.O., Bock, C. & Reipschlag, A. (2000) Modulation of the cost of pHi regulation during
575 metabolic depression: A P-31-NMR study in invertebrate (*Sipunculus nudus*) isolated muscle.
576 *Journal of Experimental Biology*, **203**, 2417-2428.

577 Pörtner, H.O., Peck, L.S. & Somero, G.N. (2007) Thermal limits and adaptation in marine Antarctic
578 ectotherms: an integrative view. *Philosophical Transactions of the Royal Society B-Biological
579 Sciences*, **362**, 2233-2258.

580 Robertson, R.F., El-Haj, A.J., Clarke, A., Peck, L.S. & Taylor, E.W. (2001) The effects of temperature on
581 metabolic rate and protein synthesis following a meal in the isopod *Glyptonotus antarcticus*
582 Eights (1852). *Polar Biology*, **24**, 677-686.

583 Sandjensen, K. & Pedersen, M.F. (1994) PHOTOSYNTHESIS BY SYMBIOTIC ALGAE IN THE FRESH-
584 WATER SPONGE, SPONGILLA-LACUSTRIS. *Limnology and Oceanography*, **39**, 551-561.

585 Schram, J.B., Schoenrock, K.M., McClintock, J.B., Amsler, C.D. & Angus, R.A. (2014) Multiple stressor
586 effects of near-future elevated seawater temperature and decreased pH on righting and
587 escape behaviors of two common Antarctic gastropods. *Journal of Experimental Marine
588 Biology and Ecology*, **457**, 90-96.

589 Sewell, M.A. & Hofmann, G.E. (2011) Antarctic echinoids and climate change: a major impact on the
590 brooding forms. *Global Change Biology*, **17**, 734-744.

591 Somero, G.N. (2010) The physiology of climate change: how potentials for acclimatization and
592 genetic adaptation will determine 'winners' and 'losers'. *J Exp Biol*, **213**, 912-920.

593 Somero, G.N. (2012) The Physiology of Global Change: Linking Patterns to Mechanisms. *Annual*
594 *Review of Marine Science, Vol 4*, **4**, 39-61.

595 Spicer, J.I., Widdicombe, S., Needham, H.R. & Berge, J.A. (2011) Impact of CO₂-acidified seawater on
596 the extracellular acid-base balance of the northern sea urchin *Strongylocentrotus*
597 *droebachiensis*. *Journal of Experimental Marine Biology and Ecology*, **407**, 19-25.

598 Stumpp, M., Trubenbach, K., Brennecke, D., Hu, M.Y. & Melzner, F. (2012) Resource allocation and
599 extracellular acid-base status in the sea urchin *Strongylocentrotus droebachiensis* in
600 response to CO₂ induced seawater acidification. *Aquat Toxicol*, **110-111**, 194-207.

601 Suckling, C.C., Clark, M.S., Richard, J., Morley, S.A., Thorne, M.A.S., Harper, E.M. & Peck, L.S. (2015)
602 Adult acclimation to the combined temperature and pH stressors significantly enhances
603 reproductive outcomes compared to short-term exposures. *Journal of animal ecology*, **84**,
604 773-784.

605 Sunday, J.M., Bates, A.E. & Dulvy, N.K. (2011) Global analysis of thermal tolerance and latitude in
606 ectotherms. *Proc Biol Sci*, **278**, 1823-1830.

607 Venables, H.J., Clarke, A. & Meredith, M.P. (2013) Wintertime controls on summer stratification and
608 productivity at the western Antarctic Peninsula. *Limnology and Oceanography*, **58**, 1035-
609 1047.

610 Watson, S.-A., Peck, L.S., Tyler, P.A., Southgate, P.C., Tan, K.S., Day, R.W. & Morley, S.A. (2012)
611 Marine invertebrate skeleton size varies with latitude, temperature and carbonate
612 saturation: implications for global change and ocean acidification. *Global Change Biology*,
613 **18**, 3026-3038.

614 Widdicombe, S. & Needham, H.R. (2007) Impact of CO₂-induced seawater acidification on the
615 burrowing activity of *Nereis virens* and sediment nutrient flux. *Marine Ecology Progress*
616 *Series*, **341**, 111-122.

617 Winberg, C.G. (1960) Rate of metabolism and food requirements of fishes. *Translated Series of the*
618 *Fisheries Research Board of Canada*, **194**.

619 Wittmann, A.C. & Pörtner, H.O. (2013) Sensitivities of extant animal taxa to ocean acidification.
620 *Nature Climate Change*, **3**, 995-1001.

621 Wood, C.M. (1993) Ammonia and urea metabolism and excretion. *The Physiology of Fishes* (ed. D.H.
622 Evans), pp. 177-230. CRC Press, Boca Raton, FL.

623 Wood, H.L., Spicer, J.I. & Widdicombe, S. (2008) Ocean acidification may increase calcification rates,
624 but at a cost. *Proceedings of the Royal Society B-Biological Sciences*, **275**, 1767-1773.

625 Zhao, C., Feng, W., Wei, J., Zhang, L., Sun, P. & Chang, Y. (2015) Effects of temperature and feeding
626 regime on food consumption, growth, gonad production and quality of the sea urchin
627 *Strongylocentrotus intermedius*. *Journal of the Marine Biological Association of the United*
628 *Kingdom*, **11**.

629

630

631

632 **Figure Legends**

633 Figure 1. The energy absorbed from food. All values are in Joules per hour per g ash free dry
634 mass. Treatments are: *Low temperature control* = -0.3 °C, pH 8.0; *High temperature control*
635 = 2 °C, pH 8.0; *pH -0.3* = 2 °C, pH 7.8; *pH -0.5* = 2 °C, pH 7.5. Filled bars = September,
636 open bars = November. ** indicates a significant difference in the energy absorbed between
637 months ($F_{(1,64)} = 35.0, P < 0.01$). Different letters indicate that low temperature controls
638 absorbed significantly less energy than other treatments (ANOVA $F_{(3,64)} = 5.8, P < 0.01$;
639 Tukey tests, *pH -0.3*, $T = 3.5$ and *pH -0.5*, $T = 3.7, P < 0.01$). Mean (\pm SE).

640

641 Figure 2. Oxygen consumption of *S. neumayeri*, pre and 6 days post feeding, in September
642 and November. Treatments are: *Low temperature control* = -0.3°C, pH 8.0; *High temperature*
643 *control* = 2 °C, pH 8.0; *pH -0.3* = 2 °C, pH 7.8; *pH -0.5* = 2 °C, pH 7.5. A, indicates that low
644 temperature controls consumed less oxygen than high temperature control and *pH -0.3*
645 treatments (ANOVA, $F_{(3,139)} = 3.8, P = 0.01$; High temperature control, $T = 2.6, P < 0.05$; *pH*
646 *-0.3* treatment, $T = 3.0, P < 0.05$). * indicates that there was a significant increase in oxygen
647 consumption post feeding (ANOVA, $F_{(1,139)} = 6.3, P < 0.05$). Mean (\pm SE).

648

649 Figure 3. Ammonia production of *S. neumayeri* in September and November before and 6
650 days after feeding. Treatments are: *Low temperature control* = -0.3 °C, pH 8.0; *High*
651 *temperature control* = 2 °C, pH 8.0; *pH -0.3* = 2 °C, pH 7.8; *pH -0.5* = 2 °C, pH 7.5. **
652 indicates a significant difference between pre and post feeding (Kruskal Wallis test: $H = 12.7,$
653 $P < 0.01$).


654 Fig. 4. Atomic O:N ratio for *S. neumayeri* in September (top panel) and November (bottom
655 panel) before and 6 days after feeding. Treatments are: Tcur = -0.3 °C, pH 7.8; pHcur = 2 °C,
656 pH 8.1; pH-0.3 = 2 °C, pH 7.8; pH-0.5 = 2 °C, pH 7.5. * indicates a significant difference
657 between pre and post feeding.

658

659 Figure 5. The scope for growth in September (filled bars) and November (open bars). **
660 indicates a significant difference between months ($H = 5.3, P < 0.01$). *Low temperature*
661 *control* = -0.3 °C, pH 8.0; *High temperature control* = 2 °C, pH 8.0; *pH -0.3* = 2 °C, pH 7.8;
662 *pH -0.5* = 2 °C, pH 7.5. Mean (\pm SE).


663

692





693

694 Fig. 1





695
696 Fig. 2
697



698

699 Fig. 3.

700




701

702

703

704



705

706 Fig. 5

	Low	High		
Seawater parameter	temperature	temperature	-0.3 pH	-0.5 pH
	control	Control		
Alkalinity	1733 ± 25	1851 ± 37	1753 ± 40	1805 ± 34
$p\text{CO}_2$ (µatm)	417 ± 15	420 ± 13	834 ± 39	1361 ± 36
pH _{NIST}	7.98 ± 0.02	8.00 ± 0.01	7.72 ± 0.01	7.52 ± 0.01
Ω calcite	1.20 ± 0.10	1.50 ± 0.03	0.76 ± 0.02	0.51 ± 0.02
Ω aragonite	0.75 ± 0.06	0.9 ± 0.02	0.48 ± 0.01	0.32 ± 0.01
Temperature (°C)	-0.3 ± 0.0	1.7 ± 0.1	1.9 ± 0.1	2.2 ± 0.1
Salinity (psu)	35 ± 0.2	35 ± 0.2	35 ± 0.2	35 ± 0.1

Table 1: Mean (\pm SE) water parameters in the adult *Sterechinus neumayeri* microcosm over the course of the experiment following the format of Barry *et al.*, (2010). Values for $p\text{CO}_2$, Ω calcite, Ω aragonite and total alkalinity were modelled from CO2SYS (Lewis & Wallace 1988) with refitted constants (Mehrbach *et al.* 1973; Dickson & Millero 1987).

Parameter		<i>Low temperature control</i>	<i>High temperature control</i>	<i>pH -0.3</i>	<i>pH -0.5</i>
A_{eff}	Sept	0.77 ± 0.06a	0.71 ± 0.05a	0.66 ± 0.06a	0.55 ± 0.08a
	Nov	0.87 ± 0.02b	0.91 ± 0.02b	0.87 ± 0.02b	0.91 ± 0.03b
C	Sept	2.4 ± 0.5a	1.3 ± 0.2ab	1.0 ± 0.3b	1.0 ± 0.2b
	Nov	4.8 ± 1.1c	3.1 ± 0.5cd	2.7 ± 0.3d	2.2 ± 0.5d

Table 2: Absorption efficiency (A_{eff}) and energy consumed (C, $\text{J}\cdot\text{h}^{-1}\text{gAFDM}^{-1}$), in September and November. Treatments are: *Low temperature control* = $-0.3\text{ }^{\circ}\text{C}$, pH 8.0; *High temperature control* = $1.7\text{ }^{\circ}\text{C}$, pH 8.0; *pH -0.3* = $1.9\text{ }^{\circ}\text{C}$, pH 7.8; *pH -0.5* = $2.2\text{ }^{\circ}\text{C}$, pH 7.5. Mean ± SE. Different lower case letters indicate that absorption efficiency was lower in September than November ($F_{(1,64)} = 40.8$, $P > 0.01$). Different lower case letters indicate that more energy was consumed in the low temperature control than other treatments (ANOVA, $F_{(3,64)} = 6.6$, $P < 0.01$; Tukey tests, *pH -0.3*, $T = 3.7$ and *pH -0.5*, $T = 4.1$, $P < 0.01$) and was less in September than November ($F_{(1,64)} = 35.7$, $P < 0.01$).

Parameter		<i>Low</i>	<i>High</i>	<i>pH -0.3</i>	<i>pH -0.5</i>
		<i>temperature</i>	<i>temperature</i>		
		<i>control</i>	<i>control</i>		
Test	Sept	734 ± 56	611 ± 72	628 ± 89	549 ± 64
AFDM,					
mg					
	Nov	791 ± 105	712 ± 80	703 ± 50	600 ± 68
Gonad	Sept	629,	634,	506,	685,
AFDM,		510-859	358-675	337-746	288-828
mg					
	Nov	728,	511,	554,	589,
		407-935	362-871	349-675	345-777
GSI	Sept	48.1 ± 1.6	46.2 ± 2.5	46.2 ± 2.3	48.9 ± 3.7
	Nov	47.2 ± 4.3	42.4 ± 3.7	43.8 ± 3.2	47.1 ± 3.7
C:N	Sept	5.4,	5.7,	5.6,	6.0,
		5.1-5.9	5.3-6.0	5.2-6.0	5.3-6.3
	Nov	5.9,	6.1,	5.8,	6.4,
		5.5-6.4	5.4-6.2	4.7-5.8	5.3-8.0
C:H	Sept	0.52,	0.53,	0.53,	0.54,
		0.51-0.53a	0.53-0.54ab	0.52-0.54bc	0.53-0.54c

Nov	0.53,	0.53,	0.54,	0.54,
	0.52-0.53a	0.52-0.53ab	0.54-0.55bc	0.53-0.55c

Table 3. The ash free dry mass (AFDM) of the test and gonad, the gonad somatic index (GSI), the carbon to nitrogen (C:N) and carbon to hydrogen (C:H) ratio in the gonad in September and November. Values are means \pm SE or median, interquartile range (the latter is used where data were not normally distributed, even after transformation. Different letters after the interquartile range indicate significantly different C:H ratios.

Duration of incubation	Trait	Control low temperature	Control high temperature	pH -0.3	pH -0.5	Ref
6 months	Egg size	+	-	-(-)	-	Suckling et al. 2015
	Fertilization success	=	+	=	=	Suckling et al. 2015
	Hatching success	+			-	Suckling et al. 2015
	Larval survival	+	+		-	Suckling et al. 2015
17 months	Egg size	-	+(-)	+	+(+)	Suckling et al. 2015
	Fertilization success		+		-	Suckling et al. 2015
	Hatching success	=	=	=	=	Suckling et al. 2015
	Larval survival	-	+		-	Suckling et al. 2015
8 to 24 months	Metabolic rate		=	=	=	Suckling et al. 2015
8 to 40 months	Test growth	=	=	=	=	Suckling et al. 2015 Current study
8 to 40 months	Gonad allocation	=	=	=	=	Suckling et al. 2015 Current Study
After 40 months	Metabolic rate	-	+	+	+	Current Study
After 40 months	Food consumption	+	-	-	-	Current Study
After 40 months	Ammonia production	=	=	=	=	Current Study
After 40 months	Scope for growth	=	=	=	=	Current Study

Table 4. Summary of effect of combined temperature and pH treatments on *S. neumayeri*