

Phosphorus sorption, supply potential and availability in soils with contrasting parent material and soil chemical properties Daly, K.; Styles, D.; Lalor, S.; Wall, D.P.

European Journal of Soil Science

DOI: 10.1111/ejss.12260

Published: 01/07/2016

Peer reviewed version

Cyswllt i'r cyhoeddiad / Link to publication

Dyfyniad o'r fersiwn a gyhoeddwyd / Citation for published version (APA): Daly, K., Styles, D., Lalor, S., & Wall, D. P. (2016). Phosphorus sorption, supply potential and availability in soils with contrasting parent material and soil chemical properties. *European Journal of Soil Science*, *66*(4), 792-801. https://doi.org/10.1111/ejss.12260

Hawliau Cyffredinol / General rights Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

• Users may download and print one copy of any publication from the public portal for the purpose of private study or research.

- You may not further distribute the material or use it for any profit-making activity or commercial gain
 You may freely distribute the URL identifying the publication in the public portal ?

Take down policy If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

1	Phosphorus sorption, supply potential and availability in soils with
2	contrasting parent material and soil chemical properties.
3	
4	K. DALY ^a , D. STYLES ^b , S. LALOR ^c &.D.P. WALL ^a
5	
6	^a Teagasc, Johnstown Castle, Environmental Research Centre, Co. Wexford, Ireland, ^b School
7	of Environment, Natural Resources and Geography. Bangor University, Bangor, Gwynedd
8	LL57 2DG, U.K., & ^c Grassland Agro, Dock Road, Limerick, Ireland.
9	
10	
11	Correspondence: K.Daly. Email: karen.daly@teagasc.ie
12	
13	
14	Running title: Phosphorus sorption and buffering
15	
16	
17	Keywords: Phosphorus, desorption, phosphorus solubility, soil fertility, buffering capacity,
18	temperate soils.
19	
20	

21 Summary

22

23 Soil phosphorus (P) management requires a more targeted and soil specific approach than is 24 currently applied for agronomic recommendations and environmental evaluation. Phosphorus buffering capacities control the supply of P in the soil solution and were 25 26 measured across Irish soils with contrasting parent material and chemical properties. Langmuir sorption buffer capacities (MBC) and binding energies (b) were strongly correlated 27 28 with soil pH and extractable aluminium (Al). A broken-line regression fitted to the 29 relationship between MBC and Al derived a change-point value for Al above which MBC 30 increased linearly. Soils above the change point were predominantly acidic to neutral with 31 non-calcareous parent material, larger buffering capacities and binding energies than 32 calcareous soils. Ratios of Mehlich3-Al and P (Al:P) were used to relate buffering capacity 33 with supply potential in non-calcareous soils. Large ratios of Al:P were associated with poor P availability, characteristic of strongly P fixing soils. Threshold values of iron-oxide paper 34 35 strip P (FeO-P) and Morgan's P revealed Al:P ratios where soils began to supply P in 36 available form. The change-point for Morgan's P fell within the current target index for P availability; however, the confidence interval was more compatible with previous agronomic 37 38 P indices used in Ireland. Relationships between Morgan's P and measures of extractable P, M3-P and Olsen P, deviated in calcareous soils at large soil P contents, indicative of P 39 40 precipitation processes dominating in these soils. Identifying differences in soil P buffering 41 capacity at laboratory scale would improve agronomic and environmental assessment at field 42 and catchment scales.

- 43
- 44
- 45

46 Introduction

47

48 Phosphorus (P) is an essential element to all known life and an adequate supply is required in 49 agricultural soils to meet the demands for crop growth and grazing animal health. The finite 50 nature of rock phosphate and Europe's dependence on imports from China and Asia and 51 North Africa, is putting its supply under pressure as reserves are diminishing (Cordell et al. 2009). The majority (79%) of the P imported is used to make fertilizer for use in agriculture 52 53 and food production and in animal feeds (11%) (Johnston & Steen, 2000). In many countries, 54 fertilizer P recommendations are based on P index systems where bands of soil test P values 55 are used to guide application rates. In Ireland the agronomic soil test for P is Morgan's 56 extractable P, and for easier management and knowledge transfer at the farm scale Morgan's 57 P levels have been categorized as indices; 1 (deficient), 2 (low), and 3 (agronomic optimum) 58 and 4 (excessive) (Lalor & Coulter, 2008). Fertilizer recommendations have been inextricably linked with soil tests since the 1950s and 1960s, following the general 59 60 philosophy of 'build up and maintenance' (Vos, 1998). In recent years the agri-environmental 61 landscape has changed, especially, since the implementation of water quality directives.

Morgan's P values are now used as the basis for risk assessment of P loss from Irish soils.
Going forward, farmers face huge challenges in managing P resources on farms including
increased P requirements for sustainable intensification to meet rising global food demands,
the maintenance of soil fertility under stringent environmental constraints and increasingly
expensive P fertiliser. The growing need for economic and environmental sustainability has
stimulated interest in efficient P fertilizer utilization and nutrient management on farms.

Future soil fertility management requires a more targeted and soil specific approach than is currently applied, to maximize the crop use efficiency of valuable P inputs, and to increase the P supplying capacities of soils under reduced P input regimes. Phosphorus buffering capacity is described as the ability to supply soluble P as a function of capacity, and
differences in P buffering and sorption capacities can significantly influence supply and
availability of P in a water soluble form (Burkitt *et al.* 2001).

74 Whilst it has been accepted that no soil test is universal (Vos, 1998), recent research incorporating P sorption and buffering with soil P tests, has provided P indices that are soil-75 76 type specific for improved critical soil test P predictions to give a value of soil test P where availability is optimized (Moody et al. 1997; Van Rotterdam et al. 2010). Similarly, 77 78 quality/intensity relationships have provided useful indices for predicting supply of slowly 79 available P in soils using easily measured soil properties with traditional soil tests (McLaren 80 et al. 2014; Van Rotterdam et al. 2012). More recently, Reijneveld et al. (2014) proposed a 81 strategy for modelling novel methods of P capacity and intensity, with historical P tests, 82 combined with field trials, for agronomic recommendations and environmental evaluations.

83 Over the past decade Irish fertiliser P recommendations have been updated for 84 grassland soils in Ireland, using results from a fertilizer response trial that did not 85 differentiate soil type, (Schulte & Herlihy, 2007). The objectives of this present study were to measure P buffering capacities across a range of Irish grassland soils with contrasting parent 86 material and chemical properties and relate these values to soil attributes, pertaining to P 87 88 availability and supply. Soil samples with contrasting parent material and chemistry were collected from grassland sites across Ireland and the relevance of contrasting soil P buffering 89 90 and sorption is discussed with regard to availability and examined in the context of our 91 current revised agronomic recommendations.

- 92
- 93
- 94
- 95

96 Materials and methods

97

98 Soil Sampling

99 Principal agricultural soils were identified from the General Soil Map of Ireland (Gardiner & 100 Radford, 1980), the mapping unit of which is the soil association. These include 101 amalgamations of Great Soil Groups and associated parent material, and for this study, 120 sampling locations were identified that best represented principal soil associations used in 102 103 grassland systems in Ireland (Table 1). The soil sampling programme also targeted sites with 104 a range of soil test P levels (Morgan's P) so that the agronomic P indices were reflected in the 105 data. Samples were taken on field-by-field basis to a depth of 100 mm with a soil corer (1.5-106 cm diameter) with approximately 30-40 cores bulked to provide a composite sample from 107 each field and prior to the spreading of fertilizers and manures. One hundred and twenty 108 samples were air-dried and sieved to remove roots and stones (2 mm mesh) and stored in air-109 tight boxes at room temperature prior to analysis.

110

111 Phosphorus sorption isotherms

Phosphorus sorption isotherms were derived using a modification of the standardized batch 112 113 technique by Nair et al. (1984). Because of the time consuming nature of this technique, a sub-set of 74 soil samples, representing each soil association, were equilibrated with P in 114 115 solution and the data was evaluated using the Langmuir model (Paulter & Sims, 2000) as follows. Six, 40-ml solutions of P concentration 0, 5, 10, 15, 20 and 25 mg l⁻¹ in KH₂PO₄ 116 were added to 2 g soil samples in 50ml centrifuge tubes in duplicate. The suspensions were 117 118 shaken at room temperature for 24 hour, centrifuged and filtered, and the concentration of P 119 in solution was measured colorimetrically (John, 1970) using the molybdate blue method. Phosphorus sorbed to the soil was calculated as the difference between initial concentration 120

121 and P concentration measured at equilibrium. Adsorption data and affinity constants were122 determined by fitting sorption data to the Langmuir isotherm equation:

123
$$C/X = 1/Xm x b + C/Xm,$$
 (1)

which represents a straight line where X is P sorbed (mg P kg⁻¹), C is the equilibrium P concentration (mg P l⁻¹) with an intercept 1/Xm x b and slope 1/Xm. The linear form of the Langmuir equation was then used to derive Xm, the sorption maximum (mg P kg⁻¹) and b, a constant related to the binding energy (l mg⁻¹). Phosphorus sorption isotherms were plotted for each soil sample collected and the Langmuir model derived sorption maximum (Xm, mg kg⁻¹), binding energy (b, l mg⁻¹). Maximum buffer capacity (MBC, l kg⁻¹) was calculated as the product of Xm and b (Indiati *et al.* 1999).

131

132 Extractable P (FeO-P, Olsen P, Mehlich3-P, Morgan's P)

133 The iron-oxide paper strip test was used as a P sink method for estimating labile inorganic P 134 on 120 soils sampled, according to the method of Menon et al. (1988). One FeO-impregnated paper strip (100 x 20 mm) was added to each sample of soil (1 g) suspended in 40 ml of 0.01 135 M CaCl₂ solution and shaken for 16 hour on an end-over-end shaker. The paper strips were 136 eluted with 40 ml 0.1 M H₂SO₄ for one hour and the P concentrations in the acidic solution 137 138 determined colorimetrically. Morgan's P was determined on 120 soils, with 6.5 ml of soil in a 139 buffered acetate-acetic acid reagent at a 1:5 (v/v) soil to solution ratio extracting for 30 minutes (Peech & English, 1944). Mehlich3-P (M3-P) was determined on 120 soils using the 140 141 modified Mehlich test (Mehlich, 1984) to extract P, aluminium (Al), iron (Fe) and calcium (Ca), at a 1:10 soil solution ratio of 2 g of soil shaken with 20 ml Mehlich3 reagent (0.2 M 142 CH₃COOH + 0.25 M NH₄NO₃ + 0.015 M NH₄F + 0.13M HNO₃ + 0.001M EDTA) for 5 143 144 minutes on a reciprocating shaker. The extracts were filtered and P in solution measured colorimetrically with a molybdate blue method (John, 1970). Olsen P was measured on 60 145

samples only because of a lack of sample availability. This method used 2-g sample shaken
with 40 ml 0.5 M NaHCO₃, buffered at pH 8.5, for 30 minutes. Phosphorus concentration in
the solution was determined colorimetrically with a molybdate blue method (John, 1970).

149 Soil pH was determined on a 2:1 soil-water ratio paste on 120 samples and percentage organic matter (% OM) measured from 120 samples as the loss-on-ignition of 5-g samples at 150 500° C. Results were analysed with SAS Enterprise Guide V5 to obtain correlation and 151 regression information. The NLMIXED procedure in SAS was used for change-point 152 153 analysis and broken-line regression. Optimization was performed using the dual quasi-154 Newton algorithm, with a dual Broyden, Fletcher, Goldfarb, and Shanno (BFGS) update of 155 the Cholesky factor of an approximate Hessian. PROC NLMIXED computes approximate 156 standard errors for estimates and predictions using the delta method of truncated expansions.

157

- 158 **Results**
- 159

160 Chemical properties of soils collected

161 Soil chemical properties of representative soil associations are presented in Table 2 and their correlations are tabulated in Table 3. Soil pH ranged from 4.7-7.1, with smallest values 162 163 measured in soil associations 14, 15 and 25 and largest values recorded in soil associations 30 and 34 on limestone parent material. Extractable Ca, ranged from 284-8660 mg kg⁻¹ across 164 all soils and was positively correlated with soil pH (r = 0.80). Extractable Al and Fe ranged 165 from 69-1730 mg kg⁻¹ and 215-666 mg kg⁻¹, respectively with largest concentrations 166 measured on soil associations 14 and 25. Extractable Al was negatively correlated with soil 167 pH (r = -0.71) and largest concentrations associated with acidic soils. Across all soil 168 associations, OM% ranged from 5.3–15.7 % and some weak but significant correlations were 169 170 observed between OM% and extractable Fe (r = 0.47) and Ca (r = 0.32) possibly indicating

171 the presence of amorphous forms of Fe and Ca among these soils. Morgan's P ranged from 172 1.5–32.9 mg 1⁻¹ and correlated with other P extraction methods, where the strongest 173 correlations were observed with M3-P (r = 0.79) and FeO-P (r = 0.73) and weakest with 174 Olsen-P (r = 0.63). Across all soils, Mehlich3 extracted of the most P (13.4–285 mg kg⁻¹), 175 followed by FeO-P (15.4–128 mg kg⁻¹) and Olsen-P (8.9–89.6 mg kg⁻¹).

176

177 Langmuir sorption properties

The Langmuir model provided a highly significant (P < 0.001) fit to soils in this study ($R^2 >$ 178 0.95) and values of Langmuir parameters Xm, b and MBC derived from the isotherms across 179 180 all soil associations are presented in Table 2. Soil pH was inversely correlated with all of the 181 Langmuir sorption parameters Xm, (r = -0.46), b (r = -0.62) and MBC (r = -0.65) whilst extractable Al positively correlated with Xm, b and MBC. These soil attributes provided the 182 strongest correlations with Langmuir sorption terms, although the scatter plots were 183 nonlinear. Soil pH accounted for 55% of the variation in log transformed MBC (Figure 1) 184 185 with buffering capacity decreasing as soil pH increased from 4.7 to 7.1. The regression 186 equation is listed in Table 4. Soil associations derived from limestone parent material recorded the smallest values of each of the Langmuir terms with typically high pH values of 187 >6.0. Visual inspection of the relationship between MBC and Al indicated small MBC 188 189 values at small Al values with a sharp increase in MBC above a threshold Al value. This 190 threshold value was derived by regression modelling of the data where a broken-line 191 regression was fitted to log transformed MBC and extractable Al in Figure 2, and a significant change-point (P < 0.001) derived at an Al concentration of 612 (+/- 104) mg kg⁻¹, 192 above which buffer capacity in soil increased linearly with increasing Al. Parameter estimates 193 194 for this relationship are presented in Table 5. Above the upper confidence limit of the change-point for Al (>716 mg kg⁻¹), soils was predominantly derived from non-calcareous 195

196 parent material. This group of soils were characterised by predominantly low pH values 197 ranging from 4.7 to 6.3 (mean = 5.3) and strong Langmuir sorption properties. Mean MBC and b values in these soils were 1250 l kg⁻¹ and 2.56 l mg⁻¹, respectively, and mean Al and Ca 198 values of 1072 and 1451 mg kg⁻¹, respectively. These soils are denoted here as non-199 200 calcareous soils (n = 76) and their summary statistics are presented in Table 6. When separated out from the overall data, Al in these soils accounted for 65% and 57% of the 201 202 variation in MBC and b, respectively. The remaining soils, were mostly soil associations 203 derived from limestone parent material, with soil pH in the range 5.2–7.1, mean extractable Ca values of 3487 mg kg⁻¹, and mean MBC and b values of 346 l kg⁻¹ and 1.01 l mg⁻¹, 204 205 respectively. These soils are denoted here as calcareous (n = 44). There were no significant 206 correlations between Al, Ca and Fe and any of the Langmuir sorption values in the calcareous soils group. Four samples with Al values of (776–779 mg kg⁻¹) that exceeded the upper value 207 208 of the 95% confidence interval around the change-point were assigned to the calcareous 209 group because of parent material, large Ca values and weak Langmuir sorption properties.

210

211 Aluminium to phosphorus ratios in non-calcareous soils.

212 There was no direct correlation between the Langmuir terms and the different measures of 213 extractable P, except for a weak correlation between Xm and FeO-P (r = 0.26). Given that 214 buffering capacity and affinity in non-calcareous soils was strongly associated with extractable Al, the ratio of M3-Al and M3-P (Al:P) was used to infer sorptive capacity and 215 216 ability to supply P, in these soils. This ratio describes capacity and affinity for P, relative to P 217 already sorbed. The weight ratios developed for Al:P were in line with other studies using Ca:P ratios (McLaren et al. 2014) with practical significance since soil testing laboratories 218 219 tend to record results on a weight basis. To relate P buffering to supply in soluble and plant available forms, ratios of Al:P were plotted against P desorption and availability as measured 220

221 by FeO-P and Morgan's P and the relationship is depicted in Figures 3(a). The smallest FeO-P values corresponded to large Al:P ratios, however, as Al:P ratios decreased towards a point 222 of inflection on the graph, below which, FeO-P increased steadily. Following the pattern in 223 224 the data, a broken line regression was fitted and a significant change-point was found (P <0.001) at the FeO-P value of 43.2 mg kg⁻¹ within a lower and upper 95% confidence interval 225 of 32.3 and 53.9 mg kg⁻¹, respectively. The value of Al:P at this change-point was 8.1 within 226 227 a confidence interval of 5.5 and 11.8, respectively. To relate buffering to agronomic P, the 228 relationship between Al:P and Morgan's P was examined and a significant change-point was found in the relationship (P < 0.001). A broken line regression fitted to the data derived a 229 change-point for Morgan's P at 7.3 mg l⁻¹ within a 95% confidence interval of 5.9 and 8.7 mg 230 231 1⁻¹(Figure 3(b)). The value of Al:P at the Morgan's P change-point was 11.7 within the confidence interval of 9.2 and 15.0. Since the estimate of 11.7 for Al:P at the Morgan's P 232 233 change-point falls within the confidence interval for value of Al:P at the FeO-P change-point, they are not significantly different. Thus the Al:P ratio at the Morgan's P change-point is 234 235 consistent, within the range of possible values estimated at the FeO-P change-point. The 236 parameter estimates for relationships in Figure 3(a) and 3(b) are presented in Table 5.

237

238 Calcium to P ratios in calcareous soils

239 Mehlich3 extractable Ca and P were expressed as a ratio (Ca:P) and plotted against values of 240 FeO-P and Morgan's P in a similar analysis for calcareous soils. One sample with an 241 extremely large Ca value of 8888 mg kg⁻¹ which biased the relationship was removed from 242 the analysis and a non-linear, negative relationship between Ca:P and FeO-P ($R^2 = 0.32$) and 243 Morgan's P ($R^2 = 0.33$), was observed. These relationships showed no change points above 244 or below which, P became more or less available, and the regressions were not strong enough 245 to be predictive. 247 Relationship between P tests in calcareous and non-calcareous soils.

The standard agronomic P test used in Ireland, Morgan's P, was significantly correlated with 248 each of the other extractable P methods FeO-P (r = 0.73), M3-P (r = 0.79) and Olsen-P (r = 0.79) 249 250 0.63) across all soils. Correlations across all soils may conceal the effect that variations in Al and Ca could have on extraction efficiency in the soils presented in this study. Figures 4(a)251 252 and 4(b) depict the relationship between Morgan's P and measures of available P M3-P, 253 Olsen-P, for non-calcareous and calcareous soil groups. The relationships between Morgan's P and M3-P, and Olsen P were broadly similar within each of the soil group, however, they 254 255 deviated as soil P concentrations increased, indicating less extractable M3-P and Olsen-P at a 256 given Morgan's P concentration in the calcareous soils. The relationship between FeO-P and Morgan's P (Figure 5) was more comparable between soil groups within the typical 257 agronomic range (Morgan's P 0–10 mg l⁻¹) but had some divergence in the relationship at 258 259 very large P values. However, only a small number of samples were present at extremely large Morgan's P values. Overall, the relationships between these variables was stronger in 260 non-calcareous ($R^2 = 0.71$) compared to calcareous soils ($R^2 = 0.42$) and the regression 261 equations between soil tests are presented in Table 4. 262

263

264 **Discussion**

265

266 Langmuir sorption properties and soil characteristics

Values of Langmuir sorption parameters in this present study are in line with those reported elsewhere on agricultural soils over a soil pH range 4.6–8.3 (Indiati *et al.* 1999). These authors reported Xm ranging from $37.2-4990 \text{ mg kg}^{-1}$, b values from $0.75-6.12 \text{ l mg}^{-1}$ and MBC values from $51.4-5260.8 \text{ l kg}^{-1}$. Quintero *et al.* (1999) reported a decrease in P retention and maximum buffering capacities in soils as pH increased from 3 to 7, and attributed a strong buffering capacity to large amounts of both Al and Fe, in line with results reported here. Calcareous soils in this present study recorded small values of Xm, b and MBC and which are in line with those reported by Amrani *et al.* (1999) on calcareous soils in Manitoba with Langmuir buffer capacities and affinity values between $146 - 808 \text{ l kg}^{-1}$ and $0.15 - 0.6 \text{ l mg}^{-1}$, respectively.

Soils grouped as non-calcareous exhibited greater Xm, b and MBC, because of their 277 278 greater amounts of extractable Al. Broggi et al. (2011) found significant influence of soil pH 279 and Al content, with smaller adsorption energies in limed soils. In this present study, soil pH 280 and Al were negatively correlated with each other and associated with strong P sorbing soils. 281 The strong association between P sorption and amounts of extractable Al in preference to Fe 282 is also in line with other international, (Burkitt et al. 2002; Paulter and Sims, 2000) and Irish 283 studies (Daly et al. 2001; Maguire et al. 2001). Consistent with results reported here, some 284 researchers reported Al to be more strongly correlated than Fe with P sorption (Villapando & 285 Graetz, 2001). Extractable Al is also strongly associated with plant growth and Ballard & 286 Fiskell (1974) concluded that the dominant nature of extractable Al provides the best indicator of P retention in soils. 287

288

289 Relating sorptive properties to P supply

To relate measures of extractable P to sorptive properties, the relationship between ratios of M3 extractable Al:P and FeO-P was described by a broken-line regression with changepoints which revealed a value for Al:P below which FeO-P concentrations increased steadily. Above this Al:P change-point, soils were characterized as strongly P sorbing, indicative of poor Al-P solubility. A similar relationship was found between Al:P and Morgan's P and the change point corresponded to a Morgan's P value of 7.3 mg Γ^{-1} . This value falls within the

upper end of the current soil test P index range for optimum P availability (5.1–8 mg l⁻¹) set 296 297 for Irish soils by Schulte & Herlihy (2007). Above the Al:P change-points for both FeO-P 298 and Morgan's P it is likely that the P sorbed on surface oxides of Al was more tightly sorbed 299 and less available for release as soluble and plant available forms. Once the amounts of P 300 increased in the soils and reduced the Al:P ratios below the change point values, the soils 301 began to supply P in soluble and plant-available forms. The Al:P values at the change-points 302 for both FeO-P and Morgan's P were consistent and not statistically different. Soils below the 303 change-point were typified by large MBC and b values, with low pH and large Al concentrations and are described here as strong 'P fixing' non-calcareous soils. Quintero et 304 305 al. (1999) reported that 70 % of added P in strongly P fixing soils remained in the non-labile 306 P pools, whereas 30-40 % of P added to soils with poor buffering capacity was non-labile, 307 and noted that greater rates of fertilizer P are required for soils with large Al and Fe contents 308 to reach an optimum P availability for plant uptake. Similarly, Fernandez & Warren (1994) 309 related different sorption properties to varying amounts of extractable Al and concluded that 310 fertiliser rates should vary according to amount of Al in soils. Our results implications for 311 fertilizer recommendations on strong P fixing soils where Al:P ratios are large although require further data to be able to revise existing recommendations. However, this work 312 313 provides easily measured soil properties (Al:P) that can be combined with conventional soil 314 testing to describe P supply potential across soils with different sorptive properties.

315

316 Interpreting the Morgan's P threshold within agronomic P indices.

Across the soils studied here the change-point value for Morgan's P of 7.3 mg 1^{-1} in Figure 3(b) represents the ability to supply P in a plant available form consistently. Accounting for some of the variability between soils a confidence interval around this change-point Morgan's P concentration of 5.9 to 8.7 mg 1^{-1} was calculated. This range is possibly more

compatible with the previous target index (Index 3) range of 6.1 to 10 mg l⁻¹ Morgan's P 321 (Coulter. 2004), before they were revised downward to $5.1-8 \text{ mg l}^{-1}$. These revised values 322 were based on grass yield response studies of a range of fertilizer P inputs over 8 soil series 323 324 (Schulte & Herlihy, 2007). However, the soils used in the latter trial comprised of four 325 limestone and four non-limestone parent material soil groups, and did not cover the range of soil chemistry found in soils in Ireland. The fertilizer rates required to reach 95% potential 326 yield and herbage P concentration of 3.0 g kg⁻¹ were not statistically related to differences 327 between soil types or parent material groups. The lack of contrast in soil chemical properties 328 329 across the eight soil series failed to provide the statistical significance required to include soil 330 type as a factor and ultimately resulted in a P index system that does not account for 331 differences in P sorption or buffering capacities that could influence fertilizer uptake and 332 critical soil P values.

333 The difference in P sorption characteristics between calcareous and non-calcareous soils in this present study results largely from contrasting soil properties such as soil pH and 334 335 extractable Al and Ca. Herlihy & McGrath (2007) found no significant difference in Langmuir sorption maxima among soil series and soil parent material groups (limestone 336 verses non-limestone) with the exception of equilibrium buffer capacity (EBC) values for the 337 range 0-3 mg l⁻¹ P where there were larger values were observed in the non-limestone soil 338 339 groups. They also reported no significant differences in pH, clay, oxalate-extractable Al, with 340 the exception of significantly larger oxalate extractable Fe in non-limestone soils. The lack of 341 contrast between their soil groups in terms of chemical properties made it difficult to 342 statistically distinguish sorption characteristics in their data.

343

344 Processes of sorption and precipitation: implications for soil P management

345 In line with previous studies on Irish soils, extractable inorganic P as measured by M3-P, 346 Olsen-P and FeO-P were strongly correlated with Morgan's P on soils collected in this study (Daly et al. 2001; Daly & Casey, 2005; Herlihy et al. 2006). However, the contrast in soils, 347 348 exhibited by differences in Langmuir sorption properties, soil pH and amounts of extractable 349 Al, Fe and Ca, influenced the relationships between extractable P and Morgan's P. These 350 relationships indicated larger Olsen and M3-P concentrations in non-calcareous soils and 351 smaller concentrations in calcareous soils, at similar Morgan's P values, particularly as soil 352 test P concentrations increased. The relationship between Olsen-P and Morgan's P was reported for 199 soils sampled in the cross-border region of Northern Ireland (Foy et al. 353 1997) and although the soil tests were strongly related ($R^2=0.70$), the authors reported the 354 355 Olsen-Morgan relationship in high pH soils and soils over carboniferous geology followed a 356 different trend in predicting the smallest Olsen-P values. The relationship between Morgan's 357 P and Olsen-P in this present study concurs with Foy et al. (1997) with the smallest Olsen-P values observed in the calcareous soils at similar Morgan's P values. Morgan's P may be 358 359 over-estimating Olsen-P and M3-P at large P values P in calcareous soils, where the process 360 of Ca-P precipitation becomes dominant at large soil P contents.

The FeO-P values were more comparable in non-calcareous and calcareous soil groups at similar Morgan's P value and the relationship showed only slight divergence at greater extremes of soil test P concentrations. The FeO-P test is reported to be a 'P sink' method, favoured over chemical extraction for estimates of P desorption and for quantity intensity relationships, as it is not confounded by chemistry (McLaren *et al.* 2014; Van Rotterdam *et al.* 2012) and suitable across soil types.

These diverging relationships indicate stronger sorptive properties in highly buffered soils (Dougherty *et al.* 2011) with a larger pool of P in non-calcareous soils that can continuously supply P to the soil solution and precipitation of Ca-P at large P values in 370 calcareous soils. Chemical fractionation on calcareous soils has recorded 98% of total 371 inorganic P as Ca-P and desorption from the soil matrix using FeO strips was related to labile P available for plant uptake (Ahmad et al. 2006). The interaction of P with soil components 372 373 in calcareous soils is controlled by both surface reactions and precipitation and at small P 374 concentrations non-carbonate clays provide surfaces for P adsorption. However, once these 375 sites have been filled, and at greater soil P concentrations, calcium carbonate plays a role in the precipitation of P and subsequent reduction of P availability to the plant and soil solution 376 377 (Von Wandruszka, 2006).

378 Precipitation of Ca-P in soils at large P values could have implications for P transport 379 at the catchment scale. Mellander et al. (2012) recorded sub-hourly P concentrations and 380 discharge delivered to spring water in a karst catchment in western Ireland characterized by 381 calcareous parent material overlain by shallow soils. Despite a large proportion of soils 382 measuring excessive Morgan's P concentrations with a good connectivity to groundwater, 383 background P concentrations in groundwater were small and posed a low risk of increasing 384 surface water P concentrations. The authors conclude that processes or P attenuation in the 385 soil, geology and hydrology in this catchment were limiting P transfers to surface waters. Smaller values of M3-P at large Morgan's P values in the calcareous soils in our study 386 supports the attenuation hypothesis in Mellander et al. (2012) that the process of desorption is 387 388 overtaken by precipitation at elevated soil P contents.

Herlihy *et al.* (2004) reported results from a P trial that included a 'zero P treatment' on soils across the full range of soil P indices (1–4) used in Ireland, for both calcareous and non-calacareous soil parent material groups. They reported more rapid rates of decline in soil test P in limestone soils at low P indices that could possibly be explained by the poorer buffering in these soils compared to the non-limestone soil group. As P is depleted from the soil solution phase from plant uptake and not replaced in added fertilizer, soils that are better buffered can utilise their larger store of labile inorganic P reserves to replenish the soil solution phase more rapidly than poorly buffered soils with smaller overall P reserves which may be more tightly held in the shorter term. Strongly buffered soil will exhibit a slower rate of decline in soil test P concentration than poorly buffered soils over time and modelling studies on Irish soils (Schulte *et al.* 2010) could consider including sorptive properties such as Al:P to improve estimates.

401 Differences in soil P supply and release between calcareous and non-calcareous soils 402 have implications for how P fertilizer application should be managed. For non-calcareous 403 soils splitting the annual P allowance using a little and often approach to P application for 404 grassland, over the annual growing season may be more appropriate for maintaining adequate 405 P in soil solution for soils with P concentrations below the change-point for Morgan's P (< 7.3 mg l^{-1}) with Al:P ratios above those at the change-point. Above the threshold Morgan's P 406 407 value, maintenance P applications could be applied at the start of the growing season. In contrast it may be less important to split the annual P application for calcareous soils as a 408 409 smaller proportion of added P would be fixed. In terms of P build-up, calcareous soils may 410 require less P fertilizer overall than strong high P fixing soils to build up to the agronomic 411 optimal range (Index 3) because the binding energies are less in these soils.

412

413 **Conclusions and Recommendations**

414

Phosphorus dynamics in grassland soils are controlled by the chemical nature of the soil and the P retained in soil. In this study, P sorptive capacity and strength to which P is bound in soil was to a large extent influenced by soil pH and amounts of extractable Al. This was illustrated by a change-point value of extractable Al in soil, above which P buffering capacities increased linearly, facilitating the classification of soils into calcareous and non420 calcareous groups. Non-calcareous soils with large buffering capacities had larger P reserves 421 at similar soil test P than calcareous soils with smaller P buffer capacities. Acknowledging 422 differences in P buffering capacities in grassland soils, combined with soil P testing, could 423 improve P management and efficiency; however, sorption isotherms are too time consuming 424 for routine soil testing. Ratios of Al:P could be used as a surrogate for Langmuir buffer 425 capacities in non-calcareous soils, as this indicator was related to supply potential and release of P in an available form. This indicates the potential for incorporating easily measured soil 426 427 chemical properties with agronomic P testing, particularly for strong P fixing soils where P availability could be improved by adjusting Al:P ratios. Diverging relationships between soil 428 429 P tests in calcareous and non-calcareous soils can be attributed to differences in P reserves in 430 strongly buffered non-calcareous soils, and P precipitation at large P contents, in calcareous 431 soils with poor P buffering capacities. Distinguishing differences in P buffering in soil, at 432 process scale, will support improved agronomic and environmental assessment at both field and catchment scale, specifically, our interpretation of P build up and release, P balances, 433 434 rates of soil P decline and P attenuation in catchments.

435

436 Acknowledgement

437

438 The authors would like to thank Dr. Jim Grant for statistical advice and support and his

439 technical assistance in data analysis.

440

441

442

443

445 **References**

- Ahmad, W., Rahmatullah, Ahmad-Gill, M. & Aziz, T. 2006. Bioavailability of Inorganic
 Phosphorus Fractions in Calcareous Soils Estimated by Neubauer Technique, IronImpregnated Filter Paper, and Chemical Test. *Communications in Soil Science & Plant Analysis*, 37, 1185-1193.
- Amrani, M., Westfall, D.G. and Mougli, L. 1999. Phosphate sorption in calcareous Moroccan
 soils as affected by soil properties. *Communications in Soil Science & Plant Analysis*, 30,
 1299-1314.
- Ballard, R. & Fiskell, J.G. 1974. Phosphorus retention in coastal plain forest soils: I.
 Relationship to soil properties. *Soil Science Society of America Journal*, **38**, 250-255.
- Broggi, F., Oliveira, A.D., Freire, F.J., Freire, M.D.S. & Nascimento, C.D. 2011. Phosphate
 capacity factor in mineralogically different soils in Pernambuco and the influence of pH
 on the maximum capacity of adsorption. *Ciência e Agrotecnologia*, **35**, 77-83.
- Burkitt, L.L., Gourley, C.J.P., Sale, P.W.G., Uren, N.C., Hannah, M.C. 2001. Factors
 affecting the change in extractable phosphorus following the application of phosphatic
 fertiliser on pasture soils in southern Victoria. *Australian Journal of Soil Research*, 39,
 759-771.
- Burkitt, L.L., Moody, P.W., Gourley, C.J.P. & Hannah, M.C. 2002. A simple phosphorus
 buffering index for Australian soils. *Australian Journal of Soil Research*, 40, 497-513.
- 464 Cordell, D., Drangert, J.O. & White, S. 2009. The story of phosphorus: Global food security
 465 and food for thought. *Global Environmental Change*, **19**, 292-305.
- 466 Coulter, B.S. (ed.) 2004. Nutrient and Trace Element Advice for Grassland, Tillage,
 467 Vegetable and Fruit Crops. 2nd edition. Teagasc Publications, Dublin, Ireland.
- 468 Daly, K. & Casey, A. 2005. Environmental aspects of soil phosphorus testing. Irish Journal
- 469 of Agricultural & Food Research, 44, 261-279.

- Daly, K., Jeffrey, D. & Tunney, H. 2001. The effect of soil type on phosphorus sorption
 capacity and desorption dynamics in Irish grassland soils. *Soil Use & Management*, 17,
 12-20.
- 473 Dougherty, W.J., Burkitt, L.L. & Corkrey, R. 2011. The effect of soil P sorption properties
 474 and phosphorus fertiliser application strategy on "incidental" phosphorus fertiliser
 475 characteristics: a laboratory study. 2011. *Nutrient Cycling in Agroecosystems*, **89**, 189476 197.
- Fernandes, M.L.V. & Warren, G.P. 1994. Exchangeable and non-exchangeable phosphate
 sorption in Portuguese soils. *Fertilizer research*, **37**, 23-34.
- 479 Foy, R.H., Tunney, H., Carroll, M.J., Byrne, E., Gately, T., Bailey, J.S. 1997. A comparison
- 480 of Olsen and Morgan soil phosphorus test results from the cross-border region of Ireland.

481 Irish Journal of Agricultural & Food Research, **36**, 185-193.

- Gardiner, M.J. & Radford, T. 1980. Soil Associations of Ireland and Their Land Use
 Potential. *Explanatory Bulletin to Soil Map of Ireland, 1980, National Soil Survey of Ireland*, Soil Survey Bulletin No.6. ISBN 0-905442-49-0.
- Herlihy, M., & McGrath, D. 2007. Phosphorus fractions and adsorption characteristics in
 grassland soils of varied soil phosphorus status. *Nutrient Cycling in Agroecosystems*, 77,
 15-27.
- Herlihy, M., McCarthy, J., Breen, J. & Moles, R. 2004. Effects over time of fertiliser P and
 soil series on P balance, soil-test P and herbage production. *Irish Journal of Agricultural & Food Research*, 43, 147-160.
- Herlihy, M., McCarthy, J. & Brennan, D. 2006. Divergent relationships of phosphorus soil
 tests in temperate grassland soils. *Communications in Soil Science & Plant Analysis*, 37,
 693-705.

- Indiati, R., Neri, U., Sharpley, A.N. & Fernandes, M.L. 1999. Extractability of Added
 Phosphorus in Short-Term Equilibration Tests of Portuguese Soils. *Communications in Soil Science & Plant Analysis*, **30**, 1807-1818.
- John, M.K. 1970. Colorimetric determination of phosphorus in soil and plant materials with
 ascorbic acid. *Soil Science*, **109**, 214-220.
- Johnston, A.E. & Steen, I. 2000. Understanding Phosphorus and its Use in Agriculture. *EFMA. European Fertilizer Manufacturers Association*, Brussels, Belgium
- 501 Lalor, L. & Coulter, S. 2008. Major and minor micronutrient advice for productive
 502 agricultural crops. Teagasc.
- Maguire, R.O., Foy, R.H., Bailey, J.S. & Sims, J.T. 2001. Estimation of the phosphorus
 sorption capacity of acidic soils in Ireland. *European Journal of Soil Science*, 52, 479-487.
- 505 McLaren, T., Guppy, C., Tighe, M., Moody, P. & Bell, M. 2014. Dilute acid extraction is a
- 506 useful indicator of the supply of slowly available phosphorus in Vertisols. *Soil Science*

507 Society of America Journal, **78**, 139-146

- Mehlich, A.1984. Mehlich-3 soil test extractant: a modification of Mehlich-2 extractant. *Communications in Soil Science & Plant Analysis.* 15, 1409-1416.
- 510 Mellander, P.E., Jordan, P., Wall, D.P., Melland, A.R., Meehan, R., Kelly, C. et al. 2012.
- 511 Delivery and impact bypass in a karst aquifer with high phosphorus source and pathway
- 512 potential. *Water Research*, **46**, 2225-2236.
- Menon, R.G., Hammond, L.L. & Sissingh, H.A. 1988. Determination of plant-available
 phosphorus by the iron hydroxide-impregnated filter paper (Pi) soil test. *Soil Science Society of America Journal*, 52, 110-115.
- 516 Moody, P.W. 2007. Interpretation of a single-point P buffering index for adjusting critical
- 517 levels of the Colwell soil P test. *Australian Journal of Soil Research*, **45**, 55-62.

- 518 Nair, P.S., Logan, T.J., Sharpley, A.N., Sommers, L.E., Tabatabai, M.A. & Yuan, T. 1984.
- 519 Interlaboratory comparison of a standardized phosphorus adsorption procedure. *Journal of*520 *Environmental Quality*, 13, 591-595.
- 521 Paulter, M.C. & Sims, J.T. 2000. Relationship between soil test P, soluble P and P saturation
 522 in Delaware soils. *Soil Science Society of America Journal*, 64, 765-733.
- 523 Peech, M. & English, L. 1944. Rapid microchemical soil test. *Soil Science*, **57**, 167-194.
- 524 Quintero, C.E., Boschetti, G.E. & Benavidez, R.A. 1999 Phosphorus retention in some soils
- of the Argentinean Mesopotamia. *Communications in Soil Science & Plant Analysis*, 30
 (9&10), 1449-1461.
- 527 Reijneveld, A., Termorshuizen, A., Vedder, H. & Oenema, O. 2014. Strategy for innovation
- in soil tests illustrated for P tests. *Communications in Soil Science & Plant Analysis*, 45,
 498-515.
- 530 Schulte, R.P.O. & Herlihy, M. 2007. Quantifying responses to phosphorus in Irish grasslands:
- Interactions of soil and fertiliser with yield and P concentration. *European Journal of Agronomy*, **26**, 144-153.
- 533 Schulte, R.P.O., Melland, A.R., Fenton, O., Herlihy, M., Richards, K. & Jordan, P. 2010.
- Modelling soil phosphorus decline: expectations of Water Framework Directive policies. *Environmental science & policy*, 13, 472-484.
- Villapando, R.R. & Graetz, D.A. 2001. Phosphorus sorption and desorption properties of the
 spodic horizon from selected Florida Spodosols. *Soil Science Society of America Journal*,
 65, 331-339.
- 539 Van Rotterdam, A.M.D., Bussink, D.W., Temminghoff, E.J.M. & Van Reimsdijk, W.K.
- 540 2012. Predicting the potential of soils to supply phosphorus by integrating soil chemical
- 541 processes and standard soil tests. *Geoderma*, **189**, 617-626.

542	Von Wandruszka, R. 2006. Phosphorus retention in calcareous soils and the effect of organic
543	matter on its mobility. Geochemical Transactions, 7, 6.
544	Vos, R. 1998. Fertility recommendations: past and present. Communications in Soil Science
545	& Plant Analysis, 29 , 1429-1440.
546	
547	
548	
549	
550	
551	
552	
553	
554	

Table 1 The locations and description of soil associations used for sample collection, describing arrangements of Great Soil Groups and
 associated parent material with texture and drainage class included (Gardiner & Radford, 1980).

551	assor
558	_
	-

Location	Soil	Great Soil Groups	Parent Material	Texture class	Drainage class
	Association				
Tipperary	34	Luvisol (70%), Gleysol (20%),	Gravelly limestone	Gravelly loam	Well drained
		Cambisol (10%)	till		
Tipperary	30	Luvisol (70%), Cambisol (20%),	Calcareous, limestone	Sandy loam	Well drained
		Gleysol (5%), Histosol (5%).	till		
Cork	15	Podzol (60%), Cambisol (20%), Gleysol	Old red sandstone,	Gravelly loam	Well drained
		(20%)	shale		
Roscommon	31	Luvisol (80%), Gleysol (10%),	Calcareous glacial till	Loam	Well drained
		Cambisol (5%), Histosol (5%).			
Tyrone	25	Gleysol (50%), Cambisol (40%),	Silurian sandstone-	Clay loam	Imperfect to poor
		Histosol (10%)	shale composition		
Meath	14	Cambisol (75%), Gleysol (15%), Podzol	Ordovician shale,	Loam	Well drained
		(10%)	glacial till.		

Table 2. Range (min-max) of soil properties measured across samples collected from soil association listed in Table 1. Soil pH, %Organic matter (% OM), Morgan's P, Iron-oxide paper-strip P (FeO-P), Mehlich3 extractable Al, Fe, Ca and P (M3-P) were measured on all soils (n =120). Across soil associations Langmuir sorption isotherm values were derived for a subset of samples (n = 74). Olsen extractable P (Olsen-P) derived for a sub-set of soils (n = 60) did not include soil associations 30 and 14.

							Langmuir Te	erms		Phosphor	us Extracts	
Soil	pН	OM	Al	Fe	Ca	Xm	b	MBC	Morgan's	M3-P	FeO-P	Olsen-P
Associatio									Р			
n												
		/ %		/ mg	g kg ⁻¹		/ 1 mg ⁻¹	$/ 1 kg^{-1}$	/ mg 1 ⁻¹		/ mg kg ⁻¹	
34	5.2-7.0	5.7-15.4	69-1160	215-642	1420-8660	263-333	0.66-4.48	213-476	1.5-32.9	18-168	21.6-38	14.6-57.6
30	5.4-7.1	5.5-11.9	349-780	254-592	1795-5543	263-400	0.63-1.29	250-417	1.6-17.5	17.8-101	19.6-69.6	
15	5.0-6.7	8.3-15.7	580-1530	454-655	894-5674	270-625	0.54-2.71	147-1429	6.6-18.2	60.8-211	39.8-83.2	30.6-77.5
31	4.9-6.9	5.9-15.5	317-1240	306-612	777-5040	303-435	0.35-2.0	119-769	1.5-11.8	13.4-110	15.4-58.4	8.9-34.2
25	4.8-6.3	5.3-14.5	901-1730	390-666	284-3566	385-588	1.43-7.67	714-3333	1.8-25.8	18.4-285	15.4-128	15.5-89.6
14	4.7-6.4	7.5-14.7	254-1730	373-556	769-4721	357-556	1.0-2.89	435-1429	2.6-9.2	30.5-114	17.4-59.2	

Table 3. Pearson product moment correlation matrix of r-values significant at P = 0.05 derived between soil pH, % organic matter (OM), Mehlich3 extractable P, Al, Fe and Ca / mg kg⁻¹, Morgan's P / mg l⁻¹, Olsen-P / mg kg⁻¹ and iron-oxide paper strip P (FeO-P / mg kg⁻¹). Sorption properties include Langmuir sorption maximum (Xm / mg kg⁻¹); binding energy, (b / 1 mg⁻¹) and maximum buffer capacity, (MBC / 1 kg⁻¹). NS denotes non-significant correlation at P = 0.05.

	Xm	В	MBC	Morgan's P	M3-P	FeO-P	Olsen-P	pН	OM	Al	Fe	Ca
Xm	1.00											
b	0.38	1.00										
MBC	0.53	0.98	1.00									
Morgan's P	NS	NS	NS	1.00								
M3-P	0.41	NS	NS	0.79	1.00							
FeO-P	0.26	NS	NS	0.73	0.78	1.00						
Olsen-P	NS	NS	NS	0.63	0.88	0.78	1.00					
pН	-0.46	-0.62	-0.65	0.23	NS	NS	NS	1.00				
ÔM	0.29	NS	NS	0.31	0.37	0.23	0.42	NS	1.00			
Al	0.53	0.76	0.80	NS	0.22	NS	NS	-0.71	NS	1.00		
Fe	0.48	0.41	0.48	NS	0.40	NS	0.43	-0.48	0.47	0.55	1.00	
Ca	NS	-0.57	-0.58	0.24	NS	NS	NS	0.80	0.32	-0.66	-0.40	1.00

Table 4 Regression equations for relationships between logMBC and soil pH across all soils, and relationships between Morgan's P and extractable measures of soil P, namely, Mehlich3-P, Olsen-P and iron-oxide paper strip P (FeO-P) for both calcareous and non-calcareous soils.

у	Equation		Coefficients	Standard errors	R^2
logMBC	$y = a + (b \times pH)$	а	4.92	0.23	0.55
		b	-0.37	0.04	
M2 P non colearnous soils	$\mathbf{v} = \mathbf{o} + (\mathbf{b} \mathbf{v} \mathbf{M} \mathbf{o} \mathbf{r} \mathbf{g} \mathbf{o} \mathbf{r}' \mathbf{c} \mathbf{P})$	0	Q 1/	5 50	0.83
M3-P non-calcaleous sons	$y = a + (0 \times Morgan S P)$	a 1	0.14	5.59	0.85
		b	11.01	0.58	
M3-P calcareous soil	$v = a + (b \times Morgan's P)$	а	22.47	4.35	0.80
	y (c	h	4 67	0.38	
		U	1.07	0.50	
Olsen-P non-calcareous soils	$y = a + (b \times Morgan's P)$	а	9.34	3.82	0.66
		b	3.52	0.38	
	(1 - 1)	_	15 64	4.10	0.62
Olsen-P calcareous soils	$y = a + (b \times Morgan \otimes P)$	a	15.64	4.19	0.62
		b	1.18	0.27	
FeO-P non-calcareous soils	v = a + (h x Morgan's P)	а	5 18	2 59	0.71
100 1 non culculcous sons	$y = a + (b \times 1001 \text{gain } 51)$	u h	3.68	0.27	0.71
		U	5.08	0.27	
FeO-P calcareous soils	$y = a + (b \times Morgan' \otimes P)$	а	21.68	3.00	0.42
		b	1.39	0.26	

Table 5 Parameter estimates derived from the broken-line regressions fitted to non-calcareous soil data with change-points (x = c) highlighted in bold, for extractable Al mg kg⁻¹, FeO-P mg kg⁻¹ and Morgan's P mg l⁻¹.

		Estimate	Standard error	t value	Р	95% confi	dence limit
						Lower	Upper
Figure 2	Intercept 1	6.1799	0.3299	18.73	< 0.0001	5.5217	6.8381
(n = 73)	Slope 1	-0.0008	0.0007	-1.22	0.2280	-0.0023	0.0005
	$\mathbf{x} = \mathbf{c}$	611.92	52.1751	11.76	<0.0001	507.83	716.01
	Value of logMBC at change-point	5.6331	0.1324	42.54	< 0.0001	5.3690	5.8973
	Slope 2	0.0029	0.0002	13.08	< 0.0001	0.0024	0.0033
Figure 3(a)	Intercept 1	3.9422	0.1433	27.51	< 0.0001	3.6568	4.2276
(n = 76)	Slope 1	-0.0402	0.0054	-7.40	< 0.0001	-0.0511	-0.0294
	$\mathbf{x} = \mathbf{c}$	43.1187	5.4313	7.94	<0.0001	32.3013	53.9361
	Value of logAl:P at change-point	2.2048	0.1692	13.03	< 0.0001	1.8678	2.5417
	Slope 2	-0.00861	0.004052	-2.12	0.0369	-0.1668	-0.0005
Figure 3(b)	Intercept 1	4.2247	0.1627	25.92	< 0.0001	3.9006	4.5487
(n = 76)	Slope 1	-0.2297	0.0311	-7.37	< 0.0001	-0.2917	-0.1676
	$\mathbf{x} = \mathbf{c}$	7.3041	0.7153	10.21	<0.0001	5.8794	8.7287
	Value of logAl:P at change-point	2.5471	0.1148	22.19	< 0.0001	2.3185	2.7756
	Slope 2	-0.06198	0.01129	-5.49	< 0.0001	-0.0844	-0.0395

Table 6 Summary statistics of general soil properties; pH, % organic matter (OM), Mehlich3 extractable Al, Fe, Ca, P, Morgan's P and ironoxide paper strip P (FeO-P) of soils divided into non-calcareous (n = 76) and calcareous (n = 44) soil groups. Langmuir sorption isotherm data, sorption maxima (Xm), maximum buffer capacity (MBC), binding energy (b) in non-calcareous soils (n = 41) and calcareous soils (n = 33).

615 Olsen-P values were measured on a subset of non-calcareous (n = 47) and calcareous soils (n = 14).

616

						Ι	angmuir T.	erms	Р	hosphoru	is Extracts	S
	Soil	OM	Al	Fe	Ca	Xm	b	MBC	Morgan's	M3-P	FeO-P	Olsen-
	pН								Р			Р
	_	/%		/m	g kg ⁻¹		/l mg ⁻¹	/l kg ⁻¹	/mg l ⁻¹		/mg kg ⁻	1
						Non-	calcareous	soils				
Min	4.7	5.3	807	369	284	303	0.80	244	1.5	13.4	3.3	8.9
Max	6.3	18.3	1730	666	4713	625	8.00	3333	25.8	285	128	89.6
Median	5.2	9.5	1072	553	1451	476	2.56	1250	7.0	85.9	30.4	37.3
Mean	5.3	10.1	1108	540	1631	467	3.04	1417	8.15	98	35.6	40
						Ca	lcareous sc	oils				
Min	5.2	5.5	68.9	215	777	244	0.35	119	1.5	17.8	4.4	16.8
Max	7.1	16.3	794	642	8660	667	2.10	1000	32.9	168	69.6	57.6
Median	6.2	9.3	658	425	3539	323	0.94	303	7.1	56.1	31	27
Mean	6.2	9.9	597	432	3457	343	1.01	346	9.0	64.7	34.3	31

617

620

621 **Figure 1.** The relationship between log transformed Langmuir maximum buffer capacity 622 (MBC) and soil pH ($R^2 = 0.55$) for the sub-set of samples for which sorption isotherms were 623 derived (n=74).

624

Figure 2. Broken-line regression fitted to the relationship between log transformed Langmuir maximum buffer capacity and extractable for the subset of samples for which sorption isotherms were derived (n = 74), with a change-point derived for Al at 615 mg kg⁻¹ within a lower and upper confidence interval of 508 and 716 mg kg⁻¹, respectively.

629

Figure 3. Ratios of Mehlich3-Al to M3-P (Al:P) plotted against (a) FeO-P and (b) Morgan's P in non-calcareous (n = 76) separated out from the full data-set. A broken-line regression fitted to both data-set calculated change-points for FeO-P at 43.2 mg kg⁻¹ within lower and upper confidence interval of 32.3 and 53.9 mg kg⁻¹, respectively. The dotted line represents change-point values of Morgan's P and Al:P on the x and y axes, respectively. Morgan's P change-point was derived at 7.3 mg l⁻¹ within lower and upper confidence interval of 5.9 and 8.7 mg l⁻¹, respectively.

637

Figure 4. The Relationship between Morgan's P and (a) M3-P for both non-calcareous (n = 76) and calcareous soils (n = 44); and (b) Olsen-P for non-calcareous (n = 47) and calcareous (n = 14) soils, with regression lines and confidence intervals showing diverging relationships in both soil groups with smaller M3-P and Olsen-P values, large Morgan's P values in calcareous soils.

- **Figure 5.** The relationship between Morgan's P and FeO-P in non-calcareous (n = 76) and
- 645 calcareous (n = 44) soils with overlapping confidence intervals showing comparable values
- 646 of FeO-P at similar Morgan's P between both soil groups.