

Relationships between reservoir water quality and catchment habitat type

Gough, Rachel; Cohen, Yvonne; Fenner, Nathalie; Cannon, Jonathan; Freeman, Christopher

International Journal of Environmental Technology and Management

DOI: 10.1504/IJETM.2016.074799

Published: 01/01/2016

Peer reviewed version

Cyswllt i'r cyhoeddiad / Link to publication

Dyfyniad o'r fersiwn a gyhoeddwyd / Citation for published version (APA): Gough, R., Cohen, Y., Fenner, N., Cannon, J., & Freeman, C. (2016). Relationships between reservoir water quality and catchment habitat type. *International Journal of Environmental Technology and Management, 19*(1), 16-39. https://doi.org/10.1504/IJETM.2016.074799

Hawliau Cyffredinol / General rights Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

· Users may download and print one copy of any publication from the public portal for the purpose of private study or research.

- You may not further distribute the material or use it for any profit-making activity or commercial gain
 You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

1 Relationships between reservoir water quality and catchment

2 habitat type

3 Rachel Gough¹*, Yvonne Cohen¹, Nathalie Fenner¹, Jonathan Cannon², Christopher Freeman¹

4 ¹ School of Biological Sciences, Bangor University, Deiniol Road, Bangor, Gwynedd, LL57 2UW, UK

5 ² Dŵr Cymru Welsh Water Head Office, Pentwyn Road, Nelson, Treharris, Mid Glamorgan, CF46 6LY,

6 *UK*

7 *Corresponding author: <u>bss408@bangor.ac.uk</u>.

- 8
- 9 Abstract
- 10 Numerous catchment characteristics including topography, geology, soil and vegetation are reported
- 11 to exert a strong influence on mean surface water properties. The present study employs a
- 12 geographical information system (GIS) approach to examine, for the first time, the relationship
- 13 between reservoir water quality (dissolved organic carbon (DOC) concentration, colour, nitrate
- 14 concentration and pH) and catchment Phase 1 Habitat coverage. Analysis was conducted on 2
- 15 occasions and at 2 different spatial scales. Numerous statistically significant correlations were
- 16 identified, suggesting the use of Phase 1 Habitat data could help improve predictive models of
- 17 surface water quality. The occurrence and strength of correlations varied seasonally in response, we
- 18 argue, to temporal variations in hydrological regime and anthropogenic activity. The data also
- 19 suggest that the proximity of habitat types to the reservoir is significant in affecting reservoir water
- 20 quality. The findings are used to recommend suitable measures for drinking water companies to
- 21 mitigate against water quality issues.
- 22 **Key words:** dissolved organic carbon; drinking water; catchment; Phase 1 Habitat; soil; geographical
- 23 information system.

24 1. Introduction

25 The biogeochemical properties of surface waters are acquired, to a large extent, during the passage 26 of water through the catchment due to the interaction of water with vegetation, soils and mineral 27 layers. Various organic and inorganic compounds will be solubilised and transported downstream 28 during runoff, influencing solute concentrations, pH and ionic strength (Stutter et al., 2006). 29 Although surface water quality exhibits temporal variations in response to weather events and 30 seasonal drivers (Gergel et al., 1999; Scott et al., 1998; Soulsby et al., 2006), physical catchment 31 characteristics including topography, geology, soil and vegetation type will, to a large extent, 32 determine mean biogeochemical characteristics (Billett and Cresser, 1992; Clair et al., 1994; Holden 33 et al., 2007; Hope et al., 1994; Sobek et al., 2007). Amongst these variables, soil type is widely 34 considered to represent a dominant control on surface water composition and quality (Aitkenhead 35 et al., 1999; Billett and Cresser, 1992; Hope et al., 1997; Soulsby et al., 2006; Stutter et al., 2006). 36 Though its development is strongly influenced by other catchment features including soil 37 characteristics, habitat type may also be an important factor affecting surface water quality. 38 Vegetation type influences catchment hydrology, primary production and organic matter inputs 39 (Ordóñez et al., 2008; Zhang et al., 2011), which affect soil composition and chemistry and in turn, 40 drainage water quality. Forested catchments for example, have been associated with the production 41 of dissolved organic carbon (DOC)-rich drainage waters (Grayson et al., 2012; Hope et al., 1994) with 42 differences in DOC concentration and flux also reported between different tree species (Chow et al., 43 2009; Fröberg et al., 2011; Gough et al., 2012). Wetland habitat coverage is also reported to be a 44 strong predictor of surface water DOC concentration (Gergel et al., 1999; Hope et al., 1994). 45 Surface water characteristics can also be strongly affected by anthropogenic activity. For example, the application of agricultural fertilisers has been associated with significant leaching of nutrients 46

47 (nitrates and phosphates) into surface waters (Badruzzaman et al., 2012). Elevated nutrient

48 concentrations may in turn result in eutrophication and algal blooms (Correll, 1998; Freeman et al.,

2009; Hecky and Kilham, 1988; Vollenweider, 1968), which are particularly problematic in drinking
water supplies (Smith, 1998). Liming of agricultural land has been associated with increased surface
water pH (Hindar et al., 2003). Drainage of wetland habitats in an attempt to improve their
economic value has also been linked to elevated colour and DOC concentrations in surface waters
(Holden et al., 2004; Wallage et al., 2006).

54 The Phase 1 Habitat Survey of Wales, completed by the Countryside Council of Wales (CCW) in 1997, provides a record of habitat coverage and land use (Howe et al., 2005). In its digitised form, using 55 56 geographical information system (GIS) software, the data offers a useful means of measuring the 57 spatial extent of different habitat types within catchments. Since the classification scheme includes 58 both natural habitats and anthropogenic features (e.g. arable land and improved grassland), the data 59 holds significant potential for researchers concerned with investigating catchment influences on 60 surface water quality. The characteristics of surface waters supplying drinking water treatment 61 works (WTWs) is important for water companies which have a responsibility to provide a safe and 62 reliable drinking water supply for their customers. The concentration of DOC in surface waters is 63 particularly important, with the removal of DOC from drinking water supplies representing the single 64 biggest treatment cost for the water treatment industry (Watts et al., 2001). Elevated DOC 65 concentrations in raw water can inflate treatment costs by increasing the coagulant and disinfectant 66 doses required (Chow et al., 2005; Edzwald, 1993) and the frequency of filter backwashes (Eikebrokk 67 et al., 2004). DOC in finished water is problematic since it can cause undesirable colour, odour and 68 taste (Davies et al., 2004; WHO, 2011), transports organic and inorganic micro-pollutants (Gao et al., 1998; Rothwell et al., 2007) and leads to bacterial regrowth in distribution systems (Prévost et al., 69 70 1998). Crucially, DOC also acts as a precursor to potentially harmful disinfection by-products (DBPs) 71 including trihalomethanes (THMs). These are formed during chlorination, a treatment necessary to 72 ensure that finished water meets microbiological safety standards (WHO, 2011).

73 Rising surface water DOC concentrations have been observed in many areas of central and northern 74 Europe and North America in the past couple of decades (Freeman et al., 2001; Hejzlar et al., 2003; 75 Monteith et al., 2007; Skjelkvåle et al., 2005; Stoddard et al., 2003; Worrall et al., 2003). In the UK, 76 measurements undertaken at 22 upland sites showed a mean increase in DOC concentration of 91% 77 between 1988 and 2003 (Evans et al., 2005). DOC concentrations tend to be highest, and rising most 78 rapidly in peat-dominated, upland catchments (Freeman et al., 2001), which in the UK, supply over 79 70% of drinking water (Watts et al., 2001). In this context of declining surface water quality, 80 developing a better understanding of catchment influences is crucial for the drinking water industry. 81 The importance of catchment characteristics in affecting the quality of drinking water supplies is 82 recognised by the UK drinking water regulator, the drinking water inspectorate (DWI) who recommend that "catchment and raw water source protection" is included in the drinking water 83 84 safety plans of drinking water providers (DWI, 2005).

85 A GIS approach, which can offer an effective means of visualising and measuring landscape features 86 is increasingly being used in the study of catchment influences on hydrochemistry. GIS software has 87 become an important tool in the modelling of hydrological processes and its use in developing 88 predictive models for various water quality parameters within catchments based on land use and 89 other catchment characteristics is particularly relevant for water treatment companies. For example 90 Foster and McDonald (2000) used GIS and spatially referenced data on pastoral farming intensity to 91 model and display sources of cryptosporidium risk in drinking water catchments. Lake et al. (2003) 92 developed a nitrate leaching model using GIS and information on a number of physical catchment 93 characteristics. This was used to identify areas of groundwater vulnerable to nitrate pollution. 94 Recently, Grayson et al. (2012) used a GIS approach, and ITE land cover data (similar to Phase 1 95 Habitat data) to identify correlations between drinking water reservoir colour and the spatial extent 96 of different land cover classes. A multicriteria evaluation approach was then used to develop a 97 predictive model for water colour production potential in the catchments and create a colorimetric,

risk-based map from the data. However, as yet, the use of Phase 1 Habitat data for predicting
catchment water quality has not been explored.

100 This study investigates potential relationships between Phase 1 Habitat classes and reservoir water 101 quality (DOC concentration, colour, nitrate concentration and pH). GIS mapping was used to 102 measure the spatial extent of Phase 1 Habitat types in 16 drinking water reservoir catchments in 103 north Wales. Correlation analysis was then used to identify statistically significant relationships 104 between these land cover classes and reservoir water quality in spring and autumn. Analyses were 105 carried out both at a whole-catchment scale, and in a 250 m buffer zone surrounding the reservoirs 106 in order to assess the importance of proximity in the occurrence and strength of the correlations. 107 Such research is important for informing future catchment management practices. Identifying 108 problematic land cover will also help water treatment companies target monitoring programmes 109 and mitigation strategies, and improved understanding of seasonality in raw water quality will also 110 enable better optimization of treatment processes.

111 2. Methods

112 2.1. Study sites and sampling regime

113 Water samples were collected on 2 occasions (in September 2007 and March 2008) from the raw 114 water (i.e. pre-treatment) supply of 16 WTWs in north Wales. Where the raw water supply was derived from more than 1 reservoir, composite samples were collected, and Phase 1 Habitat data 115 116 was also combined. The timing of sampling was chosen to correspond with the seasonal maximum 117 (autumn) and minimum (spring) in reservoir DOC concentration. 14 of the WTWs included in this 118 study are located in upland catchments, with the remaining 2 situated in lowland, agricultural areas. 119 Uplands are defined as areas more than 250 m above sea level (Mitchell, 1991). These areas are 120 typically characterised by high rainfall, low mean temperatures and acidic soils (Foster and 121 McDonald, 2000).

122 2.2. Hydrochemical analysis

123 pH was measured on un-filtered samples using a Mettler Toledo S20 pH meter (Mettler Toledo,

Leicester, UK), calibrated daily with pH 4 and pH 7 reference standards (Sigma-Aldrich, Dorset, UK).

125 Colour measurements (Hazen) were obtained from WTW data at the time of sample collection. One

126 degree Hazen (1 mg L⁻¹ Pt/Co) is defined as the colour produced by 1 mg L⁻¹ Pt (as K₂PtCl₆) in the

127 presence of 2 mg L⁻¹ cobalt (II) chloride hexahydrate (Mitchell and McDonald, 1992).

128 Before DOC measurement, samples were passed through a 0.45 µm cellulose acetate filter to 129 remove particulate organic carbon, as per the operational definition of DOC (Thurman, 1985). DOC 130 concentrations were determined using a Shimadzu Total Organic Carbon 5000 analyser (Shimadzu, 131 Milton Keynes, UK), with a carrier gas of high purity air at a flow rate of 150 mL min⁻¹ and a 33 µL 132 injection volume. Calibration was performed with a one point calibration, using a 100 mg L^{-1} KO₄H₅C₈ 133 solution (total organic carbon – TOC) and a 100 mg L⁻¹ Na₂CO₃/100 mg L⁻¹ NaHCO₃ solution (inorganic 134 carbon – IC). DOC concentrations were calculated by subtracting IC values from TOC values. Analysis 135 of TOC and IC standard solutions at 10 mg L⁻¹ intervals demonstrated that the analyser performed 136 linearly from 0 to 200 mg L^{-1} , with r^2 values > 0.9. All reagents were supplied by Sigma-Aldrich, 137 Dorset, UK.

Nitrate concentration was determined using a Dionex DX-120 ion chromatograph equipped with an
IonPac AS14A anion analytical column (both Thermo Scientific, Hertfordshire, UK). The eluent was a
1.0 mM Na₂HCO₃/8.0 mM NaCO₃ solution (reagents supplied by Sigma-Aldrich, Dorset, UK) made
with Milli Q water and the flow rate, 1 mL min⁻¹. Concentrations were determined using a five point
calibration with standard Dionex solutions.

143 2.3. Geographical information systems (GIS) analysis

Version 9.2 of the ArcGIS package (ESRI, Buckinghamshire, UK) was used to display and quantify the
 spatial extent of habitat types within each reservoir catchment. First, the watersheds associated

with each reservoir were mapped. This was achieved using the *Hydrology* functions in the *Spatial Analyst* extension and a digital elevation model downloaded from Digimap (EDINA, 2014) (10 m
resolution). Defined watersheds were then clipped to other GIS layers displaying habitat type.
Habitat information was displayed using digitised version of the Phase 1 Habitat Survey of Wales
(Howe et al., 2005). In addition to this whole-catchment analysis, habitat coverage was also
measured in a 250 m-wide zone around the perimeter of each reservoir.

152 2.4. Statistical analysis

153 For statistical analysis, Phase 1 Habitat categories were organized into more generalised groupings

154 (Table 1). Statistical analysis was performed using version 20 of the SPSS statistical package (IBM,

155 New York, USA). Depending on the conditions satisfied by the data, Pearson's correlation and

156 Spearman's correlation analyses was employed to test for significant correlations between Phase 1

157 habitat type coverage and reservoir water quality. This analysis was also performed using the subset

158 of Phase 1 Habitat data covering a zone of 250 m directly adjacent to the reservoir.

159 **3. Results and discussion**

160 *3.1. DOC and colour*

161 The absence of any statistically significant correlations between catchment woodland and scrub 162 coverage and reservoir DOC concentration and colour (Table 2 and 3) is surprising given that 163 previous research indicates a strong positive relationship between forest coverage and DOC 164 concentration (Grayson et al., 2012; Hope et al., 1994). High DOC flux from forested catchments is 165 partly due to high DOC loading as rainwater passes through above ground biomass (Kawasaki et al., 166 2005; Stevens et al., 1989) as well as the large source of leachable carbon in the litter layer (Hongve, 167 1999). However, DOC concentrations are also reported to vary significantly between different tree 168 species (Gough et al., 2012). Our habitat categories did not account for this potential variation, 169 which may explain the absence of any statistically significant correlations in this study.

170 A moderate negative correlation was observed between unimproved grassland and spring DOC 171 concentration at the whole catchment scale (p < 0.05; Table 2) and no correlations between 172 unimproved grassland and DOC or colour in the 250 m buffer zone analysis (Table 3). The negative 173 correlation corroborates the findings of previous studies. For example, Grayson et al. (2012) report a 174 significant negative correlation between water colour and moorland grass coverage across 18 175 drinking water catchments in Yorkshire. In a UK-wide study, Armstrong et al. (2007) found that 176 heather dominated, drained catchments produced the highest water colour followed by mixed 177 vegetation and grass dominated catchments. Van den Berg et al. (2012) also reported lowest mean 178 pore water DOC concentration in grassland sites compared with other vegetation categories 179 (woodlands, heathlands and moorlands) in their survey of 41 UK sites. This association may relate to 180 solubility controls since colour release in temperate grasslands is reported to be suppressed by 181 acidic conditions (Hopkins et al., 1990; Miller, 2008). However, our unimproved grassland category 182 included neutral and calcareous grassland and no correlations were identified in the present study 183 between pH and unimproved grassland coverage.

184 Negative correlations were identified between tall herb and fern habitat coverage and autumn DOC 185 concentration and colour at the whole catchment scale (both p < 0.05; Table 2) and between tall 186 herb and fern coverage and autumn colour in the 250 m buffer zone analysis (p < 0.05; Table 3). At 187 first this result seems unexpected since bracken coverage, which was dominant in this habitat class, 188 has been associated with high primary productivity and the accumulation of large amounts of litter, 189 forming a large pool of organic matter (Marrs et al., 2000). However, Potthast et al. (2012) observed 190 that, compared with pasture land (Setaria grass) the litter present in bracken habitat showed a 191 significantly lower rate of decay. In addition, they found a significant decrease in microbial biomass 192 and activity when pasture land was invaded by bracken. Therefore, if bracken coverage tends to 193 replace improved grassland habitats in drinking water catchments, its presence may reduce DOC 194 production. The occurrence of these correlations in autumn may relate to this being the litter fall

period, when the leaching of DOC from decomposing litter would normally contribute significantly to
DOC export (Kalbitz et al., 2000).

197 Negative correlations between heathland coverage and DOC concentration and colour occurred at 198 both spatial scales. In the whole catchment analysis, heathland coverage displayed a moderate 199 negative correlation with autumn DOC concentration and colour (both p < 0.05) and a strong 200 negative correlation with spring colour (p < 0.01; Table 2). In the 250 m buffer analysis a moderate 201 negative correlation was identified with autumn DOC concentration and colour (both p < 0.05), a 202 moderate negative correlation with spring DOC concentration (p < 0.05) and a strong negative 203 correlation with spring colour (p < 0.01; Table 3). These negative relationships were surprising given 204 that Calluna, a common species in heath habitats, has been reported to produce highly-coloured 205 drainage water (Grayson et al., 2012). This has been attributed to their relatively dry soil conditions 206 which confer high rates of aerobic microbial decomposition (Clutterbuck and Yallop, 2010). Many 207 heath habitats were also formed as a result of peatland drainage. This former status is likely to 208 further enhance DOC and colour release due to the large carbon stocks associated with peat 209 substrate (Fenner et al., 2009). Conversely however, moisture constraints in heath habitats are 210 reported to inhibit phenol oxidase activity (Toberman et al., 2008). According to the enzymic latch 211 theory, this can suppress DOC production by causing an accumulation of phenolic compounds which 212 inhibit the activity of hydrolase enzymes (Freeman et al., 2001). This may explain the negative 213 correlations between heathland coverage and reservoir DOC concentration and colour observed in 214 the present study. Overall, the relationship appeared stronger at the 250 m buffer scale, suggesting 215 that proximity to the reservoir affected the degree to which this habitat influenced reservoir water 216 quality.

At the whole catchment scale a moderate positive correlation was identified between fen/ mire habitat and autumn DOC concentration (p < 0.05; Table 2). This habitat also correlated positively with autumn and spring DOC concentration in the 250 m buffer analysis (both p < 0.05; Table 3).

220 Positive correlations between swamp coverage and reservoir DOC concentration and colour were 221 also identified and were striking in terms of the strength of the correlations observed and their 222 occurrence at both spatial scales and both sampling times (Table 2 and 3); all were strong positive 223 correlations (p < 0.01) except for the swamp/ autumn DOC concentration correlation at the whole 224 catchment scale which was a moderate positive trend (p < 0.05). These positive relationships are 225 likely to be linked to the wetland status of these habitats. Percentage wetland coverage has been 226 identified as an important predictor of stream water DOC concentration (Eckhardt and Moore, 1990; 227 Gergel et al., 1999; Hope et al., 1994). A combination of high primary productivity and low 228 decomposition rates causes the accumulation of deep layers of peat in wetland environments 229 (Mitsch and Gosselink, 2000). The considerable depth of organic material in such environments 230 provides a large pool of available carbon (Thurman, 1985) and the inhibitory effect of anaerobic 231 conditions on microbial metabolism promotes the formation of DOC end products (Fenner et al., 232 2009). In addition, in wetland systems, the depth of the organic horizon limits contact between 233 drainage waters and the adsorption sites within the mineral soil horizon, which also contributes to 234 high DOC loading (Tipping et al., 1999). However, our data also show an absence of statistically 235 significant correlations between DOC concentration/ colour and other habitat categories which are, 236 or include, wetlands (marsh/ marshy grassland, bog, flush and spring). This suggests that the type of 237 wetland present may be an important determinant of drainage water DOC concentration. It may be 238 significant that, of these wetland habitat types, swamp and fen/ mire habitats tend to be more 239 nutrient-rich than the other wetland habitats (Mitsch and Gosselink, 2000) which may support 240 higher rates of primary productivity and thus a larger pool of organic carbon.

Strong positive correlations between arable coverage and DOC concentration were observed at the whole catchment scale in autumn and spring (both p < 0.01; Table 2). Moderate positive correlations were also identified with colour at both sampling times (both p < 0.05). The 250 m buffer zone could not be included in this analysis since there was only 1 catchment where arable land was present in this zone (Figure 1). The interpretation of these positive correlations is not straightforward since in

246 previous studies arable land use has been associated with lower carbon content than other land use 247 types. For example, soil solution carbon concentrations for soils in northern Saskatchewan, Canada, 248 decreased in the following order: aspen forest > recently cleared forest > wheat/ fallow field (McFee 249 and Kelly, 1995). Similarly, in their review article, Chantigny (2003) reports that dissolved organic 250 matter concentrations vary as follows: forest soils > grassland soils > arable soils. This variation, it is 251 suggested, is partly due to differences in vegetation type (e.g. tree vs. herbaceous plant) (Chantigny, 252 2003) as well as the lower carbon content associated with arable soils (Zsolnay, 1996). In addition, 253 aerobic conditions, which tend to occur in arable soils encourage the complete mineralisation of 254 organic matter to CO_2 , as opposed to DOC and CO_2 end products in anaerobic decomposition (Boddy 255 et al., 2008; Fenner et al., 2009). However, water soluble carbon content in arable soils is also 256 reported to vary depending on crop plants used (Zsolnay, 1996) and temporally, during crop cycles 257 (Campbell et al., 1999) and with successive cultivations (Delprat et al., 1997). In addition, application 258 of organic fertilisers on agricultural soils is reported to substantially increase the concentration of 259 soluble organic carbon (Gregorich et al., 1998). Although it is not possible to isolate the cause of the 260 positive correlations observed here between arable land use and DOC concentration/ colour, it is 261 notable that the correlations occurred despite this land use being virtually absent in the reservoir 262 250 m buffer zone.

263 At the whole catchment scale moderate positive correlations were identified between buildings 264 coverage and autumn DOC concentration (p < 0.05) and between the "other" category and autumn 265 DOC concentration (p < 0.05) and spring colour (p < 0.05; Table 2). A moderate positive correlation 266 was also found between buildings coverage and autumn DOC concentration in the 250 m buffer 267 zone analysis (p < 0.05; Table 3). Given the rural location of the catchments in the present study it is 268 likely that farm buildings will account for a significant proportion of the buildings category. Indeed, a 269 strong positive correlation between buildings and arable coverage was identified in the whole 270 catchment data ($r_s = 0.803$, p < 0.01). This correlation may therefore explain the relationship 271 between buildings and DOC concentration, though the reason for this being confined to the autumn

analysis is unclear. The correlations between the "other" category and DOC concentration and
colour at the whole catchment scale are also difficult to interpret since this category includes
unknown habitat classes ("not accessed" land and "illegible" data inputs). It may be significant
however, that bare ground (J.4; Table 1) is included in this category, which may provide a source of
readily-leachable organic matter.

277 *3.2. Nitrate and pH*

278 Strong positive correlations were observed between arable coverage and nitrate concentration in 279 the whole catchment analysis in autumn and spring (both p < 0.01; Table 2). This is likely to be 280 caused by the leaching of organic or inorganic fertiliser (Neill, 1989). The stronger correlation in the 281 spring analysis may be due to the timing of fertiliser application, which for arable crops tends to 282 occur in late winter/spring (MAFF, 2000; Trudgill et al., 1991). As mentioned earlier, the 250 m 283 buffer zone could not be included in the analysis due to there being only 1 catchment where arable 284 was present in this zone. It is interesting therefore that strong correlations exist at the whole 285 catchment scale despite arable coverage being virtually absent in the 250 m buffer zone. The 286 application of fertiliser may also explain the strong positive correlations between improved 287 grassland coverage and nitrate concentration in both the whole catchment analysis (p < 0.01 in 288 autumn and spring; Table 2), and the 250 m buffer zone analysis (p < 0.05 and p < 0.01 in autumn 289 and spring, respectively; Table 3). Again, stronger correlations in the spring analysis at both spatial 290 scales are likely to be due to the timing of fertiliser application. The strong positive correlations 291 between woodland and scrub coverage and spring nitrate concentrations at both spatial scales (both 292 p < 0.01; Table 2 and 3) may be explained by the application of fertiliser prior to tree planting in 293 commercial forestry plantations (Drinan et al., 2013).

294 Moderate positive correlations were identified between fen/ mire coverage and reservoir nitrate 295 concentration in spring sampling at the whole catchment scale (p < 0.05; Table 2) and in both 296 autumn and spring in the 250 m buffer analysis (both p < 0.05; Table 3). These correlations are likely

to relate to the nutrient status of this habitat; fen systems are typically associated with relatively
high nutrient concentrations due to their being supplied by drainage water from surrounding
mineral soil (Mitsch and Gosselink, 2000).

300 The positive correlation between buildings and nitrate concentration at the whole catchment scale 301 in spring (p < 0.05; Table 2) may be due to the positive correlation mentioned earlier between 302 buildings and arable coverage. In addition, the urine and droppings of mammals and birds has been 303 identified as an important non-agricultural source of ammonia (DEFRA, 2002). Nitrifying bacteria in 304 the soil may then convert ammonia to nitrate. Therefore, assuming that a significant proportion of 305 the buildings in this category are farms, then the leaching of ammonia from domestic animals may 306 also account for this correlation. The leaching of nitrates from septic tanks and fertiliser stores may 307 also explain this association.

308 A moderate positive correlation was observed between marsh/ marshy grassland coverage and 309 reservoir nitrate concentration but only in the 250 m buffer analysis in spring (p < 0.05; Table 3). The 310 reason for this is not clear but may be an artefact of the positive association between marsh/ 311 marshy grassland and other habitat types displaying a positive correlation with nitrate. For example, 312 in the 250 m buffer zone analysis, marsh/ marshy grassland coverage correlates positively with 313 woodland and scrub ($r_s = 0.646$, p < 0.01), improved grassland ($r_s = 0.771$, p < 0.01) and fen/mire 314 habitat ($r_s = 0.560$, p < 0.05), all of which show a positive correlation with spring nitrate 315 concentration at this spatial scale.

The supply of nitrate and phosphate is critical in determining the growth rates of phytoplankton in freshwater systems with elevated concentrations resulting in eutrophication in some cases (Hecky and Kilham, 1988). In drinking water sources algal blooms can lead to a number of treatment issues including taste and odour problems, elevated TOC levels, increased coagulant and chlorine demand, membrane fouling and an increase in DBPs (Bernhardt et al., 1991; Li et al., 2012; Nguyen et al., 2005). Elevated reservoir nitrate concentrations may also increase the formation of nitrogenous

322 DBPs (NDBPs), produced during the disinfection stage of water treatment either directly, or
 323 indirectly *via* increased algal biomass and consequently increased concentrations of dissolved
 324 organic nitrogen in the raw water (Ritson et al., 2014).

The negative correlation between heathland coverage and reservoir pH in the 250 m buffer analysis (*p* < 0.05; Table 3) is likely to be related to the preference of heath vegetation for acidic soils (Holden et al., 2007) which has a corresponding effect on drainage water pH (Cresser and Edwards, 1987). A positive relationship has been reported between DOC solubility and pH (Lumsdon et al., 2005). Thus solubility controls may also help to explain the negative correlations between heathland coverage and reservoir DOC concentration and colour.

331 The absence of correlations between pH and some of the other habitat types which tend to be 332 associated with peat substrates (bare peat, bog, fen/ mire, flush and spring and swamp) is surprising 333 given that peatlands tend to produce acidic drainage waters. This is reported to result from the 334 accumulation of organic acids, the enhanced activity of sulphur-metabolising bacteria under 335 waterlogged conditions and high cation exchange capacity (Clymo, 1964; Urban et al., 1995). 336 Coniferous forest stands, which represent a large proportion of forest coverage in north Wales, are 337 also associated with acidic drainage waters (Eisalou et al., 2013; Gough et al., 2012). A significant 338 decrease in pH has been reported as rainwater passes through coniferous canopies and litter 339 (Eisalou et al., 2013), due to the high exchangeable acidity of coniferous foliage and litter and the 340 fact that coniferous litter is readily leached of organic acids (Alfredsson et al., 1998; Nykvist, 1963).

341 3.3. Temporal and spatial variations in correlations

At the whole catchment scale, more associations between Phase 1 Habitat categories and DOC concentration were identified in autumn than in spring. A difference in hydrological regime due to higher rainfall in September than March may explain this contrast. Higher rainfall will result in a larger contribution of surface runoff to discharge water (Horton, 1933) which is likely to enhance the

influence of surface characteristics such as vegetation/ litter characteristics. The influence of habitat
may also be enhanced by higher above ground biomass following the growing season.

348 Overall there were fewer statistically significant correlations identified between habitat types and 349 reservoir nitrate concentration in autumn compared with spring (Table 2 and 3). We have already 350 suggested that fertiliser application may be significant in explaining a number of correlations 351 between habitat type and reservoir nitrate concentration (Neill, 1989), and that its timing may 352 explain the greater number and strength of correlations in spring (MAFF, 2000; Trudgill et al., 1991). 353 However, the drivers of seasonal variations in surface water nitrate concentration are known to be 354 complex, comprising numerous biogeochemical and hydrological processes (Martin et al., 2004). 355 Stream nitrate concentrations tend to exhibit a summer minima and a winter maxima (Neill, 1989; 356 Reynolds et al., 1992). This is explained in part by variations in the supply of nitrogen. For example, 357 in the summer, the availability of leachable nitrate in the soil is limited by lower atmospheric inputs 358 and plant uptake and in streams by macrophyte uptake (Cooke and Cooper, 1988) and denitrification 359 (Hill, 1979). In winter on the other hand, plant uptake decreases, atmospheric inputs increase and in-360 stream losses decrease due to lower primary productivity (Reynolds et al., 1992). Surface water 361 nitrate levels may also be transport-limited, with a strong association reported with precipitation 362 and discharge (Neill, 1989; Trudgill et al., 1991). However, given that lower rainfall totals were 363 recorded in March than in September, it is more likely that reservoir nitrate levels in spring (which 364 were higher than in autumn), were supply-limited.

365 It is difficult to interpret the overall effect of spatial scale on relationships between habitat classes 366 and reservoir water quality since at the 250 m buffer scale, a number of habitat classes are present 367 in only a few catchments, or are absent altogether. For example, there was only 1 catchment where 368 arable land was identified in the 250 m buffer zone. However, there was an obvious similarity in the 369 occurrence of significant correlations at the 2 spatial scales. Although there was no clear difference 370 in the strength of the correlations between the 2 spatial scales, this similarity would suggest that

Phase 1 Habitat coverage in the 250 m buffer zone was more important than in the wider catchment
in affecting reservoir water quality. Previous studies have reported improved regressions between
land use and surface water quality parameters when the riparian area was included, or weighted
more heavily than other areas (Levine and Jones, 1990; Osborne and Wiley, 1988).

The relationship between reservoir water quality and catchment characteristics at different spatial scales may also vary temporally. For example, Gergel et al. (1999), in their study of wetland influence on DOC concentrations in Wisconsin lakes and rivers, found that in autumn, wetland coverage in the whole catchment was the best predictor of lake water DOC concentration whereas in summer, wetland coverage within 50 m of lakes was the best predictor. This could relate to seasonal hydrological changes since the timing of sampling in autumn and spring corresponded with base flow and peak flow conditions, respectively (Hurley et al., 1995).

382 In this study we noted the absence of a number correlations between Phase 1 Habitat classes and 383 surface water parameters that would typically be expected. For example the lack of positive 384 correlations between woodland and scrub habitat and a number of wetland habitats and DOC 385 concentration/ colour was unexpected. This may be due to the influence of various other catchment 386 features not included in the present study, but which previous studies have reported to influence 387 surface water chemistry. For example, slope will mediate the relationship between catchment 388 characteristics and surface water quality due to its influence on surface runoff (Rochelle et al., 1989) 389 as well as being a predictor of soil organic horizon depth (Rasmussen et al., 1989) and wetland 390 abundance (Eckhardt and Moore, 1990). The development of a particular soil type reflects a number 391 of factors including climate, parent material, topography and vegetation and is reported to be a 392 crucial factor in determining surface water composition and quality (Aitkenhead et al., 1999; Billett 393 and Cresser, 1992; Hope et al., 1997). Indeed information on soil chemical characteristics has formed 394 the basis of a number of predictive models for stream water solute concentrations (Billett and 395 Cresser, 1992; Christophersen and Wright, 1981; Cosby et al., 1985). Soil type influences spatial

396 patterns of water flow and storage (Grayson and Western, 2001; Weiler and Naef, 2003). In addition, 397 adsorption processes in mineral soils regulate the transport of organic carbon and the soil organic 398 pool is reported to be the main factor controlling DOC flux in streams (Aitkenhead et al., 1999). It 399 should also be noted that the topographic watershed does not necessarily correspond with 400 groundwater influence, which may also strongly impact on reservoir water quality (Garrison et al., 401 1987). In addition, the Phase 1 Habitat Survey of Wales was conducted between 1987 and 1997 and 402 it is likely that a number of land use changes have occurred in this time (Stevens et al., 2004). 403 Nonetheless, the present study has demonstrated that the data continues to be relevant to the 404 study of surface water quality and the extent of its coverage represents a significant benefit for 405 drinking water companies.

406 Various processes occurring in the water body will also affect surface water chemistry. For example, 407 as mentioned earlier, seasonal variations in the uptake of nitrate in surface waters influence nitrate 408 concentrations (Cooke and Cooper, 1988). DOC loss from reservoirs is reported to occur as a result 409 of sedimentation and mineralisation processes (Algesten et al., 2004), with precipitation also 410 affecting DOC concentrations via a dilution effect (Engstrom, 1987). Conversely, DOC may be 411 produced within the water body (autochthonous DOC), potentially supressing the relationship 412 between DOC concentration and colour and terrestrial drivers. Nonetheless, a substantial number of 413 correlations were identified between Phase 1 Habitat data and reservoir water characteristics in the 414 present study. This, we suggest, relates both to the direct influence of vegetation/ land cover on 415 runoff and drainage water quality and also to habitat classes being predictors of other physical 416 characteristics such as peat soils or certain management practices.

417 *3.4. Implications for potable water treatment*

418 Though previous research has cited the relationship between catchment wetland coverage and

419 surface water DOC and colour loading (Eckhardt and Moore, 1990; Gergel et al., 1999; Hope et al.,

420 1994), our data suggest that wetland type may significantly affect the magnitude of this relationship.

421 The identification of positive associations between swamp and, to a lesser extent, fen/ mire habitats 422 and DOC concentration/ colour, possibly the result of their nutrient supply (Mitsch and Gosselink, 423 2000), may justify monitoring the quality of drainage waters in these areas. Given that these habitat 424 types also occupy very small proportions of the catchments included in this study (Figure 2), it may 425 be that diverting drainage water from these areas would be a cost-effective strategy for improving 426 reservoir water quality. Monitoring of drainage waters and diversion of water courses may also be 427 appropriate for areas of arable land use which arguably exerted the strongest influence on surface 428 water quality. This land use class correlated with DOC, colour and nitrate concentrations at both 429 sampling times despite being virtually absent from the 250 m buffer zone of the reservoirs. The 430 apparent impact of arable land on reservoir water quality highlights the importance of excluding this 431 activity from areas close to the reservoir. In cases where the diversion of problematic drainage 432 waters is not possible, it may be appropriate to blend reservoir water with water from another 433 catchment, as has been employed previously as a strategy to reduce water discolouration from peat 434 (Grayson et al., 2012).

435 Allowing the expansion of habitat types whose coverage correlates negatively with DOC/ colour may 436 be a suitable strategy in some cases. However, the potential benefits to surface water quality may 437 be outweighed by other detrimental impacts. For example, tall herb and fern coverage, which in the 438 present study was dominated by bracken, correlates negatively with DOC and colour but bracken 439 habitat has no economic value and is associated with the leaching of carcinogenic compounds such 440 as ptaquiloside (Rasmussen et al., 2003). Heathland coverage also correlated negatively with DOC 441 concentration and colour in the present study, but this we argue, may relate to site-specific factors 442 such as soil moisture constraints since Calluna vegetation is typically associated with highly-coloured 443 waters (Grayson et al., 2012).

Given the number of statistically significant correlations identified in the present study, and thenational scale of Phase 1 Habitat data, we suggest that future research should explore integrating

Phase 1 Habitat data into predictive models for reservoir water quality. The present study has also
highlights the fact that correlations between catchment characteristics and surface water quality
may vary on a seasonal basis; an important consideration as researchers seek to develop more
sophisticated predictive models.

450 4. Conclusions

This study has considered, for the first time, the use of catchment Phase 1 Habitat data for
predicting reservoir water quality. Our analysis was conducted at two different spatial and temporal
scales, to investigate the effect of season and the proximity of habitat types to the reservoir in
affecting potential associations between habitat type and water quality parameters.

Numerous statistically significant correlations were observed between Phase 1 Habitat classes and reservoir water quality. These could be explained either by the direct impact of vegetation on drainage water or its association with other physical catchment characteristics or land management practices. Arable land cover appeared to have the most substantial impact on reservoir water quality, correlating strongly with DOC concentration, colour and nitrate concentration at both sampling times. This was despite arable land being virtually absent from the 250 m buffer zone.

The degree to which habitat classes affected reservoir water quality appeared to vary on a seasonal basis, with more correlations between habitat classes and DOC concentration in autumn, and between habitat classes and nitrate concentration in spring. However, a striking similarity was observed between correlations at the whole catchment scale and within the 250 m buffer zone. We therefore suggest that in general, the influence of habitat coverage on reservoir water quality parameters increases with proximity to the reservoir.

Although previous research has identified a link between wetland abundance and surface water
DOC/ colour loading, our findings suggest that the type of wetland habitat present is also important.
We found that swamp and fen/ mire habitats were the only wetland types which correlated with

470 reservoir DOC or colour. This specificity, we suggest, may relate to the high nutrient levels in these

471 habitats which may support higher rates of primary production than other wetland types.

472 Based on the number and strength of correlations observed, we suggest that predictive models for

473 surface water characteristics based on catchment characteristics could be improved by incorporating

474 Phase 1 Habitat data. The findings of this study are important for drinking water companies

475 concerned with maintaining finished water quality and may be of use in targeting monitoring of

drainage water in catchments and selecting appropriate mitigation strategies such as diverting or

477 blending water.

478 5. Acknowledgements

This research was part-funded by the European Social Fund (ESF) through the European Union's Convergence programme administered by the Welsh Government. Match funding and access to water samples and water quality data was provided by Dŵr Cymru Welsh Water (DCWW). Christopher Freeman and Nathalie Fenner acknowledge funding from NERC under the first EU ERA-EnvHealth call (FP7-ENV-2007- CSA-1.2.3-01) and the large NERC grant, *Characterisation of the nature, origins and ecological significance of dissolved organic matter in freshwater ecosystems* (NE/K010689/1).

486 6. References

Aitkenhead, J.A., Hope, D. and Billett, M.F. (1999) 'The relationship between dissolved organic
carbon in stream water and soil organic carbon pools at different spatial scales', *Hydrological Processes*, Vol. 13 No. 8, pp.1289 - 1302.

Alfredsson, H., Condron, L.M., Clarholm, M. and Davis, M.R. (1998) 'Changes in soil acidity and
organic matter following the establishment of conifers on former grassland in New Zealand', *Forest Ecology and Management*, Vol. 112 No. 3, pp. 245 - 252.

Algesten, G., Sobek, S., Bergström, A., Ågren, A., Tranvik, L.J. and Jansson, M. (2004) 'Role of lakes
for organic carbon cycling in the Boreal zone', *Global Change Biology*, Vol. 10 No. 1, pp. 141 - 147.

495 Armstrong, A., Holden, J., Kay, P., Chapman, P., Clements, S., Foulger, M., McDonald, A.T. and

Walker, A. (2007) *Grip-blocking in upland catchments; costs and benefits*, Tech. Rep. Final Report for
Yorkshire Water.

Badruzzaman, M., Pinzon, J., Oppenheimer, J. and Jacangelo, J.G. (2012) 'Sources of nutrients
impacting surface waters in Florida: a review', *Journal of Environmental Management*, Vol. 109, pp.
80 - 92.

Bernhardt, H., Schell, H., Hoyer, O. and Lusse, B. (1991) 'Influence of algogenic organic substances on
flocculation and filtration', *WISA*, Vol. 1, pp. 41 - 57.

Billett, M.F. and Cresser, M.S. (1992) 'Predicting stream-water quality using catchment and soil
chemical characteristics', *Environmental Pollution*, Vol. 77 No. 2-3, pp. 263 - 268.

Boddy, E., Roberts, P., Hill, P.W., Farrar, J. and Jones, D.L. (2008) 'Turnover of low molecular weight
dissolved organic C (DOC) and microbial C exhibit different temperature sensitivities in Arctic tundra
soils', *Soil Biology and Biochemistry*, Vol. 40 No.7, pp. 1557 - 1566.

Campbell, C., Lafond, G., Biederbeck, V., Wen, G., Schoenau, J. and Hahn, D. (1999) 'Seasonal trends
in soil biochemical attributes: effects of crop management on a black chernozem', *Canadian Journal*of Soil Science, Vol. 79 No. 1, pp. 85 - 97.

511 Chantigny, M.H. (2003) 'Dissolved and water-extractable organic matter in soils: a review on the 512 influence of land use and management practices', *Geoderma*, Vol. 113 No. 3-4, pp. 357 - 380.

513 Chow, C.W., Fabris, R., Drikas, M. and Holmes, M. (2005) 'A case study of treatment performance
514 and organic character', *Journal of Water Supply Research and Technology – AQUA*, Vol. 54 No. 6, pp.
515 385 - 395.

Chow, A.T., Lee, S., O'Geen, A.T., Orozco, T., Beaudette, D., Wong, P., Hernes, P.J., Tate, K.W. and
Dahlgren, R.A. (2009) 'Litter contributions to dissolved organic matter and disinfection byproduct

518 precursors in California Oak woodland watersheds', Journal of Environmental Quality, Vol. 38 No. 6,

519 pp. 2334 - 2343.

- 520 Christophersen, N. and Wright, R.F. (1981) 'Sulfate budget and a model for sulfate concentrations in
- 521 stream water at Birkenes, a small forested catchment in southernmost Norway', *Water Resources*
- 522 *Research*, Vol. 17 No. 2, pp. 377 389.
- 523 Clair, T.A., Pollock, T.L. and Ehrman, J.M. (1994) 'Exports of carbon and nitrogen from river basins in 524 Canada's Atlantic Provinces', *Global Biogeochemical Cycles*, Vol. 8 No. 4, pp. 441 - 450.
- 525 Clutterbuck, B. and Yallop, A.R. (2010) 'Land management as a factor controlling dissolved organic
- 526 carbon release from upland peat soils 2: changes in DOC productivity over four decades', Science of
- 527 *the Total Environment*, Vol. 408 No. 24, pp. 6179 6191.
- 528 Clymo, R. (1964) 'The origin of acidity in sphagnum bogs', *The Bryologist*, Vol. 67 No. 4, pp. 427 431.
- 529 Cooke, J.G. and Cooper, A.B. (1988) 'Sources and sinks of nutrients in a New Zealand hill pasture
 530 catchment III. Nitrogen', *Hydrological Processes*, Vol. 2 No. 2, pp. 135 149.
- 531 Correll, D.L. (1988) 'The role of phosphorus in the eutrophication of receiving waters: a review',
 532 *Journal of Environmental Quality*, Vol. 27 No. 2, pp. 261 266.
- 533 Cosby, B., Hornberger, G., Galloway, J. and Wright, R. (1985) 'Modeling the effects of acid
- deposition: assessment of a lumped parameter model of soil water and streamwater chemistry', *Water Resources Research*, Vol. 21 No. 1, pp. 51 63.
- 536 Cresser, M.S. and Edwards, A. (1987) *Acidification of freshwaters*, Cambridge University Press,
 537 Cambridge.
- Davies, J-M., Roxborough, M. and Mazumder, A. (2004) 'Origins and implications of drinking water
 odours in lakes and reservoirs of British Columbia, Canada', *Water Research*, Vol. 38 No. 7, pp. 1900 1910.
- 541 Delprat, L., Chassin, P., Linères, M. and Jambert, C. (1997) 'Characterization of dissolved organic
- carbon in cleared forest soils converted to maize cultivation', *European Journal of Agronomy*, Vol. 7
 No. 1-3, pp. 201 210.
- 544 DEFRA (Department of Environment, Food and Rural Affairs) (2002) Ammonia in the UK.
- 545 Drinan, T.J., Graham, C.T., O'Halloran, J. and Harrison, S.S.C. (2013) 'The impact of catchment conifer
- 546 plantation forestry on the hydrochemistry of peatland lakes', *Science of the Total Environment*, Vol.
- 547 443, pp. 608 620.
- 548 DWI (Drinking Water Inspectorate). *A brief guide to drinking water safety plans.*
- 549 http://www.sswm.info/content/brief-guide-water-safety-plans (Accessed 11 August 2014).
- Eckhardt, B. and Moore, T. (1990) 'Controls on dissolved organic carbon concentrations in streams,
 southern Quebec', *Canadian Journal of Fisheries and Aquatic Sciences*, Vol. 47 No. 8, pp. 1537 1544.
- 552 EDINA. Digimap collections. [online] http://edina.ac.uk/digimap (Accessed 11 August 2014).

- Edzwald, J. (1993) 'Coagulation in drinking water treatment: particles, organics and coagulants', *Water Science and Technology*, Vol. 27 No. 11, pp. 21 35.
- 555 Eikebrokk, B., Vogt, R. and Liltved, H. (2004) 'NOM increase in northern European source waters:
- discussion of possible causes and impacts on coagulation/contact filtration processes', *Water Supply*,
 Vol. 4 No. 4, pp. 47 54.
- 558 Eisalou, H.K., Şengönül, K., Gökbulak, F., Serengil, Y. and Uygur, B. (2013) 'Effects of forest canopy
- cover and floor on chemical quality of water in broad leaved and coniferous forests of Istanbul,
- 560 Turkey', Forest Ecology and Management, Vol. 289, pp. 371 377.
- 561 Engstrom, D.R. (1987) 'Influence of vegetation and hydrology on the humus budgets of Labrador 562 lakes', *Canadian Journal of Fisheries and Aquatic Sciences*, Vol. 44 No. 7, pp. 1306 - 1314.
- 563 Evans, C.D., Monteith, D.T. and Cooper, D.M. (2005) 'Long-term increases in surface water dissolved 564 organic carbon: observations, possible causes and environmental impacts', *Environmental Pollution*, 565 Vol. 137 No. 1, pp. 55, 71
- 565 Vol. 137 No. 1, pp. 55 71.
- 566 Fenner, N., Freeman, C. and Worrall, F. (2009) 'Hydrological controls on dissolved organic carbon 567 production and release from UK peatlands' in Baird, A.J. et al (Eds.), *Carbon cycling in northern*
- 568 *peatlands*, American Geophysical Union, Washington DC, pp. 237-249.
- Foster, J.A. and McDonald, A.T. (2000) 'Assessing pollution risks to water supply intakes using
 geographical information systems (GIS)', *Environmental Modelling & Software*, Vol. 15 No. 3, pp. 225
 234.
- Freeman, C., Ostle, N. and Kang, H. (2001) 'An enzymic 'latch' on a global carbon store a shortage of
 oxygen locks up carbon in peatlands by restraining a single enzyme', *Nature*, Vol. 409 No. 6817, pp.
 149 149.
- Freeman, A.M., Lamon, III E.C. and Stow, C.A. (2009) 'Nutrient criteria for lakes, ponds, and
 reservoirs: a Bayesian TREED model approach', Ecological Modelling, Vol. 220 No. 5, pp. 630 639.
- 577 Freeman, C., Evans, C.D., Monteith, D.T., Reynolds, B. and Fenner, N. (2001) 'Export of organic 578 carbon from peat soils', Nature, Vol. 412 No. 6849, pp. 785 - 785.
- 579 Fröberg, M., Hansson, K., Kleja, D.B. and Alavi, G. (2011) 'Dissolved organic carbon and nitrogen 580 leaching from scots pine, norway spruce and silver birch stands in southern Sweden', *Forest Ecology*
- 581 *and Management*, Vol. 262 No. 9, pp. 1742 1747.
- 582 Gao, J.P., Maguhn, J., Spitzauer, P. and Kettrup, A. (1998) 'Sorption of pesticides in the sediment of 583 the Teufelsweiher pond (southern Germany). II: competitive adsorption, desorption of aged residues 584 and effect of dissolved organic carbon', *Water Research*, Vol. 32 No. 7, pp. 2089 - 2094.
- Garrison, P.J., Greb, S.R., Knauer, D.R., Wentz, D.A., Krohelski, J.T., Bockheim, J.G., Gherini, S.A. and
 Chen, C.W. (1987) 'Application of the ILWAS model to the northern Great Lakes states', *Lake and Reservoir Management*, Vol. 3 No. 1, pp. 356 364.

- 588 Gergel, S.E., Turner, M.G. and Kratz, T.K. (1999) 'Dissolved organic carbon as an indicator of the scale 589 of watershed influence on lakes and rivers', *Ecological Applications*, Vol. 9 No. 4, pp. 1377 - 1390.
- 590 Gough, R., Holliman, P.J., Willis, N., Jones, T.G. and Freeman, C. (2012) 'Influence of habitat on the
- quantity and composition of leachable carbon in the O2 horizon: potential implications for potable

water treatment', *Lake and Reservoir Management*, Vol. 28 No. 4, pp. 282 - 292.

Grayson, R. and Western, A. (2001) 'Terrain and the distribution of soil moisture', *Hydrological Processes*, Vol. 15 No. 13, pp. 2689 - 2690.

- 595 Grayson, R., Kay, P., Foulger, M. and Gledhill, S. (2012) 'A GIS based MCE model for identifying water 596 colour generation potential in UK upland drinking water supply catchments', *Journal of Hydrology*,
- 597 Vol. 420-421, pp. 37 45.
- Gregorich, E., Rochette, P., McGuire, S., Liang, B. and Lessard, R. (1998) 'Soluble organic carbon and
 carbon dioxide fluxes in maize fields receiving spring-applied manure', *Journal of Environmental Quality*, Vol. 27 No. 1, pp. 209 214.
- 601 Hecky, R. and Kilham, P. (1988) 'Nutrient limitation of phytoplankton in freshwater and marine
- 602 environments: a review of recent evidence on the effects of enrichment', *Limnology and*603 *Oceanography*, Vol. 33 No. 4, pp. 796 822.
- Hejzlar, J., Dubrovský, M., Buchtele, J. and Ruzicka, M. (2003) 'The apparent and potential effects of
 climate change on the inferred concentration of dissolved organic matter in a temperate stream (the
 Malše river, south Bohemia)', *Science of the Total Environment*, Vol. 310 No. 1-3, pp. 143 152.
- Hill, A. (1979) 'Denitrification in the nitrogen budget of a river ecosystem', *Nature*, Vol. 281 No.
 5729, pp. 291 292.
- Hindar, A., Wright, R.F., Nilsen, P., Larssen, T. and Høgberget, R. (2003) 'Effects on stream water
 chemistry and forest vitality after whole-catchment application of dolomite to a forest ecosystem in
 southern Norway', *Forest Ecology and Management*, Vol. 180 No. 1-3, pp. 509 525.
- Holden, J., Chapman, P.J., Labadz, J.C. (2004) 'Artificial drainage of peatlands: hydrological and
 hydrochemical process and wetland restoration', *Progress in Physical Geography*, Vol. 28 No. 1, pp.
 95 123.
- Holden, J., Shotbolt, L., Bonn, A., Burt, T.P., Chapman, P.J., Dougill, A.J., Fraser, E.D.G., Hubacek, K.,
 Irvine, B., Kirkby, M.J., Reed, M.S., Prell, C., Stagl, S., Stringer, L.C., Turner, A. and Worrall, F. (2007)
- 617 'Environmental change in moorland landscapes', *Earth-Science Reviews*, Vol. 82 No. 1-2, pp. 75 100.
- Hongve, D. (1999) 'Production of dissolved organic carbon in forested catchments', *Journal of Hydrology*, Vol. 224, pp. 91 99.
- Hope, D., Billett, M.F. and Cresser, M.S. (1994) 'A review of the export of carbon in river water:
 fluxes and processes', *Environmental Pollution*, Vol. 84 No. 3, pp. 301 324.

- Hope, D., Billett, M.F. and Cresser, M.S. (1997) 'Exports of organic carbon in two river systems in NE
 Scotland', *Journal of Hydrology*, Vol. 193 No. 1-4, pp. 61-82.
- Hopkins, D., Ibrahim, D., O'donnell, A. and Shiel, R. (1990) 'Decomposition of cellulose, soil organic
 matter and plant litter in a temperate grassland soil', *Plant and Soil*, Vol. 124 No. 1, pp. 79 85.
- Horton, R.E. (1933) 'The role of infiltration in the hydrologic cycle', *Transactions American Geophysical Union*, Vol. 14, pp. 446 460.
- Howe, L., Blackstock, T., Burrows, C. and Stevens, J. (2005) 'The habitat survey of Wales', *British Wildlife*, Vol. 16 No. 3, pp. 153 162.
- Hurley, J.P., Benoit, J.M., Babiarz, C.L., Shafer, M.M., Andren, A.W., Sullivan, J.R., Hammond, R. and
 Webb, D.A. (1995) 'Influences of Watershed Characteristics on Mercury Levels in Wisconsin Rivers', *Environmental Science and Technology*, Vol. 29 No. 7, pp. 1867 1875.
- Kalbitz, K., Solinger, S., Park, J.H., Michalzik, B. and Matzner, E. (2000) 'Controls on the dynamics of
 dissolved organic matter in soils: a review', *Soil Science*, Vol. 165 No. 4, pp. 277 304.
- Kawasaki, M., Ohte, N. and Katsuyama, M. (2005) 'Biogeochemical and hydrological controls on
 carbon export from a forested catchment in central Japan', *Ecological Research*, Vol. 20 No. 3, pp.
 347 358.
- Lake, I.R., Lovett, A.A., Hiscock, K.M., Betson, M., Foley, A., Sünnenberg, G., Evers, S. and Fletcher, S.
- 639 (2003) 'Evaluating factors influencing groundwater vulnerability to nitrate pollution: developing the
- 640 potential of GIS', *Journal of Environmental Management*, Vol. 68 No. 3, pp. 315 328.
- Levine, D.A. and Jones, W.W. (1990) 'Modeling phosphorus loading to three Indiana reservoirs: a
 geographic information system approach', *Lake and Reservoir Management*, Vol. 6 No. 1, pp. 81 91.
- Li, L., Gao, N., Deng, Y., Yao, J., Zhang, K. (2012) 'Characterization of intracellular & extracellular
- algae organic matters (AOM) of microcystic aeruginosa and formation of AOM-associated
- disinfection byproducts and odor & taste compounds', *Water Research*, Vol. 46 No. 4, pp. 1233 1240.
- Lumsdon, D.G., Stutter, M.I., Cooper, R.J. and Manson, J.R. (2005) 'Model assessment of
 biogeochemical controls on dissolved organic carbon partitioning in an acid organic soil',
- 649 *Environmental Science and Technology*, Vol. 39 No. 20, pp. 8057 8063.
- Marrs, R.H., Le Duc, M.G., Mitchell, R.J., Goddard, D., Paterson, S. and Pakeman, R.J. (2000) 'The
 ecology of bracken: its role in succession and implications for control', *Annals of Botany*, Vol. 85
 (suppl B), pp. 3 15.
- Martin, C., Aquilina, L., Gascuel-Odoux, C., Molénat, J., Faucheux, M. and Ruiz, L. (2004) 'Seasonal
 and interannual variations of nitrate and chloride in stream waters related to spatial and temporal

- patterns of groundwater concentrations in agricultural catchments', *Hydrological Processes*, Vol. 18
 No. 7, pp. 1237 1254.
- McFee, W.W. and Kelly, J.M. (1995) 'Management-induced changes in the actively cycling fractions
 of soil organic matter', *Carbon*, Vol. 1, pp. 119 138.
- Miller, C.J. (2008) *Mechanisms of water colour release from organic soils and consequences for catchment management*. Unpublished PhD thesis, University of Aberdeen, Aberdeen.
- 661 MAFF (Ministry of Agriculture, Fisheries and Food) (2000) *Fertiliser recommendations for agricultural* 662 *and horticultural crops (RB209).*
- Mitchell, G. and McDonald, A. (1992) 'Relationship between different methods of colour
 measurement in potable water', *Transactions of the Institute of Measurement and Control*, pp. 306 309.
- Mitchell, G.N. (1991) 'Water quality issues in the British uplands', *Applied Geography*, Vol. 11 No. 3,
 pp. 201 214.
- 668 Mitsch, W. and Gosselink, J. (2007) *Wetlands*, 4th ed., John Wiley & Sons, New Jersey.
- Monteith, D.T., Stoddard, J.L., Evans, C.D., de Wit, H.A., Forsius, M., Hogasen, T., Wilander, A.,
- 570 Skjelkvale, B.L., Jeffries, D.S., Vuorenmaa, J., Keller, B., Kopacek, J. and Vesely, J. (2007) 'Dissolved
- organic carbon trends resulting from changes in atmospheric deposition chemistry', *Nature*, Vol.
- 672 450, pp. 537 540.
- Neill, M. (1989) 'Nitrate concentrations in river waters in the south-east of Ireland and their
 relationship with agricultural practice', *Water Research*, Vol. 23 No. 11, pp. 1339 1355.
- 675 Nguyen, M.L., Westerhoff, P., Baker, L., Hu, Q., Esparza-Soto, M. and Sommerfeld, M. (2005)
- 676 'Characteristics and reactivity of algae-produced dissolved organic carbon', *Journal of Environmental* 677 *Engineering*, Vol. 131 No. 11, pp. 1574 1582.
- Nykvist, N. (1963) 'Leaching and decomposition of water-soluble organic substances from different
 types of leaf and needle litter', *Studia Forestalia Suecica*, Vol. 3, pp. 1 31.
- 680 Ordóñez, J.A.B., de Jong, B.H.J., García-Oliva, F., Aviña, F.L., Pérez, J.V., Guerrero, G., Martínez, R. and
- Masera, O. (2008) 'Carbon content in vegetation, litter, and soil under 10 different land-use and
- 682 land-cover classes in the central highlands of Michoacan, Mexico', Forest Ecology and Management,
- 683 Vol. 255 No. 7, pp. 2074 2084.
- Osborne, L.L. and Wiley, M.J. (1988) 'Empirical relationships between land use/cover and stream
 water quality in an agricultural watershed', *Journal of Environmental Management*, Vol. 26 No. 1, pp.
 9 27.

- Potthast, K., Hamer, U. and Makeschin, F. (2012) 'Land-use change in a tropical mountain rainforest
 region of southern Ecuador affects soil microorganisms and nutrient cycling', *Biogeochemistry*, Vol.
 111 No. 1-3, pp. 151 167.
- 690 Prévost, M., Rompré, A., Coallier, J., Servais, P., Laurent, P., Clément, B. and Lafrance, P. (1998)
 691 'Suspended bacterial biomass and activity in full-scale drinking water distribution systems: impact of
- 692 water treatment', *Water Research*, Vol. 32 No. 5, pp. 1393 1406.
- Rasmussen, L.H., Kroghsbo, S., Frisvad, J.C. and Hansen, H.C.B. (2003) 'Occurrence of the
 carcinogenic bracken constituent ptaquiloside in fronds, topsoils and organic soil layers in Denmark', *Chemosphere*, Vol. 51 No. 2, pp. 117 127.
- Rasmussen, J.B., Godbout, L. and Schallenberg, M. (1989) 'The humic content of lake water and its
 relationship to watershed and lake morphometry', *Limnology and Oceanography*, Vol. 34 No. 7, pp.
 1336 1343.
- Reynolds, B., Emmett, B.A. and Woods, C. (1992) 'Variations in streamwater nitrate concentrations
 and nitrogen budgets over 10 years in a headwater catchment in mid-Wales', *Journal of Hydrology*,
 Vol. 136 No. 1-4, pp. 155 175.
- Ritson, J.P., Graham, N.J.D., Templeton, M.R., Clark, J.M., Gough, R. and Freeman, C. (2014) 'The
 impact of climate change on the treatability of dissolved organic matter (DOM) in upland water
 supplies: a UK perspective', *Science of the Total Environment*, Vol. 473–474 No. 1, pp. 714 730.
- Rochelle, B.P., Liff, C.I., Campbell, W.G., Cassell, D.L., Robbins Church, M. and Nusz, R.A. (1989)
 'Regional relationships between geomorphic/hydrologic parameters and surface water chemistry
 relative to acidic deposition', *Journal of Hydrology*, Vol. 112 No. 1, pp. 103 120.
- Rothwell, J.J., Evans, M.G., Daniels, S.M. and Allott, T.E.H. (2007) 'Baseflow and stormflow metal
 concentrations in streams draining contaminated peat moorlands in the Peak District National Park
 (UK)', *Journal of Hydrology*, Vol. 341 No. 1-2, pp. 90 104.
- Scott, M.J., Jones, M.N., Woof, C. and Tipping, E. (1998) 'Concentrations and fluxes of dissolved
 organic carbon in drainage water from an upland peat system', *Environment International*, Vol. 24
- 713 No. 5-6, pp. 537 546.
- 714 Skjelkvåle, B.L., Stoddard, J.L., Jeffries, D.S., Tørseth, K., Høgåsen, T., Bowman, J., Mannio, J.,
- 715 Monteith, D.T., Mosello, R., Rogora, M., Rzychon, D., Vesely, J., Wieting, J., Wilander, A. and
- 716 Worsztynowicz, A. (2005) 'Regional scale evidence for improvements in surface water chemistry
- 717 1990–2001', *Environmental Pollution*, Vol. 137 No. 1, pp. 165 176.
- Smith, V.H. (1998) 'Cultural eutrophication of inland, estuarine, and coastal waters' in Pace, M.L. et
 al (Eds.), *Successes, Limitations, and Frontiers in Ecosystem Science*, Springer-Verlag, New York, pp. 7
 49.

- 721 Sobek, S., Tranvik, L.J., Prairie, Y.T., Kortelainen, P. and Cole, J.J. (2007) 'Patterns and regulation of
- dissolved organic carbon: an analysis of 7,500 widely distributed lakes', Limnology and
- 723 Oceanography, Vol. 52 No. 3, pp. 1208 1219.
- Soulsby, C., Tetzlaff, D., Rodgers, P., Dunn, S. and Waldron, S. (2006) 'Runoff processes, stream water
- residence times and controlling landscape characteristics in a mesoscale catchment: an initial
 evaluation', *Journal of Hydrology*, Vol. 325 No. 1-4, pp. 197 221.
- Stevens, P.A., Hornung, M. and Hughes, S. (1989) 'Solute concentrations, fluxes and major nutrient
 cycles in a mature sitka-spruce plantation in Beddgelert forest, north Wales', *Forest Ecology and Management*, Vol. 27 No. 1, pp. 1 20.
- Stevens, J.P., Blackstock, T.H., Howe, E.A. and Stevens, D.P. (2004) 'Repeatability of Phase 1 habitat
 survey', *Journal of Environmental Management*, Vol. 73 No. 1, pp. 53-59.
- Stoddard, J.L., Kahl, J.S., Deviney, F.A., Dewalle, D.R., Driscoll, C.T. and Herlihy, A.T. (2003) *Response*of surface water chemistry to the clean air act amendments of 1990, Tech. Rep. EPA 620/R-03/001.
- 734 Stutter, M.I., Deeks, L.K., Low, D. and Billett, M.F. (2006) 'Impact of soil and groundwater
- heterogeneity on surface water chemistry in an upland catchment', *Journal of Hydrology*, Vol. 318
 No. 1-4, pp. 103 120.
- Thurman, E.M. (Ed.), (1985) Organic Geochemistry of Natural Waters, Kluwer Academic Publishers,
 Lancaster.
- 739 Tipping, E., Woof, C., Rigg, E., Harrison, A.F., Ineson, P., Taylor, K., Benham, D., Poskitt, J., Rowland,
- A.P., Bol, R. and Harkness, D.D. (1999) 'Climatic influences on the leaching of dissolved organic
- 741 matter from upland UK moorland soils, investigated by a field manipulation experiment',
- 742 Environment International, Vol. 25 No. 1, pp. 83 95.
- 743 Toberman, H., Evans, C.D., Freeman, C., Fenner, N., White, M., Emmett, B.A. and Artz, R.R.E. (2008)
- 'Summer drought effects upon soil and litter extracellular phenol oxidase activity and soluble carbon
 release in an upland *calluna* heathland', *Soil Biology and Biochemistry*, Vol. 40 No. 6, pp. 1519 1532.
- Trudgill, S., Burt, T., Heathwaite, A.L. and Arkell, B. (1991) 'Soil nitrate sources and nitrate leaching
 losses, Slapton, south Devon', *Soil Use and Management*, Vol. 7 No. 4, pp. 200 206.
- Urban, N.R., Verry, E.S. and Eisenreich, S.J. (1995) 'Retention and mobility of cations in a small
 peatland trends and mechanisms', *Water, Air, and Soil Pollution*, Vol. 79 No. 1-4, pp. 201 224.
- van den Berg, L.J.L., Shotbolt, L. and Ashmore, M.R. (2012) 'Dissolved organic carbon (DOC)
 concentrations in UK soils and the influence of soil, vegetation type and seasonality', *Science of the*
- 753 *Total Environment*, Vol. 427–428, pp. 269 276.

- Vollenweider, R.A. (1968) *The scientific fundamentals of lake and stream eutrophication, with particular reference to phosphorus and nitrogen as eutrophication factors,* Tech. Rep. DAS/
 DSI/68.27.
- Wallage, Z.E., Holden, J. and McDonald, A.T. (2006) 'Drain blocking: an effective treatment for
 reducing dissolved organic carbon loss and water discolouration in a drained peatland', *Science of the Total Environment*, Vol. 367 No. 2-3, pp. 811 821.
- 760 Watts, C.D., Naden, P.S., Machell, J. and Banks, J. (2001) 'Long term variation in water colour from
- 761 Yorkshire catchments', *Science of the Total Environment*, Vol. 278 No. 1-3, pp. 57 72.
- Weiler, M. and Naef, F. (2003) 'Simulating surface and subsurface initiation of macropore flow', *Journal of Hydrology*, Vol. 273 No. 1, pp. 139 154.
- 764 WHO (World Health Organization). (2011) *Guidelines for drinking-water quality*.
- Worrall, F., Reed, M., Warburton, J. and Burt, T. (2003) 'Carbon budget for a British upland peat
 catchment', *Science of the Total Environment*, Vol. 312 No. 1-3, pp. 133 146.
- Zhang, W., An, S., Xu, Z., Cui, J. and Xu, Q. (2011) 'The impact of vegetation and soil on runoff
 regulation in headwater streams on the east Qinghai–Tibet Plateau, China', *CATENA*, Vol. 87 No. 2,
 pp. 182 189.
- Zsolnay, A. (1996) 'Dissolved humus in soil waters', in Piccolo, A. (Ed.), Humic Substances in
- 771 *Terrestrial Ecosystems*, Elsevier Science, Amsterdam, pp. 171 223.

773 7. Tables and Figures

774

Table 1.	Categorisation	of Phase 1	habitat types.
	0		/ 1

Category used in	Phase 1 Habitat	Category used in	Phase 1 Habitat
present study	classification	present study	classification
Woodland and scrub	A.1	Fen/ mire	E.3
	A.2		
Recently-felled woodland	A.4	Bare peat	E.4
Unimproved grassland	B.1.1	Swamp	F.1
	B.3.1		F.2
Improved grassland	B.1.2	Water	G.1
	B.2.2		G.2
	B.4		
Marsh/ marshy grassland	B.5	Rock/ scree/ quarry	l.1
			1.2
Tall herb and fern	C.1	Arable	J.1.1
	C.2		
	C.3		
Heathland	D.1	Caravan site	J.3.4
	D.2		
	D.3		
	D.5		
	D.6		
Bog	E.1	Buildings	J.3.6
Flush and spring	E.2	Other	J.1.2
			J.4
			Not accessed
			Illegible

775

Table 2. Correlation coefficients (*r*) for statistically significant correlations between percentage Phase
 1 Habitat coverage and reservoir water quality (whole-catchment analysis).

I habitat coverage and reservoir water quality (whole caterinent analysis).								
	Autumn [DOC]	Spring [DOC]	Autumn Colour (Hazen)	Spring Colour (Hazen)	Autumn [NO₃ ⁻]	Spring [NO₃ ⁻]	Autumn pH	Spring pH
Woodland and scrub			· · · · ·			0.743**		
Recently-felled woodland								
Improved grassland					0.636**	0.677**		
Unimproved grassland		-0.512*						
Marsh/ marshy grassland								
Tall herb and fern	-0.499*		-0.588*					
Heathland	-0.543*		-0.543*	-0.649**				
Bog								
Flush and spring								
Fen/ mire	0.564*					0.577*		
Bare peat								
Swamp	0.612*	0.690**	0.624**	0.636**				
Water								
Rock/ scree/ quarry								
Arable	0.734**	0.651**	0.508*	0.549*	0.632**	0.721**		
Caravan site								
Buildings	0.596*					0.580*		
Other	0.548*			0.539*				

778

779

* indicates p < 0.05 and ** indicates p < 0.01. All results shown relate to Spearman's correlation

analysis.

Table 3. Correlation coefficients (*r*) for statistically significant correlations between percentage Phase
 1 Habitat coverage and reservoir water quality (250 m buffer zone analysis).

	age anu	reservon	watert	juanty (Z	50 III DU	bullel zolle allalysis).				
	Autumn	Spring	Autumn	Spring	Autumn	Spring	Autumn	Spring		
	[DOC]	[DOC]	Colour	Colour	[NO3 ⁻]	[NO3 ⁻]	рН	рН		
			(Hazen)	(Hazen)						
Woodland and scrub						0.774**				
Recently-felled woodland										
Improved grassland					0.617*	0.655**				
Unimproved grassland										
Marsh/ marshy grassland						0.502*				
Tall herb and fern			-0.559*							
Heathland	-0.560*	<u>-0.517*</u>	-0.499*	-0.652**			-0.558*			
Bog										
Flush and spring										
Fen/ mire	0.600*	0.513*			0.587*	0.578*				
Bare peat										
Swamp	0.647**	0.709**	0.624**	0.636**						
Water										
Rock/ scree/ quarry										
Arable										
Buildings	0.499*									
Other										

783 784

* indicates p < 0.05 and ** indicates p < 0.01. Underlined result relates to Pearson's correlation analysis and the remainder to Spearman's correlation analysis.

785





Figure 1. Percentage Phase 1 Habitat coverage in 250 m reservoir buffer zone.





Figure 2. Percentage Phase 1 Habitat coverage in whole reservoir catchments.