

A meta-analysis of integrated multi-trophic aquaculture

Kerrigan, Daniel; Suckling, Coleen

Reviews in Aquaculture

DOI:

[10.1111/raq.12186](https://doi.org/10.1111/raq.12186)

Published: 01/08/2018

Peer reviewed version

[Cyswllt i'r cyhoeddiad / Link to publication](#)

Dyfyniad o'r fersiwn a gyhoeddwyd / Citation for published version (APA):

Kerrigan, D., & Suckling, C. (2018). A meta-analysis of integrated multi-trophic aquaculture: Extractive species growth is most successful within close proximity to open-water fish farms. *Reviews in Aquaculture*, 10(3), 560-572. <https://doi.org/10.1111/raq.12186>

Hawliau Cyffredinol / General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

1 **A meta-analysis of integrated multi-trophic aquaculture: Extractive species growth is**
2 **most successful within close proximity to open-water fish farms**

3
4 Daniel Kerrigan; Coleen C. Suckling*

5
6 School of Ocean Sciences, Bangor University, Menai Bridge, Anglesey. LL59 5AB. UK

7
8 * Corresponding author: Coleen C. Suckling; Email: coleen.suckling@bangor.ac.uk and
9 coleen.suckling@cantab.net

10
11 **Running title**

12 Extractive species in open-water IMTA

13
14 **Abstract**

15
16 Fish farming in open water releases dissolved and particulate waste (inorganic and organic)
17 into the surrounding marine environment. To reduce this environmental impact, commercial
18 extractive species can be grown alongside to utilise and reduce this waste, a technique known
19 as integrated multi-tropic aquaculture (IMTA). Information is lacking on whether: 1) IMTA is
20 generally successful with respect to extractive species growth responses; 2) at what spatial
21 scale they can be cultivated from fish cage nutrient sources. Focussing on bivalves and
22 macroalgae as extractive species, this study uses a meta-analysis approach to summarise and
23 conclude peer reviewed data on IMTA to address these information gaps. We show that there
24 are clear benefits to integrating bivalves and macroalgae with fish farms. Bivalves grown
25 within, and relatively near, fish cages (0 m and 1-60 m distance categories respectively)
26 showed significantly higher biomass production relative to controls compared to those grown
27 at larger spatial scales (61+ m). However, biomass production of macroalgae was
28 significantly higher than controls only within close proximity to fish cages (0 m). This
29 information shows increased extractive species production is generally greatest at relatively
30 small spatial scales. It also highlights the need for more site specific information (e.g.
31 seawater parameters, hydrodynamics, food supply, farm capacity) in future studies. The
32 allocation of control sites and locating these at suitable distances (>1 km) from fish farm
33 effluent sources to avoid fish farm nutrient contamination is also recommended.

35 **Keywords:** Bivalves; extractive species; fish farm; integrated multi-trophic aquaculture
36 (IMTA); macroalgae; sustainable aquaculture.

37

38 **1. Introduction**

39

40 Large-scale increases in the intensive mariculture of high-value carnivorous
41 organisms can often result in a number of environmental and sustainability problems (Naylor
42 et al, 2000; Valiela et al, 2001; Naylor et al, 2005; Cabello, 2006; Bergqvist & Gunnarsson,
43 2013). Consequently, this has contributed towards a generally negative public perception of
44 aquaculture, particularly in Western nations (Mazur & Curtis, 2008) which could restrict the
45 potential for future growth in this much needed sector (White et al, 2004). A particular
46 concern is the release of allochthonous nutrients into the surrounding water column from the
47 rearing of carnivorous fish (e.g. Salmonids) in floating sea cages (open-water farming).
48 Nutrients are released as both particulate (organic) waste (uneaten feed pellets and fish
49 faeces) and as dissolved (inorganic) nutrients as a result of nutrient leaching from particulate
50 waste and fish excretory products (Olsen & Olsen, 2008; Wang et al, 2012). Nutrient loading
51 due to fish farming is considerable (Wang et al, 2012) and can negatively impact the benthic
52 environment due to smothering and increased organic enrichment, leading to alterations in
53 sediment chemistry with knock-on effects on benthic biodiversity (Giles, 2008; Olsen &
54 Olsen, 2008; Hargrave, 2010). Many attempts to reduce nutrient loading surrounding fish
55 farms have been made (e.g. improving the digestibility of fish feeds, computerized feed-
56 management systems), however such technological improvements have not yet eliminated the
57 problem of nutrient pollution associated with fish farming (Islam, 2005; Wang et al, 2012).

58 One solution to reducing the environmental impact of fish farming is the use of
59 integrated multi-trophic aquaculture (IMTA). IMTA can be used to potentially recycle these
60 nutrients by cultivating additional commercially relevant organisms. These ‘extractive
61 species’ are able to intercept and assimilate aquaculture derived waste (both organic and
62 inorganic) when cultivated alongside fed fish species (Edwards et al, 1988; Chopin et al,
63 2001; Neori et al, 2004; Troell et al, 2009). This IMTA approach could therefore potentially
64 bio-mitigate the negative environmental impacts of aquaculture whilst simultaneously
65 providing a secondary marketable product for the farmer with possible economical benefit
66 and improved public perception (Chopin et al, 2001; Troell et al, 2003; Ridler et al, 2007). In
67 practice IMTA can take the form of a large variety of systems particularly in Asia (e.g.
68 temporal integration of rice and shrimp or the polyculture of shrimp, fish and crabs in

69 brackish ponds; Troell, 2009) however the majority of Western IMTA operations are land-
70 based, recirculating systems successfully rearing crops of finfish, macroalgae and
71 macroalgivores (e.g. Neori et al, 1996). The majority of general Western aquaculture
72 activities are carried out at sea but at present, there are relatively few commercial examples of
73 open-water IMTA systems (Barrington et al, 2009).

74 A variety of organisms (e.g. echinoderms or crustaceans; Cook & Kelly, 2007;
75 Barrington et al., 2009; Nelson et al, 2012) have been included as part of open-water IMTA
76 trials, however the most commonly cultivated extractive groups are bivalves and macroalgae.
77 In contrast to echinoderms, bivalves and macroalgae are cultivated down current of fish
78 cages, allowing natural water currents to move farm nutrient waste towards these suspended
79 extractive species. The groups can be divided into organic (bivalves) and inorganic
80 (macroalgae) extractive species based on whether the group in question utilises the organic or
81 inorganic nutrients released from fish farms. Suspension feeding bivalves are generalist
82 consumers, able to ingest a variety of particle types and sizes, therefore particulate fish waste
83 could provide an additional food source for bivalves (Jones & Iwama, 1991; Troell et al,
84 2003). Laboratory and field studies utilizing stable isotopes and fatty acids as biomarkers
85 have confirmed that bivalves (*Mytilus edulis*, *Mytilus galloprovincialis*, *Perna viridis*) are
86 able to capture and assimilate fish farm derived organic waste (Lefebvre et al, 2000; Mazzola
87 & Sara, 2001; Gao et al, 2006; Reid et al, 2010; Redmond et al, 2010; MacDonald et al,
88 2011). Similarly, Pacific oysters (*Crassostrea gigas*) have demonstrated high growth rates
89 ($0.7\% \text{ day}^{-1}$) when used as biofilters in land-based IMTA systems (Shpigel, 2005). Although
90 mathematical models have suggested that the capacity of bivalves to assimilate farm derived
91 waste may be limited in an open-water context (Cranford et al, 2013), the high food supply
92 environment surrounding fish farms potentially provides an opportunity for increased bivalve
93 growth (Page & Hubbard, 1987; Brown & Hartwick, 1988).

94 Macroalgal species chosen for inclusion within open-water IMTA operations are
95 typically those with value either as a foodstuff (e.g. *Saccharina latissima*), for industrial
96 applications such as agar extraction (e.g. *Gracilaria spp*), or the cosmetics market (e.g.
97 beauty spas and products). Most are capable of utilising ammonium cations (NH_4^+) which is
98 the primary nitrogen species emitted by fish farms (Hanisak, 1983; Chen et al, 2003;
99 Fernandez-Jover et al, 2007). Nitrogen availability is often a major constraint limiting
100 macroalgal growth, particularly in temperate but also some tropical regions (Lobban &
101 Harrison, 1996; Larned, 1998). Therefore, in areas where macroalgal growth is nitrogen
102 limited (e.g. northern temperate regions) greater availability of inorganic nitrogen found

103 within the vicinity of fish farms could result in increased macroalgal growth rates (Chopin et
104 al, 2001; Neori et al, 2004). Ammonium levels immediately surrounding fish farms have been
105 observed to be below the saturation threshold for macroalgae such as *S. latissima* (Ahn et al,
106 1998) and *Gracilaria vermiculophylla* (Abreu et al, 2011a) suggesting that macroalgae are
107 capable of fully exploiting these available nutrients (Sanderson et al, 2012; Handå et al,
108 2013). Such uptake has been evidenced within controlled land-based IMTA systems, with 72
109 % of nitrogen removed concurrent with increased macroalgal growth (e.g. Neori et al, 2000;
110 Chopin et al, 2001; Matos et al, 2006; Abreu et al, 2011b). Based on this evidence, it can be
111 expected that macroalgal species will show high growth rates in the vicinity of fish farm
112 structures releasing high inorganic nutrient loads.

113 Land-based IMTA systems are mostly closed loop systems thus allowing control of
114 nutrient rich waste (Chopin et al, 2001). In contrast, open-water IMTA lacks this fine control
115 with the dilution of waste occurring by natural seawater movement (e.g. currents). These
116 systems are however generally sheltered within fjordic systems (e.g. Scottish sea lochs) and
117 are likely to have regular current patterns (Navas et al, 2011) leading to the general
118 assumption that the organic and inorganic nutrients will progressively disperse as distance
119 increases from the farm. This increased dilution with distance from farm effect may severely
120 affect the ability of extractive species to intercept nutrient rich waste, thus raising concerns
121 over the effectiveness of using IMTA within an open-water context (Cranford et al, 2013).
122 There is therefore a knowledge gap on the spatial scale at which extractive species can be
123 located in order to assimilate waste and increase profitability.

124 This study will focus only on extractive species growth at different spatial scales to
125 determine whether IMTA can be regarded as worthwhile. Individual trials investigating the
126 cultivation of macroalgae as an extractive species show some indication of positive results
127 whereas those using bivalves as extractive species are less clear. To date this information has
128 not been collated in an informative manner. An overview to determine the effectiveness of
129 open-water IMTA in the context of the production of extractive species is therefore required.
130 Such an overview could help stakeholders determine whether IMTA practices are worth
131 adopting, as well as contributing towards Blue growth (Whitmarsh et al, 2006; DEFRA,
132 2015).

133 This study aims to summarise from the available literature whether open-water IMTA
134 results in extractive species growth augmentation. More specifically, it will focus on the
135 growth of bivalves and macroalgae cultivated in the vicinity of open-water fish farms.
136 Growth responses at increasing distances from the fish cage will also be investigated to help

137 determine the best location to place the extractive species in relation to the farm, information
138 which could be useful for IMTA implementation. We hypothesize that extractive species will
139 show increased growth (relative to controls) when cultured alongside fish farms with growth
140 augmentation declining as distance increases from the closest fish cage.

141

142 **2. Methods**

143

144 *2.1. Data selection*

145

146 A comprehensive search of peer reviewed literature was carried out during early 2015
147 using a keyword search of the Web of Science database. Studies were located using the
148 terms; “bioremediation”, “bivalve”, “growth”, “IMTA”, “integrated aquaculture”,
149 “macroalgae”, “mussel”, “polyculture”, “salmon” and “seaweed” with those studies of direct
150 relevance to the subject matter of this study selected for use. All literature selected compared
151 the growth rate of extractive species (bivalves or macroalgae) cultivated in the vicinity of
152 commercial open-water fish farms with an expressly specified control. Studies which did not
153 include a designated control were excluded from this analysis. All studies were experimental
154 interventions except for Wallace (1980) who measured the growth of naturally occurring
155 fouling mussels on artificial structures both in the vicinity and at a distance from fish cages.
156 Only studies which provided all necessary data (growth parameters, standard deviation
157 values, sample size) were included within this analysis. Data were restricted to the use of
158 cultivated extractive species of potential commercial interest (Bivalves: *Crassostrea gigas*,
159 *Mytilus edulis*, *Mytilus galloprovincialis*, *Mytilus planulatus*, *Ostrea edulis* and *Placopecten*
160 *magellanicus*; Macroalgae: *Gracilaria chilensis*, *Palmaria palmata*, *Sargassum hemiphyllum*,
161 *Sargassum henslowianum*, *Saccharina latissima* and *Ulva spp*). Data were further restricted
162 to those studies which quantified shell length for bivalves and the parameters blade length
163 (cm), biomass production (kg fresh mass m⁻¹) and specific growth rate (SGR; % day⁻¹) for
164 macroalgae. The latter was averaged across the total length of time for each respective study.
165 Data were extracted from graphical figures within the literature using digitizing software
166 (PlotDigitizer; <http://plotdigitizer.sourceforge.net>).

167 Several of the studies included within the meta-analysis contributed more data points
168 than other studies. For example, Navarrete-Mier et al (2010) measured the growth of two
169 extractive species (*O. edulis* and *M. edulis*) at five different distances (0 m, 25 m, 120 m, 300
170 m & 600 m) from the nearest fish farm thereby contributing ten data points to the meta-

171 analysis. In contrast, measuring the growth of one extractive species (*C. gigas*) at one
172 distance (e.g. Jiang et al, 2013) contributed only a single data point towards the meta-
173 analysis. In this study the difference in growth in extractive species between experimental
174 (IMTA) and control sites for each distance was treated as a separate data point, providing the
175 selection criteria described above were met. Although using multiple observations from a
176 single study can decrease the independence of these data points, it was necessary due to the
177 limited number of studies suitable for inclusion (e.g. Kroker et al, 2010).

178

179 2.2. Data analysis

180

181 All studies used in this meta-analysis compared the growth rate of extractive species
182 grown in the vicinity of open-water fish farms with those of a designated control, therefore
183 standardized mean difference was used as the effect size. Effect size is used to quantify the
184 magnitude of difference between two groups with the difference expressed in standard
185 deviation units (Sullivan & Feinn, 2012). A positive value for effect size indicates the
186 experimental group outperformed the control group, a negative effect size indicates
187 underperformance. An effect size of zero indicates no difference between experimental and
188 control groups. For each data point, standardized mean difference was expressed as Cohens
189 d' which was calculated using Formula 1 (Gurevitch et al, 2001; Lakens, 2013) where M^E and
190 M^C represent mean extractive species size at the end of the experimental period for the
191 experimental and control groups respectively. The use of Cohens d' can give a biased
192 estimate of effect size when sample sizes are small (< 20) or differ between experimental and
193 control groups (Hedges & Olkin, 1985). This was encountered within this analysis, therefore
194 an unbiased corrected effect size (Hedges g') was used in this analysis. Hedges g' was
195 calculated from Cohens d' using Formula 2 where n_E & n_C represent sample size for the
196 experimental and control groups respectively (Gurevitch et al, 2001; Lakens, 2013). Variance
197 in effect size (V_D) can be calculated by squaring standard error in effect size (SE_D) which is
198 calculated using Formula 3 (Gurevitch et al, 2001; Lakens, 2013), where d is effect size
199 (Hedges g').

200

201 **Formula 1:** Cohens $d = (M^E - M^C) / SD_{pooled}$

202 **Formula 2:** Hedges $g' = Cohens\ d' \times [1 - (3/4)(n_E + n_C - 9)]$

203 **Formula 3:** $SE_D = \sqrt{[(n_E + n_C) / n_E n_C] + [d^2 / 2(n_E + n_C)]}$

204

205 The studies used in this analysis varied due to interspecific differences between
206 species (e.g. growth rate) and variation in site-specific conditions (e.g. temperature, salinity
207 & chlorophyll-*a* levels). Therefore, a random-effects model was used to calculate the
208 weighted mean effect size. Random-effects models account for two sources of sampling
209 error; within-study variance and between-study variance. Within study variance is given by
210 V_D (see above). Between study variance (r) was calculated by subtracting the degrees of
211 freedom ($n-1$) from total variance and then dividing by a scaling factor, using equations given
212 by Borenstein et al, 2007. Total variance (V_D^*) for each data point was calculated by adding
213 together V_D and r . The reciprocal of V_D^* , w_i was used to determine the weighting each data
214 point carried within the combined effect.

215 For each study a weighted mean effect size (T) was calculated. All data points from
216 each study were combined using Formula 4 (Borenstein et al, 2007) where T_i is effect size
217 (Hedges g'). The standard error of mean effect size (SE_T) was calculated using Formula 5.
218 The significance of weighted mean effect size was assessed by constructing 95%
219 bootstrapped confidence intervals around weighted mean effect size using equations given by
220 Borenstein et al (2007). If 95% confidence intervals do not cross zero, weighted mean effect
221 size can be considered significant (Borenstein et al, 2007). Forest plots were constructed to
222 show the results graphically (weighted mean effect size \pm 95% confidence intervals). Studies
223 which contributed a single data point (e.g. Sara et al, 2009) were presented simply as Hedges
224 g' for graphical representation of effect size. The total weighted mean effect size (T^*) was
225 then calculated by combining all of these weighted data points from all studies for bivalves
226 and macroalgae (Formula's 4 & 5). All calculations were performed on Microsoft Excel 2016
227 (version 16.0) using the framework provided by Neyeloff et al (2012) as a guide.

228

229 **Formula 4:** $T = \Sigma (w_i T_i) / \Sigma w_i$

230 **Formula 5:** $SE_T = \sqrt{(1/w_i)}$

231

232 *2.3 Distance subgroup analysis*

233

234 To determine the effect of distance on the growth of extractive species, each data
235 point was categorized into a subgroup. Bivalve data points were categorized into four
236 distance categories between the bivalves and the nearest stocked fish cage; 1) 0 m, 2) 1-60 m,
237 3) 61-299 m and 4) 300+ m. "0 m" indicates the bivalves were located inside, suspended
238 underneath or "immediately adjacent" to the fish cage. Previous studies have reported that 99

239 % of particles originating from fish farms will settle within 60 m (Coyne et al, 1994; Giles,
240 2008), therefore a distance category of “1-60 m” was also used in this analysis. Not all
241 studies reported an explicit distance between bivalves and fish cages, studies which stated
242 experimental bivalves were located “adjacent” to fish cages were presumed to be between 1
243 and 60 m of fish cages and thus were included within this distance category. Bivalves taken
244 from floats supporting fish farms (Wallace, 1980) were also included within the “1-60 m”
245 category. To maximise categorical balance, the threshold between the 3rd and 4th distance
246 categories was set at an arbitrary value of 300 m.

247 Macroalgae data points were categorized into three distance categories; 1) 0 m, 2) 1-
248 60 m and 3) 61+ m. “0 m” indicates macroalgae were cultivated “within” or “attached” to the
249 fish farm. Major nutrient enhancement is found within 60 m of fish farms (Sanderson et al,
250 2008), therefore, similar to bivalves (described above) the next category was set as “1-60 m”.
251 Not all studies reported an explicit distance between macroalgae and fish cages. Studies
252 which stated that macroalgae were located “adjacent” to fish cages were presumed to be
253 between 1 and 60 m of fish cages and thus were included within this distance category. As
254 only five data points were made at distances exceeding 60 m, to maximise categorical
255 balance the final category was therefore set as “61+ m”. A weighted mean effect size with
256 95% confidence intervals was then constructed for each individual study and subgroup in
257 addition to total weighted mean effect size using the methods described above.

258

259 *2.4. Sensitivity*

260

261 To test the robustness of our findings, a sensitivity analysis was performed using the
262 method employed by Kroeker et al (2010). To summarise, those studies with the largest effect
263 size (regardless of distance) were systematically removed from the meta-analysis, which was
264 then re-run to determine what effect removal had on the meta-analysis outcome. This step
265 was then repeated with effect sizes of decreasing magnitude to determine how many studies
266 needed to be removed to change the significance of the overall result. Similarly, if any study
267 contributed five or more data points to the meta-analysis then this study was removed and the
268 meta-analysis re-ran.

269

270 **3. Results**

271

272 *3.1. Bivalves*

273

274 Twelve studies were found which compared the growth of bivalves cultivated in the
275 vicinity of fish farms with designated controls. From these 12 studies, 43 data points were
276 extracted and incorporated within the meta-analysis. The analysis showed that IMTA had an
277 overall significantly positive effect on the growth rate of bivalves, as indicated by total
278 weighted mean effect size (T^* ; Figure. 1). However, the growth augmentation bivalves
279 experienced in open-water ITMA systems varied according to distance from the closest fish
280 farm. Bivalves within the 0 m and 1-60 m subgroups showed significantly higher growth than
281 controls whereas bivalves grown at further distance points (61 – 299 m and 300+ m) grew at
282 a similar rate to control bivalves (Figure. 2).

283 The stepwise removal of the fifteen largest effect sizes and the removal of the three
284 studies which contributed five or more data points (Jones & Iwama, 1995; Navarrete-Mier et
285 al, 2010; Lander et al, 2012) did not alter the significance of either total weighted mean effect
286 size or weighted mean effect size for each distance category. The sensitivity analysis
287 therefore indicates that the findings of this meta-analysis are robust.

288

289 3.2. *Macroalgae*

290

291 Eight studies were found which compared the growth of macroalgae cultivated in the
292 vicinity of fish farms with designated controls. From these eight studies, 24 data points were
293 extracted and incorporated within the meta-analysis. The analysis showed that IMTA had an
294 overall significantly positive effect on the growth of macroalgae, as indicated by total
295 weighted mean effect size (T^* ; Figure. 3). However, the growth augmentation macroalgae
296 experienced in open-water ITMA systems varied according to distance from the closest fish
297 farm (Figure. 4). Macroalgae within the 0 m subgroup grew significantly faster than controls
298 whereas macroalgae grown further in distance (1-60 m and 61+ m) grew at a similar rate to
299 control macroalgae.

300 A single study (Sanderson et al, 2012) contributed five or more data points to this
301 meta-analysis. The removal of Sanderson et al (2012) did not alter the significance of total
302 weighted mean effect size. However, the removal of the five largest effect sizes from the
303 database used to calculate total weighted mean effect size altered the significance of total
304 weighted mean effect size (significant to non-significant). Stepwise removal of high
305 magnitude data points from the subgroup analysis did not alter the significance of any
306 findings. As the removal of high magnitude data points altered the significance of total

307 weighted mean effect size, the findings for macroalgae are less robust than for bivalves.

308

309 **4. Discussion**

310

311 The growth of extractive species (bivalves and macroalgae) was significantly greater
312 than controls when integrated with open-water fish farms. Macroalgae cultivated within fish
313 farms (0 m) performed significantly better than those at increasing distances. Bivalves
314 cultivated within (0 m) and near (1-60 m) fish farms showed significantly greater growth than
315 those located at further distances (61+ m). Extractive species therefore show best growth
316 performances when located within close proximity to fish farms. Macroalgae generated higher
317 growth rates when integrated within the farm but bivalves showed a larger spatial scale of up to
318 60 m distance for highest growth performances. Overall, these results demonstrate that IMTA
319 is effective with respect to extractive species augmentation which could help farmers generate
320 a profit, particularly if compared to monospecific farms. Although these results lend support to
321 the implementation of open-water IMTA systems, the meta-analyses showed high variation.
322 Such variation can be attributed to many factors which we will now discuss.

323

324 *4.1. Sources of variation*

325

326 *4.1.1. Seasonality and nutrient / food supply*

327

328 The growth enhancement of macroalgae cultivated as part of open-water IMTA
329 systems can be attributed to a fertilisation effect due to increased nutrient levels found within
330 fish farms. Further evidence for the utilisation of farm derived inorganic nutrients is provided
331 by the increased nitrogen content of macroalgae cultivated 10 m from fish cages and by the
332 enrichment of *S. latissima* in a nitrogen isotope (δ^{15}) typical of fish effluent (Troell et al, 1997;
333 Sanderson et al, 2012). Many factors can influence nutrient levels surrounding fish farms
334 including feeding regime, hydrodynamics (Sanderson et al, 2008) and ambient nutrient levels.
335 Macroalgal growth is often limited during the summer due to low ambient nutrient levels
336 (Lobban & Harrison, 1996), it can therefore be expected that IMTA macroalgae will
337 experience the greatest growth enhancement during summer months due to the availability of
338 dissolved nutrients released by nearby fish farms (Chopin et al, 2001; Neori et al, 2004).
339 Studies included within this meta-analysis support this theory with Abreu et al (2009)
340 observing greatest growth rates of *G. chilensis* during the summer. Similarly, Handå et al

341 (2013) found the growth enhancement of IMTA macroalgae to be most pronounced in summer
342 and Wang et al (2014) observed the growth increase of *S. latissima* to occur at a time when
343 dissolved nitrogen levels were at their lowest. In contrast, Halling et al (2005) found no
344 significant difference in macroalgal biomass production between IMTA and control sites.
345 Although ambient nutrient levels were not expressively measured by Halling et al (2005), the
346 findings of this study may have been influenced by seasonality given that the study occurred
347 through the austral winter (when ambient nutrients may not limit macroalgal growth). Troell et
348 al (1997) who cultivated the same species (*G. chilensis*) at the same site but instead during
349 summer months reported a 40% increase in integrated macroalgae production when compared
350 to controls. These examples provide strong evidence that the seasonal timing of IMTA trials
351 may influence the results. It is therefore likely that macroalgae cultivated by Halling et al
352 (2005) may not have received the full benefits of integration due to their selected growth
353 period.

354 Other seasonality related factors may also influence the growth benefit macroalgae
355 experience when integrated with open-water fish farms. Nutrient emissions from open-water
356 fish farms vary, but generally increase during the course of the grow-out cycle (typically two
357 years) peaking (up to four-fold) in late summer of the second year when fish feeding levels are
358 highest (Strain & Hargrave, 2005; Reid et al, 2013a). Commercial macroalgal harvest would
359 have to occur during early to mid-summer as macroalgae begins to degenerate in late summer
360 (Lobban & Harrison, 1996), therefore peak nutrient emissions from fish farms would not be
361 available to IMTA extractive macroalgae. Consequently, for a large period of the year
362 (particularly during the first year of fish rearing) macroalgae would not be exposed to
363 substantially elevated nutrient levels. How macroalgal growth augmentation varies during the
364 typical two year grow-out cycle has yet to be assessed. Such information is required because
365 commercial IMTA ventures which integrate macroalgae will need to account for variations in
366 farm derived nutrient emissions e.g. by scaling back macroalgal cultivation during the first
367 year of fish growth.

368 Food availability for bivalves varies due to a variety of factors (e.g. light, nutrient
369 levels) that undergo regular spatial and temporal fluctuations (Page & Hubbard, 1987; Navarro
370 & Thompson, 1995; Litchman, 1998; Cranford & Hill, 1999). At mid to high latitudes
371 phytoplankton levels are at their seasonal minima throughout the winter, therefore natural
372 populations of bivalves often show minimal growth during this period of food limitation
373 (Malouf & Breese, 1977; Hilbish, 1986). Wallace (1980) suggested that the faster growth rates
374 of *M. edulis* observed at fish farms was likely due to mussels receiving a continuous supply of

375 farm derived waste throughout the winter, thus facilitating year round growth. Similar
376 conclusions were made by Lander et al (2012) who found that the growth advantage gained by
377 *M. edulis* in the vicinity of fish farms was most pronounced in the autumn and winter months.
378 The importance of seasonal timing could explain why some studies found bivalves showed no
379 significant growth enhancement through IMTA. Part of the study carried out by Cheshuk et al
380 (2003) was conducted on an empty farm (no fish present) during the austral winter period (3 ½
381 months from June to September). During this time these mussels (*M. planulatus*) were not
382 exposed to farm derived waste. Chlorophyll-*a* measurements showed that natural food
383 availability was at its lowest during this time, thus leading to the low growth rates reported.

384 When ambient particle concentration, often described as total particulate matter
385 (TPM), exceed a certain threshold (e.g. 5.0 mg TPM l⁻¹ for *M. edulis*) then a significant
386 proportion of ingested particles are not digested but instead are rejected as pseudofaeces
387 (Widdows et al, 1979). Saturation of mussel feeding due to high ambient TPM was suggested
388 by Cheshuk et al (2003) as a possible mechanism for why only modest enhancement of *M.*
389 *planulatus* growth was observed. The use of bivalves as extractive species in open-water IMTA
390 fish farms is dependent on bivalves directly consuming particulate fish waste. The validity of
391 such systems may therefore be compromised if ambient particle concentrations surrounding
392 fish farms were consistently higher than the pseudofaeces threshold. Therefore, bivalve growth
393 enhancement may likely be achieved in IMTA systems located in areas with seasonally or
394 consistently low ambient seston levels (Troell & Norberg, 1998). In oligotrophic waters, farm
395 derived nutrients could also stimulate local phytoplankton production thereby increasing the
396 food supply for secondary consumers e.g. bivalves (Sara et al, 2009). The relationship between
397 bivalve-fish IMTA and variations in local food supply is complex and will require more focus
398 in future IMTA studies. Quantification of the assimilation of farm derived waste at varying
399 particle concentrations would assist in elucidating the relationship between the outcome of
400 open-water IMTA and ambient seston levels.

401

402 4.1.2. Hydrodynamics

403

404 Models indicate that bivalves are best able to capture particulate fish waste when
405 cultivated in areas with slow (< 0.05 m s⁻¹) current speeds (Troell & Norberg, 1998; Cranford
406 et al, 2013). Studies included within this meta-analysis measuring faster current speeds of up to
407 0.11 m s⁻¹ (e.g. Navarette-Mier et al, 2010) found no evidence for bivalve growth
408 augmentation. Particle capture efficiency is dependent on the amount of time available to filter

409 particles from the surrounding water column which is dependent on current speed (fast current
410 speeds equal less time to extract food particles and *vice versa*). IMTA bivalves cultivated in
411 areas where currents do not regularly exceed 0.05 m s^{-1} can therefore be expected to show
412 greater growth enhancement when integrated with open-water fish farms (Troell & Norberg,
413 1998; Cranford et al, 2013).

414 Nutrient dispersal surrounding open-water fish farms is influenced by hydrodynamics,
415 subsurface geographical features and the structure of the fish cage (Sanderson et al, 2008).
416 Understanding dispersal patterns and how they change over time is a complex task requiring
417 extensive field work and advanced modelling (Olsen et al, 2008). An understanding of nutrient
418 emissions from open-water fish farms (also referred to as volumetric loading) would be of
419 importance for commercial IMTA ventures, because it would allow farmers to obtain the
420 maximum growth benefit for their crop through optimum placement of extractive species.
421 Optimum placement is likely to be highly site-specific therefore this meta-analysis cannot
422 provide detailed information on how to organise a commercial IMTA farm besides showing the
423 general distances at which extractive species can be cultivated.

424

425 *4.1.3. Species specific responses*

426

427 Species specific differences in fish faecal properties, extractive species optimal
428 growth, assimilation and feeding mechanisms and patterns will have contributed towards our
429 results. Bivalve growth rates differ intrinsically between species, as shown by comparative
430 studies (Epifanio, 1979; Laing et al, 1987; Cardoso et al, 2006). Intra-specific variation could
431 thus have contributed to the varying growth responses seen in IMTA bivalves as seen by
432 Rensel et al (2011). However, given the small number of studies suitable for inclusion in this
433 meta-analysis, the data were not used to determine the species specific level of contribution, of
434 farmed fish or extractive species, to the effect sizes reported in this study. More open-water
435 IMTA intervention studies on a range of farmed fish and extractive species types would help to
436 determine which species provide the biggest influence on IMTA responses. Despite this
437 variation, increased extractive species growth found during this study demonstrates that
438 increased extractive species production is generally achievable across a range of species.

439

440

441 *4.1.4. Control site selection*

442

443 Macroalgae have demonstrated increased production at distances as great as 800 m
444 from fish farm effluent sources (Abreu et al, 2009). In some cases, control sites have been
445 located within this range and demonstrate pronounced biomass production (Halling et al,
446 2005). It is therefore likely that these sites have been located within the dispersal range of fish
447 farm effluents and therefore could mask potential farm specific responses. To avoid
448 downstream nutrient contamination of controls in future studies, we recommend that control
449 site location considers the hydrodynamics of the area and suitably high distances (e.g. > 1 km
450 (8 km as used by Abreu et al (2009)) from fish cages. Furthermore, the selection criteria for
451 this meta-analysis outlined that literature which did not include a designated control were
452 excluded from the analysis, and as a consequence several studies were not used in the analysis.
453 We therefore recommend the use of well-placed controls in future intervention experiments to
454 help increase the body of evidence around IMTA responses.

455

456 *4.1.5. Site specific information*

457

458 The description of site specific conditions was variable or sometimes absent in the
459 literature (e.g. chlorophyll-*a* concentration, mean current speed). This information is required
460 to understand what effects (if any) site-specific conditions have on the capacity of extractive
461 species to capture and assimilate fish waste. This information is also valuable to farmers by
462 allowing identification of localities where commercial IMTA ventures are most likely to
463 succeed. Based from our experience in this meta-analysis we recommend that future studies
464 consider the following information for study areas: water temperature and salinity, mean and
465 maximum current speeds and their direction, chlorophyll-*a* concentration, particulate organic
466 matter and TPM concentrations. Additionally, details on fish-farm size and feeding protocols
467 would be beneficial towards understanding the effect of distance between fish cages and
468 extractive species.

469

470 *4.2. Meta-analysis limitations*

471

472 One of the limitations with meta-analysis is the reliance on publicly available data
473 which could create a bias on the reported effect sizes. This is a possibility for the data used
474 within this meta-analysis because the majority (80 %) of the macroalgal growth responses
475 included in this meta-analysis demonstrated strong evidence for increased macroalgal growth.
476 Half of the bivalve studies used in this meta-analysis reported positive growth responses

477 relative to controls. It is possible that data has not been made publicly available from studies
478 which were unsuccessful in extractive augmentation within IMTA. Such information will be
479 critical for future meta-analytical summaries of IMTA implementation as more literature is
480 released in this field of research. It is therefore highly recommended that researchers and
481 journals encourage publicising data which demonstrates unsuccessful extractive
482 augmentation within IMTA to prevent possible future bias.

483 The initial literature search identified 14 studies which cultivated macroalgae in the
484 vicinity of open-water fish farms. However, six of these studies were excluded from
485 subsequent analysis due to the lack of an expressly specified control or a lack of reporting of
486 data (e.g. sample size) required for the meta-analysis. Therefore, only eight macroalgal
487 studies (containing 24 data points) were included within this analysis, compared to 12 bivalve
488 studies (containing 43 data points). Given that significant bivalve growth enhancement was
489 found within the 1-60 m distance category despite the generally rapid settling velocity of
490 particulate fish waste (Law et al, 2014), the lack of significant macroalgal growth
491 enhancement at distances greater than 0 m could be deemed surprising. The paucity of
492 suitable macroalgal studies could be a potential causative factor behind the lack of significant
493 macroalgal growth enhancement at distances greater than 0 m. This therefore emphasises the
494 recommendation for future IMTA studies to include suitable controls within experimental
495 designs as it is only with reference to controls that the presence (or lack of) growth
496 enhancement can be determined.

497

498 *4.3. Logistics of extractive species in IMTA*

499

500 As the dilution of particulate fish waste increases with distance from fish cages
501 (Doglioli et al, 2004), bivalves cultivated within fish cages themselves (0 m) will be exposed to
502 higher concentrations of particulate waste (and thus increased food availability) than those at
503 greater distances. Therefore, greater growth augmentation for bivalves and macroalgae within
504 the 0 m subgroup is predictable. However, significantly increased bivalve growth was also
505 observed within a larger spatial scale, in this case up to 60 m. Although significant macroalgal
506 growth augmentation was only found for the 0 m subgroup, individual studies have found
507 significantly increased macroalgal production at distances of up to 800 m from fish farms
508 (Abreu et al, 2009). Such findings are encouraging as farmers are unlikely to adopt IMTA
509 practices if the installation of extractive species interferes with the day to day operations of the
510 fish farm (the major cash crop) e.g. restricting access to fish cages or impeding water flow

511 (thus reducing oxygen supply to fish). Therefore, extractive species must be appropriately
512 located (e.g. not inside a fish cage). The results of this meta-analysis indicate that spatial
513 constraints may not represent an impediment to widespread open-water IMTA. To maximise
514 farm waste recapture (as well as biomass production), bivalves should be cultivated close to
515 fish cages due to the rapid settling velocity of particulate fish waste (Law et al, 2014).
516 However, care should be taken in locating the macroalgal component of open-water IMTA
517 farms as excess particulate fish waste could potentially settle on macroalgal fronds thus
518 blocking light and restricting growth.

519 The evidence presented in this study shows that by adopting IMTA practices,
520 economic advantages could be gained by farmers (though increased production of extractive
521 species). By providing secondary marketable crops, IMTA farms exhibit a greater degree of
522 economic diversification compared to monoculture operations. Diversification represents a
523 form of insurance, as a marketable product will still be produced in the event of disease
524 outbreaks or infrastructure damage (e.g. net failure). Profitable markets presently exist for the
525 sale of bivalves (Lucas & Southgate, 2012) and the farmed seaweed market is likely to grow in
526 Western nations given the increasing popularity of seaweed consumption (Brownlee et al,
527 2012). Much of the infrastructure and equipment for IMTA (e.g. rope lines, boats, buoys) will
528 already be present on fish farms, therefore the start-up costs of IMTA in farms is likely to be
529 low (Lander et al, 2012). However, labour costs can be expected to increase on IMTA farms as
530 extra work hours will be required for the maintenance and harvesting of extractive species
531 (Holdt & Edwards, 2014). A model based on a Canadian farm estimates that net present value
532 (NPV) is increased by 24% when mussels and seaweed are grown alongside Atlantic salmon
533 (Ridler et al, 2007) however more transparent models containing greater detail (e.g.
534 proportions of extractive species) would be of use to farmers in determining the economies of
535 integration. Surveys have shown a positive attitude towards IMTA amongst the general public
536 indicating that consumer acceptance will not be a barrier to IMTA expansion, with 50% of
537 participants willing to pay 10% extra for IMTA labelled products (Barrington et al, 2010).
538 Therefore, a system of eco-labelling may allow IMTA farmers to charge a higher price for their
539 products and thus keep IMTA farms profitable in the face of falling fish prices (Whitmarsh et
540 al, 2006) or competition with larger firms (Ridler et al, 2007). The profitability of IMTA farms
541 may also be improved if coastal management systems legally oblige operations to pay for the
542 environmental cost of their activities via discharge taxes (“user pays” concept; Troell et al,
543 2003).

544 However, before IMTA becomes more widely implemented there are a number of

545 mitigation and biosecurity issues regarding commercial IMTA that need to be satisfactorily
546 resolved. While bivalve integration has shown generally positive growth responses in this
547 meta-analysis, the net organic loading from bivalves (released as faeces) combined with the
548 fish farm may still have a negative impact on the underlying benthos. It has been
549 recommended by the Fisheries and Oceans Canada (DFO) Science Advisory Schedule (DFO,
550 2013) to use extractive deposit feeding species (e.g. sea urchins, sea cucumbers and
551 polychaetes) located underneath the suspended bivalve extractive species to consume these
552 heavy organic solids (Cranford et al., 2013; DFO, 2013; Reid et al., 2013b). The
553 implementation of adding another trophic level into IMTA will require structural
554 considerations relating to the fish farm (e.g. oxygen supply via seawater flow and efficient
555 connection between trophic levels; DFO, 2013).

556 Previous work has found that IMTA bivalves grown in water of sufficient depth are
557 highly unlikely to act as reservoirs for fish pathogens such as infectious salmon anaemia virus
558 (ISAV) or *Vibrio anguillarum* and may assist in the control of drug-resistant pathogens and
559 parasites such as the sea louse *Lepeoptheirus salmonis* (Mortensen, 1993; Skar & Mortensen,
560 2007; Molloy et al, 2011; Pietrak et al, 2012; Molloy et al, 2014). Haya et al (2004) similarly
561 found that extractive species (*M. edulis* and *S. latissima*) cultured in the vicinity of a *S. salar*
562 farm did not accumulate hazardous therapeutants or contaminants (e.g. heavy metals) above
563 background levels. To facilitate the spread of open-water IMTA further work regarding
564 bioaccumulation within extractive species and potential disease transfer within farms requires
565 consideration to dispel any concerns farmers or regulatory bodies may have regarding IMTA.
566 If commercial IMTA is to become widespread, legislation governing aquaculture operations
567 may have to be reformed so that policy recommendations (e.g. minimum distances between
568 mussel and fish farms) do not act as barriers to commercial IMTA (Alexander, 2015).

569

570 4.4. Conclusions and future research considerations

571

572 This study demonstrates that; 1) extractive species cultivated in the vicinity of open-
573 water fish farms experience a growth benefit due to integration and 2) close proximity of
574 extractive species to the farm (0 m for macroalgae and 0-60 m for bivalves) can increase
575 performance and therefore possibly profit but there appears to be some spatial flexibility
576 around this if logistical constraints require it. Even though the extent at which nutrient
577 extraction is carried out is still quantitatively unknown, spatially extensive locations of
578 extractive species are known to significantly reduce organic loading around fish cages (Reid et

579 al, 2013a; Holdt & Edwards, 2014).

580 Future study recommendations include: 1) allocating control sites and locating these
581 at suitable distances (>1 km to 8km) from fish farm effluent sources to avoid fish farm nutrient
582 contamination; 2) including site details such as seawater parameters (e.g. temperature,
583 salinity), hydrodynamics (current speeds and direction), food supply (chlorophyll-*a*, particulate
584 organic matter and total particle matter concentrations), farm capacity (farm size and feeding
585 protocols); and 3) determining the extent to which spatially extensive extractive species
586 cultivation mitigates nutrient discharge from open-water fish farms (including consideration of
587 the organic loading from the bivalve component of IMTA farms).

588 Open-water IMTA is still in development and further research can be expected to
589 improve IMTA methodologies (e.g. optimum placings for extractive species and a wider range
590 of commercial extractive species) leading towards more sustainable IMTA systems. Although
591 complete (100%) nutrient sequestration is not practically feasible, future IMTA efforts should
592 be encouraged given the environmental and economic merits of integration.

593

594 **Acknowledgements**

595 This study was generated as part of the third year dissertation module through the host
596 University's BSc Marine Biology degree programme.

597

598 **References**

599 Abreu, M., Varela, D., Henriquez, L., Villarroel, A., Yarish, C., Sousa-Pinto, I. & Buschmann,
600 A. 2009. Traditional vs integrated multi-trophic aquaculture of *Gracilaria chilensis* C.J. Bird,
601 J.McLachlan & E.C. Olivera: Productivity and physiological performance. *Aquaculture*, **293**:
602 211-220.

603

604 Abreu, M., Pereira, R., Yarish, C., Buschmann, A. & Sousa-Pinto, I. 2011a. IMTA with
605 *Gracilaria vermiculophylla*: Productivity and nutrient removal performance of the macroalgae
606 in a land-based pilot scale system. *Aquaculture*, **312**: 77-87.

607

608 Abreu, M., Pereira, R., Buschmann, A., Sousa-Pinto, I. & Yarish, C. 2011b. Nitrogen uptake
609 responses of *Gracilaria vermiculophylla* (Ohmi) Papenfuss under combined and single
610 addition of nitrate and ammonium. *Journal of Experimental Marine Biology and Ecology*. **407**:
611 190-199.

612

613 Ahn, O., Petrell, R. & Harrison, P. 1998. Ammonium and nitrate uptake by *Laminaria*
614 *saccharina* and *Nereocystis luetkeana* originating from a salmon sea cage farm. *Journal of*
615 *Applied Phycology*, **10**: 333-340.

616

617 Alexander, K., Potts, T., Freeman, S., Israel, D., Johansen, J., Kletou, D., Meland, M.,
618 Pecorino, D., Rebours, C., Shorten, M. & Angel, D. 2015. The implications of aquaculture
619 policy and regulation for the development of integrated multi-trophic aquaculture in Europe.
620 *Aquaculture*, **443**: 16-23.

621

622 Barrington, K., Chopin, T. & Robinson, S. 2009. Integrated multi-trophic aquaculture (IMTA)
623 in marine temperate waters. In: D. Soto (ed.). *Integrated mariculture: a global review*. FAO
624 Fisheries and Aquaculture Technical Paper. No. 529. Rome, FAO. pp. 7–46

625

626 Barrington, K., Ridler, N., Chopin, T., Robinson, S. & Robinson, B. 2010. Social aspects of the
627 sustainability of integrated multi-trophic aquaculture. *Aquaculture international*, **18**: 201-211.

628

629 Bergqvist, J. & Gunnarsson, S. 2013. Finfish aquaculture: Animal welfare, the environment
630 and ethical implications. *Journal of Agricultural and Environmental Ethics*, **26**: 75-99.

631

632 Borenstein, M., Hedges, L. & Rothstein, H. 2007. Meta-analysis: Fixed effects vs. random
633 effects. Retrieved from [https://www.meta-analysis.com/downloads/Meta-](https://www.meta-analysis.com/downloads/Meta-analysis%20fixed%20effect%20vs%20random%20effects.pdf)
634 [analysis%20fixed%20effect%20vs%20random%20effects.pdf](https://www.meta-analysis.com/downloads/Meta-analysis%20fixed%20effect%20vs%20random%20effects.pdf) (30/05/2016).

635

636 Brown, J. & Hartwick, B. 1988. Influences of temperature, salinity and available food upon
637 suspended culture of the pacific oyster *Crassostrea gigas* I. Absolute and allometric growth.
638 *Aquaculture*, **70**: 231-251.

639

640 Brownlee, I., Fairclough, A., Hall, A. & Paxman, J. 2012. The potential health benefits of
641 macroalgae and macroalgae extract. In: *Macroalgae: ecology, nutrient composition and*
642 *medicinal uses. Marine Biology: Earth Sciences in the 21st Century*. Nova Science Publishers,
643 Hauppauge, New York, 119-136.

644

645 Buschmann, A., Varela, D., Hernandez-Gonzalez, M. & Huovinen, P. 2008. Opportunities and

646 challenges for the development of an integrated macroalgae-based aquaculture activity in
647 Chile: determining the physiological capabilities of *Macrocystis* and *Gracilaria* as biofilters.
648 *Journal of Applied Phycology*, **20**: 571-577.

649

650 Cabello, F. 2006. Heavy use of prophylactic antibiotics in aquaculture: A growing problem for
651 human and animal health and for the environment. *Environmental Microbiology*, **8**: 1137-1144.
652

653 Cardoso, J., Witte, J., & van der Veer, H. 2006. Intra- and interspecies comparison of energy
654 flow in bivalve species in Dutch coastal waters by means of the Dynamic Energy Budget
655 (DEB) theory. *Journal of Sea Research*, **56**: 182-197.

656

657 Chen, Y., Beveridge, M., Telfer, T. & Roy, W. 2003. Nutrient leaching and settling rate
658 characteristics of the faeces of Atlantic salmon (*Salmo salar*) and the implications for
659 modelling of solid waste dispersion. *Journal of Applied Ichthyology*, **19**: 114-117.

660

661 Cheshuk, B., Purser, G. & Quintana, R. 2003. Integrated open-water mussel (*Mytilus*
662 *planulatus*) and Atlantic Salmon (*Salmo salar*) culture in Tasmania, Australia. *Aquaculture*,
663 **218**: 357-378.

664

665 Chopin, T., Buschmann, A., Halling, C., Troell, M., Kautsky, N., Neori, A., Kraemer, G.,
666 Zertuche-Gonzalez, J., Yarish, C. & Neefus, C. 2001. Integrating macroalgae into marine
667 aquaculture systems: A key towards sustainability. *Journal of Phycology*, **37**: 975-986.
668

669

670 Cook, E. & Kelly, M. 2007. Enhanced production of the sea urchin *Paracentrotus lividus* in
671 integrated open-water cultivation with Atlantic salmon *Salmo salar*. *Aquaculture*, **273**: 573-
672 585.

673

674 Coyne, R., Hiney, M., O'Connor, B., Kerry, J., Cazabon, D. & Smith, P. 1994. Concentration
675 and persistence of oxytetracycline in sediments under a marine salmon farm. *Aquaculture*, **123**:
676 31-42.

677

678 Cranford, P. & Hill, P. 1999. Seasonal variation in food utilization by the suspension-feeding
679 bivalve molluscs *Mytilus edulis* and *Placopecten magellanicus*. *Marine Ecology Progress*

680 *Series*, **190**: 223-239.

681

682 Cranford, P., Reid, G. & Robinson, S. 2013. Open water integrated multi-trophic aquaculture:
683 Constraints on the effectiveness of mussels as an organic extractive component. *Aquaculture*
684 *Environmental Interactions*, **4**: 163-173.

685

686 DEFRA. 2015. United Kingdom multi-annual national plan for the development of sustainable
687 aquaculture. Department for Environment Food and Rural Affairs. 39 pp.

688

689 DFO. 2013. Review of the organic extractive component of integrated multi-trophic
690 aquaculture (IMTA) in Southwest New Brunswick with emphasis on the blue mussel. Fisheries
691 and Oceans Canada (DFO) Science Advisory Schedule. Report number 2013/056.

692

693 Doglioli, A., Magaldi, M., Vezzulli, L. & Tucci, S. 2004. Development of a numerical model to
694 study the dispersion of wastes coming from a marine fish farm in the Ligurian Sea (Western
695 Mediterranean). *Aquaculture*, **231**: 215-235.

696

697 Edwards, P., Pullin, R. & Gartner, J. 1988. Research and education for the development of
698 integrated crop-livestock-fish farming systems in the tropics. ICLARM Studies and Reviews,
699 vol 16. International Centre for Living Aquatic Resources Management, Manila: 53pp.

700

701 Epifanio, C. 1979. Growth in bivalve molluscs: Nutritional effects of two or more species of
702 algae in diets fed to the American oyster *Crassostrea virginica* (Gmelin) and the hard clam
703 *Mercenaria mercenaria* (L.). *Aquaculture*, **18**: 1-12.

704

705 Fernandez-Jover, D., Sanchez-Jerez, P., Bayle-Sempere, J., Carratala, A. & Leon, V. 2007.
706 Addition of dissolved nitrogen and dissolved organic carbon from wild fish faeces and food
707 around Mediterranean fish farms: Implications for waste-dispersal models. *Journal of*
708 *Experimental Marine Biology and Ecology*, **340**: 160-168.

709

710 Gao, Q., Shin, P., Lin, G., Chen, S. & Cheung, S. 2006. Stable isotope and fatty acid evidence
711 for uptake of organic waste by green lipped mussels *Perna viridis* in a polyculture fish farm
712 system. *Marine Ecology Progress Series*, **317**: 273-283.

713

714 Giles, H. 2008. Using Bayesian networks to examine consistent trends in fish farm benthic
715 impact studies. *Aquaculture*, **274**: 181-195.
716

717 Gurevitch, J., Curtis, P. & Jones, M. 2001. Meta-analysis in Ecology. *Advances in Ecological*
718 *Research*, 32: 200-247.
719

720 Halling, C., Aroca, G., Cifuentes, M., Buschmann, A. & Troell, M. 2005. Comparison of spore
721 inoculated and vegetative propagated cultivation methods of *Gracilaria chilensis* in an
722 integrated macroalgae and fish cage culture. *Aquaculture international*, **13**: 409-422.
723

724 Handå, A., Forbord, S., Wang, X., Broch, O., Dahle, S., Storseth, T., Reitan, K., Olsen, Y. &
725 Skjermo, J. 2013. Seasonal- and depth- dependent growth of cultivated kelp (*Saccharina*
726 *latissima*) in close proximity to salmon aquaculture (*Salmo salar*) in Norway. *Aquaculture*,
727 **414**: 191-201.
728

729 Hanisak, M. 1983. The nitrogen relationships of marine macroalgae. In Carpenter, E. &
730 Capone, D (eds). *Nitrogen in the marine environment*. New York Academic Press: 699-730.
731

732 Hargrave, B. 2010. Empirical relationships describing benthic impacts of Salmon aquaculture.
733 *Aquaculture Environment Interactions*, **1**: 33-46.
734

735 Haya, K., Sephton, D., Martin, J. & Chopin, T. 2004. Monitoring of therapeutants and
736 phycotoxins in Kelps and Mussels co-cultured with Atlantic salmon in an integrated multi-
737 trophic aquaculture system. *Bulletin of the Aquaculture Association of Canada*, **104**: 29-34.
738

739 Hedges, L. & Olkin, I. 1985. *Statistical methods for meta-analysis*. San Diego, CA: Academic
740 Press.
741

742 Hilbish, T. 1986. Growth trajectories of shell and soft tissue in bivalves: Seasonal variation in
743 *Mytilus edulis* L. *Journal of Experimental Marine Biology and Ecology*, **96**: 103-113.
744

745 Holdt, S. & Edwards, M. 2014. Cost-effective IMTA: A comparison of the production
746 efficiencies of mussels and macroalgae. *Journal of Applied Phycology*, **26**: 33-945.
747

748 Islam, S. 2005. Nitrogen and phosphorus budget in coastal and marine cage aquaculture and
749 impacts of effluent loading on ecosystem: review and analysis towards model development.
750 *Marine Pollution Bulletin*, **40**: 48-61.
751

752 Jiang, Z., Wang, G., Fang, J. & Mao, Y. 2013. Growth and food sources of Pacific oyster
753 *Crassostrea gigas* integrated culture with Sea bass *Lateolabrax japonicus* in Ailian Bay, China.
754 *Aquaculture International*, **21**: 45-52.
755

756 Jones, T. & Iwama, G. 1991. Polyculture of the Pacific oyster, *Crassostrea gigas* (Thunberg),
757 with Chinook salmon, *Oncorhynchus tshawytscha*. *Aquaculture*, **92**: 313-322.
758

759 Kroeker, K., Kordas, R., Crim, R. & Singh, G. 2010. Meta-analysis reveals negative yet
760 variable effects of ocean acidification on marine organisms. *Ecology Letters*, **13**: 1419-1434.
761

762 Lakens, D. 2013. Calculating and reporting effect sizes to facilitate cumulative science: a
763 practical primer for t-tests and ANOVAs. *Frontiers in Psychology*, **4**: 863.
764

765 Laing, I., Utting, S. & Kilada, R. 1987. Interactive effect of diet and temperature on the growth
766 of juvenile clams. *Journal of Experimental Marine Biology and Ecology*, **113**: 23-38.
767

768 Lander, T., Robinson, S., MacDonald, B. & Martin, J. 2012. Enhanced growth rates and
769 condition index of blue mussels (*Mytilus edulis*) held at integrated multi-trophic aquaculture
770 sites in the Bay of Fundy. *Journal of Shellfish Research*, **31**: 997-1007.
771

772 Larned, S. 1998. Nitrogen versus Phosphorus – limited growth and sources of nutrients for
773 coral reef macroalgae. *Marine Biology*, **132**: 409-421.
774

775 Law, B., Hill, P., Maier, I., Milligan, T. & Page, F. 2014. Size, settling velocity and density of
776 small suspended particles at an active salmon aquaculture site. *Aquaculture Environment*
777 *Interactions*, **6**: 29-42.
778

779 Lefebvre, S., Barille, L. & Clerc, M. 2000. Pacific oyster (*Crassostrea gigas*) feeding
780 responses to fish farm-effluent. *Aquaculture*, **187**: 185-198.
781

782 Litchman, E. 1998. Population and community responses of phytoplankton to fluctuating light.
783 *Oecologica*, **117**: 247-257.
784

785 Lobban, C. & Harrison, P. 1996. Macroalgae ecology and physiology. Cambridge University
786 Press: 376pp.
787

788 Lucas, J. & Southgate, P. 2012. Aquaculture: Farming aquatic animals and plants. Wiley-
789 Blackwell: 648pp.
790

791 MacDonald, B., Robinson, S. & Barrington, K. 2011. Feeding activity of mussels (*Mytilus*
792 *edulis*) held in the field at an integrated multi-trophic aquaculture (IMTA) site (*Salmo salar*)
793 and exposed to fish food in the laboratory. *Aquaculture*, **314**: 244-251.
794

795 Malouf, R. & Breese, W. 1977. Seasonal changes in the effects of temperature and water flow
796 rate on the growth of juvenile pacific oysters, *Crassostrea gigas* (Thunberg). *Aquaculture*, **12**:
797 1-13.
798

799 Matos, J., Costa, S., Rodrigues, A., Pereira, R. & Sousa-Pinto, I. 2006. Experimental integrated
800 aquaculture of fish and red macroalgae in Northern Portugal. *Aquaculture*, **252**: 31-42.
801

802 Mazur, N. & Curtis, A. 2008. Understanding community perceptions of aquaculture: lessons
803 from Australia. *Aquaculture International*, **16**: 601-621.
804

805 Mazzola, A. & Sara, G. 2001. The effect of fish farming organic waste on food availability for
806 bivalve molluscs (Gaeta Gulf, Central Tyrrhenian, MED): stable carbon isotope analysis.
807 *Aquaculture*, **192**: 361-379.
808

809 Molloy, S., Pietrak, M., Bouchard, D. & Bricknell, I. 2011. Ingestion of *Lepeophtheirus*
810 *salmonis* by the blue mussel *Mytilus edulis*. *Aquaculture*, **311**: 61-64.
811

812 Molloy, S., Pietrak, M., Bouchard, D. & Bricknell, I. 2014. The interaction of infectious
813 salmon anaemia virus (ISAV) with the blue mussel, *Mytilus edulis*. *Aquaculture Research*, **45**:
814 509-518.
815

816 Mortensen, S. 1993. Passage of infectious pancreatic necrosis virus (IPNV) through
817 invertebrates in an aquatic food chain. *Diseases of Aquatic Organisms*, **16**: 41-45.
818

819 Navarette-Mier, F., Sanz-Lazaro, C. & Marin, A. 2010. Does bivalve mollusc polyculture
820 reduce marine fin fish farming environmental impact? *Aquaculture*, **306**: 101-107.
821

822 Navarro, J. & Thompson, R. 1995. Seasonal fluctuations in the size spectra, biochemical
823 composition and nutritive value of the seston available to a suspension-feeding bivalve in a
824 subarctic environment. *Marine Ecology Progress Series*, **125**: 95-106.
825

826 Navas, J., Telfer, T. & Ross, L. 2011. Application of 3D hydrodynamic and particle tracking
827 models for better environmental management of finfish culture. *Continental Shelf Research*,
828 **31**: 675-684.
829

830 Naylor, R., Goldburg, R., Primavera, J., Kautsky, N., Beveridge, M., Clay, J., Folke, C.,
831 Lubchenco, J., Mooney, H. & Troell, M. 2000. Effect of aquaculture on world fish supplies.
832 *Nature*, **405**: 1017-1024.
833

834 Naylor, R., Hindar, K., Fleming, I., Goldburg, R., Williams, S., Volpe, J., Whoriskey, F., Eagle,
835 J., Kelso, D. & Mangel, M. 2005. Fugitive salmon: Assessing the risks of escaped fish from
836 net-pen aquaculture. *Bioscience*, **55**: 427-437.
837

838 Nelson, E., MacDonald, B. & Robinson, S. 2012. The absorption efficiency of the suspension-
839 feeding sea cucumber, *Cucumaria frondosa*, and its potential as an extractive integrated multi-
840 trophic aquaculture (IMTA) species. *Aquaculture*, **370-71**: 19-25.
841

842 Neori, A., Krom, M., Ellner, S., Boyd, C., Popper, D., Rabinovitch, R., Davison, P., Dvir, O.,
843 Zuber, D., Ucko, M., Angel, D. & Gordin, H. 1996. Macroalgae biofilters as regulators of
844 water quality in integrated fish-macroalgae culture units. *Aquaculture*, **141**: 183-199.
845

846 Neori, A., Shpigel, M. & Ben-Ezra, D. 2000. A sustainable integrated system for the culture of
847 fish, macroalgae and abalone. *Aquaculture*, **186**: 279-291.
848

849 Neori, A., Chopin, T., Troell, M., Bushcman, A., Kraemer, G., Halling, C., Shpigel, M. &

850 Yarish, C. 2004. Integrated aquaculture: rationale, evolution and state of the art emphasizing
851 macroalgae biofiltration in modern mariculture. *Aquaculture*, **231**: 361-391.

852

853 Neyeloff, J., Fuchs, S. & Moreira, L. 2012. Meta-analyses and Forest plots using a Microsoft
854 excel spreadsheet: step-by-step guide focusing on descriptive data analysis. *BMC Research*
855 *Notes* **5**:52.

856

857 Olsen, L., Holmer, M. & Olsen, Y. 2008. Perspective of nutrient emission from fish
858 aquaculture in coastal waters: Literature review with evaluated states of knowledge. *The*
859 *Fishery and Aquaculture Industry Fund*, FHF Project no. 542014, 87pp.

860

861 Olsen, Y. & Olsen, L. 2008. Environmental impact of aquaculture on coastal planktonic
862 ecosystems. In: Tsukamoto, K., Kawamura, T., Takeuchi, T., Beard, T. & Kaiser, M. (eds).
863 *Fisheries for Global Welfare and Environment, Proceedings of the 5th World Fisheries*
864 *Congress 2008*, Terrapub, Tokyo: 181-196.

865

866 Page, H. & Hubbard, D. 1987. Temporal and spatial patterns of growth in mussels (*Mytilus*
867 *edulis*) on an offshore platform: relationships to water temperature and food availability.
868 *Journal of Experimental Marine Biology and Ecology*, **111**: 159-179.

869

870 Peharda, M., Zupan, I., Bavcevic, L., Frankic, A. & Klanjscek, T. 2007. Growth and condition
871 index of mussel *Mytilus galloprovincialis* in experimental integrated aquaculture. *Aquaculture*
872 *Research*, **38**: 1714-1720.

873

874 Pietrak, M., Molloy, S., Bouchard, D., Singer, J. & Bricknell, I. 2012. Potential role of *Mytilus*
875 *edulis* in modulating the infectious pressure of *Vibrio anguillarum* 02 β on an integrated multi-
876 trophic aquaculture farm. *Aquaculture*, **326-329**: 36-39.

877

878 Redmond, K., Magnasen, T., Hansen, P., Strand, O. & Meier, S. 2010. Stable isotopes and fatty
879 acids as tracers of the assimilation of salmon fish feed in blue mussels (*Mytilus edulis*).
880 *Aquaculture*, **298**: 202-210.

881

882 Reid, G., Liutkus, M., Bennet, A., Robinson, S., MacDonald, B. & Page, F. 2010. Absorption
883 efficiency of blue mussels (*Mytilus edulis* and *M. trossulus*) feeding on Atlantic salmon (*Salmo*

884 *salar*) feed and fecal particulates: Implications for integrated multi-trophic aquaculture.
885 *Aquaculture*, **209**: 165-169.

886

887 Reid, G., Chopin, T., Robinson, S., Azevedo, P., Quinton, M. & Belyea, E. 2013a. Weight
888 ratios of the kelps, *Alaria esculenta* and *Saccharina latissima*, required to sequester dissolved
889 inorganic nutrients and supply oxygen for Atlantic salmon, *Salmo salar*, in integrated multi-
890 trophic aquaculture systems. *Aquaculture*, **408-409**: 34-46.

891

892 Reid, G., Robinson, S., Chopin, T. & MacDonald, B. 2013b. Dietary proportion of fish culture
893 solids required by shellfish to reduce the net organic load in open-water integrated multi-
894 trophic aquaculture: A scoping exercise with cocultured Atlantic Salmon (*Salmo salar*) and
895 Blue Mussel (*Mytilus edulis*). *Journal of Shellfish Research*, **32**: 509-517.

896

897 Rensel, J., Bright, K. & Siegrist, Z. 2011. Integrated fish-shellfish mariculture in Puget sound.
898 Final Report. NOA award – NA08OAR4170860.

899

900 Ridler, N., Wowchuk, M., Robinson, B., Barrington, K., Chopin, T., Robinson, S., Page, F.,
901 Reid, G., Szemerda, M., Sewuster, J. & Boyne-Travis, S. 2007. Integrated multi-trophic
902 aquaculture (IMTA): A potential strategic choice for farmers. *Aquaculture Economics &*
903 *Management*, **11**: 99-110.

904

905 Sanderson, J., Cromey, C., Dring, M. & Kelly, M. 2008. Distribution of nutrients for
906 macroalgae cultivation around salmon cages at farm sites in north-west Scotland. *Aquaculture*,
907 **278**: 60-68.

908

909 Sanderson, J., Dring, M., Davidson, K. & Kelly, M. 2012. Culture, yield and bioremediation
910 potential of *Palmaria palmata* (Linnaeus) Weber & Mohr and *Saccharina latissima* (Linnaeus)
911 C.E. Lane, C. Mayers, Druehl & G.W. Saunders adjacent to fish farm cages in northwest
912 Scotland. *Aquaculture*, **354-355**: 128-135.

913

914 Sara, G., Zenone, A. & Tomasello, A. 2009. Growth of *Mytilus galloprovincialis* (mollusca,
915 bivalvia) close to fish farms: a case of integrated multi-trophic aquaculture within the
916 Tyrrhenian Sea. *Hydrobiologia*, **636**: 129-136.

917

918 Shpigel, M. 2005. The use of bivalves as biofilters and valuable product in land based
919 aquaculture systems – review. In Dame, R. & Olenin, S. (Eds.), *The Comparative roles of*
920 *suspension-feeders in ecosystems*. Kluwer Academic Pub, Dordrecht, The Netherlands: 400pp.
921

922 Skar, C. & Mortensen, S. 2007. Fate of infectious salmon anaemia virus (ISAV) in
923 experimentally challenged blue mussels *Mytilus edulis*. *Diseases of Aquatic Organisms*, **74**: 1-
924 6.
925

926 Stirling, H. & Okumus, I. 1995. Growth and production of mussels (*Mytilus edulis* L.)
927 suspended at salmon cages and shellfish farms in two Scottish sea lochs. *Aquaculture*, **134**:
928 193-210.
929

930 Strain, P. & Hargrave, B. 2005. Salmon aquaculture, nutrient fluxes and ecosystem processes
931 in Southwestern New Brunswick. *Environmental Effects of Marine Finfish Aquaculture*, **5**: 29-
932 57.
933

934 Sullivan, G. & Feinn, R. 2012. Using effect size – or why the *P* value is not enough. *Journal of*
935 *Graduate Medical Education*, **4**: 279-282.
936

937 Troell, M., Halling, C., Nilsson, A., Buschmann, A., Kautsky, N. & Kautsky, L. 1997.
938 Integrated marine cultivation of *Gracilaria chilensis* (Gracilariales, Rhodophyta) and salmon
939 cages for reduced environmental impact and increased economic output. *Aquaculture*, **156**: 45-
940 61.
941

942 Troell, M. & Norberg, J. 1998. Modelling output and retention of suspended solids in an
943 integrated salmon-mussel culture. *Ecological Modelling*, **110**: 65-77.
944

945 Troell, M., Halling, C., Neori, A., Chopin, T., Buschmann, A., Kautsky, N. & Yarish, C. 2003.
946 Integrated mariculture: asking the right questions. *Aquaculture*, **226**: 69-90.
947

948 Troell, M. 2009. Integrated marine and brackish water aquaculture in tropical regions:
949 research, implementation and prospects. In D. Soto (ed.). *Integrated mariculture: a global*
950 *review. FAO Fisheries and Aquaculture Technical Paper*. No. 529. Rome, FAO. pp 47-131.
951

952 Troell, M., Joyce, A., Chopin, T., Neori, A., Buschmann, A. & Fang, J. 2009. Ecological
953 engineering in aquaculture – Potential for integrated multi-trophic aquaculture in marine
954 offshore systems. *Aquaculture*, **297**: 1-9.
955

956 Valiela, I., Bowen, J. & York, J. 2001. Mangrove forests: One of the worlds threatened major
957 tropical environments. *Bioscience*, **51**: 807-815.
958

959 Wallace, J. 1980. Growth rates of different populations of the edible mussel, *Mytilus edulis*, in
960 North Norway. *Aquaculture*, **19**: 303-311.
961

962 Wang, X., Olsen, L., Reitan, K. & Olsen, Y. 2012. Discharge of nutrient wastes from salmon
963 farms: environmental effects, and potential for integrated multi-trophic aquaculture.
964 *Aquaculture Environment Interactions*, **2**: 267-283.
965


966 Wang, X., Broch, O., Forbord, S., Handå, A., Skjermo, J., Reitan, K., Vadstein, O. & Olsen, Y.
967 2014. Assimilation of inorganic nutrients from salmon (*Salmo salar*) farming by the
968 macroalgae (*Saccharina latissima*) in an exposed coastal environment: implications for
969 integrated multi-trophic aquaculture. *Journal of Applied Phycology*, **26**: 1869-1878.
970

971 White, K., O'Neill, B. & Tzankova, Z. 2004. At a crossroads: Will aquaculture fulfil the
972 promise of a blue revolution? Report. Seaweb Aquaculture Clearinghouse Report, Washington
973 DC: 17pp.
974

975 Whitmarsh, D., Cook, E. & Black, K. 2006. Searching for sustainability in aquaculture: An
976 investigation into the economic prospects of an integrated salmon-mussel production system.
977 *Marine Policy*, **30**: 293-298.
978


979 Widdows, J., Fieth, P. & Worrall, C. 1979. Relationships between seston, available food and
980 feeding activity in the Common Mussel, *Mytilus edulis*. *Marine Biology*, **50**: 195-207.
981
982
983
984
985

986 **Figure legends**



987
 988
 989
 990
 991
 992
 993
 994
 995
 996
 997
 998
 999
 1000
 1001
 1002
 1003
 1004
 1005

Figure 1. Bivalve growth relative to controls (weighted mean effect size \pm 95% confidence intervals) when integrated with open-water fish farms presented for individual studies and for all studies collated (total weighted mean effect size (T*)). Boxed numbers represent the number of data points used to calculate weighted mean effect sizes.



1006

1007

Figure 2. Weighted mean effect size (\pm 95% confidence intervals) for bivalves cultivated at varying distances from open-water fish farms. Boxed numbers represent the number of data points used to calculate weighted mean effect sizes.

1009

1010

1011

1012

1013

1014

1015

1016

1017

1018

1019

1020


1021

1022

1023

1024

1025



1026

1027

1028

1029

1030

1031

1032

1033

1034

1035

1036

1037

1038

1039

1040

1041


1042

1043

1044

1045

Figure 3. Macroalgal growth relative to controls (weighted mean effect size \pm 95% confidence intervals) when integrated with open-water fish farms presented for individual studies and for all studies collated (total weighted mean effect size (T*)). Boxed numbers represent the number of data points used to calculate weighted mean effect sizes.



1046

1047

1048

1049

Figure 4. Weighted mean effect size (\pm 95% confidence intervals) for macroalgae cultivated at varying distances from open-water fish farms. Boxed numbers represent the number of data points used to calculate weighted mean effect sizes.