

Evaluation of Molecular Methods for the Detection and Quantification of **Pathogen-Derived Nucleic Acids in Sediment**

Farkas, Kata; Hassard, Francis; McDonald, James; Malham, Shelagh; Jones, David

Frontiers in Microbiology

DOI: 10.3389/fmicb.2017.00053

Published: 24/01/2017

Publisher's PDF, also known as Version of record

Cyswllt i'r cyhoeddiad / Link to publication

Dyfyniad o'r fersiwn a gyhoeddwyd / Citation for published version (APA): Farkas, K., Hassard, F., McDonald, J., Malham, S., & Jones, D. (2017). Evaluation of Molecular Methods for the Detection and Quantification of Pathogen-Derived Nucleic Acids in Sediment. Frontiers in Microbiology, 6, 1-12. Article 53. https://doi.org/10.3389/fmicb.2017.00053

Hawliau Cyffredinol / General rights Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

• Users may download and print one copy of any publication from the public portal for the purpose of private study or research.

- You may not further distribute the material or use it for any profit-making activity or commercial gain
 You may freely distribute the URL identifying the publication in the public portal ?

Take down policy If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.





Evaluation of Molecular Methods for the Detection and Quantification of Pathogen-Derived Nucleic Acids in Sediment

Kata Farkas^{1*}, Francis Hassard², James E. McDonald³, Shelagh K. Malham² and Davey L. Jones¹

¹ School of Environment, Natural Resources and Geography, Bangor University, Bangor, Wales, ² School of Ocean Sciences, Bangor University, Bangor, Wales, ³ School of Biological Sciences, Bangor University, Bangor, Wales

The accurate detection of pathogens in environmental matrices, such as sediment, is critical in understanding pathogen fate and behavior in the environment. In this study, we assessed the usefulness of methods for the detection and quantification of Vibrio spp. and norovirus (NoV) nucleic acids in sediment. For bacteria, a commonly used direct method using hexadecyltrimethylammonium bromide (CTAB) and phenol-chloroform-isoamyl alcohol (PCI) extraction was optimized, whereas for NoV, direct and indirect (virus elution-concentration) methods were evaluated. For quantification, commercially available quantitative PCR (qPCR) and reverse transcription qPCR (RT-qPCR) kits were tested alongside a digital PCR (dPCR) approach. CTAB-based extraction combined with 16 h polyethylene glycol 6000 (PEG6000) precipitation was found to be suitable for the direct extraction of high abundance bacterial and viral nucleic acids. For the indirect extraction of viral RNA, beef extract-based elution followed by PEG6000 precipitation and extraction using the NucliSENS® MiniMag® Nucleic Acid Purification System and the PowerViral® Environmental RNA/DNA Isolation Kit and gRT-PCR resulted in 83-112 and 63-69% recoveries of NoV, respectively. dPCR resulted in lower viral recoveries (47 and 9%) and ~ 4 orders of magnitude lower Vibrio concentrations (3.6-4.6 log₁₀ gc/100 g sediment) than was observed using qPCR. The use of internal controls during viral quantification revealed that the RT step was more affected by inhibitors than the amplification. The methods described here are suitable for the enumeration of viral and/or bacterial pathogens in sediment, however the use of internal controls to assess efficiency is recommended.

Keywords: dPCR, norovirus, Vibrio, nucleic acid extraction, pathogen detection, qRT-PCR, sediment

INTRODUCTION

Pathogenic bacteria and viruses, found in environmental water due to wastewater discharge, agricultural activities, and run-off have been shown to associate with waterborne outbreaks of human disease (Radin, 2014). Pathogens in water readily adsorb to both inorganic and organic matter (Jin and Flury, 2002) resulting in the accumulation of pathogens in sediment (Staley et al., 2012). Hence, viruses and bacteria are often found in surface sediment in significantly higher

OPEN ACCESS

Edited by:

Petr Kralik, Veterinary Research Institute, Czechia

Reviewed by:

Christine Paillard, Centre National de la Recherche Scientifique (CNRS), France Chris Whitehouse, United States Food and Drug Administration (FDA), USA

> *Correspondence: Kata Farkas fkata211@gmail.com

Specialty section:

This article was submitted to Food Microbiology, a section of the journal Frontiers in Microbiology

Received: 07 October 2016 Accepted: 09 January 2017 Published: 24 January 2017

Citation:

Farkas K, Hassard F, McDonald JE, Malham SK and Jones DL (2017) Evaluation of Molecular Methods for the Detection and Quantification of Pathogen-Derived Nucleic Acids in Sediment. Front. Microbiol. 8:53. doi: 10.3389/fmicb.2017.00053 concentrations than in the overlying water column (Rao et al., 1986; Duhamel and Jacquet, 2006; Vignaroli et al., 2013; Perkins et al., 2014). Furthermore, the association of viral and bacterial particles with sediment particles has been shown to increase the persistence of those pathogens (Gerba and McLeod, 1976; Smith et al., 1978; LaBelle and Gerba, 1980; Davies et al., 1995; Anderson et al., 2005). Pathogens may be released from sediment to water as a result of physical disturbance or variations in physico-chemical properties of water due to weather changes (An et al., 2002; Haramoto et al., 2009) and can be ingested by crustacea and shellfish destined for human consumption (Landry et al., 1983; Oliveira et al., 2011; Lowther et al., 2012). These events can result in public health threats far from the source of contamination.

In order to assess the potential health risks related to the contamination of the sediment, the development of reliable methods for the detection and accurate quantification of pathogens is essential. Bacterial contamination of effluent discharges and bathing water quality is traditionally measured by culture-dependent microbiological plating (International Organization for Standardization, ISO 9308-1:2014, 2014). Even though bacterial culturing is a simple and inexpensive approach, recent studies suggest that culture-dependent methods underrepresent the bacterial abundance, as viable but non-culturable (VBNC) bacteria cannot be reliably enumerated in this way (Pinto et al., 2015). Pathogenic viruses have also been studied in environmental matrices using cell or tissue culturing, however those methods are time-consuming, require experienced staff and the results often underestimate viral titer due to the aggregation of viral particles (Charles et al., 2009; Farkas et al., 2015). Furthermore, many pathogenic viruses (e.g., human noroviruses) cannot be propagated in vitro.

Molecular approaches such as quantitative (qPCR) and reverse transcription qPCR (RT-qPCR) are often used for the rapid and accurate enumeration of pathogen-derived nucleic acids in environmental samples (Dick and Field, 2004; Bae and Wuertz, 2009; Furet et al., 2009; Girones et al., 2010) including sediment (Green and Lewis, 1999; Miura et al., 2011; Kim et al., 2014; Staggemeier et al., 2015a). PCR-based methods have shown the most promising results in terms of viral or bacterial recovery, however some chemical and biological compounds (e.g., nucleases, humic substances) within the sediment may inhibit the enzymes of RT and/or PCR resulting in false negative results (Meschke and Sobsey, 1998; Rock et al., 2010). However, progressive development of commercially available qPCR and qRT-PCR kits has resulted in procedures more resistant to inhibition.

Digital PCR (dPCR) using water-oil emulsion droplets or chip-based technology have been frequently used in diagnostics and virus detection (Sedlak and Jerome, 2013; Ding and Mu, 2016). Recently the usefulness of dPCR-based quantification for water and fruit samples has also been investigated (Ishii et al., 2014; Coudray-Meunier et al., 2015; Fraisse et al., 2017). Results have suggested that dPCR allowed a more accurate quantification of viral nucleic acids than q(RT-)PCR and was not adversely affected by high concentrations of humic acid (Hoshino and Inagaki, 2012) and other inhibitors associated with environmental samples (Fraisse et al., 2017). Although molecular approaches are unable to address the infectivity of the target pathogen, these methods have been widely used, especially for those viruses that cannot be cultured *in vitro*.

Approaches commonly used for the extraction of pathogens and pathogen-derived nucleic acids from sediments have shown high variability in efficiency (Miura et al., 2011), particularly among different types of sediment (Johnson et al., 1984). Comparison between studies is often challenging due to the various extraction and quantification approaches used (Rames et al., 2016). To facilitate detailed studies on viruses and bacteria, further validation of the protocols currently used for quantification is necessary. The aim of this study was to assess the efficiency of different extraction and PCR-based quantification methods for the recovery of bacteria (Vibrio spp.) and NoV commonly found in coastal and marine environments. Bacterial DNA was extracted directly from sediment. For virus recovery, direct and traditional, indirect elution-concentration methods were used. Vibrio spp. and NoV nucleic acids were then quantified using q(RT-)PCR and dPCR-based approaches.

MATERIALS AND METHODS

Environmental Sediment Samples

For the bacterial enumeration experiments, sediment samples from eight different sites (Site 1: $53^{\circ}44'43.50''$ N, $2^{\circ}49'14.30''$ W, Site 2: $53^{\circ}44'36.80''$ N, $2^{\circ}49'49.10''$ W, Site 3: $53^{\circ}44'7.48''$ N, $2^{\circ}51'43.29''$ W, Site4: $53^{\circ}44'3.70''$ N, $2^{\circ}52'39.10''$ W, Site 5: $53^{\circ}43'58.50''$ N, $2^{\circ}53'20.90''$ W, Site 6: $53^{\circ}43'55.50''$ N, $2^{\circ}54'6.10''$ W, Site 7: $53^{\circ}43'43.90''$ N, $2^{\circ}58'14.00''$ W, Site 8: $53^{\circ}43'54.40''$ N, $2^{\circ}58'29.60''$ W) were collected from the River Ribble and estuary, North England, UK at high tide (**Figure 1**). The top 10 cm of sediment was collected by a mechanically operated Van Veen grab, aseptically placed in a 50 mL polypropylene centrifuge tube and stored at 4° C. Subsequently, small aliquots (\sim 1 g subsamples in triplicates) from each replicate grab were frozen at -80° C pending nucleic acid extraction.

For the NoV experiments, sediment samples were collected in the Conwy estuary (53°16′N 3°49′W), North Wales, UK during receding tide. The top 1 cm layer of sediment was collected manually in sterile plastic bottles and stored at 4°C. Preliminary findings suggested that the sediment samples contained no NoV, thus those were subject to virus spiking. Prior to seeding with viruses, sediment was aliquoted (5, 2, and 0.25 g/tube) and autoclaved in order to inactivate bacterial enzymes which could interfere with the spiking studies.

Norovirus Seeding

Human norovirus (kindly provided by Prof. Ian Goodfellow, University of Cambridge, UK) was isolated from an anonymized clinical sample, collected as part of an ethically approved study at the University of Cambridge lead by Dr. Lydia Drumwright. The viral sample was generated by the preparation of a 10% solution using phosphate-buffered saline, pH 7.4 which was subsequently filtered through a $0.2 \,\mu$ m filter. An RNase treatment was performed on $100 \,\mu$ L 10-time diluted sample as described in Topping et al. (2009). The test revealed no



significant concentration loss compared to a non-treated sample, suggesting that the sample contains predominantly intact NoV particles. NoV was added to sterilized sediment to achieve a final concentration of $\sim 2 \times 10^5$ genome copies (gc)/g sediment. Experiments were set up in triplicates and with one negative control (no virus added). Samples were incubated at room temperature on an orbital shaker at 90 rpm for 30 min to allow the attachment of viral particles to sediment.

Extraction of Bacterial DNA

Bacterial genomic DNA (gDNA) was extracted from 0.5 g sediment in triplicates using the direct extraction method based on the method of Griffiths et al. (2000). In brief, 0.5 mL glass beads, 0.5 mL of hexadecyltrimethylammonium bromide (CTAB) extraction buffer and 0.5 mL of phenol-chloroformisoamyl alcohol (25:24:1; pH 8.0; PCI) were added to samples which were then lysed at 5.5 m/s for 30 s. Samples were centrifuged at $14,000 \times g$ for 5 min and the top aqueous phase (containing nucleic acids) was transferred to a new tube. An equal volume of chloroform-isoamyl alcohol (24:1) (CI) was added, followed by centrifugation at 14,000 \times g for 5 min. Nucleic acids were precipitated from aqueous layer by the addition of 2 volumes of 30% (wt/vol) polyethylene glycol 6000 (PEG6000)-1.6 M NaCl. The mixture was incubated at room temperature for 1, 2, or 16 h at 4° C and centrifuged at 16,000 \times g for 10 min. The pellet was washed with 0.2 mL ice cold 70% ethanol and air dried prior to elution in 50 µL molecular-grade water with no subsequent pre-treatments. The concentration and quality of gDNA was checked using a Qubit fluorimeter 2.0 (Invitrogen, UK) and a Nanodrop ND-1000 (Nanodrop, USA), respectively. Results suggested high dsDNA concentrations in all samples, hence all samples were analyzed using qPCR as described below.

Direct Extraction of NoV RNA

The spiked sediment samples (0.25 g each) were mixed with 0.25 mL PBS. Viral nucleic acids were extracted directly for sediment using the CTAB-method described above. For PEG precipitation, samples were incubated at room temperature for 2 or 16 h at 4° C.

The suitability of commercial kits was also evaluated. Nucleic acids from 2 g NoV-spiked sediment were extracted using the PowerSoil[®] Total RNA Isolation Kit (MO BIO Laboratories, USA) according to manufacturer's protocol and eluted in 50 μ L molecular-grade water. The PowerViral[®] Environmental DNA/RNA Isolation Kit (MO BIO Laboratories, USA) aiming to coextract viral RNA and DNA was also used. For that purpose, the spiked samples (0.25 g each) were mixed with 0.25 mL PBS prior to extraction. Beta-mercaptoethanol and PCI was added to the lysis buffer as advised by the manufacturer. Viral RNA was eluted in 50 μ L molecular-grade water. The characteristics of each approach are summarized in Table 1.

Indirect Extraction of NoV RNA

NoV particles were eluted and concentrated as described in Lewis and Metcalf (1988). In brief, 5 g NoV-spiked sediment samples were mixed with 15 mL 3% beef extract in 2 M NaNO₃ (pH 5.5) for 30 min and the solid matter was removed by centrifugation at 2500 \times g, 10 min. The pH of the eluent was adjusted to 7.5 and incubated in 15% PEG6000 and 2% NaCl overnight at 4° C and centrifuged at 2500 \times g for 80 min. The resulting pellet was subject to nucleic acid extraction, using the CTABbased extraction (detailed above) and two commercial kits, the NucliSENS[®] MiniMag[®] Nucleic Acid Purification System (bioMérieux SA, France), and the PowerViral® Environmental RNA/DNA Isolation Kit (MO BIO Laboratories, USA). Nucleic acids were extracted according to manufacturer's protocol and eluted in 50 µL molecular biology grade water. Prior to extraction using the MiniMag[®] System, samples were incubated in 10 mg/mL proteinase K solution at 37°C for 60 min. No PCI was used when samples were extracted using the PowerViral[®] Kit. The characteristics of each approach are summarized in Table 1.

Quantitative PCR and RT-PCR

All qPCR assays were carried out in a QuantStudioTM Flex 6 Real-Time PCR System (Applied Biosystems, USA). For quantification, dilution series of a plasmid DNA carrying the target sequence were used (Primerdesign Ltd, UK). For all samples the original and a 10-times (vol:vol) diluted extract were tested. A positive NoV RNA control—extracted from fecal matter using the PowerViral Kit and 1000× diluted in molecular-grade

Extraction	Company	Cost per sample	Extraction type	Sample size	Processing time	Final volume	Additional equipment	Harmful chemicals
		-						
CTAB-based extraction	N/A	£1	Physical, PEG	0.25 g	45 min + 12–16 h incubation	25–100 μL	Beadbeater	PCI, CI
PowerSoil [®] Total RNA Isolation Kit	MoBio	£8.60	Physical, Column	2 g	3 h	50–100 μL	None	PCI
PowerViral Environmental DNA/RNA Isolation Kit	MoBio	£5	Column	0.2 mL	45 min	50–100 μL	None	BME*
NucliSENS [®] MiniMag [®] Nucleic Acid Purification System	Bio Mérieux SA	£2.80	Magnetic beads	0.5 mL	45 min	50–100 μL	MiniMag system*	None
*Optional.								

TABLE 1 | Characteristics of the extraction methods used for the direct and indirect extraction of viral nucleic acids.

TABLE 2 | Characteristics of the quantification kits used in this study.

	Kit	Company	Cost per sample	RT	Polymerase	Duration
One-step	RNA Ultrasense [®]	Invitrogen	£5.80	Superscript [®] III	Platinum [®] Taq	3 h 40 min
	Oasig	Primer design	£2	N/A	N/A	2 h
Two-step	Superscript [®] IV RT	Invitrogen	£5.30	Superscript [®] IV	-	20 min
	KAPA Force Probe qPCR	KAPA Biosystems	£1.20	-	KAPA3G	2 h
	QuantStudio TM 3D Digital PCR	Applied Biosystems	£8	-	N/A	2 h 45 min

water—was also added to qRT-PCR reactions. The characteristics of each approach are summarized in **Table 2**.

For detection of bacterial (*Vibrio* spp.) DNA by qPCR, two commercially available TaqMan qPCR mixes were tested, namely the Oasig qPCR Master Mix (Primerdesign Ltd, UK) and the KAPA Force Probe qPCR mix (KAPA Biosystems, USA). The 20 μ L reaction mixes contained 1× qPCR mix, 1 μ g bovine serum albumin (BSA), 1 μ L Vibrio spp. Primer/Probe mix (Primerdesign Ltd, UK), and 4 μ L sample or standard. Using the Oasig qPCR Master Mix, the initial denaturation was 2 min at 95°C followed by 50 cycles of amplification consisting of 95°C for 15 s and 60°C for 60 s. Using the KAPA Force Probe qPCR mix, the 5 min denaturation at 98°C was followed by 45 cycles of amplification consisting of 95°C for 30 s.

For detection of NoV RNA, three RT-qPCR approaches were tested. A single-step TaqMan-based qRT-PCR assay was used according to the method described previously by Flannery et al. (2014) using RNA Ultrasense One-step qRT-PCR kit (Invitrogen, USA). The 20 μ L qPCR reaction mix contained 1× RNA Ultrasense Reaction Mix with 1 μ L RNA Ultrasense Enzyme Mix, 10 pmol of the forward (ATG TTC AGR TGG ATG AGR TTC TCW GA), 20 pmol of the reverse (TCG ACG CCA TCT TCA TTC ACA) primers, 5 pmol of the probe (FAM-AGC ACG TGG GAG GGC GAT CG-TAMRA), 0.1× ROX reference dye, 1 μ g BSA and 4 μ L of the sample/plasmid DNA. Following a 60 min step of reverse transcription at 55°C and a 5 min step of denaturation at 95°C, the 45 cycles of amplification consisted of 95°C for 15 s, 60°C for 60 s, and 65°C for 60 s.

A single-step TaqMan-based qRT-PCR assay using the same NoV standards, primers, and probe as described above with the Oasig OneStep qRT-PCR Master Mix (Primerdesign Ltd., UK) was also used, according to the manufacturer's instructions. The 20 μ L reaction mix contained 4 μ L sample/standard and 1 μ g BSA. The reverse transcription was performed at 42°C for 10 min followed by a 2 min denaturation at 95°C and 50 cycles of amplification consisting of 95°C for 15 s and 60°C for 60 s.

The usefulness of two-step RT-qPCR for NoV RNA quantification was also investigated. For RT the Superscript IV (Invitrogen, USA) was used with oligo $(dT)_{18}$ primers according to the manufacturer's instructions. The 20 μ L reaction mix contained 4 μ L RNA extract. Following RT, the cDNA was quantified by qPCR using the KAPA Force Probe qPCR mix (KAPA Biosystems, USA) and the standards, primers and probe for NoV as detailed above. The 20 μ L reaction mix contained 4 μ L sample/standard and 1 μ g BSA. The 5 min denaturation at 98°C was followed by 45 cycles of amplification consisting of 95°C for 15 s, 60°C for 60 s, and 65°C for 30 s.

Nucleic Acid Quantification Using dPCR

Digital PCR was performed using the QuantStudioTM 3D Digital PCR System using sealed chip technology (Applied Biosystems, USA). The 14.5 μ L PCR mixture contained 7.5 μ L QuantStudioTM 3D Digital PCR Master Mix v2, 0.8 μ L of Norovirus genogroup 2 or *Vibrio* spp. primer/probe mix (Primerdesign Ltd, UK) and 5 μ L sample. The reaction mixture was loaded onto a QuantStudioTM 3D Digital PCR 20 K Chip using an automatic chip loader. Amplification was carried



out using the following thermal cycling conditions: $96^{\circ}C$ for 10 min, then 39 cycles of $60^{\circ}C$ for 2 min and $96^{\circ}C$ for 30 s followed by $60^{\circ}C$ for 2 min. After amplification the chips were allowed to cool to room temperature and then read on the QuantStudioTM 3D Instrument. The results were further analyzed using the QuantStudio 3D AnalysisSuiteTM Cloud Software.

Internal Control for NoV Quantification

The level of inhibition for each extract type was estimated using plasmid DNA incorporating the target sequence of the NoV quantification assays described above. The plasmid DNA was added to each negative RNA extract to reach a concentration of ${\sim}10^3$ copies/ μL .

Statistical Analysis

Paired *t*-tests were used to analyze the qPCR results, the data was found to have no significant outliers between the two related groups. Next a Shapiro-Wilk test of normality revealed approximately normal distributions between each paired observation. Finally, a Welch's *t*-test was undertaken on SPSS version 22 (IBM, USA).

RESULTS AND DISCUSSION

Comparison of PEG Incubation Times on Quantification of Bacterial DNA from Sediment

The original protocol for direct extraction of bacterial nucleic acids from soil samples suggested the use of a 2 h incubation time to enable DNA binding to PEG (Griffiths et al., 2000) resulting in sufficient DNA precipitation. However, many other studies use longer PEG incubation times (Krsek and Wellington, 1999; Roose-Amsaleg et al., 2001). In order to investigate whether incubation times have an effect on bacterial DNA extraction efficiency, we incubated the samples in PEG solution for 1, 2, and 16 h. The results suggested that increased incubation times resulted in higher DNA concentrations when qPCR quantification was used (Figure 2). Between 1 and 16 h incubation time, the median Vibrio spp. abundance increased by 1.3 and 1.2 log₁₀ gc/100 g sediment for the Oasig and KAPA qPCR method, respectively. In contrast, dPCR had a 0.7 and $0.1 \log_{10} \text{gc}/100 \text{g}$ sediment reduction in Vibrio spp. abundance between 1-2 and 2-16h of PEG incubation. This implies that prolonged PEG incubation resulted in inhibited quantification by dPCR. It has been suggested that PCR inhibitors can be concentrated along with nucleic acids during PEG precipitation (LaMontagne et al., 2002; Miura et al., 2011). Although, dPCR has been shown to be resistant to humic acid, a common inhibitor in environmental samples (Hoshino and Inagaki, 2012), our findings suggest that these sediment samples may have contained other organic substances that also interfere with dPCR amplification.

Comparison of Different Methods for Quantifying Bacteria in Sediment

Vibrio spp. quantification was significantly greater in all PEG incubation times with KAPA mastermix (range 7.9-9.2 log10 gc/100 g sediment) compared to Oasig mastermix $(0-6.56 \log_{10})$ gc/100 g sediment) and dPCR (3.6-4.6 log₁₀ gc/100 g sediment; Figure 2). Overall the highest concentrations were achieved for the samples incubated in PEG solution overnight using the KAPA mastermix for quantification resulting in 1.7×10^9 gc/100 g sediment on average. Dilution, 10× (vol:vol) of DNA extracts did not improve the recovery suggesting limited inhibition using this mastermix. Significantly lower DNA concentrations or negative results were observed using the Oasig qPCR mastermix. Recovery of Vibrio spp. targets increased for the 1 and 2 h PEG treatments on diluted samples, suggesting inhibition of the mastermix. The observed average Vibrio spp. abundance increased by 0.5 log₁₀ gc/100 g sediment for the 1 h PEG treatment (p < 0.05; average fold increase 1×10^6 gc/100 g sediment). Dilution of the 2 h PEG samples allowed recovery of Vibrio spp. at three sites which gave a negative qPCR result using undiluted samples; this resulted in an average fold increase of 3×10^7 gc/100 g sediment. However, the remaining five sites remained negative using qPCR result when testing on the diluted samples. In contrast, Vibrio spp. target recoveries decreased by 1 log₁₀ gc/100 g sediment for the 16 h treatment with dilution, principally due to low initial nucleic acid concentration (typically <0.1 ng/ μ L). This highlights the role of PEG incubation on inhibitor carryover.

Interestingly very low *Vibrio* concentrations were observed using dPCR for quantification regardless of the PEG incubation times. It has been suggested that qPCR-based quantification may overestimate the actual DNA concentrations, whereas dPCR gives a more accurate estimate on DNA concentrations (Yang et al., 2014). However, the five orders of magnitude difference between the dPCR and qPCR results is higher than expected and may be due to the inhibition of the dPCR. Based on the high variation observed among samples analyzed by dPCR, the low concentrations may be due to analytical error.

The observed *Vibrio* concentrations using the KAPA mastermix correlated with previous findings on the concentration of *Vibrio* spp. in estuarine sediment. Using qPCR for quantification the *Vibrio* concentrations were 8–13 log10 gc/100 g estuarine sediment (Givens et al., 2014) and 4–9 log10 gc/100 g costal sediment (Vezzulli et al., 2009). Therefore, 16 h PEG incubation times when extracting nucleic acids and use of inhibitor resistant mastermix is recommended when quantifying bacteria/viruses from sediment samples.

Direct Extraction of Viral RNA from Sediment

The results of direct extractions of NoV RNA from sediment are summarized in Table 3. Typically, when samples were analyzed directly by qRT-PCR, very low, or no recovery was noted. However, when samples were diluted 10-fold prior to quantification, viral RNA could be recovered. As shown previously, PEG has the capacity to enrich inhibitors affecting the reverse transcriptase and DNA polymerase enzymes (Miura et al., 2011). Our results suggest that with sample dilution, the effect of inhibitors was eliminated and 82% recovery was achieved in samples incubated for 16 h in PEG solution. In contrast, NoV was not recovered with the original CTAB-based method of Griffiths et al. (2000) with 2 h PEG incubation regardless of sample dilution. In accordance with our findings on Vibrio spp. recovery (detailed above), these results suggest that the short incubation time did not allow for good levels of nucleic acid binding to PEG.

The PowerSoil[®] Total RNA Isolation Kit also performed well for the direct recovery of viral RNA, however the recoveries were slightly lower than those achieved using the CTAB-based extraction with 16 h PEG incubation (21 vs. 82%). Even though the kit uses inhibitor removal technology and column-based purification after the PCI extraction, the dilution of the sample prior to qRT-PCR was necessary to accurately quantify viral RNA.

In contrast to the PowerSoil[®] and CTAB protocols, low recoveries (<0.1%) were observed in samples extracted using the PowerViral[®] Environmental DNA/RNA Isolation Kit. Although previous research has demonstrated successful recovery of viruses from environmental samples, such as biosolids and concentrated surface water, using this method (Iker et al., 2013), our results suggest that this approach is not adequate for sediment samples.

TABLE 3 | Genome copy numbers (gc) and recoveries (Rec%) of NoV RNA from spiked sediment samples using direct extraction methods.

Sample	Dilution	Concentration (gc/sample)	SD	Rec%	SD
CTAB 16 h	Undiluted	0	0	0	0
CTAB 16 h	10× diluted	1.2×10^{5}	1.7×10^{4}	82.1	11.4
CTAB 2 h	Undiluted	0	0	0	0
CTAB 2 h	10× diluted	0	0	0	0
PowerSoil	Undiluted	4.5×10^{1}	7.5×10^{0}	<0.1	<0.1
PowerSoil	10× diluted	3.0×10^{4}	7.7×10^{1}	20.5	0.1
PowerViral	Undiluted	3.5×10^{1}	5.1 × 10 ⁰	<0.1	<0.1
PowerViral	$10 \times \text{diluted}$	1.6×10^{2}	6.3 × 10 ¹	<0.1	<0.1

Number of copies added to each sample: 1.47×10^5 . Diluted samples were ten times diluted prior to qRT-PCR. 2 and 16 h indicates the duration of PEG6000 incubation. SD represent standard deviations of independent experiments (n = 2).

Direct extraction has been shown to be suitable for the detection of viral nucleic acids in environmental samples however, recoveries depend on the type of the sample and the target viruses (Miura et al., 2011; Honjo et al., 2012; Iker et al., 2013). The CTAB-based extraction has been widely used for studying microbial communities in soil (Amos et al., 2015; Mayer et al., 2015; Santos et al., 2015), and this study supports its usefulness for the recovery of high abundance viruses and bacteria from sediment. It is a cost-effective approach and requires less preparation time than the PowerSoil[®] Kit (**Table 1**). The major disadvantage of direct extraction is the use of hazardous chemicals and physical extraction in order to separate nucleic acids. These treatments may affect the integrity of the viral and bacterial genomes, although this is unlikely to impact qPCR-based quantification directly.

Effect of Beef Extract-Based Elution on Nucleic Acid Extraction and RT-qPCR

Beef extract-based solutions are routinely used for the elution of infectious viral particles from environmental samples, e.g., sediment and water (Pepper and Gerba, 2000; Ikner et al., 2012). However, the residual beef extract may interfere with enzymes used for molecular quantification (Iker et al., 2016). In order to determine beef extraction-related inhibition, 3% beef extract in 2 M NaNO₃ pH 5.5 solution was spiked with a known concentration of NoV and concentrated using PEG solution. Viral nucleic acids were extracted from the PEG pellet and quantified using PCR-based approaches.

Results suggested that beef extract solution did not significantly inhibit the extraction and qRT-PCR (**Table 4**). Using the RNA Ultrasense One-step qRT-PCR system for quantification, results suggested no inhibition on quantification using the MiniMag[®] System or the PowerViral[®] Kit. Using these extraction approaches, the NoV RNA was fully recovered, suggesting that the residual beef extract and the PEG had no effect on the extraction or quantification. As the CTAB-based extraction gave the best recoveries for the direct extraction of viral nucleic acids from sediment, its usefulness for indirect extraction was investigated. Results showed only 1% recovery, TABLE 4 | Genome copy numbers (gc) and recoveries (Rec%) of NoV from 3% beef extract in 2 M NaNO₃, pH 5.5–PEG precipitation, followed by RNA extraction: NucliSENS[®] MiniMag[®] Nucleic Acid Purification System (MM), the PowerViralTM Environmental RNA/DNA Isolation Kit (PV), and the CTAB-based method adapted from Griffiths et al. (2000; CTAB).

Quantification	Extraction	Undiluted nucleic acid extract				Diluted nucleic acid extract			
		Control		Sample		Control		Sample	
		gc	Gc	Rec%	SD	gc	gc	Rec%	SD
RNA Ultrasense	MM	2.3 × 10 ⁵	2.8 × 10 ⁵	123.9	15.9	2.2 × 10 ⁵	2.3 × 10 ⁵	100.9	28.5
One-step	PV	2.3×10^{5}	2.8×10^{5}	112.6	54.4	2.1 × 10 ⁵	3.3×10^{5}	133.0	44.3
qRT-PCR	CTAB	6.6×10^{4}	1.7×10^{3}	0.9	1.5	2.3×10^{5}	2.2×10^{3}	1.1	2.0
Oasig	MM	4.5×10^{2}	3.1 × 10 ²	_*	_*	4.7 × 10 ²	3.4×10^{2}	_*	_*
One-step	PV	5.6×10^{1}	5.3×10^{1}	_*	_*	6.3×10^{1}	1.3×10^{2}	_*	_*
qRT-PCR	CTAB	5.9×10^{4}	0	0	0	0	0	0	0
Superscript IV RT	MM	1.7 × 10 ⁵	3.5×10^{4}	12.4	3.3	2.6 × 10 ⁵	2.8 × 10 ⁵	97.9	3.0
KAPA force	PV	1.6×10^{4}	5.3×10^{4}	20.7	2.7	2.5×10^{4}	2.3×10^{5}	89.1	3.3
Probe qPCR	CTAB	2.7×10^{3}	2.7×10^{1}	<0.1	0	2.5×10^3	9.0×10^{3}	3.9	1.8

Undiluted and 10 times diluted nucleic acid extracts were subject to quantification. SD represent standard deviations of experiments (n = 3). *Recoveries were not calculated due to the low RNA concentration observed in the controls.

TABLE 5 | Genome copy numbers (gc) and recoveries (Rec%) of NoV from sediment using elution—concentration followed by RNA extraction: NucliSENS[®] MiniMag[®] Nucleic Acid Purification System (MM) and the PowerViralTM Environmental RNA/DNA Isolation Kit (PV).

Quantification Extrac	Extraction	Control	Undiluted nucleic acid extract			Diluted nucleic acid extract			n
		gc	gc	Rec%	SD	gc	Rec%	SD	
RNA Ultrasense	MM	3.8 × 10 ⁵	2.7 × 10 ⁵	70.9	14.6	4.3 × 10 ⁵	112.0	1.5	3
One-step qRT-PCR	PV	2.8 × 10 ⁵	2.8×10^{4}	10.1	13.6	1.7 × 10 ⁵	62.8	19.7	3
Superscript IV RT KAPA Force	MM	3.4×10^{5}				2.9×10^{5}	83.1	1.6	2
Probe qPCR	PV					2.4×10^{5}	68.6	3.7	2
Superscript IV RT	MM	4.0 × 10 ⁵				1.9 × 10 ⁵	46.7	_	1
dPCR	PV					3.6×10^{4}	8.8	12.3	2

Undiluted and 10 times diluted nucleic acid extracts were subject to quantification. SD represent standard deviations of independent experiments.

and sample dilution prior to qRT-PCR did not improve recovery efficiency. This suggested that the extraction was inhibited by residual beef extract and therefore this approach was not used in subsequent indirect extractions.

In order to further investigate the inhibitory effect of beef extract on quantification, different qPCR-based methods were tested. The Oasig OneStep qRT-PCR system did not provide the accurate quantification of NoV RNA in diluted fecal matter and in the beef extract samples (**Table 4**). The concentrations detected in both sample types were 3–4 orders of magnitude lower than those detected using the RNA Ultrasense One-step qRT-PCR system, regardless of the extraction method used (**Table 4**). The controls and dilution also showed lower concentrations than expected suggesting that the qRT-PCR conditions were suboptimal. Nonetheless, increased RT-step and extension times did not improve the results (data not shown). Therefore, the Oasig OneStep qRT-PCR system was not deemed appropriate for the sample types/conditions examined in this study.

Indirect Extraction of Viral Nucleic Acids from Sediment

The results of virus recovery from sediment spiked with NoV using beef extract elution followed by PEG precipitation are summarized in **Table 5**. The method was shown to extract 80–100% of rotavirus and hepatitis A virus from sediment (Lewis and Metcalf, 1988) and used to quantify enterovirus, rotavirus, and hepatitis A virus (Le Guyader et al., 1994) and adenovirus (Staggemeier et al., 2015b) in sediment.

In this study, different extraction and quantification methods for the recovery of NoV RNA were compared. The best recoveries were achieved using the MiniMag[®] System followed by qRT-PCR using the RNA Ultrasense system, however sample dilution prior to quantification was necessary to achieve full NoV recovery. Using the MiniMag[®] System followed by two-step RTqPCR also resulted in high recoveries (83%) together with the PowerViral[®] Kit extraction followed by one-step or two-step RT-qPCR (63–69%). Using the two-step quantification approach, $4\,\mu L$ of the RNA extract was the subject of the RT step, resulting in a five-time diluted sample. No further dilution was necessary prior to qPCR.

The cDNA sample derived from RT was additionally quantified using dPCR. For samples extracted using the MiniMag[®] System, the recovery rate (47%) was comparable with qPCR-based quantification, whereas recoveries observed in samples extracted using the PowerViral[®] Kit were significantly lower and showed high variation (0.2–17.5%) suggesting this extraction approach was not appropriate for dPCR quantification from sediments.

Previous reports have also shown that the MiniMag[®] System is suitable for the extraction of nucleic acids of enteric viruses from concentrated water samples treated with beef extract solution (Rutjes et al., 2005; Baert et al., 2011; Sano et al., 2011). The system has also been used for viral nucleic acid extraction from meat, fruits and vegetables (Butot et al., 2007; Summa et al., 2012). Furthermore, a method using the MiniMag[®] System for the extraction of hepatitis A virus and NoV for the detection of enteric viruses in shellfish has attained accreditation by the International Organization for Standardization (ISO; Lees and CEN WG6 TAG4, 2010) suggesting that the system is suitable for the extraction of viral nucleic acids from difficult environmental samples. The PowerViral® Kit has also been shown to outperform other commercial extraction kits when viral nucleic acids were extracted from fecal matter, biosolids, and concentrated water samples with residual beef extract (Iker et al., 2013). The extractions can be performed within 45 min and the use of non-standard equipment and hazardous chemicals is not a requirement of successful extraction (Table 1).

Assessment of Inhibition

In order to understand the inhibitory effect of environmental matrices on PCR amplification, the negative controls of direct and indirect extractions were spiked with known concentrations of plasmid DNA incorporating the target sequence of NoV qRT-PCR. Results suggested that the inhibitors extracted along with nucleic acids had no or little effect on the polymerase enzymes of the four quantification types when indirect extraction was used combined with MiniMag[®] or PowerViral[®] Kit extraction (Table 6). However, the negative results suggested that those samples where direct extraction was used contained high concentrations of inhibitors affecting the polymerase activity of the Oasig OneStep qRT-PCR and the KAPA Force Probe qPCR systems. Nonetheless, DNA could be precisely quantified using the RNA Ultrasense and dPCR systems regardless of extraction type. Results imply that the observed inhibition described in previous sections had a significant effect on reverse transcriptase but little impact on the polymerase enzymes.

q(RT)-PCR vs. Digital PCR

In this study, we aimed to evaluate the usefulness of 3D dPCR for the detection and quantification of viral and bacterial nucleic acids. However, while the plasmid spiking experiments suggested no significant inhibition from environmental samples (**Table 6**), the results of the *Vibrio* experiments suggested some obstruction may be due to the presence of inhibitors or analytical

TABLE 6 | Copy numbers and recoveries (Rec%) of plasmid DNA from negative NoV RNA extracts of: indirect extraction of sediment using NucliSENS[®] MiniMag[®] Nucleic Acid Purification System (MM), indirect extraction using PowerViralTM Environmental RNA/DNA Isolation Kit (PV), and direct extraction using CTAB-based method adapted from Griffiths et al. (2000; CTAB).

Control	3.1 × 10 ³		
MM	3.3×10^{3}	108.4	14.4
PV	1.7×10^{3}	56.5	17.0
CTAB	2.1×10^{3}	69.0	5.7
Control	3.0 × 10 ³		
MM	2.6×10^{3}	88.2	9.5
PV	2.9×10^{3}	97.1	18.0
CTAB	0	0	0
Control	3.0 × 10 ³		
MM	2.8×10^{3}	91.6	8.6
PV	2.8×10^{3}	93.7	3.2
CTAB	0	0	0
Control	2.0 × 10 ³		
MM	2.0×10^{3}	96.1	15.4
PV	1.7×10^{3}	85.7	7.8
CTAB	2.8×10^{3}	136.1	3.8
	Control MM PV CTAB Control MM PV CTAB Control MM PV CTAB Control MM PV CTAB	$\begin{array}{c c} {\rm Control} & 3.1 \times 10^3 \\ {\rm MM} & 3.3 \times 10^3 \\ {\rm PV} & 1.7 \times 10^3 \\ {\rm CTAB} & 2.1 \times 10^3 \\ {\rm Control} & 3.0 \times 10^3 \\ {\rm MM} & 2.6 \times 10^3 \\ {\rm PV} & 2.9 \times 10^3 \\ {\rm CTAB} & 0 \\ {\rm Control} & 3.0 \times 10^3 \\ {\rm MM} & 2.8 \times 10^3 \\ {\rm PV} & 2.0 \times 10^3 \\ {\rm MM} & 2.0 \times 10^3 \\ {\rm MM} & 2.0 \times 10^3 \\ {\rm PV} & 1.7 \times 10^3 \\ {\rm CTAB} & 2.8 \times 10^3 \\ \end{array}$	$\begin{array}{c c c c c c c c } & 3.1 \times 10^3 & & & & \\ & MM & 3.3 \times 10^3 & 108.4 & \\ & PV & 1.7 \times 10^3 & 56.5 & \\ & CTAB & 2.1 \times 10^3 & 69.0 & \\ \hline & Control & 3.0 \times 10^3 & & \\ & MM & 2.6 \times 10^3 & 88.2 & \\ & PV & 2.9 \times 10^3 & 97.1 & \\ & CTAB & 0 & 0 & \\ \hline & Control & 3.0 \times 10^3 & & \\ & MM & 2.8 \times 10^3 & 91.6 & \\ & PV & 2.8 \times 10^3 & 91.6 & \\ & PV & 2.8 \times 10^3 & 93.7 & \\ & CTAB & 0 & 0 & \\ \hline & Control & 2.0 \times 10^3 & \\ & MM & 2.0 \times 10^3 & 96.1 & \\ & PV & 1.7 \times 10^3 & 85.7 & \\ & CTAB & 2.8 \times 10^3 & 136.1 & \\ \hline \end{array}$

SD represent standard deviations of independent experiments (n = 2).

limitations of the technology (Figure 2). Furthermore, dPCR only gives accurate quantification in a narrow concentration range ($\sim 10^4 - 10^1$ DNA copies/µL), whereas qPCR covers a much broader range $(10^6 - 10^1 \text{ DNA copies}/\mu\text{L})$. Both approaches failed to give accurate quantification for low concentration samples (<30 copies/reaction), however using q(RT-)PCR, a clear amplification curve was observed suggesting that those samples were positive. On the contrary, dPCR results for those samples were either considered negative or results were ambiguous due to background noise. Previous studies have achieved more reliable results on environmental samples using dPCR than with qPCR (Hoshino and Inagaki, 2012; Coudray-Meunier et al., 2015), however the water-oil emulsion droplet technology used by the equipment validated in those studies may give different recoveries from 3D dPCR system. Overall, based on our results, further validation and method development is essential in order to routinely use 3D dPCR approaches in environmental studies.

Molecular Detection of Pathogen-Derived Nucleic Acids

Nucleic acid extraction followed by qPCR-based quantification is the most commonly used method for the rapid detection of pathogens in environmental samples, allowing prompt response in case of a public health threat. However, false negative results may occur as a result of PCR inhibition or poor efficiency of nucleic acid extraction



(Miura et al., 2011; Iker et al., 2013; Staggemeier et al., 2015b). Hence, the methods used for the detection of pathogens should be thoroughly validated. Sediments have the ability to accumulate bacterial and viral pathogens, however, they also accumulate organic matter that may inhibit extraction and qPCR quantification and therefore special measures are needed for method development.

Ideally nucleic acid extraction methods are suitable for the co-extraction of RNA and DNA originated from a wide range of microorganisms and viruses. Our results suggested that using CTAB and beadbeating for cell/viral lysis efficiently released bacterial and viral nucleic acids. The elution-concentration method using beef extract for elution and PEG6000 for precipitation described here and in Lewis and Metcalf (1988) has been shown to be suitable for the detection and quantification of enteric viruses in sediment (**Figure 3**).

Regardless of the nucleic acid extraction method used a ten times sample dilution was necessary prior to qPCR. The only exception was when the KAPA Robust qPCR mastermix was used to quantify Vibrio DNA. Sample dilution may not be needed if post-extract purification procedures are used, however these additional steps may enhance the degradation of viral and bacterial genomes. Nonetheless, the detection limit of qPCRbased approaches is generally very low (≤ 10 gc) and sample dilution prior to q(RT-)PCR would still allow the detection of 10-100 gc. Hence, even when sample dilution is necessary, the assay is far more sensitive than other available quantification methods e.g., culturing. Regardless of sample dilution prior to qPCR, quantification showed great variation depending on the approach used (Figure 3). The best recoveries were achieved using the RNA Ultrasense One-step gRT-PCR kit for viral RNA and the KAPA Robust qPCR mastermix for both bacterial DNA and viral cDNA. Interestingly, the dPCR approach we used throughout this study did not efficiently quantify the target DNA/cDNA sequences. As discussed in Section q(RT)-PCR vs. Digital PCR, this may have been a result of analytical error. Further, evaluation of dPCR is necessary in order to assess the usefulness of dPCR for the quantification of DNA in sediment samples.

CONCLUSIONS AND RECOMMENDATIONS

Direct extraction of nucleic acids using the CTAB method has been shown to be suitable for the co-extraction of high abundance viral and bacterial nucleic acids, and further analysis of the sample would help to better understand the composition of microbial and viral communities and to estimate health risks.

For NoV, indirect extraction combined with RNA extraction using the MiniMag^(R) System and with quantification using the RNA Ultrasense kit gave the best recoveries (**Figure 3**), however sample dilution prior to qRT-PCR was necessary for accurate quantification. Furthermore, the samples obtained via PEG precipitation may be used for the evaluation of viral infectivity for viruses that can be cultured *in vitro* (Lewis and Metcalf, 1988) and viral integrity assays. Therefore, we recommend the use of this elution—concentration protocol for different types of sediment, however, as suggested by others (Le Guyader et al., 2009; Mattison et al., 2009), extraction and quantification controls should be used to address assay performance.

The two-step RT-qPCR method described here also gave adequate results. Our results suggest that better adjusted conditions for the RT and/or the more advanced reverse transcriptase used during the separate RT step enabled full cDNA synthesis in the sample. In addition, twostep approaches may be a useful and cost-efficient tool when it is desirable to study different RNA viruses in a single sample. Currently, we do not recommend the application of 3D dPCR for nucleic acid quantification in environmental studies, however it may be a valuable tool for precise quantification when used along with q(RT-)PCR.

AUTHOR CONTRIBUTIONS

KF and FH wrote the manuscript and undertook the experiments and data analysis. SM, JM, and DJ advised on experiment design. KF, FH, SM, JM, and DJ prepared manuscript for submission. All authors have approved the final version to be published.

FUNDING

This work was funded by the Natural Environment Research Council (NERC) and the Food Standards Agency (FSA) under

REFERENCES

- Amos, G. C., Borsetto, C., Laskaris, P., Krsek, M., Berry, A. E., Newsham, K. K., et al. (2015). Designing and implementing an assay for the detection of rare and divergent NRPS and PKS clones in European, Antarctic and Cuban soils. *PLoS ONE* 10:e01 38327. doi: 10.1371/journal.pone.0138327
- An, Y. J., Kampbell, D. H., and Breidenbach, G. P. (2002). Escherichia coli and total coliforms in water and sediments at lake marinas. Environ. Pollut. 120, 771–778. doi: 10.1016/S0269-7491(02)00173-2
- Anderson, K. L., Whitlock, J. E., and Harwood, V. J. (2005). Persistence and differential survival of fecal indicator bacteria in subtropical waters and sediments. *Appl. Environ. Microbiol.* 71, 3041–3048. doi: 10.1128/AEM.71.6.3041-3048.2005
- Bae, S., and Wuertz, S. (2009). Rapid decay of host-specific fecal Bacteroidales cells in seawater as measured by quantitative PCR with propidium monoazide. *Water Res.* 43, 4850–4859. doi: 10.1016/j.watres.2009.06.053
- Baert, L., Mattison, K., Loisy-Hamon, F., Harlow, J., Martyres, A., Lebeau, B., et al. (2011). Review: norovirus prevalence in Belgian, Canadian and French fresh produce: a threat to human health? *Int. J. Food Microbiol.* 151, 261–269. doi: 10.1016/j.ijfoodmicro.2011.09.013
- Butot, S., Putallaz, T., and Sanchez, G. (2007). Procedure for rapid concentration and detection of enteric viruses from berries and vegetables. *Appl. Environ. Microbiol.* 73, 186–192. doi: 10.1128/AEM.01248-06
- Charles, K. J., Shore, J., Sellwood, J., Laverick, M., Hart, A., and Pedley, S. (2009). Assessment of the stability of human viruses and coliphage in groundwater by PCR and infectivity methods. *J. Appl. Microbiol.* 106, 1827–1837. doi: 10.1111/j.1365-2672.2009.04150.x
- Coudray-Meunier, C., Fraisse, A., Martin-Latil, S., Guillier, L., Delannoy, S., Fach, P., et al. (2015). A comparative study of digital RT-PCR and RT-qPCR for quantification of hepatitis A virus and norovirus in lettuce and water samples. *Int. J. Food Microbiol.* 201, 17–26. doi: 10.1016/j.ijfoodmicro.2015. 02.006
- Davies, C. M., Long, J. A., Donald, M., and Ashbolt, N. J. (1995). Survival of fecal microorganisms in marine and freshwater sediments. *Appl. Environ. Microbiol.* 61, 1888–1896.
- Dick, L. K., and Field, K. G. (2004). Rapid estimation of numbers of fecal Bacteroidetes by use of a quantitative PCR assay for 16S rRNA genes. *Appl. Environ. Microbiol.* 70, 5695–5697. doi: 10.1128/AEM.70.9.5695-5697. 2004
- Ding, X., and Mu, Y. (2016). Research and application progress of digital nucleic acid amplification detection techniques. *Chin. J. Anal. Chem.* 44, 512–521. doi: 10.1016/S1872-2040(16)60918-0
- Duhamel, S., and Jacquet, S. (2006). Flow cytometric analysis of bacteria- and virus-like particles in lake sediments. J. Microbiol. Methods 64, 316–332. doi: 10.1016/j.mimet.2005.05.008
- Farkas, K., Varsani, A., Marjoshi, D., Easingwood, R., McGill, E., and Pang, L. (2015). Size exclusion-based purification and PCR-based quantitation of MS2 bacteriophage particles for environmental applications. *J. Virol. Methods* 213, 135–138. doi: 10.1016/j.jviromet.2014.11.024
- Flannery, J., Rajko-Nenow, P., Winterbourn, J. B., Malham, S. K., and Jones, D. L. (2014). Effectiveness of cooking to reduce norovirus and infectious F-specific

the Environmental Microbiology and Human Health (EMHH) Programme (NE/M010996/1).

ACKNOWLEDGMENTS

The authors would like to thank Prof. Ian Goodfellow and Dr. Lydia Drumwright (University of Cambridge, UK) for providing virus sample, Life Technologies Ltd., for use of the QuantStudioTM 3D Digital PCR System and Prof. Alan McCarthy (University of Liverpool) for his helpful review comments on the manuscript.

RNA bacteriophage concentrations in *Mytilus edulis. J. Appl. Microbiol.* 117, 564–571. doi: 10.1111/jam.12534

- Fraisse, A., Coudray-Meunier, C., Martin-Latil, S., Hennechart-Collette, C., Delannoy, S., Fach, P., et al. (2017). Digital RT-PCR method for hepatitis A virus and norovirus quantification in soft berries. *Int. J. Food Microbiol.* 243, 36–45. doi: 10.1016/j.ijfoodmicro.2016.11.022
- Furet, J. P., Firmesse, O., Gourmelon, M., Bridonneau, C., Tap, J., Mondot, S., et al. (2009). Comparative assessment of human and farm animal faecal microbiota using real-time quantitative PCR. *FEMS Microbiol. Ecol.* 68, 351–362. doi: 10.1111/j.1574-6941.2009.00671.x
- Gerba, C. P., and McLeod, J. S. (1976). Effect of sediments on the survival of *Escherichia coli* in marine waters. *Appl. Environ. Microbiol.* 32, 114–120.
- Girones, R., Ferrús, M. A., Alonso, J. L., Rodriguez-Manzano, J., Calgua, B., de Abreu Corrêa, A., et al. (2010). Molecular detection of pathogens in water-the pros and cons of molecular techniques. *Water Res.* 44, 4325–4339. doi: 10.1016/j.watres.2010.06.030
- Givens, C. E., Bowers, J., DePaola, A., Hollibaugh, J., and Jones, J. (2014). Occurrence and distribution of *Vibrio vulnificus* and *Vibrio parahaemolyticus*potential roles for fish, oyster, sediment and water. *Lett. Appl. Microbiol.* 58, 503–510. doi: 10.1111/lam.12226
- Green, D. H., and Lewis, G. D. (1999). Comparative detection of enteric viruses in wastewaters, sediments and oysters by reverse transcription-PCR and cell culture. *Water Res.* 33, 1195–1200. doi: 10.1016/S0043-1354(98)00313-3
- Griffiths, R. I., Whiteley, A. S., O'Donnell, A. G., and Bailey, M. J. (2000). Rapid method for coextraction of DNA and RNA from natural environments for analysis of ribosomal DNA- and rRNA-based microbial community composition. *Appl. Environ. Microbiol.* 66, 5488–5491. doi: 10.1128/AEM.66.12.5488-5491.2000
- Haramoto, E., Kitajima, M., Katayama, H., Asami, M., Akiba, M., and Kunikane, S. (2009). Application of real-time PCR assays to genotyping of F-specific phages in river water and sediments in Japan. *Water Res.* 43, 3759–3764. doi: 10.1016/j.watres.2009.05.043
- Honjo, M. N., Minamoto, T., and Kawabata, Z. (2012). Reservoirs of Cyprinid herpesvirus 3 (CyHV-3) DNA in sediments of natural lakes and ponds. *Vet. Microbiol.* 155, 183–190. doi: 10.1016/j.vetmic.2011.09.005
- Hoshino, T., and Inagaki, F. (2012). Molecular quantification of environmental DNA using microfluidics and digital PCR. Syst. Appl. Microbiol. 35, 390–395. doi: 10.1016/j.syapm.2012.06.006
- Iker, B. C., Bright, K. R., Pepper, I. L., Gerba, C. P., and Kitajima, M. (2013). Evaluation of commercial kits for the extraction and purification of viral nucleic acids from environmental and fecal samples. *J. Virol. Methods* 191, 24–30. doi: 10.1016/j.jviromet.2013.03.011
- Iker, B. C., Kitajima, M., and Gerba, C. P. (2016). "Extraction and purification of viral nucleic acids from environmental samples," in *Sample Preparation Techniques for Soil, Plant, and Animal Samples*, ed M. Micic (New York, NY: Springer), 315–324.
- Ikner, L. A., Gerba, C. P., and Bright, K. R. (2012). Concentration and recovery of viruses from water: a comprehensive review. *Food Environ. Virol.* 4, 41–67. doi: 10.1007/s12560-012-9080-2
- International Organization for Standardization, ISO 9308-1:2014 (2014). Water Quality – Enumeration of Escherichia coli and Coliform Bacteria – Part 1:

Membrane Filtration Method for Waters with Low Bacterial Background Flora. Geneva.

- Ishii, S., Kitamura, G., Segawa, T., Kobayashi, A., Miura, T., Sano, D., et al. (2014). Microfluidic quantitative PCR for simultaneous quantification of multiple viruses in environmental water samples. *Appl. Environ. Microbiol.* 80, 7505–7511. doi: 10.1128/AEM.02578-14
- Jin, Y., and Flury, M. (2002). Fate and transport of viruses in porous media. *Adv. Agron.* 77, 39–102. doi: 10.1016/S0065-2113(02)77013-2
- Johnson, R. A., Ellender, R. D., and Tsai, S. C. (1984). Elution of enteric viruses from Mississippi estuarine sediments with lecithin-supplemented eluents. *Appl. Environ. Microbiol.* 48, 581–585.
- Kim, M., Gutiérrez-Cacciabue, D., Schriewer, A., Rajal, V., and Wuertz, S. (2014). Evaluation of detachment methods for the enumeration of *Bacteroides fragilis* in sediments via propidium monoazide quantitative PCR, in comparison with *Enterococcus faecalis* and *Escherichia coli. J. Appl. Microbiol.* 117, 1513–1522. doi: 10.1111/jam.12630
- Krsek, M., and Wellington, E. (1999). Comparison of different methods for the isolation and purification of total community DNA from soil. J. Microbiol. Methods 39, 1–16. doi: 10.1016/S0167-7012(99)00093-7
- LaBelle, R. L., and Gerba, C. P. (1980). Influence of estuarine sediment on virus survival under field conditions. *Appl. Environ. Microbiol.* 39, 749–755.
- LaMontagne, M., Michel, F., Holden, P., and Reddy, C. (2002). Evaluation of extraction and purification methods for obtaining PCR-amplifiable DNA from compost for microbial community analysis. J. Microbiol. Methods 49, 255–264. doi: 10.1016/S0167-7012(01)00377-3
- Landry, E. F., Vaughn, J. M., Vicale, T. J., and Mann, R. (1983). Accumulation of sediment-associated viruses in shellfish. Appl. Environ. Microbiol. 45, 238–247.
- Le Guyader, F. S., Parnaudeau, S., Schaeffer, J., Bosch, A., Loisy, F., Pommepuy, M., et al. (2009). Detection and quantification of noroviruses in shellfish. *Appl. Environ. Microbiol.* 75, 618–624. doi: 10.1128/AEM. 01507-08
- Le Guyader, F., Dubois, E., Menard, D., and Pommepuy, M. (1994). Detection of hepatitis A virus, rotavirus, and enterovirus in naturally contaminated shellfish and sediment by reverse transcription-seminested PCR. *Appl. Environ. Microbiol.* 60, 3665–3671.
- Lees, D., and CEN WG6 TAG4 (2010). International standardisation of a method for detection of human pathogenic viruses in molluscan shellfish. *Food Environ. Virol.* 2, 146–155. doi: 10.1007/s12560-010-9042-5
- Lewis, G. D., and Metcalf, T. G. (1988). Polyethylene glycol precipitation for recovery of pathogenic viruses, including hepatitis A virus and human rotavirus, from oyster, water, and sediment samples. *Appl. Environ. Microbiol.* 54, 1983–1988.
- Lowther, J. A., Gustar, N. E., Powell, A. L., Hartnell, R. E., and Lees, D. N. (2012). Two-year systematic study to assess norovirus contamination in oysters from commercial harvesting areas in the United Kingdom. *Appl. Environ. Microbiol.* 78, 5812–5817. doi: 10.1128/AEM.01046-12
- Mattison, K., Brassard, J., Gagné, M., Ward, P., Houde, A., Lessard, L., et al. (2009). The feline calicivirus as a sample process control for the detection of food and waterborne RNA viruses. *Int. J. Food Microbiol.* 132, 73–77. doi: 10.1016/j.ijfoodmicro.2009.04.002
- Mayer, R. E., Vierheilig, J., Egle, L., Reischer, G. H., Saracevic, E., Mach, R. L., et al. (2015). Automated sampling procedures supported by high persistence of bacterial fecal indicators and Bacteroidetes genetic microbial source tracking markers in municipal wastewater during short-term storage at 5°C. Appl. Environ. Microbiol. 81, 5134–5143. doi: 10.1128/AEM. 00998-15
- Meschke, J., and Sobsey, M. (1998). Comparative adsorption of Norwalk virus, poliovirus 1 and F RNA coliphage MS2 to soils suspended in treated wastewater. *Water Sci. Technol.* 38, 187–189. doi: 10.1016/S0273-1223(98)00 823-3
- Miura, T., Masago, Y., Sano, D., and Omura, T. (2011). Development of an effective method for recovery of viral genomic RNA from environmental silty sediments for quantitative molecular detection. *Appl. Environ. Microbiol.* 77, 3975–3981. doi: 10.1128/AEM.02692-10
- Oliveira, J., Cunha, A., Castilho, F., Romalde, J. L., and Pereira, M. J. (2011). Microbial contamination and purification of bivalve shellfish: crucial aspects in monitoring and future perspectives – A mini-review. *Food Control* 22, 805–816. doi: 10.1016/j.foodcont.2010.11.032

- Pepper, I. L., and Gerba, C. P. (2000). "Chapter 8 Environmental sample collection and processing," in *Environmental Microbiology, 3rd Edn.*, eds I. L. Pepper, C. P. Gerba, T. J. Gentry (San Diego, CA: Academic Press), 157–175.
- Perkins, T. L., Clements, K., Baas, J. H., Jago, C. F., Jones, D. L., Malham, S. K., et al. (2014). Sediment composition influences spatial variation in the abundance of human pathogen indicator bacteria within an estuarine environment. *PLoS ONE* 9:e112951. doi: 10.1371/journal.pone.0112951
- Pinto, D., Santos, M. A., and Chambel, L. (2015). Thirty years of viable but nonculturable state research: unsolved molecular mechanisms. *Crit. Rev. Microbiol.* 41, 61–76. doi: 10.3109/1040841X.2013.794127
- Radin, D. (2014). New trends in food-and waterborne viral outbreaks. Arch. Biol. Sci. 66, 1–9. doi: 10.2298/ABS1401001R
- Rames, E., Roiko, A., Stratton, H., and MacDonald, J. (2016). Technical aspects of using human adenovirus as a viral water quality indicator. *Water Res.* 96, 308–326. doi: 10.1016/j.watres.2016.03.042
- Rao, V. C., Metcalf, T. G., and Melnick, J. L. (1986). Human viruses in sediments, sludges, and soils. *Bull. World Health Organ.* 64, 1–13.
- Rock, C., Alum, A., and Abbaszadegan, M. (2010). PCR inhibitor levels in concentrates of biosolid samples predicted by a new method based on excitation-emission matrix spectroscopy. *Appl. Environ. Microbiol.* 76, 8102–8109. doi: 10.1128/AEM.02339-09
- Roose-Amsaleg, C., Garnier-Sillam, E., and Harry, M. (2001). Extraction and purification of microbial DNA from soil and sediment samples. *Appl. Soil Ecol.* 18, 47–60. doi: 10.1016/S0929-1393(01)00149-4
- Rutjes, S. A., Italiaander, R., van den Berg, H. H., Lodder, W. J., and de Roda Husman, A. M. (2005). Isolation and detection of enterovirus RNA from large-volume water samples by using the NucliSens miniMAG system and real-time nucleic acid sequence-based amplification. *Appl. Environ. Microbiol.* 71, 3734–3740. doi: 10.1128/AEM.71.7.3734-3740. 2005
- Sano, D., Pérez-Sautu, U., Guix, S., Pinto, R. M., Miura, T., Okabe, S., et al. (2011). Quantification and genotyping of human sapoviruses in the Llobregat river catchment, Spain. *Appl. Environ. Microbiol.* 77, 1111–1114. doi: 10.1128/AEM.01721-10
- Santos, N., Santos, C., Valente, T., Gortázar, C., Almeida, V., and Correia-Neves, M. (2015). Widespread environmental contamination with *Mycobacterium tuberculosis* complex revealed by a molecular detection protocol. *PLoS ONE* 10:e0142079. doi: 10.1371/journal.pone.0142079
- Sedlak, R. H., and Jerome, K. R. (2013). Viral diagnostics in the era of digital polymerase chain reaction. *Diagn. Microbiol. Infect. Dis.* 75, 1–4. doi: 10.1016/j.diagmicrobio.2012.10.009
- Smith, E. M., Gerba, C. P., and Melnick, J. L. (1978). Role of sediment in the persistence of enteroviruses in the estuarine environment. *Appl. Environ. Microbiol.* 35, 685–689.
- Staggemeier, R., Bortoluzzi, M., da Silva Heck, T. M., da Luz, R. B., Fabres, R. B., Soliman, M. C., et al. (2015a). Animal and human enteric viruses in water and sediment samples from dairy farms. *Agric. Water Manage* 152, 135–141. doi: 10.1016/j.agwat.2015.01.010
- Staggemeier, R., Bortoluzzi, M., da Silva Heck, T. M., da Silva, T., Spilki, F. R., and de Matos Almeida, S. E. (2015b). Molecular detection of human adenovirus in sediment using a direct detection method compared to the classical polyethylene glycol precipitation. *J. Virol. Methods* 213, 65–67. doi: 10.1016/j.jviromet.2014.11.019
- Staley, C., Reckhow, K. H., Lukasik, J., and Harwood, V. J. (2012). Assessment of sources of human pathogens and fecal contamination in a Florida freshwater lake. *Water Res.* 46, 5799–5812. doi: 10.1016/j.watres.2012.08.012
- Summa, M., von Bonsdorff, C., and Maunula, L. (2012). Evaluation of four virus recovery methods for detecting noroviruses on fresh lettuce, sliced ham, and frozen raspberries. J. Virol. Methods 183, 154–160. doi: 10.1016/j.jviromet.2012.04.006
- Topping, J., Schnerr, H., Haines, J., Scott, M., Carter, M., Willcocks, M., et al. (2009). Temperature inactivation of feline calicivirus vaccine strain FCV F-9 in comparison with human noroviruses using an RNA exposure assay and reverse transcribed quantitative real-time polymerase chain reaction—a novel method for predicting virus infectivity. *J. Virol. Methods* 156, 89–95. doi: 10.1016/j.jviromet.2008.10.024
- Vezzulli, L., Pezzati, E., Moreno, M., Fabiano, M., Pane, L., Pruzzo, C., et al. (2009). Benthic ecology of *Vibrio* spp. and pathogenic

Vibrio species in a coastal Mediterranean environment (La Spezia Gulf, Italy). *Microb. Ecol.* 58, 808–818. doi: 10.1007/s00248-009-9542-8

- Vignaroli, C., Luna, G., Pasquaroli, S., Di Cesare, A., Petruzzella, R., Paroncini, P., et al. (2013). Epidemic *Escherichia coli* ST131 and *Enterococcus faecium* ST17 in coastal marine sediments from an Italian beach. *Environ. Sci. Technol.* 47, 13772–13780. doi: 10.1021/es4019139
- Yang, R., Paparini, A., Monis, P., and Ryan, U. (2014). Comparison of nextgeneration droplet digital PCR (ddPCR) with quantitative PCR (qPCR) for enumeration of Cryptosporidium oocysts in faecal samples. *Int. J. Parasitol.* 44, 1105–1113. doi: 10.1016/j.ijpara.2014.08.004

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2017 Farkas, Hassard, McDonald, Malham and Jones. This is an openaccess article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.