



## **Anxiety and perceived psychological stress play an important role in the immune response after exercise**

Edwards, Jason; Walsh, Neil; Diment, Bethany; Roberts, Ross

### **Exercise Immunology Review**

Published: 31/01/2018

Peer reviewed version

[Cyswllt i'r cyhoeddiad / Link to publication](#)

*Dyfyniad o'r fersiwn a gyhoeddwyd / Citation for published version (APA):*

Edwards, J., Walsh, N., Diment, B., & Roberts, R. (2018). Anxiety and perceived psychological stress play an important role in the immune response after exercise. *Exercise Immunology Review*, 24.

#### **Hawliau Cyffredinol / General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

#### **Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

**Anxiety and perceived psychological stress play an important role in the immune response after exercise**

**RUNNING HEAD:** Psychological stress, exercise and immunity

Jason P. Edwards<sup>1</sup>, Neil P. Walsh<sup>1</sup>, Bethany C. Diment<sup>1,2</sup> and Ross Roberts<sup>3</sup>

<sup>1</sup>College of Health and Behavioural Sciences, Bangor University, Bangor, Gwynedd, UK.

<sup>2</sup>Norwich Medical School, University of East Anglia, Norwich, Norfolk, UK.

<sup>3</sup>Institute for the Psychology of Elite Performance, Bangor University, Bangor, Gwynedd, UK.

*Corresponding Author:*

Prof. Neil P. Walsh FACSM

College of Health and Behavioural Sciences,

Bangor University,

Bangor,

LL57 2PZ,

UK.

Email: n.walsh@bangor.ac.uk

Telephone: + 44 1248 383480

## ABSTRACT

There are common pathways by which psychological stress and exercise stress alter immunity. However, it remains unknown whether psychological stress plays a role in the *in vivo* immune response to exercise. We examined the relationship between anxiety and perceived psychological stress reported before exercise and *in vivo* immunity after exercise using skin sensitisation with Diphenylcyclopropanone (DPCP). In a randomised design, sixty four, thoroughly familiarised, males completed widely used psychological instruments to assess state-anxiety and perceived psychological stress before exercise, and ran either 30 minutes at 60% (30MI) or 80% (30HI)  $\dot{V}O_{2peak}$ , 120 minutes at 60% (120MI)  $\dot{V}O_{2peak}$  or rested (CON) before DPCP sensitisation. Cutaneous recall to DPCP was measured **as the dermal thickening response to a low-dose series DPCP challenge 4-weeks after sensitisation.** After accounting for exercise ( $R^2 = 0.20$ ;  $P < 0.01$ ), multiple-regression showed that pre-exercise state-anxiety (STAI-S;  $\Delta R^2 = 0.19$ ;  $P < 0.01$ ) and perceived psychological stress ( $\Delta R^2 = 0.13$ ;  $P < 0.05$ ) **were moderately associated** with the DPCP response after exercise. The STAI-S scores before exercise were considered low-to-moderate in these familiarised individuals (median split; mean STAI-S of low 25 and moderate 34). Further examination showed that the DPCP response after exercise (30MI, 30HI or 120MI) was 62% lower in those reporting low *vs.* moderate state-anxiety before exercise (**mean difference in dermal thickening: -2.6 mm; 95% CI: -0.8 to -4.4 mm;  $P < 0.01$** ). As such, the results indicate a beneficial effect of moderate (*vs.* low) state-anxiety and perceived psychological stress on *in vivo* immunity after exercise. Moreover, correlations were of comparable strength for the relationship between physiological stress (heart rate training impulse) and the summed dermal response to DPCP ( $r = -0.37$ ; **95% CI: -0.05 to -0.62;  $P = 0.01$** ), and state-anxiety and the summed dermal response to DPCP ( $r = 0.39$ ; **95% CI: 0.08 to 0.63;  $P < 0.01$** ). In conclusion, state-anxiety and perceived psychological stress levels before exercise play an

important role in determining the strength of the *in vivo* immune response after exercise. These findings indicate a similar strength relationship for the level of state-anxiety prior to exercise and the level of physiological stress during exercise with the *in vivo* immune response after exercise. Future research is required to investigate exercise-immune responses in athletes, military personnel and others in physically demanding occupations experiencing higher levels of psychological stress than those reported in this study e.g. related to important competition, military operations and major life events. Nevertheless, the present findings support the recommendation that exercise scientists should account for anxiety and psychological stress when examining the immune response to exercise.

**KEYWORDS:** Running, Immunity, In vivo, Diphenhydramine, STAI

## INTRODUCTION

Numerous studies report an increase in upper respiratory tract infection (URTI) symptoms following a bout of strenuous exercise and during periods of heavy training in athletes (25, 33, 37), and there is widespread agreement that a transient suppression of immune function is at least partly responsible (48). A multitude of training and lifestyle stressors are thought to be involved in the observed decrease in immune function in athletes and military personnel; including, prolonged training sessions, exposure to environmental extremes (e.g. heat, cold and high altitude), poor nutrition and poor sleep (41-43, 47, 48). For example, prolonged heavy exercise ( $\geq 2$  h) transiently decreases *in vitro* measures of immunity in isolated blood samples (48) and more clinically meaningful *in vivo* measures of immunity instigated at the skin, including delayed type hypersensitivity (DTH) and contact hypersensitivity (CHS) (6, 16, 24). Indeed, recent work highlights the immunosuppressive effect of prolonged exercise (2 h) on the induction of CHS using the novel antigen Diphenylcyclopropenone (DPCP) (16, 24). Besides the immunosuppressive effects of prolonged heavy training sessions, the training environment and lifestyle stressors such as nutritional deficits (e.g. energy, macro- and micro- nutrients) and poor sleep (e.g. total deprivation and disruption) have long been implicated in the decrease in immune function in athletes and military personnel (41-43). Somewhat surprisingly, field studies (multi-stressor environment) and laboratory studies mimicking real-world athletic and military scenarios by exposing participants to these stressors, either separately or combined, demonstrate only subtle and short-lived modulation of immunity at rest and in response to exercise (5, 28, 31, 40). Rather than decrease immunity, some studies actually show a beneficial ‘priming’ effect of stressors such as short-term sleep disruption (1 night) (28), intermittent cold exposure (29) and intermittent hypoxic exposure on immunity (50). As such, there is a pressing need for research investigating other

likely **behavioural, environmental and lifestyle** candidates involved in the observed decrease in immune function in athletes and military personnel.

Given the well-known and marked influence of psychological stress on immunity and infection resistance (10, 13), and the likely shared mechanisms by which psychological stress and exercise stress alter immunity (36); i.e. principally through activation of the hypothalamic-pituitary-adrenal (HPA) axis and sympathetic-adrenal-medullary (SAM) axis and subsequent immunomodulatory hormones, it has been hypothesised that psychological stress **can** play a role in the decrease in immunity with prolonged heavy exercise and heavy training (8, 36, 49). Unfortunately, exercise immunologists rarely report measures of psychological stress in their studies and so there is little by way of empirical evidence to support this hypothesis (38). That there are striking similarities in the way acute and chronic psychological stress and acute and chronic exercise stress influence immunity provides indirect support for this hypothesis. For example, although chronic psychological stress is widely accepted to decrease immunity and increase infection risk (10, 13), short-lasting, moderate-intensity psychological stress can enhance *in vivo* immunity (21) and is considered a fundamental adaptive response to help us survive (13). Similarly, prolonged heavy exercise and heavy training are widely accepted to decrease immunity and increase infection risk (48), but short-lasting, moderate-intensity exercise stress can enhance *in vivo* immunity (34).

With this information in mind, using a multiple linear regression model, we tested, and provide evidence supporting our hypothesis that the level of anxiety and perceived psychological stress reported by **an** individual prior to exercise play an important role in determining the strength of the *in vivo* immune response to DPCP after exercise.

## METHODS

Using the CHS responses to exercise from a previous study (16), here we present previously unpublished and novel insights regarding the influence of anxiety and perceived psychological stress on *in vivo* immunity after exercise.

### Participants

Sixty four healthy, non-smoking, recreationally active males (age  $22 \pm 3$  years; height  $180 \pm 6$  cm; body mass  $76.7 \pm 11.5$  kg;  $\dot{V}O_{2peak}$   $57 \pm 6$  mL/kg/min) gave written informed consent to participate in the study. Participants had no previous history of exposure to DPCP and were excluded if they were taking any medication or dietary supplements, or had a history of atopy or any other immune-related or inflammatory dermatological **condition**. Participants were required to abstain from alcohol and exercise for 24 h before and 48 h after the experimental trials. The study received local ethics committee approval, and all protocols were conducted in accordance with the Declaration of Helsinki (2013).

Participants were matched for age and aerobic fitness (gas exchange threshold and  $\dot{V}O_{2peak}$ ) before being randomly assigned to one of four groups. Groups were 1) 120 min of seated rest (CON); 2) 30 min of moderate-intensity ( $60\% \dot{V}O_{2peak}$ ) exercise (30MI); 3) 30 min of high-intensity ( $80\% \dot{V}O_{2peak}$ ) exercise (30HI); or 4) 120 min of moderate-intensity ( $60\% \dot{V}O_{2peak}$ ) exercise (120MI).

### Preliminary measures and familiarisation

$\dot{V}O_{2peak}$  was estimated by means of a ramped exercise test on a treadmill (h/p/cosmos Mercury 4.0, Nussdorf-Traunstein, Germany) as described (16). At least 24 h after the preliminary measures and approximately 7 days before the experimental trial, participants

were informed of their group allocation and attended the laboratory for familiarisation. For exercising participants, the calculated exercise intensity was verified, and the participant was familiarised by running for 50% of their allocated exercise duration. During this visit, all participants were familiarised with blood sampling and other relevant procedures.

### **Experimental procedures**

On the day of the experimental trial, participants were transported to the laboratory at 0730 h and provided with a standard breakfast (0.03 MJ/kg) before completing widely used, validated psychological instruments. The level of anxiety was assessed using the state aspect of the State Trait Anxiety Inventory (STAI-S): the STAI-S is one of the most commonly used scales to measure anxiety, which has been defined as an unpleasant emotional state that exists at a given moment in time and at a particular level of intensity, and is characterised by subjective feelings of tension, apprehension, nervousness, and worry (45). The STAI-S consists of 20-items, with responses being measured on a four-point Likert scale (from 1 'not at all' to 4 'very much so') and a range of scores from 20–80 (composite reliability = 0.94). Perceived psychological stress was assessed using the Perceived Stress Scale (PSS): the PSS is a widely used psychological instrument for measuring the perception of stress, and measures the degree to which life situations are considered stressful by the individual during the previous month (11). The PSS is a 14-item inventory, with responses measured on a five-point Likert scale (from 0 'never' to 4 'very often') and a range of scores from 0–56 (composite reliability = 0.73). Average PSS score for young adults has been reported as  $21 \pm 7$  and high PSS score in posttraumatic stress disorder patients as  $34 \pm 8$  (11, 26). Participants assigned to 120MI began running on a treadmill at 1100 h, and those assigned to 30HI and 30MI began at 1230 h, so that all participants completed the exercise at the same time of day (1300 h). Heart rate was monitored continuously during the experimental trials (Polar FT1,



Polar Electro, Kempele, Finland). Immediately after the exercise, participants showered and returned to the laboratory within 15 min of completion before being sensitised to DPCP at 1320 h, exactly 20 min after exercise cessation. This **short standardised** delay in sensitisation allowed cutaneous blood flow to return to baseline (16).

### **Blood collection and analysis**

Blood samples were collected before, immediately after, and 1 h after exercise or seated rest by venepuncture into two separate vacutainer tubes (Becton Dickinson, Oxford, UK), one containing K<sub>3</sub>EDTA, and one containing lithium heparin. The samples were spun at 1500 g for 10 minutes in a refrigerated centrifuge. Plasma was aliquoted into Eppendorf tubes, and immediately frozen at -80 °C for later analysis. Plasma epinephrine and norepinephrine were determined on K<sub>3</sub>EDTA plasma, and plasma cortisol was determined on lithium heparin plasma using commercially available ELISA kits (CatCombi, IBL International, Hamburg, Germany and DRG Instruments, Marburg, Germany, respectively). The intra-assay coefficient of variation for plasma epinephrine, norepinephrine and cortisol was 4.1%, 4.1% and 4.4%, respectively.

### **Induction of CHS**

The sensitising exposure to the novel antigen DPCP involved application of an occluded patch, constituting a 12-mm aluminium Finn chamber (Epitest Oy, Tuusula, Finland) on scanpor hypoallergenic tape containing an 11-mm filter paper disc (16). The paper disc was soaked in 22.8 µL of 0.125% DPCP in acetone (patch = 30 µg/cm<sup>2</sup> DPCP) and allowed to dry for 5 min before being applied to the skin on the lower back for exactly 48 h.

## **Elicitation**

The magnitude of *in vivo* immune responsiveness was quantified by measuring the responses elicited by secondary exposure to DPCP. Twenty eight days after the initial sensitisation to DPCP, all participants received a challenge with a low-concentration dose-series of DPCP on individual patches, each comprising an 8-mm aluminium Finn chamber on scanpor hypoallergenic tape containing a 7-mm filter paper disc. Patches were applied to the volar aspect of the upper arm in the following concentrations: 10  $\mu\text{L}$  of DPCP: 0.0048%, 1.24  $\mu\text{g}/\text{cm}^2$ ; 0.0076%, 1.98  $\mu\text{g}/\text{cm}^2$ ; 0.0122%, 3.17  $\mu\text{g}/\text{cm}^2$ ; 0.0195%, 5.08  $\mu\text{g}/\text{cm}^2$ ; 0.0313%, 8.12  $\mu\text{g}/\text{cm}^2$ ; and, 10  $\mu\text{L}$  of 100% acetone served as a control patch for background subtraction. Patches were applied in randomly allocated order at the local site to minimise any anatomical variability in responses. Elicitation patches were removed after 6 h, and the strength of immune reactivity was assessed as the cutaneous responses 48 h after application (16).

## **Assessment of CHS responses**

Dermal thickness was determined at each elicitation site using a high-frequency ultrasound scanner (Episcan, Longport Inc, Reading, UK). The ultrasound probe was placed over the centre of each patch site together with ultrasound gel. The mean of three measurements was taken from each 12-mm scan image assessed at a later time by a blinded investigator. Mean skinfold thickness was determined from triplicate measurements at each elicitation site using modified spring-loaded skin callipers (Harpenden Skinfold Calliper, British Indicators, England, UK). As previously described (24), this method provides an objective measure of skin oedema (inflammatory swelling). Skinfold thickness was recorded to the nearest 0.1 mm by placing the jaws of the calliper at the outer diameter of the response site and measuring skin thickness only (no subcutaneous fat). Skinfold thickness assessed using skinfold

callipers has previously been shown to be strongly related ( $r = 0.93$ ) with high-frequency ultrasound readings of dermal thickness (16). Mean skin erythema was determined from triplicate measurements at each elicitation site using an erythema meter (ColorMeter DSM11, Cortex Technology, Hadsund, Denmark) which provides an objective measure of skin redness (24). Mean background values were determined from triplicate measurements at the acetone patch site for both thickness and redness. To determine the increase in thickness and redness, the value from the acetone-only site was subtracted from each elicitation site value. The values for increase in dermal thickness, skinfold thickness and erythema over all the doses were summed to give an approximation of the area under the dose–response curve, representative of the overall reactivity of each participant to DPCP (24).

### **Statistical analyses**

Hierarchical linear regression analysis was used to examine the relationship between STAI-S and PSS (in 2 separate models) and *in vivo* immunity after exercise. In step 1 of each model, the influence of exercise on the summed dermal thickening response to DPCP was accounted for by calculating the training impulse (TRIMP) to reflect the level of physiological stress, as described (2). In step 2, the influence of each psychological measure on the summed dermal thickening response to DPCP was assessed. Sample size was deemed appropriate for the multiple linear regression analysis with 2 steps, in line with recommendations (46). To further illustrate the influence of anxiety on *in vivo* immunity after exercise, we performed additional analyses by categorising the population based on STAI-S scores using a median split; whereby, the levels before exercise were defined as low anxiety (LOW: STAI-S  $\leq 29$ ; mean 25) and moderate anxiety (MOD: STAI-S  $\geq 30$ ; mean 34): the STAI-S ranges for LOW and MOD are in line with those reported in the literature (30, 45). Independent *t*-tests were used to compare the summed dermal responses to DPCP in LOW and MOD in each group

(30MI, 30HI, 120MI and CON). Comparisons of psychological measures between groups (30MI, 30HI, 120MI and CON) were made using one-way ANOVA. A two-way, mixed-model ANOVA was used to analyse DPCP responses across the full dose-series challenge (anxiety level x dose) and circulating stress hormones (anxiety level x time) with significant differences identified using *post hoc* Tukey HSD, where appropriate. Pearson correlation coefficients were also calculated between physiological stress (TRIMP) and the DPCP response, and anxiety and the DPCP response. To determine the influence of anxiety on the threshold DPCP dose that elicits a response, logarithmic transformation was performed on the DPCP data (LOW vs. MOD). This enabled the calculation of the  $x$ -intercept when  $y = 0$ , using linear regression on the linear portion of the dose-response curve. A threshold dose for a response to DPCP was then calculated by back transformation (antilog). Data are presented as mean  $\pm$  SD, unless otherwise stated and statistical significance was accepted at  $P < 0.05$ . Data were checked for normality and where appropriate natural log transformation was performed before analysis. Statistical analyses were performed using common statistical software packages (SPSS 22; IBM, Chicago, IL, and GraphPad Prism 5.0, San Diego, CA). Cohen's  $d$  effect sizes ( $d$ ) are presented to indicate the meaningfulness of group differences for DPCP responses; whereby, values greater than 0.2, 0.5, and 0.8 represent small, medium, and large effects, respectively (9).

## RESULTS

### STAI-S Anxiety

Prior to exercise, there were no differences in psychological measures between groups (e.g. STAI-S scores for 30MI, 30HI, 120MI and CON) and participants reported low-to-moderate STAI-S scores (Fig. 1A). In step 1 of the regression model (Table 1), exercise (TRIMP;  $78 \pm 60$  AU) was a significant predictor accounting for 20% of the variance in the summed dermal thickening response to DPCP ( $P < 0.01$ ); whereby, greater physiological stress was associated with a lower DPCP response following exercise. In step 2, STAI-S score was a significant predictor over and above exercise, accounting for an additional 19% of the variance in DPCP response ( $P < 0.01$ ); together, exercise and anxiety accounted for 39% of the variance in the dermal thickening response to DPCP (Table 1). Pearson correlation coefficients were of comparable, **moderate** strength for the relationship between physiological stress and the summed dermal response to DPCP (TRIMP;  $r = -0.37$ ,  $R^2 = 0.13$ ,  $P = 0.01$ ), and anxiety and the summed dermal response to DPCP (STAI-S score;  $r = 0.39$ ,  $R^2 = 0.15$ ,  $P < 0.01$ ). This association between anxiety before exercise and *in vivo* immunity after an exercise challenge indicates that LOW were more likely to have a lower DPCP response following exercise stress than MOD (Fig. 1B). When reported as the summed response to the five DPCP challenge doses, dermal thickening response was 62% lower in LOW than MOD (LOW  $1.6 \pm 2.3$  and MOD  $4.2 \pm 3.1$  mm;  $P < 0.01$ ;  $d = 1.0$ ).

\*\*\*Table 1 near here\*\*\*

\*\*\*Fig. 1 near here\*\*\*

The ubiquitous influence of anxiety on *in vivo* immunity after exercise challenge (but not rested CON) is further illustrated in the comparisons between LOW and MOD in each group

(30MI, 30HI, 120MI and CON; Fig. 2A-D). Responses to DPCP assessed as skinfold thickness and erythema (data not shown for brevity), were smaller in LOW *vs.* MOD for 30MI ( $P < 0.01$ ) and 30HI ( $P < 0.05$ ; Fig. 2A-B), but not CON. The suppressive effect of LOW *vs.* MOD was also apparent in 120MI ( $P = 0.05$ ;  $d = 0.9$ ; Fig. 2C) which is particularly striking given that the suppressive effect of prolonged exercise on the induction of DPCP immune memory has been reported (16). The lower CHS response to exercise in LOW *vs.* MOD is also illustrated in the smaller dermal thickening response across the full dose-series of DPCP in LOW *vs.* MOD ( $F(1, 35) = 11.1$ ,  $P < 0.01$ ; Fig. 1B for 30MI and 30HI). Furthermore, the threshold dose for a positive response to DPCP was calculated using the linear part of the dose-response curves. Compared with MOD, LOW required a 4-times greater DPCP dose ( $1.5 \mu\text{g}/\text{cm}^2$ ) to elicit a positive response.

\*\*\*Fig. 2 near here\*\*\*

### **Perceived Stress Scale**

Participants reported low-to-moderate PSS scores ( $16.5 \pm 5.3$ ). After accounting for the influence of exercise in step 1 of the regression model (Table 1), PSS score was a significant, **moderate** predictor (in step 2), accounting for an additional 13% of the variance in DPCP response ( $P < 0.05$ ); together, exercise and PSS score accounted for 33% of the variance in the dermal thickening response to DPCP (Table 1). This association between the perception of psychological stress in the last month (i.e. the degree to which life situations are considered stressful) and *in vivo* immunity after exercise challenge indicates that participants reporting lower life stress were more likely to have a lower DPCP response following an exercise challenge than participants reporting moderate life stress.

### **Circulating stress hormones**

When comparing LOW and MOD, a significant anxiety level x time interaction was observed for circulating epinephrine concentration ( $F(2, 88) = 5.9; P < 0.01$ ); whereby, epinephrine was lower in LOW than MOD at pre-exercise (LOW  $0.25 \pm 0.17$  vs. MOD  $0.58 \pm 0.46$  nmol/L;  $P < 0.01$ ), but not different at post or 1 h post-exercise. Similarly, an independent *t*-test showed that circulating cortisol concentration was also lower pre-exercise in LOW than MOD (LOW  $545 \pm 190$  vs. MOD  $699 \pm 289$  nmol/L;  $P < 0.05$ ); albeit, there was no significant interaction. Nevertheless, the lower circulating epinephrine and cortisol concentration in LOW than MOD before exercise represent large ( $d = 0.94$ ) and medium ( $d = 0.63$ ) effects, respectively. Circulating norepinephrine was not different between LOW and MOD.

## DISCUSSION

The aim of this work was to investigate the influence of anxiety and perceived psychological stress on the *in vivo* immune response after exercise. The findings support our hypothesis that the level of anxiety and perceived psychological stress reported by the individual prior to exercise play an important role in determining the strength of the subsequent *in vivo* immune response after exercise (Table 1 and Fig. 1): *in vivo* immunity was assessed by DPCP sensitisation after exercise and recall responses measured 28 d later. Moreover, the findings indicate a similar, moderate strength relationship for the level of anxiety prior to exercise (STAI-S;  $r = 0.39$ ) and the level of physiological stress during exercise (TRIMP;  $r = -0.37$ ) with the *in vivo* immune response after exercise challenge. The ubiquitous influence of anxiety on the immune response after exercise is further evidenced by a lower *in vivo* immune response to DPCP in individuals reporting low compared with moderate anxiety, regardless of the intensity and duration of the exercise challenge (30MI, 30HI and 120MI, Fig. 2A–C). These findings support the recommendation that exercise scientists should account for anxiety and psychological stress when examining the immune response to exercise.

The findings of the present study demonstrate an important interaction between the a priori level of anxiety and perceived psychological stress and the subsequent immune response after an exercise challenge. We previously showed no significant influence of 30MI or 30HI on *in vivo* immunity (16), but these new insights show a lower *in vivo* immune response in individuals reporting low compared with moderate anxiety in 30MI and 30HI (Fig. 2A–B). Moreover, although we have previously shown a suppressive effect of 120MI compared with rested control on *in vivo* immunity (16), particularly striking is the 50% lower *in vivo* immune



response in individuals reporting low compared with moderate anxiety on 120MI (Fig. 2C). Given that DPCP is benign, determining the clinical significance of these findings, with specific regard to infection (skin and other) is an important avenue for future research. Preferably, the strength of the cutaneous recall response to DPCP could be generalised beyond skin immunity to indicate the immune system's general ability to respond to an infectious challenge. The available evidence in this regard is supportive as cutaneous immune measures are impaired in individuals with acute infectious illness (3, 22), diabetes and psoriasis (1) and predict mortality in critically ill HIV-infected patients (17). That we show lower pre-exercise circulating cortisol and epinephrine in the low compared with moderate anxiety group raises the possibility that stress hormones may modulate the immune response to subsequent exercise; indeed, stress hormones are considered to play important roles in preparing the immune system for challenge (13, 15). For example, administration of physiological doses of corticosterone and epinephrine increased T-cell drainage away from the site of DTH challenge to lymph nodes, which in-turn enhanced the DTH response in rats (15). In addition, adrenalectomy has been shown to eliminate stress-induced immune-enhancement in rats, likely by reducing the glucocorticoid and epinephrine response (15). Nevertheless, post-exercise circulating cortisol and epinephrine were not different between individuals reporting low and moderate anxiety in the present study; as such, further research is required into the underlying mechanisms.

Regarding the timing of the psychological measurements, the findings were unlikely due to an acute anticipatory effect prior to exercise as our participants underwent thorough familiarisation to all procedures, including running 50% of their allocated exercise duration; indeed, the success of familiarisation is shown as similar STAI-S scores prior to exercise and rested CON (Fig. 1A). In addition, our findings for the relationship between STAI-S score

and the *in vivo* immune response after exercise are further supported by the relationship between PSS score and the *in vivo* immune response after exercise: PSS assesses the perception of stress, and measures the degree to which life situations spanning the last month are considered stressful (whereas STAI-S provides an acute measure of anxiety) (11). As such, the PSS findings provide added confidence regarding the observed association between psychological stress and the *in vivo* immune response after exercise challenge. It remains to be shown whether individuals are predisposed to respond to stressful situations, such as competitive sport or military scenarios, in a predictable manner with regards to neuro-endocrine-immune responses. In support of this notion, there is some evidence that personality traits predict endocrine-stress-reactivity (4, 19); nevertheless, further research is required to investigate this novel concept in exercise immunology, and to establish whether the findings of the present study extend to other immune measures e.g. vaccination responses (7) and mucosal immunity (23). Further research is also required to disentangle the influence of psychological and physiological strain during prolonged exercise (e.g. during endurance and ultra-endurance events) on *in vivo* immunity. Psychological stress measurements were made before exercise in the present study and it is reasonable to assume that psychological stress during more prolonged exercise (e.g. 120MI) might also play a role in the observed decrease in the *in vivo* immune response (Fig. 2C).

### **Bridging the gap between exercise immunology and psycho-neuro-immunology**

Research investigators have long since acknowledged a role for psychological stress in the decrease in immunity associated with heavy exercise and training but there is little empirical research to support this hypothesis (8, 36). Since Clow and Hucklebridge's Exercise Immunology Review article highlighting this working hypothesis in 2001 (8) there have been

> 3,000 peer-reviewed publications in exercise immunology (using the search terms ‘exercise’ and ‘immune’, Web of Science™) yet < 5% of these publications include the search terms ‘psychological stress’ or ‘anxiety’. Closer inspection of this small subset of exercise immunology publications reveals that the large majority mention a putative role for psychological stress or anxiety in exercise-immune modulation; however, only a small handful of original investigations either attempt to manipulate psychological stress or include objective measures of psychological stress (27, 32, 38, 39). The present study answers the recent calls to physiologists (51) and exercise immunologists (49) to incorporate objective psychological measurements in their human studies.

The findings herein support the recommendation that exercise immunologists should include aspects of mental health (e.g. psychological stress and others), in a broader conceptual framework of exercise-immune interactions alongside other factors thought to decrease immunity in athletes and military personnel (e.g. prolonged training sessions, poor nutrition etc.). This will inform and direct research questions and experimental designs with the aim of improving our understanding of the complicated exercise-immune interactions and with the potential to provide effective countermeasures to immune impairment in those concerned. To this end, the exercise immunologist’s toolkit will be enhanced by joining forces with experts in the ever expanding field of psycho-neuro-immunology to begin to disentangle the psychosocial and physiological underpinning of decreased immunity and increased infection risk in high level athletes, military personnel **and others in physically demanding occupations**. Our finding that pre-exercise anxiety and perceived psychological stress accounted for additional variance in post-exercise *in vivo* immunity after accounting for exercise (using TRIMP) emphasises the importance of incorporating psychological measurements in studies investigating the immune response to exercise. As do the similar strength correlations for pre-

exercise anxiety (STAI-S;  $r = 0.39$ ) and physiological stress during exercise (TRIMP;  $r = -0.37$ ) with *in vivo* immunity after exercise. These findings indicate a beneficial effect of moderate (*vs.* low) anxiety and perceived psychological stress on *in vivo* immunity after exercise (Fig. 2A and D); as such, the findings accord with the immune-enhancement theory of moderate stress (13, 20, 21). Further research is required to investigate exercise-immune responses in athletes, military personnel and others in physically demanding occupations (e.g. firefighters and mountain rescue workers) experiencing higher levels of psychological stress than those reported in this study e.g. as might occur in relation to important competition, major life events etc. The immuno-suppressive effects of chronic high stress in rats (3 weeks of restraint and shaking stress) (14) and humans (examination period) (44) are widely acknowledged (13). As such, research is required to test the hypothesis that chronic high levels of psychological stress exacerbate the decrease in *in vivo* immunity after exercise. Irrespective, the present findings support the recommendation that exercise scientists should account for anxiety and psychological stress when examining the immune response to exercise, and for coaches and support staff to monitor anxiety and psychological stress alongside more traditional physiological measures of training stress. Accordingly, recent evidence highlights that aspects of mental health such as psychological stress and depression are important risk factors for illness in Olympic athletes (18). In time, studies may demonstrate the utility of interventions to alter psychological stress in order to optimise immunity and host defence in athletes, military personnel and those in physically demanding occupations. There is good reason for optimism as an 8-week mindfulness meditation programme increased the antibody response to influenza vaccine in employees working in a highly stressful environment (*vs.* waiting-list controls) (12). Also, although somewhat limited methodologically, preliminary work in competitive athletes showed that a 3-week stress management intervention reduced the number of days out due to illness and injury (35).

## CONCLUSIONS

In conclusion, these findings show that anxiety and perceived psychological stress levels prior to exercise play an important role in determining the strength of the *in vivo* immune response after exercise. Moreover, these findings indicate a similar, moderate strength relationship for the level of state-anxiety prior to exercise and the level of physiological stress during exercise with the *in vivo* immune response after exercise. Future research is required to investigate exercise-immune responses in athletes and others in physically demanding occupations experiencing higher levels of psychological stress than those reported in this study e.g. related to important competition and major life events. Nevertheless, these findings support the recommendation that exercise scientists should account for anxiety and psychological stress when examining the immune response to exercise.

## ACKNOWLEDGEMENTS

Bethany Diment received a 125<sup>th</sup> anniversary research scholarship from Bangor University.

This study received no external funding. None of the authors had a conflict of interest.

## REFERENCES

1. Bangsgaard N, Engkilde K, Menne T, Lovendorf M, Jacobsen GK, Olsen J, and Skov L. Impaired hapten sensitization in patients with autoimmune disease. *Clin Exp Immunol* 165: 310-317, 2011.
2. Banister EW. Modelling elite athletic performance. In: *Physiological testing of elite athletes*, edited by Green H, McDougal J, and Wenger H. Champaign, IL: Human Kinetics, 1991, p. 403-424.
3. Bennett BK, Hickie IB, Vollmer-Conna US, Quigley B, Brennan CM, Wakefield D, Douglas MP, Hansen GR, Tahmindjis AJ, and Lloyd AR. The relationship between fatigue, psychological and immunological variables in acute infectious illness. *Aust N Z J Psychiatry* 32: 180-186, 1998.
4. Bibbey A, Carroll D, Roseboom TJ, Phillips AC, and de Rooij SR. Personality and physiological reactions to acute psychological stress. *Int J Psychophysiol* 90: 28-36, 2013.
5. Booth CK, Coad RA, Forbes-Ewan CH, Thomson GF, and Niro PJ. The physiological and psychological effects of combat ration feeding during a 12-day training exercise in the tropics. *Mil Med* 168: 63-70, 2003.
6. Bruunsgaard H, Hartkopp A, Mohr T, Konradsen H, Heron I, Mordhorst CH, and Pedersen BK. In vivo cell-mediated immunity and vaccination response following prolonged, intense exercise. *Med Sci Sports Exerc* 29: 1176-1181, 1997.
7. Burns VE. Using vaccinations to assess in vivo immune function in psychoneuroimmunology. *Methods Mol Biol* 934: 371-381, 2012.
8. Clow A, and Hucklebridge F. The impact of psychological stress on immune function in the athletic population. *Exerc Immunol Rev* 7: 5-17, 2001.
9. Cohen J. *Statistical power analysis for the behavioral sciences*. Hillsdale, NJ: Erlbaum, 1988, p. 79-80.
10. Cohen S, Tyrrell DA, and Smith AP. Psychological stress and susceptibility to the common cold. *N Engl J Med* 325: 606-612, 1991.
11. Cohen S, and Williamson G. Psychological stress in a probability sample of the United States. In: *The social psychology of health*, edited by Spacapan S, and Oskamp S. Newbury Park, CA: Sage, 1988, p. 31-67.
12. Davidson RJ, Kabat-Zinn J, Schumacher J, Rosenkranz M, Muller D, Santorelli SF, Urbanowski F, Harrington A, Bonus K, and Sheridan JF. Alterations in brain and immune function produced by mindfulness meditation. *Psychosom Med* 65: 564-570, 2003.
13. Dhabhar FS. Effects of stress on immune function: the good, the bad, and the beautiful. *Immunol Res* 58: 193-210, 2014.
14. Dhabhar FS, and McEwen BS. Acute stress enhances while chronic stress suppresses cell-mediated immunity in vivo: a potential role for leukocyte trafficking. *Brain Behav Immun* 11: 286-306, 1997.
15. Dhabhar FS, and McEwen BS. Enhancing versus suppressive effects of stress hormones on skin immune function. *Proc Natl Acad Sci U S A* 96: 1059-1064, 1999.
16. Diment BC, Fortes MB, Edwards JP, Hanstock HG, Ward MD, Dunstall HM, Friedmann PS, and Walsh NP. Exercise intensity and duration effects on in vivo immunity. *Med Sci Sports Exerc* 47: 1390-1398, 2015.
17. Dolan MJ, Clerici M, Blatt SP, Hendrix CW, Melcher GP, Boswell RN, Freeman TM, Ward W, Hensley R, and Shearer GM. In vitro T cell function, delayed-type hypersensitivity skin testing, and CD4+ T cell subset phenotyping independently

- predict survival time in patients infected with human immunodeficiency virus. *J Infect Dis* 172: 79-87, 1995.
18. Drew MK, Vlahovich N, Hughes D, Appaneal R, Peterson K, Burke L, Lundy B, Toomey M, Watts D, Lovell G, Praet S, Halson S, Colbey C, Manzanero S, Welvaert M, West N, Pyne DB, and Waddington G. A multifactorial evaluation of illness risk factors in athletes preparing for the Summer Olympic Games. *J Sci Med Sport* 20: 745-750, 2017.
  19. Edelstein RS, Yim IS, and Quas JA. Narcissism predicts heightened cortisol reactivity to a psychosocial stressor in men. *J Res Pers* 44: 565-572, 2010.
  20. Edwards KM, Burns VE, Carroll D, Drayson M, and Ring C. The acute stress-induced immunoenhancement hypothesis. *Exerc Sport Sci Rev* 35: 150-155, 2007.
  21. Edwards KM, Burns VE, Reynolds T, Carroll D, Drayson M, and Ring C. Acute stress exposure prior to influenza vaccination enhances antibody response in women. *Brain Behav Immun* 20: 159-168, 2006.
  22. Haider S, Coutinho Mde L, Emond RT, and Sutton RN. Tuberculin anergy and infectious mononucleosis. *Lancet* 2: 74, 1973.
  23. Hanstock HG, Walsh NP, Edwards JP, Fortes MB, Cosby SL, Nugent A, Curran T, Coyle PV, Ward MD, and Yong XH. Tear fluid SIgA as a noninvasive biomarker of mucosal immunity and common cold risk. *Med Sci Sport Exer* 48: 569-577, 2016.
  24. Harper Smith AD, Coakley SL, Ward MD, Macfarlane AW, Friedmann PS, and Walsh NP. Exercise-induced stress inhibits both the induction and elicitation phases of in vivo T-cell-mediated immune responses in humans. *Brain Behav Immun* 25: 1136-1142, 2011.
  25. Hellard P, Avalos M, Guimaraes F, Toussaint JF, and Pyne DB. Training-related risk of common illnesses in elite swimmers over a 4-yr period. *Med Sci Sports Exerc* 47: 698-707, 2015.
  26. Hu E, Koucky EM, Brown WJ, Bruce SE, and Sheline YI. The role of rumination in elevating perceived stress in posttraumatic stress disorder. *J Interpers Violence* 29: 1953-1962, 2014.
  27. Huang CJ, Webb HE, Garten RS, Kamimori GH, and Acevedo EO. Psychological stress during exercise: lymphocyte subset redistribution in firefighters. *Physiol Behav* 101: 320-326, 2010.
  28. Ingram LA, Simpson RJ, Malone E, and Florida-James GD. Sleep disruption and its effect on lymphocyte redeployment following an acute bout of exercise. *Brain Behav Immun* 47: 100-108, 2015.
  29. Jansky L, Pospisilova D, Honzova S, Ulicny B, Sramek P, Zeman V, and Kaminkova J. Immune system of cold-exposed and cold-adapted humans. *Eur J Appl Physiol Occup Physiol* 72: 445-450, 1996.
  30. Julian LJ. Measures of anxiety: State-Trait Anxiety Inventory (STAI), Beck Anxiety Inventory (BAI), and Hospital Anxiety and Depression Scale-Anxiety (HADS-A). *Arthritis Care Res (Hoboken)* 63 Suppl 11: S467-472, 2011.
  31. Laing SJ, Oliver SJ, Wilson S, Walters R, Bilzon JL, and Walsh NP. Neutrophil-degranulation and lymphocyte-subset response after 48 hr of fluid and/or energy restriction. *Int J Sport Nutr Exerc Metab* 18: 443-456, 2008.
  32. Moreira A, Arsati F, Lima-Arsati YB, Simoes AC, and De Araujo VC. Monitoring stress tolerance and occurrences of upper respiratory illness in basketball players by means of psychometric tools and salivary biomarkers. *Stress Health* 27: e166-e172, 2011.

33. Nieman DC, Johanssen LM, Lee JW, and Arabatzis K. Infectious episodes in runners before and after the Los Angeles Marathon. *J Sports Med Phys Fitness* 30: 316-328, 1990.
34. Pascoe AR, Singh MAF, and Edwards KM. The effects of exercise on vaccination responses: a review of chronic and acute exercise interventions in humans. *Brain Behav Immun* 39: 33-41, 2014.
35. Perna FM, Antoni MH, Baum A, Gordon P, and Schneiderman N. Cognitive behavioral stress management effects on injury and illness among competitive athletes: a randomized clinical trial. *Ann Behav Med* 25: 66-73, 2003.
36. Perna FM, Schneiderman N, and LaPerriere A. Psychological stress, exercise and immunity. *Int J Sports Med* 18 Suppl 1: S78-83, 1997.
37. Peters EM, and Bateman ED. Ultramarathon running and upper respiratory tract infections. An epidemiological survey. *S Afr Med J* 64: 582-584, 1983.
38. Rehm KE, Elci OU, Hahn K, and Marshall GD. The impact of self-reported psychological stress levels on changes to peripheral blood immune biomarkers in recreational marathon runners during training and recovery. *Neuroimmunomodulation* 20: 164-176, 2013.
39. Rehm KE, Sunesara I, Tull MT, and Marshall GD. Psychological stress moderates the relationship between running volume and cd4+ T cell subpopulations. *J Biol Reg Homeos Ag* 30: 449-457, 2016.
40. Severs Y, Brenner I, Shek PN, and Shephard RJ. Effects of heat and intermittent exercise on leukocyte and sub-population cell counts. *Eur J Appl Physiol Occup Physiol* 74: 234-245, 1996.
41. Shephard RJ. Immune changes induced by exercise in an adverse environment. *Can J Physiol Pharmacol* 76: 539-546, 1998.
42. Shephard RJ, and Shek PN. Heavy exercise, nutrition and immune function: is there a connection? *Int J Sports Med* 16: 491-497, 1995.
43. Shephard RJ, and Shek PN. Interactions between sleep, other body rhythms, immune responses, and exercise. *Can J Appl Physiol* 22: 95-116, 1997.
44. Smith A, Vollmer-Conna U, Bennett B, Wakefield D, Hickie I, and Lloyd A. The relationship between distress and the development of a primary immune response to a novel antigen. *Brain Behav Immun* 18: 65-75, 2004.
45. Spielberger CD. *Manual for the State-Trait Anxiety Inventory*. Palo Alto, CA: Consulting Psychologists Press, 1983, p. 12-14.
46. Tabachnick BG, and Fidell LS. *Using multivariate statistics*. Essex, UK: Pearson Education, 2013, p. 159-160.
47. Walsh NP, Gleeson M, Pyne DB, Nieman DC, Dhabhar FS, Shephard RJ, Oliver SJ, Bermon S, and Kajeniene A. Position statement. Part two: maintaining immune health. *Exerc Immunol Rev* 17: 64-103, 2011.
48. Walsh NP, Gleeson M, Shephard RJ, Gleeson M, Woods JA, Bishop NC, Fleshner M, Green C, Pedersen BK, Hoffman-Goetz L, Rogers CJ, Northoff H, Abbasi A, and Simon P. Position statement. Part one: immune function and exercise. *Exerc Immunol Rev* 17: 6-63, 2011.
49. Walsh NP, and Oliver SJ. Exercise, immune function and respiratory infection: an update on the influence of training and environmental stress. *Immunol Cell Biol* 94: 132-139, 2016.
50. Wang JS, Chen WL, and Weng TP. Hypoxic exercise training reduces senescent T-lymphocyte subsets in blood. *Brain Behav Immun* 25: 270-278, 2011.
51. Wehrwein EA, and Carter JR. The mind matters: psychology as an overlooked variable within physiology studies. *Physiology (Bethesda)* 31: 74-75, 2016.



## FIGURE LEGENDS

**FIGURE 1.** Effect of state-anxiety prior to exercise on the *in vivo* immune response after exercise. (A) Low (LOW) and moderate (MOD) levels of anxiety. Data are Mean  $\pm$  SD. (B) Contact hypersensitivity (CHS) assessed as elicitation challenge 28 d after DPCP induction. Dermal thickening response to the full dose-series challenge with DPCP is shown (30MI and 30HI). Data are Mean  $\pm$  SEM for clarity. <sup>1</sup>Shown for comparison.

**FIGURE 2.** Effect of state-anxiety prior to exercise on the *in vivo* immune response after exercise of varying intensity and duration. (A–D) Summed increase in skinfold thickening response to DPCP challenge for each exercise group (30MI, 30HI and 120MI) and rested CON. Data are Mean  $\pm$  SD.

**TABLE 1.** Multiple linear regression analysis examining the influence of state-anxiety and perceived psychological stress level prior to exercise on the subsequent *in vivo* immune response after exercise. Contact hypersensitivity (CHS) assessed as the summed dermal thickening response to the full dose-series elicitation challenge with DPCP 28 d after DPCP induction. After accounting for the **negative** influence of exercise in step 1, separate models show the **positive** influence of anxiety (from low to moderate levels), assessed using STAI-S in step 2 (A) and perceived psychological stress (from low to moderate levels) over the last month, assessed using PSS in step 2 (B), respectively.

<i>Dependent variable: CHS</i>	<b>B</b>	<b>SE</b>	$\beta$	<i>t</i>	$\Delta F$	$R^2$	$\Delta R^2$
<b>A. Step 1</b>							
Exercise (TRIMP) <sup>1</sup>	-0.005	0.002	-0.44	-2.93	8.56	0.20**	0.20**
<b>Step 2</b>							
STAI-S	0.06	0.02	0.44	3.24	10.50	0.39**	0.19**
<b>B. Step 1</b>							
Exercise (TRIMP) <sup>1</sup>	-0.005	0.002	-0.44	-2.93	8.56	0.20**	0.20**
<b>Step 2</b>							
PSS	0.06	0.02	0.36	2.54	6.45	0.33**	0.13*

<sup>1</sup>TRIMP = training impulse; STAI-S = State Trait Anxiety Inventory; PSS = Perceived Stress Scale; \*  $P < 0.05$ ; \*\*  $P < 0.01$ .

FIGURE 1.




FIGURE 2.

