Human parahippocampal cortex supports spatial binding in visual working memory

Dundon, Neil; Katshu, Mohammad Zia Ul; Harry, Bronson; Roberts, Daniel; Leek, Charles; Downing, Paul; Sapir, Ayelet; Roberts, Craig; D-Avossa, Giovanni

Cerebral Cortex

DOI:
[10.1093/cercor/bhx231](https://doi.org/10.1093/cercor/bhx231)

Published: 15/09/2017

Peer reviewed version

[Cyswllt i'r cyhoeddiad / Link to publication](#)

Dyfyniad o'r fersiwn a gyhoeddwyd / Citation for published version (APA):

Dundon, N., Katshu, M. Z. U., Harry, B., Roberts, D., Leek, C., Downing, P., Sapir, A., Roberts, C., & D-Avossa, G. (2017). Human parahippocampal cortex supports spatial binding in visual working memory. *Cerebral Cortex*, 2017, 1-11. <https://doi.org/10.1093/cercor/bhx231>

Hawliau Cyffredinol / General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

26

Abstract

27 Studies investigating the functional organisation of the medial temporal lobe (MTL)
28 suggest that parahippocampal cortex (PHC) generates representations of spatial and
29 contextual information used by the hippocampus in the formation of episodic
30 memories. However, evidence from animal studies also implicates PHC in spatial
31 binding of visual information held in short term, working memory. Here we examined
32 a 46-year-old man (PJ), after he had recovered from bilateral medial occipitotemporal
33 cortex strokes resulting in ischemic lesions of PHC and hippocampal atrophy, and a
34 group of age-matched healthy controls. When recalling the colour of one of two
35 objects, PJ misidentified the target when cued by its location, but not shape. When
36 recalling the position of one of three objects, he frequently misidentified the target,
37 which was cued by its colour. Increasing the duration of the memory delay had no
38 impact on the proportion of binding errors, but did significantly worsen recall
39 precision in both PJ and controls. We conclude that PHC may play a crucial role in
40 spatial binding during encoding of visual information in working memory.

41

42 **Keywords:** Feature binding; Medial temporal lobe; Parahippocampal cortex; Spatial
43 Memory; Visual working memory

44

Introduction

45 The medial temporal lobe (MTL) comprises the hippocampus and parahippocampal
46 regions, i.e., entorhinal cortex, perirhinal cortex (PRC) and parahippocampal cortex
47 (PHC). These structures play a prominent role in episodic memory, as evidenced by
48 the dense anterograde amnesia, which follows damage to MTL (Scoville and Milner
49 1957; Corkin 1984; Corkin et al. 1997). Modular accounts of MTL function have
50 suggested that the hippocampus synthesises episodic memories by binding
51 information about the identity and location of objects carried respectively by two
52 different streams (Eichenbaum et al. 2007; Diana et al. 2007).

53

54 MTL structures have also been implicated in short term memory processes
55 (Ranganath and Blumenfeld 2005; Graham et al. 2010; Yonelinas, 2013). First,
56 animal models have pointed to specific molecular mechanisms in the mammalian
57 MTL dedicated to the storage of short term memories, and separate from those
58 involved in long term memory (Deacon et al. 2002; Reisel et al. 2002). Single unit
59 recordings and lesion studies in non-human primates have further demonstrated that
60 the hippocampus (Friedman and Goldman-Rakic 1988), entorhinal cortex (Suzuki et
61 al. 1997), PRC (Davachi and Goldman-Rakic 2001) and PHC (Bachevalier and
62 Nemanic 2008) contribute to the encoding and recall of information from short term
63 memory. These animal findings complement neuropsychological studies of patients
64 with amnesia resulting from Korsakoff's Syndrome, encephalitis and colloid cysts
65 (Holdstock et al. 1995), and patients with surgical (Aggleton 1992; Owen et al. 1995)
66 or ischemic (Holdstock et al. 2002) lesions to the MTL, demonstrating retention
67 deficits for novel stimuli over delay intervals as short as two seconds (Ranganath
68 and Blumenfeld 2005).

69

70 An increasing body of evidence further suggests that short term memory exploits the
71 same MTL modules as episodic memory; that is, PRC codes information about an
72 object's identity and PHC codes an object's location and its context, and these two
73 streams are bound in the hippocampus (Pertsov et al. 2013; Watson et al. 2013; Yee et
74 al. 2014; Libby et al. 2014). Consistent with the idea that in short term memory
75 identity and location information are processed separately and then bound, patients
76 with hippocampal damage can exhibit deficits recalling object-location conjunctions
77 after 1.0s delays, even when unimpaired recalling either object identities or locations
78 (Olson et al. 2006a; 2006b). However, other studies report that patients with damage
79 to the hippocampus do not necessarily show deficits in recalling object-location
80 conjunctions, suggesting that spatial binding is preserved (e.g. Jeneson et al. 2010; see
81 Yonelinas 2013 for a review).

82

83 An alternative possibility is that spatial binding in short term memory occurs in
84 parahippocampal regions, rather than the hippocampus proper. In support of this
85 view, data in both rats (Burwell and Amaral 1998) and monkeys (Suzuki and Amaral
86 1994) indicate that PRC and PHC are reciprocally connected, suggesting that the
87 parcellation of identity and spatial information is not absolute, and that there may
88 already be substantial cross-talk between object and spatial/context related
89 information in parahippocampal regions. Further, recordings in rats have
90 demonstrated single unit responses for object-location conjunctions in the PHC
91 homologue (Barker and Warburton 2011).

92

Behavioural studies in monkeys have provided crucial evidence for the role of PHC in spatial binding. Rhesus monkeys with PHC lesions are impaired in both simple location and object-location conjunction tasks (Malkova and Mishkin, 2003). This short term memory impairment was observed in a delayed match-to-sample task, where the sample contained two non-identical objects. After a six-second delay, the test array contained one of the objects in its original location (the target), and an identical item either at the location of the sample foil (object-place condition), or at a novel location not previously occupied by either sample object (location condition). Monkeys with PHC lesions were impaired identifying the target in both conditions, while monkeys with lesions in the hippocampus showed no impairment in either task (Malkova and Mishkin 2003). Hippocampectomised monkeys were likewise unimpaired in a later study, using a more difficult task with an increased number of objects and locations (Belcher et al. 2006).

A cross-species homology in the short term memory functionality of PHC is partly supported by the observation that patients with PHC lesions also exhibit a decrement in spatial recall (Ploner et al. 2000), although this impairment is only observed using delays greater (i.e. >15.0s) than those used by Malkova and Mishkin (2003). In addition, functional imaging data in healthy subjects demonstrate heightened right PHC activation during both encoding and maintenance of object-location conjunctions, relative to trials where objects or locations are memorised separately (Luck et al. 2010). However, no neuropsychological study has so far demonstrated that PHC contributes to spatial binding in human short term memory.

117 In the present study, we examined the nature and extent of spatial and short term
118 memory deficits associated with focal PHC lesions, by testing a middle-aged man (PJ)
119 with bilateral posterior circulation strokes involving the PHC, but sparing the
120 hippocampus and PRC. Our experiments were driven by three specific research
121 questions: 1) does damage to PHC produce binding difficulties and if so, are the
122 binding problems specifically spatial or do they generalise to other visual dimensions;
123 2) do binding impairments reflect deficits in memory encoding or maintenance; and
124 3) is the binding impairment secondary to a loss of positional information either in
125 memory or perception?

126
127 Both PJ and controls showed dependent decrements in the precision of spatial recall,
128 however PJ's recall precision was significantly worse than controls at longer delays
129 (5.0s). PJ also showed impaired spatial binding. This impairment was unaffected by
130 the duration of the memory delay. Finally, PJ's binding deficits did not generalise
131 across visual dimensions, since he performed normally when recall involved the
132 conjunction of non-spatial features. We conclude that PHC serves a spatially specific
133 binding function in short term memory, and that this function appears to be
134 independent of PHC's role in recall precision.

135

Methods

PJ: history and clinical assessment

PJ was first seen by one of the authors (CR), four months after he had suffered a cerebrovascular accident. PJ was 45 years old when he developed headaches, visual and mental status changes over the course of a few hours. Two days after the onset of these symptoms, he was admitted to a stroke-unit at a regional hospital. During the admission, he continued to be confused and agitated. The diagnostic work-up revealed bilateral posterior circulation strokes involving the occipito-temporal cortex. No cause for the stroke was identified. PJ had no significant medical history, except for cluster headaches, which responded well to standard treatment.

Upon returning home, he was not able to resume his full-time occupation as an animal breeder, because of difficulties finding his way around the house and farm, where he had moved two years prior. He also relinquished driving, because he could not find his way around familiar streets. He was able to sketch the overall layout of his home, but frequently misidentified rooms and the family resorted to placing signs on internal doors to help him find his way around. His ability to repair equipment around the farm was also diminished, because of difficulty identifying the correct tool in a cluttered environment.

PJ's visual perimetry was formally assessed three and five months following the ischemic injury, with a binocular field test (Esterman, 1982). He showed strict upper quadrantanopias, worse on the left than on the right. There was evidence of partial recovery on the second assessment (see figure S3).

Formal clinical psychometric testing was conducted approximately 6 months following his stroke. The standardised scores are presented in supplementary table 1. His general intellectual functioning fell within the average range, as measured with the Wechsler Adult Intelligence scale, fourth edition (WAIS-IV). This was affected negatively by slowed processing speed on visual tasks. He performed similarly on the verbal (Verbal Comprehension Index) and non-verbal scale (Perceptual Reasoning Index) of the WAIS-IV. His expressive and receptive language functions were grossly intact. He did however often require verbal instructions to be repeated. His information-processing speed was in the borderline range on the WAIS-IV. Memory function was significantly impaired for both visual and verbal material. He had difficulties with learning and acquisition of new material and also with delayed recall. Performance was not improved for recognition memory. His errors on a visual memory task were primarily misplacement errors. He demonstrated set-loss errors on a word generation task and also required reminding of rules on a problem-solving task. Performance on executive functioning tasks was mixed; he performed at the expected level on a planning and problem-solving task. His performance on a verbal fluency task was within normal limits. His score on an attention-shifting and inhibition task was in the impaired range of ability. PJ passed on all subtests of object perception from the Visual Object and Space Perception Battery (Warrington and James 1991), except for progressive silhouettes, where he had a raw score of 11, indicating mild impairment. He was also faultless in all subtests of space perception.

PJ was scanned using a research MRI protocol and tested behaviourally at the Bangor University School of Psychology approximately one year and ten months following the ischemic event, when he was 47 years of age. Testing took place on two

consecutive days.

Control Participants

Behavioural comparison: Ten right-handed, healthy male participants were recruited from the local community. Controls were screened for any history of major neuropsychiatric disorders and visual impairments. IQ was measured with the 2-subtest (vocabulary and matrix reasoning) version of the Wechsler Abbreviated Scale of Intelligence (WASI; Wechsler 1999). Supplementary Table 2 summarises the characteristics of the control group. The mean age was 48.2 years (sd: 6.4), the mean IQ was 101.1 (sd: 7.6) and the mean age leaving school was 16.6 (sd: 0.7). On all these variables, PJ and controls were matched; all p-values were above .095 using a modified t-test (Crawford and Howell 1998).

Anatomical comparison: A convenience sample of 10 healthy male participants was drawn from a Bangor University image register. The mean age was 43.3 years (sd: 4.9).

All participants were compensated for their time and travel expenses. All participants gave written, informed consent prior to initiating any experimental procedure. The testing procedures had been reviewed and approved by the Betsi Cadwaladr University Health Board and the Bangor University School Psychology Ethics committees.

Behavioural testing: overview and material

210 PJ and controls performed three computer-based behavioural experiments. Testing
211 took place in a dark room; participants sat comfortably, unrestrained, approximately
212 85cm from an LCD screen (NEC LCD3210). Participants were encouraged to actively
213 scan the display and foveate individual stimuli. Custom-coded Matlab scripts
214 (Mathworks 2014a), using a set of freely available routines designed to facilitate the
215 coding of visual experiments (Brainard 1997), controlled the experiments and
216 generated the displays. Matlab scripts were run on an Apple iMac 10.

217

218 Statistical comparison of PJ and controls

219 We computed the significance of performance differences between PJ and the control
220 group in all experiments using a modified t-test (Crawford and Howell 1998). Where
221 performance was measured with a percentage or ratio, we conducted the t-test on
222 logarithmically transformed values.

223

224 **Imaging**

225 Imaging – image acquisition and analysis

226 PJ and the anatomical comparison controls were scanned on a Phillips Achieva 3T
227 MR scanner with a 32-channel head coil. T1 weighted images (TE = 4.32ms; 8° flip
228 angle) were acquired axially with a 0.7mm isotropic voxel-size. PJ's T1 weighted
229 anatomical volume was bias corrected and normalised to the atlas representative
230 MNI152 template using SPM12 (Ashburner and Friston 2003). The mapping included
231 a 12-degrees-of-freedom affine transform followed by a local deformation, computed
232 after the lesion had been masked using a hand-drawn region. The normalised anatomy
233 was obtained by interpolation via a 4th degree B-spline, and resampled using a 0.7mm
234 linear voxel size. Skull stripped anatomy was obtained using a modified version of

FSL's BET, which is optimised for tissue segmentation in the presence of brain pathology (Lutkenhoff et al. 2014). To determine whether PJ's stroke encroached onto perirhinal and entorhinal cortex, probabilistic maps of these regions were superimposed on his brain anatomy (Hindy and Turk-Browne 2016). Lesion boundaries were drawn by a board-certified adult neurologist, using the co-registered T1 and FLAIR images.

Lesion anatomy results

Figure 1 shows axial and coronal slices from the MNI Atlas co-registered T1-weighted scan of PJ's brain. In the left hemisphere the lesion volume is 6.25 cm³, in the right hemisphere 10.71 cm³. Figure 1A shows that the ischemic lesions in medial occipitotemporal cortex (mOTC) of the left and right hemisphere lie posterior to the location of entorhinal and perirhinal cortex (marked respectively in red and green), identified in a previous group study (Hindy and Turk-Browne 2016). Figure S1 provides additional anatomical information about the relationship between lesion and entorhinal and perirhinal cortex. The coronal slices in figure 1B demonstrate that the fornix is intact, however sections -23 to -32 suggest hippocampal volume loss on the right. Also, retrosplenial cortex and the adjacent precuneus are spared in both hemispheres. Figure S2 shows sagittal slices through medial brain structures, which highlights the extent of the damage to PHC and lingual gyrus. Given the apparent hippocampal volume loss, we compared PJ's left and right hippocampal volumes to those of the anatomical comparison controls. A stereological procedure was used to estimate hippocampal volumes in all participants (Keller and Roberts 2009). The input images were the T1 weighted brain volumes in native scanner space. A regular cubic grid with a step of 3 pixels was superimposed on coronal slices, with a random

starting position. The senior author, a board-certified neurologist, outlined the hippocampal formation to determine the number of overlaying grid points. The hippocampal formation included the hippocampus, dentate gyrus and subiculum. The anterior border of the hippocampal formation was the alveus, the posterior border was the crux of the fornix. The hippocampal borders were also identified in axial and sagittal slices. The procedure was implemented using ImageJ (Schneider et al. 2012) and a stereology dedicated plugin (Merzin 2008). This analysis indicated that PJ's left (3931mm^3) and right (2530mm^3) hippocampi were not significantly smaller than controls (left: mean = 3561mm^3 ; $t(9) = 0.516$, $p = 0.618$; right: mean = 3816mm^3 $t(9) = -1.79$, $p = 0.108$). However, the volumetric difference between the left and right hippocampi was significantly greater for PJ than for controls ($t(9) = 2.641$, $p = 0.027$), suggesting that PJ's right hippocampus may have been atrophied.

Experiment 1: spatial vs. non-spatial binding in working memory

Experiment 1 – Rationale

Primate studies (Malkova and Mishkin 2003; Belcher et al. 2006) have suggested that PHC is involved in remembering locations in close peri-personal space as well as spatial binding in working memory. In this first experiment, we examined visual working memory spatial and feature binding in PJ, a man with PHC lesions, and a group of age-matched controls. On each trial, participants had to remember the colour, shape and location of two objects. After a short delay, participants were cued to recall the colour of one of the objects, identified either by its location on the screen, or by its shape. We reasoned that if human PHC is involved in spatial binding, then PJ's recall performance should be worse than controls, specifically on location trials.

Experiment 1 – Methods

Figure 2A shows a schematic representation of Experiment 1's trial structure. In each trial, an equilateral triangle and a square, whose side lengths were 2.42° and 1.72° respectively, appeared side-to-side in the lower half of the screen, at an eccentricity of 4.25° along the main diagonal, for 2.0s. The shapes were either red, blue or green. A 200ms pattern mask, and then a 2.0s blank screen, followed the sample display. The recall screen contained three coloured rectangles, 1.0° wide and 3.0° high, whose lower edges were aligned 2.5° above the screen center and spaced horizontally 9.0° apart. A bright cross (location cue) or the outline of one of the two shapes (shape cue) identified the target. The location cues, which also included a dark cross, appeared at the locations occupied by the two shapes. The shape cue appeared 3.0° below the screen center. Participants reported the target colour by placing a cursor over the corresponding coloured rectangle and clicking the mouse button. The mouse click prompted the beginning of a new trial, after a 1.0s delay, during which the screen was blank. Participants practiced the task over ten trials and then completed ninety trials, including both shape and location cued recalls. Trial order was randomised, minimising participants' ability to predict whether a shape or location cue would follow the sample display. To ensure that PJ had not forgotten the task instructions, we asked him to describe what he had been doing after each block. In each instance he correctly reported that he had been recalling either the probed shape colour, or the colour at the location of the white cross.

Experiment 1 – Data analysis

We scored trials based on whether participants reported (a) the correct target colour (correct response), (b) the colour of the non-target shape (binding error), or (c) neither

the target nor the non-target colour, i.e., dummy colour (generic error). We then calculated the proportion of binding (BE) and generic errors (GE) for each cue condition (location and shape) and compared PJ and the control group's recall accuracy using odds ratios. We computed two odds ratios: the first was the ratio of the proportion of binding errors in location vs. shape cued trials (i.e., $[BE_{\text{location}} / BE_{\text{shape}}]$). The second was the ratio of binding errors over generic errors in location vs. shape cued trials (i.e., $[BE_{\text{location}} / GE_{\text{location}}] / [BE_{\text{shape}} / GE_{\text{shape}}]$). If a participant's data cells contained zero counts, a value of 0.5 was added to all cells prior to computing the ratios (Gart and Zweifel 1967).

Experiment 1 – Results: impaired spatial binding in visual working memory

The left-hand panels of figures 2B and 2C report the proportion of generic errors following location and shape cues, while the right-hand panels show the proportion of binding errors. PJ made more binding errors when the target was identified by a location than a shape cue ($p < 0.001$; Fisher exact test). PJ was also much more likely to make a binding than a generic error following a location ($p < 0.001$, two-tailed binomial test), but not a shape cue ($p = 0.5$), suggesting that his difficulties did not reflect a problem remembering which colours had been shown. For PJ, the odds ratio of making a binding error in the location vs. shape cue trials was 60.7, which was significantly greater than the control group average of 0.501 (95% CI: [0.23 - 1.06], $t(9) = 3.72$, $p = 0.005$), suggesting that he was much more likely to make a binding error on location than shape cue trials, while controls were modestly more accurate following a location than a shape cue. Moreover, PJ's odds ratio of making a binding rather than a generic error in the location vs shape cued trials was 29.0 which was again significantly greater than the control group average of 0.421 (95% CI: [0.21 -

0.83], $t(9) = 3.46$, $p = 0.007$), confirming that he was much more likely to make a binding than a generic error on location rather than shape cued trials, while controls were more likely to make a binding than a generic error on shape rather than location trials.

Experiment 1: Interim discussion

PJ showed a remarkable deficit binding objects to their location in a working memory task. When he reported the colour of one of two objects, he was able to do so accurately for targets cued by their shape. However, when a target was identified by its location, his performance was greatly diminished because of numerous binding errors. Control participants, on the other hand, showed comparable recall accuracy irrespective of the cue type. These findings strongly suggest that PJ's impairment cannot be attributed to either diminished memory for the report feature, i.e. the target's colour, or a binding deficit that generalises across visual dimensions. Rather, PJ shows a binding impairment that is specifically spatial.

Experiment 2: delayed spatial recall

Experiment 2 – Rationale

In the previous experiment, we demonstrated that PJ suffers a specific spatial binding impairment in a working memory task. In experiment 2, we examined whether spatial binding impairments reflect diminished resolution of spatial data in working memory, or rather disruption of spatial binding. To this end we assessed the effects of the duration of the memory delay on both the precision of spatial recall and the proportion of binding errors.

Experiment 2 – Methods

Figure 3A summarises Experiment 2's trial structure. The sample stimulus consisted of three coloured discs, 0.8° in diameter. The discs were red, green and blue, and remained visible for 2.0s. A 1.0s long pattern mask followed the sample. A central colour cue (a 0.3° wide square) appeared either immediately after the pattern mask, or after an additional 4.0s interval, during which only a white central fixation point was visible. The cue identified the target of the same colour. The participants placed the cursor at the recalled target location and clicked the mouse to record their response and initiate the next trial. The location of the discs included the center of the screen and the vertices of a virtual square, at an eccentricity of 6.0° . 2D Gaussian displacement (s.d. = 0.9°) jittered the position of each disc. Each participant completed two blocks of one hundred and twenty trials each.

Experiment 2 – Data analysis

First, we identified trials in which participants had made a binding error, i.e. when the recalled position was closer to the one of the non-target items than the target, and the distance from the non-target item was no greater than half the minimum distance between canonical locations, i.e. 3.0° (Pertzov et al. 2013). After tabulating and removing binding errors, we estimated recall accuracy and precision. Accuracy reflects how close a participant's average reported location is to the true target position. Precision reflects the magnitude of trial-to-trial deviations from a participant's average reported location. Accuracy is diminished by systematic errors, which depend on factors such as display size and memory load (Katshu and d'Avossa 2014), while precision is thought to reflect the resolution of spatial memory (Bays et al. 2009). These two variables were computed using linear regressions. We computed

two regressions whose dependent variables were the azimuth and elevation of the reported target location, respectively. The regressors in each case included a constant and the target's azimuth and elevation. The results of the regression analysis were used to estimate the systematic biases reporting the target location. The scaling factor was the divergence of the error field, which we previously found to be the main linear component of the systematic error (Katshu and d'Avossa 2014). We quantified recall precision using the standard deviation of the residuals from the model fits. The variance and standard deviations of the variable errors were computed using the same procedure employed in a previous study (Katshu and d'Avossa 2014), and averaged over azimuth and elevation. Precision changes between short and long delays were quantified using an efficiency measure, namely a ratio whose numerator was the recall variance following 1.0s delays and denominator was recall variance following 5.0s delays.

Experiment 2 – Results: recall precision, but not binding errors, affected by memory delay

PJ made more binding errors than controls, following both 1.0s and 5.0s delays. Otherwise, both PJ and controls performed similarly in terms of accuracy and precision.

The proportion of binding errors are shown in the left-hand panels of figure 3B and 3C. Overall, PJ made a binding error on 9.44% of trials, which was significantly greater than the control group average of 3.21% (95% CI: [2.24 - 4.18]; $t(9) = 4.02$; $p = 0.003$). Increasing the duration of the memory delay had no effect on the proportion of PJ's relative binding errors; PJ's odds ratio for making a binding error following

1.0s vs. 5.0s delays was 1.27, which was not significantly different to the control group average of 1.0 (95% CI: [0.72 - 1.38]; $t(9) = 0.462$; $p = 0.655$), and suggested a non-significant tendency for more binding errors following short than long memory delays. Further, 40% (6/15) of PJ's binding errors on short delay trials, and 50% (6/12) of his binding errors on long delay trials, occurred when the target appeared in the upper portion of the screen; a goodness of fit test reported that his binding errors were not biased toward the target appearing in either the upper or lower half of the screen following either delay ($\chi^2(3) = 1$, $p = .801$). We can therefore conclude that his binding issues are unlikely due to his upper visual field deficit impacting the encoding of the entire sample stimulus.

Both PJ and controls showed systematic distortions. Following both short and long memory delays, PJ reported targets displaced leftward (1.0s: -0.24° ; 5.0s: -0.23°) and upward (1.0s: 0.15° ; 5.0s: 0.09°). In contrast, controls' group mean displacement was rightward (1.0s: 0.09° , 95% CI: $[-0.09 - 0.26]$; 5.0s: 0.07° , 95% CI: $[-0.12 - 0.27]$;) and downward (1.0s: -0.37° , 95% CI: $[-0.55 - -0.19]$; 5.0s: -0.28° , 95% CI: $[-0.45 - -0.11]$). However, PJ's displacements were not significantly different from controls for both delays (all p -values > 0.100). PJ also tended to overestimate the position of targets relative to the screen center, indicated by an error divergence of 0.04 following 1.0s delays and 0.16 following 5.0s delays. In contrast, controls underestimated targets relative to the screen center, as indicated by a group average error divergence of -0.26 (95% CI: $[-0.36 - -0.15]$) following 1.0s delays and -0.29 (95% CI: $[-0.41 - -0.16]$) following 5.0s delays. However, PJ and controls did not differ significantly (both p -values > 0.055).

Recall precision data are summarised in the right-hand panel of figure 3B and 3C. In contrast to binding errors, increasing the delay had a significant effect on recall precision. PJ's error standard deviation was 1.33° following 1.0s delays, which was not statistically different from the control group average of 1.01° (95% CI: $[0.91 - 1.10]$; $t(9) = 2.11$; $p = 0.064$). PJ's error standard deviation following 5.0s delays (1.78°) was statistically larger than the control group average of 1.18° (95% CI: $[1.09 - 1.27]$; $t(9) = 4.23$; $p = 0.002$). However, PJ's efficiency after a 5.0s delay compared to a 1.0s delay was 0.56, which was not significantly smaller than the control group average of 0.73 (95% CI: $[0.65 - 0.82]$; $t(9) = -1.37$; $p = 0.203$).

Experiment 2: Interim discussion

The experiment yielded a number of findings. First PJ made more binding errors than controls, confirming that he exhibited an impairment of spatial binding using a task in which the target location was the report rather than the cue variable. Secondly, following 1.0s delay the precision recalling the target location was not appreciably different between PJ and controls, suggesting that his binding impairment did not reflect a problem recalling the target location precisely. Moreover, while increasing the memory delay did not increase the proportion of binding errors, it did significantly diminish both PJ and controls' spatial recall precision, providing additional evidence that recall precision did not account for binding errors. In summary, PJ shows frequent binding errors, but spatial recall precision which is comparable to that of controls. Crucially, changing the duration of the memory delay produces dissociable effects on recall precision and binding.

Experiment 3: centroid estimation

Experiment 3 – Rationale

In experiment 3 we ascertained whether PJ's diminished recall of a target position may reflect a sensory impairment. While this seems unlikely given the finding that PJ's recall precision was not significantly diminished compared to controls (with 1.0s delay), it was important to establish the extent to which sensory difficulties may have limited his performance. We therefore assessed participants' spatial accuracy and precision in a perceptual task.

Experiment 3 – Methods

This experiment assessed participants' ability to localise the centroid, namely the average location, of three white discs. The discs' diameter was 0.5° (see figure 4A for a schematic representation of the trial structure). The discs remained visible until participants had positioned a crosshair shaped cursor at the desired location and clicked the mouse. Following a blank, 1.0s-long interval, a novel set of discs appeared and the procedure was repeated. Discs could occupy any of seven canonical locations. These included the screen center and the vertices of a virtual concentric hexagon, with a side length of 6.87° . All permutations of three out of seven canonical target locations, less any resulting in a collinear configuration, were used as sample arrays. Each possible permutation appeared twice, for a total of sixty-four trials. A pseudorandom, zero mean, circular Gaussian distribution, with a standard deviation of 0.6° , was used to jitter each disc's position independently. Prior to testing, instructions were read to the participants. The centroid was defined as the point in space where the triangle, whose vertices coincided with the discs' locations, would balance in the horizontal plane (Baud-Bovy and Soechting 2001). One of the experimenters also provided a visual demonstration, using a cut-out triangular shape.

Prior to testing, participants completed twenty-five practice trials. At the end of each practice trial, the reported and actual positions of the centroid were shown for 2.0s.

Experiment 3 – Data analysis

We estimated the systematic and variable error of participants' centroid estimations, by fitting a linear model to the azimuth and elevation of the reported centroid location. The model regressors included a constant and the centroid azimuth and elevation. Two metrics were used to characterise the systematic error: 1) the constant displacement, that is the tendency to report the centroid above, below, right or left of its true location, and 2) scaling factor, measuring the linear relationship between reported and actual centroid positions. These are, respectively, the estimated intercept and beta parameters of the linear model. We computed precision as the standard deviation of the variable error, i.e., residuals from the model, using the same methods used in Experiment 2.

Experiment 3 – Results: accuracy and precision of centroid estimation

The left-hand panels of figure 4B and 4C illustrate the direction of systematic biases in centroid estimates. PJ and controls respectively reported the centroid -0.07° and -0.10° (95% CI: $[-0.15^\circ - -0.04^\circ]$) left of its veridical position, suggesting that both showed a similarly small leftward bias, ($t(9) = 0.322$, $p = 0.755$). However, PJ reported the centroid 0.56° above its veridical position. This bias was significantly larger than controls, who showed a group average upward bias of 0.06° (95% CI: $[-0.02^\circ - 0.14^\circ]$; $t(9) = 3.69$, $p = 0.005$). The middle panel of figure 4B and 4C summarise the linear scaling for centroid estimates. PJ varied the reported centroid azimuth by a factor of 0.97, and elevation by a factor of 1.00, in both cases reflecting

an almost perfect linear relationship between reported and actual centroid positions. These values were comparable to those shown by controls, namely 0.99 for azimuth (95% CI: [0.94 – 1.03]; $t(9) = -0.263$, $p = 0.799$), and 0.97 for elevation (95%CI: [0.93 – 1.01]; $t(9) = 0.443$, $p = 0.668$). Finally, PJ's azimuth variable error standard deviation, 0.67° , was not significantly different from the control average of 0.69° (95%CI = [0.56° - 0.82°]; $t(9) = -0.091$, $p = 0.931$), nor was his elevation variable error standard deviation, 0.77° , significantly different from the control average of 0.59° (95%CI = [0.47° - 0.70°]; $t(9) = 0.925$, $p = 0.380$), suggesting that both the vertical and horizontal precision of his centroid judgements was relatively spared.

Experiment 3 – Interim discussion

PJ showed a strong tendency to report the centroid above its true location. This probably represents a compensatory strategy for his upper visual field defect. In fact, hemianopic patients display a bias toward their blind field when judging the midpoint of horizontal line (Barton and Black 1998; Kerkhoff and Buchers 2008). However, both PJs accuracy and precision estimating the centroid position were within the control group's range. We conclude that aside from compensatory visual defect biases, PJ's ability to localise perceptually is largely spared and unlikely to account for his diminished recall precision.

530

Discussion

531 We tested a middle-aged man (PJ) with bilateral mOTC strokes involving the PHC.
532 Acutely, PJ had developed a derangement of attention and short-term memory
533 (Horenstein et al. 1967; Medina et al. 1977; Shih et al. 2007). At the time of testing,
534 PJ was no longer delirious, but continued to have difficulties with his memory as well
535 as navigating familiar environments. The latter is a form of spatial disorientation
536 previously attributed to PHC lesions in humans (Zola-Morgan et al. 1989; Epstein et
537 al. 2001). Animal studies have demonstrated additional deficits in spatial working
538 memory following PHC lesions in non-human primates (Malkova and Mishkin 2003;
539 Bachevalier and Nemanic 2008). Whether the same deficits characterise human
540 patients with PHC lesions is not yet known.

541

542 We found that PJ had a profound deficit binding an object to its location in a working
543 memory task. When he recalled the colour of one of two objects, after a short memory
544 delay, he could accurately do so when the target was cued by its shape. However,
545 when the target was cued by its location, his accuracy was greatly diminished because
546 he made numerous binding errors, frequently reporting the colour of the non-target
547 item instead of the colour of the target. Control participants, on the other hand, were
548 accurate whether the target was identified by the location or shape cue. These findings
549 strongly suggest that PJ was impaired only when using a location cue and that this
550 impairment could not be attributed to either diminished memory for the report feature,
551 i.e. the target's colour, or a binding deficit that generalises across spatial and non-
552 spatial visual dimensions. According to a recent study, generalised binding difficulties
553 may instead characterise recall performance in individuals with autoimmune temporal
554 encephalitis, which mainly affects the hippocampal formation (Pertzov et al. 2013).

555

556 Some animal and imaging studies have indeed shown that both anterior PHC and
557 hippocampus contribute to object-in-place associations in short-term memory (Milner
558 et al. 1997; Bachevalier and Nemanic 2008). However, animal data suggest that
559 hippocampal involvement in spatial binding is restricted to tasks where spatial
560 relations are incidentally encoded (Bachevalier and Nemanic 2008). These findings,
561 together with ours, suggest that in tasks where spatial information is intentionally
562 encoded and recalled, the role of PHC goes beyond simply providing spatial data to
563 the hippocampus, where general purpose processes bind visual features in working
564 memory. Moreover, our findings confirm that binding in visual working memory is
565 liable to be disrupted by focal brain lesions (Gorgoraptis et al. 2011), supporting the
566 idea that it is a neural function independent from those underpinning the
567 representations of individual features (Wheeler and Treisman 2002; Smyrnis et al.
568 2005).

569

570 Binding errors do not reflect the resolution of spatial information

571 When PJ reported the location of one of three objects held in memory he erroneously
572 reported the location of one of the non-target items more frequently than controls.
573 This finding suggests that PJ had difficulties with spatial binding, whether space was
574 the cue or report dimension. One might argue that PJ's spatial binding impairment
575 simply reflects degraded spatial representations. In other words, diminished ability
576 recalling the location of an object might explain his difficulties using spatial
577 information to identify targets in memory. However, this hypothesis is not supported
578 by our data. PJ was able to estimate the centroid of simple dot configurations as
579 precisely as controls, indicating that despite the presence of an upper visual field

defect, the spatial resolution of visual data was not prominently affected in this perceptual task. Moreover, PJ's precision recalling the location of visual targets was not appreciably different from that of controls, even though his proportion of spatial binding errors was much greater. Finally, binding errors did not become more frequent when the delay interval was increased, although the precision of spatial recall did decrease. We conclude that binding errors do not reflect the temporal decay of a memory trace, contrary to previous suggestions (Zhang and Luck 2009). Moreover, our findings are consistent with observations that binding errors are not affected by the duration of the memory delay in either patients with hippocampal pathology (Pertsov et al. 2013) or healthy controls (Gorgoraptis et al. 2011), although whether binding errors may be effected by longer (e.g., >20.0s) delays remains to be established. Finally, varying the spatial memory demands at the time of recall in a spatial version of the Sternberg working memory task does not change the likelihood of committing a binding error, confirming that binding errors do not reflect confusion among features of the probe dimension (Smyrnis et al. 2005). Taken together, the available evidence in healthy controls and patients instead suggests that binding errors reflect interference with early processes, engaged at the time when visual information is encoded in working memory. However, a recent high-resolution fMRI study has suggested that load dependent signals in PHC during the delay period of a match-to-sample-task may reflect on-going binding processes (Schon et al. 2016).

Delays affect the precision of spatial recall

PJ's spatial recall precision was similar to that of controls when the memory delay lasted 1.0s. When the memory delay was 5.0s long, both he and controls suffered a decrement in recall precision. These are not entirely novel findings. Recall precision

is known to decrease with longer memory delays in healthy controls (Sheth and Shimojo 2001; Zhang and Luck 2009). Moreover, recall precision disproportionately decreases in patients with PHC lesions, although significantly so only following memory delays greater than 20s (Ploner et al. 2000). This finding is in keeping with our own: recall efficiency following 5.0s vs 1.0s delays was lower in PJ than in controls, however this difference was not significant. Combined, these data are consistent with the idea that following PHC lesions, spatial recall precision decays more quickly than in healthy controls, as opposed to declining abruptly. More generally, our findings are in keeping with the view that spatial recall draws information from a limited capacity resource (Bays et al. 2009), whose resolution diminishes over time. Therefore, delay dependent changes in spatial recall precision most likely reflect a limited ability to maintain information in working memory rather than impaired encoding, in contrast to the binding deficits discussed above. Finally, PJ's performance in our experiments is consistent with his neuropsychological profile, which is principally characterised by impairment on various memory tasks, including those that do not have a spatial binding component, such as the Logical Memory test and the Rey Auditory Verbal Learning Test. However we do not yet know the extent to which diminished recall precision and spatial binding account for the broad memory deficits observed following lesions to PHC.

Could the hippocampus be the site for short term memory spatial binding?

In the present study we identified impairments resulting from focal lesions to PHC, and found a spatial binding deficit in short term memory. Our data cannot rule out the possibility that binding takes place outside PHC, for example, in the hippocampus. Indeed, comparison of hippocampal volumes in PJ and age and gender matched

controls suggest hippocampal atrophy in PJ. Lateralised hippocampal atrophy commonly follows distal, ipsilateral stroke, even in young patients unlikely to harbour neurodegenerative processes (Schaapsmeeders et al. 2015a, 2015b), suggesting that the hippocampus may be particularly vulnerable to the effects of deafferentation. PJ's hippocampal atrophy raises the possibility that spatial binding deficits reflect diminished function within the hippocampus. Our data cannot refute this alternative hypothesis. As mentioned in the introduction, previous studies in patients with inflammatory and anoxic damage involving the hippocampus (e.g. Pertzov et al. 2013; Watson et al. 2013; Yee et al. 2014) have also demonstrated spatial binding impairments, lending support to the hippocampus' role in feature binding. Nonetheless, the specific spatial nature of PJ's binding impairment, which did not generalise to other visual dimensions (i.e., shape), is inconsistent with the proposal that the hippocampus provides a general purpose binding mechanism. Therefore, we conclude that spatial binding is either carried out in hippocampus, using inputs from PHC, or that PHC itself initiates spatial binding processes.

Concluding remarks

This study provides novel information on the role of MTL, by showing that a man with a lesion involving PHC, hippocampal atrophy, but spared PRC, has a selective deficit in short term spatial binding. This deficit is not explained by diminished resolution of spatial information. Our findings are consistent with the idea that spatial binding processes in short term memory may be initiated in the PHC even before visual information reaches the hippocampus.

655

Acknowledgments

656 This work was supported in part by the Biotechnology and Biological Sciences
657 Research Council grant BB/1007091/1. The authors thank Paul Mullins for his
658 assistance with MRI data acquisition, and for providing the anatomical control data.

References

- Aggleton JP. 1992. The functional effects of amygdala lesions in humans: A comparison with findings from monkeys. New York (NY): Wiley-Liss.
- Ashburner J, Friston KJ. 2003. Spatial normalization using basis functions. In: Frackowiak RS, Friston KJ, Frith CD, Dolan RJ, Price CJ, Ashburner J, Penny WD, Zeki S, editors. Human brain function. Oxford: Academic Press. p. 655-672.
- Bachevalier J, Nemanic S. 2008. Memory for spatial location and object-place associations are differently processed by the hippocampal formation, parahippocampal areas TH/TF and perirhinal cortex. *Hippocampus*. 18(1):64-80.
- Barker GR, Warburton EC. 2011. When is the hippocampus involved in recognition memory? *J Neurosci*. 31(29):10721-10731.
- Barton JJ, Black SE. 1998. Line bisection in hemianopia. *J Neurol Neurosurg Psychiatry*. 64(5):660-662.
- Baud-Bovy G, Soechting J. 2001. Visual localization of the center of mass of compact, asymmetric, two-dimensional shapes. *J Exp Psychol Hum Percept Perform*. 27(3):692-706.
- Bays PM, Catalao RF, Husain M. 2009. The precision of visual working memory is

685 set by allocation of a shared resource. J Vision. 9(10):7-7.

686

687 Belcher AM, Harrington, RA, Malkova, L, Mishkin, M. 2006. Effects of hippocampal

688 lesions on the monkey's ability to learn large sets of object-place associations.

689 Hippocampus. 16(4):361-367.

690

691 Brainard DH. 1997. The psychophysics toolbox. Spatial vision. 10:433-436.

692

693 Burwell RD, Amaral DG. 1998. Perirhinal and postrhinal cortices of the rat:

694 interconnectivity and connections with the entorhinal cortex. J Comp Neurol.

695 391(3):293-321.

696

697 Corkin S. 1984, June. Lasting consequences of bilateral medial temporal lobectomy:

698 Clinical course and experimental findings in HM. Semin Neurol. 4(2):249-259.

699

700 Corkin S, Amaral DG, González RG, Johnson KA, Hyman, BT. 1997. HM's medial

701 temporal lobe lesion: findings from magnetic resonance imaging. J Neurosci.

702 17(10):3964-3979.

703

704 Crawford JR, Howell DC. 1998. Comparing an individual's test score against norms

705 derived from small samples. Clin Neuropsychol. 12(4):482-486.

706

707 Davachi L, Goldman-Rakic PS. 2001. Primate rhinal cortex participates in both visual

708 recognition and working memory tasks: functional mapping with 2-DG. J

709 Neurophys. 85(6):2590-2601.

710

711 Deacon RM, Bannerman DM, Kirby BP, Croucher A, Rawlins JNP. 2002. Effects of
712 cytotoxic hippocampal lesions in mice on a cognitive test battery. *Behav Brain*
713 *Res.* 133(1):57-68.

714

715 Diana RA, Yonelinas AP, Ranganath C. 2007. Imaging recollection and familiarity in
716 the medial temporal lobe: a three-component model. *Trends Cogn Sci.*
717 11(9):379-386.

718

719 Eichenbaum H, Yonelinas AR, Ranganath C. 2007. The medial temporal lobe and
720 recognition memory. *Annu Rev Neurosci.* 30:123.

721

722 Epstein R, DeYoe EA, Press DZ, Rosen AC, Kanwisher N. 2001. Neuropsychological
723 evidence for a topographical learning mechanism in parahippocampal cortex.
724 *Cognitive Neuropsychol.* 18(6):481-508.

725

726 Esterman B. (1982). Functional scoring of the binocular field. *Ophthalmology.*
727 89:1226-1234.

728

729 Friedman HR, Goldman-Rakic PS. 1988. Activation of the hippocampus and dentate
730 gyrus by working-memory: a 2-deoxyglucose study of behaving rhesus
731 monkeys. *J Neurosci.* 8(12):4693-4706.

732

733 Gorgoraptis N, Catalao RF, Bays PM, Husain M. 2011. Dynamic updating of working
734 memory resources for visual objects. *J Neurosci.* 31(23):8502-8511.

735

736 Graham KS, Barense MD, Lee AC. 2010. Going beyond LTM in the MTL: a
737 synthesis of neuropsychological and neuroimaging findings on the role of the
738 medial temporal lobe in memory and perception. *Neuropsychologia*. 48(4):831-
739 853.

740

741 Habib M, Sirigu A. 1987. Pure topographical disorientation: a definition and
742 anatomical basis. *Cortex*. 23(1):73-85.

743

744 Hindy NC, Turk-Browne NB. 2016. Action-based learning of multistate objects in the
745 medial temporal lobe. *Cereb Cortex*. 26(5):1853-1865.

746

747 Holdstock JS, Shaw C, Aggleton JP. 1995. The performance of amnesic subjects on
748 tests of delayed matching-to-sample and delayed matching-to-position.
749 *Neuropsychologia*. 33(12):1583-1596.

750

751 Holdstock JS, Mayes AR, Roberts N, Cezayirli E, Isaac CL, O'Reilly RC, Norman
752 KA. 2002. Under what conditions is recognition spared relative to recall after
753 selective hippocampal damage in humans?. *Hippocampus*. 12(3):341-351.

754

755 Horenstein S, Chamberlin W, Conomy J. 1967. Infarction of the fusiform and
756 calcarine regions: agitated delirium and hemianopia. *T Am Neurol Assoc*.
757 92:85.

758

759 Jeneson A, Mauldin KN, Squire LR. 2010. Intact working memory for relational

760 information after medial temporal lobe damage. J Neurosci. 30(41):13624-
761 13629.
762
763 Katshu MZUH, d'Avossa G. 2014. Fine-grained, local maps and coarse, global
764 representations support human spatial working memory. PloS one.
765 9(9):e107969.
766
767 Keller SS, Roberts N. 2009. Measurement of brain volume using MRI: software,
768 techniques, choices and prerequisites. J Anthropol Sci. 87:127-51.
769
770 Kerkhoff G, Bucher L. 2008. Line bisection as an early method to assess
771 homonymous hemianopia. Cortex. 44(2):200-205.
772
773 Libby LA, Hannula DE, Ranganath C. 2014. Medial temporal lobe coding of item and
774 spatial information during relational binding in working memory. J Neurosci.
775 34(43):14233-14242.
776
777 Luck D, Danion JM, Marrer C, Pham BT, Gounot D, Foucher J. 2010. The right
778 parahippocampal gyrus contributes to the formation and maintenance of bound
779 information in working memory. Brain Cognition. 72(2):255-263.
780
781 Lutkenhoff ES, Rosenberg M, Chiang J, Zhang K, Pickard JD, Owen AM, Monti
782 MM. 2014. Optimized brain extraction for pathological brains (optiBET). PLoS
783 One. 9(12):e115551.
784

785 Malkova L, Mishkin M. 2003. One-trial memory for object-place associations after
786 separate lesions of hippocampus and posterior parahippocampal region in the
787 monkey. *J Neurosci.* 23(5):1956-1965.
788

789 Medina JL, Chokroverty S, Rubino FA. 1977. Syndrome of agitated delirium and
790 visual impairment: a manifestation of medial temporo-occipital infarction. *J*
791 *Neurol Neurosurg Psychiatry.* 40(9):861-864.
792

793 Merzin M. 2008. Applying stereological method in radiology. Volume measurement.
794 Bachelor's thesis. University of Tartu.
795

796 Olson IR, Page K, Moore KS, Chatterjee A, Verfaellie M. 2006a. Working memory
797 for conjunctions relies on the medial temporal lobe. *J Neurosci.* 26(17):4596-
798 4601.
799

800 Olson IR, Moore KS, Stark M, Chatterjee A. 2006b. Visual working memory is
801 impaired when the medial temporal lobe is damaged. *J Cog Neurosci.*
802 18(7):1087-1097.
803

804 Owen AM, Sahakian BJ, Semple J, Polkey CE, Robbins TW. 1995. Visuo-spatial
805 short-term recognition memory and learning after temporal lobe excisions,
806 frontal lobe excisions or amygdalo-hippocampectomy in man.
807 *Neuropsychologia.* 33(1):1-24.
808

809 Pertzov Y, Miller TD, Gorgoraptis N, Caine D, Schott JM, Butler C, Husain M. 2013.

810 Binding deficits in memory following medial temporal lobe damage in patients
811 with voltage-gated potassium channel complex antibody-associated limbic
812 encephalitis. *Brain*. awt129.
813
814 Ploner CJ, Gaymard BM, Rivaud-Péchoux S, Baulac M, Clémenceau S, Samson S,
815 Pierrot-Deseilligny C. 2000. Lesions affecting the parahippocampal cortex yield
816 spatial memory deficits in humans. *Cereb Cortex*. 10(12):1211-1216.
817
818 Ranganath C, Blumenfeld RS. 2005. Doubts about double dissociations between
819 short-and long-term memory. *Trends Cogn Sci*. 9(8):374-380.
820
821 Reisel D, Bannerman DM, Schmitt WB, Deacon RM, Flint J, Borchardt T, Seeburg
822 PH, Rawlins JNP. 2002. Spatial memory dissociations in mice lacking GluR1.
823 *Nat Neurosci*. 5(9):868-873.
824
825 Schaapsmeeders P, van Uden IW, Tuladhar AM, Maaijwee NA, van Dijk EJ, Rutten-
826 Jacobs LC, Arntz RM, Schoonderwaldt HC, Dorresteyn LD, de Leeuw FE,
827 Kessels RP. 2015. Ipsilateral hippocampal atrophy is associated with long -
828 term memory dysfunction after ischemic stroke in young adults. *Hum Brain*
829 *Mapp*. 36(7):2432-2442.
830
831 Schaapsmeeders P, Tuladhar AM, Maaijwee NA, Rutten-Jacobs LC, Arntz RM,
832 Schoonderwaldt HC, Dorresteyn LD, van Dijk EJ, Kessels RP, de Leeuw FE.
833 2015. Lower ipsilateral hippocampal integrity after ischemic stroke in young
834 adults: a long-term follow-up study. *PloS One*. 10(10):p.e0139772.

835

836 Schneider CA, Rasband WS, Eliceiri KW. 2012. NIH Image to ImageJ: 25 years of
837 image analysis. *Nat Methods*. 9:671-675.

838

839 Schon K, Newmark RE, Ross RS, Stern CE. 2016. A working memory buffer in
840 parahippocampal regions: evidence from a load effect during the delay period.
841 *Cereb Cortex*. 2016. 26(5):1965-74.

842

843 Scoville WB, Milner B. 1957. Loss of recent memory after bilateral hippocampal
844 lesions. *J Neurol Neurosurg Psychiatry*. 20(1):11-21.

845

846 Sheth BR, Shimojo S. 2001. Compression of space in visual memory. *Vision Res*.
847 41(3):329-341.

848

849 Shih H, Huang W, Liu C, Tsai T, Lu C, Lu M, Chen P, Tseng C, Jou S, Tsai C, Lee
850 CC. 2007. Confusion or delirium in patients with posterior cerebral arterial
851 infarction. *Acta Neurol Taiwanica*. 16(3):136-142.

852

853 Smyrnis N, d'Avossa G, Theleritis C, Mantas A, Ozcan A, Evdokimidis I. 2005.
854 Parallel processing of spatial and serial order information before moving to a
855 remembered target. *J Neurophysiol*. 93(6):3703-3708.

856

857 Suzuki WA, Miller EK, Desimone R. 1997. Object and place memory in the macaque
858 entorhinal cortex. *J Neurophysiol*. 78(2):1062-1081.

859

860 Suzuki WL, Amaral DG. 1994. Perirhinal and parahippocampal cortices of the
861 macaque monkey: cortical afferents. *J Comp Neurol.* 350(4):497-533.
862

863 Warrington EK, James M. 1991. The visual object and space perception battery. Bury
864 St Edmunds (United Kingdom): Thames Valley Test Company.
865

866 Watson PD, Voss JL, Warren DE, Tranel D, Cohen NJ. 2013. Spatial reconstruction
867 by patients with hippocampal damage is dominated by relational memory
868 errors. *Hippocampus.* 23(7):570-580.
869

870 Wechsler D. 1999. Wechsler abbreviated scale of intelligence. Psychological
871 Corporation.
872

873 Wheeler ME, Treisman AM. 2002. Binding in short-term visual memory. *J Exp*
874 *Psychol Gen.* 131(1):48-64.
875

876 Yee LT, Hannula DE, Tranel D, Cohen NJ. 2014. Short-term retention of relational
877 memory in amnesia revisited: accurate performance depends on hippocampal
878 integrity. *Front Human Neurosci.* 8(16).
879

880 Yonelinas AP. 2013. The hippocampus supports high-resolution binding in the
881 service of perception, working memory and long-term memory. *Behav Brain*
882 *Res.* 254:34-44.
883

884 Zhang W, Luck SJ. 2009. Sudden death and gradual decay in visual working memory.

885 Psychol Sci. 20(4):423-428.

886

887 Zola-Morgan S, Squire LR, Amaral DG, Suzuki WA. 1989. Lesions of perirhinal and

888 parahippocampal cortex that spare the amygdala and hippocampal formation

889 produce severe memory impairment. J Neurosci. 9(12):4355-4370.

890

891

892

893

894

895

896

897

898

899

900

901

902

903

904

905

906

907

908

909

Figure 1. Lesion anatomy. T1 weighted, MNI atlas registered axial (panel A) and coronal (panel B) slices are displayed in neurological coordinates, and illustrate the extent of ischemic damage in the left and right mOTC. In panel A, the axial slices also highlight the location of entorhinal and perirhinal cortex, in red and green respectively. These regions lay anteriorly and laterally to the boundaries of the ischemic lesions. In panel B, coronal slices highlight parahippocampal and hippocampal structures, including the fornix. The ischemic lesions lay inferiorly and posteriorly to the hippocampus and spare the fornix and the retrosplenial cingulate cortex. The hippocampi appear diminished in volume, more so on the right.

Figure 2. Spatial vs. non-spatial binding in working memory. Panel A shows the trial structure. The sample display for all participants (including PJ) contained a square and a triangle, placed side by side in the bottom half of the screen. The two objects were red, blue or green and never had the same colour. After a brief pattern mask and blank delay, three vertical coloured bars appeared as well as a cursor, which the participant used to report the colour of the memory target. In shape trials, targets were identified by a probe whose outline matched the target shape. In location trials, the

location of targets were identified by a white cross. Panel B shows each individual participants' error rate on a greyscale, with lighter colours representing a higher proportion of errors; the left panel shows generic errors, the right panel shows binding errors. On each panel, the upper row shows errors following shape probes, while the lower row shows errors following location probes, for PJ (blue outline) and each of the controls (red outline). Panel C shows PJ's and the group averaged proportion of generic and binding errors. Error bars are standard error of the mean.

937

938 Figure 3. Delayed spatial recall. Panel A shows the structure of immediate and
 939 delayed, spatial recall trials. The sample display for all participants (including PJ)
 940 contained three coloured discs, which could appear in both the upper and lower
 941 portion of the screen. The participants had to reproduce the position of one of the
 942 discs (the target) using a mouse cursor after either a 1.0s pattern mask or an additional

4.0s delay. The target was identified by its colour, indicated by a visual probe
displayed at the center of the screen. Panel B (left) shows PJ's (blue outline) and
controls' (red outline) individual percentage of binding errors on a greyscale,
following 1.0s (upper row) and 5.0s (lower row) delays, with lighter colours
representing a higher proportion of errors. Panel B (right) shows recall precision (95%
error ellipses) in 1.0s and 5.0s delayed recall trials for PJ (blue) and controls (red).
Panel C shows PJ's and the group averaged proportion of binding errors and
precision. Error bars are standard error of the mean.

Figure 4. Centroid estimation. Panel A shows the trial structure. The participants
placed a cursor at the centroid of the configuration formed by three bright discs. The

discs remained visible until the participant made a response by clicking the mouse. Panel B shows each participant's constant displacement (arrow vectors), scaling (diamond plot) and precision (uncertainty ellipses) in locating the centroid. The length of the diamond plot's hemi-axes corresponds to 1.0 scaling factor. Panel C shows PJ's and group averaged values of the constant displacement and scaling factor, separately for azimuth (X) and elevation (Y). The precision measure shown is the square root of the mean error variance for azimuth and elevation. Error bars in all cases are standard error of the mean.