

**Bangor University**

## **PROFESSIONAL DOCTORATES**

Choice in Life (transgender older adults' mental health) and in Death (UK GPs' euthanasia discourses)

Maddock, Emily

*Award date:*  
2017

*Awarding institution:*  
Bangor University

[Link to publication](#)

### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

### **Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



PRIFYSGOL  
**BANGOR**  
UNIVERSITY

**CHOICE IN LIFE (transgender older adults' mental health)  
AND IN DEATH (UK GPs' euthanasia discourses)**

**Emily Maddock**

**Bangor University**

Submitted in part fulfilment of the final degree award

Doctorate in Clinical Psychology

June 2017

## Contents

<b>Declarations.....</b>	<b>4</b>
<b>Acknowledgements.....</b>	<b>5</b>
<b>Thesis abstract.....</b>	<b>6</b>
<b>Literature review: Guidance for Authors.....</b>	<b>7</b>
<b>Factors that impact on the mental health of transgender older adults: A systematic review of the literature.....</b>	<b>20</b>
Abstract.....	21
Introduction.....	22
Method.....	24
Results .....	26
Socioeconomic Factors.....	26
Employment, education and income.....	26
Military service.....	29
Psychosocial Factors.....	30
Relationship status.....	30
Victimisation and discrimination.....	31
Gender identity disclosure.....	32
Social support and community belonging.....	34
Internalised stigma.....	35
Development of a cohesive self and self-acceptance.....	36
Spiritual/religious activity.....	37
Biomedical Factors.....	38
Physical health and health related behaviours.....	38
Transition related health care.....	38
Discussion.....	40
Limitations.....	41
Implications and directions for future research.....	42
References.....	44
<b>Empirical paper: Guidance for Authors.....</b>	<b>53</b>
<b>General Practitioners and the euthanasia and assisted suicide debate in the United Kingdom: A Foucauldian exploration of their discourses.....</b>	<b>70</b>
Abstract.....	71
Introduction.....	72
Method.....	75
Methodology.....	75
Research ethics.....	77
Sampling.....	77
Participants.....	78
Interviews.....	78

Analysis.....	79
Results.....	80
First author's declaration.....	80
Interpretations.....	80
Palliative care.....	82
Medicine is limited, is EU/AS the answer?.....	84
Patients have the right to EU/AS, but do not count on me.....	86
Discussion.....	91
References.....	96
<b>Contributions to personal development, clinical practice and theory.....</b>	<b>100</b>
Personal reflections.....	101
Choosing topics.....	101
Doing qualitative research.....	103
The interface between 'real life' and academic research.....	106
Implications for clinical practice.....	107
Implications for the person and clinicians/professionals.....	108
Training implications.....	109
Service level implications.....	110
On a societal level.....	111
Implications for theory development and future research.....	112
Research philosophy.....	112
Future research.....	114
References.....	116
<b>Appendices.....</b>	<b>119</b>
Appendix A - School of Psychology ethics application.....	119
Appendix B - School of Psychology ethical approval letter.....	132
Appendix C - Betsi Cadwaladr University Health Board Research and Development West Department R&D approval application.....	134
Appendix D - Betsi Cadwaladr University Health Board Research and Development West Department approval letter.....	172
Appendix E - Advert.....	175
Appendix F - Information sheet.....	177
Appendix G - Informed consent form.....	182
Appendix H - Interview schedule.....	185
Appendix I - Example transcript.....	187
Appendix J - Trans Health Factsheet on Ageing – Rising to the challenge.....	205
Appendix K - Word count statement.....	208

### Declarations

This work has not been previously accepted in substance for any degree and is not being concurrently submitted in candidature for any degree.

Signed: .....

Date: .....

#### Statement 1

This thesis is the result of my own investigations, except where otherwise stated. Other sources are acknowledged by footnotes giving explicit references. A list of references is appended.

Signed: .....

Date: .....

#### Statement 2

I hereby give consent for my thesis, if accepted, to be available using:

a) I agree to deposit an electronic copy of my thesis (the Work) in the Bangor University (BU) Institutional Digital Repository, the British Library ETHOS system, and /or in any other repository authorized for use by Bangor University and where necessary have gained the required permissions for the use of third party material.

Signed: .....

Date: .....

## **Acknowledgements**

Firstly, a huge thank you to Carolien, whose energy and enthusiasm has been inspiring. Your support has been invaluable, especially when I could no longer see the wood for the trees!

Danny, thank you for your patience, encouragement and grounding. I could not have come this far without you.

And finally, thank you to the research team, who provided guidance along the way.

**CHOICE IN LIFE (transgender older adults' mental health)  
AND IN DEATH (UK GPs' euthanasia discourses)**

**Literature review:** Transgender older adults' mental health needs are poorly understood. A systematic review of the literature revealed that transgender older adults' mental health is affected by three types of factors; socioeconomic, psychosocial and biomedical.

Transgender older adults may have unique mental health needs that differ from the general older adult population, transgender adults and youth. Fear of victimisation and

discrimination, and a lack of social support result in transgender older adults being less likely

to engage in end of life planning. **Empirical paper:** The debate in the United Kingdom (UK)

regarding the legalisation of euthanasia (EU) and assisted suicide (AS) is ongoing. However,

the views of UK GPs regarding the legalisation of EU/AS are not well understood. Six white

British GPs participated in semi-structured interviews. Their discourses were analysed using

Foucauldian discourse analysis. While the GPs' factual knowledge regarding EU/AS was

limited, they revealed three discourses: 'palliative', 'medicine is limited, is EU/AS the

answer?' and 'Patients have the right to EU/AS, but do not count on me'. **Implications for**

**theory and practice:** Both papers highlighted the influence of societal norms and related

discourses on individuals and on services, the need for open communication, staff training

and for services to be sensitive to personal choice, both for patients and staff. Many

influential models of aging are underpinned by normative assumptions. The use of these

models by services and in research may be perpetuating these assumptions through the lack

of consideration of those who do not meet such assumptions, including transgender older

adults.

## Literature Review: Guidance for Authors



[Skip Navigation](#)

## [Oxford Journals](#)

- [Contact Us](#)
- [My Basket](#)
- [My Account](#)

# [The Gerontologist](#)

- [About This Journal](#)
- [Contact This Journal](#)
- [Subscriptions](#)
- [Current Issue](#)
- [Archive](#)
- [Search](#)
- [Oxford Journals](#)
- [Medicine & Health](#) & [Social Sciences](#)
- [The Gerontologist](#)
- [For Authors](#)
- [Instructions to Authors](#)

## Instructions to Authors

**On this page:**

[Introduction](#)

[Aims and Scope of the Journal](#)

[Types of Manuscripts Considered](#)

[Formatting](#)

[Components of the Manuscript](#)

[Additional Policies and Considerations](#)

[How to Submit Online Using ScholarOne](#)

[Review Process](#)

[Paper Acceptance Information](#)

[Author Rights and Benefits](#)

[Authors Self-Archiving/Public Access Policy](#)

[Open Access](#)

[Opportunities for New Reviewers/Mentors](#)

[Editorial Correspondence](#)

### INTRODUCTION

The Gerontological Society of America (GSA), the publisher of The Gerontologist, was founded in 1945 to promote the scientific study of aging, to encourage exchanges among researchers and practitioners from the various disciplines related to gerontology, and to foster the use of gerontological research in forming public policy. The Society's 5,500+ members

include leading gerontological researchers, educators, and practitioners in biological, medical, behavioral, and social sciences and the humanities. For more information about the Gerontological Society of America, visit [geron.org](http://geron.org).

The organization fosters collaboration between physicians, nurses, biologists, behavioral and social scientists, psychologists, social workers, economists, policy experts, those who study the humanities and arts, and many other scholars and researchers in aging. This stems from the belief that the intersection of research from diverse areas is the best way to achieve the greatest impact and promote healthy aging. Through networking and mentorship opportunities, GSA provides a professional “home” for career gerontologists and students at all levels.

## AIMS AND SCOPE OF THE JOURNAL

*The Gerontologist* is a bimonthly journal of The Gerontological Society of America that provides a multidisciplinary perspective on human aging primarily through the publication of research that is relevant to social policy, program development, and service delivery. It reflects and informs the broad community of disciplines and professions involved in understanding the aging process and providing service to older people. Articles, including those in applied research, should include a conceptual framework and testable hypotheses, and report research finding with implications for policy or practice. Contributions from social and psychological sciences, biomedical and health sciences, political science and public policy, economics, education, law, and the arts and humanities are welcome. Brief descriptions of innovative practices and programs are appropriate in the Practice Concepts section. Please refer below to the Types of Manuscripts Considered for additional information about all types of manuscripts.

Due to the high volume of submissions, we are unable to offer pre-screening advice. Instead, please refer to the aims and scope of the journal to determine if *The Gerontologist* is a suitable journal for your work.

## TYPES OF MANUSCRIPTS CONSIDERED BY THE GERONTOLOGIST

All manuscripts submitted to *The Gerontologist* should address practice and/or policy implications. **\*The word limits listed below include abstract, text, and references. Tables and figures must not exceed an additional 5 pages (Literature Review submissions must not exceed an additional 10 pages of tables and figures). To manage the word and page counts, authors are encouraged to submit detailed methodology, tables, and/or figures as appendices. If your manuscript is accepted, appendices are available to readers online only.**

a. **Research Articles** present the results of original research. These manuscripts may be no longer than 6,000\* (7,000\* for qualitative studies) words. The word count includes; abstract, text and references. Tables and figures must not exceed an additional 5 pages. The text is usually divided into sections with the headings: Introduction, Design and Methods, Results, and Discussion. Subheads may also be needed to clarify content. Research design and analysis procedures as well as implications for practice or policy must be clearly described (See [Schoenberg, N. & McAuley, W. J. \(2007\). Promoting qualitative research. \*The Gerontologist\*, 47\(5\), 576–577.](#) and [Schoenberg, N.E., Miller, E.A. and Pruchno, R., \*The Qualitative Portfolio at The Gerontologist: Strong and Getting Stronger. The Gerontologist\*](#)

[\(2011\) 51\(3\): 281–284](#) ). Qualitative manuscripts should avoid the subheading “A Qualitative Study.”

Humanities and Arts: Please refer to the following editorial for additional detail with these types of submission: [Kivnick, H.Q. & Pruchno, R. \(2011\). Bridges and Boundaries: Humanities and Arts Enhance Gerontology, \*The Gerontologist\*, 51\(2\), 142-144.](#)

**b. Literature Review or Measurement Online Only Articles** *The Gerontologist* has launched a new policy welcoming submissions using sophisticated scale development procedures (6,000\* words) and/or literature reviews (8,000\* words) for consideration for publication. These articles will be published online only (title would appear in a print issue Table of Contents for the journal, but the article would appear online only). Articles will go through our usual peer review and editing processes. They will receive a DOI, be searchable, and will be available electronically.

**c. Brief Reports** Brief reports are encouraged for significant and innovative papers that are not as long as full research articles, but are equivalent in quality. Manuscripts should be no more than 2,500\* words. The word count includes the abstract, text and references.

**d. The Forum** Timely scholarly review articles or well-documented arguments presenting a viewpoint on a topical issue are published in this section. Total length should be no more than 5,000\* words. The word count includes the abstract, text and references.

**e. Practice Concepts** A Practice Concepts manuscript describes, in 6,000\* words or fewer, an innovative practice amenable to replication OR the translation of a practice that already has an evidence base but has not yet been trialed in the "real world." Authors must clearly specify the following information about the practice: (a) uniqueness or innovation, (b) theoretical or conceptual basis, (c) essential components or features, (d) evidence that supports replication or suggests modifications, and (e) challenges to feasibility and/or barriers to implementation in real world settings. The article should be structured to highlight these points (the structure of a research article does not necessarily apply). An important goal is to provide enough information about the practice to allow its replication (in a research context) or translation (in a real world context). The word count includes the abstract, text and references.

Please refer to the following editorials for further information about Practice Concept submissions: [Morrow-Howell, N. & Noelker L. \(2006\). Raising the Bar to Enhance the Research–Practice Link. \*The Gerontologist\*, 46, 315-316](#) and [Kimberly S. Van Haitsma, K.S., Williamson, J.B., & Pruchno, R. \(2011\). Practice Concepts and Policy Studies: The New Divide, \*The Gerontologist\*, 51\(6\), 731-733.](#)

**f. Policy Studies** The aim of this section to highlight policy research; to share some of the best research on aging issues with policymakers, senior advocates, and others involved in aging policy issues; and to bring to bear on the policy process the knowledge and expertise of gerontological researchers heralding from diverse academic backgrounds. We are particularly interested in articles that make use of theories, conceptual frameworks, and the like that help readers interpret the policy analysis presented. Policy studies manuscripts deal with aging-related public policy programs and proposed changes in those programs in the United States, as well as similar programs in other countries. The tone and style of the manuscript should be scholarly, but also easily understood and of interest to a broad audience that includes not only

GSA members but journalists, Congressional staff members, and others who are involved in formulating and reforming at the local, state, and federal level social policies that impact the older population. The content should tell a new and interesting story that is supported by empirical data. (Essays linked to important public policy debates that are not grounded on original empirical research would be best submitted to the Forum section of the journal.) Manuscripts should be no longer than 7,000\* words.

Suitable topics include, but are not limited to, the following:

- Major aging-related federal social programs such as Social Security, Medicare, Medicaid, the Affordable Care Act, and the numerous programs funded by the Administration on Aging;
- Federal, state, and local policies dealing with assisted living, long-term care, elder abuse, illegal scams affecting the older population, late-life healthcare rationing, end-of-life care, assisted suicide, age-restricted communities, aging in place, migration patterns among the older population, federal and state taxes, and estates and trusts;
- Nongovernmental or corporate policies that foster or adversely affect the well-being of the older population, such as outsourcing, workplace flexibility, inclusion and exclusion in retention programs; and
- Cross-national policy studies—of particular interest are studies about other developed countries that are grappling with aging policy-related challenges that are similar to those in the United States.

g. **On Film and Digital Media.** Please refer to the editorial "[Launching 'On Film and Digital Media.'](#)"

h. **Book Reviews.** Book reviews are published in essay form. Reviews are prepared at the request of the Book Review Editor and are not guaranteed for acceptance prior to submission. Unsolicited book review essays are not accepted. Books for review should be sent to Judith G. Gonyea, PhD, Book Review Editor, Social Research Department, School of Social Work, Boston University, 264 Bay State Road, Boston, MA 02115.

i. **Guest Editorials** Upon occasion, the Editor-in-Chief will invite guest editorials. Unsolicited editorials are not accepted.

*The Gerontologist* does not publish obituaries, speeches, announcements of programs, or new product information.

Supplement issues of *The Gerontologist* are additional and externally funded issues. Please contact the editorial office at [tg@geron.org](mailto:tg@geron.org) for further information. The Gerontologist also publishes special issues, developed by the editors of *The Gerontologist*, within our regularly scheduled bimonthly issues.

## FORMATTING

Manuscripts are to be submitted in Microsoft Word or a Word-compatible program at [ScholarOne](#). Manuscripts submitted in other formats will be unsubmitted and returned to the corresponding author for correction prior to editor review. Please DO NOT submit PDF versions of your manuscript submission materials. A peer-review title page will be created by the system and will be combined with the main document file into a single PDF document. This document will be used for the peer review process.

*The Gerontologist* uses **APA style**. General guidelines follow; for more detailed information, consult the [Publication Manual of the American Psychological Association](#) (6th ed.). Please see section [TYPES OF MANUSCRIPTS CONSIDERED BY THE GERONTOLOGIST](#) above for additional information about the types of submissions and word counts. Please read [“Editorial: Science or Fishing?”](#) for valuable information about manuscript preparation.

This journal has a double-blind review policy. Therefore, we require authors to submit **TWO** versions of the article file, anonymous and non-anonymous (see below for additional information). Please upload **ONLY** these two files, the anonymous and non-anonymous manuscript files. Include your tables and figures in these documents, following APA guidelines.

**Abbreviations:** Ensure that the use of abbreviations is clear and that each one is defined in the text at its first mention only.

**In-text References and Citations** Refer to the [Publication Manual of the American Psychological Association](#) (6th ed.) for style and see the [FORMATTING](#) section above. References in text are shown by citing in parentheses the author’s surname and the year of publication. Example: “. . . a recent study (Jones, 1987) has shown. . . .” If a reference has two authors, the citation includes the surnames of both authors each time the citation appears in the text. When a reference has more than two authors and fewer than six authors, cite all authors the first time the reference occurs. In subsequent citations, and for all citations having six or more authors, include only the surname of the first author followed by “et al.” Multiple references cited at the same point in the text are in alphabetical order by author’s surname.

**Instructions for Anonymous Files** Upload one anonymous version of your manuscript with no author names or contact information on the title page, blinded acknowledgment details, and removed other self-identifying author information. Be sure to check there are no self references in the text itself that would reveal the identity of the authors. Please remove those instances and insert the term "blinded for review" in its place. This includes self-references to your name and University/Institute. This version may be seen by peer-reviewers.

**Instructions for Non-Anonymous Files** Upload a complete version of the manuscript with all of the author and acknowledgment details. This version will be seen by the editors and will be the version published, IF accepted.

## COMPONENTS OF THE MANUSCRIPT

**Cover Letter (Optional).** A cover letter is not required and is optional. It should explain how the manuscript is innovative, provocative, timely, and of interest to a broad audience, and other information authors wish to share with editors. Note: The cover letter for manuscripts will NOT be shared with reviewers.

**Title page.** A title page should be a completely separate page that includes the following:

- (1) Title of the manuscript, APA recommends that a title be no more than 12 words.
- (2) **All** authors' full name(s), affiliations, and email addresses.
- (3) The corresponding author should be clearly designated.

**Abstract and Key Words.** On a separate page, each manuscript must include a brief

abstract. Structured abstracts for research articles, Brief Reports, and Practice Concepts submissions should be approximately 250 words (the web-based system will not accept an abstract of more than 250 words), and must include the following headings: **Purpose of the study, Design and Methods, Results, and Implications**. Forum and Policy Studies manuscripts must also include an abstract of about 200 words, but may be without structured headings.

Below the abstract, authors should supply three to five key words that are NOT in the title. (Please avoid elders, older adults, or other words that would apply to all manuscripts submitted to *The Gerontologist*.) Please note three key words must be entered in order to move forward in the online submission process.

**Text.** The text of research articles, Brief Reports, and Practice Concepts submissions should follow the headings included in the structured abstract (see above Abstract and Key Words). Forum and Policy Studies manuscript should also be divided into headings, as appropriate for the submission. Articles may need subheadings within some sections to clarify their content. The Implications should not merely restate the results but should interpret the results and specify the policy and/or practice implications.

(1) The word counts for the different kinds of publications considered by the Journal are presented above and are inclusive of the abstract, text and references.

(2) If manuscripts greatly exceed these word count limits, your manuscript may be returned to you for correction BEFORE the peer review process can begin. If you would like to appeal the word count limit for the text of the manuscript, permission must be granted by the Editor in Chief prior to submission. When submitting, please indicate in your cover letter that permission has been granted.

**Acknowledgment (Optional).** If the authors choose to include acknowledgments recognizing funders or other individuals, they should be placed on a separate page immediately following the title page. The self-identifying acknowledgments should be removed from the anonymous version of the manuscript.

### **Conflict of Interest**

Please declare any conflict of interest, if applicable.

### **FUNDING**

Details of all funding sources for the work in question should be given in a separate section entitled Funding. This should appear before the Acknowledgements section.

The following rules should be followed:

- The sentence should begin: ‘This work was supported by ...’
- The full official funding agency name should be given, i.e. ‘National Institutes of Health’, not ‘NIH’ (full RIN-approved list of UK funding agencies) Grant numbers should be given in brackets as follows: ‘[grant number xxxx]’
- Multiple grant numbers should be separated by a comma as follows: ‘[grant numbers xxxx, yyyy]’

- Agencies should be separated by a semi-colon (plus 'and' before the last funding agency)
- Where individuals need to be specified for certain sources of funding the following text should be added after the relevant agency or grant number 'to [author initials]'.

An example is given here: 'This work was supported by the National Institutes of Health [AA123456 to C.S., BB765432 to M.H.]; and the Alcohol & Education Research Council [hfygr667789].'

Oxford Journals will deposit all NIH-funded articles in PubMed Central. See [http://www.oxfordjournals.org/for\\_authors/repositories.html](http://www.oxfordjournals.org/for_authors/repositories.html) for details. Authors must ensure that manuscripts are clearly indicated as NIH-funded using the guidelines above.

### *Crossref Funding Data Registry*

In order to meet your funding requirements authors are required to name their funding sources, or state if there are none, during the submission process. For further information on this process or to find out more about the CHORUS initiative please click [here](#).

### **Reference List**

Arrange alphabetically by author's surname; do not number. The reference list includes only references cited in the text. Do not include references to private communications or submitted work. Consult the [\*Publication Manual of the American Psychological Association\*](#) (6th ed.) for correct form.

Examples:

Journals: Kaskie, B., Imhof, S., Cavanaugh, J., & Culp, K. (2008). Civic engagement as a retirement role for aging Americans. *The Gerontologist*, 48, 368–377.

doi:10.1093/geront/48.3.368

Books: Quadagno, J. S. (1982). *Aging in early industrial societies*. New York: Academic Press.

### **Tables**

Tables are to be numbered consecutively with Arabic numbers and have a brief title for each. Place table footnotes immediately below the table, using superscript letters (a, b, c) as reference marks. Asterisks are used only for probability levels of tests of significance (\* $p < .05$ ). Tables should be embedded in the anonymous and non-anonymous files.

### **Figures/Illustrations**

Please include your figures embedded in the anonymous and non-anonymous word processing file, per APA style guidelines. The journal reserves the right to reduce the size of illustrative material. Each figure should be at sufficient resolution (i.e., 300 dpi at 5 in.) to be clear, sharp images when reduced to print size. Figures must be professionally lettered in a sans-serif type (e.g., Arial or Helvetica). All labels used in figures should be in upper case in both the figure and the caption.

### **Color figures**

Figures may appear in color online, but will only appear in color in print when deemed

necessary. Please contact the editorial office for further information about color figures at [tg@geron.org](mailto:tg@geron.org).

### **Captions for Tables and Illustrations**

Type captions on a separate page with numbers corresponding to the illustrations. Figure captions should provide with sufficient information so that the reader can understand it with minimal reference to the text. Explain symbols, arrows, numbers, or letters used in illustrations. Explain internal scale and identify staining method in photomicrographs. Captions for tables and illustrations should be adequate to interpret the content of the table or figure without reference to the text.

### **Supplementary Material**

Supplementary material can be made available by the publisher online only and linked to the published article. This material includes supporting material that is not essential for inclusion in the full text to understand the conclusions of the paper but contains data that is additional or complementary and directly relevant to the article content and therefore may benefit the reader. Such information might include more detailed methods, extended data sets/data analysis, or additional figures.

It is standard practice for appendices to be made available online only as supplementary material. All text and figures must be provided in suitable electronic formats. All material to be considered as supplementary material must be submitted for peer review at the same time as the manuscript and included in the anonymous and nonanonymous versions of the manuscript (do not upload the material as separate files) and indicated clearly as supplementary material. Also ensure that the supplementary material is cited in the main manuscript where necessary, for example, “(see Supplementary Material)” or “(see Supplementary Figure 1).” The material cannot be altered or replaced after the paper has been accepted for publication, and **it will not be edited**.

### **Appendices**

All appendices will be published online only as supplementary material (please see [FORMATTING](#) and [SUPPLEMENTARY MATERIAL](#) instructions above).

## **ADDITIONAL POLICIES AND CONSIDERATIONS**

### **Permissions for Illustrations and Figures**

Permission to reproduce copyright material, for print and online publication in perpetuity, must be cleared and if necessary paid for by the author; this includes applications and payments to DACS, ARS and similar licensing agencies where appropriate. Evidence in writing that such permissions have been secured from the rights-holder must be made available to the editors. It is also the author's responsibility to include acknowledgements as stipulated by the particular institutions. Oxford Journals can offer information and documentation to assist authors in securing print and online permissions: please see Guidelines for Authors at [http://www.oxfordjournals.org/for\\_authors/](http://www.oxfordjournals.org/for_authors/). Information on permissions contacts for a number of main galleries and museums can also be provided. Should you require copies of this, please contact the editorial office of the journal in question or the Oxford Journals Rights department.



## **Ethics**

*The Gerontologist* expects that authors will observe high standards with respect to publication ethics. For example, the following practices are unacceptable: (1) falsification or fabrication of data, (2) plagiarism, including duplicate publication of the authors' own work, in whole or in part without proper citation, (3) misappropriation of the work of others such as omission of qualified authors or of information regarding financial support. Allegations of unethical conduct will be discussed initially with the corresponding author. In the event of continued dispute the matter will be referred to the author's institution and funding agencies for investigation and adjudication.

Oxford Journals, publisher of *The Gerontologist*, is a member of the Committee on Publication Ethics (COPE), and the journal strives to adhere to the COPE code of conduct and guidelines. For further information see <http://publicationethics.org/>.

Submission of a manuscript to *The Gerontologist* implies that it has not been published or is not under consideration elsewhere. If accepted for this journal, it is not to be published elsewhere without permission. As a further condition of publication, the corresponding author will be responsible, where appropriate, for certifying that permission has been received to use copyrighted instruments or software employed in the research and that human or animal subjects approval has been obtained.

In the case of coauthored manuscripts, the corresponding author will also be responsible for signing a license to publish form on behalf of his or her co-authors indicating that the manuscript is original work, has not been previously published, will not be submitted for publication to any other journal following acceptance in *The Gerontologist*, and all necessary permissions have been received.

## **HOW TO SUBMIT ONLINE USING SCHOLARONE**

Manuscripts must be submitted online at *The Gerontologist's* online submission site. Instructions on submitting your manuscript online can be viewed [here](#). After reading the instructions, if you still have questions on how to submit your manuscript online, contact the Managing Editor, Megan McCutcheon, at [tg@geron.org](mailto:tg@geron.org).

## **REVIEW PROCESS**

All papers published in *The Gerontologist* are subject to peer review. Papers that are outside the scope of the journal, that do not comply with the guidelines below or are otherwise judged to be unsuitable by the editor will be rejected without peer review. Peer-reviewed manuscripts are sent to at least two independent referees for evaluation. Authors are encouraged to suggest reviewers of international standing. Reviewers advise on the originality and scientific merit of the paper; the editors make all final decisions on publication. The corresponding author will be notified via email with a decision and revision requirements, if applicable. Please ensure that email addresses are current for notification purposes.

## **PAPER ACCEPTANCE INFORMATION**

### **Licenses and Copyright**

It is a condition of publication in *The Gerontologist* that authors grant an exclusive license to

The Gerontological Society of America. This ensures that requests from third parties to reproduce articles are handled efficiently and consistently and will also allow the article to be as widely disseminated as possible. As part of the license agreement, authors may use their own material in other publications provided that The Gerontologist is acknowledged as the original place of publication and Oxford University Press as the Publisher.

Upon receipt of accepted papers at Oxford Journals, authors will be required to complete an online copyright license to publish form. This form will be sent to the corresponding author via email.

Please note that by submitting a manuscript for consideration for publication, you confirm that you are the corresponding/submitting author and that Oxford University Press ("OUP") may retain your email address for the purpose of communicating with you about the submission. You agree to notify OUP immediately if your details change. If your article is accepted for publication OUP will contact you using the email address you designated in the submission process. Please note that OUP does not retain copies of rejected articles.

Information about the Creative Commons licensee can be found at <http://creativecommons.org/>.

## **AUTHOR RIGHTS AND BENEFITS**

### **Advance Access**

Advance Access allows for papers to be published online soon after they have been accepted for publication--reducing the time between submission and publication. Articles posted for Advance Access have been copyedited and typeset but not yet paginated for inclusion in a specific issue of the journal. Appearance in Advance Access constitutes official publication, with full-text functionality, and the Advance Access version can be cited by a unique DOI (Digital Object Identifier). The final manuscript is then paginated into an issue, at which point it is removed from the Advance Access page. Both versions of the paper continue to be accessible and citable.

### **Authors Self-Archiving/Advance Access Policy**

For information about this journal's policy, please visit our [Author Self-Archiving policy page](#).

### **Open Access**

The Gerontologist authors have the option to publish their paper under the [Oxford Open initiative](#); whereby, for a charge, their paper will be made freely available online immediately upon publication. After your manuscript is accepted the corresponding author will be required to accept a mandatory license to publish agreement.

As part of the licensing process you will be asked to indicate whether or not you wish to pay for open access.

If you do not select the open access option, your paper will be published with standard subscription-based access and you will not be charged.

Oxford Open articles are published under Creative Commons licenses.

RCUK/Wellcome Trust funded authors publishing in The Gerontologist can use the Creative

Commons Attribution license (CC-BY) for their articles.

All other authors may use the following Creative Commons licenses:

- Creative Commons Attribution Non-Commercial license (CC-BY-NC)
- Creative Commons Attribution Non-Commercial No Derivatives license (CC-BY-NC-ND)

Please click [here](#) for more information about the Creative Commons licenses.

You can pay Open Access charges using our Author Services site. This will enable you to pay online with a credit/debit card, or request an invoice by email or post. The applicable open access charges vary according to which Creative Commons license you select. The open access charges are as follows.

Charges for CC-BY:

- Regular charge: £2125/ \$3400 / €2763
- List B Developing country charge\*: £1063/ \$1700 / €1382
- List A Developing country charge\*: £0 /\$0 / €0

Charges for CC-BY-NC/CC-BY-NC-ND:

- Regular charge: £1875/ \$3000 / €2438
- List B Developing country charge\*: £938/ \$1500 / €1219
- List A Developing country charge\*: £0 /\$0 / €0

\*Visit our developing countries page (click [here](#) for a list of qualifying countries).

Please note that these charges are in addition to any color charges that may apply.

Orders from the UK will be subject to the current UK VAT charge. For orders from the rest of the European Union, OUP will assume that the service is provided for business purposes. Please provide a VAT number for yourself or your institution, and ensure you account for your own local VAT correctly.

### **Offprints**

Authors will receive electronic access to their paper free of charge. Printed offprints may be purchased in multiples of 50. Authors may purchase offprints through the author services site where they sign their online license forms.

You can pay Open Access charges using our Author Services site. This will enable you to pay online with a credit/debit card, or request an invoice by email or post. The applicable open access charges vary according to which Creative Commons licence you select. The open access charges are as follows.

Charges for CC-BY:

- Regular charge: £2125/ \$3400 / €2763
- List B Developing country charge\*: £1063/ \$1700 / €1382
- List A Developing country charge\*: £0 /\$0 / €0

Charges for CC-BY-NC/CC-BY-NC-ND:

- Regular charge: £1875/ \$3000 / €2438
- List B Developing country charge\*: £938/ \$1500 / €1219
- List A Developing country charge\*: £0 /\$0 / €0

\*Visit our developing countries page (click [here](#) for a list of qualifying countries).

Please note that these charges are in addition to any colour charges that may apply.

Orders from the UK will be subject to the current UK VAT charge. For orders from the rest of the European Union, OUP will assume that the service is provided for business purposes. Please provide a VAT number for yourself or your institution, and ensure you account for your own local VAT correctly.

## **OPPORTUNITIES FOR NEW REVIEWERS/MENTORS**

For students or emerging scholars interested in participating in the review process for *The Gerontologist*, we provide an opportunity for novice reviewers. Novice reviewers are added as a third reviewer to manuscripts that we send out for peer review. These reviews are given full consideration in the editorial decision. This process gives new professionals an opportunity not only to review a manuscript, but also to experience the review process, editorial decisions, and manuscript revision process. This also helps the novice reviewer learn how to publish in *The Gerontologist*. After individuals successfully complete two or more reviews, novice reviewers will become regular reviewers. All reviewers are asked to return their reviews within two weeks.

We also encourage individuals to serve as a mentor to new professionals who are interested in the review process for *The Gerontologist*. The current review form allows for this designation so that all invited reviews can be completed with a reviewer-in-training, if interested.

## **Editorial Correspondence**

Questions? Contact the Editorial Office at [tg@geron.org](mailto:tg@geron.org).

**Factors that impact on the mental health of transgender older adults: A systematic review  
of the literature**

Emily L. Maddock (Corresponding author)  
North Wales Clinical Psychology Programme  
Bangor University  
Bangor, LL57 2DG  
Psp4f4@bangor.ac.uk

Carolien Lamers  
North Wales Clinical Psychology Programme  
Bangor University  
Bangor, LL57 2DG  
c.lamers@bangor.ac.uk

This paper is part of the Doctorate in Clinical Psychology  
North Wales Clinical Psychology Programme  
Bangor University

## Abstract

**Purpose:** This review provides an update on the limited body of research examining the factors that impact on the mental health of transgender older adults. Factors reported in the transgender aging literature are identified and considered in relation to the broader adult and youth transgender literature, taking into consideration the shifting social climate.

**Design and Methods:** The databases PsycINFO, ProQuest Social Sciences Premium Collection, Social Science Database, Political Science Database and Science Direct were searched for studies published between 2000 and 2017. The following terms were used: (Transgender\*), male-to-female, female-to-male, gender identity, gender expression, (old\*) adults, elder, mental health and successful aging. Eight articles were identified that met the criteria for inclusion in the review. **Results:** Three factors were identified. Socioeconomic factors included employment, education and income, and military service. Psychosocial factors included relationship status, victimisation and discrimination, gender identity disclosure, social support and community belonging, internalised stigma, development of a cohesive self and self-acceptance, and spiritual/religious activity. Biomedical factors included physical health and health related behaviours, and transition related health care.

**Implications:** Transgender older adults may have unique mental health needs that differ from the general older adult population, transgender adults and youth. These needs and cohort effects should be taken into consideration by health and social care organisations involved in the mental health care provision for transgender older adults.

**Key Words:** *Lesbian, Gay, Bisexual and Transgender (LGBT), Quality of life (QOL), Gender identity*

Transgender older adults' mental health needs continue to be poorly understood and are understudied. Although there has been a recent increase in research focusing on the mental health of transgender individuals, much of this focus has been on transgender youth, health care access, socio-demographic characteristics and rates of mental health difficulties. Little research has attempted to understand the factors that influence the mental health of transgender individuals, and even fewer studies focus on older adults who identify as transgender. In this review, mental health is conceptualised as a subjective state of cognitive and emotional wellbeing that encompasses, quality of life (QOL), satisfaction and contentment. Psychological distress, suicidal ideation and mental health diagnoses such as depression and anxiety are referred to as indicators of mental health difficulties (Fredriksen-Goldsen, Kim, Shiu, Goldsen, & Emler, 2015; Hoy-Ellis et al., 2017; McCann et al., 2013).

The term 'transgender' is used to describe individuals whose gender identity, expression or behaviours do not match those usually associated with their biological sex (Persson, 2009). Biological sex is categorised as either female, male or intersexed according to an individual's anatomy. Traditionally, gender has been dichotomised as either male or female and assigned at birth according to biological sex. However, gender identity is increasingly being understood as a more complex and multidimensional construct that refers to an individual's 'sense of self' as being male, female or a sense of self that does not conform to this traditional dichotomy.

Population estimates are difficult both because many official documents do not include data on gender identity (e.g. the United States [US] & United Kingdom [UK] Census) and because many transgender individuals prefer not to disclose their gender identity (Meerwijk & Sevelius, 2017). There is currently no validated estimate of the population of transgender adults in the UK. Based on survey data from 1998, Reed, Rhodes, Schofield, and Wylie

(2009) estimated that in 2007 there was likely to be a minimum of 10,000 transgender individuals over the age of sixteen in the UK, who had sought transition related health care. Transgender individuals are estimated to make up 0.5% of the American adult population (Conron, Scott, Stowell, & Landers, 2012). Slightly higher prevalence rates have been reported for transgender youth. In a school based survey, 1.2% of 8,166 students in New Zealand reported being transgender and 2.5% were not sure about their gender (Clark et al., 2014). Similar rates of 1.3% of 2,730 middle school students reported identifying as transgender in an American school based survey (Shields et al., 2013).

Transgender individuals report higher rates of anxiety (40.4% for transgender women & 47.5% for transgender men; Budge & Adelson, 2013) and depression (48-62%; Nemoto, Bodeker, & Iwamoto, 2011) than the general population (28.8% & 16.6% respectively; Kessler, Berglund, Demler, Jin, & Walters, 2005). The National Transgender Discrimination Survey (Hass, Rodgers, & Herman, 2014) conducted in America reported lifetime suicide attempts of 45% for 18-44 year olds, 33% for 55-64 year olds and 16% for over 65s. Like other minority groups, transgender individuals can find themselves in a social context that does not provide adequate support or health care to meet their physical and mental health needs.

Stigma-related discrimination and victimisation have long been recognised as having a profoundly negative impact on the mental health and wellbeing of minority groups. Transgender individuals are at greater risk of experiencing discrimination and victimisation than the general population (Bradford, Reisner, Honnold, & Xavier, 2013). Prevalence estimates of transgender-related discrimination have been reported to exceed 60% among adults (Clements-Nolle, Marx, & Katz, 2006).



It is clear that transgender individuals across the life span are at higher risk of experiencing mental health issues. Yet, although the mental health needs of transgender individuals are increasingly being recognised, there continues to be a lack of understanding of the factors that impact on mental health, particularly for transgender older adults. In addition, understanding transgender older adults' mental health is complicated by the fact that what constitutes an "older adult" is culturally defined. In most western societies it is generally accepted to be retirement age (63 for women and 65 for men in the United Kingdom [UK]) but can include people as young as 50 (Department of Health, 2001). The term older adult will be used in this review to reflect this common definition and will refer to individuals aged 50 and over. This paper reviews the limited research on the factors that impact the mental health of transgender older adults. Factors are discussed in comparison with the experience of non-transgender older adults, transgender adults and youth as reported in the literature.


### **Method**

The databases PsycINFO, ProQuest Social Sciences Premium Collection, Social Science Database, Political Science Database and Science Direct were searched using the following terms: (Transgender\*), male-to-female, female-to-male, gender identity, gender expression, (old\*) adults, elder, mental health and successful aging. As the transgender older adult population has been 'invisible' and the focus of little research until approximately the year 2000 (Witten, 2016), the search was limited to studies published between 2000 and 2017. The search and selection process was completed by the first author.

The search produced 272 articles of potential relevance. The resulting abstracts were assessed for relevance and excluded if they were duplicates, not available in English, were not exclusive to older transgender adults or did not include factors specific to mental health

or wellbeing (see Figure 1). Qualitative and quantitative studies that assessed mental health factors as variables in statistical models and questionnaires were included.

Eight articles were identified that met the criteria to be included in the review (see Table 1). Although these studies ranged in quality from rigorous large scale quantitative designs (e.g. Fredriksen-Goldsen et al., 2015) to discussions of case vignettes (e.g. Ettner & Wylie, 2013), all eight studies were included due to the limited amount of studies available.



Factors that affect the health and wellbeing of transgender older adults have been categorised in the literature as biomedical, psychosocial and socioeconomic (Ettner & Wylie, 2013; Witten 2016). Therefore, each article was reviewed and factors that impacted on the mental health of transgender older adults were identified (see Table 1 for summary) and presented under the following headings (1) socioeconomic factors, (2) psychosocial factors and (3) biomedical factors.

## **Results**

Although the literature base is currently limited, the available studies provide an initial indication of the factors that impact on transgender older adult's mental health. Under the socioeconomic category, the following factors were identified: employment, education and income, and military service. Under the psychosocial category, the following factors were identified: relationship status, victimisation and discrimination, gender identity disclosure, social support and community belonging, internalised stigma, development of a cohesive self and self-acceptance, and spiritual/religious activity. Under the biomedical category, the following factors were identified: physical health and health related behaviours, and transition related health care. Each of the factors is summarised and reviewed in comparison to the general older adult, transgender adult and youth literature where appropriate.

### **Socioeconomic Factors**

#### *Employment, education and income*

In the general population, employment and higher levels of income are associated with lower levels of psychological distress, depression and anxiety compared to unemployment

Table 1  
 Summary of studies included in the review and factors identified.

Study	Year	Data collection	N	Age group	% transgender	Mental health outcomes	Factors identified	
							Positive	Negative
Elder	2016	Semi-structured interviews	10	60+	100%	Subjective reports of experiences (suicidal ideation, depression, loneliness & low self-esteem)	Gender transition, advocating/teaching & raising awareness of transgender issues, community connectedness	Historical victimisation/discrimination, rejection/loss & discrimination in religious community
Ettner & Wylie	2013	Case vignettes	5	60+	100%	Subjective reports of depression, anxiety & wellbeing	Self-acceptance	Hormone deficiency
Fabbre	2015	Interviews & participant observations	22	50+	100%	Subjective reports of wellbeing & experiences of emotions	Prior military service, development of a cohesive self & self-acceptance	Heteronormative societal gender expectations & internalized stigma
Fredriksen-Goldsen et al.	2013	Cross-sectional survey	2,560 LGBT	50+	7% (174)	Depressive symptomatology & perceived stress	Social support & community belonging	Internalised stigma, victimisation, fear of accessing health services, obesity, lack of physical activity & concealment of gender identity
Fredriksen-Goldsen et al.	2015	Cross-sectional survey	2,560 LGBT	50+	7% (174)	Physical & mental health related QOL	Positive sense of sexual identity, social network size, social support, income, employment, health promoting behaviours	Victimisation & discrimination, gender identity disclosure & chronic health conditions

Hoy-Ellis et al.	2017	Cross-sectional survey	183	50+	100%	Depressive symptomatology & psychological health related quality of life	Prior military service	Identity stigma
McCann et al.	2013	Survey & semi-structured interviews	144 LGBT	55+	7% (10)	Mental health status, suicide & self-harm & substance misuse	-	Victimisation & discrimination (physical violence/verbal insults), loss, rejection, gender identity disclosure & substance misuse
Porter et al.	2013	Cross-sectional survey	289	51+	100%	Subjective experience of successful aging	Religious affiliation	Discrimination by religious groups

and lower levels of income (<\$20,000; Frech & Damaske, 2012; Hergenrather, Zeglin, McGuire-Kuletz, & Rhodes, 2015; Prause & Dooley, 2001; Robert et al., 2009; Van Gorp et al., 2007). The positive impact of employment and income on mental health is also consistently demonstrated in the transgender adult population (Rotondi et al., 2011). Retirement has been associated with reduced levels of depression, including when compared to those in employment in the general population (Mandal & Roe, 2008).

Employment and income were reported to have a positive impact on the mental health QOL of transgender older adults (Fredriksen-Goldsen et al., 2015). However, an elevated rate of suicide attempts has been shown for those who have experienced discrimination in the work environment, particularly those who had experienced physical violence ( $n = 209$ ; 65%; Hass et al., 2014). The Fredriksen-Goldsen et al. (2015) study was of high quality and used multivariate linear regression to assess the independent contributions of variables, including age (*Young-old* = 50-64 years,  $n = 1,078$ ; *Middle-old* = 65-79 years,  $n = 1,138$ ; & *Old-old* = 80+ years,  $n = 247$ ), on mental health QOL. While the positive impact of income was consistent across age groups but strongest in the young-old group, education was not reported to have a positive effect on mental health QOL. This finding is consistent with the general older adult literature (Robert et al., 2009) which suggests that income is a stronger predictor of mental health QOL than education for older people, regardless of gender orientation.

### *Military service*

Within the general older adult population, military service has been associated with poorer mental health in the short term following service compared with those without prior military service (Wilmoth, London, & Parker, 2010). However, recent research shows

positive impacts of prior military service on older adults' mental health in the longer term (Spiro, Settersten, & Aldwin, 2016). In a recent study, the negative impact of identity related stigma on the mental health of transgender older adults was attenuated by prior military service. Prior military service was a significant predictor of lower depressive symptomatology and higher psychological health related QOL in transgender older adults (Hoy-Ellis et al., 2017).

Furthermore, in a study by Fabbre (2015), male-to-female transgender older adults reported that the desire to fulfil societal gender expectations during youth and adulthood led them to consider or enrol in military service. Participants hoped that the masculine military environment would reduce their transgender feelings and reduce taunting for being feminine. Although military experience was reported to constrain gender identity, in later life it was viewed as a positive experience which contributed to the process of self-acceptance.

The Hoy-Ellis et al. (2017) study provides initial evidence that the negative impact of identity related stigma on the mental health of transgender older adults can be attenuated by prior military service. This was a good quality study that employed a rigorous method and analysis, using weighted multivariate linear models to evaluate the relationship between health related QOL, depressive symptomatology, identity stigma and prior military service. Prior military service is a potentially protective factor that warrants further investigation, especially considering 41% of transgender older adults reported previous military service in an American national community based survey (Fredriksen-Goldsen et al., 2013).

### **Psychosocial Factors**

### *Relationship status*

A negative impact of being partnered (married or in a relationship) was reported for transgender older adults in the old-old group (80+ years), but not for the middle-old or young-old groups (Fredriksen-Goldsen et al., 2015). This finding is not consistent with the positive impact of being partnered in the general older adult population (Namkee & Jung-Hwa, 2011) and the lesbian, gay, bisexual and transgender (LGBT) older adult population, when age group distinctions are not made (Williams & Fredriksen-Goldsen, 2014).

### *Victimisation and discrimination*

The studies that examine the mental health of transgender older adults consistently identify victimisation, particularly physical violence, as negatively impacting on mental health (Elder, 2016; Fredriksen-Goldsen et al., 2013; Fredriksen-Goldsen et al., 2015; McCann, Sharek, Higgins, Sheerin, & Glacken, 2013). The rates of victimisation and discrimination were highest among the young-old and lowest among the old-old participants. However, the negative impact on mental health was stronger in the young-old and old-old than in the middle-old (Fredriksen-Goldsen et al., 2015).

Those who had experienced victimisation were at higher risk of attempting suicide, according to survey data and qualitative interviews. In the McCann et al. (2013) study, 11% of the survey respondents had considered suicide in the previous year and 4% had self-harmed. Nineteen percent of survey respondents reported experiencing physical violence and 47% had experienced verbal insults related to their LGBT identity. Interview participants discussed suicidal ideation and self-harm as being related to victimisation and discrimination.



Victimisation and discrimination, as well as the fear of these experiences, was specifically linked to the social context of the 1940-50s in which many transgender older adults grew up (Elder, 2016). In this good quality thematic qualitative study, participants reported fearing being committed to an institution, being beaten or killed, throughout their adolescence/adulthood as society was not accepting of transgender people. Some participants spoke about attending vigils for transgender friends who died violent deaths, and linked these experiences to trauma and suicidal feelings.

The negative impact of victimisation and discrimination on transgender older adults' mental health is consistent with the broader literature relating to transgender adults and youth. Transgender individuals across the lifespan experience wide spread discrimination (housing, work, education & health care; Bradford, Reisner, Honnold, & Xavier, 2013).

#### *Gender identity disclosure*

Gender identity disclosure was consistently identified as negatively impacting on the mental health of transgender older adults, the impact was highest among the young-old and old-old age ranges (Elder, 2016; Fredriksen-Goldsen et al., 2015; McCann et al., 2013).

Findings from qualitative interviews seem to suggest that this could be linked to the potential or actual rejection and loss experienced following disclosure (McCann et al., 2013). Rejection by family members was also associated with higher rates (57%) of lifetime suicide attempts compared to those who did not experience rejection by family members (33%; Hass et al., 2014). Furthermore, gender identity concealment was shown to be associated with increased depressive symptomatology and perceived stress for transgender participants when compared to LGB older adults (Fredriksen-Goldsen et al., 2013). The emerging literature is suggestive of a 'catch 22' situation, where disclosing gender identity

could result in rejection and negatively impact on mental health, while at the same time concealing gender identity may be stressful and increase depressive symptoms. Gender identity disclosure might be further complicated by perceived recognition by others as transgender. Higher rates (45%) of suicide attempts were reported among transgender females (male to female) who perceived others could 'most of the time' tell they are transgender compared to those who perceived others could 'never' tell they are transgender (36%; Hass et al., 2014).

For transgender adults, the literature is less suggestive of a consistently negative impact of gender identity disclosure on mental health. Reports suggest a mixture of both positive and negative experiences, primarily influenced by the type of response received when the disclosure is made. In Galupo, Krum, Hagen, Gonzalez, and Bauerband's (2014) study, 55% of adult participants reported positive and supportive experiences of gender identity disclosure to friends that had a self-affirming effect. Negative experiences were reported by 25% of the participants, responses included believing the transgender identity was not genuine and refusing to use the chosen pronoun or name. Twenty three percent of adult participants reported indifferent responses to their disclosure. Positive impacts on mental health such as increased self-acceptance, self-confidence and higher levels of well-being have been reported when the disclosure was received positively (Lepore, Ragan, & Jones, 2000; Rodriguez & Kelly, 2006).

Awareness of transgender identity is developing at increasingly younger ages (Floyd & Bakeman, 2006). However, many studies show that the younger youth are less accepting of LGBT peers (Heinze & Horn, 2009; Poteat, Espelage, & Koenig, 2009). Despite this, gender disclosure during adolescence has been shown to be associated with lower levels of depression and higher levels of satisfaction and self-esteem. Although many transgender

youth experience school based victimisation that has a negative impact on their mental health, this may be outweighed by not having to conceal gender identity, which was also associated with higher levels of depression (Russell, Toomey, Ryan, & Diaz, 2014). For today's transgender youth, disclosing transgender identity during adolescence is associated with better mental health in early adulthood (Russell et al., 2014).

#### *Social support and community belonging*

Social support and a sense of community belonging were consistently found to have a positive impact on the mental health of transgender older adults and were significantly associated with lower levels of depressive symptomatology and perceived stress (Elder, 2016; Fredriksen-Goldsen et al. 2013; Fredriksen-Goldsen et al., 2015). Further insight was provided by the Fredriksen-Goldsen et al. (2013) study that compared transgender older adults to LGB older adults. Transgender older adults were shown to report lower levels of social support and community belonging than the LGB older adults, despite reporting larger social networks. There were inconsistent findings on the impact of social network size on mental health. Fredriksen-Goldsen et al. (2015) reported a positive impact associated with larger social network size while others found no effect. One possible explanation for the inconsistent findings is that while social network size is considered a measure for social support, it is more likely to be the quality and support provided by the social relationship that is important.

Active involvement in advocacy and teaching others in a variety of settings (churches, courts, hospitals, schools & conferences) about transgender related topics and issues was reported to be linked to resilience and had a positive impact on mental health (Elder, 2016). This linked to a sense of community involvement, especially when transgender older adults

were integrated with younger transgender individuals at events such as transgender marches.

Family, peer and community support have also been shown to be protective factors against suicide attempts and associated with better mental health outcomes for both transgender adults and youth (Budge & Adelson, 2013; Liu & Mustanski, 2012; Moody & Smith, 2013; Snapp, Watson, Russell, Diaz, & Ryan, 2015). Family acceptance has been shown to be a particularly important predictor of self-esteem for youth (Snapp et al., 2015). In a longitudinal study (McConnell, Birkett, & Mustanski, 2016), younger people who received support from their family and peers reported significantly lower levels of psychological distress than those who did not receive support from family or peers. Youth who received support from peers but not family, reported higher levels of psychological distress than those who did receive family support. However, they also reported increasing levels of family support throughout adolescence into early adulthood, and reduced psychological distress associated with the increase in family support.

Additionally, in a study by Witten (2014) transgender individuals over the age of 18 reported concerns and worries about lacking social support in later life. Concerns included reduced independence and that if later life care was required their gender related choices and wishes would not be respected; for example, their preferred pronoun and name would not be put on their death certificate or gravestone. Other concerns included not being able to live the final years of their lives as their true selves and suicide or euthanasia was spoken about as preferable to returning to a non-gender affirming identity. These concerns resulted in a lack of end of life preparation, such as the use of advance directives, wills and power of attorney options.

### *Internalised stigma*

Although only addressed in two studies, higher levels of internalised stigma were found to be significantly associated with higher depressive symptomatology and perceived stress (Fabbre, 2015; Fredriksen-Goldsen et al., 2013). In one study (Fredriksen-Goldsen et al., 2013) transgender older adults were compared to non-transgender LGB older adults. While internalised stigma had a negative impact on the mental health of LGB older adults, the rates of internalised stigma were higher among transgender older adults.

Internalised stigma has been conceptualised as resulting from minority stressors, such as exposure to high levels of victimisation and discrimination due to transgender identity. Given that transgender older adults may have been exposed to victimisation and discrimination for many years, and have lived during a time when being transgender was less accepted than it is today, it seems likely that internalised stigma is a relevant factor that warrants further exploration.

### *Development of a cohesive self and self-acceptance*

The development of a cohesive self and self-acceptance has been reported to have a positive impact on the mental health of transgender older adults (Ettner & Wylie, 2013; Fabbre, 2015). In a good quality extended case qualitative study by Fabbre (2015), transgender older adults reported being acutely aware of how their gender identities did not match the heteronormative expectations of Western society. They reported spending many years trying to 'fit' into expected gender roles and feeling as though they had failed when this was not achieved. This was associated with feelings of anger, and frustration and reported to negatively impact on mental health. However, 'failing' to fulfil society's expectations was seen as an important step towards self-acceptance and living authentically

according to their own expectations. This self-acceptance was associated with improved mental health, self-confidence and a sense of peace. Transgender older adults reported the development of a cohesive self that integrated previous attempts to 'fit' societal gender expectations with their self-accepted gender identity. As well as self-acceptance of gender identity, a positive sense of sexual identity was reported to be associated with better mental health QOL among transgender older adults (Fredriksen-Goldsen et al., 2015).

### *Spiritual/religious activity*

Spiritual/religious activity was not found to have a significant impact on depressive symptomatology, perceived stress or successful aging for transgender older adults (Fredriksen-Goldsen et al., 2013; Porter, Ronneberg & Witten, 2013).

Although little research has addressed religion and spirituality among transgender adults and youth, the available literature suggests both benefits and risks to mental health. In a qualitative study by Westerfield (2013), transgender adults reported a connection with a higher power and viewing themselves as being made as transgender by God. They also reported concern regarding how others with religious beliefs would perceive them and had experienced both supportive and discriminatory responses. In another qualitative study (Dahl & Galliher, 2012), transgender youths reported mixed experiences of religious involvement. Negative experiences included feelings of inadequacy, religious-based guilt, depressive symptoms and social strain. Positive experiences included building resilience, increased sense of self, enhanced acceptance of others, religious values and social support.

Similarly, religion has been reported to be both protective and a risk factor among LGB individuals. Religion has been associated with higher levels of internalised homophobia, resulting in a negative impact on mental health and wellbeing (Kralovec, Fartacek, Fartacek,

& Plöderl, 2014). Two-thirds of sexual minorities have reported experiencing conflict between their religious and sexual identities (Dahl & Galliher, 2009).

It is widely accepted that religion and spirituality have a positive impact on the wellbeing, physical and mental health of the general older adult population. Religion and spirituality have been reported to impact on relationship satisfaction (Cohen & Koenig, 2003), to bring meaning to life (Emery & Pargament, 2004) and to help adjustment to changes associated with aging (Sadler & Biggs, 2006). Religion has also been shown to be a protective factor against suicide in the general population (Colucci & Martin, 2008; Gearing & Lizardi, 2009).

### **Biomedical Factors**

#### *Physical health and health related behaviours*

Chronic health conditions were shown to negatively impact on mental health QOL (Fredriksen-Goldsen et al., 2015). This negative impact was highest among the young-old and old-old participants. This is consistent with the general older adult population (Braden et al., 2008).

Fear of accessing services and lack of physical activity were shown to mediate the relationship between gender identity and depressive symptomatology and perceived stress in transgender older adults (Fredriksen-Goldsen et al., 2013). This was consistent with a follow up study by Fredriksen-Goldsen et al. (2015) that showed physical activity and routine health check-ups were positively associated with mental health QOL. Leisure activity and no substance misuse were also shown to positively impact on mental health QOL which

was consistent with qualitative interviews that suggested substance misuse negatively impacted on the mental health of transgender older adults (McCann et al., 2013).

#### *Transition related health care*

Higher rates of lifetime suicide attempts were reported by older participants who had received or wanted to receive transition related health care like hormone treatment and surgery (Hass et al., 2014). However, this study had several methodological limitations and it was unclear what percentage of attempts occurred before or after transition related health care. This finding is not consistent with the broader literature that suggests the use of transition related health care is associated with improved mental health and reduced suicide risk for transgender adults (Bailey, Ellis, & McNeil, 2014; Wilsen, Chen, Arayasirikul, Winzel, & Raymond, 2015). Furthermore, in a qualitative study by Elder (2016), transgender older adults reported waiting until they were in their 50s before pursuing transition related health care due to the extreme levels of stigma, discrimination and victimisation that they experienced in the 1940-50s. Older participants associated delaying, having to wait to transition and the increased physical complications with transitioning later in life with poorer mental health (Elder, 2016).

Ettner and Wylie (2013) discussed that male-to-female transgender older adults who are not receiving hormone treatment may be at risk of untreated testosterone deficiency, which can lead to depression. Although testosterone decline is a typical part of aging, testosterone deficiency in male-to-female transgender older adults may be missed by health care professionals. This bio-medical factor warrants further exploration.



## Discussion

Today's transgender older adults aged in a social context that viewed them as mentally ill and subjected them to brutal treatments such as castration, lobotomies and involuntary institutionalization. Extreme victimisation and discrimination began to reduce in the 1970's and society has gradually become more supportive and accepting over the last few decades. This historical social context has had primarily negative implications for today's transgender older adults. This is key for understanding the factors that may impact on their mental health, and how this may differ from transgender adults and youth developing in a more accepting society. This review highlights the complex nature of the mental health of transgender older adults. There are likely to be many factors that directly or indirectly affect the wellbeing and mental health of this group. There is a complex web of interactions between factors for individuals identifying as transgender and this review has aimed to begin to identify some of these factors rather than explore interactions in great detail.

The finding that being partnered had a negative impact only on the old-old transgender older adults is perhaps better understood when considered in relation to historical social context. It may be that the social context in which the old-old group grew up in, had an influence on their decisions to conceal their gender identity and get married or have a relationship that fitted with the social expectations associated with their birth sex. This may have been less likely to happen for younger generations who grew up in a more accepting social context and were more able to have gender affirming relationships. The lower rates of victimisation and discrimination among the old-old group but higher levels of negative impact on mental health might be related to this. There are numerous potential explanations that require further investigation. This finding perhaps indicates differences in

coping between the age groups, different types of victimisation or different support availability. Another possible explanation is that reluctance to disclose gender identity in the old-old group due to historic social context, has protected them from some victimisation.

A negative impact of gender identity disclosure was consistently identified for transgender older adults, which was at odds with the mixed but generally more positive experiences reported by transgender adults and youth. This important difference is again better understood by considering the less accepting historic social context. It could be that due to the extreme levels of discrimination and victimisation experienced by transgender persons in the 1940-50s, transgender older adults' experiences of disclosing their gender identity have been generally more negative.

The positive impact of social support and a sense of community belonging on the mental health of transgender older adults was consistent with the positive impact of social support reported in the transgender adult and youth literature. However, the more extensive transgender adult and youth literature includes more specific findings, such as the positive impact of family acceptance and support that exceeds the positive impact of peer support. The impact of different forms of support (e.g. family, peer, partner, community & combinations of support) on the mental health of transgender older adults has yet to be investigated.

### **Limitations**

This review includes only a small number of studies that use a range of methods, including selective recruitment methods from which the results cannot be generalized. Study samples may be biased to transgender older adults who are more confident and feel able to disclose their gender identity. Those who are the most at risk or who are

experiencing mental health difficulties may be under represented in study samples, particularly the old-old age group. Although the factors have been categorised as biomedical, psychosocial and socioeconomic according to conceptualisations in the literature (Ettner & Wylie, 2013; Witten 2016), some of the factors are complex and interlinking and could be conceptualised as fitting into more than one of the categories.

It was beyond the scope of this review to critique the theoretical underpinnings of the research that addresses the mental health of transgender older adults, such as heteronormative models of aging and social theories that pathologise non-normative gender identities. However, it is likely that the structuring of services, health interventions and research based on these heteronormative models also impacts on the mental health of transgender individuals.

The broad umbrella term 'lesbian, gay, bisexual and transgender' (LGBT) is used to describe a diverse group of individuals whose sexual and/or gender identity does not match that which is usually associated with the individual's birth sex. Some of the studies included in this review used LGBT samples without distinguishing between gender identity and sexual orientation. Although the more extensive LGB literature has proved to be a valuable starting place for research focusing on transgender related issues, it is increasingly becoming apparent that there are important differences between transgender individuals' and LGB individuals' mental health. These differences warrant independent research and potentially have implications for clinical practice.

### **Clinical implications and directions for future research**

This review identifies and summarizes some of the factors identified in a small number of studies that appear to impact on the mental health of transgender older adults. It

highlights that transgender older adults may have unique mental health needs that differ from the general older adult population and transgender adults and youth, such as the impact of delaying transition until later in life due to the experience of extreme discrimination and victimisation in the 1940-50s. These needs and cohort effects should be taken into consideration by health and social care organisations involved in the mental health care provision of transgender older adults. Staff working with transgender older adults need to be aware that their views on transitioning might differ from their clients and that a common ground needs to be found for developing a therapeutic relationship. Factors such as religion and being in a partnership, which are considered to be protective in most general older adult research (Emery & Pargament, 2004; Sadler & Biggs, 2006), might not fulfil this role in transgender older adults. Specific issues raised by transgender adults about end of life care require attention in health care provision as well as care settings.

Identifying factors that have a positive impact on the mental health of transgender older adults specifically is crucial for tailoring interventions that promote good mental health and successful aging. Potential positive factors i.e. a positive sense of sexual identity, advocacy, and prior military service, require further exploration. Additional research to verify the impact of these factors on the mental health of transgender older adults (and transgender individuals of all ages) would be beneficial and could have useful implications; for example, health and social services may be able to facilitate their involvement in advocacy.

## References

- Bailey, L., Ellis, S. J., & McNeil, J. (2014). Suicide risk in the UK trans population and the role of gender transition in decreasing suicidal ideation and suicide attempt. *Mental Health Review Journal, 19*(4), 209-220. doi:10.1108/MHRJ-05-2014-0015
- Braden, J. B., Zhang, L., Fan, M., Unützer, J., Edlund, M. J., & Sullivan, M. D. (2008). Mental health service use by older adults: The role of chronic pain. *The American Journal of Geriatric Psychiatry, 16*(2), 156-167. doi: 10.1097/JGP.0b013e31815a3ea9
- Bradford, J., Reisner, S. L., Honnold, J. A., & Xavier, J. (2013). Experiences of transgender-related discrimination and implications for health: Results from the Virginia Transgender Health Initiative Study. *American Journal of Public Health, 103*, 1820–1829. doi:10.2105/AJPH.2012.300796
- Budge, S. L., & Adelson, J. L. (2013). Anxiety and depression in transgender individuals: The roles of transition status, loss, social support, and coping. *Journal of Consulting and Clinical Psychology, 81*(3), 545-557. doi:10.1037/a0031774
- Clark, T. C., Lucassen, M. F. G., Bullen, P., Denny, S. J., Fleming, T. M., Robinson, E. M., & Rossen, F. V. (2014). The health and well-being of transgender high school students: Results from the New Zealand adolescent health survey (Youth'12). *Journal of Adolescent Health, 55*, 93-99. doi.org/10.1016/j.jadohealth.2013.11.008
- Clements-Nolle, K., Marx, R., & Katz, M. (2006). Attempted suicide among transgender persons: The influence of gender-based discrimination and victimization. *Journal of Homosexuality, 51*(3), 53–69. doi:10.1300/J082v51n03\_04

- Cohen, A. B., & Koenig, H. G. (2003). Religion, religiosity and spirituality in the biopsychosocial model of health and ageing. *Ageing International*, 28(3), 215–241. doi:10.1007/s12126-002-1005-1.
- Colucci, E., & Martin, G. (2008). Religion and spirituality along the suicidal path. *Suicide and Life-Threatening Behavior*, 38(2), 229–244. doi:10.1521/suli.2008.38.2.229
- Conron, K. J., Scott, G., Stowell, G. S., & Landers, S. J. (2012). Transgender health in Massachusetts: Results from a household probability sample of adults. *American Journal of Public Health*, 102, 118-122. doi:10.2105/AJPH.2011.300315
- Dahl, A., & Galliher, R. V. (2009). LGBTQ young adult experiences of religious and sexual identity integration. *Journal of LGBT Issues in Counseling*, 3, 92–112. doi:10.1080/15538600903005268
- Dahl, A. L., & Galliher, R. V. (2012). LGBTQ adolescents and young adults raised within a Christian religious context: Positive and negative outcomes. *Journal of Adolescence*, 35, 1611-1618. doi:10.1016/j.adolescence.2012.07.003
- Department of Health. (2001). *National Service Framework for Older people*. Retrieved from [https://www.gov.uk/government/uploads/system/uploads/attachment\\_data/file/198033/National\\_Service\\_Framework\\_for\\_Older\\_People.pdf](https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/198033/National_Service_Framework_for_Older_People.pdf)
- Elder, A. B. (2016). Experiences of older transgender and gender nonconforming adults in psychotherapy: A qualitative study. *Psychology of Sexual Orientation and Gender Diversity*, 3(2), 180–186. doi:10.1037/sgd0000154

- Emery, E. E., & Pargament, K. I. (2004). The many faces of religious coping in late life: Conceptualization, measurement, and links to well-being. *Ageing International*, 29(1), 3–27. doi:10.1007/s12126-004-1007-2
- Ettner, R., & Wylie, K. (2013). Psychological and social adjustment in older transsexual people. *Maturitas*, 74, 226-229. doi:10.1016/j.maturitas.2012.11.011
- Fabbre, V. D. (2015). Gender transitions in later life: A queer perspective on successful aging. *The Gerontologist*, 55, 144-153. doi:10.1093/geront/gnu079
- Floyd, F. J., & Bakeman, R. (2006). Coming-out across the life course: Implications of age and historical context. *Archives of Sexual Behavior*, 35(3), 287-296. doi:10.1007/s10508-006-9022-x
- Frech, A., & Damaske, S. (2012). The relationships between mothers' work pathways and physical and mental health. *Journal of Health and Social Behavior*, 53(4), 396–412. doi:org/10.1177/0022146512453929
- Fredriksen-Goldsen, K. I., Cook-Daniels, L., Kim, H., Erosheva, E. A., Emler, C. A., Hoy-Ellis, C. P., Goldsen, J., & Muraco, A. (2013). Physical and mental health of transgender older adults: An at-risk and underserved population. *The Gerontologist*, 54(3), 488-500. doi: 10.1093/geront/gnt021
- Fredriksen-Goldsen, K. I., Kim, H., Shiu, C., Goldsen, J., & Emler, C. A. (2015). Successful aging among LGBT older adults: Physical and mental health-related quality of life by age group. *The Gerontologist*, 55(1), 154-168. doi:10.1093/geront/gnu081

- Galupo, M. P., Krum, T. E., Hagen, D. B., Gonzalez, K. A., & Bauerband, L. A. (2014). Disclosure of transgender identity and status in the context of friendship. *Journal of LGBT issues in counseling, 8*, 25-42. doi:10.1080/15538605.2014.853638
- Gearing, R. E., & Lizardi, D. (2009). Religion and suicide. *Journal of Religion and Health, 48*(2), 332-341. doi:10.1007/s10943-008-9181-2
- Hass, A. P., Rodgers, P. L., & Herman, J. L. (2014). *Suicide attempts among transgender and gender non-conforming adults; Findings of the national transgender discrimination survey*. Retrieved from <http://williamsinstitute.law.ucla.edu/wp-content/uploads/AFSP-Williams-Suicide-Report-Final.pdf>
- Heinze, J. E., & Horn, S. S. (2009). Intergroup contact and beliefs about homosexuality in adolescence. *Journal of Youth and Adolescence, 38*(7), 937-951. doi:10.1007/s10964-009-9408-x
- Hergenrather, K. C., Zeglin, R. J., McGuire-Kuletz, M., & Rhodes, S. D. (2015). Employment as a social determinant of health: A review of longitudinal studies exploring the relationship between employment status and mental health. *Rehabilitation Research, Policy, and Education, 29*(3), 261-290. doi.org/10.1891/2168-6653.29.3.261
- Hoy-Ellis, C. P., Shiu, C., Sullivan, K. M., Kim, H., Sturges, A. M., & Fredriksen-Goldsen, K. I. (2017). Prior military service, identity stigma, and mental health among transgender older adults. *The Gerontologist, 57*, 63-71. doi:10.1093/geront/gnw173
- Kessler, R. C., Berglund, P., Demler, O., Jin, R., Merikangas, K. R., & Walters, E. E. (2005). Lifetime prevalence and age-of-onset distributions of *DSM-IV* disorders in the



- National Comorbidity Survey Replication. *Archives of General Psychiatry*, 62, 593–602. doi:10.1001/archpsyc.62.6.593
- Kralovec, K., Fartacek, C., Fartacek, R., & Plöderl, M. (2014). Religion and suicide risk in lesbian, gay and bisexual Austrians. *Journal of Religion and Health*, 53, 413–423. doi:10.1007/s10943-012-9645-2
- Lepore, S. J., Ragan, J. D., & Jones, S. (2000). Talking facilitates cognitive-emotional processes of adaptation to an acute stressor. *Journal of Personality and Social Psychology*, 78, 499–508. doi:10.1037//0022-3514.78.3.499
- Liu, R. T., & Mustanski, B. (2012). Suicidal ideation and self-harm in lesbian, gay, bisexual, and transgender youth. *American Journal of Preventive Medicine*, 42(3), 221–228. doi:10.1016/j.amepre.2011.10.023
- Mandal, B., & Roe, B. (2008). Job loss, retirement and the mental health of older Americans. *Journal of Mental Health Policy and Economics*, 11(4), 167–176. doi:10.2139/ssrn.991134
- McCann, E., Sharek, D., Higgins, A., Sheerin, F., & Glacken, M. (2013). Lesbian, gay, bisexual and transgender older people in Ireland: Mental health issues. *Aging and Mental Health*, 17(3), 358–365. doi:10.1080/13607863.2012.751583
- McConnell, E. A., Birkett, M., & Mustanski, B. (2016). Families matter: Social support and mental health trajectories among lesbian, gay, bisexual, and transgender youth. *Journal of Adolescent Health*, 59(6), 674–680. doi:10.1016/j.jadohealth.2016.07.026

- Meerwijk, E. L., & Sevelius, J. M. (2017). Transgender population size in the United States: A meta-regression of population-based probability samples. *American Journal of Public Health, 107*(2), 216. doi:10.2105/AJPH.2016.303578a.
- Moody, C., & Smith, N. G. (2013). Suicide protective factors among trans adults. *Archives of Sexual Behavior, 42*(5), 739–752. doi:10.1007/s10508-013-0099-8
- Namkee, G. C., & Jung-Hwa, H. (2011). Relationship between spouse/partner support and depressive symptoms in older adults: Gender difference. *Aging & Mental Health, 15*(3), 307-317. doi:10.1080/13607863.2010.513042
- Nemoto, T., Bödeker, B., & Iwamoto, M. (2011). Social support, exposure to violence and transphobia, and correlates of depression among male-to-female transgender women with a history of sex work. *American Journal of Public Health, 101*, 1980–1988. doi:10.2105/AJPH.2010.197285
- Persson, D. I. (2009). Unique challenges of transgender aging: Implications from the literature. *Journal of Gerontological Social Work, 52*(6), 633-646.  
doi:10.1080/01634370802609056
- Porter, K. E., Ronneberg, C. R., & Witten, T. M. (2013). Religious affiliation and successful aging among transgender older adults: Findings from the Trans Metlife survey. *Journal of religion, Spirituality & Aging, 25*, 112-138.  
doi:10.1080/15528030.2012.739988
- Poteat, V. P., Espelage, D. L., & Koenig, B. W. (2009). Willingness to remain friends and attend school with lesbian and gay peers: Relational expressions of prejudice among

heterosexual youth. *Journal of Youth and Adolescence*, 38(7), 952–962.

doi:10.1007/s10964-009-9416-x

Prause, J., & Dooley, D. (2001). Favourable employment status change and psychological depression: A two-year follow-up analysis of the National Longitudinal Survey of Youth. *Applied Psychology: An International Review*, 50(2), 282–304. doi .org/10.1111/1464-0597.00059

Reed, B., Rhodes, S., Schofield, P., & Wylie, K. (2009). *Gender variance in the UK: Prevalence, incidence, growth and geographic distribution*. Retrieved from Gender Identity Research and Education Society website: <http://gires.org.uk/assets/Medpro-Assets/GenderVarianceUK-report.pdf>

Robert, S. A., Cherepanov, D., Palta, M., Dunham, N. C., Feeny, D., & Fryback, D. G. (2009). Socioeconomic status and age variations in health-related quality of life: Results from the National Health Measurement Study. *Journals of Gerontology Series B- Psychological Sciences & Social Sciences*, 64B(3), 378-389. doi:10.1093/geronb/gbp012

Rodriguez, R. R., & Kelly, A. E. (2006). Health effects of disclosing secrets to imagined accepting versus nonaccepting confidants. *Journal of Social & Clinical Psychology*, 25(9), 1023–1047. doi: 10.1521/jscp.2006.25.9.1023

Rotondi, N. K., Bauer, G. R., Travers, R., Travers, A., Scanlon, K., Kaay, M. (2011). Depression in male-to-female transgender Ontarians: Results from the Trans PULSE project. *Canadian Journal of Community Mental Health*, 30(2), 113-133. doi:10.7870/cjcmh-2011-0020

- Russell, S. T., Toomey, R. B., Ryan, C., & Diaz, R. M. (2014). Being out at school: The implications for school victimization and young adult adjustment. *American Journal of Orthopsychiatry, 84*(6), 635-643. doi:10.1037/ort0000037
- Sadler, E., & Biggs, S. (2006). Exploring the links between spirituality and 'successful ageing'. *Journal of Social Work Practice, 20*(3), 267–280. doi:10.1080/02650530600931757.
- Shields, J.P., Cohen, R., Glassman, J. R., Whitaker, K., Franks, H., & Bertolini, I. (2013). Estimating population size and demographic characteristics of lesbian, gay, bisexual, and transgender youth in middle school. *Journal of Adolescent Health, 52*(2), 248-50. doi:10.1016/j.jadohealth.2012.06.016
- Snapp, S. D., Watson, R. J., Russell, S. T., Diaz, R. M., & Ryan, C. (2015). Social support networks for LGBT young adults: Low cost strategies for positive adjustment. *Family Relations, 64*(3), 420-430. doi:10.1111/fare.12124
- Spiro, A., Settersten, R. A., & Aldwin, C. M. (2016). Long-term outcomes of military service in aging and the life course: A positive re-envisioning. *The Gerontologist, 56*, 1–13. doi:10.1093/geront/gnw038
- Van Gorp, W. G., Rabkin, J. G., Ferrando, S. J., Mintz, J., Ryan, E., Borkowski, T., & Mcelhiney, M. (2007). Neuropsychiatric predictors of return to work in HIV/AIDS. *Journal of the International Neuropsychological Society, 13*(1), 80–89. doi:org/10.1017/S1355617707070117
- Westerfield, E. M. (2013). *Transgender peoples' experiences of religion and spirituality* (Master's thesis). Retrieved from ProQuest, Ann Arbor MI (AAI1514854).

Williams, M. E., & Fredriksen-Goldsen, K. I. (2014). Same-sex partnerships and the health of older adults. *Journal of Community Psychology, 42*, 558–570.

doi:10.1002/jcop.21637

Wilmoth, J. M., London, A. S., & Parker, W. M. (2010). Military service and men's health trajectories in later life. *The Journals of Gerontology, Series B: Psychological Sciences and Social Sciences, 65*, 744–755. doi:10.1093/geronb/gbq072

Wilson, E. C., Chen, Y. H., Arayasirikul, S., Wenzel, C., & Raymond, H. F. (2015). Connecting the dots: Examining transgender women's utilization of transition-related medical care and associations with mental health, substance use, and HIV. *Journal of Urban Health, 92*(1), 182–92. doi:10.1007/s11524-014-9921-4.

Witten, T. M. (2014). End of life, chronic illness, and trans-identities. *Journal of Social Work in End-of-Life & Palliative Care, 10*, 1–26. doi:10.1080/15524256.2013.877864

Witten, T. M. (2016). The intersectional challenges of aging and of being a gender non-conforming adult. *Journal of the American Society of Aging, 40*(2), 63–70. Retrieved from <http://search.proquest.com/openview/2d7da3f7d27ccbbae50e12a18e7ad2fc/1?pq-origsite=gscholar&cbl=30306>

## **Empirical Paper: Guidance for Authors**

## Your Paper Your Way

We now differentiate between the requirements for new and revised submissions. You may choose to submit your manuscript as a single Word or PDF file to be used in the refereeing process. Only when your paper is at the revision stage, will you be requested to put your paper in to a 'correct format' for acceptance and provide the items required for the publication of your article.

**To find out more, please visit the Preparation section below.**



### Introduction

Click [here](#) for guidelines on Special Issues.

Click [here](#) for guidelines on Qualitative methods.

*Social Science & Medicine* provides an international and interdisciplinary forum for the dissemination of social science research on health. We publish original research articles (both empirical and theoretical), reviews, position papers and commentaries on health issues, to inform current research, policy and practice in all areas of common interest to social scientists, health practitioners, and policy makers. The journal publishes material relevant to any aspect of health and healthcare from a wide range of social science disciplines (anthropology, economics, epidemiology, geography, policy, psychology, and sociology), and material relevant to the social sciences from any of the professions concerned with physical and mental health, health care, clinical practice, and health policy and the organization of healthcare. We encourage material which is of general interest to an international readership.

### *Journal Policies*

The journal publishes the following types of contribution:

- 1) Peer-reviewed original research articles and critical analytical reviews in any area of social science research relevant to health and healthcare. These papers may be up to 8000 words including abstract, tables, figures, references and (printed) appendices as well as the main text. Papers below this limit are preferred.
- 2) Peer-reviewed short communications of findings on topical issues or published articles of between 2000 and 4000 words.
- 3) Submitted or invited commentaries and responses debating, and published alongside, selected articles (please select the article type 'Discussion' when submitting a Commentary).
- 4) Special Issues bringing together collections of papers on a particular theme, and usually [guest edited](#).

Due to the high number of submissions received by *Social Science & Medicine*, Editorial Offices are not able to respond to questions regarding the appropriateness of new papers for the journal. If you are unsure whether or not your paper is within scope, please take some time to review previous issues of the journal and the Aims and Scope at

<https://www.journals.elsevier.com/social-science-and-medicine/>.

## Submission checklist

You can use this list to carry out a final check of your submission before you send it to the journal for review. Please check the relevant section in this Guide for Authors for more details.

### Ensure that the following items are present:

One author has been designated as the corresponding author with contact details:

- E-mail address
- Full postal address

All necessary files have been uploaded:

*Manuscript:*

- Include keywords
- All figures (include relevant captions)
- All tables (including titles, description, footnotes)
- Ensure all figure and table citations in the text match the files provided
- Indicate clearly if color should be used for any figures in print

*Graphical Abstracts / Highlights files* (where applicable)

*Supplemental files* (where applicable)

Further considerations

- Manuscript has been 'spell checked' and 'grammar checked'
- All references mentioned in the Reference List are cited in the text, and vice versa
- Manuscript does not exceed the word limit
- All identifying information has been removed from the manuscript, including the file name itself
- Permission has been obtained for use of copyrighted material from other sources (including the Internet)
- Relevant declarations of interest have been made
- Journal policies detailed in this guide have been reviewed
- Referee suggestions and contact details provided, based on journal requirements

For further information, visit our [Support Center](#).



## Before You Begin

### Ethics in Publishing

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/ethicalguidelines>.

Please note that any submission that has data collected from human subjects requires ethics approval. If your manuscript does not include ethics approval, your paper will not be sent out for review.

### Declaration of interest



All authors are requested to disclose any actual or potential conflict of interest including any financial, personal or other relationships with other people or organizations within three years of beginning the submitted work that could inappropriately influence, or be perceived to influence, their work. [More information](#).

### **Submission declaration and verification**

Submission of an article implies that the work described has not been published previously (except in the form of a conference abstract or as part of a published lecture or thesis for an academic qualification), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. To verify originality, your article may be checked by the originality detection software iThenticate. See also <http://www.elsevier.com/editors/plagdetect>.

### **Changes to authorship**

Authors are expected to consider carefully the list and order of authors **before** submitting their manuscript and provide the definitive list of authors at the time of the original submission. Any addition, deletion or rearrangement of author names in the authorship list should be made only **before** the manuscript has been accepted and only if approved by the journal Editor. To request such a change, the Editor must receive the following from the **corresponding author**: (a) the reason for the change in author list and (b) written confirmation (e-mail, letter) from all authors that they agree with the addition, removal or rearrangement. In the case of addition or removal of authors, this includes confirmation from the author being added or removed.

Only in exceptional circumstances will the Editor consider the addition, deletion or rearrangement of authors **after** the manuscript has been accepted. While the Editor considers the request, publication of the manuscript will be suspended. If the manuscript has already been published in an online issue, any requests approved by the Editor will result in a corrigendum.

### ***Article transfer service***

This journal is part of our Article Transfer Service. This means that if the Editor feels your article is more suitable in one of our other participating journals, then you may be asked to consider transferring the article to one of those. If you agree, your article will be transferred automatically on your behalf with no need to reformat. Please note that your article will be reviewed again by the new journal. [More information](#).

### **Copyright**

Upon acceptance of an article, authors will be asked to complete a 'Journal Publishing Agreement' (see [more information](#) on this). An e-mail will be sent to the corresponding author confirming receipt of the manuscript together with a 'Journal Publishing Agreement' form or a link to the online version of this agreement.

Subscribers may reproduce tables of contents or prepare lists of articles including abstracts for internal circulation within their institutions. [Permission](#) of the Publisher is required for

resale or distribution outside the institution and for all other derivative works, including compilations and translations. If excerpts from other copyrighted works are included, the author(s) must obtain written permission from the copyright owners and credit the source(s) in the article. Elsevier has [preprinted forms](#) for use by authors in these cases.

For open access articles: Upon acceptance of an article, authors will be asked to complete an 'Exclusive License Agreement' ([more information](#)). Permitted third party reuse of open access articles is determined by the author's choice of [user license](#).

### ***Author rights***

As an author you (or your employer or institution) have certain rights to reuse your work. [More information](#).

### ***Elsevier supports responsible sharing***

Find out how you can [share your research](#) published in Elsevier journals.

### **Role of the funding source**

You are requested to identify who provided financial support for the conduct of the research and/or preparation of the article and to briefly describe the role of the sponsor(s), if any, in study design; in the collection, analysis and interpretation of data; in the writing of the articles; and in the decision to submit it for publication. If the funding source(s) had no such involvement then this should be stated. Please see <http://www.elsevier.com/funding>.

### ***Funding body agreements and policies***

Elsevier has established a number of agreements with funding bodies which allow authors to comply with their funder's open access policies. Some funding bodies will reimburse the author for the Open Access Publication Fee. Details of [existing agreements](#) are available online.

### **Open access**

This journal offers authors a choice in publishing their research:

#### ***Open access***

- Articles are freely available to both subscribers and the wider public with permitted reuse.
- An open access publication fee is payable by authors or on their behalf, e.g. by their research funder or institution.

#### ***Subscription***

- Articles are made available to subscribers as well as developing countries and patient groups through our [universal access programs](#).
- No open access publication fee payable by authors.

Regardless of how you choose to publish your article, the journal will apply the same peer review criteria and acceptance standards.

For open access articles, permitted third party (re)use is defined by the following [Creative Commons user licenses](#):

***Creative Commons Attribution (CC BY)***

Lets others distribute and copy the article, create extracts, abstracts, and other revised versions, adaptations or derivative works of or from an article (such as a translation), include in a collective work (such as an anthology), text or data mine the article, even for commercial purposes, as long as they credit the author(s), do not represent the author as endorsing their adaptation of the article, and do not modify the article in such a way as to damage the author's honor or reputation.

***Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)***

For non-commercial purposes, lets others distribute and copy the article, and to include in a collective work (such as an anthology), as long as they credit the author(s) and provided they do not alter or modify the article.

The open access publication fee for this journal is **USD 3200**, excluding taxes. Learn more about Elsevier's pricing policy: <https://www.elsevier.com/openaccesspricing>.

***Green open access***

Authors can share their research in a variety of different ways and Elsevier has a number of green open access options available. We recommend authors see our [green open access page](#) for further information. Authors can also self-archive their manuscripts immediately and enable public access from their institution's repository after an embargo period. This is the version that has been accepted for publication and which typically includes author-incorporated changes suggested during submission, peer review and in editor-author communications. Embargo period: For subscription articles, an appropriate amount of time is needed for journals to deliver value to subscribing customers before an article becomes freely available to the public. This is the embargo period and it begins from the date the article is formally published online in its final and fully citable form. [Find out more](#).

This journal has an embargo period of 36 months.

***Elsevier Publishing Campus***

The Elsevier Publishing Campus ([www.publishingcampus.com](http://www.publishingcampus.com)) is an online platform offering free lectures, interactive training and professional advice to support you in publishing your research. The College of Skills training offers modules on how to prepare, write and structure your article and explains how editors will look at your paper when it is submitted for publication. Use these resources, and more, to ensure that your submission will be the best that you can make it.

***Language (usage and editing services)***

Please write your text in good English (American or British usage is accepted, but not a mixture of these). Authors who feel their English language manuscript may require editing to eliminate possible grammatical or spelling errors and to conform to correct scientific English may wish to use the [English Language Editing service](#) available from Elsevier's WebShop.

**Submission**

Submission to this journal occurs online and you will be guided step by step through the creation and uploading of your files. Please submit your article via <http://ees.elsevier.com/ssm>. The system automatically converts source files to a single PDF file of the article, which is used in the peer-review process. Please note that even though

manuscript source files are converted to PDF files at submission for the review process, these source files are needed for further processing after acceptance. All correspondence, including notification of the Editor's decision and requests for revision, takes place by e-mail.

## Reviewers

Please provide the names and email addresses of 3 potential reviewers and state the reason for each suggestion. Colleagues within the same institution and co-authors within the last 5 years should not be included in the suggestions. Note that the editor retains the sole right to decide whether or not the suggested reviewers are used.

## Additional information

Please note author information is entered into the online editorial system (EES) during submission and must *not* be included in the manuscript itself.

*Social Science & Medicine* does not normally list more than six authors to a paper, and special justification must be provided for doing so. Further information on criteria for authorship can be found in *Social Science & Medicine*, 2007, 64(1), 1-4.

Authors should approach the Editors in Chief if they wish to submit companion articles.

Information about our peer-review policy can be found [here](#).

Please note that we may suggest accepted papers for legal review if it is deemed necessary.



## Preparation

### NEW SUBMISSIONS

Submission to this journal proceeds totally online and you will be guided stepwise through the creation and uploading of your files. The system automatically converts your files to a single PDF file, which is used in the peer-review process.

As part of the Your Paper Your Way service, you may choose to submit your manuscript as a single file to be used in the refereeing process. This can be a PDF file or a Word document, in any format or lay-out that can be used by referees to evaluate your manuscript. It should contain high enough quality figures for refereeing. If you prefer to do so, you may still provide all or some of the source files at the initial submission. Please note that individual figure files larger than 10 MB must be uploaded separately.

### References

There are no strict requirements on reference formatting at submission. References can be in any style or format as long as the style is consistent. Where applicable, author(s) name(s), journal title/book title, chapter title/article title, year of publication, volume number/book chapter and the pagination must be present. Use of DOI is highly encouraged. The reference style used by the journal will be applied to the accepted article by Elsevier at the proof stage. Note that missing data will be highlighted at proof stage for the author to correct.

### ***Formatting Requirements***

The journal operates a double blind peer review policy. For guidelines on how to prepare your paper to meet these criteria please see the [attached guidelines](#). The journal requires that your manuscript is submitted with double spacing applied. There are no other strict formatting requirements but all manuscripts must contain the essential elements needed to convey your manuscript, for example Abstract, Keywords, Introduction, Materials and Methods, Results, Conclusions, Artwork and Tables with Captions.

If your article includes any Videos and/or other Supplementary material, this should be included in your initial submission for peer review purposes.

Divide the article into clearly defined sections.

## **REVISED SUBMISSIONS**

### ***Use of word processing software***

Regardless of the file format of the original submission, at revision you must provide us with an editable file of the entire article. Keep the layout of the text as simple as possible. Most formatting codes will be removed and replaced on processing the article. The electronic text should be prepared in a way very similar to that of conventional manuscripts (see also the [Guide to Publishing with Elsevier](#)). See also the section on Electronic artwork.

To avoid unnecessary errors you are strongly advised to use the 'spell-check' and 'grammar-check' functions of your word processor.

### **Essential cover page information**

The Cover Page should **only** include the following information:

- ***Title.*** Concise and informative. Titles are often used in information-retrieval systems. Avoid abbreviations and formulae where possible and make clear the article's aim and health relevance.
- ***Author names and affiliations in the correct order.*** Where the family name may be ambiguous (e.g., a double name), please indicate this clearly. Present the authors' affiliation addresses (where the actual work was done) below the names. Indicate all affiliations with a lower-case superscript letter immediately after the author's name and in front of the appropriate address. Provide the full postal address of each affiliation, including the country name and, if available, the e-mail address of each author.
- ***Corresponding author.*** Clearly indicate who will handle correspondence at all stages of refereeing and publication, also post-publication. **Ensure that telephone and fax numbers (with country and area code) are provided in addition to the e-mail address and the complete postal address. Contact details must be kept up to date by the corresponding author.**
- ***Present/permanent address.*** If an author has moved since the work described in the article was done, or was visiting at the time, a 'Present address' (or 'Permanent address') may be indicated as a footnote to that author's name. The address at which the author actually did the work must be retained as the main, affiliation address. Superscript Arabic numerals are used for such footnotes.

### ***Text***

In the main body of the submitted manuscript this order should be followed: abstract, main text, references, appendix, figure captions, tables and figures. Author details, keywords and acknowledgements are entered separately during the online submission process, as is the

abstract, though this is to be included in the manuscript as well. During submission authors are asked to provide a word count; this is to include ALL text, including that in tables, figures, references etc.

### *Title*

Please consider the title very carefully, as these are often used in information-retrieval systems. Please use a concise and informative title (avoiding abbreviations where possible). Make sure that the health or healthcare focus is clear.

### **Abstract**

An abstract of up to 300 words must be included in the submitted manuscript. An abstract is often presented separately from the article, so it must be able to stand alone. It should state briefly and clearly the purpose and setting of the research, the principal findings and major conclusions, and the paper's contribution to knowledge. For empirical papers the country/countries/locations of the study should be clearly stated, as should the methods and nature of the sample, the dates, and a summary of the findings/conclusion. Please note that excessive statistical details should be avoided, abbreviations/acronyms used only if essential or firmly established, and that the abstract should not be structured into subsections. Any references cited in the abstract must be given in full at the end of the abstract.

### **Research highlights**

Research highlights are a short collection of 3 to 5 bullet points that convey an article's **unique contribution to knowledge** and are placed online with the final article. We allow 85 characters per bullet point including spaces. They should be supplied as a separate file in the online submission system (further instructions will be provided there). You should pay very close attention to the formulation of the Research Highlights for your article. Make sure that they are **clear, concise and capture the reader's attention**. If your research highlights do not meet these criteria we may need to return your article to you leading to a delay in the review process.

### **Keywords**

Up to 8 keywords are entered separately into the online editorial system during submission, and should accurately reflect the content of the article. Again abbreviations/acronyms should be used only if essential or firmly established. For empirical papers the country/countries/locations of the research should be included. The keywords will be used for indexing purposes.

### **Methods**

Authors of empirical papers are expected to provide full details of the research methods used, including study location(s), sampling procedures, the date(s) when data were collected, research instruments, and techniques of data analysis. Specific guidance on the reporting of qualitative studies are provided [here](#).

Systematic reviews and meta-analyses must be reported according to [PRISMA](#) guidelines.

### **Footnotes**

There should be no footnotes or endnotes in the manuscript.

## Artwork

### *Electronic artwork*

#### *General points*

- Make sure you use uniform lettering and sizing of your original artwork.
- Preferred fonts: Arial (or Helvetica), Times New Roman (or Times), Symbol, Courier.
- Number the illustrations according to their sequence in the text.
- Use a logical naming convention for your artwork files.
- Indicate per figure if it is a single, 1.5 or 2-column fitting image.
- For Word submissions only, you may still provide figures and their captions, and tables within a single file at the revision stage.
- Please note that individual figure files larger than 10 MB must be provided in separate source files.

A detailed [guide on electronic artwork](#) is available.

**You are urged to visit this site; some excerpts from the detailed information are given here.**

#### *Formats*

Regardless of the application used, when your electronic artwork is finalized, please 'save as' or convert the images to one of the following formats (note the resolution requirements for line drawings, halftones, and line/halftone combinations given below):

EPS (or PDF): Vector drawings. Embed the font or save the text as 'graphics'.

TIFF (or JPG): Color or grayscale photographs (halftones): always use a minimum of 300 dpi.

TIFF (or JPG): Bitmapped line drawings: use a minimum of 1000 dpi.

TIFF (or JPG): Combinations bitmapped line/half-tone (color or grayscale): a minimum of 500 dpi is required.

#### **Please do not:**

- Supply files that are optimized for screen use (e.g., GIF, BMP, PICT, WPG); the resolution is too low.
- Supply files that are too low in resolution.
- Submit graphics that are disproportionately large for the content.

### *Color artwork*

Please make sure that artwork files are in an acceptable format (TIFF (or JPEG), EPS (or PDF), or MS Office files) and with the correct resolution. If, together with your accepted article, you submit usable color figures then Elsevier will ensure, at no additional charge, that these figures will appear in color online (e.g., ScienceDirect and other sites) regardless of whether or not these illustrations are reproduced in color in the printed version. **For color reproduction in print, you will receive information regarding the costs from Elsevier after receipt of your accepted article.** Please indicate your preference for color: in print or online only. [Further information on the preparation of electronic artwork](#).

### *Figure captions*

Ensure that each illustration has a caption. A caption should comprise a brief title (**not** on the figure itself) and a description of the illustration. Keep text in the illustrations themselves to a minimum but explain all symbols and abbreviations used.

## Tables

Please submit tables as editable text and not as images. Tables can be placed either next to the

relevant text in the article, or on separate page(s) at the end. Number tables consecutively in accordance with their appearance in the text and place any table notes below the table body. Be sparing in the use of tables and ensure that the data presented in them do not duplicate results described elsewhere in the article. Please avoid using vertical rules and shading in table cells.

## References

### *Citation in text*

Please ensure that every reference cited in the text is also present in the reference list (and vice versa). Any references cited in the abstract must be given in full at the end of the abstract. Unpublished results and personal communications are not recommended in the reference list, but may be mentioned in the text. If these references are included in the reference list they should follow the standard reference style of the journal (see below) and should include a substitution of the publication date with either "Unpublished results" or "Personal communication" Citation of a reference as "in press" implies that the item has been accepted for publication.

### *Web references*

As a minimum, the full URL should be given and the date when the reference was last accessed. Any further information, if known (DOI, author names, dates, reference to a source publication, etc.), should also be given. Web references can be listed separately (e.g., after the reference list) under a different heading if desired, or can be included in the reference list.

### *Data references*

This journal encourages you to cite underlying or relevant datasets in your manuscript by citing them in your text and including a data reference in your Reference List. Data references should include the following elements: author name(s), dataset title, data repository, version (where available), year, and global persistent identifier. Add [dataset] immediately before the reference so we can properly identify it as a data reference. The [dataset] identifier will not appear in your published article.

### *References in special issue articles, commentaries and responses to commentaries*

Please ensure that the words 'this issue' are added to any references in the reference list (and any citations in the text) to other articles which are referred to in the same issue.

### *Reference management software*

Most Elsevier journals have their reference template available in many of the most popular reference management software products. These include all products that support [Citation Style Language styles](#), such as [Mendeley](#) and [Zotero](#), as well as [EndNote](#). Using the word processor plug-ins from these products, authors only need to select the appropriate journal template when preparing their article, after which citations and bibliographies will be automatically formatted in the journal's style. If no template is yet available for this journal, please follow the format of the sample references and citations as shown in this Guide.

The current *Social Science & Medicine* EndNote file can be directly accessed by clicking [here](#).

Users of Mendeley Desktop can easily install the reference style for this journal by clicking the following link:



<http://open.mendeley.com/use-citation-style/social-science-and-medicine>

When preparing your manuscript, you will then be able to select this style using the Mendeley plug-ins for Microsoft Word or LibreOffice.

### **Reference formatting**

There are no strict requirements on reference formatting at submission. References can be in any style or format as long as the style is consistent. Where applicable, author(s) name(s), journal title/book title, chapter title/article title, year of publication, volume number/book chapter and the pagination must be present. Use of DOI is highly encouraged. The reference style used by the journal will be applied to the accepted article by Elsevier at the proof stage. Note that missing data will be highlighted at proof stage for the author to correct. If you do wish to format the references yourself they should be arranged according to the following examples:

### **Reference style**

*Text:* All citations in the text should refer to:

1. *Single author:* the author's name (without initials, unless there is ambiguity) and the year of publication;
2. *Two authors:* both authors' names and the year of publication;
3. *Three or more authors:* first author's name followed by 'et al.' and the year of publication. Citations may be made directly (or parenthetically). Groups of references should be listed first alphabetically, then chronologically.

Examples: 'as demonstrated (Allan, 2000a, 2000b, 1999; Allan and Jones, 1999). Kramer et al. (2010) have recently shown ....'

*List:* References should be arranged first alphabetically and then further sorted chronologically if necessary. More than one reference from the same author(s) in the same year must be identified by the letters 'a', 'b', 'c', etc., placed after the year of publication.

*Examples:*

Reference to a journal publication:

Van der Geer, J., Hanraads, J.A.J., Lupton, R.A., 2010. The art of writing a scientific article. *J. Sci. Commun.* 163, 51–59.

Reference to a book:

Strunk Jr., W., White, E.B., 2000. *The Elements of Style*, fourth ed. Longman, New York.

Reference to a chapter in an edited book:

Mettam, G.R., Adams, L.B., 2009. How to prepare an electronic version of your article, in: Jones, B.S., Smith, R.Z. (Eds.), *Introduction to the Electronic Age*. E-Publishing Inc., New York, pp. 281–304.

Reference to a website:

Cancer Research UK, 1975. Cancer statistics reports for the UK.

<http://www.cancerresearchuk.org/aboutcancer/statistics/cancerstatsreport/> (accessed 13.03.03).

Reference to a dataset:

[dataset] Oguro, M., Imahiro, S., Saito, S., Nakashizuka, T., 2015. Mortality data for Japanese oak wilt disease and surrounding forest compositions. *Mendeley Data*, v1. <https://doi.org/10.17632/xwj98nb39r.1>.

### **Video data**

Elsevier accepts video material and animation sequences to support and enhance your scientific research. Authors who have video or animation files that they wish to submit with

their article may do so during online submission. Where relevant, authors are strongly encouraged to include a video still within the body of the article. This can be done in the same way as a figure or table by referring to the video or animation content and noting in the body text where it should be placed. These will be used instead of standard icons and will personalize the link to your video data. All submitted files should be properly labeled so that they directly relate to the video file's content. In order to ensure that your video or animation material is directly usable, please provide the files in one of our recommended file formats with a maximum size of 10 MB. Video and animation files supplied will be published online in the electronic version of your article in Elsevier Web products, including ScienceDirect: <http://www.sciencedirect.com>. For more detailed instructions please visit our video instruction pages at <http://www.elsevier.com/artworkinstructions>. Note: since video and animation cannot be embedded in the print version of the journal, please provide text for both the electronic and the print version for the portions of the article that refer to this content.

### **Supplementary data**

Elsevier accepts electronic supplementary material to support and enhance your research. Supplementary files offer the author additional possibilities to publish supporting applications, accompanying videos describing the research, more detailed tables, background datasets, sound clips and more. Supplementary files supplied will be published online alongside the electronic version of your article in Elsevier Web products, including ScienceDirect: <http://www.sciencedirect.com>. In order to ensure that your submitted material is directly usable, please provide the data in one of our recommended file formats. Authors should submit the material in electronic format together with the article and supply a concise and descriptive caption for each file. For more detailed instructions please visit our artwork instruction pages at <http://www.elsevier.com/artworkinstructions>.

### ***Data linking***

If you have made your research data available in a data repository, you can link your article directly to the dataset. Elsevier collaborates with a number of repositories to link articles on ScienceDirect with relevant repositories, giving readers access to underlying data that give them a better understanding of the research described.

There are different ways to link your datasets to your article. When available, you can directly link your dataset to your article by providing the relevant information in the submission system. For more information, visit the [database linking page](#).

For [supported data repositories](#) a repository banner will automatically appear next to your published article on ScienceDirect.

In addition, you can link to relevant data or entities through identifiers within the text of your manuscript, using the following format: Database: xxxx (e.g., TAIR: AT1G01020; CCDC: 734053; PDB: 1XFN).

### **AudioSlides**

The journal encourages authors to create an AudioSlides presentation with their published article. AudioSlides are brief, webinar-style presentations that are shown next to the online article on ScienceDirect. This gives authors the opportunity to summarize their research in their own words and to help readers understand what the paper is about. [More information](#)

[and examples are available](#). Authors of this journal will automatically receive an invitation e-mail to create an AudioSlides presentation after acceptance of their paper.

### Interactive plots

This journal enables you to show an Interactive Plot with your article by simply submitting a data file. [Full instructions](#).



**After Acceptance**

### Online proof correction

Corresponding authors will receive an e-mail with a link to our online proofing system, allowing annotation and correction of proofs online. The environment is similar to MS Word: in addition to editing text, you can also comment on figures/tables and answer questions from the Copy Editor. Web-based proofing provides a faster and less error-prone process by allowing you to directly type your corrections, eliminating the potential introduction of errors.

If preferred, you can still choose to annotate and upload your edits on the PDF version. All instructions for proofing will be given in the e-mail we send to authors, including alternative methods to the online version and PDF.

We will do everything possible to get your article published quickly and accurately. Please use this proof only for checking the typesetting, editing, completeness and correctness of the text, tables and figures. Significant changes to the article as accepted for publication will only be considered at this stage with permission from the Editor. It is important to ensure that all corrections are sent back to us in one communication. Please check carefully before replying, as inclusion of any subsequent corrections cannot be guaranteed. Proofreading is solely your responsibility.

### Offprints

The corresponding author will, at no cost, receive a customized [Share Link](#) providing 50 days free access to the final published version of the article on [ScienceDirect](#). The Share Link can be used for sharing the article via any communication channel, including email and social media. For an extra charge, paper offprints can be ordered via the offprint order form which is sent once the article is accepted for publication. Both corresponding and co-authors may order offprints at any time via Elsevier's [Webshop](#). Corresponding authors who have published their article open access do not receive a Share Link as their final published version of the article is available open access on ScienceDirect and can be shared through the article DOI link.

## Guidelines for Qualitative Papers

There is no one qualitative method, but rather a number of research approaches which fall under the umbrella of 'qualitative methods'. The various social science disciplines tend to have different conventions on best practice in qualitative research. However SS&M has prepared the following

general guidance for the writing and assessment of papers which present qualitative data (either alone or in combination with quantitative methods). General principles of good practice for all research will also apply.

## **Fitness for purpose**

Are the methods of the research appropriate to the nature of the question(s) being asked, i.e.

- Does the research seek to understand social processes or social structures &/or to illuminate subjective experiences or meanings?
- Are the settings, groups or individuals being examined of a type which cannot be pre-selected, or the possible outcomes not specified (or hypothesised) in advance?

## **Methodology and methods**

- All papers must include a dedicated methods section which specifies, as appropriate, the sample recruitment strategy, sample size, and analytical strategy.

## **Principles of selection**

Qualitative research is often based on or includes non-probability sampling. The unit(s) of research may include one or a combination of people, events, institutions, samples of natural behaviour, conversations, written and visual material, etc.

- The selection of these should be theoretically justified e.g. it should be made clear how respondents were selected
- There should be a rationale for the sources of the data (e.g respondents/participants, settings, documents)
- Consideration should be given to whether the sources of data (e.g people, organisations, documents) were unusual in some important way
- Any limitations of the data should be discussed (such as non response, refusal to take part)

## **The research process**

In most papers there should be consideration of

- The access process
- How data were collected and recorded
- Who collected the data
- When the data were collected
- How the research was explained to respondents/participants

## **Research ethics**

- Details of formal ethical approval (i.e. IRB, Research Ethics Committee) should be stated in the main body of the paper. If authors were not required to obtain ethical approval (as is the case in some countries) or unable to obtain ethical approval (as sometimes occurs in resource-poor settings) they should explain this. Please anonymise this information as appropriate in the manuscript, and give the information when asked during submission.
- Procedures for securing informed consent should be provided

Any ethical concerns that arose during the research should be discussed.

## **Analysis**

The process of analysis should be made as transparent as possible (notwithstanding the conceptual and theoretical creativity that typically characterises qualitative research). For example

- How was the analysis conducted
  - How were themes, concepts and categories generated from the data
  - Whether analysis was computer assisted (and, if so, how)
  - Who was involved in the analysis and in what manner
- Assurance of analytic rigour. For example
  - Steps taken to guard against selectivity in the use of data
  - Triangulation
  - Inter-rater reliability
  - Member and expert checking
  - The researcher's own position should clearly be stated. For example, have they examined their own role, possible bias, and influence on the research (reflexivity)?

## **Presentation of findings**

### **Consideration of context**

The research should be clearly contextualised. For example

- Relevant information about the settings and respondents/participants should be supplied
- The phenomena under study should be integrated into their social context (rather than being abstracted or de-contextualised)
- Any particular/unique influences should be identified and discussed

### **Presentation of data:**

- Quotations, field notes, and other data where appropriate should be identified in a way which enables the reader to judge the range of evidence being used
- Distinctions between the data and their interpretation should be clear
- The iteration between data and explanations of the data (theory generation) should be clear
- Sufficient original evidence should be presented to satisfy the reader of the relationship between the evidence and the conclusions (validity)

- There should be adequate consideration of cases or evidence which might refute the conclusions

**General Practitioners and the euthanasia and assisted suicide debate in the United Kingdom: A Foucauldian exploration of their discourses**

Emily L. Maddock (Corresponding author)

North Wales Clinical Psychology Programme

Bangor University

Bangor, LL57 2DG

Psp4f4@bangor.ac.uk

Carolien Lamers

North Wales Clinical Psychology Programme

Bangor University

Bangor, LL57 2DG

c.lamers@bangor.ac.uk

This paper is part of the Doctorate in Clinical Psychology

North Wales Clinical Psychology Programme

Bangor University

## Abstract

The debate in the United Kingdom (UK) regarding the legalisation of euthanasia (EU) and assisted suicide (AS) is ongoing. While EU/AS are not legal in the UK, doctors use palliative sedation and high doses of painkillers to manage the end of life stages. However, were EU/AS to become end of life choice options, UK General Practitioners (GPs) are likely to be involved in assessing and assisting in these choices. The purpose of this study was to gain an understanding of the discourses of UK GPs regarding EU/AS. Six white British GPs participated in semi-structured interviews. Their discourses were analysed using Foucauldian discourse analysis (FDA); a form of discourse analysis based on the philosophy of Michel Foucault, which is primarily concerned with revealing the knowledge and power relationships in discourse. While the GPs' factual knowledge regarding EU/AS was limited, they revealed a comfortable 'palliative' discourse, in which GPs hold a powerful 'expert' position but in which EU/AS did not feature. However, a less comfortable 'medicine is limited, is EU/AS the answer?' discourse emerged, where GPs were no longer the 'experts'. From the 'medicine is limited, is EU/AS the answer?' discourse, GPs moved to a further discourse where EU/AS could be a patient's right, however most GPs settled on a 'someone else should do it' discourse. In this discourse where EU/AS was legalised, the GPs returned to the comfortable 'palliative' discourse and 'expert' position, and considered EU/AS not necessarily as part of the role of a GP. Increased factual knowledge of EU/AS is necessary for UK GPs to engage in an informed debate, and would allow GPs to develop new, alternative discourses. Future research should take their discourses and gaps in knowledge into consideration.

*Keywords:* Euthanasia, Assisted dying, Foucault, Palliative care, Physician assisted dying



The way that death and dying are thought about, spoken about and subsequently responded to is specific to the cultural context at any given time (Strazzari, 2005). As medical knowledge and technology advanced throughout the latter half of the twentieth century in the United Kingdom (UK), dying shifted from an accepted, natural part of life (Callahan, 2009) to a medical challenge, with the aim to prolong life as long as possible (Clark, 2002). Dying increasingly took place in hospices, hospitals and other medical/care settings and people could be kept alive beyond the point at which death would naturally have occurred. These advances have contributed to an increase in end of life care options, such as palliative sedation and the use of high doses of pain killers to control physical symptoms. This is often referred to as the 'medicalization of dying' (Karsoho, Fishman, Wright, & Macdonald, 2016).

The medicalization of death and the prolonging of life beyond its natural end point has contributed to debates about the medical decision making process regarding the point at which it is decided that interventions are stopped, treatment is withdrawn or life supporting equipment is turned off. These decisions ultimately determine when death occurs (Karsoho, Fishman, Wright, & Macdonald, 2016). Several public and patient groups in the UK have been campaigning for the legalisation of euthanasia (EU) and assisted suicide (AS) so that the dying person can be given the option to determine the endpoint of their life rather than a doctor.

While EU and AS are not legal in the UK, the options available to doctors to manage the end of life stages are palliative sedation and high doses of painkillers. Palliative sedation is the intentional administration of sedative drugs to a terminally ill patient with the aim of reducing consciousness to a level that provides relief from physical symptoms and pain that

cannot otherwise be managed (Broeckaert & Nunez Olarte, 2002). The use of high doses of pain killers to control otherwise unmanageable physical symptoms may have as an unintended side effect the shortening of life, often referred to by medics as 'the double effect' (Marcoux, Mishara, & Durand, 2007). Both practices have been hotly debated in the UK, with critics claiming that they are forms of 'soft euthanasia' (Ten Have & Welie, 2014; Howland, 2005).

In contrast, EU is the deliberate ending of a competent person's life, at their request, for the purpose of relieving suffering (Deliens & Van der Wal, 2003). In assisted suicide, a competent patient ends their own life using means (e.g. lethal drugs) provided by a medical practitioner (Wilson et al., 2007). Legislation supporting EU and AS has been in place in the Netherlands and Belgium since 2002, in Luxembourg since 2009 and in Colombia since 2015. Several other countries and states of America also have legislation permitting either EU or AS, with Canada passing legislation legalising Physician assisted dying in June 2016.

In the Netherlands, GPs perform EU more frequently than any other medical specialists (Onwuteaka-Philipsen et al., 2003). Research into the views and experience of Dutch GPs has revealed both positive and negative impacts of being involved in the EU/AS process. A mixture of anxiety, loss and relief were reported by GPs in a study by Van Marwijk, Haverkate, Van Royen, and The (2007). Open patient-GP communication regarding EU has been highlighted as particularly important (Borgsteede et al., 2007). Opinions about who should initiate EU discussions differed between GPs and many reported finding it difficult to judge the 'right' time to discuss EU. Having a good patient-doctor relationship has also been shown to be important, with many GPs saying they are only willing to perform EU if they have a good relationship with the patient (Van Marwijk, Haverkate, Van Royen, & The,

2007). Norwood (2007) interviewed patients and GPs revealing that conversations about EU changed the patient-GP relationship. Open discussions resulted in a deepening and strengthening of the relationship.

If EU/AS were to be legalised in the UK, UK doctors, including GPs, would be directly affected by this ruling. Doctors are currently involved in decisions about the use of palliative sedation and pain management, which acknowledges the risk of 'the double effect'. However, the views of UK GPs regarding the legalisation of EU/AS are not well understood (Seale, 2009b). The available research suggests that the majority (32-63%) of GPs are opposed to the legalisation of EU/AS in the UK, but less so than geriatricians (80%; Market Research Services, 2004). A survey by Seale (2009b) that included 1073 GPs, also showed UK doctors (79.9%) to be opposed to EU/AS. Those who were in support of EU/AS stated a strong need for safeguards, that GPs should have the right to 'opt out' of involvement with EU/AS and that non-medical practitioners should be able to be involved. Of those who opposed EU/AS, 35% agreed that withholding life sustaining treatment in the interest of relieving suffering, or providing treatment that might also have a life shortening effect, was acceptable.

Given that there have been (and are likely to be future) attempts to pass EU/AS legislation in the UK, a better understanding of the views of GPs regarding EU/AS is essential, as the implementation could be hampered by a lack of willingness of GPs to be involved in carrying out EU/AS. To gain a deeper insight into the discourses of UK GPs regarding EU/AS, the same methodology to Lamers and Williams (2015) will be used, who explored the discourses of older people in the UK regarding EU/AS. Foucauldian discourse analysis (FDA) will be used to address two research questions:

1. What knowledge and power underlies general practitioners discourses in relation to the EU/AS debate?
2. What is the position of general practitioners in the EU/AS debate?

## **Method**

### **Methodology**

Discourse analysis (DA) is a social constructionist approach that is concerned with the relationship between language and social processes (Bryman, 2001). A discourse is the narrative of a phenomenon constructed through speech, writing and practice (Wetherell, Taylor & Yates, 2015). FDA is a form of DA based on the theories of the French philosopher Michel Foucault and is primarily concerned with power relationships in society as revealed through discourses. As the EU/AS debate centres around the questions as to who has the power to decide when and how to end a life, this methodology seemed suitable to answer the posed questions.

Foucault conceptualised discourses as constitutive, constructing what is considered to be 'real', 'correct' or the 'truth' at any given time in a particular culture (Gutting, 2005). This 'truth' is constructed through the process of normalization, when a particular discourse becomes dominant and considered 'correct'. Through this discourse, implicit and explicit 'rules of behaviour' (e.g. social norms and legislation) are created and this influences how individuals and institutions think and act. Existing discourses can be consolidated or indeed challenged. When changes in the knowledge of a particular phenomenon occur, this can produce new conceptualisations of a phenomenon and an alternative discourse gets created. With a change in knowledge as the catalyst for the development and evolution of discourses, knowledge is also fundamental to power relationships (Wetherell, Taylor &

Yates, 2004). Power relationships are developed when people who have different aspects of knowledge, are divided into social positions of 'experts', those who have knowledge about the 'objects', who in turn become the 'subjects' based on knowledge as it is expressed through discourse. Foucault theorised that the position of 'subject' is itself produced and defined by the discourse. The 'subject' is an individual who personifies the knowledge created by a particular discourse.

As discourses are ever evolving; what is considered to be 'true' or 'correct' evolves over time. Thus, FDA does not seek to reveal one single definitive 'truth' relating to a phenomenon, instead FDA seeks to reveal the power relationships created by discourses at any given time in a particular culture.

Medical, illness and end of life care discourses position medical practitioners, such as GPs, in the 'expert' position. They hold specialist knowledge regarding illnesses and treatments, and use associated medical and diagnostic language. Subsequently, the experience of an individual becomes the 'object' of the GP's knowledge and their discourse. Through these discourses, the position of 'patient' (subject position) is created and associated behavioural 'norms' are established. The patient (subject) conforms to the 'rules' defined by the medical discourses and behaves accordingly, a process Foucault referred to as 'subjectification' (Wetherell, Taylor & Yates, 2004). Institutions, such as hospitals, also hold authority and power, as through and in these institutions the 'experts' (medical practitioners) monitor the 'objects' (patients) treatment and progress. This experience of surveillance leads to the 'subject' (patients) engaging in self-surveillance (self-monitoring), a process that Foucault believed to be the most powerful mechanism of control.

In the UK, through the frequent legal challenges and media discussions, the dominant discourse that EU/AS is unacceptable and could lead to a 'slippery slope' is being challenged repeatedly. Discourses about possible inadequacies of current end of life care and a cultural shift to self-determination, contribute to an emerging discourse that EU/AS is needed and beneficial (Lamers & Williams, 2015). Nevertheless, the discourse to maintain EU/AS as an illegal act remains dominant, and is supported by a legal framework that sets out how GPs and other medical professionals are allowed to act when providing end of life care, and as such operates as a surveillance system for the doctors. These six Foucauldian concepts; knowledge, power, surveillance, expert, object and subject will be used to understand the GPs' discourses around EU/AS.

### **Research Ethics**

Ethical approval for this research was obtained from both the School of Psychology Ethics Committee at Bangor University and from the Betsi Cadwaladr University Health Board Research and Development West Department.

### **Sampling**

FDA is a social constructionist approach and is therefore concerned with gaining an in depth understanding of participants' construction of reality through language. As it does not aim to generalise results, random sampling is not required. Therefore, recruitment was opportunistic and via adverts (see Appendix E) distributed through GP meetings to GPs currently employed by Betsi Cadwaladr University Health Board and those who have retired. The information available to prospective participants included the advert, an information sheet (see Appendix F) detailing the purpose of the research, the process, data protection

procedures and researcher information and contact details. Initial contact with prospective participants was via e-mail or telephone, and consent to participate was discussed.

## Participants

Participants were six white British GPs aged between 35-73 years, three were currently practising and three had retired. Length of time in practice ranged between 3-41 years.

Demographic details were recorded (see Table 1).

Table 1  
*Participant Demographic Details*

Participant	Gender	Age	Professional status	Years in practice	Length of retirement
Grace	Female	60	Retired	35	1 month
John	Male	73	Retired	41	8 years
Peter	Male	67	Retired	32	8 years
Sarah	Female	39	Practicing	15	-
William	Male	58	Practicing	30	-
Jane	Female	35	Practicing	3	-

*Note.* Participant names are pseudonyms.

## Interviews

Interviews were conducted by the first author at Bangor University or at the GPs' home. The interview process began with confirmation that the GP had read and understood the information sheet. The information sheet addressed the action that would be undertaken were a GP to describe a clinical practice that might cause concern. If this were to happen, the supervisory team would be informed and appropriate action taken. The opportunity was given for the GPs to ask questions or raise concerns. The informed consent form (see

Appendix G) was completed at the beginning of the interview. The interview was recorded using a Dictaphone.

The interviews were semi-structured and followed an interview schedule (see Appendix H) that was based on that used by Lamers and Williams (2015). The interview schedule was altered to incorporate questions that would distinguish between the GPs' professional and personal views. The interview began with externalising questions with the aim of establishing GPs' knowledge and understanding of EU/AS, e.g. "Can you tell me in your own words what euthanasia means?" This was followed by personalising questions with the aim of accessing the language used by GPs to describe their views and personal meaning of EU/AS, and finally, specifying questions regarding the circumstances under which they would consider EU/AS. GPs were given the opportunity to discuss or add comments they felt were relevant or important that had not been addressed in the interview. Explorative prompts were used throughout the interviews to facilitate naturally occurring speech as much as possible. Interviews lasted 30-45 minutes and ended with a debrief.

### **Analysis**

As there is no single agreed method for conducting FDA, in keeping with Lamers and Williams (2015), the analysis followed the procedure outlined by Georgaca and Avdi (2011).

The analysis consisted of the following stages:

1. The interviews were transcribed by the first author (see Appendix I for example transcript). Transcription included vocal tones and pauses (bold = said loudly/with emphasis; underlined = said quietly/softly; (.) = pause for 1-2 seconds; (..) = pause for 3+ seconds).



2. Transcripts were read and reread to identify the main discourses relating to each of the two research questions. This included identifying and examining implied or explicit expressions of the GPs' views on EU/AS, and how they positioned themselves in the debate, keeping in mind the Foucauldian concepts of knowledge, power, surveillance, expert, object and subject. In addition to the main discourses, other interrelated discourses were identified and examined.
3. The dynamics of the interaction were examined, focusing on the GPs' discursive agenda and interaction with the interviewer (e.g. seeking agreement).
4. The GPs' position (expert, object or subject) within each discourse was established.
5. The way in which each discourse maintained or challenged current practices and power structures was identified.
6. The impact of the discourses on the GPs behaviour was explored.

## **Results**

### **First Author's Declaration**

The first author has no previous research, clinical or personal experiences with EU/AS and does not hold strong views either way related to EU/AS. The first author's interest in EU/AS came from the increase in media coverage of the EU/AS debate in the UK.

### **Interpretations**

As FDA seeks to reveal the power relationships associated with discourses, rather than seeking to reveal one single definitive truth, the results are presented as interpretations. It should be acknowledged that both the author's and the reader's own discourses influence

the interpretation of the discourses revealed by the GPs. All identifying details have been anonymised and GPs' quotes are presented under pseudonyms.

Throughout the interviews, GPs' discourses regarding EU/AS drew on three types of knowledge; their knowledge about what EU/AS is and the law in the UK, professional knowledge about illness and treatment, and personal knowledge about societal norms, expectations and their own values. As Foucault theorised, this knowledge and the associated discourses have developed and evolved over a number of years.

*Sarah: "I think God my views have changed so much from perhaps 20 years ago, I think perhaps 20 years ago I was far more kind of **yes**, absolutely, everyone should be given the right to do it and I feel so much more uncomfortable about it now".*

GPs' factual knowledge regarding EU/AS was confused. Although they described EU as the intentional ending of a patient's life with the aim of relieving suffering, there was confusion around who made this decision, the patient or the medical practitioner. No GP referred explicitly to the fact that the patient had to be competent. Most GPs described EU as being primarily a doctor initiated decision, and several of the GPs engaged in checking or seeking agreement from the interviewer: *"We talk about euthanizing animals, don't we?"* (Jane).

There was also some confusion regarding the difference between EU and AS *"Erm (..) I'm not sure whether I've thought of there being a difference (..) erm (..)"* (Sarah). Most GPs described AS as helping someone to die by providing the means, e.g. a lethal drug that the person takes themselves. In contrast with EU, all GPs described AS as being initiated by, and a decision made by, the patient not the medical practitioner. However, GPs were unable to accurately differentiate between EU and AS, being confused about who makes the decision

and who carries out the procedure. With this confusion around the nature of the acts, their discourses were based on inaccurate knowledge.

*Grace: "I suppose they are slightly different in that assisted suicide suggests that the person is making the plans and simply requires help with the actual process of **doing**, whereas euthanasia would suggest that others have more of a say in the initiation and carrying out of the act. That for me is the difference".*

### **Palliative care**

The GPs revealed a dominant 'palliative' discourse that was underpinned by the GPs' professional knowledge of illness, illness trajectory, treatment options and symptom management. They talked about managing end of life situations effectively with medication in such a way that patients were able to have 'a good death' and EU/AS was not viewed as necessary or part of the GPs' treatment repertoire.

*Sarah: "At the moment with things like syringe drivers we write up the drugs but we don't actually administer (.) it's district nurses generally that do it, erm so yes you might alter doses and say yes they need more at the moment, they're more distressed or restless or this needs to be added to the driver".*

Here, GPs were in the 'expert' position and held the power to make medical decisions that might have the side effect of hastening somebody's death: "*what you do all the time as a doctor is make judgments about things that aren't black and white*" (Jane). It is worth noting that it is not usually the GP who implements their own decision, it is often the district nurses. The patient's experience was the 'object' of the 'palliative' discourse: "*I don't think there has been one where we've just left them without, where it hasn't been sorted out*"

(Sarah). Within this discourse, the GP 'sorted out' the 'object' (i.e. 'one') by using their expert knowledge and medical decisions, thus creating a 'subject' that gets 'sorted out'.

The 'palliative' discourse was comfortable and safe for the GPs, who spoke confidently and fluently. Within this discourse the GP makes decisions for the patient, thus taking the expert position and creating the 'subject' who complies accordingly.

*Jane: "When it's palliative care and its people who are dying anyway, then there is a massive overlap between (.) treatments that are going to kill someone and treatments that are going to help someone. So I feel much more comfortable discussing in those situations 'well this is what we are doing with your medication'".*

*Sarah: "Erm, and I think we are all well aware that by increasing pain medication (.) that potentially can shorten life because of the side effects but it's a very, they are in those last few days anyway so by increasing, actually it's more important that their pain is controlled".*

The 'palliative' discourse also has a 'covert' strand, where the GPs acknowledge the risk of the 'double effect' and where medical decisions might begin to move closer to the acts of EU/AS. GPs quote medical colleagues who are seen as more knowledgeable and therefore more powerful, implying an internal level of surveillance.

*Jane: "A couple of senior doctors I've worked with have said things like (.) they talk about a palliative patient and you say at least they aren't in any pain and you sort of see them go but mmm but if they were at least we could then (.) offer some morphine that might then cut things down a bit. Because actually it's dragging on for*

*them and it's dragging on for the family. But actually if someone isn't in any pain (..)".*

While palliative care creates a safe discourse, with the GPs using their expert knowledge with the aim of alleviating pain and suffering, they allude to an awareness that at times their intervention might come close to the 'double effect' and the challenge of surveillance emerges.

### **Medicine is limited, is EU/AS the answer?**

However, GPs began to acknowledge that for a minority of patients a 'good death' was not always possible; palliative treatment and pain management options had been exhausted or were ineffective at relieving symptoms. In describing the limitations of medicine, their own knowledge and skills, a much less comfortable discourse emerged.

*Jane: "Erm, we love to think we've got loads of options in medicine and actually sometimes you just don't. And actually for some people dying is going to be bad. So I think it's naive isn't the right word but I think to say oh you know, modern medicine can make it better, I don't think that is true".*

GPs were themselves beginning to challenge the previous 'palliative' discourse. They described these situations as *"difficult because they [GPs] are used to being in control of a situation like that"* (Peter), and themselves as feeling *"helpless"* (William). These words are inferring a shift from being and feeling powerful to becoming powerless. In these situations, the knowledge and power had shifted away from the GP, with the patient experiencing the limitations of the medical knowledge. When the patient and GP were confronted with the 'medicine is limited' discourse, EU/AS was considered to be an alternative. William was

aware that he might have different views about EU/AS from his colleagues, and he senses some level of surveillance.

*William: "I've got fairly (.) well I think positive views but other people would disagree. Strong views on euthanasia from my experience of seeing people suffering and I don't think that's because (..) I don't think that's because we don't, haven't made reasonable attempts to relieve suffering I think some things are just (..) beyond science and society to help individuals and so I think euthanasia and assisted suicide should be an option".*

When confronted with the discourse 'Medicine is limited, is EU/AS the answer?', GPs described engaging in two behaviours that were interpreted as attempts to avoid the vulnerable position that this discourse would put them in, as this could potentially lead to 'risky' EU/AS conversations. Jane commented that *"the words [EU/AS] (..) make everyone very edgy"*. The first behaviour was to attempt to shut down any conversations the patients may initiate regarding EU/AS.

*Jane: "I think for me a conversation like that would always start off with 'wow ok, this is medically a really difficult area because this is not something that I can help you with'".*

The second behaviour was to call in the experts, as the GPs questioned their own knowledge and power.

*Peter: "You could call an expert in when things were getting out of (.) getting slightly out of control, you weren't sure what to do. And we had a good one round here and I'd just ring him up and say look er, I've got this problem and der der der*

*der and you don't have to come and see it, but erm, the patient, but can you tell me what to do, or very often I'd ring him up and say look, erm I'm getting out of control with this and I need you to actually come and talk to these people and say look these are the options, I don't think they are believing me because I'm only a general practitioner so to speak. And we just need a little more authority or something and that would work in that way, er, erm and things would settle down".*

In this strand of the 'medicine is limited' discourse, Peter moved out of the 'expert' position, and into a subject position. Peter's comment "*I'm getting out of control with this*" reflects a level of self-surveillance regarding his expertise and ensuring that his own behaviour was in line with the law. By inviting an 'expert', Peter also invited in external surveillance. By gaining the perspective of an authoritative expert, he was able to regain control and re-establish his position in a 'palliative' discourse.

Despite attempts to remain in the 'palliative' discourse, the 'medicine is limited' discourse frequently re-emerged as GPs knew that there would be situations when a good death was not possible. The discourse was underpinned by the GPs' knowledge of the limitations of treatments and medicines as well as their experience of end of life care not always being able to meet the needs of patients. Based on this professional knowledge, the 'medicine is limited' discourse was unavoidable and the possibility of EU/AS as an alternative discourse emerged.

### **Patients have the right to EU/AS, but do not count on me**

In this "*vulnerable position*" (Grace) where the dominant 'palliative' discourse was challenged by the 'medicine is limited, is EU/AS the answer?' discourse, the possibility of a 'patients should have the right to EU/AS' discourse was considered.

Grace: *“At the moment it’s kind of by the back door by let’s put in DNAR [do not attempt resuscitation] orders on people, well what are we really saying in that situation? You know, in the event of something happening we are not actually going to give you active care, well that’s one step along the road to actually having that conversation with relatives which would be (.) I really do think we are reaching the point where quality of life is really dreadful here and we now say to them I think it would be counterproductive in the event of something happening to try to resuscitate this person, if their heart stopped or whatever. Maybe we would need to be more brave and say, you know do you feel that we should be looking at the option of ending this person’s life. I don’t think it’s a massive step because if we really looked inside what we are doing we are probably not that far away now”.*

In Grace’s description, the confusion in the knowledge about the exact nature of EU/AS is noted, where she proposes to engage in conversations with family and friends about the patient’s quality of life and the futility of treatment, rather than the patient themselves. Grace alludes again to the ‘covert’ strand of the ‘palliative’ discourse, and makes reference to the ‘double effect’.

Peter: *“We just said helping people to die and that sort of thing. And we’d talk about it, but in the time I was there, there was no doctor who would say yeah I do it or I would do it because it’s actually illegal. Maximum 14 years in prison?! It’s something like that, it’s something fairly heavy. So, we would touch on it and this sort of thing and we would talk about erm, again, this er, the ethics of the double effect, so in fact you’re actually talking about euthanasia without actually mentioning it maybe or calling it by the name”.*



Here, Peter also refers to a 'covert' strand known amongst doctors, reflecting an awareness of the power they exert over medical decisions, while also exhibiting a shared self-surveillance, and an awareness of the surveillance of the legal system. The GP moved between the position of 'expert' and being powerful, to the 'subject' under surveillance. Sarah expressed her concerns about the risks related to naming an act in terms of EU/AS.

*Sarah: "Erm, and I think we are all well aware that by increasing pain medication (.) that potentially can shorten life because of the side effects but it's a very, they are in those last few days anyway so by increasing, actually it's more important that their pain is controlled. But to actually say, right, to purposefully, yes, we are actually going to make sure that is that. That is obviously life ending".*

The discourse that 'patients should have the right to EU/AS' was generally described in positive and collaborative language. While the patient was seen in the 'expert' role with knowledge about their own life, there was also a tension noted with the patient in a position of power and control, leading to a reversal of positions held under a 'palliative' discourse.

*Grace: "We give choice in medicine on so many issues, we involve people in choice more. And therefore my view now is that people probably have the right to choose for themselves whether or not their life should be ended".*

*Jane: "I think by the time you are having conversations about it [EU/AS], they [patients] know what position they are putting you in".*

However, when GPs attempted to define when EU/AS could be seen as a potential option, the 'patients should have the right to EU/AS' discourse seemed to be challenged. All GPs

spoke about EU/AS as only being acceptable “*if somebody just had a few months to go and was suffering a lot*” (John) and for “*a clear physical condition not a mental health problem, clear physical condition (.) that is (.) unbearable for the individual (.) that is very resistant to any sort of palliative treatment*” (William). Their attempt to define the criteria for when EU/AS could be considered, brings the GPs’ knowledge about health and illness back as a central player in the decision making, thus reinstating their position of power. They challenge the ‘subject’s’ knowledge and expertise to judge when their own life would no longer be bearable.

Sarah: “*That’s what sits really uncomfortably, the what if, because I don’t want to be at a point where I’m going to be incontinent or I’m gonna be this or this and therefore on this day I want my life to end. Erm, well what if it never happens or what if when you get there it’s not how you think it might be, or it might be*”.

Sarah starts to introduce the word ‘I’, beginning to bring her personal ideas and experiences to the EU/AS debate. This third type of personal knowledge regarding one’s own values and experiences, informed this discourse further, as well as societal discourses about when death is acceptable.

Grace: “*As society becomes less comfortable with the concept of death and wants to keep people as long as possible and we are not supposed to be age discriminatory, the care that we’re now offering people is probably worse than it was many years ago when we were rather more pragmatic about not treating chest infections or other things*”.

Possibly, Grace was referring to a discourse around an aged death (Lamers & Williams, 2015), which might be valued differently to a death in younger people or from non-age

related conditions. However, when exploring the consequences of a 'patients should have the right to EU/AS' discourse, GPs found themselves not only no longer positioned as the 'expert', their actions related to EU/AS, would also become the 'subject' of strict legal surveillance as well as scrutiny by their colleagues, patients and the public. GPs expressed concerns about being known as "*the death doctor*" (Jane) and no longer being trusted by their communities; "*You know, it's not great to have a reputation as the one who will bump you off if you get a bit too problematic*" (Grace).

In exploring the implications of the 'patients should have the right to EU/AS' discourse, GPs developed a 'someone else should do it' strand as a response. They excluded themselves from an active involvement with EU/AS, if this end of life choice were to be legalised in the UK. They felt that EU/AS required a legislative framework which should offer GPs the option to 'opt out', with the proviso that they would refer on to a colleague who was prepared to undertake EU/AS. They drew on a familiar pregnancy termination discourse, where a similar practice exists. It is worth noting that in the 'palliative' discourse, perhaps GPs had also 'opted out' when they described the practice of prescribing an increase in medication, but other professionals administering it.

*Grace: "I don't think the Swiss model is a bad one that it is not doctors administering the (.) treatment. I actually think there is an important distinction between doctors supporting euthanasia and assisted suicide and being the agents there of. Erm, I suppose it could work if you had the equivalent that we have in termination clinics. Where you have doctors who simply work there (..) or nurses who work there. Erm, equally, (..) in terminations you know, doctors have the right to opt out. Erm, and I'm not sure that that should necessarily be doctors or whether it should be practitioners*

*who are simply there to administer the means. I'm not even sure that I would want to be the person writing the prescription".*

Although all GPs stated that patients should have the right to access EU/AS, only William anticipated being willing to have an active role in EU/AS, should it be legalised. This strand of the 'patients should have the right to EU/AS' discourse provided a comfortable position for GPs regarding EU/AS, resolving the vulnerabilities and issues evident in the 'medicine is limited' discourse, but that GPs 'shouldn't be the ones to do it'. The GPs returned to the 'palliative' discourse, as the most viable alternative, and one which also allowed them to remain in the 'expert' position.

### **Discussion**

The GPs engaged in multiple and conflicting discourses before settling on a 'Patients should have the right to EU/AS, but someone else should do it' discourse. This discourse positioned the GPs as supporting the legalisation of EU/AS in the UK, provided that they did not have an active role in the EU/AS process. However, the question remains whether their lack of accurate knowledge and understanding of EU/AS has disadvantaged them in engaging with alternative discourses.

It is possible that GPs were referring to other concepts associated with EU/AS, which not only confuses discourse, but might have made the discrepancy between palliative care and EU/AS more nebulous; the 'covert' strand. When the law talks about EU/AS carried out in a competent person, this is also referred to as 'voluntary'. If the decision is made by someone other than the patient and the patient is incapable of making the decision themselves, this is referred to as 'non-voluntary' (Strazzari, 2005). The way in which EU/AS is achieved can be classified as either 'active', e.g. the administration of a lethal drug, or

'passive', the withholding of life sustaining treatment (Strazzari, 2005). The latter might resonate with the palliative care practice of withdrawing treatments or interventions.

A more accurate understanding of EU/AS, the distinctions between other end of life practices and the 'double effect', is necessary in order for UK GPs to engage in an informed debate. The impact that this lack of knowledge regarding EU/AS had on their discourses and ability to engage in an informed debate, highlights the need for improved training and education on end of life options, EU/AS and the associated laws.

In the 'palliative' discourse, GPs held medical and illness related knowledge which positioned them in the 'expert' position, from which they made legal end of life care decisions, such as when to increase painkilling medication. Patients were able to have 'good deaths', as unpleasant symptoms and pain were successfully controlled. The knowledge, 'expert' position and power of GPs expressed in this discourse, reflects the medicalization of dying.

When confronted with the 'medicine is limited' discourse, one of the behaviours that the GPs described engaging in was to attempt to shut down any conversations the patients may initiate regarding EU/AS, thus limiting the patient's access to the discourse. The positioning of the GPs in this debate mirrors the findings in Lamers and Williams' (2015) study, where older people stated that the medical profession seemed not willing to engage with them in the EU/AS debate.

Although the GPs' reluctance to engage in discussions regarding EU/AS is understandable given that EU/AS remains illegal in the UK, this behaviour is problematic in that it may negatively impact the patient-GP relationship and inhibit the patient from expressing concerns or difficulties regarding their end of life care. The importance of

communication in healthcare, and in palliative care specifically, has long been recognised (Royal College of Nursing, 2014). Open patient-doctor discussions regarding a number of end-of-life topics has been shown to result in greater concordance between patient end of life care goals and those documented by medical practitioners, as well as increased patient satisfaction with care (You et al., 2014).

Decisions about end of life care can be influenced by patient-doctor end of life care discussions (Mack et al., 2012). In countries where EU/AS is legal, open patient-GP communication regarding EU has been highlighted as particularly important (Borgsteede et al., 2007), even when the patient does not ultimately choose an EU/AS death. Norwood (2009) interviewed patients and GPs in the Netherlands (where EU is legal) and reported that conversations about EU had a positive impact on the patient-GP relationship. Open discussions resulted in a deepening and strengthening of the relationship. The GPs' concerns about actively engaging in EU/AS might be a reflection of their lack of knowledge regarding EU/AS and are potentially amplified by their reluctance to engage in discussions with their patients and colleagues regarding EU/AS.

However, with patient-GP communication essential and impacting not only on end of life decisions but also the patient-GP relationship, it will be beneficial for GPs to begin to engage in end of life care discussions with their patients, regardless whether they have raised EU/AS. This communication will enhance the GPs' understanding of the patients' concerns and reasons for raising EU/AS, help to protect the patient-GP relationship whilst also being able to address the patient concerns or difficulties.

It is possible that the six GPs have felt under 'surveillance' by the interviewer, who had indicated at the start of the interview that any concerns regarding the GPs' practice would

be reported to the relevant authorities and regulatory bodies. This surveillance, along with the GPs' knowledge of legal prohibitions, could have led to the 'covert' strand noted in the 'palliative' and 'medicine is limited' discourses. Therefore, it should be acknowledged that the context may have influenced the GPs' discourses. Also, GPs whose practices might be harder to distinguish from EU/AS would have been unlikely to have volunteered to take part in the study. As discourses are ever evolving, the results can only be considered within the socio-political context at the time of the research. Furthermore, the interpretation of the GPs' discourses by the authors will have been influenced by the authors' own knowledge, views and discourses.

The discourses may have been influenced by demographic factors. All of the GPs were white British, practising in the UK and they will have reflected western discourses regarding illness, care of the dying, EU/AS and the role of a GP. The GPs varied in age and length of time in practice and will have experienced different training and practice approaches that have changed and developed over time. This study does not address the possibility that younger, newly qualified GPs may hold different discourses from older, more experienced or retired GPs. Indeed, some of the GPs discussed that length of time in practice was a factor that had shaped their views regarding EU/AS.

The methodology itself brings limitations, as FDA assumes that reality is discursively constructed and contingent, it can minimise the non-semantic aspects of reality, particularly the material structuring of economic and political reality. FDA does not take into consideration the impact of such non-semantic factors on the construction of the 'object', which is conceptualised as being entirely discourse constructed (Reed, 2000). FDA has also

been criticised for minimising the significance of human agency in the construction, evolution and function of discourse.

This study has revealed that the GPs' discourses regarding EU/AS are complex, at times conflicting and based on three different types of knowledge. Future research that aims to understand GPs' views on EU/AS may need to take these complexities and deficits in knowledge into account.



## References

- Borgsteede, S. D., Deliens, L., Graafland-Riedstra, C., Francke, A. L., Van der Wal, G., & Willems, D. L. (2007). Communication about euthanasia in general practice: Opinions and experiences of patients and their general practitioners. *Patient Education and Counselling, 66*, 156-161. doi:10.1016/j.pec.2006.10.006
- Broeckeaert, B., & Nunez Olarte, J. M. (2002). Sedation in palliative care: facts and concepts. In H. Ten Have, & D. Clark (Eds.), *The ethics of palliative care: European perspectives* (pp. 166-180). Buckingham: Open University Press.
- Bryman, A. (2001). *Social Research Methods*. Oxford: Oxford University Press.
- Callahan, D. (2009). Death, mourning, and medical progress. *Perspectives in Biology and Medicine, 52*(1), 103-15. doi:10.1353/pbm.0.0067
- Clark, D. (2002). Between hope and acceptance: The medicalization of dying. *The British Medical Journal, 324*, 905-907. doi:doi.org/10.1136/bmj.324.7342.905
- Deliens, L., & Van der Wal, G. (2003). The euthanasia law in Belgium and the Netherlands. *The Lancet, 362*, 1239-40. doi:10.1016/S0140-6736(03)14520-5
- Georgaca, E., & Avdi, E. (2011). Discourse analysis. In D. Harper & A. R. Thompson (Eds), *Qualitative research methods in mental health and psychotherapy* (pp. 147-161). Oxford, England: Wiley-Blackwell. doi:10.1002/9781119973249.ch11
- Gutting, G. (2005). *FOUCAULT. A Very Short Introduction*. Oxford: Oxford University Press.

- Hagelin, J., Nilstun, T., Hau, J., & Carlsson, H-E. (2004). Surveys on attitudes towards legalisation of euthanasia: Importance of question phrasing. *Journal of Medical Ethics, 30*(6), 521-23. doi:10.1136/jme.2002.002543
- Howland, J. (2005). Questions about palliative sedation: An act of mercy or mercy killing? *Ethics and Medics, 30*(8), 1-2.
- Karsoho, H., Fishman, J. R., Wright, D. K., & Macdonald, M. E. (2016). Suffering and medicalization at the end of life: The case of physician-assisted dying. *Social Science & Medicine, 170*, 188-196. doi:10.1016/j.socscimed.2016.10.010
- Lamers, C. P. T., & Williams, R. R. (2015). Older people's discourses about euthanasia and assisted suicide: A Foucauldian exploration. *The Gerontologist, 56*(6), 1072-1081. doi:10.1093/geront/gnv102
- Mack, J. W., Cronin, A., Keating, N, L., Taback, N., Huskamp, H. A., Malin, J. L., ... Weeks, J. C. (2012). Associations between end-of-life discussion characteristics and care received near death: A prospective cohort study. *Journal of Clinical Oncology, 30*(35), 4387–95. doi:10.1200/JCO.2012.43.6055.
- Marcoux, I., Mishara, B. L., & Durand, C. (2007). Confusion between euthanasia and other end-of-life decisions: Influences on public opinion poll results. *Canadian Journal of Public Health, 98*(3), 235-239. doi:10.17269/cjph.98.820
- Market Research Services. (2004). *Public opinion on the legalisation of euthanasia/PAS. Appendix 7*. In: House of Lords. Assisted Dying for the Terminally Ill Bill – First Report. London: House of Lords. Retrieved from <https://www.publications.parliament.uk/pa/ld200405/ldselect/ldasdy/86/8617.htm>

Norwood, F. (2007). Nothing more to do: Euthanasia, general practice, and end-of-life discourse in the Netherlands. *Medical Anthropology*, 26, 139-174.

doi:10.1080/014597407012833165

Onwuteaka-Philipsen, B. D., Van der Heide, A., Koper, D., Keij-Deerenberg, I., Rietjens, J. A., Rurup, M. L., ... Van der Maas, P. J. (2003). Euthanasia and other end-of-life decisions in the Netherlands in 1990, 1995, and 2001. *The Lancet*, 362, 395–9.

doi:10.1016/S0140-6736(03)14029-9

Reed, M. (2000). The limits of discourse analysis in organizational analysis. *SAGE Journals*, 7(3), 524-530. doi:10.1177/135050840073011

Royal College of Nursing. (2014). *Defining nursing*. Retrieved from

<https://www.rcn.org.uk/professional-development/publications/pub-004768>

Seale, C. (2006). National survey of end-of-life decisions made by UK medical practitioners.

*Palliative Medicine*, 20, 3-10. doi:10.1191/0269216306pm1094oa

Seale, C. (2009a). End-of-life decisions in the UK involving medical practitioners. *Palliative*

*Medicine*, 23, 198-204. doi:10.1177/0269216308102042

Seale, C. (2009b). Legalisation of euthanasia or physician-assisted suicide: survey of doctors'

attitudes. *Palliative Medicine*, 23, 205-212. doi:10.1177/0269216308102041

Strazzari, M. (2005). Ageing, Dying, and Death in the Twenty-first Century. In J. Germov

(Eds.), *Second opinion: an introduction to health sociology* (pp. 244-264). Retrieved

from [http://lib.oup.com.au/he/health/germov5e/ch13\\_3.pdf](http://lib.oup.com.au/he/health/germov5e/ch13_3.pdf)

Ten Have, H. T., & Welie, J. V. M. (2014). Palliative sedation versus euthanasia: An ethical assessment. *Journal of Pain and Symptom Management*, 47(1), 123-136.

Doi:10.1016/j.jpainsymman.2013.03.008

Van Marwijk, H., Haverkate, L., Van Royen, P., & The, A. (2007). Impact of euthanasia on primary care physicians in the Netherlands. *Palliative Medicine*, 21, 609-614.

doi:10.1177/0269216307082475

Wetherell, M., Taylor, S., & Yates, S. J. (2004). *Discourse theory and practice: A reader*.

London: SAGE Publications.

Wetherell, M., Taylor, S., & Yates, S. J. (2015). *Discourse as Data: A guide for analysis*. Milton

Keynes: The Open University.

Wilson, K. G., Chochinov, H. M., McPherson, C. J., Skirko, M. G., Allard, P., Chary, S., ... Clinch, J. J. (2007). Desire for euthanasia or physician assisted suicide in palliative cancer care. *Health Psychology*, 26(3), 314-323.

You, J. J., Dodek, P., Lamontagne, F., Downar, J., Sinuff, T., Jiang, X., ... Heyland, K. (2014).

What really matters in end-of-life discussions? Perspectives of patients in hospital with serious illness and their families. *Canadian Medical Association Journal*,

186(18), 679–87. doi:10.1503/cmaj.140673

**Contributions to personal development, clinical practice and theory**

Emily L. Maddock

Bangor University, Bangor, LL57 2DG

Psp4f4@bangor.ac.uk

This final paper focuses on three areas. Firstly, my personal reflections on the project are discussed, followed by the clinical implications of the literature review and empirical paper, and finally the contributions of both papers to theory development and future research.

### **Personal reflections**

#### ***Choosing topics***

My interest in euthanasia (EU) and assisted suicide (AS) has developed over a number of years with the growing debate and media coverage in the United Kingdom (UK). Lamers and Williams (2015) used Foucauldian Discourse Analysis (FDA) to explore the discourses of older adults in the UK regarding EU/AS. This study provided the starting point for the development of the ideas for both the literature review and the empirical paper. Given that older adults' discourses revealed General Practitioners (GPs) as being in a position of power when it comes to making end of life decisions (Lamers & Williams, 2015), exploring the discourses of UK GPs themselves regarding EU/AS was the next logical step. This previous study, along with the fact that the views of UK GPs regarding EU/AS are not well understood (Seale, 2009), led to the development of the research questions addressed in the empirical paper.

My initial ideas for the literature review developed from considering Foucauldian power concepts in relation to my own clinical practice. One particular experience that had evoked strong feelings for me stood out. I had been left feeling very frustrated when I attempted to utilise the gender reassignment referral pathway for a service user, but was met with resistance from another professional who, despite clear guidelines, felt it was not part of their role to make the referral and engaged in obstructive behaviours such as not responding to my correspondence. I was struck by the amount of power that this

professional held and how it left me feeling frustrated, although not as frustrated as the service user. Given that in the discourses of older adults, medical professionals were seen to be in the 'expert' position and holding the power in medical and end of life decisions (Lamers & Williams, 2015), I began to think about the position of transgender older adults and how, as a minority group, they related to powerful medical professionals.

This thinking led me to explore the literature on transgender older adults, however, it quickly became apparent that there was very little research available and none that addressed their discourses or relationships with professionals (perhaps reflecting a lack of power). From this, the idea developed to focus on what factors are implicated in the mental health of transgender older adults.

Having now completed the research project, I have considered how the findings might influence my perception of the experience that initially led me to consider the mental health of transgender older adults. I feel that there are several similarities between the power relationships revealed in the empirical study and those that were at play in my experience.

GPs' discourses around EU/AS revealed their position as the powerful 'experts' and the patient as the 'subject'. This indeed also seemed to be the case in the experience that I had, the service user and I were both in the 'subject' position, lacking power over the referral process. Similar to the GPs who shut down uncomfortable EU/AS discussions, the professional concerned in my experience also engaged in shutting down communication, perhaps indicating a lack of knowledge regarding transgender related issues and the referral process. Interestingly, the professional also engaged in a 'someone else should do it' discourse and did not see the gender clinic referral as part of their role.

For me, these similarities highlight how a possible lack of knowledge that many professionals have around transgender related issues, can lead to certain discourses and related behaviours, which can impact on the service user and their mental health. Indeed, this lack of knowledge amongst professionals was apparent in the discussions I had with Sophie<sup>1</sup>, a 70 year old woman who identifies as transgender. She agreed to meet with me to discuss my literature review. Although these comments are my own reflections, and no research has been done that investigates the discourses of transgender older adults or medical professionals regarding transgender related issues, I do feel that these reflections have been beneficial in raising my awareness of potential power relationships, related discourses and the way this impacts on my behaviour and role, as well as the impact of these discourses on service users.

### ***Doing qualitative research***

Having previously only used quantitative research methods, I made the decision to undertake a qualitative research project, based in part on the idea that the experience would be more beneficial to my professional development. Although I feel that this has indeed been the case, the process of understanding and using an unfamiliar methodology such as FDA, whilst juggling the pressures of being a Trainee, has been challenging.

I began by reading studies that had used FDA and researching Foucauldian philosophies in a hope to 'get my head around' FDA reasonably quickly. However, the more I researched the more it became apparent that Foucault's writings are extensive, complex, and at times contradictory. My understanding of FDA seemed to consolidate through the process of

---

<sup>1</sup>Sophie is a pseudonym



actually applying it (and realising that there is no agreed 'it') to my own data. Through this process I was able to understand FDA as bringing a unique perspective to interpret data.

The experience of applying FDA to my data has developed my ability to think flexibly whilst maintaining a particular 'stance' based on the philosophy of Foucault. Taking a Foucauldian perspective locates the experience of the individual in power relationships as they exist in institutions and society as a whole. However, the Foucauldian philosophies that underpin FDA can be criticised for the absence of an agreed methodology, and potentially being based on Foucault's own personal life experiences.

While it is widely accepted that too little is known about Foucault's private life to be able to say with any certainty to what extent it influenced his philosophical work and writings (Gutting, 2005), it is conceivable that his early exposure to powerful academics, his authoritarian father and receiving psychiatric care could have influenced his ideas about power and knowledge.

It has been speculated that Foucault attempted suicide in his early adulthood due to struggling with his homosexuality, resulting in him receiving psychiatric care. It is possible that this experience contributed to his ideas about sexuality, the medical profession holding power, surveillance and control. Foucault posed that with changes in knowledge, came new power relationships that were played out in peoples' behaviour. Thus ever evolving discourses created new realities in which power was executed through institutions such as prisons, schools and hospitals through modes such as surveillance. Foucault can be criticised for taking a negative stance that focuses on how institutions can use power to control the 'subject' and not acknowledging the positive aspects of knowledge and power within

institutions such as hospitals e.g. the power to keep somebody safe. Perhaps this negative stance reflects his perception of his own life experiences.

The use of FDA to analyse the GPs' discourses regarding EU/AS was appropriate given that power relationships are prominent in EU/AS and the ongoing debate in the UK. However, the potential impact of my power as the researcher in relation to data creation, analysis and interpretation should be acknowledged. Discourse analysis is primarily applied to naturally occurring discourses, but due to the lack of availability of any records or texts reflecting the discourses of UK GPs regarding EU/AS, it was necessary to generate data through semi-structured interviews. Through developing and using an interview schedule I created an artificial structure that may have influenced the GPs' discourses. Furthermore, the GPs may have felt under 'surveillance' by me, as the researcher, with the power to report any concerning practice to the supervisory team. This may have influenced the GPs' discourses.

Although my own views and power could potentially have influenced the GPs' discourses, discourse analysis is not concerned with revealing one definitive truth but instead provides an account of a reality, as constructed by discourse at any given time. Therefore, the potential impact of my own views and power on the GPs' discourses, and the interpretation of these discourses, must be acknowledged and considered. Through this acknowledgment and consideration, the potential impact can be minimised and contextualised.

Although I do not hold strong views regarding EU/AS and do not take a position of either supporting or opposing its legalisation in the UK, it was noted that several of the GPs engaged in seeking my approval when giving their views in the interview. Furthermore, one

GP commented at the end of the interview “I’ve used terminology to you today because you’re a Psychologist that most lay people wouldn’t have understood” (Peter), indicating an assumed level of knowledge and power associated with my profession that influenced Peter’s choice of language. Despite these limitations, FDA provides an alternative way of understanding and conceptualising a phenomenon such as EU/AS through discourses, power and knowledge as they are executed in a particular social context.

### ***The interface between ‘real life’ and academic research***

My experience as a ‘scientist-practitioner’ has been that there is sometimes a mismatch between the ‘real world’ and the academic literature. Therefore, having reviewed the literature on the factors that impact on the mental health of transgender older adults, I was fortunate enough to be able to meet with Sophie, who has identified as female since her teens and transitioned from male to female 11 years ago. I met with Sophie to discuss how the literature review matched her lived experience of identifying as transgender. Sophie has extensive knowledge and experience of transgender related issues and is involved in running transgender support groups, networks, delivering diversity training to health boards, schools, police services and is involved in large scale research projects across the UK. Although Sophie’s experience cannot be considered to be representative of other transgender older adults, she provided some very useful insights and contributed greatly to my learning experience and contextualising of the literature.

Sophie felt strongly that academic research generally focuses on the negative experiences of transgender individuals, such as being victimised and experiencing stigma, and neglects the positive experiences. She highlighted that quantitative questionnaires as well as qualitative studies often ask questions regarding the experience of victimisation

without clarifying or reporting when this experience happened. She explained that in her experience, most transgender adults, and particularly older adults, will have experienced some form of victimisation, but for most this will have happened 20 or maybe 30 years ago. Sophie felt that the lack of consideration in research of when the victimisation took place results in a disproportionately negative representation of the experience of being transgender in current society. Furthermore, she felt that this has a negative, fear inducing impact on those who are experiencing confusion regarding their gender identity.

Sophie spoke about the difficulties she, and others she knows, have experienced when accessing physical and mental health care, and that it is not uncommon for people who identify as transgender to experience additional end of life difficulties. This included the impact of dementia on gender identity, where people might forget that they have transitioned from one gender to another and become confused and distressed. She spoke about how families, who have been reluctant to accept the gender identity of a loved one, often revert to calling the person by their birth name and sex once they have died, including at their funeral. Thus disregarding the person's wishes about how they are remembered. Sophie felt the literature review was important, especially as there are many aspects of transgender aging and mental health that are not yet sufficiently addressed in the literature.

### **Implications for clinical practice**

The clinical implications of both papers are discussed in relation to the person, clinicians, training, service level, and on a societal level.

### ***Implications for the person and clinicians/professionals***

Both papers highlight the influence and impact of societal norms as reflected in discourses on individuals and on services. For GPs, the discourses around EU/AS led to a fear of being labelled the 'doctor of death' if EU/AS were to be legalised in the UK. This led GPs to respond with a 'someone else should do it' discourse, disengaging from the consequences of a possible change in legal status of EU/AS. For some transgender older adults, the discourses and stigma associated with not conforming to societal gender norms influenced their decisions to disclose their gender identity, and even to join military service in the hope that it would reduce their transgender feelings (Fabbre, 2015).

It is important for health and social care professionals who may be working with transgender older adults to be aware of how societal norms/discourses, such as gender expectations, may impact on transgender older adults; for example, they may be less willing to disclose their gender identity for fear of rejection (McCann, Sharek, Higgins, Sheerin, & Glacken, 2013). It is also important for professionals to be aware of how such discourses may influence their own behaviour, e.g. making assumptions or decisions based on biological sex.

Both papers have emphasised the challenges in communication and bringing views into the open. Foucault proposed that this type of behaviour is directly influenced by the related discourses (Wetherell, Taylor & Yates, 2004). This was evident in the empirical study when the GPs were confronted with the uncomfortable 'medicine is limited' discourse. One of the behaviours that GPs described engaging in, to avoid the 'medicine is limited' discourse that highlighted their lack of knowledge, was shutting down any conversations the patients tried to initiate regarding EU/AS. This can be problematic given the importance of patient-GP

communication in developing a good patient-GP relationship and patient satisfaction (You et al., 2014). It may be more helpful for GPs to explore patient concerns regarding end of life issues, despite limited options, rather than shutting down communication. This may help to protect the patient-GP relationship and address patient worries.

### ***Training implications***

Both papers highlight that a lack of knowledge about a particular issue can result in reluctance to engage in discussions. Therefore, offering staff training that addresses end of life issues may be an effective way of enhancing and encouraging patient-professional communication. GPs and other staff who are involved in delivering palliative and end of life care also need to be aware of the specific needs of minority groups, such as transgender older adults. Awareness of gender related issues may contribute to staff feeling more confident in discussing and delivering appropriate care to transgender older adults. Training on transgender terminology may also facilitate communication around transgender related issues. Although not specific to older adults, the importance of staff training on transgender related issues and mental health has been recognised in the 'LGBT People & Mental Health: Guidance for Services and Practitioners' produced by the National LGB&T Partnership (2015a).

Incorporating research findings, such as that the factors that impact on the mental health of transgender older adults may differ from transgender adults and youth need to be incorporated in psycho-education for multi-disciplinary teams who are directly involved in the care and support of transgender older adults, and for service users themselves. Regular supervision for staff who are delivering mental health and end of life services may also be a useful way to ensure diversity needs are being met.

### ***Service level implications***

GPs' discourses regarding EU/AS revealed that GPs are usually in the 'expert' position and have the power to make end of life decisions, such as when to increase pain killing medication. However, in the 'medicine is limited' discourse GPs had reached the limits of their knowledge and skills to manage end of life situations. Therefore, they 'called in' colleagues who were seen to have more expertise and medical authority. This highlights the importance of multi-disciplinary team meetings in palliative care, where responsibility and knowledge can be shared. Furthermore, there is a need for services to be transparent and openly discuss end of life options, what is and what is not available, with patients, without fear of repercussions for the professionals. Services involved in end of life care need to promote the use of end of life planning and advance directives, as this can give patients a sense of control over the end of their life. Indeed, in the GPs' discourses, patients raised the possibility of EU/AS, and this may have been an attempt at gaining control over the end of their life.

Services also need to be aware that transgender older adults are more likely to identify friends as potential care givers and for the role of the next of kin, than the general older adult population (De Vries & Gutman, 2016). This can be problematic as friends are often not recognised in service policies that are based on normative family structures and care provision (Barker, 2002). Indeed, it was noted in the empirical paper that GPs often addressed family members regarding end of life decisions. Transgender older adults may lack important social support networks (Fredriksen-Goldsen et al., 2013), which makes them less likely to have discussed or made end of life plans, such as living wills (De Vries &

Gutman, 2016). Services may need to consider the role of patients' friends when developing service policies related to end of life care.

Although information sheets, such as the 'Trans Health Factsheet on Ageing – Rising to the challenge' (The National LGB&T Partnership, 2015b) have recently become available (see Appendix J), the literature reviewed and Sophie's comments would suggest this information is not yet being used routinely. Furthermore, services need to be sensitive to and facilitate personal choice, both for patients and for staff. This was highlighted by GPs in the 'patients have the right to EU/AS, but do not count on me' discourse, where GPs expressed that they should have the option to opt out of practices that are not in keeping with their personal values.

### ***On a societal level***

The debate regarding the legalisation of EU/AS in the UK is ongoing. However, as reported in the empirical paper, GPs have confused and limited knowledge regarding EU/AS which may hinder their ability to successfully engage in the debate. Opinions of the medical profession have previously been considered in parliament during debates (The Commission on Assisted Dying, 2012), and have contributed to decisions not to legalise EU/AS in the UK. Therefore, there may be a need, on a societal level, for a critical evaluation of the opinions that contribute to this decision making process. Both papers highlight the need for the recognition of individuals' values and rights to make life choices, whether that is related to gender expression or the right to opt out of providing a service that is not in line with personal values.



## **Implications for theory development and future research**

### ***Research philosophy***

Many psychological and societal theories and models of aging, such as the successful aging paradigm, are underpinned by normative assumptions, including heteronormativity (Fabre, 2015); the assumption in western cultures that biological sex, gender and sexuality have a 'natural' alignment (e.g. the expectation that females are feminine).

Models that are based on normative assumptions, such as heteronormativity, are perpetuating these assumptions through researching and describing people against these assumptions, and through the association of successful aging with people who meet these assumptions. The successful aging paradigm has been criticised for lacking consideration of inequality, gender and race (Fabre, 2015). This criticism is particularly relevant for transgender people who might not meet the criteria of heteronormativity. Transgender older adults are more likely to experience victimisation, discrimination, a lack of social support and physical health problems associated with transition. They are therefore more likely to experience poor mental health, be reluctant to engage with professionals and are less likely to make end of life plans. Given these difficulties, heteronormative models would conceptualise transgender older adults as not aging successfully. This can also lead to a lack of specialist service provision for minority groups such as transgender older adults, as the services are aimed at people who meet the normative assumptions.

This perpetuating of normative assumptions and lack of service provision can further contribute to an internalised stigma (Maylon, 1982) of not meeting the expected behaviours and outcomes for one's grouping. Therefore, it is important to critically evaluate the use of these models in underpinning research and service provision.

While it must be acknowledged that transgender older adults are more likely to experience additional difficulties than the general older adult population, the question must also be asked; how helpful is it for research and services to be underpinned by models that are based on, and perpetuate, normative assumptions that may contribute to internalised stigma and that fail to consider positive aspects of transgender aging such as the development of a cohesive self? Indeed, Sophie stated that in her opinion academic research has a tendency to focus on the negative aspects of transgender aging and that this can be fear inducing.

The GPs' discourses regarding EU/AS also seem to be influenced by dominant research and care models that have normative assumptions. For example, within the palliative discourse, GPs spoke about the need for patients to have a 'good death'. GPs' descriptions of what a 'good death' would involve, appeared to be based on normative assumptions such as heteronormativity, the importance of independence and productivity. There was no reference made to gender, race or cohort effects. Furthermore, the impact of such normative assumptions was apparent in Lamers and Williams (2015) study of the discourses of older adults regarding EU/AS, who spoke about becoming dependent on others as an undesirable position.

Both studies, and Sophie's comments, suggest the need for increased development and use of models and theories of aging, communication, identity and social support that are not based exclusively on normative assumptions such as heteronormativity, independence and productivity. The development and use of such models and theories in research and service provision could in itself contribute to reducing (internalised) stigma as well as improving servicing provision for all, including minority groups.

### ***Future research***

While the 'palliative' discourse was consistent with the current UK legal position on EU/AS, the GPs' discourse that 'patients should have the right to EU/AS' is not consistent with the views of the majority of the GPs in other studies. One possible explanation for this inconsistency is that previous research, primarily large scale questionnaires, asks questions regarding GPs' views on EU/AS from the perspective that the GPs holds an active role in EU/AS. For example, a question in Seale's (2009b) study asks: 'A person with an incurable and a painful illness, from which they will die, for example, someone dying of cancer. Do you think that, if they ask for it, a doctor should ever be allowed by law to end their life, or not?' Such questions do not allow for the complexity of GPs' views.

When a survey included the option to make qualitative comments, (Seale, 2009b) those GPs who were in support of EU/AS stated that GPs should have the right to 'opt out' of involvement with EU/AS and that non-medical practitioners should be able to be involved. These added qualitative comments are consistent with the 'someone else should do it' discourse revealed by the GPs.

Furthermore, the wording used in questionnaires and the respondents' knowledge of EU/AS can bias responses (Hagelin, Nilstun, Hau, & Carlsson, 2004; Marcoux, Mishara, & Durand, 2007; Seale, 2009a). The finding that the GPs were not able to accurately define or distinguish between EU/AS, and that they were not opposed to EU/AS per se but rather opposed to their role in it, might bring a different perspective to the UK GPs' responses to such questionnaires regarding EU/AS.

When GPs considered that patients might have EU/AS as a potential end of life choice option, they felt that EU/AS was only acceptable if somebody was close to death, was

suffering a lot, and had a physical illness. They did not explicitly refer to the age at which death would occur nor the type of illness the patient would have. Future research investigating GPs' discourses regarding EU/AS when considering age or health condition might be worth exploring, as the older adults revealed an 'aged death' discourse (Lamers & Williams, 2015), where death was considered as the natural order and uneventful, and distinguished from untimely deaths in younger people. The discourses of other medical specialisms (e.g. oncology) and health professionals such as nurses, might reveal different discourses. The latter group might hold a very different discourse regarding EU/AS, as they seem to be the ones carrying out the decisions of their medical colleagues and are likely to be present when death occurs.

GPs described dementia as a 'tricky area' regarding EU/AS, due to patients' lacking the capacity to make end of life decisions. Dementia is also a concern for transgender older adults, who fear they will become confused about their gender and that professionals will deliver care based on their previous gender. Both studies highlight the need for further research into dementia and its impact on end of life decisions.

## References

- Barker, J. C. (2002). Neighbours, friends and other non-kin caregivers of community-living dependent elders. *Journals of Gerontology, Series B: Psychological Sciences and Social Science*, 57(3), 158-167. doi.org/10.1093/geronb/57.3.S158
- De Vries, B., & Gutman, G. (2016). End-of-life preparations among LGBT older adults. *Journal of the American Society on Aging*, 40(2), 46-48. Retrieved from <http://search.proquest.com.ezproxy.bangor.ac.uk/docview/1812643835/fulltext/7C69446E66CC4426PQ/1?accountid=14874>
- Fabbre, V. D. (2015). Gender transitions in later life: A queer perspective on successful aging. *The Gerontologist*, 55, 144-153. doi:10.1093/geront/gnu079
- Fredriksen-Goldsen, K. I., Cook-Daniels, L., Kim, H., Erosheva, E. A., Emllet, C. A., Hoy-Ellis, C. P., Goldsen, J., & Muraco, A. (2013). Physical and mental health of transgender older adults: An at-risk and underserved population. *The Gerontologist*, 54(3), 488-500. doi: 10.1093/geront/gnt021
- Fredriksen-Goldsen, K. I., Kim, H., Shiu, C., Goldsen, J., & Emllet, C. A. (2015). Successful aging among LGBT older adults: Physical and mental health-related quality of life by age group. *The Gerontologist*, 55(1), 154-168. doi:10.1093/geront/gnu081
- Gutting, G. (2005). *FOUCAULT. A Very Short Introduction*. Oxford: Oxford University Press.
- Lamers, C. P. T., & Williams, R. R. (2015). Older people's discourses about euthanasia and assisted suicide: A Foucauldian exploration. *The Gerontologist*, 56(6), 1072-1081. doi:10.1093/geront/gnv102

- Maylon, A. K. (1982). Psychotherapeutic implications of internalized homophobia in gay men. *Journal of Homosexuality*, 7, 59-69. doi.org/10.1300/J082v07n02\_08
- McCann, E., Sharek, D., Higgins, A., Sheerin, F., & Glacken, M. (2013). Lesbian, gay, bisexual and transgender older people in Ireland: Mental health issues. *Aging and Mental Health*, 17(3), 358-365. doi:10.1080/13607863.2012.751583
- Seale, C. (2009). Legalisation of euthanasia or physician-assisted suicide: survey of doctors' attitudes. *Palliative Medicine*, 23, 205-212. doi:10.1177/0269216308102041
- The Commission on Assisted dying. (2012). *The Falconer Report*. Retrieved from [https://www.demos.co.uk/files/476\\_CoAD\\_FinalReport\\_158x240\\_I\\_web\\_single-NEW\\_.pdf?1328113363](https://www.demos.co.uk/files/476_CoAD_FinalReport_158x240_I_web_single-NEW_.pdf?1328113363)
- The National LGB&T Partnership. (2015a). *LGBT People & Mental Health: Guidance for Services and Practitioners*. Retrieved from <https://nationallgbtpartnershipdotorg.files.wordpress.com/2015/10/np-mental-health-service-practitioners.pdf>
- The National LGB&T Partnership. (2015b). *Trans Health Factsheet on Ageing – Rising to the challenge*. Retrieved from <https://nationallgbtpartnershipdotorg.files.wordpress.com/2012/07/np-trans-health-factsheet-ageing-final.pdf>
- Wetherell, M., Taylor, S., & Yates, S. J. (2004). *Discourse theory and practice: A reader*. London: SAGE Publications.
- You, J. J., Dodek, P., Lamontagne, F., Downar, J., Sinuff, T., Jiang, X., ... Heyland, K. (2014). What really matters in end-of-life discussions? Perspectives of patients in hospital

with serious illness and their families. *Canadian Medical Association Journal*,  
186(18), 679–87. doi:10.1503/cmaj.140673

**Appendix A**

**School of Psychology Ethics Application**



**Appendix B**

**School of Psychology Ethical Approval Letter**

Dear Emily,

2016-15670 General Practitioners and the euthanasia and assisted suicide debate in the United Kingdom: A Foucauldian exploration of their discourses

Your research proposal number 2016-15670 has been reviewed by the Psychology Ethics and Research Committee and the committee are now able to confirm ethical and governance approval for the above research on the basis described in the application form, protocol and supporting documentation. This approval lasts for a maximum of three years from this date.

Ethical approval is granted for the study as it was explicitly described in the application

If you wish to make any non-trivial modifications to the research project, please submit an amendment form to the committee, and copies of any of the original documents reviewed which have been altered as a result of the amendment. Please also inform the committee immediately if participants experience any unanticipated harm as a result of taking part in your research, or if any adverse reactions are reported in subsequent literature using the same technique elsewhere.

---

**Appendix C**

**Betsi Cadwaladr University Health Board Research and Development West**

**Department R&D approval application**

**Appendix D**

**Betsi Cadwaladr University Health Board Research and Development West**

**Department approval letter**

**Appendix E**

**Advert**

**Appendix F**  
**Information Sheet**

**Appendix G**

**Informed Consent Form**

**Appendix H**

**Interview Schedule**



**Appendix I**

**Example Transcript**

**Appendix J**

**Trans Health Factsheet on Ageing – Rising to the challenge**

**Appendix K**

**Word count statement**