

**Bangor University**

## **DOCTOR OF PHILOSOPHY**

### **Sheep urine patch nitrous oxide emissions: measurement and mitigation**

Marsden, Karina

*Award date:*  
2015

*Awarding institution:*  
Bangor University

[Link to publication](#)

#### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

#### **Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

**Sheep urine patch nitrous oxide emissions:  
Measurement and mitigation**

Karina A. Marsden

November (2015)



PRIFYSGOL  
**BANGOR**  
UNIVERSITY

A thesis submitted to Bangor University in candidature for the degree  
Philosophiae Doctor

School of Environment Natural Resources and Geography,  
Bangor University,  
Bangor,  
Gwynedd,  
LL57 2UW

## Declaration and Consent

### Details of the Work

I hereby agree to deposit the following item in the digital repository maintained by Bangor University and/or in any other repository authorized for use by Bangor University.

**Author Name:** Karina Anne Marsden

**Title:** Laughing gas (N<sub>2</sub>O), livestock urine and nitrification inhibitors: Spatial and temporal investigations

**Supervisor/Department:** Dave Chadwick / SENRGy

**Funding body (if any):** Bangor University

**Qualification/Degree obtained:** PhD

This item is a product of my own research endeavours and is covered by the agreement below in which the item is referred to as “the Work”. It is identical in content to that deposited in the Library, subject to point 4 below.

### Non-exclusive Rights

Rights granted to the digital repository through this agreement are entirely non-exclusive. I am free to publish the Work in its present version or future versions elsewhere.

I agree that Bangor University may electronically store, copy or translate the Work to any approved medium or format for the purpose of future preservation and accessibility. Bangor University is not under any obligation to reproduce or display the Work in the same formats or resolutions in which it was originally deposited.

### Bangor University Digital Repository

I understand that work deposited in the digital repository will be accessible to a wide variety of people and institutions, including automated agents and search engines via the World Wide Web.

I understand that once the Work is deposited, the item and its metadata may be incorporated into public access catalogues or services, national databases of electronic theses and dissertations such as the British Library’s EThOS or any service provided by the National Library of Wales.

I understand that the Work may be made available via the National Library of Wales Online Electronic Theses Service under the declared terms and conditions of use (<http://www.llgc.org.uk/index.php?id=4676>). I agree that as part of this service the National Library of Wales may electronically store, copy or convert the Work to any approved medium or format for the purpose of future preservation and accessibility. The National Library of Wales is not under any obligation to reproduce or display the Work in the same formats or resolutions in which it was originally deposited.

**Statement 1:**

This work has not previously been accepted in substance for any degree and is not being concurrently submitted in candidature for any degree unless as agreed by the University for approved dual awards.

Signed ..... (candidate)

Date .....

**Statement 2:**

This thesis is the result of my own investigations, except where otherwise stated. Where correction services have been used, the extent and nature of the correction is clearly marked in a footnote(s).

All other sources are acknowledged by footnotes and/or a bibliography.

Signed ..... (candidate)

Date .....

**Statement 3:**

I hereby give consent for my thesis, if accepted, to be available for photocopying, for inter-library loan and for electronic storage (subject to any constraints as defined in statement 4), and for the title and summary to be made available to outside organisations.

Signed ..... (candidate)

Date .....

**NB:** Candidates on whose behalf a bar on access has been approved by the Academic Registry should use the following version of **Statement 3:**

**Statement 3 (bar):**

I hereby give consent for my thesis, if accepted, to be available for photocopying, for inter-library loans and for electronic storage (subject to any constraints as defined in statement 4), after expiry of a bar on access.

Signed ..... (candidate)

Date .....

**Statement 4:**

Choose **one** of the following options

a) I agree to deposit an electronic copy of my thesis (the Work) in the Bangor University (BU) Institutional Digital Repository, the British Library ETHOS	
--	--

system, and/or in any other repository authorized for use by Bangor University and where necessary have gained the required permissions for the use of third party material.	
b) I agree to deposit an electronic copy of my thesis (the Work) in the Bangor University (BU) Institutional Digital Repository, the British Library ETHOS system, and/or in any other repository authorized for use by Bangor University when the approved <b>bar on access</b> has been lifted.	
c) I agree to submit my thesis (the Work) electronically via Bangor University's e-submission system, however I <b>opt-out</b> of the electronic deposit to the Bangor University (BU) Institutional Digital Repository, the British Library ETHOS system, and/or in any other repository authorized for use by Bangor University, due to lack of permissions for use of third party material.	

*Options B should only be used if a bar on access has been approved by the University.*

**In addition to the above I also agree to the following:**

1. That I am the author or have the authority of the author(s) to make this agreement and do hereby give Bangor University the right to make available the Work in the way described above.
2. That the electronic copy of the Work deposited in the digital repository and covered by this agreement, is identical in content to the paper copy of the Work deposited in the Bangor University Library, subject to point 4 below.
3. That I have exercised reasonable care to ensure that the Work is original and, to the best of my knowledge, does not breach any laws – including those relating to defamation, libel and copyright.
4. That I have, in instances where the intellectual property of other authors or copyright holders is included in the Work, and where appropriate, gained explicit permission for the inclusion of that material in the Work, and in the electronic form of the Work as accessed through the open access digital repository, *or* that I have identified and removed that material for which adequate and appropriate permission has not been obtained and which will be inaccessible via the digital repository.
5. That Bangor University does not hold any obligation to take legal action on behalf of the Depositor, or other rights holders, in the event of a breach of intellectual property rights, or any other right, in the material deposited.
6. That I will indemnify and keep indemnified Bangor University and the National Library of Wales from and against any loss, liability, claim or damage, including without limitation any related legal fees and court costs (on a full indemnity bases), related to any breach by myself of any term of this agreement.

Signature: .....

Date : .....

---

## Thesis Summary

---

Demand for livestock products are expected to rise due to increasing global population, urbanisation and affluence. Sustainably managing livestock excreta will be central to achieving an expansion in production whilst minimising environmental damage. The deposition of excreta to soil by livestock accounts for ca. 21% of the UK agricultural N<sub>2</sub>O emissions. Accurate quantification of N<sub>2</sub>O and assessments of the efficacy of mitigation technologies are key research areas for progressing toward enhanced sustainability and productivity in grazed grasslands. The overall thesis aims are to enhance understanding of N cycling and losses in sheep urine patches, as ‘hot-spots’ and ‘hot-moments’ for rapid nutrient cycling. Objectives were (i) to determine how sheep urine patch and environmental parameters influence N<sub>2</sub>O emissions, (ii) to determine the optimal way to accurately measure N<sub>2</sub>O emissions from sheep urine patches via the static chamber technique, and (iii) to assess the efficacy of synthetic nitrification inhibitors as a mitigation strategy for urine patch N<sub>2</sub>O emissions. Sheep-grazed grasslands were selected for study, based on the lack of current available evidence for these agroecosystems. N<sub>2</sub>O emissions were monitored from sheep urine-influenced soil in small incubation vessels, or by the static chamber technique in the field (manual and automated campaigns). The use of <sup>14</sup>C-labelled inhibitors were also employed in laboratory studies, to trace the fate of nitrification inhibitors in the plant-soil-microbe system and provide a better understanding of the factors that affect the efficacy of inhibitors to reduce N<sub>2</sub>O emissions. Urine patch size and N concentration were found to be important parameters influencing emissions of N<sub>2</sub>O from sheep urine patches. Emissions of N<sub>2</sub>O were generally lower than the IPCC default of 1% of the N applied in sheep excreta, where peaks in emissions occurred alongside rainfall events. Total extractable N, oxidation reduction potential and soil water-filled pore space were determined to be key drivers of N<sub>2</sub>O emissions from sheep urine under controlled conditions. The urine patch diffusional area was shown to be important for accurate quantification of N<sub>2</sub>O emissions when using the chamber technique; the importance of daily sampling of emissions, an assessment of the diurnal nature of N<sub>2</sub>O emissions and having a high number of replicate chambers to adequately represent the large spatial variability in N<sub>2</sub>O emissions was also confirmed. The nitrification inhibitors DCD and DMPP had contrasting behaviours in differing soil types. DCD had a greater sorption in comparison to DMPP and microbial uptake and degradation were concluded to be important parameters influencing their effective period in the soil. A graminaceous plant was shown to be able to acquire DCD intact through its roots and translocate the compound to shoots which raises concerns about contamination of food products. A liquid application of DMPP was not effective in reducing cumulative N<sub>2</sub>O emissions from sheep urine patches in the field. The efficacy of nitrification inhibitors to reduce N<sub>2</sub>O appears to vary widely, nevertheless they are a mitigation strategy that could be implemented in the short term. Achieving enhanced sustainability and productivity in grazed grasslands is a complex problem, requiring an interdisciplinary approach and the involvement of policy-makers and farmers to resolve. There are several mitigation strategies available or being developed, and some which require more research before being practicable. Advances in technologies to measure and mitigate N<sub>2</sub>O emissions will greatly enhance our knowledge of N cycling and losses, and the potential to alleviate such losses from the urine patch environment in the near future.

---

# Thesis Contents

---

Declaration and consent.....	i
Thesis summary.....	iv
Acknowledgements.....	ix
Abbreviations.....	x
List of tables.....	xi
List of figures.....	xiii
<b>Chapter 1: Introduction.....</b>	<b>1</b>
1.1 Introduction to the challenges facing the livestock sector.....	2
1.2 Thesis background and rationale.....	2
1.3 Thesis aims and objectives.....	5
1.4 Experimental chapter information.....	8
1.5 Discussion points arising from objectives.....	9
1.6 References.....	10
<b>Chapter 2: Literature Review.....</b>	<b>15</b>
2.1 Introduction.....	16
2.2 The reactive N cascade in the context of livestock production.....	17
2.3 Sheep urine patch characteristics.....	22
2.4 Changes in soil-microbe-plant properties following urine deposition.....	32
2.5 Terrestrial transformations of N leading to N <sub>2</sub> O formation.....	37
2.6 Measurement of N <sub>2</sub> O emissions from urine patches.....	42
2.7 Nitrification inhibitors as tools to mitigate sheep urine patch N <sub>2</sub> O emissions .....	44
2.8 Conclusions.....	49
2.9 References.....	51
<b>Chapter 3: The urine patch diffusional area: an important N<sub>2</sub>O source?.....</b>	<b>71</b>
Abstract.....	72
3.1 Introduction.....	73
3.2 Materials and methods.....	75

3.3 Results.....	82
3.4 Discussion.....	90
3.5 Conclusions.....	97
3.6 Acknowledgements.....	98
3.7 References.....	99
<b>Chapter 4: Disentangling the effect of sheep urine patch size and N loading rate on cumulative N<sub>2</sub>O emissions .....</b>	<b>107</b>
Abstract.....	108
4.1 Introduction.....	109
4.2 Materials and Methods.....	111
4.3 Results.....	118
4.4 Discussion.....	126
4.5 Conclusions.....	131
4.6 Acknowledgements.....	132
4.7 References.....	133
<b>Chapter 5: The mobility of nitrification inhibitors under simulated ruminant urine deposition and rainfall: a comparison between DCD and DMPP.....</b>	<b>139</b>
Abstract.....	140
5.1 Introduction.....	141
5.2 Materials and methods.....	144
5.3 Results.....	151
5.4 Discussion.....	161
5.5 Conclusions.....	166
5.6 Acknowledgements.....	167
5.7 References.....	168
<b>Chapter 6: Plant acquisition and metabolism of the synthetic nitrification inhibitor dicyandiamide and naturally-occurring guanidine from agricultural soils.....</b>	<b>175</b>
Abstract.....	176
6.1 Introduction.....	177



6.2 Materials and methods.....	179
6.3 Results.....	186
6.4 Discussion.....	195
6.5 Acknowledgements.....	201
6.6 References.....	202
<b>Chapter 7: Effect of time-since DMPP application on mitigating N<sub>2</sub>O emissions from sheep urine patches.....</b>	<b>209</b>
Abstract.....	210
7.1 Introduction.....	211
7.2 Materials and methods.....	213
7.3 Results.....	220
7.4 Discussion.....	231
7.5 Conclusions.....	238
7.6 Acknowledgements.....	239
7.7 References.....	240
<b>Chapter 8: Discussion and recommendations for future research.....</b>	<b>247</b>
8.1 Introduction.....	248
8.2 General discussion.....	248
8.3 What implications do variations in urine patch and environmental parameters have on the upscaling of N <sub>2</sub> O emissions from grazed grasslands?.....	252
8.4 How should the chamber technique be deployed, to monitor N <sub>2</sub> O emissions from urine patches?.....	255
8.5 Are nitrification inhibitors a suitable strategy for reducing N <sub>2</sub> O emissions from grazed grasslands?.....	256
8.6 What alternative strategies are there for the mitigation of N <sub>2</sub> O emissions from urine patches?.....	258
8.7 How should future research be prioritised, in order to achieve enhanced productivity and sustainability in grazed grasslands?.....	267
8.8 Conclusions.....	269
8.9 References.....	271

<b>Appendix 1:</b> Supplementary material for Chapter 3.....	283
<b>Appendix 2:</b> Supplementary material for Chapter 4.....	286
<b>Appendix 3:</b> Details of an attempted study utilising lux marked <i>Nitrosomonas europaea</i> ...	287

---

## Acknowledgements

---

Firstly, I would like to thank my supervisors Dave Chadwick and Davey Jones for their excellent supervision throughout my PhD study. Dave - I hope I have done you proud as your first 'Bangor PhD', thanks for always having your door open and being available for helpful discussions. I appreciate the opportunities you have presented to me, not just for my PhD, but for my future career too. Davey – I would also extend my sincere thanks for the opportunities you have presented to me all the way from undergraduate level. Working for you prior to my PhD provided me with many of the experiences, resources and faculties required to make it this far. Thanks also for reading manuscripts in time for strict deadlines! In addition, thanks go to Paul Hill, who has always been happy to discuss my work and for providing helpful comments on the co-authored manuscripts.

I would like to thank all the technical staff: John Evans, Sarah Chesworth, Jonathan Roberts, Gordon Turner, Helen Simpson and Al Jones who have all been very helpful. A special thanks to Mark Hughes and Llinos Hughes for assistance setting up field and urine collection trials and to Will and Ruth for allowing me access to their flock at the University farm.

So many peers and colleagues have provided help and assistance along the way - too many to mention here! I appreciate from the bottom of my heart the people that spared some time for me, no matter how short. A particular thanks to Antonio Martinez for working with me and being so helpful during his stay at ECW and to Jairo Cardoso for helping me with sheep urine collection – I wasn't expecting anyone to do that! A massive thanks to Helen Glanville and Rory Shaw, when disaster struck you both really cheered me up with afternoon tea on the pier.

Thanks go to my family for their endless love, support and encouragement and just for listening to me whinge! I can't wait to spend a bit of quality time with my 'sibs' and 'nibs'.

Last but definitely not least thanks to Tali, for always being there for me, believing in me and encouraging me. You have definitely done more than your fair share of cooking and cleaning these past few years! Here's to better times ahead.

---

## Abbreviations

---

<b>AMO</b> - Ammonia monooxygenase	<b>IPCC</b> - Intergovernmental Panel on Climate Change
<b>ANOVA</b> - Analysis of variance	<b>K</b> - Potassium
<b>C</b> - Carbon	<b>LSD</b> - Least squares difference
<b>Ca</b> - Calcium	<b>N<sub>2</sub></b> - Nitrogen molecule
<b>CEC</b> - Cation exchange capacity	<b>Na</b> - Sodium
<b>CH<sub>4</sub></b> - Methane	<b>NAR</b> - Nitrate reductase
<b>CO<sub>2</sub></b> - Carbon dioxide	<b>NH<sub>3</sub></b> - Ammonia
<b>DCD</b> - Dicyandiamide	<b>NH<sub>4</sub><sup>+</sup></b> - Ammonium
<b>DEFRA</b> – Department of Environment Food and Rural Affairs	<b>NI</b> - Nitrification inhibitors
<b>DMPP</b> - 3,4-Dimethylpyrazole phosphate	<b>NIR</b> - Nitrite reductase
<b>DMSO</b> - Dimethylsulfoxide	<b>NO<sub>2</sub><sup>-</sup></b> - Nitrite
<b>DNRA</b> - Dissimilatory nitrate reduction to ammonia	<b>NO<sub>3</sub><sup>-</sup></b> - Nitrate
<b>DOC</b> - Dissolved organic carbon	<b>N<sub>2</sub>O</b> - Nitrous oxide
<b>DON</b> - Dissolved organic nitrogen	<b>NO</b> - Nitric oxide
<b>DOP</b> – Dissolved organic phosphorus	<b>NO<sub>2</sub></b> - Nitrogen dioxide
<b>DW</b> - Dry weight	<b>NO<sub>x</sub></b> - Nitric oxide and/or nitrogen dioxide
<b>EC</b> - Electrical conductivity	<b>NOR</b> - Nitric oxide reductase
<b>ECD</b> - Electron capture detector	<b>NOS</b> - Nitrous oxide reductase
<b>EF</b> - Emission factor	<b>NUE</b> - Nitrogen use efficiency
<b>FAO</b> - Food and Agriculture Organization	<b>O<sub>2</sub></b> - Oxygen
<b>FID</b> - Flame ionization detector	<b>OCT</b> - Organic cation transporter
<b>FW</b> - Fresh weight	<b>ORP</b> - Oxidation reduction potential
<b>GHG</b> - Greenhouse gas	<b>P</b> - Phosphorus
<b>GC</b> - Gas chromatograph	<b>PLFA</b> - Phospholipid fatty acid
<b>GPS</b> - Global positioning system	<b>PO<sub>4</sub><sup>3-</sup></b> - Phosphate
<b>HAO</b> - Hydroxylamine oxidoreductase	<b>SEM</b> – Standard error of the mean
<b>HPLC</b> - High performance liquid chromatography	<b>UK</b> - United Kingdom
	<b>UV</b> - Ultraviolet
	<b>WFPS</b> – Water-filled pore space

---

## List of Tables

---

**Table 3.1** Properties of the Eutric Cambisol used to fill soil mesocosms. Values represent means  $\pm$  SEM ( $n = 4$ ) and results are reported on a dry weight basis.

**Table 3.2** Properties of sheep urine, applied to Eutric Cambisol mesocosms. Values represent means  $\pm$  SEM ( $n = 3$ ), where replicates are analytical replicates of urine combined from 5 individual sheep.

**Table 3.3** Multiple regression analysis with  $\ln N_2O$  as the dependent variable,  $\ln$  total extractable soil nitrogen (TN) and  $\ln$  oxidation reduction potential (ORP) as predictor variables and water-filled pore space (WFPS; 50% and 70%) as a categorical predictor variable.

**Table 4.1** Properties of the Eutric Cambisol at the field site receiving artificial urine application. Values represent means  $\pm$  SEM ( $n = 3$ ) and results are reported on a dry weight basis.

**Table 4.2** Pasture biomass ( $t DM ha^{-1}$ ) and foliar N content (%) 2 weeks and 4 weeks following artificial sheep urine application to a Eutric Cambisol. Values represent means  $\pm$  SEM ( $n = 3$ ), and letters indicate significant differences ( $p < 0.05$ ).

**Table 5.1** Properties of soils used to fill the soil columns, values represent means  $\pm$  SEM,  $n = 3$ , letters indicate significant differences between the soils, results are reported on a dry soil weight basis and N.D. stands for not determined.

**Table 5.2** Soil-to-solution partition coefficient ( $K_d$ ) for the nitrification inhibitors DCD and DMPP at field relevant concentrations (1 and 10  $mg l^{-1}$ ) with and without the presence of sheep urine. Values represent means ( $n = 3$ )  $\pm$  SEM.

**Table 6.1** Soil properties of Eutric Cambisol and Sapric Histosol used in soil rhizosphere microcosms. Values represent means  $\pm$  SEM,  $n = 4$ , letters indicate significant differences between the two soils and results are reported on a dry soil weight basis.

**Table 6.2** Summary of results for wheat uptake rates and tissue localization of  $^{14}\text{C}$ -DCD and  $^{14}\text{C}$ -guanidine (0.01, 0.1 and 1 mM) under sterile conditions and from Eutric Cambisol and Sapric Histosol rhizosphere microcosms following a 2 h chase period. Values represent means  $\pm$  SEM ( $n = 4$ ).

**Table 6.3** Summary of results for microbial mineralization, microbial uptake and sorption of  $^{14}\text{C}$ -DCD and  $^{14}\text{C}$ -guanidine (0.01, 0.1 and 1 mM) in a Eutric Cambisol and a Sapric Histosol after 2 h. Values represent means  $\pm$  SEM ( $n = 4$ ).

**Table 7.1** Properties of the Eutric Cambisol at the field site receiving urine and DMPP applications. Values represent means  $\pm$  SEM ( $n = 3$ ) and results are reported on a dry weight basis.

---

## List of Figures

---

- Figure 1.1** The nitrogen cycle of grazed pastures, adapted from Saggar et al. (2013).
- Figure 2.1** Schematic of the N cycle. Source: Stark and Richards (2008).
- Figure 2.2** Diagrammatic representation of the urine patch area, where  $r_w$  represents the wetted area radius,  $r_p$  represents the pasture response area radius and  $d$  represents the depth of urine infiltration. Source: Li et al. (2012).
- Figure 2.3** Distribution of  $\text{Br}^-$  from simulated sheep and cattle urine, using potassium bromide solution as a tracer. Source: Williams and Haynes (1994).
- Figure 2.4** The microbial pathways leading to  $\text{N}_2\text{O}$ , with environmental conditions favourable for each process and enzymes regulating each pathway. Source: Baggs (2011).
- Figure 3.1** Aerial schematic view of the Eutric Cambisol mesocosms, repacked to a depth of 5 cm. The shaded region represents the area of direct urine application, labelled rows display sampling regions and T1 to T6 represent successive sampling time points, where further time points were sampled from parallel mesocosms.
- Figure 3.2** Changes in soil pH (panels a, b, c and d), electrical conductivity (EC; panels e, f, g and h) and oxidation reduction potential (ORP; panels i, j, k and l) following sheep urine application to a Eutric Cambisol, maintained at either 50% or 70% water-filled pore space (WFPS), and sampled at increasing distances away from the direct area of application (Zone A, 0-3 cm: panels a, e and i; Zone B, 3-6 cm: panels b, f and j; Zone C, 6-9 cm: panels c, g and k; Zone F, control: panels d, h and l). Symbols represent means  $\pm$  SEM ( $n = 4$ ). Figure legend applies to all panels and text on the top row of panels applies to each respective column.
- Figure 3.3** Soil extractable total dissolved nitrogen (panels a, b, c, d), ammonium (panels e, f, g, h), nitrate (panels i, j, k and l), nitrous oxide emissions (panels m, n, o and p)

and extractable dissolved organic carbon (panels q, r, s and t) following sheep urine application to a Eutric Cambisol, maintained at either 50% or 70% water-filled pore space (WFPS), and sampled at increasing distances away from the direct area of application. Symbols represent means  $\pm$  SEM ( $n = 4$ ). Figure legend applies to all panels and text on the top row of panels applies to each respective column.

**Figure 3.4** Cumulative nitrous oxide emissions following sheep urine application to a Eutric Cambisol, maintained at either 50% or 70% water-filled pore space (WFPS) and sampled at increasing distances away from direct area of urine application. Bars represent means  $\pm$  SEM ( $n = 4$ ) and different letters indicate significant differences (Fisher's LSD,  $p < 0.05$ ).

**Figure 4.1** Mineral N dynamics following artificial sheep urine application of varying N concentration and patch size to a Eutric Cambisol, panel a) extractable soil ammonium-N panel b) extractable soil nitrate-N, panel c) soil solution nitrate-N and panel d) soil solution ammonium-N. Figure legend applies to all panels, symbols and error bars denote means  $\pm$  SEM, where  $n = 3$  for panels a) and b), however,  $n$  varies (1-3) for panels c) and d), due to variability in successful soil solution collection by Rhizon samplers.

**Figure 4.2** Dynamics of total extractable nitrogen and organic carbon (panels a and b, respectively) and soil solution nitrogen and dissolved organic carbon (panels c and d, respectively) following artificial sheep urine application (of varying patch size and N content) to a Eutric Cambisol. Figure legend applies to all panels, symbols and error bars denote means  $\pm$  SEM, where  $n = 3$  for panels a) and b), however,  $n$  varies (1-3) for panels c) and d), due to variability in successful soil solution collection by Rhizon samplers.



**Figure 4.3** Weather station data (panel a), nitrous oxide emissions (panel b) and water-filled pore space (WFPS; panel c) following application of artificial sheep urine of varying N concentration and patch size to a Eutric Cambisol. Figure legends apply to each respective panel, for both panels b and c, where the traces represent the treatment mean ( $n = 3$ ).

**Figure 4.4** Cumulative  $N_2O$ -N emissions over the duration of the field trial (40 days), where artificial urine of varying N content and patch size was applied to a Eutric Cambisol. Bars represent means  $\pm$  SEM ( $n = 3$ ).

**Figure 5.1** The percentage of extractable DCD- $^{14}C$  or DMPP- $^{14}C$  at depth fractions of sandy loam (a and d), sandy clay loam (b and e) and Sapric Histosol (c and f) columns, following a simulated 40 mm rainfall event (a, b and c) and a sheep urine deposition plus a 40 mm rainfall event (d, e and f). Soil type labels apply to each column and legends apply to each row of panels. Bars represent means ( $n = 3$ ) and error bars represent SEM.

**Figure 5.2** The percentage of extractable DCD- $^{14}C$  (a, b and c) or DMPP- $^{14}C$  (d, e and f) compared to  $NH_4^+$  at differing soil depth fractions following a simulated urine deposition plus 40 mm rainfall event applied to sandy loam (a and d), sandy clay loam (b and d) and Sapric Histosol (c and f) columns. Soil type labels apply to each column and legends apply to each row of panels. Bars represent means ( $n = 3$ ) and error bars denote SEM.

**Figure 5.3** Linear sorption isotherms for  $^{14}C$ -DCD (a, c and e) and  $^{14}C$ -DMPP (b, d and f) in either a 0.01 M  $CaCl_2$  or sheep urine matrix, in a sandy loam (a and b), sandy clay loam (c and d) and Sapric Histosol (e and f). Symbols represents means ( $n = 3$ ), bi-directional error bars represent SEM for sorption and equilibrium solution

concentrations, legends apply to each column of panels and soil type labels apply to each row of panels.

**Figure 5.4** Linear sorption isotherms of  $\text{NH}_4^+$  in 0.01 M  $\text{CaCl}_2$  matrix, in three soils (sandy loam and sandy clay loam textured Eutric Cambisol and a Sapric Histosol). Symbols represents means ( $n = 3$ ) and bi-directional error bars represent SEM for sorption and equilibrium solution concentrations.

**Figure 5.5** Cumulative desorption of  $^{14}\text{C}$ -DCD and  $^{14}\text{C}$ -DMPP in a sandy loam (a, b, c and d), sandy clay loam (e, f, g and h) and a Sapric Histosol (i, j, k and l) at 1 mg DCD  $\text{l}^{-1}$  (a, e and i), 10 mg DCD  $\text{l}^{-1}$  (b, f and j), 1 mg DMPP  $\text{l}^{-1}$  (c, g and k) and 10 mg DMPP  $\text{l}^{-1}$  (d, h and l) in either a 0.01 M  $\text{CaCl}_2$  or sheep urine matrix. Text in the top row of panels applies to each respective column of panels and legend applies to all panels. Symbols represents means ( $n = 3$ ) and error bars represent SEM for sorption and equilibrium solution concentrations.

**Figure 6.1** Molecular structure of the nitrification inhibitor, dicyandiamide (DCD; Panel A), and its naturally occurring analogue guanidine (Panel B), which are used in this study.

**Figure 6.2** Rates of  $^{14}\text{C}$ -dicyandiamide (DCD) and  $^{14}\text{C}$ -guanidine uptake by wheat (*Triticum aestivum*) roots grown in sterile hydroponic culture at three concentrations (0.01, 0.1 and 1 mM). Values represent means  $\pm$  SEM ( $n = 4$ ) and different letters indicate significant differences between means of compounds and concentrations (Fisher's LSD;  $p < 0.05$ ).

**Figure 6.3** Phosphorimages showing the distribution of  $^{14}\text{C}$ -dicyandiamide (DCD; Panel A) and  $^{14}\text{C}$ -guanidine (Panel B) within sterile wheat plants. The plants were exposed to each substrate (0.01 mM) for 2 h prior to imaging. The dark colour in the top

images represents the distribution of  $^{14}\text{C}$ -label within the plant tissue while, the images underneath are the corresponding photographs of the same plants.

**Figure 6.4** Cumulative mineralization of  $^{14}\text{C}$ -dicyandiamide (DCD) or  $^{14}\text{C}$ -guanidine in sterile wheat plants. Plants were exogenously supplied with each substrate (0.01 mM) for the whole 2 h monitoring period. Values represent mean  $\pm$  SEM ( $n = 4$ ).

**Figure 6.5** Uptake of  $^{14}\text{C}$ -dicyandiamide (DCD) or  $^{14}\text{C}$ -guanidine (0.01, 0.1 and 1 mM) by wheat plants grown in either a Eutric Cambisol (Panel A) or Sapric Histosol (Panel B), legend applies to both panels. Values represent means  $\pm$  SEM ( $n = 4$ ) and different letters indicate significant differences between means of different compounds, concentrations and between each soil type (Fisher's LSD;  $p < 0.05$ ).

**Figure 6.6** Mineralization of a)  $^{14}\text{C}$ -dicyandiamide (DCD) in a Eutric Cambisol b)  $^{14}\text{C}$ -DCD in Sapric Histosol c)  $^{14}\text{C}$ -guanidine in Eutric Cambisol and d)  $^{14}\text{C}$ -guanidine in Sapric Histosol at three concentrations (0.01, 0.1 and 1 mM). Legend applies to all panels and symbols represent means  $\pm$  SEM ( $n = 4$ ).

**Figure 6.7** Amount of added  $^{14}\text{C}$  label recovered in the soil solution pool and in the 0.5 M  $\text{K}_2\text{SO}_4$ -extractable pool from a)  $^{14}\text{C}$ -dicyandiamide (DCD; 0.01 mM) added to Eutric Cambisol b)  $^{14}\text{C}$ -DCD (0.01 mM) added to Sapric Histosol c)  $^{14}\text{C}$ -guanidine (0.01 mM) in Eutric Cambisol and d)  $^{14}\text{C}$ -guanidine (0.01 mM) in Sapric Histosol. Legend applies to all panels and symbols represent means  $\pm$  SEM ( $n = 4$ ).

**Figure 7.1** Rainfall, soil temperature and soil water-filled pore space during the course of the field trial. Hourly rainfall and soil (0-10 cm) temperature are shown in panel a) and water-filled pore space within the automated chambers are displayed in panel b). The date of measurements are displayed in dd/mm/yy format and legends apply to each respective panel only.

**Figure 7.2** N<sub>2</sub>O fluxes following sheep urine and DMPP applications to a Eutric Cambisol, where panel a) contains fluxes from treatments SU and NU and panels b), c) and d) display fluxes from SU + DMPP, SU + DMPP 2 and SU + DMPP 4, respectively. In each panel the black line represents the mean flux ( $n = 3$ ) and the grey shaded area represents the upper and lower bounds of the SEM. In treatment NU, error bars denote SEM. Arrows indicate the timing of either sheep urine (SU) or DMPP applications and the dates along the x-axis are in dd/mm/yy format.

**Figure 7.3** Cumulative nitrous oxide emissions, over 9 weeks post urine and DMPP applications to a Eutric Cambisol. Light grey bars represent means where  $n = 3$ , dark grey bars represent means where  $n = 2$ , with the high emitting replicates removed. Error bars denote SEM and different lower case letters represent significant differences (Welch's test, with Games-Howell pairwise comparisons) between treatments, where the same groupings were found both with and without outliers removed.

**Figure 7.4** Diurnal nature of N<sub>2</sub>O emissions and soil temperature over 6 days. The solid line represents the measured N<sub>2</sub>O flux from a single high emitting replicate from SU + DMPP 2 treatment, and the dashed line is soil temperature as measured by the Acclima SDI-12 combined temperature and moisture sensors in the chamber. The date/time format along the x-axis is expressed in dd/mm/yy hh:mm.

**Figure 7.5** Mineral nitrogen dynamics during the course of the field trial. Extractable ammonium-N can be seen in panel a) and extractable nitrate-N in panel b), which were measured on extracted soils from replicated plots (outside chambers). The soil solution ammonium-N is shown in panel c) and soil solution nitrate-N in panel d), which were measured from Rhizon samples located within the chambers. The figure legend applies to all panels, symbols represent means ( $n = 3$ ), and error bars

denote SEM. Arrows in panel d) represent timings of sheep urine (SU) and/or DMPP applications to the respective treatments. Asterisks above symbols represent level of significant differences (ANOVA) and was conducted for data in panels a) and b) only. The date format along the x-axis is dd/mm/yy.

**Figure 7.6** Total extractable and soil solution dissolved N and C dynamics. Total extractable dissolved N is displayed in panel a) and total extractable dissolved organic C is displayed in panel b), which were measured on extracted soils from replicated plots (outside chambers). Soil solution dissolved N is displayed in panel c) and soil solution dissolved organic C is displayed in panel d), which were measured using Rhizon samplers located within the chambers. The figure legend applies to all panels, symbols represent means ( $n = 3$ ), and error bars denote SEM. Arrows in panel d) represent timings of sheep urine (SU) and/or DMPP applications to the respective treatments. Asterisks above symbols represent level of significant differences (ANOVA) and was conducted for data in panels a) and b) only. The date along the x-axis is expressed in dd/mm/yy.

**Figure 8.1** Strategies to reduce N losses from grazed pastures.

---

# Chapter 1

---

## Introduction

Sheep urine patch nitrous oxide emissions: Measurement and mitigation

## **1.1 Introduction to the challenges facing the livestock industry**


Global demand for livestock products are set to increase in the future, due to projected increases in global population to 9 billion by 2050, increased income and urbanization (Smith et al., 2010; Thornton, 2010). Yet, the livestock sector is already estimated to contribute 14.5% of the global anthropogenic greenhouse gas (GHG) emissions (Gerber et al., 2013), in the form of carbon dioxide (CO<sub>2</sub>), methane (CH<sub>4</sub>) and nitrous oxide (N<sub>2</sub>O). Emissions of these three GHG's reflect losses of energy, organic matter and nitrogen (N) from agricultural systems (Eckard et al., 2010; Gerber et al., 2013). The major challenge facing the livestock sector is to meet increased demand efficiently and sustainably, whilst curbing its associated environmental impacts. In recent years, the potential for mitigation of non-CO<sub>2</sub> GHG's have been increasingly recognised (Reay et al., 2012).

## **1.2 Thesis background and rationale**

The research presented herein, is situated within the biogeochemical N cycle of grazed pastures (Figure 1), and specifically pertains to urine patches deposited by sheep. These sites are recognised as 'hot-spots' and 'hot-moments' for N<sub>2</sub>O emissions and other N loss processes (see Figure 1; van Groenigen et al., 2005; Cowan et al., 2015), undermining the efficiency of grazed grasslands. For a detailed analysis of N cycling within urine patches, readers are referred to Chapter 2 (a review of the literature).

A biogeochemical 'hot-spot' is defined as patches with disproportionately high reaction rates in comparison to the surrounding area (McClain et al., 2003) e.g. a small volume of soil with accelerated microbial processing rates compared to surrounding soil conditions. The concept does not solely apply to one biogeochemical process (Hagedorn and Bellamy, 2011), however, the large spatial variability in GHG emissions from soil are often attributed to hot-spots. Such GHG emission hot-spots require a supply of reactants and favourable conditions

for GHG production (McClain et al., 2003; Hagedorn and Bellamy, 2011; Kuzyakov and Blagodatskaya, 2015). A ‘hot-moment’ is defined as time periods with higher microbial processing rates in comparison to intervening periods (McClain et al., 2003). Hot-spots and hot-moments can occur separately, or at the same time. The urine patch supplies additions of reactive N, C and moisture to soil, removing many of the limitations to microbial activity and can stimulate the major  $N_2O$  producing pathways of nitrification and denitrification (Firestone and Davidson, 1989; Wrage et al., 2001; Bateman and Baggs, 2005). Thus, urine patches in grasslands can be described as hot-spots and hot-moments for  $N_2O$  emissions.



**Figure 1** The nitrogen cycle of grazed pastures, adapted from Saggar et al. (2013).

Due to the myriad of edaphic, environmental and management practices which can influence N cycling in the animal-plant-soil-atmosphere system, there are large uncertainties associated with broad-scale estimates of  $N_2O$  emissions from grazed grasslands (Hagedorn and



Bellamy, 2011; Butterbach-bahl et al., 2013). Nevertheless, recent estimates of the N<sub>2</sub>O emissions arising from grazing returns to grasslands are ca. 40% of the global anthropogenic N<sub>2</sub>O emissions arising from animal production systems (Oenema et al., 2005) and ca. 21% of the emissions of N<sub>2</sub>O from UK agriculture (UNFCCC, 2015). This represents one of the three largest sources of N<sub>2</sub>O from agriculture, alongside leaching and fertiliser applications, representing a potentially large emission reduction potential from the livestock sector.

This thesis broadly focuses on the measurement and mitigation of N<sub>2</sub>O from grassland soils, at the scale of the urine patch. Both accurate quantification of N<sub>2</sub>O emissions and assessments of suitable mitigation technologies are key research areas for making progress toward combatting the current challenges identified for the livestock sector, to reduce its GHG emissions. Sheep-grazed grasslands were selected for study, based on a need for specific data pertaining to such systems. This contrasts with urine patches in cattle based systems for which a large amount of scientific literature already exists.

For N<sub>2</sub>O emission measurements undertaken in this thesis, the use of gas chromatography was employed, where headspace gas samples were taken from small incubation vessels in laboratory studies, and from static-chambers in field studies. The standard chamber technique (Hutchinson and Mosier, 1980) is a widely employed technique for measuring N<sub>2</sub>O fluxes from field soils. Although several new advances in analytical technology for the measurement of N<sub>2</sub>O from soils are being made (this will be discussed in Chapter 2), the chamber technique is still likely to be used for the foreseeable future, due to its low cost. Therefore, methodologies for the accurate quantification of N<sub>2</sub>O emissions from urine patches, using this strategy, are warranted. An automated static chamber based system was also used, allowing measurements of N<sub>2</sub>O emissions from urine patches at a high temporal resolution (8 flux measurement per 24 h). This allowed an assessment of different sampling strategies based on varying the frequency of measurements.

There are a wide range of techniques which could be employed to mitigate N<sub>2</sub>O emissions from ruminant animals and urine patches, and this thesis focuses on only one potential strategy - the use of nitrification inhibitors. These compounds can be potentially utilised by application to the soil (Di and Cameron, 2003), by incorporating into livestock diet (Ledgard et al., 2008) or via addition to slurry and manure (Dittert et al., 2001; Hatch et al., 2005). The mechanisms and use of this technology in ruminant urine-influenced soil is reviewed in Chapter 2. Briefly, this mitigation technology was selected for research due to its potential to provide multiple benefits including improvements in nitrogen use efficiency, reductions in nitrate leaching and losses of gaseous N (Di and Cameron, 2002; Di and Cameron, 2007; Ruser and Schulz, 2015). Such techniques which have multiple benefits are likely to be important tools in any future mitigation strategies. However, the efficacy of nitrification inhibitors to provide such benefits varies widely under different edaphic and climatic conditions (Barth et al., 2001; Kelliher et al., 2008; Menéndez et al., 2012; Abalos et al., 2014). Further research is also required for determining their practicality, particularly in sheep urine-influenced soils. This thesis explores some of the factors influencing nitrification inhibitor efficacy, and focuses on two of the most commonly used nitrification inhibitors, dicyandiamide (DCD) and 3,4-dimethylpyrazole phosphate (DMPP) (Liu et al., 2013). Here, the use of <sup>14</sup>C-labelled nitrification inhibitors was employed in laboratory studies, to trace the fate of these inhibitors in the soil-microbe-plant system.

### **1.3 Thesis aims and objectives**

This section details the aims and objectives of the thesis, followed by a brief description of the relevant chapters which include experimental work pertaining to each objective. A list of the experimental chapter titles can be found in section 1.8; for individual experimental chapter hypotheses, please refer to the prepared manuscripts. Discussion points arising from

the overall objectives are included as bullet points in section 1.9. The purpose of these are to relate the insights gained from the current research, with recommendations for the direction of future research. This forms the basic structure of the final discussion section of the thesis (Chapter 8).

### *1.3.1 Thesis aims*

The overall aims of the thesis are to enhance understanding of N cycling and losses in urine patches, as hot-spots and hot-moments for rapid nutrient cycling; to determine how to accurately measure N<sub>2</sub>O emissions from sheep urine patches via the chamber technique, and to assess the efficacy of synthetic nitrification inhibitors as a strategy to reduce N<sub>2</sub>O emissions from sheep urine patches.

### *1.3.2 Objective 1*

*Determine how a) sheep urine patch parameters and b) environmental parameters influence N<sub>2</sub>O emissions from grassland soils.*

#### a) Sheep urine patch parameters

Urine patches vary widely in terms of N content, composition, patch size, volume and frequency (Betteridge et al., 2010). In Chapter 3, the horizontal diffusion of solutes out of the urine patch is assessed, investigating how this influences N<sub>2</sub>O emissions. In Chapter 4, the effects of urine patch size, N content and volume on N<sub>2</sub>O emissions are investigated within the typical ranges reported for sheep. In Chapter 7, average sheep urine volumes for Welsh Mountain ewes are determined and N<sub>2</sub>O emissions are quantified under field conditions.

#### b) Environmental parameters

Other than reactive N inputs, the main drivers of N<sub>2</sub>O emission from soil include moisture status (related to the soil oxygen status), temperature and the availability of labile C

(Smith, 2010; Butterbach-Bahl et al., 2013). The influence of soil water-filled pore space on emissions of N<sub>2</sub>O from urine-influenced soil is investigated under controlled conditions in Chapter 3. In Chapters 4 and 7, urine N is applied under different seasons including spring and late summer, respectively. This allows some assessment of N<sub>2</sub>O emissions under differing weather patterns, due to seasonality. The effect of spatial heterogeneity in soil conditions on N<sub>2</sub>O emissions from urine patches is also discussed in Chapter 7.

### *1.3.3 Objective 2*

*Determine how to accurately monitor N<sub>2</sub>O emissions from sheep urine patches, using the closed chamber technique.*

In Chapter 3, the importance of including the urine patch diffusional area for monitoring N<sub>2</sub>O emissions is assessed, which has implications for urine application methods within chambers, under experimental conditions. The high frequency automated chamber system was used in Chapters 4 and 7. In Chapter 4, the data are used to determine how different sampling strategies can influence the accuracy of N<sub>2</sub>O emission measurements from sheep urine patches. In Chapter 7, the diurnal nature of GHG fluxes are assessed, and insights gained are considered in the context of reduced frequency chamber measurements. Additionally, Chapter 7 considers how automated chamber based systems can be optimised for capturing both spatial and temporal variability in N<sub>2</sub>O emissions.

### *1.3.4 Objective 3*

*Investigate the factors which influence nitrification inhibitor efficacy in ruminant urine-influenced soils and assess their potential as an N<sub>2</sub>O mitigation strategy for sheep-grazed grasslands.*

Factors which influence the efficacy of nitrification inhibitors in soils are assessed in Chapters 5, 6 and 7. In Chapter 5, a comparative study was conducted for DCD and DMPP to determine their vertical mobility, and potential dis-location from urine derived ammonium, in a range of soils contrasting in texture and organic matter content. Chapter 6 assesses the magnitude of plant acquisition of DCD, both as a DCD depletion pathway and as a potential vector for contamination of food products. In Chapter 7, the direct measurement of the efficacy of DMPP in reducing N<sub>2</sub>O emissions from real sheep-derived urine patches, under summer conditions are investigated, varying the time-since DMPP application.

#### **1.4 Experimental chapter information**

The experimental chapters of the current thesis have been prepared in the style of journal article manuscripts. The title page of each experimental chapter contains details of the authors, author contributions to the manuscript, the current progress of each manuscript (e.g. published / accepted / submitted / not yet submitted) and any additional information, such as whether the work has / will be presented at a conference. The thesis consists of five experimental chapters, located in Chapters 3-7 of the current document. For continuity and clarity, the experimental chapters will be referred to as they appear in this thesis. The titles of each experimental chapters are as follows:

**Chapter 3:** The urine patch diffusional area: an important N<sub>2</sub>O source?

**Chapter 4:** Disentangling the effect of sheep urine patch size and N loading rate on cumulative N<sub>2</sub>O emissions.

**Chapter 5:** The mobility of nitrification inhibitors under simulated ruminant urine deposition and rainfall: a comparison between DCD and DMPP.

**Chapter 6:** Plant acquisition and metabolism of the synthetic nitrification inhibitor dicyandiamide and naturally-occurring guanidine from agricultural soils.

**Chapter 7:** Effect of time-since DMPP application on mitigating N<sub>2</sub>O emissions from sheep urine patches.

### **1.5 Discussion points arising from objectives**

The results and conclusions of the objectives of the thesis will be discussed in Chapter 8, where the following questions will form the basis of the structure of the discussion:

- What implications do variations in urine patch and environmental parameters have on the upscaling of N<sub>2</sub>O emissions from grazed grasslands?
- How should the chamber technique be deployed, to monitor N<sub>2</sub>O emissions from urine patches?
- Are nitrification inhibitors a suitable strategy for reducing N<sub>2</sub>O emissions from grazed grasslands?
- What alternative strategies are there for the mitigation of N<sub>2</sub>O emissions from urine patches?
- How should future research be prioritised, in order to achieve enhanced productivity and sustainability in grazed grasslands?

## 1.6 References

- Abalos, D., Jeffery, S., Sanz-Cobena, A., Guardia, G., Vallejo, A. 2014. Meta-analysis of the effect of urease and nitrification inhibitors on crop productivity and nitrogen use efficiency. *Agriculture, Ecosystems and Environment* 1, 136-144.
- Barth, G., Von Tucher, S., Schmidhalter, U. 2001. Influence of soil parameters on the effect of 3,4-dimethylpyrazole phosphate as a nitrification inhibitor. *Biology and Fertility of Soils* 24, 98-102.
- Betteridge, K., Hoogendorn, C., Costall, D., Carter, M., Griffiths, W. 2010. Sensors for detecting and logging spatial distribution of urine patches of grazing sheep and cattle. *Computers and Electronics in Agriculture* 73, 66-73.
- Butterbach-Bahl, K., Baggs, E.M., Dannenmann, M., Kiese, R., Zechmeister-Boltenstern, S. 2013. Nitrous oxide emissions from soils: how well do we understand the processes and their controls? *Philosophical Transactions of the Royal Society of London B: Biological Sciences* 368, 20130122.
- Cowan, N.J., Norman, P., Famulari, D., Levy, P.E., Reay, D.S., Skiba, U.M. 2015. Spatial variability and hotspots of soil N<sub>2</sub>O fluxes from intensively grazed grassland. *Biogeosciences* 12, 1585-1596.
- Di, H., Cameron, K.C. 2002. The use of a nitrification inhibitor, dicyandiamide (DCD), to decrease nitrate leaching and nitrous oxide emission in a simulated grazed and irrigated grassland. *Soil Use and Management* 4, 395-403.
- Di, H., Cameron, K.C. 2003. Mitigation of nitrous oxide emissions in spray-irrigated grazed grassland by treating the soil with dicyandiamide, a nitrification inhibitor. *Soil Use and Management* 19, 284-290.

- Di, H., Cameron, K.C. 2007. Effects of the nitrification inhibitor dicyandiamide on soil mineral N, pasture yield, nutrient uptake and pasture quality in a grazed pasture system. *Soil Use and Management* 23, 111-120.
- Dittert, K., Bol, R., King, R., Chadwick, D., Hatch, D. 2001. Use of a novel nitrification inhibitor to reduce nitrous oxide emission from <sup>15</sup>N-labelled dairy slurry injected into soil. *Rapid Communications in Mass Spectrometry* 15, 1291-1296.
- Eckard, R.J., Grainger, C., de Klein, C.A.M. 2010. Options for the abatement of methane and nitrous oxide from ruminant production: A review. *Livestock Science* 130, 47-56.
- Firestone, M.K, Davidson, E.A. 1989. Microbiological basis of NO and N<sub>2</sub>O production and consumption in soil. *Exchange of Trace Gases between Terrestrial Ecosystems and the Atmosphere* 47, 7-21.
- Gerber, P. J., Steinfeld, H., Henderson, B., Mottet, A., Opio, C., Dijkman, J., Tempio, G. 2013. Tackling climate change through livestock: a global assessment of emissions and mitigation opportunities. Food and Agriculture Organization of the United Nations (FAO), Rome.
- Hagedorn, F., Bellamy, P. 2011. Hot spots and hot moments for greenhouse gas emissions from soils. In *Soil Carbon in Sensitive European Ecosystems: From Science to Land Management*. John Wiley & Sons, Ltd. Chichester, UK.
- Hatch, D., Trindade, H., Cardenas, L., Carneiro, J., Hawkins, J., Scholefield, D., Chadwick, D. 2005. Laboratory study of the effects of two nitrification inhibitors on greenhouse gas emissions from a slurry-treated arable soil: impact of diurnal temperature cycle. *Biology and Fertility of Soils* 41, 225-232.
- Hutchinson, G.L., Mosier, A.R. 1980. Improved soil cover method for field measurement of nitrous oxide fluxes. *Soil Science Society of America Journal* 45, 311-316.



- Kelliher, F.M., Clough, T.J., Clark, H., Rys, G., Sedcole, J.R. 2008. The temperature dependence of dicyandiamide (DCD) degradation in soils: A data synthesis. *Soil Biology and Biochemistry* 40, 1878-1882.
- Ledgard, S.F., Menneer, J.C., Dexter, M.M., Kear, M.J., Lindsey, S., Peterse, J.S., Pacheco, D. 2008. A novel concept to reduce nitrogen losses from grazed pastures by administering soil nitrogen process inhibitors to ruminant animals: A study with sheep. *Agriculture, Ecosystems and Environment* 125, 148-158.
- Liu, C., Wang, K., Zheng, X. 2013. Effects of nitrification inhibitors (DCD and DMPP) on nitrous oxide emission, crop yield and nitrogen uptake in a wheat maize cropping system. *Biogeosciences* 10, 2427-2437.
- McClain, M.E., Boyer, E.W., Dent, C.L., Gergel, S.E., Grimm, N.B., Groffman, P.M., Hart, S.C., Harvey, J.W., Johnston, C.A., Mayorga, E., McDowell, W.H., Pinay, G. 2003. Biogeochemical hot spots and hot moments at the interface of terrestrial and aquatic systems. *Ecosystems* 6, 301-312.
- Menéndez, S., Barrena, I., Setien, I., González-Murua, C., Estavillo, J. M. 2012. Efficiency of nitrification inhibitor DMPP to reduce nitrous oxide emissions under different temperature and moisture conditions. *Soil Biology and Biochemistry* 53, 82-89.
- Oenema, O., Wrage, N., Velthof, G.L., van Groenigen, J.W., Dolfing, J., Kuikman, P.J. 2005. Trends in global nitrous oxide emissions from animal production systems. *Nutrient Cycling in Agroecosystems* 72, 51-56.
- Reay, D.S., Davidson, E.A., Smith, K.A., Smith, P., Melillo, J.M., Dentener, F., Crutzen, P.J. 2012. Global agriculture and nitrous oxide emissions. *Nature Climate Change* 2, 410-416.

- Ruser, R., Schulz, R. 2015. The effect of nitrification inhibitors on the nitrous oxide (N<sub>2</sub>O) release from agricultural soils – a review. *Journal of Plant Nutrition and Soil Science* 178, 171-188.
- Saggar, S., Jha, N., Deslippe, J., Bolan, N.S., Luo, J., Giltrap, D.L., Kim, D.-G., Zaman, M., Tillman, R.W. 2013. Denitrification and N<sub>2</sub>O:N<sub>2</sub> production in temperate grasslands: Processes, measurements, modelling and mitigating negative environmental impacts. *Science of the Total Environment* 465, 173-195.
- Smith, K. (2010). *Nitrous Oxide and Climate Change*. London, Earthscan.
- Thornton, P. K. 2010. Livestock production: recent trends, future prospects. *Philosophical Transactions of the Royal Society of London B: Biological Sciences* 365, 2853-2867.
- UNFCCC, 2015. Greenhouse Gas Inventory Data. United Nations Framework Convention on Climate Change.
- van Groenigen, J.W., Kuikman, P.J., de Groot, W.J.M., Velthof, G.L. 2005. Nitrous oxide emission from urine influenced soil as influenced by urine composition and soil physical conditions. *Soil Biology and Biochemistry* 37, 463-473.
- Wrage, N., Velthof, G.L., Van Beusichem, M.L., Oenema, O. 2001. Role of nitrifier denitrification in the production of nitrous oxide. *Soil Biology and Biochemistry* 33, 1723-1732.



---

## **Chapter 2**

---

### **Literature Review**

## 2.1 Introduction


Over the past few decades, there has been a large worldwide increase in the number of domestic animals, due to increasing consumption of meat protein *per capita* (Oenema et al., 2005). It is projected that the global population will reach 9 billion by 2050 (Smith, 2010), and the demand for meat and milk-based products is projected to rise by 68% and 57%, respectively, by 2030 compared to the year 2000 (FAO, 2006). Sustainably managing livestock excreta will be central to achieving an expansion in production with minimal environmental damage (Davidson, 2009; Chadwick et al., 2015). This literature review focuses on the urine patch of grazing animals as sites of accelerated nitrogen (N) cycling processes and as discrete sources of N losses from grazed pastures. The review will demonstrate that these sites can contribute to environmental, economic and human health issues as the reactive N cascades through different chemical forms in the biosphere, hydrosphere and atmosphere (Bolan et al., 2004; Sutton et al., 2011). The literature relating to urine patch research is largely dominated by cattle-based systems, therefore, this review aims to assess current understanding of N cycling and losses with specific reference to sheep urine patches, although comparisons to cattle based systems are made where necessary.

This review focusses initially on the reactive N cascade (the ability of reactive N to transition through several chemical forms, causing a series of negative effects on ecosystems and human health; Galloway et al. (2010)), as a necessary background to understanding the complexity of N cycling in the environment and from livestock production systems. Current knowledge of sheep urine composition and urine patch characteristics are then reviewed, followed by terrestrial N transformations and drivers of N losses. Techniques to measure nitrous oxide (N<sub>2</sub>O) emissions from urine patches are assessed and the use of nitrification inhibitors as a means to reduce N losses from the grazed pasture system are also considered in detail.

## 2.2 The reactive N cascade in the context of livestock production

Nitrogen limits primary productivity in most ecosystems (LaBauer and Treseder, 2008), despite being the most abundant element in the atmosphere (Pelletier and Tyedmers, 2010). This is because N mostly resides as the non-reactive  $N_2$  molecule, and the forces required to break the strong triple bond of the  $N_2$  molecule renders it unusable to most living organisms (Galloway et al., 2003). Reactive N has been defined by Galloway et al. (2003) as all inorganic reduced forms of N ( $NH_3$ ,  $NH_4^+$ ), inorganic oxidized forms of N ( $NO_x$ ,  $HNO_3$ ,  $N_2O$ ,  $NO_2^-$ ,  $NO_3^-$ ) and organic forms of N (e.g. urea, amino acids, proteins, nucleic acids, amines), which are essential components of the building blocks of life (Galloway and Cowling, 2002). Prior to human intervention, the amounts of reactive N in circulation was limited, due to biological N fixation by specialised microorganisms (e.g. free-living bacteria, blue-green algae and N-fixing bacteria associated with the roots of leguminous plants) being the dominant source of reactive N creation (Galloway and Cowling, 2002). In the nineteenth century the development of the Haber-Bosch process, an industrial process capable of producing reactive N for fertilizer, was fundamental for being able to produce enough food for a growing population (Galloway et al., 2010). However the concurrent rise in the use of fertilizers, cultivation of legumes and fossil fuel consumption, have contributed to a dramatic increase in reactive N creation (Galloway et al., 2004; Bouwman et al., 2009).

Livestock systems fit into the global N cycle (see Fig. 2.1) by converting reactive N stored in plant resources, which are of no immediate other use, into human-edible protein (Steinfeld and Gerber, 2010; Dijkstra et al., 2013). Grasslands cover 20-40% of the land surface of earth, depending on the definition used (Panunzi, 2008), and fodder produced to feed animals accounts for ca. 30% of all arable land (Steinfeld et al., 2006; Bouwman et al., 2013). Both plants and animals associated with livestock production are highly inefficient at incorporating applied N into their tissues. Grazing animals excrete a large proportion (70-95%) of the N they



- |   |   |   |   |
|---|---|---|---|
| A | Symbiotic and asymbiotic N fixation             | K | dissolved organic N ( $J_{\text{DON}}$ )  |
| B | Wet and dry depositions                         | L | Denitrification (biotic and abiotic)  |
| C | Mineral fertiliser inputs                       | M | Volatilisation  |
| D | Organic matter inputs                           | N | Anammox (anaerobic ammonium oxidation): $\text{NH}_4^+ + \text{NO}_2^- \rightarrow \text{N}_2 + 2 \text{H}_2\text{O}$ |
| E | Nitrification                                   | O | Ammonium fixation and desorption  |
| F | Mineralisation                                  | P | Dissimilatory nitrate reduction to ammonium (DNRA)  |
| G | Ammonification                                  | Q | Leaching of dissolved organic N   |
| H | Immobilisation                                  |   |   |
| I | Leaching of mineral N                           |   |   |
| J | Crop uptake of mineral ( $J_{\text{MIN}}$ ) and |   |   |

**Figure 2.1** Schematic of the N cycle. Source: Stark and Richards (2008).

consume (Oenema et al., 2005), mainly because the amount of N required for the optimal growth of vegetative material exceeds the amount of N that grazing animals require for protein synthesis (Ussiri and Lal, 2012). As a result, labile nutrients from the consumed, digested and excreted plant material are returned to soil, at a greater rate than the comparative degradation of plant litter (Harrison and Bardgett, 2008). The urine fraction is considered more labile than the faeces, where up to 60% (Haynes and Williams, 1993) of consumed N is excreted. In addition, there is estimated to be more N stored in managed manure produced by livestock, than there is in global N fertilizer use (Bouwman et al., 2009; Bouwman et al., 2013), highlighting the importance of livestock systems as contributors to elevated levels of reactive N in the environment. In the UK, the N input from excreta N and inorganic N are similar e.g. input from manure: 323 kt N year<sup>-1</sup>, N excretion on pasture range and paddock: 613 kt N year<sup>-1</sup> and N input from inorganic fertilizers applied to cropland and grassland: 1150 kt N year<sup>-1</sup> (Laura Cárdenas, personal communication, 2016).

The pathways of N removal processes (Fig. 2.1) from agricultural systems include N offtake in the form of crops/livestock products, volatilisation of NH<sub>3</sub>, losses of N to water by leaching of NO<sub>3</sub><sup>-</sup> and lateral flows (e.g. run off and erosion), denitrification and release of gaseous N compounds (NO, NO<sub>2</sub>, N<sub>2</sub>O, N<sub>2</sub>) and the anaerobic oxidation of NH<sub>4</sub><sup>+</sup> to N<sub>2</sub> (anammox). Some of these N removal processes are environmentally benign e.g. conversion of reactive N to N<sub>2</sub>, yet others have far-reaching consequences for human health and the environment. Atmospheric N species can be re-deposited to land through wet (e.g. by dissolving in water droplets) or dry deposition (e.g. deposition of particulate matter). Atmospheric NH<sub>3</sub> is present in the atmosphere in gaseous form, but can readily dissolve in atmospheric water or react with acidic gases or particles to form hygroscopic salts, atmospheric NH<sub>4</sub><sup>+</sup> is generally in particulate form (Mosier, 2001). Atmospheric NO<sub>3</sub><sup>-</sup> occurs as particulates



(Orel and Seinfeld, 1977), whereas  $\text{NO}_x$ ,  $\text{N}_2$  and  $\text{HNO}_3$  exist in gaseous form, and the latter can dissolve in water droplets (Hanke et al., 2003).

Gaseous losses of N which are of environmental concern include  $\text{N}_2\text{O}$  and nitric oxides ( $\text{NO}$  and  $\text{NO}_2$ ; collectively known as  $\text{NO}_x$ ).  $\text{N}_2\text{O}$  emissions are one of the major environmental implications associated with livestock production (Ambus et al., 2007; Kool et al., 2006), and the urine patches of grazing animals are recognised hot spots for  $\text{N}_2\text{O}$  emissions from grazed grasslands. In literature reviews conducted by van Groenigen (2005) and Selbie et al. (2015), the average fraction of applied ruminant urine N emitted as  $\text{N}_2\text{O}$  was found to be 1.7% ( $n = 31$ ) and 2.1% ( $n = 40$ ). These are similar to the default IPCC emission factor used for cattle excreta (2% of applied N) in Tier 1 inventories, and likely reflects the greater proportion of studies on cattle urine patches compared to sheep urine patches. The default IPCC emission factor for sheep urine is 1% of the applied urinary N, due to lower urine volumes and lower compaction in sheep grazed grassland (IPCC, 2006).  $\text{N}_2\text{O}$  is a potent greenhouse gas (Orwin et al., 2010), with an atmospheric lifetime of over 100 years and a global warming potential ca. 300 times that of carbon dioxide ( $\text{CO}_2$ ; Wrage et al., 2001). The increase in  $\text{N}_2\text{O}$  since the industrial period contributes an additional radiative forcing of  $0.16 \pm 0.02 \text{ W m}^{-2}$  and is primarily anthropogenic in origin (IPCC, 2007). Of the global total  $\text{N}_2\text{O}$  emissions from animal manure in 2000, 41% was attributed to grazing, where the highest emissions on a per animal basis were from non-dairy cattle, followed by sheep (Oenema et al., 2005). Indirect emissions of  $\text{N}_2\text{O}$  can also occur when N compounds are lost via volatilisation (followed by N deposition), runoff and leaching. The lost N can subsequently be transformed to  $\text{N}_2\text{O}$  in a different location (e.g. freshwater sediments, drainage ditches; Mosier et al., 1998; Oenema et al., 2005), as part of the reactive N cascade.

In addition to the global warming potential of  $\text{N}_2\text{O}$  residing in the troposphere, stratospheric depletion of ozone is another side effect.  $\text{N}_2\text{O}$  can react with atomic oxygen to

form nitric oxide (NO) in the stratosphere, inducing ozone destruction (Crutzen, 1970; Davidson et al, 2000; Wrage et al., 2001). This gas is currently the dominant ozone-depleting substance in the atmosphere, surpassing chlorofluorocarbons, which have seen a successful reduction in emissions following the Montreal Protocol (Ravishankara et al., 2009). A depletion of ozone in the stratosphere is associated with increased risk of skin cancer and damage to plants. In contrast to N<sub>2</sub>O causing the reduction of stratospheric ozone, NO<sub>x</sub> emissions result in the production of ozone in the troposphere (Davidson et al., 2000). Tropospheric ozone is a GHG, an air pollutant effecting human health (e.g. respiratory diseases), and it can damage vegetation (Aneja et al., 2001).

Nitrate leaching is the ‘vertical’ transport of NO<sub>3</sub><sup>-</sup> through soil by water, which can often reach surface and ground water. Increasing NO<sub>3</sub><sup>-</sup> levels in drinking water can result in methaemoglobinemia (Di and Cameron, 2002) or ‘blue-baby syndrome’, although the documented cases of this are extremely rare. This high ingestion of NO<sub>3</sub><sup>-</sup> impairs the ability of blood to transport oxygen around the body, where infants are particularly susceptible. It should also be noted, however, that NO<sub>3</sub><sup>-</sup> at lower concentrations can also have a beneficial impact on human health (Lundberg et al., 2004). In addition to its human impact, NO<sub>3</sub><sup>-</sup> leaching into aquatic environments can lead to eutrophication, algal blooms and death of fish (Di and Cameron, 2002). In a review of 22 studies, average losses of NO<sub>3</sub><sup>-</sup> from urine patches amounted to 20% of the applied urine N, although this varies with N loading rate and season of application (Selbie et al., 2015).

The volatilisation of NH<sub>3</sub> can result in aerosol formation, acid deposition and soil acidification (ApSimon et al., 1987; DEFRA, 2002). Deposition of NH<sub>4</sub><sup>+</sup>-N can also enrich N-poor habitats, resulting in eutrophication (DEFRA, 2002). In a recent review by Selbie et al. (2015), a mean and range of 12.9 (1-38)% of applied urine-N was volatilised as NH<sub>3</sub> from

ruminant urine patches ( $n = 59$ ). Sherlock and Goh (1984) measured an  $\text{NH}_3$  volatilisation loss of 12-38% of sheep urine N, when applied at a rate of  $500 \text{ kg N ha}^{-1}$ .

As can be seen from this section, reducing the losses of N from livestock production could yield multiple environmental benefits, the protection of ecosystem services (e.g. clean air and drinking water), greenhouse gas emission reductions and ozone layer protection, as well as conserving N that could fertilise pasture. Due to the cost of applying fertilizers or purchasing animal feed, improvements in N use efficiency can additionally provide an economic benefit for farmers, providing any improvement measures are cost-effective.

## **2.3 Sheep urine patch characteristics**

The simulation of urine events are often required when studying N losses from grazed pastures. This section aims to provide an overview of characteristics relating to sheep urine patches ranging from urine chemical composition and storage of collected urine, urine patch area, urine volume and frequency, and urine-to-soil application methods, in order to understand how best to re-create urine patches for experimental monitoring.

### *2.3.1 Urine chemical composition*

Natural ruminant urine composition can be highly variable due to individual animal's physiology, diet and water intake (Shand et al., 2002; Dijkstra et al., 2013). Sheep urine generally consists of  $5\text{-}16.9 \text{ g N l}^{-1}$  (Sherlock and Goh, 1984; Bristow et al., 1992), and in a recent meta-analysis conducted by Selbie et al. (2015), the average N concentration of sheep urine, from sheep fed a grass-based diet, was reported to be  $8.7 \text{ g N l}^{-1}$ . The predominant nitrogenous constituent of sheep urine is urea ( $\text{CO}(\text{NH}_2)_2$ ), which is formed in the liver as a means of detoxifying ammonia in systemic circulation (Spek et al., 2013). Urea comprises the largest fraction of urinary N at 80-90% of the total urinary N excreted (Sherlock and Goh,

1984; Dijkstra et al., 2013). Other nitrogenous constituents at smaller quantities in sheep urine include purine derivatives (allantoin, hippuric acid, uric acid, xanthine and hypoxanthine) creatine, creatinine, amino acids, and  $\text{NH}_3$  (Verbic et al., 1990; Bristow et al., 1992).

Purine derivatives arise from the breakdown product of purines (Verbic et al., 1990), as ruminant feed is generally low in purine content, the majority of purine derivatives in ruminant urine originates from rumen microorganisms (Chen and Gomes, 1992). The excretion of purine derivatives has been used as an index of microbial protein supply in ruminants (Chen et al., 1995). The purine bases adenine and guanine, are deaminated to xanthine and hypoxanthine, and mostly converted to uric acid, due to the activity of xanthine oxidase present in the intestine and liver (Tas and Susanbeth, 2007; Dijkstra et al., 2013). Uric acid is subsequently oxidized to allantoin in the liver, and excreted at ca. 4% of urinary-N. Hippuric acid is a conjugate molecule of benzoic acid and glycine, formed for the removal of benzoic acid consumed via aromatic and phenolic constituents within fodder (van Groenigen et al., 2005). In the sheep urine studied by Bristow et al. (1992), ca. 4% of urine N was comprised of hippuric acid. Xanthine, hypoxanthine and uric acid are generally excreted in urine in smaller quantities, where all compounds were present at < 1% of the total excreted N in Bristow et al. (1992). The metabolism of glycine, methionine and arginine produces creatine (ca. 4% of sheep urine N), which can be converted to the cyclic anhydride creatinine (< 1% of sheep urine N) for excretion (Bristow et al., 1992).

Bathurst (1952) investigated the amino acid distribution within the urine of cattle and sheep, where glycine was found to be the major amino acid present. In sheep and cattle urine, total glycine (including bound glycine within hippuric acid and peptides) represented 94% and 98% of the total amino-N respectively. The much lower concentration of other amino acids within the urine suggested that if the bound glycine was in peptide form, it would probably be composed of glycine residues. In contrast, Bristow et al. (1992) found glycine to be prevalent

in sheep urine, however, other amino acids were also found in similar quantities e.g.  $\beta$ -alanine and taurine. In cattle and goat urine, also investigated by Bristow et al. (1992), a dominance of glycine and taurine was observed. Expressed as a percentage of total N, the amino-N fraction has been found to comprise 9.3% (Bathurst, 1952) and 10.5-15.9% (Doak, 1952), however, Bristow et al. (1992) found a much lower value of 0.08-3.67%. As with the other urinary N constituents, the amino fraction also seems highly variable, which is likely to be due to differences in animal diet and physiological status.


The quantity of  $\text{NH}_4^+$  in sheep urine, expressed as a percentage of the total urine N, is generally less than 0.5% (Bristow et al., 1992). Nevertheless, urea can be rapidly hydrolysed via the enzyme urease, to yield  $\text{NH}_3$  and carbamate as follows (Mobley and Hausinger, 1989):



The carbamate then spontaneously hydrolyses forming carbonic acid and another  $\text{NH}_3$  molecule:



The carbonic acid then dissociates forming  $\text{H}^+$  and  $\text{HCO}_3^-$ , and the resulting  $\text{NH}_3$  equilibrates with water forming  $\text{NH}_4^+$  as displayed by the following equations (Mobley and Hausinger, 1989):



Initially, urine does not contain urease (Whitehead and Raistrick, 1993) but it can soon become present within urine samples, which can result in decreasing urea contents and increasing  $\text{NH}_4^+$  contents if left unfrozen or unacidified. This is because urease is ubiquitous in terrestrial and aquatic environments (Mobley and Hausinger, 1989) and the rate of hydrolysis of urea is dependent on factors which influence the urease-producing bacteria and the activity of the enzyme (Whitehead and Raistrick, 1993). Notably, temperature is an important factor, where higher temperatures would speed up urea hydrolysis due to more favourable conditions for bacterial multiplication and more optimal conditions for enzymatic activity. For these reasons, many authors describe rapidly returning to laboratories with urine samples and advocate freezing samples as soon as possible (Bristow et al., 1992; Whitehead and Raistrick, 1993), or analysing immediately (Shand et al., 2002). Freezing should halt population increase and/or activity of the urease producing microbes (Whitehead and Raistrick, 1993) and analysing immediately would restrict the time taken for microbes to proliferate.

Whitehead and Raistrick (1993) incubated urine from cows at 5, 10, 20 and 35 °C over a period of 21 days to gain information on the factors influencing urea hydrolysis in stored urine. With incubation temperatures over 5 °C, the sum of urea-N and ammoniacal-N were greater than the initial urea-N concentration, indicating that other N containing components were also converted to ammonium (Whitehead and Raistrick, 1993). Knowlton et al. (2010) investigated three collection methods for Holstein heifer urine (chilled, acidified or acidified 6 h after collection) fed either a high or low protein diet. It was found that all three methods performed similarly with regards to N retention, regardless of the protein content of diet (and therefore higher concentrations of urinary N). In order to study the effects of urine deposition on soil, urine should be collected and frozen immediately. Acidifying the urine would alter the pH and would, therefore, not be suitable for investigating transformations of urine compounds within the soil.

In addition to the major nitrogenous constituents of sheep urine, it is also comprised of aqueous salts. Haynes and Williams (1992) analysed the cations and anions within the urine of sheep which were fed either grass/clover pasture or supplements. In both diets,  $K^+$  and  $Na^+$  were the most prevalent cations in the urine, however, sheep on supplements had higher  $Na^+$  than  $K^+$  content and the sheep fed grass had higher  $K^+$  than  $Na^+$  content. This is likely to be due to the initial concentrations of these elements in the feed. Cations present at a lesser extent include  $Mg^{2+}$ ,  $Ca^{2+}$  and  $NH_4^+$ . The major anions found within the urine were  $HCO_3^-$  and  $Cl^-$ , with  $SO_4^{2-}$  and  $NO_3^-$  in smaller quantities (Haynes and Williams, 1992).

Synthetic urine has been used within experiments on several occasions (e.g. Shand et al., 2000; Shand et al., 2002; Kool et al., 2006; Lucas and Jones, 2006; Hoogendoorn et al., 2008), and is comprised of various quantities of the compounds present in natural urine. Advantages of using synthetic over natural urine include the fact that urine can be difficult to collect (especially if large volumes are required), while synthetic urine provides a convenient medium for experimental use (Shand et al., 2002). Natural urine can be highly variable in terms of N content and caution should be taken when interpreting results from experiments using synthetic urine and microcosm studies, especially if scaling up to pasture scale. Shand et al. (2002) noticed using synthetic sheep urine resulted in an overestimation of results in comparison to natural sheep urine, for example soil pH, dissolved organic C, total dissolved phosphate, dissolved organic N and  $NH_4^+$  and  $NO_3^-$  concentrations were all overestimated in this study. Similarly, Kool et al. (2006) compared urine from cattle with three synthetic versions (urea + glycine, urea + hippuric acid and artificial urine with the same nitrogenous composition as the real cow urine) in a soil incubation study. None of the artificial urine studies adequately represented the real urine across all measured variables ( $N_2O$  flux, pH, soil respiration and mineral N dynamics) and the use of real urine was advocated wherever possible,

although some artificial urine treatments had similar cumulative N<sub>2</sub>O emissions to real urine in the study by Bell et al. (2015).

Sheep urine composition can vary widely as a function of diet, water and physiological status of the animal. Therefore, wherever possible, real sheep urine should be collected for experimental purposes. Ideally the urine collected should reflect the grazing systems and management practices relevant to the area of study e.g. relevant sheep breeds, diets and applying at representative times of the year (to match diet). When collecting urine for field application, samples should be frozen as soon as possible to avoid losses of N. In the case of urine application to soil, urine should not be acidified, as this would change the pH of the urine, and may alter downstream N processing. Artificial urine can provide a useful medium where alterations in N composition are to be examined, but should be made as close in composition to real urine as possible. Further work is required to understand how urine composition influences downstream N<sub>2</sub>O producing processes.


### *2.3.2 Sheep urine patch area*

The urine patch is the area where certain nutrients are returned and recycled, or lost, from pastures and often show up in pastures as areas of grass which are greener than others. The area and volume of soil influenced by a urine patch are important parameters to consider when recreating urine patches in the field. The urine patch area has been considered in terms of both the wetted area and the effective area, and when used to model N leaching losses, the volume of soil influenced by urine has been considered as an inverted cone (Fig. 2.2; Li et al., 2012). The wetted area has been defined as the area of soil where urine has been directly voided, and the effective area has been defined as the area where plants can access the nutrient pool within the patch (Lantinga et al., 1987; Selbie et al., 2015). Solutes within the urine patch could move via mass flow (the movement of water and solutes together due to a pressure




gradient) or via diffusion (the movement of solutes due to a concentration gradient), where solute movement via diffusion can occur independently of water (Oyewole et al., 2013).

The effective area is generally larger than the wetted area due to horizontal and vertical diffusion of solutes, and extension of roots into the nutrient rich patch. If a soil has a large infiltration capacity then the wetted area would be expected to decrease, as the urine would tend to travel down the soil profile with minimal lateral spread (Li et al., 2012). What is less clear about this definition of the urine patch area, is the extent of horizontal diffusion of solutes in the urine patch over time and what impact this has on soil processes, N<sub>2</sub>O emissions and the measurement of such emissions from urine patches (this is investigated in Chapter 3).



**Figure 2.2** Diagrammatic representation of the urine patch area, where  $r_w$  represents the wetted area radius,  $r_p$  represents the pasture response area radius and  $d$  represents the depth of urine infiltration. Source: Li et al. (2012).

Williams and Haynes (1994) used a bromide recovery technique to determine the potential size and distribution of typical sheep (200 ml) and cattle (2000 ml) urination events applied by pouring onto soil, at relevant heights for each animal. As shown in figure 2.3, the depth is shallower and the surface area covered is smaller for the sheep compared to cattle urine, due to the lower volume applied.



**Figure 2.3** Distribution of  $\text{Br}^-$  from simulated sheep and cattle urine, using potassium bromide solution as a tracer. Source: Williams and Haynes (1994).

In this study, the wetted area of the sheep urine patch applied to a silt loam soil with a *Lolium perenne* L. dominated sward ranged from 0.043-0.055  $\text{m}^2$ , and the maximum soil depth reached was 15 cm, with the amount and concentration decreasing with depth. Doak (1952) also provided a value for sheep urine patch area of 300  $\text{cm}^2$  based on a 150 ml urine application to a . The urine patch area is likely to vary due to differences in urinary volume slope, soil type and condition (e.g. compaction) and soil moisture conditions, and differences relating to these factors are relatively understudied (Selbie et al., 2015).

In addition to the urine patch area, the total area of pasture covered by urine events has also been studied. Betteridge et al. (2010b) describe the use of a urine detecting sensor for use on female cattle or sheep, where urine flows over a thermistor, and any urination event is logged as the temperature changes. A GPS unit is also attached to the animal so the location of the urine patch can be recorded. From these data the distribution of urine can be mapped, providing useful information on the spatial distribution of urine patches at pasture scale. Betteridge et al.

(2010b) determined that cattle and sheep deposited ca. 41% and 30% of urine over 5% and 7.5% of the pasture area, respectively. Areas of more intense urine deposition were found in livestock ‘camping areas’. This supports the idea of targeting e.g.  $\text{NO}_3^-$  leaching or  $\text{N}_2\text{O}$  mitigation strategies to certain areas within the field, where urine frequency is highest and soil conditions are more conducive to  $\text{N}_2\text{O}$  production.

Betteridge et al. (2013) describe advances in such sensors, where urine volume, frequency and N concentration can all be measured. These are important parameters to consider as the concentration of N and the volume of each urine event causes urine patches of differing N loading rates. Consideration of the mean urine patch volume and frequency causes differences in modelled  $\text{NO}_3^-$  leaching losses of 5-10%, when compared to using mean values for urine concentration and volume (Li et al., 2012; Betteridge et al., 2013). A similar phenomenon may also occur when using average urine volumes and N concentrations for predicting  $\text{N}_2\text{O}$  emissions from grazed grasslands.

### *2.3.3 Sheep urine volume and frequency*

Urine volume is mainly influenced by the mineral load and water ingested by the animal (Selbie et al., 2015). Urine volume has been reported to be higher when herbage leaves are wet with rainwater or dew (Doak, 1952). Urine volume can also vary as a response to coping with changes in ambient temperatures (Betteridge et al., 2010a). Typically, individual sheep urine events range between 0.1-0.2 l (Doak, 1952; Haynes and Williams, 1993). Daily sheep urine volume ranged from 0.5-3.0 l in Ledgard et al. (2008), and sheep urine frequency has been reported to range between 13-20 times per day (Doak, 1952; Betteridge et al. 2010a; Betteridge et al., 2010b). More research is required on typical urine volumes and frequencies of sheep urine events, particularly in relation to different breeds, diets and management strategies. An assessment of the individual urine volumes from Welsh Mountain ewes is made in Chapter 7.

#### *2.3.4 Sheep urine-to-soil application methods*

The most common method for re-creating urine patches for experimental use is applying urine on an area basis within replicated plots (e.g. the equivalent of 4 l m<sup>-2</sup>; Kelliher et al., 2014). Urine has been applied to soil evenly using watering cans, where the height of urine deposition has even been considered in some studies e.g. Williams and Haynes (1994). Representing sheep urine patches by applying urine over a large area e.g. 1 m<sup>2</sup>, has advantages in that there is a large area of soil which can be utilised for soil sampling and instalment of static chambers for monitoring gas fluxes. Nevertheless, in reality, the size of sheep urine patches are much smaller (0.03-0.05 m<sup>2</sup>) and the size of urine patches has been shown to influence N losses (e.g. greater NO<sub>3</sub><sup>-</sup> leaching losses from cattle-sized compared to sheep-sized urine patches (McDowell and Houlbrooke, 2009), and lower N<sub>2</sub>O emissions due to a more even spread across the pasture deposited in smaller sheep-sized patches compared to larger cattle-sized patches (IPCC, 2006)). Many studies apply urine evenly within the chamber basal area or in lysimeter studies (Koops et al., 1997), yet this may confound measurements by the restriction of diffusion of solutes into surrounding soil and the prevention of root extension into the patch. Applying urine over a fixed area, however, is a sensible approach, as it allows the N loading rate to be calculated. It is recommended that the size of urine patches used in field studies should reflect, as closely as possible, the size of a representative urine patch. Other factors which may influence urine N processing is matching the temperature of urine application, although this has not been studied to our knowledge. It is suggested, that static chambers used for monitoring greenhouse gas emissions from urine patches should be larger than the size of the urine patch, to prevent chamber walls restricting diffusion of soluble N and C into surrounding soil (this is investigated in Chapter 3).

## **2.4 Changes in soil-microbe-plant properties following urine deposition**

### *2.4.1 Changes in soil properties*

This section describes the changes in soil properties that occur as a direct result of urine deposition. Shand et al. (2002) added synthetic sheep urine to soils under field conditions and in sward boxes. The applied urea was rapidly hydrolysed within the soil, following first order kinetics, with a half-life of 16.1 hours. The hydrolysis of urea within soil is generally a fast process, which can be stimulated by hippuric acid, which is also present in urine (Monaghan and Barraclough, 1992). The alkaline products generated during urea hydrolysis generally increase soil pH by up to 3 units (Carter, 2007; van Groenigen et al. 2005). This 1000-fold increase in  $\text{OH}^-$  is associated with increased organic matter solubility and desorption of anions held on exchange sites (Shand et al., 2002). Following nitrification or  $\text{NH}_3$  volatilisation, the pH decreases again (Clough et al., 2010). In the experiment conducted by Shand et al. (2002) the pH had returned to that of control soils by 56 days, however, the maximum pH occurred 2 days following urine addition in the sward boxes, but 7 days following urine addition to the field. The solubilisation of dissolved organic N (DON) and consequences for leaching of DON is also an understudied area for urine-influenced soils (van Kessel et al., 2008; Selbie et al., 2015)

As urea in urine hydrolyses, a large amount of  $\text{NH}_4^+$  is generated, increasing as the process progresses. If nitrification proceeds quickly, a sharp rise in concentrations of  $\text{NO}_3^-$  within the soil will occur (with the associated decrease in  $\text{NH}_4^+$ ). The rate of nitrification is important in determining the potential for loss via processes such as leaching and denitrification (Monaghan and Barraclough, 1992). This is due to the greater mobility of  $\text{NO}_3^-$  within soils in comparison to  $\text{NH}_4^+$ , therefore enhancing the probability of the N being leached. Generally, in grassland soils, nitrification proceeds only slowly relative to the rate of urea hydrolysis (Monaghan and Barraclough, 1992).

In addition to a rise in pH, the electrical conductivity (EC) of soil also increases following urine addition (Orwin et al., 2010). This is due to the addition of urinary salts, which lower the osmotic potential of the soil water. Increased levels of P have been demonstrated in soils following urine application. Generally ruminant urine does not contain high amounts of P, therefore, the increase must be soil based (Bertram et al., 2012). Bertram et al. (2012) displayed a decrease in total phospholipid-derived fatty acids (PLFA) count (a structural component of cellular membranes), combined with an increase in soluble P in soils treated with urine. The PLFA count is representative of the living microbial biomass, as upon cell death the PLFAs will rapidly degrade. The release of P-containing cell constituents following cell lysis was posed as a possible explanation for the observed soluble P increase and PLFA decrease. Another explanation for higher concentrations of available P is the increased organic matter solubility (i.e. DOP), following the associated pH increase after urine addition to soil.

Soil dissolved organic C (DOC) increases following urine addition, and subsequently declines back to original values over time (Orwin et al., 2010). Within the urine patch, possible sources of C include the C within the urine itself, increased solubilisation of the soil C due to urine addition, lysis of soil microbial cells due to stress induced changes following urine addition or from the leakage of C from scorched plant roots (Ambus et al., 2007). Carter et al. (2006) and Ambus et al. (2007) tested the hypothesis that urine deposition causes an increase in the release of root-derived C compounds, resulting in more denitrification and N<sub>2</sub>O emissions. Both studies concluded that urine did not cause a rapid increase in C release from roots, possibly due to urine-induced root death reducing respiration. Lambie et al. (2012) found that C solubilisation from soil cores amended with cow urine was ten times that solubilised with water alone. The evolution of CO<sub>2</sub> following urine addition to soils exceeded the amount initially present in the urine in studies by Clough et al. (2003) and Kool et al. (2006). This

indicated that priming of soil C release had occurred due to increased soil microbial activity (Kool et al., 2006).

#### *2.4.2 Urine patch and microorganisms*

Microorganisms control many of the processes involved with C and N cycling in soils, therefore the effect urine has on the microbial community is an important area of study. The microbial community are responsible for N<sub>2</sub>O producing and reducing pathways, and understanding how the individual processes relate to changes in environmental conditions and agricultural management practices is key for the development of mitigation strategies (Smith, 2010). Orwin et al. (2010) highlight the fact that our understanding of the biotic responses to urine addition is still limited. Features of the urine patch can be both stimulatory and inhibitory for microbial activity. Large inputs of N stimulate nitrifier and denitrifier activity (Orwin et al., 2010), and further factors such as the addition of water, labile C and other nutrients could have a stimulatory effect. Hostile conditions of the urine patch include a high pH, high osmotic stress and a build-up of toxic substances / metabolites (e.g. NO<sub>2</sub><sup>-</sup> and NH<sub>3</sub>; Petersen et al., 2004).

Orwin et al. (2010) conducted an experiment adding bovine urine to soil cores at two differing moisture treatments (35% and 75% water-filled pore space; WFPS). At 75% WFPS, the microbial biomass was often significantly lower than the dry treatment; however, rates of nitrification and denitrification were higher in the wet treatment. Nitrous oxide fluxes were also significantly higher in the 75 % WFPS treatment in comparison to 35 %. It was concluded that moisture content and aeration status had a strong effect on the way soil microbes processed the urine-N (Orwin et al., 2010). Lovell and Jarvis (1996) found the microbial biomass increased in size following urine application; however, this was not significantly different to the water control. Although microbial biomass size was similar, there was an increase in

activity (CO<sub>2</sub> production) and functioning of the soil microbial biomass after urine treatment (Lovell and Jarvis, 1996).

An increase in PLFA yields were observed in an experiment conducted by Petersen et al. (2004), where cattle urine was applied to confined field plots. This increase in the size of the microbial pool occurred over four days, with a shift in community composition. More anteiso fatty acids were produced, which belong to the family of branched-chain fatty acids (Kaneda, 1991). It was suggested that the presence of these fatty acids resulted in an increase in membrane permeability, which may be a stress response to hyperosmotic conditions. Bertram et al. (2012) found little evidence for a direct effect of increasing urinary salt concentrations on the stress of the soil microbial community, however, microbial stress was induced by the addition of the urine in association with a high moisture content. A decrease in PLFAs were observed between days 3 and 8 following bovine urine addition to soil cores in the study conducted by Bertram et al. (2012). This decline occurred at the same time as an increase in available P and was attributed to stress induced lysis of microbial cells, releasing P-containing constituents.

Di et al. (2010) investigated nitrification rates and growth of ammonia-oxidizing archaea and ammonia-oxidizing bacteria in differing soil depths, following urine amendment. The ammonia-oxidizing bacteria were more abundant and had a higher activity in comparison to the ammonia-oxidizing archaea, especially in the surface horizon. The rate of nitrification was attributed to the ammonia-oxidizing bacteria in these New Zealand pasture soils. This has consequences for the design of nitrification inhibition strategies, where this study supports targeted surface application. In this study, additions of urine also stimulated growth of ammonia oxidizing bacteria but not archaea.

The effect of urine deposition on fungal communities has been less well examined than other soil microorganisms. A shift in fungal community structure was observed by Rooney and



Clipson (2009) following the addition of synthetic sheep urine to soil microcosms. Shifts in fungal populations may be observed due to a variety of factors including changes in pH, nutrient availability, and changes in rhizodeposition (Rooney and Clipson, 2009). Jirout et al. (2013) isolated 36 fungal species from cattle overwintering areas, under laboratory conditions 64% of these species were able to produce N<sub>2</sub>O. The contribution of N<sub>2</sub>O via fungal species under field conditions is not well understood and requires further research.


#### 2.4.3 Urine and herbage

A reduction in root biomass has been reported following urine applications, for example the root biomass of *Agrostis capillaris* and *Lolium perenne* significantly decreased in a pot experiment examining the addition of synthetic sheep urine over time (Rooney and Clipson, 2009). Vegetation may be scorched (necrosis of tissues) following urine additions to soil, and botanical composition may be altered (Shand et al., 2002). Rooney and Clipson (2009) reported scorching becoming apparent 8-10 days following synthetic urine addition to soil microcosms. Ledgard et al. (1982) reported a decrease in clover growth and N fixation due to cow urine addition to pasture.

Herbage biomass production is generally found to increase following urine deposition e.g. Williams and Haynes (1994) reported an increase in herbage dry matter of 63-75% in urine treated plots compared to non-urine treated plots. The N and K content of herbage was found to increase following sheep urine application, due to the high concentrations of these elements in the initial urine samples (Haynes and Williams, 1992). The increased N supply within pastures can result in dark green coloured vegetation above urine patches. The response of vegetation will be larger than the initial wetted area following deposition, due to the roots and stolons of pasture plants which are able to access this pool of nutrients (Li et al., 2012), and due to some diffusion of NO<sub>3</sub><sup>-</sup>.

## 2.5 Terrestrial transformations of N leading to N<sub>2</sub>O formation

As the measurement and mitigation of N<sub>2</sub>O emissions from sheep urine patches is the major topic of this thesis, this section will discuss the terrestrial N transformations that may occur in urine-influenced soils. Current understanding of the microbial pathways leading to N<sub>2</sub>O production have been described in detail in Wrage et al. (2001), Baggs (2008) and Baggs and Philippot (2010). A schematic of the microbial processes leading to N<sub>2</sub>O can be seen in Figure 2.4, where ammonia oxidation, denitrification and nitrate ammonification are the microbial N<sub>2</sub>O-genic processes in soils. The majority of urine N enters the soil as urea and other small organic N molecules, which mineralize to NH<sub>4</sub><sup>+</sup> in the soil. Mineralisation is the process where soil organic N is converted to NH<sub>4</sub><sup>+</sup>/NH<sub>3</sub> and immobilisation occurs when NH<sub>4</sub><sup>+</sup>/NH<sub>3</sub> is converted back to organic N. These processes are likely to regulate the supply/removal of reactants for N<sub>2</sub>O-producing pathways, however, this section will focus on N transformations which directly lead to N<sub>2</sub>O production, therefore mineralisation and immobilisation will not be considered in detail here.



**Figure 2.4** The microbial pathways leading to N<sub>2</sub>O, with environmental conditions favourable for each process and enzymes regulating each pathway. Source: Baggs (2011).

### 2.5.1 Nitrification

Biological nitrification is the oxidation of  $\text{NH}_4^+$  or  $\text{NH}_3$  to  $\text{NO}_3^-$ , involving two intermediates, hydroxylamine ( $\text{NH}_2\text{OH}$ ) and nitrite ( $\text{NO}_2^-$ ). In Fig. 2.4, nitrification is represented by the combination of ammonia oxidation and nitrite oxidation. The first step in nitrification involves oxidation of  $\text{NH}_3$  to  $\text{NH}_2\text{OH}$ , which is carried out by primary nitrifiers and catalysed by the enzyme ammonia monooxygenase (AMO; Fig. 2.4; Wrage et al., 2001). Subsequently, the  $\text{NH}_2\text{OH}$  is oxidised to  $\text{NO}_2^-$ , which is catalysed by the enzyme hydroxylamine oxidoreductase (HAO; Fig. 2.4). Oxidation of  $\text{NO}_2^-$  to  $\text{NO}_3^-$  is carried out by the genera *Nitrobacter* (Haynes and Williams, 1993) and catalysed by nitrite oxidoreductase (Wrage et al., 2001). Autotrophic nitrifiers i.e. the process of ammonia oxidation, can be a significant source of NO and  $\text{N}_2\text{O}$  (Lipschultz et al., 1981) e.g. under anaerobic conditions  $\text{NO}_2^-$ , instead of oxygen, can be used as a terminal electron acceptor where  $\text{N}_2\text{O}$  and NO are produced (FAO, 2001).  $\text{N}_2\text{O}$  can also be formed via the chemical decomposition of the nitrification pathway intermediates ( $\text{NH}_2\text{OH}$  or  $\text{NO}_2^-$ ; Wrage et al., 2001).

Urine-N would undergo nitrification following urea hydrolysis, or the breakdown of other urinary N constituents into  $\text{NH}_3$  (e.g. amino acids, purine derivatives). Ammonia oxidation is carried out by Nitrobacteriaceae under aerobic conditions, where maximum activity is observed at 30-60% WFPS (FAO, 2001; Wrage et al., 2001). The known bacterial autotrophs that oxidise  $\text{NH}_3$  belong to the genera *Nitrosomonas* and *Nitrospira* (Smith, 2010). As inputs of N via urine are generally in forms other than  $\text{NO}_3^-$ , nitrification is an important step to provide the substrate for other microbial processes which utilise  $\text{NO}_3^-$  (Fig. 2.4; Yamulki et al., 1998). Nitrification was generally thought to be carried out by autotrophic bacteria (Pedersen et al., 1999), yet a range of heterotrophic bacterial and fungal nitrifiers are also known (Wrage et al., 2001; Smith et al., 2010). Heterotrophic nitrifiers can use inorganic

or organic substrates for nitrification (Barraclough and Puri, 1995) and are also able to denitrify under aerobic conditions (Robertson et al., 1989; Wrage et al., 2001).

The process of nitrifier denitrification is described as the reduction of  $\text{NO}_2^-$  to  $\text{N}_2\text{O}$  by ammonia oxidizers (Kool et al., 2010). Therefore,  $\text{N}_2\text{O}$  arising from ammonia oxidizers can occur during ammonia oxidation or denitrification by the ammonia oxidizers themselves. Baggs (2011) questions the inclusion of this process as a separate  $\text{N}_2\text{O}$  producing process, as it essentially involves the same enzymes used in denitrification. The ability of ammonia-oxidizing bacteria to denitrify appears widespread, therefore, a redefinition of the functioning of this microbial group may be required (Shaw et al., 2006; Baggs, 2011). Wrage et al. (2004) assessed the contribution of nitrifier denitrification in incubation studies under varying grassland conditions, including artificial urine patches. Nitrifier denitrification was found to be a potentially important pathway for  $\text{N}_2\text{O}$  production, as the greater the concentration of artificial urine-N applied the greater the concentration of  $\text{NO}_2^-$  in the soil. The increased  $\text{NO}_2^-$  in the soil would increase the substrate available for nitrifier denitrification. The environmental factors which influence this pathway are poorly understood (Smith, 2010), yet increased  $\text{NO}_2^-$  concentrations and mildly anaerobic conditions have been identified as potential rate regulating processes (Monaghan and Barraclough, 1993). The high pH and  $\text{NH}_3$  concentrations within urine patches slow down the oxidation of  $\text{NO}_2^-$  to  $\text{NO}_3^-$ , allowing accumulation of  $\text{NO}_2^-$  (Monaghan and Barraclough, 1993). It is therefore concluded that nitrifier denitrification is a potentially important source of  $\text{N}_2\text{O}$  in urine patches, but further work utilising natural urine and field conditions will be required to determine its importance under a variety of ecological conditions.

### 2.5.3 Biological denitrification

Denitrification involves the reduction of  $\text{NO}_3^-$ , to  $\text{NO}_2^-$ ,  $\text{NO}$ ,  $\text{N}_2\text{O}$  and  $\text{N}_2$  (Fig. 2.4). The process is mainly carried out by diverse genera of aerobic heterotrophic microorganisms, which are abundant in soils (Lloyd, 1993). The enzymes (and genes coding for these enzymes) which catalyse the denitrification pathway include nitrate reductase (NAR in Fig 2.4; *napA* and *narG* genes), nitrite reductase (NIR in Fig 2.4), nitric oxide reductase (NOR in Fig 2.4; *nirS* and *nirK* genes), and nitrous oxide reductase (NOS in Fig 2.4; *nosZ* gene; Hochstein and Tomlinson, 1988; Smith, 2010). The genes coding for these enzymes have been used in studies to determine the environmental factors which influence denitrifier community composition (Smith, 2010).

The facultative anaerobes involved in denitrification are able to utilise  $\text{NO}_3^-$  as a terminal electron acceptor under low  $\text{O}_2$  conditions (Wrage et al., 2001). Provided there is sufficient  $\text{NO}_3^-$  and readily available C, under low  $\text{O}_2$  conditions,  $\text{N}_2\text{O}$  can be released as an intermediate (Wrage et al., 2001). Smith (2010) identifies the main controlling influences on denitrifier activity as availability of C,  $\text{O}_2$  status of the soil (influenced by WFPS), availability of N, temperature and pH. The addition of urine to soil influences all of the factors which affect denitrifier activity. Lloyd (1993) highlights the fact that initially, denitrifiers require aerobic conditions for efficient biomass production, however, rapid denitrification occurs under low  $\text{O}_2$  conditions. This may be important upon consideration of grazing livestock urination events, which may increase anaerobic conditions in localised areas previously unaffected by low  $\text{O}_2$  levels. Denitrification has been found to be highly spatially variable, due to the variable nature of identified controlling factors of denitrification, including anaerobic soil microsites (FAO, 2001). The denitrification process occurs at particularly high rates in soil which has 50-90% WFPS (FAO, 2001).

The process of co-denitrification (Fig. 2.4) involves the utilisation of one molecule of N from a co-substrate (e.g.  $\text{NH}_3$ , amino acids, aniline, azide) combined with one molecule of

N from NO or N<sub>2</sub>O, potentially forming two molecules of N<sub>2</sub>O compared to one via the normal denitrification process (Su et al., 2004; Baggs, 2011). The significance of this pathway in terms of N<sub>2</sub>O emissions from soil is yet to be established.

#### 2.5.4 Nitrate ammonification

Nitrate ammonification (Fig. 2.4), also termed dissimilatory nitrate reduction to ammonia (DNRA), is the process where NO<sub>3</sub><sup>-</sup> is reduced to NO<sub>2</sub><sup>-</sup>, followed by NH<sub>4</sub><sup>+</sup> (Fig. 2.4; Smith, 2010). This process is carried out by fermentative bacteria and is less well characterised in soils than other N transformations. The major environmental drivers governing its occurrence in soils include the redox status, C/NO<sub>3</sub><sup>-</sup> ratio, pH, bulk density, sand content and NO<sub>2</sub><sup>-</sup> concentration (Wan et al., 2009; Rütting et al., 2011). The process is thought to conserve N, as NO<sub>3</sub><sup>-</sup> is converted to the less mobile NH<sub>4</sub><sup>+</sup>. N<sub>2</sub>O can also form during this anaerobic process, which may occur concurrently with denitrification (Stevens et al., 1998). Nitrate ammonification is favoured under high pH levels, which would occur initially under a urine patch scenario. Following urea hydrolysis and nitrification in a urine patch, NO<sub>3</sub><sup>-</sup> would not be limiting, therefore, competition for C would determine whether denitrification or nitrate ammonification would occur (Tiedje, 1988).

#### 2.5.5 Chemodenitrification

Chemodenitrification is attributed to any non-biological source of NO, N<sub>2</sub>O or N<sub>2</sub> (Smith, 2010), therefore, is not displayed in Fig. 2.4. This process is generally favoured under high NO<sub>2</sub><sup>-</sup> concentrations and a pH of < 5.5 (Haynes and Williams, 1993). An alkaline pH is usually present initially within a urine patch, therefore, chemodenitrification reactions are expected to be of minor importance for N<sub>2</sub>O emissions under this scenario, unless dealing with very acidic soils.

## 2.6 Measurement of N<sub>2</sub>O emissions from urine patches

The major challenges associated with measuring N<sub>2</sub>O includes the low atmospheric concentration of the gas (320 ppb), and the episodic nature of N<sub>2</sub>O emissions, with large flux variations observed both spatially and temporally (Rapson and Dacres, 2014). This section explores methodological, analytical and modelling approaches to quantifying N<sub>2</sub>O emissions from soils. The static chamber technique, combined with analysis of headspace gas samples via gas chromatography (GC) equipped with an electron capture detector (ECD) is one of the most widely used, versatile and low-cost technologies used for N<sub>2</sub>O flux measurements (Hutchinson and Mosier, 1980; de Klein and Harvey, 2012), which has been in use for over 40 years (Hensen et al., 2013), resulting in a wide range of largely comparable data sets. This technique was utilised in the experimental chapters of this thesis, via both manual and automated campaigns.

The optimisation and need for standardisation of the static chamber technique for monitoring N<sub>2</sub>O fluxes has been recognised (de Klein and Harvey, 2012), where important considerations when using this technique include chamber design, deployment, sampling frequency and time of day to sample. The calculation of fluxes can either be conducted by assuming linearity in the increase of chamber headspace concentrations, or by using non-linear or best-fit model approaches (Kroon et al., 2008; Pederson et al., 2010; Hensen et al., 2013). Prioritising increased chamber numbers to account for spatial variability in N<sub>2</sub>O fluxes, rather than prioritising increased number of samples per chamber closure period has been suggested by Chadwick et al. (2014), due to the far greater uncertainty introduced from spatial variability in fluxes, compared with either reducing the number of headspace samples or the assumption of linearity of headspace concentrations. The final strategy used, however, is likely to be a compromise between the necessary requirements of the experiment, and time and resources available for specific projects. Automated chamber systems reduce manual labour and provide

enhanced temporal resolution, although chamber numbers and spatial area coverage are generally limited.

In order to improve and develop current or new mitigation strategies for N<sub>2</sub>O emissions from soils, quantifying the microbial sources of N<sub>2</sub>O production will be necessary (Köster et al., 2013). Previous techniques which have been utilised to apportion N<sub>2</sub>O production to different sources (e.g. nitrification or denitrification) include the use of process-specific inhibitors such as C<sub>2</sub>H<sub>2</sub>, or isotopic labelling techniques and natural abundance techniques ( $\delta^{15}\text{N}$  and  $\delta^{18}\text{O}$ ; Baggs et al., 2008; Decock and Six, 2013; Wolf et al., 2015). Novel approaches to source partition N<sub>2</sub>O emissions from soils include the measurement of N<sub>2</sub>O isotopomers i.e. the distribution of <sup>15</sup>N in N<sub>2</sub>O as either <sup>14</sup>N<sup>15</sup>N<sup>16</sup>O (<sup>15</sup>N<sub>α</sub>) or <sup>15</sup>N<sup>14</sup>N<sup>16</sup>O (<sup>15</sup>N<sub>β</sub>; Toyoda and Yoshida, 1999; Decock and Six, 2013). The site preference (SP =  $\delta^{15}\text{N}_\alpha - \delta^{15}\text{N}_\beta$ ) of N<sub>2</sub>O has been shown to differ during differing N<sub>2</sub>O production pathways, therefore, this technique is gaining interest as a tool for identifying the source of N<sub>2</sub>O from soils. Iotopomers of N<sub>2</sub>O have been analysed using Fourier transform infrared mass spectroscopy (FTIR-MS) and, more recently, quantum cascade laser based absorption spectroscopy (QCLAS; Waechter et al., 2008; Wolf et al., 2015). The latter is more applicable for continuous measurements under field conditions and has been found to have a higher precision for SP values in comparison to FTIR-MS (Köster et al., 2013).

Tower-based micrometeorological techniques for measuring N<sub>2</sub>O emissions are conducted over larger areas (>10 m<sup>2</sup> to field and regional scale) and, therefore, avoid some of the issues related to spatial heterogeneity when utilising the chamber technique (Pattey et al., 2007; Phillips et al., 2007; Hensen et al., 2013). The range of micrometeorological techniques is beyond the scope of this review, however, it is extensively reviewed in Skinner and Wagner-Riddle (2012). Advances in optical techniques for measuring N<sub>2</sub>O (e.g. QCLAS) are providing excellent performance, sensitivity and reliability for tower-based N<sub>2</sub>O flux measurements,



however, such instrumentation is much more costly than a GC-based system (Hensen et al., 2013). Modelling techniques have also been used to estimate N<sub>2</sub>O emissions and NO<sub>3</sub><sup>-</sup> leaching from grazed pastures (Saggar et al., 2007; Giltrap et al., 2010; Cichota et al., 2013). Process based models are important for estimating N<sub>2</sub>O emissions from soils under differing land use types and future climate change scenarios, where micrometeorological techniques are useful for validating process based models (Eugster and Zeemun, 2006; Pattey et al., 2007).

The Intergovernmental Panel on Climate Change (IPCC) provide a three tier methodological approach to estimating N<sub>2</sub>O emissions, for national inventory purposes. The Tier 1 approach uses a default emission factor (1% of applied N for sheep and 2% for cattle) in combination with activity data (e.g. number of grazing animals; IPCC, 2006). This is a simplistic approach, which does not consider confounding factors such as climatic conditions, soil type or differences in management (IPCC, 2006). The Tier 2 approach uses country specific data and the Tier 3 approach is based on modelling. Estimates of N<sub>2</sub>O emissions from varying sectors are important in determining where mitigation options should be focused, however, individual source estimates of N<sub>2</sub>O emissions are still considered to have a high uncertainty (Ambus et al., 2007; IPCC, 2007). The accuracy of such estimates may decrease if C and N interactions are not adequately represented under changing climatic conditions (Gärdenäs et al. 2011). Increasing regional field-scale measurements of direct and indirect N<sub>2</sub>O emissions will improve the accuracy of country specific emission estimates. In order to provide more accurate emission data, a move to the Tier 2 and Tier 3 approach is warranted.

## **2.7 Nitrification inhibitors as tools to mitigate sheep urine patch N<sub>2</sub>O emissions**

Several differing mitigation strategies have been proposed for reducing N losses from grazed pastures, which have been reviewed extensively in Eckard et al. (2010) and Luo et al. (2010). In this Thesis, the use of nitrification inhibitors were selected for study, as in principal,

they appear to be an attractive strategy to reduce both  $\text{NO}_3^-$  leaching and  $\text{N}_2\text{O}$  emissions from urine patches or fertilizer applications, whilst also potentially providing biomass yield increases (Di and Cameron, 2007; Abalos et al., 2014). Increasing the residence time of  $\text{NH}_4^+$  increases opportunity for plant acquisition, immobilization and adsorption (Di and Cameron 2007), thus reducing the  $\text{NO}_3^-$  supply available for leaching and denitrification processes. If more of the added N remains as  $\text{NH}_4^+$  plant use efficiency can increase and losses via leaching and denitrification may be reduced (Cookson and Cornforth, 2002). Such additional benefits may become important in terms of reducing the amount of N lost as  $\text{N}_2\text{O}$  from agricultural soils (e.g. 1% of that applied in sheep excreta), as this does not represent a significant economic loss to farmers. However, combined with the reduction of other N losses (e.g.  $\text{NO}_3^-$  leaching), nitrification inhibitors become a more attractive strategy for improving agricultural N use efficiency. In addition, nitrification inhibitors may reduce  $\text{N}_2\text{O}$  emissions from two biological  $\text{N}_2\text{O}$  production pathways, from the process of nitrification itself, and by reducing the amount of substrate available for denitrification.

The first and rate-limiting step of nitrification, the oxidation of  $\text{NH}_4^+$  to  $\text{NO}_2^-$  (Zerulla et al., 2001; Moir et al., 2007), is inhibited by a wide variety of compounds of biological (e.g. root exudates; Subbarao et al., 2007; Zakir et al., 2008; Subbarao et al., 2009) and synthetic origin (e.g. dicyandiamide (DCD), 3,4-dimethylpyrazole-phosphate (DMPP), several heterocyclic N compounds; McCarty and Bremner, 1989; Zerulla et al., 2001). Ammonia monooxygenase has a broad range of substrates which it can catalytically oxidize, therefore, many nitrification inhibitors work on the basis of competing for the active site of this enzyme (McCarty, 1999). Although several compounds which inhibit nitrification have been identified, this thesis focuses on two of the most widely used nitrification inhibitors in agriculture, DCD and DMPP (Liu et al. 2013). DCD blocks the electron transport chain in the cytochrome of ammonia monooxygenase (AMO), whereas DMPP binds indiscriminately to the membrane-

bound AMO (Chaves et al. 2006; Fiencke and Bocke 2006; Benckiser et al. 2013). No significant toxicological effect on soil bacterial communities have been found for DCD applied to soils (O'Callaghan et al., 2010; Wakelin et al., 2013), and any effects have been shown to be minimal and transient (Morales et al., 2015). DMPP has also been the subject of extensive ecotoxicological tests, required for European Law, and is not recognized as a hazardous substance (Zerulla et al., 2001). Studies indicate that both DCD and DMPP effect ammonia-oxidizing bacteria, but have a limited effect of ammonia-oxidizing archaea, suggesting that ammonia oxidation could still take place from this consortia of soil microorganisms, following nitrification inhibitor addition to soils (O'Callaghan et al., 2010; Di et al., 2010; Kleineidam et al., 2011).

The nitrification inhibitor DCD has been researched extensively for use within livestock urine patches, and has even been used in commercial pastures in New Zealand (O'Callaghan et al., 2010; Liu et al., 2013). Different application methods of DCD have been studied for use in urine patches, typically DCD is sprayed directly onto pasture soil (Moir et al., 2007), however, oral administration to cattle (O'Connor et al. 2013; Welten et al. 2013), infusion of DCD into the rumen or abomasum of sheep (Ledgard et al. 2008), incorporation into fertiliser granules and addition of DCD within a biodegradable hydrogel to slow its release in soil (Minet et al. 2013) have also been investigated. The blanket approach to DCD application is generally inefficient, as the majority of N within older urine patches would have already been converted to  $\text{NO}_3^-$  (Zaman and Nguyen, 2012). Targeting urine patches within space and time e.g. infusing DCD into the gastrointestinal tract of sheep (Ledgard et al., 2008), or applications to livestock camping areas, would allow enhanced efficiency of nitrification inhibitors and would reduce the overall cost in comparison to broadcast applications (Luo et al., 2015).

Typical DCD application rates are  $10 \text{ kg DCD ha}^{-1}$  (Moir et al., 2007; O'Connor et al., 2012a; O'Connor et al., 2012b), although higher application rates (up to  $60 \text{ kg DCD ha}^{-1}$ ) have

been suggested for increased efficacy (Luo et al., 2015). In intensive sheep winter grazing systems in New Zealand, DCD (10 kg ha<sup>-1</sup>) was found to reduce N<sub>2</sub>O emissions and NO<sub>3</sub><sup>-</sup> leaching from late autumn applied urine patches (300 kg N ha<sup>-1</sup>) by 72% and 70%, respectively (Moir et al., 2010). In hill-grazed pastures, DCD reduced sheep urine patch N<sub>2</sub>O emissions by 40-80% (Hoogendoorn et al., 2008). No significant effect of DCD was found for NO<sub>3</sub><sup>-</sup> leaching from sheep urine patches applied to a winter forage crop, although the amount leached was generally three times lower than that from cattle urine patches (McDowell and Houlbrooke, 2009). Moir et al. (2007) demonstrated that a fine particle suspension of DCD, applied at a rate of 10 kg ha<sup>-1</sup>, was effective at reducing NO<sub>3</sub><sup>-</sup> levels from cattle urine patches (1000 kg N ha<sup>-1</sup>) from 40-80 kg N ha<sup>-1</sup> to 10 kg N ha<sup>-1</sup>. Forage dry matter production was also found to increase by at least 20% annually in the DCD treatments, over the course of the field trial. The results of O'Connor et al. (2012a; 2012b), however, suggest that applying DCD (10 kg ha<sup>-1</sup>) does not consistently increase herbage production in Irish grassland systems.

DMPP has been less well studied for use within urine patches, in comparison to DCD. This is likely to be due to the development of this product as a fertilizer coating, marketed as ENTEC (developed by BASF; Pasda et al., 2001; Zerulla et al., 2001). Typically, DMPP has been applied at application rates ten times lower than that of DCD, at 0.5-1 kg DMPP ha<sup>-1</sup>, which may result in a lower cost of application for farmers. It has also been demonstrated to have a lower phytotoxicity in comparison to DCD (Zerulla et al., 2001). Di and Cameron (2011) found a significant reduction in the ammonium oxidation rate constant in cattle urine-influenced (1000 kg N ha<sup>-1</sup>) pastures in New Zealand, following a liquid application of DMPP. In a lysimeter study conducted by Di and Cameron (2012), a liquid DMPP application (5 kg DMPP ha<sup>-1</sup>) was found to be just as effective as DCD (10 kg DCD ha<sup>-1</sup>), at reducing N<sub>2</sub>O emissions (by 62-66%) and NO<sub>3</sub><sup>-</sup> leaching from a 1000 kg N ha<sup>-1</sup> cattle urine application. Misselbrook et al. (2014) and Barneze et al. (2015), however, found that pyrazole derivatives

(similar in structure to DMPP) did not significantly reduce N<sub>2</sub>O emissions from cattle urine, applied under UK summer conditions.

The fate of nitrification inhibitors applied to soil is important, as this will regulate the amount of active compound available in the soil, and the length of time they are effective in reducing N losses (O'Connor et al., 2012a). The effect of temperature and moisture on the persistence of both nitrification inhibitors is fairly well described. DCD is most effective when applied at temperatures of 10°C or less (Kelliher et al., 2008) and a decrease in efficiency of DCD may also be observed under high rainfall, where the compound may be leached below the grassland rooting zone (O'Connor et al., 2012a). DMPP performance has also been found to be best under cold and wet conditions (Menéndez et al., 2012). Differences in efficacy between these two nitrification inhibitors has been attributed to a lower mobility of DMPP in comparison to DCD, due to a greater sorption of DMPP (Wissemeier et al., 2001; Zerulla et al., 2001; Di and Cameron, 2012). The potential for spatial separation of the nitrification inhibitor from NH<sub>4</sub><sup>+</sup> is another suggestion for explaining differences in their efficacy (Ruser and Schulz, 2015). However, few studies directly compare these processes for DCD and DMPP, especially for urine-influenced soil. In Chapter 5, a comparative assessment of the mobility and co-location of DCD and DMPP with urine-derived NH<sub>4</sub><sup>+</sup> is made, in a range of soils of contrasting texture and organic matter. The factors which may influence persistence of the compounds within the soil are also compared, e.g. sorption, microbial uptake and microbial degradation.

Despite the promising reductions in N<sub>2</sub>O emissions and NO<sub>3</sub><sup>-</sup> leaching observed by application of DCD to urine patches, in 2012, the use of DCD was suspended for sale and use in New Zealand (Ministry for Primary Industries, 2013). This was due to the finding of milk products contaminated with DCD, when nitrification inhibitors were applied to commercial dairy pastures (Kim et al., 2012; Chen et al., 2014). There is limited information, however,

regarding the mechanism of DCD entry into milk products. In Chapter 6, the potential for direct plant acquisition of this nitrification inhibitor is assessed.

Of the limited assessments of DMPP, and similar compounds, applied to ruminant urine patches, its effectiveness appears inconclusive. Further research is required to determine the efficacy of DMPP under differing edaphic and environmental conditions. Whether DMPP is effective at reducing N<sub>2</sub>O emissions from sheep urine patches has also not been studied. In Chapter 7, the effectiveness of DMPP applied to sheep urine patches under UK summer conditions is assessed, varying the time-since DMPP application.

## **2.8 Conclusions**

Urine patches in grazed grasslands can be sites of significant N loss (as N<sub>2</sub>O, NO<sub>x</sub>, NO<sub>3</sub><sup>-</sup> and NH<sub>3</sub>), resulting in the contribution to detrimental environmental, human health and economic impacts. Reducing agricultural inputs of reactive N may pose a difficult challenge under increasing intensity, yet it could ultimately lead to a wide range of benefits spanning resource efficiency and human and ecological health. Understanding the spatial and temporal dynamics of C and N interactions in the urine patch could feed into modelling systems, enabling accurate quantification of N loss pathways, under a range of environmental parameters. The best mitigation options are likely to take into account the spatial and temporal variability of N<sub>2</sub>O emissions from urine patches, therefore, understanding these processes are important for the development of optimal mitigation strategies.

Further work is required to understand how urine patch parameters can influence N<sub>2</sub>O production e.g. determining the importance of including urine patch diffusional areas within chambers (addressed in Chapter 3) and determining how urine patch size and N concentration influence N<sub>2</sub>O emissions, within typical ranges reported for sheep (addressed in Chapter 4). The efficacy of nitrification inhibitors in reducing N<sub>2</sub>O emissions has been found to vary

widely. Further research is required to determine factors influencing nitrification inhibitor removal/degradation processes in contrasting soils (addressed in Chapters 5 and 6) and to determine whether DMPP is effective at reducing N<sub>2</sub>O emissions from sheep urine patches under summer conditions in the UK (addressed in Chapter 7).

## 2.9 References

- Abalos, D., Jefferey, S., Sanz-Cobena, A., Guardia, G., Vallejo, A. 2014. Meta-analysis of the effect of urease and nitrification inhibitors on crop productivity and nitrogen use efficiency. *Agriculture, Ecosystems and Environment* 189, 136-144.
- Ambus, P., Petersen, S.O., Soussana, J.F. 2007. Short-term carbon and nitrogen cycling in urine patches assessed by combined carbon-13 and nitrogen-15 labelling. *Agriculture, Ecosystems and Environment* 121, 84-92.
- Aneja, V.P., Roelle, P.A., Murray, G.C., Southerland, J., Erisman, J.W., Fowler, D., Asman, W.A.H., Patni, N. 2001. Atmospheric nitrogen compounds II: emissions, transport, transformation, deposition and assessment. *Atmospheric Environment* 35, 1903-1911.
- ApSimon, H.M., Kruse, M., Bell, J.N.B. 1987. Ammonia emissions and their role in acid deposition. *Atmospheric Environment*, 21, 1939-1946.
- Baggs, E.M. 2008. A review of stable isotope techniques for N<sub>2</sub>O source partitioning in soils: recent progress, remaining challenges and future considerations. *Rapid Communications in Mass Spectrometry* 22, 1664-1672.
- Baggs, E.M. 2011. Soil microbial sources of nitrous oxide: recent advances in knowledge, emerging challenges and future direction. *Current Opinion in Environmental Sustainability* 3, 321-327.
- Baggs, E.M., Philippot, L. 2010. Microbial terrestrial pathways to N<sub>2</sub>O. In Ed: Smith, K.A. *Nitrous Oxide and Climate Change*, Earthscan Publications.
- Barneze, A.S., Minet, E.P., Cerri, C.C., Misselbrook, T. 2015. The effect of nitrification inhibitors on nitrous oxide emissions from cattle urine depositions to grassland under summer conditions in the UK. *Chemosphere* 119, 122-129.



- Barraclough, D., Puri, G. 1995. The use of  $^{15}\text{N}$  pool dilution and enrichment to separate the heterotrophic and autotrophic pathways of nitrification. *Soil Biology & Biochemistry* 27, 17-22.
- Bathurst, N.O. 1952. The amino acids of sheep and cow urine. *Journal of Agricultural Science* 42, 476-478.
- Bell, M.J., Rees, R.M., Cloy, J.M., Topp, C.F.E., Bagnall, A., Chadwick, D.R. 2015. Nitrous oxide emissions from cattle excreta applied to a Scottish grassland: Effects of soil and climatic conditions and a nitrification inhibitor. *Science of the Total Environment* 508, 343-353.
- Benckiser, G., Christ, E., Herbert, T., Weiske, A., Blome, J., Hardt, M. 2013. The nitrification inhibitor 3,4-dimethylpyrazole-phosphate (DMPP) – quantification and effects on soil metabolism. *Plant and Soil* 371, 257-266.
- Bertram, J.E., Orwin, K.H., Clough, T.J., Condon, L.M., Sherlock, R.R., O’Callaghan, M. 2012. Effect of soil moisture and bovine urine on microbial stress. *Pedobiologia* 55, 211-218.
- Betteridge, K., Costall, D., Balladur, S., Upsdell, M., Umemura, K. 2010a. Urine distribution and grazing behaviour of female sheep and cattle grazing a sheep New Zealand hill pasture. *Animal Production Science* 50, 624-629.
- Betteridge, K., Costall, D.A., Li, F.Y., Luo, D., Ganesh, S. 2013. Why we need to know what and where cows are urinating – a urine sensor to improve nitrogen models. *Proceedings of the New Zealand Grassland Association* 75, 119-124.
- Betteridge, K., Hoogendoorn, C., Costall, D., Carter, M. & Griffiths, W. 2010b. Sensors for detecting and logging spatial distribution of urine patches of grazing female sheep and cattle. *Computers and Electronics in Agriculture* 73, 66-73.

- Bolan, N.S., Saggar, S., Luo, J., Bhandral, R., Singh, J. 2004. Gaseous emissions of nitrogen from grazed pastures: processes, measurements and modelling, environmental implications and mitigation. *Advances in Agronomy* 84, 37-120.
- Bouwman, A.F., Beusen, A.H.W., Billen, G. 2009. Human alteration of the global nitrogen and phosphorus soil balances for the period 1970-2005. *Global Biogeochemical Cycles* 23, 1-16.
- Bouwman, L., Goldewijk, K.K., Van Der Hoek, K.W., Beusen, A.H.W., VanVuuren, D.P., Willems, J., Rufino, M.C., Stehfest, E. 2013. Exploring global changes in nitrogen and phosphorus cycles in agriculture induced by livestock production over the 1900-2050 period. *Proceedings of the National Academy of Sciences* 110, 20882-20887.
- Bristow, A.W., Whitehead, D.C., Cockburn, J.E. 1992. Nitrogenous constituents in the urine of cattle, sheep and goats. *Journal of the Science of Food and Agriculture* 59, 387-394.
- Carter, M.S. 2007. Contribution of nitrification and denitrification to N<sub>2</sub>O emissions from urine patches. *Soil Biology and Biochemistry* 39, 2091-2102.
- Carter, M.S., Klumpp, K., Le Roux, X. 2006. Lack of increased availability of root derived C may explain the low N<sub>2</sub>O emission from low N-urine patches. *Nutrient Cycling in Agroecosystems* 75, 91-100.
- Chadwick, D.R., Cardenas, L., Misselbrook, T.H., Smith, K.A., Rees, R.M., Watson, C.J., McGeough, K.L., Williams, J.R., Cloy, J.M., Thorman, R.E., Dhanoa, M.S. 2014. Optimizing chamber methods for measuring nitrous oxide emissions from plot-based agricultural experiments. *European Journal of Soil Science* 65, 295-307.
- Chadwick, D., Jia, W., Tong, Y., Yu, G., Shen, Q., Chen, Q. 2015. Improving manure nutrient management towards sustainable agricultural intensification in China. *Agriculture, Ecosystems and Environment* 209, 34-46.

- Chaves, B., Opoku, A., De Neve, S., Boeckx, P., Van Cleemput, O., Hofman, G. 2006. Influence of DCD and DMPP on soil N dynamics after incorporation of vegetable crop residues. *Biology and Fertility of Soils* 43, 62-68.
- Chen, X.B., Gomes, M.J. 1992. Estimation of microbial protein supply to sheep and cattle based on urinary excretion of purine derivatives – an overview of the technical details. International Feed Resources Unit, Rowett Research Institute, Aberdeen, Occasional publication.
- Chen, X.B., Mejia, A.T., Kyle, D.J., Ørskov, E.R. 1995. Evaluation of the use of the purine derivative: creatinine ratio in spot urine and plasma samples as an index of microbial protein supply to ruminants: studies in sheep. *Journal of Agricultural Science* 125, 137-143.
- Chen, X-H., Zhou, L-X., Zhao, Y-G., Pan, S-D., Jin, M-C., 2014. Application of nanoring amino-functionalized magnetic polymer dispersive micro-solid-phase extraction and ultra fast liquid chromatography-tandem mass spectrometry in dicyandiamide residue analysis of powdered milk. *Talanta* 119, 187-192.
- Cichota, R., Snow, V.O., Vogeler, I. 2013. Modelling nitrogen leaching from overlapping urine patches. *Environmental Modelling and Software* 41, 15-26.
- Clough, T.J., Bertram, J.E., Ray, J.L., Condon, L.M., O’Callaghan, M., Sherlock, R.R., Wells, N.S. 2010. Unweathered wood biochar impact on nitrous oxide emissions from a bovine-urine-amended pasture soil. *Soil Science Society of America* 74, 852-860.
- Clough, T.J., Sherlock, R.R., Kelliher, F.M. 2003. Can liming mitigate N<sub>2</sub>O fluxes from urine-amended soils? *Australian Journal of Soil Research* 41, 439-457.
- Cookson, W.R., Cornforth, I.S. 2002. Dicyandiamide slows nitrification in dairy cattle urine patches: effects on soil solution composition, soil pH and pasture yield. *Soil Biology and Biochemistry* 34, 1461-1465.

- Crutzen, P.J. 1970. The influence of nitrogen oxide on the atmospheric ozone content. *Quarterly Journal of the Royal Meteorological Society* 96, 320-325.
- Davidson, E.A. 2009. The contribution of manure and fertilizer nitrogen to atmospheric nitrous oxide since 1860. *Nature Geoscience* 2, 659-662.
- Davidson, E.A., Keller, M., Erickson, H.E., Verchot, L.V., Veldkamp, E. 2000. Testing a conceptual model of soil emissions of nitrous and nitric oxides. *BioScience* 50, 667-680.
- Decock, C., Six, J. 2013. How reliable is the intramolecular distribution of  $^{15}\text{N}$  in  $\text{N}_2\text{O}$  to source partition  $\text{N}_2\text{O}$  emitted from soil? *Soil Biology and Biochemistry* 65, 114-127.
- De Klein, C.A.M., Harvey, M.J. 2012. Nitrous Oxide Chamber Methodology Guidelines. Global Research Alliance on Agricultural Greenhouse Gases, Ministry for Primary Industries, New Zealand.
- Di, H.J., Cameron, K.C. 2002. Nitrate leaching in temperate agroecosystems: sources, factors and mitigating strategies. *Nutrient Cycling in Agroecosystems* 46, 237-256.
- Di, H.J., Cameron, K.C. 2005. Reducing environmental impacts of agriculture by using a fine particle suspension nitrification inhibitor to decrease nitrate leaching from grazed pastures. *Agriculture, Ecosystems and Environment* 109, 202-212.
- Di, H.J., Cameron, K.C. 2007. Nitrate leaching losses and pasture yields as affected by different rates of animal urine nitrogen returns and application of a nitrification inhibitor – a lysimeter study. *Nutrient Cycling in Agroecosystems* 79, 281-290.
- Di, H.J., Cameron, K.C., 2011. Inhibition of ammonium oxidation by a liquid formulation of 3,4-dimethylpyrazole phosphate (DMPP) compared with a dicyandiamide (DCD) solution in six new Zealand grazed grassland soils. *Journal of Soils and Sediments* 11, 1032-1039.

- Di, H.J., Cameron, K.C. 2012. How does the application of different nitrification inhibitors affect nitrous oxide emissions and nitrate leaching from cow urine in grazed pastures? *Soil Use and Management* 28, 54-61.
- Di, H.J., Cameron, K.C., Shen, J-P., Winefield, C.S., O'Callaghan, M., Bowatte, S., He, J-Z. 2010. Ammonia-oxidizing bacteria and archaea grow under contrasting soil nitrogen conditions. *FEMS Microbiology Ecology* 72, 386-394.
- DEFRA (2002). Ammonia in the UK. <http://archive.defra.gov.uk/environment/quality/air/airquality/publications/ammonia/documents/ammonia-in-uk.pdf> Accessed 11.12.12.
- Dijkstra, J., Oenema, O., van Groenigen, J.W., Spek, J.W., van Vuuren, A.M., Bannink, A. 2013. Diet effects on urine composition of cattle and N<sub>2</sub>O emissions. *Animal* 7, 292-302.
- Doak, B.W. 1952. Some chemical changes in the nitrogenous constituents of urine when voided on pasture. *The Journal of Agricultural Science* 42, 162-171.
- Eckard, R.J., Grainger, C., de Klein, C.A.M. 2010. Options for the abatement of methane and nitrous oxide from ruminant production: a review. *Livestock Science* 130, 47-56.
- Eugster, W., Zeeman, M. 2006. Micrometeorological techniques to measure ecosystem-scale greenhouse gas fluxes for model validation and improvement. *International Congress Series* 1293, 66-75.
- Fiencke, C., Bock, E. 2006. Immunocytochemical localization of membrane-bound ammonia monooxygenase in cells of ammonia oxidizing bacteria. *Archives of Microbiology* 185, 99-106.
- Food and Agriculture Organization (FAO). 2001. Global estimates of gaseous emissions of NH<sub>3</sub>, NO and N<sub>2</sub>O from agricultural land. <ftp://ftp.fao.org/agl/agll/docs/globest.pdf> Accessed 14.05.12.

- Food and Agriculture Organisation (FAO). 2006. World Agriculture: Towards 2030/2050 – Interim Report. Food and Agriculture Organisation, Rome.
- Galloway, J.N., Aber, J.D., Erisman, J.W., Setzinger, S.P., Howarth, R.W., Cowling, E.B., Cosby, J.B. 2003. The nitrogen cascade. *BioScience* 53, 341-356.
- Galloway, J.N., Cowling, E.B. 2002. Reactive nitrogen and the world: 200 years of change. *AMBIO: A Journal of the Human Environment* 31, 64-71.
- Galloway, J., Dentener, F., Burke, M., Durmont, E., Bouwman, A.F., Kohn, R.A., Mooney, H.A., Seitzinger, S., Kroeze, C. 2010. The impact of animal production systems on the nitrogen cycle. In: Steinfeld, H., Mooney, H.A., Schneider, F., Neville, L.E. (Eds.) *Livestock in a Changing Landscape* (pp 83-96). Island Press, Washington DC, USA.
- Galloway, J.N., Dentener, F.J., Capone, D.G., Boyer, E.W., Howarth, R.W., Seitzinger, S.P., Asner, G.P., Cleveland, C.C., Green, P.A., Holland, E.A., Karl, D.M., Michaels, A.F., Porter, J.F., Townsend, A.R., Vörösmarty, C.J. 2004. Nitrogen cycles: past, present, and future. *Biogeochemistry* 70, 153-226.
- Giltrap, D.L., Singh, J., Saggar, S., Zaman, M. 2010. A preliminary study to model the effects of a nitrification inhibitor on nitrous oxide emissions from urine-amended pasture. *Agriculture, Ecosystems and Environment* 136, 310-317.
- Hanke, M., Umann, B., Uecker, L., Arnold, F., Bunz, H. 2003. Atmospheric measurements of gas-phase HNO<sub>3</sub> and SO<sub>2</sub> using chemical ionization mass spectrometry during the MINATROC field campaign 2000 on Monte Cimone. *Atmospheric Chemistry and Physics* 3, 417-436.
- Harrison, K. A., Bardgett, R. D. 2008. Impacts of grazing and browsing by large herbivores on soils and soil biological properties. In: *The ecology of browsing and grazing* (pp. 201-216). Springer Berlin Heidelberg.

- Haynes, R.J., Williams, P.H. 1992. Changes in soil solution composition and pH in urine-affected areas of pasture. *Journal of Soil Science* 43, 323-334.
- Haynes, R.J., Williams, P.H. 1993. Nutrient cycling and soil fertility in the grazed pasture ecosystem. *Advances in Agronomy* 49, 119-199.
- Hensen, A., Skiba, U., Famulari, D. 2013. Low cost and state of the art methods to measure nitrous oxide emissions. *Environmental Research Letters* 8, 025022.
- Hochstein, L.I., Tomlinson, G.A. 1988. The enzymes associated with denitrification. *Annual Reviews of Microbiology* 42, 231-261.
- Hoogendoorn, C.J., de Klein, C.A.M., Rutherford, A.J., Letica, S., Devantier, B.P. 2008. The effect of increasing rates of nitrogen fertiliser and a nitrification inhibitor on nitrous oxide emissions from urine patches on sheep grazed hill country pasture. *Animal Production Science* 48, 147-151.
- Hutchinson, G.L., Mosier, A.R. 1980. Improved soil cover method for field measurement of nitrous oxide fluxes. *Soil Science Society of America Journal* 45, 311-316.
- IPCC. 2007. Fourth Assessment Report: Climate Change 2007. TS.2.1.1 Changes in Atmospheric Carbon Dioxide, Methane and Nitrous Oxide. [http://www.ipcc.ch/publications\\_and\\_data/ar4/wg1/en/tssts-2-1-1.html](http://www.ipcc.ch/publications_and_data/ar4/wg1/en/tssts-2-1-1.html) Accessed 05/11/12.
- Jirout, J., Šimek, M., Elhottová, D. 2013. Fungal contribution to nitrous oxide emissions from cattle impacted soils. *Chemosphere* 90, 565-572.
- Kaneda, T. 1991. Iso and anteiso-fatty acids in bacteria: biosynthesis, function, and taxonomic significance. *Microbiology and Molecular Biology Reviews* 55, 288-302.
- Kelliher, F.M., Cox, N., van der Weerden, T.J., de Klein, C.A.M., Luo, J., Cameron, K.C., Di, H.J., Giltrap, D., Rys, G. 2014. Statistical analysis of nitrous oxide emission factors

- from pastoral agriculture field trials conducted in New Zealand. *Environmental Pollution* 186, 63-66.
- Kim, D-G., Giltrap, D., Saggar, S., Palmada, T., Berben, P., Drysdale, D. 2012. Fate of the nitrification inhibitor dicyandiamide (DCD) sprayed on a grazed pasture: effect of rate and time of application. *Soil Research* 50, 337-347.
- Kleineidam, K., Košmrlj, K., Kublik, S., Palmer, I., Pfab, H., Ruser, R., Fiedler, S., Schloter, M. 2011. Influence of the nitrification inhibitor 3,4-dimethylpyrazole phosphate (DMPP) on ammonia-oxidizing bacteria and archaea in rhizosphere and bulk soil. *Chemosphere* 84, 182-186.
- Knowlton, K.F., McGilliard, M.L., Zhao, Z., Hall, K.G., Mims, W., Hanigan, M.D. 2010. Effective nitrogen preservation during urine collection from Holstein heifers fed diets with high or low protein content. *Journal of Dairy Science* 93, 323-329.
- Kool, D.M., Hoffland, E., Hummelink, E.W.J., van Groenigen, J.W. 2006. Increased hippuric acid content of urine can reduce soil N<sub>2</sub>O fluxes. *Soil Biology and Biochemistry* 38, 1021-1027.
- Kool, D.M., Wrage, N., Zechmeister-Boltenstern, S., Pfeffer, M., Brus, D., Oenema, O. & van-Groenigen, J-W. 2010. Nitrifier denitrification can be a source of N<sub>2</sub>O from soil: a revised approach to the dual-isotope labelling method. *European Journal of Soil Science* 61, 759-772.
- Koops, J.G., Van Beusichem, Oenema, O. 1997. Nitrous oxide production its source and distribution in urine patches on grassland on peat soil. *Plant and Soil* 19, 57-65.
- Köster, J.R., Well, R., Tuzson, B., Bol, R., Dittert, K., Giesemann, A., Emmenegger, L., Manninen, A., Cárdenas, L., Mohn, J. 2013. Novel laser spectroscopic techniques for continuous analysis of N<sub>2</sub>O isotopomers – application and intercomparison with



- isotope ratio mass spectrometry. *Rapid Communications in Mass Spectrometry* 27, 216-222.
- Kroon, P.S., Hensen, A., van dem Bulk, W.C.M., Jongejan, P.A.C., Vermeulen, A.T. 2008. The importance of reducing the systematic error due to non-linearity in N<sub>2</sub>O flux measurements by static chambers. *Nutrient Cycling in Agroecosystems* 82, 175-186.
- LaBauer, D.S., Treseder, K.K. 2008. Nitrogen limitation of net primary productivity in terrestrial ecosystems is globally distributed. *Ecology* 89, 371-379.
- Lambie, S.M., Schipper, L.A., Balks, M.R., Baisden, W.T. 2012. Solubilisation of soil C following treatment with cow urine under laboratory conditions. *Soil Research* 50, 50-57.
- Lantinga, E.A., Keuning, J.A., Groenwold, J., Deenen, P.J.A.G. 1987. Distribution of excreted nitrogen by grazing cattle and its effects on sward quality, herbage production and utilization. In *Animal Manure on Grassland and Fodder Crops. Fertilizer or Waste?* (pp. 103-117). Springer Netherlands.
- Ledgard, S.F., Steele, K.W., Saunders, W.H.M. 1982. Effects of cow urine and its major constituents on pasture properties. *New Zealand Journal of Agricultural Research* 25, 61-68.
- Li, F.Y., Betteridge, K., Cichota, R., Hoogendoorn, C.J., Jolly, B.H. 2012. Effects of nitrogen load variation in animal urination events on nitrogen leaching from grazed pasture. *Agriculture Ecosystems and Environment* 159, 81-89.
- Lipschultz, F., Zafiriou, O.C., Wofsy, S.C., Mcelroy, M.B., Valois, F.W., Watson, S.W. 1981. Production of NO and N<sub>2</sub>O by soil nitrifying bacteria. *Nature* 294, 641-643.
- Liu, C., Wang, K., Zheng, X. 2013. Effects of nitrification inhibitors (DCD and DMPP) on nitrous oxide emission, crop yield and nitrogen uptake in a wheat maize cropping system. *Biogeosciences* 10, 2427-2437.

- Lloyd, D. 1993. Aerobic denitrification in soils and sediments: from fallacies to facts. *Trends in Ecology and Evolution* 8, 352-356.
- Lovell, R. D., Jarvis, S.C. 1996. Effects of urine on soil microbial biomass, methanogenesis, nitrification and denitrification in grassland soils. *Plant and Soil* 186, 265-273.
- Lucas, S.D., Jones, D.L. 2006. Biodegradation of estrone and 17  $\beta$ -estradiol in grassland soils amended with animal wastes. *Soil Biology and Biochemistry* 38, 2803-2815.
- Lundberg, J.O., Weitzberg, E., Cole, J.A., Benjamin, N. 2004. Nitrate, bacteria and human health. *Nature Reviews Microbiology* 2, 593-602.
- Luo, J., de Klein, C.A.M., Ledgard, S.F., Saggart, S. 2010. Management options to reduce nitrous oxide emissions from intensively grazed pastures: a review. *Agriculture, Ecosystems and Environment* 136, 282-291.
- Luo, J., Ledgard, S., Wise, B., Welten, B., Lindsey, S., Judge, A., Sprosen, M. 2015. Effect of dicyandiamide (DCD) delivery method, application rate, and season on pasture urine patch nitrous oxide emissions. *Biology and Fertility of Soils* 51, 453-464.
- McCarty, G.W. 1999. Modes of action of nitrification inhibitors. *Biology and Fertility of Soils* 29, 1-9.
- McCarty, G.W., Bremner, J.M. 1989. Inhibition of nitrification in soil by heterocyclic nitrogen compounds. *Biology and Fertility of Soils* 8, 204-211.
- McDowell, R.W., Houlbrooke, D.J. 2009. The effect of DCD on nitrate leaching losses from a winter forage crop receiving applications of sheep or cattle urine. *Proceedings of the New Zealand Grassland Association* 17, 117-120.
- Menéndez, S., Barrena, I., Setien, I., González-Murua, C., Estavillo, J.M. 2012. Efficiency of nitrification inhibitor DMPP to reduce nitrous oxide emissions under different temperature and moisture conditions. *Soil Biology and Biochemistry* 53, 82-89.

- Minet, E.P., O'Carroll, C., Rooney, D., Breslin, C., McCarthy, C.P., Gallagher, L., Richards, K.G. 2013. Slow delivery of a nitrification inhibitor (dicyandiamide) to soil using a biodegradable hydrogel of chitosan. *Chemosphere* 93, 2854-2858.
- Ministry for Primary Industries (2013) [WWW document] URL <http://www.mpi.govt.nz/Portals/0/Documents/news-resources/news/letter-of-assurance.pdf> Accessed 01/09/14.
- Misselbrook, T.H., Cardenas, L.M., Camp, V., Thorman, R.E., Williams, J.R., Rollett, A.J., Chambers, B.J. 2014. An assessment of nitrification inhibitors to reduce nitrous oxide emissions from UK agriculture. *Environmental Research Letters* 9, 11506.
- Mobley, H.L.T., Hausinger, R.P. 1989. Microbial ureases: significance, regulation and molecular characterization. *Microbiological Reviews* 53, 85-108.
- Moir, J.L., Cameron, K.C., Di, H.J. 2007. Effects of the nitrification inhibitor dicyandiamide on soil mineral N, pasture yield, nutrient uptake and pasture quality in a grazed pasture system. *Soil Use and Management* 23, 111-120.
- Moir, J.L., Wild, M.A., Cameron, K.C., Di, H.J. 2010. The effect of DCD on nitrogen losses from sheep urine patches applied to lysimeters in autumn. *Proceedings of the New Zealand Grassland Association* 72, 197-202.
- Monaghan, R.M., Barraclough, D. 1992. Some chemical and physical factors affecting the rate and dynamics of nitrification in urine-affected soils. *Plant and Soil* 143, 11-18.
- Monaghan, R.M., Barraclough, D. 1993. Nitrous oxide and dinitrogen emissions from urine-affected soil under controlled conditions. *Plant and Soil* 151, 127-138.
- Morales, S.E., Jha, N., Saggari, S. 2015. Impact of urine and the application of the nitrification inhibitor DCD on microbial communities in dairy-grazed pasture soils. *Soil Biology and Biochemistry* 88, 344-353.

- Mosier, A.R. 2001. Exchange of gaseous nitrogen compounds between agricultural systems and the atmosphere. *Plant and Soil* 228, 17-27.
- Mosier, A., Kroeze, C., Nevison, C., Oenema, O., Seitzinger, S., van Cleemput, O. 1998. Closing the global N<sub>2</sub>O budget: nitrous oxide emissions through the agricultural nitrogen cycle. *Nutrient Cycling in Agroecosystems*, 52, 225-248.
- O'Callaghan, M., Gerard, E.M., Carter, P.E., Lardner, R., Sarathchandra, U., Burch, G., Ghani, A., Bell, N. 2010. Effects of the nitrification inhibitor dicyandiamide (DCD) on microbial communities in a pasture soil amended with bovine urine. *Soil Biology and Biochemistry* 42, 1425-1436.
- O'Connor, P.J., Hennessy, D., Brophy, C., O'Donovan, M., Lynch, M.B. 2012a. The effect of the nitrification inhibitor dicyandiamide (DCD) on herbage production when applied at different times and rates in the autumn and winter. *Agriculture, Ecosystems and Environment* 152, 79-89.
- O'Connor, P.J., Hennessy, D., Lynch, M.B., Slettery, H., Lewis, E. 2013. The effect of dicyandiamide on rumen and blood metabolites, diet digestibility and urinary excretion. *Livestock Science* 155, 35-37.
- O'Connor, P.J., Lynch, M.B., Cahalan, E., O'Donovan, M., Hennessy, D. 2012b. The effect of the nitrification inhibitor dicyandiamide (DCD) on spring and annual herbage production in urine patches when applied in late summer or early autumn. *Grass and Forage Science*, doi: 10.1111/gfs.12011.
- Oenema, O., Wrage, N., Velthof, G.L., van Groenigen, J.W., Dolfing, J., Kuikman, P.J. 2005. Trends in global nitrous oxide emissions from animal production systems. *Nitrogen Cycling in Agroecosystems* 72, 51-65.
- Orel, A.E., Seinfeld, J.H. 1977. Nitrate formation in atmospheric aerosols. *Environmental Science and Technology* 11, 1000-1007.

- Orwin, K.H., Bertram, J.E., Clough, T.J., Condon, L.M., Sherlock, R.R., O'Callaghan, M., Ray, J., Baird, D.B. 2010. Impact of bovine urine deposition on soil microbial activity, biomass, and community structure. *Applied Soil Ecology* 44, 89-100.
- Oyewole, O.A., Inselsbacher, E., Näsholm, T. 2014. Direct estimation of mass flow and diffusion of nitrogen compounds in solution and soil. *New Phytologist* 201, 1056-1064.
- Panunzi, E. 2008. Are grasslands under threat? Brief analysis of FAO statistical data on pasture and fodder crops. UN Food and Agriculture Organization, Rome.
- Pasda, G., Hähndel, R., Zerulla, W. 2001. Effect of fertilizers with the new nitrification inhibitor DMPP (3,4-dimethylpyrazole phosphate) on yield and quality of agricultural and horticultural crops. *Biology and Fertility of Soils* 34, 85-97.
- Pattey, E., Edwards, G.C., Desjardins, R.L., Pennock, D.J., Smith, W., Grant, B., MacPherson, J.I. 2007. Tools for quantifying N<sub>2</sub>O emissions from agroecosystems. *Agricultural and Forest Meteorology* 142, 103-119.
- Pedersen, A.R., Petersen, S.O., Schelde, K. 2010. A comprehensive approach to soil atmosphere trace-gas flux estimation with static chambers. *European Journal of Soil Science* 61, 888-902.
- Pedersen, H., Dunkin, K.A., Firestone, M.K. 1999. The relative importance of autotrophic and heterotrophic nitrification in a conifer forest soil as measured by <sup>15</sup>N tracer and pool dilution techniques. *Biogeochemistry* 44, 135-150.
- Petersen, S.O., Roslev, P., Bol, R. 2004. Dynamics of a pasture soil microbial community after deposition of cattle urine amended with [<sup>13</sup>C]urea. *Applied and Environmental Microbiology* 70, 6363-6369.
- Pelletier, N., Tyedmers, P. 2010. Forecasting potential global environmental costs of livestock production 2000-2050. *Proceedings of the National Academy of Sciences* 107, 18371-18374.

- Phillips, F.A., Leuning, R., Baigent, R., Kelly, K.B., Denmead, O.T. 2007. Nitrous oxide flux measurements from an intensively managed irrigated pasture using micrometeorological techniques. *Agricultural and Forest Meteorology* 143, 92-105.
- Rapson, T.D., Dacres, H. 2014. Analytical techniques for measuring nitrous oxide. *TrAC Trends in Analytical Chemistry* 54, 65-74.
- Ravishankara, A.R., Daniel, J.S., Portmann, R.W. 2009. Nitrous oxide (N<sub>2</sub>O): the dominant ozone-depleting substance emitted in the 21<sup>st</sup> century. *Science* 326, 123-125.
- Robertson, L.A., Cornelisse, R., De Vos, P., Hadjioetomo, R., Kuenen, J.G. 1989. Aerobic denitrification in various heterotrophic nitrifiers. *Antonie Van Leeuwenhoek* 56, 289-299.
- Rooney, D.C., Clipson, N.J.W. 2009. Synthetic sheep urine alters fungal community structure in an upland grassland soil. *Fungal Ecology* 2, 36-43.
- Rütting, T., Boeckx, P., Müller, C., Klemmedtsson, L. 2011. Assessment of the importance of dissimilatory nitrate reduction to ammonium for the terrestrial nitrogen cycle. *Biogeosciences* 8, 1779-1791.
- Saggar, S., Hedley, C.B., Giltrap, D.L., Lambie, S.M. 2007. Measured and modelled estimates of nitrous oxide emission and methane consumption from a sheep-grazed pasture. *Agriculture, Ecosystems and Environment* 122, 357-365.
- Selbie, D.R., Buckthought, L.E., Shepherd, M.A. 2015. The challenge of the urine patch for managing nitrogen cycling in grazed pasture systems. *Advances in Agronomy* 129, 229-292.
- Shand, C.A., Williams, B.L., Smith, S., Young, M.E. 2000. Temporal changes in C, P and N concentrations in soil solution following application of synthetic sheep urine to soil under grass. *Plant and Soil* 222, 1-13.

- Shand, C.A., Williams, B.L., Dawson, L.A., Smith, S., Young, M.E. 2002. Sheep urine affects soil solution nutrient composition and roots: differences between field and sward box soils and the effects of synthetic and natural sheep urine. *Soil Biology and Biochemistry* 34, 163-171.
- Shaw, L.J., Nicol, G.W., Smith, Z., Fear, J., Prosser, J.I., Baggs, E.M. 2006. *Nitrosospira* spp. can produce nitrous oxide *via* a nitrifier denitrification pathway. *Environmental Microbiology* 8, 214-222.
- Sherlock, R.R., Goh, K.M. 1984. Dynamics of ammonia volatilization from simulated urine patches and aqueous urea applied to pasture I. Field experiments. *Nutrient Cycling in Agroecosystems* 5, 181-195.
- Skinner, H.R., Wagner-Riddle, C. 2012. Micrometeorological methods for assessing greenhouse gas fluxes. *Managing Agricultural Greenhouse Gases: Coordinated Agricultural Research through GRACEnet to Address our Changing Climate*, pp 367-383.
- Smith, K. 2010. *Nitrous oxide and Climate Change*. London: Earthscan.
- Spek, J.W., Dijkstra, J., Van Duinkerken, G., Bannink, A. 2013. A review of factors influencing milk urea concentration and its relationship with urinary urea excretion in lactating dairy cattle. *Journal of Agricultural Science* 151, 407-423.
- Stark, C.H., Richards, K.G. 2008. The continuing challenge of agricultural nitrogen loss to the environment in the context of global change and advancing research. *Dynamic Soil, Dynamic Plant* 2, 1-12.
- Steinfeld, H., Gerber, P. 2010. Livestock production and the global environment: Consume less or produce better? *Proceedings of the National Academy of Sciences* 107, 18237-18238.

- Steinfeld, H., Gerber, P., Wassenaar, T., Castel, V., Rosales, M., De Haan, C. 2006. *Livestock's Long Shadow*. Rome: FAO, 2006.
- Stevens, R.J., Laughlin, R.J., Malone, J.P. 1998. Soil pH affects the processes reducing nitrate to nitrous oxide and di-nitrogen. *Soil Biology and Biochemistry* 30, 1119-1126.
- Su, F., Takaya, N., Shoun, H. 2004. Nitrous oxide-forming codenitrification catalyzed by cytochrome P450nor. *Bioscience, Biotechnology and Biochemistry* 68, 473-475.
- Subbarao, G.V., Nakahara, K., Hurtado, M.P., Ono, H., Moreta, D.E., Salcedo, A.F., Yoshihashi, A.T., Ishikawa, T., Ishitani, M., Ohnishi-Kameyama, M., Yoshida, M., Rondon, M., Rao, I.M., Lascano, C.E., Berry, W.L., Ito, O. 2009. Evidence for biological nitrification inhibition in Brachiaria pastures. *Proceedings of the National Academy of Sciences* 106, 17302-17307.
- Subbarao, G.V., Rondon, M., Ito, O., Ishikawa, T., Rao, I.M., Nakahara, K., Lascano, C., Berry, W.L. 2007. Biological nitrification inhibition (BNI) – is it a widespread phenomenon? *Plant and Soil* 294, 5-18.
- Sutton, M.A., Oenema, O., Erisman, J.W., Leip, A., van Grinsven, H., Winiwarter, W. 2011. Too much of a good thing. *Nature* 472, 159-161.
- Tas, B.M., Susanbeth, A. 2007. Urinary purine derivatives excretion as an indicator of in vivo microbial N flow in cattle: a review. *Livestock Science* 111, 181-192.
- Tiedje, J. M. 1988. Ecology of denitrification and dissimilatory nitrate reduction to ammonium. In *Biology of Anaerobic Microorganisms*, Ed: A. J. B. Zehnder, 179–244. Wiley, New York.
- Toyoda, S., Yoshida, N. 1999. Determination of nitrogen isotopomers of nitrous oxide on a modified isotope ratio mass spectrometer. *Analytical Chemistry* 71, 4711-4718.



- Ussiri, D., Lal, R. 2012. Soil emission of nitrous oxide and its mitigation. Springer Science & Business Media.
- van Groenigen, J.W., Kuikman, P.J., de Groot, W.J.M., Velthof, G.L. 2005. Nitrous oxide emissions from urine-treated soil as influenced by urine composition and soil physical conditions. *Soil Biology and Biochemistry* 37, 463-473.
- van Kessel, C., Clough, T., van Groenigen, J.W. 2008. Dissolved organic nitrogen: an overlooked pathway of nitrogen loss from agricultural systems? *Journal of Environmental Quality* 38, 393-401.
- Verbic, J., Chen, X.B., MacLeod, N.A., Ørskov, E.R. 1990. Excretion of purine derivatives by ruminants. Effect of microbial nucleic acid infusion on purine derivative excretion by steers. *The Journal of Agricultural Science* 114, 243-248.
- Waechter, H., Mohn, J., Tuzson, B., Emmenegger, L., Sigrist, M.W. 2008. Determination of N<sub>2</sub>O isotopomers with quantum cascade laser based absorption spectroscopy. *Optics Express* 16, 9239-9244.
- Wakelin, S.A., Clough, T.J., Gerard, E.M., O'Callaghan, M. 2013. Impact of short-interval repeat applications of dicyandiamide on soil N transformation in urine patches. *Agriculture, Ecosystems and Environment* 167, 60-70.
- Wan, Y.J., Ju, X.T., Ingwersen, J., Schwarz, U., Stange, C.F., Zhang, F.S., Streck, T. 2009. Gross nitrogen transformations and related nitrous oxide emissions in an intensively used calcareous soil. *Soil Science Society of America Journal* 73, 102-112.
- Wolf, B., Merbold, L., Decock, C., Tuzson, B., Harris, E., Six, J., Emmenegger, L., Mohn, J. 2015. First on-line isotopic characterization of N<sub>2</sub>O above intensively managed grassland. *Biogeosciences* 12, 2517-2531.

- Welten, B.G., Ledgard, S.F., Schipper, L.A., Judge, A.A. 2013. Effect of amending cattle urine with dicyandiamide on soil nitrogen dynamics and leaching of urinary-nitrogen. *Agriculture, Ecosystems and Environment* 167, 12-22.
- Whitehead, D.C., Raistrick, N. 1993. Nitrogen in the excreta of dairy cattle: changes during short-term storage. *Journal of Agricultural Science* 121, 73-81.
- Williams, P.H., Haynes, R.J. 1994. Comparison of initial wetting pattern, nutrient concentrations in soil solution and the fate of <sup>15</sup>N-labelled urine in sheep and cattle urine patch areas of pasture soil. *Plant and Soil* 162, 49-59.
- Wissemeier, A.H., Linzmeier, W., Gutser, R., Weigelt, W., Schmidhalter, U. 2001. The new nitrification inhibitor DMPP (ENTEC®) – comparisons with DCD in model studies and field applications. Food security and sustainability. In: *Plant nutrition* (Horst WJ et al. eds.). Kluwer Academic Publishers, Dordrecht, the Netherlands, pp. 702-703 doi: 10.1007/0-306-47624-X
- Wrage, N., Velthof, G.L., van Beusichem, M.L., Oenema, O. 2001. Role of nitrifier denitrification in the production of nitrous oxide. *Soil Biology and Biochemistry* 33, 1723-1732.
- Wrage, N., Velthof, G.L., Laanbroek, H.J. & Oenema, O. 2004. Nitrous oxide production in grassland soils: assessing the contribution of nitrifier denitrification. *Soil Biology and Biochemistry* 36, 229-236.
- Yamulki, S., Jarvis, S.C., Owen, P. 1998. Nitrous oxide emissions from excreta applied in a simulated grazing pattern. *Soil Biology and Biochemistry* 30, 491-500.
- Zakir, H.A.K.M., Subbarao, G.V., Pearse, S.J., Gopalakrishnan, S., Ito, O., Ishikawa, T., Kawano, N., Nakahara, K., Yoshihashi, T., Ono, H., Yoshida, M. 2008. Detection, isolation and characterization of root-exuded compound, methyl 3-(4-hydroxyphenyl)

propionate, responsible for biological nitrification inhibition by sorghum (*Sorghum bicolor*). *New Phytologist* 180, 442-451.

Zaman, M., Nguyen, M.L. 2012. How application timings of urease and nitrification inhibitors affect N losses from urine patches in pastoral systems. *Agriculture, Ecosystems and Environment* 156, 37-48.

Zerulla, W., Barth, T., Dressel, J., Erhardt, K., von Locquenghien, K.H., Pasda, G., Rädle, M., Wissemeier, A.H. 2001. 3,4-Dimethylpyrazole phosphate (DMPP) – a new nitrification inhibitor for agriculture and horticulture. *Biology and Fertility of Soils* 34, 79-84.

---

## Chapter 3

---

### **The urine patch diffusional area: an important N<sub>2</sub>O source?**

Karina A. Marsden\*, Davey L. Jones, David R. Chadwick

This manuscript has been published in Soil Biology and Biochemistry

Marsden, K.A., Jones, D.L., Chadwick, D.R. 2016. The urine patch diffusional area: An important N<sub>2</sub>O source? Soil Biology and Biochemistry 92, 161-170.

KAM, DLJ and DRC designed and conceived the experiment, KAM conducted the practical work, analysed the data and prepared the manuscript.

## **Abstract**

Urine patches contribute greatly to greenhouse gas emissions within livestock-grazed ecosystems. The effective area of a ruminant urine patch comprises the wetted area, the diffusional area and the pasture response area. This study specifically assesses the importance of considering the diffusional area for monitoring urine patch N<sub>2</sub>O emissions. Spatial and temporal changes in N<sub>2</sub>O emissions and potential drivers of emissions (soil pH, EC, redox potential, dissolved organic carbon and nitrogen, NO<sub>3</sub><sup>-</sup> and NH<sub>4</sub><sup>+</sup>) were measured in sheep urine amended Eutric Cambisol mesocosms, maintained at 50% or 70% water-filled pore space (WFPS). At 70% WFPS, over 10 weeks, the emission factor (EF) was greater when considering the wetted area plus a 9 cm diffusional area (EF = 2.75 ± 0.72% of applied N) than when considering the wetted area alone (EF = 1.44 ± 0.30% of applied N); differences were not statistically significant at 50% WFPS. Redox potential, total extractable N and WFPS contributed significantly to the observed variation in daily N<sub>2</sub>O fluxes from the urine patch. We conclude that the urine patch diffusional area is an extremely important source of emissions from urine patches. This has implications when measuring EFs, as the lateral diffusion of solutes may be restricted by chamber walls resulting in an underestimate of N<sub>2</sub>O emissions, particularly at higher soil moisture contents. Site-specific assessments of the urine patch diffusional area should be made, and accounted for, prior to monitoring emissions and calculating emission factors from urine patches applied within chambers.

**Key Words:** animal waste; denitrification; DOC; DON; nitrous oxide; grazing returns

### 3.1 Introduction

Grazing returns of excreta to pasture soils are estimated to account for 40% of the total (direct and indirect) nitrous oxide (N<sub>2</sub>O) emissions from animal production systems, globally (Oenema et al., 2005). Additions of labile carbon (C) and nitrogen (N) to soil in the form of urine (van Groenigen et al., 2005) fuel the major microbial N<sub>2</sub>O producing processes of nitrification and denitrification, creating “hot spots” and “hot moments” for emissions within pastures (McClain et al., 2003; Groffman et al., 2009). The current default IPCC emission factor used in national inventories for excretal deposition to soils are 1% and 2% of deposited N for sheep and cattle, respectively (IPCC, 2006); yet, this Tier 1 approach lacks accuracy as it fails to account for variation in N<sub>2</sub>O emissions due to environmental, edaphic or management related factors (Skiba and Smith, 2000; Skiba et al., 2012; Buckingham et al., 2014).

Variability in N<sub>2</sub>O emissions from urine patches can arise due to differences in urine composition, the amount of N excreted and the volume and frequency of urine events (Dijkstra et al., 2013). Additionally, microbial N<sub>2</sub>O production and consumption processes depend on several interacting environmental controls (Bouwman et al., 2013) such as N supply, soil temperature, soil moisture, oxidation-reduction potential (ORP), the availability of labile organic compounds, soil type, soil pH and climate (Skiba and Smith, 2000; Butterbach-bahl et al., 2013). The ORP conditions can affect the chemical status of soil nutrients and affect the activity of microorganisms, and this parameter is generally negatively correlated with pH and EC and can be used to indicate the aeration status of soils (Husson, 2013). Urine patches offer potential for emission reductions and improvements of nitrogen use efficiency (NUE) within the agricultural sector, yet a greater understanding of the spatial and temporal variability in N<sub>2</sub>O emissions from urine patches (at several scales of magnitude) is required to improve emission estimates and provide information for emission reduction strategies, such as the use of nitrification inhibitors.

The urine patch “wetted area”, where urine is directly voided, has been distinguished from the “effective area” which incorporates the diffusive edge of solutes and the plants able to access this pool of nutrients via root extension (Selbie et al., 2015). It is suggested that the effective area actually comprises the “wetted area”, the “diffusional area” and the “pasture response area” in order to distinguish between these regions. The pasture response area can extend to twice the initial wetted area (Doak, 1952), however, the diffusive edge of urinary N has been shown not to exceed 20 cm beyond the initial wetted area in three soil types (granitic Brunisol, Neoluvisol and Calcisol; Decau et al., 2003). The diffusional area may vary with urinary volume, solute concentrations, soil texture (relating to tortuosity and cation exchange capacity), soil moisture content, topography and time. The pasture response area is likely to be dependent on the magnitude of the diffusional area, the vegetation type and the corresponding root architecture.

Dennis et al. (2011) maintain that for investigating soil nutrient cycling processes, the wetted area is more important than the pasture response area, however, the diffusive edge of solutes may also be important to consider. An overestimation of  $\text{NO}_3^-$  leaching losses from urine applied to lysimeters may occur if no room is allowed for the diffusional area (Selbie et al., 2015). Similarly, underestimations of  $\text{N}_2\text{O}$  emissions may occur if the urine patch diffusional area is not considered (e.g. applying urine uniformly to the entire area beneath a static chamber for gas flux measurements), due to chamber walls preventing the lateral movement of solutes into surrounding soil. Koops et al. (1997) have demonstrated that  $\text{N}_2\text{O}$  losses from the diffusive zone of an artificial urine patch, applied to a peat grassland, can reach the same order of magnitude as the area where urine was directly applied.

This experiment was predicated on the need to assess the importance of the urine patch diffusional area for different solutes,  $\text{N}_2\text{O}$ -regulating soil properties (e.g. dissolved organic C and ORP) and the accuracy of  $\text{N}_2\text{O}$  emission measurements. Eutric Cambisol mesocosms,

amended with sheep urine, were established in order to 1) assess the spatial and temporal changes in soil properties in the wetted and diffusional area, 2) identify those soil properties which are key drivers of N<sub>2</sub>O emissions under two moisture regimes, and 3) compare the N<sub>2</sub>O emission factor from the wetted area with that of the wetted and diffusional areas combined. Two soil moisture regimes (50% and 70% WFPS) were included, as this was considered important with regards to both N<sub>2</sub>O production processes, emissions and the spatial distribution of solutes within the urine patch.

## **3.2 Materials and Methods**

### *3.2.1. Soil sampling and analysis*

Independent replicate ( $n = 4$ ) samples of a Eutric Cambisol (0–10 cm) were collected from a sheep-grazed, fertilised grassland located at Henfaes Agricultural Research Station, Abergwyngregyn, North Wales (53°14'N, 4°01'W). After collection, the soil was sieved through a 10 mm mesh. Soil moisture content was determined by oven drying (105 °C, 24 h), and organic matter was determined by loss-on-ignition (450 °C, 16 h; Ball, 1964). Soil pH and electrical conductivity (EC) were measured using standard electrodes submerged in 1:2.5 (w/v) soil-to-distilled water suspensions. The oxidation-reduction potential (ORP) was measured directly in the soil using an ELIT 31C ORP combination electrode (EA Instruments Ltd., London, UK) connected to a mV reader.

Total soil C and N were determined on oven-dried, ground soil using a TruSpec® Analyzer (Leco Corp., St. Joseph, MI). Within 24 h of soil collection, 1:5 (w/v) soil-to-0.5 M K<sub>2</sub>SO<sub>4</sub> extractions were performed; the total dissolved C and N (mineral and organic) in the resulting extracts were determined with a Multi N/C 2100S Analyzer (AnalytikJena, Jena, Germany). Nitrate (NO<sub>3</sub><sup>-</sup>), ammonium (NH<sub>4</sub><sup>+</sup>) and phosphate (P) within the 0.5 M K<sub>2</sub>SO<sub>4</sub> extracts were measured by the colorimetric methods of Miranda et al. (2001), Mulvaney (1996)



and Murphy and Riley (1962), respectively. The cations (Na, K and Ca) within 1:5 (w/v) soil-to-1 M NH<sub>4</sub>Cl soil extracts were measured using a Sherwood Model 410 Flame Photometer (Sherwood Scientific Ltd, Cambridge, UK). A summary of the soil characteristics is provided in Table 3.1.

**Table 3.1** Properties of the Eutric Cambisol used to fill soil mesocosms. Values represent means  $\pm$  SEM ( $n = 4$ ) and results are reported on a dry weight basis.

Eutric Cambisol properties	
Texture	Sandy Clay Loam
Field wet bulk density (g cm <sup>-3</sup> )	1.57 $\pm$ 0.05
Moisture content (%)	21.9 $\pm$ 1.00
pH	6.91 $\pm$ 0.17
EC ( $\mu$ S cm <sup>-1</sup> )	65.4 $\pm$ 5.14
ORP (mV)	368 $\pm$ 10.3
Total C (%)	3.29 $\pm$ 0.22
Total N (%)	0.26 $\pm$ 0.15
C:N ratio	13.0 $\pm$ 0.99
Dissolved organic C (mg C kg <sup>-1</sup> )	102 $\pm$ 8.66
Total dissolved N (mg N kg <sup>-1</sup> )	13.8 $\pm$ 1.89
Extractable NO <sub>3</sub> <sup>-</sup> (mg N kg <sup>-1</sup> )	2.28 $\pm$ 0.32
Extractable NH <sub>4</sub> <sup>+</sup> (mg N kg <sup>-1</sup> )	0.41 $\pm$ 0.24
Extractable P (mg P kg <sup>-1</sup> )	9.27 $\pm$ 0.91
Exchangeable Na (mg kg <sup>-1</sup> )	54.0 $\pm$ 5.98
Exchangeable K (mg kg <sup>-1</sup> )	181 $\pm$ 21.5
Exchangeable Ca (g kg <sup>-1</sup> )	1.09 $\pm$ 0.05

### 3.2.2. Urine collection and analysis

Welsh Mountain ewes ( $n = 5$ ) were fed a diet of freshly cut grass (*Lolium perenne* L.; 80%) and white clover (*Trifolium repens* L.; 20%). Ewes were housed in individual pens on plastic slatted flooring designed for sheep (Rimco Ltd., Yorkshire, UK), with collection trays located beneath the flooring for urine collection. Urine samples were centrifuged (4000 g; 10 min) and immediately frozen (-20°C) until required, to minimize losses of N. The urine

collected from five replicate sheep was bulked, in order to provide sufficient urine of a homogenous composition for experimental use. The total dissolved C, total dissolved N, pH, EC,  $\text{NO}_3^-$ ,  $\text{NH}_4^+$ , P and cations were determined directly within the urine samples, as described for the soil samples. The urea content of the urine was determined via the method of Orsonneau et al. (1992). A summary of the urine characteristics is provided in Table 3.2.


**Table 3.2** Properties of sheep urine, applied to Eutric Cambisol mesocosms. Values represent means  $\pm$  SEM ( $n = 3$ ), where replicates are analytical replicates of urine combined from 5 individual sheep.

Urine properties	
pH	9.15 $\pm$ 0.01
EC (mS $\text{cm}^{-1}$ )	14.1 $\pm$ 0.20
Dissolved organic C (g C $\text{l}^{-1}$ )	6.03
Total N (g N $\text{l}^{-1}$ )	3.86
Urea (g N $\text{l}^{-1}$ )	2.71 $\pm$ 0.61
$\text{NH}_4^+$ (mg N $\text{l}^{-1}$ )	129 $\pm$ 5.30
$\text{NO}_3^-$ (mg N $\text{l}^{-1}$ )	1.08 $\pm$ 0.02
P (mg P $\text{l}^{-1}$ )	11.6 $\pm$ 0.34
Na (mg $\text{l}^{-1}$ )	692 $\pm$ 1.59
K (g $\text{l}^{-1}$ )	4.00 $\pm$ 0.05
Ca (mg $\text{l}^{-1}$ )	48.4 $\pm$ 0.54

### 3.2.3 Soil mesocosm preparation, treatment application and sampling regime

Briefly, 5 kg of fresh soil ( $n = 4$ ) was weighed into polypropylene trays (internal height: 11 cm, internal length: 35.5 cm, internal width: 26.5 cm; see Appendix 1, Fig. 1), and repacked to a depth of 5 cm, resulting in a fresh bulk density of 1.10  $\text{g cm}^{-3}$ . Bare pasture soil mesocosms were used, in order to gain a mechanistic understanding of processes that occur at the soil-urine interface, in the absence of competing factors (e.g. plant removal of nutrients,  $\text{NO}_3^-$  leaching). The soil mesocosms were wetted with distilled water using a fine mist sprayer to facilitate even

coverage to achieve 50% and 70% water-filled pore space (WFPS), where the initial starting weights were recorded. The mesocosms were pre-incubated in a greenhouse maintained at 20°C for 1 week before application of urine, to ensure any observed effects were not due to soil disturbance (e.g. sieving). Soil mesocosms were maintained under these conditions for the duration of the experiment, and rewetted weekly with distilled water to achieve initial starting weights using a fine mist sprayer. Urine (36 ml) was applied in a strip (24 cm × 3 cm) across the width of the mesocosms, resulting in an equivalent urine-to-soil surface area for an average sheep urine deposition (150 ml over 300 cm<sup>2</sup>; Doak, 1952). This urine application resulted in an equivalent total N loading rate of ca. 200 kg N ha<sup>-1</sup>, where other studies investigating sheep urine patches have used N application rates of ca. 300 kg N ha<sup>-1</sup> (Haynes and Williams, 1993; Moir et al., 2003). Soil was sampled at increasing distances away from the centre of the urine patch, along a horizontal diffusional gradient. Briefly, 0-3 cm represents the centre of the urine patch, with further sampling conducted at 3-6, 6-9, 9-12, 15-18 and 27-30 cm away from the direct area of urine application, hereafter referred to as zone A, B, C, D, E and F, respectively (see Fig. 3.1). The final sampling distance (27–30 cm; Zone F) was considered to be the control, as we hypothesised that this zone would receive no effect from the urine application. Samples were taken from three parallel mesocosms for each replicate ( $n = 4$ ), to provide enough soil sampling points for the duration of the experiment (10 weeks). Sampling was conducted three times a week for the first two weeks and once a week thereafter, until the end of the experiment.



**Figure 3.1** Aerial schematic view of the Eutric Cambisol mesocosms, repacked to a depth of 5 cm. The shaded region represents the area of direct urine application, labelled rows display sampling regions and T1 to T6 represent successive sampling time points, where further time points were sampled from parallel mesocosms.

### *3.2.4 Monitoring nitrous oxide emissions and changes in soil properties*

Soil from each sampling zone (see Fig. 3.1; ca. 53 g) was removed, weighed and placed into gas-tight polypropylene containers (total volume 700 cm<sup>3</sup>) fitted with a silicone Suba Seal<sup>®</sup> (VWR International, Lutterworth, UK). Gas samples (20 ml) were taken at 0 and 60 min following container lid closure and were stored in pre-evacuated 20 ml glass vials. The linearity of gas build up within the containers was checked by taking four gas samples (0, 20, 40 and 60 min) on each sampling day in zone A, as this was expected to have the highest emissions. Linearity of gas build-up within containers was met ( $R^2 > 0.95$ ) on 29 and 39% of occasions at 50% and 70% WFPS, respectively. Chadwick et al. (2014) state that non-linear fluxes can arise during occasions of no significant net flux. Our data supports this in that where the linear assumption was violated, the fluxes tended to be minimal and during periods of high emissions the data fitted well to the linear model. We, therefore, consider this acceptable as a poorly fitted linear model at periods of non-significant fluxes is unlikely to cause excessive bias in overall emission estimates. In addition, a high amount of sampled headspace air (during the linearity checks), may have resulted in poorly linear data.

Gas samples were analysed for N<sub>2</sub>O with a Varian 450 GC (Agilent Technologies, Santa Clara, CA), fitted with a <sup>63</sup>Ni electron capture detector (ECD), where the column, injector and detector temperatures were 50, 100 and 330°C, respectively. After gas sampling, the soil pH, EC and ORP were measured using the methods described previously. Total dissolved organic C, total dissolved N, NH<sub>4</sub><sup>+</sup> and NO<sub>3</sub><sup>-</sup> were measured following extraction of the excavated soils with 0.5 M K<sub>2</sub>SO<sub>4</sub>, as described previously (Section 2.1).

### *3.2.5 Statistical analysis*

All statistical analyses were performed in either SPSS Statistics 20.0 (IBM UK Ltd, Portsmouth, UK) or Minitab 17.1.0 (Minitab Inc., State College, PA). Spatial and temporal

differences between soil pH, EC and ORP in Zones A to E were compared to the control (zone F), and differences between soil incubated at 50% and 70% WFPS were determined via one-way ANOVA, followed by Fisher's LSD post-hoc test. Normality and homogeneity of variance assumptions were tested on log-transformed data using Shapiro Wilk and Levene's test, respectively and significance was determined at the  $p < 0.05$  level. Cumulative N<sub>2</sub>O emissions were determined by integration using the trapezoidal rule, and differences between the cumulative emissions in each zone were compared via one-way ANOVA, as above. The N<sub>2</sub>O emission factor (EF) for each treatment was calculated via the following equation:

$$EF = \text{treatment N}_2\text{O-N} - \text{control N}_2\text{O-N} / \text{Total N applied} \times 100\% \quad (\text{Eqn. 3.1})$$

Differences in emission factors between Zone A and the sum of Zones A-D were compared via one-way ANOVA, as above.

In order to determine the amount of variation in N<sub>2</sub>O emissions explained by measured soil parameters, multiple linear regression was used. Data were ln transformed where the distribution was improved by the transformation, in order to approximate normality. During exploratory data analysis, best subset's regression was used; this procedure compares all models for a given set of predictor variables (pH, EC, ORP, Total N, DOC, NO<sub>3</sub><sup>-</sup>, NH<sub>4</sub><sup>+</sup>, and WFPS), and provides summary statistics (R<sup>2</sup>, adjusted R<sup>2</sup>, predicted R<sup>2</sup>, S and Mallows' Cp) for the best two candidate models with increasing numbers of fixed predictor variables. The number and type of predictor variables were chosen based on the criteria of having a high R<sup>2</sup>, adjusted R<sup>2</sup> and predicted R<sup>2</sup>, a low value of S (which represents the standard deviation of the error term) and Mallows' Cp values close to the number of terms in the model. The best candidate models were then inputted into the normal multiple regression regime in Minitab.

### 3.3 Results

#### 3.3.1 Urine patch pH, EC and ORP


Spatial and temporal variation in soil pH, EC and ORP was observed in the Eutric Cambisol following sheep urine application (Fig. 3.2). In Figure 3.2, data are only displayed for the zones A (Fig. 3.2a, e and i), B (Fig. 3.2b, f and j), C (Fig. 3.2c, g and k) and F (Fig. 3.2d, h and l), as data for zones D (see Appendix 1, Fig. 2a, c and e) and E (see Appendix 1, Fig. 2b, d and f) were similar to zone F. The initial soil pH was  $6.87 \pm 0.15$  (50% WFPS) and  $6.84 \pm 0.16$  (70% WFPS; Fig. 3.2d), which increased immediately to pH  $8.58 \pm 0.08$  (50% WFPS) and  $8.21 \pm 0.14$  (70% WFPS) following urine deposition to zone A (Fig. 3.2a). During the first 3 days of incubation, the pH in zone A (50% WFPS) was more alkaline ( $p < 0.01$ ) than that of the control (zone F; Fig. 3.2d). By day 7 the pH had returned to a similar ( $p > 0.05$ ) value to zone F, however, after 10 days the pH was more acidic ( $p < 0.01$ ) than soil previously unaffected by urine, and remained so for the duration of the experiment. Differences in pH, in comparison to the control, only extended to zone B (Fig. 3.2b); this zone was more alkaline ( $p < 0.05$ ) in comparison to zone F (Fig. 3.2d) immediately after urine deposition and returned to the control value faster (after 2 days) than the immediate area of application (Fig. 3.2a). After 4 days, zone B was more acidic ( $p < 0.05$ ) than that of the control for the duration of the experiment. The spatial and temporal changes in pH were generally very similar at 50% and 70% WFPS.

The Eutric Cambisol had an EC of  $68.7 \pm 6.1$  and  $74.8 \pm 7.2 \mu\text{S cm}^{-1}$  (50% and 70% WFPS, respectively) without urine application (Fig. 3.2h), which increased to  $367 \pm 8.5$  and  $360 \pm 30.8 \mu\text{S cm}^{-1}$  (50% and 70% WFPS, respectively) immediately following urine application (Fig. 3.2e). The EC of Zone A and B (Fig. 3.2e and f, respectively) of the 50% WFPS treatment was greater ( $p < 0.001$ ) for all sample points in comparison to zone F (Fig. 3.2h). A greater EC was also observed in zones C (Fig. 3.2g) and D (see Appendix 1, Fig. 2c),

however, these only became significant ( $p < 0.05$ ) after 4 and 7 days, respectively, indicating a temporal delay in the lateral movement of solutes. No difference ( $p < 0.05$ ) in EC was observed in the 50% WFPS soil from zone E (see Appendix 1, Fig. 2d) in comparison to the control. At both 50% and 70% WFPS in zones A and B (Fig. 3.2e and f, respectively), the EC was immediately higher than the control (Fig. 3.2h), remaining so for the duration of the experiment; the EC in zone C (Fig. 3.2g) was immediately higher than the control (Fig. 3.2h) at 70% WFPS, however, at 50% WFPS it took 4 days for the EC in zone C to be greater than the control. After 10 days, the EC in zone D at 70% WFPS was greater ( $p < 0.001$ ) than the control (Fig. 3.2h), and remained so for the duration of the experiment.

Following application of urine the ORP in zone A at 50% and 70% WFPS ( $163 \pm 14$  and  $160 \pm 27$  mV, respectively; Fig. 3.2i) was lower ( $p < 0.001$ ) than zone F ( $382 \pm 6$  and  $383 \pm 6$  mV at 50% and 70% WFPS, respectively; Fig. 3.2l) and increased to control levels after 13 days. The ORP in zone B (Fig. 3.2j) at 50% and 70% WFPS ( $224 \pm 28$  and  $148 \pm 21$  mV, respectively) was also lower ( $p < 0.01$ ) than zone F (Fig. 3.2l), and increased to that of the control after one week following urine deposition. The ORP was lower ( $p < 0.05$ ) at 50% WFPS in zone A as opposed to the 70% WFPS treatment (Fig. 3.2i), during days 13-48 after urine application.






**Figure 3.2** Changes in soil pH (panels a, b, c and d), electrical conductivity (EC; panels e, f, g and h) and oxidation reduction potential (ORP; panels i, j, k and l) following sheep urine application to a Eutric Cambisol, maintained at either 50% or 70% water-filled pore space (WFPS), and sampled at increasing distances away from the direct area of application (Zone A, 0-3 cm: panels a, e and i; Zone B, 3-6 cm: panels b, f and j; Zone C, 6-9 cm: panels c, g and k; Zone F, control: panels d, h and l). Symbols represent means  $\pm$  SEM ( $n = 4$ ). Figure legend applies to all panels and text on the top row of panels applies to each respective column.

### 3.3.2 Spatial and temporal dynamics of nitrogen and carbon in the urine patch

Nitrogen and carbon dynamics following sheep urine deposition to a Eutric Cambisol are shown in Figure 3.3, which includes results for total extractable N (Fig. 3.3a, b, c and d),

$\text{NH}_4^+$  (Fig. 3.3e, f, g and h),  $\text{NO}_3^-$  (Fig. 3.3i, j, k and l), total DOC (Fig. 3.3m, n, o and p) and  $\text{N}_2\text{O}$  emissions (Fig. 3.3q, r, s and t), as no major differences were observed in comparison to the control for Zones D (Appendix 1, Fig. 3a, c, e, g and i) and E (Appendix 1, Fig. 3b, d, f, h and j), only Zones A (Fig. 3.3a, e, i, m and q), B (Fig. 3.3b, f, j, n and r), C (Fig. 3.3c, g, k, o and s) and F (Fig. 3.3d, h, l, p and t) are displayed. Most of the applied urine-N was in the form of urea (Table 3.2), which quickly hydrolysed in the soil. This resulted in immediately high soil  $\text{NH}_4^+$  concentrations in zone A (50% WFPS; Fig. 3.3e), which peaked at the first sample point at  $240 \pm 44 \text{ mg NH}_4^+\text{-N kg}^{-1}$  soil DW. In Zone A of the 70% WFPS soil, the  $\text{NH}_4^+$  was high at the first sample point ( $93 \pm 25 \text{ mg NH}_4^+\text{-N kg}^{-1}$  soil DW) but peaked 3 days following urine application at  $140 \pm 45 \text{ mg NH}_4^+\text{-N kg}^{-1}$  soil DW. The  $\text{NH}_4^+$  did not diffuse far in the soil and only minor amounts were measured further than zone B.

As nitrification proceeded, the  $\text{NH}_4^+$  concentration decreased and a concomitant increase in  $\text{NO}_3^-$  was observed. In zone A, the  $\text{NO}_3^-$  concentration peaked 19 days following urine application (Fig. 3.3i), where concentrations were higher in the soil incubated at 50% WFPS ( $268 \pm 9 \text{ mg NO}_3^-\text{-N kg}^{-1}$  soil DW) than the soil incubated at 70% WFPS ( $207 \pm 5 \text{ mg NO}_3^-\text{-N kg}^{-1}$  soil DW). Following the rapid increase in  $\text{NO}_3^-$  concentration, a decreasing trend was observed over 19–41 days following urine application. After 41 days, the  $\text{NO}_3^-$  concentration increased at similar rates to that of the control (zone F; Fig. 3.3l) in all zones, but the concentration remained higher than the control in zones A–D. The  $\text{NO}_3^-$  diffused further than the  $\text{NH}_4^+$ , and a temporal delay in the diffusion of  $\text{NO}_3^-$  into outer zones was observed. The major peaks in  $\text{N}_2\text{O}$  emission occurred in zones A and B during the first 20 days following urine application (Fig. 3.3m and n, respectively), while  $\text{NO}_3^-$  concentrations were still increasing. Emissions peaked immediately following urine deposition, where  $882 \pm 190$  and



**Figure 3.3** Soil extractable total dissolved nitrogen (panels a, b, c, d), ammonium (panels e, f, g, h), nitrate (panels i, j, k and l), nitrous oxide emissions (panels m, n, o and p) and extractable dissolved organic carbon (panels q, r, s and t) following sheep urine application to a Eutric Cambisol, maintained at either 50% or 70% water-filled pore space (WFPS), and sampled at increasing distances away from the direct area of application. Symbols represent means  $\pm$  SEM ( $n = 4$ ). Figure legend applies to all panels and text on the top row of panels applies to each respective column.


1825 ± 774 µg N<sub>2</sub>O-N m<sup>-2</sup> h<sup>-1</sup> were emitted from zone A at 50% and 70% WFPS, respectively. The greatest emissions observed in zone B were also on the day of urine application, and were lower than that of zone A at 431 ± 146 and 1048 ± 531 µg N<sub>2</sub>O-N m<sup>-2</sup> h<sup>-1</sup> at 50% and 70% WFPS, respectively. Another peak in emissions was observed 13 days following urine deposition in zones A and B, and this was more pronounced in the soil incubated at 70% WFPS. After 20 days following urine application, no major N<sub>2</sub>O emissions were measured, yet NO<sub>3</sub><sup>-</sup> levels decreased beyond this point.

During the first day of urine application, the concentration of total DOC in the control soil was 66.6 ± 8.6 and 79.1 ± 9.6 mg C kg<sup>-1</sup> soil DW at 50% and 70% WFPS, respectively (Fig. 3.3t). Due to the presence of labile C within the sheep urine and the potential for urine to solubilise soil organic matter, the concentration of DOC in soil within zone A was 161 ± 12.0 and 169 ± 27.0 mg C kg<sup>-1</sup> soil DW at 50% and 70% WFPS, respectively (Fig. 3.3q). This rapidly decreased over the course of one week following urine application to 54.0 ± 15.8 and 60.4 ± 10.4 mg C kg<sup>-1</sup> soil DW in zone A at 50% and 70% WFPS, respectively. A similar trend was observed in zone B (Fig. 3.3r), but at lower initial concentrations (106 ± 1.9 and 131 ± 7.6 mg C kg<sup>-1</sup> soil DW at 50% and 70% WFPS, respectively) indicating rapid movement and/or solubilisation of DOC into this zone.

### 3.3.3 Cumulative N<sub>2</sub>O emissions and urine patch emission factors

The cumulative N<sub>2</sub>O emissions within each zone are displayed in Figure 3.4. Greater cumulative N<sub>2</sub>O emissions (p < 0.01) were only observed within zone A at 50% WFPS, with respect to the control treatment. At 70% WFPS, both zone A (p < 0.01) and zone B (p < 0.05) emitted greater amounts of N<sub>2</sub>O in comparison to the control. Greater cumulative emissions were observed in the 70% WFPS treatment in comparison to the 50% WFPS in zone A, B and C (p < 0.01) but not D, E and F (p > 0.05). The emission factor when only considering zone A

was greater ( $p < 0.01$ ) in the soil maintained at 70% WFPS ( $1.44 \pm 0.30\%$  of applied N over 69 days) as opposed to the same soil maintained at 50% WFPS ( $0.44 \pm 0.06\%$  of applied N over 69 days). The  $N_2O$  emission factor at 70% WFPS was greater ( $p < 0.05$ ) when summing zones A – D ( $2.75 \pm 0.72\%$  of N applied over 69 days) than when only considering zone A ( $1.44 \pm 0.30\%$  of applied N over 69 days); this was not the case in the 50% WFPS soil, where accounting for the diffusive area had no effect on the  $N_2O$  emission factor.



**Figure 3.4** Cumulative nitrous oxide emissions following sheep urine application to a Eutric Cambisol, maintained at either 50% or 70% water-filled pore space (WFPS) and sampled at increasing distances away from direct area of urine application. Bars represent means  $\pm$  SEM ( $n = 4$ ) and different letters indicate significant differences (Fisher’s LSD,  $p < 0.05$ ).

### 3.3.4 Multiple regression analysis

The data included in the multiple regression analysis were those from zones A and B as these regions emitted most  $N_2O$ ; in addition, the control data (zone F) were also included. The results of the best subset’s regression (see Table 3.3) revealed a potential model containing

three predictor variables for individual ‘daily’ fluxes which fitted the selection criteria well. Increasing the number of variables beyond this did not substantially improve the predictive power of the model, therefore, in order to avoid over-fitting, only three variables were used. The model contained total extractable soil N, ORP and WFPS as predictor variables ( $R^2$ : 0.56; adjusted  $R^2$ : 0.55, predicted  $R^2$ : 0.53; Mallows’ Cp: 4.2; S: 0.72). The parameters were then entered separately into the least squares multiple regression model. The results of the regression are presented in Table 3.3 and the regression equations for 50% and 70% WFPS were:

$$50\% \text{ WFPS: } \ln \text{ N}_2\text{O 'daily' flux} = 14.65 - 1.879 \ln \text{ ORP} + 0.2364 \ln \text{ Total N} \quad (\text{Eqn. 3.2})$$

$$70\% \text{ WFPS: } \ln \text{ N}_2\text{O 'daily' flux} = 15.11 - 1.879 \ln \text{ ORP} + 0.2364 \ln \text{ Total N} \quad (\text{Eqn. 3.3})$$

**Table 3.3** Multiple regression analysis with  $\ln \text{ N}_2\text{O}$  as the dependent variable,  $\ln$  total extractable soil nitrogen (TN) and  $\ln$  oxidation reduction potential (ORP) as predictor variables and water-filled pore space (WFPS; 50% and 70%) as a categorical predictor variable.

Term	Unstandardized coefficients		Standardized coefficients	<i>T</i>	<i>p</i>	VIF <sup>a</sup>
	<b>B</b>	<b>SEM</b>	<b>Beta</b>			
(Constant)	5.05	0.09	-	57.0	0.00	-
TN	0.24	0.08	0.19	2.89	0.00*	1.31
ORP	-1.88	0.21	-0.61	-9.09	0.00*	1.32
WFPS (70% vs. 50%)	0.46	0.13	0.21	3.65	0.00*	1.00
<b>Model Summary</b>	<b><i>F</i></b>	<b><i>R</i><sup>2</sup></b>	<b><i>R</i><sup>2</sup> (adj)<sup>b</sup></b>	<b><i>R</i><sup>2</sup> (pred)<sup>c</sup></b>		
	54.6*	0.56	0.55	0.53		

<sup>a</sup> VIF = variance inflation factor, values close to 1 indicate predictors are not correlated.

<sup>b</sup>  $R^2$  (adj) =  $R^2$  adjusted for number of terms in the model.

<sup>c</sup>  $R^2$  (pred) = Predicted  $R^2$ , a measure of how well the model predicts the dependent variable for new observations.

\*  $p < 0.05$

### 3.4 Discussion

#### 3.4.1 Within-urine patch spatial and temporal variability

The first objective of this study was to determine changes in soil chemical properties of a sheep urine patch both spatially and temporally, and assess how this may influence N<sub>2</sub>O production. Within the first few hours of sheep urine application, the soil pH and EC, total extractable N and NH<sub>4</sub><sup>+</sup>, N<sub>2</sub>O emissions and total DOC had increased while the ORP had decreased in soil directly wetted by the urine. The same trend was observed in zone B, but to a lesser extent, indicating the spread of solutes by mass flow into the region of soil adjacent to the wetted area. The addition of urine to soil may have increased DOC within the soil, due to solubilisation of soil organic matter or lysis of microbial cells by the applied urine (Monaghan and Barraclough, 1993; Ambus et al., 2007; Lambie et al., 2012). The greatest spatial effects were observed for extractable total N and NO<sub>3</sub><sup>-</sup>, and EC, whereas all other urine induced soil changes mainly occurred in zones A and B. This is likely to be due to rapid diffusion of NO<sub>3</sub><sup>-</sup> and other ions present within ruminant urine through the soil.

In this study the soil pH increased by ca. 2–2.5 pH units, which can be attributed to the high carbonate content of the urine and to alkaline products generated during urea hydrolysis (van Groenigen et al., 2005; Carter, 2007). Soil pH returned to control levels after 7 days, following which it remained more acidic than the control due to the acidifying processes of ammonification, nitrification and urea hydrolysis (Bolan et al., 1991). The spatial changes in pH within a urine patch may be important for within-patch variability and source partitioning of N<sub>2</sub>O, as the N<sub>2</sub>O product ratios of nitrification, denitrification and dissimilatory NO<sub>3</sub><sup>-</sup> reduction to NH<sub>4</sub><sup>+</sup> (DNRA) are all influenced by soil pH (Stevens et al., 1998; Šimek and Cooper, 2002; Mørkved et al., 2007). The pH optimum of nitrification is 6.5–8.0 (Šimek and Cooper, 2002), and these conditions are generally met when ruminant urine is deposited to agricultural soils. In our study the pH dropped below the optimum for nitrification, to values

as low as pH 5.7. Denitrifying enzyme activity has been shown to be highest at, or near, the soils natural pH (Šimek et al., 2002), however, reductions in pH (from  $6.82 \pm 0.40$  to  $5.52 \pm 0.48$ ) over a 10 month period increased the  $N_2O/N_2$  product ratio of denitrification (Čuhel et al., 2010). In either case, denitrification activity and  $N_2O$  release via denitrification are likely to increase once the initial high pH within the urine patch has subsided, whereas  $N_2O$  release from nitrification may occur immediately following urine deposition. This may explain the split peak observed in  $N_2O$  emissions in this (see Fig. 3.3m and n) and other studies involving urine deposition (e.g. Di and Cameron, 2012; Boon et al., 2014). Alternatively, the second peak may reflect emissions from a more recalcitrant N-containing urine constituent. Advances in the use of stable isotopes, molecular techniques (Bateman and Baggs, 2005; Wrage et al., 2005; Baggs, 2008; Baggs, 2011) and quantum cascade laser based absorption spectroscopy for the measurement of  $N_2O$  isotopomers (Waechter et al., 2008; Decock and Six, 2013) will facilitate our understanding of the source partitioning of  $N_2O$  following urine deposition to soils.

The differences observed between EC in the different zones revealed a faster lateral movement of solutes at 70% compared to 50% WFPS. This indicates that dilution, mixing and diffusion within soils of a high moisture content may lead to a faster movement of  $NO_3^-$  to anaerobic denitrifying microsites within the soil (Luo et al., 1999), where diffusion of soluble carbon may then become limiting (Myrold and Tiedje, 1985). For the majority of the incubation, the soils could be described as moderately oxidized, however, the urine-influenced soil was poorly oxidised at the beginning of the study; this may be due to a localised increase in biological oxygen demand for degradation of the added urinary C (Azam et al., 2002; Baral et al., 2014). Interestingly, the ORP was lower ( $p < 0.05$ ) at 50% WFPS in zone A as opposed to the 70% WFPS treatment, during days 13-48 after urine application. Due to a reduction in the oxygen content of the soils, it may have been expected that the 70% WFPS treatment would have a lower ORP than the 50% WFPS treatment; however, pH and the abundance of oxidizing



and reducing agents can also influence ORP. Here, it is postulated that a greater dilution of oxidizing and reducing agents may have occurred at 70% WFPS, which resulted in a lower ORP at this moisture content. The majority of N<sub>2</sub>O emissions occurred when the ORP was between 160 – 350 mV, which is in line with results from studies of paddy and arable soils (Patrick and Jugsujinda, 1992; Yu et al., 2001).

The extractable soil NH<sub>4</sub><sup>+</sup> concentration was initially higher than the control in zone A, indicating rapid urea hydrolysis. The NH<sub>4</sub><sup>+</sup> concentration peaked at 50% WFPS on the day of urine application, however, at 70% WFPS the NH<sub>4</sub><sup>+</sup> concentration peaked 3 days after urine deposition. Increasing moisture increases urease activity up to field capacity, following which it decreases (Dharmakheerthi and Thenabadu, 1996). By using the soil water characteristics estimator of Saxton and Rawls (2006), the WFPS at field capacity was estimated to be 53% which may explain the slight delay in NH<sub>4</sub><sup>+</sup> generation at 70% WFPS as this was above field capacity. The time taken for completion of urea hydrolysis at both moisture contents is similar to that of other studies (e.g. Yadav et al., 1987). As the NH<sub>4</sub><sup>+</sup> was oxidised, the NO<sub>3</sub><sup>-</sup> concentration in the urine-influenced soil increased. The major emission period of N<sub>2</sub>O took place during the first 20 days after urine application, whilst nitrification was still taking place. As the soils were not completely saturated it is suggested that both nitrification and denitrification contributed to the overall N<sub>2</sub>O emissions, due to a combination of aerobic and anaerobic microsites within the soil. Less N<sub>2</sub>O emissions at 50% WFPS are consistent with an inhibitory effect of a greater oxygen content upon denitrification. The magnitude of N<sub>2</sub>O fluxes were similar to that measured by Allen et al. (1996), where dairy cow urine was applied to pasture blocks in a laboratory incubation.

Minimal N<sub>2</sub>O fluxes were observed beyond 20 days of incubation, even though NO<sub>3</sub><sup>-</sup> concentrations remained higher than the control, suggesting another factor may have been limiting N<sub>2</sub>O production. As temperature and moisture were controlled in this study, it is

suggested that labile C limitation (or DOC being present in a low quality form) prevented N<sub>2</sub>O emissions from denitrification, and an NH<sub>4</sub><sup>+</sup> limitation prevented N<sub>2</sub>O production via nitrification. Interestingly, NO<sub>3</sub><sup>-</sup> concentrations continued to decrease following the major N<sub>2</sub>O peak in the absence of plants; possible removal mechanisms are immobilization and diffusion into surrounding soil. Studies investigating the effect of increasing DOC on denitrification rates commonly use glucose as a readily available C source e.g. Weier et al. (1993), however, further work is required to understand how differences in DOC molecular weight and concentration, may influence denitrification rates. Some studies have demonstrated more readily available C compounds stimulate denitrification more than complex molecules (Bremner and Shaw, 1958; deCatanzaro and Beauchamp, 1985) and therefore, determining the effect of DOC species specifically found within ruminant urine on N<sub>2</sub>O emissions may explain some of the variability associated with emissions from urine patches related to urine composition.

### *3.4.2 Predicting N<sub>2</sub>O emission by multiple regression analysis*

The second objective of this study was to determine the amount of variation in N<sub>2</sub>O emissions which could be predicted by the measured soil parameters. In the final multiple regression model changes in total extractable soil N, WFPS and ORP all contributed significantly ( $p < 0.001$ ) to the variation in N<sub>2</sub>O emissions, explaining 55.6% of the total variation. Provided all other variables are held constant, increasing the total N by 1% resulted in a 0.24% increase in N<sub>2</sub>O emissions and decreasing ORP by 1% resulted in a 1.88% increase in N<sub>2</sub>O emissions, under these experimental conditions. N<sub>2</sub>O emissions were, on average, 58% higher in soil incubated at 70% in comparison to 50% WFPS, when holding ORP and total extractable N constant. Model parameters which contributed the most new information to the model followed the sequence ORP > WFPS > total extractable N. Low amounts of organic N were extracted from soils following urea hydrolysis and, therefore, the inclusion of total

extractable N in the best subset's regression, as opposed to individual  $\text{NO}_3^-$  or  $\text{NH}_4^+$  concentrations, supports the tenet that the majority of  $\text{N}_2\text{O}$  emissions were due to a combination of nitrification and denitrification.

### *3.4.3 The importance of urine patch edge effects*

The third objective of this study was to determine how important considering urine patch edge effects are when calculating  $\text{N}_2\text{O}$  emission factors. The marked difference between the lateral distribution of  $\text{NH}_4^+$  and  $\text{NO}_3^-$  within the urine patch highlights the importance of considering the urine patch diffusional area when monitoring  $\text{N}_2\text{O}$  emissions via chambers, or studies monitoring N loss via lysimeters. Under field conditions the mass of mineral N available for lateral diffusion may be influenced by plant uptake, the extent of vertical diffusion, leaching and preferential flow through soil macropores. The mesocosms used in this study used homogenised soil (i.e. preferential flow unlikely) and did not include effects of plant uptake, leaching, drainage and vertical diffusion beyond 5 cm. It would be expected that accounting for these processes would result in a lower mass of mineral N available for lateral diffusion than observed in this study. Nevertheless, the lateral diffusion of N in our study was 11 cm less than that observed by Decau et al. (2003), where cattle urine (3 l over 0.4 m<sup>2</sup>) was applied to 1 m deep lysimeters with a cross-sectional area of 2 m<sup>2</sup>. However, the experimental conditions in Decau et al. (2003) were not similar to those used in this study.

The  $\text{NH}_4^+$  derived from the urine application remained central to the urine patch, with only small amounts diffusing up to 3 cm away from the initial wetted area. Conversely, the highly mobile  $\text{NO}_3^-$  diffused ca. three times as far from the centre of the urine patch and persisted in the soil for a longer period. These results suggest that  $\text{N}_2\text{O}$  production via nitrification would be limited by the lateral diffusion of  $\text{NH}_4^+$ , and are therefore only likely to occur in the initial wetted area and the area of soil influenced by mass flow of urine through

soil immediately after deposition. On the other hand, denitrification of urinary nitrogen may occur both centrally and within a larger diffusional area of soil around the urine patch.

This suggests that in order for mitigation strategies such as synthetic or biological nitrification inhibitors to be effective at reducing  $\text{N}_2\text{O}$  emissions, it would be beneficial for the inhibitors to possess a similar charge and diffusion coefficient to  $\text{NH}_4^+$ . As roots can undergo death and decomposition in the direct urine deposition zone (Shand et al., 2002), it is likely that the biological (i.e. plant) delivery of nitrification inhibitors will be of most significance in the diffusive zone. Research regarding biological denitrification inhibition is still in its infancy (Bardon et al., 2014) and further research is required, yet, an effective denitrification inhibitor would ideally match the diffusive speed of  $\text{NO}_3^-$ .

In this study, emissions were ca. 1.5 and 2 (50% and 70% WFPS, respectively) times greater when considering the wetted and diffusional area (sum of zones A-D) in comparison to the wetted area only (zone A). Under field conditions this figure may be expected to be lower due to some removal of  $\text{NO}_3^-$  via plant uptake, draining, leaching and vertical diffusion however, it may be more representative of times where plant uptake is low or urine is deposited to areas of bare soil in the field. The walls of chambers for measuring gaseous emissions from soil are generally inserted to a depth of 5 cm. If a urine patch is applied uniformly throughout a chamber, then the chamber walls may prevent lateral diffusion of  $\text{NO}_3^-$  and DOC into the surrounding soil, resulting in greater concentrations than would have been present otherwise or a deeper infiltration of urine. This could potentially overestimate denitrification, however, due to the limited diffusive speed of  $\text{NH}_4^+$  perturbation to  $\text{N}_2\text{O}$  emissions from nitrification may be minimal. On the other hand, not considering the urine patch diffusional area may underestimate  $\text{N}_2\text{O}$  emissions, due to the smaller zone of soil influenced by the urine, resulting in fewer microbes exposed to the addition of N, DOC and moisture. Similarly, due to fewer microbes at soil depth, a greater vertical movement of urine may reduce direct emissions. It

cannot be excluded that the opposing effects could cancel each other out. Further work is required to assess these potential processes, which could be investigated via the use of  $^{15}\text{N}$ -labelled urine applied to larger and deeper pasture mesocosms with an intact sward, comparing source-partitioned emissions with and without a chamber wall to restrict diffusion.

#### 3.4.4 Theoretical diffusion of $\text{NO}_3^-$ and $\text{NH}_4^+$ through soil

A calculation of the theoretical diffusive speed of  $\text{NO}_3^-$  and  $\text{NH}_4^+$  through soil may be useful for estimating the urine patch diffusional area in differing soils (and hence the additional area required within chambers to improve accuracy of emission measurements). To assess this we compared the theoretical linear distance of diffusive movement to the observed diffusive movement in the mesocosms. The effective diffusion coefficient ( $D_e$ ) can be calculated using the equation

$$D_e = \frac{D_1 \times \theta f \times dC_1}{dC_s} \quad (\text{Eqn. 3.4})$$

where  $D_1$  is the diffusion coefficient in pure water,  $\theta$  is the soil volumetric moisture content,  $f$  is the impedance or pore tortuosity factor and  $dC_1/dC_s$  is the reciprocal of the buffer power (Nye and Tinker, 2000). To calculate  $D_e$  in the mesocosms we used  $D_1$  values of 1.60 and 1.64  $\text{cm}^2 \text{d}^{-1}$  for  $\text{NO}_3^-$  and  $\text{NH}_4^+$ , respectively (Lide, 2004), an  $f$  value of 0.3 (Jones et al., 2005) and the moisture contents of the mesocosms. Values for the buffer power in the same soil, were obtained from (Jones et al., 2012). Further, the linear distance ( $L$ ) of diffusive movement of  $\text{NO}_3^-$  and  $\text{NH}_4^+$  through time can be calculated as

$$L = (2D_e t)^{1/2} \quad (\text{Eqn. 3.5})$$

where  $t$  is time. Using these parameters, the linear diffusive distance of  $\text{NH}_4^+$  over 10 weeks was calculated as 1.50 and 1.72 cm at 50% and 70% WFPS, respectively. The calculated diffusive distance of  $\text{NO}_3^-$  was greater at 4.18 and 4.94 cm at 50% and 70% WFPS, respectively. In the soil mesocosms, increased  $\text{NH}_4^+$  was observed up to 3 cm from the urine patch edge, whereas increased  $\text{NO}_3^-$  concentrations were observed up to 9 cm from the urine patch edge. Some disparity between observed and theoretical values may be due to the coarser scale of measurement (3 cm fractions) in the mesocosms and saturation of the exchange phase with urine derived  $\text{K}^+$  (and other ions) which could lower the sorption of  $\text{NH}_4^+$ . The formation and subsequent diffusion of  $\text{NO}_3^-$  may have occurred after the  $\text{NH}_4^+$  had diffused 1.50–1.72 cm, which may be the reason for the greater observed compared to theoretical diffusive distance of  $\text{NO}_3^-$ . Further validation of this method by comparison to measured urine patch diffusional areas in the field, across varying soil types, soil moisture contents, microtopography and urine patch N concentrations and volume need to be investigated prior to utilising this equation as a method for determining the chamber size required for an experimental urine patch.

### **3.5 Conclusions**

The results of our study show that  $\text{N}_2\text{O}$  emissions can extend beyond the initial wetted area of a urine patch, and that this effect is greater under a high soil moisture content. For a typical sheep urine application to a Eutric Cambisol with an even surface, an additional 9 cm around the initial wetted area would have been required to capture the majority of  $\text{N}_2\text{O}$  emissions via a chamber based system. The additional area required around a urine patch may also vary alongside urine volume, patch area, the concentration of N applied, the soil type beneath the patch and the underlying microtopography. These conditions are likely to be highly site specific, therefore, preliminary assessments should be conducted in order to assess the

magnitude of the urine patch diffusional area, and additional area inside chambers should be allowed for, prior to monitoring emissions.

### **3.6 Acknowledgments**

Thanks to Jonathan Roberts, Rory Shaw, John Hyland and Matthew Scowen for their assistance sieving soil in the field. KAM would also like to thank the School of Environment Natural Resources and Geography, Bangor University, for sponsoring this study.

### 3.7 References

- Ambus, P., Petersen, S.O., Soussana, J.F., 2007. Short-term carbon and nitrogen cycling in urine patches assessed by combined carbon-13 and nitrogen-15 labelling. *Agriculture, Ecosystems and Environment* **121**, 84-92.
- Allen, A.G., Jarvis, S.C., Headon, D.M., 1996. Nitrous oxide emissions from soils due to inputs of nitrogen from excreta return by livestock on grazed grassland in the UK. *Soil Biology and Biochemistry* **28**, 597-607.
- Azam, F., Müller, C., Weiske, A., Benckiser, G., Ottow, J.C.G., 2002. Nitrification and denitrification as sources of atmospheric nitrous oxide – role of oxidizable organic carbon and nitrogen. *Biology and Fertility of Soils* **35**, 54-61.
- Baggs, E.M., 2008. A review of stable isotope techniques for N<sub>2</sub>O source partitioning in soils: recent progress, remaining challenges and future considerations. *Rapid Communications in Mass Spectrometry* **22**, 1664-1672.
- Baggs, E.M., 2011. Soil microbial sources of nitrous oxide: recent advances in knowledge, emerging challenges and future direction. *Current Opinion in Environmental Sustainability* **3**, 321-327.
- Ball, D.F. 1964. Loss-on-ignition as an estimate of organic matter and organic carbon in non-calcareous soils. *Journal of Soil Science* **15**, 84-92.
- Baral, K.R., Thomsen, A.G., Olesen, J.E., Peterson, S.E., 2014. Controls of nitrous oxide emission after simulated cattle urine deposition. *Agriculture, Ecosystems and Environment* **188**, 103-110.
- Bardon, C., Piola, F., Bellvert, F., Haichar, F. el Z., Comte, G., Meiffren, G., Pommier, T., Puijalón, S., Tsafack, N., Poly, F., 2014. Evidence for biological denitrification inhibition (BDI) by plant secondary metabolites. *New Phytologist* **204**, 620-630.



- Bateman, E.J., Baggs, E.M., 2005. Contributions of nitrification and denitrification to N<sub>2</sub>O emissions from soils at different water-filled pore space. *Biology and Fertility of Soils* **41**, 379-388.
- Bolan, N.S., Hedley, M.J., White, R.E., 1991. Processes of soil acidification during nitrogen cycling with emphasis on legume based pastures. *Plant and Soil* **134**, 53-63.
- Boon, A., Robinson, J.S., Chadwick, D.R., Cardenas, L.M., 2014. Effect of cattle urine addition on the surface emissions and subsurface concentrations of greenhouse gases in a UK peat grassland. *Agriculture, Ecosystems and Environment* **186**, 23-32.
- Bouwman, A.F., Beusen, A.H.W., Griffioen, J., van Groenigen, J.W., Hefting, M.M., Oenema, O., van Puijenbroek, P.J.T.M., Seitzinger, S., Slomp, C.P., Stehfest, E., 2013. Global trends and uncertainties in terrestrial denitrification and N<sub>2</sub>O emissions. *Philosophical Transactions of the Royal Society B* **368**, DOI: 10.1098/rstb.2013.0112.
- Bremner, J.M., Shaw, K. 1958. Denitrification in soil. II. Factors affecting denitrification. *The Journal of Agricultural Science* **51**, 40-52.
- Buckingham, S., Anthony, S., Bellamy, P.H., Cardenas, L.M., Higgins, S., McGeough, K., Topp, C.F.E., 2014. Review and analysis of global agricultural N<sub>2</sub>O emissions relevant to the UK. *Science of the Total Environment* **487**, 164-172.
- Butterbach-bahl, K., Baggs, E.M., Dannenmann, M., Kiese, R., Zechmeister-Boltenstern, S., 2013. Nitrous oxide emissions from soils: how well do we understand the processes and their controls? *Philosophical Transactions of the Royal Society B* **368**, 20130122.
- Carter, M.S., 2007. Contribution of nitrification and denitrification to N<sub>2</sub>O emissions from urine patches. *Soil Biology and Biochemistry* **39**, 2091-2102.
- Chadwick, D.R., Cardenas, L., Misselbrook, T.H., Smith, K.A., Rees, R.M., Watson, C.J., Mcgeough, K.L., Williams, J.R., Cloy, J.M., Thorman, R.E., Dhanoa, M.S., 2014.

- Optimizing chamber methods for measuring nitrous oxide emissions from plot-based agricultural experiments. *European Journal of Soil Science* **65**, 295–307.
- Čuhel, J., Šimek, M., Laughlin, R.J., Bru, D., Chèneby, D., Watson, C.J., Philippot, L., 2010. Insights into the effects of soil pH on N<sub>2</sub>O and N<sub>2</sub> emissions and denitrifier community size and activity. *Applied and Environmental Microbiology* **76**, 1870-1878.
- de Catanzaro, J.B., Beauchamp, E.G. 1985. The effect of some carbon substrates on denitrification rates and carbon utilization in soil. *Biology and Fertility of Soils*, **1**, 183-187.
- Decau, M.L., Simon, J.C., Jacquet, A., 2003. Fate of urine nitrogen in three soils throughout a grazing season. *Journal of Environmental Quality* **32**, 1405-1403.
- Decock, C., Six, J., 2013. How reliable is the intramolecular distribution of <sup>15</sup>N in N<sub>2</sub>O to source partition N<sub>2</sub>O emitted from soil? *Soil Biology and Biochemistry* **65**, 114-127.
- Dennis, S.J., Moir, J.L., Cameron, K.C., Di, H.J., Hennessy, D., Richards, K.G., 2011. Urine patch distribution under dairy grazing at three stocking rates in Ireland. *Irish Journal of Agricultural and Food Research* **50**, 149-160.
- Dharmakheerthi, R.S., Thernabadu, M.W., 1996. Urease activity in soils: a review. *Journal of the National Science Foundation of Sri Lanka* **24**, 159-195.
- Di, H.J., Cameron, K.C., 2012. How does the application of different nitrification inhibitors affect nitrous oxide emissions and nitrate leaching from cow urine in grazed pastures? *Soil Use and Management* **28**, 54-61.
- Dijkstra, J., Oenema, O., van Groenigen, J.W., Spek, J.W., van Vuuren, A.M., Bannink, A., 2013. Diet effects on urine composition of cattle and N<sub>2</sub>O emissions. *Animal* **7**, 292-302.
- Doak, B.W., 1952. Some chemical changes in the nitrogenous constituents of urine when voided on pasture. *The Journal of Agricultural Science* **42**, 162-171.

- Groffman, P.M., Butterbach-Bahl, K., Fulweile, R.W., Gold, A.J., Morse, J.L., Stander, E.K., Tague, C., Tonitto, C., Viddon, P., 2009. Challenges to incorporating spatially and temporally explicit phenomena (hotspots and hot moments) in denitrification models. *Biogeochemistry* **93**, 49-47.
- Haynes, R.J., Williams, P.H., 1993. Nutrient cycling and soil fertility in the grazed pasture ecosystem. *Advances in Agronomy* **46**, 119-199.
- Husson, O. 2013. Redox potential (Eh) and pH as drivers of soil/plant/microorganism systems: a transdisciplinary overview pointing to integrative opportunities for agronomy. *Plant and Soil* **362**, 389-417.
- IPCC 2006, 2006 IPCC Guidelines for National Greenhouse Gas Inventories, Prepared by the National Greenhouse Gas Inventories Programme, Eggleston H.S., Buendia L., Miwa K., Ngara T., Tanabe K. (Eds). Published: IGES, Japan.
- Jones, D.L., Healey, J.R., Willett, V.B., Farrar, J.F., Hodge, A., 2005. Dissolved organic nitrogen uptake by plants-an important N uptake pathway? *Soil Biology and Biochemistry* **37**, 413-423.
- Jones, D.L., Rousk, J., Edwards-Jones, G., DeLuca, T.H., Murphy, D.V., 2012. Biochar mediated changes in soil quality and plant growth in a three year field trial. *Soil Biology and Biochemistry* **45**, 113-124.
- Koops, J.G., van Beusichem, M.L., Oenema, O., 1997. Nitrous oxide production, its source and distribution in urine patches on grassland on peat soil. *Plant and Soil* **191**, 57-65.
- Lambie, S.M., Schipper, L.A., Balks, M.R., Baisden, W.T., 2012. Solubilisation of soil carbon following treatment with cow urine under laboratory conditions. *Soil Research* **50**, 50-57.
- Lide, D.R. (Ed.), 2004. *CRC Handbook of Chemistry and Physics, Student Edition*. CRC Press.

- Luo, J., Tillman, R.W., Ball, P.R., 1999. Factors regulating denitrification in a soil under pasture. *Soil Biology and Biochemistry* **31**, 913-927.
- McClain, M.E., Boyer, E.W., Dent, L., Gergel, S.E., Grimm, N.B., Groffman, P.M., Hart, S.C., Harvey, J.W., Johnston, C.A., Mayorga, E., McDowell, W.H., Pinay, G., 2003. Biogeochemical hot spots and hot moments at the interface of terrestrial and aquatic systems. *Ecosystems* **6**, 301-312.
- Miranda, K.M., Epsey, M.G., Wink, D.A., 2001. A rapid, simple, spectrophotometric method for simultaneous detection of nitrate and nitrite. *Nitric Oxide* **5**, 62-71.
- Moir, J.L. Edwards, G.R., Berry, E., 2012. Nitrogen uptake and leaching loss of thirteen temperate grass species under high N loading. *Grass and Forage Science* **68**, 313-325.
- Monaghan, R.M., Barraclough, D., 1993. Nitrous oxide and dinitrogen emissions from urine-affected soil under controlled conditions. *Plant and Soil* **151**, 127-138.
- Mørkved, P.T., Dörsch, P., Bakken, L.R., 2007. The N<sub>2</sub>O product ratio of nitrification and its dependence on long-term changes in soil pH. *Soil Biology and Biochemistry* **39**, 2048-2057.
- Mulvaney, R.L., 1996. Nitrogen – inorganic forms. In: Sparks, D.L. (Eds.), *Methods of Soil Analysis. Part 3*. Madison, WI, USA: Soil Science Society of America Inc., pp. 1123-1184.
- Murphy, J., Riley, J.P., 1962. A modified single solution method for determination of phosphate in natural waters. *Analytica Chimica Acta* **27**, 31-36.
- Myrold, D.D., Tiedje, M., 1985. Diffusional constraints on denitrification in soil. *Soil Science Society of America Journal* **49**, 651-657.
- Nye, P.H., Tinker, P.B., 2000. *Solute Movement in the Rhizosphere*. Oxford University Press, Oxford.

- Oenema, O., Wrage, N., Velthof, G.L., van Groenigen, J.W., Dolfing, J., Kuikman, P.J., 2005. Trends in global nitrous oxide emissions from animal production systems. *Nutrient Cycling in Agroecosystems* **72**, 51-56.
- Orsonneau, J-L., Massoubre, C., Cabanes, M., Lustenberger, P., 1992. Simple and sensitive determination of urea in serum and urine. *Clinical Chemistry* **38**, 619-623.
- Patrick, W.H., Jugsujinda, A., 1992. Sequential reduction and oxidation of inorganic nitrogen, manganese and iron in flooded soils. *Soil Science Society of America Journal* **56**, 1071-1073.
- Saxton, K.E., Rawls, W.J. 2006. Soil water characteristic estimates by texture and organic matter for hydrologic solutions. *Soil Science Society of America Journal* **70**, 1569-1578.
- Selbie, D.R., Buckthought, L.E., Shepherd, M.A., 2015. The challenge of the urine patch for managing nitrogen in grazed pasture systems. *Advances in Agronomy*, doi:10.1016/bs.agron.2014.09.004
- Shand, C.A, Williams, B.L., Dawson, L.A., Smith, S., Young, M.E., 2002. Sheep urine affects soil solution nutrient composition and roots: differences between field and sward box soils and the effects of synthetic and natural sheep urine. *Soil Biology and Biochemistry* **34**, 163-171.
- Šimek, M., Cooper, J.E., 2002. The influence of soil pH on denitrification: progress towards the understanding of this interaction over the last 50 years. *European Journal of Soil Science* **53**, 345-354.
- Šimek, M., Jišová, L., Hopkins, D.W., 2002. What is the so-called pH optimum for denitrification in soil? *Soil Biology and Biochemistry* **9**, 1227-1234.
- Skiba, U., Smith, K.A., 2000. The controls of nitrous oxide emissions from agricultural and natural soils. *Chemosphere – Global Change Science* **2**, 379-386.

- Skiba, U., Jones, S.K., Dragosits, U., Drewer, J., Fowler, D., Rees, R.M., Pappa, V.A., Cardenas, L., Chadwick, D., Yamulki, S., Manning, A.J., 2012. UK emissions of the greenhouse gas nitrous oxide. *Philosophical Transactions B* **367**, 1175-1185.
- van Groenigen, J.W., Kuikman, P.J., de Groot, W.J.M, Velthof, G.L., 2005. Nitrous oxide emissions from urine-treated soil as influenced by urine composition and soil physical conditions. *Soil Biology and Biochemistry* **37**, 463-473.
- Waechter, H., Mohn, J., Tuzson, B., Emmenegger, L., Sigrist, M.W., 2008. Determination of N<sub>2</sub>O isotopomers with quantum cascade laser based absorption spectroscopy. *Optics Express* **16**, 9239-9244.
- Weier, K.L., Doran, J.W., Power, J.F., Walters, D.T., 1993. Denitrification and the dinitrogen/nitrous oxide ratio as affected by soil water, available carbon and nitrate. *American Society of Agronomy* **57**, 66-72.
- Wrage, N., van Groenigen, J.W., Oenema, O., Baggs, E., 2005. A novel dual-isotope labelling method for distinguishing between soil sources of N<sub>2</sub>O. *Rapid Communications in Mass Spectrometry* **19**, 3298-3306.
- Yadav, D.S., Kumar, V., Singh, M., Relan, P.S., 1987. Effect of temperature and moisture on kinetics of urea hydrolysis and nitrification. *Australian Journal of Soil Research* **25**, 185-191.
- Yu, K.W., Wang, Z.P., Vermoesen, A., Patrick Jr., W.H., Van Cleemput, O., 2001. Nitrous oxide and methane emissions from different soil suspensions: effect of soil redox status. *Biology and Fertility of Soils* **34**, 25-30.



---

## Chapter 4

---

### **Disentangling the effect of sheep urine patch size and N loading rate on cumulative N<sub>2</sub>O emissions**

Karina A. Marsden\*, Davey L. Jones, David R. Chadwick

This experimental chapter forms the basis of a publication which has been accepted for a special edition of *Animal Production Science*, and accepted for an oral presentation in the Greenhouse Gases and Animal Agriculture 2016 conference in Melbourne, Australia.

Marsden, K.A., Jones, D.L., Chadwick, D.R. 2016. Disentangling the effect of sheep urine patch size and N loading rate on cumulative N<sub>2</sub>O emissions. *Animal Production Science*, In Press.

KAM, DLJ and DRC designed and conceived the experiment; KAM conducted the experimental work, analysed the data and prepared the manuscript.



## **Abstract**

Ruminant urine N concentration and volume are important parameters influencing the size and N loading rate of urine patches deposited to soil. Such parameters can influence N cycling and emissions of the greenhouse gas, nitrous oxide (N<sub>2</sub>O) from grazed grassland, yet, there is limited information on the effect of these parameters within typical ranges reported for sheep. We used an automated, high frequency gas monitoring system to investigate N<sub>2</sub>O emissions from varying urine N application rates and patch sizes under field conditions. Using artificial sheep urine, we manipulated urine N concentration to provide two urine N application rates (4 and 16 g N l<sup>-1</sup>; equivalent to 200 and 800 kg N ha<sup>-1</sup>). We investigated the effect of urine patch size with equal N application rates (4 × 125 cm<sup>2</sup> vs. 500 cm<sup>2</sup>, at 200 and 800 kg N ha<sup>-1</sup>) and the effect of patch size with unequal N application rates, but the same total amount of N applied (62.5 ml over 125 cm<sup>2</sup> at 800 kg N ha<sup>-1</sup> and 250 ml over 500 cm<sup>2</sup> at 200 kg N ha<sup>-1</sup>). Cumulative emissions of N<sub>2</sub>O generally increased with N loading rate, whether applied as one large urine patch or four smaller ones. Cumulative N<sub>2</sub>O emissions increased when the N was applied in four smaller urine patches compared to one large patch; this difference was significant at 800 kg N ha<sup>-1</sup>, but not at 200 kg N ha<sup>-1</sup>. When the total amount of N applied was held constant (1 g of N), the amount of N<sub>2</sub>O released was similar when urine was applied as a high N concentration small patch (800 kg N ha<sup>-1</sup>) compared to a low N concentration large patch (200 kg N ha<sup>-1</sup>). Urine N<sub>2</sub>O emission factors in this study were, on average, ten times lower than the IPCC default of 1% for sheep excreta. This research clearly demonstrates that the chemical and physical nature of the urine patch influences N<sub>2</sub>O emissions, yet further research is required to gather more data on typical sheep urine volumes (individual and daily), urination frequency, urine N concentrations and the typical volumes of soil influenced by urine deposition, to provide more accurate estimates of emissions from sheep grazed pastures.

**Key Words:** agricultural systems; global climate change; microbial processes; ruminants.

## 4.1 Introduction

Ruminant urine N concentration, volume and frequency vary widely, influencing the size and N loading rate of urine patches deposited to pasture soils. These parameters can influence the fate of urinary N in grazed pastures, including emissions of the powerful greenhouse gas nitrous oxide (N<sub>2</sub>O) and leaching of NO<sub>3</sub><sup>-</sup> (Li et al., 2012). The default IPCC N<sub>2</sub>O emission factor for both cattle and sheep was 2% of applied N, however, due to the lower volume and higher frequency of sheep urine events, and more moderate levels of soil compaction under sheep-grazed pastures, the default emission factor has been lowered for sheep from 2% to 1% of the applied N (IPCC, 2006). The range in urine volumes deposited by sheep and cattle (0.13-2 l) has been shown to influence N transformations and processes (e.g. longer retention of mineral N, greater plant growth and N uptake in larger patches) even when the total N applied is constant (Orwin et al., 2009). Whilst these results display a clear difference between sheep and cattle urine patches, where large differences in urinary volume are produced between the two species, less work has been conducted on N cycling and N<sub>2</sub>O emissions within the typical ranges of urine patch sizes reported for sheep.

The concentration of N excreted in urine is a function of the amount of surplus metabolized N to be excreted, the volume of urine produced and the frequency of urine events (Hoogendoorn et al., 2010) and can range from 1-18 g N l<sup>-1</sup> (Bristow et al., 1992; Oenema et al., 1997; Hoogendoorn et al., 2010). Urine volume is mainly influenced by water intake and the mineral load ingested by the animal (Selbie et al., 2015), and can be high when the moisture content of the diet is high, or when the herbage leaves are wet with rain water or dew (Doak, 1952). Urine volume can also vary as a response to coping with changes in ambient temperatures (Betteridge et al., 2010a). Data for sheep urine volume and frequency are scarce, but typically individual sheep urine events range between 100-200 ml (Doak, 1952; Haynes and Williams, 1993). However, variation within individual sheep urine volumes can be high

e.g. the mean and range of 40 individual urine events between 6 Welsh Mountain ewes fed (*ad libitum*) a *Lolium perenne* L. dominated sward was 104 (18-397) ml (Marsden, unpubl.). Daily sheep urine volume ranged from 0.5-3 l in Ledgard et al. (2008) and the frequency of urination events is reported to range between 13-20 times per day (Doak, 1952; Betteridge et al. 2010a; Betteridge et al., 2010b).

Urine volume, soil moisture status, soil type and topography can all influence the urine patch wetted area and volume of soil influenced by a urine patch. The wetted area of a urine patch has been defined as the surface area covered by urine following deposition to soil, as opposed to the pasture response area which can extend beyond the wetted area (Lantinga et al., 1987; Li et al., 2012). Typical wetted areas for a sheep urine event with a volume of 150 ml is reported to be 300 cm<sup>2</sup> (Doak, 1952) and for 200 ml is reported to be 430-550 cm<sup>2</sup> (Williams and Haynes, 1994). Typical urine applications of 4 l m<sup>-2</sup> were reported to be utilised in Kelliher et al. (2014).

Due to the wide range in sheep urine volume and frequency, and the interactive effects of dietary N content, energy use, and ambient temperature fluctuations it is evident that sheep urine patches vary widely in N concentration and volume, resulting in patches of different sizes and of different N loading rates. It can be envisaged that at times where sheep may be fed a diet low in moisture content, or when ambient temperatures are high, that their urine will be more concentrated, having a higher N concentration deposited over a small patch size. On the other hand, when the diet is high in moisture and the sheep is fully hydrated, urine may be deposited in larger sized urine patches with a lower N concentration. This experiment was designed to test these two extremes, to determine how differences in sheep urine patch size and N concentration may influence N cycling and cumulative N<sub>2</sub>O emissions arising from such urine patches.

Using a high frequency automated greenhouse gas monitoring system we assessed whether: 1) cumulative N<sub>2</sub>O emissions will increase with increasing urine N loading rate, when keeping the urine patch size constant, 2) cumulative emissions will be higher from four small urine patches compared to one large urine patch, when keeping the total N loading rate constant, and 3) cumulative emissions will be higher from a low N concentration large urine patch compared to high N concentration small urine patch, where N loading rate differs, but the total amount of N applied remains constant. The total amount of applied N equates to the total amount of N in the volume of applied urine. The N loading rate is a function of the urine N concentration and the surface area the urine is applied to (Selbie et al., 2015). We also examined the effect of reduced sampling frequency (i.e. reflecting the typical static chamber measurement approach) on measured cumulative N<sub>2</sub>O emissions.

## **4.2 Materials and methods**

### *4.2.1 Field site*

The field site was established at Henfaes Research Station, Abergwyngregyn, North Wales (53°14'N, 4°01'W) in March, 2015. The soil at the site is classified as a Eutric Cambisol and is of mixed glacial till in origin, deposited ca. 10,000 years ago. The field was reseeded in 1990 with a *Lolium perenne* L. and *Trifolium repens* L. mix. The field has since received moderate fertilizer applications (ca. 120 kg N ha<sup>-1</sup> y<sup>-1</sup> in years 1990-2002 and ca. 60-80 kg N ha<sup>-1</sup> y<sup>-1</sup> in subsequent years) and is regularly grazed by Welsh Mountain ewes at a moderate stocking density (2-3 livestock units / ha). The area was fenced off just over 3 months prior to treatment application, to prevent the effect of recent livestock urine patches on monitored gas fluxes from the soil.

#### *4.2.2 Soil sampling and analysis*

Soil (0-10 cm) was sampled in triplicate at block level ( $n = 3$ ) to determine soil characteristics (Table 4.1); this was conducted four days prior to treatment application. Soil moisture content was determined by oven drying (105°C; 24 h) and organic matter was determined by the loss-on-ignition (450 °C; 16 h; Ball, 1964). Soil pH and electrical conductivity (EC) were determined using standard electrodes in 1:2.5 (w/v) soil-to-distilled water suspensions. Total C and N were determined on oven-dried and ground samples in a TruSpec® Analyzer (Leco Corp., St. Joseph, MI). Dissolved C and dissolved N were determined in 1:5 (w/v) soil-to-0.5 M K<sub>2</sub>SO<sub>4</sub> extracts, according to Jones and Willet (2006). Microbial biomass C and N were determined via the chloroform fumigation-extraction method of Voroney et al. (2008), using K<sub>EC</sub> and K<sub>EN</sub> correction factors of 0.35 and 0.5, respectively. Total extractable P, NO<sub>3</sub><sup>-</sup>, and NH<sub>4</sub><sup>+</sup> were determined in 0.5 M K<sub>2</sub>SO<sub>4</sub> extracts via the methods of Murphy and Riley (1962), Miranda et al. (2001) and Mulvaney (1996), respectively. Exchangeable cations (Na, K, Ca) within 1:5 (w/v) soil-to-1 M NH<sub>4</sub>Cl extracts were measured using a model 410 flame photometer (Sherwood Scientific Ltd., Cambridge, UK).

#### *4.2.3 Experimental design and treatment application*

Artificial sheep urine was made up according to Lucas and Jones (2006), the composition of which is in line with the suggestion of Kool et al. (2006) in containing at least urea and hippuric acid for a realistic simulation of N<sub>2</sub>O emissions from artificial urine patches. The N concentration of the artificial urine was modified by adjusting the proportion of urea to obtain a total of 4 g N l<sup>-1</sup> and 16 g N l<sup>-1</sup>, where all other urine constituents were held constant. The two N concentrations were chosen to reflect low and high N concentrations in the urine, within typical reported ranges for N concentration within sheep urine. The effect of urine patch size was investigated by comparing a single large urine patch with four smaller patches, while

holding total N loading rate constant. The effect of patch size with unequal N loading rates, but the same total amount of applied N, was also assessed, by comparing a low N concentration large urine patch to a high N concentration small urine patch.

**Table 4.1** Properties of the Eutric Cambisol at the field site receiving artificial urine application. Values represent means  $\pm$  SEM ( $n = 3$ ) and results are reported on a dry weight basis.

Eutric Cambisol properties	
Texture	Sandy Clay Loam
Bulk density ( $\text{g cm}^{-3}$ )	$1.08 \pm 0.05$
Gravimetric moisture content (%)	$42.8 \pm 0.80$
Organic matter (%)	$11.1 \pm 0.68$
pH	$6.08 \pm 0.04$
EC ( $\mu\text{S cm}^{-1}$ )	$41.7 \pm 8.1$
Total C (%)	$5.47 \pm 0.77$
Total N (%)	$0.42 \pm 0.05$
C:N ratio	$12.8 \pm 0.65$
Dissolved organic C ( $\text{mg C kg}^{-1}$ )	$109 \pm 13.7$
Total dissolved N ( $\text{mg N kg}^{-1}$ )	$16.2 \pm 0.82$
Microbial biomass C ( $\text{g C kg}^{-1}$ )	$2.32 \pm 0.09$
Microbial biomass N ( $\text{mg N kg}^{-1}$ )	$186 \pm 2.60$
Extractable $\text{NO}_3^-$ ( $\text{mg N kg}^{-1}$ )	$0.05 \pm 0.01$
Extractable $\text{NH}_4^+$ ( $\text{mg N kg}^{-1}$ )	$1.47 \pm 0.15$
Extractable P ( $\text{mg P kg}^{-1}$ )	$6.56 \pm 1.21$
Exchangeable Na ( $\text{mg kg}^{-1}$ )	$13.4 \pm 1.74$
Exchangeable K ( $\text{mg kg}^{-1}$ )	$123 \pm 37.5$
Exchangeable Ca ( $\text{g kg}^{-1}$ )	$0.46 \pm 0.07$

The study was set up in a randomised block design, with 5 treatments ( $n = 3$ ), as shown in Appendix 2 (Fig. 1). The treatments were as follows: 1) control (no urine application), 2) large urine patch with a low N concentration ( $4 \text{ g N l}^{-1}$ ; 1 g of N applied in 250 ml covering  $500 \text{ cm}^2$ ; N loading rate =  $200 \text{ kg N ha}^{-1}$ ), 3) large urine patch with a high N concentration ( $16 \text{ g N l}^{-1}$ ; 4 g of N applied in 250 ml, covering  $500 \text{ cm}^2$ ; N loading rate =  $800 \text{ kg N ha}^{-1}$ ), 4) four

small urine patches with a low N concentration ( $4 \text{ g N l}^{-1}$ ;  $4 \times 0.25 \text{ g}$  of N applied in  $4 \times 62.5 \text{ ml}$ , covering  $125 \text{ cm}^2$  each; N loading rate =  $200 \text{ kg N ha}^{-1}$ ) and 5) four small urine patches with a high N concentration ( $16 \text{ g N l}^{-1}$ ;  $4 \times 1 \text{ g}$  of N applied in  $4 \times 62.5 \text{ ml}$ , covering  $125 \text{ cm}^2$  each; N loading rate =  $800 \text{ kg N ha}^{-1}$ ).

Duplicate plots were established, one to allow for chamber gas flux measurements and the other for soil sampling during the study. Artificial urine was applied by evenly pouring onto the soil, using a fixed template as an area guide. All urine patches were smaller than the chamber basal area ( $2500 \text{ cm}^2$ ), ensuring additional room inside the chamber for the urine patch diffusional area, which is important for accurately monitoring urine patch  $\text{N}_2\text{O}$  emissions (Marsden et al., 2016). Soil cores (0-5 cm) were taken from the area of immediate urine application, during the course of the study. Soil sampling was conducted four days prior to treatment application (for measurement of background soil characteristics), on the day of a urine application, and 2, 4, 7, 10, 14, 17, 21, 28 and 35 days following treatment application. The soil sampling was conducted more frequently at the beginning of the study (three times in the first week) in order to provide information on soil N transformation processes which occur relatively quickly following organic N additions to soil (e.g. urea hydrolysis and nitrification). Soil sampling reduced in frequency throughout the remainder of the study (twice per week in the second and third week, and once per week, thereafter). When soil cores were taken, they were timed to match manual flux measurements taken from the control plots. Soil cores were returned immediately to the laboratory, where post processing took place within 24 h of sampling. Prior to extraction, soils were homogenised by gently mixing within the sample bags.

Soil moisture was monitored within each individual chamber by using Acclima SDI-12 digital TDT® sensors (Acclima Inc., ID, USA), monitoring at a 0.5 h measurement frequency. The sensors were inserted diagonally through the urine patch, and were in situ 3 weeks prior to urine application. Rhizon suction samplers (2.5 mm diameter, 5 cm porous tube length, 12 cm

tubing length; Rhizosphere Research Products, Wageningen, Netherlands) were also inserted at an angle of 45°, into the centre of individual urine patches, 3 weeks prior to urine application. This allowed sampling of soil solution within the chamber in a non-destructive manner when the soil was wet enough for sample collection. Successful soil solution samples were collected 4 and 3 days prior to urine application, on the day of urine application and 2, 4, 7, 10, 14 and 17 days following urine application. Beyond this point soil conditions were too dry, so samples were not collected. Additionally, air temperature was monitored both inside and outside chambers using Thermochron iButtons® (iButtonLink, LLC, WI, USA) logging temperature every 1 h. A weather station was located near to the field site, where rainfall, soil (0-10 cm) and air temperature were monitored hourly.

#### *4.2.4 Pasture biomass and N content*

The pasture within chambers was cut 2 and 4 weeks following urine application, to a height of 5 cm. Samples were oven dried (80°C; 24 h), weighed for biomass, and ground before analysis of N content as described previously.

#### *4.2.5 Automated and manual greenhouse gas measurements*

Greenhouse gas fluxes were measured from twelve non-steady-state, non-through flow chambers connected to a mobile, automated high frequency measurement system (Queensland University of Technology, Institute for Future Environments, Brisbane, Australia) as detailed in Scheer et al. (2014). Stainless steel chamber bases were inserted into the ground (10 cm depth) for 4 weeks prior to treatment application, to ensure no effects from soil disturbance were observed during the study. Chambers (50 cm x 50 cm x 15 cm) were fixed to the bases, which opened and closed during sampling via pneumatic actuators. Chamber headspace samples were automatically pumped (ca. 200 ml min<sup>-1</sup>) from the chambers, through Teflon



tubing to the sampling unit, which housed a LI-COR LI-820 non-dispersive infrared gas analyser (LI-COR, St Joseph, MI, USA) to measure the CO<sub>2</sub> concentration, and a gas chromatograph (SRI 8610C, Torrance, USA), equipped with a <sup>63</sup>Ni electron capture detector (ECD) and flame ionization detector (FID) to measure N<sub>2</sub>O and CH<sub>4</sub> concentrations in the chamber headspace, respectively. The samples passed through an Ascarite (sodium hydroxide coated silica) column prior to entering the GC, in order to remove moisture and CO<sub>2</sub>, which was changed periodically in order to protect the ECD cell.

The sampling routine consisted of three blocks of four chambers, where each block of chambers close sequentially for a period of 1 h, while the other two sets are open allowing restoration of ambient conditions. During the 1 h closure period each chamber is sampled for 3 min, followed by a calibration standard (500 ppb N<sub>2</sub>O; 880 ppm CO<sub>2</sub>; 3 ppm CH<sub>4</sub>; ± 2 % of the certified value, BOC gases, Liverpool, UK). The 15 min cycle repeats four times, before the next set of chambers are sampled. One entire cycle takes 3 h to complete, allowing up to 8 flux measurements per 24 h period.

The automated system consisted of twelve chambers only, therefore, control treatments were sampled manually from static chambers ( $n = 3$ ), as N<sub>2</sub>O fluxes were expected to be minimal from this treatment. Manual chamber flux measurements were taken once per day, between 10 am and 12 noon (de Klein and Harvey, 2012) as close to daily as possible, for the duration of the experiment. The gas samples were taken by placing polypropylene upturned buckets (ca. 26 cm in height) onto collars (26 cm diameter), inserted to a depth of 10 cm. The chambers were fitted with a re-sealable vent, to allow pressure equalisation when placing chambers onto bases, and were fitted with Suba-Seals ® (Sigma, Gillingham, UK) to allow sampling of the headspace. Samples were taken with a syringe every 15 min, over the period of 1 h, to match the automated system. Samples were stored in pre evacuated 20 ml glass vials,

before being analysed on a Clarus 500 GC with a TurboMatrix headspace autoanalyser (Perkin Elmer, CT, USA).

#### 4.2.6 Statistical analysis

Cumulative N<sub>2</sub>O emissions were determined by integration using the trapezoidal rule. The resulting cumulative emissions were corrected for the chamber area assumed to be unaffected by urine application (i.e. total chamber area = 0.25 m<sup>2</sup>, urine patch area = 0.05 m<sup>2</sup> and control area unaffected by urine = 0.2 m<sup>2</sup>), by deducting the cumulative emissions arising from the control treatment over an area of 0.2 m<sup>2</sup>. Further calculations e.g. emission factors were based on the control-corrected values for cumulative emissions. To compare whether cumulative emissions or emission factors increased with increasing N loading rate, t-tests were conducted, comparing the results from treatments 2 with 3 and treatment 4 with treatment 5. In order to determine whether emissions are greater from small or large urine patches (with the same N loading rate) the same procedure was used, comparing treatment 2 with treatment 4 and treatment 3 with treatment 5. To test the third hypothesis that cumulative emissions would be higher from a low N concentration large urine patch (200 kg N ha<sup>-1</sup>) compared with a high N concentration small urine patch (800 kg N ha<sup>-1</sup>) with the same total amount of N applied, the results from treatment 5 were divided by 4 (assuming that cumulative emissions from each individual urine patch within the chamber were equal) and compared with the results from treatment 2. Differences in pasture biomass and foliar N content at 2 and 4 weeks following urine application were assessed via ANOVA with Tukey's post-hoc test, after assessing normality and homogeneity of variance.

To assess how the accuracy of cumulative N<sub>2</sub>O emissions from urine patches is affected by the frequency of sampling, we compared the high frequency N<sub>2</sub>O dataset with the N<sub>2</sub>O dataset containing (where available): 1) daily measurements between 10 a.m. and 2 p.m., 2) weekly measurements between 10 a.m. and 2 p.m., and 3) a sampling campaign where 3 flux


measurements were taken in the first week after urine application, 2 flux measurements in the second week after urine application and once a week thereafter. These different (infrequent sampling) regimes reflected those typically used in the literature for determining emissions from urine patches. Differences were reported as the mean across all replicates and treatments. All statistical analysis was conducted in Minitab 17.0 (Minitab Inc., State College, PA, USA).

## 4.3 Results

### 4.3.1 Effect of urine N content and patch size on soil C and N dynamics

The dynamics of total (free and exchangeable)  $\text{NH}_4^+$  and  $\text{NO}_3^-$  concentrations in soil can be seen in Fig. 4.1a and b, respectively, while the concentration of  $\text{NH}_4^+$  and  $\text{NO}_3^-$  in soil solution (extracted with Rhizon samplers within the chambers) is presented in Fig. 4.1c and d. During the second half of the field trial the lack of rainfall prevented the acquisition of soil water with the Rhizon samplers and thus, there are no results to present. Extractable  $\text{NH}_4^+$  increased up to 4 days following urine application in the treatments receiving a high N loading rate, and then followed a declining trend. As expected, extractable  $\text{NH}_4^+$  peaked at a lower concentration in the lower N containing treatments, and began to decline two days following urine application. There were no apparent differences in the urine-induced increase of soil  $\text{NH}_4^+$  concentration due to patch size. The  $\text{NO}_3^-$  concentration steadily increased in the soil extracts and soil solution, with lower amounts of  $\text{NO}_3^-$  produced in the low N containing treatments. Differences in  $\text{NO}_3^-$  concentration between patch sizes were evident, with less accumulation of  $\text{NO}_3^-$  in smaller patches, and with concentrations returning to background values faster than in large urine patches. In the high N containing large urine patch, the  $\text{NO}_3^-$  concentration was still higher than the control at the end of the experiment, and it cannot be ruled out that further  $\text{N}_2\text{O}$  emissions would have occurred after this time.


The total extractable dissolved N and organic C present in soil during the course of the study can be seen in Fig. 4.2a and b, respectively, while the soil solution total dissolved N and C can be seen in Fig. 4.2c and d, respectively. Concentrations of extractable and soil solution



**Figure 4.1** Mineral N dynamics following artificial sheep urine application of varying N concentration and patch size to a Eutric Cambisol, panel a) extractable soil ammonium-N panel b) extractable soil nitrate-N, panel c) soil solution nitrate-N and panel d) soil solution ammonium-N. Figure legend applies to all panels, symbols and error bars denote means  $\pm$  SEM, where  $n = 3$  for panels a) and b), however,  $n$  varies (1-3) for panels c) and d), due to variability in successful soil solution collection by Rhizon samplers.

N were high at the beginning of the study, due to the large amounts of dissolved organic N deposited into the soil within the urine. The N persisted in the soil for longer periods in the high N containing urine patches. The soil solution C was immediately high following urine application, and rapidly declined to control values. Dissolved organic C may have been higher

in the high N containing treatments, due to the increased proportion of urea in these treatments, and/or it may have been caused by a priming effect, which has been demonstrated for ruminant urine-influenced soils for both C (Lambie et al., 2013) and N (Di and Cameron, 2008).



**Figure 4.2** Dynamics of total extractable nitrogen and organic carbon (panels a and b, respectively) and soil solution nitrogen and dissolved organic carbon (panels c and d, respectively) following artificial sheep urine application (of varying patch size and N content) to a Eutric Cambisol. Figure legend applies to all panels, symbols and error bars denote means  $\pm$  SEM, where  $n = 3$  for panels a) and b), however,  $n$  varies (1-3) for panels c) and d), due to variability in successful soil solution collection by Rhizon samplers.

### 4.3.2 Pasture biomass and foliar N content

The pasture biomass and N content after 2 and 4 weeks following urine application can be seen in Table 4.2. As expected, the pasture biomass and foliar N content were generally larger in the urine treatments compared to the control (except pasture biomass 4 weeks after urine application). Increasing the N loading rate in a single large urine patch did not increase ( $p > 0.05$ ) pasture biomass 2 or 4 weeks after urine application, nor did it increase foliage N content in the first 2 weeks after urine application, however, foliage N did increase after 4 weeks ( $p < 0.05$ ).


**Table 4.2** Pasture biomass (g DM m<sup>-2</sup>) and foliar N content (%) at 2 and 4 weeks following artificial sheep urine application to a Eutric Cambisol. Values represent means  $\pm$  SEM ( $n = 3$ ), and letters indicate significant differences ( $p < 0.05$ ).

Treatment	2 weeks after urine application		4 weeks after urine application	
	Biomass (t DM ha <sup>-1</sup> )	Foliar N content (%)	Biomass (t DM ha <sup>-1</sup> )	Foliar N content (%)
Control	43.7 $\pm$ 3.8 a	2.96 $\pm$ 0.14 a	74.1 $\pm$ 5.7 a	2.18 $\pm$ 0.14 a
Low N concentration large urine patch	73.0 $\pm$ 2.9 b	4.05 $\pm$ 0.13 bc	114.6 $\pm$ 14.9 a	3.22 $\pm$ 0.06 b
High N concentration large urine patch	74.4 $\pm$ 2.1 b	4.32 $\pm$ 0.10 cd	127.7 $\pm$ 17.3 a	3.97 $\pm$ 0.19 c
Four low N concentration small urine patches	71.4 $\pm$ 1.6 b	3.82 $\pm$ 0.10 b	106.4 $\pm$ 1.1 a	3.05 $\pm$ 0.05 b
Four high N concentration small urine patches	92.0 $\pm$ 4.0 c	4.71 $\pm$ 0.03 d	107.3 $\pm$ 16.6 a	4.26 $\pm$ 0.16 c
ANOVA $P$ value	< 0.001	< 0.001	0.12	< 0.001

Increasing the N loading rate in four smaller urine patches increased pasture biomass ( $p < 0.01$ ) 2 weeks after urine application and increased foliar N content ( $p < 0.01$ ) on both sampling occasions. Patch size had no effect ( $p > 0.05$ ) on pasture biomass or foliar N content when applied at 200 kg N ha<sup>-1</sup>, either 2 or 4 weeks following urine deposition. At 800 kg N ha<sup>-1</sup>, the biomass was greater ( $p < 0.05$ ) in four small urine patches compared to one large urine patch after 2 weeks, however, no differences ( $p > 0.05$ ) were observed after 4 weeks. No differences ( $p > 0.05$ ) were observed between the foliar N content of four small patches compared to one large urine patch, when urine was applied at 800 kg N ha<sup>-1</sup>.

#### *4.3.3 N<sub>2</sub>O fluxes from artificial urine-influenced soil*

The results for the N<sub>2</sub>O emissions over the course of the experiment can be seen in Fig. 4.3b, alongside the weather station data in Fig. 4.3a and the soil water-filled pore space (WFPS; calculated from the SDI-12 soil moisture sensors) in Fig. 4.3c. The initial peak in N<sub>2</sub>O emissions were similar across all treatments where urine was applied, indicating initially the N content was not limiting N<sub>2</sub>O production. Following the maxima of this initial peak, the two high N treatments begin to branch off from the low N treatments, and generally remained higher than the low N containing treatments for ca. 2 weeks. The major peaks in N<sub>2</sub>O emissions occurred just after rainfall events (Fig. 4.3a and b). The largest N<sub>2</sub>O peaks were observed for the high N containing small urine patches, which peaked at around 150 µg N<sub>2</sub>O-N m<sup>-2</sup> h<sup>-1</sup>.




**Figure 4.3** Weather station data (panel a), nitrous oxide emissions (panel b) and water-filled pore space (WFPS; panel c) following application of artificial sheep urine of varying N concentration and patch size to a Eutric Cambisol. Figure legends apply to each respective panel, for both panels b and c, where the traces represent the treatment mean ( $n = 3$ ).

#### 4.3.4 Cumulative N<sub>2</sub>O emissions

Cumulative N<sub>2</sub>O emissions (mg N<sub>2</sub>O-N 0.05 m<sup>-2</sup>) over the 39 day measuring period (Fig. 4.4) followed the numerical trend high N small urine patches ( $6.75 \pm 0.54$ ) > high N large patch ( $4.01 \pm 0.57$ ) > low N small patches ( $2.56 \pm 2.40$ ) > low N large patch ( $1.36 \pm 0.44$ ) > control ( $0.36 \pm 0.07$ ). The variability in cumulative N<sub>2</sub>O emissions was greater in the low N small urine patches treatment, which arose from one ‘high emitting’ replicate.





**Figure 4.4** Cumulative N<sub>2</sub>O-N emissions over the duration of the field trial (40 days), where artificial urine of varying N content and patch size was applied to a Eutric Cambisol. Bars represent means  $\pm$  SEM ( $n = 3$ ).

Increasing the N loading rate from 200 to 800 kg N ha<sup>-1</sup> within a large urine patch increased ( $p < 0.05$ ) cumulative N<sub>2</sub>O emissions by a factor of 3, from  $1.36 \pm 0.44$  to  $4.01 \pm 0.57$  mg N<sub>2</sub>O-N 0.05 m<sup>-2</sup> over 39 days. No difference ( $p > 0.05$ ) was observed between cumulative N<sub>2</sub>O emissions when increasing N concentration in the four small sized urine patches. The non-significant effect of increasing N concentration on cumulative N<sub>2</sub>O emissions in the small urine patches is likely to be the result of the high N<sub>2</sub>O emitting replicate; if this replicate is removed from the analysis the difference then becomes significant ( $p < 0.01$ ), although the potential for such high emissions cannot be ruled out from smaller urine patches.

Increasing the number of replicates per treatment is recommended in order to capture spatial variability of N<sub>2</sub>O emissions under field conditions.

The second hypothesis was to determine whether cumulative N<sub>2</sub>O emissions were higher from four small compared to one large urine patch (with the same N loading rate). No effect of patch size ( $p > 0.05$ ) was observed between the cumulative emissions at the lower N loading rate (200 kg N ha<sup>-1</sup>), regardless of whether the high-emitting replicate was included or removed. However, under a higher N loading rate (800 kg N ha<sup>-1</sup>), cumulative N<sub>2</sub>O emissions were greater ( $p < 0.05$ ) from the four small patches ( $6.75 \pm 0.54$  mg N<sub>2</sub>O-N 0.05 m<sup>-2</sup> over 40 days) compared to a single large urine patch ( $4.01 \pm 0.57$  mg N<sub>2</sub>O-N 0.05 m<sup>-2</sup> over 40 days).

The third hypothesis was that cumulative emissions would be higher from a large urine patch with a low N concentration, as opposed to a small urine patch with a high N concentration (manipulating N loading rate, but keeping total N applied constant). The cumulative emissions from a high N concentration single small urine patch was  $1.69 \pm 0.14$  mg N<sub>2</sub>O-N 0.05 m<sup>-2</sup>, over the duration of the study, where no differences were found ( $p > 0.05$ ) compared to the low N concentration large urine patch, at  $1.36 \pm 0.44$  mg N<sub>2</sub>O-N 0.05 m<sup>-2</sup>.

#### 4.3.5 Emission factors

The N<sub>2</sub>O emission factor for the low N concentration large urine patch over 39 days was  $0.10 \pm 0.05\%$  of the N applied and  $0.09 \pm 0.02\%$  from the high N concentration large urine patch. The emission factors from the four low N concentration small urine patches was  $0.22 \pm 0.24\%$  of the applied N, where one replicate had the highest emission factor of 0.69%, highlighting the large variability observed in this treatment. The N<sub>2</sub>O emission factor from the four high N concentration small urine patches was  $0.16 \pm 0.01\%$  of the total N applied and the emission factor from the high N concentration large urine patch was  $0.09 \pm 0.02\%$  of the N applied. No effect of patch size on N<sub>2</sub>O emission factors was observed in the low N

concentration treatments, however, the emission factor was significantly greater ( $p < 0.05$ ) from the four small urine patches compared to a single large urine patch, at  $800 \text{ kg N ha}^{-1}$ . No significant differences in  $\text{N}_2\text{O}$  emission factors were observed when increasing N concentration from  $4\text{--}16 \text{ g N l}^{-1}$  either from four small patches, or from a single large patch.

#### *4.3.6 Effect of reduced sampling frequency on cumulative $\text{N}_2\text{O}$ emissions*

When sampling frequency was reduced to one flux measurement per day (compared to 8 flux measurements per day), between the hours of 10 a.m. and 2 p.m., the mean difference in cumulative emissions was within  $1 \pm 6\%$  of the high frequency measurement value, although differences between individual replicates ranged from an underestimate of 39% and an overestimate of 50% of the value obtained by taking 8 flux measurements per day. Sampling  $\text{N}_2\text{O}$  fluxes once a week resulted in a mean underestimation of the cumulative emissions of  $54 \pm 24\%$ , and individual replicates ranged from an underestimate of 268% and an overestimate of 37%. The last sampling regime comparison was to measure  $\text{N}_2\text{O}$  fluxes three times in the first week following urine application, twice in the second week, and once per week thereafter. Across all treatments and replicates, this sampling regime would have underestimated emissions by  $22 \pm 10\%$  of the high frequency value, and individual replicates ranged from an underestimate of 90% to an overestimate of 38%.

## **4.4 Discussion**

When urine was applied at the same N loading rate,  $\text{NO}_3^-$  concentrations did not peak as high, and returned to control values faster in four smaller urine patches compared to a single large patch. The observed effect could have been due to the larger wetted perimeter created by the four small urine patches (179 cm) in comparison to one large urine patch (89 cm). This may have allowed increased plant access to the nutrient rich patch via roots and stolons, which may

explain the greater plant biomass and foliar N content when increasing urine N loading rate in smaller urine patches. Diffusion away from the urine patch edge has been identified as an important mechanism for N processing and transformations in urine patches of different sizes (Orwin et al., 2009). The difference in patch sizes may have allowed a greater lateral, as opposed to vertical, diffusion of soluble N and C into surrounding soil, potentially influencing differing proportions of nitrifying and denitrifying microorganisms. A greater diffusion of  $\text{NO}_3^-$  beyond the area of immediate urine application may have also resulted in the lower peak of  $\text{NO}_3^-$  found in smaller urine patches, as soil cores were only taken where urine had been directly applied, and Rhizon samples were taken at the centre of the urine patch. The soil mineral N concentration remained higher for a longer period with increasing urine N concentration, which means N losses can occur under partially differing environmental conditions from the high N treatment compared to the lower N treatment (Dijkstra et al., 2013).

Concentrations of  $\text{NO}_3^-$ ,  $\text{NH}_4^+$ , total N and DOC in the urine patches (measured in both the soil extracts and the soil solution) tended to vary widely, having large standard errors. This large spatial variability in factors which can drive  $\text{N}_2\text{O}$  emissions, is likely to have contributed to the large observed variation in  $\text{N}_2\text{O}$  emissions in this study. This could also be problematic when comparing urine patch soil properties from duplicated plots with  $\text{N}_2\text{O}$  emissions from chambers, as processes may be occurring at different rates in the different locations. We suggest the development of non-destructive sampling techniques for monitoring soil conditions within chambers to ensure monitored soil properties accurately reflect conditions within chambers.

In this study, increasing urine N loading rate from  $200 \text{ kg N ha}^{-1}$  to  $800 \text{ kg N ha}^{-1}$  generally increased cumulative  $\text{N}_2\text{O}$  emissions, but the emission factors were not significantly different. This is similar to the finding of Selbie et al. (2014), who found a curvilinear increase in cumulative  $\text{N}_2\text{O}$  emissions upon increasing N loading rate. Increasing urine N loading rate,

however, had no effect on the fraction of N<sub>2</sub>O produced under laboratory (van Groenigen et al., 2005a) and field conditions (van Groenigen, 2005b; Dai et al., 2013; Luo et al., 2013). However, urine derived N<sub>2</sub>O emission factors have also been shown to decrease (Selbie et al., 2014) or increase alongside N loading rate (Singh et al., 2009). There are contradictory results for the response of urine patch N<sub>2</sub>O emission factors to increasing N loading rates and results to date are contradictory and uncertain (Luo et al., 2013; Selbie et al., 2014).

When manipulating urine patch size and N loading rate, but maintaining the same total amount of N applied, no differences were found in cumulative N<sub>2</sub>O emissions or emission factors, which is similar to the findings of van Groenigen et al. (2005a). Cumulative N<sub>2</sub>O emissions and emission factors, however, were significantly greater from four small urine patches compared to one large urine patch, when applied at 800 kg N ha<sup>-1</sup>. It is difficult to determine a management strategy which would reduce N<sub>2</sub>O emissions based on the results of this treatment, as it assumes a difference in the frequency of urine events, but with the same N concentration and total volume of urine excreted. In reality, ruminant urine N concentration and volume varies widely between and within days, and between individuals of the same species (Betteridge et al., 1986; Hoogendoorn et al., 2010; Betteridge et al., 2013). Nevertheless, based on the results of this study and the sheep urine patch sizes used here, it would seem that sheep which urinate more frequently in smaller volumes would emit more N<sub>2</sub>O than a sheep urinating less frequently in larger volumes, given the same high urinary N concentration. Further work is required to determine how this effects other N loss processes such as NO<sub>3</sub><sup>-</sup> leaching and NH<sub>3</sub> volatilisation, as given the range of patch sizes excreted by cattle, increasing patch size has been shown to logarithmically increase NO<sub>3</sub><sup>-</sup> leaching (Li et al., 2012).

It has been recognised that more information is required for values of ruminant urine volume, urine N concentration and urination frequency, where the development of sensor based

technology which can measure all three of these values (e.g. Betteridge et al., 2013), will be important for improving models to predict N losses from grazed grassland. An increase in our knowledge of how ruminant urine N concentration is affected by dietary factors will also be important, e.g. Pacheco et al. (2010) found urinary N concentration of cattle urine to be related to the moisture content, dietary cation-anion difference and soluble sugar content of the forage. Increasing knowledge in these areas will also assist in a better understanding on the effect of differing management and mitigation strategies to reduce N losses (e.g. feed type, grazing intensity, and inclusion of diuretics in the diet).

Increasing urine volume e.g. by the inclusion of NaCl in the diet, has been suggested as a way to reduce N losses as it would reduce N loading rate per urination, potentially promoting a better dispersion of N across the pasture (Costall and Betteridge, 2010; Pacheco et al., 2010), although this will depend on livestock movement and behaviour. The results of our study, however, have shown that N<sub>2</sub>O emission were no different in patches of differing N loading rates, for the same given amount of N applied. Therefore, for a reduction in N<sub>2</sub>O emissions from sheep urine patches, it is suggested that a reduction in dietary N would need to occur concurrently with any mitigation strategies based on increasing total urine volume.

Both Ledgard et al. (2007) and Liu and Zhou (2014) show that increasing NaCl in the diet of cattle and sheep, respectively, results in increased urine volume and frequency, but the average volume of individual urine events remained similar. Both Li et al. (2012) and Betteridge et al. (2013) have shown that considering mean values (rather than varied values) for cattle urine volume, frequency and N concentration causes differences of ca. 5-10% in modelled NO<sub>3</sub><sup>-</sup> leaching losses. The use of mean values for urine volume and N concentration are also often used in plot based studies of urine patch N<sub>2</sub>O emissions, which may also cause inaccuracies when upscaling such emissions.

Across all urine treatments, N<sub>2</sub>O emission factors were on average ten times lower than that of the 1% default IPCC emission factor for sheep excreta (IPCC, 2006). In a large study based on numerous field trials conducted across New Zealand, the mean sheep urine patch emission factors from lowland pasture soils was also found to be lower than the default IPCC emission factor for excreted N at pasture, at 0.55% of the applied N (Kelliher et al., 2014). The low emission factors found in this study may have been caused by the relatively dry soil conditions which prevailed during the study, or due to the short period of measurement (i.e. a year is the normal requirement for calculation of emission factors). Low emission factors have been previously observed above sheep grazed pastures under dry summer months, with maximum emissions occurring under cooler and wetter soil conditions (Allen et al., 1996; Saggar et al., 2007). Rainfall and soil drainage class have been identified as key variables influencing N<sub>2</sub>O emission factors from urine returns (de Klein et al., 2003), and a consideration of environmental and management factors (which are not considered in the IPCC Tier 1 approach) will undoubtedly be important for better constraining N<sub>2</sub>O emission estimates from grazed soils.

The values for emission factors in this study should be considered with care, as they were derived from artificial urine and overall emissions were fairly low and variable, which may have masked some effects. It is suggested that similar studies should be conducted under differing environmental conditions to determine if the effects hold true. Real urine is generally advocated for the calculation of emission factors (deKlein et al., 2003), however, the use of artificial urine was beneficial in this study as it allowed the manipulation of N concentration. The artificial urine contained the recommended constituents for adequate representations of urine for N<sub>2</sub>O emissions (Kool et al., 2006) and the proportion of urea was increased to manipulate the N content, which is consistent with an increase in the proportion of urea

excreted by ruminants occurring as a result of increased dietary crude protein (Dijkstra et al., 2013).

Our results show that daily sampling of gases between 10 a.m. and 2 p.m. resulted in a good approximation of mean N<sub>2</sub>O emissions. Sampling more often directly after urine application and tapering the sampling towards the end of the experiment resulted in a better approximation of emissions than weekly sampling, although both sampling strategies had a high variability and resulted in a mean underestimation of emissions. Additional sources of error which may occur during manual gas sampling, which were not assessed here, include errors associated with vial evacuation, filling and storage prior to analysis of the gases. Using an automated system, with samples being immediately pumped to a GC, eliminates the potential for such errors.

#### **4.5 Conclusions**

Cumulative N<sub>2</sub>O emissions generally increased, but emission factors remained similar, with increasing N loading rate (200–800 kg N ha<sup>-1</sup>). Emission factors were similar from a low N concentration large urine patch (200 kg N ha<sup>-1</sup>) compared to a high N concentration small urine patch (800 kg N ha<sup>-1</sup>), when applying the same total amount of N. However, emissions were larger from four smaller patches compared to one large urine patch under the same N loading rate (800 kg N ha<sup>-1</sup>). These conclusions suggest a reduction in overall dietary N would need to occur alongside any mitigation strategies which manipulate urine volume, in order to be effective at reducing N<sub>2</sub>O emissions from sheep urine patches. For the same total volume and concentration of N excreted in the urine, sheep which urinate little and often may be causing greater emissions than sheep which urinate less frequently with a larger volume and patch size. With this in mind, further research is required to gather more data on typical sheep urine volume (both individual urine events and daily volumes), frequency and N



concentrations. Such parameters may vary due to sheep breed, edaphic and climatic conditions, diet and management or mitigation practices (e.g. inclusion of salt in diet). Improving our knowledge on the links between the plant-animal-soil–atmosphere systems will be important for determining more accurate emission estimates from sheep grazed grasslands, and for determining the efficacy of N<sub>2</sub>O mitigation strategies.

#### **4.6 Acknowledgements**

We would like to thank Christian Brunk at QUT for his assistance and support during the set up phase of the automated GHG system, Lara Jane Pritchard for assistance with field sampling and John Evans, Mark Hughes and Llinos Hughes for their assistance setting up the field trial. KAM would also like to thank the School of Environment, Natural Resources and Geography, Bangor University, for sponsoring this study. This work was supported by the Natural Environment Research Council under grant award NE/MO15351/1.

## 4.7 References

- Allen AG, Jarvis SC, Headon DM (1996). Nitrous oxide emissions from soils due to inputs of nitrogen from excreta return by livestock on grazed grassland in the U.K. *Soil Biology and Biochemistry* **28**, 597-607.
- Ball DF (1964). Loss-on-ignition as an estimate of organic matter and organic carbon in non-calcareous soils. *Journal of Soil Science* **15**, 84-92.
- Betteridge K, Andrews WGK, Sedcole JR (1986). Intake and excretion of nitrogen, potassium and phosphorus by grazing steers. *The Journal of Agricultural Science* **106**, 393-404.
- Betteridge K, Costall D, Balladur S, Upsdell M, Umemura K (2010a). Urine distribution and grazing behaviour of female sheep and cattle grazing a sheep New Zealand hill pasture. *Animal Production Science* **50**, 624-629.
- Betteridge K, Costall DA, Li FY, Luo D, Ganesh S (2013). Why we need to know what and where cows are urinating – a urine sensor to improve nitrogen models. In *Proceedings of the New Zealand Grassland Association* **75**, 119-124.
- Betteridge K, Hoogendoorn C, Costall D, Carter M, Griffiths W (2010b). Sensors for detecting and logging spatial distribution of urine patches of grazing female sheep and cattle. *Computers and Electronics in Agriculture* **73**, 66-73.
- Bristow A, Whitehead DC, Cockburn JE (1992). Nitrogenous constituents in the urine of cattle, sheep and goats. *Journal of the Science of Food and Agriculture* **59**, 387-394.
- Costall K, Betteridge K (2010). Methods of delivering salt to cattle. *Proceedings of the New Zealand Society of Animal Production* **70**, 296-298.
- Dai Y, Di HJ, Cameron KC, He J-Z (2013). Effects of nitrogen application rate and a nitrification inhibitor dicyandiamide on ammonia oxidizers and N<sub>2</sub>O emissions in a grazed pasture soil. *Science of the Total Environment* **465**, 125-135.

- de Klein CAM, Barton L, Sherlock RR, Li Z, Littlejohn RP (2003). Estimating a nitrous oxide emission factor for animal urine from some New Zealand pasture soils. *Australian Journal of Soil Research* **41**, 381-399.
- De Klein CAM, Harvey, MJ. 2012. Nitrous Oxide Chamber Methodology Guidelines. Global Research Alliance on Agricultural Greenhouse Gases, Ministry for Primary Industries, New Zealand.
- de Klein CAM, Luo J, Woodward KB, Styles T, Wise B, Lindsey S, Cox N (2014). The effect of nitrogen concentration in synthetic cattle urine on nitrous oxide emissions. *Agriculture, Ecosystems and Environment* **188**, 85-92.
- Di HJ, Cameron KC (2008). Sources of nitrous oxide from <sup>15</sup>N-labelled animal urine and urea fertiliser with and without a nitrification inhibitor, dicyandiamide (DCD). *Australian Journal of Soil Research* **46**, 76-82.
- Dijkstra J, Oenema O, van Groenigen JW, Spek JW, van Vuuren AM, Bannink A (2013). Diet effects on urine composition of cattle and N<sub>2</sub>O emissions. *Animal* **7**(S2), 292-302.
- Doak BW (1952). Some chemical changes in the nitrogenous constituents of urine when voided on pasture. *The Journal of Agricultural Science* **42**, 162-171.
- Haynes RJ, Williams PH (1993). Nutrient cycling and soil fertility in the grazed pasture ecosystem. *Advances in Agronomy* **49**, 119-199.
- Hoogendoorn CJ, Betteridge K, Costall DA, Ledgard SF (2010). Nitrogen concentration in the urine of cattle, sheep and deer grazing a common ryegrass/cocksfoot/white clover pasture. *New Zealand Journal of Agricultural Research* **53**, 235-243.
- IPCC (2006). IPCC Guidelines for National Greenhouse Gas Inventories Volume 4 - Agriculture, Forestry and Other Land Use. <http://www.ipcc-nggip.iges.or.jp/public/2006gl/vol4.html>

- Jones DL, Willett VB (2006). Experimental evaluation methods to quantify dissolved organic nitrogen (DON) and dissolved organic carbon (DOC) in soil. *Soil Biology and Biochemistry* **5**, 991-999.
- Kelliher FM, Cox N, van der Weerden TJ, de Klein CAM, Luo J, Cameron KC, Di HJ, Giltrap D, Rys G (2014). Statistical analysis of nitrous oxide emission factors from pastoral agriculture field trials conducted in New Zealand. *Environmental Pollution* **186**, 63-66.
- Kool DM, Hoffland E, Abrahamse SPA, van Groenigen JW (2006). What artificial urine composition is adequate for simulating soil N<sub>2</sub>O fluxes and mineral N dynamics? *Soil Biology and Biochemistry* **38**, 1757-1763.
- Lambie SM, Schipper LA, Balks MR, Baisden WT (2013). Priming of soil decomposition leads to losses of carbon in soil treated with cow urine. *Soil Research* **51**, 513-520.
- Lantinga EA, Keuning JA, Groenwold J, Deenen PJAG (1987). Distribution of excreted nitrogen by grazing cattle and its effects on sward quality, herbage production and utilization. In *Animal Manure on Grassland and Fodder Crops, Fertilizer or Waste?* pp 103-117, Springer, Netherlands.
- Ledgard, SF, Menneer JC, Dexter MM, Kear MJ, Lindsey S, Peters JS, Pacheco DS (2008). A novel concept to reduce nitrogen losses from grazed pastures by administering soil nitrogen process inhibitors to ruminant animals: a study with sheep. *Agriculture, Ecosystems and Environment* **125**, 148-158.
- Ledgard SF, Welten BG, Menneer JC, Betteridge K, Crush JR, Barton MD (2007). New nitrogen mitigation technologies for evaluation in the Lake Taupo catchment. *Proceedings of the New Zealand Grassland Association* **69**, 117-121.
- Li FY, Batteridge K, Cichota R, Hoogendoorn CJ, Jolly BH (2012). Effects of nitrogen load variation in animal urination events on nitrogen leaching from grazed pastures. *Agriculture, Ecosystems and Environment* **159**, 81-89.

- Liu H, Zhou D (2014). Mitigation of ammonia and nitrous oxide emissions from pasture treated with urine of sheep fed diets supplemented with NaCl. *Animal Feed Science and Technology* **192**, 39-47.
- Lucas SD, Jones DL (2006). Biodegradation of estrone and 17  $\beta$ -estradiol in grassland soils amended with animal wastes. *Soil Biology and Biochemistry* **38**, 2803-2815.
- Luo J, Hoogendoorn C, van der Weerden T, Saggar S, de Klein C, Giltrap D, Rollo M, Rys G (2013). Nitrous oxide emissions from grazed hill land in New Zealand. *Agriculture, Ecosystems and Environment* **181**, 58-68.
- Marsden KA, Jones DL, Chadwick DR (2016). The urine patch diffusional area: an important N<sub>2</sub>O source? *Soil Biology and Biochemistry* **92**, 161-170.
- Miranda KM, Epsey MG, Wink DA (2001). A rapid, simple, spectrophotometric method for simultaneous detection of nitrate and nitrite. *Nitric Oxide* **5**, 62-71.
- Oenema O, Velthof GL, Yamulki S, Jarvis SC (1997). Nitrous oxide emissions from grazed grassland. *Soil Use and Management* **13**, 288-295.
- Orwin KH, Bertram JE, Clough TJ, Condon LM, Sherlock RR, O'Callaghan M (2009). Short-term consequences of spatial heterogeneity in soil nitrogen concentrations caused by urine patches of different sizes. *Applied Soil Ecology* **42**, 271-278.
- Pacheco D, Lowe K, Hickey M, Burke JL, Cosgrove GP (2010). Seasonal and dietary effects on the concentration of urinary N from grazing cows. *Proceedings of the 4<sup>th</sup> Australasian Science Symposium*, 68-73.
- Saggar S, Hedley CB, Giltrap DL, Lambie SM (2007). Measured and modelled estimates of nitrous oxide emission and methane consumption from a sheep-grazed pasture. *Agriculture, Ecosystems and Environment* **122**, 357-365.

- Scheer C, Rowlings DW, Firrel M, Deuter P, Morris S, Grace PR (2014). Impact of nitrification inhibitor (DMPP) on soil nitrous oxide emissions from an intensive broccoli production system in sub-tropical Australia. *Soil Biology and Biochemistry* **77**, 243-251.
- Selbie DR, Buckthought LE, Shepherd MA (2015). The challenge of the urine patch for managing nitrogen in grazed pasture systems. *Advances in Agronomy* **129**, 229-292.
- Selbie DR, Cameron KC, Di HJ, Moir JL, Lanigan JG, Richards KG (2014). The effect of urinary nitrogen loading rate and a nitrification inhibitor on nitrous oxide emissions from a temperate grassland soil. *Journal of Agricultural Science* **152**, S159-S171.
- Singh J., Saggarr S., Bolan NS (2009). Influence of dicyandiamide on nitrogen transformation and losses in cow-urine-amended soil cores from grazed pasture. *Animal Production Science* **49**, 253-261.
- van Groenigen JW, Kuikman PJ, de Groot WJM, Velthof GL (2005a). Nitrous oxide emission from urine-treated soil as influenced by urine composition and soil physical conditions. *Soil Biology and Biochemistry* **37**, 463-473.
- van Groenigen JW, Velthof GL, van der Bolt FJE, Vos A, Kuikman PJ (2005b). Seasonal variation in N<sub>2</sub>O emissions from urine patches: effects of urine concentration, compaction and dung. *Plant and Soil* **273**, 15-27.
- Voroney RP, Brookes PC, Beyaert RP (2008). Soil microbial biomass C, N, P and S. In: Carter, M.R and Gregorich, E.G., eds. *Soil sampling and methods of analysis*, 2<sup>nd</sup> edn. Boca Raton, FL, USA: CRC Press, pp. 637-651.
- Williams PH, Haynes RJ (1994). Comparison of initial wetting pattern, nutrient concentrations in soil solution and the fate of <sup>15</sup>N-labelled urine in sheep and cattle urine patch areas of pasture soil. *Plant and Soil* **162**, 49-59.



---

## Chapter 5

---

### **The mobility of nitrification inhibitors under simulated ruminant urine deposition and rainfall: a comparison between DCD and DMPP**

Karina A. Marsden<sup>\*a</sup>, Antonio J. Marín-Martínez<sup>b</sup>, Antonio Vallejo<sup>c</sup>, Paul W. Hill<sup>a</sup>, Davey L. Jones<sup>a</sup> & David R. Chadwick<sup>a</sup>.

This manuscript has been accepted for publication in *Biology and Fertility of Soils*.

Marsden, K.A., Marín-Martínez, A.J., Vallejo, A., Hill, P.W., Jones, D.L., Chadwick, D.R. 2016. The mobility of nitrification inhibitors under simulated urine deposition and rainfall: a comparison between DCD and DMPP. *Biology and Fertility of Soils*, In Press.

KAM, AJM-M, AV, PWH, DLJ and DRC designed and conceived the experiment; AV conducted preliminary trials; KAM and AJM-M conducted experimental work and KAM analysed results and prepared the manuscript.



## Abstract

Urine patches within pasture soils are hotspots for N cycling and losses, where nitrification inhibitors (NI) offer a means of reducing such losses. Within urine influenced soil, more research has been conducted on the effectiveness of dicyandiamide (DCD) than 3,4-dimethylpyrazole phosphate (DMPP) as NI. Differences in the efficacy of these NI are often ascribed to a greater mobility of DCD, which may lead to spatial separation from  $\text{NH}_4^+$  and nitrifying microorganisms. We tested the mobility of  $^{14}\text{C}$ -labelled DCD and DMPP relative to sheep urine derived  $\text{NH}_4^+$  in soil columns of contrasting texture and organic matter content, following simulated rainfall. We also assessed factors influencing the vertical mobility of these NI in soils, including solubility, sorption/desorption processes and microbial degradation and uptake. Following 40 mm rainfall, without the presence of sheep urine, the distribution of both NI were similar in the soil columns, however, there was a greater retention of DCD compared to DMPP in the top 1 cm. Both NI appeared to co-locate well with urine-derived  $\text{NH}_4^+$ , and the presence of sheep urine altered the leaching profile of the NI (compared to rainfall application alone), but this effect was inhibitor and soil-type dependent. A greater sorption to the soil matrix was observed for DCD in comparison to DMPP in all three studied soils, and the presence of urine generally increased desorption processes. Of the NI applied to the soil columns, 18-66% was taken up within 30 min by the microbial community. However, only small amounts (<1%) were mineralized during this period. In conclusion, due to the greater adsorption of DCD as opposed to DMPP and similarity in the degree of co-location of both NI with urine  $\text{NH}_4^+$ , the results of this study suggest that differences in microbial uptake and degradation may be more important parameters for explaining differences in the efficacy of reducing nitrification. Further work is required to determine the comparative efficacy of both NI in reducing nitrification rates under field conditions in a range of soil types and environmental conditions.

**Key Words:** grazed grassland; livestock ecosystems; nitrogen use efficiency; nutrient dynamics

## 5.1 Introduction

In pasture soils, high loadings of nitrogen (N) are deposited within ruminant urine patches and these sites are particularly vulnerable to losses of N to the environment. Typically, 20% of deposited urinary-N is leached as  $\text{NO}_3^-$ , 13% is volatilised as  $\text{NH}_3$  and 2% is emitted as the greenhouse gas  $\text{N}_2\text{O}$  (Selbie et al. 2015). While  $\text{N}_2\text{O}$  constitutes a small agronomic loss in terms of magnitude of N, having nearly 300 times the global warming potential of  $\text{CO}_2$  (IPCC 2007), it accounts for 46% of agricultural greenhouse gas emissions (Smith et al. 2007). The agricultural sector will need to contribute to decreasing emissions (Misselbrook et al. 2014) in order to achieve targets (80% reduction from 1990 baseline levels by 2050) set by the UK, and other governments. Reducing N loss via  $\text{NO}_3^-$  leaching and improving N use efficiency would translate to a direct economic benefit for farmers and reducing  $\text{N}_2\text{O}$  emissions from grasslands could contribute to decreasing emissions from the livestock sector.

Nitrification inhibitors (NI) are a potential mitigation strategy which can reduce losses of N from urine patches deposited to grassland soils (Di and Cameron 2012; Ledgard et al. 2014; Luo et al. 2015). By delaying the conversion of  $\text{NH}_4^+$  to  $\text{NO}_3^-$ , the opportunity for plant acquisition, immobilization, fixation and adsorption of  $\text{NH}_4^+$  is increased (Di and Cameron 2007). This can potentially reduce emissions of  $\text{N}_2\text{O}$  from both nitrification and denitrification (Gilsanz et al., 2016), where nitrification has been shown to be the dominant  $\text{N}_2\text{O}$  producing process in soils with a WFPS of 35-60%, but at a higher WFPS (70%), denitrification becomes the dominant  $\text{N}_2\text{O}$  producing process (Bateman and Baggs 2005). By the same processes, NI can also reduce the amount of  $\text{NO}_3^-$  available for leaching (Di and Cameron 2004).

Two of the most widely used NI are dicyandiamide (DCD) and dimethylpyrazole-phosphate (DMPP) (Liu et al. 2013). DCD blocks the electron transport chain in the cytochrome of ammonia monooxygenase (AMO), whereas DMPP binds indiscriminately to the membrane-bound AMO (Chaves et al. 2006; Fiencke and Bocke 2006; Benckiser et al. 2013), delaying the first and rate-limiting step of nitrification (the oxidation of  $\text{NH}_4^+$  to  $\text{NO}_2^-$ ; Zerulla et al. 2001). In comparison to DCD, DMPP has been shown to be less phytotoxic, and lower application rates are required (Wissemeier et al. 2001; Zerulla et al. 2001; Di and Cameron 2012). Both NI have demonstrable efficacy in reducing N losses from fertilizer applications (Weiske et al. 2001; Liu et al. 2013) and livestock slurry (Fangueiro et al. 2009; Pereira et al. 2010), however, DCD applications to urine patches have been more widely researched (e.g. Di and Cameron 2003; O'Callaghan et al. 2010; de Klein et al. 2011) in comparison to DMPP (e.g. Di and Cameron 2011; Di and Cameron 2012). Some authors have found a difference between the efficacy of DCD and DMPP e.g. Weiske et al. (2001) found DMPP to reduce  $\text{N}_2\text{O}$  emissions from fertilizer by an average of 49%, whereas DCD reduced emissions by an average of 26%. Di and Cameron (2012), however, found that DCD and DMPP reduced  $\text{N}_2\text{O}$  emissions by a similar amount from cattle urine (62 and 66% reduction, respectively). The efficacy of NI in reducing  $\text{N}_2\text{O}$  emissions and  $\text{NO}_3^-$  leaching can vary widely. In a laboratory study of nine contrasting UK soils, the efficacy of DCD to inhibit  $\text{NH}_4^+$  oxidation, net  $\text{NO}_3^-$  production and emissions of  $\text{N}_2\text{O}$  was lower in soils of high temperature, clay content and organic matter content (McGeough et al. 2016). Differences in efficacy have been attributed to a lower mobility of DMPP in comparison to DCD, due to a greater sorption of DMPP (Wissemeier et al. 2001; Zerulla et al. 2001; Di and Cameron 2012). Having a high mobility may lead to the spatial separation of NI from  $\text{NH}_4^+$  and nitrifying microorganisms (Ruser and Schulz 2015).

Physicochemical characteristics which can influence mobility within soil include solubility and sorption/desorption processes (Carrillo-González et al. 2006). The greater the solubility of a chemical in water, the greater the potential for vertical transport. The sorption/desorption of chemicals within the soil is mainly influenced by the organic matter content of soil, where charges associated with the chemical influences the types of bond established. DCD is a net neutrally charged molecule and binding takes place on the surface of organic matter, through hydrogen bonding of the  $-NH_2$  and  $=NH$  functional groups to negative carboxyl groups of organic matter (Zhang et al. 2004; Singh et al. 2008; Shepherd et al. 2012). Conversely, DMPP is positively charged and adsorption is thought to occur to the mineral fraction of soils, such as clays and silts (Barth et al. 2001, 2008). As chemicals are transported through soil, further biological transformations may influence the quantity of substance available for vertical movement e.g. microbial uptake and mineralization. As DMPP is a heterocyclic compound, it is not readily degradable, resulting in a slower degradation rates in soil in comparison to DCD (Weiske et al. 2001; Chaves et al. 2006). In the specific conditions of urine influenced soils, the high concentration of  $NH_4^+$ ,  $K^+$  and other cations, may saturate cation exchange sites leading to further movement of NI down the soil profile.

The objective of this study was to obtain information on how the combination of NI characteristics and soil conditions can affect the mobility and co-location of  $NH_4^+$  and NI in soils, with ruminant urine as the source of  $NH_4^+$ . We investigated physicochemical (solubility and sorption/desorption) and biological (microbial uptake and degradation) factors influencing the vertical mobility of DCD and DMPP, in soil columns of contrasting texture and organic matter following a 40 mm rainfall event, with and without the presence of sheep urine. We hypothesised that 1) DCD would move further down the soil profile than DMPP following simulated rainfall, due to a greater sorption of DMPP, 2) a greater co-location would be observed for DMPP with urine  $NH_4^+$ , due to the lower mobility of DMPP, 3) the presence of

sheep urine would increase vertical movement and desorption of both NI, due to saturation of soil exchange sites by ions within sheep urine and 4) a greater microbial uptake and mineralization would occur for DCD in comparison to DMPP, due to a greater bioavailability of DCD in comparison to DMPP.

## **5.2 Materials and methods**

### *5.2.1 Soil and sheep urine analysis*

Three soil types were selected for experimentation, on the basis of contrasting texture and organic matter content: a sandy loam textured Eutric Cambisol (53°24'N, 4°02'W), a sandy clay loam textured Eutric Cambisol (53°14'N, 4°01'W) and a high organic matter containing Sapric Histosol (52°52'N, 0°47'W). Both Eutric Cambisol samples were collected from beneath moderately sheep grazed and fertilised pasture, and the Sapric Histosol was collected from a eutrophic lowland peat used in intensive arable agriculture. The soil types used in this study had not been previously exposed to either DCD or DMPP. A summary of soil properties is presented in Table 1.

Soil was sampled in triplicate (0–10 cm depth), sieved (< 2 mm) in order to reduce sample heterogeneity and stored at 4°C until required. Soil moisture content was determined by weight difference after oven drying (105°C), and organic matter was determined on dry soil by loss-on-ignition in a muffle furnace (450°C; Ball 1964). Soil C:N ratio was determined on oven-dried, ground soil samples using a TruSpec® Analyzer (Leco Corp., St. Joseph, MI). The cation exchange capacity (CEC) of soils was determined using an unbuffered salt extraction method of Schofield (1949). Soil pH and electrical conductivity (EC) were measured using standard electrodes (1: 2.5 (w/v) soil-to-distilled water). Soluble C and N were determined in 1:5 soil-to-0.5 M K<sub>2</sub>SO<sub>4</sub> extracts using a Multi N/C 2100S Analyzer (AnalytikJena, Jena, Germany), within 24 h of sample collection, according to Jones and Willett (2006). Microbial

C and N were determined using the chloroform fumigation-extraction method of Voroney et al. (2008) ( $K_{EC} = 0.35$  and  $K_{EN} = 0.5$ ). Total available nitrate ( $\text{NO}_3^-$ ), ammonium ( $\text{NH}_4^+$ ) and phosphate (P) were determined within 0.5 M  $\text{K}_2\text{SO}_4$  extracts via the colorimetric procedures of Miranda et al. (2001), Mulvaney (1996) and Murphy and Riley (1962), respectively. Cations ( $\text{Na}^+$ ,  $\text{K}^+$  and  $\text{Ca}^{2+}$ ) were determined within 1:5 (w/v) soil-to-1 M  $\text{NH}_4\text{Cl}$  extracts using a Sherwood Model 410 Flame Photometer (Sherwood Scientific Ltd, Cambridge, UK).

Sheep urine was collected from Welsh Mountain ewes fed a diet of 80% *Lolium perenne* L. and 20% *Trifolium repens* L., where urine was collected between the hours of 10 a.m. and 4 p.m. Several urine samples from a single sheep were pooled for experimental use. The urine was frozen (unacidified) before use to avoid losses of N. The sheep urine had a pH of 8.99 and an EC of 22  $\text{mS cm}^{-1}$ ; the urine contained a total of 2.27  $\text{g N l}^{-1}$ , 3.00  $\text{g organic C l}^{-1}$ , 1.71  $\text{g urea N l}^{-1}$ , 44.9  $\text{mg NH}_4^+\text{-N l}^{-1}$ , 0.44  $\text{mg NO}_3^-\text{-N l}^{-1}$ , 0.92  $\text{mg P l}^{-1}$ , 7.16  $\text{g K l}^{-1}$ , 1.11  $\text{g Na l}^{-1}$  and 73.3  $\text{mg Ca l}^{-1}$ . Properties were measured directly on the urine via the methods described above and urea was measured using the method of Orsonneau et al. (1992).

**Table 5.1** Properties of soils used to fill the soil columns, values represent means  $\pm$  SEM,  $n = 3$ , letters indicate significant differences between the soils, results are reported on a dry soil weight basis and N.D. stands for not determined.

<b>Soil Property</b>	<b>Sandy Loam</b>	<b>Sandy Clay Loam</b>	<b>Sapric Histosol</b>
Texture: Sand (%)	80.0 $\pm$ 2.52 a	49.0 $\pm$ 2.00 b	N.D.
Silt (%)	11.7 $\pm$ 2.19 a	31.3 $\pm$ 0.88 b	N.D.
Clay (%)	8.33 $\pm$ 0.67 a	19.7 $\pm$ 1.20 b	N.D.
Organic matter (%)	5.40 $\pm$ 0.29 a	5.10 $\pm$ 0.44 a	77.6 $\pm$ 0.36 b
Gravimetric moisture content (%)	14.2 $\pm$ 0.12 a	19.5 $\pm$ 0.33 b	61.2 $\pm$ 0.10 c
Cation exchange capacity (meq 100 g <sup>-1</sup> )	14.0 $\pm$ 0.29 a	14.8 $\pm$ 0.68 a	80.8 $\pm$ 0.80 b
pH	6.64 $\pm$ 0.15 a	6.42 $\pm$ 0.04 a	6.37 $\pm$ 0.06 a
Electrical conductivity ( $\mu$ S cm <sup>-1</sup> )	51.1 $\pm$ 13.5 a	67.8 $\pm$ 7.34 ab	107 $\pm$ 11.8 b
Total C (g kg <sup>-1</sup> )	24.1 $\pm$ 2.50 a	19.1 $\pm$ 0.76 a	392 $\pm$ 1.33 b
Total N (g kg <sup>-1</sup> )	3.29 $\pm$ 0.14 a	3.12 $\pm$ 0.13 a	26.6 $\pm$ 0.09 b
Dissolved organic C (mg C kg <sup>-1</sup> )	52.7 $\pm$ 2.57 a	64.7 $\pm$ 13.4 a	959 $\pm$ 79.1 b
Total dissolved N (mg N kg <sup>-1</sup> )	7.51 $\pm$ 0.78 a	9.92 $\pm$ 0.66 a	71.4 $\pm$ 6.98 b
Microbial C (g kg <sup>-1</sup> )	0.35 $\pm$ 0.01 a	0.38 $\pm$ 0.04 a	4.41 $\pm$ 0.20 b
Microbial N (mg kg <sup>-1</sup> )	27.2 $\pm$ 0.79 a	26.1 $\pm$ 1.33 a	777 $\pm$ 42.3 b
Extractable NH <sub>4</sub> <sup>+</sup> (mg N kg <sup>-1</sup> )	0.20 $\pm$ 0.01 a	0.24 $\pm$ 0.03 a	6.50 $\pm$ 0.46 b
Extractable NO <sub>3</sub> <sup>-</sup> (mg N kg <sup>-1</sup> )	0.61 $\pm$ 0.14 a	0.89 $\pm$ 0.12 a	29.3 $\pm$ 2.73 b
Extractable P (mg kg <sup>-1</sup> )	0.32 $\pm$ 0.004 a	0.08 $\pm$ 0.04 a	26.1 $\pm$ 3.59 b
Exchangeable K (meq kg <sup>-1</sup> )	2.39 $\pm$ 0.43 a	1.61 $\pm$ 0.31 a	8.95 $\pm$ 3.79 a
Exchangeable Na (meq kg <sup>-1</sup> )	0.77 $\pm$ 0.08 a	1.37 $\pm$ 0.08 a	5.17 $\pm$ 0.51 b
Exchangeable Ca (meq kg <sup>-1</sup> )	51.8 $\pm$ 1.51 a	53.3 $\pm$ 2.12 a	617 $\pm$ 17.4 b

### 5.2.2 Comparative mobility of $^{14}\text{C}$ -DCD and $^{14}\text{C}$ -DMPP under simulated rainfall

To compare the mobility of  $[\text{U}]^{14}\text{C}$ -DCD and  $5\text{-}^{14}\text{C}$ -DMPP (American Radiolabelled Chemicals, St Louis, MO, USA) in contrasting soils under a simulated rainfall event, polypropylene tubes ( $n = 3$ ; 15 cm depth; 0.8 cm diameter) were repacked with sieved, field moist soil (soil sampled at 0-10 cm depth) to approximate field bulk density values (9, 8 and 5 g for the sandy loam, sandy clay loam and Sapric Histosol). This resulted in bulk densities of 1.0, 0.9 and 0.4 g cm<sup>-3</sup> and porosities of 0.60, 0.67 and 0.71 in the sandy loam, sandy clay loam and Sapric Histosol soil columns, respectively (particle density was assumed to be 2.65 g cm<sup>-3</sup> in the mineral soils and 1.4 g cm<sup>-3</sup> in the organic soils; Rowell 1994). In order to pack the soils evenly, additions of soil to the columns were made at increments, followed by lightly tapping the columns at the base to create an even density. The bottom of the tubes contained nylon mesh, to allow for drainage of leachate and to prevent any loss of soil. Nevertheless, no leachate was present following the rainfall simulations. Field relevant application rates of either  $^{14}\text{C}$ -DCD (1 g l<sup>-1</sup>; 50  $\mu\text{l}$ ; *ca.* 1 kBq) or  $^{14}\text{C}$ -DMPP (0.1 g l<sup>-1</sup>; 50  $\mu\text{l}$ ; *ca.* 1 kBq) were applied to the top of the column and a 40 mm rainfall event was simulated by adding 2 ml of distilled water drop-wise to the soil columns, *ca.* 5 minutes after the application of the NI. This rainfall event was chosen to simulate UK storm conditions which promote rapid water movement down the soil profile. It should also be noted that these leaching rates also approximate rates of water movement down preferential flow pathways in the soil profile under lower rainfall events. Preliminary studies indicated that the wetting front generally reached, but did not exceed the soil column length (15 cm).

The soil columns were incubated for 0.5 h, at laboratory temperature, after which the tubes were cut into the following depth fractions with a scalpel: 0-1, 1-2, 2-3, 3-5, 5-7, 7-9, 9-12 and 12-15 cm. The entire cut sections (including tube, to extract soil adhered to inner edge) were extracted with 0.5 M K<sub>2</sub>SO<sub>4</sub> (1:5 w/v; 0.5 h; 150 rpm). An aliquot (1.5 ml) of the soil



solution was centrifuged (14 000 g; 5 min) and the resulting supernatant was mixed with HiSafe 3 scintillant (PerkinElmer, Llantrisant, UK) and the activity measured using a Wallac 1404 Liquid Scintillation Counter (Wallac EG&G, Milton Keynes, UK).

### *5.2.3 Effect of ruminant urine on NI mobility and co-location of NI with urine ammonium*

To determine if the presence of urine influences the vertical movement of DCD or DMPP, soil columns ( $n = 3$ ) were prepared and processed as above. A sheep urine deposition event was simulated *ca.* 2 minutes following application of the  $^{14}\text{C}$ -DCD or  $^{14}\text{C}$ -DMPP and preceding the simulated rainfall event, by applying 250  $\mu\text{l}$  of sheep urine to the top of the soil column. The vertical distribution of the NI was compared to soil columns incubated without urine.

The  $\text{NH}_4^+$  concentration of the 0.5 M  $\text{K}_2\text{SO}_4$  extracts of each depth fraction was also determined on the urine treated soil columns, via the method described previously, to determine the co-location of  $^{14}\text{C}$ -DCD and  $^{14}\text{C}$ -DMPP with urine-derived  $\text{NH}_4^+$ .

### *5.2.4 Solubility of DCD and DMPP in water*

To determine the water solubility of DCD and DMPP, the OECD (1995) flask method was used. Briefly, 5 g of NI was added to 10 ml of water ( $n = 3$ ) and incubated at 30°C on a rotary shaker for 24 h. One replicate was then removed and incubated at 20°C for 24 h with occasional shaking, before centrifuging at 10 000 g. Samples were syringe filtered (0.2  $\mu\text{m}$ ) and analysed for total dissolved C, as described above, and the amount of NI dissolved in the water calculated. One of the remaining replicates was incubated for another 24 h at 30°C and the final replicate was incubated for an additional 48 h, before incubation at 20°C for a further 24 h and preparation of samples for analysis of dissolved C. This was conducted to ensure additional time had no effect on the amount of NI dissolved.

### 5.2.5 Sorption and desorption

Sorption isotherms were determined for  $^{14}\text{C}$ -DCD and  $^{14}\text{C}$ -DMPP in the three contrasting soils in the presence and absence of sheep urine, using similar methods as Jones and Brassington (2003). Briefly,  $^{14}\text{C}$ -DCD or  $^{14}\text{C}$ -DMPP was applied (50  $\mu\text{l}$ ; *ca.* 1 kBq) to 0.5 g ( $n = 3$ ) of field moist soil, following which 2.5 ml of either 0.01 M  $\text{CaCl}_2$  or sheep urine was added to the soils. A total of 8 concentrations of  $^{14}\text{C}$ -DCD and  $^{14}\text{C}$ -DMPP were used, ranging from 0.08-10 mg NI  $\text{l}^{-1}$ . The soil suspensions were shaken for 0.5 h at 150 *rpm* on a rotary shaker, subsequently an aliquot (1.5 ml) was centrifuged (10 000 *g*; 5 min) and the  $^{14}\text{C}$  activity in the supernatant determined by liquid scintillation counting, as above. Sorption of  $\text{NH}_4^+$  was also assessed as described above, using 8 concentrations ranging from 2.3-300 mg  $\text{NH}_4^+\text{-N}$   $\text{l}^{-1}$ , in 0.01 M  $\text{CaCl}_2$ . The  $\text{NH}_4^+$  concentration in the supernatant was determined as above. The partition coefficient ( $K_d$ ) for the NI (with and without the presence of urine) and  $\text{NH}_4^+$  within soil, was determined via Equation 1, where  $C_{\text{ads}}$  ( $\mu\text{mol kg}^{-1}$ ) is the concentration adsorbed to the soil solid phase at equilibrium and  $C_{\text{sol}}$  ( $\mu\text{mol l}^{-1}$ ) is the adsorbate concentration remaining in solution at equilibrium.

$$K_d = C_{\text{ads}} / C_{\text{sol}} \quad (\text{Eqn. 5.1})$$

Desorption, with and without the presence of sheep urine, was determined by adding either  $^{14}\text{C}$ -DCD (25  $\mu\text{l}$ ; *ca.* 0.5 kBq) or  $^{14}\text{C}$ -DMPP (25  $\mu\text{l}$ ; *ca.* 0.5 kBq) at two concentrations (1 and 10 mg  $\text{l}^{-1}$ ) to 0.2 g of soil ( $n = 3$ ). Four successive washes of the soil were conducted by adding either 1 ml of 0.01 M  $\text{CaCl}_2$  or sheep urine and by conducting a final wash with 0.5 M  $\text{K}_2\text{SO}_4$ . Samples were shaken for 0.5 h at 150 *rpm* on a rotary shaker between the additions of fresh wash solution. At the end of each wash period the samples were centrifuged (10 000 *g*; 5 min) and the supernatant removed prior to adding fresh 0.01 M  $\text{CaCl}_2$  or sheep urine. The

activity in the supernatant was determined as described previously, where the activity residing in the entrained solution trapped within the soil matrix was accounted for. After the final wash, soils were dried (105 °C; 24 h) and ground before combustion in an OX400 biological oxidizer (RJ Harvey, Hillsdale, NJ, USA), where evolved  $^{14}\text{CO}_2$  was captured in Oxysolve C-400 (Zinsser analytic, Frankfurt, Germany) to quantify  $^{14}\text{C}$  remaining bound to soils following washes.

#### 5.2.6 Substrate mineralization and microbial uptake

Mineralization and microbial uptake of  $^{14}\text{C}$ -DCD and  $^{14}\text{C}$ -DMPP in the three soils was determined to quantify the degradation of the NI during the course of the incubation, via the methods of Hill et al (2007). 0.3 ml of  $^{14}\text{C}$ -DCD or  $^{14}\text{C}$ -DMPP (*ca.* 0.5 kBq ml $^{-1}$ ; 0.1 and 1 g l $^{-1}$ ) were added to 1 cm $^3$  of soil ( $n = 3$ ), contained in 10 cm $^3$  glass vessels. Evolved  $^{14}\text{CO}_2$  was captured by flowing (*ca.* 100 ml min $^{-1}$ ) moist air over the soil surface, with the outflow passing through two consecutive 0.1 M NaOH traps (capture efficiency > 95%; Hill et al. 2007). Traps were changed after 0.05, 0.12, 0.25, 0.5, 1, 2, 4 and 8 h, and the activity in the solution determined by liquid scintillation counting as above.

To enable calculation of the  $^{14}\text{C}$ -NI pool taken up by soil microbes,  $^{14}\text{C}$ -DCD or  $^{14}\text{C}$ -DMPP (0.3 ml; *ca.* 0.5 kBq; 0.1 and 1 g l $^{-1}$ ) was pipetted evenly onto the soil surface (1 g;  $n = 3$ ) and extractions using ice-cold 0.5 M K $_2$ SO $_4$  (1:5 w/v) were conducted at 0.05, 0.12, 0.25, 0.5 and 1 h following addition of the substrate. The soils were shaken (150 rpm; 0.5 h) and subsequently centrifuged (10 000 g, 10 min). The  $^{14}\text{C}$  in the resulting supernatant was determined via liquid scintillation counting as described above. Uptake of the substrate by soil microbes was calculated by deducting the 0.5 M K $_2$ SO $_4$  extractable pool from the starting  $^{14}\text{C}$  pool. This is assuming the extraction procedure removed all exchangeable  $^{14}\text{C}$ -NI and the remainder was taken up into the microbial biomass.

### 5.2.7 Statistical analysis

To compare the vertical mobility of  $^{14}\text{C}$ -DCD and  $^{14}\text{C}$ -DMPP under rainfall, to determine how urine influences the vertical mobility of the NI and to compare the co-location of the NI with that of urine- $\text{NH}_4^+$ , a one-way ANOVA with Tukey's post-hoc test was used to compare each section depth, following testing for normality (Ryan-Joiner test) and homogeneity of variance (Levene's test). The same analysis was conducted on the slope of the linear sorption isotherms, following log transformation of data. A one-way ANOVA was also conducted for desorption after the fourth wash in either urine or  $\text{CaCl}_2$ , for mineralization at the 8 h time point and microbial uptake after 1 h. All statistical analyses were performed in Minitab 17.1.0 (Minitab Inc., State College, PA).


## 5.3 Results

### 5.3.1 Comparative vertical mobility of NI following simulated rainfall

The distribution of extractable DCD- $^{14}\text{C}$  and DMPP- $^{14}\text{C}$  following simulated rainfall was generally similar within each soil type (Fig. 1). However, a greater retention of  $^{14}\text{C}$ -DCD was observed in comparison to  $^{14}\text{C}$ -DMPP in the top 0-1 cm depth fraction of the sandy loam (Fig. 1a) and sandy clay loam columns (Fig. 1b). The total amount of DCD- $^{14}\text{C}$  extracted from the columns with 0.5 M  $\text{K}_2\text{SO}_4$  after 0.5 h was  $66.6 \pm 1.29$ ,  $63.9 \pm 0.65$  and  $38.8 \pm 0.53\%$  of that applied to the sandy loam, sandy clay loam and the Sapric Histosol, respectively. In comparison, the percentage of DMPP- $^{14}\text{C}$  extracted from the columns was  $79.4 \pm 2.14$ ,  $72.5 \pm 1.42$  and  $39.1 \pm 1.39\%$  of that applied to the sandy loam, sandy clay loam and Sapric Histosol, respectively.

The presence of sheep urine reduced ( $p < 0.01$ ) the quantity of extractable DCD- $^{14}\text{C}$  and DMPP- $^{14}\text{C}$  in the top 1 cm of the sandy loam columns (Fig. 1a and d), increased ( $p < 0.01$ ) the amount of extractable DCD- $^{14}\text{C}$  in the bottom 12-15 cm depth fraction and had no effect ( $p$

> 0.05) on the extractable amount of DCD-<sup>14</sup>C and DMPP-<sup>14</sup>C in each remaining depth fraction. The presence of sheep urine did not influence the extractable amount of DCD-<sup>14</sup>C or DMPP-<sup>14</sup>C in any studied depth fraction of the sandy clay loam columns (Fig. 1b and e). The presence of urine had no effect ( $p > 0.05$ ) on the amount of extractable DCD-<sup>14</sup>C in each depth fraction of the Sapric Histosol (Fig. 1c and f). However, it decreased ( $p < 0.001$ ) the extractable amount of DMPP-<sup>14</sup>C from the top 1 cm and increased ( $p < 0.001$ ) the extractable amount in the 12-15 cm depth fraction.



**Figure 5.1** The percentage of extractable DCD-<sup>14</sup>C or DMPP-<sup>14</sup>C at depth fractions of sandy lam (a and d), sandy clay loam (b and e) and Sapric Histosol (c and f) columns, following a simulated 40 mm rainfall event (a, b and c) and a sheep urine deposition plus a 40 mm rainfall event (d, e and f). Soil type labels apply to each column and legends apply to each row of panels. Bars represent means ( $n = 3$ ) and error bars represent SEM.

The percentage of applied DCD-<sup>14</sup>C extracted from the soil columns with applied sheep urine plus rainfall was  $79.6 \pm 6.58$ ,  $72.9 \pm 1.92$  and  $43.6 \pm 0.73\%$  of the added label applied to the sandy loam, the sandy clay loam and Sapric Histosol, respectively. The total amount of DMPP-<sup>14</sup>C extracted from the soil columns under the same conditions was  $79.4 \pm 0.77$ ,  $71.5 \pm 3.54$  and  $47.9 \pm 0.01\%$  in the sandy loam, sandy clay loam and Sapric Histosol, respectively. In conclusion, urine increased the total amount of DCD extracted from soils, but had no effect on DMPP.


### 5.3.2 Co-location of NI with urine-derived ammonium

In general, the distribution of both DCD-<sup>14</sup>C and DMPP-<sup>14</sup>C within the soil profile coincided well with the urine-derived NH<sub>4</sub><sup>+</sup> (Fig. 2). A greater ( $p < 0.001$ ) percentage of total column extractable DCD-<sup>14</sup>C in comparison to NH<sub>4</sub><sup>+</sup> was found in the top 1 cm in all three soil types (Fig. 2a, b, and c), indicating a retention of DCD at the soil surface. A greater ( $p < 0.001$ ) percentage of total extractable NH<sub>4</sub><sup>+</sup> in comparison to DMPP-<sup>14</sup>C was found in the 9-12 cm depth fraction of the sandy loam columns (Fig. 2d). Greater ( $p < 0.001$ ) amounts of total extractable NH<sub>4</sub><sup>+</sup> in comparison to DCD-<sup>14</sup>C were also found in the 9-12 cm depth fraction of the sandy clay loam columns (Fig. 2e), indicating some dis-location of NI with NH<sub>4</sub><sup>+</sup> at depth. No differences were observed at any depth fraction for DMPP-<sup>14</sup>C and urine-NH<sub>4</sub><sup>+</sup> in the sandy clay loam or the Sapric Histosol columns (Fig. 2e and f, respectively), indicating similar vertical distributions under conditions of mass flow.

### 5.3.3 Solubility of DCD and DMPP in water

The solubility of DMPP at 20°C was significantly higher ( $p < 0.001$ ) at  $125 \pm 2.4 \text{ g l}^{-1}$  in comparison to that of DCD at  $73.2 \pm 2.0 \text{ g l}^{-1}$ . An increasing trend was not observed in the


replicates maintained for 48 and 72 h, indicating that saturation of the NI within the matrix had occurred after 24 h.



**Figure 5.2** The percentage of extractable DCD-<sup>14</sup>C (a, b and c) or DMPP-<sup>14</sup>C (d, e and f) compared to NH<sub>4</sub><sup>+</sup> at differing soil depth fractions following a simulated urine deposition plus 40 mm rainfall event applied to sandy loam (a and d), sandy clay loam (b and d) and Sapric Histosol (c and f) columns. Soil type labels apply to each column and legends apply to each row of panels. Bars represent means (*n* = 3) and error bars denote SEM.

#### 5.3.4 Sorption

Sorption isotherms for DCD-<sup>14</sup>C (Fig. 3a, c and e), DMPP-<sup>14</sup>C (Fig. 3b, d and f) and NH<sub>4</sub><sup>+</sup> (Fig. 4) were linear, where all R<sup>2</sup> values were greater than 0.95. The gradient of the linear sorption isotherms were steeper (*p* < 0.001) in the Sapric Histosol compared to the other soil types for both DCD-<sup>14</sup>C, DMPP-<sup>14</sup>C and NH<sub>4</sub><sup>+</sup> indicating greater amounts of sorption in this




**Figure 5.3** Linear sorption isotherms for <sup>14</sup>C-DCD (a, c and e) and <sup>14</sup>C-DMPP (b, d and f) in either a 0.01 M CaCl<sub>2</sub> or sheep urine matrix, in a sandy loam (a and b), sandy clay loam (c and d) and Sapric Histosol (e and f). Symbols represents means ( $n = 3$ ), bi-directional error bars represent SEM for sorption and equilibrium solution concentrations, legends apply to each column of panels and soil type labels apply to each row of panels.

soil type. In comparison to DMPP-<sup>14</sup>C, greater sorption occurred for DCD-<sup>14</sup>C in the Sapric Histosol in both matrices (0.01 M CaCl<sub>2</sub> and urine). However, no differences were observed between DCD-<sup>14</sup>C and DMPP-<sup>14</sup>C sorption in the other two soil types ( $p > 0.05$ ). In the Sapric Histosol the gradient of the DMPP-<sup>14</sup>C sorption isotherm was steeper ( $p < 0.001$ ) in the 0.01



M CaCl<sub>2</sub> matrix as opposed to the urine matrix. The calculated soil-to-solution partition coefficients ( $K_d$ ; Table 2) followed the trend sandy loam < sandy clay loam < Sapric Histosol for DCD-<sup>14</sup>C, DMPP-<sup>14</sup>C and NH<sub>4</sub><sup>+</sup>; for the NI this trend was observed at both concentrations and within both matrices.



**Figure 5.4** Linear sorption isotherms of NH<sub>4</sub><sup>+</sup> in 0.01 M CaCl<sub>2</sub> matrix, in three soils (sandy loam and sandy clay loam textured Eutric Cambisol and a Sapric Histosol). Symbols represents means ( $n = 3$ ) and bi-directional error bars represent SEM for sorption and equilibrium solution concentrations.

**Table 5.2** Soil-to-solution partition coefficient ( $K_d$ ) for the nitrification inhibitors DCD and DMPP at field relevant concentrations (1 and 10 mg l<sup>-1</sup>) with and without the presence of sheep urine. Values represent means ( $n = 3$ )  $\pm$  SEM.


Soil Type	Matrix	DCD		DMPP		NH <sub>4</sub> <sup>+</sup> 300 mg N l <sup>-1</sup>
		1 mg l <sup>-1</sup>	10 mg l <sup>-1</sup>	1 mg l <sup>-1</sup>	10 mg l <sup>-1</sup>	
Sandy loam textured Eutric Cambisol	0.01 M	1.25 $\pm$	1.34 $\pm$	0.87 $\pm$	0.63 $\pm$	1.27 $\pm$
	CaCl <sub>2</sub>	0.10	0.05	0.01	0.05	0.35
Sandy clay loam textured Eutric Cambisol	0.01 M	1.88 $\pm$	1.48 $\pm$	1.15 $\pm$	0.93 $\pm$	1.56 $\pm$
	CaCl <sub>2</sub>	0.05	0.09	0.05	0.05	0.18
Sapric Histosol	0.01 M	15.6 $\pm$	14.3 $\pm$	12.8 $\pm$	11.0 $\pm$	3.86 $\pm$
	CaCl <sub>2</sub>	0.28	0.31	0.33	0.44	0.11
Sandy loam textured Eutric Cambisol	Sheep urine	0.64 $\pm$ 0.10	0.64 $\pm$ 0.18	0.96 $\pm$ 0.05	0.84 $\pm$ 0.08	-
	Sheep urine	1.47 $\pm$ 0.15	1.17 $\pm$ 0.18	1.01 $\pm$ 0.04	1.11 $\pm$ 0.07	-
Sapric Histosol	Sheep urine	15.0 $\pm$ 0.36	13.4 $\pm$ 0.66	7.73 $\pm$ 0.43	6.99 $\pm$ 0.35	-

#### 5.3.4 Desorption

Generally, the presence of urine increased total desorption (Fig. 5) of both DCD-<sup>14</sup>C and DMPP-<sup>14</sup>C numerically (although not statistically) at the fourth consecutive wash, in all soil types. The presence of urine increased desorption of DMPP-<sup>14</sup>C in the Sapric Histosol at 1 mg l<sup>-1</sup> ( $p < 0.001$ ; Fig. 5k) and 10 mg l<sup>-1</sup> ( $p < 0.05$ ; Fig. 5l). Interestingly, the same trend was not observed for DCD-<sup>14</sup>C in the same soil type (Fig. 5i and j). Desorption of DCD-<sup>14</sup>C in the CaCl<sub>2</sub> matrix was greater ( $p < 0.05$ ) in the sandy loam soil (Fig. 5a and b) compared to the

Sapric Histosol (Fig. 5i and j), but desorption was no greater ( $p > 0.05$ ) in the sandy clay loam soil (Fig 5e and f). In the urine matrix desorption of DCD- $^{14}\text{C}$  was lower ( $p < 0.01$ ) in the Sapric Histosol (Fig. 5i and j) compared to the sandy loam (Fig. 5a and b) and sandy clay loam (Fig. 5e and f) textured Eutric Cambisol at both studied concentrations. For both studied concentrations of DMPP- $^{14}\text{C}$  in the  $\text{CaCl}_2$  matrix, desorption was lower ( $p < 0.01$ ) in the Sapric Histosol (Fig. 5k and l) in comparison to either the sandy loam (Fig. 5c and d) or the sandy clay loam (Fig. 5g and h) Eutric Cambisol. In the urine matrix, however, no differences in desorption of DMPP- $^{14}\text{C}$  was observed at either concentration between the soil types.

When comparing between the applied  $^{14}\text{C}$ -NI in the  $\text{CaCl}_2$  matrix, there was a greater desorption ( $p < 0.05$ ) of DMPP- $^{14}\text{C}$  (Fig. 5e and f) in comparison to DCD- $^{14}\text{C}$  (Fig. 5g and h) in the sandy clay loam. The same trend was true for the urine matrix, except no differences were observed between DCD- $^{14}\text{C}$  and DMPP- $^{14}\text{C}$  at  $1 \text{ mg compound l}^{-1}$ . In the urine matrix, greater ( $p < 0.001$ ) amounts of DMPP- $^{14}\text{C}$  (Fig. 5k and l) desorbed in comparison to DCD- $^{14}\text{C}$  (Fig. 5i and j) in the Sapric Histosol, at both studied concentrations. The final wash conducted with  $0.5 \text{ M K}_2\text{SO}_4$  typically only increased desorption by minor amounts (ranging from 1.4 to 6.0%), indicating that the previous washes had removed the majority of the extractable NI- $^{14}\text{C}$  from the soils. The mass balance for the total recovered DCD- $^{14}\text{C}$  and DMPP- $^{14}\text{C}$  following biological oxidation was  $101 \pm 0.63\%$  and  $101 \pm 0.95\%$  respectively, for all soil types, applied concentrations and matrices.



**Figure 5.5** Cumulative desorption of  $^{14}\text{C}$ -DCD and  $^{14}\text{C}$ -DMPP in a sandy loam (a, b, c and d), sandy clay loam (e, f, g and h) and a Sapric Histosol (i, j, k and l) at  $1 \text{ mg DCD l}^{-1}$  (a, e and i),  $10 \text{ mg DCD l}^{-1}$  (b, f and j),  $1 \text{ mg DMPP l}^{-1}$  (c, g and k) and  $10 \text{ mg DMPP l}^{-1}$  (d, h and l) in either a  $0.01 \text{ M CaCl}_2$  or sheep urine matrix. Text in the top row of panels applies to each respective column of panels and legend applies to all panels. Symbols represents means ( $n = 3$ ) and error bars represent SEM for sorption and equilibrium solution concentration.

### 5.3.5 Mineralization

The results of the mineralization assay confirmed that only minor degradation of the  $^{14}\text{C}$ -NI would have occurred under the conditions and duration (0.5 h) of the column study. Mineralization of  $^{14}\text{C}$ -DCD within all three soil types ranged from 0.10 to 0.35% of added  $^{14}\text{C}$  label, over 0.5 h. For  $^{14}\text{C}$ -DMPP the amount degraded over the period of the column incubation was even lower, ranging from 0.03 to 0.16% of the added  $^{14}\text{C}$  label. After 8 h, the amount of  $^{14}\text{C}$ -DCD mineralized was still low, ranging from 0.41 to 1.67% of the  $^{14}\text{C}$ -label applied in the three soil types and at both studied concentrations; the amount of  $^{14}\text{C}$ -DMPP mineralized after 8 h ranged from 0.05 to 0.25% of the applied label. Greater amounts ( $p < 0.01$ ;  $0.84 \pm 0.15$  and  $1.53 \pm 0.30\%$  more at 0.1 and 1 g NI l<sup>-1</sup>, respectively) of  $^{14}\text{C}$ -DCD mineralized in the Sapric Histosol in comparison to  $^{14}\text{C}$ -DMPP, at both studied concentrations. The same pattern was also seen for the sandy clay loam textured Eutric Cambisol at 0.1 g NI l<sup>-1</sup>, where  $0.83 \pm 0.12\%$  more  $^{14}\text{C}$ -DCD was mineralized in comparison to  $^{14}\text{C}$ -DMPP. No differences were observed in the amount of  $^{14}\text{C}$ -DMPP mineralized between all soil types at either studied concentration. For  $^{14}\text{C}$ -DCD at 0.1 g l<sup>-1</sup>,  $1.17 \pm 0.21\%$  and  $0.69 \pm 0.18\%$  more  $^{14}\text{C}$ -DCD mineralized in the Sapric Histosol in comparison to either the sandy loam or sandy clay loam textured Eutric Cambisol, respectively. No differences ( $p > 0.05$ ) were observed in the amount of  $^{14}\text{C}$ -DCD mineralized between the different soils at the higher studied concentration.

### 5.3.6 Microbial uptake

After 1 h in the sandy loam textured Eutric Cambisol, no difference ( $p < 0.05$ ) was observed between the amount of DCD- $^{14}\text{C}$  or DMPP- $^{14}\text{C}$  acquired by the soil microbes, which ranged between 20 and 23% of that applied, at both studied concentrations. The same trend was observed in the sandy clay loam textured Eutric Cambisol, where uptake ranged from 18 to 28% of that applied. In the Sapric Histosol, greater amounts ( $p < 0.001$ ) of DCD- $^{14}\text{C}$  ( $66 \pm$

0.36%) was acquired by soil microbes in comparison to DMPP-<sup>14</sup>C ( $51 \pm 2.67\%$ ) at the higher application rate, however, no differences ( $p > 0.05$ ) between DCD-<sup>14</sup>C ( $56 \pm 0.32\%$ ) and DMPP-<sup>14</sup>C ( $61 \pm 1.01\%$ ) were observed at the lower concentration. The microbial uptake was two to three-fold greater ( $p < 0.001$ ) in the Sapric Histosol compared to the mineral soils for both NI and at both studied concentrations. The results of the microbial uptake study correspond well with that of the soil column studies, indicating that the deficit in the amount of <sup>14</sup>C-NI recovered from the soil columns is that which was immobilised into microbial biomass and degraded within the soils.

#### **5.4 Discussion**

Our first hypothesis was that DCD would be more mobile and translocate further down the soil profile than DMPP, due to the positive charge and rapid sorption of DMPP to soil colloids (Azam et al. 2001). The results of the column study investigating the vertical movement of DCD and DMPP over 0-15 cm under a 40 mm rainfall event, revealed that the mobility of both NI were similar, and DCD did not appear to be more mobile than DMPP. A greater sorption for DCD in comparison to DMPP was found in the organic and mineral soils, contradicting our hypothesis that a greater adsorption of DMPP would occur. A greater sorption was found in the Sapric Histosol compared to the mineral soils for both NI, suggesting negatively charged domains within organic matter are important for adsorption processes. However, if the results of Fig.5 are expressed on a per g of organic C basis, the results between the soils are more similar, suggesting a partition phenomenon rather than charged-based sorption. Compounds possessing a greater octanol-water partition coefficient than others will show slower sorption; the octanol-water partition coefficient (Log P) of DCD and DMPP is predicted to be -1.03 and 0.92 for DCD and DMPP, respectively (Chemicalize.org 2016). In addition, DCD is hydrophilic (Turowski and Deshmukh 2004), which in combination with the

low value for the octanol-water partition coefficient, suggests strong absorption and permeation into organic matter. Protection and occlusion of NI from nitrifiers and other microbes due to sorption, may reduce the effectiveness of NI, at least in the short term. Sorption may also protect against some microbial degradation, and if remobilisation of NI occurs, it may prolong the inhibitory effect (Barth et al. 2001).

Under some circumstances DMPP has been found to be more effective than DCD at inhibiting nitrification and reducing N<sub>2</sub>O losses (Weiske et al. 2001; Chaves et al. 2006; Irigoyen et al. 2006) and this difference in efficacy is often attributed to the lower mobility of DMPP in comparison to DCD and hence a greater spatial separation of DCD with NH<sub>4</sub><sup>+</sup>. Nevertheless, the inhibition of the oxidation of NH<sub>4</sub><sup>+</sup> to NO<sub>3</sub><sup>-</sup> only occurs when the nitrifying population have taken up the NI. This study only examined the mobility of NI and NH<sub>4</sub><sup>+</sup>, however, a consideration of the distribution of nitrifying microorganisms and their acquisition of the NI is an important aspect for future research. Our second hypothesis was that DMPP would co-locate with urine-NH<sub>4</sub><sup>+</sup> more than DCD. In our study, both NI appeared to coincide well with urine-derived NH<sub>4</sub><sup>+</sup>, with only few incidences of the percentage of extractable NH<sub>4</sub><sup>+</sup> being higher than the extractable NI-<sup>14</sup>C label at depth. Nevertheless, this study only focused on the short-term co-location of NI with urine-NH<sub>4</sub><sup>+</sup>, and further generation of urine-NH<sub>4</sub><sup>+</sup> would occur post urea hydrolysis. Being a neutral compound, urea may also be susceptible to vertical transport (Dawar et al. 2011). However, urea hydrolysis is normally complete within ca. 2 days, reducing the time available for vertical movement. Comparing the soil-to-solution partition coefficients of the NI and NH<sub>4</sub><sup>+</sup>, it appears that the *K<sub>d</sub>* of NH<sub>4</sub><sup>+</sup> and DCD are more similar than that of DMPP and NH<sub>4</sub><sup>+</sup> in the sandy loam and the sandy clay loam soils, but large differences were found for both NI compared to NH<sub>4</sub><sup>+</sup> in the Sapric Histosol. It may be possible to use NI and NH<sub>4</sub><sup>+</sup> *K<sub>d</sub>* values from differing soil types in order to assess which (if any) inhibitor may be more effective, where similar *K<sub>d</sub>* values may result in a greater co-location of the two

chemicals. Further work is required to assess whether this would be a useful proxy for reducing N<sub>2</sub>O emissions and improving NI use in order to maximise efficacy.

The third objective was to determine if the presence of sheep urine resulted in a greater vertical movement of both NI. Without the presence of urine (rainfall only), a retention of DCD was observed in the top 1 cm of the sandy loam and sandy clay loam textured soils, and a retention of both NI was found in the top 1 cm of the Sapric Histosol. A retention of NI at the surface may be beneficial in that nitrification decreases with pasture soil depth (Young et al. 2002), which may result in a greater coincidence of NI with nitrifiers, nevertheless, the use of sieved soils in this study would have altered any natural depth distribution of nitrifiers in the incubated soil profiles. The addition of sheep urine to the soil columns had a mixed effect on the depth distribution of the NI, depending on the soil type and inhibitor. Relative to simulated rainfall alone, the presence of sheep urine resulted in a greater amount of extractable DCD-<sup>14</sup>C at the bottom of the sandy loam soil columns, and reduced the extractable amount of both DCD-<sup>14</sup>C and DMPP-<sup>14</sup>C in the top 1 cm of the sandy loam soil columns, however, no differences in depth distribution of either NI were observed following sheep urine and rainfall application to the sandy clay loam soil. The presence of sheep urine resulted in lower amounts of DMPP-<sup>14</sup>C at the top 1 cm and greater amounts of extractable DMPP-<sup>14</sup>C at 12-15 cm in the Sapric Histosol, but no such trend was observed for DCD-<sup>14</sup>C. To consider the reasons behind these results, a consideration of the soil, NI and urine properties are required.

The soils used in this study were all of a similar pH (Table 1), however the addition of urine would have altered the soil pH and made conditions in the soil columns more alkaline. As DCD is amphipathic, sorption has been shown to be pH dependent, where increases in pH above pH 5 lead to increased sorption (Zhang et al. 2004). This may partially explain why the vertical distribution of DCD-<sup>14</sup>C was similar whether urine was present or not in the Sapric Histosol. In the solubility assay, DMPP was found to be over 1.5 times more soluble in water



than DCD. However, a saturated solution ( $125 \text{ g l}^{-1}$ ) of DMPP is acidic (ca. pH 3), whereas dissolving DCD results in a near neutral pH. The solubility of DMPP at pH 7, is reported to be  $46 \text{ g l}^{-1}$  (Zerulla et al. 2001) - considering this value results in DCD having a greater solubility than DMPP. Thus, DMPP solubility may vary widely as a function of soil pH and buffer capacity, but whether this influences the mobility relative to DCD is unclear. As the soils used in this study had a similar pH, NI solubility would not have varied much due to the effect of soil type.

The sandy loam and sandy clay loam soils had a similar CEC and organic matter content (Table 1), however, both parameters were greater in the Sapric Histosol. The NI sorption and partitioning into organic matter, and the availability of cation exchange sites may have all contributed to the differences in the distribution of the NI, with and without the presence of urine. The low CEC in the mineral soils and saturation of these exchange sites with cations present within the urine, may explain the higher amounts of DCD- $^{14}\text{C}$  and DMPP- $^{14}\text{C}$  at the bottom of the sandy loam soil columns, compared to the rainfall only treatment. This trend did not hold true, however, for the sandy clay loam columns despite the similarity in soil properties.

The results of our desorption assays revealed that even after one wash with  $0.01 \text{ M CaCl}_2$ , a large proportion of DCD and DMPP was remobilised into solution. This supports the theory that remobilisation of these NI may occur e.g. following urine deposition and/or heavy rainfall events. In the case of ruminant urine events this effect may be important, as urine is generally deposited at varying times following NI application to pasture. While the presence of urine generally increased desorption, the effect was strongest for DMPP- $^{14}\text{C}$  in the Sapric Histosol, and weakest for DCD- $^{14}\text{C}$  in the same soil type. This again shows a clear contrast in the binding mechanisms and behaviour of these two NI. In the Sapric Histosol, the vertical movement of DMPP was enhanced due to the presence of urine. This trend was not observed for DCD, which may be due to differences in the physico-chemical properties of these NI. The

partitioning and adsorption of DCD in the organic soil may have prevented its vertical movement due to urine addition.

The short-term microbial mineralization of DCD was faster in comparison to DMPP in all soil types, which supports our fourth hypothesis. This is consistent with results of other studies, where DMPP has been found to have a longer residence time in comparison to DCD (Chaves et al. 2006). DCD degrades to CO<sub>2</sub> and NH<sub>4</sub><sup>+</sup> via guanilyc urea, guanidine, and urea (Amberger 1986; Kelliher et al. 2008). The first stage of DCD degradation can occur on the surface of metal oxides, which catalyse the reaction of DCD and water to guanylurea (Hallinger et al. 1990). Biological degradation of DCD by common soil microorganisms has also been demonstrated, where DCD is supplied as the sole N source in pure culture (Hauser and Haselwandter 1990; Schwarzer et al. 1998). As DMPP is a heterocyclic compound, it is not readily degradable (Chaves et al. 2006), although information on the mechanism and degradation pathways of this inhibitor are still lacking. Results from this study indicate that the microbial community were better able to degrade DCD in comparison to DMPP, where DMPP degrading bacteria may take longer time periods to establish in comparison to DCD. DMPP has also been found to have a longer effect on the abundance of ammonium oxidizing bacteria in comparison to DCD (Kuo et al. 2015) and there is evidence that DMPP has an inhibitory effect on both ammonium oxidizing bacteria and archaea (Florio et al. 2014).

In this study, even after 1 h, the microbial uptake of both inhibitors accounted for a large proportion of that applied. Approximately 20% was taken up by soil microbes in the mineral soils and > 50% of that applied was taken up by soil microbes in the Sapric Histosol, which was likely to be a function of the greater microbial biomass in this soil. To be effective, the NI would need to be acquired by the target microbial communities (ammonium oxidizing bacteria and archaea). Immobilisation into non-target microbial communities could, therefore, equate to a fairly large removal mechanism for these NI and this requires further investigation.

No difference was observed between DCD and DMPP in the proportion acquired by soil microbes in the mineral soils. However, uptake was greater for DCD compared to DMPP in the Sapric Histosol at the higher studied concentration. This suggests a slight preference of, or bioavailability of DCD to the soil microbial community in the short-term.

The results of this study should be considered with care, as repacked soils were used and there were no preferential flow pathways as would occur under field conditions, which would potentially enhance the vertical movement of either NI. The soils were also sieved, which removed any natural variation of soil properties which can occur with depth. The use of soil columns could have also promoted vertical movement of NI or urine, by restricting lateral diffusion. This is in contrast to the study by Azam et al (2001), where shallow petri dishes were used and NI and  $\text{NH}_4^+$  were applied in the centre. This approach may have promoted lateral movement of solutes. The soils used to repack the soil columns in this study were field moist and the initial starting moisture content of the soil could also influence the movement of inhibitors through the soil. Further work should attempt to establish the comparative efficacy of both NI, their uptake by nitrifying and non-nitrifying microorganisms and co-location with  $\text{NH}_4^+$  over longer time scales and under field conditions with differing soil moisture contents.

## **5.5 Conclusions**

A similar distribution of DCD and DMPP was observed up to a depth of 15 cm following a simulated rainfall event in one organic and two mineral soils. The presence of sheep urine did not influence the depth distribution of DCD following rainfall, but enhanced the movement of DMPP down the profile, especially in the organic soil. A greater sorption was found for DCD in comparison to DMPP in the soil types studied here and the presence of urine generally increased desorption of both NI. The results of our study suggest that the efficacy of

NI are influenced more by differences in microbial uptake and degradation rates than by differences in sorption and desorption rates to the soil matrix.

## **5.6 Acknowledgements**

KAM would like to thank the School of Environment, Natural Resources and Geography, Bangor University, for sponsoring this study. This work was supported by the UK Natural Environment Research Council, under grant award (NE/M015351/1).

## 5.7 References

- Amberger A (1986) Potentials of nitrification inhibitors in modern N-fertilizer management. *J Plant Nutr Soil Sc* 149:469-484
- Azam F, Benckiser G, Müller C, Ottow JCG (2001) Release, movement and recovery of 3,4-dimethylpyrazole phosphate (DMPP), ammonium, and nitrate from stabilized nitrogen fertilizer granules in a silty clay soil under laboratory conditions. *Biol Fertil Soils* 34: 118-125
- Ball DF (1964) Loss-on-ignition estimate of organic matter and organic carbon in calcareous soils. *J Soil Sci* 15:84-92
- Barth G, von Tucher S, Schmidhalter U (2001) Influence of soil parameters on the effect of 3,4-dimethylpyrazole-phosphate as a nitrification inhibitor. *Biol Fertil Soils* 34:98-102
- Barth G, von Tucher S, Schmidhalter U (2008) Effectiveness of 3,4-dimethylpyrazole phosphate as nitrification inhibitor in soil as influenced by nitrification inhibitor concentration, application form, and soil matric potential. *Pedosphere* 18:378-385
- Bateman EJ, Baggs EM (2005) Contributions of nitrification and denitrification to N<sub>2</sub>O emissions from soils at different water-filled pore space. *Biol Fertil Soils* 41:379-388
- Benckiser G, Christ E, Herbert T, Weiske A, Blome J, Hardt M (2013) The nitrification inhibitor 3,4-dimethylpyrazole-phosphate (DMPP) – quantification and effects on soil metabolism. *Plant Soil* 371: 257-266
- Carrillo-González R, Šimůnek J, Sauve S, Adriano D (2006) Mechanisms and pathways of trace element mobility in soils. *Adv Agron* 91:111-178
- Chaves B, Opoku A, De Neve S, Boeckx P, Van Cleemput O, Hofman G (2006) Influence of DCD and DMPP on soil N dynamics after incorporation of vegetable crop residues. *Biol Fertil Soils* 43:62-68

Chemicalize.org (2016). Used for name to structure generation/prediction of LogP properties.

January 2016, chemicalize.org and Chemaxon (<http://www.chemaxon.com>)

Dawar K, Zaman M, Rowarth JS, Blennerhassett J, Turnbull MH (2011) Urea hydrolysis and lateral and vertical movement in the soil: effects of urease inhibitor and irrigation. *Biol Fertil Soils* 47:139-146.

de Klein CAM, Cameron KC, Di HJ, Rys G, Monaghan RM, Sherlock RR (2011) Repeated annual use of the nitrification inhibitor dicyandiamide (DCD) does not alter its effectiveness in reducing N<sub>2</sub>O emissions from cow urine. *Anim Feed Sci Tech* 166:480-491

Di HJ, Cameron KC (2003) Mitigation of nitrous oxide emissions in spray-irrigated grazed grassland by treating the soil with dicyandiamide, a nitrification inhibitor. *Soil Use Manage* 19:284-290

Di HJ, Cameron KC (2004) Treating grazed pasture soil with a nitrification inhibitor eco-n<sup>TM</sup>, to decrease nitrate leaching in a deep sandy soil under spray irrigation – a lysimeter study. *New Zeal J Agr Res* 47: 351-361.

Di HJ, Cameron KC (2007) Nitrate leaching losses and pasture yields as affected by different rates of animal urine nitrogen returns and application of a nitrification inhibitor – a lysimeter study. *Nutr Cycle Agroecosys* 79:281-290

Di HJ, Cameron KC (2011) Inhibition of ammonium oxidation by a liquid formulation of 3,4-Dimethylpyrazole phosphate (DMPP) compared with a dicyandiamide (DCD) solution in six New Zealand grazed grassland soils. *J Soil Sediment* 11:1032-1039

Di HJ, Cameron KC (2012) How does the application of different nitrification inhibitors affect nitrous oxide emissions and nitrate leaching from cow urine in grazed pastures. *Soil Use Manage* 28:54-61

- Fangueiro D, Fernandes A, Coutinho J, Mreira N, Trindade H (2009) Influence of two nitrification inhibitors (DCD and DMPP) on annual ryegrass yield and soil mineral N dynamics after incorporation with cattle slurry. *Commun Soil Sci Plan* 40:3387-3398
- Fiencke C, Bock E (2006) Immunocytochemical localization of membrane-bound ammonia monooxygenase in cells of ammonia oxidizing bacteria. *Arch Microbiol* 185:99-106
- Florio A, Clark AM, Hirsch PR, Jhurrea D, Benedetti A (2014) Effects of the nitrification inhibitor 3,4-dimethylpyrazole phosphate (DMPP) on abundance and activity of ammonia oxidizers in soil. *Biol Fertil Soils* 50:795-807
- Gilsanz C, Báez D, Misselbrook, TH, Dhanoa MS, Cárdenas LM (2016). Development of emission factor and efficiency of two nitrification inhibitors, DCD and DMPP. *Agri Ecosyst Environ* 216:1-8.
- Hallinger S, Wallnöfer PR, Goldbach H, Amberger A (1990) Several aspects of bacterial dicyandiamide degradation. *Naturwissenschaften* 77:332-334
- Hauser M, Haselwandter K (1990) Degradation of dicyandiamide by soil bacteria. *Soil Biol Biochem* 22:113-114
- Hill PW, Kusyakov Y, Jones D, Farrar J (2007) Response of root respiration and root exudation to alterations in root C supply and demand in wheat. *Plant Soil* 291:131-141
- IPCC (2007). *Climate change 2007 The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change.* Solomon S, Qin D, Manning M, Chen Z, Marquis M, Averyt KB, Tignor M and Miller HL (Eds) Cambridge University Press, Cambridge 996 pp
- Irigoyen I, Lamsfus C, Aparico-Tejo P, Muro J (2006) The influence of 3,4-dimethylpyrazole phosphate and dicyandiamide on reducing nitrate accumulation in spinach under Mediterranean conditions. *J Agr Sci* 144:555-562

- Jones DL, Brassington DS. 2003. Sorption of organic acids in acid soils and its implications in the rhizosphere. *Eur J Soil Sci* **49**:447-455.
- Jones DL, Willett VB (2006) Experimental evaluation methods to quantify dissolved organic nitrogen (DON) and dissolved organic carbon (DOC) in soil. *Soil Biol Biochem* **5**:991-999
- Kelliher FM, Clough TJ, Clark H, Rys G, Sedcole JR (2008) The temperature dependence of dicyandiamide (DCD) degradation in soils: A data synthesis. *Soil Biol Biochem* **40**: 1878-1882
- Kou YP, Wei K, Chen GX, Wang ZY, Xu H (2015) Effects of 3,4-dimethylpyrazole phosphate and dicyandiamide on nitrous oxide emission in a greenhouse vegetable soil. *Plant Soil Environ* **61**:29-35
- Ledgard SF, Luo J, Sprosen MS, Wyatt JB, Balvert SF, Lindsey SB (2014). Effects of the nitrification inhibitor dicyandiamide (DCD) on pasture production, nitrous oxide emissions and nitrate leaching in Waikato, New Zealand. *New Zeal J Agr Res* **57**:294-315.
- Liu C, Wang K, Zheng X (2013) Effects of nitrification inhibitors (DCD and DMPP) on nitrous oxide emission, crop yield and nitrogen uptake in a wheat maize cropping system. *Biogeosciences* **10**:2427-2437
- Luo J, Ledgard S, Wise B, Welten B, Lindsey S, Judge A, Sprose M (2015). Effects of dicyandiamide (DCD) delivery method, application rate, and season on pasture urine patch nitrous oxide emissions. *Biol Fertil Soils* **51**:453-464.
- McGeough KL, Watson CJ, Müller C, Laughlin JR, Chadwick DR (2016) Evidence that the efficacy of the nitrification inhibitor dicyandiamide (DCD) is affected by soil properties in UK soils. *Soil Biol Biochem* **94**:222-232



- Miranda KM, Epsey MG, Wink DA (2001) A rapid, simple, spectrophotometric method for simultaneous detection of nitrate and nitrite. *Nitric Oxide* 5:62-71
- Misselbrook TH, Cardenas LM, Camp V, Thorman RE, Williams JR, Rollett AJ, Chambers BJ (2014) An assessment of nitrification inhibitors to reduce nitrous oxide emissions from UK agriculture. *Environ Res Lett* 9:115006.
- Mulvaney RL (1996) Nitrogen - inorganic forms. In: Sparks DL (Eds) *Methods of Soil Analysis. Part 3*. Soil Science Society of America Inc., Madison, WI, pp 1123-1184
- Murphy J, Riley JP (1962) A modified single solution method for the determination of phosphate in natural waters. *Anal Chim Acta* 27:31-36
- O'Callaghan M, Gerard EM, Carter PE, Lardner R, Sarathchandra U, Burch G, Ghani A, Bell N (2010) Effect of the nitrification inhibitor dicyandiamide (DCD) on microbial communities in a pasture soil amended with bovine urine. *Soil Biol Biochem* 42:1425-1436
- OECD (1995) OECD guideline for the testing of chemicals – water solubility. No. 105. OECD, Paris.
- Orsonneau J-L, Massoubre C, Cabanes M, Lustenberger P (1992) Simple and sensitive determination of urea in serum and urine. *Clin Chem* 38:619-623
- Pereira J, Fanguero D, Chadwick DR, Misselbrook TH, Coutinho J, Trindade H (2010) Effect of cattle slurry pre-treatment by separation and addition of nitrification inhibitors on gaseous emissions and N dynamics: a laboratory study. *Chemosphere* 79:620-627
- Rowell DL (2014) *Soil Science: Methods and Applications*. Routledge, New York, USA
- Ruser R, Schulz R (2015) The effect of nitrification inhibitors on the nitrous oxide (N<sub>2</sub>O) release from agricultural soils – a review. *J Plant Nutr Soil Sc* 178:171-188

- Schofield RK (1949) Effect of pH on electric charges carried by clay particles. *J Soil Sci* 1:1-8
- Schwarzer C, Auer B, Klima J, Haselwandter K. (1998) Physiological and electron microscopical investigations on syntrophic dicyandiamide degradation by soil bacteria. *Soil Biol Biochem* 3:385-391
- Selbie DR, Buckthought LE, Shepherd MA (2015) The challenge of the urine patch for managing nitrogen in grazed pasture systems. *Adv Agron* 129:229-292
- Shepherd M, Wyatt J, Welten B (2012) Effect of soil type and rainfall on dicyandiamide concentrations in drainage from lysimeters. *Soil Res* 50:67-75
- Singh J, Sagggar S, Giltrap DL, Bolan NS (2008) Decomposition of dicyandiamide (DCD) in three contrasting soils, and its effects on nitrous oxide emissions, soil respiratory activity, and microbial biomass – an incubation study. *Soil Res* 46:517-525
- Smith P, Martino D, Cai Z, Gwary D, Janzen H, Kumar P, McCarl B, Ogle S, O'Mara F, Rice C, Scholes B, Sirotenko O (2007) Agriculture In: Metz B, Davidson OR, Bosch PR, Dave R, Meyer LA (Eds) In *Climate Change 2007: Mitigation*. Cambridge University Press, Cambridge, UK and New York. .
- Turowski M, Deshmukh B (2004) Direct chromatographic method for determination of hydrogen cyanamide and dicyandiamide in aqueous solutions. *Anal Lett* 37:1981:1989
- Voroney RP, Brookes PC, Beyaert RP (2008) Soil microbial biomass C, N, P and S. In: Carter MR, Gregorich EG (Eds) *Soil sampling and methods of analysis*, 2<sup>nd</sup> edn. CRC Press, Boca Raton, FL, pp 637-651
- Weiske, A, Benckiser G, Herbert T, Ottow J (2001) Influence of the nitrification inhibitor 3,4-dimethylpyrazole phosphate (DMPP) in comparison to dicyandiamide (DCD) on nitrous oxide emissions, carbon dioxide fluxes and methane oxidation during three years of repeated application in field experiments. *Biol Fertil Soils* 34:109-117

- Wissemeier AH, Linzmeier W, Gutser R, Weigelt W, Schmidhalter U (2001) The new nitrification inhibitor DMPP (ENTECC®) – comparisons with DCD in model studies and field applications. Food security and sustainability. In: Horst WJ, Schenk MK, Bürket A, Claassen N, Flessa H, Frommer WB, Golbach H, Olf H-W, Römheld B, Sattelmacher B, Schmidhalter U, Schubert S, Wirén NV, Wittenmayer L (Eds) Plant Nutrition. Kluwer Academic Publishers, Dordrecht, the Netherlands, pp. 702-703 doi: 10.1007/0-306-47624-X
- Young SR, Black AS, Conyers MK (2002) Distribution of nitrification within surface soils under pasture. *Commun Soil Sci Plan* 33:1507-1518
- Zerulla W, Barth T, Dressel J, Erhardt K, von Locquenghien KH, Pasda G, Rädle M, Wissemeier AH (2001) 3,4-Dimethylpyrazole phosphate (DMPP) – a new nitrification inhibitor for agriculture and horticulture. *Biol Fertil Soils* 34:79-84
- Zhang H-J, Wu Z-J, Zhou Q-X (2004) Dicyandiamide sorption-desorption behaviour on soils and peat humus. *Pedosphere* 14:395-399

---

## Chapter 6

---

### **Plant acquisition and metabolism of the synthetic nitrification inhibitor dicyandiamide and naturally-occurring guanidine from agricultural soils**

Karina A. Marsden\*, Matthew Scowen, Paul W. Hill, Davey L. Jones & David R. Chadwick

This work has been published in *Plant and Soil*, and was presented at the BASF Top Science-Symposium 2015 on Nitrification Inhibitor Usage in Organic and Mineral Fertilizers in Dortmund, Germany.

Marsden, K.A., Scowen, M., Hill, P.W., Jones, D.L., Chadwick, D.R. 2015. Plant acquisition and metabolism of the synthetic nitrification inhibitor dicyandiamide and naturally-occurring guanidine from agricultural soils. *Plant and Soil* 395, 201-214.

KAM, MS, PWH, DLJ and DRC designed and conceived the experiment; KAM conducted the practical work; KAM analysed the results and prepared the manuscript.

## Abstract

There is increasing interest and use of nitrification inhibitors (NI) in agroecosystems, yet little is known of their fate *in planta*. Residues of the organic, N-rich NI, dicyandiamide (DCD), have been found in milk products following commercial application to pasture. We investigated whether plant acquisition and metabolism of DCD were consistent with plant-mediated transmission from soil to agricultural food products. Uptake rates, translocation to the shoot, degradation of the label within wheat tissue and availability within two soils of DCD and the structurally similar naturally occurring N-rich molecule, guanidine, were measured using  $^{14}\text{C}$  labelling. Under sterile conditions, over 2 h wheat took up (34 and 14  $\mu\text{mol g}^{-1}$  root DW  $\text{h}^{-1}$  at 1 mM: DCD and guanidine, respectively), translocated (7-15 and 19-22%) and metabolised (0.4 and 0.9% of uptake) DCD- and guanidine- $^{14}\text{C}$ . Both molecules were also acquired from soil by wheat despite concurrent soil sorption and microbial uptake. Both DCD and guanidine can be acquired and metabolised by graminaceous plants. Although probably not a significant route of N acquisition, plant uptake provides a direct route of DCD entry into the food chain.

**Key Words:** Bioavailability; Dicyandiamide (DCD); Guanidine; Mineralization; Nitrification inhibitor; Nitrogen cycle.

## 6.1 Introduction

Within agricultural soils, nitrification represents one of the dominant nitrogen (N) flow pathways and is responsible for generating  $\text{NO}_3^-$  which can be lost to the wider environment via leaching and denitrification (Zerulla et al. 2001). To reduce these N losses, effective management strategies are required to improve N use efficiency (NUE) within most agroecosystems. One potential solution to the problem is the application of synthetic or natural nitrification inhibitors (NI) to the soil to slow the conversion of  $\text{NH}_4^+$  to  $\text{NO}_3^-$  (Subbarao et al. 2012; Abalos et al. 2014).


Among the many identified NI, synthetic dicyandiamide (DCD;  $\text{C}_2\text{H}_4\text{N}_4$ ) is one of the most widely researched and one of the few used at a commercial scale (O'Callaghan et al. 2010; Liu et al. 2013). DCD has been investigated in a wide range of arable and livestock-based agroecosystems, where applications of DCD ( $10\text{-}30 \text{ kg ha}^{-1}$ ) have been shown to be effective in reducing nitrous oxide ( $\text{N}_2\text{O}$ ) emissions following spreading of either N fertilisers (Weiske et al. 2001; Di and Cameron 2006; Cui et al. 2011), livestock slurry (Hatch et al. 2005) or ruminant urine (Di and Cameron 2006; Dai et al. 2013; Barneze et al. 2015). DCD application to soil has also been shown to reduce  $\text{NO}_3^-$  leaching after the application of inorganic N fertilisers and from livestock urine patches (Di et al. 2009; Cui et al. 2011). A recent meta-analysis investigating the effect of soil-applied NI (including DCD), indicated that, on average, they result in an 8% increase in crop yield and a 13% increase in NUE (Abalos et al. 2014).

A range of application routes for DCD have been investigated including introduction to urine patches by oral administration to cattle (O'Connor et al. 2013; Welten et al. 2013), infusion of DCD into the rumen or abomasum of sheep (Ledgard et al. 2008), incorporation into fertiliser granules or addition of DCD in a biodegradable hydrogel to slow its release in soil (Minet et al. 2013). However, the simplest, least controversial and consequently most

widespread route is direct application to soil. Although direct application to soil is both practical and has demonstrable efficacy, this method leaves DCD susceptible to degradation by soil microbes, which may reduce persistence and increases in NUE. There is also the potential for removal from soil due to uptake by plant roots.

In 2012, low level residues of DCD were found in New Zealand dairy products from which plant interception and uptake of DCD with subsequent transfer to ruminant milk have been hypothesised to be key vectors in its contamination (Kim et al. 2012; Chen et al. 2014). Although DCD is not currently known to pose a significant risk to human health, the discovery of DCD in milk has led to a voluntary suspension of sale and use of DCD in New Zealand until international acceptable limits for its presence in milk products can be agreed (Ministry for Primary Industries 2013).

Whether plants are able to take up DCD is currently uncertain. Similarly, whether (if taken up) DCD persists in plant tissues or is rapidly degraded is unknown. Historically, DCD has been suggested for use as an organic fertiliser, although as opposed to intact plant uptake of this molecule, prior degradation to  $\text{NH}_4^+$  (and subsequent nitrification to  $\text{NO}_3^-$ ) was the presumed route of N supply to plants (Reddy 1964). Limited evidence from some >100 year old investigations (Hutchinson and Miller 1912) show increased soil residence of DCD in the absence of plants, and reported phytotoxicity and yield reduction at high rates of DCD addition ( $25 \text{ kg ha}^{-1}$ ) to clover, suggest that plant uptake does take place (Amberger 1986; Di and Cameron 2004; Macadam et al. 2003; Kelliher et al. 2008). The occurrence of the structurally similar molecule guanidine ( $\text{CH}_5\text{N}_3$ ; Fig. 6.1) and derivatives in a wide range of organisms including soil, plants and associated microbes may also suggest the existence of capacity for acquisition and assimilation of DCD by plants (Kato et al. 1986; Prescott and John 1996; Schulten and Schnitzer 1998; Kawano and Hwang 2010; G uthner et al. 2014).



**Figure 6.1** Molecular structure of the nitrification inhibitor, dicyandiamide (DCD; Panel A), and its naturally occurring analogue guanidine (Panel B), which are used in this study.

This investigation was predicated on the need to determine whether 1) the capacity of plants to take up DCD through their roots and translocate it to shoots is great enough, and 2) that the rate of subsequent degradation *in planta* is slow enough, to make plants a realistic route for the transmission of DCD from soil to agricultural food products. It also presents an opportunity to investigate earlier speculation on the potential of guanidine and similar compounds to be added to the rapidly growing list of routes via which plants have the capacity to obtain N from soil (Hutchinson and Miller 1912; Lewis 1936; Bollard 1966; Warren 2014).

We aimed to test the following hypotheses: 1) graminaceous plants have the capacity to take up both DCD and guanidine with their roots; 2) both DCD and guanidine can be metabolised by plants; 3) plants can take up both DCD and guanidine from soil; 4) the magnitude of competing substrate removal processes from soil (microbial uptake, mineralization and sorption) will be regulated by soil type.

## 6.2 Materials and methods

### 6.2.1 Soil properties

Two contrasting UK agricultural soils were used in this study (Table 6.1). The first was a mineral sandy loam textured Eutric Cambisol collected from a sheep-grazed fertilised grassland in North Wales (53°14'N, 4°01'W), while the second was an organic Sapric Histosol



**Table 6.1** Soil properties of Eutric Cambisol and Sapric Histosol used in soil rhizosphere microcosms. Values represent means  $\pm$  SEM,  $n = 4$ , letters indicate significant differences between the two soils and results are reported on a dry soil weight basis.

<b>Soil Property</b>	<b>Eutric Cambisol</b>	<b>Sapric Histosol</b>
Moisture Content (%)	25.0 $\pm$ 0.27 a	61.2 $\pm$ 0.09 b
Organic Matter (%)	7.70 $\pm$ 1.41 a	77.2 $\pm$ 0.47 b
Cation Exchange Capacity (meq 100 g <sup>-1</sup> )	14.8 $\pm$ 0.68* a	80.8 $\pm$ 0.80* b
pH	6.67 $\pm$ 0.14 a	6.37 $\pm$ 0.06 a
Electrical Conductivity ( $\mu$ S cm <sup>-1</sup> )	50.1 $\pm$ 1.85 a	103 $\pm$ 9.56 b
Total Carbon (g kg <sup>-1</sup> )	23.8 $\pm$ 1.50 a	391 $\pm$ 1.15 b
Total Nitrogen (g kg <sup>-1</sup> )	2.92 $\pm$ 0.07 a	26.5 $\pm$ 0.07 b
Total Organic Carbon (mg C kg <sup>-1</sup> )	88.2 $\pm$ 6.29 a	959 $\pm$ 79.1 b
Total Organic Nitrogen (mg N kg <sup>-1</sup> )	10.6 $\pm$ 3.70 a	37.8 $\pm$ 5.35 b
Microbial C (g kg <sup>-1</sup> )	1.58 $\pm$ 0.05 a	4.41 $\pm$ 0.20 b
Microbial N (g kg <sup>-1</sup> )	0.33 $\pm$ 0.03 a	0.96 $\pm$ 0.08 b
NH <sub>4</sub> <sup>+</sup> (mg N kg <sup>-1</sup> )	4.29 $\pm$ 0.32 a	5.90 $\pm$ 0.69 a
NO <sub>3</sub> <sup>-</sup> (mg N kg <sup>-1</sup> )	4.34 $\pm$ 0.20 a	27.8 $\pm$ 2.45 b
PO <sub>4</sub> <sup>3-</sup> (mg P kg <sup>-1</sup> )	9.98 $\pm$ 0.39 a	32.0 $\pm$ 6.68 b
K <sup>+</sup> (meq kg <sup>-1</sup> )	12.3 $\pm$ 2.36 a	8.46 $\pm$ 2.72 a
Na <sup>+</sup> (meq kg <sup>-1</sup> )	0.95 $\pm$ 0.05 a	5.08 $\pm$ 0.37 b
Ca <sup>2+</sup> (meq kg <sup>-1</sup> )	71.5 $\pm$ 11.0 a	619 $\pm$ 12.5 b

\*  $n = 3$

collected from an intensive arable production area in East Anglia (52°52'N, 0°47'W). At each site, four independent replicate soil samples (0-10 cm) were collected, sieved to pass 2 mm, and stored at 4°C in gas permeable polythene bags until the start of the experiment, where field

moist soils were utilised for experimentation. Soil moisture content was determined by oven drying (105°C, 24 h), and soil organic matter content by loss-on-ignition (450°C, 16 h; Ball, 1964). Total soil C and N content was determined using a CHN2000 Analyzer (Leco Corp., St. Joseph, MI). Soil pH and electrical conductivity (EC) were measured using standard electrodes in 1:2.5 (w/v) soil: distilled water extracts. Available C and N was determined using a 1:5 (w/v) soil-to-0.5 M K<sub>2</sub>SO<sub>4</sub> extract within 24 h of sample collection, according to Jones and Willett (2006). Total dissolved C and N in the extracts were determined with a Multi N/C 2100S (AnalytikJena, Jena, Germany). Microbial biomass C and N were determined by CHCl<sub>3</sub> fumigation-extraction according to Voroney et al. (2008) using K<sub>EC</sub> and K<sub>EN</sub> correction factors of 0.35 and 0.50, respectively. K<sub>2</sub>SO<sub>4</sub>-extractable P, NO<sub>3</sub><sup>-</sup> and NH<sub>4</sub><sup>+</sup> were determined using the colorimetric methods of Murphy and Riley (1962), Miranda et al. (2001) and Mulvaney (1996), respectively. Exchangeable cations were measured within 1:5 w/v soil-to-1 M NH<sub>4</sub>Cl extracts using a model 410 flame photometer (Sherwood Scientific Ltd., Cambridge, UK).

### 6.2.2 Plant uptake and translocation of DCD and guanidine under sterile conditions

DCD and guanidine uptake rates were determined under sterile conditions to determine if they were taken up intact (i.e. without prior microbial cleavage). Wheat seeds (*Triticum aestivum* var. Granary) were surface sterilised with 14% (v/v) NaClO and 80% ethanol and grown aseptically according to Hill et al. (2011). Briefly, surface sterilised seeds were germinated on agar containing 50% Murishage and Skoog basal medium, to screen for microbial contamination; after which they were transferred aseptically to Phytatrays (Sigma Aldrich, Gillingham, UK) containing sterile perlite and 50% Murishage and Skoog basal medium, supplemented with 10 mg l<sup>-1</sup> Na-metasilicate. Wheat plants were grown at 20°C, with a 16 h photoperiod and light intensity (PAR) of 500 μmol m<sup>-2</sup> s<sup>-1</sup>, until they had reached the third leaf stage. The roots of individual intact wheat plants ( $n = 4$ ) were placed in 12 ml of

sterile (0.2 µm-filtered) solution containing either  $^{14}\text{C}$ -DCD or  $^{14}\text{C}$ -guanidine (ca. 1 kBq ml<sup>-1</sup>; American radiolabelled Chemicals, St Louis, MO, USA), for a period of 2 h. A solution concentration of 1 mM DCD was chosen to reflect the DCD concentration found within soil solution in response to typical field application rates (10 kg ha<sup>-1</sup>) and two lower concentrations of 0.01 and 0.1 mM, were chosen to represent subsequent dilution of field applied DCD by diffusive and mass flow processes. Subsequently, the plant roots were thoroughly rinsed in 0.01 M CaCl<sub>2</sub>, followed by deionised water and the roots and shoots oven-dried (80°C, 24 h). To quantify the  $^{14}\text{C}$  content of the plants, the roots and shoots were separately combusted in an OX400 biological oxidizer (RJ Harvey, Hillsdale, NJ, USA), the  $^{14}\text{CO}_2$  captured in Oxosol scintillant (National Diagnostics, Atlanta, GA, USA) and  $^{14}\text{C}$  measured using a Wallac 1404 Liquid Scintillation Counter (Wallac EG&G, Milton Keynes, UK). To visualise the location of  $^{14}\text{C}$ -DCD and  $^{14}\text{C}$ -guanidine in the root and shoot tissues, the  $^{14}\text{C}$  distribution within intact plants was imaged using a Cyclone Plus phosphor-imaging system (PerkinElmer, Waltham, MA, USA) using an exposure time of 1 h.

### *6.2.3 Plant metabolism of DCD and guanidine*

To determine whether DCD or guanidine could be mineralized within the plant, sterile wheat seeds were prepared as described previously. The roots of intact plants were then placed in sterile (0.2 µm-filtered) solutions containing either  $^{14}\text{C}$ -DCD or  $^{14}\text{C}$ -guanidine (4 ml; 1 kBq ml<sup>-1</sup>; 0.01 mM). The plants were then placed in sterile 250 cm<sup>3</sup> polypropylene vessels through which moist air was passed at a rate of ca. 600 ml min<sup>-1</sup>. The outflow was bubbled through Oxosol scintillant to capture any respired  $^{14}\text{CO}_2$ . The Oxosol was changed after 1, 5, 10, 20, 40, 60, 90 and 120 min and captured  $^{14}\text{C}$  measured as described above.

#### 6.2.4 Plant uptake of DCD and guanidine from soil

To determine the uptake rates of DCD and guanidine from soil, wheat seeds ( $n = 4$ ) were individually sown into Eutric Cambisol or Sapric Histosol rhizosphere microcosms (240 mm long; internal diameter 8 mm) as described in Owen and Jones (2001). Each microcosm contained approximately 12 g FW soil. Plants were grown under the same conditions used for the sterile uptake study (Section 2.2), until they had reached the third leaf stage. At this point, solutions of  $^{14}\text{C}$ -DCD or  $^{14}\text{C}$ -guanidine (ca. 1 kBq ml<sup>-1</sup>; 0.01, 0.1 and 1 mM) were injected directly into the rhizosphere soil. A total of 4 injections (0.25 ml each) using 1 ml polypropylene syringes and 18 gauge needles at four depths (3, 9, 15 and 21 cm) were made, to facilitate an even distribution of solution within the microcosms. After 2 h, plants were removed from the microcosms and washed thoroughly in 0.01 mM CaCl<sub>2</sub>, followed by distilled water. After drying (80°C, 24 h), the  $^{14}\text{C}$  content of the root and shoot material was determined as described above. To estimate the quantity of root in contact with the injected solution, blue ink was injected as above into another set of microcosms ( $n = 4$ ). The sections of root exposed to the ink were removed, washed, dried (80 °C) and weighed.

This experiment was repeated under similar conditions, utilising unlabelled DCD (1 mM), in order to establish whether the intact DCD molecule could be detected in wheat shoot extracts via HPLC. After injection of the substrate and washing of the root material, the shoot was separated from the root. The shoots of four wheat plants were bulked ( $n = 3$ ) in order to increase the likelihood of ascertaining a measurable peak on the HPLC, and ground in 2 ml of DMSO in a borosilicate Griffiths tube; 1 ml of the DMSO extract was then evaporated to dryness under vacuum in a rotary evaporator. The pellet was resuspended in 0.5 ml of HPLC grade water, centrifuged (10 000 g), and analysed for DCD using an adapted method of Turowski and Deshmukh (2004) on a Varian ProStar HPLC, with UV detection at 215 nm. The column used was a Luna 5u SCX (250 x 4.6 mm; 5µm; 100 Å), DCD eluted after ca. four

minutes in this system. Chromatograms were compared to standards and the control plants (no injected DCD).

#### *6.2.5 DCD and guanidine mineralization within soils*

DCD and guanidine mineralization were determined, over the same time course as the rhizosphere uptake study (Section 2.5), by measuring the rate of  $^{14}\text{CO}_2$  evolution after the addition of 0.2 ml of  $^{14}\text{C}$ -DCD or  $^{14}\text{C}$ -guanidine (ca. 5 kBq ml $^{-1}$ ; 0.01, 0.1 and 1 mM) to 1 cm $^3$  of each soil ( $n = 4$ ). Soils were contained in a 10 cm $^3$  sealed glass vessel, with moist air flowing (ca. 100 ml min $^{-1}$ ) over the soil surface. Evolved  $^{14}\text{CO}_2$  was captured by passing the outflow through two consecutive 0.1 M NaOH traps (capture efficiency > 95%; Hill et al. 2007). Traps were changed after 1, 5, 10, 20, 40, 60, 90 and 120 min, and the activity in the solution determined by liquid scintillation counting after mixing with HiSafe 3 scintillant (PerkinElmer, Llantrisant, UK).

#### *6.2.6 DCD and guanidine microbial uptake and sorption within soils*

The amount of substrate remaining in soil solution (i.e. that remaining after microbial uptake and abiotic removal processes) was determined according to the centrifugal-drainage procedure of Hill et al. (2008). Briefly,  $^{14}\text{C}$ -DCD or  $^{14}\text{C}$ -guanidine (0.2 ml; ca. 5 kBq ml $^{-1}$ ; 0.01, 0.1 or 1 mM) was pipetted evenly onto the soil surface. After 1, 5, 10, 20, 40, 60, 90 and 120 min, soil solution was recovered by centrifugation (4000 g, 1 min, 20°C) and the amount of  $^{14}\text{C}$ -DCD or  $^{14}\text{C}$ -guanidine in the recovered soil solution determined by liquid scintillation counting as described above.

The amount of DCD or guanidine sorbed to the solid phase and present in soil solution was determined by performing 0.5 M K $_2$ SO $_4$  extracts over time. Briefly,  $^{14}\text{C}$ -DCD or  $^{14}\text{C}$ -guanidine was mixed with soil and incubated as described above. After 5, 10, 20, 40, 60, 90 or

120 min the soil was extracted with ice-cold 0.5 M K<sub>2</sub>SO<sub>4</sub> (1:5 w/v) by shaking (150 rev min<sup>-1</sup>) for 20 min. After shaking, the extracts were centrifuged (10,000 g, 10 min) and <sup>14</sup>C in the supernatant determined by liquid scintillation counting. The amount of substrate sorbed to the soils was calculated by deducting the soil solution pool from the K<sub>2</sub>SO<sub>4</sub> extractable pool; the amount of substrate taken up by soil microbes was determined by deducting the soil solution pool and the sorbed pool from the total substrate added.


#### 6.2.7 Statistical analysis

All measurements were performed in quadruplicate unless otherwise stated. Log-transformation of data, followed by a one-way ANOVA with Fisher's LSD post-hoc test, was used to compare the following: wheat root uptake rates of DCD and guanidine under sterile conditions, the percentage of substrate translocation to wheat shoot biomass under sterile conditions, wheat root uptake rates from soil rhizosphere microcosms (after correcting for root weight exposed to injected solutions), percentage of substrate translocation to shoot biomass from wheat grown in soil, microbial uptake and sorption of the substrate at the 2 h time point. Data were tested for normality (Shapiro-Wilk test) and homogeneity of variance assumptions (Levene's test) prior to conducting the ANOVA. Microbial mineralization data were analysed by one-way ANOVA as above, except data were not transformed. An independent samples t-test was used to compare plant metabolism of assimilated compounds at the 2 h time point, after testing the normality and homogeneity of variance assumptions. All statistical analyses were performed in IBM SPSS Statistics 20.0.

## 6.3 Results

### 6.3.1 DCD and guanidine uptake and tissue localization in sterile wheat plants

Rates of uptake of  $^{14}\text{C}$ -DCD and  $^{14}\text{C}$ -guanidine by wheat roots increased ( $p < 0.001$ ) with increasing concentration (from 0.01 to 1 mM) in sterile solution (Table 6.2; Fig. 6.2). Phosphorimaging revealed a fairly even distribution of the  $^{14}\text{C}$  label added as DCD within the shoot biomass (Table 6.2; Fig. 6.3), with the majority remaining in the roots. Phosphorimaging of  $^{14}\text{C}$  added as guanidine in the wheat plant showed an even distribution throughout the root system, but it was present predominantly in the lower regions of the shoot (Table 6.2; Fig.6.3). Wheat roots acquired 2.37% less ( $p < 0.01$ ) of the  $^{14}\text{C}$  label added as DCD in comparison to guanidine at a concentration of 0.01 mM. No differences ( $p > 0.05$ ) were observed at 0.1 mM, while 1.71% more ( $p < 0.01$ ) DCD was acquired in comparison to guanidine at 1 mM.



**Figure 6.2** Rates of  $^{14}\text{C}$ -dicyandiamide (DCD) and  $^{14}\text{C}$ -guanidine uptake by wheat (*Triticum aestivum*) roots grown in sterile hydroponic culture at three concentrations (0.01, 0.1 and 1 mM). Values represent means  $\pm$  SEM ( $n = 4$ ) and different letters indicate significant differences between means of compounds and concentrations (Fisher's LSD;  $p < 0.05$ ).

**Table 6.2** Summary of results for wheat uptake rates and tissue localization of  $^{14}\text{C}$ -DCD and  $^{14}\text{C}$ -guanidine (0.01, 0.1 and 1 mM) under sterile conditions and from Eutric Cambisol and Saprific Histosol rhizosphere microcosms following a 2 h chase period. Values represent means  $\pm$  SEM ( $n = 4$ ).

	Experimental condition	$^{14}\text{C}$ -DCD			$^{14}\text{C}$ -Guanidine		
		0.01 mM	0.1 mM	1 mM	0.01 mM	0.1 mM	1 mM
Uptake rate ( $\mu\text{mol g}^{-1}$ root DW $\text{h}^{-1}$ )	Sterile	0.23 $\pm$ 0.04	2.25 $\pm$ 0.35	34.0 $\pm$ 6.29	0.60 $\pm$ 0.13	2.51 $\pm$ 0.32	14.3 $\pm$ 4.72
Uptake rate ( $\text{nmol g}^{-1}$ root DW $\text{h}^{-1}$ )		Eutric Cambisol	3.02 $\pm$ 0.92	30.9 $\pm$ 6.17	353 $\pm$ 94.3	7.13 $\pm$ 1.88	49.8 $\pm$ 13.9
Uptake rate ( $\text{nmol g}^{-1}$ root DW $\text{h}^{-1}$ )	Saprific Histosol		2.18 $\pm$ 0.57	21.1 $\pm$ 1.72	174 $\pm$ 13.4	5.25 $\pm$ 0.77	47.2 $\pm$ 14.7
Shoot translocation (% of acquired label)	Sterile	7.46 $\pm$ 1.18	7.76 $\pm$ 0.50	15.0 $\pm$ 4.05	22.4 $\pm$ 3.96	19.0 $\pm$ 6.58	22.0 $\pm$ 3.18
Shoot translocation (% of acquired label)		Eutric Cambisol	9.13 $\pm$ 2.28	5.42 $\pm$ 0.51	7.37 $\pm$ 2.36	0.71 $\pm$ 0.25	1.58 $\pm$ 0.40
Shoot translocation (% of acquired label)	Saprific Histosol		7.80 $\pm$ 1.72	7.55 $\pm$ 1.48	7.16 $\pm$ 1.54	0.94 $\pm$ 0.13	1.10 $\pm$ 0.64






**Figure 6.3** Phosphorimages showing the distribution of  $^{14}\text{C}$ -dicyandiamide (DCD; Panel A) and  $^{14}\text{C}$ -guanidine (Panel B) within sterile wheat plants. The plants were exposed to each substrate (0.01 mM) for 2 h prior to imaging. The dark colour in the top images represents the distribution of  $^{14}\text{C}$ -label within the plant tissue while, the images underneath are the corresponding photographs of the same plants.

### 6.3.2 Plant mineralization of assimilated DCD and guanidine

Of the added  $^{14}\text{C}$ -DCD and  $^{14}\text{C}$ -guanidine taken up by the plant, only small amounts were mineralized to  $^{14}\text{CO}_2$  during the experiment. After 2 h,  $0.44 \pm 0.07\%$  of the acquired  $^{14}\text{C}$ -DCD and  $0.90 \pm 0.22\%$  of acquired  $^{14}\text{C}$ -guanidine had been metabolised to  $^{14}\text{CO}_2$  within the plant with no differences observed between the two substrates ( $p > 0.05$ ). For both compounds, respiration rates tended to be faster in the first 20 minutes of the incubation (Fig. 6.4).



**Figure 6.4** Cumulative mineralization of  $^{14}\text{C}$ -dicyandiamide (DCD) or  $^{14}\text{C}$ -guanidine in sterile wheat plants. Plants were exogenously supplied with each substrate (0.01 mM) for the whole 2 h monitoring period. Values represent mean  $\pm$  SEM ( $n = 4$ ).

### 6.3.3 DCD and guanidine uptake from soil rhizosphere microcosms

Wheat plants took up DCD- $^{14}\text{C}$  and guanidine- $^{14}\text{C}$  when grown in soil-filled microcosms (Fig. 6.5; Table 6.2), however, the rates were lower than those grown under sterile conditions. To evaluate the quantity of roots which were exposed to the injected  $^{14}\text{C}$  labelled substrates a blue ink tracer was injected into the microcosms. From this, we estimated that  $41 \pm 8$  and  $32 \pm 5\%$  of the total root biomass was exposed to the injected  $^{14}\text{C}$ -substrates in the Eutric Cambisol and Sapric Histosol soil, respectively.


Increasing the concentration of injected  $^{14}\text{C}$ -DCD into soil rhizosphere microcosms increased ( $p < 0.001$ ) rates of root uptake of the label in both soil types (Table 6.2). Although a consistently higher mean root uptake rate of DCD was observed from the Eutric Cambisol in

comparison to the Sapric Histosol at each concentration, differences were not significant ( $p > 0.05$ ). After 2 h, wheat roots had acquired  $0.88 \pm 0.12$ ,  $0.96 \pm 0.12$  and  $1.09 \pm 0.21\%$  of total applied DCD- $^{14}\text{C}$  (0.01, 0.1 and 1 mM respectively) from the Eutric Cambisol and translocated 5.42 - 9.13% to the shoot material (Table 6.2); the amount acquired by wheat from the Sapric Histosol was ca. half this:  $0.50 \pm 0.08$ ,  $0.53 \pm 0.06$  and  $0.44 \pm 0.06\%$  (0.01, 0.1 and 1 mM, respectively), with 7.16 - 7.80% translocated to the shoot (Table 6.2).

Increasing the concentration of injected  $^{14}\text{C}$ -guanidine into the soil-filled microcosms increased rates of root uptake of guanidine- $^{14}\text{C}$  in both soil types ( $p < 0.001$ ; Fig. 6.5). In the Eutric Cambisol, wheat roots acquired less ( $p < 0.01$ ) of the  $^{14}\text{C}$  label added as DCD in comparison to guanidine at 0.01 mM, yet uptake rates were similar ( $p > 0.05$ ) at 0.1 and 1 mM. In the Sapric Histosol, wheat roots acquired less ( $p < 0.05$ ) of the  $^{14}\text{C}$  label added as DCD in comparison to guanidine at 0.01 and 1 mM, however, uptake rates were similar at 1 mM ( $p > 0.05$ ; Table 6.2). After 2 h, wheat roots had acquired  $2.21 \pm 0.38$ ,  $1.54 \pm 0.33$  and  $1.00 \pm 0.09\%$  of total applied guanidine- $^{14}\text{C}$  (0.01, 0.1 and 1 mM respectively) from the Eutric Cambisol and translocated 0.71 – 2.09% to the shoot material (Table 6.2); the total amount acquired by wheat from the Sapric Histosol was  $1.26 \pm 0.03$ ,  $1.05 \pm 0.14$  and  $0.50 \pm 0.08\%$  (0.01, 0.1 and 1 mM, respectively), with 0.94 – 2.24% translocated to the shoot (Table 6.2). The percentage of acquired guanidine- $^{14}\text{C}$  which was translocated to the shoot was lower than that of DCD- $^{14}\text{C}$  in both soil types ( $p < 0.05$ ; Table 6.2).

Unlabelled DCD was detected within the wheat shoot extracts by HPLC, confirming intact uptake and translocation of the molecule to shoot material. The concentrations of DCD recovered in shoots following injection of 1 ml of 1 mM DCD and a 2 h incubation period were  $79.7 \pm 19.2$  and  $43.7 \pm 3.75$  nmol DCD  $\text{g}^{-1}$  shoot DW in the Eutric Cambisol and Sapric Histosol, respectively. DCD concentrations in wheat shoots calculated from the  $^{14}\text{C}$  data showed the same higher shoot recovery in the mineral soil and were of the same order at 25.8

$\pm 4.71$  and  $9.48 \pm 2.31$  nmol DCD  $\text{g}^{-1}$  shoot DW in the Eutric Cambisol and Sapric Histosol, respectively. Some disparity in the absolute values may have resulted from the different groups of plants used for the two methods of measurement.




**Figure 6.5** Uptake of  $^{14}\text{C}$ -dicyandiamide (DCD) or  $^{14}\text{C}$ -guanidine (0.01, 0.1 and 1 mM) by wheat plants grown in either a Eutric Cambisol (Panel A) or Sapric Histosol (Panel B), legend applies to both panels. Values represent means  $\pm$  SEM ( $n = 4$ ) and different letters indicate significant differences between means of different compounds, concentrations and between each soil type (Fisher's LSD;  $p < 0.05$ ).

#### 6.3.4 Microbial mineralization of DCD and guanidine within soil

Over 2 h, the amount of  $^{14}\text{C}$ -DCD and  $^{14}\text{C}$ -guanidine mineralized in soil was small (Fig. 6.6; Table 6.3). After 2 h in the Eutric Cambisol, less ( $p < 0.01$ )  $^{14}\text{C}$ -DCD was mineralized in comparison to  $^{14}\text{C}$ -guanidine, at 0.01 and 0.1 mM. Less  $^{14}\text{C}$ -DCD was also mineralized at 1 mM, however, differences were not significant ( $p > 0.05$ ). In the Sapric Histosol, more ( $p < 0.01$ )  $^{14}\text{C}$ -guanidine was mineralized in comparison to  $^{14}\text{C}$ -DCD at the lowest concentration (0.01 mM); at 0.1 and 1 mM less ( $p < 0.05$ )  $^{14}\text{C}$ -guanidine was mineralized in comparison to  $^{14}\text{C}$ -DCD. After 2 h, greater amounts of  $^{14}\text{C}$ -DCD were degraded in the Sapric Histosol in

comparison to the Eutric Cambisol at all concentrations ( $p < 0.05$ ); conversely, greater amounts of  $^{14}\text{C}$ -guanidine were degraded in the Eutric Cambisol as opposed to the Sapric Histosol at all studied concentrations ( $p < 0.05$ ).



**Figure 6.6** Mineralization of a)  $^{14}\text{C}$ -dicyandiamide (DCD) in a Eutric Cambisol b)  $^{14}\text{C}$ -DCD in Sapric Histosol c)  $^{14}\text{C}$ -guanidine in Eutric Cambisol and d)  $^{14}\text{C}$ -guanidine in Sapric Histosol at three concentrations (0.01, 0.1 and 1 mM). Legend applies to all panels and symbols represent means  $\pm$  SEM ( $n = 4$ ).


**Table 6.3** Summary of results for microbial mineralization, microbial uptake and sorption of  $^{14}\text{C}$ -DCD and  $^{14}\text{C}$ -guanidine (0.01, 0.1 and 1 mM) in a Eutric Cambisol and a Sapric Histosol after 2 h. Values represent means  $\pm$  SEM ( $n = 4$ ).

Soil Type		$^{14}\text{C}$ -DCD			$^{14}\text{C}$ -Guanidine		
		0.01 mM	0.1 mM	1 mM	0.01 mM	0.1 mM	1 mM
Mineralization (% of applied label)	Eutric Cambisol	1.58 $\pm$ 0.08	1.09 $\pm$ 0.08	0.89 $\pm$ 0.37	6.50 $\pm$ 0.45	2.13 $\pm$ 0.24	1.46 $\pm$ 0.21
Mineralization (% of applied label)	Sapric Histosol	2.82 $\pm$ 0.21	1.91 $\pm$ 0.08	2.40 $\pm$ 0.31	4.02 $\pm$ 0.63	1.12 $\pm$ 0.23	0.64 $\pm$ 0.04
Microbial uptake (nmol g <sup>-1</sup> soil DW)	Eutric Cambisol	0.19 $\pm$ 0.004	1.68 $\pm$ 0.14	15.2 $\pm$ 0.75	0.24 $\pm$ 0.02	1.48 $\pm$ 0.09	10.8 $\pm$ 0.98
Microbial uptake (nmol g <sup>-1</sup> soil DW)	Sapric Histosol	0.63 $\pm$ 0.01	5.97 $\pm$ 0.18	60.6 $\pm$ 1.64	0.68 $\pm$ 0.01	5.09 $\pm$ 0.14	50.1 $\pm$ 1.36
Sorption (nmol g <sup>-1</sup> soil DW)	Eutric Cambisol	0.28 $\pm$ 0.002	3.03 $\pm$ 0.14	32.0 $\pm$ 0.57	0.28 $\pm$ 0.02	3.78 $\pm$ 0.07	41.3 $\pm$ 0.04
Sorption (nmol g <sup>-1</sup> soil DW)	Sapric Histosol	0.60 $\pm$ 0.01	6.31 $\pm$ 0.16	62.2 $\pm$ 1.68	0.56 $\pm$ 0.01	7.33 $\pm$ 0.12	74.1 $\pm$ 1.06

### 6.3.5 Microbial uptake and sorption of DCD and guanidine within soil

The amount of  $^{14}\text{C}$ -substrate added as DCD or guanidine present in the soil solution pool and the  $\text{K}_2\text{SO}_4$ -extractable pool is shown in Figure 6.7. As the trends in the data were similar across all concentrations, only the 0.01 mM data are presented. Increasing applied DCD or guanidine concentration from 0.01 to 1 mM increased ( $p < 0.001$ ) the amount of DCD taken up by microbes in both soil types, at all concentrations (Table 6.3). Greater amounts of DCD and guanidine were taken up by microbes in the Sapric Histosol in comparison to the Eutric Cambisol at all studied concentrations ( $p < 0.001$ ). In the Eutric Cambisol a greater amount of added guanidine was taken up by microbes in comparison to DCD at 0.01 mM ( $p < 0.001$ ), and less guanidine was taken up in comparison to DCD at 0.1 and 1 mM ( $p < 0.001$ ). In the Sapric Histosol no difference was observed in the amount of substrate taken up by microbes as DCD or guanidine at 0.01 mM ( $p > 0.05$ ) and a lower amount of guanidine was taken up by microbes at 0.1 and 1 mM ( $p < 0.01$ ), in comparison to DCD (Table 6.3).

Increasing either DCD or guanidine concentration from 0.01 to 1 mM increased the amount of substrate sorbed to both soil types at all studied concentrations ( $p < 0.001$ ; Table 6.3). A greater amount of either DCD or guanidine sorbed to the Sapric Histosol in comparison to the Eutric Cambisol at all studied concentrations ( $p < 0.001$ ). No difference was found in the amount of DCD or guanidine sorbed in either soil type at 0.01 mM ( $p > 0.05$ ), but greater amounts of guanidine sorbed in comparison to DCD in both soil types at concentrations of 0.1 and 1 mM ( $p < 0.001$ ; Table 6.3).



**Figure 6.7** Amount of added <sup>14</sup>C label recovered in the soil solution pool and in the 0.5 M K<sub>2</sub>SO<sub>4</sub>-extractable pool from a) <sup>14</sup>C-dicyandiamide (DCD; 0.01 mM) added to Eutric Cambisol b) <sup>14</sup>C-DCD (0.01 mM) added to Sapric Histosol c) <sup>14</sup>C-guanidine (0.01 mM) in Eutric Cambisol and d) <sup>14</sup>C-guanidine (0.01 mM) in Sapric Histosol. Legend applies to all panels and symbols represent means ± SEM (*n* = 4).

## 6.4 Discussion

### 6.4.1 Plant uptake

Our results clearly demonstrate that in the absence of competing physical and biological soil processes (e.g. sorption, microbial uptake and microbial degradation), wheat roots can acquire DCD and guanidine from solution. Interestingly, rates of uptake ( $34.0 \pm 6.29 \mu\text{mol}$



DCD  $\text{g}^{-1}$  root DW  $\text{h}^{-1}$ ;  $14.3 \pm 4.72 \mu\text{mol}$  guanidine  $\text{g}^{-1}$  root DW  $\text{h}^{-1}$ ) at 1 mM under similar experimental conditions, were of a similar magnitude to other small N and C containing molecules found in soil (e.g. ca.  $25 \mu\text{mol}$   $\text{NO}_3^-$   $\text{g}^{-1}$  root DW  $\text{h}^{-1}$ ; ca.  $23 \mu\text{mol}$  alanine  $\text{g}^{-1}$  root DW  $\text{h}^{-1}$ ; Hill et al. 2013); however, rates of uptake were greater for DCD and guanidine when considering moles of N acquired ( $136 \mu\text{mol}$  DCD-N  $\text{g}^{-1}$  root DW  $\text{h}^{-1}$  and  $42.9 \mu\text{mol}$  guanidine-N  $\text{g}^{-1}$  root DW  $\text{h}^{-1}$  compared to  $25 \mu\text{mol}$   $\text{NO}_3^-$ -N  $\text{g}^{-1}$  root DW  $\text{h}^{-1}$  and  $23 \mu\text{mol}$  alanine-N  $\text{g}^{-1}$  root DW  $\text{h}^{-1}$ ; Hill et al. 2013).

The mechanisms of DCD uptake and subsequent translocation within plants remain unknown. Being a synthetic compound, we hypothesise that no DCD-specific membrane transporters exist, however, it is possible that uptake could be facilitated by transporter proteins for structurally similar, naturally occurring molecules. A recent study by Eggen and Lillo (2012), found that a pharmaceutical drug used for diabetes II, metformin, had a high bioconcentration factor within seeds of *Brassica napus*. Like DCD, the drug metformin is structurally similar to the naturally occurring molecule, guanidine. The proposed mechanism of metformin entry to plant cells was via organic cation transporters (OCT), which transport naturally occurring N-containing compounds across the cell membrane. Expression of OCT has been demonstrated within several *Arabidopsis* tissues, including root tissue (Lelandais-Brière et al. 2007; Kűfner and Koch 2008; Eggen and Lillo 2012). Substrates for plant OCT have not been well characterised, however, guanidine has been identified as a substrate for mammalian OCT (Cova et al. 2002). Assuming that DCD is an analogue of guanidine and acquired by the same mechanism, we would expect similar rates of uptake for both compounds. However, under sterile conditions we observed lower uptake rates of DCD compared to guanidine at 0.01 mM, similar rates at 0.1 mM and higher rates in comparison to guanidine at 1 mM. This suggests that the transporter affinity differs for the two compounds, and may also suggest there are other alternative transport pathways which need to be further investigated.

#### 6.4.2 *Plant translocation and assimilation*

Lower amounts of DCD-<sup>14</sup>C were recovered in the shoot in comparison to guanidine-<sup>14</sup>C. However, the DCD-<sup>14</sup>C appeared to be more evenly distributed within the shoot tissue. The low rates of mineralization of DCD- and guanidine-<sup>14</sup>C within the plant over the short incubation time employed here may suggest saturation of metabolic pathways and thus that plant capacity to use the C and N acquired in these molecules is low. Recovery of intact DCD in wheat shoots by HPLC perhaps supports this view. However, preferential accumulation of structurally similar metformin, in seeds of *Brassica napus* and the occurrence of other guanidine derivatives, including arginine, in seeds may suggest storage of DCD and guanidine i.e. a lack of metabolic pathways which are directly connected to respiration (Ngamga et al. 2007; Eggen and Lillo 2012).

#### 6.4.3 *Competition for DCD and guanidine in the rhizosphere*

Wheat was able to acquire ca. 0.5-1.0% of the <sup>14</sup>C applied as DCD and ca. 0.5-2% of that applied as guanidine from soil within 2 h when in competition with the rhizosphere microbial community and sorption processes. Although realistic DCD soil solution concentrations were chosen for this study, direct injection into the rhizosphere may have resulted in a greater amount of root surface area exposed to NI-containing soil solution than may be expected under field conditions. Actual DCD concentrations in the rhizosphere may vary according to NI application method, weather conditions, crop type, soil temperature and moisture and time since application. As only the C within molecules was isotopically labelled and detected within plant tissue, the results of the <sup>14</sup>C-DCD uptake study do not unequivocally demonstrate that DCD was taken up intact from soil without prior lysis by soil microbes, or whether it remained intact once inside the plant without further degradation or transformation

of the DCD molecule. Detection of unlabelled DCD within wheat shoots via HPLC, however, shows that intact uptake of DCD by plants took place.

Our results suggest that soil type is a regulator of DCD and guanidine bioavailability, with wheat acquiring consistent numerically (although not statistically) greater amounts of DCD-<sup>14</sup>C and guanidine-<sup>14</sup>C from the mineral Eutric Cambisol in comparison to the organic Sapric Histosol. A combination of a greater microbial uptake and sorption in the Sapric Histosol similarly suggests a lower availability for plant acquisition when compared to the Eutric Cambisol. The Sapric Histosol has a greater cation exchange capacity and more soil organic matter (which has been identified as important source of DCD binding domains (Jacinthe and Pichtel, 1992; Zhang et al. 2004)) in comparison to the Eutric Cambisol, which may have led to greater sorption in this soil. The Sapric Histosol also had a greater microbial biomass compared to the Eutric Cambisol on a soil weight basis, which may have caused greater amounts of DCD and guanidine to be taken up by microbes in this soil.

In comparison to some other simple C substrates (e.g. amino acids, sugars) and the level of sorption to soil particles, the mineralization of DCD and guanidine by soil microbes was very slow (Hill et al. 2008; Wilkinson et al. 2014). This might be expected considering that neither soil has previously been exposed to the synthetic nitrification inhibitor, DCD. The dissimilar pattern of DCD and guanidine mineralization, particularly at low concentrations in the Eutric Cambisol supports the tenet that DCD is not rapidly extracellularly degraded to guanidine. The greater degradation of DCD in the Sapric Histosol in comparison to the Eutric Cambisol may be explained by the greater microbial biomass on a soil weight for weight basis, although reasons for the lower rates of guanidine mineralization in the organic soil compared to the mineral soil are unclear. The low rates of guanidine uptake and mineralization at higher guanidine application rates ( $\geq 0.1$  mM) may also suggest that although naturally found in the microbial community, the capacity to internally assimilate high amounts of guanidine limits its

metabolic conversion to CO<sub>2</sub>. This supports the conclusions of Rajbanshi et al. (1992) in that the microbial community requires longer time periods than 2 h to adapt to utilising DCD in soil. However, we cannot exclude the possibility that both DCD and guanidine are metabolised largely by pathways which do not feed into respiration.

Our results support the suggestion of Kelliher et al. (2014) that plants may play a role in reducing the half-life of DCD within soils, and this may be more pronounced in mineral soils as opposed to organic soils where plant uptake rates were greater due to lower amounts being sorbed and taken up by microbes. Interestingly, they also suggest that plants are able to derive some additional N from soils where DCD has been applied, and that guanidine (and perhaps other similar N-rich molecules) can be added to the growing list of naturally occurring N forms which plants are able to acquire from soil and metabolise. However, we hesitate to suggest that this has a significant role in plant N nutrition.

#### *6.4.4 DCD entry into the food chain*

There are three obvious routes of entry for DCD into meat or milk products: by i) consumption of pasture which has intercepted DCD on the foliage during spray application of the NI ii) direct livestock consumption of pasture or forage (which has acquired DCD from the soil as demonstrated here) and iii) ingestion of soil particles containing DCD.

Based on a dairy cow consuming 15 kg DM day<sup>-1</sup> of grass (McDonald et al. 1996), with a standing biomass of 2000 kg DM ha<sup>-1</sup> (O'Donovan and Dillon 1999), following a spray application of DCD (10 kg ha<sup>-1</sup>), with 5% of the total applied DCD intercepted via the canopy (Kim et al. 2012) we estimate that 3.5 g of DCD cow<sup>-1</sup> day<sup>-1</sup> could be consumed if allowed to graze immediately following application. Assuming that wheat is representative of other grasses and extrapolation of uptake and respiration rates to 24 h (with 50% translocation to shoots over this period), we estimate that 0.43 mg DCD cow<sup>-1</sup> day<sup>-1</sup> and 0.15 mg DCD cow<sup>-1</sup>

day<sup>-1</sup> could be ingested following consumption of grass grown on the Eutric Cambisol and the Sapric Histosol, respectively. To estimate the amount of DCD consumed via soil ingestion we assume an even distribution of DCD in the top 10 cm of soil (soil bulk density of 1.10 and 0.31 g cm<sup>-3</sup> for the Eutric Cambisol and Sapric Histosol, respectively) with removal due to plant uptake as above, and microbial mineralization at 1.30% and 5.93% (Eutric Cambisol and Sapric Histosol, respectively) of applied DCD over 24 h (measured over this period by Scowen, M; unpublished). Based on 2% of the cow's DM intake being soil (Thornton and Abrahams, 1983), 2.68 and 9.1 mg DCD cow<sup>-1</sup> day<sup>-1</sup> could be ingested with soil in the Eutric Cambisol and Sapric Histosol, respectively. Our calculations are based on a number of assumptions and upscaling laboratory data to field conditions, therefore, care should be taken when considering these estimations. Further research is required to elucidate how application methods (e.g. liquid vs. granular formulations) and environmental conditions (e.g. movement of NI into the root zone due to irrigation or rainfall) may influence pasture plant acquisition of this NI under field conditions.

Based on our estimates the magnitude of risk for DCD entry into the food chain via cattle follows the trend: pasture interception of DCD > soil ingestion of DCD > plant acquisition of DCD. However, the greatest risk pathway (canopy interception) would be transient (e.g. under high rainfall) and easily controlled by preventing grazing immediately following DCD application to pasture. Controlling the amount of DCD ingested beyond this point represents more of a challenge for out-grazing livestock and plant uptake may be a more significant pathway over longer periods. The potential for DCD to enter raw foods destined for direct human consumption (e.g. salad vegetables) also requires further investigation as the concentrations of DCD may be higher than in dairy products. Further, if like some other guanidine derivatives DCD accumulates in seeds, application of DCD to arable grain, oilseed

or pulse crops could result in further direct DCD entry into the food chain due to the capacity of plants to acquire DCD from soil through roots (Ngamga et al. 2007; Eggen and Lillo, 2012).

## **6.5 Acknowledgments**

This work was supported by the UK Natural Environment Research Council under grant award NE/IO12303/1. Partial funding for the research was also provided by an EU Knowledge Economy Skills Scholarship.

## 6.6 References

- Abalos D, Jefferey S, Sanz-Cobena A, Guardia G, Vallejo A, (2014) Meta-analysis of the effect of urease and nitrification inhibitors on crop productivity and nitrogen use efficiency. *Agr Ecosyst Environ* 189: 136-144
- Amberger A, (1986) Potentials of nitrification inhibitors in modern N-fertilizer management. *J Plant Nutr Soil Sci* 149: 469-484
- Ball D F, (1964) Loss-on-ignition estimate of organic matter and organic carbon in calcareous soils. *J Soil Sci* 15: 84-92
- Barneze A S, Minet A P, Cerri C C, Misselbrook T, (2015) The effect of nitrification inhibitors on nitrous oxide emissions from cattle urine depositions to grassland under summer conditions in the UK. *Chemosphere* 119: 122-129
- Bollard E G, (1966) A comparative study of the ability of organic nitrogenous compounds to serve as sole sources of nitrogen for the growth of plants. *Plant Soil* 25: 153-166
- Cova E, Laforenza U, Gastaldi G, Sambuy Y, Tritto S, Faelli A, Ventura U, (2002) Guanidine transport across the apical and basolateral membranes of human intestinal Caco-2 cells is mediated by two different mechanisms. *J Nutr* 132: 1995-2003
- Cui M, Sun X, Hu C, Di H J, Tan Q, Zhao C, (2011) Effective mitigation of nitrate leaching and nitrous oxide emissions in intensive vegetable production systems using a nitrification inhibitor, dicyandiamide. *J Soils Sediments* 11: 722-730
- Chen X-H, Zhou L-X, Zhao Y-G, Pan S-D, Jin M-C, (2014) Application of nanoring amino-functionalized magnetic polymer dispersive micro-solid-phase extraction and ultra fast liquid chromatography-tandem mass spectrometry in dicyandiamide residue analysis of powdered milk. *Talanta* 119: 187-192

- Dai Y, Di H J, Cameron K C, He J-Z, (2013) Effects of nitrogen application rate and a nitrification inhibitor dicyandiamide on ammonia oxidizers and N<sub>2</sub>O emissions in a grazed pasture soil. *Sci Total Environ* 465: 125-135
- Di H, Cameron K C, (2004) Effects of temperature and application rate of a nitrification inhibitor, dicyandiamide (DCD), on nitrification rate and microbial biomass in a grazed pasture soil. *Aust J Soil Res* 42: 927-932
- Di H, Cameron K C, (2006) Nitrous oxide emissions from two dairy pasture soils as affected by different rates of a fine particle suspension nitrification inhibitor, dicyandiamide. *Biol Fertil Soils* 42: 472-480
- Di H, Cameron K C, Shen J P, He J Z, Winefield C S, (2009) A lysimeter study of nitrate leaching from grazed grassland as affected by a nitrification inhibitor, dicyandiamide, and relationships with ammonia oxidizing bacteria and archaea. *Soil Use Manage* 25: 454-461
- Eggen T, Lillo C, (2012) Antidiabetic II drug metformin in plants: uptake and translocation to edible parts of cereals, oily seeds, beans, tomato, squash, carrot and potatoes. *J Agric Food Chem* 60: 6929-6325
- Güthner T, Mertschenk B, Schulz B, (2014) Guanidine and Derivatives. In Wiley-VCH Ed. Ullmann's Fine Chemicals, Wiley-VCH Verlag GmbH & Co. KGaA, Boschstr. 12, 69469, Weinheim, Germany.
- Hatch D, Trinidad H, Cardenas L, Carneiro J, Hawkins J, Scholefield D, Chadwick D, (2005) Laboratory study of the effects of two nitrification inhibitors on greenhouse gas emissions from a slurry-treated arable soil: impact of diurnal temperature cycle. *Biol Fertil Soils* 41: 225-232
- Hill P W, Farrar J F, Jones D L, (2008) Decoupling of microbial glucose uptake and mineralization in soil. *Soil Biol Biochem* 40: 616-624



- Hill P W, Kuzyakov Y, Jones D, Farrar J, (2007) Response of root respiration and root exudation to alterations in root C supply and demand in wheat. *Plant Soil* 291: 131-141
- Hill P W, Marsden K A, Jones D L, (2013) How significant to plant N nutrition is the direct consumption of soil microbes by roots? *New Phytol* 199: 948-955
- Hill P W, Quilliam R S, DeLuca T H, Farrar J, Farrell M, Roberts P, Newsham K K, Hopkins D W, Bardgett R D, Jones D L, (2011) Acquisition and assimilation of nitrogen as peptide-bound and D-enantiomers of amino acids by wheat. *PLoS ONE*, 6, e19220. doi:10.1371/journal.pone.0019220
- Hutchinson H B, Miller N H J, (1912) The direct assimilation of inorganic and organic forms of nitrogen by higher plants. *J Agric Sci* 4: 282-302
- Jacinthe P A, Pichtel J R, (1992) Interaction of nitrapyrin and dicyandiamide with soil humic compounds. *Soil Sci Soc Am J* 56: 465-470
- Jones D L, Willett V B, (2006) Experimental evaluation methods to quantify dissolved organic nitrogen (DON) and dissolved organic carbon (DOC) in soil. *Soil Biol Biochem* 5: 991-999
- Kato T, Kondo T, Mizuno K, (1986) Occurrence of guanidino compounds in several plants. *Soil Sci Plant Nutr* 32: 487-491
- Kawano M, Hwang J, (2010) Influence of guanidine, imidazole, and some heterocyclic compounds on dissolution rates of amorphous silica. *Clays Clay Miner* 58: 757-765
- Kelliher F M, Clough T J, Clark H, Rys G, Sedcole J R, (2008) The temperature dependence of dicyandiamide (DCD) degradation in soils: A data synthesis. *Soil Biol Biochem* 40: 1878-1882

- Kelliher F M, van Koten C, Kear M J, Sprosen M S, Ledgard F S, de Klein C A M, Leticia S A, Luo J, Rys G, (2014) Effect of temperature on dicyandiamide (DCD) longevity in pastoral soils under field conditions. *Agric Ecosyst Environ* 186: 201-204
- Kim D-G, Giltrap D, Saggar S, Palmada T, Berben P, Drysdale D, (2012) Fate of the nitrification inhibitor dicyandiamide (DCD) sprayed on a grazed pasture: effect of rate and time of application. *Soil Res* 50: 337-347
- Küfner L, Koch W, (2008) Stress regulated members of the plant organic cation transporter family are localized to the vacuolar membrane. *BMC Res Notes*, 1: doi:10.1186/1756-0500-1-43.
- Ledgard S F, Menneer J C, Dexter M M, Kear M J, Lindsey S, Peters J S, Pacheco D, (2008) A novel concept to reduce nitrogen losses from grazed pastures by administering soil nitrogen process inhibitors to ruminant animals: A study with sheep. *Agric Ecosyst Environ* 125: 148-158
- Lelandais-Brière C, Jovanovic M, Torres G A M, Perrin Y, Lemoine R, Corre-Menguy F, Hartmann C, (2007) Disruption of AtOCT1, an organic cation transporter gene, affects root development and carnitine-related responses in Arabidopsis. *Plant J* 51: 154-164
- Lewis A H, (1936) The fertilizer value of some concentrated materials, particularly urea and guanidine and their nitrates and phosphates. *J Agric Sci* 26: 509-526
- Liu C, Wang K, Zheng X, (2013) Effects of nitrification inhibitors (DCD and DMPP) on nitrous oxide emission, crop yield and nitrogen uptake in a wheat-maize cropping system. *Biogeosciences Discuss* 10: 711-737
- Macadam X M B, del Prado A, Merino P, Esatvillo J M, Pinto M, González-Murua C, (2003) Dicyandiamide and 3, 4-dimethyl pyrazole phosphate decrease N<sub>2</sub>O emissions from

- grasslands but dicyandiamide produces deleterious effects in clover. *J Plant Physiol* 160: 1517-1523
- McDonald P, Edwards R, Greenhalgh J F D, (1996) *Animal Nutrition*. Essex, England: Longman Group Limited.
- Minet E P, O'Carroll C, Rooney D, Breslin C, McCarthy CP, Gallagher L, Richards K G, (2013) Slow delivery of a nitrification inhibitor (dicyandiamide) to soil using a biodegradable hydrogel of chitosan. *Chemosphere* 93: 2854-2858
- Ministry for Primary Industries (2013) [WWW document] URL <http://www.mpi.govt.nz/Portals/0/Documents/news-resources/news/letter-of-assurance.pdf> [accessed 1 September 2014].
- Miranda K M, Epsey M G, Wink D A, (2001) A rapid, simple, spectrophotometric method for simultaneous detection of nitrate and nitrite. *Nitric Oxide-Biol Ch* 5: 62-71
- Mulvaney R L, (1996) Nitrogen - inorganic forms. In: Sparks, D.L. ed. *Methods of Soil Analysis. Part 3*. Madison, WI, USA: Soil Science Society of America Inc., pp 1123-1184
- Murphy J, Riley J P, (1962) A modified single solution method for the determination of phosphate in natural waters. *Anal Chim Acta* 27: 31-36
- Ngamga D, Fanzo Free S N Y, Tane P, Fomum Z T, (2007) Millaurine A, a new guanidine alkaloid from seeds of *Millettia laurentii*. *Fitoterapia* 78: 276-277
- Owen A G, Jones D L, (2001) Competition for amino acids between wheat roots and rhizosphere microorganisms and the role of amino acids in plant N acquisition. *Soil Biol Biochem* 33: 651-657
- O'Callaghan M, Gerard E M, Carter P E, Lardner R, Sarathchandra U, Burch G, Ghani A, Bell N, (2010) Effect of the nitrification inhibitor dicyandiamide (DCD) on

- microbial communities in a pasture soil amended with bovine urine. *Soil Biol Biochem* 42: 1425-1436
- O'Connor P J, Hennessy D, Lynch M B, Slettery H, Lewis E, (2013) The effect of dicyandiamide on rumen and blood metabolites, diet digestibility and urinary excretion. *Livest Sci* 155: 35-37
- O'Donovan M O, Dillon P, (1999) Measurement of grassland management practice on commercial dairy farms: TEAGASC: Co. Cork, Ireland. 4351:1
- Prescott A G, John P, (1996) Dioxygenases: molecular structure and role in plant metabolism. *Annu Rev Plant Phys* 47: 245-271
- Rajbanshi S S, Benckiser G, Ottow J C G, (1992) Mineralization kinetics and utilization as an N source of dicyandiamide (DCD) in soil. *Naturwissenschaften* 79: 26-27
- Reddy G R, (1963) Effect of varying quantities of dicyandiamide on the utilization of nitrogen by several crops from sodium nitrate and ammonium sulphate. *J Agr Sci* 62: 35-38
- Schulten H R, Schnitzer M, (1998) The chemistry of soil organic nitrogen: a review. *Biol Fert Soils* 26: 1-15
- Subbarao G V, Sahrawat K L, Nakahara K, Ishikawa T, Kishii M, Rao I M, Hash C T, George T S, Rao P S, Nardi P, Bonnett D, Berry W, Suenaga K, Lata J C, (2012) Chapter six - Biological nitrification inhibition - a novel strategy to regulate nitrification in agricultural systems. In: Sparks, D.L., eds. *Advances in Agronomy*, Academic Press 114: 249-302
- Thornton I, Abrahams P, (1983) Soil ingestion – a major pathway of heavy metals into livestock grazing contaminated land. *Sci Total Environ* 28: 287-294
- Turowski M, Deshmukh B, (2004) Direct chromatographic method for determination of hydrogen cyanamide and dicyandiamide in aqueous solutions. *Anal Lett* 9: 1981-1989

- Voroney R P, Brookes P C, Beyaert R P, (2008) Soil microbial biomass C, N, P and S. In: Carter, M.R and Gregorich, E.G., eds. Soil sampling and methods of analysis, 2<sup>nd</sup> edn. Boca Raton, FL, USA: CRC Press, pp. 637-651.
- Warren C R, (2014) Organic N molecules in the soil solution: what is known, what is unknown and the path forwards. *Plant Soil* 375: 1-19.
- Weiske A, Benckiser G, Herbert T, Ottow J, (2001) Influence of the nitrification inhibitor 3,4-dimethylpyrazole phosphate (DMPP) in comparison to dicyandiamide (DCD) on nitrous oxide emissions, carbon dioxide fluxes and methane oxidation during 3 years of repeated application in field experiments. *Biol Fert Soils* 34: 109-117.
- Welten B G, Ledgard S F, Schipper L A, Judge A A, (2013) Effect of amending cattle urine with dicyandiamide on soil nitrogen dynamics and leaching of urinary-nitrogen. *Agr Ecosyst Environ* 167: 12-22
- Wilkinson A, Hill P W, Farrar J F, Jones D L, Bardgett R D, (2014) Rapid microbial uptake and mineralization of amino acid and peptide-N along a grassland productivity gradient. *Soil Biol Biochem* 72: 75-73
- Zerulla W, Barth T, Dressel J, Erhardt K, von Locquenghien K H, Pasda G, Rädle M, Wissemeier A H, (2001) 3,4-Dimethylpyrazole phosphate (DMPP) – a new nitrification inhibitor for agriculture and horticulture. *Biol Fert Soils* 34: 79-84.
- Zhang H J, Wu Z J, Zhou Q X, (2004) Dicyandiamide sorption-desorption behaviour on soils and peat humus. *Pedosphere* 14: 395-399

---

## Chapter 7

---

### **Effect of time-since DMPP application on mitigating N<sub>2</sub>O emissions from sheep urine patches**

Karina A. Marsden\*, Davey L. Jones, David R. Chadwick

KAM, DLJ and DRC designed and conceived the experiment, KAM conducted the experimental work, analysed the results and prepared the manuscript.

## Abstract

Nitrification inhibitors are a potential technology to mitigate N<sub>2</sub>O emissions from the urine patches of grazing animals. At present there is limited information regarding the efficacy of the nitrification inhibitor 3,4-dimethylpyrazole phosphate (DMPP) in reducing N<sub>2</sub>O emissions from ruminant urine patches, as opposed to the well-studied nitrification inhibitor, dicyandiamide. In practical terms, urine patches would be deposited to soil at various times following the application of a nitrification inhibitor to soil. We hypothesised that the effectiveness of DMPP in reducing cumulative N<sub>2</sub>O emissions would decrease the longer the time since DMPP application. This study utilised an automated closed chamber technique, to monitor fluxes of N<sub>2</sub>O from sheep urine patches (725 kg N ha<sup>-1</sup>; 150 ml; 300 cm<sup>2</sup>) deposited to a Eutric Cambisol, where DMPP was applied (1 kg ha<sup>-1</sup>) on the same day, 2 weeks before and 4 weeks before urine application. Fluxes were monitored continuously from 4 weeks before to 9 weeks after urine application. DMPP was found to be ineffective in reducing cumulative N<sub>2</sub>O emissions when applied prior to, or even when applied at the same time as the sheep urine. Some effect of DMPP in delaying the accumulation of NO<sub>3</sub><sup>-</sup> was observed, with effects being greater the shorter the time since DMPP application. The temporal dynamics of N<sub>2</sub>O fluxes were also altered where DMPP was applied on the same day as the urine. The N<sub>2</sub>O emission factors ranged from ca. 0.33-2.49% of the applied N, with an average of 0.81 ± 0.22% across all chambers. This was lower than the default IPCC emission factor of 1% for sheep, despite the fairly high N application. Heterogeneity in soil conditions were deemed responsible for the large spatial variability of N<sub>2</sub>O emissions observed in this study. Further work is required to determine whether DMPP can reduce emissions from urine patches at a range of N concentrations, under different climatic conditions and with a range of nitrification inhibitor application methods.

**Key words:** diurnal; grassland; greenhouse gases; nitrogen cycle; sustainable intensification.

## 7.1 Introduction

The agricultural sector is the second largest source of greenhouse gases in the UK (following the energy sector; Webb et al., 2014), contributing ca. 8% of the UK's total (direct and indirect) anthropogenic emissions (Gill et al., 2010). The agricultural sector will need to contribute to the ambitious GHG emissions reduction target set by the UK Climate Change Act, i.e. to reduce emissions by 80% from base year levels, by 2050. The sustainable management of livestock excreta will form an important component of reducing agricultural emissions, especially under projected increases in global meat consumption (Davidson, 2009). Pastures cover a greater portion of the earth's surface compared to arable soils (7% and 3%, respectively; Oenema et al., 2005) and urine patches deposited to pastures are known hotspots for N cycling, where microbial transformations (e.g. nitrification and denitrification) can result in elevated N<sub>2</sub>O emissions (de Klein and van Logtestijn, 1994; Carter, 2007). Excreta (dung and urine) deposited to pasture soils account for ca. 40% of the global N<sub>2</sub>O emissions arising from animal production systems (Oenema et al., 2005).

The use of nitrification inhibitors within grazed pastures offers a potential means of reducing N losses whilst promoting nitrogen use efficiency (Di and Cameron, 2002; Di and Cameron, 2007). By delaying the first and rate-limiting step of nitrification (the oxidation of NH<sub>4</sub><sup>+</sup> to NO<sub>2</sub><sup>-</sup>; Chaves et al., 2006; Fiencke and Bockel, 2006; Benckiser et al., 2013), nitrification inhibitors enable nitrogen to persist in the soil in the ammoniacal form for longer. This increases opportunity for plant acquisition of applied N, reduces NO<sub>3</sub><sup>-</sup> leaching and can reduce N<sub>2</sub>O emissions from both nitrification and denitrification processes. Two of the most widely researched nitrification inhibitors include dicyandiamide (DCD) and 3,4-dimethylpyrazole phosphate (DMPP; Liu et al., 2013), where the latter is a newer product, developed by BASF (Ludwigshafen, Germany; Barth et al., 2001; Zerulla et al., 2001). In addition, reviews on the efficacy of nitrification inhibitors have revealed a greater effectiveness



in reducing N<sub>2</sub>O emissions from grasslands, in comparison to other land use types (Akiyama et al., 2010; Gilsanz et al., 2016).

There have been a greater number of studies exploring the effect of DCD in reducing N<sub>2</sub>O emissions from urine patches compared to DMPP (e.g. Di and Cameron, 2003; deKlein et al., 2011; Misselbrook et al., 2014; Bell et al., 2015). The use of an alternative nitrification inhibitor to DCD within pasture soils may be necessary, due to evidence of plant acquisition of DCD, which may lead to contamination of milk products from dairy pasture soils (Marsden et al., 2015). DMPP may be a suitable alternative to DCD for use in pasture soils as it has a lower phytotoxicity and slower degradation rates in soil (Weiske et al., 2001; Zerulla et al., 2001).

DMPP has been shown to be effective in reducing N<sub>2</sub>O emissions from fertilisers (Linzmeier et al., 2001; Menéndez et al., 2012; Bell et al., 2015) and livestock slurry (Chadwick et al., 2001; Hatch et al., 2005; Merino et al., 2005) but there are few studies investigating DMPP efficacy where ruminant urine is the nitrogen source (e.g. Di and Cameron, 2011; Di and Cameron, 2012; Gilsanz et al., 2016). When applied at the same time as a cattle urine application (1000 kg N ha<sup>-1</sup>), a liquid DMPP (5 kg ha<sup>-1</sup>) application reduced N<sub>2</sub>O emissions by 66% (Di and Cameron, 2012). Misselbrook et al. (2014) and Barneze et al. (2015) investigated the N<sub>2</sub>O reduction efficacy of pyrazole derivatives (which are structurally similar to DMPP) applied to cattle urine, however, no significant reduction in N<sub>2</sub>O emissions were observed. DMPP performance has been found to be best under cold and wet conditions (Menéndez et al., 2012), therefore, whether DMPP is effective under warmer soil conditions is still unclear.

This study was carried out to determine whether DMPP is effective in reducing N<sub>2</sub>O emissions arising from pasture soils influenced by sheep urine, under UK summer conditions. Given that sheep will deposit urine at various times following an application of DMPP, and that cost and labour associated with application will increase with the need for repeated

applications, we wanted to assess the duration of DMPP efficacy in reducing N<sub>2</sub>O emissions. We used a high frequency automated GHG monitoring system to evaluate the efficacy of a liquid DMPP application in reducing N<sub>2</sub>O emissions from sheep urine applied immediately after, 2 weeks after and 4 weeks after DMPP application. We hypothesised that N<sub>2</sub>O reduction efficacy would reduce with the greater the time-since application, due to degradation and immobilization of the nitrification inhibitor within the soil.

## **7.2 Materials and Methods**

### *7.2.1 Field site*

The study site was established on a lowland grazed grassland at Henfaes Research station, Abergwyngregyn, North Wales (53°14'N, 4°01'W). The soil at the site is classified as a Eutric Cambisol, and is of glacial till (deposited ca. 10 000 years ago) in origin. The recent history includes moderate fertilizer applications, light grazing by Welsh Mountain ewes and reseeded in 1990 with a *Lolium perenne* L. and *Trifolium repens* L. mix. To prevent the effects of recent livestock excretal depositions on monitored gas fluxes, livestock were excluded from the study site five months prior to treatment application.

### *7.2.2 Soil sampling and analysis*

To provide the background soil characteristics (Table 7.1) of the study site, soil was sampled (0-10 cm, 1.5 cm diameter) in triplicate, at the block level ( $n = 3$ ). The bulk density was measured by inserting a 100 cm<sup>3</sup> metal ring into the ground, oven-drying the removed soil core (105°C; 24 h), and sieving (< 2 mm) the dry soil to remove and weigh stones. Gravimetric moisture content was determined by oven drying soil (105°C; 24 h) and organic matter was determined by loss-on-ignition (450°C; 16 h; Ball, 1964). Soil pH and electrical conductivity (EC) were measured with standard electrodes, submerged in 1:2.5 (w/v) soil-to-distilled water

suspensions. The total soil C and N content of oven-dried and ground samples were determined using a TruSpec® Analyzer (Leco Corp., St. Joseph, MI); pasture foliar C and N contents, before and during the study, were also analysed using the same instrument. Dissolved (1:5; w/v; soil-to-0.5M K<sub>2</sub>SO<sub>4</sub>) total C and N concentrations in soil were determined by the method of Jones and Willet (2006), using a multi N/C 2100S analyser (AnalytikJena, Jena, Germany). The CHCl<sub>3</sub>-fumigation-extraction method of Voroney et al. (2008), was used to determine microbial biomass C and N, where the resulting C and N in the extracts were analysed as described previously, using K<sub>EC</sub> and K<sub>EN</sub> correction factors of 0.35 and 0.5, respectively. Concentrations of P, NO<sub>3</sub><sup>-</sup> and NH<sub>4</sub><sup>+</sup> were determined in 0.5 M K<sub>2</sub>SO<sub>4</sub> soil (0-5 cm) extracts using colorimetric methods described in Murphy and Riley (1962), Miranda et al. (2001) and Mulvaney (1996), respectively. Exchangeable cations (Na, K, Ca) were measured using a Sherwood Model 410 flame photometer (Sherwood Scientific Ltd., Cambridge, UK) in 1 M NH<sub>4</sub>Cl extracts (1:5, w/v, soil-to-1 M NH<sub>4</sub>Cl).

### 7.2.3 Sheep urine collection and analysis

Welsh Mountain ewes ( $n = 6$ ) were housed in individual pens, containing slatted plastic flooring specifically designed for sheep (Rimco Ltd., Yorkshire, UK). The pen flooring was slightly raised above the ground, allowing room for urine collection trays. The sheep were allowed free access to water and provided with a diet (*ad libitum*) of freshly cut pasture similar in composition to the field site. During the urine collection period individual urine volumes were recorded, the samples were filtered (0.45 µm) and frozen (unacidified). The urine collected from all sheep were bulked (to provide sufficient volume of homogenous composition) and thoroughly mixed before application to the soil. Total dissolved organic C, total N, P, NO<sub>3</sub><sup>-</sup>, NH<sub>4</sub><sup>+</sup> and cations in the sheep urine were analysed as described for the soil extracts, and the urea content was determined via the method of Orsonneau et al. (1992).

**Table 7.1** Properties of the Eutric Cambisol at the field site receiving urine and DMPP applications. Values represent means  $\pm$  SEM ( $n = 3$ ) and results are reported on a dry weight basis.

Eutric Cambisol properties (0-10cm)	
Texture	Sandy Clay Loam
Bulk density ( $\text{g cm}^{-3}$ )	$1.08 \pm 0.05$
Gravimetric moisture content (%)	$21.5 \pm 0.9$
Organic matter (%)	$9.99 \pm 0.32$
pH	$5.91 \pm 0.03$
EC ( $\mu\text{S cm}^{-1}$ )	$32.7 \pm 0.9$
Total C (%)	$3.99 \pm 0.01$
Total N (%)	$0.34 \pm 0.01$
C:N ratio	$11.6 \pm 0.2$
Dissolved organic C ( $\text{mg C kg}^{-1}$ )	$170 \pm 25$
Total dissolved N ( $\text{mg N kg}^{-1}$ )	$23.1 \pm 2.8$
Microbial biomass C ( $\text{g C kg}^{-1}$ )	$2.44 \pm 0.30$
Microbial biomass N ( $\text{mg N kg}^{-1}$ )	$171 \pm 10$
Extractable $\text{NO}_3^-$ ( $\text{mg N kg}^{-1}$ )	$3.97 \pm 0.45$
Extractable $\text{NH}_4^+$ ( $\text{mg N kg}^{-1}$ )	$3.23 \pm 0.99$
Extractable P ( $\text{mg P kg}^{-1}$ )	$3.37 \pm 0.44$
Exchangeable Na ( $\text{mg kg}^{-1}$ )	$43.8 \pm 1.1$
Exchangeable K ( $\text{mg kg}^{-1}$ )	$140 \pm 12$
Exchangeable Ca ( $\text{g kg}^{-1}$ )	$1.07 \pm 0.09$

#### 7.2.4 Experimental design, treatments and application dates

The study was laid out in a randomised block design, consisting of five treatments ( $n = 3$ ) as follows: 1) no urine application (control), 2) sheep urine only, 3) sheep urine plus DMPP applied at the same time, 4) sheep urine plus DMPP applied 2 weeks before urine application, and 5) sheep urine + DMPP applied 4 weeks before urine application, hereafter referred to as NU, SU, SU + DMPP, SU + DMPP 2, and SU + DMPP 4, respectively. For each treatment, urine patches were duplicated next to each flux chamber, allowing one patch for monitoring

greenhouse gases from undisturbed soils and another for soil sampling. Sheep urine (150 ml) was poured onto respective plots using a fixed template as an area guide (300 cm<sup>2</sup>), to represent an average sheep urine volume and patch size (Doak, 1952). Liquid applications of DMPP (250 ml; 0.1 g DMPP l<sup>-1</sup>) were sprayed (a normal nitrification inhibitor application method for grazed grasslands; Di and Cameron, 2011) by hand onto respective plots (2500 cm<sup>2</sup> across the chamber basal area), at a frequently used equivalent rate of 1 kg DMPP ha<sup>-1</sup> (Zerulla et al., 2001). The first DMPP treatment application to respective plots (4 weeks before urine application; SU + DMPP 4) took place on 29/6/2015, the second DMPP treatment application (2 weeks before urine application; SU + DMPP 2) was applied on 13/7/2015, and the final DMPP treatment application (SU + DMPP) took place on 27/7/2015. Following this, sheep urine was applied to all plots except for the control on the same date as the final DMPP application.

#### *7.2.5 Greenhouse gas emission monitoring*

An automated high frequency (8 flux measurements per day/night cycle) greenhouse gas monitoring system (Queensland University of Technology, Institute for Future Environments, Brisbane, Australia), as described in Scheer et al. (2014), was used to measure fluxes of N<sub>2</sub>O from the urine treated soils. As the automated system consisted of 12 chambers only (i.e. enough for 4 treatments, where  $n = 3$ ), fluxes of N<sub>2</sub>O from the NU treatment were measured manually using the conventional static chamber technique (1 flux measurement as close to daily as possible); fluxes of N<sub>2</sub>O were expected to be lower and less variable from the NU treatment, compared to urine treatments, as no N was applied. We, therefore, expected the difference in measurement techniques to cause minimal bias in the results.

For the automated chambers, stainless steel chamber bases (2500 cm<sup>2</sup> basal area) were inserted into the soil (10 cm depth), and chambers (50 cm × 50 cm × 15 cm) were clamped to

the bases. The chambers opened and closed via pneumatic actuators, where one block of chambers would close sequentially every 1 h. While one block of chambers were closed, the other two blocks of chambers were open, allowing ambient conditions to be restored within the chambers. Chamber headspace gas samples were pumped (ca. 200 ml min<sup>-1</sup>) to a sampling unit, through Teflon tubing. The sampling unit housed a LI-COR LI-820 non-dispersive infrared gas analyser (LI-COR, St Joseph, MI, USA) to measure CO<sub>2</sub>. Samples were then passed through an Ascarite (sodium hydroxide coated silica) filter before being pumped to a gas chromatograph (SRI 8610C, Torrance, USA), equipped with a <sup>63</sup>Ni electron capture detector (ECD) and flame ionization detector (FID) to measure N<sub>2</sub>O and CH<sub>4</sub> concentrations, respectively. Over the 1 h chamber closure period, each chamber ( $n = 4$ ) was sampled at 3 minute intervals followed by a calibration standard (500 ppb N<sub>2</sub>O; 880 ppm CO<sub>2</sub>; 3 ppm CH<sub>4</sub>;  $\pm 2$  % of the certified value, BOC gases, Liverpool, UK). For each chamber, this results in headspace concentration measurements once every 15 min, over the course of 1 h. A full cycle takes 3 hours to complete, resulting in (where uninterrupted measurement occurs) 8 flux measurements per 24 h.

The manual gas samples from the NU treatment were taken by placing polypropylene upturned buckets (ca. 26 cm in height) onto collars (26 cm diameter), inserted to a depth of 10 cm. The chambers were fitted with a re-sealable vent to allow pressure equalisation when placing chambers onto bases, and were fitted with Suba-Seals ® (Sigma, Gillingham, UK) to allow headspace gas sampling. Headspace samples were taken with a syringe every 15 min over the period of 1 h, to match the automated system. Samples were stored in pre evacuated 20 ml glass vials, before being analysed on a Perkin Elmer 580 Gas Chromatograph served by a TurboMatrix 110 auto sampler (Perkin Elmer, CT, USA). The system contained two Elite-Q PLOT megabore capillary columns, with split injection. One column was linked to an ECD (375°C), and the other to an FID (350°C) with methaniser, and the oven temperature was 50°C.

Fluxes (from both methods) were measured for a total of 13 weeks, (beginning 4 weeks before urine application and ending 9 weeks after urine application). Prior to the start of the experiment, standard gases were analysed by both GCs (laboratory and field system) and shown to be not significantly different (data not shown).

#### *7.2.6 Ancillary measurements*

To support the emission measurements, several ancillary measurements were made. Soil cores (0-5 cm) were taken from replicated plots on several dates throughout the experiment. Gravimetric moisture content and 1:5 (w/v) soil-to-0.5 M K<sub>2</sub>SO<sub>4</sub> extractions were conducted on the resulting soil samples. The extracts were analysed for NO<sub>3</sub><sup>-</sup>, NH<sub>4</sub><sup>+</sup>, extractable dissolved organic C and total N, as described previously. In order to allow sampling of the soil solution from within the chambers in a non-destructive manner, Rhizon suction samplers (2.5 cm diameter, 5 cm porous part, 12 cm length tubing; Rhizosphere Research Products, Wageningen, Netherlands) were inserted into the centre of the chamber plots. Soil solution samples were collected periodically from the Rhizons through a needle inserted into an evacuated 9 ml plastic container, when soil conditions were moist enough for successful sample collection. The soil solution samples were also analysed for NO<sub>3</sub><sup>-</sup>, NH<sub>4</sub><sup>+</sup>, dissolved organic C and dissolved total N, as described previously. Soil moisture sensors (Acclima SDI-12 digital TDT® sensors; Acclima Inc., ID, USA) were inserted diagonally through the urine patch (but were in situ several weeks prior to urine application) within the chambers, in order to measure soil moisture non-destructively. The pasture foliar N content was measured within the chambers at block level prior to urine application, and pasture biomass and foliar N content was measured in each chamber at 3, 6 and 9 weeks following urine application. The grass was also cut in the duplicated plots at the same time as the chambers to ensure consistency. Air temperature was monitored inside and outside the chambers using ThermoChron iButtons®

(iButtonLink, LLC, WI, USA) logging temperature every 1 h. Rainfall, soil (0-10 cm) and air temperature were monitored hourly at a weather station near to the field site.

### 7.2.7 Data processing and statistical analysis

Fluxes of N<sub>2</sub>O were calculated from each chamber as described in Scheer et al. (2014). The combination of three or four data points on the slope of N<sub>2</sub>O increase or decrease, over the 1 h chamber closure period, which provided the best R<sup>2</sup> value was used in the final reported fluxes. Cumulative N<sub>2</sub>O emissions were calculated by trapezoidal integration (from the point of urine application onwards) and emission factors were calculated using the following equation:

$$EF = (\text{treatment N}_2\text{O-N} - \text{control N}_2\text{O-N}) / \text{Total N applied} \times 100\% \quad (\text{Eqn. 1})$$

Differences in cumulative N<sub>2</sub>O emissions were analysed by Welch's test followed by Games-Howell pairwise comparisons, due to the data not conforming to homogeneity of variance assumptions required for a parametric statistical procedure, even after data transformation. After assessing the normality (Shapiro Wilk test) and homogeneity of variance (Levene's test) assumptions, the N<sub>2</sub>O emission factors were compared via one way ANOVA, with Tukey's post-hoc test, following log transformation of the data. The same analysis was conducted for extractable NH<sub>4</sub><sup>+</sup>, NO<sub>3</sub><sup>-</sup>, N and dissolved organic C at each sampling point. The soil solution NH<sub>4</sub><sup>+</sup>, NO<sub>3</sub><sup>-</sup>, N and dissolved organic C had a variable *n*, ranging from 1-3, due to successful sample collection being dependent on soil conditions being moist enough. Therefore, numerical trends in the soil solution NH<sub>4</sub><sup>+</sup>, NO<sub>3</sub><sup>-</sup>, total N and dissolved organic C data were reported rather than statistical differences. The soil solution mineral N data allowed determination of whether N processing was occurring at similar rates in the chambers and the



duplicated plots used for the soil coring, as the Rhizon samplers were located directly inside the chambers. Pasture biomass and foliar N content were also analysed by ANOVA, as above, on each of the three biomass cutting dates. All statistical analyses were conducted in Minitab 17.0 (Minitab Inc., State College, PA, USA).


## **7.3 Results**

### *7.3.1 Urine characteristics*

The composite sheep urine sample contained 14.5 g N l<sup>-1</sup> and 28.7 g C l<sup>-1</sup>, which resulted in an equivalent N application rate of 725 kg N ha<sup>-1</sup> in all applied urine patches. The urine contained 12.4 g urea-N l<sup>-1</sup>, 0.4 mg NO<sub>3</sub><sup>-</sup>-N l<sup>-1</sup>, 211 mg NH<sub>4</sub><sup>+</sup>-N l<sup>-1</sup>, 5.5 mg P l<sup>-1</sup>, 1.85 g Na l<sup>-1</sup>, 13.9 g K l<sup>-1</sup> and 284 mg Ca l<sup>-1</sup>. The urine had a pH and EC of 8.48 and 25.3 mS cm<sup>-1</sup>, respectively. Out of a total of 40 urine events between the 6 sheep, the mean and range of individual urine volumes were 104 (18-397) ml per urination event.

### *7.3.2 Weather data and soil moisture*


The hourly rainfall, soil temperature (0-10 cm) and water-filled pore space data throughout the study period can be seen in Fig. 7.1. The first DMPP application on the 29/06/15 for treatment SU + DMPP 4, was applied under warm and dry conditions. The second DMPP application on the 13/07/15, for treatment SU + DMPP 2, was applied during a period of rainfall. The final DMPP and urine applications (27/07/15) were also under periods of rainfall. The total rainfall over the course of the study was 137 mm, and the average soil temperature (0-10 cm) was 16 °C, ranging from 11-20 °C. The average WFPS over the course of the study was 27%, and ranged from 5-69%, with minimum values at the beginning of the experiment following an extended dry period, and maximum values occurring after rainfall events.



**Figure 7.1.** Rainfall, soil temperature and soil water-filled pore space during the course of the field trial. Hourly rainfall and soil (0-10 cm) temperature are shown in panel a) and water-filled pore space within the automated chambers are displayed in panel b). The date of measurements are displayed in dd/mm/yy format and legends apply to each respective panel only.

### 7.3.2 Nitrous oxide emissions


The fluxes of N<sub>2</sub>O from all treatments can be seen in Fig. 7.2. When considering the whole automated chamber data set (8420 flux measurements), 79% of the headspace concentration data had an  $R^2 > 0.95$ , 88% had an  $R^2 > 0.90$  and 97% of the data had an  $R^2 > 0.80$ , indicating the increase/decrease in N<sub>2</sub>O within the chamber headspace over time fitted well to the assumption of linearity. In the vast majority of cases three, rather than four, data points of increase in chamber headspace concentration over time resulted in a higher  $R^2$ . As can be seen in Fig. 7.2a, the main N<sub>2</sub>O peak from the SU treatment occurred within the first nine days following urine application, with a maxima of  $264 \pm 179 \mu\text{g N}_2\text{O-N m}^{-2} \text{ h}^{-1}$ . DMPP was effective in reducing this initial N<sub>2</sub>O peak when applied at the same time as the urine (Fig. 7.2b), however, a broader, shallower peak began around the 30/08/15, which was not evident in the SU treatment. In both treatments, SU + DMPP 2 and SU + DMPP 4, this initial peak was similar in size (Fig. 7.2c and d, respectively) with maximum peaks in the first nine days of  $192 \pm 151$  and  $231 \pm 89 \mu\text{g N}_2\text{O-N m}^{-2} \text{ h}^{-1}$ , respectively. Notably the variation in N<sub>2</sub>O emissions is higher in Fig. 7.2c and 2d compared to 7.2a and 7.2b, and a strong diurnal trend in N<sub>2</sub>O fluxes was observed.



**Figure 7.2.**  $\text{N}_2\text{O}$  fluxes following sheep urine and DMPP applications to a Eutric Cambisol, where panel a) contains fluxes from treatments SU and NU and panels b), c) and d) display fluxes from SU + DMPP, SU + DMPP 2 and SU + DMPP 4, respectively. In each panel the black line represents the mean flux ( $n = 3$ ) and the grey shaded area represents the upper and lower bounds of the SEM. In treatment NU, error bars denote SEM. Arrows indicate the timing of either sheep urine (SU) or DMPP applications and the dates along the x-axis are in dd/mm/yy format.

### 7.3.3 Cumulative N<sub>2</sub>O emissions and emission factors

Cumulative N<sub>2</sub>O emissions were calculated from the point of urine application onwards, and can be seen in Fig. 7.3. Mean cumulative emissions ranged from 0.4-1.0 kg N<sub>2</sub>O-N ha<sup>-1</sup>. Due to the high variability of the SU + DMPP 2 and SU + DMPP 4, homogeneity of variance was not achieved across the treatments, therefore, Welch's test was used to assess differences. The results of the Welch's test were positive ( $p < 0.05$ ) both with and without the removed 'high emitting' replicates; the statistical groupings from the Games-Howell test can be seen in Fig. 7.4. Here it can be seen that the SU and SU + DMPP treatments emitted significantly more N<sub>2</sub>O in comparison to NU. Due to the large variability in the SU + DMPP 2 and SU + DMPP 4 treatments (and the low value for  $n$  when one replicate is removed), no differences were observed between these treatments and NU. In contrast to our hypotheses, DMPP did not reduce the cumulative N<sub>2</sub>O emissions from sheep urine patches when applied at the same time as the urine, or when applied 2 and 4 weeks before urine application. The N<sub>2</sub>O emission factors were  $0.41 \pm 0.05$ ,  $0.43 \pm 0.04$ ,  $1.22 \pm 0.56$  and  $1.19 \pm 0.65\%$  of the applied N for SU, SU + DMPP, SU + DMPP 2 and SU + DMPP 4 treatments, respectively (without removal of outliers). No significant differences ( $p > 0.05$ ) were found between the treatments.




**Figure 7.3.** Cumulative nitrous oxide emissions, over 9 weeks post urine and DMPP applications to a Eutric Cambisol. Light grey bars represent means where  $n = 3$ , dark grey bars represent means where  $n = 2$ , with the high emitting replicates removed. Error bars denote SEM and different lower case letters represent significant differences (Welch's test, with Games-Howell pairwise comparisons) between treatments, where the same groupings were found both with and without outliers removed.

#### 7.3.4 Diurnal nature of N<sub>2</sub>O emissions

In order to investigate the diurnal nature of N<sub>2</sub>O emissions, data were selected over the period from 08/08/2015 to 14/08/15. The 'high emitting' replicate from SU + DMPP 2 treatment was chosen for investigation, as it is unlikely that other factors e.g. available N and C, were limiting N<sub>2</sub>O production during this period. The diurnal nature of N<sub>2</sub>O emissions can be seen in Fig. 7.4, where N<sub>2</sub>O emissions are shown to be out of phase with soil temperature, which may indicate a lag phase between soil temperature and the activity of N<sub>2</sub>O producing microorganisms. The maximum N<sub>2</sub>O emissions over this diurnal cycle generally occurred

between ca. 00:30 and 06:30 (hh:mm), and the minimum emissions in the diurnal cycle occurred between ca. 15:30 and 21:30 (hh:mm). The maximum soil temperature over the period of observation shown in Fig. 7.3 was 26 °C, and the minimum soil temperature was 12 °C. The highest soil temperatures generally occurred at around 13:30 (hh:mm), however the minimum temperature occurred anywhere between 00:00 and 06:00 (hh:mm).



**Figure 7.4.** Diurnal nature of N<sub>2</sub>O emissions and soil temperature over 6 days. The solid line represents the measured N<sub>2</sub>O flux from a single high emitting replicate from SU + DMPP 2 treatment, and the dashed line is soil temperature as measured by the Acclima SDI-12 combined temperature and moisture sensors in the chamber. The date/time format along the x-axis is expressed in dd/mm/yy hh:mm.

### 7.3.5 Mineral N dynamics


The extractable NH<sub>4</sub><sup>+</sup> and NO<sub>3</sub><sup>-</sup> from soil cores in replicated plots can be seen in Figure 7.5a and b, respectively, and the soil solution NH<sub>4</sub><sup>+</sup> and NO<sub>3</sub><sup>-</sup> from Rhizon samples can be seen in Figure 7.5c and d, respectively. Prior to urine application, no significant differences ( $p >$

0.05) in extractable  $\text{NH}_4^+$  concentrations were observed between treatments. On the day of urine application (27/07/15), all extractable  $\text{NH}_4^+$  concentrations were greater than the control ( $p < 0.01$ ), with no further differences between treatments. On the second day following urine application, extractable  $\text{NH}_4^+$  was only higher ( $p < 0.05$ ) in the SU treatment in comparison to the control. Four days following urine application the extractable  $\text{NH}_4^+$  was greater in all DMPP containing treatments ( $p < 0.05$ ) compared to the control, but the SU treatment had returned to control levels ( $p > 0.05$ ), indicating DMPP applications were delaying nitrification. One week following urine application the SU and SU + DMPP 4 treatments had returned to control levels ( $p > 0.05$ ), however, the SU + DMPP and SU + DMPP 2 treatments had higher extractable  $\text{NH}_4^+$  concentrations in comparison to the control ( $p < 0.05$ ). This indicates that the effect of DMPP was shorter lived in the SU + DMPP 4, compared to SU + DMPP 2 and SU + DMPP treatments. No differences were observed in extractable  $\text{NH}_4^+$  between treatments 10 days after urine application. Two weeks following urine application SU + DMPP had a higher extractable  $\text{NH}_4^+$  content compared to the control ( $p < 0.05$ ), and after 3 weeks extractable  $\text{NH}_4^+$  was greater in SU + DMPP and the SU + DMPP 2 compared to the control ( $p < 0.05$ ). No further differences in extractable  $\text{NH}_4^+$  concentrations were observed beyond three weeks following urine application ( $p > 0.05$ ).

No differences were observed in extractable  $\text{NO}_3^-$  concentrations between treatments prior to, on the day of urine application or two and four days following urine application ( $p > 0.05$ ). After one week, however, all urine containing treatments had a higher extractable  $\text{NO}_3^-$  content in comparison to NU ( $p < 0.001$ ). Ten days following urine application the extractable  $\text{NO}_3^-$  concentrations were still greater than NU in all urine containing treatments. Here, the SU had a higher extractable  $\text{NO}_3^-$  concentration than the SU + DMPP treatment, but not the SU + DMPP 2 or SU + DMPP 4 treatments, indicating a delay in nitrification when DMPP was applied at the same time as the urine only. Two weeks after urine application, extractable  $\text{NO}_3^-$



was greater in all urine containing treatments compared to NU, but no other differences were detected beyond this date ( $p > 0.05$ ).




**Figure 7.5.** Mineral nitrogen dynamics during the course of the field trial. Extractable ammonium-N can be seen in panel a) and extractable nitrate-N in panel b), which were measured on extracted soils from replicated plots (outside chambers). The soil solution ammonium-N is shown in panel c) and soil solution nitrate-N in panel d), which were measured from Rhizon samples located within the chambers. The figure legend applies to all panels, symbols represent means ( $n = 3$ ), and error bars denote SEM. Arrows in panel d) represent timings of sheep urine (SU) and/or DMPP applications to the respective treatments.

Asterisks above symbols represent level of significant differences (ANOVA) and was conducted for data in panels a) and b) only. The date format along the x-axis is dd/mm/yy.

Rhizon soil solution samples were only obtained when soil conditions were moist enough, therefore n varied one each sample date, depending on whether sufficient sample was obtained for analysis. Soil solution  $\text{NH}_4^+$  concentrations in the treatments containing sheep urine appeared to return to control values at a similar time to the soil extracts. The soil solution  $\text{NO}_3^-$  concentrations increased steadily during the 9 days following urine application. At the point of 9 days following urine application, the soil solution  $\text{NO}_3^-$  followed the numerical trend  $\text{C} < \text{SU} + \text{DMPP} < \text{SU} + \text{DMPP 2} < \text{SU} + \text{DMPP 4} < \text{SU}$ , which indicates DMPP delayed nitrification rates compared to sheep urine alone, but was less effective with the longer the time since DMPP application. Notably, the soil solution  $\text{NO}_3^-$  concentrations were numerically higher than NU for a longer period in comparison to the extractable  $\text{NO}_3^-$  data, indicating a difference in N processing rates between duplicated plots and the chambers.

#### 7.3.5 Dissolved organic C and N dynamics

The total extractable N and dissolved organic C during the course of the field trial can be seen in Fig. 7.6a and b, respectively; the total N and dissolved organic C measured in the soil solution samples can be seen in Fig. 7.6c and d, respectively. Differences between total extractable N were only found at one sampling point following urine application to the plots, this was at two weeks following urine application where the SU + DMPP extractable N was greater than the control ( $p < 0.05$ ). Similarly, not many differences in extractable dissolved organic C were found between the treatments at the different sampling points. Here, the only differences found were two days following urine application, where SU and SU + DMPP 2 had higher levels of extractable dissolved organic C compared to the control ( $p < 0.05$ ).



**Figure 7.6.** Total extractable and soil solution dissolved N and C dynamics. Total extractable dissolved N is displayed in panel a) and total extractable dissolved organic C is displayed in panel b), which were measured on extracted soils from replicated plots (outside chambers). Soil solution dissolved N is displayed in panel c) and soil solution dissolved organic C is displayed in panel d), which were measured using Rhizon samplers located within the chambers. The figure legend applies to all panels, symbols represent means ( $n = 3$ ), and error bars denote SEM. Arrows in panel d) represent timings of sheep urine (SU) and/or DMPP applications to the respective treatments. Asterisks above symbols represent level of significant differences (ANOVA) and was conducted for data in panels a) and b) only. The date along the x-axis is expressed in dd/mm/yy.

The soil solution N increased following urine application, and remained numerically higher than the control until the end of the study, which reflected the remaining  $\text{NO}_3^-$  in the soil solution of the chamber plots. In the two chambers which emitted the highest amounts of

N<sub>2</sub>O, it was found that the soil solution dissolved organic C concentration was ca. 2 and 4 times higher than the other samples collected across the plots, prior to urine application. The soil solution dissolved organic C concentration increased following urine application and treatments appeared to decline at similar rates. No observable differences in soil solution dissolved organic C were found between treatments after ca. 3 weeks following urine deposition.

#### *7.3.6 Pasture biomass and foliar N content*

The foliar N content measured at block level ( $n = 3$ ), was  $3.5 \pm 0.5\%$  prior to urine application. No significant differences ( $p > 0.05$ ) in foliar N content were observed between treatments 3 weeks following urine application, where the mean foliar N content across all treatments was  $3.4 \pm 0.2\%$ . The urine containing treatments all had a significantly greater pasture foliar N content in comparison to the control ( $p < 0.01$ ) at 6 and 9 weeks following urine application, where the mean foliar N content of all the urine containing treatments was  $4.3 \pm 0.1\%$  and  $4.6 \pm 0.2\%$ , compared to the control value of  $3.2 \pm 0.2\%$  and  $3.0 \pm 0.4\%$ , respectively. No significant differences ( $p > 0.05$ ) were found for pasture biomass in the plots at 3, 6 or 9 weeks following urine application, where the mean pasture biomass across all samples were  $0.47 \pm 0.2$ ,  $0.51 \pm 0.03$  and  $0.29 \pm 0.04$  t DM ha<sup>-1</sup>, respectively.

## **7.4 Discussion**

### *7.4.1 Effect of DMPP on sheep urine patch N<sub>2</sub>O emissions and mineral N dynamics*

In this study, a liquid DMPP application (1 kg ha<sup>-1</sup>) did not reduce cumulative N<sub>2</sub>O emissions from sheep urine patches under UK summer conditions; this was true when DMPP was applied 4 weeks before, 2 weeks before and even when applied on the same day as the sheep urine. Whilst no significant effect of DMPP was found for cumulative N<sub>2</sub>O emissions,

there was an observable effect of the nitrification inhibitor based on temporal trends in the N<sub>2</sub>O fluxes and mineral N data. When considering the SU + DMPP treatment, the initial N<sub>2</sub>O peak following urine application was much reduced in comparison to the SU only treatment, however, a broader, shallower peak towards the end of the measurement period was observed. In addition, the extractable NH<sub>4</sub><sup>+</sup> concentrations were higher in the SU + DMPP and SU + DMPP 2 treatments up to three weeks following urine application. This suggests that nitrification rates were delayed, with the effect being greater the shorter the time since DMPP application. Whilst some effect of DMPP was observable, this tended to delay, rather than reduce overall N<sub>2</sub>O emissions in this study. The N application in the sheep urine patches was relatively high at 725 kg N ha<sup>-1</sup>, yet Di et al. (2012) found DMPP reduced N<sub>2</sub>O emissions by 66% from cattle urine N applications as high as 1000 kg N ha<sup>-1</sup>, although a higher DMPP application of 5 kg ha<sup>-1</sup> was used in this study.

Reasons for not observing an effect of DMPP on cumulative N<sub>2</sub>O emissions could be because of the high soil temperatures, which may have resulted in rapid DMPP degradation by microbial activity. Menéndez et al. (2012) and Vasquez et al. (unpublished), however, found no effect of increasing temperature between 10 and 20 °C on DMPP degradation, which represents the typical range of soil temperatures observed in this field trial. The soil water content has also been identified as an important parameter influencing the persistence of this molecule in soil, where a lower water content (and hence greater oxygen content) may increase oxidation of the molecule (Chen et al., 2010; Menéndez et al., 2012). In this study the maximum observed water-filled pore space was 69%, but more often it remained between 20 and 40%, which renders depletion of DMPP via oxidation a possible degradation pathway. Further depletion of DMPP from the soil solution could have occurred due to immobilisation within non-target microbial biomass or canopy interception of the applied inhibitor.

Another possible reason for the lack of effect of DMPP on cumulative N<sub>2</sub>O emissions could be inefficient mixing with the soil solution and urine-N. Given that the DMPP was applied under rainfall conditions in the treatments SU + DMPP and SU + DMPP 2, and that there was an observable effect of DMPP on nitrification rates and the temporal nature of N<sub>2</sub>O fluxes in SU + DMPP treatment, it would suggest that the inhibitor at least partially mixed with the urine derived N. The DMPP applied 4 weeks before urine application, however, may have experienced more degradation or pasture canopy interception during application, as it was not applied during a rainfall event. Further research based on different DMPP application methods is required, e.g. investigating whether DMPP would have been more effective if mixed with the urine prior to application to soil. Comparatively more research has been conducted on different application methods of DCD in comparison to DMPP (e.g. Ledgard et al., 2008; Minet et al., 2013; Wakelin et al., 2013; Luo et al., 2015). This may be due to the original commercial development of DMPP as a fertiliser coating (Pasda et al., 2001; Zerulla et al., 2001), rather than for its use on urine-influenced soils.

#### *7.4.3 Spatial variability in N<sub>2</sub>O fluxes*

N<sub>2</sub>O emissions from agricultural soils are notoriously variable, both in time and space (Hénault et al., 2012), and this study proved to be no exception. Notably the SU and the SU + DMPP treatments did not have as high a variability in N<sub>2</sub>O emissions as the SU + DMPP 2 and SU + DMPP 4 treatments. Due to this we can conclude with more confidence that DMPP had no effect on cumulative N<sub>2</sub>O emissions when applied at the same time as the urine under these experimental conditions. Bearing this in mind, it is unlikely that the DMPP would have had an appreciable effect on emissions when DMPP was applied weeks before the urine, even if the data had not suffered from the large spatial variability.

Whilst a high temporal variability was observed, the automated chamber system captured this variability, thereby reducing bias in emission calculations due to sampling time and frequency, which can be an artefact of manually sampled static chamber based campaigns e.g. Yamulki et al. (2001), Scheer et al. (2014). The N<sub>2</sub>O emissions in this study were also, however, highly spatially variable. Here, one replicate in each of the treatments SU + DMPP 2 and SU + DMPP 4 demonstrated a marked difference to their respective replicates. Future studies should consider maximising the number of replicates in order to capture spatial variability in N<sub>2</sub>O emissions, when using an automated chamber based system. This will potentially result in a having to select fewer treatments for investigation, however, would be appropriate where emission factors are to be measured for inventory purposes.

Prior to urine application, the soil solution samples from inside the ‘high emitting’ chambers had 2 (242 mg C l<sup>-1</sup>) and 4 (489 mg C l<sup>-1</sup>) times the amount of dissolved organic C compared to the mean of the remaining soil solution samples (125 ± 10 mg C l<sup>-1</sup>; mean ± SEM; *n* = 10). The spatial heterogeneity in dissolved organic C may have been caused by several reasons, including root exudation in the ‘rhizosphere’, litter decomposition and root death in the ‘detritosphere’ and labile and recalcitrant organic material passed through the hind gut of soil organisms or faeces of larger organisms in ‘biopores’, which are recognised as hotspots of microbial activity (Kuzyakov and Blagodatskaya, 2015). Urine patches themselves are also considered hotspots of microbial activity. As the plots in this study were fenced off several months prior to the simulated urine applications, the possibility of the high dissolved organic C arising from a previous urine deposition is unlikely. Whatever the origin of the labile C in these plots, we believe the overlapping of two unique hotspots occurred. The presence of greater amounts of labile C in the two high emitting chambers prior to urine application may have resulted in an increased abundance and activity of soil microorganisms (e.g. nitrifiers and denitrifiers), compared to the soil enclosed by the other chambers. The addition of more labile

C and N in the form of urine, may then have led to rapid N turnover and increased N<sub>2</sub>O emissions from these plots. Alternatively, DMPP could have provided a C source in itself, where uneven applications within the chambers may have resulted in a large variability in emissions.

#### 7.4.2 N<sub>2</sub>O emission factors

No differences were found for N<sub>2</sub>O emission factors between treatments. The mean N<sub>2</sub>O emission factor across all treatments ( $n = 12$ ) was  $0.81 \pm 0.22\%$  of the applied N. The lowest emission factor recorded was 0.30% of the applied N which occurred in the SU + DMPP 2 treatment, and the highest emission factor was 2.49% of the applied N which occurred in the SU + DMPP 4 treatment. In general, emission factors were lower than the current default IPCC emission factor of 1% of the applied N for sheep excreta (IPCC, 2007), which is consistent with reported emission factors being lower under spring-summer conditions in comparison to autumn-winter (Allen et al., 1996). Nevertheless, the overlapping of nutrient hotspots more than doubled the default IPCC emission factor for sheep urine, where emission factors of 2.24 and 2.49% of the applied N occurred. The measurement period of this study was shorter than the recommended emission factor measurement period of 1 year and it did not include a measurement of emissions from the dung fraction of excreta, therefore, care must be taken when considering these emission factor values.

#### 7.4.4 Diurnal nature of N<sub>2</sub>O fluxes

The existence of diurnal trends in N<sub>2</sub>O flux data has been identified in several studies (e.g. Thomson et al., 1997; Yamulki et al., 2000; Yamulki et al., 2001; Hyde et al. 2005a; Hyde et al., 2005b; Hatch et al. 2005; Alves et al., 2012; Scheer et al., 2014). Scheer et al. (2014) found the diurnal nature of N<sub>2</sub>O fluxes appeared to be greatest when the mineral N and moisture



were non-limiting. This appeared to be true in our study, and the diurnal trend was also strongest in the chambers where urine was applied to a dissolved organic C hotspot. Our study shows that maximum N<sub>2</sub>O fluxes occurred at night time and early morning, where the diurnal nature of fluxes was out of phase with soil temperature. A similar phenomenon was observed in other temperate grassland systems where cattle urine (Yamulki et al., 2000; Yamulki et al., 2001) and ammonium-nitrate fertilizer (Hyde et al., 2005) were the N sources. In the studies by Yamulki et al. (2000) and Yamulki et al. (2001), maximum N<sub>2</sub>O fluxes occurred between the late afternoon and early morning and Hyde et al. (2005a) observed minimum N<sub>2</sub>O fluxes at mid-day and maximum fluxes between 20:00 and 22:00 (hh:mm). However, other studies have also shown soil N<sub>2</sub>O flux and soil temperature to be closer in phase with each other, e.g. Hatch et al. (2005) showed a lag time of only 30 minutes between maximum temperature and N<sub>2</sub>O fluxes, although this study was conducted under controlled conditions, on repacked soil cores, in the absence of plants and with slurry as the N source. In the study by Scheer et al. (2014), maximum emissions occurred between 12:00 and 15:00 (hh:mm) and minimum emissions occurred between 04:00 and 07:00 (hh:mm); in this study, the diurnal fluxes were assessed in a treatment containing fertilizer application, after the incorporation of broccoli residues in an intensive broccoli production site in a sub-tropical climate.

The observed variation could be due to the fact that N<sub>2</sub>O emissions are dependent on several factors (not just soil temperature), which may interact in different ways to produce different timings of minimum and maximum flux. In addition, the solubility and microbial consumption of O<sub>2</sub> increases alongside temperature, which may lead to reductions in N<sub>2</sub>O due to the occurrence of complete denitrification (Yamulki et al., 2000). The pattern of N<sub>2</sub>O production and consumption across soil depth may also vary with temperature, where maximum and minimum temperatures would occur later at soil depth. Another possible reason is that greater compaction could result in slower gas diffusivity (Ball et al., 1999; Sitaula et al.,

2000) from the source of N<sub>2</sub>O to the chamber headspace i.e. a shorter lag between maximum temperature and maximum N<sub>2</sub>O flux was found in the sieved and repacked soils of Hatch et al. (2005), as well as soils which had been subjected to rotary hoeing during incorporation of vegetable residues in Scheer et al. (2014). This area of study requires further research as it has implications for recommendations of the best time to sample N<sub>2</sub>O emissions over a 24 h period to reflect the average daily fluxes. It also supports the use of methods with a high temporal resolution for monitoring N<sub>2</sub>O emissions, where possible.

#### *7.4.5 Destructive vs. non-destructive soil sampling for measuring soil mineral N pools*

In this study differences were found between the mineral N dynamics as measured by taking soil cores from duplicate urine patches (destructive) and as measured in the soil solution of the chambers (non-destructive). The soil solution NO<sub>3</sub><sup>-</sup> concentrations in the urine treated chambers remained higher than the control for a longer period than the extractable NO<sub>3</sub><sup>-</sup>, therefore, further N<sub>2</sub>O emissions could have occurred beyond the duration of this study. The trends in soil solution and soil extractable NO<sub>3</sub><sup>-</sup> concentrations are more indicative of whether N processing rates were occurring at similar rates in the chambers, than comparing soil solution NH<sub>4</sub><sup>+</sup> and soil extractable NH<sub>4</sub><sup>+</sup>. Provided N processing rates are occurring equally, the soil solution NO<sub>3</sub><sup>-</sup> should follow similar temporal trends to the extractable NO<sub>3</sub><sup>-</sup>, due to limited sorption occurring for NO<sub>3</sub><sup>-</sup>. Conversely, the soil solution NH<sub>4</sub><sup>+</sup> data represents the pool of NH<sub>4</sub><sup>+</sup> that was not bound to exchange sites, therefore, comparisons between the processing rates of NH<sub>4</sub><sup>+</sup> in the soil solution and the soil extracts cannot be made, as they represent different pools. Nevertheless, soil solution NH<sub>4</sub><sup>+</sup> concentrations in the treatments containing sheep urine appeared to return to control values at a similar time to the soil extracts.

The differences in N processing found between the duplicated urine patches may reflect artefacts of coring the duplicated urine patches. This may have influenced the aeration and

water infiltration dynamics in these areas. Conversely, the chambers may have caused more shading and/or interception of rainfall in comparison to the duplicated plots. The difference in N processing rates highlights the need for the development and use of non-destructive sampling techniques for use inside chambers. Rhizon samplers were useful in this study for determining differences, which would not have been observed from only taking soil cores from duplicated urine patches. Limitations of the Rhizons were that successful sample collection depended on sufficient soil moisture, which sometimes resulted in a variable number of replicates on different sample dates. Deploying multiple Rhizons within a urine patch could potentially overcome the issue of the occasional small sample volume. Nevertheless, Rhizon samplers would be ineffective under extended dry periods unless the soil water content is modified artificially, which is not recommended as modification of the soil water content may also modify N processing rates.

## **7.5 Conclusions**

A recent review by Gilsanz et al. (2016), highlights the paucity of studies regarding DMPP efficacy in reducing N<sub>2</sub>O emissions, in comparison to the well-researched inhibitor DCD. This is especially true of urine-influenced soils, where this study contributes new information for DMPP efficacy in sheep urine patches, under UK summer conditions. In this study, DMPP influenced the dynamics of N processing rates, but had no effect on overall cumulative N<sub>2</sub>O emissions or emission factors. Further research should be conducted to determine whether DMPP is effective in livestock grazed pastures at a range of urine-N applications, under different seasonal conditions and with a range of nitrification inhibitor delivery methods to the soil.

## **7.6 Acknowledgements**

KAM would like to thank the School of Environment, Natural Resources and Geography, Bangor University, for sponsoring this study. In addition, thanks go to Neil Donovan at Rothamsted Research for analysing gas samples, Sarah Chesworth, Terry Cox, Rory Shaw, Jonathan Roberts, Gordon Turner, Mark Hughes, Llinos Hughes and Jairo Cardoso Abreu at Bangor University for all the assistance with setting up the field trial and sample processing.

## 7.7 References

- Akiyama, H., Yan, X., Yagi, K. 2010. Evaluation of effectiveness of enhanced-efficiency fertilizers as mitigation options for N<sub>2</sub>O and NO emissions from agricultural soils: meta-analysis. *Global Change Biology*, 16, 1837-1846.
- Allen, A.G., Jarvis, S.C., Headon, D.M. 1996. Nitrous oxide emissions from soils due to inputs of nitrogen from excreta return by livestock on grazed grassland in the UK. *Soil Biology and Biochemistry* 28, 597-607.
- Alves, B.J.R., Smith, K.A., Flores, R.A., Cardoso, A.S., Oliviera, W.R.D., Jantalia, C.P., Urquiaga, S., Boddey, R.M. 2012. Selection of the most suitable sampling time for static chambers for the estimation of daily mean N<sub>2</sub>O flux from soils. *Soil Biology and Biochemistry* 46, 129-135.
- Ball, D.F. 1964. Loss-on-ignition as an estimate of organic matter and organic carbon in non-calcareous soils. *Journal of Soil Science* 15, 84-92.
- Ball, B.C., Scott, A., Parker, J.P. 1999. Field N<sub>2</sub>O, CO<sub>2</sub> and CH<sub>4</sub> fluxes in relation to tillage, compaction and soil quality in Scotland. *Soil and Tillage Research* 53, 29-39.
- Barneze, A.S., Minet, E.P., Cerri, C.C., Misselbrook, T. 2015. The effect of nitrification inhibitors on nitrous oxide emissions from cattle urine depositions to grassland under summer conditions in the UK. *Chemosphere* 119, 122-129.
- Barth, G., von Tucher, S., Schmidhalter, U. 2001. Influence of soil parameters on the effect of 3,4-dimethylpyrazole-phosphate as a nitrification inhibitor. *Biology and Fertility of Soils* 34, 98-102.
- Bell, M.J., Rees, R.M., Cloy, J.M., Topp, C.F.E., Bagnall, A., Chadwick, D.R. 2015. Nitrous oxide emissions from cattle excreta applied to a Scottish grassland: Effects of soil and climatic conditions and a nitrification inhibitor. *Science of the Total Environment* 508, 343-353.

- Benckiser, G., Christ, E., Herbert, T., Weiske, A., Blome, J., Hardt, M. 2013. The nitrification inhibitor 3,4-dimethylpyrazole-phosphate (DMPP) – quantification and effects on soil metabolism. *Plant and Soil* 371, 257-266.
- Carter, M.S. 2007. Contribution of nitrification and denitrification to N<sub>2</sub>O emissions from urine patches. *Soil Biology and Biochemistry* 39, 2091-2102.
- Chaves B., Opoku, A., De Neve, S., Boeckx, P., Van Cleemput, O., Hofman, G. 2006. Influence of DCD and DMPP on soil N dynamics after incorporation of vegetable crop residues. *Biology and Fertility of Soils* 43, 62-68.
- Chen, D., Suter, H.C., Islam, A., Edis, R. 2010. Influence of nitrification inhibitors on nitrification and nitrous oxide (N<sub>2</sub>O) emissions from a clay loam soil fertilized with urea. *Soil Biology and Biochemistry*, 42, 660-664.
- Davidson, E.A. 2009. The contribution of manure and fertilizer nitrogen to atmospheric nitrous oxide since 1860. *Nature Geoscience* 2, 659-662.
- De Klein, C.A.M., Cameron, K.C., Di, H.J., Rys, G., Monaghan, R.M., Sherlock, R.R. 2011. Repeated annual use of the nitrification inhibitor dicyandiamide (DCD) does not alter its effectiveness in reducing N<sub>2</sub>O emissions from cow urine. *Animal Feed Science and Technology* 166, 480-491.
- De Klein, C.A.M., van Logtestijn, R.S.P. 1994. Denitrification and N<sub>2</sub>O emission from urine-affected grassland soil. *Plant and Soil* 163, 235-242.
- Di, H., Cameron, K.C. 2002. The use of a nitrification inhibitor, dicyandiamide (DCD), to decrease nitrate leaching and nitrous oxide emission in a simulated grazed and irrigated grassland. *Soil Use and Management* 4, 395-403.
- Di, H.J., Cameron, K.C. 2003. Mitigation of nitrous oxide emissions in a spray-irrigated grazed grassland by treating the soil with dicyandiamide, a nitrification inhibitor. *Soil Use and Management* 19, 284-290.

- Di, H., Cameron, K.C. 2007. Effects of the nitrification inhibitor dicyandiamide on soil mineral N, pasture yield, nutrient uptake and pasture quality in a grazed pasture system. *Soil Use and Management* 23, 111-120.
- Di, H.J., Cameron, K.C. 2011. Inhibition of ammonium oxidation by a liquid formulation of 3,4-dimethylpyrazole phosphate (DMPP) compared with dicyandiamide (DCD) in six new Zealand grazed grassland soils. *Journal of Soils and Sediments* 11, 1032–1039.
- Di, H.J., Cameron, K.C. 2012. How does the application of different nitrification inhibitors affect nitrous oxide emissions and nitrate leaching from cow urine in grazed pastures? *Soil Use and Management* 28, 54-61.
- Doak, B.W., 1952. Some chemical changes in the nitrogenous constituents of urine when voided on pasture. *The Journal of Agricultural Science* 42, 162-171.
- Fiencke, C., Bock, E. 2006. Immunocytochemical localization of membrane-bound ammonia monooxygenase in cells of ammonia oxidizing bacteria. *Archives of Microbiology* 185, 99-106.
- Gill, M., Smith, P., Wilkinson, J.M. 2010. Mitigating climate change: the role of domestic livestock. *Animal* 4, 323-333.
- Gilsanz, C., Báez, D., Misselbrook, T.H., Dhanoa, M.S., Cárdenas, L.M. 2016. Development of emission factors and efficiency of two nitrification inhibitors, DCD and DMPP. *Agriculture, Ecosystems and Environment*, 216, 1-8.
- Hatch, D., Trindade, H., Cardenas, L., Carneiro, J., Hawkins, J., Scholefield, D., Chadwick, D. 2005. Laboratory study of the effects of two nitrification inhibitors on greenhouse gas emissions from a slurry-treated arable soil: impact of diurnal temperature cycle. *Biology and Fertility of Soils* 41, 225-232.

- Hénault, C., Gossel, A., Mary, B., Roussel, M., Léonard, J. 2012. Nitrous oxide emission by agricultural soils: a review of spatial and temporal variability for mitigation. *Pedosphere*, 22, 426-433.
- Hyde, B., Ryan, M., Hawkins, M., Connolly, J., Carton, O.T. 2005a. Nitrous Oxide Emissions, Teagasc Environment Research Centre.
- Hyde, B.P., Fanning, A.F., Ryan, M., Carton, O.T. 2005b. An examination of the diurnal variability in nitrous oxide emissions. In: O'Mara, F.P. (Eds.) 20<sup>th</sup> International Grassland Congress, Wageningen Academic Pub.
- Jones, D.L., Willett, V.B. 2006. Experimental evaluation methods to quantify dissolved organic nitrogen (DON) and dissolved organic carbon (DOC) in soil. *Soil Biology and Biochemistry* 5, 991-999.
- Kuzyakov, Y., Blagodatskaya, E. 2015. Microbial hotspots and hot moments in soil: Concept & review. *Soil Biology and Biochemistry* 83, 184-199.
- Ledgard, S.F., Menneer, J.C., Dexter, M.M., Kear, M.J., Lindsey, S., Peters, J.S., Pacheco, D. 2008. A novel concept to reduce nitrogen losses from grazed pastures by administering soil nitrogen process inhibitors to ruminant animals: a study with sheep. *Agriculture, Ecosystems and Environment* 125, 148-158.
- Linzmeier, W., Gutser, R., Schmidhalter, U. 2001. Nitrous oxide emission from soil and from a nitrogen-15-labelled fertilizer with the new nitrification inhibitor 3,4-dimethylpyrazole phosphate (DMPP). *Biology and Fertility of Soils* 34, 103-108.
- Liu, C., Wang, K., Zheng, X. 2013. Effects of nitrification inhibitors (DCD and DMPP) on nitrous oxide emission, crop yield and nitrogen uptake in a wheat-maize cropping system. *Biogeosciences Discussions* 10, 711-737.



- Luo, J., Ledgard, S., Wise, B., Welten, B., Lindsey, S., Judge, A., Sprosen, M. 2015. Effect of dicyandiamide (DCD) delivery method, application rate and season on pasture urine patch nitrous oxide emissions. *Biology and Fertility of Soils* 51, 453-464.
- Marsden, K.A., Scowen, M., Hill, P.W., Jones, D.L., Chadwick, D.R. 2015. Plant acquisition and metabolism of the synthetic nitrification inhibitor dicyandiamide and naturally-occurring guanidine from agricultural soils. *Plant and Soil* 395, 201-214.
- Menéndez, S., Barrena, I., Setien, I., González-Murua, C., Estavillo, J.M. 2012. Efficiency of nitrification inhibitor DMPP to reduce nitrous oxide emissions under different temperature and moisture conditions. *Soil Biology and Biochemistry* 53, 82-89.
- Merino, P., Menéndez, S., Pinto, M., González-Murua, C., Estavillo, J.M. 2005. 3,4-Dimethylpyrazole phosphate reduces nitrous oxide emissions from grassland after slurry application. *Soil Use and Management* 21, 53-57.
- Minet, E. P., O'Carroll, C., Rooney, D., Breslin, C., McCarthy, C.P., Gallagher, L., Richards, K.G. 2013. Slow delivery of a nitrification inhibitor (dicyandiamide) to soil using a biodegradable hydrogel of chitosan. *Chemosphere* 93, 2854-2858.
- Miranda, K.M., Epsey, M.G., Wink, D.A., 2001. A rapid, simple, spectrophotometric method for simultaneous detection of nitrate and nitrite. *Nitric Oxide* 5, 62-71.
- Misselbrook, T.H., Cardenas, L.M., Camp, V., Thorman, R.E., Williams, J.R., Rollett, A.J., Chambers, B.J. 2014. An assessment of nitrification inhibitors to reduce nitrous oxide emissions from UK agriculture. *Environmental Research Letters* 9, 11506.
- Mulvaney, R.L., 1996. Nitrogen – inorganic forms. In: Sparks, D.L. (Eds.), *Methods of Soil Analysis. Part 3*. Madison, WI, USA: Soil Science Society of America Inc., pp. 1123-1184.
- Murphy, J., Riley, J.P., 1962. A modified single solution method for determination of phosphate in natural waters. *Analytica Chimica Acta* 27, 31-36.

- Oenema, O., Wrage, N., Velthof, G.L., van Groenigen, J.W., Dolfing, J., Kuikman, P.J. 2005. Trends in global nitrous oxide emissions from animal production systems. *Nutrient Cycling in Agroecosystems* 72, 51-65.
- Orsonneau, J-L., Massoubre, C., Cabanes, M., Lustenberger, P., 1992. Simple and sensitive determination of urea in serum and urine. *Clinical Chemistry* 38, 619-623.
- Pasda, G., Hähndel, R., Zerulla, W. 2001. Effect of fertilizers with the new nitrification inhibitor DMPP (3,4-dimethylpyrazole phosphate) on yield and quality of agricultural and horticultural crops. *Biology and Fertility of Soils* 34, 85-97.
- Scheer, C., Rowlings, D.W., Firrel, M., Deuter, P., Morris, S., Grace, P.R. 2014. Impact of nitrification inhibitor (DMPP) on soil nitrous oxide emissions from an intensive broccoli production system in sub-tropical Australia. *Soil Biology and Biochemistry* 77, 243-251.
- Sitaula, B.K., Hansen, S., Sitaula, J.I.B., Bakken, L.R. 2000. Effects of soil compaction on N<sub>2</sub>O emission in agricultural soil. *Chemosphere – Global Change Science* 2, 367-371.
- Thomson, P.E., Parker, J.P., Arah, J.R., Clayton, H., Smith, K.A. 1997. Automated soil monolith-flux chamber system for the study of trace gas fluxes. *Soil Science Society of America Journal* 61, 1323-1330.
- Voroney, R.P., Brookes, P.C., Beyaert, R.P. 2008. Soil microbial biomass C, N, P and S. In: Carter, M.R and Gregorich, E.G., eds. *Soil sampling and methods of analysis*, 2<sup>nd</sup> edn. Boca Raton, FL, USA: CRC Press, pp. 637-651.
- Wakelin, S.A., Clough, T.J., Gerard, E.M., O’Callaghan, G.M. 2013. Impact of short-interval, repeat application of dicyandiamide on soil N transformation in urine patches. *Agriculture, Ecosystems and Environment* 167, 60-70.
- Webb, N., Broomfield, M., Buys, G., Cardenas, L., Murrels, T., Pang, Y., Passant, N., Thistlewaite, G., Watterson, J. 2014. UK Greenhouse Gas Inventory, 1990 to 2012.

Annual report for submission under the Framework Convention on Climate Change.  
Oxfordshire, UK: AEA. Report number: Ricardo-AEA/R/3407.

Weiske, A., Benckiser, G., Herbert, T., Ottow, J.G.C. 2001. Influence of the nitrification inhibitor 3,4-dimethylpyrazole phosphate (DMPP) in comparison to dicyandiamide (DCD) on nitrous oxide emissions, carbon dioxide fluxes and methane oxidation during three years of repeated application in field experiments. *Biology and Fertility of Soils* 34, 109-117.

Yamulki, S., Toyoda, S., Yoshida, N., Veldkamp, E., Grant, B., Bol, R. 2001. Diurnal fluxes and the isotopomer ratios of N<sub>2</sub>O in a temperate grassland following urine amendment. *Rapid Communications in Mass Spectrometry* 15, 1263-1269.

Yamulki, S., Wolf, I., Bol, R., Grant, B., Brumme, R., Veldkamp, E., Jarvis, S.C. 2000. Effects of dung and urine amendments on the isotopic content of N<sub>2</sub>O released from grasslands. *Rapid Communications in Mass Spectrometry* 14, 1356-1360.

Zerulla, W., Barth, T., Dressel, J., Erhardt, K., von Locquenghien, K.H., Pasda, G., Rädle, M., Wissemeier, A.H. 2001. 3,4-dimethylpyrazole phosphate (DMPP) – a new nitrification inhibitor for agriculture and horticulture. *Biology and Fertility of Soils* 34, 79-84.

Zhang, H-J., Wu, Z-J., Zhou, Q-X. 2004. Dicyandiamide sorption-desorption behaviour on soils and peat humus. *Pedosphere* 14, 395-399.

---

## Chapter 8

---

### Discussion and recommendations for future research

## 8.1 Introduction

Detailed discussion of the results from the experiments in this Thesis are described in each experimental Chapter. In this Chapter, the experimental work presented in Chapters 3-7 is summarised and discussed in relation to the common themes within the Thesis, the results are then discussed in relation to the initial objectives, and in terms of the wider implications of the results. An assessment is made of how future mitigation strategies and research should be prioritised in order to achieve enhanced sustainability and productivity in livestock grazed grasslands.

## 8.2 General discussion

This section discusses the results in terms of common themes within the Thesis. This Thesis consists of five experimental chapters where sheep urine (real or artificial) and/or nitrification inhibitors (DCD and DMPP) are applied to differing soils, with or without the presence of plants, in either laboratory or field settings.

Real sheep urine was applied to soils within Chapters 3, 5 and 7, and artificial urine was used in Chapter 4. The collected sheep urine samples used in Chapter 3 had a lower N concentration ( $3.6 \text{ g N l}^{-1}$ ) than that of the urine samples collected and used for Chapter 7 ( $14.5 \text{ g N l}^{-1}$ ). The urine patch N loading rate used in Chapter 3 equated to around  $200 \text{ kg N ha}^{-1}$ , whereas the N loading rate in the field trial of Chapter 7 was  $725 \text{ kg N ha}^{-1}$ . Artificial urine was made up at two concentrations ( $4 \text{ g N l}^{-1}$  and  $16 \text{ g N l}^{-1}$ ) and applied at two N loading rates (equivalent to  $200 \text{ kg N ha}^{-1}$  and  $800 \text{ kg N ha}^{-1}$ ) for use in Chapter 4, which were similar to the measured range of urine N concentrations found in Chapters 3 and 7. The observed range in urine N concentration supports other findings of a wide range in urinary N concentrations of grazing animals, despite similarities in the diet (*Lolium perenne* L. dominated sward) fed to the animals and the breed under investigation (Welsh Mountain ewes).

The nitrification inhibitor DCD was utilised in Chapters 5 and 6 and DMPP was utilised in Chapters 5 and 7. In Chapters 5 and 6 it was shown that DCD was bioavailable to both soil microorganisms and plants, respectively. While DMPP was also acquired by soil microorganisms, it appeared to be less available to soil microbes in comparison to DCD, which may explain the slower degradation rates of DMPP in comparison to DCD. Direct comparisons of the efficacy of DCD and DMPP were not made, although factors influencing their vertical mobilities were compared in Chapter 5. Both nitrification inhibitors behaved in different ways in contrasting soils, suggesting that differences in their efficacy in contrasting soils may also occur. The nitrification inhibitor DMPP was chosen for use in the final field trial (Chapter 7) as fewer studies had been conducted using this inhibitor to target urine patches in the soil compared to DCD, and due to the fact that the recent suspension of the sale and use of DCD may prompt the use of alternative nitrification inhibitors within pasture soils. DMPP was found to be ineffective at reducing emissions of  $\text{N}_2\text{O}$  from the Eutric Cambisol, under the conditions of field trial in Chapter 7. The results of the studies on nitrification inhibitors suggest that they are not a one-stop solution for reducing  $\text{N}_2\text{O}$  emissions from urine patches. Further work is required to optimise their efficacy in a cost-effective manner e.g. by using at high risk times, in high risk areas or by using technologies to apply directly to urine patches in the soil.

A common soil type (Eutric Cambisol; sandy clay loam) was utilised in Chapters 3, 4, 5 and 7; a sandy loam textured Eutric Cambisol was used in Chapter 5; and a Histosol was used in Chapters 5 and 6, to collectively provide some contrasting soil conditions. The Cambisols are one of the most common soil types based on land area (covering some 1.5 billion ha), and are widely used for food production (FAO, 2001), therefore, these soils represent an important area of study. Sheep urine was applied to a Eutric Cambisol in Chapters 3, 4 and 7. The maximum extractable  $\text{NO}_3^-$  following urine application to these soils were similar in the field trials (Chapter 4 and 7; ca.  $100 \text{ mg NO}_3^- \text{-N kg}^{-1} \text{ soil DW}$ ), but was higher in the laboratory

study (Chapter 3 ca. 270 mg NO<sub>3</sub><sup>-</sup>-N kg<sup>-1</sup> soil DW) which is likely to be an effect of the absence of plants in this study. The time taken for NO<sub>3</sub><sup>-</sup> to peak in these soils following urine application was similar, ranging from ca. 20-28 days. The extractable dissolved organic C following urine application to the Eutric Cambisol was similar in these three studies, ranging between 100-200 mg C kg<sup>-1</sup> soil DW. This suggests that C availability is not limited in these soils, however, this was measuring the extractable pool, where the organic C may be bound to soil particles and potentially unavailable to the soil microbes. The soil solution dissolved organic C is more indicative of what is potentially bioavailable following urine deposition, and this was measured in both Chapters 4 and 7. These results both show a sharp increase and decline of dissolved organic C immediately after urine application, which also peaked at similar values (ca. 2000 mg C l<sup>-1</sup>), despite the fact that artificial and real urine were used in the different studies. This suggests that the artificial urine may have adequately represented the composition of real urine. This peak of organic C was short-lived, however, lasting a period of just under a week in Chapter 4 and just over a week in Chapter 7 (this generally coincided with peak in N<sub>2</sub>O emissions following urine application). This suggests that the labile C applied to or created (e.g. priming or solubilisation of organic matter) within the urine patch is rapidly consumed, where conditions then return to similar values as measured prior to urine application. The N<sub>2</sub>O emissions varied quite widely from this soil type, with the highest emissions occurring under the laboratory study. This may have been expected as conditions of moisture were maintained at high values (70% WFPS) throughout the incubation and there were no plants competing for the mineral N within the soil. The major N<sub>2</sub>O emission period lasted for around 20 days after urine application in Chapters 3 and 4, but lasted for a longer period in Chapter 7 (ca. 60 days after urine application, mainly in the replicates which emitted high amounts of N<sub>2</sub>O). It is recognised that emissions could have been monitored for longer time periods e.g. to calculate emission factors 1 year of measurements are recommended. Levels of NO<sub>3</sub><sup>-</sup> had returned to

control values by the end of the study period in Chapter 3 and 7, however, there was still some  $\text{NO}_3^-$  available at the end of the study in Chapter 4, where  $\text{N}_2\text{O}$  emissions may have still occurred following rainfall events.

In Chapters 4, 6 and 7, pasture plants were present within the studies, however, no plants were present in Chapters 3 and 5. The inclusion of plants within studies of the urine patch is important due to the role plants play in regulating substrate availability in the rhizosphere. The absence of plants in Chapter 3 is likely to have resulted in an over-estimation of  $\text{N}_2\text{O}$  emissions, as it is likely that more N would have been available in the soil for  $\text{N}_2\text{O}$  producing processes. It may have also increased lateral diffusion of N, due to higher concentrations within the centre of the urine patch, creating a larger concentration gradient for diffusion. While Chapter 3 examined the urine patch diffusional area, a consideration of the pasture response area could also be important for accurately monitoring  $\text{N}_2\text{O}$  emissions from urine patches within chambers. The chamber walls may prevent the growth of adjacent plant roots into the urine patch, resulting in less of the available N being incorporated into plant tissues. The presence of plant roots may have altered the flow pathways of urine, DCD and DMPP in the soil columns of Chapter 5. Rhizosphere microorganisms may have played a greater role in the immobilisation of DCD and DMPP, if plants roots and their associated microbial communities were represented in the soil columns. This may have overestimated the vertical movement of the inhibitors in the studied soils.

Three of the experimental Chapters were laboratory studies (Chapters 3, 5 and 6), which are useful for understanding mechanistic processes; nevertheless, these studies are limited in what can be concluded from them, due to dissimilarities between laboratory and field conditions. In Chapter 3, 5 and 6 sieved soils were used; while this creates homogeneous conditions, it may misrepresent conditions in the field. Conditions which are modified include the breakdown of aggregate structures affecting soil porosity and bulk density, the removal of



stones and roots which can affect preferential flow pathways, homogenisation of the soil which removes any natural depth distribution of microorganisms and the potential breakage of fungal hyphae which may stimulate the release of nutrients into the soil. This has to be considered when interpreting the results of the laboratory studies in this Thesis, and extrapolations to field conditions need to be considered with care. The sieved soils were re-packed to a depth of 5 cm in Chapter 3 and to a soil column depth of 15 cm in Chapter 5. The use of shallow trays in Chapter 3 may have promoted lateral diffusion of solutes, while the use of small soil columns in Chapter 5 may have promoted vertical movement of urine and NI. Using intact pasture blocks or pasture soil cores, with a greater soil depth than 5 cm would be preferential for future laboratory studies, as many of the soil conditions would remain unchanged, yet environmental parameters such as moisture and temperature could be controlled.

Two of the experimental chapters were field based studies (Chapters 4 and 7). The data gathered from field trials are generally more realistic, although trends can be difficult to interpret due to confounding factors which can be present under natural conditions. A large variability was observed in many of the measured soil parameters and N<sub>2</sub>O emissions, which reflects the inherent spatial variability present within pasture soils. While the soil type and urinary N concentrations deposited were fairly similar in the two field trials, N<sub>2</sub>O emissions differed largely on the two separate occasions. This is likely to be due to the different climatic regimes observed during the trials, where one was comparatively warmer and drier (Chapter 4) and the other was cooler and wetter (Chapter 7).

### **8.3 What implications do variations in urine patch and environmental parameters have on the upscaling of N<sub>2</sub>O emissions from grazed grasslands?**

The first objective was to determine how a) sheep urine patch parameters and b) environmental parameters influence N<sub>2</sub>O emissions from grassland soils. In Chapter 4, the

effect of urine patch size and N concentration within urine patches was assessed. From this study it was concluded that increasing urine N concentration increased cumulative N<sub>2</sub>O emissions, but not the N<sub>2</sub>O emission factor. An effect of patch size was also observed, when holding N application rate constant, where four smaller patches emitted more N<sub>2</sub>O than a single large urine patch, at an equivalent N loading rate of 800 kg N ha<sup>-1</sup>. There is a paucity of data regarding urination frequency, volume and N concentration under differing diets, management strategies and environmental conditions. The importance of considering the frequency and concentration distribution of urine events for modelling N losses from grazed pastures is becoming more apparent (Li et al., 2012). Advances in sensor-based technology for monitoring urine events *in situ* will be most useful in the future (e.g. Betteridge et al., 2010, 2013), to feed into models, and to test management and mitigation measures under more natural conditions (i.e. compared to housing animals and subsequent urine collection, which may influence the urine composition).

In Chapter 3, changes in N<sub>2</sub>O emissions at the urine patch-soil interface were assessed, under controlled conditions. The most important driving variables for N<sub>2</sub>O emissions under controlled conditions were water-filled pore space (WFPS), redox potential and total extractable N content. In Chapters 4 and 7, peaks in N<sub>2</sub>O emissions also appeared alongside rainfall events. Under the warm and dry conditions in the field trial of Chapter 4, the N<sub>2</sub>O emissions from sheep urine patches over 10 weeks were ten times lower than the IPCC default emission factor for N<sub>2</sub>O from sheep excreta. In the late summer urine application there was more rainfall and emission factors were generally closer to that of the default (0.80% of applied N), however, emissions were still not measured for a whole year, and further emissions could have occurred beyond the period measured in this study. Scaling up from individual chamber-based urine patch fluxes to field scale fluxes can cause uncertainty for livestock based systems. Typically, livestock have to be excluded from plots with chambers, yet, the presence of

livestock is likely to alter emissions e.g. no consideration of compaction caused by hooves or overlapping of urine patches (de Klein et al., 2006; Pleasants et al., 2007; Cichota et al., 2013). The use of collars which are low lying to the ground would allow livestock to be present within a chamber study site. Upscaling of emissions would also use mean values for urine volume and frequency per head of livestock, yet, this may cause uncertainties due to differences in N processing rates of patches of different sizes.

The IPCC Tier 1 approach to calculating N<sub>2</sub>O emission factors is too simplistic and does not consider the effect of soil type, management strategies or climatic variables on excretal emissions (de Klein et al., 2001; Skiba et al., 2012). A move to IPCC Tier 2 and 3 approaches is necessary to better constrain N<sub>2</sub>O emission estimates from grasslands, where far more regional measurements of N<sub>2</sub>O emissions from sheep excreta will be required. Disaggregation of N<sub>2</sub>O emissions based on rainfall data and soil drainage class may be warranted, as variation between years may result in large biases of emission estimates (van der Weerden et al., 2012). In New Zealand, disaggregation of emission factors based on hill slopes has also been suggested to reflect the spatial difference in soil conditions and livestock resting behaviour in these areas, as most emission factor measurements are based on flat or lowland areas, which could represent an overestimation of N<sub>2</sub>O emissions (Luo et al., 2013; Saggar et al., 2015). The effect of future climate scenarios are also important to consider, such as the expected increase in extreme weather events e.g. drought and flooding. Both could significantly affect N<sub>2</sub>O emissions estimates, where emissions would be expected to be lower under drought conditions and higher under periods of heavy rainfall.

#### **8.4 How should the chamber technique be deployed, to monitor N<sub>2</sub>O emissions from urine patches?**

The second objective was to determine how to accurately monitor N<sub>2</sub>O emissions from urine patches, using the closed chamber technique. In Chapter 3, the importance of considering the urine patch diffusional area whilst monitoring N<sub>2</sub>O emissions from urine patches was assessed. From the results of this study, it is recommended that, when calculating urine patch N<sub>2</sub>O emission factors, extra room inside static chambers should be allowed for the diffusion of solutes into surrounding soil. Further work could be conducted here, by comparing the actual restriction effect posed by chamber walls in the field and assessing whether considering the pasture response area also influences N<sub>2</sub>O emission measurements. Methods of measurements which do not disturb soil conditions or impact on the diffusion of solutes could be utilised, such as eddy covariance methods, or the use of sub-surface gas probes (Li and Kelliher, 2005) which do not rely on excluding stock from an area, which can influence N<sub>2</sub>O emission e.g. by compaction with hooves. In addition, field validation of diffusion coefficient calculations for urine constituents should be conducted, where this approach could be used to estimate chamber size requirements to cover a urine patch.

In Chapter 4, an assessment of the headspace sampling frequency from static chambers following urine application was made. From this, it is recommended that where resources are available, sampling as close to daily as possible between 10 am and 12 noon, would closely resemble the fluxes as measured by an automated system. In Chapter 7, the diurnal nature of N<sub>2</sub>O fluxes were examined, and maximum fluxes were observed during the night or early morning. It is likely that the diurnal pattern of N<sub>2</sub>O emissions depends on land-use type, and site specific characteristics e.g. gas diffusivity through the soil profile. Therefore, where possible, an assessment of the diurnal variation in N<sub>2</sub>O fluxes in the area of study should be

conducted prior to a manual sampling campaign. This will allow an assessment of the best time of day to sample to represent the average daily flux.

A large spatial variability in N<sub>2</sub>O emissions was observed in the field trials of Chapters 4 and 7, despite the exclusion of livestock several months prior to treatment application. It is advised that replicate chambers should be maximised as much as resources allow, in order to capture this spatial variability. Automated chamber systems are generally limited by the number of available chambers, therefore, treatment numbers would need to be minimized to ensure sufficient replication, and hence an adequate spatial representation of N<sub>2</sub>O emissions. It is also noted that this spatial variability can influence ancillary measurements taken to support N<sub>2</sub>O emission factors. When taking ancillary measurements from duplicated urine patches, N processing rates can differ. Advances in non-destructive techniques to monitor urine patch characteristics will be important for future urine patch research.

### **8.5 Are nitrification inhibitors a suitable strategy for reducing N<sub>2</sub>O emissions from grazed grasslands?**

The third objective was to investigate the factors which influence nitrification inhibitor efficacy in ruminant urine-influenced soils and assess their potential as an N<sub>2</sub>O mitigation strategy for sheep grazed grasslands. In Chapter 5, the vertical mobility and factors influencing vertical mobility (solubility, sorption/desorption, microbial uptake and degradation) of the nitrification inhibitors DCD and DMPP were assessed in three soils contrasting in texture and organic matter content, with and without the presence of sheep urine. It was found that the presence of sheep urine altered the depth distribution of both nitrification inhibitors, and increased desorption of the nitrification inhibitors, where this effect was soil and inhibitor type dependent. Sorption by soil was greater for DCD in comparison to DMPP in all tested soils and a large proportion of the applied inhibitors were immobilized by the microbial biomass.

Immobilization of nitrification inhibitors into non-target microbial biomass could equate to a large depletion of the bioavailable nitrification inhibitor pool. In this Chapter, it was shown that soil type is a regulator of the factors which can influence nitrification inhibitor efficacy. Further work is required to determine the comparative efficacy of DCD and DMPP in reducing nitrification rates and N<sub>2</sub>O emissions from sheep urine patches, across a similar range of soil types. This may allow recommendations of which nitrification inhibitor, if any, are more effective under contrasting soil types.

In Chapter 6, the capacity for graminaceous plants to directly acquire DCD through roots, and to translocate this to shoots, was demonstrated. It was concluded that this is a potential vector for DCD entry into the food chain, and is one of the likely causes of the contamination of New Zealand milk products with DCD. The contamination of milk products from DCD-treated pasture has resulted in some consumer scepticism of nitrification inhibitor use, and alternatives are currently being sought (Bates et al., 2015). While DMPP is a different compound, and may not be as phytotoxic as DCD (e.g. leaf necrosis at 8 times the recommended application rate; Zerulla et al., 2001), it is still a synthetic compound which may also attract similar scepticism. A direct comparison of plant acquisition of both DCD and DMPP should be made, to confirm the same outcome will not occur with DMPP. Screening of plants capable of exuding biological nitrification inhibitors, through plant breeding, may prove to be a more fruitful approach (Subbarao et al., 2006).

The use of nitrification inhibitors in grazed pastures would be fairly quick to implement if a cost-effective delivery mechanisms can be developed, and they represent a useful strategy for improving nitrogen use efficiency in grazed pastures. However, the benefits associated with nitrification inhibitor use varies widely, suggesting benefits may not always outweigh the cost of application. This has implications for the widespread adoption of any mitigation strategy. In Chapter 7, DMPP applied to sheep urine patches under summer conditions was completely

ineffective at reducing cumulative N<sub>2</sub>O emissions, even when applied at the same time as the urine. Further work is required to assess the reasons why it may not have been effective, where potentially higher application rates may have been required (which would also influence the cost of application) or temperatures may have been too high to render it effective.

Targeting urine patches within space and time is another major challenge associated with the effective use of nitrification inhibitors, and differing application methods will be a key area of research to bring this technology from a tried and tested concept to a widely adopted mitigation strategy. While orally administering synthetic nitrification inhibitors to ruminants has been examined, where large amounts of consumed inhibitor were excreted alongside the urine, this approach needs extensive ethical considerations. In particular, producers and consumers need assurance that there are no adverse effects of the inhibitors on grazing animal health and to ensure that no contamination would be present in milk or meat tissues destined for human consumption (Ledgard et al., 2008). As a wide range of compounds are able to inhibit nitrification, there may be compounds naturally present within urine or forage plants which could be utilised for targeted inhibitor-to-urine applications. Here, the screening of urine chemical constituents or feed constituents for nitrification inhibitor potential e.g. attempted study shown in Appendix 2, using a lux marked strain of *Nitrosomonas europaea*, could be a useful approach for novel nitrification inhibitor identification (Subbarao et al., 2006).

## **8.6 What alternative strategies are there for the mitigation of N<sub>2</sub>O emissions from urine patches?**

The mitigation of N<sub>2</sub>O from urine patches is likely to require a combination of different strategies, to be most effective, which will need to be assessed by life cycle analysis and whole-farm system modelling approaches, in order to ensure the avoidance of pollution swapping (Eckard et al., 2010; Gerber et al., 2013; Schils et al., 2013). On-farm N<sub>2</sub>O mitigation strategies


which have been utilised, tested or conceived can be broadly divided into plant, animal, urine, soil and management-based interventions, as shown in Figure 8.1. Nevertheless, some of these options would fit under several of the categories suggested here.

#### *8.6.1 Plant-based mitigation strategies*

Plant-based strategies to reduce urine patch N<sub>2</sub>O emissions include the use of high-sugar ryegrass varieties in pastures. Such grasses have been shown to improve rumen N utilisation, potentially reducing N excreted in the urine (Moir et al., 2001; Edward et al., 2007). Pacheco et al. (2010) found urine N concentration to be inversely related to dietary cation-anion intake and soluble sugar content of differing forages, potentially reducing the N loading rate of individual urine events. Further field validation of potential effects on N<sub>2</sub>O emissions and N utilisation by livestock are still required.

Condensed tannins are another potential plant-based strategy to reduce urine derived N<sub>2</sub>O. Condensed tannin extracts are currently expensive, however, plant breeding may offer an approach to increase tannin levels in forage plants (Smith et al., 2010). Supplementation of condensed tannin extract has been shown to increase the proportion of N excreted in the faeces, and lower the N in the urine of both sheep (Carulla et al., 2005) and cattle (Misselbrook et al., 2005; Grainger et al., 2009). This is due to the complexation of condensed tannins with proteins in the rumen, which results in some protection against microbial degradation and can serve to





**Figure 8.1** Strategies to reduce N losses from grazed pastures.

increase the absorption of amino acids in the abomasum or lower intestine, or the complex can be excreted in the faeces (Min et al., 2003; de Klein and Eckard, 2008; Smith, 2010). Other plant-based strategies which might reduce total N losses from grazed pastures is breeding plants which can intercept and acquire more urine or fertilizer N through differing root architecture, pasture with high yields but lower N contents and the inclusion of clover in pastures to increase biologically fixed N (Ledgard et al., 2009; de Klein et al., 2010; Malcolm et al., 2014; Selbie et al., 2015).

The biological exudation of N process inhibitors is also an emerging option for reducing N<sub>2</sub>O emissions from grazed grasslands. Compounds could also potentially enter the soil via root death, or, if present in shoot material it could potentially be consumed by livestock and if protected from degradation in the rumen, excreted in the urine. Several plants have been identified to exude nitrification inhibiting compounds including the tropical grass *Brachiaria* spp. and Sorghum, however, the capacity varies between species and further establishment of the effects under field conditions is warranted (Subbarao et al., 2007; Subbarao et al., 2009; Zeng et al., 2015). Recent studies have also utilised biochar plant-extract complexes, where biological nitrification inhibitors are adsorbed onto the biochar, prolonging their effective period in the soil (Muñoz et al., 2014; Reyes-Escobar et al., 2015). The potential for biological denitrification inhibition to occur via plant secondary metabolites has been shown in Bardon et al. (2014), yet this is still an understudied area. There is also current interest in understanding the physiological and genetic plant traits which regulate N<sub>2</sub>O emissions, in order to potentially improve agronomic performance alongside environmental sustainability (Thomson et al., 2012).

### 8.6.2 Soil-based mitigation strategies

In addition to the application of synthetic nitrification inhibitors, inhibiting or promoting other N cycle processes could also reduce N<sub>2</sub>O emissions. The nitrification inhibitor DCD has been shown to be more effective in reducing N<sub>2</sub>O emissions, NH<sub>3</sub> volatilisation and pasture N uptake from cow urine when applied in combination with a urease inhibitor e.g. N-(n-butyl) thiophosphoric triamide (nBTPT), than when applied alone (Zaman et al., 2009). Although urease inhibitors are not particularly effective at reducing N<sub>2</sub>O emissions when used solely, the combined use with nitrification inhibitors is a sensible approach, as the use of nitrification inhibitors alone can increase ammonia volatilisation, as the N remains in the soil as ammonium for longer (Davies and Williams, 1995).

Saggar et al. (2013) describes denitrification as both an environmental threat and an opportunity. The loss of N as N<sub>2</sub>O via denitrification could be prevented, or on the other hand, promotion of complete denitrification to N<sub>2</sub> via nitrous oxide reductase is another approach to reducing N<sub>2</sub>O (Thomson et al., 2012). Factors which influence the product ratio of N<sub>2</sub>O/N<sub>2</sub> includes soil pH, which has led to the suggestion of liming as an N<sub>2</sub>O mitigation strategy (Clough et al., 2003; Zaman et al., 2007; Bakken et al., 2012; McMillan et al., 2015). The use of lime has been studied for potential effects on encouraging complete denitrification to N<sub>2</sub>, and the use of the soil amendment zeolite has also been considered for reducing the NH<sub>4</sub><sup>+</sup> pool available for nitrification (Zaman et al., 2007; Zaman et al., 2008; Zaman and Nguyen, 2010; McMillan et al., 2015). Zeolite can absorb NH<sub>4</sub><sup>+</sup>, as it is a negatively charged natural aluminosilicate (Zaman et al., 2007), which may reduce the available supply of NH<sub>4</sub><sup>+</sup> for nitrification and the amount of available downstream NO<sub>3</sub><sup>-</sup> for denitrification. Zeolite has been found to reduce N<sub>2</sub>O emissions from urine-influenced soils, and liming has been shown to increase the amount of N<sub>2</sub>, as opposed to N<sub>2</sub>O from urine-influenced soils (Zaman et al., 2007; Zaman et al., 2010).

Biochar is an additional soil amendment, which, alongside increasing C sequestration in grassland may also have benefits for N cycling in grazed pastures (Taghizadeh-Toosi et al., 2011; Taghizadeh-Toosi et al., 2012). Biochar has been increasingly recognised for its interaction with the N cycle, where the  $\text{NH}_3$  and  $\text{NH}_4^+$  can absorb into the porous structure (Clough et al., 2009a). Recently, biochar has also been found to reduce the  $\text{N}_2\text{O}/\text{N}_2$  product ratio, facilitating the last step of denitrification, possibly by facilitating the transfer of electron to denitrifiers (Cayuela et al., 2013). In urine-influenced soil the effect of biochar on denitrification appears to be mixed, where some trials have shown a reduction (Taghizadeh-Toosi et al., 2011) and others have shown no change in nitrous oxide emissions (Clough et al., 2009a; Anderson et al., 2014).

Copper has been shown to down regulate the nitrous oxide reductase gene, *NosZ* (Richardson et al., 2009; Sullivan et al., 2013), therefore, when copper is in limited supply the fraction of N released as  $\text{N}_2\text{O}$  from soils may increase. Ensuring copper in soil is non-limiting is a potential strategy to avoid this, via the use of copper-based fertilizers, although to our knowledge, this has not been assessed in any field trials. As denitrification is inhibited by oxygen, techniques which promote soil aeration and reduce soil compaction could be important to consider (Eckard et al., 2010). Moving the location of water and feed containers in the field would prevent several urine patches being deposited to the same area, which can exacerbate N losses (Shepherd and Chambers, 2007; Matthews et al., 2010). Other areas such as sheep tracks and gateways could potentially have greater  $\text{N}_2\text{O}$  emissions, which could be targeted within  $\text{N}_2\text{O}$  emission reduction strategies (Matthews et al., 2010).

### 8.6.3 *Animal-based mitigation strategies*

Animal-based strategies to reduce  $\text{N}_2\text{O}$  emissions would involve the selective breeding of ruminants for ‘environmentally friendly’ traits. While there are no such breeding

interventions taking place to our knowledge, the suggestion of breeding livestock which walk while they urinate or urinate more frequently have been made (de Klein and Eckard, 2008; de Klein et al., 2010), in order to improve the spread of urine N across the pasture. Additional improvements may also be made by improving the N conversion efficiency in the rumen of livestock i.e. the amount of N converted to meat or milk (de Klein et al., 2010).

#### *8.6.4 Management strategies*

Several management strategies could be employed for tightening the N cycle in grazed pastures. The rate, source, timing and method of fertilizer applications should be carefully considered to ensure it matches the peak N demand of plants and it should not be applied under conditions of heavy rainfall (Eckard et al., 2010). The interactive effect of fertilizer application and time of urine deposition has also been shown to influence N<sub>2</sub>O emissions and NO<sub>3</sub><sup>-</sup> leaching (Anger et al., 2003; Buckthought et al., 2011). Grazing could be restricted for a period of time following fertilizer application as a potential solution to avoid any fertilizer-urine interaction effects. Restricting grazing at other risk periods e.g. winter periods where rainfall is high and compaction by hooves is more likely (de Klein et al., 2006), has also been suggested as an N<sub>2</sub>O mitigation strategy. Effluent from livestock can then be collected and re-applied when N demand and soil conditions are appropriate, which can also minimise N losses. While this is an option for certain cattle systems, sheep grazing is often extensive, where animals graze all year round. Further research is required to determine mitigation strategies specifically for extensive systems.

The avoidance of camping areas through grazing management may reduce N<sub>2</sub>O emissions, such as through grazing management e.g. rotational grazing, as opposed to continual grazing (Petersen et al., 2004; Dahlin et al., 2005). On the other hand, knowledge of camping areas could be viewed as an opportunity for the use of targeted mitigation strategies e.g.

nitrification inhibitors. Further studies are required on rotational grazing systems to determine how differences in management strategies effect N<sub>2</sub>O emissions. Generally, increasing stocking rates will increase N<sub>2</sub>O emissions, however, rotational grazing can cause less or more N<sub>2</sub>O emissions depending on management, climatic and regional factors (Saggar et al., 2007). Silvopastoral systems may increase shading/shelter areas for livestock, where tree roots may help acquire N leached beyond the pasture root zone, which could be emitted as N<sub>2</sub>O indirectly (Schoeneberger et al., 2012).

Balancing the protein-to-energy ratio of ruminant diets can reduce the amount of urinary N excreted. As discussed previously, this could be achieved through the use of high-sugar fodder varieties; alternatively the provision of high-energy supplements can be provided where animals are grazing high protein grass (Eckard et al., 2010). The inclusion of salt, or other diuretic compounds may reduce the N load per urination by increasing total urinary volume and increasing the spread of urine across the pasture (Ledgard et al., 2007; Costall and Betteridge, 2010; Pacheco et al., 2010; Liu and Zhou, 2014).

#### 8.6.5 Demand-led strategies

The simplest strategy to reducing N<sub>2</sub>O emissions from urine patches would be to decrease livestock numbers, and hence the total amount of excreta returned to land (Luo et al., 2010; Ussiri and Lal, 2012). This could be achieved through a reduction in meat protein consumption *per capita*, or even a complete switch to plant-based protein consumption, which under simulated modelled scenarios have been shown to cause large reductions in agricultural non-CO<sub>2</sub> greenhouse gases (Stehfest et al., 2009; Popp et al., 2010; Westhoek et al., 2014). Nevertheless, changes in the supply and demand for animal products and the change in land-use associated with any reduction in animal product consumption are highly uncertain (Westhoek et al., 2014; Decock et al., 2015). In addition, the effects on livestock producers and

downstream processing and supply chains, which provide income to rural economies, are likely to be negative and have to be considered (Westhoek et al., 2014). Livestock production can also occur on marginal land, which is not suitable for growing crops (Kingston-Smith et al., 2010). Encouraging a reduction in animal product consumption, alongside reductions in food wastage and better transport and supply chains, is a sensible strategy to reduce agricultural GHG emissions, nevertheless, it is considered unlikely that widespread vegan/vegetarianism would be adopted and with increasing population, global projections point in the opposite direction (Smith, 2010; Thornton et al., 2010). Direct mitigation strategies at the farm-scale will inevitably have an important role to play in emission reductions from the livestock sector in the short term.

Despite a wide range of potential mitigation strategies for N<sub>2</sub>O, implementation of techniques are lagging, and the development of N<sub>2</sub>O specific policies may be required (Schils et al., 2013). Nitrification inhibitors, fertilizer management and managing animal diets have been considered to show the best potential for N<sub>2</sub>O emission reductions in the short term (de Klein and Eckard, 2008). Whilst some of the techniques described here need further research and development, the solutions adopted by farmers will more than likely depend on the practicality and economics of individual farms (Luo et al., 2010). Strategies to reduce N<sub>2</sub>O emissions from urine patches could be incorporated into agri-environmental schemes, in order to encourage adoption. Farmers may need assistance in an advisory capacity, to manage N efficiently and to ensure the avoidance of pollution swapping. Several of the suggested mitigation strategies are more suited to intensive livestock systems, and further novel strategies may be required under more extensive systems (Eckard et al., 2010).

## **8.7 How should future research be prioritised, in order to achieve enhanced productivity and sustainability in grazed grasslands?**

Achieving enhanced productivity and sustainability in grazed grassland is a complex challenge, which will require a multidisciplinary approach to resolve. In order to understand the spatial and temporal variability of soil processes and N transformations at the scale of the urine patch, advances will need to be made in order to monitor urine patches in as non-destructive a manner as possible. The use of planar optodes to measure changes in soil pH, O<sub>2</sub>, CO<sub>2</sub> and NH<sub>4</sub><sup>+</sup> have been described for use in the rhizosphere (Blossfeld et al., 2011; Blossfeld, 2013), where such technology may also be useful for studying urine-influenced soil, although the technique is confined to two dimensions. The use of N<sub>2</sub>O microelectrodes (Elberling et al., 2010) and the development of *in-situ* electrodes to monitor soil mineral N (Shaw et al., 2013), could provide a method of studying the urine patches without causing excessive soil disturbance. Fluorescence *in situ* hybridization (FISH) based approaches have been reported for use in identifying microbial hotspots and microbial groups within hotspots in either 2D or 3D surfaces (e.g. soil root interface; Musat et al., 2008; Schmidt et al., 2012; Schmidt and Eickhorst, 2013; Schmidt and Eickhorst, 2014; Kuzyakov and Blagodatskaya, 2015). Knowledge of wider N cycling in the urine patch is lacking, where several process rates need further study, such as determining nitrification, denitrification and mineralization in differing zones of the urine patch. There is limited research regarding the influence of urine on DOC and DON solubilisation, and whether the increase in DOC fuels denitrification. Modelling N process rates in the urine patch using <sup>15</sup>N tracing models would provide useful insight into the fate of urinary N (Müller et al., 2004; Müller et al., 2007; McGeough et al., 2016). Recently, N<sub>2</sub>O has also been shown to be produced by ammonia-oxidising bacteria present on the leaves of pasture plants influenced by urine, where further investigations should be conducted to quantify the magnitude of this N<sub>2</sub>O source (Bowatte et al., 2015).



Understanding the source partitioning (major soil process; nitrification or denitrification) of N<sub>2</sub>O release from soil will be fundamental for improving N<sub>2</sub>O mitigation strategies (Köster et al., 2013), as some techniques may prevent one microbial N<sub>2</sub>O pathway but promote another. It may also improve understanding of the environmental regulators of N<sub>2</sub>O production pathways, aside from nitrification and denitrification. Advances in technology able to continuously monitor the isotopomers and site preference of N<sub>2</sub>O in the field, promise a greater understanding of the source partitioning of N<sub>2</sub>O production, where this technique is currently known to distinguish between N<sub>2</sub>O produced via hydroxylamine reduction, nitrate reduction, and between bacterial and fungal denitrification (Thomson et al., 2012). The ‘omics’ (genomics, proteomics, metabolomics) era is also another important research area, which could advance understanding of N cycling and losses in the urine patch. Additionally, omics of urine itself may provide better knowledge of urinary compounds, some of which are known to influence N<sub>2</sub>O production and emission from soil e.g. hippuric acid and benzoic acid (Kool et al., 2006; Bertram et al., 2009; Clough et al., 2009b). There is a general need for more research relating the microbial composition and functioning in urine-influenced soils, in relation to N<sub>2</sub>O emissions. Process-based models will continually need to be developed and updated to incorporate advances in our knowledge of N<sub>2</sub>O production in soils.

In terms of mitigating N losses from the urine patch efficiently, targeting the areas of the pasture in space and time is another challenge which needs to be met. A few novel technologies have been made and are continuing to be developed for cattle-based systems, including the use of a tail mounted nitrification inhibitor dispensing device, which dispenses a given amount of inhibitor following urination (Bates and Quin, 2013). The use of robotised vehicles have also been developed (Bates et al., 2015), sensing where the urine patches are located via fluorescence, temperature or chlorophyll measurements. Nevertheless, the time taken for the pasture to respond to a urine application can be long after the N loss risk period,

and temperature can dissipate too quickly for these strategies to be useful (Bates and Quin, 2013; Bates et al., 2015). This has led to the more recent development of a towable or automated technique, of measuring the electrical conductivity and moisture content of the pasture soil, which resulted in a good detection of urine patches (Bates et al., 2015). This method of sensing the urine patch has been described for use with a combination of a urease inhibitor, gibberellic acid and a dissolved organic C supply (a fulvic acid polymer), in order to delay urea hydrolysis, promote plant growth in the urine patch and reduce N losses, where preliminary trials showed a comparative efficacy to DCD (Bates et al., 2015). Further advances in such technologies will be important for providing practical and cost-effective measures to reduce N losses at the farm scale.

Care should be taken when selecting mitigation strategies, to ensure that ‘pollution swapping’ does not occur. Some strategies which reduce N<sub>2</sub>O emissions may exacerbate losses of other N forms, or other greenhouse gases e.g. housing livestock would result in manure piles which generate ammonia, and would require bought in concentrates, which is more carbon intensive. When considering mitigation strategies, a whole life cycle approach needs to be considered, in order to consider the potential trade-offs which may result due to implementation of such measures.

## **8.8 Conclusions**

The N concentration and size of sheep urine patches have been shown to influence N<sub>2</sub>O emissions, therefore, further data is required on urine volume, frequency and N concentrations under a range of management techniques and climatic variables. Environmental parameters exert a strong influence on N<sub>2</sub>O emissions from urine patches, where measured N<sub>2</sub>O emissions from sheep urine patches under summer conditions were lower than the default IPCC Tier 1 approach to calculating N losses from sheep excreta. A move to Tier 2 and 3 approaches will

be fundamental to better constrain N<sub>2</sub>O emission estimates from grasslands, based on variability in soil type, hill slopes, climate and management related factors.

When utilising chambers to calculate N<sub>2</sub>O emission factors from sheep urine patches, sufficient room should be allowed for the urine patch diffusional area (and potentially the pasture response area) in order to reflect emissions as closely as possible. If using a manual sampling campaign sampling should be conducted as close to daily as possible, an assessment of the diurnal variability of N<sub>2</sub>O emissions is recommended and chamber number should be maximized under both manual and automated campaigns, in order to adequately capture spatial variation in N<sub>2</sub>O emissions.

The capacity for plants to directly acquire DCD and translocate this compound to the shoots, as demonstrated in this thesis, represents a potential vector for the contamination of livestock products when using inhibitor, where the use of synthetic nitrification may attract livestock product consumer scepticism. This combined with the apparent lack of effectiveness of DMPP in reducing N<sub>2</sub>O emissions from sheep urine patches, suggests the use of nitrification inhibitors alone will not be sufficient to entirely reduce N<sub>2</sub>O emissions from urine patches. The efficacy of nitrification inhibitors does vary widely, yet, they represent a mitigation strategy which could be implemented in the short term.

Achieving enhanced sustainability and productivity in livestock systems is a challenging and complex problem, which will require expertise and collaboration across conventional scientific disciplines, policy-makers and farmers. There are, however, several mitigation strategies available, being developed, or some which require much more research before being practicable. Advances in technologies to measure and mitigate N<sub>2</sub>O emissions will greatly enhance our knowledge of N cycling in the urine patch environment in the near future.

## 8.9 References

- Anderson, C.R., Hamonts, K., Clough, T.J., Condrón, L.M. 2014. Biochar does not affect soil N-transformations or microbial community structure under ruminant urine patches but does alter relative proportions of nitrogen cycling bacteria. *Agriculture, Ecosystems and Environment* 191, 63-72.
- Anger, M., Hoffman, C., Kühbauch, W. 2003. Nitrous oxide emissions from artificial urine patches applied to different N-fertilized swards and estimated annual N<sub>2</sub>O emissions for differently fertilized pastures in an upland location in Germany. *Soil Use and Management* 19, 104-111.
- Bakken, L.R., Berkaust, L., Liu, B., Frostegard, A. 2012. Regulation of denitrification at the cellular level: a clue to the understanding of N<sub>2</sub>O emissions from soils. *Philosophical Transactions of the Royal Society B: Biological Sciences* 367, 1226-1234.
- Bardon, C., Piola, R., Bellvert, F., Haichar, F.Z., Comte, G., Meiffren, G., Pommier, T., Puijalón, S., Tsafak, N., Poly, F. 2014. Evidence for biological denitrification inhibition (BDI) by plant secondary metabolites. *New Phytologist* 204, 620-630.
- Bates, G., Quin, B. 2013. Low-cost detection and treatment of fresh cow urine patches. *Proceedings of the New Zealand Grassland Association* 75, 125-130.
- Bates, G., Quin, B., Bishop, P. 2015. Low-cost detection and treatment of fresh cow urine patches. Pastoral Robotics Ltd., Auckland, New Zealand.
- Bertram, J.E., Clough, T.J., Sherlock, R.R., Condrón, L.M., O'Callaghan, M., Wells, N.S., Ray, J.L. 2009. Hippuric acid and benzoic acid inhibition of urine derived N<sub>2</sub>O emissions from soil. *Global Change Biology* 15, 2067-2077.
- Betteridge, K., Costall, D., Balladur, S., Upsdell, M., Umemura, K. 2010. Urine distribution and grazing behaviour of female sheep and cattle grazing a sheep New Zealand hill pasture. *Animal Production Science* 50, 624-629.

- Betteridge, K., Costall, D.A., Li, F.Y., Luo, D., Ganesh, S. 2013. Why we need to know what and where cows are urinating – a urine sensor to improve nitrogen models. In Proceedings of the New Zealand Grassland Association 75, 119-124.
- Blossfeld, S. 2013. Light for the dark side of plant life:- planar optodes visualizing rhizosphere processes. Plant and Soil, 369, 29-32.
- Blossfeld, S., Gansert, D., Thiele, B., Kuhn, A.J., Lösch, R. 2011. The dynamics of oxygen concentration, pH value and organic acids in the rhizosphere of *Juncus* spp. Soil Biology and Biochemistry 43, 1186-1197.
- Bowatte, S., Newton, P.C.D., Brock, S., Theobald, P., Luo, D. 2015. Bacteria on leaves: a previously unrecognised source of N<sub>2</sub>O in grazed pastures. The ISME Journal 9, 265-267.
- Buckthought, L., Snow, V.O., Shepherd, M.A., Clough, T.J., Cameron, K.C., Di, H. 2011. Nitrogen fertiliser and urine patch interaction – use of APSIM to aid experimental design. Fertilizer and Lime Research Centre, Massey University: Palmerston North, New Zealand. Available at <http://flrc.massey.ac.nz/publications.html> [accessed 23 November 2015].
- Carulla, J.E., Kreuzer, M., Machmüller, A., Hess, H.D. 2005. Supplementation of *Acacia mearnsii* tannins decreases methanogenesis and urinary nitrogen in forage-fed sheep. Crop and Pasture Science 56, 961-970.
- Cayuela, M.L., Sánchez-Monedero, M.A., Roig, A., Hanley, K., Enders, A., Lehmann, J. 2013. Biochar and denitrification in soils: when, how much and why does biochar reduce N<sub>2</sub>O emissions? Scientific Reports 3, 1-7.
- Cichota, R., Snow, V.O., Vogeler, I. 2013. Modelling nitrogen leaching losses from overlapping urine patches. Environmental Modelling and Software 41, 15-26.

- Clough, T.J., Bertram, J.E., Ray, J.L., Condon, L.M., O'Callaghan, M., Sherlock, R.R., Wells, N.S. 2009a. Unweathered wood biochar impact on nitrous oxide emissions from a bovine-urine-amended pasture soil. *Soil Science Society of America Journal* 74, 852-860.
- Clough, T.J., Ray, J.L., Buckthought, L.E., Calder, J., Baird, D., O'Callaghan, M., Sherlock, R.R., Condon, L.M. 2009b. *Soil Biology and Biochemistry* 41, 2222-2229.
- Clough, T.J., Sherlock, R.R., Kelliher, F.M. 2003. Can liming mitigate N<sub>2</sub>O fluxes from urine-amended soil? *Soil Research* 41, 439-457.
- Costall, K., Betteridge, K. 2010. Methods of delivering salt to cattle. *Proceedings of the New Zealand Society of Animal Production* 70, 296-298.
- Dahlin, A.S., Emanuelsson, U., McAdam, J.H. 2005. Nutrient management in low input grazing-based systems of meat production. *Soil Use and Management* 21, 122-131.
- Davies, D.M., Williams, P.J. 1995. The effect of the nitrification inhibitor dicyandiamide on nitrate leaching and ammonia volatilization: A U.K. nitrate sensitive areas perspective. *Journal of Environmental Management* 45, 263-272.
- Decock, C., Lee, J., Necpalova, M., Pereira, E.I.P., Tendall, D.M., Six, J. 2015. Mitigating N<sub>2</sub>O emissions from soil: from patching leaks to transformative action. *Soil Discussions* 2, 903-915.
- de Klein, C.A.M., Eckard, R.J. 2008. Targeted technologies for nitrous oxide abatement from animal agriculture. *Animal Production Science* 48, 14-20.
- de Klein, C.A.M., Monaghan, R.M., Ledgard, S.F., Shepherd, M. 2010. A system's perspective on the effectiveness of measures to mitigate the environmental impacts of nitrogen losses from pastoral dairy farming. *Proceedings of the Australasian Dairy Science Symposium* 4, 14-28.

- de Klein, C.A.M., Sherlock, R.R., Cameron, K.C., van der Weerden, T.J. 2001. Nitrous oxide emissions from agricultural soils in New Zealand – a review of current knowledge and directions for future research. *Journal of the Royal Society of New Zealand* 31, 543-574.
- de Klein, C.A.M., Smith, L.C., Monaghan, R.M. 2006. Restricted autumn grazing to reduce nitrous oxide emissions from dairy pastures in Southland, New Zealand. *Agriculture, Ecosystems and Environment* 112, 192-199.
- Eckard, R.J., Grainger, C., de Klein, C.A.M. 2010. Options for the abatement of methane and nitrous oxide from ruminant production: A review. *Livestock Science* 130, 47-56.
- Edwards, G.R., Parsons, A.J., Rasmussen, S., Bryant, R.H. 2007. High sugar ryegrasses for livestock systems in New Zealand. *New Zealand Grassland Association*, 161-171.
- Elberling, B., Christiansen, H.H., Hansen, B.U. 2010. High nitrous oxide production from thawing permafrost. *Nature Geoscience* 3, 332-335.
- FAO, 2001. "World Soil Resources Reports" In: Driessen, P. Deckers, J. (Eds), *Lecture Notes on the Major Soils of the World*. Rome, pp 1 -10.  
<http://www.fao.org/docrep/003/Y1899E/y1899e00.htm#toc>
- Grainger, C., Clarke, T., Auld, M.J., Beauchemin, K.A., McGinn, S.M., Waghorn, G.C., Eckard, R.J. 2009. Potential use of *Acacia mearnsii* condensed tannins to reduce methane emissions and nitrogen excretion from dairy cows. *Canadian Journal of Animal Science* 89, 241-251.
- Gerber, P.J., Hristov, A.N., Henderson, B., Makkar, H., Oh, J., Lee, C., Meinen, R., Montes, F., Ott, T., Firkins, J., Rotz, A., Dell, C., Adesogan, A.T., Yang, W.Z., Tricarico, J.M., Kebreab, E., Waghorn, G., Dijkstra, J., Oosting, S. Technical options for the mitigation of direct methane and nitrous oxide emissions from livestock: a review. *Animal* 7, 220-234.

- Kingston-Smith, A.H., Edwards, J.E., Huws, S.A., Kim, E.J., Abberton, M. 2010. Plant-based strategies towards minimising 'livestock's long shadow'. *Proceedings of the Nutrition Society* 69, 613-620.
- Kool, D.M., Hoffland, E., Hummelink, E.W.J., van Groenigen, J.W. 2006. Increased hippuric acid content of urine can reduce soil N<sub>2</sub>O fluxes. *Soil Biology and Biochemistry* 38, 1021-1027.
- Köster, J.R., Well, R., Tuzson, B., Bol, R., Dittert, K., Gieseemann, A., Emmenegger, L., Manninen, A., Cárdenas, L., Mohn, J. 2013. Novel laser spectroscopic techniques for continuous analysis of N<sub>2</sub>O isotopomers – application and intercomparison with isotope ratio mass spectrometry. *Rapid Communications in Mass Spectrometry* 27, 216-222.
- Kuzyakov, Y., Blagodatskaya, E. 2015. Microbial hotspots and hot moments in soil: Concept & review. *Soil Biology and Biochemistry* 83, 184-199.
- Ledgard, S.F., Menneer, J.C., Dexter, M.M., Kear, M.J., Lindse, S., Peters, J.S., Pacheco, D. 2008. A novel concept to reduce nitrogen losses from grazed pastures by administering soil nitrogen process inhibitors to ruminant animals: a study with sheep. *Agriculture, Ecosystems and Environment* 125, 148-158.
- Ledgard, S., Schils, R., Eriksen, J., Luo, J. 2009. Environmental impacts of grazed clover/grass pastures. *Irish Journal of Agricultural and Food Research* 48, 209-226.
- Ledgard, S.F., Welten, B.G., Menneer, J.C., Betteridge, K., Crush, J.R., Barton, M.D. 2007. New nitrogen mitigation technologies for evaluation in the Lake Taupo catchment. *Proceedings of the New Zealand Grassland Association* 69, 117-121.
- Li, F.Y., Betteridge, K., Cichota, R., Hoogendoorn, C.J., Jolly, B.H. 2012. Effects of nitrogen load variation in animal urination events on nitrogen leaching from grazed pastures. *Agriculture, Ecosystems and Environment* 159, 81-89.



- Li, Z., Kelliher, F.M. 2005. Determining nitrous oxide emissions from subsurface measurements in grazed pasture: a field trial of alternative technology. *Australian Journal of Soil Research* 43, 677-687.
- Liu, H., Zhou, D. 2014. Mitigation of ammonia and nitrous oxide emissions from pasture treated with urine of sheep fed diets supplemented with NaCl. *Animal Feed Science and Technology* 192, 39-47.
- Luo, J., de Klein, C.A.M., Ledgard, S.F., Saggar, S. 2010. Management options to reduce nitrous oxide emissions from intensively grazed pastures: a review. *Agriculture, Ecosystems and Environment* 136, 282-291.
- Luo, J., Hoogendoorn, C., van der Weerden, T., Saggar, S., de Klein, C., Giltrap, D., Rollo, M., Rys, G. 2013. Nitrous oxide emissions from grazed hill land in New Zealand. *Agriculture, Ecosystems and Environment* 181, 58-68.
- Malcolm, B.J., Cameron, K.C., Di, H.J., Edwards, G.R., Moir, J.L. 2014. The effect of four different pasture species compositions on nitrate leaching losses under high N loading rate. *Soil Use and Management* 30, 58-68.
- Matthews, R.A., Chadwick, D.R., Retter, A.L., Blackwell, M.S.A., Yamulki, S. 2010. Nitrous oxide emissions from small-scale farmland features of UK livestock farming systems. *Agriculture, Ecosystems and Environment* 136, 192-198.
- McGeough, K.L., Watson, C.J., Müller, C., Laughlin, R.J. and Chadwick, D.R. (2016). Evidence that the efficacy of the nitrification inhibitor dicyandiamide (DCD) is affected by soil properties in UK soils. *Soil Biology and Biochemistry* (accepted).
- McMillan, A.M.S., Pal, P., Phillips, R.L., Palmada, T., Berben, P., Jha, N., Saggar, S., Luo, J. 2015. Can pH amendments in grazed pastures help reduce N<sub>2</sub>O emissions from denitrification? – The effects of liming and urine addition on the completion of denitrification in fluvial and volcanic soils. *Soil Biology and Biochemistry*,

- Miller, L.A., Moorby, J.M., Davies, D.R., Humphreys, M.O., Scollan, N.D., MacRae, J.C., Theodorou, M.K. 2001. Increased concentration of water-soluble carbohydrate in perennial ryegrass (*Lolium perenne* L.): milk production from late-lactation dairy cows. *Grass and Forage Science* 56, 383-394.
- Min, B.R., Barry, T.N., Attwood, G.T., McNabb, W.C. 2003. The effect of condensed tannins on the nutrition and health of ruminants fed fresh temperate forages: a review. *Animal Feed Science and Technology*. 106, 3-19.
- Misselbrook, T.H., Powell, J.M., Broderick, G.A., Grabber, J.H. 2005. Dietary manipulation in dairy cattle: laboratory experiments to assess the influence on ammonia emissions. *Journal of Dairy Science* 88, 1765-1777.
- Müller, C., Rütting, T., Kattge, J., Laughlin, R.J., Stevens, R.J. 2007. Estimation of parameters in complex <sup>15</sup>N tracing models by Monte Carlo sampling. *Soil Biology and Biochemistry* 39, 715-726.
- Müller, C., Stevens, R.J., Laughlin, R.J., 2004. A <sup>15</sup>N tracing model to analyse N transformations in old grassland soil. *Soil Biology and Biochemistry* 36, 619-632.
- Muñoz, C., Quilodrán, C., Navia, R. 2014. Evaluation of biochar-plant-extracts complexes on soil nitrogen dynamics. *Journal of Biobased Materials and Bioenergy* 8, 377-385.
- Musat, N., Halm, H., Winterholler, B., Hoppe, P., Peduzzi, S., Hillion, F., Horreard, F., Amann, R., Jorgensen, B.B., Kuypers, M.M.M. 2008. A single-cell view on the ecophysiology of anaerobic phototrophic bacteria. *Proceedings of the National Academy of Sciences*, 105, 17861-17866.
- Pacheco, D., Lowe, K., Hickey, M.J., Burke, J.L., Cosgrove, G.P. 2010. Seasonal and dietary effects on the concentration of urinary N from grazing dairy cows. In: Edwards, G.R.,

- Bryant, R.H. (Eds.) Proceedings of the 4th Australasian Dairy Science Symposium, Caxton Press, Lincoln University, New Zealand, pp. 68–73.
- Petersen, S.O., Simek, M., Stamatiadis, S., Yamulki, S. 2004. Nitrous oxide emissions from grazed grassland: effects of cattle management and soil conditions. In: Weiske, Achim (Ed.) Greenhouse Gas Emissions from Agriculture. Mitigation options and Strategies, Institute for Energy and Environment. Leipzig, Germany, pp. 75-78.
- Pleasants, A.B., Shorten, P.R., Wake, G.C. 2007. The distribution of urine deposited on a pasture from grazing animals. *The Journal of Agricultural Science* 145, 81-86.
- Popp, A., Lotze-Campen, H., Bodirsky B. 2010. Food consumption, diet shifts and associated non-CO<sub>2</sub> greenhouse gases from agricultural production. *Global Environmental Change* 20, 451-462.
- Reyes-Escobar, J., Zagal, E., Sandoval, M., Navia, R., Muñoz, C. 2015. Development of a biochar-plant-extract-based nitrification inhibitor and its application in field conditions. *Sustainability* 7, 13585-13596.
- Richardson, D., Felgate, H., Watmough, N., Thomson, A., Baggs, E. 2009. Mitigating release of the potent greenhouse gas N<sub>2</sub>O from the nitrogen cycle – could enzymic regulation hold the key? *Trends in Biotechnology* 27, 388-397.
- Saggar, S., Giltrap, D.L., Davison, R., Gibson, R., de Klein, C.A.M., Rollo, M., Ettema, P., Rys, G. 2015. Estimating direct N<sub>2</sub>O emissions from sheep, beef and deer grazed pastures in New Zealand hill country: accounting for the effect of land slope on the N<sub>2</sub>O emission factors from urine and dung. *Agriculture, Ecosystems and Environment* 205, 70-78.
- Saggar, S., Giltrap, D.L., Li, C., Tate, K.R. 2007. Modelling nitrous oxide emissions from grazed grassland in New Zealand. *Agriculture, Ecosystems and Environment* 119, 205-216.

- Saggar, S., Jha, N., Deslippe, J., Bolan, N.S., Luo, J., Giltrap, D.L., Kim, D.-G., Zaman, M., Tillman, R.W. 2013. Denitrification and N<sub>2</sub>O:N<sub>2</sub> production in temperate grasslands: Processes, measurements, modelling and mitigating negative impacts. *Science of the Total Environment* 465, 173-1795.
- Schils, R.L.M., Eriksen, J., Ledgard, S.F., Vellinga, Th.V., Kuikman, P.J., Luo, J., Petersen, S.O., Velthof, G.L. 2013. Strategies to mitigate nitrous oxide emissions from herbivore production systems. *Animal* 7, 29-40.
- Schmidt, H., Eickhorst, T. 2014. Detection and quantification of native microbial populations on soil-grown rice roots by catalyzed reporter deposition-fluorescence in situ hybridization. *FEMS Microbiology Ecology* 87, 390-402.
- Schmidt, H., Eickhorst, J. 2013. Spatio-temporal variability of microbial abundance and community structure in the puddled layer of a paddy soil cultivated with wetland rice (*Oryza sativa* L.). *Applied Soil Ecology* 72, 93-102.
- Schmidt, H., Eickhorst, T., Mussmann, M. 2012. Gold-FISH: a new approach to the in situ detection of single microbial cells combining fluorescence and scanning electron microscopy. *Systematic and Applied Microbiology* 35, 518-525.
- Schoeneberger, M., Gary B., de Gooijer, H., Soolanayakanahally, R., Sauer, T., Brandle, J., Zhou, X., Current, D. 2012. Branching out: agroforestry as a climate change mitigation and adaptation tool for agriculture. *Journal of Soil and Water Conservation* 67, 128A-136A.
- Selbie, D.R., Buckthought, L.E., Shepherd, M.A. 2015. The challenge of the urine patch for managing nitrogen in grazed pasture systems. *Advances in Agronomy* 129, 229-292.
- Shaw, R., Williams, A.P., Miller, A., Jones, D.L. 2013. Assessing the potential for ion selective electrodes and dual wavelength UV spectroscopy as a rapid on-farm measurement of soil nitrate concentration. *Agriculture* 3, 327-341.

- Shepherd, M., Chambers, B. 2007. Managing nitrogen on the farm: the devil's in the detail. *Journal of the Science of Food and Agriculture* 87, 558-568.
- Skiba, U., Jones, S.K., Dragostis, U., Drewer, J., Fowler, D., Rees, R.M., Pappa, V.A., Cardenas, L., Chadwick, D., Yamulki, S., Manning, A.J. 2012. UK emissions of the greenhouse gas nitrous oxide. *Philosophical Transactions of the Royal Society B: Biological Sciences* 367, 1175-1185.
- Smith, K. (2010). *Nitrous Oxide and Climate Change*. London, Earthscan.
- Stehfest, E., Bouwman, L., van Vuuren, D.P., den Elzen, M.G.J., Eickhout, P.K. 2009. Climate benefits of a changing diet. *Climatic Change* 95, 83-102.
- Subbarao, G.V., Ishikawa, O., Ito, K., Nakahara, K., Wang, H.Y., Berry, W.L. 2006. A bioluminescence assay to detect nitrification inhibitors released from plant roots: a case study with *Brachiaria humidicola*. *Plant and Soil* 288, 101-112.
- Subbarao, G.V., Nakahara, G.V., Hurtado, M.D.P., Ono, H., Moreta, D.E., Salcedo, A.F., Yoshihashi, A.T., Ishikawa, T., Ishitani, M., Ohnishi-Kameyama, M., Yoshida, M., Rondon, M., Rao, I.M., Lascano, C.E., Berry, W.L., Ito, O. 2009. Evidence for biological nitrification inhibition in *Brachiaria* pastures. *Proceedings of the National Academy of Sciences* 106, 17302-17307.
- Subbarao, G.V., Rondon, M., Ito, O., Ishikawa, T., Rao, I.M., Nakahara, K., Lascano, C., Berry, W.L. 2007. Biological nitrification inhibition (BNI) - is it a widespread phenomenon? *Plant and Soil* 294, 5-18.
- Sullivan, M.J., Gates, A.J., Appia-Ayme, C., Rowley, G., Richardson, D.J. 2013. Copper control of bacterial nitrous oxide emission and its impact on Vitamin B12-dependent metabolism. *Proceedings of the National Academy of Sciences* 110, 19926-19931.
- Taghizadeh-Toosi, A., Clough, T.J., Condon, L.M., Sherlock, R.R., Anderson, C.R., Craigie, R.A. 2011. Biochar incorporation into pasture soil suppresses in situ nitrous oxide

- emissions from ruminant urine patches. *Atmospheric Pollutants and Trace Gases* 40, 468-476.
- Taghizadeh-Toosi, A., Clough, T.J., Sherlock, R.R., Condon, L.M. 2011. A wood based biochar captures  $\text{NH}_3\text{-N}$  generated from ruminant urine-N, retaining its bioavailability. *Plant and Soil* 353, 73-84.
- Thomson, A.J., Giannopoulos, G., Pretty, J., Baggs, E.M., Richardson, D.R. 2012. Biological sources and sinks of nitrous oxide and strategies to mitigate emissions. *Philosophical Transactions of the Royal Society of London B: Biological Sciences* 367, 1157-1168.
- Thornton, P. K. 2010. Livestock production: recent trends, future prospects. *Philosophical Transactions of the Royal Society of London B: Biological Sciences* 365, 2853-2867.
- Ussiri, D., Lal, R. 2012. Soil emission of nitrous oxide and its mitigation. Springer Science and Business Media.
- van der Weerden, T.J., Kelliher, F.M., de Klein, C.A.M. 2012. Influence of pore size distribution and soil water content on nitrous oxide emissions. *Soil Research* 50, 125-135.
- Westhoek, H., Lesschen, J.P., Rood, T., Wagner, S., De Marco, A., Murphy-Bokern, D., Leip, A., van Grinsven, H., Sutton, M.A., Oenema, O. 2014. Food choices, health and the environment: Effects of cutting Europe's meat and dairy intake. *Global Environmental Change* 26, 196-205.
- Zaman, M., Nguyen, M.L. 2010. Effect of lime and zeolite on  $\text{N}_2\text{O}$  and  $\text{N}_2$  emissions from a pastoral soil treated with urine or nitrate-N fertilizer under field conditions. *Agriculture, Ecosystems and Environment* 136, 254-261.
- Zaman, M., Nguyen, M.L., Matheson, F., Blennerhasset, J.D., Quin, B.F. 2007. Can soil amendments (zeolite or lime) shift the balance between nitrous oxide and dinitrogen

- emissions from pasture and wetland soils receiving urine or urea-N? *Soil Research* 45, 543-553.
- Zaman, M., Nguyen, M.L., Saggar, S. 2008. N<sub>2</sub>O and N<sub>2</sub> emissions from pasture and wetland soils with and without amendments of nitrate, lime and zeolite under laboratory conditions. *Soil Research* 46, 526-534.
- Zaman, M., Saggar, S., Blennerhasset, J.D., Singh, J. 2009. Effect of urease and nitrification inhibitors on N transformation, gaseous emissions of ammonia and nitrous oxide, pasture yield and N uptake in grazed pasture system. *Soil Biology and Biochemistry* 41, 1270-1280.
- Zeng, H., Di, T., Zhu, Y., Subbarao, G.V. 2015. Transcriptional response of plasma membrane H<sup>+</sup>-ATPase genes to ammonium nutrition and its functional link to the release of biological nitrification inhibitors from sorghum roots. *Plant and Soil* 398, 301-312.
- Zerulla, W., Barth, T., Dressel, J., Erhardt, K., von Locquenghien, K.H., Pasda, G., Rädle, M., Wissemeier, A. 2001. 3,4-Dimethylpyrazole phosphate (DMPP) – a new nitrification inhibitor for agriculture and horticulture. *Biology and Fertility of Soils* 34, 79-84.

---

## Appendix 1


---

### Supplementary material for Chapter 3




**Figure 1** Eutric Cambisol mesocosms maintained at 50% and 70% WFPS.





**Figure 2** Changes in soil pH (panels a and b), electrical conductivity (EC; panels c and d) and oxidation reduction potential (ORP; panels e and f) following sheep urine application to a Eutric Cambisol, maintained at either 50% or 70% water-filled pore space (WFPS), and sampled at 9-12 cm (panels a, c and e) and 15-18 cm (panels b, d and f) away from the direct area of application. Symbols represent means  $\pm$  SEM ( $n = 4$ ). Figure legend applies to all panels and text on the top row of panels applies to each respective column.




**Figure 3** Extractable total dissolved nitrogen (panels a and b), ammonium (panels c and d), nitrate (panels e and f), nitrous oxide emissions (panels g and h) and total dissolved organic carbon (panels i and j) following sheep urine application to a Eutric Cambisol, maintained at either 50% or 70% water-filled pore space (WFPS), and sampled at 9-12 cm (panels a, c, e, g and i) and 15-18 cm (panels b, d, f, h and j) away from the direct area of application.

Symbols represent means  $\pm$  SEM ( $n = 4$ ). Figure legend applies to all panels and text on the top row of panels applies to each respective column.

## Appendix 2

### Supplementary material for Chapter 4



Key:



Automated chamber



Manual chamber



Duplicated soil sampling plots



Low N concentration large urine patch ( $4 \text{ g N l}^{-1}$ ;  $1 \text{ g N total}$ ;  $200 \text{ kg N ha}^{-1}$ )



High N concentration large urine patch ( $16 \text{ g N l}^{-1}$ ;  $4 \text{ g N total}$ ;  $800 \text{ kg N ha}^{-1}$ ;  $500 \text{ cm}^2$ )



Four low N concentration small urine patches ( $4 \text{ g N l}^{-1}$ ;  $4 \times 0.25 \text{ g of N total}$ , applied in  $4 \times 62.5 \text{ ml}$ , covering  $125 \text{ cm}^2$  each;  $200 \text{ kg N ha}^{-1}$ )



Four high N concentration small urine patches ( $16 \text{ g N l}^{-1}$ ;  $4 \times 1 \text{ g of N total}$ , applied in  $4 \times 62.5 \text{ ml}$ , covering  $125 \text{ cm}^2$  each;  $800 \text{ kg N ha}^{-1}$ )

**Figure 1** Plot layout for the field trial conducted in Chapter 4.

---

## Appendix 3

---

### **Details of an attempted study using lux marked *Nitrosomonas europaea***

This study utilised a lux marked strain of the nitrifying bacterium *Nitrosomonas europaea* ATCC19718 (pHLUX20). Unfortunately, some problems occurred during experimentation, and the long culture time of this microorganism made it difficult to troubleshoot problems with the bio assay. Due to time constraints, it was decided to discontinue the study. A summary of the main objectives and some photographs of culturing this microorganism are shown below (Fig 1).

The objectives of this experiment were to test whether any compounds commonly excreted in the urine of ruminants, are able to inhibit nitrification. If any compounds within urine do inhibit nitrification, then it may be possible to increase excreted inhibitor concentrations through diet. Excretion of the inhibitor would then theoretically be deposited in the exact location it is required – which may save money/labour costs in comparison to e.g. blanket applications of nitrification inhibitors.

The following objectives were planned:

- 1) Monitor growth of *Nitrosomonas* culture by measuring OD<sub>600</sub>
- 2) Confirm that inhibition of fluorescence corresponds to inhibition of nitrification (i.e. % nitrite production in assay)
- 3) Screen the following urine compounds for their inhibitory effect on nitrification: allantoin, uric acid, creatinine, creatine, hippuric acid, benzoic acid and compare with chemical nitrification inhibitors e.g. DCD, DMPP and the standard Allylthiourea

- 4) Narrow down compounds which have an inhibitory effect, and then test the mode of action of inhibition i.e. test whether the ammonium oxidation (AMO) or the hydroxylamine pathway (HAO) pathway is interfered with.
- 5) Test inhibition of nitrification in soils in the lab, determine the inhibitory potential of a concentration range of the nitrification inhibitors, a concentration range within artificial urine and finally by spiking urine with increased amounts of inhibitor.
- 6) Test the potential for N<sub>2</sub>O emission reduction in soils – pot trial.



**Figure 1** Culture of a lux marked strain of the nitrifying bacterium *Nitrosomonas europaea* ATCC19718 (pHLUX20).