



Sensitivity of peatland litter decomposition to changes in temperature and rainfall

Bell, M.; Ritson, J.P.; Verhoef, A.; Brazier, R.E.; Templeton, M.R.; Graham, N.; Freeman, Christopher; Clark, J.M.

Geoderma

DOI:

[10.1016/j.geoderma.2018.06.002](https://doi.org/10.1016/j.geoderma.2018.06.002)

Published: 01/12/2018

Peer reviewed version

[Cyswllt i'r cyhoeddiad / Link to publication](#)

Dyfyniad o'r fersiwn a gyhoeddwyd / Citation for published version (APA):

Bell, M., Ritson, J. P., Verhoef, A., Brazier, R. E., Templeton, M. R., Graham, N., Freeman, C., & Clark, J. M. (2018). Sensitivity of peatland litter decomposition to changes in temperature and rainfall. *Geoderma*, 331, 29-37. <https://doi.org/10.1016/j.geoderma.2018.06.002>

Hawliau Cyffredinol / General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Sensitivity of peatland litter decomposition to changes in temperature and rainfall

AUTHORS AND AFFILIATIONS

Michael C. Bell^{a,*}, Jonathan P. Ritson^{b,c}, Anne Verhoef^a, Richard E. Brazier^d, Michael R. Templeton^c, Nigel J.D. Graham^c, Chris Freeman^e, Joanna M. Clark^a

^a Soil Research Centre, Department of Geography and Environmental Science, School of Archaeology, Geography and Environmental Science, The University of Reading, Whiteknights, PO Box 227, Reading RG6 6AB, UK

^b Grantham Institute for Climate Change, Imperial College London, South Kensington, London SW7 2AZ, UK

^c Department of Civil and Environmental Engineering, Imperial College London, South Kensington, London SW7 2AZ, UK

^d Geography, College of Life and Environmental Sciences, University of Exeter, EX4 4RJ, UK

^e Wolfson Carbon Capture Laboratory, School of Biological Sciences, Bangor University, Bangor, Gwynedd LL57 2UW, UK

*Corresponding author: Soil Research Centre, Department of Geography and Environmental Science, School of Archaeology, Geography and Environmental Science, The University of Reading, Whiteknights, PO Box 227, Reading RG6 6AB, UK. E-mail address: m.c.bell@pgr.reading.ac.uk

Abstract

Changes to climate are projected over the next 50 years for many peatland areas. As decomposition of peat-forming vegetation is likely to be intrinsically linked to these changes in climate, a clear understanding of climate-peat dynamics is required. There is concern that increased temperature and decreased precipitation could increase the rate of decomposition and put the carbon sink status of many peatlands at risk, yet few studies have examined the impact of both climatic factors together. To better understand the sensitivity of peatland decomposition to changes in both temperature and precipitation and their interaction, we conducted a short-term laboratory experiment in which plant litters and peat soil were incubated, in isolation, in a factorial design. Treatments simulated baseline and projected climate averages derived from the latest UK climate change projections (UKCP09) for Exmoor, a climatically marginal peatland in SW England. Regular carbon dioxide flux measurements were made throughout the simulation, as well as total mass loss and total dissolved organic carbon (DOC) leached. The largest effect on carbon loss in this multifactor experiment was from substrate, with *Sphagnum*/peat releasing significantly less C in total during the experiment than dwarf shrubs/graminoids. Climate effects were substrate specific, with the drier rainfall treatment increasing the DOC leaching from *Calluna*, but decreasing it from *Sphagnum*. Partitioning between CO₂ and DOC was also affected by climate, but only for the peat and *Sphagnum* samples, where the future climate scenarios (warmer and drier) resulted in a greater proportion of C lost in gaseous form. These results suggest that indirect effects of climate through changes in species composition in peatlands could ultimately turn out to be more important for litter decomposition than direct effects of climate change from increased temperatures and decreased rainfall.

Keywords: decomposition; Exmoor; blanket peat; uplands; DOC; UK

1. Introduction

Northern peatlands are an important carbon store, holding around one third of the global soil carbon stock (Gorham, 1991). For peatlands to accumulate organic matter, and thus sequester carbon from the atmosphere, the overall loss of carbon from the system through the combined decomposition of plant litter and peat must be lower than C input via litter production and vascular plant root exudation (Frolking et al., 2010; Limpens et al., 2008). Decomposition is the breakdown of organic matter. It can be due to one or a number of physical, chemical or biological processes. Carbon dioxide (CO₂), dissolved organic carbon (DOC), nutrients and stable hummus are among the principal final products of decomposition (Bragazza et al., 2009). Carbon is primarily lost as CO₂ in peatlands (Billett et al., 2010). However, additional losses in the form of methane (CH₄), produced during anaerobic decomposition, and DOC to aquatic systems can be an important component of the peatland carbon balance in some areas (Köehler et al., 2011). The speed and final product of the decomposition process is dependent on the chemistry of the organic matter, and the environment in which it decomposes (Limpens et al., 2008).

Understanding how peat and plant litter decomposition could be affected by climate change, specifically changes in temperature and rainfall, is key to an improved understanding of how the peatland carbon source/sink status could change during the 21st century. Northern peatlands are typically dominated by remains of mosses of the genus *Sphagnum*, a bryophyte (Turetsky, 2003). *Sphagnum* has played an important role in peatlands becoming such a large carbon sink. Not only does the recalcitrance of its litter mean that it keeps C in the system for longer, but the presence of *Sphagnum* maintains acidic, nutrient poor conditions which make it difficult for plants with more labile litter to establish themselves (van Breemen, 1995).

Climate change may alter vegetation types in peatlands with vascular plants becoming more dominant (Buttler et al., 2015; Dieleman et al., 2015; Fenner et al., 2007; Weltzin et al., 2003). However, many ongoing catchment management programmes aim to encourage *Sphagnum* species over vascular plants as part of peatland restoration programmes (Grand-Clement et al., 2013).

While some authors have studied the effects of climate variables on the treatability of water from different peatland plant species (e.g. Ritson et al., 2014; Tang et al., 2013), little work has so far been conducted on the total flux of carbon and the partitioning between gaseous and fluvial losses from different peatland plant species. This partitioning is an important parameter to include in carbon cycle models, particularly for climate change modelling as there is likely to be a lag between DOC release from the soil and its incorporation into the atmospheric pool, as in-stream processing can lead to the temporary storage of C within the aquatic system (Hope et al., 1994). An understanding of partitioning of carbon losses between aquatic and gaseous fractions from different vegetation sources is also important for predicting whether catchment management programmes aimed at restoring certain species have the desired holistic effects of improving drinking water quality (including enhanced C sequestration) and do not result in environmental problem-shifting.

Decomposition of litters in the field is traditionally measured using the litter bag technique (Bragazza et al., 2009; Wieder and Lang, 1982). Litter of a known mass is enclosed in mesh bags with openings large enough for decomposers to access the food source, but small enough to prevent the physical loss of litters. Samples are incubated in the field, either on, or just below the soil surface (Johnson and Damman, 1991; Moore et al., 2007) and decomposition rate is quantified via mass loss or changing nutrient quotients of the litter (Keuskamp et al., 2013). *In situ* litter decomposition studies, using this technique, have

suggested multiple drivers for increased decomposition rates in peatland systems including: elevated nutrient additions (Bubier et al., 2007), water-table drawdown (Straková et al., 2012), litter quality (Limpens and Berendse, 2003) and temperature (Moore et al., 2007). However, these field-based studies are limited to a single measure of decomposition; mass loss. Mass loss gives a valuable measure of net decomposition, which can be useful for comparing between litters, but does not provide information on the final product of the decomposition process. Carbon dioxide and DOC not only represent different flux pathways (gaseous versus aquatic), but can also be indicative of different processes and stages of decomposition. CO₂ is in effect representative of fully utilised carbon, whereas DOC could be stored in the peat column or utilised by microbes and exported to the atmosphere via respiration (Pastor et al., 2003; Turetsky, 2003).

Laboratory incubations of peatland litters to examine the relationship between decomposition and climate have been limited to date. Studies that have been conducted have tended to concentrate on a single measure of decomposition, have focussed solely on peat rather than vegetation, or have just looked at one environmental variable, such as temperature (e.g. Neff and Hooper 2002; Moore et al. 2008) or water table (e.g. Freeman et al. 1993), in isolation. The effect of interactions between changes in temperature and rainfall on the decomposition of peatland litter is not currently well understood (Breeuwer et al., 2008). A gradient based mesocosm study measured increased C soil respiration in mesocosms transplanted to a warmer, drier location (Bragazza et al., 2016). However, isolating the effects of climatic variables such as temperature and rainfall from other confounding variables can be difficult in the field as these factors naturally co-vary. It is also difficult to separate the effects on different litters and measure the relative decomposition from different sources in the field or in intact mesocosms. To address this, previous studies have incubated soils or

litters from wetland sites in isolation, notably Moore and Dalva (2001), Neff and Hooper (2002) and Wickland et al. (2007). This was the approach taken here.

A better understanding of the sensitivity of different peatland litters to changes in temperature and rainfall regime is needed to enable predictions of the impact of climate change on the stability of peatland carbon stocks, and to support better parameterisations of models that simulate past, current and future climate (Frolking et al., 2010; Smith et al., 2010). Therefore, the objectives of this research were as follows: (1) assess how litter decomposition rates varied between common ombrotrophic peatland plant species (*Calluna vulgaris*, *Molinia caerulea*, *Sphagnum* moss, and mixed litter) compared to peat; (2) determine the relative importance (or partitioning) of gaseous (i.e. CO₂) versus aquatic (DOC) fluxes during decomposition with respect to plant species and peat; (3) evaluate the importance of simulated temperature and rainfall changes on controlling total decomposition, C fluxes and partitioning between CO₂ and DOC. This final objective links to another study (Ritson et al., 2014), which used the same experimental treatments to examine the impact of climate change on the treatability (for potable water supply) of dissolved organic matter.

2. Methods

2.1. *Field site and sample collection*

In the south-west UK, blanket peat covers large parts of the upland areas and constitute an important store of C (Parry and Charman, 2013). These blanket peatlands represent the southern-most blanket peat in the UK and are considered climatically marginal (Clark et al., 2010). The shallow peat of Exmoor was heavily damaged by intensive drainage during the 19th and 20th centuries, and recent efforts have been made to restore large areas

through a programme of ditch blocking (Grand-Clement et al., 2014, 2013). The field sites are dominated by *Molinia caerulea* (purple moor grass) (Gatis et al., 2015) and are classified as UK National Vegetation Classification M25: *Molinia caerulea* - *Polentilla erecta* mires (Rodwell, 1991). Paleoecological studies indicate that purple moor grass expansion corresponds with the industrial revolution and prior to this, *Sphagnum* was more dominant species within this region (Chambers et al., 1999). Exmoor typically receives precipitation of around 1800–2600 mm yr⁻¹ with mean winter and summer temperatures of between 4.5–5.5 °C and 10–12 °C, respectively (Met Office, 2014). Exmoor receives a relatively high input of nutrients through atmospheric deposition, and the mean deposition of total nitrogen and oxidised sulphur for the period 2008–2012 was estimated to be 15.5–16.16 kg N h⁻¹ yr⁻¹ and 4.41–4.86 kg S h⁻¹ yr⁻¹, respectively (DEFRA, 2015).

Samples of vegetation and peat were collected from two catchments, Aclands (51° 07'54.2" N 3° 48'43.3" W) and Spooners (51° 07'23.3" N 3° 45'11.8" W), within Exmoor National Park, UK, during July 2013. A further description of these field sites, including maps, can be found in Grand-Clement et al. (2014) and Luscombe et al. (2015). Five different substrates were collected: *Calluna*, *Molina* and *Sphagnum*, mixed litter and peat. Fresh leaves and branches were collected from *Calluna vulgaris* and *Molinia caerulea*, avoiding any plants which had been partially eaten by herbivores. Mixed litter (predominantly *Molinia caerulea* but also some *Eriophorum vaginatum*, both at the early stages of decomposition) was collected from the bog surface. Intact *Sphagnum* spp. sods (entire plants) were collected from the centre of stands and peat was collected from the top ca. 30 cm of the soil profile using a screw auger. The samples were transported back to the laboratory in cool boxes within 8 hours of collection, and stored at 4 °C prior to their preparation. Sample collection is described fully in Ritson et al. (2014).

Table 1. Organic matter properties for peatland vegetation and peat classes used in the experiment. Values are means ($n = 5$) of subsample values, numbers in brackets are the standard error of the mean. Aromaticity is a qualitative measure and was calculated from cross polarization-magic angle spinning nuclear magnetic resonance (CP-MAS NMR).

Class	Dominant substrate	C (%)	N (%)	C:N ratio	Moisture content (g g dry wt ⁻¹)	Aromaticity (%)
<i>Calluna</i>	<i>Calluna vulgaris</i>	49.3 (0.02)	1.35 (0.01)	36.4 (0.22)	1.90 (0.01)	8
Mixed litter	Senesced <i>Molinia c</i>	45.9 (0.01)	1.29 (0.01)	35.6 (0.28)	5.38 (0.11)	5
<i>Molinia</i>	<i>Molinia caerulea</i>	45.5 (0.02)	2.34 (0.02)	19.5 (0.13)	3.56 (0.05)	5
Peat	Peat	29.1 (0.02)	1.64 (0.01)	17.7 (0.07)	6.06 (0.06)	11
<i>Sphagnum</i>	<i>Sphagnum</i> spp.	41.8 (0.03)	1.00 (0.01)	42.1 (0.28)	15.020.15)	7

2.2. Sample preparation

Samples from each vegetation class (*Calluna vulgaris*, *Molinia caerulea*, mixed litter and *Sphagnum* spp.), were cut into 2-3 cm sections and mixed thoroughly within their class. Any material not from the target group was removed. Once sorted, cut and homogenised, the vegetation samples were left to air-dry in the laboratory to constant mass (Moore et al., 2007). The structure of peat is permanently altered when dried, becoming hydrophobic (Holden and Burt, 2002). For this reason, the peat was treated differently to the other substrates and was kept refrigerated and at field moist conditions prior to the experiment.

Air-dried sub-samples of each plant species and field moist peat ($n = 5$) were oven dried at 70 °C for 24 h to determine an air-dry to oven-dry conversion factor. This conversion factor was

applied to estimate the oven-dry weight of samples at the beginning of the simulation. These oven dry samples were analysed for litter quality (see methods below; Table 1).

2.3. *Experimental design*

2.3.1. *Experimental set up*

Polypropylene Buchner funnels (Thermo Scientific, USA) were filled with approximately 2 g dry-weight of air-dry vegetation or 10 g homogenised peat. The stem of the funnel was packed with acid washed glass wool to ensure no loss of particulates, and each Buchner funnel was placed above an amber-glass bottle to collect leachate, for separate analysis (see Ritson et al., 2014). The climate control facilities comprise two versatile environmental test chambers (Panasonic MLR-352H). The incubators were set to cycle between 12 h at the mean daily maximum and 12 h at the mean daily minimum temperature. Deionised water was applied in amounts equivalent to the monthly rainfall, with a dispenser set to the required volume based on the area of the Buchner funnel top. This was applied gradually and evenly, over approximately 30 seconds, taking care not to disturb the samples.

2.3.2. *Experimental design*

The experiment was set up as a factorial ANOVA design, where substrate, temperature and rainfall were the treatment factors in a 5 x 2 x 2 design. Five replicates of each treatment were conducted, giving a total of 100 samples (see Ritson et al., 2014). Climate treatments were based on (i) baseline (1961-1990; Met Office, 2013) mean minimum and mean maximum daily temperature and mean rainfall totals for July and August for Exmoor National Park, UK, and (ii) a projection from the UKCP09 high emissions scenario for 2080s (2070-2100). All four

possible combinations of baseline and projected temperature and rainfall were applied as treatments. Mean minimum daily temperatures were set to 11.0 °C (± 0.3 °C for test chamber error) for the baseline, and 15.4 °C (± 0.3 °C) for the UKCP09 treatment. Mean maximum daily temperatures were set to 18.2 °C (± 0.3 °C) for the baseline treatment and 23.7 °C (± 0.3 °C) for the UKCP09 treatment. Rainfall totalled 100.9 and 117.5 mm for the baseline treatment for July and August, respectively, and 75.4 and 67.2 mm for the UKCP09 treatment. Overall, this was equivalent to a ca. 5.0 °C increase in temperature and ca. 35 % decrease in rainfall for the UKCP09 treatments. The relative humidity (to program the test chambers) and number of rain days were set at mean values from the baseline period for all samples (for relative humidity: 80 % in July, 79 % in August; for rain days: 12 in July and 14 in August). The rainfall total for each month was split evenly between the rain days (resulting in irrigation amounts ranging between 4.2 mm and 8.4 mm per event, or 19 and 38 ml, depending on treatment and month), and the amounts were applied every 2-4 days. The conversion from mm to ml was based on the area of the funnel.

2.4. *Measurements*

2.4.1. *Carbon dioxide evolution*

At eight points during the experiment (on days 2, 7, 10, 17, 31, 45, 50, and 60 after the start of the incubation) samples were analysed for CO₂ efflux. Samples were taken out of the incubators individually and the top part of the Buchner funnel, containing the sample, was removed and placed into an airtight container (Lock&Lock, South Korea) which was connected to an infra-red gas analyser, IRGA (8100A, LI-COR, USA). Samples were removed for measurement in a random order to avoid the introduction of systematic measurement error.

Observations with the IRGA lasted for 2 minutes during which time the concentration of CO₂ in the chamber (water vapour compensated) was measured every second, and the samples were out of the incubators for a maximum of 5 minutes. The rate of change in CO₂ concentration in the headspace during the 2-minute measurement period was used to calculate the instantaneous flux by applying a linear regression. The IRGA system was allowed to purge for a minute between samples, and a 30 second deadband (a period where the IRGA is running without the measurements being used in the flux estimate) was introduced at the beginning of each measurement to allow for adequate mixing of air in the headspace. The CO₂ time series for each sample was analysed separately, and the deadband was adjusted where necessary. All flux estimates were derived from a minimum of 60 consecutive CO₂ concentration measurements (i.e. 60 seconds). Individual flux estimates were converted to mg CO₂-C g C⁻¹ day⁻¹. Cumulative CO₂ flux over the course of the two month experiment was calculated for each sample based on a linear interpolation of fluxes between the successive measurements (Neff and Hooper, 2002), expressed as mg CO₂-C g C⁻¹.

2.4.2. *Dissolved organic carbon*

DOC analysis was performed on the cumulative leachate collected at the end of the simulation period. Samples were filtered through a GF/F filter (Whatman, UK) with a nominal pore size of 0.7 µm, acidified to ~pH 2 with HCl and stored at 4°C prior to analysis. Organic carbon content was measured as non-purgeable organic carbon (NPOC) using a total organic carbon analyser (Shimadzu TOC-V, Japan). Calibration and quality control checks were run daily from freshly prepared standards and samples were analysed within one week of the end of the simulation. Organic carbon concentration (mg L⁻¹) was converted to DOC leached (mg) based on the volume of leachate collected. As temperature and rainfall treatment affected

the volume of sample collected, this was then reported as mg DOC-C g C⁻¹, based on the starting dry mass and initial C content of each sample.

2.4.3. *Mass loss*

At the end of the two-month incubation period, samples were removed from the incubators, dried at 70 °C for 48 h and weighed. C loss was estimated based on the starting mass and starting C content, as well as the end mass and end C content. This was expressed as mgC g C⁻¹.

2.5. *Litter quality analysis*

The oven-dry subsamples from the beginning, and all samples from the end of the experiment were ground using a laboratory disc mill with an agate barrel (Tema Machinery Ltd, UK) and analysed for total C and N (Thermo FLASH 2000 elemental analyser). The handling and analysis of samples were undertaken together with reference material (Pahokee Peat, International Humic Substances Society) in order to verify the calibration and quality of the results. Subsamples from the start of the experiment were also analysed for C character using nuclear magnetic resonance (NMR) analysis. Cross polarisation-magic angle spinning (CP-MAS) ¹³C NMR analysis was undertaken at the University of Reading chemical analysis facility (CAF) on a Bruker AV500 instrument based on the method of Wong et al. (2002). This technique is used widely in environmental studies and is good for looking at relative changes in carbon distribution across a series of samples (Simpson et al., 2011). For the vegetation samples 1024 transients (executions of the pulse sequence) were taken, however due to the presence of paramagnetic material in the peat soil transients were increased to 4096 to

improve the signal to noise ratio. Magic angle spinning (MAS) spin rate was set at 10 kHz to shift spinning side bands out of the region of interest and all signals were externally referenced to adamant. The aromatic peak between 110-140 ppm was integrated and its percentage of the total signal calculated to give an indication of the differences between relative aromaticity between samples (Peuravuori et al., 2003), which can be used as an alternative qualitative measure of the recalcitrance of the litter (see Table 1). For further reading on using NMR analysis for environmental research, see Simpson et al. (2011).

2.6. *Statistical analysis*

All statistical analyses were carried out using R version 3.0.2 (R Core Team, 2015) and plots were generated using the ggplot2 package (Wickham, 2009). The experiment involved a three-way analysis of variance (ANOVA) design where temperature, rainfall and substrate were the experimental factors. The effect of these factors, and their interaction, on response variables of CO₂ flux, mass loss, DOC leached and CO₂:DOC ratio production was analysed (Table 2). A Fligner-Killeen test was performed to determine that there was equal variance between groups, and residuals were checked for normality. Carbon dioxide flux, DOC flux and CO₂:DOC ratio were all square-root transformed prior to inclusion, in order to meet the assumption of normally distributed residuals. A critical value of $P = 0.05$ was used for statistical significance. For multiple comparisons, the Tukey honest significant differences (HSD) test was used. The variance explained by each factor was estimated using omega-squared (ω^2), a test which is suitable for estimating effect size from small sample sizes (Keselman, 1975).

3. Results

3.1. Variability of decomposition rates between different substrates

Decomposition was greatest for the *Molinia* samples and slowest for the peat samples (Fig. 1). The same substrate rankings were evident in terms of the following: the estimated cumulative CO₂ flux (means ranged from 942.87 mgC gC⁻¹ for *Molinia* to 41.26 mgC gC⁻¹ for peat; Fig. 1); the DOC flux (means ranged from 20.95 mgC gC⁻¹ for *Molinia* to 1.07 mgC gC⁻¹ for peat; Fig. 1); and the overall total measured mass loss (means ranged from 511.21 mgC gC⁻¹ for *Molinia* to 27.43 mgC gC⁻¹ for peat; Fig. 2). Substrate was a significant explanatory factor in the ANOVA for all of these response variables ($P < 0.001$; Table 2). Post-hoc tests suggested significant differences in the order *Molina* > *Sphagnum* = Mixed litter > *Calluna* > peat for CO₂ flux; *Molinia* > *Calluna* = Mixed litter > *Sphagnum* > peat for DOC flux; and *Molinia* > *Calluna* > *Sphagnum* = Mixed litter > peat for mass loss.

There was a steady decrease in CO₂ release over the entire incubation period, for all substrates and treatments. The decrease in CO₂ flux was greatest for the *Molinia* samples, with a reduction in estimated flux between the first (day 1 to 30) and second half (day 31 to 60) of the experiment of 64.27%, compared to peat where the difference was 25.13% – the lowest decline recorded.

In terms of total mass balance, the overall loss of C measured directly through mass loss of the substrate and by the total fluxes (DOC + CO₂) were not equal. Whilst DOC fluxes were less than the total mass loss, the cumulative CO₂ flux estimated by integration across the experimental period (*sensu* Neff and Hooper, 2002) was greater than the total mass loss.

Table 2. ANOVA model results, where the model fitted is $response = Temperature * Rainfall * Substrate$ (Cumulative CO₂, DOC and CO₂:DOC ratio are square-root transformed data).

Significant differences ($P < 0.05$) are highlighted in bold and effect sizes (ω^2) are stated in brackets. “Temp”, “Rain” and “Sub” are short for Temperature, Rainfall and Substrate, respectively).

Variable	Factor	F	df	P value
Cumulative CO ₂ (mgCO ₂ -C gC ⁻¹)	Temperature	1.04	1	0.311
	Rainfall	3.25	1	0.075
	Substrate	631.08	4	< 0.001 (0.960)
	Temp:Rain	0.06	1	0.804
	Temp:Sub	1.27	4	0.288
	Rain:Sub	1.68	4	0.164
	Temp:Rain:Sub	1.27	4	0.290
	Mass loss (mgCO ₂ -C gC ⁻¹)	Temperature	20.10	1
Rainfall		0.01	1	0.927
Substrate		946.23	4	< 0.001 (0.966)
Temp:Rain		0.01	1	0.915
Temp:Sub		3.80	4	0.007 (0.003)
Rain:Sub		2.72	4	0.035 (0.002)
Temp:Rain:Sub		0.56	4	0.690
DOC (mgCO ₂ -C gC ⁻¹)		Temperature	18.19	1
	Rainfall	3.38	1	0.070
	Substrate	247.14	4	< 0.001 (0.838)
	Temp:Rain	0.04	1	0.837
	Temp:Sub	5.12	4	0.001 (0.014)
	Rain:Sub	11.34	4	< 0.001 (0.035)
	Temp:Rain:Sub	4.43	4	0.003 (0.012)
	CO ₂ -C:DOC-C	Temperature	15.57	1
Rainfall		26.16	1	< 0.001 (0.097)
Substrate		19.96	4	< 0.001 (0.292)
Temp:Rain		0.01	1	0.921
Temp:Sub		3.27	4	0.015 (0.035)
Rain:Sub		8.71	4	< 0.001 (0.119)
Temp:Rain:Sub		2.24	4	0.072

3.2. Carbon dioxide to DOC partitioning

The cumulative flux of CO₂ was greater than DOC for all of the samples (Fig. 1). Mean CO₂:DOC quotients ranged from 31.56 for *Molinia* to 68.67 for *Sphagnum* (all climate

treatments pooled), indicating that proportionally more carbon was lost as DOC from *Molinia* than *Sphagnum*. Substrate was again the strongest factor explaining the variation in partitioning between gaseous and aquatic fluxes ($P < 0.001$). There was a significant relationship between total CO₂ flux and DOC leached between groups when all data were pooled, indicating that DOC leaching increases with CO₂ flux. However, this relationship was not observed within substrate groups.

The general trend was for an increase in the CO₂:DOC quotient under UKCP09 future climate scenarios (Fig. 1). There was a statistically significant interaction between rainfall and substrate ($P < 0.001$), with the proportion of C lost as CO₂ generally increasing under the UKCP09 (drier) rainfall scenario. Post-hoc tests revealed that drier conditions were only significant in changing the partitioning between gas and aquatic carbon fluxes for the peat ($P < 0.001$) and *Sphagnum* ($P < 0.001$) samples, the mean of which increased from 24.91 (± 5.74) to 78.64 (± 13.85) and from 47.34 (± 2.41) to 101.78 (± 16.25) when comparing the baseline temperature and rainfall to the UKCP09 temperature and rainfall treatments. Partitioning between CO₂ and DOC in other substrates, *Molinia*, *Calluna* and mixed litter, was unaffected by temperature and rainfall treatments ($P > 0.05$).




Figure 1. Treatment effects on measures of carbon fluxes, and their ratio; (a) Cumulative CO₂ flux, (b) DOC flux, (c) Cumulative CO₂ to DOC ratio (Error bars indicate one standard error).

Different letters denote statistically significant differences in means between substrates and

asterisks denote significant differences between treatments for substrates based on Tukey HSD test ($P < 0.05$).

3.3. *Relative importance and interaction between temperature and rainfall in controlling decomposition*

Increasing temperature had a significant effect on decreasing DOC flux ($P < 0.001$) and increasing mass loss ($P < 0.001$). In addition, significant interactions between temperature and substrate ($P < 0.001$) and rainfall and substrate ($P = 0.001$) were seen for DOC flux and for mass loss ($P = 0.007$ and $P = 0.035$, respectively), indicating that the effect of simulated climate change depended on substrate type. For DOC, post hoc tests suggested that the interaction was only significant between temperature and *Calluna*, whereas for rainfall, there was a significant interaction between rainfall treatment and *Calluna* and rainfall treatment and *Sphagnum*. This interaction between rainfall and substrate was not unidirectional, however, with the drier rainfall treatment increasing DOC in *Calluna* samples, but decreasing it in *Sphagnum*. With mass loss, only *Sphagnum* was close to being significant on its own. Differences in cumulative CO₂ flux between climate treatments (both temperature and rainfall) were non-significant ($P > 0.05$), though rainfall was close to the critical threshold ($P = 0.075$). No significant interaction between temperature and rainfall for any of the measures of decomposition was observed (CO₂ flux, DOC flux or mass loss).




Figure 2. Treatment effects on C loss estimated from mass and C quotient measurements (Error bars indicate one standard error). Different letters denote statistically significant differences in means between substrates and asterisks denote significant differences between treatments for substrates based on Tukey HSD test ($P < 0.05$).

4. Discussion

4.1. Carbon fluxes and decomposition rates between different substrates

The fluxes of CO₂ and DOC, and the mass loss, were lowest from the peat samples compared to litter sources, consistent with previous work (Moore and Dalva, 2001; Tang et al., 2013). Decomposition in peatland ecosystems progresses along a continuum from freshly senesced plant litter to peat. Part of the reason that peat decomposes more slowly than fresh litter is because more labile compounds are preferentially utilised by decomposers (Wieder and Lang, 1982). This results in older peat being composed of progressively more recalcitrant materials (Broder et al., 2012). In the field, these differences are likely to be even greater as decomposition of peat is further inhibited by anoxic and acidic conditions which prevail for longer at greater depths. Here, we used measures of C:N ratio and aromaticity to characterise the quality of the substrates. A high C:N ratio usually suggests recalcitrant litter, as N is limiting

for microbes and can also stimulate the activity of certain enzymes (Bragazza et al., 2012). Here peat had the lowest C:N ratio, which could be due to retention of N containing compounds during the decomposition process (Kalbitz and Geyer, 2002). Aromaticity is another potential measure of decay resistance as aromatic polymers are characterised by structural and chemical recalcitrance. In this instance, peat had the highest estimated proportion of aromatic compounds (Table 1), which could in part explain its low decomposition rate relative to the other substrates.

In terms of specific fluxes, *Sphagnum* produced the least DOC in line with previous observations of high retention of carbon in this species (Fenner et al., 2004). *Sphagnum* also produced slightly more CO₂ than *Calluna*, which was unexpected given the well documented recalcitrance of *Sphagnum* litter which has been shown to decompose much more slowly than other peat forming vegetation (van Breemen, 1995). This could have been a result of environmental conditions, specifically water content, as the poor water holding capacity of *Calluna* litter meant that lack of water supply could have limited its decomposition during laboratory simulation experiments (Blok et al., 2015; Toberman et al., 2008).

The estimates of cumulative CO₂ flux were approximately double the amount estimated from C mass loss, suggesting an overestimation of cumulative CO₂ flux by integrating point measurements recorded here. This could partly be due to the measurement of some autotrophic respiration from peat roots in the earlier stages of the experiment, or an inadequate representation of night-time respiration rates. However, given that all samples were treated in the same way, and were well mixed to begin with, it is reasonable to assume that the observed differences between groups are robust.

4.2. *Relative importance of gaseous and fluvial fluxes*

The release of DOC during organic matter decomposition to surface waters is an important component of the peatland carbon budget (Billett et al., 2010). Little is known about partitioning between CO₂ and DOC fluxes in peatlands (Bonnett et al., 2006). Herein, it was shown that *Molinia* produced the most DOC as a proportion of its starting mass, and peat produced the least. However, in terms of the ratio of CO₂ to DOC, *Calluna* and mixed litter had the lowest ratios, signalling that a higher proportion of mass is lost as DOC than CO₂ for these substrates, when compared with the others. Again, absolute values for CO₂:DOC may not be accurate due to overestimation of measured CO₂ fluxes, but comparisons between groups should still be robust.

Previous studies have reported DOC fluxes equivalent to over 20% of net ecosystem exchange (NEE) (e.g. Dinsmore et al., 2010; Koehler et al., 2011). In some instances, DOC exported to streams has been estimated to exceed NEE (Billett et al., 2004), highlighting the importance of including this flux when estimating source/sink status of a peatland. Strong correlations have been found between gross primary productivity (GPP) and DOC concentrations in peatland catchments (Dinsmore et al., 2013; Harrison et al., 2008), suggesting a large amount of DOC is produced directly by living plants, indirectly by rhizosphere priming or by the decay of fresh plant litter. The difference in the amount of DOC released from the peat and litter samples in this study during decomposition supports the theory that the decay of fresh litter is a substantive DOC source (Evans et al., 2007; Palmer et al., 2001). In the field, a lot will depend on the regulating effect of water table depth on decomposition, flow pathways of runoff (Wallage and Holden, 2011), as well as substrate.

We observed a lower DOC flux but no significant change in CO₂ flux under the UKCP09 (warmer) temperature scenario. Combined with the lower DOC flux under UKCP09 (drier)

rainfall scenario for some of the substrates (*Sphagnum*), this resulted in an increase in the partitioning coefficient between CO₂ and DOC flux for some of the substrates (*Sphagnum* and peat), which is in accordance with the findings of Neff and Hooper (2002) in their arctic soil samples. This implies that a greater proportion of C released from *Sphagnum* and peat would be lost in the form of CO₂ than DOC under warmer and drier conditions, with the effect of rainfall being slightly stronger than temperature. The strength of the rainfall effect suggests that it is due more to the decreased flux of DOC than due to a reduction in total decomposition. The total DOC measured in this study was in effect net DOC production; a function of release and transport, as well as its mineralisation. The decreased DOC flux could be explained by a suppression of DOC mobility by drought-induced acidification (Clark et al., 2006; Clark et al., 2012). DOC production could be affected by both microbial enzymatic activity and physical leaching. It could be that DOC was produced, but lack of water supply limited physical leaching and so it was subsequently consumed by microbes present in the samples (Moore and Dalva, 2001). The findings presented here, that CO₂:DOC partitioning from *Sphagnum* and peat could be affected by climate change, but that the partitioning from other substrates was not affected, will be of interest to those modelling the response of peatland C balance as it indicates that DOC flux cannot be estimated from ecosystem respiration alone. It is also useful information for water treatment engineers who are trying to plan future climate-proof investments in treatment works, as these results imply that DOC could increase or decrease in the future, depending on the proportion of different plant species in the catchments, and the pattern of rainfall.

4.3. *Effects and interactions between temperature and rainfall*

Results indicate that simulated temperature changes have a greater effect on the loss of DOC (generally decreasing the flux) and on mass loss (generally increasing the flux) than on the production of CO₂. The effect varied between substrates, for example, drier conditions increased the DOC flux from *Calluna*, but decreased it from *Sphagnum*. When looking at the main drivers of increased decomposition in peatlands, other laboratory based studies have also found mixed results. Higher temperatures led to increased mass loss (between 5, 10 and 15 °C) in a one year incubation of Scots pine needles below a living moss layer (Domisch et al., 2006), supporting the results found here. Neff and Hooper (2002) also found that temperature controlled the CO₂ flux from their incubated soil samples, but the response depended largely on vegetation community. However, in their study on DOC production across a range of spatial scales, Preston et al. (2011) found that temperature had no effect on DOC production in their peat microcosm experiment, regardless of moisture treatment. This contrasts with Clark et al. (2006) who found increased DOC release at 20 °C relative to 10 °C in another laboratory study incubating peat mesocosms.

Whilst measured mass loss did increase positively with temperature, measured CO₂ fluxes did not. Investigations at lower temperatures than those used here have suggested *Sphagnum* becomes more productive, growing faster, with increases in temperature from a low base (Breeuwer et al., 2008), but that the opposite can occur for warmer temperatures (17.5 and 21.7 °C) (Breeuwer et al., 2009). Results herein indicate that decomposition, measured through mass loss, increased with temperature, suggesting that the strength of the peatland C sink could be reduced during summer months.

Rainfall treatment was just above the $P = 0.05$ critical threshold for cumulative CO₂ ($P = 0.075$) and DOC flux ($P = 0.070$), with lower values seen under the UKCP09 (lower rainfall)

conditions. For DOC in peat, this agrees with the findings of Tang et al. (2013) who found that DOC export was greater in the wetter treatments, and declined in the drought treatments. Observed responses of DOC export to drought or experimentally lowered water tables have been varied, with suggestions that DOC export could increase due to drought (Worrall et al., 2006). However, there are mechanisms which could explain the decrease in DOC observed here. Decreased DOC during droughts could be due to a suppression of DOC transport by drought-induced decrease in DOC solubility (Clark et al., 2012, 2006), or an increase in hydrophobicity (Worrall et al., 2008) rather than a change in the production of DOC through organic matter decomposition. A difference here is that this study simulated rainfall whereas those examining the mechanisms have tended to focus on water table effects, either through manipulations in the laboratory or event-based analysis in the field.

4.4. *Limitations of the study*

The data presented here are a useful indication of the relative importance of different substrates as sources of DOC and CO₂ fluxes, decomposition and the potential for changes in partitioning between CO₂ and DOC under future climate conditions. However, these are laboratory based simulation experiments and are subject to constraints that must be taken into account when interpreting the results. Samples may have been exposed to more extreme drying conditions than under field conditions as moisture levels were not regulated by living vegetation and/or underlying water table, and the lack of a peat substrate beneath the samples from the vegetation groups could also have increased the loss of water and so negated any rainfall treatment effect. Ritson et al. (2017) found that any exposure to oxygen (during laboratory experiments) increased DOC from peat and changed the quality of the C. The differences with the results presented here suggest a high sensitivity to the amount and

frequency of the rainfall treatments. The findings of these laboratory experiments now need to be tested under field conditions.

4.5. *Wider implications for our understanding of peatland carbon balances*

While predictions about future rainfall patterns are variable, with both increases and decreases projected for northern latitudes (IPCC, 2014), models are in strong agreement that temperatures at mid-to-high latitudes will increase due to climate change (Murphy et al., 2009), and that drought events are likely to become more frequent (Meehl and Tebaldi, 2004). While there are differences in both magnitude and direction of change between the substrates, the data here suggest that warmer summer temperatures and decreased rainfall could increase overall decomposition, reduce the flux of DOC, and subsequently alter the partitioning of carbon released during decomposition from DOC to CO₂.

Other studies have suggested that the effect of climate change will be more through indirect effects on the composition of vegetation, with graminoids and dwarf shrubs being favoured over *Sphagnum* mosses in warmer, drier conditions (Kapfer et al., 2011; Weltzin et al., 2003). Given the greater lability of the litter produced by these plant functional types, seen here in the greater decomposition of the *Molinia* samples, this could be a concern for climate change mitigation. However, in SW England, where these samples were collected, graminoids (*Molinia*) are already highly prevalent (Chambers et al., 1999) and so the effect may not be as severe, as climate-induced vegetation change may already have occurred. The shallow peat in this area suggests the area could have been sequestering C at a lower rate than other blanket peatlands, or even losing C, though further work looking at the entire C balance is needed to verify this.

In the field, it is likely that climate change may increase productivity, and therefore litter production, alongside any increase in decomposition rates. Studies have already provided evidence that warmer temperatures can increase rates of carbon sequestration at some northern latitude peatlands, due in part to longer and warmer growing seasons enhancing plant growth (Loisel and Yu, 2013). Therefore, when interpreting the differences in decomposition rates of the different substrates in the context of *in situ* actual conditions, it is also important to consider their respective litter production rates and substrate supply (Ritson et al., 2016). While *Molinia* showed the fastest decomposition in this experiment, it also tends to produce more litter than *Calluna* (e.g. Aerts, 1989), so is likely to be a more significant source of C at the catchment scale. Also, while both substrates generally produce less “litter” than *Sphagnum* (Bragazza et al., 2012), this may change in a warmer, drier climate (Breeuwer et al., 2009). Further work is needed to take into account the effect of climate change on productivity and litter production, as well as on decomposition, to assess the likely impacts of climate change on the future carbon balance of peatlands.

5. Conclusions

This study has used short-term laboratory incubations of peat and peat-forming vegetation from a climatically marginal blanket peatland in SW England in order to assess the relative importance of different temperature and rainfall scenarios for the main C fluxes (CO₂ and DOC) and their partitioning during short-term litter decay. The largest variation in C fluxes and total mass loss was seen between substrates, with smaller fluxes (cumulative CO₂ and DOC) and mass lost from *Sphagnum*/peat compared to higher fluxes and mass loss from dwarf shrub/graminoids. The CO₂:DOC ratio was lowest for *Molinia* and highest for *Sphagnum*. Climate factors were important too, but the effect depended largely on the substrate. For

example, while the future (drier) rainfall scenario increased the DOC flux from *Calluna*, it had the opposite effect for *Sphagnum*. Therefore, indirect effects on species composition in peatlands could ultimately turn out to be more important than direct effects of climate change from increased temperatures and decreased rainfall.

Upscaling of this work from laboratory to field is needed in order to confirm these laboratory-based findings and improve understanding of the likely impacts of climate change on C fluxes and decomposition in peatlands. Further work across natural climatic gradients is one way of addressing this research need, deploying a space for time substitution to understand future climate change.

Acknowledgements

This research was funded by a studentship received by Michael Bell from the Natural Environment Research Council (grant number 1227376). Jonathan Ritson would like to thank the Grantham Institute: Climate and Environment and Climate-KIC for financial support. Brazier acknowledges the support of the South West Water-funded Mires-on-the-Moors project. The authors would also like to thank South West Water and especially the Mires project staff for access to sites as well as the Exmoor National Park Authority. The help of Dr Radek Kowalczyk of University of Reading for running the NMR experiments and advising on their experimental design is also greatly appreciated.

References

1. Aerts, R., 1989. Aboveground Biomass and Nutrient Dynamics of *Calluna vulgaris* and *Molinia caerulea* in a Aboveground biomass and nutrient dynamics of *Calluna vulgaris*

and *Molinia caerulea* in a dry heathland. *OIKOS* 56, 31–38.

2. Billett, M., Charman, D., Clark, J., Evans, C., Evans, M., Ostle, N., Worrall, F., Burden, A., Dinsmore, K., Jones, T., McNamara, N., Parry, L., Rowson, J., Rose, R., 2010. Carbon balance of UK peatlands: current state of knowledge and future research challenges. *Clim. Res.* 45, 13–29. doi:10.3354/cr00903
3. Billett, M.F., Palmer, S.M., Hope, D., Deacon, C., Storeton-West, R., Hargreaves, K.J., Flechard, C., Fowler, D., 2004. Linking land-atmosphere-stream carbon fluxes in a lowland peatland system. *Global Biogeochem. Cycles* 18, 1–12. doi:10.1029/2003GB002058
4. Blok, D., Elberling, B., Michelsen, A., 2015. Initial Stages of Tundra Shrub Litter Decomposition May Be Accelerated by Deeper Winter Snow But Slowed Down by Spring Warming. *Ecosystems* 19, 155–169. doi:10.1007/s10021-015-9924-3
5. Bonnett, S.A.F., Ostle, N., Freeman, C., 2006. Seasonal variations in decomposition processes in a valley-bottom riparian peatland. *Sci. Total Environ.*, 370, 561–573.
6. Bragazza, L., Buttler, A., Habermacher, J., Brancaleoni, L., Gerdol, R., Fritze, H., Hanajík, P., Laiho, R., Johnson, D., 2012. High nitrogen deposition alters the decomposition of bog plant litter and reduces carbon accumulation. *Glob. Chang. Biol.* 18, 1163–1172. doi:10.1111/j.1365-2486.2011.02585.x
7. Bragazza, L., Buttler, A., Siegenthaler, A., Mitchell, E.A.D., 2009. Plant Litter Decomposition and Nutrient Release in Peatlands, in: Baird, A.J., Belyea, L.R., Comas, X., Reeve, A.S., Slater, L. (Eds.), *Carbon Cycling in Northern Peatlands*. American Geophysical Union, pp. 99–110.
8. Breeuwer, A., Heijmans, M.M.P.D., Gleichman, M., Robroek, B.J.M., Berendse, F., 2009. Response of *Sphagnum* species mixtures to increased temperature and

- nitrogen availability. *Plant Ecol.* 204, 97–111. doi:10.1007/s11258-009-9571-x
9. Breeuwer, A., Heijmans, M.M.P.D., Robroek, B.J.M., Berendse, F., 2008. The effect of temperature on growth and competition between *Sphagnum* species. *Oecologia* 156, 155–67. doi:10.1007/s00442-008-0963-8
 10. Broder, T., Blodau, C., Biester, H., Knorr, K.H., 2012. Peat decomposition records in three pristine ombrotrophic bogs in southern Patagonia. *Biogeosciences* 9, 1479–1491. doi:10.5194/bg-9-1479-2012
 11. Bubier, J.L., Moore, T.R., Bledzki, L. a., 2007. Effects of nutrient addition on vegetation and carbon cycling in an ombrotrophic bog. *Glob. Chang. Biol.* 13, 1168–1186. doi:10.1111/j.1365-2486.2007.01346.x
 12. Chambers, F.M., Mauquoy, D., Todd, P.A., 1999. Recent rise to dominance of *Molinia caerulea* in environmentally sensitive areas : new perspectives from palaeoecological data. *J. Appl. Ecol.* 36, 719–733.
 13. Clark, J., Gallego-Sala, A., Allott, T., Chapman, S., Farewell, T., Freeman, C., House, J., Orr, H., Prentice, I., Smith, P., 2010. Assessing the vulnerability of blanket peat to climate change using an ensemble of statistical bioclimatic envelope models. *Clim. Res.* 45, 131–150. doi:10.3354/cr00929
 14. Clark, J.M., Chapman, P.J., Heathwaite, A.L., Adamson, J.K., 2006. Suppression of Dissolved Organic Carbon by Sulfate Induced Acidification during Simulated Droughts. *Environ. Sci. Technol.* 40, 1776–1783. doi:10.1021/es051488c
 15. Clark, J.M., Heinemeyer, A., Martin, P., Bottrell, S.H., 2012. Processes controlling DOC in pore water during simulated drought cycles in six different UK peats. *Biogeochemistry* 109, 253–270. doi:10.1007/s10533-011-9624-9
 16. Dinsmore, K.J., Billett, M.F., Dyson, K.E., 2013. Temperature and precipitation drive

- temporal variability in aquatic carbon and GHG concentrations and fluxes in a peatland catchment. *Glob. Chang. Biol.* 19, 2133–48. doi:10.1111/gcb.12209
17. Dinsmore, K.J., Billett, M.F., Skiba, U.M., Rees, R.M., Drewer, J., Helfter, C., 2010. Role of the aquatic pathway in the carbon and greenhouse gas budgets of a peatland catchment. *Glob. Chang. Biol.* 16, 2750–2762. doi:10.1111/j.1365-2486.2009.02119.x
18. Domisch, T., Finér, L., Laine, J., Laiho, R., 2006. Decomposition and nitrogen dynamics of litter in peat soils from two climatic regions under different temperature regimes. *Eur. J. Soil Biol.* 42, 74–81. doi:10.1016/j.ejsobi.2005.09.017
19. Evans, C.D., Freeman, C., Cork, L.G., Thomas, D.N., Reynolds, B., Billett, M.F., Garnett, M.H., Norris, D., 2007. Evidence against recent climate-induced destabilisation of soil carbon from ¹⁴C analysis of riverine dissolved organic matter. *Geophys. Res. Lett.* 34, L07407. doi:10.1029/2007GL029431
20. Fenner, N., Freeman, C., Ostle, N.J. and Reynolds, B., 2004. Peatland carbon efflux partitioning reveals that sphagnum photosynthate contributes to the DOC pool. *Plant and Soil.* 259, 345–354.
21. Fenner, N., Ostle, N.J., McNamara, N., Sparks, T., Harmens, H., Reynolds, B., Freeman, C., 2007. Elevated CO₂ Effects on Peatland Plant Community Carbon Dynamics and DOC Production. *Ecosystems* 10, 635–647. doi:10.1007/s10021-007-9051-x
22. Frohking, S., Roulet, N.T., Tuittila, E., Bubier, J.L., Quillet, A., Talbot, J., Richard, P.J.H., 2010. A new model of Holocene peatland net primary production, decomposition, water balance, and peat accumulation. *Earth Syst. Dyn.* 1, 1–21. doi:10.5194/esd-1-1-2010
23. Freeman, C., Lock, M.A., Reynolds, B., 1993. Fluxes of carbon dioxide, methane and nitrous oxide from a Welsh peatland following simulation of water table draw-down:

- Potential feedback to climatic change. *Biogeochemistry*. 19, 51–60.
24. Gatis, N., Luscombe, D.J., Grand-Clement, E., Hartley, I.P., Anderson, K., Smith, D., Brazier, R.E., 2015. The effect of drainage ditches on vegetation diversity and CO₂ fluxes in a *Molinia caerulea*-dominated peatland. *Ecohydrology* n/a-n/a. doi:10.1002/eco.1643
25. Gorham, E., 1991. Northern Peatlands: Role in the carbon cycle and probable responses to climatic warming. *Ecol. Appl.* 1, 182–195.
26. Grand-Clement, E., Anderson, K., Smith, D., Luscombe, D., Gatis, N., Ross, M., Brazier, R.E., 2013. Evaluating ecosystem goods and services after restoration of marginal upland peatlands in South-West England. *J. Appl. Ecol.* n/a-n/a. doi:10.1111/1365-2664.12039
27. Grand-Clement, E., Luscombe, D.J., Anderson, K., Gatis, N., Benaud, P., Brazier, R.E., 2014. Antecedent conditions control carbon loss and downstream water quality from shallow, damaged peatlands. *Sci. Total Environ.* 493, 961–73. doi:10.1016/j.scitotenv.2014.06.091
28. Harrison, A.F., Taylor, K., Scott, A., Poskitt, J., Benham, D., Grace, J., Chaplow, J., Rowland, P., 2008. Potential effects of climate change on DOC release from three different soil types on the Northern Pennines UK: examination using field manipulation experiments. *Glob. Chang. Biol.* 14, 687–702. doi:10.1111/j.1365-2486.2007.01504.x
29. Holden, J., Burt, T.P., 2002. Infiltration, runoff and sediment production in blanket peat catchments: implications of field rainfall simulation experiments. *Hydrol. Process.* 16, 2537–2557. doi:10.1002/hyp.1014
30. Hope, D., Billett, M.F., Cresser, M.S., 1994. A review of the export of carbon in river

water: Fluxes and processes. *Environ. Pollut.* 84, 301–324.

31. IPCC, 2014. IPCC, 2014: Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change [Core Writing Team, R.K. Pachauri and L.A. Meyer (eds.)]. Geneva, Switzerland.
32. Johnson, L.C., Damman, A.W.H., 1991. Species Controlled Sphagnum Decay on a South Swedish raised bog Sphagnum decay. *Oikos* 61, 234–242.
33. Kapfer, J., Grytnes, J.-A., Gunnarsson, U., Birks, H.J.B., 2011. Fine-scale changes in vegetation composition in a boreal mire over 50 years. *J. Ecol.* 99, 1179–1189. doi:10.1111/j.1365-2745.2011.01847.x
34. Keselman, H.J., 1975. A Monte Carlo investigation of three estimates of treatment magnitude: Epsilon squared, eta squared, and omega squared. *Can. Psychol. Rev.* 16, 44–48.
35. Keuskamp, J.A., Dingemans, B.J.J., Lehtinen, T., Sarneel, J.M., Hefting, M.M., 2013. Tea Bag Index: a novel approach to collect uniform decomposition data across ecosystems. *Methods Ecol. Evol.* 4, 1070–1075. doi:10.1111/2041-210X.12097
36. Koehler, A.-K., Sottocornola, M., Kiely, G., 2011. How strong is the current carbon sequestration of an Atlantic blanket bog? *Glob. Chang. Biol.* 17, 309–319. doi:10.1111/j.1365-2486.2010.02180.x
37. Laine, A., Sottocornola, M., Kiely, G., Byrne, K.A., Wilson, D., Tuittila, E.-S., 2006. Estimating net ecosystem exchange in a patterned ecosystem: Example from blanket bog. *Agric. For. Meteorol.* 138, 231–243.
38. Limpens, J., Berendse, F., 2003. How litter quality affects mass loss and N loss from decomposing Sphagnum. *Oikos* 103, 537–547. doi:10.1034/j.1600-

0706.2003.12707.x

39. Loisel, J., Yu, Z., 2013. Recent acceleration of carbon accumulation in a boreal peatland, south central Alaska. *J. Geophys. Res. Biogeosciences* 118, 41–53.
doi:10.1029/2012JG001978
40. Luscombe, D.J., Anderson, K., Gatis, N., Grand-Clement, E., Brazier, R.E., 2015. Using airborne thermal imaging data to measure near-surface hydrology in upland ecosystems. *Hydrol. Process.* 29, 1656–1668. doi:10.1002/hyp.10285
41. Meehl, G.A., Tebaldi, C., 2004. More intense, more frequent, and longer lasting heat waves in the 21st century. *Science* 305, 994–7. doi:10.1126/science.1098704
42. Moore, T.R., Bubier, J.L., Bledzki, L., 2007. Litter Decomposition in Temperate Peatland Ecosystems: The Effect of Substrate and Site. *Ecosystems* 10, 949–963.
doi:10.1007/s10021-007-9064-5
43. Moore, T.R., Dalva, M., 2001. Some controls on the release of dissolved organic carbon by plant tissues and soils. *Soil Sci.* 166, 38–47.
44. Moore, T.R., Paré, D., Boutin, R., 2008. Production of Dissolved Organic Carbon in Canadian Forest Soils. *Ecosystems* 11, 740–751. doi:10.1007/s10021-008-9156-x
45. Murphy, J.M., Sexton, D.M.H., Jenkins, G.J., Booth, B.B.B., Brown, C.C., Clark, R.T., Collins, M., Harris, G.R., Kendon, E.J., Betts, R.A., Brown, S.J., Humphrey, K.A., McCarthy, M.P., McDonald, R.E., Stephens, A., Wallace, C., Warren, R., Wilby, R., Wood, R.A., 2009. UK Climate Projections Science Report: Climate Change Projections.
46. Neff, J.C., Hooper, D.U., 2002. Vegetation and climate controls on potential CO₂, DOC and DON production in northern latitude soils. *Glob. Chang. Biol.* 8, 872–884.
doi:10.1046/j.1365-2486.2002.00517.x

47. Palmer, S.M., Hope, D., Billett, M.F., Dawson, J.J.C., Bryant, C.L., 2001. Sources of organic and inorganic carbon in a headwater stream: Evidence from carbon isotope studies. *Biogeochemistry* 52, 321–338. doi:10.1023/A:1006447706565
48. Parry, L.E., Charman, D.J., 2013. Modelling soil organic carbon distribution in blanket peatlands at a landscape scale. *Geoderma* 211–212, 75–84.
doi:10.1016/J.GEODERMA.2013.07.006
49. Peuravuori, J., Ingman, P. and Pihlaja, K., 2003. Critical comments on accuracy of quantitative determination of natural humic matter by solid state (13)C NMR spectroscopy. *Talanta* 59(1), 177–89.
50. Preston, M.D., Eimers, M.C., Watmough, S.A., 2011. Effect of moisture and temperature variation on DOC release from a peatland: conflicting results from laboratory, field and historical data analysis. *Sci. Total Environ.* 409, 1235–42.
doi:10.1016/j.scitotenv.2010.12.027
51. R Core Team, 2015. R: A language and environment for statistical computing.
52. Ritson, J.P., Bell, M., Graham, N.J.D., Templeton, M.R., Brazier, R.E., Verhoef, A., Freeman, C., Clark, J.M., 2014. Simulated climate change impact on summer dissolved organic carbon release from peat and surface vegetation: Implications for drinking water treatment. *Water Res.* 67, 66–76. doi:10.1016/j.watres.2014.09.015
53. Ritson, J.P., Bell, M., Brazier, R.E., Grand-Clement, E., Graham, N.J.D., Freeman, C., Smith, D., Templeton, M.R., Clark, J.M., 2016. Managing peatland vegetation for drinking water treatment. *Scientific Reports.* 6, 36751.
54. Rodwell, J., 1991. *British plant communities, Vol 2: mires and heaths.* Cambridge University Press, Cambridge.
55. Smith, J., Gottschalk, P., Bellarby, J., Chapman, S., Lilly, a, Towers, W., Bell, J.,

- Coleman, K., Nayak, D., Richards, M., Hillier, J., Flynn, H., Wattenbach, M., Aitkenhead, M., Yeluripati, J., Farmer, J., Milne, R., Thomson, a, Evans, C., Whitmore, a, Falloon, P., Smith, P., 2010. Estimating changes in Scottish soil carbon stocks using ECOSSE. I. Model description and uncertainties. *Clim. Res.* 45, 179–192.
doi:10.3354/cr00899
56. Straková, P., Penttilä, T., Laine, J., Laiho, R., 2012. Disentangling direct and indirect effects of water table drawdown on above- and belowground plant litter decomposition: consequences for accumulation of organic matter in boreal peatlands. *Glob. Chang. Biol.* 18, 322–335. doi:10.1111/j.1365-2486.2011.02503.x
57. Tang, R., Clark, J.M., Bond, T., Graham, N., Hughes, D., Freeman, C., 2013. Assessment of potential climate change impacts on peatland dissolved organic carbon release and drinking water treatment from laboratory experiments. *Environ. Pollut.* 173, 270–277.
58. Toberman, H., Evans, C.D., Freeman, C., Fenner, N., White, M., Emmett, B.A., Artz, R.R.E., 2008. Summer drought effects upon soil and litter extracellular phenol oxidase activity and soluble carbon release in an upland *Calluna* heathland. *Soil Biology & Biochemistry* 40, 1519–1532.
59. Turetsky, M.R., 2003. The Role of Bryophytes in Carbon and Nitrogen Cycling. *Bryologist* 106, 395–409.
60. van Breemen, N., 1995. How Sphagnum bogs down other plants. *Trends Ecol. Evol.* 10, 270–275.
61. Wallage, Z.E., Holden, J., 2011. Near-surface macropore flow and saturated hydraulic conductivity in drained and restored blanket peatlands. *Soil Use Manag.* 27, 247–254.
doi:10.1111/j.1475-2743.2011.00336.x

62. Weltzin, J.F., Bridgham, S.D., Pastor, J., Chen, J., Harth, C., 2003. Potential effects of warming and drying on peatland plant community composition. *Glob. Chang. Biol.* 9, 141–151. doi:10.1046/j.1365-2486.2003.00571.x
63. Wickham, H., 2009. *ggplot2: elegant graphics for data analysis*. Springer, New York.
64. Wieder, R.K., Lang, G.E., 1982. A Critique of the Analytical Methods Used in Examining Decomposition Data Obtained From Litter Bags. *Ecology* 63, 1636–1642.
65. Wong, S., Hanna, J. V, King, S., Carroll, T.J., Eldridge, R.J., Dixon, D.R., Bolto, B.A., Hesse, S., Abbt-Braun, G., Frimmel, F.H., 2002. Fractionation of natural organic matter in drinking water and characterization by ¹³C cross-polarization magic-angle spinning NMR spectroscopy and size exclusion chromatography. *Environ. Sci. Technol.* 36, 3497–503. doi:10.1021/es010975z
66. Worrall, F., Gibson, H.S., Burt, T.P., 2008. Production vs. solubility in controlling runoff of DOC from peat soils – The use of an event analysis. *J. Hydrol.* 358, 84–95. doi:10.1016/j.jhydrol.2008.05.037