

Harnessing positive species interactions as a tool against climate-driven loss of coastal biodiversity

Bulleri, Fabio; Eriksson, Britas Klemens ; Queirós, Ana ; Airolidi, Laura ; Arenas, Francisco; Arvanitidis, Christos ; Bouma, Tjeerd; Crowe, Tasman; Davoult, Dominique; Guizien, Katell ; Iveša, Ljiljana ; Jenkins, Stuart; Michalet, Richard; Olabarria, Celia ; Procaccini, Gabriele ; Serrão, Ester A. ; Wahl, Martin ; Benedetti-Cecchi, Lisandro

PLoS Biology

DOI:
[10.1371/journal.pbio.2006852](https://doi.org/10.1371/journal.pbio.2006852)

Published: 04/09/2018

Peer reviewed version

[Cyswllt i'r cyhoeddiad / Link to publication](#)

Dyfyniad o'r fersiwn a gyhoeddwyd / Citation for published version (APA):

Bulleri, F., Eriksson, B. K., Queirós, A., Airolidi, L., Arenas, F., Arvanitidis, C., Bouma, T., Crowe, T., Davoult, D., Guizien, K., Iveša, L., Jenkins, S., Michalet, R., Olabarria, C., Procaccini, G., Serrão, E. A., Wahl, M., & Benedetti-Cecchi, L. (2018). Harnessing positive species interactions as a tool against climate-driven loss of coastal biodiversity. *PLoS Biology*, 6(9), Article e2006852. <https://doi.org/10.1371/journal.pbio.2006852>

Hawliau Cyffredinol / General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Full title: **Harnessing positive species interactions as a tool against climate-driven loss of coastal biodiversity**

Short title: Positive species interactions in changing climates

Fabio Bulleri^{1*}, Britas Klemens Eriksson², Ana Queirós³, Laura Airoidi⁴, Francisco Arenas⁵, Christos Arvanitidis⁶, Tjeerd J. Bouma⁷, Tasman P. Crowe⁸, Dominique Davoult⁹, Katell Guizien¹⁰, Ljiljana Iveša¹¹, Stuart R. Jenkins¹², Richard Michalet¹³, Celia Olabarria¹⁴, Gabriele Procaccini¹⁵, Ester A. Serrão¹⁶, Martin Wahl¹⁷, Lisandro Benedetti-Cecchi¹

¹Dipartimento di Biologia, Università di Pisa, CoNISMa, Via Derna 1, 56126, Pisa, Italy

²Groningen Institute for Evolutionary Life-Sciences, University of Groningen, Nijenborgh 7, 9747 AG Groningen, The Netherlands

³Plymouth Marine Laboratory, Prospect Place, PL1 3DH Plymouth, UK

⁴University of Bologna, Dipartimento di Scienze Biologiche, Geologiche ed Ambientali (BiGeA), & Centro Interdipartimentale di Ricerca per le Scienze Ambientali (CIRSA), UO CoNISMa, Italy

⁵CIIMAR - Interdisciplinary Center of Marine and Environmental Research, Av. General Norton de Matos S/N 4450-208, Matosinhos, Portugal

⁶Institute of Marine Biology, Biotechnology and Aquaculture, Hellenic Centre for Marine Research, Thalassokosmos, Former US Base at Gournes, 71003, Heraklion, Crete, Greece

⁷NIOZ Royal Netherlands Institute for Sea Research, Department of Estuarine and Delta Systems and Utrecht University, P.O. Box 140, 4400 AC Yerseke, The Netherlands

⁸Earth Institute and School of Biology and Environmental Science, University College Dublin

⁹Sorbonne Université, CNRS, UMR 7144, Station Biologique, Place Georges Teissier, F. 29680 Roscoff, France

¹⁰Sorbonne Universités, UPMC Univ Paris 06, CNRS, Laboratoire d'Ecogéochimie des Environnements Benthiques (LECOB), Observatoire Océanologique, F-66650, Banyuls/Mer, France

¹¹Ruder Bošković Institute, Center for Marine Research, G. Paliaga, Rovinj, Croatia

¹²School of Ocean Sciences, Bangor University, Menai Bridge, Anglesey, LL59 5AB, UK

¹³UMR CNRS 5805 EPOC, University of Bordeaux, 33405 Talence, France

¹⁴Departamento de Ecoloxía e Bioloxía Animal, Facultade de Ciencias del Mar, Campus Lagoas-Marcosende, Universidade de Vigo, 36701, Vigo-Pontevedra, Spain.

¹⁵Stazione Zoologica Anton Dohrn, Villa Comunale, 80121, Napoli, Italy

¹⁶CCMAR, CIMAR, University of Algarve, Campus de Gambelas, 8005-139, Faro, Portugal

¹⁷GEOMAR Helmholtz Centre for Ocean Research, D-24105 Kiel, Germany

* Corresponding author: email: fabio.bulleri@unipi.it

Abstract

40 Habitat-forming species sustain biodiversity and ecosystem functioning in harsh
environments through the amelioration of physical stress. Nonetheless, their role in shaping patterns
of species distribution under future climate scenarios is generally overlooked. Focusing on coastal
systems, we assess how habitat-forming species can influence the ability of stress-sensitive species
to exhibit plastic responses, adapt to novel environmental conditions or track suitable climates.
45 Here, we argue that habitat-former populations could be managed as a nature-based solution against
climate-driven loss of biodiversity. Drawing from different ecological and biological disciplines, we
identify a series of actions to sustain the resilience of marine habitat-forming species to climate
change, as well as their effectiveness and reliability in rescuing stress-sensitive species from
increasingly adverse environmental conditions.

50

55

Positive species interactions under climate change

60 Anthropogenic climate change is causing unprecedented alterations to Earth's ecosystems [1,2]. Modifications in species distribution and abundance as a consequence of altered environmental conditions can be the direct result of physiological and/or phenological responses [3]. More often, climate-induced modifications in individual physiology, phenology and behavior scale up to the community level through the filter of species interactions [4]. Nonetheless, species
65 interactions are still seldom incorporated into models aiming to forecast species distribution under future climate scenarios [5,6].

Although terrestrial and marine studies have started addressing the effects of climate change on the balance between negative and positive species interactions (Box 1) [7-11], the role of habitat-formers (Box 1) in shaping future patterns of species distribution is yet to be fully explored. This is
70 at odds with compelling evidence showing that habitat-formers frequently facilitate other species in otherwise hostile environments [5,8,12-15] and can enhance conservation and restoration success [16-18]. Habitat-formers have allowed species to persist under dramatic changes in climate in the past and acted as important evolutionary forces. For instance, environmental stress amelioration by canopy-forming Quaternary plants has allowed Tertiary plant lineages adapted to moist conditions
75 to persist despite the onset of an unfavorable climate [19]. Indeed, biogenic modification of abiotic conditions (Box 1) underpins pivotal chapters in the evolution of life on Earth: in the Cambrian, the development of biomineralised skeletons (e.g. trilobites and other arthropods), a response to the advent of predation, caused reworking and oxygenation of ocean sediments (i.e. the burrowing revolution), giving rise to the ancestors of many modern groups of animals [20]. Milder conditions
80 due to warming may reduce the reliance of extant species on habitat-formers in some extreme environments, such as alpine and arctic tundra [7]. There is, however, undisputable evidence that increasingly harsher physical conditions are a major driver of the current biodiversity crisis across ecosystems on Earth [1,2], suggesting that the importance of physical stress amelioration by habitat-formers is set to increase under future climate scenarios.

Box 1. Glossary

Positive species interactions: interactions among species, also referred to as facilitative interactions or facilitation, in which at least one of the participants benefits from the presence of the other, while neither is disadvantaged. These include interactions between co-evolved, mutually obligate organisms, as well as looser, facultative interactions between species that did not co-evolve.

Habitat-former: A species able to support the persistence of other species by providing suitable environmental conditions, enhancing the availability of or access to limiting resources or reducing the effects of negative species interactions, such as competition, predation and diseases. Habitat-formers include ecosystem engineers which are defined as organisms that affect other species through the creation, modification and maintenance of habitat. Biotic and abiotic conditions are not necessarily optimal (relative to other habitats) for all the species found in the presence of a habitat-former.

Biogenic modification of environmental conditions: Modification of environmental conditions operated by a living organism (i.e. a habitat-former). Similarly, biogenic amelioration or buffering of environmental stress refers to the case in which the presence of a living organism reduces the intensity of stressful environmental conditions for other species.

Biogenic refugia: habitats formed by living organisms and of limited spatial extent that allow other species to escape adverse environmental or biological conditions and from which they can subsequently expand when suitability of external conditions is restored.

Benefactor and beneficiary species: The benefactor is a species able to deliver benefits to other species, defined as beneficiary species. A species may behave as a benefactor under some environmental conditions or resource availability levels, but not under others. For example, an intertidal canopy-forming macroalga (i.e. the benefactor) can benefit understory (i.e. the beneficiaries) reducing heat and desiccation at high-shore levels. By contrast, it can negatively influence understory species lower on the shore, where heat and desiccation stress are less severe.

Epigenetic mechanisms: Mechanisms which form the basis of the dynamic regulation of gene expression through chromatin remodeling, DNA methylation, non-coding RNA-associated genes and histone modification. Epigenetic changes can be inherited but do not involve changes in the underlying DNA sequence.

Assortative mating: Non-random mating model in which the frequency of mating between individuals with a similar genotype and/or phenotype is higher than that expected by chance.

Climate rescuer: A habitat-former resistant/resilient to climate change providing suitable environmental conditions to species that would otherwise be unable to maintain viable population under future climate scenarios.

Habitat-formers are key in shaping community structure and ecosystem functioning in marine environments, through both local and long-distance positive interactions that extend across coastal landscapes [12,13,21]. In transitional and shallow-water environments, the habitat-former concept has traditionally been applied to sessile species, such as mangroves, salt-marsh plants, seagrasses, macroalgae, bivalves and corals [22] (Fig. 1A-E). However, mobile species that modify the characteristics of sediments through their burrowing or feeding activity (i.e. bioturbators; Fig. 1F), such as holothurians, crustaceans and polychaetes, could play a similar role from tidal flats to abyssal plains [23]. Here, we assess the circumstances under which biogenic amelioration of environmental stress may sustain coastal biodiversity and ecosystem functioning in the face of climate change and, hence, be used as a nature-based solution for coastal conservation and restoration.

FIGURE 1. Habitat-formers in intertidal and subtidal environments. (A) clumps of the mussel *Mytilus edulis* on a tidal flat in the Wadden sea, the Netherlands (Photo credit: B.K.E. Eriksson); (B) mangrove trees of the species *Avicennia marina* along the central coasts of the Red Sea (Photo credit: T. Dailianis); (C) fronds of the brown seaweed *Fucus vesiculosus* at low tide on a rocky

shore of the Iberian Peninsula (Photo credit: E. Serrão); (D) the seagrass *Posidonia oceanica* in shallow waters of Crete, Aegean Sea (Photo credit: T. Dailianis); (E) multi-specific canopy stands formed by the brown seaweeds *Cystoseira barbata*, *C. compressa* and *C. crinita* on shallow rocky reefs of Croatia, NE Adriatic Sea (Photo credit: L. Iveša); (F) burrowing by the sea cucumber *Holothuria scabra* exposes anoxic sediments on a reef flat in Fiji (Photo credit: S. Lee).

Biogenic refugia against climate change

Biogenic buffering of environmental stress has been documented in harsh, transitional habitats, such as intertidal rocky and sandy shores, mudflats and salt marshes [12]. For example, intertidal macroalgal canopies or mussels beds reduce heat and desiccation stress during emersion, sustaining diversity and productivity of benthic communities [24,25]. However, while the role of geo-morphological refugia for species persistence in the face of past and current changes in climate is recognized [26], that of biogenic refugia (Box 1) remains unexplored. Benefactors (Box 1) may provide climatically suitable habitat for stress-susceptible species, increasing their survival during acute climate-driven disturbance events, such as heat-waves or sea-storms. For example, intertidal mussel clumps enhance cordgrass survival during severe drought events and function as nuclei for vegetative recovery in the aftermath [11]. In subtidal environments, macrophyte photosynthetic activity buffers calcifying organisms from ocean acidification by increasing pH [27,28]. Daily uptake of CO₂ by plants increases pH within the surrounding diffusive boundary layer and these effects can scale up to adjacent habitats, such as stony corals or mussel beds [28,29]. Subtidal canopies also attenuate wave-action and, at shallow depths, light stress [30,31]. Below the sediment surface, biogenic activity can reduce the impacts of seasonal hypoxia driven by heat-waves. Seawater flushing and particle mixing by large burrowing marine invertebrates (i.e. bioturbation and bioirrigation) facilitate oxygenation of sedimentary pore water spaces and the burial of organic matter, ameliorating biogeochemical conditions within sediments [32,33]. Indeed, reduction of physical stress by bioturbators (e.g. temperature-driven hypoxia) may explain why the proportion of

benthic species on soft-sediments shifting their trailing edge at the pace predicted by seawater warming rates is lower than expected [34].

165 Facilitation can expand the distribution of beneficiary species beyond the range predicted from their physiological tolerance matrices [35-37]. The magnitude of the biogenic reduction of thermal stress may exceed - by far - the increment expected under warming climates. For example, intertidal canopies of the seaweed *Ascophyllum nodosum* reduced summer maximum rock temperatures in New England by as much as $\sim 8^{\circ}\text{C}$ [24] and mussels and algal turfs ameliorated
170 lethal and sub-lethal thermal stress over 14° of latitude [35].

Reliance of beneficiaries on biogenic amelioration of environmental conditions may increase under future climates, at least until beneficiary species possibly adapt to the new conditions. Thus, a large proportion of species in a community might become obligate associates with habitat-formers. The survival of beneficiary species would depend, first, upon the spatial and temporal extent of the
175 biogenic refugia and, second, their fitness therein. Refugia might be too small to allow beneficiaries to maintain viable populations. In addition, life in biogenic habitats can entail costs due to competition either with the benefactor itself or other associated species [37].

Adapt, move or perish: the role of biogenic habitat

180 A species that is currently neither resistant (unaffected) nor resilient (able to recover) to climate change must either adapt or move to persist. Can habitat-formers influence the mechanisms underpinning species potential to i) exhibit plastic responses, ii) genetically adapt to novel environmental conditions or iii) track suitable climates?

i) Pre-existing phenotypic plasticity, allowing individuals to acclimate, may sustain short-term
185 population persistence, before evolutionary adaptation can take place [3]. Rapid adaptation to novel environmental conditions, through the activation of alternative metabolic pathways or the modification of gene expression levels by epigenetic mechanisms (Box 1), has been demonstrated in marine organisms [38,39]. Acclimation can also influence subsequent generations and biogenic

habitats may facilitate species acclimation via developmental or transgenerational plasticity

190 exposing individuals to sub-lethal temperatures during extreme events, such as heat-waves [40, 41].

ii) Adaptation to changing climate by selection of individual traits across generations can require time, especially for long-lived organisms. Body mass, reproduction type (e.g. sexual versus vegetative) and generation time influence local adaptation rates [42]. By virtue of their smaller body mass and shorter generation times, adaptation can be expected to be generally more rapid in
 195 beneficiary species than in habitat-formers. However, given that current climate-driven changes may modify marine habitats at rates fast exceeding the potential for adaptive, genetic change within populations, habitat-formers may buy population persistence time for stress-sensitive species. The evolutionary potential of positive interactions remains unquantified [43], but small-scale variation in the intensity of negative biotic interactions (e.g. predation) has been shown to promote rapid
 200 adaptive differentiation [44].

Several lines of argument do suggest that biogenic habitats may influence fine-scale genetic structure of associated species. First, at the seascape scale, patches of habitat-formers alternating with open surfaces - a common configuration of transitional coastal environments - increase spatial heterogeneity in selective pressures, thus sustaining genetic polymorphism. This may explain the
 205 inverted dominance of two alleles found homozygous in barnacles living in exposed sites versus underneath a canopy-forming macroalga [45]. Second, enhanced aggregation of individuals seeking shelter in biogenic habitats, in association with limited dispersal and occurrence of within-habitat environmental gradients [10], can influence genetic structuring through isolation by distance [46]. Third, habitat-formers can elicit phenotypic variations in beneficiary species that, when involving
 210 reproductive traits, may enhance fine-scale genetic structuring through assortative mating (Box 1) [46,47]. Biogenic enhancement of genetic variation would be particularly important in populations at range edges since they may have lower genetic variability compared to central populations [48].

iii) Under lethal climate-driven stress, the synchrony of migration capacities determines species interaction outcomes at the leading edge of range shifts. Three different scenarios describe

215 how climate change can alter species interactions [49]. In the first, all the species within a community migrate synchronously to track climate change without noteworthy modification of the interaction environment. Thus, facilitative effects of habitat-formers could be maintained in newly colonized areas (Fig. 2, scenario 1).

220 **FIGURE 2. Alternative scenarios of interaction between benefactor and beneficiary species after climate change.** Under the current climate, southern and northern canopy-forming macroalgae facilitate different species of barnacles (in the northern hemisphere in this example). Under scenario 1, species migrate synchronously to track suitable climate resulting in no significant modifications of the interacting environment and no generation of novel interactions: extant positive
 225 interactions are maintained. Under scenario 2, all species exhibit the same time lag in migration and interact in harsher environmental conditions, resulting in either (a) a strengthening of positive interactions or, (b) in the case in which levels of stress become excessive, in the collapse of facilitation. Under scenario 3, species migration is asynchronous, generating novel interactions. In this example, southern species migrate poleward and start interacting with extant, non-migrating
 230 species. Positive interactions between each original pair of canopy-forming macroalgae and barnacles are likely to be maintained (green arrows). Novel interactions (red arrow) between canopy-formers and barnacles can be either positive or negative, while novel interactions between canopy-formers and between barnacles are likely to be negative.

235 In the second scenario, all species exhibit the same migration lag, thus interacting under changing environmental conditions. Enhanced levels of environmental stress may increase the frequency and/or intensity of positive interactions [15] (Fig. 2, scenario 2a). For example, along the US east coast, intertidal macroalgal canopies fostered cirriped survival at thermally stressful southern sites [24]. By contrast, at northern cooler sites, benefits were overridden by increased
 240 whelk predation. Progressive warming may strengthen stress mitigation benefits, shifting the net

effect of canopies from negative to positive also at northern sites. Alternatively, facilitation may collapse if environmental stress becomes extreme and impairs the ability of the benefactor to deliver benefits [50] (Fig. 2, scenario 2b).

In the third scenario, some species migrate towards cooler climates and start interacting with resident, non-migrating species (Fig. 2, scenario 3). Such novel interactions can be either positive or negative. Recruitment through seeds, spores or larvae represents a critical stage of range shifts. Juvenile stages are often less tolerant to stressful conditions than adults and biogenic stress amelioration might be crucial to enable their recruitment outside their current distributional range. For instance, on the US east coast, salt marsh vegetation facilitates recruits of the black mangrove, *Avicennia germinans*, at its northward distributional limit [51]. Positive effects do not necessarily stem from environmental stress reduction, but might be generated by alleviation of resource limitation, competition or predation pressure. For example, reefs formed by the Pacific oyster north of its former range provide native mussels with shelter from crab predation [52].

What makes a habitat-former a climate rescuer species?

Ecosystem-wide effects of environmental stress buffering. The first requisite of a climate rescuer (Box 1) is the ability to sustain biodiversity and ecosystem functioning through stress alleviation (Fig. 3). This effect is not limited to temperature or desiccation, but extends to other climate-related stressors, such as ocean acidification, hypoxia, increased UV radiation and changing hydrodynamic regimes. Ideally, positive effects should not be limited to single habitats, but should propagate to other ecosystems. Primary habitat-formers can provide substrate for other habitat-formers (facilitation cascades: [53]) or promote other species across the landscape (habitat cascades: [54]) through long-distance interactions [55]. Within this context, stress-tolerant species that facilitate other species both within and across habitats should be considered standout climate rescuers.

FIGURE 3. Climate rescuer identification and management. The diagram describes sequential steps towards the identification of a climate-rescuer species and possible management actions aimed to sustain i) life-traits underpinning its resistance to future environmental stress and ii) population traits that determine the strength and reliability of its positive effects on stress-sensitive species.

270

Resilient morphology and phenology under changing climates. Climate rescuer species should be able to persist in increasingly stressful environments without facing morphological or phenological modifications that undermine their facilitative functionality (Fig. 3). Climate rescuing would be supported if the benefactor can withstand a greater magnitude of change in a given climate-driven stressor than its beneficiaries whilst still sustaining function. Thus, the success of the benefactor-beneficiary relationship hinges on the relationship between the (climate) response traits of the benefactor relative to its ability to express the (ameliorating) functional effect trait supporting the beneficiary under a changing environment.

275

Functional effect traits of a habitat-former are often related to morphology and can be altered by climate change. Known changes in species morphology associated with climate changes include reduced average body size in ectotherms [56]. Likewise, calcifying organisms, including important habitat-formers such as bivalves, may reduce their growth to compensate for increased metabolic costs incurred in acidified seawater [57]. Reduced size may confer weaker ability to deliver benefits to other species. Calcifiers may also experience changes in the chemical make up of their shells under ocean acidification and warming. This may render them less structurally robust to physical forcing, reducing their ability to serve as anchoring structures for marine diversity [58].

285

Modifications in phenology may also reduce stress buffering capacity. For example, on the coasts of British Columbia, experimental warming delayed the development of annual intertidal algae [59], potentially exposing associated species to desiccation and heat stress during spring low tides.

290

Viable populations under changing climates. The ability of a habitat-former to maintain viable populations at the edges of its distribution or within warming hotspots determines its potential to act as a critical refugium (Fig. 3). In some cases, habitat-former populations have collapsed at the warmer limit of their distribution [60,61]. In other cases, poleward shifts have occurred without
 295 changes at the equatorial range edge. For example, reduced risk of winter freezing has promoted poleward migration of some mangroves, at the expense of salt-marshes, but with no significant equatorial edge contraction [62]. In the southern hemisphere, tropical corals and seagrasses have expanded towards higher latitudes without modifying their northernmost boundaries [63,64].

In addition, climate rescuer populations should not undergo thinning during hot seasons or
 300 extreme atmospheric events (i.e. exhibit large temporal fluctuations) because they might become too sparse to buffer environmental stress. Since habitat modification is often density dependent [65], assessing whether there is a minimal (threshold) population density or size that is needed for benefits to accrue seems crucial.

305 **Active management of habitat-formers to mitigate biodiversity loss**

By virtue of their potential to ameliorate environmental stress, habitat-former populations could be managed as a tool against climate-driven loss of biodiversity (Fig. 3). Major threats to marine habitat-formers and approaches to their conservation have been thoroughly reviewed elsewhere [66] and will be not re-iterated here. Instead, we outline a number of actions to sustain
 310 habitat-formers facing novel climatic conditions, as well as population traits enhancing their effectiveness and reliability in rescuing stress-sensitive species.

Enhancing habitat-former tolerance to novel climatic conditions

Genotype selection. Persistence of target habitat-former populations can be enhanced by selecting stress-tolerant genotypes. Genetic variation in traits relevant under global change seems high in
 315 coastal biota [67] and novel quantitative genetic analyses can provide accurate estimates of persistence probability of wild populations [68]. High genetic variation occurs among populations

with reduced gene flow, but also within the same population. For example, resilience to heat-waves differs between shallow and deep genotypes of the same populations of the seagrass *Posidonia oceanica* [39]. If this is caused by inherited genetic adaptation rather than acclimation to different developmental depths, then assisted relocation of such stress-tolerant genotypes - reared either in the lab or in the field - could rescue declining populations and enhance sub-population connectivity.

Synthetic biology. Although its application in the field of conservation is still in its infancy, synthetic biology is moving fast and may represent a strategic tool under future climates if accompanied by thorough risk-assessment and complying with environmental ethics [69].

Organisms have been genetically modified to enhance their resistance to biotic (e.g. disease) and abiotic (e.g. drought, salinity, heat) stressors, both in terrestrial and marine environments [70,71]. Gene editing of a single habitat-forming species may indirectly enhance the persistence of an entire suite of stress-susceptible species under adverse climates. The molecular basis for tolerance to environmental stress has been identified in key habitat-forming species, such as oysters and corals [72,73]. New genome editing techniques, such as CRISPR/Cas9, may rapidly advance this field.

Assisted evolution. Tolerance to stress can be enhanced through human-assisted acceleration of natural processes [74]. Short-term variance in biotic or abiotic pressures is critical to build stress tolerance [75]. For example, rapid fluctuations between benign and severe conditions accelerated adaptation to warming in the marine diatom *Thalassiosira pseudonana* since population size expansion during favorable periods increased the probability of fixing beneficial mutations [76]. Thus, controlled alternation of high and low stress phases in mesocosms - climate incubators - may act as an accelerator for adaptation to climate change, as high stress phases cause selective mortality of sensitive genotypes while stress relaxation phases allow surviving genotypes to recover and, possibly, reproduce [77].

The microbiome. Microbial symbionts influence host physiology, behavior and resistance to disease [78]. High genetic diversity and fast generational turn-over of symbionts can allow rapid adaptation to novel climatic conditions, potentially raising host fitness [79]. Laboratory thermal

selection could expand the temperature tolerance range of the coral-dinoflagellate *Symbiodinium* after ~80 asexual generations, corresponding to just 2.5 years [80]. Although the mechanisms
 345 regulating property transfer from the microbiome to the host (i.e. emergence of stress tolerance at the holobiont level) are yet to be fully understood, assisted microbiome evolution might be a formidable tool for raising habitat-former tolerance to novel climatic conditions.

Enhancing habitat-former population traits under novel climatic conditions

350 **Conservation biology.** By drawing on conservation and restoration knowledge, population viability of potential climate rescuers can be actively sustained (Fig. 3). Habitat-former population size and resilience can be enhanced by supporting connectivity through protection of source populations, restoration of natural migration corridors or the creation of new ones [81]. In some cases, managed relocation (or assisted migration) of habitat-formers at strategic sites might enhance connectivity
 355 among their populations, as well as among populations of beneficiary species. Likewise, herbivore release from predation can result in the overgrazing of habitat-forming macrophytes and trophic cascade restoration could be necessary to foster their persistence [82].

Mitigation of other anthropogenic stressors. Control of local/regional anthropogenic perturbations potentially exacerbating the impact of climate stressors will likely enhance habitat-
 360 former population resilience to climate and non-climate stressors [66,83]. For example, removal of excess nutrients enhances the tolerance of canopy-forming macroalgae to increased temperature [83].

Biodiversity. A large body of literature suggests a positive relationship between biodiversity and both resilience and temporal stability [84]. Thus, promoting multi-species assemblages of habitat-
 365 formers that are, to some degree, functionally interchangeable, may increase the reliability of their positive effects on other species under changing environmental conditions. In addition, greater micro-habitat availability in multi-species assemblages of habitat-formers may enhance the coexistence among beneficiary species and, hence, broaden the number of species sheltered from

adverse climatic conditions [84]. When desirable, the formation and maintenance of multi-species
 370 assemblages could be pursued through active control of competitively dominant species that would
 otherwise form mono-specific stands or through the seeding of subordinate species. Similar actions
 could be implemented to enhance genotype diversity, although they would require better
 understanding of competitive hierarchies between clonal genotypes.

Eco-engineering. Maritime infrastructures, off-shore installations and hard coastal-defences
 375 (breakwaters, seawalls) significantly change species distributions and ecological connectivity [85].
 Eco-engineering designs of artificial habitats including conservation or restoration objectives have
 the potential to turn these changes into an opportunity to sustain climate rescuer populations by
 supplying suitable habitat or providing new dispersal routes facilitating their migrations and that of
 beneficiary species. As previously demonstrated in the fields of restoration and conservation
 380 [16,17], engineering man-made structures for sustaining target habitat-forming species would be
 sufficient for attracting a suite of facultative and obligate associated species and represents,
 therefore, a cost-effective approach.

Non-native species. Where native habitat-formers are lacking, non-native species might be
 considered as alternative climate rescuers as they may revitalize functionalities that would be
 385 otherwise lost, including the support of diverse communities and the provision of climate refuges.
 The use of non-native species in conservation is still highly debated, but, in extreme cases, they
 may be the only chance of avoiding massive species loss when key habitat-formers decline due to
 global and regional human-driven changes (Box 2).

390 **Box 2. The role of non-native species as climate rescuers.** The view that all non-native species
 represent a threat to native biodiversity has been challenged on the grounds that some of them cause
 no harm and can contribute to achieve conservation and restoration goals [86,87].
 Climate change is predicted to foster invasions via enhanced propagule dispersal and decreased
 biotic resistance of native communities [86,88]. In addition, poleward shifts of coastal species have

395 been documented throughout the globe [88]. By virtue of their better adaptation to novel climate
 conditions, non-native species may be the primary cause of native species decline or local
 extinction. On the other hand, non-natives may replace natives when they decline as a consequence
 of other anthropogenic stressors. Although the effects of non-native habitat-formers on marine
 biodiversity are often complex and variable [89,90], there are examples of non-native species
 400 compensating, to some extent, for native habitat-former loss. For example, in areas of Chesapeake
 Bay where native eelgrass beds have retreated, the macroalga, *Gracilaria vermiculophylla*, provides
 suitable habitat for the native blue crab, *Callinectes sapidus*, a highly valued recreational and
 commercial species [91]. Positive effects of non-native habitat-formers can scale up to whole
 communities and influence ecosystem functioning. For example, long-term bioirrigation by the non-
 405 native polychaetes *Marenzelleria* spp. alleviates soft-sediment hypoxia in the Baltic Sea [92].
 Likewise, the non-native seaweed *Sargassum muticum* confers benthic assemblages greater
 resistance to warming and acidification than native macroalgal canopies [93].
 Of course, the benefits and risks of using non-native species as climate rescuers do not differ from
 those already described for restoration or conservation practice [94]. Many aspects of biological
 410 invasions, including their perception and management, are still highly controversial [95,96]. By no
 means, do we negate the capacity of non-native species to alter native biodiversity and to impair
 ecosystem functioning; rather, we suggest that their potential to rescue native species from changing
 climates should be not discarded a priori, but benefits and risks fully evaluated on a case-by-case
 basis.

415 **Concluding remarks**

Amelioration of physical stress by habitat-formers sustains species persistence in harsh
 environments [14,15]. This service might become increasingly important under future climates. The
 potential of habitat-formers to act as climate rescuers relies on their ability to maintain key
 420 individual and population traits in the face of climate changes. Likewise, the strength of rescuing

effects depends upon source-sink dynamics and the interplay of stabilizing and destabilizing forces regulating the co-existence between the benefactor and the beneficiaries, as well as among beneficiaries. Thus, current ability to ameliorate environmental conditions is not sufficient in itself to make a habitat-former a climate rescuer species. Nonetheless, some habitat-forming species

425 display the right individual and population traits (Box 3). Drawing from different ecological and biological disciplines, a series of management actions can sustain the strength and reliability of their climate rescuing effects. Within a multi-disciplinary framework (Fig. 3), understanding how biogenic habitats influence evolutionary adaptation of beneficiary species to changing conditions and their ability to track suitable climates should be considered a priority. Developing the concept

430 of sustaining habitat-former populations as a nature-based solution to climate change will likely depend on our ability and willingness to address ethical issues in modern conservation, such as those related to the use of synthetic biology, non-native species, assisted species evolution and species relocation. Finally, the general features of one or a few species that reduce climate-driven abiotic stress for other species that we describe in coastal systems are likely to be found also in

435 other types of ecosystems. For example, heat tolerance of freshwater gastropods is lowered in hypoxic conditions [97] and may be sustained by macrophyte oxygen production. In high-alpine systems, some cushion plants mitigate the effects of warming on native grasses [9]. Likewise, during drought events, canopy-forming mosses enhance the survival of smaller mosses and hepatics in their understory [98]. Thus, the broad conclusions we derive for coastal ecosystems under climate

440 change may also apply to other ecosystems.

Box 3. Examples of potential climate rescuers

Climate rescuer on the sand

445 Sea cucumbers play an important role in coastal environments since they bioturbate sediments and recycle nutrients, sustaining the diversity and functioning of benthic communities [99]. The sea

cucumber *Holothuria scabra* (the ‘sandfish’; Fig. 1F) is distributed throughout the Indo-Pacific region, between 30° N and 30° S of latitude. It is an active burrower and enhances sediment oxygenation, buffering negative effects of hypoxia caused by eutrophication and warming [33]. In addition, it can foster seagrass growth and productivity via re-mineralization of nutrients and/or their release from sediment pore water [99], potentially triggering a facilitation cascade. This species is cultured and it seems able to rapidly adapt to variable environmental conditions (e.g. salinity, temperature) through behavioral and molecular mechanisms [100,101]. For instance, in aquaculture facilities, extreme water temperatures, exceeding 31° C, caused no mortality of juveniles and, indeed, fostered their growth [102]. Finally, the entire mitochondrial genome of this species has been sequenced [103]. For the reasons above, this species may offer a nature-based solution for alleviating the impact of temperature-driven hypoxia.

Climate rescuer on the rocks

The brown macroalga *Fucus vesiculosus* (Fig. 1C) occupies wide ecological and geographical ranges. Presently, it spans latitudes from above 70° N (Norway) to near 30° N (Morocco) withstanding, at low tide, extreme freezing (e.g., Labrador Sea), extreme heat (e.g., above 40° C in Iberia) and variable salinities (estuaries, the Baltic Sea). It can function as climate rescuer for taxa beyond the southern limits of most intertidal fucoid seaweeds of the NE Atlantic, which can be vertically compressed and geographically restricted beyond the NW Iberian climate refugium [104]. In contrast, *F. vesiculosus* extends further south, persisting in more extreme conditions. Although it suffered the loss of many populations of a southern genetic lineage [105], reciprocal transplants showed that populations that persisted from this southern lineage have better adaptive traits for their habitat [106]. In this species, the costs of thermal stress to cellular metabolism (recorded as molecular heat shock response) can be escaped when high temperatures co-occur with rapid extreme desiccation [36]. Producing large quantities of recruits of *F. vesiculosus* is a standard procedure because this species has been for decades widely used as a model in developmental biology, reproductive ecology, ecophysiology, including in experimental field outplants [107].

475

Because the species is easily propagated and the southern populations have the capacity to withstand heat stress and maintain large canopies in areas where few other large intertidal canopies exist, this species may offer a nature-based solution for alleviating the impact of multiple stressors on intertidal community diversity and abundance, along its warm range limits.

Acknowledgements

480

Ideas and concepts presented in this paper were developed during the foresight workshop POSTCLIMA, funded by EuroMarine (European Marine Research Network; <http://euromarinenetwork.eu/>). We sincerely thank three anonymous reviewers for providing comments and constructive criticism on an earlier draft. F.B. wish to thank the University of Pisa for providing access to facilities during the workshop.

485

References

- 490 1. Pecl GT, Araújo MB, Bell JD, Blanchard J, Bonebrake TC, Chen I-C, et al. Biodiversity
redistribution under climate change: Impacts on ecosystems and human well-being. *Science*. 2017;
355(6332): eaai9214. doi: 10.1126/science.aai9214.
2. Scheffers BR, De Meester L, Bridge TCL, Hoffmann AA, Pandolfi JM, Corlett RT, et al. The
495 broad footprint of climate change from genes to biomes to people. *Science*. 2016; 354(6313):
aaf7671. doi: 10.1126/science.aaf7671.
3. Somero GN. The physiology of climate change: how potentials for acclimatization and genetic
adaptation will determine ‘winners’ and ‘losers’. *J Exp Biol* 2010; 213(6): 912-20. doi:
10.1242/jeb.037473.
- 500 4. Ockendon N, Baker DJ, Carr JA, White EC, Almond REA, Amano T, et al. Mechanisms
underpinning climatic impacts on natural populations: altered species interactions are more
important than direct effects. *Global Change Biol*. 2014; 20(7): 2221-9. doi: 10.1111/gcb.12559.
- 505 5. Cavieres LA, Brooker RW, Butterfield BJ, Cook BJ, Kikvidze Z, Lortie CJ, et al. Facilitative
plant interactions and climate simultaneously drive alpine plant diversity. *Ecol Lett*. 2014; 17(2):
193-202. doi: 10.1111/ele.12217.
6. Wisz MS, Pottier J, Kissling WD, Pellissier L, Lenoir J, Damgaard CF, et al. The role of biotic
510 interactions in shaping distributions and realised assemblages of species: implications for species
distribution modelling. *Biol Rev*. 2013; 88(1): 15-30. doi: 10.1111/j.1469-185X.2012.00235.x.
7. Anthelme F, Cavieres LA, Dangles O. Facilitation among plants in alpine environments in the
face of climate change. *Front Plant Sci*. 2014; 5(387). doi: 10.3389/fpls.2014.00387.
- 515 8. Brooker RW, Maestre FT, Callaway RM, Lortie CL, Cavieres LA, Kunstler G, et al. Facilitation
in plant communities: the past, the present, and the future. *J Ecol*. 2008; 96(1): 18-34. doi:
10.1111/j.1365-2745.2007.01295.x.
- 520 9. Cavieres LA, Sierra-Almeida A. Facilitative interactions do not wane with warming at high
elevations in the Andes. *Oecologia*. 2012; 170(2): 575-84. doi: 10.1007/s00442-012-2316-x.
10. Jurgens LJ, Gaylord B. Edge effects reverse facilitation by a widespread foundation species. *Sci*
Rep. 2016; 6: 37573. doi: 10.1038/srep37573.
- 525 11. Angelini C, Griffin JN, van de Koppel J, Lamers LPM, Smolders AJP, Derksen-Hooijberg M, et
al. A keystone mutualism underpins resilience of a coastal ecosystem to drought. *Nat Commun*.
2016; 7: 12473. doi: 10.1038/ncomms12473.
12. Bulleri F. Facilitation research in marine systems: state of the art, emerging patterns and insights
for future developments. *J Ecol*. 2009; 97(6): 1121-30. doi: 10.1111/j.1365-2745.2009.01567.x.
- 530 13. Donadi S, van der Heide T, van der Zee EM, Eklöf JS, de Koppel Jv, Weerman EJ, et al. Cross-
habitat interactions among bivalve species control community structure on intertidal flats. *Ecology*.
2013; 94(2): 489-98. doi: 10.1890/12-0048.1.

- 535 14. He Q, Bertness MD, Altieri AH. Global shifts towards positive species interactions with increasing environmental stress. *Ecol Lett.* 2013; 16(5): 695-706. doi: 10.1111/ele.12080.
15. Bertness MD, Callaway R. Positive interactions in communities. *Trends Ecol Evol.* 1994; 9(5): 191-3. doi: 10.1016/0169-5347(94)90088-4.
- 540 16. Gómez-Aparicio L, Zamora R, M.Gómez J, A.Hódar J, Castro J, Baraza E. Applying plant facilitation to forest restoration: a meta-analysis of the use of shrubs as nurse plants. *Ecol Appl.* 2004; 14(4): 1128-38. doi: doi:10.1890/03-5084.
- 545 17. Halpern BS, Silliman BR, Olden JD, Bruno JP, Bertness MD. Incorporating positive interactions in aquatic restoration and conservation. *Front Ecol Environ.* 2007; 5(3): 153-60. doi: 10.1890/1540-9295(2007)5[153:ipiiar]2.0.co;2.
- 550 18. Byers JE, Cuddington K, Jones CG, Talley TS, Hastings A, Lambrinos JG, et al. Using ecosystem engineers to restore ecological systems. *Trends Ecol Evol.* 2006; 21(9): 493-500. doi: 10.1016/j.tree.2006.06.002.
19. Valiente-Banuet A, Rumebe AV, Verdú M, Callaway RM. Modern Quaternary plant lineages promote diversity through facilitation of ancient Tertiary lineages. *Proc Natl Acad Sci U S A.* 2006; 103(45): 16812-17. doi: 10.1073/pnas.0604933103.
- 555 20. Meysman FJR, Middelburg JJ, Heip CHR. Bioturbation: a fresh look at Darwin's last idea. *Trends Ecol Evol.* 2006; 21(12): 688-95. doi: 10.1016/j.tree.2006.08.002.
- 560 21. Koppel Jvd, Heide Tvd, Altieri AH, Eriksson BK, Bouma TJ, Olff H, et al. Long-distance interactions regulate the structure and resilience of coastal ecosystems. *Annu Rev Mar Sci.* 2015; 7(1): 139-58. doi: 10.1146/annurev-marine-010814-015805.
- 565 22. Altieri AH, van de Koppel J, Foundation species in marine environments. In: Bertness MD, Bruno JF, Silliman BR, Stachowicz JJ editors. *Marine Community Ecology and Conservation.* Sinauer Associates, Inc., Sunderland; 2014. pp. 37-56.
23. Solan M, Cardinale BJ, Downing AL, Engelhardt KAM, Ruesink JL, Srivastava DS. Extinction and ecosystem function in the marine benthos. *Science.* 2004; 306(5699): 1177-80. doi: 10.1126/science.1103960.
- 570 24. Leonard GH. Latitudinal variation in species interactions: a test in the New England rocky intertidal zone. *Ecology.* 2000; 81(4): 1015-30. doi: 10.1890/0012-9658(2000)081[1015:lvisia]2.0.co;2.
- 575 25. Silliman BR, Bertness MD, Altieri AH, Griffin JN, Bazterrica MC, Hidalgo FJ, et al. Whole-community facilitation regulates biodiversity on Patagonian rocky shores. *Plos One.* 2011; 6(10): e24502. doi: 10.1371/journal.pone.0024502.
- 580 26. Keppel G, Van Niel KP, Wardell-Johnson GW, Yates CJ, Byrne M, Mucina L, et al. Refugia: identifying and understanding safe havens for biodiversity under climate change. *Global Ecol Biogeogr.* 2012; 21(4): 393-404. doi: 10.1111/j.1466-8238.2011.00686.x.
- 585 27. Hendriks IE, Olsen YS, Ramajo L, Basso L, Steckbauer A, Moore TS, et al. Photosynthetic activity buffers ocean acidification in seagrass meadows. *Biogeosciences.* 2014; 11(2): 333-46. doi: 10.5194/bg-11-333-2014.

28. Wahl M, Schneider Covachã S, Saderne V, Hiebenthal C, Müller JD, Pansch C, et al. Macroalgae may mitigate ocean acidification effects on mussel calcification by increasing pH and its fluctuations. *Limnol. Oceanogr.* doi: 10.1002/lno.10608.
- 590 29. Unsworth RKF, Collier CJ, Henderson GM, McKenzie LJ. Tropical seagrass meadows modify seawater carbon chemistry: implications for coral reefs impacted by ocean acidification. *Environ Res Lett.* 2012; 7(2): 024026. doi:10.1088/1748-9326/7/2/024026.
- 595 30. Pinsky ML, Guannel G, Arkema KK. Quantifying wave attenuation to inform coastal habitat conservation. *Ecosphere.* 2013; 4(8): 1-16. doi: 10.1890/es13-00080.1.
- 600 31. Bennett S, Wernberg T, de Bettignies T, Kendrick GA, Anderson RJ, Bolton JJ, et al. Canopy interactions and physical stress gradients in subtidal communities. *Ecol Lett.* 2015; 18(7): 677-86. doi: 10.1111/ele.12446.
- 605 32. Gutiérrez JL, Jones CG, Byers JE, Arkema KK, Berkenbusch K, Commito JA, et al. Physical ecosystem engineers and the functioning of estuaries and coasts. In: Wolanski E, McLusky D editors. *Treatise on Estuarine and Coastal Science.* Elsevier, Amsterdam, The Netherlands; 2011. pp. 53-81.
33. Lee S, Ford AK, Mangubhai S, Wild C, Ferse SCA. Effects of sandfish (*Holothuria scabra*) removal on shallow-water sediments in Fiji. *PeerJ.* 2018; 6: e4773. doi: 10.7717/peerj.4773.
- 610 34. Hiddink JG, Burrows MT, Molinos JG. Temperature tracking by North Sea benthic invertebrates in response to climate change. *Global Change Biol.* 2015; 21(1): 117-29. doi: 10.1111/gcb.12726.
- 615 35. Jurgens LJ, Gaylord B. Physical effects of habitat-forming species override latitudinal trends in temperature. *Ecol Lett.* 21(2): 190-6. doi: 10.1111/ele.12881.
- 620 36. Mota CF, Engelen AH, Serrao EA, Pearson GA. Some don't like it hot: microhabitat-dependent thermal and water stresses in a trailing edge population. *Funct Ecol.* 2015; 29(5): 640-9. doi: 10.1111/1365-2435.12373.
37. Bulleri F, Bruno JF, Silliman BR, Stachowicz JJ. Facilitation and the niche: implications for coexistence, range shifts and ecosystem functioning. *Funct Ecol.* 2016; 30(1): 70-8. doi: 10.1111/1365-2435.12528.
- 625 38. Torda G, Donelson JM, Aranda M, Barshis DJ, Bay L, Berumen ML, et al. Rapid adaptive responses to climate change in corals. *Nat Clim Change.* 2017; 7(9): 627-36. doi: 10.1038/nclimate3374.
- 630 39. Marín-Guirao L, Entrambasaguas L, Dattolo E, Ruiz JM, Procaccini G. Molecular mechanisms behind the physiological resistance to intense transient warming in an iconic marine plant. *Front Plant Sci.* 2017; 8(1142). doi: 10.3389/fpls.2017.01142.
- 635 40. Donelson JM, Munday PL, McCormick MI, Nilsson GE. Acclimation to predicted ocean warming through developmental plasticity in a tropical reef fish. *Global Change Biol.* 2011; 17(4): 1712-9. doi: 10.1111/j.1365-2486.2010.02339.x.

41. Putnam HM, Gates RD. Preconditioning in the reef-building coral *Pocillopora damicornis* and the potential for trans-generational acclimatization in coral larvae under future climate change conditions. *J Exp Biol.* 2015; 218(15): 2365-72. doi: 10.1242/jeb.123018.
- 640 42. Stillwell RC, Fox CW. Geographic variation in body size, sexual size dimorphism and fitness components of a seed beetle: local adaptation versus phenotypic plasticity. *Oikos.* 2009; 118(5): 703-12. doi: 10.1111/j.1600-0706.2008.17327.x.
- 645 43. Soliveres S, Smit C, Maestre FT. Moving forward on facilitation research: response to changing environments and effects on the diversity, functioning and evolution of plant communities. *Biol Rev.* 2015; 90(1): 297-313. doi: 10.1111/brv.12110.
- 650 44. Sanford E, Kelly MW. Local adaptation in marine invertebrates. *Annu Rev Mar Sci.* 2011; 3: 509-35. doi: 10.1146/annurev-marine-120709-142756
- 655 45. Schmidt PS, Rand DM. Adaptive maintenance of genetic polymorphism in an intertidal barnacle: Habitat- and life-stage-specific survivorship of *Mpi* genotypes. *Evolution.* 2001; 55(7): 1336-44.
46. Castellanos MC, Donat-Caerols S, Gonzalez-Martinez SC, Verdu M. Can facilitation influence the spatial genetics of the beneficiary plant population? *J Ecol.* 2014; 102(5): 1214-21. doi: 10.1111/1365-2745.12278.
- 660 47. Liancourt P, Choler P, Gross N, Thibert-Plante X, Tielbörger K. How facilitation may interfere with ecological speciation. *Internat. J Ecol.* 2012; 725487. doi: 10.1155/2012/725487.
48. Diekmann OE, Serrao EA. Range-edge genetic diversity: locally poor extant southern patches maintain a regionally diverse hotspot in the seagrass *Zostera marina*. *Mol Ecol.* 2012; 21(7): 1647-57. doi: 10.1111/j.1365-294X.2012.05500.x.
- 665 49. Alexander JM, Diez JM, Hart SP, Levine JM. When climate reshuffles competitors: A call for experimental macroecology. *Trends Ecol Evol* 2016; 31(11): 831-41. doi: 10.1016/j.tree.2016.08.003.
- 670 50. Michalet R, Brooker RW, Cavieres LA, Kikvidze Z, Lortie CJ, Pugnaire FI, et al. Do biotic interactions shape both sides of the humped-back model of species richness in plant communities? *Ecol Lett.* 2006; 9(7): 767-73. doi: 10.1111/j.1461-0248.2006.00935.x.
- 675 51. Guo HY, Zhang YH, Lan ZJ, Pennings SC. Biotic interactions mediate the expansion of black mangrove (*Avicennia germinans*) into salt marshes under climate change. *Global Change Biol.* 2013; 19(9): 2765-74. doi: 10.1111/gcb.12221.
- 680 52. Reise K, Buschbaum C, Büttger H, Wegner KM. Invading oysters and native mussels: from hostile takeover to compatible bedfellows. *Ecosphere.* 2017; 8(9). doi: 10.1002/ecs2.1949.
53. Altieri AH, Silliman BR, Bertness MD. Hierarchical organization via a facilitation cascade in intertidal cordgrass bed communities. *Am Nat.* 2007; 169(2): 195-206. doi: 10.1086/510603.
- 685 54. Thomsen MS, Wernberg T, Altieri AH, Tuya F, Gulbransen D, McGlathery KJ, et al. Habitat cascades: The conceptual context and global relevance of facilitation cascades via habitat formation and modification. *Integr Comp Biol.* 2010; 50(2): 158-75. doi: 10.1093/icb/icq042.

55. van de Koppel J, van der Heide T, Altieri AH, Eriksson BK, Bouma TJ, Olff H, et al. Long-distance interactions regulate the structure and resilience of coastal ecosystems. *Annu Rev Mar Sci*. 2015; 7: 139-58. doi: 10.1146/annurev-marine-010814-015805.
56. Daufresne M, Lengfellner K, Sommer U. Global warming benefits the small in aquatic ecosystems. *Proc Natl Acad Sci U S A*. 2009; 106(31): 12788-93. doi: 10.1073/pnas.0902080106.
57. Garilli V, Rodolfo-Metalpa R, Scuderi D, Brusca L, Parrinello D, Rastrick SPS, et al. Physiological advantages of dwarfing in surviving extinctions in high-CO₂ oceans. *Nat Clim Change*. 2015; 5: 678. doi: 10.1038/nclimate2616.
58. Gaylord B, Hill TM, Sanford E, Lenz EA, Jacobs LA, Sato KN, et al. Functional impacts of ocean acidification in an ecologically critical foundation species. *J Exp Biol* 2011; 214(15): 2586-94. doi: 10.1242/jeb.055939.
59. Kordas RL, Dudgeon S, Storey S, Harley CDG. Intertidal community responses to field-based experimental warming. *Oikos*. 2015; 124(7): 888-98. doi: 10.1111/oik.00806.
60. Thomson JA, Burkholder DA, Heithaus MR, Fourqurean JW, Fraser MW, Statton J, et al. Extreme temperatures, foundation species, and abrupt ecosystem change: an example from an iconic seagrass ecosystem. *Global Change Biol*. 2015; 21(4): 1463-74. doi: 10.1111/gcb.12694.
61. Rilov G. Multi-species collapses at the warm edge of a warming sea. *Sci Rep*. 2016; 6: 36897. doi: 10.1038/srep36897.
62. Osland MJ, Day RH, Hall CT, Brumfield MD, Dugas JL, Jones WR. Mangrove expansion and contraction at a poleward range limit: climate extremes and land-ocean temperature gradients. *Ecology*. 2017; 98(1): 125-37. doi: 10.1002/ecy.1625.
63. Wernberg T, Bennett S, Babcock RC, de Bettignies T, Cure K, Depczynski M, et al. Climate-driven regime shift of a temperate marine ecosystem. *Science*. 2016; 353(6295): 169-72. doi: 10.1126/science.aad8745.
64. Gorman D, Turra A, Bergstrom ER, Horta PA. Population expansion of a tropical seagrass (*Halophila decipiens*) in the southwest Atlantic (Brazil). *Aquat Bot*. 2016; 132: 30-6. doi: 10.1016/j.aquabot.2016.04.002.
65. Bouma TJ, Friedrichs M, Van Wesenbeeck BK, Temmerman S, Graf G, Herman PMJ. Density-dependent linkage of scale-dependent feedbacks: a flume study on the intertidal macrophyte *Spartina anglica*. *Oikos*. 2009; 118(2): 260-8. doi: 10.1111/j.1600-0706.2008.16892.x.
66. O'Leary JK, Micheli F, Airoidi L, Boch C, De Leo G, Elahi R, et al. The resilience of marine ecosystems to climatic disturbances. *Bioscience*. 2017; 67(3): 208-20. doi: 10.1093/biosci/biw161.
67. Reusch TBH. Climate change in the oceans: evolutionary versus phenotypically plastic responses of marine animals and plants. *Evol Appl*. 2014; 7(1): 104-22. doi: 10.1111/eva.12109.
68. Gienapp P, Fior S, Guillaume F, Lasky JR, Sork VL, Csilléry K. Genomic quantitative genetics to study evolution in the wild. *Trends Ecol Evol*. 32(12): 897-908. doi: 10.1016/j.tree.2017.09.004.

69. Piaggio AJ, Segelbacher G, Seddon PJ, Alphey L, Bennett EL, Carlson RH, et al. Is it time for synthetic biodiversity conservation? *Trends Ecol Evol* 32(2): 97-107. doi: 10.1016/j.tree.2016.10.016.
70. Harfouche A, Meilan R, Altman A. Tree genetic engineering and applications to sustainable forestry and biomass production. *Trends Biotechnol.* 2011; 29(1): 9-17. doi: 10.1016/j.tibtech.2010.09.003.
71. Levin RA, Voolstra CR, Agrawal S, Steinberg PD, Suggett DJ, van Oppen MJH. Engineering strategies to decode and enhance the genomes of coral symbionts. *Front Microbiol* 2017; 8: 1220. doi:10.3389/fmicb.2017.01220.
72. Barshis DJ, Ladner JT, Oliver TA, Seneca FO, Traylor-Knowles N, Palumbi SR. Genomic basis for coral resilience to climate change. *Proc Natl Acad Sci U S A.* 2013; 110(4): 1387-92. doi: 10.1073/pnas.1210224110.
73. Zhang GF, Li L, Meng J, Qi HG, Qu T, Xu F, et al. Molecular basis for adaptation of oysters to stressful marine intertidal environments. *Annu Rev Anim Biosci.* 2016; 4: 357-81. doi: 10.1146/annurev-animal-022114-110903.
74. van Oppen MJH, Oliver JK, Putnam HM, Gates RD. Building coral reef resilience through assisted evolution. *Proc Natl Acad Sci U S A.* 2015; 112(8): 2307-13. doi: 10.1073/pnas.1422301112.
75. Carpenter SR, Brock WA, Folke C, van Nes EH, Scheffer M. Allowing variance may enlarge the safe operating space for exploited ecosystems. *Proc Natl Acad Sci U S A.* 2015; 112(46): 14384-9. doi: 10.1073/pnas.1511804112.
76. Schaum CE, Buckling A, Smirnoff N, Studholme DJ, Yvon-Durocher G. Environmental fluctuations accelerate molecular evolution of thermal tolerance in a marine diatom. *Nat Comm* 2018; 9(1): 1719. doi: 10.1038/s41467-018-03906-5.
77. Wahl M, Buchholz B, Winde V, Golomb D, Guy-Haim T, Muller J, et al. A mesocosm concept for the simulation of near-natural shallow underwater climates: The Kiel Outdoor Benthocosms (KOB). *Limnol Oceanogr-Meth.* 2015; 13(11): 651-63. doi: 10.1002/lom3.10055.
78. Costello EK, Stagaman K, Dethlefsen L, Bohannan BJM, Relman DA. The application of ecological theory toward an understanding of the human microbiome. *Science.* 2012; 336(6086): 1255-62. doi: 10.1126/science.1224203.
79. Zilber-Rosenberg I, Rosenberg E. Role of microorganisms in the evolution of animals and plants: the hologenome theory of evolution. *FEMS Microbiol Rev.* 2008; 32(5): 723-35. doi: 10.1111/j.1574-6976.2008.00123.x.
80. Chakravarti LJ, Beltran VH, van Oppen MJH. Rapid thermal adaptation in photosymbionts of reef-building corals. *Global Change Biol.* 2017; 23(11): 4675-88. doi: 10.1111/gcb.13702
81. Rudnick DA, Ryan SJ, Beier P, Cushman SA, Dieffenbach F, Epps CW, et al. The role of landscape connectivity in planning and implementing conservation and restoration priorities. *Issues Ecol* 2012; 16:1-23.

82. Östman Ö, Eklöf J, Eriksson BK, Olsson J, Moksnes P-O, Bergström U. Top-down control as important as nutrient enrichment for eutrophication effects in North Atlantic coastal ecosystems. *J Appl Ecol.* 2016; 53(4): 1138-47. doi: 10.1111/1365-2664.12654.
83. Strain EMA, van Belzen J, van Dalen J, Bouma TJ, Airolidi L. Management of local stressors can improve the resilience of marine canopy algae to global stressors. *Plos One.* 2015; 10(3): e0120837. doi: 10.1371/journal.pone.0120837.
84. Cardinale BJ, Duffy JE, Gonzalez A, Hooper DU, Perrings C, Venail P, et al. Biodiversity loss and its impact on humanity. *Nature.* 2012; 486: 59-67. doi: 10.1038/nature11148.
85. Bishop MJ, Mayer-Pinto M, Airolidi L, Firth LB, Morris RL, Loke LHL, et al. Effects of ocean sprawl on ecological connectivity: impacts and solutions. *J Exp Mar Biol Ecol.* 2017; 492: 7-30. doi: 10.1016/j.jembe.2017.01.021.
86. Walther GR, Roques A, Hulme PE, Sykes MT, Pysek P, Kuhn I, et al. Alien species in a warmer world: risks and opportunities. *Trends Ecol Evol.* 2009; 24(12): 686-93. doi: 10.1016/j.tree.2009.06.008.
87. Schlaepfer MA. Do non-native species contribute to biodiversity? *PLoS Biol.* 2018; 16(4): e2005568. doi: 10.1371/journal.pbio.2005568.
88. Sorte CJB, Williams SL, Carlton JT. Marine range shifts and species introductions: comparative spread rates and community impacts. *Global Ecol Biogeogr* 2010; 19(3): 303-16. doi: 10.1111/j.1466-8238.2009.00519.
89. Maggi E, Benedetti-Cecchi L, Castelli A, Chatzinikolaou E, Crowe TP, Ghedini G, et al. Ecological impacts of invading seaweeds: a meta-analysis of their effects at different trophic levels. *Divers Distrib.* 2015; 21(1): 1-12. doi: 10.1111/ddi.12264.
90. Guy-Haim T, Lyons DA, Kotta J, Ojaveer H, Queiros AM, Chatzinikolaou E, et al. Diverse effects of invasive ecosystem engineers on marine biodiversity and ecosystem functions: A global review and meta-analysis. *Global Change Biol.* 2018; 24(3): 906-24. doi: 10.1111/gcb.14007.
91. Johnston CA, Lipcius RN. Exotic macroalga *Gracilaria vermiculophylla* provides superior nursery habitat for native blue crab in Chesapeake Bay. *Mar Ecol Prog Ser.* 2012; 467: 137-46. doi: 10.3354/meps09935.
92. Norkko J, Reed DC, Timmermann K, Norkko A, Gustafsson BG, Bonsdorff E, et al. A welcome can of worms? Hypoxia mitigation by an invasive species. *Global Change Biol.* 2012; 18(2): 422-34. doi: 10.1111/j.1365-2486.2011.02513.x.
93. Olabarria C, Arenas F, Viejo RM, Gestoso I, Vaz-Pinto F, Incera M, et al. Response of macroalgal assemblages from rockpools to climate change: effects of persistent increase in temperature and CO₂. *Oikos.* 2013; 122(7): 1065-79. doi: 10.1111/j.1600-0706.2012.20825.x.
94. Schlaepfer MA, Sax DF, Olden JD. The potential conservation value of non-native species. *Conserv Biol.* 2011; 25(3): 428-37. doi: 10.1111/j.1523-1739.2010.01646.x.
95. Davis MA, Chew MK, Hobbs RJ, Lugo AE, Ewel JJ, Vermeij GJ, et al. Don't judge species on their origins. *Nature.* 2011; 474: 153. doi: 10.1038/474153a.

96. Simberloff D. Non-natives: 141 scientists object. *Nature*. 2011; 475: 36. doi: 10.1038/475036a.
- 840 97. Koopman KR, Collas FPL, van der Velde G, Verberk W. Oxygen can limit heat tolerance in freshwater gastropods: differences between gill and lung breathers. *Hydrobiologia*. 2016; 763(1): 301-12. doi: 10.1007/s10750-015-2386-y.
- 845 98. Mulder CPH, Uliassi DD, Doak DF. Physical stress and diversity-productivity relationships: The role of positive interactions. *Proc Natl Acad Sci U S A*. 2001; 98(12): 6704-8. doi: 10.1073/pnas.111055298.
99. Purcell SW, Conand C, Uthicke S, Byrne M. Ecological roles of exploited sea cucumbers. *Oceanogr Mar Biol Annu Rev*. 2016; 54: 367-86.
- 850 100. Kamyab E, Kuhnhold H, Novais SC, Alves LMF, Indriana L, Kunzmann A, et al. Effects of thermal stress on the immune and oxidative stress responses of juvenile sea cucumber *Holothuria scabra*. *J Comp Physiol B*. 2017; 187(1): 51-61. doi: 10.1007/s00360-016-1015-z.
- 855 101. Mercier A, Battaglione SC, Hamel JF. Daily burrowing cycle and feeding activity of juvenile sea cucumbers *Holothuria scabra* in response to environmental factors. *J Exp Mar Biol Ecol*. 1999; 239(1): 125-56. doi: 10.1016/s0022-0981(99)00034-9.
- 860 102. Lavitra T, Fohy N, Gestin PG, Rasolofonirina R, Eeckhaut I. Effect of water temperature on the survival and growth of endobenthic *Holothuria scabra* (Echinodermata: Holothuroidea) juveniles reared in outdoor ponds. *SPC Beche-de-Mer Inform Bull*. 2010; 30: 25-8.
- 865 103. Xia JJ, Ren CH, Yu ZH, Wu XY, Qian J, Hu CQ. Complete mitochondrial genome of the sandfish *Holothuria scabra* (Holothuroidea, Holothuriidae). *Mitochondrial DNA Part A*. 2016; 27(6): 4174-5. doi: 10.3109/19401736.2014.1003899.
- 870 104. Neiva J SE, Assis J, Pearson GA, Coyer JA, Olsen JL, et al. Climate oscillations, range shifts and phylogeographic patterns of North Atlantic Fucaceae. In: Hu Z-M, Fraser C, editors. *Seaweed Phylogeography. Adaptation and Evolution of Seaweeds under Environmental Change*. Springer, Netherlands; 2016. p. 279-308.
- 875 105. Nicastro KR, Zardi GI, Teixeira S, Neiva J, Serrao EA, Pearson GA. Shift happens: trailing edge contraction associated with recent warming trends threatens a distinct genetic lineage in the marine macroalga *Fucus vesiculosus*. *BMC Biol*. 2013; 11: 6. doi: 10.1186/1741-7007-11-6.
106. Saada G, Nicastro KR, Jacinto R, McQuaid CD, Serrao EA, Pearson GA, et al. Taking the heat: distinct vulnerability to thermal stress of central and threatened peripheral lineages of a marine macroalga. *Divers Distrib*. 2016; 22(10): 1060-8. doi: 10.1111/ddi.12474.
- 880 107. Ladah L, Bermudez R, Pearson G, Serrao E. Fertilization success and recruitment of dioecious and hermaphroditic furoid seaweeds with contrasting distributions near their southern limit. *Mar Ecol Prog Ser* 2003; 262: 173-83. doi: 10.3354/meps262173.





Figure 1






Figure 3