

# Extent and retreat history of the Barra Fan Ice Stream offshore western Scotland and northern Ireland during the last glaciation

Callard, Sarah Louise; Ó Cofaigh, C.; Benetti, Sara; Chiverrell, R.C.; Van Landeghem, Katrien; Saher, Margot; Gales, Jenny; Small, David; Clark, Chris; Livingstone, S.J.; Fabel, Derek; Moreton, Steven

#### **Quaternary Science Reviews**

DOI: 10.1016/j.quascirev.2018.10.002

Published: 01/12/2018

Peer reviewed version

Cyswllt i'r cyhoeddiad / Link to publication

Dyfyniad o'r fersiwn a gyhoeddwyd / Citation for published version (APA): Callard, S. L., O Cofaigh, C., Benetti, S., Chiverrell, R. C., Van Landeghem, K., Saher, M., Gales, J., Small, D., Clark, C., Livingstone, S. J., Fabel, D., & Moreton, S. (2018). Extent and retreat history of the Barra Fan Ice Stream offshore western Scotland and northern Ireland during the last glaciation. Quaternary Science Reviews, 280-302. https://doi.org/10.1016/j.quascirev.2018.10.002

#### Hawliau Cyffredinol / General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

· Users may download and print one copy of any publication from the public portal for the purpose of private study or research.

- You may not further distribute the material or use it for any profit-making activity or commercial gain
   You may freely distribute the URL identifying the publication in the public portal ?

Take down policy If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

- 1 Extent and retreat history of the Barra Fan Ice Stream offshore western Scotland and
- 2 northern Ireland during the last glaciation.
- 3 S. Louise Callard<sup>1,\*</sup>, Colm Ó Cofaigh<sup>1</sup>, Sara Benetti<sup>2</sup>, Richard C. Chiverrell<sup>3</sup>, Katrien J.J. Van
- 4 Landeghem<sup>4</sup>, Margot H. Saher<sup>4</sup>, Jenny A. Gales<sup>5</sup>, David Small<sup>1</sup>, Chris D. Clark<sup>6</sup>, Stephen, J.
- 5 Livingstone<sup>6</sup>, Derek Fabel<sup>7</sup>, Steven G. Moreton<sup>8</sup>
- <sup>1</sup> Department of Geography, Durham University, Durham, DH1 3LE, UK
- <sup>7</sup> <sup>2</sup> School of Geography and Environmental Sciences, Ulster University, Coleraine, BT52 1SA,
- 8 Northern Ireland
- 9 <sup>3</sup>Department f Geography, University of Liverpool, Liverpool, UK
- <sup>4</sup> School of Ocean Sciences, Bangor University, Menai Bridge, UK
- <sup>5</sup>University of Plymouth, School of Biological and Marine Sciences, Drake Circus, Plymouth,
   PL4 8AA
- <sup>6</sup>Department of Geography, University of Sheffield, Sheffield, S10 2TN, UK
- <sup>7</sup>Scottish Universities Environmental Research Centre, University of Glasgow, East Kilbride,
   G75 0QF
- <sup>8</sup> Natural Environment Research Council, Radiocarbon facility, East Kilbride, Scotland, G75
   OQF, UK
- 18
- 19 Abstract
- 20 During the Last Glacial Maximum (LGM) the marine-terminating Barra Fan Ice Stream (BFIS), a major conduit of the British Irish Ice Sheet (BIIS), drained much of western Scotland 21 22 and northwest Ireland with ice streaming onto the continental shelf of the Malin Sea. The extent and retreat history of this ice stream across the shelf, until now, is not well known. In particular, 23 geochronological constraints on the history of this ice stream have thus far been restricted to 24 25 deep-sea cores or terrestrial cosmogenic nuclide dating onshore, with ages across the shelf absent. To understand the possible external forcing factors acting on this marine terminating 26 27 ice stream during retreat, improved geochronological constraint on its deglaciation is necessary. Here, we present new geophysical data, marine sediment cores and over forty 28 29 radiocarbon dates to provide important constraints on maximum extent of the BFIS, as well as the timing and pattern of retreat back across the Malin Shelf. Dated moraines and grounding-30 zone wedges (GZW) seen in seafloor sub-bottom profiles provide evidence that the BFIS 31 reached the Malin Shelf edge during the LGM and was at its maximum extent around 26.7 ka 32 BP. The presence of two sets of GZWs suggests that the style of retreat was episodic. The new 33 radiocarbon chronology shows that retreat from the shelf edge was underway by 25.9 ka BP, 34 with the majority of the continental shelf ice free by 23.2 ka BP, and that glacimarine conditions 35 were present in the Sea of Hebrides by 20.2 ka BP at the latest. Collectively, these results 36 37 indicate that the majority of the Malin Shelf was free of grounded ice by ~21.5-20 ka BP, which is up to 4,000 years earlier than previously reconstructed. We attribute this early deglaciation 38

to high relative sea level caused by glacial isostatic depression when the BIIS reached its maximum extent promoting ice shelf and grounding line instability. Two deep troughs, forming reverse bed slopes, aided the continued retreat of the BFIS. This suggests that local ice loading and bed morphology can be significant controls on the destabilisation of a marine-terminating ice stream and can override the influence of ocean and atmospheric temperatures.

44

Keywords: British-Irish Ice Sheet; glacimarine; Last Glacial Maximum; radiocarbon dating;
ice-sheet retreat; grounding-zone wedges.

47

# 48 **1. Introduction**

During the Last Glacial Maximum (LGM) defined here as ca. 26.5-19 ka (P.U. Clark et al., 49 2009) the British-Irish Ice Sheet (BIIS) extended to the continental shelf break offshore of 50 western Ireland and Scotland (Bradwell et al., 2008; Clark et al., 2012; Ó Cofaigh et al., 2012, 51 2016; Peters et al., 2015, 2016). Despite the global eustatic sea level being 134 m lower than 52 present (Lambeck et al., 2014), up to two thirds of the BIIS was marine influenced and was 53 drained by ice streams (Clark et al., 2012). It is therefore likely that the BIIS was sensitive to 54 both oceanic and atmospheric changes, both major forcing factors in deglaciation (Clark et al., 55 56 2012). The BIIS is therefore a good analogue for the West Antarctic Ice Sheet (WAIS) and Greenland Ice Sheet, which are currently undergoing rapid thinning and retreat of marine-based 57 58 sectors (Joughin et al., 2014; Mouginot et al., 2015). Predicting the impact of these changes, particularly with respect to sea level, requires modelling informed by empirical data on the 59 60 processes, rates and controls of palaeo-marine based ice stream retreat (Hindmarsh, 2018). Reconstructing the deglaciation of the BIIS will therefore provide valuable empirical data to 61 62 test and inform ice sheet models. Although, recent research offshore Britain and Ireland, using marine geophysical data, has significantly enhanced our understanding of the glacial 63 geomorphology of the shelf (e.g. Benetti et al., 2010; Dunlop et al., 2010; Ó Cofaigh et al., 64 2012), the extent and dynamic retreat of most of the marine-based sectors of the BIIS are still 65 not well constrained chronologically. 66

67

During the LGM, the Barra Fan Ice Stream (BFIS; Dove et al., 2015) drained at least 5-10 % of the BIIS and proxy evidence from deep water marine cores adjacent to the ice stream (Knutz et al., 2001) suggests a maximum extent reaching the shelf edge. This ice stream fed the Donegal Barra Fan (DBF), the largest glacimarine depocentre of the BIIS (Fig. 1). Irish and near-shore Scottish waters provide most of the direct evidence for glaciation of the Malin Shelf, 73 with the majority of the shelf remaining unmapped. There, streamlined bedforms and over-74 deepened troughs provide evidence of grounded, fast-flowing ice that streamed southwestwards from the Inner Hebrides and then westwards around the Outer Hebrides across the 75 shelf (Howe et al., 2012; Dove et al., 2015). Undated moraine complexes at the shelf edge mark 76 the western limit of the BFIS (Selby, 1989; Dunlop et al., 2010). A series of smaller recessional 77 78 moraines step back eastwards and become increasingly abundant on the inner shelf, marking 79 the pattern and direction of retreating ice (Dunlop et al., 2010; Dove et al., 2015, Small et al., 2016). Dating control for the timing of advance and retreat of the BFIS relies on ice rafted 80 81 debris (IRD) records. These records suggest that ice extended onto the Malin Shelf by 29 ka BP (Scourse et al., 2009) with grounded ice reaching the shelf edge by ~27 ka BP (Wilson et 82 al., 2002; Peck et al. 2006; Scourse et al., 2009). A marked reduction in IRD flux to the DBF 83 at 23 ka BP potentially represents the onset of deglaciation for this sector of the BIIS margin. 84 Recent terrestrial cosmogenic nuclide (TCN) dating on Tiree constrains deglaciation of much 85 of the mid and outer sector of the BFIS to  $20.6 \pm 1.2$  ka and complete deglaciation of the inner 86 sector to between  $20.6 \pm 1.2$  and  $17.5 \pm 1.0$  ka (Small et al., 2017). 87

88

Here, we combine new geophysical, sedimentary and radiocarbon data from across the Malin Shelf to reconstruct, for the first time, the deglacial history of the BFIS at and after the LGM. The aims of this paper are threefold: 1. to describe the glacial geomorphology, seismic stratigraphy, the lithology and geochronology of glacial sediments on the Malin Shelf; 2. to reconstruct spatially and temporally the BFIS at its maximum extent and during its retreat across the Malin Shelf, and 3. to discuss the underlying controls of BFIS dynamics in the wider context of the stability of the former BIIS.

96

# 97 2. Regional setting and background

The Malin Sea, known also as the Malin Shelf (Fig. 1), describes the continental shelf north of 98 Ireland and west of Scotland, and is adjacent to a major flow pathway for Atlantic Ocean 99 circulation (Ellett., 1979). The Slope Current transports warm saline Atlantic waters 100 northwards around 500 m water depth (Dooley and Crease, 1978). Not always confined to the 101 102 continental slope, today these waters also inundate the continental shelf along the northwest coast of Scotland, including the Malin Shelf (Huthnance, 1986, Reid et al., 1997). Troughs and 103 basins bounded by the SW-NE orientated Caledonian fault system extend and interlink from 104 the Sea of Hebrides to the mid-shelf and were major flow paths for streaming ice across the 105 Malin Shelf (Davies et al., 1984; Dobson et al., 1992). Two basins, the Malin Deep in the south 106

and a basin in the north, are separated by a central bedrock high, the Stanton Banks (Dobsonet al., 1992, Fig. 7).

109

# 110 2.1. Previous work on the glacial history of the Malin Shelf

Glaciation of the continental shelf to the shelf edge did not occur until the mid-Pleistocene, 111 ~440 ka (Stoker et al., 1993), with localised ice extending to the shelf edge during two further 112 glaciations in Marine Isotope Stage (MIS) 4-2 and likely including the LGM (Stoker et al., 113 1993). Ice draining western Scotland, the North Channel and northwest Ireland likely merged 114 115 with Hebridean ice to form the BFIS (Dunlop et al., 2010; Howe et al., 2012; Finlayson et al., 2014, Dove et al., 2015), possibly reaching a shelf edge position at ~27 ka BP (Wilson et al. 116 2002; Peck et al. 2006; Scourse et al., 2009). The sea floor geomorphology (Dunlop et al., 117 2010, Finlayson et al., 2014, Dove et al., 2015) provide prima facie evidence for ice streaming 118 (e.g. elongated bedforms, convergent flows) in this sector of the BIIS. Initial retreat was likely 119 rapid given prevalence of over deepened troughs (Dunlop et al., 2010, Dove et al., 2015), with 120 potentially slower, more episodic retreat on the inner shelf (Dove et al., 2015) accompanied by 121 substantial reorganisation of ice flow directions (Finlayson et al., 2014). Iceberg furrow marks 122 on the outer shelf, west of the Malin Deep, indicate that calving accompanied the initial retreat 123 124 phase (Dunlop et al., 2010). Equivalent evidence for calving is lacking for the inner shelf, which is likely a function of deeper water in the Malin Deep and burial by post-glacial sediment 125 126 further east (Dunlop et al., 2010). Sea-level studies (e.g. Brooks et al., 2008; Bradley et al., 2011) suggest that the Malin Shelf region retained a calving margin throughout deglaciation, 127 128 which is supported by uninterrupted IRD flux into the DBF to 16 ka BP (Kroon et al., 2000).

129

130 There is no existing geochronology from the outer and mid Malin Shelf to constrain the glacial/deglacial chronology of the BFIS. However, there is limited chronological control for 131 the adjacent Outer Hebrides Ice Cap. Glacimarine sediments from the Hebridean Shelf, dated 132 to 26.4 ka (Hedges et al., 1988), provides either a minimum date for the retreat of ice from the 133 shelf edge or it merely dates some glacimarine sediment distant to a more restricted maximum 134 ice extent (Peacock et al., 1992). Dating of marine material from the mid-Hebridean Shelf 135 suggests that ice did not retreat from the mid-shelf until after 18.5 ka (Peacock et al., 1992). 136 TCN and radiocarbon ages from the eastern periphery of the Malin Sea indicates that initial 137 deglaciation occurred before 20 ka (McCabe and Clark, 2003; Clark et al., 2009, Small et al., 138 2017), earlier than expected from the limited radiocarbon dates collected offshore and 139 suggestive of a more rapid pattern of retreat across the Malin Shelf. In summary, the BFIS 140

141 deglacial chronology for the outer and mid Malin Shelf is limited and a more comprehensive

- 142 dataset is needed to test hypothesised patterns, styles and rates of ice marginal retreat.
- 143

# 144 **3. Methodology**

Cruise JC106 of the RRS James Cook in August 2014 collected sub-bottom profiler (chirp) data 145 and sediment cores (Fig. 1). Chirp data were collected using a hull mounted Kongsberg SBP 146 120 sub-bottom profiler that operated a sweep frequency between 2500 to 6500 kHz, with a 147 depth resolution of 0.3 ms. The profiles were interpreted using the IHS Kingdom<sup>™</sup> software. 148 149 To convert the two-way travel time to depth estimates an average p-wave velocity of 1620 ms<sup>-</sup> <sup>1</sup>, measure from sediment cores using a Multi-Sensor Core Logger (MSCL) was used We 150 identify eight acoustic facies (AFA-AFH) from the sub-bottom profiler data (Fig. 2), and 151 present five regional acoustic profiles from the outer (MS1-3, Fig. 3) and inner Malin Sea 152 (MS4-5, Fig. 4). 153

154

Thirty-six vibrocores were retrieved (Table 1) using the British Geological Society vibrocorer 155 with a 6 m barrel and 8 cm core diameter. The underwater position of each core was collected 156 using a Sonardyne Ranger USBL beacon attached to the vibrocorer. The on-board GEOTEK 157 158 MSCL measured the gamma density, p-wave velocity and magnetic susceptibility of each core at two centimetre resolution. All cores were described visually, noting the lithology, texture, 159 160 contacts, sedimentary structures and Munsell colour, and shear vane measurements were made using a hand held Torvane at 10 cm intervals. A GEOTEK XCT scanner provided X-161 162 radiographs at a 92 µm resolution to further refine the core lithofacies and to count the clasts larger than 2 mm, classified as IRD, in 2 cm windows down the core. The X-radiographs, visual 163 164 logs and physical properties were used to identify six lithofacies associations (LFA1-LFA6) from the shelf cores and a further five lithofacies associations from the upper continental slope 165 cores (LFA7-LFA11). Figure 5 shows examples of these and they are described in section 4.2. 166 Within the lithofacies associations, 20 individual lithofacies were identified (Table 3: after 167 Eyles et al., 1983). 168

169

From the cores a mixture of paired bivalves, mixed benthic foraminifera samples and shell fragments were collected and cleaned for dating. All dated material is assumed to have lived in a benthic and marine environment. Only whole, unabraded foraminifera specimens were picked from sieve (500, 180 and 63  $\mu$ m) residues, with the assemblage dominated by the cold water species *Elphidium clavatum*, *Cassidulina reniforme*, *Nonionella labradorica* and

*Cibicides lobatulus.* Forty-nine samples for radiocarbon dating (Table 2) were collected from 175 cores on the shelf and upper slope and submitted to the NERC radiocarbon facility in East 176 Kilbride where they were prepared to graphite and passed to the SUERC AMS laboratory 177 (SUERC publication codes) or the Keck C Cycle AMS laboratory, University of California, 178 Irvine (UCIAMS publication codes) for 14C measurement. The primary aim of the dating was 179 to constrain the timing of ice-sheet advance and retreat across the shelf. We dated 32 reworked 180 shell fragments, one fragment per date, and two mixed benthic foraminifera samples from the 181 over-consolidated diamicton units. These ages constrain the onset of ice advance across the 182 183 shelf and the timing of sediment reworking by icebergs when the core site was free of grounded ice. These ages are treated as a maximum age of reworking. A further 15 samples were taken 184 from above the over-consolidated diamicton in glacimarine sediments and constrain the timing 185 when the core site was free from grounded ice and thus a minimum age of retreat. . Their 186 stratigraphic positions are shown and are discussed in context with the lithofacies 187 interpretations (Fig. 6a-e). 188

189

The conventional <sup>14</sup>C ages were calibrated using the Marine13 curve with an inbuilt marine 190 191 reservoir correction of 400 years and a  $\Delta R$  of 0 years (OxCal 4.2 software; Reimer et al., 2013). 192 The ages are reported in the text as the calibrated  $2\sigma$  median result in the form ka BP (see Table 2). The radiocarbon reservoir age in the North Atlantic is known to vary both temporally and 193 spatial (Austin et al., 1995; Peck et al., 2006; Singarayer et al., 2008). As a sensitivity test to 194 account for the temporal and spatial variability we have applied two further age calibrations 195 196 using the  $\Delta R$  +300 and +700 yrs as exemplified by Small et al. (2013a). These results are reported in Table 2. The different  $\Delta R$  values have a modest impact and due to the large 197 uncertainties on the correct  $\Delta R$  to use (Wanamaker et al., 2012), and for clarity in the text, we 198 refer only to ages calibrated with a  $\Delta R$  of 0 unless otherwise stated. 199

200

# 201 **4. Results and interpretation**

- 202 4.1 Acoustic facies
- 203

# 204 4.1.1 Acoustic facies A (AFA)

AFA lacks internal structure and forms the acoustic basement in the northeast of the study area when visible. It typically has a weak to moderate strength, impenetrable and discontinuous upper surface reflector of high amplitude. This surface reflector generally forms rugged and variable topography that occasionally crops out at the sea bed (see MS4, Fig. 4a and examples in Fig. 2). In seismic line MS5 (Fig. 4b), AFA forms a large smooth ridge that has at least 100
m of relief. The depressions on either side of this ridge are infilled with acoustic facies AFC to
AFF.

212

#### 213 *4.1.2 Acoustic facies B (AFB)*

AFB has a transparent to faintly stratified internal acoustic return and a very strong upper 214 reflector. On the outer shelf this upper reflector exhibits a corrugated surface pattern with 215 hyperbolic reflections (e.g., MS1; Fig. 3), and a smooth upper reflector surface in the inner 216 217 shelf (e.g., MS5; Fig. 4). The basal reflector is moderate to diffuse in strength, smooth and occasionally undulating (e.g., on the outer shelf, seismic profile MS1; Fig. 3a), discontinuous 218 and often obscured by the overlying AFB. AFB exhibits a range of morphologies including 219 mounds, asymmetric wedges, ridges and sheets that reach a maximum of ca. 60 m in thickness 220 with the larger features cropping out and locally influencing the seabed topography (Fig. 3). 221 On the inner shelf, these ridges often interdigitate with AFE, and here the lower reflector is 222 diffuse and either very faint or not visible. 223

224

For the outer shelf sector, in the north (MS1: Fig. 3) AFB comprises two substantial ridges 225 226 (GZW1a and GZW1b) that are 23-25 m thick and have outer edges at the shelf break and 20 km inshore of it. The inner slopes of both ridges are ornamented with smaller mounds ranging 227 228 in height from between 15-17 m with some onlapping the adjacent fronting mound to the west. The small mounds reduce in size moving east with the easternmost mound forming a thin 229 230 horizontal lens 4 km in width (Fig. 3). Further south, AFB is set slightly back from the shelf break. On MS2 (Fig. 3), AFB forms a 58-m high wedge (TS1a) extending from 3.5 to 22 km, 231 232 which is buried to the east by an onlapping package of AFB (TS1b) that combine to form a broad, low-relief, ridge outlined in Figure 7. On the southern line (MS3), AFB comprises two 233 234 small ridges 1 km from the shelf break, and as with MS2, there is also a thick sheet (TS2) of AFB that lies in a depression 5 km from the shelf edge (Fig. 3) that also forms part of the broad, 235 subdued, ridge mapped on Figure 7. On the mid-shelf, east of the Stanton Banks and on the 236 landward slope of the Malin Deep, AFB forms a series of distinct asymmetric, acoustically 237 transparent wedge-like features on seismic line MIS5 (GZW2a-c); Fig. 5b). The surface 238 topography of these asymmetric wedges is flat and smooth. Across the study area AFB was 239 sampled in 20 cores all of which recovered a stiff shelly diamicton. 240

241

242 4.1.4 Acoustic Facies C (AFC)

AFC is acoustically transparent with no internal reflectors. It forms isolated lenses 3 to 11 m thick in topographic lows of AFA and AFB (see Fig. 2). It is bounded at its surface by a smooth, undulating and weak reflector, and it often has a hummocky lower bounding reflector. This facies occasionally forms acoustically transparent wedges that interdigitate with the glacimarine basin fills of AFE.

248

249 4.1.5 Acoustic Facies D (AFD)

AFD is characterised by chaotic, contorted and sub-parallel internal acoustic stratifications and hyperbolic diffractions that can be strong in amplitude. This facies forms a discontinuous drape that reaches a maximum thickness of 12 m, but is generally 3-4 m thick, and overlies topographic irregularities in AFB. The upper surface can be hummocky, but more often is flat due to truncation by the overlying facies. The upper bounding reflector ranges from diffuse to moderate strength.

256

# 257 4.1.6 Acoustic Facies E (AFE)

AFE (Fig. 2) is acoustically well stratified and is characterised by continuous, evenly spaced, 258 parallel to sub-parallel internal reflectors. This facies often forms an on-lapping fill (max 40 259 260 m) in the topographic lows of AFA and AFB (Fig. 4b) and also overlies and truncates AFC and AFD. On the outer shelf (MS1, Fig. 3a) and within the Inner Hebrides Trough (Fig. 4) this 261 262 facies forms relatively thin drapes that contain some acoustic blanking. In the deeper basins (MS5, Fig. 4b), which lie east of the Stanton Bank and to the south of the study area, AFE 263 264 comprises a series of stacked stratified units with each successive unit truncating and on-265 lapping the previous.

266

# 267 4.1.7 Acoustic Facies F (AFF)

AFF (Fig. 2) is an acoustically transparent to faintly laminated drape that typically forms the uppermost facies. Characterised by a diffuse lower reflector on the outer shelf, AFF forms a very thin, continuous drape of < 1 m thickness that overlies AFB and AFE. On the inner shelf this unit can be very thick (greater than 40 m thick) and directly drapes bedrock (AFA: MS4, Fig. 4a). At its thickest, no underlying facies are visible due to pulse attenuation

273

#### 274 4.1.8 Acoustic Facies G (AFG)

AFG is characterised by disrupted and chaotic internal reflectors that are contained within small

v-shaped depressions cut into the GZW diamictons (AFB: Fig. 4).

# 278 4.1.9 Acoustic Facies H (AFH)

AFH is restricted to the upper continental slope and form a series of stacked, smooth but hummocky lenses that are internally structureless and acoustically homogeneous. AFH often contains isolated point-source diffracted hyperbole concentrated at the basal reflector.

282

283 *4.1.9 Acoustic interpretation* 

AFA, is interpreted as bedrock and is restricted to the Inner Hebrides Trough and the ridge behind the Stanton Banks (Figs 4a and 4b). Elsewhere the Malin Sea shelf contains very thick sequences of Quaternary sediments that overlie and obscure the bedrock (AFA) in the seismic data (Stoker et al., 1993; Fyfe et al., 1999).

288

AFB is interpreted as a sub-glacial till that has been formed into a series of grounding-zone 289 wedges (GZWs), till sheets and moraines marking the grounding line positions of the BFIS at 290 the shelf edge and during subsequent retreat. These ice-marginal features identified in the 291 seismic lines have a strong signature in the bathymetric data (Fig. 7), but previous descriptions 292 293 were restricted to the south of the study area (Dunlop et al., 2010). Together, these data identify 294 two sets of grounding line positions, separated by two deep troughs (~170 m-bsl). The outer shelf is characterised by a series of small moraines, GZWs (GWZ1a-b) and broad subdued 295 296 wedges interpreted as till sheets (TS1-2) that combine to form the broad sediment ridge mapped in Figure 7. This marks a grounding line position that extends the width of the Malin Shelf. No 297 298 GZWs or moraines were visible in the seismic or bathymetry data from the deep troughs. Directly east of the Stanton Banks, a second set of GZWs (2a-c) are also visible in the 299 300 bathymetry data and were mapped in part by Dunlop et al. (2010). The morphology of the 301 Malin Shelf features is consistent with GZWs described from other glaciated margins (e.g., 302 Batchelor et al., 2015; Anderson and Jakobsson, 2016; Evans and Hogan, 2016). These features form subdued features on the Malin Shelf, wide relative to their height. The size and 303 morphology of these wedges are characteristic of a grounding zone, where the thickness of the 304 GZW is constrained by the sub-ice shelf cavity as opposed to moraine ridges formed at a 305 grounded tidewater cliff (Dowdeswell and Fugelli, 2012). The smaller mounds located on the 306 edge of the large GZWs in seismic line MS1 (Fig. 3a) are likely moraine banks produced by 307 308 minor ice stream re-advances during a period of prolonged retreat (Batchelor and Dowdeswell, 309 2015).

AFC is consistent with an interpretation of as debris flows (e.g. Dowdeswell et al., 2010;
Batchelor et al. 2011). AFC was not sampled during the coring.

313

AFD was captured in two cores (150VC and 153VC), both containing a massive, soft, pebbly rich mud. Thus, AFD is interpreted as an ice proximal, IRD-rich deposit. The contorted internal structure may represent modification of the sediment from either post-deposition settling or minor oscillation of the ice margin near to the area of sediment deposition. The hyperbolic diffractions are likely caused by the presence of clasts, that are either large or occur in high concentrations.

320

AFE is interpreted as glacimarine basin fills that are more distal to the ice margin. Winkelmann 321 et al. (2008) suggest that this type of stratification could be a result of changing clast 322 concentrations relating to variations in IRD and suspension dominated depositional processes. 323 AFE was sampled in eight cores that contained alternating massive pebbly rich mud with 324 laminated silts and clays. Similar acoustic facies and lithofacies are documented in the Arctic 325 (e.g., Dowdeswell et al., 2010; Batchelor et al., 2011; Hogan et al., 2016). The stratification is 326 327 often conformable in the sub-bottom profile data, consistent with suspension settling from turbid meltwater plumes and IRD rainout (Hogan et al., 2012; Ó Cofaigh et al., 2016). In MS5 328 (Fig. 4) there is evidence of truncation of the acoustically stratified sediment and the presence 329 330 of transparent wedges (AFC) within AFE. This is consistent with downslope remobilisation and erosional events characteristic of these glacimarine settings (c.f., Ó Cofaigh et al., 2016). 331

332

AFF was captured in nearly all the cores collected in this study and comprised coarse-grained sand to gravel or shell hash. On the inner Malin Shelf and in the Inner Hebrides Trough AFF consists of medium, fine or muddy-fine sand. The abundance of shell hash and whole shells, the coarse sand and gravels combined with the stratigraphy is consistent with post-glacial bottom-current winnowing (Howe et al., 2012).

338

AFG is interpreted as furrow infill, with the furrows formed by sea-bed scour by the keels of floating icebergs calved from the BFIS. Dunlop et al. (2010) noted that the outer Malin shelf bathymetry in particular contains high concentrations of v-shaped elongated furrows down to ca. 190 m water depth. The furrow infill (Core 145VC) consisted of coarse sandy-gravel overlying a stiff diamicton (Fig. 6c).

AFH is interpreted as the deposits of glacigenic debris flows on the continental slope likely 345 sourced when BFIS ice was at the shelf edge or by postglacial downslope mobilisation or 346 iceberg disturbance. The stacked lenses are characteristic of episodic subaqueous glacigenic 347 debris flows and gravity deposits sourced from ice streams crossing the continental shelf (e.g. 348 Laberg and Vorren, 1995; Elverhøi et al., 1997). AFH was recovered in three cores (124VC, 349 142VC and 143VC) and consists of a stiff basal diamicton representing the debris flows 350 overlain by laminated silts and clays, massive pebbly mud and/or massive bioturbated muds 351 that represents the transition to an ice distal setting once ice had retreated from the shelf edge. 352

353

#### 354 4.2 Sedimentology

355 4.2.1 Shelf Lithofacies Associations (LFA1-LFA4)

356 *LFA1: Stiff diamicton (Dmmc, Dms)* 

The basal lithofacies of 19 cores on the shelf and particularly the outer shelf are highly 357 consolidated diamicton (Dmmc) comprising predominantly massive, dark grey (Munsell 358 colour 5Y4-1), matrix-supported sub-rounded to sub-angular gravel to pebble-sized clasts in a 359 silty-clay to sandy-silt matrix. Shell fragments are frequent to rare, and when present they are 360 typically highly abraded although occasionally are in a good condition. In some cores LFA1 361 362 contains internal structures such as sand lenses (1-2 cm thick), crude stratification (Dms), dewatering structures and preferential clast alignments that are evident only in the X-363 364 radiographs (Fig. 5). LFA1 is typically over-consolidated with shear strengths averaging 100 kPa, varying between 45-200 kPa, and increasing with depth. LFA1 has high wet bulk densities 365 366 and magnetic susceptibilities (Fig. 6). LFA1 is constrained by 25 radiocarbon ages (Table 2), twelve of which are finite. These ages are not in stratigraphical order within individual cores 367 368 reflecting the prevalence of reworking and mixing in the sedimentary processes (Fig. 6). The oldest ages,  $26719 \pm 318$  and  $26825 \pm 302$  cal yrs BP, are from the shelf edge (125VC), with a 369 370 further five ages ranging between  $24179 \pm 212$  cal yrs BP to  $18251 \pm 169$  cal yrs BP from these outer GZWs and Till sheets (136VC, 145VC, 139VC; Table 1, Fig. 6b and 6c). 139VC 371 contains most diverse set of radiocarbon ages, ranging from infinite to six measurements 372 between  $15,031 \pm 204$  cal yrs BP and  $12,662 \pm 88$  cal yrs BP. The uppermost age in core 373 125VC, at 69.5 cm down the core also provides a young age of  $12,826 \pm 129$  cal yrs BP. 374

375

# 376 *LFA2: Soft diamicton or massive pebbly mud (LFA2: Dmm, Dms, Fmd)*

377 LFA2 is a matrix-supported diamicton with a sandy-silt matrix and gravel to pebble size clasts

378 (Dmm) and/or massive silts containing abundant pebble-size clasts (Fmd) present on the outer

shelf (146VC, 147VC) and inner shelf (148VC, 149VC, 151VC, 153VC and 154VC) (Fig. 6d 379 and 6e). These diamictons often form discrete beds (6-55 cm thick) and can repeat 2 to 5 times 380 in a core often alternating with LFA3 laminated mud. The diamicton in LFA2 varies from 381 massive (Dmm) to stratified (Dms) and have low shear strengths averaging 24.6 kPa with a 37 382 kPa maximum (149VC). The pebbly muds (Fmd) also form multiple beds ranging from 7-86 383 cm thick within individual cores, and contain fewer pebble-gravel sized clasts only visible in 384 the X-radiographs and in a muddy-matrix that has low shear strengths of no more than 10 kPa. 385 If present together (cores 146VC, 147VC, 148VC, 149VC, and 153VC), the Fmd is always 386 387 stratigraphically higher than Dmm. The Fmd is often massive, but with occasional water escape structures and preferential near vertical grain alignment (Fig. 5). The upper and lower contacts 388 of LFA2 are often sharp and can form intruded/loaded contacts with the underlying LFA3. 389 Occasionally the lower boundary can be gradational over 5 cm and consist of laminated fine 390 mud and diamicton (Fig. 5). The sediment matrix varies from dark grey to very dark grey (5Y 391 4/1, 5Y 3/1) or olive brown/greyish brown (5Y 4/2, 2.5Y 5/2) with the Dmm darker in colour 392 than the Fmd. LFA2 is characterised by medium wet bulk densities and variable high magnetic 393 394 susceptibilities (Fig. 6) due to the clast content. Seven radiocabron ages were obtained from LFA2 targeting mixed benthic foraminifera, with the three oldest ages from core 146VC on the 395 396 outer shelf  $(23,828 \pm 251, 25,897 \pm 178 \text{ and } 24,424 \pm 331 \text{ cal yrs BP}$ ; Fig. 6c) again not in stratigraphic order. A chronologically tight cluster of three radiocarbon ages come from the 397 398 mid-shelf (151VC, 153VC, 154VC;  $23,232 \pm 292, 22,679 \pm 270$  and  $22,129 \pm 261$  cal yrs BP, respectively), along with an additional 14C age for a shell fragment in a soft diamicton (149VC, 399 400  $20,204 \pm 190$  cal yrs BP; Fig. 6d and 6e).

401

# 402 *LFA3: Laminated mud (Fl, Fldef)*

LFA3 consists of greyish brown to very dark grey (2.5Y 5/2 to 5Y 3/1) fine (1cm to >1 mm 403 404 thick) horizontal laminated clays and silt with no evidence of bioturbation, although deformation structures, including convoluted bedding, ball and pillow structures or bending of 405 the underlying laminae, coincident with isolated dropstones, are visible. High angle conjugate 406 to normal brittle faults are also visible and are more prevalent towards the core base. Although 407 408 >2 mm clast are present in the Fl, the clast count is consistently less than 10 grains (per 2 cm window). Laminations vary from: (i) continuous horizontal to sub-horizontal laminae with 409 sharp upper and lower contacts; to (ii) deformed, convoluted and discontinuous laminae. Basal 410 sections of seven cores (146VC, 147VC, 148VC, 149VC, 151VC, 153VC, 154VC: Fig. 6d and 411 6e) consist of LFA3 alternating with LFA2 with sharp upper and lower contacts (Fig. 5c). Fl is 412

- 413 firmer (shear strength average 16 kPa, range 4-34 kPa) when overlain by LFA2. Towards the
  414 upper contact of Fl the bedding often becomes deformed, with convoluted structures (Fldef).
- 415
- 416 LFA4: Laminated, stratified and bioturbated mud containing frequent clasts (Fld, Flb,
  417 Fldef(d), Fs, Fsb)
- LFA4 (148-151VC: Fig. 6d and 6e) gradually transitions from a clearly defined laminated mud with no bioturbation, but frequent >2 mm clasts (Fld), to a bioturbated laminated mud (Flb) to a bioturbated and shell rich stratified mud (Fsb). Flb often consists of wispy, convoluted and/or deformed laminae that contain abundant shell fragments and gravel-sized clasts. Core 148VC contains high angle (>40°) laminae and convoluted bedding with clasts aligning in the coarser silt laminae (Fldef(d): Fig. 6d). A 14C age of 16,860  $\pm$  221 cal yrs BP came from a whole bivalve of *Yoldiella sp.* from the Flb facies (151VC: Fig. 6e).
- 425
- 426 *LFA5: Bioturbated massive mud (Fm, Fmb, Fmg):*
- LFA5 is dark grey (7.5YR 4/1) massive muds (Fm) that are commonly bioturbated (Fmb) and 427 can contain abundant shell fragments (140VC: Fig. 6b). Occasionally beds of gravelly mud 428 429 (Fmg) interbed with Fs (LFA4). Bioturbation is visually characterised by darker mottles and 430 streaks, but burrows are evident in the X-radiographs. Occasional gravel-sized clasts can be present in this facies, but it is comparatively clast free. Fm is often restricted to a single bed (2-431 432 146 cm thick) and is buried by massive shelly sand or gravel (LFA6) with a sharp but convoluted contact. Frequent lenses, pods and burrows filled with the overlying material 433 434 (LFA6) occur in the upper 10-30 cm of LFA5. LFA5 has a low average shear strength (6 kPa), low wet bulk densities and low magnetic susceptibilities (Fig. 6). Occasionally Fm interbeds 435 436 with massive, gravelly mud (Fmg) that contains abundant shell fragments and have gradational 437 contacts.
- 438
- 439 LFA6: Muddy sand and shelly sand and gravel (Sfm, Sfmb, Sm, Sl, Suf, Gm, Gms)

LFA6 is the surface unit across all cores and consists of: massive muddy fine sand (Sfm), bioturbated at times (Sfmb); massive coarse to fine sand (Sm); laminated sand (Sl); finingupward sand (Suf); gravel (Gm) and matrix-supported gravel (Gms). The physical properties of these units are typically soft (6 kPa), a low bulk densities and variable magnetic susceptibilities peaking in the gravels (Fig. 6). LFA6 is often rich in shell fragments and variable in thickness.

#### 447 *4.2.2. Lithofacies Association (LFA) Interpretation:*

LFA1 is the dominant lithofacies of the outer moraines, GZWs and till sheets that are identified 448 as AFB. They form the broad sediment ridge that spans the outer shelf marking the former 449 extent of the BFIS (Fig. 7). Multiple process interpretations for LFA1 are possible spanning 450 subglacial diamictons over-compacted by grounded ice (Ó Cofaigh et al., 2007, 2013; Evans 451 et al., 2006) to sub-ice shelf diamictons affected by severe iceberg turbation (Domack et al., 452 1999). IRD rain-out is another possible interpretation, but the over-consolidation of LFA1 453 indicates that this is not the dominant process. Morphological evidence in the bathymetric data 454 455 confirm pervasive iceberg scouring of the outer shelf down to water depths of 450 m (Dunlop et al., 2010; Stoker, 1995). The radiocarbon dating sheds some light on the process environment 456 with mixed ages and sequences showing stratigraphic age inversion. The spread of radiocarbon 457 ages between  $24.2 \pm 0.2$  and  $12.7 \pm 0.1$  ka alongside infinite measurements (: Fig. 6b), suggest 458 the mixing and compacting of subglacial and ice-marginal diamicton by icebergs. A similar 459 process of iceberg turbation of both subglacial and glacimarine diamictons occurs in Antarctica 460 and the resultant diamictons are impossible to differentiate (c.f. Powell et al., 1996). Locations 461 protected from iceberg activity, for example by a bounding moraine negate this problem, and 462 the oldest ages in subglacial diamicton from core 125VC (Fig. 6b) provide a maximum 463 464 constraint on the advance of grounded ice to the shelf edge to after  $26.7 \pm 0.3$  ka BP although this could be as young as  $26 \pm 0.1$  ka BP if using a  $\Delta R$  of 700 yrs. 465

466

The alternating beds of LFA2 (Dmm/Fmd) and LFA3 (Fl) reflect a switch to more clearly 467 468 discernible glacimarine sedimentation in an ice-proximal setting. One explanation for this pattern is a seasonal signal capturing the switch in meltwater plume dominated sedimentation 469 470 during the summer and deposition of ice rafted diamicton from icebergs calving in the winter when meltwater production is lower (Syvitski et al., 1989, Cowan et al., 1997, Cai et al., 1997). 471 472 An alternative, and favoured, interpretation is that these sediments represent a change from IRD dominated deposition forming sediments of LFA2, to a period of supressed IRD rainout 473 and meltwater plume dominated sedimentation caused by multiyear sea-ice build-up depositing 474 LFA3 (Ó Cofaigh and Dowdeswell, 2001). Similar interpretations have been made from cores 475 collected in east Greenland (e.g. Jennings and Weiner, 1996, Dowdeswell et al., 2000). The 476 LFA2 beds contain abundant foraminifera for dating and follow a time transgressive pattern 477 with distance from the shelf edge with the oldest deglacial ages from core 146VC, located on 478 the outer shelf, and the youngest from core 149VC located within the Sea of Hebrides. The 479 three ages from core 146VC do not lie in stratigraphic order indicating some downslope 480

remobilisation occurred that is evident in the X-radiographs where a change in clast alignment is visible (Fig. 5). Nevertheless, the dates of  $25.9 \pm 0.2$  and  $24.5 \pm 0.3$  cal ka BP from glacimarine sediments indicate that this section of the Malin Shelf was free from grounded ice by this time. The oldest age from the inner shelf,  $23.2 \pm 0.3$  ka BP in core 151VC (Fig. 6e), lies directly above AFB and dates the early deglaciation of this part of the Malin Shelf. Finally, a basal age from core 149VC constrains the timing of deglaciation of the Inner Hebrides Trough and indicates that the entire Malin Shelf was free by grounded ice by  $20.2 \pm 0.2$  ka BP.

488 The muds of LFA4 and LFA5 are interpreted as hemipelagic ice-distal sediments deposited in an open (glaci)marine environment through the process of suspension settling and occasional 489 490 IRD contribution. In ice-distal environments, laminated muds often grade into massive bioturbated muds (cf. Jennings, 1993; Svendsen et al., 1992). Only one radiocarbon age dated 491 492 LFA4 (151VC; Fig. 6e), providing a calibrated age of  $16.8 \pm 0.2$  ka BP indicating that the inner shelf close to Stanton Banks was an ice-distal environment by this time. Lithofacies Fldef in 493 494 core 148VC, with near vertical laminations, likely represents mass movement of sediment downslope at the core site due to its location in a steep sided bedrock basin. 495

496

The sands and gravels of LFA6 are interpreted as postglacial deposits reworked by bottom current activity across the shelf and upper slope (cf. Anderson et al., 1984; Howe et al., 2001). The differing particle size of the matrix and fining upward sequences may represent a weakening of the bottom currents with time (Viana et al., 1998). The shell hash, gravels and massive sands likely represent lag deposits (Dowdeswell et al., 1998). There are no fine glacimarine sediments preserved on the outer shelf indicating that bottom-current winnowing has been pervasive across the shelf since its deglaciation (Howe et al., 2012).

504

505 *4.2.3 Lithofacies on the upper continental slope (LFA7- LFA11)* 

506 LFA7 Matrix-supported diamicton (Dmmc)

LFA7 forms the basal lithofacies in five cores (118VC, 124VC, 142VC, 143VC, 144VC) and consists of very dark grey (5Y 3/1) sub-rounded to sub-angular gravel to pebble-sized clasts in a very stiff, silty-clay to sandy-silt matrix (Dmmc). Dmmc also contains occasional shell fragments. Core 144VC consists of a 6 m thick Dmmc that has four distinct colour changes with sharp boundaries down core that oscillate from grey to very dark grey from one bed to the next (5Y 4/1 and 5Y 3/1). LFA7 is typically over-consolidated with shear strengths averaging 82 kPa, but can be variable within a core (e.g. 142VC, Fig. 6). LFA7 has also medium wet bulk densities and magnetic susceptibilities (Fig. 6b). LFA7 is dated by nine radiocarbon ages, five of which are finite. The remaining four ages, from core 142VC, are not in stratigraphical order and range between  $22,402 \pm 123$  to  $18,138 \pm 174$  cal yrs BP. (Fig. 6)

517

# 518 LFA8: Compact stratified, laminated and deformed fine sands and muds (Sl, Ss, Fldef)

LFA8 is a dark greyish brown stratified medium to fine silty-sand occurring in two cores 519 (116VC and 124VC). The laminations vary from finely laminated medium sand and silty-sand 520 that form wispy, convoluted lamina to coarser sandier layers that form discontinuous pods and 521 522 ball and pillow structures (124VC: Fig. 5). Occasional clasts are present towards the base of LFA8. Shear strengths are high, averaging 42 kPa and varying between 13-85 kPa. LFA8 has 523 medium wet bulk densities and low magnetic susceptibilities. LFA8 is constrained by three 524 radiocarbon ages, two lying in stratigraphic order from core 116VC ( $22,750 \pm 221$  and 18,326525  $\pm$  165 cal yrs BP) and one from the base of core 124VC (18,862  $\pm$  106 cal yrs BP; Fig. 6a). 526

527

# 528 LFA9: Alternating laminated and massive pebbly muds (Fl, Fmd, Fldef)

LFA9 only occurs in core 143VC and consists of three lithofacies; finely laminated clay silt with dipping and convoluted laminae (Fldef) as well as planar, horizontal laminations (Fl), which alternates with a massive clast-rich, silty-mud with some grain alignment (Fmd; Fig. 6c). There is no evidence of bioturbation. LFA9 is dark grey (5Y 4/1) and firm with average shear strengths of 23 kPa. A radiocarbon age of 18,090  $\pm$  174 cal yrs BP dates the base of the Fl facies.

535

# 536 *LFA10: Bioturbated laminated to massive muds (Flb, Fldef, Fmb)*

537 LFA10 consists of three lithofacies; laminated muds that are either bioturbated (Flb) or deformed (Fldef) and massive bioturbated muds (Fmb). Laminated silts and clays transition 538 from continuous parallel laminae with individual laminae less than 5 mm thick to wispy, 539 discontinuous and more diffuse laminae up-core. Bioturbation is visible as black mottles on the 540 sediment surface that increase in abundance down core and as a dense collection of small 541 irregular burrows to large vertical burrows on the X-radiographs (see Fig. 5). LFA10 is olive 542 grey (5Y 4/2) and has very low shear strengths, wet bulk densities and magnetic 543 susceptibilities. LFA10 also contains low >2 mm clast content (Fig. 6). The oldest age of 544  $22,751 \pm 229$  cal yrs BP dates the Flb facies at the base of core 115VC. A shell age from Fldef 545 in core 116VC produces a younger age of  $18,326 \pm 165$  cal yrs BP. 546

548 LFA11: Massive to laminated sands and sandy-gravels (Sl, Sm, Smb, Smg, Gs)

LFA11 forms the uppermost lithofacies in seven cores. It is predominantly composed of laminated fine-medium sand (SI) or massive fine-medium sand (Sm) with gravel to pebblesized clasts (Smg) and occasional gravel layers (Gs). LFA11 if often bioturbated (Smb), as shown by burrow in the X-radiographs, and contains high concentrations of shell fragments. LFA11 is variable in colour ranging from light olive to dark grey (5Y 4/3, 2.5Y 4/1), has low shear strength, low magnetic susceptibilities and medium wet bulk densities measurements (Fig. 6)

556

# 557 4.2.4. LFA7-LFA11 interpretation

LFA7 is interpreted to be both ice marginal debris flows deposited when the BFIS was at the 558 shelf edge feeding the DBF and postglacial reworking caused by both iceberg turbation and 559 sediment gravity flows. The series of stacked diamicton beds in core 144bVC, and the finite 560 ages collected from this core, are likely examples of stacked ice marginal debris flows (cf. 561 Laberg and Vorren 1995). Whereas, the variable shear strength measurements combined with 562 the range of ages spanning  $22.4 \pm 0.1$  to  $18.1 \pm 0.2$  ka BP in core 142VC is interpreted as debris 563 flows that where subsequently reworked by icebergs once the BFIS had retreated from the shelf 564 565 edge.

566

LFA8 is interpreted as coarse-grained sandy turbidites interbedded with finer-grained muddy turbidites. Although the shear strength in both cores is high we rule out overriding by grounded ice as a cause for this due to the position of the cores on the upper continental slope at ~240 m water depth. Instead, we attribute the high shear strength to compaction during downslope flow and removal of pore water as evident from the water escape structures visible in the Xradiographs (e.g., Fig. 5). We cannot rule out compaction by icebergs but it is likely that icebergs calving from the BFIS would ground on the shallower shelf before reaching the slope.

LFA9 is similar to the alternating LFA2 and LFA3 lithofacies associations except there is only one glacimarine couplet captured. LFA9 is interpreted to represent a period dominated by suspension settling of turbid water plumes with limited IRD followed by period of IRD delivery combined with suspension settling as described in section 4.3.1. The age of  $18.1 \pm 0.2$  ka BP indicates that IRD was still deposited on the shelf slope at and after this time.

581 LFA10, characterised by bioturbated laminated muds transitioning to massive bioturbated muds, is interpreted as suspension settling from meltwater plumes. Knutz et al. (2001) noted a 582 similar lithofacies and interpretation in core MD95-2006 taken from the DBF, and dated to 583 after 22 14C ka BP. The basal ages from core 114VC and 116VC correlate with this timing. 584 The lack of sand and coarse sediment and the presence of bioturbation and marine fauna 585 suggest that the grounding line was not marginal at the time of deposition (cf. Ó Cofaigh and 586 Dowdeswell, 2001). The vertical burrows in core 115VC indicate sedimentation was rapid (cf. 587 Jennings et al., 2014). The transition from laminated to massive bioturbated mud is likely a 588 589 result of increasing distance from the ice margin and therefore a slowing in the rate of sediment 590 accumulation (Ó Cofaigh and Dowdeswell, 2001).

591

LFA11 is similar to LFA6 and likely represents postglacial sedimentation. On the slope, cross
slope and bottom current reworking are the processes responsible for the deposition of massive
and laminated sand, which are a result of both reworking and winnowing of the fine sediments
(Faugères and Stow, 1993; Stow et al., 2002).

596

# 597 5. Discussion

598 5.1. Timing of the maximum BFIS on the Malin Shelf

599 Our new geophysical data demonstrate that the BFIS reached the shelf edge of over the majority of the Malin Sea. This maximum extent is more widespread than previously figured (Dunlop 600 et al., 2010). A wide and often compound ridge is visible in the bathymetric data (Fig. 7), with 601 GZWs, till sheets and moraines occurring across the shelf edge in the north and south, and near 602 to the shelf edge in the centre of the Malin Shelf. Constraining the timing for this maximum 603 extent has proven challenging with reworked older fauna ubiquitous and iceberg turbation 604 introducing younger specimens to the glacial sediments. On the southern Malin Shelf, the shelf 605 edge moraine provided a protected setting from iceberg turbation, and the youngest radiocarbon 606 dated shell fragments in subglacial diamictons (Core 125VC) constrain shelf edge glaciation 607 to after  $26.7 \pm 0.3$  ka BP. A minimum age is provided by the oldest ages from retreat stage soft 608 609 diamictons (146VC), thus delimiting shelf edge glaciation to between 26.7  $\pm$  0.3 and 25.9  $\pm$ 610 0.2 ka BP (as late as  $25.3 \pm 0.3$  ka BP if using a  $\Delta R$  700 yrs). This timing is very similar to chronologies obtained for shelf edge glaciation elsewhere on the western marine margin of the 611 last British Irish Ice Sheet, with timings of after 26.3 ka BP for the Donegal Ice Stream (Ó 612 Cofaigh et al., submitted), and before 24.3 ka BP in the Celtic Sea (Praeg et al., 2015). A shelf 613

edge position for the BFIS between  $26.7 \pm 0.3$  to  $25.9 \pm 0.2$  ka corresponds with peak IRD flux in core MD95-2006 from the adjacent DBF that also supports a shelf-edge position at ~27 ka BP (Knutz et al., 2001; Scourse et al., 2009).

617 5.2. Iceberg calving history

Radiocarbon dates from iceberg-turbated diamicton on the outer Malin Shelf lie between 24-618 18 ka BP, and point to period of iceberg calving likely associated with the breakup of the BFIS 619 that is equivalent to evidence from other marine-based sectors of the BIIS (Benetti et al., 2010; 620 Dunlop et al., 2010; Ó Cofaigh et al., 2012; Peters et al., 2016). Dunlop et al. (2010) have 621 622 argued that the lack of iceberg furrows to the east of the Malin Deep suggest that no major 623 calving event took place after the initial breakup of ice on the outer Malin Shelf. Whereas, our 624 chronology for iceberg turbate indicates a period of consistent iceberg activity spanning 6 ka and that the inner Malin Shelf was ice free by 23 ka with iceberg turbation an important process 625 626 as late as 18 ka. Two of the outer shelf cores (139VC and 125VC) provide a very different 627 chronology for sediments with physical characteristics identical to iceberg-turbated diamicton 628 elsewhere. These sediments (139VC) contain younger samples ranging  $15 \pm 0.2$  to  $12.7 \pm 0.1$ ka BP intermixed with older materials in the range  $22.4 \pm 0.2$  to  $24.2 \pm 0.2$  ka BP (Table 1 and 629 630 Fig. 6b). Similarly, the youngest age from a reworked shell fragment in these diamictons (125VC) is  $12.8 \pm 0.1$  ka BP. If iceberg turbation was responsible for reworking shell materials 631 into diamictons after  $12.7 \pm 0.1$  ka BP, this supports a view put forward by Small et al. (2013b) 632 for the existence of a marine and calving ice margin in western Scotland during the Younger 633 Dryas Stadial, a cold event that interrupted the warming and deglaciation of the Northern 634 Hemisphere between 12.9-11.7 ka BP (Alley, 2000). If the larger  $\Delta R$  of 700 yrs is applied to 635 these ages then the youngest age from 139VC would shift from  $12662 \pm 88$  cal yrs BP to 11591 636  $\pm$  267 cal yrs BP. This is still broadly coeval to the Younger Dryas Stadial and for icebergs 637 638 reworking the shelf sediments at this time.

639

To date, understanding of the dynamics and iceberg calving history for the BFIS has relied on IRD records in the DBF (Scourse et al., 2009), which showed: (1) peaks in IRD flux attributed to grounded ice at the shelf edge by 27 ka BP and a maximum mass at 24 ka BP (Scourse et al., 2009); (2) decline in IRD reflecting a rapid ice retreat ~23 ka BP; and (3) further peaks in IRD attributed to a shelf wide BFIS readvance between 22-16 ka BP. Our retreat history differs showing an early retreat of some 19 km from the shelf edge by 25.9 ka BP, a further ~65 km by 23.2 ka BP and with the entire Malin Sea shelf ice free by 20.1 ka BP (Fig. 7 and 8) and 647 even if the larger  $\Delta R$  of 700 yrs is applied to our chronology, the timing of retreat still occurs earlier (see table 2) than previously thought. There is no evidence in the acoustic profiles or 648 sediment data for a major readvance back across the shelf between 22-16 ka BP. The glacial 649 dynamics associated with IRD flux, e.g. retreat or advance stage, have proven difficult to 650 resolve (Scourse et al., 2009), whereas our sequence of events suggests that greater BFIS-651 sourced IRD flux to the DBF relates to phases of ice retreat, and as a process, calving dominated 652 the retreat dynamics of the Malin Sea area from 24 to 18 ka BP, with renewed activity during 653 654 the Younger Dryas Stadial.

655

# 5.3. Rates and controls on ice marginal retreat

GZW are associated with temporary pauses in ice-stream retreat or readvances over 10-100s 657 years and are regarded as representing episodic rather than catastrophic retreat (Dowdeswell 658 and Fugelli, 2012). Two areas of GZW occur on the Malin Shelf, the first at the outer shelf 659 edge and the second at the landward margin of two deep (>170 m) west to east trending troughs 660 661 landward of the Stanton Banks (Figs 9 and 5b). Both troughs continue into the Inner Hebrides Trough where water depths increase to >200 m (Fig 1) and are interpreted as glacial 662 663 overdeepenings (Dunlop et al., 2010). The GZWs are relatively small, 25 m thick and >10 km wide, but similar-sized GZW occur in other previously glaciated regions e.g., Antarctica (e.g. 664 665 Anderson and Jakobsson, 2016; Evans and Hogan, 2016; c.f. Table 2 in Batchelor and 666 Dowdeswell, 2015). In those cases their relatively small size is interpreted to reflect either a brief still-stands or a low sediment flux to the grounding zone (Dowdeswell and Fugelli, 2012). 667 668

Retreat of ice to the inner of these shelf edge GZWs, GZW1b, occurred after the oldest 669 deglacial age of 25.9  $\pm$  0.2 ka BP (146VC) and a halt in retreat at this GZW likely occurred 670  $25.9 \pm 0.2$  to  $24.8 \pm 0.3$  ka BP (146VC) based on two mixed benthic foraminifera ages taken 671 from IRD-rich sediments overlying consolidated diamictons. Landward of the overdeepenings 672 and inshore of the Stanton Banks topographical high (Fig. 4b) a series of GZWs step back in a 673 southeast direction and glacimarine sediments in front of these marginal positions have basal 674 ages ranging  $23.2 \pm 0.3$  to  $22.1 \pm 0.3$  ka BP indicating ice-free conditions by this time on the 675 inner shelf. There are no obvious moraines or GZWs visible in the available regional 676 bathymetry (Fig 2), which reflects rapid retreat of the BFIS across these deeper waters and their 677 reverse slopes. Previous research has pointed to early and rapid deglaciation of the inner Malin 678 Shelf (Binns et al., 1974), but that radiocarbon dating targeted bulk carbon and were dismissed 679

680 as too old (Harkness and Wilson, 1974). Later attempts to constrain ice-free conditions in other marine cores from the southern Inner Hebrides and western Scotland were significantly 681 younger (Peacock, 2008; Peacock et al., 2012). Our eastern most deglacial age (149VC; Fig. 682 6a) confirms this early deglaciation of the inner shelf by  $20.2 \pm 0.2$  ka BP. This timing is 683 supported by terrestrial evidence that deglaciation was already underway by 20.6 ka on Tiree 684 ~20 km to the east (Small et al., 2017).  $^{36}$ Cl exposure ages from southern Skye show complete 685 deglaciation by 17.6 ka (Small et al., 2016) and so a fully deglaciated Malin Shelf by 20.2  $\pm$ 686 0.2 ka BP fits the existing regional onshore geochronology. Radiocarbon and cosmogenic 687 688 nuclide dates from onshore the north coast of Ireland suggest that ice stepped back on land by at least ~20 ka in this region (McCabe and Clark, 2003; Ballantyne, 2007; Clark et al., 2009). 689

690

# 691 5.4. Forcing and conditioning of the retreat dynamics of the BFIS

The deglaciation of the Malin Shelf coincides with a period of global sea-level lowering 692 (Lambeck et al., 2014) and local cooling of sea-surface temperature (Peck et al., 2008; Hibbert 693 et al., 2010). A prevailing hypothesis for the 'early' deglaciation of the BIIS is the influence of 694 Heinrich Event 2 (H2), which is associated with a 15-30 m global sea-level rise caused by the 695 collapse of the Laurentide Ice Sheet (Chappell, 2002; Siddall et al., 2003). This sea-level rise 696 697 would have destabilised the marine-based portions of the BIIS calving front (Scourse et al., 2009). Mg/Ca sea-surface temperature estimates 698 Furthermore, and percent of 699 Neogloboquadrina pachyderma sinistral coiling (NPS %), a proxy for the polar front, show a northward migration of the polar font at the same time (Peck et al., 2008). This migration of 700 701 the polar front allowed warmer waters to reach the calving margin of the BIIS during Greenland Stadial 2 (GS2) at 23 ka BP (Scourse et al., 2009). However, our data indicate that the BFIS 702 703 had already retreated from its shelf-edge position and was located ~100 km further inland at the time of H2. Therefore, H2 and the migration of the polar front cannot explain the timing of 704 705 initial retreat from the shelf edge, although both likely had an influence on the continued retreat of the BFIS. 706

707

An alternative mechanism for early retreat is an increase in relative sea level (RSL) caused by local isostatic loading and crustal deformation of the Malin Shelf by the BFIS. Subsequently, this RSL rise would have led to the destabilisation and retreat of the ice margin (cf. Eyles and McCabe, 1989). The majority of the Malin Shelf was ice free by  $20.2 \pm 0.2$  ka BP (at latest  $19.4 \pm 0.2$  ka BP if using a  $\Delta R$  700 yrs), and our shallowest core (153VC) is ~20 m above the global sea level minima of 134 m between 29-21 ka BP (Lambeck et al., 2014). Given this core 714 has glacimarine material dated at  $22.6 \pm 0.3$  ka BP it indicates that the glacial isostatic loading of this portion of the Malin Shelf was at least 20 m and likely significantly more for the 715 laminated glacimarine muds to be deposited without wave based erosion or iceberg turbation 716 of the sediments. This suggests relative sea level was at least ~15 m higher than is currently 717 modelled for this region at 21 ka BP (Bradley et al., 2011). Thus, although global sea levels at 718 the LGM were ~134 m lower than present (Lambeck et al. 2014), the outer Malin Shelf (120-719 720 150 m and 160 m at the shelf break) would have still been below sea level. The effect of glacial isostatic adjustment (GIA) on the outer shelf between 27-21 ka BP has not been modelled with 721 722 any degree of certainty, but existing models extending back to 21 ka BP suggest that the outer shelf was still responding to ice loading at this time (Bradley et al., 2007). Theoretical 723 treatments and modelled simulations show that the stability of marine-based ice sheets is 724 closely coupled with the depth of water at the grounding line (Schoof, 2007). A reduction in 725 sea level can be enough to stop retreat, and an increase in sea level can cause instability and 726 enhance retreat (Gomez et al., 2012). We argue that the RSL rise associated with this isostatic 727 loading would be significant enough to cause the destabilisation of the BFIS, a conclusion 728 729 consistent with those put forward by Eyles and McCabe (1989).

730

731 Topographical controls leave a clear signature in the retreat history of the BFIS, with episodic retreat and minor re-advances on the topographical highs of the outer shelf. Furthermore, we 732 733 hypothesise that the more rapid and continued retreat across the mid shelf was aided by the presence of the two east-to-west trending troughs, the Malin Deep south of the Stanton Banks 734 735 and the second trough to the north that connects to the Inner Hebrides Trough. These overdeepened troughs were likely formed by ice streaming over several glacial periods (Davies et 736 737 al., 1984; Dunlop et al., 2010) across the softer bedrock (Permian to Cretaceous strata) cropping out in that sector. Much more resistant Dalradian basement bedrock crops out on the outer shelf 738 739 (Fyfe et al., 1993) and likely contributed in the formation a reverse slope, with greater erosion of the softer lithologies (c.f. Cook and Swift, 2012). This geometry of the ice-stream bed is 740 likely to focus and enhance calving of the ice margin as it retreated back into deeper waters, 741 thus providing a mechanism for rapid and irreversible retreat (c.f. Jamieson et al., 2012; Smith 742 et al., 2016). The ice-bed topography of the Malin Shelf was, therefore, a significant factor in 743 the early demise and pace of ice marginal retreat of the BFIS. 744

745

746 6. Conclusion

- By directly dating glacial landforms and sediments on the shelf, the data presented 747 here provides the best constraint on the extent and timing of retreat of the last BFIS, a 748 major draining system of the BIIS across the Malin Shelf. 749
- Glacial geomorphic features evident on the sub-bottom profiles confirm that the BFIS 750 reached the shelf edge during the last glacial period and its maximum extent 751 sometime before  $25.9 \pm 0.2$  ka BP 752
- Retreat across the shelf is marked by a series of large GZWs and smaller moraines 753 and at least one still-stand that occurred sometime  $\sim$ 25.9 ± 0.2 -24.8 ± 0.3 ka BP. The 754 outer and mid Malin Shelf was free of grounded ice by  $23.2 \pm 0.3$  ka BP, with the 755 inner shelf ice free by  $20.2 \pm 0.2$  ka BP. 756
- We conclude that the initial retreat of BFIS was caused by the growth of the BIIS that 757 increased isostatic loading on the Malin Sea shelf leading to a relative sea-level rise. 758 Additionally, topographic control in the form of over-deepened basins likely 759
- facilitated enhanced calving of the BFIS leading to rapid and irreversible retreat of 760 this ice stream once it started to retreat into deeper water. It is likely that the impact of 761 H2 and the northward migration of the Polar front at 23 ka BP aided further retreat of 762 the BFIS; however they cannot be the cause of its initial demise.

#### 764 Acknowledgments

This research was funded by the UK Natural Environment Research Council grant; BRITICE-765 CHRONO NE/J009768/1. The work was supported by the NERC Radiocarbon Facility 766 (allocation numbers 1722.0613 and 1878.1014). Thanks are due to the staff at the NERC AMS 767 Laboratory, East Kilbride for carbon isotope measurements. We thank the officers and crew of 768 the RRS James Cook for their help with acquisition and the British Geological Survey for 769 vibrocore collection during the cruise JC106. We also thank Kasper Weilbach, Riccardo 770 Arosio, Catriona Purcell, Zoe Roseby and Elke Hanenkamp for their scientific support on the 771 JC106 leg 2 cruise. The two anonymous reviewers are thanks for their detailed and constructive 772 773 comments.

774

763

#### 775 References

776 Alley, R.B, 2000. The Younger Dryas cold interval as viewed from central Greenland.

Quaternary Science Reviews, 19, 213-226. 777

- Anderson, J.B., Brake, C.F., Myers, N.C., 1984. Sedimentation on the Ross Sea continental 778
- shelf, Antarctica. Marine Geology, 57, 295-333. 779

- Anderson, J.B., Jakobsson, M., 2016.Grounding-zone wedges on Antarctic continental
   shelves. Geological Society, London, Memoirs, 46, 243-244.
- Austin, W.E.N., Bard, E., Hunt, J.B., Kroon, D., Peacock, J.D., 1995. The <sup>14</sup>C age of the
   Icelandic Vedde Ash: implications for Younger Dryas marine reservoir age corrections.
   Radiocarbon, 37, 53-62.
- Batchelor, C.L., Dowdeswell, J.A., Hogan, K.A., 2011. Late Quaternary ice flow and
  sediment delivery through Hinlopen Trough, Northern Svalbard margin: Submarine
  landforms and depositional fan. Marine Geology, 284, 13-27.
- Batchelor, C.L., Dowdeswell, J.A., 2015. Ice-sheet grounding-zone wedges (GZWs) on highlatitude continental margins. Marine Geology, 363, 65-92.
- Benetti, S., Dunlop, P., Ó Cofaigh, C., 2010. Glacial and glacially-related features on the
  continental margin of northwest Ireland mapped from marine geophysical data. Journal of
  Maps, 6 (1), 14-29.
- Binns, P.E., Harland, R., Hughes, M.J., 1974. Glacial and post-glacial sedimentation in the
  Sea of Hebrides. Nature, 48, 751–754.
- Bradwell, T., Stoker, M.S., Golledge, N.R., Wilson, C.K., Merritt, J.W., Long, D., Everest,
  J.D., Hestvik, O.B., Stevenson, A.G., Hubbard, A.L., Finlayson, A.G., Mathers, H.E.,
- 797 2008. The northern sector of the last British Ice Sheet: Maximum extent and demise. Earth
  798 Science Reviews, 88, 207-226.
- Bradley, S.L., Milne, G.A., Shennan, I., Edwards, R., 2011. An improved glacial isostatic
  adjustment model for the British Isles. Journal of Quaternary Science, 26, 541-552.
- Brooks, A.J., Bradley, S.L., Edwards, R.J., Milne, G.A., Horton, B., Shennan, I., 2008.
  Postglacial relative sea-level observations from Ireland and their role in glacial rebound
  modelling. Journal of Quaternary Science, 23, 175-192.
- Cai, J., Powell, R.D., Cowan, E.A., Carlson, P., 1997. Lithofacies and seismic-reflection
  interpretation of temperate glacimarine sedimentation in Tarr Inlet, Glacier Bay, Alaska.
  Marine Geology, 143, 5-37.
- Chappell, J., 2002. Sea level changes forced ice breakouts in the Last Glacial cycle: new
  results from coral terraces. Quaternary Science Reviews, 21, 1229–1240.
- 809 Clark, P. U., A. S. Dyke, J. D. Shakun, A. E. Carlson, J. Clark, B. Wohlfarth, J. X. Mitrovica, 810 S. W. Hostatlar, and A. M. McCaba (2000). The Last Classical Maximum, Science
- S. W. Hostetler, and A. M. McCabe (2009), The Last Glacial Maximum, Science,
  325(5941), 710–714.
- Clark, C.D., Hughes, A.L., Greenwood, S.L., Jordan, C. and Sejrup, H.P., 2012. Pattern and
  timing of retreat of the last British-Irish Ice Sheet. Quaternary Science Reviews, 44,112146.
- Clark, C.D., Ely, J.C., Greenwood, S.L., Hughes, A.L.C., Meehan, R., Barr, I.D., Bateman,
  M.D., Bradwell, T., Doole, J., Evans, D.J.A., Jorden, C.J., Monteys, X., Pellicer, X.M.,
  Sheehy, M.,2017. BRITICE Glacial Map, version 2: a map and GIS database of glacial
- 818 landforms of the last British–Irish Ice Sheet. https://doi.org/10.1111/bor.12273. ISSN
  819 0300-9483.
- 820 Cook, S.J., Swift, D.A., 2012. Subglacial basins: Their origin and importance in glacial
- systems and landscapes. Earth-Science Reviews, 115, 332-372

- Cowan, E.A., Cai, J., Powell, R.D., Clark, J.D., Pitcher, J.N., 1997. Temperate glacimarine
  varves: an example from Disenchantment Bay, southern Alaska. Journal of Sedimentary
  Research 67,536-549.
- Davies, H.C., Dobson, M.R., Whittington, R.J., 1984. A revised seismic stratigraphy for
  Quaternary deposits on the inner continental shelf west of Scotland between 55°30'N and
  57°30'N. Boreas, 13, 49-66.
- B28 Dobson, M.R., Whittington, R.J., 1992. Aspects of the geology of the Malin Sea area.
  B29 Geological Society Special Publication, 62, 291-311.
- Bomack, E.W., Jacobson, E.A., Shipp, S., Anderson, J.B., 1999. Late Pleistocene–Holocene
  retreat of the West Antarctic Ice-Sheet system in the Ross Sea: Part 2—sedimentologic
  and stratigraphic signature. Geological Society of America Bulletin 111, 1517-1536.
- Booley, H. D., and Crease, J. 1978. Observed and geostrophic currents south and east of
  Faroe during Overflow '73. ICES CM 1978/C: 53.
- Bove, D., Arosio, R., Finlayson, A., Bradwell, T., Howe, J.A., 2015. Submarine glacial
  landforms record Late Pleistocene ice-sheet dynamics, Inner Hebrides, Scotland.
  Quaternary Science Reviews, 123, 76-90.
- Bowdeswell, J., Fugelli, E., 2012. The seismic architecture and geometry of grounding-zone
  wedges formed at the marine margins of past ice sheets. Geological Society of America
  Bulletin, 124, 1750-1761.
- B41 Dowdeswell, J.A., Elverhøi, A., Spielhagen, R., 1998. Glacimarine sedimentary processes
  and facies on the Polar North Atlantic Margins. Quaternary Science Reviews, 17, 243-272.
- B43 Dowdeswell, J.A., Whittington, R.J., Jennings, A.E., Andrews, J.T., Mackensen, A.,
- Marienfeld, P., 2000. An origin for laminated glacimarine sediments through sea-ice
  build-up and suppressed iceberg rafting. Sedimentology, 47, 557-576.
- B46 Dowdeswell, J.A., Evans, J., O Cofaigh, C., 2010. Submarine landforms and shallow acoustic
  stratigraphy of a 400 km-long fjord-shelf-slope transect, Kangerlussuaq margin, East
  Greenland. Quaternary Science Reviews, 29, 3359-3369.
- Bunlop, P., Shannon, R., McCabe, M., Quinn, R., Doyle, E., 2010. Marine geophysical
  evidence for ice sheet extension and recession on the Malin Shelf: New evidence for the
  western limits of the British Irish Ice Sheet. Marine Geology, 276, 86-99.
- Ellett, D.J., 1979. Some oceanographic features of Hebridean waters. Proceedings of the
  Royal Society of Edinburgh, 77B, 61–74.
- Elverhøi, A., Norem, H., Andersen, E.S., Dowdeswell, J.A., Fossen, I., Haflidason, H.,
  Kenyon, N.H., Laberg, J.S., King, E.L., Sejrup, H.P., Solheim, A., Vorren, T., 1997. On
  the origin and flow behaviour of submarine slides on deep-sea fans on the Norwegian–
- Barents Sea continental margin. Geo-Marine Letters, 17, 119–125.
- Evans, D.J.A., Phillips, E.R., Hiemstra, J.F. and Auton, C.A., 2006. Subglacial till:
  formation, sedimentary characteristics and classification. Earth Science Reviews, 78, 115176.
- 861 Evans, J., Hogan, K.A., 2016. Grounding-zone wedges on the northern Larsen shelf,
- Antarctic Peninsula. Geological Society, London, Memoirs, 46, 237-238.
- 863 Eyles, N., Eyles, C.H., Miall, A.D., 1983. Lithofacies types and vertical profile models; an
- alternative approach to the description and environmental interpretation of glacial diamict
- and diamictite sequences. Sedimentology, 30, 393-410.

- Eyles, N., McCabe, A.M., 1989. The Late Devensian (< 22,000 BP) Irish Sea Basin: the</li>
  sedimentary record of a collapsed ice sheet margin. Quaternary Science Reviews, 8, 307351.
- Faugères, J-C., Stow, D.A., 1993. Bottom-current-controlled sedimentation: a synthesis on
  the contourite problem. Sedimentary Geology, 82, 287-297.
- Finlayson, A., Fabel, D., Bradwell, T., Sugden, D., 2014. Growth and decay of a marine
  terminating sector of the last British-Irish Ice Sheet: a geomorphological reconstruction.

873 Quaternary Science Reviews, 83, 28-45.

- Fyfe, J.A., Long, D., Evans, D., 1993. The Geology of the Malin- Hebrides Sea Area. HMSO
  for the British Geological Survey, London.
- Gomez, N., Pollard, D., Mitrovica, J.X., Huybers, P., Clark, P.U., 2012. Evolution of a
  coupled marine ice sheet-sea level model. Journal of Geophysical Research, 117, F01013
- Harkness, D. D. and Wilson, H. W. 1979. Scottish Universities Research and Reactor Centre
  Radiocarbon Measurements III. Radiocarbon, 21, 203-56.
- Hedges, R. E. M., Housley, R. A., Law, I. A., Perry, C. and Hendy, E. 1988. Radiocarbon
  dates from the Oxford AMS System: Archaeometry Datelist 9. Archaeometry, 30, 291305.
- Hibbert, F.D., Austin, W.E.N., Leng, M.J., Gatliff, R.W., 2010. British ice sheet dynamics
  inferred from north Atlantic ice-rafted debris records spanning the last 175,000 years.
  Journal of Quaternary Science 25, 461-482.
- Hindmarsh, R.C.A., 2018. Ice Sheets and Glacier Modelling, in: Menzies, J., van der Meer,
  J.J.M. (Eds.), Past Glacial Environments. Elsevier Ltd., Amsterdam, pp. 605-661.
- Hogan, K.A., Dowdeswell, J., O Cofaigh, C., 2012. Glacimarine sedimentary processes and
  depositional environments in an embayment fed by West Greenland ice streams. Marine
  Geology, 311, 1-16.
- Hogan, K.A., Ó Cofaigh, C., Jennings, A.E., Dowdeswell, J.A., Hiemstra, J.F., 2016.
  Deglaciation of a major palaeo-ice stream in Disko Trough, West Greenland, Quaternary
  Science Reviews, 147, 5-26.
- Howe, J.A., Dove, D., Bradwell, T. and Gafeira, J., 2012. Submarine geomorphology and
  glacial history of the Sea of the Hebrides, UK. Marine Geology, 315, pp.64-76.
- Howe, J.A., Stoker, M.S., Woolfe, K.J., 2001. Deep-marine seabed erosion and gravel lags in
  the northwestern Rockall Trough, North Atlantic Ocean. Journal of the Geological
  Society, 158, 427.
- Huthnance, J.M., 1986. Rockall slope current and shelf edge processes. Proc. R. Soc.
  Edinburgh 88B, 83-101.
- Jamieson, S.S.R., Viela, A., Livingstone, S.J., Ó Cofaigh, C., Stokes, C., Hillenbrand, C-D.,
  Dowdeswell, J.A., 2012. Ice-stream stability on a reverse bed slope. Nature Geoscience, 5,
  799-802.
- Jennings, A.E., 1993. The Quaternary history of Cumberland Sound, southeastern Baffin
  Island: the marine evidence. GeHographie Physique et Quaternaire, 47, 21-42.
- Jennings, A.E., Weiner, N.J., 1996. Environmental change in eastern Greenland during the
- 907last 1300 years: evidence from foraminifera and lithofacies in Nansen Fjord 683N. The
- 908 Holocene, 6, 179-191.

- Jennings, A.E., Walton, M.E., Ó Cofaigh, C., Kilfeather, A., Andrews, J.T., Ortiz, J.D., De
- 910 Vernal, A., Dowedeswell, J.A., 2014. Paleoenvironments during Younger Dryas-Early
  911 Holocene retreat of Greenland Ice Sheet from outer Disko Trough, central west Greenland.
  912 Journal of Quaternary Science, 29, 27-40.
- Joughin, I., Smith, B.E., Medley, B., 2014. Marine ice sheet collapse potentially under way
  for the Thwaites Glacier Basin, West Antarctica. Science, 344, 735-738.
- 915 Knutz, P.C., Austin, W.E. and Jones, E.J.W., 2001. Millennial- scale depositional cycles
- related to British Ice Sheet variability and North Atlantic paleocirculation since 45 kyr BP,
  Barra Fan, UK margin. Paleoceanography, 16, 53-64.
- Kroon, D., Simmield, G., Austin, W.E.N., Derrick, S., Knutz, P., Shimmield, T., 2000.
  Century- to millennial-scale sedimentological–geochemical records of glacial-Holocene
  sediment variations from the Barra Fan (NE Atlantic). Journal of the Geological Society,
  157, 643–653.
- Laberg, J.S., Vorren, T.O., 1995. Late Weichselian submarine debris flow deposits on the
  Bear Island Trough Mouth Fan. Marine Geology 127, 45–72.
- Lambeck, K., Rouby, H., Purcell, A., Sun, Y., Sambridge, M., 2014. Sea level and global ice
  volumes from the last glacial maximum to the Holocene. Proc. Natl. Acad. Sci. U. S. A.,
  111, 15296e15303.
- McCabe, A.M. and Clark, P.U., 2003. Deglacial chronology from County Donegal, Ireland:
  implications for deglaciation of the British–Irish ice sheet. Journal of the Geological
  Society, 160,847-855.
- Mouginot, J., Rignot, E., Scheuchl, B., Fenty, I., Khazendar, A., Morlighem, M., Buzzi, A.,
  Paden, J., 2015. Fast retreat of Zachariæ Isstrøm, northeast Greenland. Science 350, 13571361.
- Ó Cofaigh, C., Dowdeswell, J.A., 2001. Laminated sediments in glacimarine environments:
  diagnostic criteria for their interpretation. Quaternary Science Reviews 20, 1411-1436.
- Ó Cofaigh, C., Evans, J., Dowdeswell, J.A., Larter, R.D., 2007. Till characteristics, genesis
   and transport beneath Antarctic paleo-ice streams. Journal of Geophysical Research, 112
- 937 Ó Cofaigh, C., Dunlop, P., Benetti, S., 2012. Marine geophysical evidence for Late
  938 Pleistocene ice sheet extent and recession off northwest Ireland. Quaternary Science
  939 Reviews, 44, 147-159.
- 940 Ó Cofaigh, C., Andrews, J., Jennings, A., Dowdeswell, J., Hogan, K., Kilfeather, A.,
- Sheldon, C., 2013. Glacimarine lithofacies, provenance and depositional processes on a
  West Greenland trough-mouth fan. Journal of. Quaternary. Science, 28, 13-26.
- 943 Ó Cofaigh, C., Hogan, K.A., Dowdeswell, J.A., Streuff, K., 2016. Stratified glacimarine
- basin-fills in West Greenland fjords. Geological Society, London, Memoirs, 46, 99-100.
- Peacock, J.D., 2008. Late Devensian palaeoenvironmental changes in the sea area adjacent to
  Islay, SW Scotland: implications for the deglacial history of the island. Scottish Journal of
  Geology, 44, 183-190.
- 948 Peacock, J.D., Austin, W.E.N., Selby, I., Graham, D.K., Harland, R., Wilkinson, I.P., 1992.
- 949 Late Devensian and Flandrian palaeoenvironmental changes on the Scottish continental
- shelf west of the Outer Hebrides. Journal of Quaternary Science, 7, 145-161.

- 951 Peacock, J.D., Horne, D.J., Whittaker, J.E., 2012. Late Devensian evolution of the marine offshore environment of western Scotland. Proceedings of the Geologists' Association, 952 123, 419-437. 953 Peck, V.L., Hall, I.R., Zahn, R., Elderfield, H., Grousset, F., Hemming, S.R., Scourse, J.D., 954 955 2006. High resolution evidence for linkages between NW European ice sheet instability 956 and Atlantic meridional overturning circulation. Earth and Planetary Science Letters, 243, 476-488. 957 Peck, V.L., Hall, I.R., Zahn, R., Elderfield, H., 2008. Millennial-scale surface and subsurface 958 paleothermometry from the northeast Atlantic, 55e8 ka BP. Paleoceanography, 23, 959 PA3221. 960 Peters, J.L., Benetti, S., Dunlop, P., Ó Cofaigh, C., 2015. Maximum extent and dynamic 961 behaviour of the last British Irish ice sheet west of Ireland. Quaternary Science Reviews, 962 128, 48-68. 963 Peters, J.L., Benetti, S., Dunlop, P., Ó Cofaigh, C., Moreton, S.G., Wheeler, A.J., Clark, 964 C.D., 2016. Sedimentology and chronology of the advance and retreat of the last British-965 Irish Ice Sheet on the continental shelf west of Ireland. Quaternary Science Reviews, 140, 966 101-124. 967 968 Powell, R.D., Dawber, M., McInnes, J.N., Pyne, A.R., 1996. Observations of the groundingline area at a floating glacier terminus. Annals of Glaciology, 22, 217-223. 969 Praeg, D., McCarron, S., Dove, D., Ó Cofaigh, C., Scott, G., Monteys, X., Facchin, L., 970 Romeo, R., Coxon., 2015. Ice sheet extension to the Celtic Sea shelf edge at the Last 971 Glacial Maximum. Quaternary Science Reviews, 111, 107-112. 972 Reid, D.G., Turrell, W.R., Walsh, M., Corten, A., 1997. Cross-shelf processes north of 973 Scotland in relation to the southerly migration of Western mackerel. ICES Journal of 974 Marine Science, 54, 168-178. 975 Reimer, P.J., Bard, E., Bayliss, A., Beck, J.W., Blackwell, P.G., Ramsey, C.B., Buck, H., 976 977 Cheng, H., Edwards, R.L., Friedrich, M., Grootes, P.M., Guilderson, T.P., Hafildason, H., 978 Hajdas, I., Hatté, C., Heaton, T.J., Hoffman, D.L., Hogg, A.G., Hughen, K.A., Kaiser, K.F., Kromer, B., McCormac, F.G., Manning, S.W., Nui, M., Reimer, R.W., Richards, 979 D.A., Scott, E.M., Southon, J.R., Staff, R.A., Turney, C.S.M., van der Plicht, J., 2013. 980 981 Intcal13 and Marine13 radiocarbon age calibration curves, 0-50,000 years Cal Bp. Radiocarbon, 55, 1869-1887. 982 Sacchetti, F., Benetti, S., Georgiopoulou, A., Shannon, P.M., O'Reilly, B.M., Dunlop, P., 983 Quinn, R., Ó Cofaigh, C., 2012. Deep-water geomorphology of the glaciated Irish margin 984 from high-resolution marine geophysical data. Marine Geology, 291-294, 113-131. 985 Schoof, C., 2007. Ice sheet grounding line dynamics: Steady states, stability, and hysteresis. 986 Journal of Geophysical Research, 112, F03S28, doi:10.1029/2006JF000664. 987 Scourse, J.D., Haapaniemi, A.I., Colmenero-Hidalgo, E., Peck, V.L., Hall, I.R., Austin, W.E., 988 989 Knutz, P.C. and Zahn, R., 2009. Growth, dynamics and deglaciation of the last British-990 Irish ice sheet: the deep-sea ice-rafted detritus record. Quaternary Science Reviews, 28, 3066-3084. 991 Selby, I. 1989. Quaternary Geology of the Hebridian Continental Margin. Unpublished PhD 992
- 993 thesis, Nottingham University.

- 994 Siddall, M., Rohling, E.J., Almogi-Labin, A., Hemleben, C., Meischner, D., Schmelzer, I., Smeed, D.A., 2003. Sea-level fluctuations of the last glacial cycle. Nature, 423, 853–858. 995 Singarayer, J.S., Richards, D.A., Ridgwell, A., Valdes, P.J., Austin, W.E.N., Beck, J.W., 996 2008. An oceanic origin for the increase of atmospheric radiocarbon during the Younger 997 998 Drvas. Geophysical Research Letters, 35, L14707. 999 Small, D., Austin, W., Rinterknecht, V., 2013a. Freshwater influx, hydrographic 1000 reorganization and the dispersal of ice-rafted detritus in the sub-polar North Atlantic Ocean during the last deglaciation. Journal of Quaternary Science, 28, 527-535. 1001 Small, D., Parrish, R.R., Austin, W.E.N.A., Cawood, P.A., Rinterknecht, V., 2013b. 1002 Provenance of North Atlantic ice rafted debris during the last deglaciation - A new 1003 application of U-Pb rutile and zircon geochronology. Geology, 41, 155-158. 1004 1005 Small, D., Rinterknecht, V., Austin, W.E.N., Bates, C.R., Benn, D.I., Scourse, J.D., Bourlès, D.L., ASTER Team and Hibbert, F.D., 2016. Implications of 36Cl exposure ages from 1006 Skye, northwest Scotland for the timing of ice stream deglaciation and deglacial ice 1007 dynamics. Quaternary Science Reviews, 150, 130-145. 1008 Small, D., Benetti, S., Dove, D., Ballantyne, C.K., Fabel, D., Clark, C.D., Gheorghiu, D.M., 1009 1010 Newall, J., Xu, S., 2017. Cosmogenic exposure age constraints on deglaciation and flow 1011 behaviour of a marine-based ice stream in western Scotland, 21-16 ka. Quaternary Science Reviews, 167, 30-46. 1012 Stoker, M.S., 1995. The influence of glacigenic sedimentation on slope-apron development 1013 1014 on the continental margin off NW Britain. In: Scrutton, R.A., Stoker, M.S., Shimmield, G.B., Tudhope, A.W (Eds.), The Tectonics, Sedimentation and Palaeoceanography of the 1015 North Atlantic Region. Geological Society London, Special Publication, 90, 159-177. 1016 Stoker, M.S., Hitchen, K., Graham, C.C., 1993. United Kingdom Offshore regional report: 1017 the geology of the Hebrides and West Shetland shelves and adjacent deep-water areas. 1018 HMSO for the British Geological Survey, London. 1019 1020 Stow, D.A.V., Faugères, J-C., Howe, J.A., Pudsey, C.J., Viana, A., 2002. Bottom currents, 1021 contourites and deep sea sediment drifts: current state-of-the-art. Geological Society of London Memoirs, 22, 7-20. 1022 Svendsen, J.I., Mangerud, J., Elverhm, A., Solheim, A. and Schiillenhelm, R.T.E., 1992. The 1023 1024 Late Weichselian glacial maximum on western Spitsbergen inferred from offshore 1025 sediment cores. Mar. Geol., 104, 1-17. Syvitski, J.P.M., Farrow, G.E., Atkinson, R.J.A., Moore, P.G., Andrews, J.T., 1989. Baffin 1026 1027 Island fjord macrobenthos: bottom communities and environmental significance. Arctic, 1028 42, 232-247. Viana, A.R., Faugères, J-C., Stow, D.A.V., 1998. Bottom-current-controlled sand deposits -1029 a review of modern shallow-to deep-water environments. Sedimentary Geology, 115, 53-1030 1031 80. 1032 Wanamaker Jr, A.D., Butler, P.G., Scourse, J.D., Heinemeier, J., Eiríksson, J., Knudsen, 1033 K.L., Richardson, C.A., 2012. Surface changes in the North Atlantic meridional overturning circulation during the last millennium. Nature communications, 3, 899. 1034
- 1035 Wilson, L.J., Austin, W.E.N., Jansen, E., 2002. The last British Ice Sheet: growth, maximum
- extent and deglaciation. Polar Research, 21, 243–250.

1037 Winkelmann, D., Scháfer, C., Stein, R., Mackensen, A., 2008. Terrigenous events and

climate history of the Sophia Basin, Arctic Ocean. Geochemistry, Geophysics, Geosystems, 1038

1039 9, Q07023.

1040

1041

#### 1042 **Figures and Tables**

1043 Fig. 1 (a) Regional schematic map showing the maximum extent of the British Irish Ice Sheet during the last glacial, modified from Peters et al. (2015) with the proposed flow-line 1044 1045 directions of the major ice streams with ice-marginal and Donegal Barra Fan positions previously published by Bennetti et al (2010), Dunlop et al. (2010) Ó Cofaigh et al. (2012), 1046 Sacchetti et al. (2012) and Clark et al. (2017) and locations mentioned in the text (OH: Outer 1047 Hebrides, KB: St Kilda Basin, S: Skye, NC: North Channel, DBF: Donegal Barra Fan); Black 1048 box marks the study area; (b) The Malin Sea continental shelf showing the labelled core 1049 1050 locations (red circles) and seismic profiles shown in Figs 3 and 4 (black lines labelled MS1-1051 MS5).

1052 1053

Fig. 2. Acoustic facies examples identified in the sub-bottom profile data on the Malin Shelf 1054 1055 and upper continental slope

1056

Fig. 3. Acoustic profiles of MS1-MS3 with schematic illustration of each acoustic profile 1057 1058 below the original data where a) is MS1, b) MS2 and c) MS3. Note the different scales used for each seismic profile. Core locations are marked with a red circle. 1059

1060

1061 Fig. 4. Acoustic profiles of MS4-MS5 with schematic illustration of each acoustic profile below the original data where a) is MS4, b) MS5. Note the different scales used for each 1062 1063 seismic profile. Core locations are marked with a red circle.

1064

Fig. 5. Representative X-radiographs of lithofacies associations LFA1-6 and LFA7-11. 1065 1066

1067 Fig. 6. Core logs of key cores mentioned in the text where a) cores located on MS3, b) cores 1068 located on MS2, c) cores located on MS1, d) cores located on MS4, e) cores located on MS5. Shear strength measurements, >2 mm clast counts, magnetic susceptibility and bulk density 1069 results are added to these logs along with the median radiocarbon ages. 1070

1071

Fig. 7. Greyscale hillshade image of the Malin Sea shelf, with previously published glacial 1072 geomorphology shaded in brown (Dunlop and Benetti) and new features identified by the 1073 1074 author, and ground truthed using acoustic sub-bottom profiles and core data, in blue. The location of the seismic lines shown in Fig. 3-4 and the cores used to constrain the chronology 1075 of the BFIS are also added, along with proposed isochrones (dashed black lines) of the BFIS 1076 margins. Source of hillshade image from EMODnet Bathymetry portal (www.emodnet-

1077

1078 bathymetry.eu).

1079	
1080	Fig. 8. Schematic diagram of the retreating BFIS from (a) its last glacial maximum extent
1081	position at 26.7 $\pm$ 0.3 ka BP, (b) early deglaciation and second grounding line position on the
1082	outer shelf between 26.7 $\pm$ 0.3-25.9 $\pm$ 0.2 ka BP, (c) rapid retreat back thought the deep
1083	troughs, (d) mid-shelf grounding position at $23.2 \pm 0.3$ ka BP and (e) retreat onto the
1084	coastline by 20 ka BP.
1085	
1086	Table 1. Location, water depth and core recovery of cores collected from the Malin Sea
1087	
1088	Table 2. Radiocarbon results for cores discussed in this study
1089	
1090	Table 3. Lithofacies identified in the cores from the Malin Sea (After Eyles et al., 1983)
1091	
1092	
1093	
1094	
1095	
1096	
1097	
1098	
1099	
1100	
1101	
1102	
1103	
1104	
1105	
1106	
1107	
1108	
1109	
1110	
1111	
1112	
1113	
1114	
1115	
1116	
1117	
1118	
1119	
1120	
1121	
1122	



# 1134 Figure 2.

Acoustic Facies	Example	Description	Interpretation
AFA	15m	Basal reflector that is smooth to rugged (see arrowed reflector in AFC and AFF below) and is acoustically structureless below this reflector.	Bedrock
AFB		Strong upper reflector. Internally transparent to faintly stratified. Ranges in morphology and includes large ridges, small mounds, asymmetric wedges and stacked sheets.	Moraines and grounding zone wedges that mark ice marginal positions
AFC	a) 15m 15m 20m 2km	Internally acoustically transparent lenses (a) or wedges (b) geometries with weak boundary reflectors.	Debris flows
AFD	5m 500m	Chaotic to contorted internal acoustic stratification. Upper surface hummocky to truncated. Tends to directly overlie AFB.	Ice proximal glacimarine sediments containing a large IRD component. Possibly contorted by an osciliating ice margin nearby.
AFE	20m 2km	Acoustically stratified facies with continuous parallel reflectors. Often contains a stacked sequence of stratified units that truncate the underlying unit.	Ice distall glacimarine basin infills.
AFF	+ + + + + + + + + + + + + + + + + + +	Acoustically semi-transparent facies, forming a conformable drape. Forms the uppermost acoustics facies across the study area.	Post-glacial sand and mud, likely deposited during the Holocene.
AFG	10m 	Discrete v-shaped cut and fill geometries restricted to the surface of AFB	Iceberg furrows that have been subsequently infilled by younger sediment.
AFH	10m 	Stacked lenses of internal structureless facies with some point-source diffracted hyperbole. Occurrence restricted to the upper continental shelf.	Glacigenic debris flows.





























Table 1. Location, water depth and core recovery of cores collected from the Malin Sea							
Core name	Location	Water depth (m)	Recovery (m)				
091VC	55.4958 N, 8.0456 W	76	0.62				
114VC	55.6814 N, 9.4269 W	568.6	4.42				
115VC	55.6802 N, 9.3696 W	384	3.67				
116VC	55.6796 N, 9.3392 W	258	2.36				
118VC	55.6799 N, 9.3075 W	178	1.7				
119VC	55.6800 N, 9.2613 W	131	1.32				
120VC	55.6805 N, 9.0756 W	110.8	0.95				
121VC	55.6808 N, 8.9909 W	114	0.6				
122VC	55.6505 N, 8.4030 W	95	0.6				
123VC	55.6255 N, 8.4613 W	91	1.435				
124VC	55.7110 N, 9.3432 W	240	5.84				
125VC	55.7337 N, 9.2515 W	129	1.08				
129VC	56.1894 N, 8.5530 W	130	1.33				
130VC	56.1853 N, 8.5800 W	126	0.5				
131VC	56.1842 N, 8.8305 W	135	1.69				
135VC	56.4447 N, 9.1864 W	783	3.38				
136VC	56.1837 N, 8.8578 W	124	3.92				
137VC	56.1825 N, 8.9184 W	141	0.34				
138VC	56.1770 N, 8.9797 W	148.5	0.275				
139VC	56.1771 N, 8.9797 W	148.8	3.67				
140VC	56.1693 N, 9.0662 W	167.12	4.235				
141VC	56.1683 N, 9.1694 W	190	0.61				
142VC	56.1684 N, 9.1804 W	201	2.93				
143VC	56.4517 N, 9.0611 W	272	3.63				
144bVC	56.4521 N, 9.0562 W	243	6.08				
145VC	56.4669 N, 8.8744 W	147	3.98				
146VC	56.4730 N, 8.7070 W	150	4.14				
147VC	56.4840 N, 8.4464 W	158	6				
148VC	56.3867 N, 7.4274 W	121	5.02				
149VC	56.3973 N, 7.4488 W	136	4.5				
150VC	56.3997 N, 7.4537 W	140	5.27				
151VC	56.1405 N, 7.5377 W	122	4.1				
153VC	56.2517 N, 7.5874 W	113.5	3.35				
154VC	56.3253 N, 7.6181 W	138	2.7				

----

Table 2. Radiocarbon results for cores discussed in this study									
Core	Depth (cm bsf)	Sample material	$\delta^{13}C_{VPDB}$ %0 ± 0.1	Carbon content (% by wt.)	<sup>14</sup> C age (yrs BP)	Calibrated age range with 0 AR (yrs BP)	Calibrated age range with 300 ΔR (yrs BP)	Calibrated age range with 700 ΔR (yrs BP)	Laboratory code
115VC	365-367	Mixed benthic foraminifera	-1.425	13.8	19299±66	22751±229	22464±145	22058±231	SUERC-63566
116VC	140	Shell fragment	0.492	11.6	15487±41	18326±165	17981±153	17528±169	SUERC-59493
116VC	231-233	Mixed benthic foraminifera	-0.993	13.1	19299±58	22750±221	22463±133	22056±220	SUERC-63567
124VC	303-304	Shell fragment	1.233	10.7	12733±39	14312±272	13914±135	13472±125	SUERC-58376
124VC	305-307	Shell fragment	1.834	11	12675±39	14189±190	13854±129	13417±108	SUERC-58377
124VC	522-524	Mixed benthic foraminifera	-1.811	11.2	16025±44	18862±106	18598±129	18131±178	SUERC-63568
125VC	69.5	Shell fragment	1.043	11.44	11373±39	12826±129	12612±80	12042±204	SUERC-72872
125VC	115	Shell fragment	1.272	11.41	22813±61	26719±318	26310±235	25955±135	SUERC-72873
125VC	117	Shell fragment	1.007	11.52	22906±62	26825±302	26408±255	26018±147	SUERC-72874
130VC	45	Shell fragment	0.875	11.8	49782±1271	Out of range	Out of range	Out of range	SUERC-59482
136VC	232-233	Shell fragment	1.94	11.5	19854±52	23418±251	23078±249	22601±188	SUERC-58378
139VC	151-154	Mixed benthic foraminifera	-2.274	10.2	18920±80	22389±184	22082±249	21583±269	UCIAMS-176381
139VC	172	Shell fragment	1.288	11.7	50780±2520	Out of range	Out of range	Out of range	SUERC-59488
139VC	208	Bryozoa	0.123	9.1	11160±39	12662±88	12379±206	11591±267	SUERC-59489
139VC	305.5	Shell fragment	0.698	11.08	13017±39	14978±221	14276±253	13789±137	SUERC-58379
139VC	312	Shell fragments	0.878	10.6	Indistinguishable from background				SUERC-59483
139VC	331.5	Shell fragment	0.657	9	13048±38	15031±204	14345±284	13825±133	SUERC-59490
139VC	332.5	Shell fragment	1.512	10.4	11280±38	12749±114	12542±104	11866±202	SUERC-59491
139VC	348.5	Shell fragment	1.158	8.3	20397±77	24179±212	23850±225	23371±265	SUERC-59487
139VC	339	Shell fragment	0.675	10.7	12729±37	14299±263	13910±131	13467±120	SUERC-59486
139VC	Shoe	Shell fragment	0.415	11.5	13023±38	14989±216	14287±259	13796±135	SUERC-58382
142VC	150-151	Mixed benthic foraminifera	-0.647	10.9	18922±50	22402±123	22078±212	21593±216	SUERC-58325
142VC	160	Shell fragment	-0.391	9.6	50975±1485	Out of range	Out of range	Out of range	SUERC-59496
142VC	189.5	Shell fragment	-0.686	11.3	41614±474	44711±864	44443±892	44063±886	SUERC-59497

142VC	223	Shell fragment	0.857	11.74	15331±41	18138±174	17810±167	17324±188	SUERC-59498
142VC	269.5	Shell fragment	1.738	11.1	17269±51	20228±198	19983±194	19510±180	SUERC-59499
142VC	210	Shell fragment	-0.174	10.6	15496±41	18339±163	17991±153	17539±169	SUERC-59503
142VC	272	Shell fragment	0.405	11.3	Indistinguishable from background				SUERC-59500
142VC	285	Shell fragment	-0.845	11.5	Indistinguishable from background				SUERC-59501
143VC	323-325	Mixed benthic foraminifera	-0.993	11.9	15287±46	18090±174	17766±174	17271±196	SUERC-63565
143VC	330	Shell fragment	0.799	11.7	44474±689	Out of range	Out of range	Out of range	SUERC-59506
143VC	363	Shell fragment	-0.304	11.5	>53893	Out of range	Out of range	Out of range	SUERC-59508
143VC	Shoe	Shell fragment	1.321	11.8	Indistinguishable from background				SUERC-59507
144bVC	517-518	Shell fragment	0.067	11.6	Indistinguishable from background				SUERC-58383
145VC	256-257	Shell fragment	0.07	11.5	15431±41	18251±169	17919±155	$17455 \pm 174$	SUERC-58384
145VC	318-320	Shell fragment	0.196	10.4	18423±51	21824±196	21450±233	20886±202	SUERC-58385
146VC	223-225	Mixed benthic foraminifera	-7.491	9.8	20200±80	23828±251	23478±288	23007±290	UCIAMS-176382
146VC	369-372	Mixed benthic foraminifera	-3.325	10.5	22030±100	25897±178	25662±225	25265±302	UCIAMS-176383
146VC	389-392	Mixed benthic foraminifera	5.401	10.6	20730±100	24424±331	24089±270	23646±300	UCIAMS-164440
147VC	293	Shell fragment	2.942	11.8	Indistinguishable from background				SUERC-58386
147VC	422	Shell fragment	0.986	10.64	Indistinguishable from background				SUERC-72875
147VC	449-450	Shell fragment	5.507	11.91	Indistinguishable from background				UCIAMS-186925
147VC	489	Shell fragment	1.245	10.78	>49150	Out of range	Out of range	Out of range	SUERC-72876
147VC	492-493	Shell fragment	3.24	7.37	Indistinguishable from background				UCIAMS-186923
149VC	421	Shell fragment	-0.491	11.8	17155±47	20204±190	19846±202	19375±180	SUERC-59509
151VC	300-302	Yodiella sp	0.194	11.3	14320±41	16860±221	16411±200	15886±174	SUERC-67939
151VC	389	Mixed benthic foraminifera	-1.673	7.7	19690±90	23232±292	22853±286	22460±194	UCIAMS-179841
153VC	277-279	Mixed benthic foraminifera	1.158	16.1	19210±110	22679±270	22365±261	21956±321	UCIAMS-164432

	154VC	211-214	Mixed benthic foraminifera	3.744	12.7	18670±90	22129±261	21757±265	21229±304	UCIAMS-164433
1276										
1277										
1278										
1279										
1280										
1281										
1282										
1283										
1284										
1285										
1286										
1287										
1288										
1289										
1290										
1291										
1292										
1293										
1294										
1295										
1296										
1297										
1298										
1299										
1201										
1202										
1302										

Table 3. Lithofacies identified in the cores from the Malin Sea (After Eyles et al., 1983)					
Lithofacies	Description				
Dmm	Diamict, matrix supported and massive				
Dmmc	Diamict, matrix supported, consolidated (>40 kPa)				
Dms	Diamict, matrix supported and stratified				
Gm	Gravel, clast supported and massive				
Gms	Gravel, matrix supported				
Sfm	Muddy fine sand, massive				
Sm	Sand, massive				
Smg	Sand, massive with some gravel				
Ss	Sand, stratified				
Sl	Sand, laminated				
Suf	Sand, upward fining				
Fm	Mud/muddy sand, massive				
Fmd	Mud/muddy sand, with clast				
Fmb	Mud/muddy sand, with bioturbation				
Fmg	Mud/muddy sand, massive with gravel				
Fs	Mud/muddy sand, stratified				
Fsb	Mud/muddy sand, stratified with bioturbation				
Fl	Mud/muddy sand, laminated				
Fldef	Mud/muddy sand, deformed laminations				
Flb	Mud/muddy sand, laminated with bioturbation				