

## Rapid growth in greenhouse gas emissions from the adoption of industrial-scale aquaculture

Yuan, J.J.; Xiang, J.; Liu, Deyan; Kang, Hojeong; He, Teihu; Kim, S.; Lin, Y.; Freeman, Christopher; Ding, W.X.

### Nature Climate Change

DOI:

[10.1038/s41558-019-0425-9](https://doi.org/10.1038/s41558-019-0425-9)

Published: 01/04/2019

Peer reviewed version

[Cyswllt i'r cyhoeddiad / Link to publication](#)

*Dyfyniad o'r fersiwn a gyhoeddwyd / Citation for published version (APA):*

Yuan, J. J., Xiang, J., Liu, D., Kang, H., He, T., Kim, S., Lin, Y., Freeman, C., & Ding, W. X. (2019). Rapid growth in greenhouse gas emissions from the adoption of industrial-scale aquaculture. *Nature Climate Change*, 9(4), 318-322. <https://doi.org/10.1038/s41558-019-0425-9>

#### Hawliau Cyffredinol / General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

#### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

1 **Rapid growth in greenhouse gas emissions from the adoption of industrial-scale aquaculture**

2

3 **Running head:** Industrial-scale aquaculture increases global warming

4 Junji Yuan<sup>1†</sup>, Jian Xiang<sup>1†</sup>, Deyan Liu<sup>1</sup>, Hojeong Kang<sup>2</sup>, Tiehu He<sup>1,3</sup>, Sunghyun Kim<sup>1‡</sup>, Yongxin

5 Lin<sup>1</sup>, Chris Freeman<sup>4\*</sup>, Weixin Ding<sup>1\*</sup>

6 <sup>1</sup> *State Key Laboratory of Soil and Sustainable Agriculture, Institute of Soil Science, Chinese*  
7 *Academy of Sciences, Nanjing 210008, China.*

8 <sup>2</sup> *School of Civil and Environmental Engineering, Yonsei University, Seoul 120–749, Korea*

9 <sup>3</sup> *University of Chinese Academy of Sciences, Beijing 100049, China*

10 <sup>4</sup> *School of Natural Sciences, Bangor University, Gwynedd LL57 2UW, UK*

11 <sup>†</sup> These authors contributed equally to this work.

12 <sup>‡</sup> Present address: *Smithsonian Environmental Research Center, Edgewater, MD 21037, USA*

13

14 **\*Author for correspondence:**

15 **Weixin Ding:** Tel: 0086-25-86881527; Fax: 0086-25-86881000; Email: [wxding@issas.ac.cn](mailto:wxding@issas.ac.cn)

16 **Chris Freeman:** Tel: 44-1248-382353; Fax: 44-1248370731; Email: [c.freeman@bangor.ac.uk](mailto:c.freeman@bangor.ac.uk)

17

18 **Article type:** Primary Research Article

19 **Length of the text, methods and legends:** 2,099, 2,466 and 229 words, respectively

20 **The number of references:** 30

21 **The number of figures and tables:** 2 figures and 2 tables

22 **Final size of figures and tables:** Fig. 1, 31 Kb, 2-column width; Fig. 2, 52 Kb, 1-column width

23 **Abstract**

24 Fisheries capture has plateaued, creating ever-greater reliance on aquaculture to feed growing  
25 populations. Aquaculture volumes now exceed those of capture fisheries globally<sup>1,2</sup>, with China  
26 dominating production through major land-use change; more than half of Chinese freshwater  
27 aquaculture systems having been converted from paddy fields<sup>1,3</sup>. However, the greenhouse gas  
28 (GHG) implications of this expansion have yet to be effectively quantified. Here we measure  
29 year-round methane (CH<sub>4</sub>), nitrous oxide (N<sub>2</sub>O) and carbon dioxide (CO<sub>2</sub>) emissions from paddy  
30 fields and new, extensively managed crab aquaculture ponds. The conversion increased associated  
31 global warming potentials (GWP) from  $8.15 \pm 0.43$  to  $28.0 \pm 4.1$  Mg CO<sub>2</sub> eq ha<sup>-1</sup>, primarily due to  
32 increased CH<sub>4</sub> emission. After compiling a worldwide database of different freshwater aquaculture  
33 systems, the top 21 producers were estimated to release  $6.04 \pm 1.17$  Tg CH<sub>4</sub> and  $36.7 \pm 6.1$  Gg N<sub>2</sub>O  
34 in 2014. We found that 80.3% of total CH<sub>4</sub> emitted originated in shallow earthen aquaculture  
35 systems, with far lower emissions from intensified systems with continuous aeration<sup>4</sup>. We therefore  
36 propose greater adoption of aerated systems is urgently required to address globally significant rises  
37 in CH<sub>4</sub> emission from the conversion of paddy fields to aquaculture.

38 With increasing demand for animal proteins due to rising populations and a leveling off in capture  
39 fisheries, global aquaculture production has increased by 500% since the late-1980s, and now  
40 represents a major global industry<sup>1</sup>. In 2014, aquaculture volume amounted to 101 million tons (Mt)  
41 and is projected to reach 230 Mt by 2030, accounting for 62% of global fish and shellfish supply for  
42 human consumption<sup>1,2</sup>. This ever-expanding aquaculture sector relies heavily on application of  
43 aquafeeds<sup>5,6</sup> which increase nutrient loadings and carbon (C) burial in aquaculture systems and  
44 adjacent water bodies<sup>7,8</sup>. Only 25% (11–36%) of the nitrogen (N) consumed by fish was converted  
45 to biomass with the remainder excreted into water as un-ionized ammonia<sup>9,10</sup>. Likewise, a  
46 substantial proportion of feed C was transformed to CO<sub>2</sub> and CH<sub>4</sub> by animals and microbes<sup>11</sup> or  
47 buried in aquaculture systems<sup>7</sup>. In 2016, about 10.9 Tg C and 1.82 Tg N from the 39.9 Mt aquafeeds  
48 were estimated to be discharged to environments in global aquaculture<sup>12</sup>. Moreover, fertilizers are  
49 widely used in the extensive and semi-intensive aquaculture systems to stimulate phytoplankton  
50 production<sup>13</sup>. These intensive C and N loadings have the potential to drive aquaculture systems to  
51 become major anthropogenic sources of CH<sub>4</sub> and N<sub>2</sub>O emissions.

52 Williams & Crutzen<sup>14</sup> tentatively estimated N<sub>2</sub>O emission from the aquaculture sector at 0.09 Tg  
53 in 2008, accounting for 0.33% of global N<sub>2</sub>O emission. Using the N<sub>2</sub>O emission factor of influent N  
54 (EF<sub>N</sub> = 1.80%) in wastewater treatment plants<sup>15</sup>, global N<sub>2</sub>O emission from aquaculture was  
55 estimated to increase from 0.15 Tg in 2009 to 0.60 Tg in 2030, which could contribute 5.72% of  
56 global anthropogenic N<sub>2</sub>O emission<sup>10</sup>. However, large uncertainties in these estimates may arise  
57 from differences in management levels<sup>16,17</sup> and yield difference between species<sup>17,18</sup>. Besides N<sub>2</sub>O,  
58 aquaculture ponds could be important anthropogenic CH<sub>4</sub> sources with characteristics of intensive C  
59 loading, shallow water and frequent mixing<sup>19</sup>. To date, >40% of worldwide aquaculture production  
60 has been carried out in earthen ponds, while estimates of overall CH<sub>4</sub> budgets in global aquaculture

61 remain scarce.

62 China is the world's largest aquaculture producer, contributing ~60% of global volume<sup>1</sup>;  
63 furthermore the volume and area of that aquaculture is steadily rising<sup>3</sup>. Above 70% of Chinese  
64 freshwater aquaculture production is carried out in extensive and semi-intensive earthen ponds<sup>3</sup>.  
65 More and more paddy fields have been, and will continue to be, converted to aquaculture ponds.  
66 They currently account for 51.3% of Chinese inland fish ponds<sup>3,18</sup>. There is clearly an urgent need  
67 for greater appreciation of the costs associated with GHG emissions incurred during the ongoing  
68 unprecedented levels of conversion of paddy fields towards industrial-scale aquaculture.

69

#### 70 **Effect of conversion of paddy field to aquaculture on GHG emission**

71 We measured year-round fluxes of CH<sub>4</sub>, N<sub>2</sub>O and CO<sub>2</sub> from three adjacent crab aquaculture ponds  
72 converted from paddy fields 12 years ago and neighboring paddy fields (PF) in the Tai Lake basin  
73 (31°02'N, 120°25'E; Supplementary Figs. 1 and 2) during 2013–2014. Wheat-rice rotation is the  
74 typical cropping system in this region. Urea was applied in PF at 150 and 280 kg N ha<sup>-1</sup> during  
75 wheat and rice seasons, respectively. Crab ponds differed in size and water depth (Supplementary  
76 Table 1). They were not equipped with aerators but were fertilized during culturing. Chinese mitten  
77 crab (*Eriocheir sinensis*) were fed with commercial feed pellets, trash fish and corn seeds at the  
78 same rate in each pond during crab production period from March to October. Annual C and N  
79 inputs in crab ponds were 1.20 Mg C ha<sup>-1</sup> and 244 kg N ha<sup>-1</sup>, respectively (Supplementary Tables  
80 2–4).

81 Annual CH<sub>4</sub> emission in PF was 218 ± 7.28 kg CH<sub>4</sub> ha<sup>-1</sup> (Fig. 1a), which was located in the  
82 upper end of the previously reported ranges (98.3–240 kg CH<sub>4</sub> ha<sup>-1</sup>) for paddy fields without  
83 organic amendment in this area<sup>20</sup>. However, conversion from PF to crab ponds sharply increased

84 CH<sub>4</sub> emission to 962 ± 62 kg CH<sub>4</sub> ha<sup>-1</sup>; this value was higher than the summarized amount of 572  
85 kg CH<sub>4</sub> ha<sup>-1</sup> in permanently inundated temperate wetlands and the default emission factor (900 kg  
86 CH<sub>4</sub> ha<sup>-1</sup>) for tropical inland freshwater wetlands proposed by the Intergovernmental Panel on  
87 Climate Change (IPCC; ref. 21).

88 The CH<sub>4</sub> EF<sub>C</sub> of C inputs from feeds and fertilizer in crab ponds was estimated at up to 60.0%  
89 (Table 1), which may be attributed to the enhanced availability of labile organic substrates and  
90 highly anaerobic environment in crab ponds. The C output as harvested crab was 0.19 Mg C ha<sup>-1</sup>  
91 (Supplementary Table 4), accounting for 16.1% of the C inputs excluding the photosynthates by  
92 submerged macrophytes. The remaining 1.04 Mg C ha<sup>-1</sup> was deposited into sediments as  
93 unconsumed feed and feces, which led mean dissolved organic C (DOC) concentrations in pond  
94 sediments to reach 7.97-fold greater than that of PF (Supplementary Table 1). Additionally, organic  
95 compounds in feed remnants and feces such as starch and protein can be more easily decomposed<sup>22</sup>  
96 to methanogenic substrates than crop residues in PF. Moreover, pond sediments were permanently  
97 inundated, thereby creating anaerobic environments ideal for methanogenesis.

98 Annual N<sub>2</sub>O emission in PF was 7.11 ± 0.23 kg N<sub>2</sub>O ha<sup>-1</sup> (Fig. 1b). The EF<sub>N</sub> of fertilizer-N  
99 applied was 1.05%, closing to the IPCC default value (1.00%) for agricultural soils<sup>23</sup>. Conversion  
100 from PF to crab ponds significantly decreased annual N<sub>2</sub>O emission by 95.4% to 0.33 ± 0.07 kg  
101 N<sub>2</sub>O ha<sup>-1</sup>, with an EF<sub>N</sub> of 0.09 ± 0.02% (Table 1). The lower N<sub>2</sub>O emission in crab ponds was  
102 disproportionate to the differences in N application rates (244 vs 430 kg N ha<sup>-1</sup>), let alone the  
103 relatively higher total inorganic N content in pond sediments (Supplementary Tables 1). Nitrous  
104 oxide is derived from both nitrification and denitrification, although denitrification produces more  
105 N<sub>2</sub>O (ref. 24). It is likely that the much lower redox potential (-124 to -160 mV) suppressed  
106 nitrification in pond sediments, which reduced overall NO<sub>3</sub><sup>-</sup> concentrations to <1 mg N kg<sup>-1</sup>. This

107 concentration was lower than the threshold value of  $5 \text{ mg N kg}^{-1}$  for active denitrification<sup>25</sup>.  
108 Moreover, the high DOC concentrations and anaerobic conditions permit  $\text{N}_2\text{O}$  to be further reduced  
109 to  $\text{N}_2$  through denitrification<sup>26</sup>.

110 Using the net ecosystem C balance method, annual loss of soil organic C (SOC) in PF was  
111 estimated to be  $0.04 \pm 0.05 \text{ Mg C ha}^{-1}$  (Table 1), which fell in the range of  $-0.27$  to  $0.67 \text{ Mg C ha}^{-1}$   
112 estimated previously in paddy fields of Tai Lake basin<sup>27</sup>. The  $\text{CO}_2$  fluxes measured by transparent  
113 chambers in crab ponds were regarded as net ecosystem exchange. On an annual basis, crab ponds  
114 were weak net  $\text{CO}_2$  sources, releasing  $0.13\text{--}1.99 \text{ Mg CO}_2 \text{ ha}^{-1}$  (Fig. 1c).

115 Conversion from PF to extensive crab ponds increased the 100-yr GWP from  $8.15 \pm 0.43$  to  
116  $28.0 \pm 4.1 \text{ Mg CO}_2 \text{ eq ha}^{-1}$ , mainly due to increased  $\text{CH}_4$  emission with a contribution of 96.3%  
117 (Table 1). Our results contrast with those of Liu et al.<sup>18</sup>, who reported such conversion significantly  
118 reduced  $\text{CH}_4$  and  $\text{N}_2\text{O}$  emissions by 48% and 56%, respectively. Annual  $\text{CH}_4$  emission in Liu's  
119 ponds (equipped with aerators and classified as semi-intensive, see below) was just  $32.6 \text{ kg CH}_4 \text{ ha}^{-1}$   
120 despite the much greater feeding rate and higher sediment DOC concentration compared to test  
121 extensive ponds. Hence, substrate availability was not the limiting factor for  $\text{CH}_4$  emissions in  
122 feeding aquaculture systems, while oxygen exposure by aeration was the key factor affecting  $\text{CH}_4$   
123 emissions. Our results highlight that GHG emissions clearly differ from one aquaculture system to  
124 another, greatly depending on the intensity of operational management. This observation illustrates  
125 the potential for mitigating the effects of future paddy field conversion through careful  
126 management.

127

## 128 **Global $\text{CH}_4$ and $\text{N}_2\text{O}$ budgets of freshwater aquaculture**

129 Here, we classified aquaculture into four systems: rice-fish, extensive, semi-intensive and intensive

130 based on local conditions and aquaculture facilities especially whether aerators are used or not (see  
131 Methods). We compiled a worldwide database of CH<sub>4</sub> and/or N<sub>2</sub>O emissions that were measured in  
132 45 inland freshwater aquaculture systems during 2003–2015 (Supplementary Table 5). Land-use  
133 and production statistics were also compiled for different aquaculture systems of top 21 freshwater  
134 aquaculture producers (Supplementary Table 6); however, data from extensive and semi-intensive  
135 systems were pooled because of unavailability of aerator-use data for separate classification. In  
136 2014, the top 21 producers contributed 97.5% of global freshwater aquaculture volume<sup>1</sup>.

137 The synthesized data show that CH<sub>4</sub> fluxes ranged from –0.03 to 37.0 mg CH<sub>4</sub> m<sup>-2</sup> h<sup>-1</sup> in  
138 rice-fish, extensive, and semi-intensive systems. Mean CH<sub>4</sub> flux in rice-fish system was the highest  
139 at 12.6 ± 3.9 mg CH<sub>4</sub> m<sup>-2</sup> h<sup>-1</sup>, followed by extensive and semi-intensive systems (Fig. 2a). The  
140 absence of CH<sub>4</sub> flux in intensive systems can be attributed to a combination of continuous aeration,  
141 water exchange and a lack of habitats for methanogens<sup>4</sup>. The rice-fish system also had the highest  
142 mean N<sub>2</sub>O flux followed by semi-intensive and extensive systems (28.4 ± 9.8 and 7.56 ± 3.02 µg  
143 N<sub>2</sub>O m<sup>-2</sup> h<sup>-1</sup>, respectively; Fig. 2b). The EF<sub>N</sub> and yield-scale N<sub>2</sub>O EF (EF<sub>Y</sub>) in intensive system  
144 were 1.16 ± 0.18% and 2.48 ± 0.42 g N<sub>2</sub>O kg<sup>-1</sup> yield, respectively, which were significantly higher  
145 than the corresponding values in extensive (0.24 ± 0.10% and 0.66 ± 0.22 g N<sub>2</sub>O kg<sup>-1</sup> yield) and  
146 semi-intensive (0.35 ± 0.16% and 0.88 ± 0.41 g N<sub>2</sub>O kg<sup>-1</sup> yield) systems. The EF<sub>Y</sub> in intensive  
147 systems was close to the IPCC default EF<sub>Y</sub> (2.66 g N<sub>2</sub>O kg<sup>-1</sup> yield) that is widely used in model  
148 estimate for aquaculture<sup>10,21</sup>, but was 2.75- and 1.82-fold greater than that for extensive and  
149 semi-intensive systems, respectively. Considering the large volume of extensive and semi-intensive  
150 aquaculture (Supplementary Table 6), previous estimates<sup>10,14</sup> of global aquaculture N<sub>2</sub>O emission  
151 may have been overestimated because of the higher default EF<sub>Y</sub> mentioned above.

152 The estimated CH<sub>4</sub> and N<sub>2</sub>O emissions from the top 21 producers in 2014 were 6.04 ± 1.17 Tg



153 CH<sub>4</sub> and 36.7 ± 6.1 Gg N<sub>2</sub>O, respectively (Table 2), which accounted for 1.82% and 0.34% of  
154 global anthropogenic CH<sub>4</sub> and N<sub>2</sub>O emissions, respectively. Methane was a key contributor (94.6%;  
155 Fig. 2e) to GWP in freshwater aquaculture, of which 1.19 ± 0.27 Tg CH<sub>4</sub> was emitted from rice-fish  
156 system and 4.85 ± 1.04 Tg CH<sub>4</sub> from extensive plus semi-intensive systems. To our knowledge, this  
157 is the first global estimate of CH<sub>4</sub> emission from freshwater aquaculture. Our estimated total N<sub>2</sub>O  
158 emission was much lower than the previous estimates of 90 Gg N<sub>2</sub>O (ref. 14) and 146 Gg N<sub>2</sub>O (ref.  
159 10) of global aquaculture. Extensive plus semi-intensive systems contributed 87.0% of global  
160 volume of freshwater aquaculture, meanwhile, were the largest CH<sub>4</sub> and N<sub>2</sub>O emitter (80.3% and  
161 45.2%, respectively) from this sector. Intensive systems accounted for 8.89% of the production,  
162 27.0% of total N<sub>2</sub>O emissions but negligible CH<sub>4</sub> emissions. Rice-fish systems represented only  
163 4.30% of aquaculture volume, yet they accounted for 19.7% and 27.8% of CH<sub>4</sub> and N<sub>2</sub>O budgets,  
164 respectively.

165 The greenhouse gas intensity (GHGI, GWP/yield) was 3.59 ± 0.74 kg CO<sub>2</sub> eq kg<sup>-1</sup> yield in  
166 extensive plus semi-intensive systems, which was 4.46-fold greater than that in intensive systems  
167 (0.66 ± 0.11 kg CO<sub>2</sub> eq kg<sup>-1</sup> yield; Fig. 2f). Therefore, if half of the current productions from  
168 extensive plus semi-intensive systems (19.5 Mt) are replaced by intensive systems, the GWP of CH<sub>4</sub>  
169 and N<sub>2</sub>O emissions from freshwater aquaculture (excluding rice-fish) will be reduced by 40.1%  
170 from 143 Tg CO<sub>2</sub> eq to 85.6 Tg CO<sub>2</sub> eq.

171 China has emerged as the world's largest freshwater aquaculture emitter of CH<sub>4</sub> (4.10 ± 0.10 Tg  
172 yr<sup>-1</sup>) and N<sub>2</sub>O (22.8 ± 7.1 Gg yr<sup>-1</sup>), contributing 68.0% and 62.1% of global budgets from the sector,  
173 respectively. In China, CH<sub>4</sub> emissions from freshwater aquaculture with 7.57 × 10<sup>6</sup> ha equates to  
174 36.5% of total CH<sub>4</sub> emissions from paddy fields, natural wetlands and lakes (11.3 Tg CH<sub>4</sub> yr<sup>-1</sup>; ref.  
175 28). Since 83.0% of Chinese freshwater aquaculture CH<sub>4</sub> emissions originate from extensive plus

176 semi-intensive systems, a substantial reduction in emissions could be achieved through improved  
177 management practices, such as installing more efficient aerators in earthen ponds and implementing  
178 optimized feeding strategies for reducing feed waste.

179 In conclusion, the conversion of paddy fields to extensive crab aquaculture ponds sharply  
180 increased GWP, primarily through a drastic increase in CH<sub>4</sub> release. Our findings emphasize the  
181 need to assess the climatic impacts of land-use shifts towards industrial-scale aquaculture. Methane  
182 is the most important GHG in freshwater aquaculture compared with N<sub>2</sub>O, and it was primarily  
183 sourced from extensive plus semi-intensive systems. Our findings indicate that effective  
184 management of extensive and semi-intensive systems through conversion to intensive systems is  
185 urgently required to mitigate GHG emissions from the unprecedented growth of aquaculture.

186

## 187 **Methods**

188 Methods, including statements of data availability and any associated accession codes and  
189 references, are available in the online version of this paper.

190

## 191 **Data availability**

192 The authors declare that the data supporting the findings of this study are available within the article  
193 and its supplementary information files.

194

## 195 **References**

- 196 1. FAO. *The State of World Fisheries and Aquaculture 2016. Contributing to food security and*  
197 *nutrition for all* (FAO, 2016).
- 198 2. FAO. *The State of World Fisheries and Aquaculture 2014. Opportunities and challenges* (FAO,

- 199 2014).
- 200 3. Ministry of Agriculture of the People's Republic of China (MoA) *China Fisheries Yearbook*  
201 *2013* (MoA of China, Chinese Agric. Press, 2014) (in Chinese).
- 202 4. Hu, Z. et al. Influence of carbohydrate addition on nitrogen transformations and greenhouse gas  
203 emissions of intensive aquaculture system. *Sci. Total Environ.* **470**, 193–200 (2014).
- 204 5. Naylor, R. L. et al. Effect of aquaculture on world fish supplies. *Nature* **405**, 1017–1024 (2000).
- 205 6. Cao, L. et al. China's aquaculture and the world's wild fisheries. *Science* **347**, 133–135 (2015).
- 206 7. Boyd, C. E., Wood, C. W., Chaney, P. L. & Queiroz, J. F. Role of aquaculture pond sediments in  
207 sequestration of annual global carbon emissions. *Environ. Pollut.* **158**, 2537–2540 (2010).
- 208 8. Chatvijitkul, S., Boyd, C. E., Davis, D. A. & McNevin, A. A. Pollution potential indicators for  
209 feed-based fish and shrimp culture. *Aquaculture* **477**, 43–49 (2017).
- 210 9. Hargreaves, J. A. Nitrogen biogeochemistry of aquaculture ponds. *Aquaculture* **166**, 181–212  
211 (1998).
- 212 10. Hu, Z., Lee, J. W., Chandran, K., Kim, S. & Khanal, S. K. Nitrous oxide (N<sub>2</sub>O) emission from  
213 aquaculture: a review. *Environ. Sci. Technol.* **46**, 6470–6480 (2012).
- 214 11. Boyd, C. E. & Tucker, C. S. *Handbook for Aquaculture Water Quality* (Craftmaster Printers,  
215 2014).
- 216 12. Alltech. *2017 Alltech Global Feed Survey* (Alltech, 2017).
- 217 13. Green, B. W. in *Feed and Feeding Practices in Aquaculture* (ed. Davis, A. D.) 27–52  
218 (Woodhead Publishing, 2015).
- 219 14. Williams, J. & Crutzen, P. Nitrous oxide from aquaculture. *Nat. Geosci.* **3**, 143–143 (2010).
- 220 15. Ahn, J. H. et al. N<sub>2</sub>O emissions from activated sludge processes, 2008–2009: results of a  
221 national monitoring survey in the United States. *Environ. Sci. Technol.* **44**, 4505–4511 (2010).

- 222 16. Paudel, S. R. et al. Effects of temperature on nitrous oxide (N<sub>2</sub>O) emission from intensive  
223 aquaculture system. *Sci. Total Environ.* **518–519**, 16–23 (2015).
- 224 17. Hu Z. *A comparison of methane and nitrous oxide emissions between paddy fields and crab/fish*  
225 *farming wetlands in southeast China*. (Doctoral thesis, Nanjing Agri. Univ., 2015) (in Chinese).
- 226 18. Liu, S. et al. Methane and nitrous oxide emissions reduced following conversion of rice paddies  
227 to inland crab-fish aquaculture in southeast China. *Environ. Sci. Technol.* **50**, 633–642 (2016).
- 228 19. Holgerson, M. A. & Raymond, P. A. Large contribution to inland water CO<sub>2</sub> and CH<sub>4</sub> emissions  
229 from very small ponds. *Nat. Geosci.* **9**, 222–226 (2016).
- 230 20. Cai, Z., Tsuruta, H. & Minami, K. Methane emission from rice fields in China: measurements  
231 and influencing factors. *J. Geophys. Res. Atmos.* **105**, 17231–17242 (2000).
- 232 21. IPCC. *2013 Supplement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories:*  
233 *Wetlands* (eds Hiraishi, T. et al.) (IPCC, 2014).
- 234 22. Burford, M. A. & Williams, K. C. The fate of nitrogenous waste from shrimp feeding.  
235 *Aquaculture* **198**, 79–93 (2001).
- 236 23. IPCC. *2006 IPCC Guidelines for National Greenhouse Gas Inventories* Vol. 4 (eds Eggleston,  
237 H. S., Buendia, L., Miwa, K., Ngara, T. & Tanabe, K.) Ch. 11 (IGES, 2006).
- 238 24. Freeman, C., Lock, M. A., Reynolds, B. & Hudson, J. A. Nitrous oxide emissions and the use of  
239 wetlands for water quality amelioration. *Environ. Sci. Technol.* **50**, 2438–2440 (1997).
- 240 25. Dobbie, K. E. & Smith, K. A. Impact of different forms of N fertilizer on N<sub>2</sub>O emissions from  
241 intensive grassland. *Nutr. Cycl. Agroecosys.* **67**, 37–46 (2003).
- 242 26. Miller, M. N. et al. Crop residue influence on denitrification, N<sub>2</sub>O emissions and denitrifier  
243 community abundance in soil. *Soil Biol. Biochem.* **40**, 2553–2562 (2008).
- 244 27. Ma, Y. et al. Net global warming potential and greenhouse gas intensity of annual rice-wheat

245 rotations with integrated soil-crop system management. *Agric. Ecosyst. Environ.* **164**, 209–219  
246 (2013).28. Chen, H. et al. Methane emissions from rice paddies natural wetlands, lakes in  
247 China: synthesis new estimate. *Global Change Biol.* **19**, 19–32 (2013).  
248 29. Xie, Z. et al. CO<sub>2</sub> mitigation potential in farmland of China by altering current organic matter  
249 amendment pattern. *Sci. China Earth Sci.* **53**, 1351–1357 (2010).  
250 30. Frei, M. et al. Methane emissions and related physicochemical soil and water parameters in  
251 rice-fish systems in Bangladesh. *Agric. Ecosyst. Environ.* **120**, 391–398 (2007).

252

### 253 **Additional information**

254 **Correspondence and requests for materials** should be addressed to W.D. or C.F.

255

### 256 **Acknowledgements**

257 This work was supported by grants from the Chinese Academy of Sciences (XDB15020100), the  
258 Natural Science Foundation of Jiangsu Province (BK20151056) and China (41501274, 41471077),  
259 and the Special Fund for Forest Scientific Research in Public Welfare (201404210). H.K. is grateful  
260 to NRF (2016R1D1A1A02937049) and KFS (2017096A001719BB01).

261

### 262 **Author contributions**

263 W.D., J.Y., and D.L. designed the study; J.X. led the GHG fluxes and auxiliary measurements with  
264 the support of T.H., S.K., and Y.L. Site selection and set-up was carried out by J.Y., and D.L. H.K.  
265 and C.F. were the key international collaborators during this research. The manuscript was drafted  
266 by J.Y., H.K., W.D. and C.F with all authors contributing to the final version.

267

268 **Competing financial interests**

269 The authors declare no competing financial interests.

270 **Methods**

271 **Site description** Field experiments were carried out in a conventional paddy field (PF) and three  
272 adjacent crab ponds in the Tai Lake basin (31°02'N, 120°25'E), Suzhou City, Jiangsu Province,  
273 China (**Supplementary Fig. 1**). This region is characterized by a subtropical monsoon climate with  
274 the long-term (1981–2010) mean annual air temperature of 16.5°C and precipitation of 1176 mm  
275 (<http://cdc.nmic.cn/home.do>). Paddy fields accounted for 65% of total cropland in this region,  
276 however, they are being rapidly converted to aquaculture ponds due to the greater economic  
277 benefits from the latter since 1980s ([ref. 31](#)).

278 Soil was developed from alluvial sediments of the Yangtze River, and classified as stagnic  
279 Anthrosols based on the USDA soil taxonomy. The surface soil (0–20 cm) had a pH (H<sub>2</sub>O) of 5.95,  
280 bulk density of 1.25 g cm<sup>-3</sup> and a loam texture with 40% sand, 34% silt and 26% clay, and  
281 contained 20.3 g kg<sup>-1</sup> organic C and 1.81 g N kg<sup>-1</sup> total N. Three neighboring aquaculture ponds for  
282 Chinese mitten crab (*Eriocheir sinensis*) cultivation were converted from paddy fields in 2001.

283

284 **Experimental design and field management** Four independent 3 × 8 m<sup>2</sup> plots were established in  
285 PF in November 2012. Winter wheat (*Triticum aestivum* L., Yangfumai 4) and summer rice (*Oryza*  
286 *sativa* L., Wuyunjing 23) was rotated during the period from November 2012 to May 2014. During  
287 wheat season, urea was applied at the rate of 150 kg N ha<sup>-1</sup>, with basal and supplemental  
288 fertilization ratio of 40%:60%. During rice season, urea was applied at the rate of 280 kg N ha<sup>-1</sup>,  
289 with the basal and supplemental fertilizer ratio of 50%:50%. Calcium superphosphate (40 and 125  
290 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> for wheat and rice, respectively) and potassium sulfate (60 and 125 kg K<sub>2</sub>O ha<sup>-1</sup> for  
291 wheat and rice, respectively) were applied as basal fertilizers (**Supplementary Table 2**). The row  
292 distance was 25 cm for rice and wheat, and the hill distance was 15 cm for rice. No irrigation was

293 performed during wheat season, while rice was managed under a typical water regime mode of  
294 flooding-midseason drainage-reflooding-moist irrigation (F-D-F-M). Crop grain and straw were  
295 harvested and oven dried at 60°C until a constant weight.

296 Parallel field experiments were conducted in three neighboring crab ponds with different size  
297 (CP1, 1.71 ha; CP2, 0.71 ha; CP3, 0.09 ha), from March 2013 to March 2014. Monoculture of  
298 Chinese mitten crab was employed at the same stocking density of 15000 ind ha<sup>-1</sup> for each pond.  
299 The submerged western waterweed (*Elodea nuttallii*) naturally vegetated in ponds and provided  
300 molting shelters and foods for crabs. Feeds and fertilizers were applied at the same rates in each  
301 pond. Snails (*Bellamya quadrata*) were introduced into the ponds twice at the rates of 600 and 400  
302 kg ha<sup>-1</sup> on April 4 and June 20, respectively, to filter feed residue and provide supplementary foods  
303 for crabs (Supplementary Tables 2 and 3). Crabs were fed with commercial feed pellets (2050 kg  
304 ha<sup>-1</sup>) (Purina Co. Ltd., Jiaxing, China), trash fish (1250 kg ha<sup>-1</sup>) and corn seeds (1150 kg ha<sup>-1</sup>),  
305 twice per day on 9:00 a.m. and 17:00 p.m. until the crabs were harvested. In order to stimulate  
306 phytoplankton and waterweed production, cake manure (residue of de-oiled oil seeds) at 40 kg ha<sup>-1</sup>  
307 was applied as basal fertilizer while urea, compound fertilizer and calcium superphosphate were  
308 applied at the rate of 130, 100 and 200 kg ha<sup>-1</sup>, respectively, with four splits of 25%:25%:25%:25%  
309 on March 29, June 5, August 12 and September 27. Annual inputs of C and N to crab ponds were  
310 1.20 Mg C ha<sup>-1</sup> and 244 kg N ha<sup>-1</sup>, respectively. Water was constantly maintained all-year round,  
311 while the water depth differed between ponds. Crab harvest started from 1 to 30 October 2013,  
312 depending on crab maturity. Crab yield was expressed as fresh weight (Supplementary Table 4).  
313 Details management practices in the two systems are shown in Supplementary Table 3.

314

315 **Measurement of GHG fluxes** Wooden boardwalks were installed in each plot to facilitate



316 collecting gas samples and measuring the auxiliary parameters (Supplementary Fig. 2). The static  
317 closed chamber technique was used to measure GHG fluxes; in PF, PVC chamber collars (50 cm ×  
318 50 cm × 20 cm) with a water-filled channel were inserted into the soil at a depth of 15 cm. In crab  
319 ponds, a specially designed system, which included four stainless steel pegs for fixing the system  
320 and two adjustable crossbars for elevating or lowering the chamber collars with the fluctuation of  
321 water level, were installed along the boardwalks to minimize water wave impact on gas sampling.  
322 Three PVC chamber collars were placed on the crossbars in each pond. If necessary, the crossbars  
323 together with the chamber collars were adjusted to the best position one day before sampling. The  
324 transparent Plexiglass chambers (50 cm × 50 cm × 15 cm) in crab ponds and the stainless steel  
325 chambers (50 cm × 50 cm × 50 cm) insulated with white foam in PF were used. See Yuan et al.<sup>32</sup> for  
326 further detailed information of the devices.

327 The GHG fluxes were measured twice weekly in crab ponds during crab production period  
328 from March to October and weekly during period without crab production from November to  
329 February (Supplementary Fig. 3). In PF, GHG fluxes were measured twice weekly from April to  
330 November and weekly from December to March. Gas sampling was conducted at 08:00–10:00 local  
331 time to minimize diurnal variation in the flux pattern. During sampling, the chamber was fitted into  
332 the water trough of the chamber collars. Each time, four gas samples of the chamber headspace  
333 were drawn using a 50-mL syringes at 0, 10, 20, and 30 min after closure and injected into 22-mL  
334 pre-evacuated glass vials. Air temperature inside the chamber was simultaneously measured with a  
335 mercury thermometer. Concentrations of CH<sub>4</sub>, N<sub>2</sub>O and CO<sub>2</sub> were determined by a gas  
336 chromatograph (Agilent 7890, Santa Clara, CA, USA) equipped with a flame ionization detector for  
337 CO<sub>2</sub> and CH<sub>4</sub> and a <sup>63</sup>Ni electron capture detector for N<sub>2</sub>O. The gas standards were provided by the  
338 National Research Center for Certified Reference Materials, Beijing, China. The precision for GHG

339 concentrations was  $\pm 0.5\%$  based on repeated measurements of gas standards. The GHG fluxes were  
340 calculated using a linear least squares fit to the four points in the time series of concentration for  
341 each plot. Data were omitted if the slope of the linear fitting had  $R^2 < 0.90$ . Since the opaque  
342 chambers were used in PF, the measured  $\text{CO}_2$  fluxes were ecosystem respiration ( $R_e$ ); in contrast,  
343  $\text{CO}_2$  fluxes in crab ponds measured by transparent chambers were net ecosystem exchange<sup>32</sup>. The  
344 dataset of GHG fluxes were supplied as **Supplementary Table 7**.

345 Annual or seasonal cumulative  $\text{CH}_4$  ( $\text{kg CH}_4 \text{ ha}^{-1}$ ),  $\text{N}_2\text{O}$  ( $\text{kg N}_2\text{O ha}^{-1}$ ) and  $\text{CO}_2$  ( $\text{kg CO}_2 \text{ ha}^{-1}$ )  
346 emissions ( $E$ ) were calculated using the following equation:

$$347 \quad E = \sum_{i=1}^n (f_i + f_{i+1}) / 2 \times (t_{i+1} - t_i) \times 24 \times 10^{-2}$$

348 where  $f$  represents the flux of  $\text{CH}_4$  ( $\text{mg CH}_4 \text{ m}^{-2} \text{ h}^{-1}$ ) or  $\text{N}_2\text{O}$  ( $\text{mg N}_2\text{O m}^{-2} \text{ h}^{-1}$ ) or  $\text{CO}_2$  ( $\text{mg CO}_2 \text{ m}^{-2}$   
349  $\text{h}^{-1}$ );  $i$  is the  $i$ th measurement;  $(t_{i+1} - t_i)$  is the days between two adjacent measurements; and  $24 \times$   
350  $10^{-2}$  was used for unit conversion.

351

352 **Auxiliary measurements** Redox potential of the intact soil and sediment at 10 cm depth was  
353 measured *in situ* using a PHB-6 pH/mV meter (Jiaoyuan Instrument, Yancheng, China). The soil of  
354 PF or sediment of ponds at 10 cm depth was collected weekly using a Russian corer for mineral N  
355 and dissolved organic C measuring. The  $\text{NH}_4^+$  and  $\text{NO}_3^-$  were extracted with 2 M KCl solution  
356 (shaken for 1 h and then filtered); extracts were filtered and analyzed on a continuous-flow analyzer  
357 (SAN++, Skalar, Breda, the Netherlands). Dissolved organic C was extracted with deionized water  
358 (shaken for 30 min at  $25^\circ\text{C}$ , centrifuged for 25 min at 4000 rpm, and filtered through 0.45- $\mu\text{m}$   
359 membrane filter) and measured on a TOC analyzer (TOC Vcph, Shimadzu, Kyoto, Japan). Soil  
360 organic C (SOC) and total N contents were determined by the wet-oxidation redox method and the

361 Kjeldahl procedure, respectively<sup>33</sup>.

362 **Estimates of SOC change in paddy field and GWP** The SOC change ( $\delta$ SOC) in PF was estimated  
363 from the net ecosystem C balance (NECB) using a coefficient of 0.213 for paddy soils in this  
364 study<sup>29</sup>, namely, the conversion rate of organic C gain to SOC is 213 g C kg<sup>-1</sup>. The NECB of the  
365 short-plant croplands was calculated according to Ma et al.<sup>27</sup>:

$$366 \text{ NECB} = \text{GPP} - \text{Re} - \text{Harvest} - \text{CH}_4 + \text{Manure}$$

367 where GPP (gross primary production) is inferred from NPP (net primary production) via the  
368 NPP/GPP ratio of 0.58 in this region deduced by Zhang et al.<sup>34</sup>; Re, CH<sub>4</sub> and manure are the C  
369 exchange through ecosystem respiration, CH<sub>4</sub> emission, and manure application, respectively;  
370 Harvest is the C of removed straw and grain, which was calculated based on biomass yields, and C  
371 and N contents in straw and grain (Supplementary Table 4). The NPP includes net primary  
372 productions of grain, straw, root, litter and rhizodeposit, according to Ma et al.<sup>27</sup>.

373 The GWP (Mg CO<sub>2</sub>eq ha<sup>-1</sup>) in PF is calculated by the following equation<sup>35</sup>:

$$374 \text{ GWP} = 28 \times \text{CH}_4 + 265 \times \text{N}_2\text{O} - 44/12 \times \delta\text{SOC}$$

375 and for crab ponds:

$$376 \text{ GWP} = 28 \times \text{CH}_4 + 265 \times \text{N}_2\text{O} + 1 \times \text{CO}_2$$

377 where CH<sub>4</sub>, N<sub>2</sub>O and CO<sub>2</sub> denote annual emissions of CH<sub>4</sub> (Mg CH<sub>4</sub> ha<sup>-1</sup>), N<sub>2</sub>O (Mg N<sub>2</sub>O ha<sup>-1</sup>) and  
378 CO<sub>2</sub> (Mg CO<sub>2</sub> ha<sup>-1</sup>), respectively.

379

380 **Data collection and classification of global freshwater aquaculture** As mentioned above, there  
381 are large uncertainties in previous model estimates of global aquaculture N<sub>2</sub>O emissions by using  
382 EFs of applied N and fish yields: First, the N<sub>2</sub>O EFs are highly dependent on management levels in  
383 the aquaculture system. For example, yield-scale N<sub>2</sub>O EF (EF<sub>Y</sub>) of carp was 1.07 g N<sub>2</sub>O kg<sup>-1</sup> yield

384 in an intensive rearing system<sup>16</sup> but was only 0.28 g N<sub>2</sub>O kg<sup>-1</sup> yield in a semi-intensive earthen  
385 pond<sup>17</sup>; Secondly, the EF<sub>Y</sub> can be biased by the major yield difference between species. For instance,  
386 although the direct N<sub>2</sub>O emission rates in two adjacent semi-intensive aquaculture ponds were  
387 comparable, the EF<sub>Y</sub> measured in crab ponds was 8.11-fold greater than that in carp ponds due to  
388 the magnitude difference in yields<sup>17,18</sup>.

389 Here, we compiled a worldwide database of GHG emissions measured in the inland freshwater  
390 aquaculture systems (Supplementary Table 5). We identified potential published studies for  
391 inclusion in the database using Web of Science with the keywords ‘greenhouse gases or CH<sub>4</sub> or N<sub>2</sub>O’  
392 and ‘aquaculture or fish farming or rice fish or aquaponics’. Twenty-four studies fell within the  
393 inland freshwater aquaculture and met the following criteria: (i) field measurement of CH<sub>4</sub> and/or  
394 N<sub>2</sub>O emissions was carried out on a per hectare or per fish yield basis; (ii) type of aquaculture  
395 system with or without aerator use was reported; (iii) the N input and yield in intensive systems  
396 were listed (see below). The dataset include 45 CH<sub>4</sub> and/or N<sub>2</sub>O emission measurements across 19  
397 sites between 2003 and 2015.

398 Generally, the aquaculture systems are classified based on production per unit volume or per  
399 unit area<sup>36</sup>; however, when estimating the regional or global GHG emissions, such classification  
400 might be unfit due to lack of the available production data counted by volume or area and  
401 deficiency of the cross-species classification criteria for big differences in production performance  
402 between culture species. Here, we classified four systems: rice-fish, extensive, semi-intensive and  
403 intensive based on the local conditions and aquaculture facilities especially aerator use or not.  
404 Actually, the stocking density and production are associated with investment on infrastructure  
405 especially aeration equipment<sup>36</sup>, because the dissolved oxygen in fish ponds should be  
406 maintained >5.0 mg L<sup>-1</sup>, theoretically<sup>37</sup>.

407       • Rice-fish systems include integrated rice field or rice field-pond complex and are used to  
408 produce fish and other aquatic animals.

409       • Extensive aquaculture systems involve excavated earthen ponds, irrigation canals and ditches,  
410 small lakes and reservoirs used for fish farming. Extensive systems have low stocking density, with  
411 natural productivity or limited supplemental feeds and no aerator system.

412       • Semi-intensive aquaculture systems include excavated earthen ponds, irrigation canals and  
413 ditches, small lakes and reservoirs, have higher stocking densities than extensive systems, and are  
414 equipped with aerators and managed with artificial feeds and intermittent aeration.

415       • Intensive aquaculture systems, which utilize man-made rearing units such as concrete/canvas  
416 tanks, raceways recirculating systems, have high stocking rates and complete diet management,  
417 intensive and continuous aeration, and frequent or continuous water exchange. The cage and pen  
418 culture performed in open water bodies like rivers, lakes and reservoirs are also classified as  
419 intensive aquaculture because of the high stocking rates and sufficient dissolved oxygen supply  
420 from the constant water exchange.

421       Global inventory of the land use and production statistics are also compiled in different  
422 aquaculture systems of the major freshwater aquaculture producers ([Supplementary Table 6](#)),  
423 however, data of extensive and semi-intensive systems were pooled because of lack of aerator use  
424 data to classify each other. Data were derived from the official fisheries statistics for 2014. In case  
425 2014 data were not available, the most recent data were used. If the national official statistical data  
426 were not available, the FAO estimate (National Aquaculture Sector Overview) or private survey  
427 data were used. Further details on the statistics used are provided in the Supplementary materials.

428

429       **Estimation of global CH<sub>4</sub> and N<sub>2</sub>O budgets** We estimated N<sub>2</sub>O emissions from intensive systems

430 by multiplying  $EF_Y$  by production. Methane emission from intensive systems was recognized as  
431 negligible because the aerobic condition limited  $CH_4$  production in such systems<sup>4</sup>. While  $CH_4$  and  
432  $N_2O$  emissions from rice-fish, extensive, and semi-intensive aquaculture systems were estimated by  
433 multiplying mean emission rates by area, because (i) the yield EF for  $CH_4$  was generally  
434 unavailable in literature; and (ii) the  $EF_Y$  would be biased by the huge yield difference between  
435 species in extensive and semi-intensive systems. Additionally, when estimating  $CH_4$  emission from  
436 rice-fish systems, the  $CH_4$  fluxes ( $32\text{--}37\text{ mg } CH_4\text{ m}^{-2}\text{ h}^{-1}$ ) measured in Bangladesh<sup>30</sup> were excluded  
437 from mean emission rates, because of the extremely high emission rates and relatively small area of  
438 rice-fish in Bangladesh ( $\sim 3.97\%$  of global rice-fish area).

439 It should be noted that our preliminary estimates possess some uncertainties. First, field  
440 measurements of  $CH_4$  and  $N_2O$  fluxes were mainly conducted during the feeding period, may result  
441 in overestimation of  $CH_4$  and  $N_2O$  emissions; secondly, only averaged  $CH_4$  and  $N_2O$  fluxes in  
442 extensive and semi-intensive systems were set up due to the absence of detailed aquaculture  
443 facilities data; thirdly, there was no detailed information relative to land use and production in  
444 aquaculture in many main producers (e.g. Brazil, Nigeria). More field measurements along with  
445 detailed national aquaculture information in those countries are required to obtain more reliable  
446 estimates. Moreover, our estimates only focused on the direct  $CH_4$  and  $N_2O$  emissions, however,  
447 GHG emission from adjacent water bodies can also be enhanced by the nutrients loading caused by  
448 water exchange in some aquaculture systems (especially intensive systems). Hence, these potential  
449 indirect emissions should be considered in future estimates.

450

## 451 **References**

452 31. Zhang, F., Xing, Y., Pu, L. & Peng, B. Study on the eco-environmental effect of land use change

- 453 in Suzhou. *Res. Soil Water Conserv.* **16**, 98–103 (2009) (in Chinese).
- 454 32. Yuan, J. et al. Exotic *Spartina alterniflora* invasion alters ecosystem-atmosphere exchange of  
455 CH<sub>4</sub> and N<sub>2</sub>O and carbon sequestration in a coastal salt marsh in China. *Glob. Change Biol.* **21**,  
456 1567–1580 (2015).
- 457 33. Carter, M. R. *Soil Sampling and Methods of Analysis* (Lewis Publishers, 1993).
- 458 34. Zhang, Y., Xu, M., Chen, H. & Adams, J. Global pattern of NPP to GPP ratio derived from  
459 MODIS data: effects of ecosystem type, geographical location and climate. *Global Ecol.*  
460 *Biogeogr.* **18**, 280–290 (2009).
- 461 35. Ciais, P. et al. in *Climate change 2013: The Physical Science Basis* (eds Stocker T. F. et al.) Ch.  
462 6, 465–570 (IPCC, Cambridge Univ. Press, 2014).
- 463 36. Lekang, O-I. *Aquaculture engineering* (John Wiley & Sons, 2008).
- 464 37. Losordo, T. M., Masser, M. P & Rakocy, J. *Recirculating Aquaculture Tank Production Systems:*  
465 *Management of Recirculating Systems* (SRAC Publication, 1998).

466 **Figure legends:**

467 **Figure 1. Annual CH<sub>4</sub>, N<sub>2</sub>O and CO<sub>2</sub> emissions from the paddy field (PF) and crab ponds (CP)**

468 **during 2013–2014. a, CH<sub>4</sub>, b, N<sub>2</sub>O, c, CO<sub>2</sub>.** Vertical bars represent standard errors of the means (*n*

469 = 4 for PF and *n* = 3 for crab ponds). Three crab ponds had different size and water depth

470 (Supplementary Table 1). ‘A’, ‘B’, and ‘C’ denote significant differences between sites (*P* < 0.05,

471 ANOVA, Tukey’s HSD test) during the entire year; ‘a’, ‘b’, and ‘c’ denote significant differences

472 between crab ponds during the crab production period or during the period without crab production.

473 CO<sub>2</sub> release from PF was calculated from soil organic C change estimates using the net ecosystem

474 C balance method.

475 **Figure 2. Literature-sourced greenhouse gas emission factors of different aquaculture. a, mean**

476 CH<sub>4</sub> emission rate, **b, mean N<sub>2</sub>O emission rate, c, N<sub>2</sub>O emission factor of applied N (EF<sub>N</sub>), d, yield**

477 based N<sub>2</sub>O emission (EF<sub>Y</sub>). Boundaries of the boxes indicate the first and third quartiles, line within

478 the box and the white square represent the median and average, respectively. Whiskers mark the




479 10th and 90th percentiles, and the outliers are shown as dots. **e, global warming potential (GWP), f,**







480 greenhouse gas intensity (GHGI, GWP/yield). Vertical bars represent standard errors of the means.

481 Aquaculture systems are classified based on the local conditions and aquaculture facilities

482 especially whether aerators were used or not.







1 **Table 1 Annual GHG emissions, net GWP and emission factors of CH<sub>4</sub> and N<sub>2</sub>O in paddy field and crab ponds**

Systems	CH <sub>4</sub> (kg CH <sub>4</sub> ha <sup>-1</sup> )	N <sub>2</sub> O (kg N <sub>2</sub> O ha <sup>-1</sup> )	CO <sub>2</sub> * (Mg CO <sub>2</sub> ha <sup>-1</sup> )	C input† (Mg C ha <sup>-1</sup> )	N input† (kg N ha <sup>-1</sup> )	δSOC‡ (Mg C ha <sup>-1</sup> )	Net GWP§ (Mg CO <sub>2</sub> eq ha <sup>-1</sup> )	EF <sub>C</sub> †† (%)	EF <sub>N</sub> ¶ (%)	EF <sub>Y</sub> ¶ (g N <sub>2</sub> O kg <sup>-1</sup> yield)
Paddy field	218 ± 7b	7.11 ± 0.23a	50.6 ± 0.9a	–	430	-0.04 ± 0.05	8.15 ± 0.43b	–	1.05 ± 0.03a	0.56 ± 0.02a
Crab ponds	962 ± 149a	0.33 ± 0.07b	0.93 ± 0.55b	1.20	244	–	28.0 ± 4.1a	60.0 ± 9.3	0.09 ± 0.02b	0.30 ± 0.07b

2 \* The value is ecosystem respiration in paddy field and net ecosystem CO<sub>2</sub> exchange for crab ponds. † Calculated by application rates and C and  
3 N contents of the fertilizers and feeds (see Supplementary Tables 2–4). ‡ Estimated from the net ecosystem carbon balance (NECB) using a  
4 coefficient of 0.213 for paddy soils<sup>29</sup>. § Net GWP = 28×CH<sub>4</sub>+265×N<sub>2</sub>O–44/12×δSOC for paddy field, and net GWP =  
5 28×CH<sub>4</sub>+265×N<sub>2</sub>O+1×CO<sub>2</sub> for crab ponds. †† The direct emission factor of C for CH<sub>4</sub> (EF<sub>C</sub>) is calculated by dividing annual CH<sub>4</sub> emission by  
6 total C input<sup>21</sup>. ¶ The direct emission factor of N for N<sub>2</sub>O (EF<sub>N</sub>) and yield-scaled emission factor for N<sub>2</sub>O (EF<sub>Y</sub>) are calculated by dividing  
7 annual N<sub>2</sub>O emission by total N input and grain/crab yield, respectively. Values are means ± standard errors.

8 **Table 2 Direct CH<sub>4</sub> (Gg CH<sub>4</sub> yr<sup>-1</sup>) and N<sub>2</sub>O (Mg N<sub>2</sub>O yr<sup>-1</sup>) emissions from**  
 9 **different freshwater aquaculture systems in global top 21 producers in 2014**

Country/region	Rice-fish systems*		Extensive plus semi-intensive systems*		Intensive systems†	Total‡	
	CH <sub>4</sub>	N <sub>2</sub> O	CH <sub>4</sub>	N <sub>2</sub> O	N <sub>2</sub> O	CH <sub>4</sub>	N <sub>2</sub> O
	China	696	5,988	3,408	11,653	5,152	3,524
India	108	925	487	1,667	–	512	2,591
Indonesia	66	571	91	313	1,955	142	2,839
Vietnam	19	161	173	590	344	162	1,095
Bangladesh	–	–	323	1,106	4	268	1,109
Myanmar	–	–	50	172	0	42	172
Brazil	–	–	45	153	430	37	584
Thailand	2	15	71	244	91	61	350
Nigeria‡	–	–	–	–	–	–	–
Philippines	–	–	8	28	373	7	401
Iran	0	2	28	97	316	24	415
USA	15	128	35	119	76	44	323
Egypt	268	2,306	1	3	442	269	2,752
Pakistan	–	–	8	28	0	7	28
Taiwan Province of China	0	0	34	116	0	28	116
Russia	–	–	57	194	71	47	265
Cambodia	0	2	1	3	208	1	213

Uganda	–	–	6	19	67	5	86
Lao PDR	2	20	21	71	55	20	146
Turkey	–	–	0	0	268	0	268
Malaysia	12	101	3	12	50	15	162
Top 21 subtotal	1,188	10,219	4,851	16,586	9,903	6,039	36,709

---

10 \* Calculated by mean CH<sub>4</sub> and N<sub>2</sub>O emission rates (Fig. 2) and the area for aquaculture  
11 (Supplementary Table 6) collected from the literature. Rates of CH<sub>4</sub> emission from rice-fish  
12 system in Bangladesh were excluded when calculating<sup>30</sup>. † Calculated by averaged  
13 yield-scaled emission factor for N<sub>2</sub>O (EF<sub>Y</sub>) (Fig. 2d) and volume of production from intensive  
14 aquaculture. The direct emission rate of CH<sub>4</sub> from intensive system was estimated at 0  
15 according to Hu et al.<sup>4</sup>. ‡ No official or private statistics is available about area and  
16 production from different systems in Nigeria.