

Integrating field and laboratory approaches for ripple development in mixed sand-clay-EPS

Baas, Jaco H.; Baker, Megan; Malarkey, Jonathan; Bass, S.J.; Manning, A.J.; Hope, J.A.; Peakall, Jeff; Lichtman, Ian; Ye, L.; Davies, Alan; Parsons, D.R.; Paterson, D.M.; Thorne, P.D.

Sedimentology

DOI:

[10.1111/sed.12611](https://doi.org/10.1111/sed.12611)

Published: 01/12/2019

Peer reviewed version

[Cyswllt i'r cyhoeddiad / Link to publication](#)

Dyfyniad o'r fersiwn a gyhoeddwyd / Citation for published version (APA):

Baas, J. H., Baker, M., Malarkey, J., Bass, S. J., Manning, A. J., Hope, J. A., Peakall, J., Lichtman, I., Ye, L., Davies, A., Parsons, D. R., Paterson, D. M., & Thorne, P. D. (2019). Integrating field and laboratory approaches for ripple development in mixed sand-clay-EPS. *Sedimentology*, 66(7), 2749-2768. <https://doi.org/10.1111/sed.12611>

Hawliau Cyffredinol / General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

SEDIMENTOLOGY

the journal of the
International Association of Sedimentologists

INTEGRATING FIELD AND LABORATORY APPROACHES FOR RIPPLE DEVELOPMENT IN MIXED SAND-CLAY-EPS

Journal:	<i>Sedimentology</i>
Manuscript ID	SED-2018-OM-310
Manuscript Type:	Original Manuscript
Date Submitted by the Author:	20-Dec-2018
Complete List of Authors:	<p>Baas, Jaco; Bangor University, School of Ocean Sciences Baker, Megan; Bangor University, School of Ocean Sciences Malarkey, J.; University of Wales Bangor, School of ocean Sciences Bass, Sarah; Plymouth University, School of Marine Science and Engineering Manning, Andrew; Plymouth University, School of Marine Science and Engineering Hope, Julie; University of Auckland, Institute of Marine Science Peakall, Jeff; University of Leeds, Earth and Environment Lichtman, Ian; National Oceanography Centre Liverpool Ye, Leiping; University of Hull, Department of Geography, Environment and Earth Sciences Davies, Alan; Bangor University, School of Ocean Sciences Parsons, Daniel; University of Hull, Geography, Environment and Earth Sciences Paterson, David; University of St. Andrews, Scottish Oceans Institute, School of Biology Thorne, Peter; National Oceanography Centre Liverpool, Proudman Oceanography Laboratory</p>
Keywords:	Current Ripples, Mixed Sand-Clay-EPS, Cohesion, Estuary, Intertidal flat

SCHOLARONE™
Manuscripts

Reviewer 1

Question 1.1 - This paper looks at the formation of bedforms on a natural mixed sand–mud–EPS intertidal flat in a macrotidal estuary. Both physical and biological origin of cohesive forces are considered to affect bedform dynamics in sandy sediment that should be redefined from the widely used as ‘clean’ sand, based on integrated field hydrodynamic and bed morphological data. These analyses are also integrated with information on experimental small-scale current ripples, to elucidate the underlying processes that control the differences between them in the rate of change of current ripple length with changing bed clay and EPS content. This is a piece of important work in the area of sedimentology, for predictive sediment transport models, bedform size predictors, and sedimentary facies analysis. I recommend the manuscript to be accepted with the following minor comments:

Reply 1.1 - We thank this reviewer for their positive assessment of the importance of our work.

Q. 1.2 - (1) In the part of METHODS, please add more details on the field campaign in May 2013. For example, how was the weight percentage of EPS measured? It should not be cited from an unpublished work (Lichtman et al., in review).

R. 1.2 - Lichtman et al. (2018) is now published, and we have expanded the Methods somewhat to address the comments of all reviewers.

Q. 1.3 - (2) Some recent publications in the areas could be cited as reference. For example: Zgheib, N., Fedele, J.J., Hoyal, D.C.J.D., Perillo, M.M., & Balachandar, S. (2018). Direct Numerical Simulation of Transverse Ripples: 1. Pattern Initiation and Bedform Interactions, 123, (3), 105-122. doi: 10.1002/2017JF004398
Liu, X., Jia, Y., Zheng, J., Wen, M., & Shan, H. (2017). An experimental investigation of wave-induced sediment responses in a natural silty seabed: New insights into seabed stratification. Sedimentology, 64(2), 508-529. doi: 10.1111/sed.12312

R. 1.3 - After careful consideration, we have decided against citing these interesting papers in our revised manuscript. The first paper is somewhat relevant in that it discusses planform patterns of bedforms, but these patterns do not include 3D shapes and the paper does not discuss how the properties of the bed might be important. The second paper is less relevant for our manuscript because it discusses wave-generated instead of current-generated bedforms.

Reviewer 2: Iris Verhagen (main comments)

Q. 2.1 - Thank you for the opportunity to review the manuscript SED-2018-OM-053 “Redefining ‘clean’ sand by integrating field and laboratory data on mixed sand-clay-EPS rippled-bed transport”, by Baas et al.

In short, my opinion is that the work is well written, has clear aims and purpose, and shows interesting new results about bed form formation in mixed sediments (clay-sand).

R. 2.1 - We thank this reviewer for their positive assessment of our work.

Q. 2.2 - However, I recommend more detail is added about the statistical methods used to (further) validate the trends that are shown.

R. 2.2 - Our statistical method is primarily based on linear regression and we take an objective approach by including both R^2 -values and p -values. R^2 -values alone are usually not sufficient, because low values do not necessarily mean that a linear trend is statistically insignificant. The probability, p , is more objective and a better measure for goodness of fit, as it compares the calculated probability to a reference value, in our case $p_{ref} = 0.05 = 5\%$. If $p < p_{ref}$, there is lower than 5% probability that the fit is statistically insignificant. Likewise, $p > p_{ref}$ means that the fit is insignificant with a probability of at least 5%. This is a standard method in statistics that objectively validates the linear relationships between our data. We therefore think that the statistical method does not require further explanation in our manuscript.

Reviewer 2: Iris Verhagen (further comments in text)

p. 3

(1) Recommend to insert reference(s) here.

Done.

(2) Recommend to insert reference(s) here.

Done.

p. 4 - I am not sure about the placement of this section. Perhaps better at the end of this paragraph or in the interpretation/discussion section?

We believe that hypotheses belong in the introduction, so we did not move this section.

p. 5 - Same comment regarding statistical methods (below on page 7) applies here, although this might be covered in the work by Lichtman et al.

This comment was covered in R. 2.2.

p. 6

(1) I understand now but found this confusing when I first read it, therefore I would consider rephrasing this section, for example including "removing the non-cohesive fraction from the mud fraction" or similar.

This section has been reworded (see page 6 in revised manuscript).

(2) See comment on p15

See reply p. 15.

(3) Consider removing this or moving it to the discussion section (p11-12).

Done.

p. 7

(1) I think that there is probably a trend here (Fig 2), but the number of data points is quite low with 12 and especially the subsurface trend (solid circles Fig 2) seems to be biased by two of the points. So I suggest that the authors need to expand on their statistical testing to back this up more (perhaps as an appendix or supplement material), e.g.:

- What statistical test(s) and data were used to obtain the p-values (T-test + hypotheses, linear regression?)

- Provide an indication of the errors (as error-bars) in the data set associated with measuring the ripple length and mud content in the bed.

[This comment was covered in R. 2.2.](#)

(2) This is not true for all beds, bed 10 and 15 have slightly higher bed mud % in the ripples compared to the beds.

[The text has been changed to reflect this.](#)

Also, if the bed mud % difference (subsurface - ripple) is plotted against the ripple length, no clear trend is visible, even though that is implied here. Or can the data not be plotted against each other in this way, i.e. was height of the bed forms different and therefore the winnowing depth/efficiency (p8) or is it possible something else influences this?

[It is not clear to us why the reviewer would expect the difference to show a relationship.](#)

(3) These are large ranges with quite a bit of overlap: 24-38% subsurface, 12-34% surficial (with only one standard deviation from the mean).

Also, beds 4-6 have surficial bed mud content within this range, bed 7 same for the subsurface.

I recommend more detail is added about this observed variation.

[This is just an order of magnitude argument here. The subsurface cohesive clay content is now discussed in more detail in a later section and includes a figure.](#)

p. 8 - Recommend to insert reference(s) here. Also a typo (, instead of .)

[A reference to Decho & Gutierrez \(2017\) has been added.](#)

p. 9

(1) This is already briefly mentioned on page 6, perhaps consider rewording there to avoid repetition.

[The earlier section has been reworded, see reply p. 6\(1\)](#)

(2) Link this to the ranges discussed on page 7 perhaps? If $c = 14\%$ then $m \sim 39\%$, quite high compared to $31 \pm 7\%$

[We cannot do this because the experiments did not have non-cohesive fines.](#)

p. 11

(1) I am assuming this is bed clay% in the ripples? Not bed mud%?

[Yes, it is bed clay %, but in the substrate, not the ripples.](#)

(2) This 'Flow energy' section becomes somewhat confusing by taking this approach. I think the content is OK but perhaps consider rewording part of it.

[We think the wording is ok here.](#)

p. 12 - Additionally, the fact that the ripple heights were recorded at a different time compared to the sampling for clay content (p5, lines 40-41) should be considered.

[Ripple heights were not recorded. Since only clay content \(which is not time sensitive\) and not EPS was determined from the sediment samples, this is not a concern. The fact that EPS was not measured directly for the samples is now clearer in the text.](#)

p. 13 - How was this measured? (see also comment in Table 1)

[The new manuscript now explains that this was determined from Equation 1.](#)

p. 15 - Yes, I wanted to make this point at the start of the paper when the spring tide conditions are mentioned, perhaps comment there (p6, line 30) that these conditions were favorable due to higher flow velocities?

The spring tide is not the reason we chose this particular day to conduct the ripple data collection.

p. 16 - Consider moving this to “Water salinity” section

We cannot do this, because this part of the section sums up the various influences of the different factors described in the previous section

p. 18

(1) These are new results discussed here, I would consider adding the experiments to the main part of this manuscript (see comment on Appendix A)

We considered doing this but feel it breaks up the text too much. However, the discussion of enhanced thresholds has been moved forward to the *Bed EPS content* section.

(2) (EPS proxy)

This has been done where Xanthan gum is first mentioned for these experiments (p. 18, 2nd para 2, line 5)

‘... θ_c for kaolin clay and the Xanthan gum (an EPS proxy), in ...’

(3) I would suggest to change this to "sedimentary rock classification" (or similar) rather than facies analysis here, when linked directly to the sandstone classifications.

This sentence does not refer to sedimentary rock classifications specifically.

p. 19

(1) Are these the results in Appendix A or the experiments by Baas et al. (2013)?

They are the results of Baas et al. (2013), which is partly the reason for keeping the description of the vertical shear flume in an appendix. The word ‘bedform’ has been inserted in this sentence for clarification: ‘... limits of the present comparative study of laboratory and field bedforms.’

(2) See comment p18 (line 47), but facies more applicable here.

We believe this is ok as it is.

p. 20 - This only discussed very briefly (p16).

We feel this is covered by using the adjective ‘possibly’.

p. 26 - typo

Done.

p. 28 - I would consider including these results in the main manuscript, especially since the title suggests laboratory data are a key part of the paper.

See reply p. 18(1).

Table 1 - Are these values calculated with Eq. 1? Or was EPS measured? If so, this is not mentioned in the methods.

The Methods section now explains this.

Fig. 2 - Only two data points seem to control the slope of the solid regression line. Are you sure this is not sample bias (low n)?

Possibly, but there is no denying the downward trend and now Λ includes the three flat-bed cases as well in Fig. 4. Also a comment has been added about the increased scatter of the field data compare to the laboratory data.

Fig. 5 - Perhaps more effective in hours?

We prefer to keep the present format, because this allows readers to match our data to tide tables.

Reviewer 3

Q. 3.1 - This is a potentially very interesting paper that sets out to examine the relationship between flume and field data sets examining ripple development in an estuary and under estuarine conditions, and seeks to examine the various influences on ripple size and development as function of flow and wave ‘energy’, clay content, water salinity and EPS concentrations. I was excited to read the paper to see how the authors had sorted this out and examine the database that would allow these factors to be discerned. There is, as the authors highlight later in their paper, a clear need for more data and analysis to examine these issues.

However, I was disappointed to find that the paper presents a very limited dataset, and that many of the factors perhaps of importance, such as EPS concentration, wave energy, salinity and even clay content further down within the bed, have not been measured or quantitatively assessed whatsoever. The result is a paper rather short on data, drawing heavily on past experimental work for comparisons, and presenting often vague and unsubstantiated discussion on the brief results. The paper then has a Discussion section where it attempts to discuss these possible findings with respect to modern and ancient sediments, arriving at the conclusion that our terminology for clean and dirty sands needs change...or we need a ‘paradigm shift in sedimentary facies analysis’ for recognizing the importance of cohesive forces for bedform dynamics. This may be true, but such a shift must be based on a full characterization of the processes that control such bedform dynamics – and this paper does not do this as it does not quantify and measure many of the factors that it identifies as important. Thus, the reader is left having to trust a broad and rather vague Discussion section where the authors speculate on the importance of these factors (see the very vague Table 2) that seems inappropriate for Sedimentology where I would judge readers would expect evidence for these contentions and not just speculation. I have marked many comments on the pdf, but the principal points of concern are:

1) There is a lack of measurement in many of the supposed vital factors – EPS, salinity, wave velocities, clay contents not just in the first few centimeters but deeper down, and assessment of 2D-vs-3D bedforms from a quantitative perspective. Many of these thus result in the paper being too speculative in many parts, and often unsupported by data.

R. 3.1 - Unfortunately, it was not possible to collect samples for EPS content analysis, in addition to clay content and ripple size analysis, during the short slack water period at the field site, as explained in the revised manuscript. We therefore have to rely on a more general relationship between bed clay content and bed EPS content based on samples that were collected independently by our team in the same fieldwork period (for details, see Lichtman *et al.*, 2018). The exact location at which these samples were collected was described somewhat confusingly in the original

manuscript. We were unclear in mentioning that the 140 m refers to the maximum distance from the sampling area for this study. The Methods section of the revised manuscript makes clear that the vast majority of the samples were collected within or in the immediate vicinity of the sampling area for this manuscript.

We agree with the reviewer that salinity may play a role in the comparison between the field and laboratory data, but it seems unrealistic to us to expect that a full comparison of complex field conditions with simplified laboratory conditions is possible. Until laboratory data on the effect of salinity on the dynamics of current ripples in mixed sand-mud and sand-EPS become available, we can only rely on less satisfactory data from literature. Unfortunately, these data are inconclusive, which the reviewer may have missed, yet lean towards a negative correlation between salinity and bed stability (and therefore ripple size). Rather than ignoring salinity, we use this information to the best of our ability in the revised manuscript, thereby encouraging further laboratory and field research on the effect of salinity.

The revised manuscript is now more explicit in mentioning that waves were absent in the tidal periods immediately preceding the ripple measurements (see also Lichtman *et al.*, 2018). This renders the section on wave velocity redundant, and it has therefore been removed.

We may have been unclear in mentioning that we took bulk samples from the ripples and from 0 to 20 mm below the base of the ripples for comparison. To address this issue, we have slightly reworded the related text in the Methods section of the revised manuscript. We do not see a reason to extend the sampling to older sediment deeper into the subsurface.

Quantitative assessment ripple three-dimensionality: See R. 3.5 below.

Q. 3.2 - 2) The authors suggest new limits for how we classify muddy or clean sediments, and yet these original limits were presumably not drawn up with ripple process mechanics in mind but rather as simple descriptors of the percentage of fines vs sands...these are surely different things.

In addition, if you may not be able to even assess the role of EPS in ancient sediments, yet it may be absolutely key, how does changing the limit even make sense here – a clean EPS-rich sand could, presumably, significantly influence the bedforms but leave no trace – so how would a new classification help here?

Since the effect of fines must also depend to some extent on the applied bed shear stress, how would such limits apply to sand/muds deposited at higher bed shear stresses than ripple bedforms...and thus would you need different limits there?

I am just unclear as to what is gained here and judge the authors rather oversell this new classification angle. Surely ‘wacke’ refers to the grain size of a rock we can classify easily in the field...could you even judge a clay content of 3%, especially given likely diagenetic alterations as well?

In addition, I feel the authors rather oversell their pitch regarding reclassifying what we mean by clean and muddy sediments, and fail to acknowledge the differing aims of the older classifications against what is presented here. As such, I do not feel the paper is acceptable for Sedimentology.

R. 3.2 – We firmly believe that many earth scientists are unaware that a few percent of clay in sand are sufficient to change bedform dynamics, and thus erroneously treat such sand as ‘clean’ from a process sedimentological perspective (hence the reference to a “paradigm shift”). Surely, sedimentologists should make informed decisions when classifying sediment, rather than setting arbitrary boundaries. However, we agree with the reviewer that there are practical limitations in redefining the boundary between clean and dirty sand to 3% clay and even lower percentages of

EPS. Instead of redefining the boundaries, we now state in the *Implication for geological studies* of the revised manuscript: “These textural classifications have practical use for categorising rocks in outcrop and core, but the boundary percentages should not be used to differentiate non-cohesive from cohesive dynamic behaviour of mixed sand–mud”. Also, the emphasis of a change in the ‘clean sand’ definition (including in the title) has been reduced.

Q. 3.3 - 3) The paper refers many times to the manuscript of Lichtmann et al. (in review, *Geomorphology*) and I would judge it essential to view this paper as well to assess any overlap between the two. The title suggests it also looks at bedform migration and maybe morphology – so how does this paper differ? The present paper also refers to results from this other paper and the paper would need to be accepted for this to be a feasible citation.

R. 3.3 – The reviewer is correct that this is a paper that they ought to have had access to in order to be able to independently assess these aspects. We hope that the reviewer will forgive us our eagerness to publish the present work. Lichtman *et al.*’s (2018) paper is now published in *Geomorphology*. There is no overlap, except that we use some of their hydrodynamic data and their relationship between bed clay and EPS content. Their paper is now also referred to in the discussion of the modification to the bed material transport rate and in quantifying the trapping of clay and EPS (see R. 3.6). See also R. 3.1.

Q. 3.4 - 4) Bed clay and EPS are assessed through a correlation from a different site!..how do we know this holds and why not measure EPS at the field site for this paper?

8) It would seem essential to have measured EPS concentrations at this site.

R. 3.4 - We do not have EPS data from the ripples and the subsurface, because there was not enough time around low slack water to take samples of high enough quality for EPS analysis. EPS cannot be sampled afterwards, because environmental conditions will be different from when the samples for clay analysis were collected. We therefore had to use Lichtman et al.’s (2018) statistically significant relationship between bed clay and EPS content, which covers the field site. See R. 3.1.

Q. 3.5 - 5) No mention at all, or quantification, is given of how 2D vs 3D are assessed..these thus leave this ‘data’, and speculations from it, purely subjective.

R. 3.5 - The revised manuscript includes a simple method for distinguishing between 2D, 2D-3D, and 3D bedforms. We applied this to the ripples in the field and to the ripples of Baas et al. (2013) for which high-quality pictures of their planform were available.

Q. 3.6 - 6) The paper refers to winnowing of fines and this has been shown in the author’s previous lab experiments. But is there any role for the advective pumping of fines into the bed through hyporheic flow that has been shown by the work of, amongst others, Aaron Packman, Winnowing may influence removal of fines from the bed, but does hyporheic flow enhance the injection of fines into the bed, especially where there are pressure gradients created across bedforms, that may induce such effects and enrich the percentage of fines in the regions underneath the maximum depth of bedform scour? I feel this issue also needs to be raised and discussed and really calls for deeper sampling of clay concentration within the bed. Could such enrichment with fines within the bed then further limit the evolution of ripples by producing an enhanced clay rich zone?

R. 3.6 – We agree. Appendix A now provides calculations on the trapping time for clay or EPS entering the ripple from the water column above, thus counteracting winnowing. These calculations show that it takes a minimum of four inundation periods for the concentration of clay/EPS to reach what it is in the water column. It is basically a slow process compared to the turn-over of sediment as a ripple migrates, and too slow to expect a significant amount of fine sediment to be stored underneath the ripples. Deeper sampling than 20 mm below the active ripple layer, as is made more clear in the revised text, is therefore not needed

Q. 3.7 - 7) It would be good to clarify if the percentages mud were volume or weight percentages?

R. 3.7 – Done, these are volume percentages.

Q. 3.8 - 9) Page 14 states salinity is postulated to make a big difference – but no data whatsoever is given to support this! The authors refer to their ‘unpublished pilot experiments’ as support but I can’t believe they expect this statement would hold up in any reputable scientific journal! Authors must present the data to support such speculations.

R. 3.8 – The reviewer may have misread the paper where salinity is concerned. We did not state that salinity makes a ‘big difference’. On the contrary, the effect of salinity is equivocal, yet leans towards a negative correlation with bed stability and therefore ripple size (see also R.3.1). In the revised manuscript, we no longer refer to the unpublished dataset.

Q. 3.9 - 10) The ‘Wider Implications’ section has just too many vague and unsubstantiated provisos as to make the conclusions far too speculative – this really seems to just highlight more of these parameters need quantification to explain these trends....this needs doing here and not in future studies.

R. 3.9 - Our objective was to take the comparison between field and lab data as far as possible, incorporating as many variables as possible. Inevitably, this requires some speculation, not in the least because this is a new field of research for which many relationships have not been quantified yet. Rather than trying to provide definitive answers to all questions, which is impossible on the basis of one field dataset and simplified laboratory experiments, the Discussion assesses the broader importance of the data that should stimulate future research in the laboratory and beyond the specific characteristics of the Dee Estuary, including the geological past. We believe that this is a justifiable and widely used approach in *Sedimentology*. Where possible, we have provided further evidence for our interpretations (see R.3.1, R.3.5, R.3.6, R.3.10)

R. 3.10 - In conclusion, this paper promised a lot but failed to deliver a full dataset on which the contentions concerning the controlling parameters of bedform evolution depend. Whilst these are probably the principal factors to account for, the paper fails to measure or fully quantify many of them, resulting in far too much speculation and speculation that cannot be supported by the data presented.

R. 3.10 – See R.3.9. The revised manuscript contains a new section that makes the effect of EPS more quantitative, based on a comparison with the work of Malarkey *et al.* (2015) and following a similar approach to the quantification of the effect of clay. These quantifications are summarised in a new Fig. 6. We have also linked the increase in the threshold of motion (Fig. 7) more directly to this section. This should help reduce the amount of perceived speculation in our manuscript.

Associate Editor: Charlie Bristow

We have received three reviews of your paper, two of which recommend publication with minor revision and a third that recommends rejection. Given the potential significance of some of the claims made within this paper it required a thorough review. The review that recommends rejection provides a series of reasons why this paper cannot be published in its present form. Chief amongst these is the lack of data. There is simply not enough data presented to properly test the hypothesis and it appears that the data sets are incomplete. This paper appears to be immature. It needs more data. This does not mean that your hypothesis is wrong, but its just not supported by the data presented. This is not something that can be fixed with a revision and therefore I agree with reviewer three that this paper should be rejected.

We have found the points raised by the reviewers to be helpful, and we have looked carefully at the manuscript in order to address the key points that they raise. Our revised manuscript contains more data and it also provides more detailed analysis of some of the datasets and a reduced emphasis on the need for a new 'clean sand' definition. See earlier comments. However, it should be emphasised that a full assessment of all parameters that control mixed sand-clay-EPS ripple dynamics in natural estuaries is not possible at present. Despite this shortcoming, the new field data are in agreement with published laboratory data and support our main conclusion that smaller amounts of clay and EPS than traditionally deemed to be the case impact on ripple development and stability, to a far greater extent than has been previously recognised. This outcome is important in terms of bedload transport and for the processes of sediment entrainment and suspended load, both of which are strongly linked to bedforms. The fieldwork also clearly demonstrates the enhanced complexity of real mixed-load sediments and the importance of additional factors that have not been examined in the simplified experiments, which presently mark the state-of-the-art in this field. Thus the present work provides much needed comparison and ground-truthing of these experimental results and an exploration of the strengths and limitations of the present laboratory based understanding in the field. Additionally, the techniques and approaches developed herein for the examination of these different processes also offer guidance for future studies in this field. Lastly, we think that the publication of the Lichtman *et al.*, (2018) paper also addresses a number of the issues raised by Reviewer 3, providing background and methodological detail, and the opportunity for the reader to independently examine these data.

INTEGRATING FIELD AND LABORATORY APPROACHES FOR RIPPLE DEVELOPMENT IN MIXED SAND–CLAY–EPS

**Baas, Jaco H.^{1*}, Baker, Megan L.¹, Malarkey, Jonathan^{1,2}, Bass, Sarah, J.³, Manning, Andrew
J.^{2,3,4}, Hope, Julie A.⁵, Peakall, Jeffrey ⁶, Lichtman, Ian D.⁷, Ye, Leiping ², Davies, Alan. G.¹,
Parsons, Daniel. R.², Paterson, David M.⁸ and Thorne, Peter D.⁷**

¹ School of Ocean Sciences, Bangor University, Menai Bridge, Isle of Anglesey, LL59 5AB, United Kingdom

² Department of Geography, Environment and Earth Sciences, University of Hull, Hull, HU6 7RX, United Kingdom

³ School of Marine Science and Engineering, Plymouth University, PL4 8AA, United Kingdom

⁴ HR Wallingford, Howbery Park, Wallingford, OX10 8BA, United Kingdom

⁵ Institute of Marine Science, University of Auckland, Private Bag 92019, Auckland, New Zealand

⁶ School of Earth and Environment, University of Leeds, LS2 9JT, United Kingdom

⁷ National Oceanography Centre, Joseph Proudman Building, 6 Brownlow Street, Liverpool, L3 5DA, United Kingdom

⁸ Scottish Oceans Institute, School of Biology, University of St. Andrews, St. Andrews, KY16 8LB, United Kingdom

* Email: j.baas@bangor.ac.uk

ABSTRACT

The shape and size of sedimentary bedforms play a key role in the reconstruction of sedimentary processes in modern and ancient environments. Recent laboratory experiments have shown that bedforms in mixed sand–clay develop at a slower rate and often have smaller heights and wavelengths than equivalent bedforms in pure sand. This is generally attributed to cohesive forces that can be of physical origin, caused by electrostatic forces of attraction between clay minerals, and of biological origin, caused by ‘sticky’ extracellular polymeric substances (EPS) produced by micro-organisms, such as microalgae (microphytobenthos) and bacteria. In the present study, we demonstrate, for the first time, that these laboratory experiments are a suitable analogue for current ripples formed by tidal currents on a natural mixed sand–mud–EPS intertidal flat in a macrotidal estuary. Integrated hydrodynamic and bed morphological measurements, collected during a spring tide under weak wave conditions near Hilbre Island (Dee estuary, NW England), reveal a statistically significant decrease in current ripple wavelength for progressively higher bed mud and EPS contents, and a concurrent change from three-dimensional linguoid to two-dimensional straight-crested ripple planform morphology. These results agree well with observations in laboratory flumes, but the rate of decrease of ripple wavelength as mud content increased was found to be substantially greater for the field than the laboratory. Since the formation of ripples under natural conditions is inherently more complex than in the laboratory, four additional factors that might affect current ripple development in estuaries, but which were not accounted for in laboratory experiments, were explored. These were current forcing, clay type, pore water salinity, and bed EPS content. Our data illustrate that clay type alone cannot explain the difference in the rate of decrease in ripple wavelength, as the bed clay contents were too low for clay type to have had a measurable effect on bedform development. Accounting for the difference in current forcing between the field and experiments, and therefore the relative stage of development with respect to equilibrium ripples, increases the difference between the ripple wavelengths. The presence of strongly cohesive EPS in the current ripples on the natural intertidal flat might explain the majority of the difference in the rate of decrease in ripple wavelength between the field and the laboratory. The effect of pore water salinity on the rate of bedform development cannot be quantified at present, but salinity is postulated herein to have had a smaller influence on the ripple wavelength than bed EPS content. The common presence of clay and EPS in many aqueous sedimentary environments implies that a re-assessment of the role of current ripples and their primary current lamination in predicting and reconstructing flow regimes is necessary, and that models that are valid for pure sand are an inappropriate descriptor for more complex mixed sediment. We propose that this re-assessment is necessary at all bed clay contents above 3%.

Running title: Mixed sand bed-clay-EPS transport

Keywords: Current Ripples, Mixed Sand–Clay–EPS, Cohesion, Estuary, Intertidal Flat

INTRODUCTION

In estuaries, the interaction of river flows, tides, and waves leads to more complex particle movement and resulting spatiotemporal distribution of deposits than in many other environments (*e.g.*, Dalrymple & Choi, 2007; van den Berg *et al.*, 2007). In particular, the common presence of sand mixed with cohesive clay and non-cohesive silt – with silt and clay collectively referred to as mud herein – renders the reconstruction of sedimentary processes from bedforms and primary current lamination in estuarine sedimentary facies a challenge, because cohesive forces can have a large influence on the erosion, transport, and deposition of sediment (*e.g.*, Mehta, 2013; Chen *et al.*, 2017). Even small volumes of mud-sized particles in sand – well within ‘clean sand’ and ‘mature sandstone’ (arenite), defined as sand with less than 25% mud by Shepard (1954) and less than 10-15% mud by Folk (1951) and Dott (1964) – are able to bind sediment particles via electrostatic van der Waals forces. This binding increases the threshold stress for sediment entrainment from the seabed, promotes flocculation of suspended sediment, and changes depositional properties, compared to non-cohesive silt and sand (*e.g.*, Mehta, 2013). In addition to this physical cohesion, estuaries are also major sites of primary and secondary production, which promotes the development of biological cohesion by organic molecules produced through biological activity (*e.g.*, extracellular polymeric substances, EPS), acting to bind particles through electrostatic interactions, hydrogen bond formation, and cation effects (*e.g.*, Underwood & Paterson, 2003). EPS is produced in sediment by microphytobenthos, chiefly diatoms and cyanobacteria (Underwood & Paterson, 2003), and bacteria.

At present, cohesion is not sufficiently well-incorporated either into engineering models for estuarine sediment transport or into geological facies models of estuaries (*e.g.*, Le Hir *et al.*, 2007; Wang & Sturm, 2016). Recent experimental work has shown that further fundamental physical sedimentological research is needed to close the knowledge gap with non-cohesive sedimentary dynamics (Baas *et al.*, 2013, Malarkey *et al.*, 2015, Schindler *et al.*, 2015; Parsons *et al.*, 2016; Baas *et al.*, 2016). These laboratory studies have highlighted that sedimentary bedforms and their primary current lamination in mixed sand–clay are significantly different from those in pure sand. Fractions of 1-10% of cohesive clay and less than 0.1% of EPS within a sand bed are sufficient to increase the development time of current ripples on a flat bed (Baas *et al.*, 2013, Malarkey *et al.*, 2015), and reduce the equilibrium height and wavelength of subaqueous dunes in mixed sand-clay (Schindler *et al.*, 2015)

and in mixed sand-clay-EPS (Parsons *et al.*, 2016). This decrease in bedform size, as clay and EPS content are increased, is particularly pronounced for subaqueous dunes (Schindler *et al.*, 2015; Parsons *et al.*, 2016). However, we hypothesise that current ripples in mixed sand–clay–EPS will also be smaller than in pure sand when developed within the time frame of a semidiurnal tide, because the additional time required to reach equilibrium dimensions compared to pure sand ranges from hours to tens of hours (Baas *et al.*, 2013; Malarkey *et al.*, 2015). A key process during bedform development in mixed sediment is the entrainment of clay, silt, and EPS into suspension from a mixed sand–mud bed, *i.e.* winnowing (McCrone, 1962), as it may facilitate the bedforms eventually reaching heights and wavelengths that are similar to their mud-free equivalents. This facilitative process appeared more important for current ripples (Baas *et al.*, 2013; Malarkey *et al.*, 2015) than for dunes (Schindler *et al.*, 2015; Parsons *et al.*, 2016). The above results were based exclusively on controlled laboratory experiments conducted with steady, uniform flow, constant flow depth, well-sorted sand, a single type of clay mineral (kaolinite) and EPS (xanthan gum), and in fresh (Baas *et al.*, 2013; Malarkey *et al.*, 2015) or brackish (Schindler *et al.*, 2015; Parsons *et al.*, 2016) water.

The present paper compares, for the first time, the dynamics of experimental small-scale bedforms in mixed sand–clay with those on an intertidal flat (Dee Estuary, NW England), based on integrated morphological and hydrodynamical data. Firstly, evidence is provided that current ripples found at low slack water at this field site and current ripples in laboratory experiments are influenced by physical and biological cohesion in a similar way. Thereafter, the inherently more complex field conditions are explored to elucidate the underlying processes that control the differences between the field data and the experimental data in the rate of change of current ripple wavelength with changing bed clay and EPS content. Finally, the wider implications of this validation study for predictive sediment transport models, bedform size predictors, and sedimentary facies analysis are discussed.

METHODS

The data described herein integrate laboratory experiments with a comprehensive set of field measurements in an intertidal environment near Hilbre Island in the Dee Estuary, NW England (Fig. 1A). The Dee Estuary is funnel-shaped and macrotidal, with a 7–8 m mean spring tidal range at Hilbre Island. Hilbre Island separates Hilbre Channel from intertidal flats west of the town of West Kirby (Fig. 1A). These tidal flats are flood-dominated and rich in fine-grained sediment (Moore *et al.*, 2009). Waves are mainly generated locally within Liverpool Bay, with northwesterly waves having the largest influence on sedimentary processes in the Dee Estuary (Brown & Wolf, 2009; Villaret *et al.*, 2011).

The intertidal flats to the northwest of Little Eye (Fig. 1A) have proven valuable as a natural laboratory for studying bedform dynamics in mixed sand–mud, owing to the large variety in sand–mud ratio, ranging from pure sand to sandy mud. During the field campaign in May 2013, the intertidal flats also contained significant amounts of EPS, ranging from 0.02% to 0.21% by weight (Lichtman *et al.*, 2018). The present paper uses textural and morphological data collected from a 1400 m² area on May 26th in 2013, where conditions had been essentially wave-free in the previous two tidal inundations. The bed was either flat, *i.e.* locally below movement threshold, or rippled with different sizes and planform patterns during exposure at low slack water (Fig. 1B). In order to study the relationship between these bedform properties and bed clay and EPS content, three plane beds and twelve rippled beds were described visually. Subsequently, using the method described in Appendix A, the 3D nature of the plan morphology of the ripples was characterised by the ripple planform index, I , where $I > 0.50$ for 2D ripples, $0.39 \leq I \leq 0.5$ for 2D-3D ripples and $I < 0.39$ for 3D ripples. The bedform wavelengths were measured by hand (collecting bedform heights within the same low slack water period was not possible), and bulk sediment samples were collected between the crest and base of individual bedforms, and down to 20 mm below the base of these bedforms, for subsequent grain size analysis using a Malvern 2000 laser particle sizer. This approach allowed comparison of surface sediment subject to winnowing (*i.e.* loss of fine particles and EPS to the water column) with deeper sediment unaffected by winnowing. The bed sampling also included flat, mud-rich, surfaces without current ripples. For these flat beds, the subsurface samples were collected in the same way as for the rippled beds, *i.e.* down to 20 mm below the surface. These samples were supplemented by surface scrapes, several millimetres deep, to distinguish undisturbed sediment from sediment that may have been subjected to winnowing at the sediment–water interface. The sand fraction in the mixed sand–mud at the study site had a median grain diameter of 0.228 mm, and the mud fraction, encompassing all particles finer than 0.063 mm, ranged from 3.8% to 37.1% by volume. The mud fraction of the ripples at the field site contained 36% clay minerals by volume (standard deviation: 4%, $n = 7$), based on X-ray powder diffraction (XRD) data, obtained using the standard methodology for bulk sediment analysis (Moore & Reynolds, 1997). In decreasing order of abundance, the clay mineral assemblage comprised illite (68%), chlorite (16%), kaolinite (13%), and mixed layer illite/smectite (3%). This 36% is therefore inferred to represent the cohesive fraction within the mud fraction; the remaining 64% of the mud fraction was dominated by non-cohesive quartz and feldspar. Following Lichtman *et al.* (2018), the cohesive clay fraction, c , can therefore be expressed in terms of the mud content, m , as:

$$c = 0.36m \quad (1)$$

thus yielding cohesive clay fractions ranging from 1.4% to 13.4% by volume. Because of time constraints during low slack water, it was not possible to collect EPS samples that exactly

corresponded to these clay samples, since the former needed to be frozen in liquid nitrogen in the field (Hope, 2016). It was thus necessary to rely on the relationship between clay content and EPS content by weight, e , determined nearby (Lichtman *et al.*, 2018):

$$e = 0.0105c + 0.0302 \quad (2)$$

($R^2 = 0.41$, $p < 0.05$, $n = 20$), yielding EPS contents ranging from 0.04% to 0.17% by weight. Adjacent to the area where the bedforms were measured, near-bed flow velocities were measured continuously during water cover using two frame-mounted acoustic Doppler velocimeters (ADV) located at 0.25 m and 0.40 m above the sediment surface. The ADV data, with an acquisition rate of 0.2 Hz, was bin-averaged over 30 s to determine Reynolds-averaged velocities in the absence of waves. Immediately before the bedforms were studied on May 26th, the field site experienced a spring tide with a maximum depth-averaged flow velocity of 0.67 m s^{-1} (based on a total period of submergence of both probes of 4.7 hours) and southeastward- and northwestward-directed currents during flood and ebb, respectively. The maximum flow depth was 2.73 m close to high slack water. The depth-averaged velocities were calculated from the vectorially added horizontal components of the ADV data and the flow depth, using the standard logarithmic law for wall-bounded shear flow (*e.g.*, van Rijn, 1990).

RELATIONSHIP BETWEEN RIPPLE PROPERTIES AND BED MUD CONTENT

Results

Table 1 provides the current ripple wavelength and planform index data and the corresponding bed mud and cohesive clay measured at the study site. The mean ripple wavelength, L , is plotted against bed mud percentage within the ripples and in the subsurface underneath the ripples in Fig. 2. Figure 2 also provides information on the planform properties of these bedforms, and photographic imagery of the various plan morphologies are shown in Fig. 3.

A statistically significant decreasing linear relationship between the mean ripple wavelength and the bed mud content, m , was found for the samples collected from the ripples and the subsurface (Fig. 2). These relationships are described by a linear trend with R^2 values of 0.44 ($p = 0.027 < 0.05$, $n = 12$) and 0.75 ($p = 0.00058 < 0.05$, $n = 12$), respectively. The mud content in the subsurface was mostly greater than the mud content within the ripples, and this difference rapidly increased as the ripple wavelength decreased (Fig. 2). For the beds with the lowest mud content, the ripple wavelength was close to the equilibrium wavelength of current ripples in pure sand with a median grain size, D_{50} , similar to the present study ($D_{50} = 0.238 \text{ mm}$ and $L = 141 \text{ mm}$ (Baas, 1999) versus $D_{50} = 0.228 \text{ mm}$ and $L = 142 \text{ mm}$

herein). The smallest ripples, formed at the highest bed mud contents, were approximately 20 mm longer than incipient current ripples in the experimental study of Baas (1999) (Fig. 2).

Figure 3A-B shows examples of flat surfaces without ripples, which were found in concert with high subsurface mud contents of 23.4%, 33.8%, and 37.1% (Table 1). As observed for the rippled beds, the mud content in the surface scrapes of the flat beds was consistently lower than in the subsurface samples down to 20 mm (Table 1). The average mud content in this surficial layer was 23%, which was higher than any of the mud contents within the ripples. The current ripples with $L \leq 95$ mm (Table 1) had straight to weakly sinuous crest lines (Fig. 3C-D); these bedforms are referred to as two-dimensional (2D) ripples in the remainder of this paper (Fig. 2). The ripples with a wavelength greater than 110 mm were either three-dimensional (3D) ripples with linguoid crests (Fig. 3G-H), or transitional between the 2D and 3D ripples (Fig. 3E-F). The beds covered in 2D-3D ripples consisted of patches of linguoid ripples next to ripples with more continuous crests. With one exception, the 2D-3D ripples were shorter in wavelength than the 3D ripples (Fig. 2).

Interpretation

These field data reveal a decreasing linear relationship between bed mud content and ripple wavelength, suggesting that clay minerals within the bed hindered the near-bed movement of non-cohesive grains (*e.g.*, Mitchener & Torfs, 1996; Baas *et al.*, 2013; Wang & Sturm, 2016). Consequently, the tidal currents were able to form 3D linguoid current ripples only in beds with $m < 11\%$ in the subsurface and $m < 6\%$ within the ripples. At progressively higher bed mud contents, cohesive forces restricted bedform development to smaller 2D-3D and 2D current ripples. The boundary between these ripple types was at bed mud contents in the subsurface and within the ripples of 18% and 10%, respectively. The beds were too cohesive for any ripple development at subsurface bed mud contents of $31 \pm 7\%$ and surficial mud contents of $23 \pm 11\%$, where 7% and 11% denote standard deviations of the mean.

The wavelength and planform morphology of the field ripples correspond remarkably well with the laboratory ripples in 0.238 mm sand of Baas (1999). The field ripples were too large to be classified as the incipient ripples of Baas (1999), which were up to 67 mm long. The wavelength of the 2D field ripples was well within the limits of the straight-crested and sinuous ripples of Baas (1999) (Fig. 2), and the 2D-3D ripples and most 3D ripples at the field site correspond in wavelength to the non-equilibrium linguoid ripples of Baas (1999). Only at the lowest bed mud contents did the 3D ripples reach wavelengths that were similar to the equilibrium linguoid ripples in the clay-free sand of Baas (1999).

COMPARISON WITH EXPERIMENTAL MIXED SAND–CLAY RIPPLES

To facilitate a direct comparison of the ripples at the field site with the ripples in the mixed sand–clay experiments of Baas *et al.* (2013), the bed mud contents in the subsurface samples collected at the field site, shown in Fig. 2, were converted to cohesive clay contents using Equation 1. The difference in cohesive clay content in the samples from the subsurface and the ripples agrees with previous studies that associated the development of bedforms with the winnowing of fine particles (*e.g.*, Baas *et al.*, 2013; Schindler *et al.*, 2015; Parsons, *et al.*, 2016; Lichtman *et al.*, 2018). The winnowing depth is limited by the height of the bedforms, which explains why the clay content in the subsurface is up to 3 times higher than the clay content in the ripples on the surface (Table 1). However, as shown in Fig. 4A, the winnowing of fine sediment in the field was not as efficient as in the laboratory experiments. Defining the boundaries of winnowing efficiency as 100% for full removal of bed clay and as 0% for full retention of clay during bedform development, the average winnowing efficiency at the field site was 35%, compared to 93% in the experiments of Baas *et al.* (2013). This low winnowing efficiency may be associated with physical and biological processes that counteract winnowing and add and store clay in the bed under field conditions, such as advection of suspended clay particles, deposition of mud flocs during high slack tide, mixing of clay and silt into the bed by benthic organisms, and hyporheic pumping (Packman *et al.*, 2000; Blois *et al.*, 2014). Following the approach of Malarkey *et al.* (2015), described in Appendix B, the time scale of hyporheic pumping into the bed – a minimum of 19.7 h equivalent to four inundation periods – is far slower than the winnowing associated with ripple overturning (0.2 to 0.4 h). Winnowing appears to have taken place also on the strongly cohesive flat beds, although this process might have been limited to the uppermost millimetres, and the reduction of bed mud content was insufficient to enable ripple formation within the time available. In addition, the production, accumulation, breakdown, and loss of EPS is complex given its multiple sources (bacteria, microphytobenthos and infauna; Decho & Gutierrez, 2017) compared to the rather simple situation represented in the laboratory with known initial concentrations of EPS, no active production, and xanthan gum being used as a proxy for EPS.

In order to further facilitate comparisons between the field and laboratory data, the ripple wavelengths were non-dimensionalised to exclude the influence of grain size on current ripple dimensions (Raudkivi, 1997; Baas, 1999; Soulsby & Whitehouse, 2005):

$$\Lambda = \frac{L - L_0}{L_E - L_0} \quad (3)$$

where Λ is the dimensionless ripple wavelength and L_E and L_0 are the wavelengths of equilibrium ripples and incipient ripples, first appearing on a flat bed, in pure sand. Since $\Lambda = 0$ for $L = L_0$ and $\Lambda = 1$ for $L = L_E$, Λ can also be thought of as the relative age of the ripple. L_E and L_0 are a function of grain size. Herein, $L_E = 142$ mm and $L_0 = 60$ mm for the 0.228 mm sand at the field site, where L_0 is taken from the experimental data for 0.238 mm sand of Baas (1999), and $L_E = 115.9$ mm and $L_0 = 28.5$ mm for the 0.143 mm sand used in the experiments of Baas *et al.* (2013). The experiments of Baas *et al.* (2013) were conducted with steady, uniform, fresh-water flow at a depth-averaged velocity of 0.36 m s^{-1} and the current ripples were given 2 hours to develop on a flat bed consisting of mixed sand and kaolin clay.

Figure 4B demonstrates that the dimensionless wavelength of the current ripples decreased linearly with increasing initial cohesive clay content in the bed for both the field data and the laboratory data. Moreover, both datasets show a change from 3D ripples via 2D-3D ripples to 2D ripples, as the initial cohesive clay content in the bed was increased (Baas *et al.*, 2013), and these planforms are demarcated by similar Λ values and I values (Fig. 4B; Table 1). The flat-bed data points from the field site, which were excluded from Fig. 2, can now be included, because they correspond to $\Lambda = 0$. The laboratory experiments thus provide an appropriate analogue for the effect of cohesive forces on current ripple size and planform morphology under estuarine conditions. However, the rate of decrease in ripple wavelength was significantly greater for the field ripples than for the experimental ripples. The field data also show more scatter than the laboratory data, particularly for high clay contents. Linear extrapolation predicts a total absence of ripple development at 13% cohesive clay at the field site (Fig. 4B).

This difference in the rate of change in ripple wavelength may be caused by physical and biological variables that were omitted from the flume experiments of Baas *et al.* (2013). Below, the possible influence on bedform dynamics of the following main variables is assessed: (i) clay mineral type; (ii) current shear stress; (iii) bed EPS content; and (iv) pore water salinity. Increased current shear stresses are expected to promote the development of current ripples on the intertidal flat, whereas strongly cohesive clay minerals, the presence of biological cohesion in the form of EPS, and seawater salinity, which may modify biological and physical cohesion, should hinder the development of the current ripples. Each of these variables is discussed in detail below, and, if possible, its relative contribution to the difference in the rate of change of ripple wavelength is determined (Fig. 4B).

DISCUSSION OF SPECIFIC CONTROLS ON RIPPLE DYNAMICS AT THE FIELD SITE

Clay type

Baas *et al.* (2013) used kaolinite clay in their experiments, whereas illite clay was dominant at the field site. Illite is more cohesive than kaolinite because illite particles have a larger specific surface area and a larger cation exchange capacity than kaolinite particles (Hillel, 2004; Yong *et al.*, 2012). Mixtures of sand and illite should, therefore, have a higher yield strength than mixtures of sand and kaolinite, and bedforms might develop more readily in mixed sand–kaolinite. However, Baker *et al.* (2017) showed that the difference in yield strength between kaolinite and montmorillonite (incl. bentonite) is small for volumetric clay concentrations below 10%. This result agrees with an experimental study by Torfs (1995, in van Ledden, 2001), in which the critical clay content for cohesive behaviour was 3–4% for both kaolinite and montmorillonite. Since illite is less cohesive than montmorillonite, all the current ripples at the field site were formed at bed clay fractions below 13%, and the subsidiary clay minerals kaolinite and chlorite in the field samples have lower yield strength than the illite, it is inferred that the effect of clay type on the difference in the rate of change of ripple wavelength between the field data and Baas *et al.*'s (2013) laboratory data was small.

Current shear stresses

The experiments of Baas *et al.* (2013) were conducted at a constant depth-averaged flow velocity of 0.36 m s^{-1} , whereas the field site was subjected to unsteady flow with a maximum depth-averaged flow velocity of 0.67 m s^{-1} (Fig. 5B). As the rate of bedform development increases non-linearly with flow velocity (Baas, 1994, 1999), the current ripples at the field site could have been preserved at the end of ebb flow in a different development stage than the mixed sand–clay ripples of Baas *et al.* (2013). This development stage would render these ripples *non-equilibrium* bedforms *sensu* Baas (1994). Before testing this hypothesis, it is important to consider the alternative that the ripples on the intertidal flat were smaller than ripples in the equivalent clay-free sand because they are *equilibrium* bedforms (Baas, 1994) with wavelengths that decrease with increasing bed clay content. Baas *et al.* (2013) discussed both possibilities without stating a preference, but, subsequently, Malarkey *et al.* (2015) provided experimental evidence that current ripples in cohesive mixtures of sand and EPS reach equilibrium heights and wavelengths that are similar to those in non-cohesive sand, provided that the flow duration is sufficiently long. Since the Baas *et al.* (2013) ripples were essentially free of clay (Fig. 4A), and probably still developing at the end of their experiments, it can be assumed that this evidence can be extrapolated from biological cohesion to physical cohesion. Thus the mathematical model of Oost & Baas (1994) and Baas *et al.* (2000) for current ripple development in unsteady flow was used to compare the development stage of the ripples at the field site and in the experiments of Baas *et al.* (2013):

$$\frac{L_{SL} - L_0}{L_E - L_0} = 1 - 0.01^{S_L} \quad \wedge \quad S_L = \int_0^T \frac{dt}{T_E(t)} \quad (4)$$

$$T_E = \left(\frac{\theta' - \theta_c}{a} \right)^{\frac{1}{b}} \quad (5)$$

In Equation 4, L_{SL} is the ripple wavelength at time T (in hours), T is the flow duration in hours, S_L is the cumulative development stage of ripple wavelength, and $T_E(t)$ is the equilibrium time in hours as a function of time t , *i.e.* the time needed to reach equilibrium ripple wavelength at the acting current strength. $S_L = 0$ for a flat bed, and $S_L \geq 1$ for equilibrium ripples. In Equation 5, T_E is related to the inverse of the flow forcing:

$$\theta' = \frac{U^2}{(\rho_s / \rho - 1) C'^2 D_{50}} \quad (6)$$

$$C' = 18 \log \left(\frac{12h}{3D_{90}} \right) \quad (7)$$

where θ' is the grain-related mobility parameter, U is the depth-averaged flow velocity, $\rho_s = 2650 \text{ kg m}^{-3}$ is the sediment density, $\rho = 1027 \text{ kg m}^{-3}$ is the density of seawater, D_{50} is the median grain size of the bed sand fraction, C' is the grain-related Chézy coefficient, h is the flow depth, and D_{90} is the 90-percentile of the grain-size distribution of the sand fraction (herein, $D_{90} = 0.364 \text{ mm}$).

In Equation 5, θ_c represents the critical mobility parameter for particle entrainment, parameterised from the Shields curve (Shields, 1936) by Soulsby (1997):

$$\theta_c = \frac{0.3}{1 + 1.2D_*} + 0.055(1 - e^{-0.02D_*}) \quad (8)$$

$$D_* = \left(\frac{(\rho_s / \rho - 1)g}{\nu^2} \right)^{1/3} D_{50} \quad (9)$$

where D_* is the dimensionless grain size parameter, $g = 9.81 \text{ m s}^{-2}$ is the constant due to gravity, and $\nu = 1.37 \times 10^{-6} \text{ m}^2 \text{ s}^{-1}$ is the kinematic viscosity of seawater at 10°C . The coefficients a and b in Equation 5 are grain-size dependent. For the field site with 0.228 mm sand herein, $a = 0.112$ and $b = 0.473$, which are linearly interpolated from values for 0.095 mm and 0.238 mm sand (*cf.*, Baas *et al.*, 2000).

Equations 4 to 9 were used to predict the final wavelength of clay-free current ripples developing on a flat bed during the flood and ebb tides immediately preceding the data collection (Fig. 5). The grain-related mobility parameter (Fig. 5C) was calculated from the water depth (Fig. 5A) and the depth-averaged flow velocity (Fig. 5B). In turn, θ' was used to predict the cumulative ripple development

stage (Fig. 5D) and finally the temporal development of ripple wavelength (Fig. 5E). As expected, the highest rates of ripple wavelength development occurred around maximum flood and maximum ebb flow. The ripples were predicted to attain their full equilibrium wavelength after 4.4 h of the available 4.7 hours of tidal flow (Fig. 5D-E). The model predicts $S_L = 1.263$ at time T , with $L_{SL} = 141.8$ mm. However, the age of the sandiest ripples at the field site might extend further back in time, as it is unknown how long these bedforms had been at equilibrium. Yet, it is unlikely that these bedforms formed more than three or four principal lunar semi-diurnal tidal cycles before the ripple wavelength data were collected because strong waves on May 24th generated wave ripples, combined flow ripples, and upper-stage plane beds (Lichtman *et al.*, 2018). The model predictions for S_L and L_{SL} in Fig. 5D-E should, therefore, be considered minimum values.

In the experiments of Baas *et al.* (2013), the equilibrium wavelength of the current ripples in pure sand, based on $L_{SL} = 0.99 L_E$ (*cf.*, Equation 4), was reached at 2.7 h. Since the experiments with mixed sand–clay were halted after 2 h, this equilibrium time corresponds to $S_L = 2/2.7 = 0.74$, which can be compared with the S_L -value in the field of 1.26 by adjusting the laboratory data of Baas *et al.* (2013) according to:

$$\Lambda = 1 - \left(\frac{L_E - L_{0.74}}{L_E - L_0} \right)^{\frac{1.26}{0.74}} \quad (10)$$

where $L_{0.74}$ is the ripple wavelength after 2 h. Equation 10 is represented by the upper curve in Fig. 6. This curve is above the curve for the laboratory data (Fig. 6) because the experiments have been adjusted as though they were run over a longer period: $1.26 \times 2.7 = 3.4$ h. Forcing the relative ages of the laboratory and field data to be the same in this way has resulted in an increase in the difference between these datasets. Hence, current shear stress cannot explain the observed difference in the rate of change of ripple wavelength between the field data and the laboratory data.

Bed EPS content

An important difference between the field data and the laboratory data of Baas *et al.* (2013) is the lack of biological cohesion in the flume experiments. The EPS content in the bed samples from the field site was not measured directly, as explained earlier, but Equation 2 can be used to make a qualitative assessment of the effect of EPS on the remaining difference in the rate of change of dimensionless ripple wavelength with increasing bed clay content (Fig. 6). The subsurface samples contained up to 0.194% EPS (Equation 2), which is comparable to the bed EPS content in the laboratory data of Malarkey *et al.* (2015). According to Malarkey *et al.* (2015), the time needed for incipient ripples to appear on the flat bed, t_i , is 0.1, 0.4, 1.0, 2.7 and 7.9 h for values of e of 0%, 0.015625%,

0.03125%, 0.0625% and 0.125%, respectively. This can be expressed as $t_i = 0.1 + 204.1e^{1.57}$, where e is given by Equation 2, such that an effective S_L can be defined as $(3.4 - t_i)/2.7 = 1.26 - 0.37t_i$, where S_L is forced to be greater than 0, and gives Λ as:

$$\Lambda = 1 - \left(\frac{L_E - L_{0.74}}{L_E - L_0} \right)^{\frac{1.26 - 0.37t_i}{0.74}} \quad (11)$$

which is the curve labelled “Equation 11” in Fig. 6. This curve represents a conservative estimate of the effect of EPS, since Equation 11 does not include the considerably longer time needed to develop from incipient to equilibrium linguoid current ripples in mixed sand–EPS as opposed to pure sand (Malarkey *et al.*, 2015). Despite this underprediction, the calculated effect of xanthan gum on ripple wavelength development rate is still considerably stronger than the observed effect (Fig. 6). Yet, Equation 11 captures the three flat-bed cases reasonably well. It is likely that the effect of xanthan gum on bedform development is stronger than that of naturally occurring EPS (Tolhurst *et al.*, 2002; van de Lageweg, 2017). Xanthan gum is strongly cohesive, and has a constant distribution in the vertical, so the experiments may not be representative of estuarine conditions, where EPS typically reaches a maximum concentration in biofilms at the sediment surface (Taylor & Paterson, 1998). We thus conclude the difference between the field and laboratory can reasonably be explained by the presence of EPS in the bed samples from the Dee Estuary.

The long delay in the initiation, as well as the long development time, of the current ripples in the EPS–sand experiments (Malarkey *et al.*, 2015), are likely to be related to increases in the threshold of motion. Thus, the flows need to first lower the threshold of motion through winnowing of EPS before ripples begin to form. Based on a separate series of flume experiments (see Appendix C for further details), Fig. 7 provides an estimation of the increase of θ_c for kaolin clay and the xanthan gum, in which the critical Shields parameter was determined for mixed sand–clay–EPS by gradually increasing the bed shear stress imposed by a steady, uniform flow until a downstream bedload trap started to collect sediment particles. For up to 1% xanthan gum and up to 30% kaolin, θ_c was up to 73% and 83% higher than for the non-cohesive reference sand. The combined effect of biological and physical cohesion led to an increase in θ_c by up to 123%. For the field site, where the current ripples formed only below 13% clay and below 0.2% EPS, Fig. 7 suggests a maximum increase in θ_c by 27% for the mixed sand–clay–EPS. It should be emphasised, however, that this percentage increase is merely an approximation of the combined effect of physical and biological cohesion on θ_c because the experiments were conducted with types of sand, clay, and EPS that were different from those at the field site.

Water salinity

The laboratory experiments of Baas *et al.* (2013) were conducted with fresh water, whereas the current ripples at the field site were formed in seawater. Salinity is well known to promote the formation of cohesive bonds between clay particles through the availability of cations that neutralise the negatively charged edges of clay platelets and therewith stimulate the attraction of these platelets by van der Waals forces (*e.g.*, Manning *et al.*, 2007; Mehta, 2013). This physicochemical attraction leads to the development of clay aggregates, *i.e.* floccules, that often have higher settling velocities than their constituent particles. The effect of salinity is larger for suspended sediments than for erosion thresholds. Experimental studies typically find a positive correlation between erosion threshold and salinity for muddy sediment (*e.g.*, Kandiah, 1974; Parchure & Mehta, 1985), but this is not always the case. For instance, no significant statistical relationship between salinity and critical shear stress existed for mixed sand—EPS in the laboratory experiments of van de Lageweg *et al.* (2017). Furthermore, the relationship between salinity and critical shear stress for sediment motion in field studies is ambiguous. Amos *et al.* (2003) found no difference in the threshold of erosion for natural muds in marine and freshwater environments, whereas Debnath *et al.* (2007) obtained a negative correlation between salinity and erosion rate for natural muds, and Spears *et al.* (2008) found that the erosion threshold increased seaward with salinity in the Eden Estuary (south-eastern Scotland). These differences between the field studies, in comparison to theoretical and experimental approaches, suggest that at least in some cases the effect of salinity on bed erosion is masked by other parameters in the natural environment, such as bed density and EPS content (*e.g.*, Grabowski *et al.*, 2011). For the present study, it is postulated that the difference in the rate of decrease of dimensionless ripple wavelength, as cohesive clay content is increased, between the field data and the experimental data (Fig. 6) is partly caused by the difference in pore water salinity. However, based on the available literature, we expect this effect to be secondary. Some support for salinity having a secondary effect is the consistency in the development of ripples in fresh water (Baas *et al.*, 2013; Malarkey *et al.*, 2015) and ripples and dunes in brackish water (Schindler *et al.*, 2015; Parsons *et al.*, 2016).

WIDER IMPLICATIONS

Frequency of occurrence of equilibrium and non-equilibrium current ripples

At the field site, the presence of cohesive clay and EPS was found to have had the greatest influence on the growth rate and size of the current ripples, followed in decreasing order of influence by flow

forcing and salinity, whereas the influence of clay type was small to negligible. However, the relative contribution of these controlling variables is likely to be unique to the measurement period at the study site.

Physical and biological cohesion are expected to have a strong influence on current ripple dynamics because small fractions of clay and EPS, akin to most aqueous environments, are sufficient to reduce current ripple size, promote two-dimensionality of ripples, and delay ripple growth. In environments where the currents and waves are stronger than at the field site, *e.g.*, in tidal channels and on wave-facing beaches, sand-rich equilibrium bedforms are probably more common, because stronger hydrodynamic forcing promotes the development rate of the bedforms, assisted by a faster rate of winnowing of clay and EPS from the bed. This raises the question if non-equilibrium mixed sand–clay–EPS ripples will dominate the bed when the currents are weaker than at the field site. In fact, this weaker hydrodynamic forcing should be the more likely scenario because the mean spring tidal range of 7–8 m in the Dee estuary is amongst the largest in Europe, and the field data were collected at spring tide in this macrotidal estuary. Under such circumstances, maximum tidal velocities are higher than in many microtidal, mesotidal and other macrotidal environments. However, since the formation of the current ripples in this study was restricted to bed clay and EPS fractions of up to only 13% and 0.2%, we infer that mixed sand–clay–EPS ripples form less readily on sand–mud tidal flats in these lower-energy environments, even under the consideration that these percentages might vary with clay type, EPS type, and flow forcing. Assuming that it is possible to form bedforms at all in such conditions, many tidal cycles would be needed to form recognisable mixed sand–clay–EPS ripples, and their preservation would depend on the frequency of occurrence of processes that destroy these bedforms, such as bioturbation and storm waves. Mixed sand–clay–EPS ripples might be restricted to a narrow strip at the boundary between sandy tidal channels and muddy tidal flats in estuaries with a smaller tidal range than the study site. Likewise, it is anticipated that during neap tides in the Dee Estuary, and in other macrotidal estuaries with a similar tidal range, the mixed sand–clay–EPS bedforms shift from intertidal flats towards areas of confined tidal flow, in particular subtidal channels.

Although the effect of pore water salinity on current ripple growth rate and size is not fully understood at present, the concentration of cations in brackish water and seawater are expected to be large enough to cause longer delays in the development of mixed sand–clay–EPS current ripples in fully marine and estuarine environments than in freshwater environments, such as rivers and lakes. It is therefore hypothesised for the benefit of future studies that equilibrium mixed sand–clay–EPS bedforms are more common in fresh water. At the field site, the influence of clay type was found to be small. However, tidal flows with significantly larger velocities than at the field site, and possibly helped by surface water waves, might be able to form current ripples at $c > 13\%$ and $e > 0.2\%$, for

which differences in yield strength between clay types start to play a more significant role in causing differences in bed cohesion (Baker *et al.*, 2017), and therefore bedform development rate.

Implications for studies in modern environments

Accurate knowledge of the geotechnical, morphological, and biological properties of the seabed is essential for the quantification of near-bed sediment transport in modern environments. These properties include the sediment size, the cohesive strength of the bed, the critical shear stress for sediment motion, and the size and migration rate of bedforms. The rate of bedload transport per unit width, q_b , can be calculated based on the volumetric sediment concentration, C_b , the particle velocity, u_p , and the saltation height of the particles, δ_b (van Rijn, 1993):

$$q_b = C_b u_p \delta_b \quad (12)$$

or on the current ripple height, H , and the current ripple migration rate, u_r (van den Berg, 1987):

$$q_b = 0.6 (1 - P) u_r H \quad (13)$$

where P is the bed porosity. Equation 12 was expressed by van Rijn (1993) also in terms of bed shear stress and sediment size:

$$q_b = 0.053 \sqrt{O g D_{50}^3} D_*^{-0.3} \left(\frac{\tau_b - \tau_{b,c}}{\tau_{b,c}} \right)^{2.1} \quad (14)$$

where τ_b is the bed shear stress, and $\tau_{b,c}$ is the critical bed shear stress according to Soulsby (1997). Baas *et al.* (2000) found a power-law relationship between flow forcing and ripple migration rate for Equation 13, which in modified form can be written as:

$$u_r = \alpha (\theta' - \theta_c)^\beta \quad (15)$$

where α and β are coefficients dependent on sediment size.

Equations 12-15, as well as other bedload transport equations, have been used successfully for cohesionless sand, albeit with large degrees of uncertainty. The fact that “it is hardly possible to predict the transport rate with an inaccuracy of less than factor 2” (van Rijn, 1993) might result in part from the lack of parameterisation of the effect of physical and biological cohesion in these equations. Even for quite modest amounts of clay and EPS, Lichtman *et al.* (2018) demonstrated large effects on the bed material transport rate associated with ripple migration. Lichtman *et al.*'s (2018) study and the present study, combined with the recent laboratory studies, on bedform development in mixed sand–clay (*e.g.*, Baas *et al.*, 2013; Malarkey *et al.*, 2015; Parsons *et al.*, 2016) suggests that predictions of q_b could be vastly improved if cohesive forces were included in bedload transport equations.

Insufficient data are available at present to achieve this, but these improvements would require the modification of $\tau_{b,c}$ in Equation 14 and θ_c in Equation 15 for cohesive forces in the bed, even at several percent of clay and hundredths of a percent of EPS. An improved parameterisation of ripple height in Equation 13 is also essential, because existing mathematical predictors for the size of current ripples (Raudkivi, 1997; Baas, 1999; Soulsby & Whitehouse, 2005), which relate the equilibrium height (and wavelength) of the ripples to the median sediment size within the ripples, have been developed only for non-cohesive sand. Hence, these predictors are unlikely to be sufficiently accurate for mixed sand–clay–EPS.

The present study strengthens the evidence collected in previous studies (*e.g.*, Baas 1994, 1999) that current ripples are often smaller in size, and more two-dimensional in planform, than those predicted from the size of the sand particles in the bed alone. In other words, the common presence of cohesive clay and EPS in the natural environment increases the likelihood of finding smaller ripples with non-equilibrium planforms. By using an equation for the development of ripple height from a flat bed in clay-free 0.228 mm sand (*e.g.*, Baas, 1999), similar to Equations 4 and 5 but with an initial height, H_0 , equal to zero, it can be shown that the smallest ripples in the study area were approximately 4 mm high, and the equilibrium height, H_E , was 17 mm. For this 13 mm difference in ripple height alone Equation 13 would overpredict the bedload transport rate by a factor 4, if the effect of cohesion is ignored. The presence of clay and EPS also increases θ_c , which is likely to further worsen this overprediction (Fig. 7).

Implication for geological studies

Whilst not a strong indicator of depositional environment, current ripples and their primary current lamination have been used extensively for the reconstruction of depositional processes in the geological record (*e.g.*, Stow, 2005). The equilibrium height and wavelength of current ripples increase, as the median size of the sediment particles in the bed increases, but their equilibrium dimensions are independent of flow velocity (Baas, 1994). Instead, flow velocity accelerates the development towards equilibrium height and wavelength. On their way to linguoid equilibrium shape, current ripples go through distinct plan morphologies, *i.e.* incipient, straight-crested, sinuous, and non-equilibrium linguoid (Baas, 1994). This progressive increase in three-dimensionality has been used to determine the development stage of current ripples in deep-marine and shallow-marine environments (*e.g.*, Baas, 1993, Oost & Baas, 1994). However, reconstructing flow properties, such as flow velocity and bed shear stress, from the size and shape of non-equilibrium current ripples in clean sand has been challenging, because Equations 4 and 5 show that the formation of non-equilibrium ripples also includes a time factor. For example, the presence of underdeveloped straight-crested

ripples and associated tabular cross-lamination may signify weak flow of relatively long duration, but also strong flow of very short duration. The present study shows that physical and biological cohesion push current ripples further away from 3D towards 2D shapes, unless the flow is strong enough to successfully winnow the clay and EPS from the bed. Above all, a full recognition of the importance of cohesive forces for bedform dynamics, even at low bed clay and EPS fractions, requires a paradigm shift in sedimentary facies analysis. Shepard (1954) put the boundary between clean sand and dirty sand at 25% clay, whereas Folk (1951) and Dott (1964) used 10-15% sediment <0.030 mm to distinguish between mature sandstone ('arenite') and immature sandstone ('wacke'). These textural classifications have practical use for categorising rocks in outcrop and core, but the boundary percentages should not be used to differentiate non-cohesive from cohesive dynamic behaviour of mixed sand–mud. Based on the laboratory and field data presented herein, a boundary between mature and immature sand of 3% detrital clay more accurately distinguishes non-cohesive from cohesive mixed sand. This boundary limits the effect of cohesion to a 20% reduction in Λ (Fig. 6) and an 8% increase in θ_c (Fig. 7). At the study site, 3% detrital clay (Fig. 4A) was equivalent to 8% mud (<0.063 mm; Equation 1), a size fraction <0.030 mm of 7%, and 0.06% EPS. Further work is needed to verify if these boundaries are applicable beyond the limits of the present comparative study of laboratory and field bedforms.

As the development of current ripples in mixed sand–clay involves loss of clay and EPS by winnowing, facies analysis should not use the clay content within current ripples and ripple cross-laminated sand to relate current ripple size and shape to bed cohesion and hydrodynamic forcing. However, the clay content in the sediment immediately below the base of current ripples and sets of cross-lamination may in many cases be representative for the original bed clay content from which the ripples were generated, especially if the textural properties of the sediment – other than the clay content – are similar. On intertidal flats in estuaries, the field data show that time is an important limiting factor for the formation of current ripples. Following on from the above interpretations for mixed sand–clay in modern estuaries, the presence of 3D current ripples and trough cross-lamination in intertidal sandstone with more than 13% clay requires extraordinary processes of formation, such as combinations of spring tides within the macrotidal range, promotion of bedload transport by near-bed wave stress, close proximity to sandy tidal channels, and a large number of consecutive tidal cycles of slow ripple growth without destructive processes. Under normal circumstances, however, such clay-rich sandstone is expected to be associated with a total absence of current ripples or with small, 2D, straight-crested current ripples and tabular cross-lamination, whereas fully developed linguoid current ripples and their cross-lamination should dominate sandstone that is poor in cohesive clay. Based on the data presented (Figs 4 and 6), the 3% detrital clay defined above for the boundary

between mature and immature sand in the subsurface might also be an appropriate upper boundary for the presence of 3D ripples and their cross-lamination in sedimentary facies associated with tidal flats in macrotidal estuaries. This maximum in clay content is inferred to get progressively lower with decreasing tidal range in mesotidal and microtidal estuaries, with an increasing probability of finding sedimentary facies with 2D current ripples and their cross-lamination or no ripples at all. Moreover, clay-poor 3D current ripples are expected to be more common in tidal flat facies that are close to tidal channel facies, where tides accelerate owing to ebb and flood flow confinement.

CONCLUSIONS

This study shows that laboratory experiments on current ripple development in mixed sand–clay–EPS are a suitable analogue for current ripple development on mixed sand–clay–EPS intertidal flats, and possibly also in other environments where physical and biological cohesion influence sediment erosion, transport, and deposition. The comparison between the field and laboratory data has led to the following main conclusions:

- Current ripples in sand develop at an increasingly slower rate, as progressively larger volumes of cohesive clay are mixed into the sand.
- These current ripples tend to become smaller and change from three-dimensional linguoid to two-dimensional straight-crested, as bed clay content is increased.
- These findings demonstrate that clay starts to affect bedform dynamics in sandy sediment at much lower concentrations than for the ‘clean sand’ and ‘arenite’ of Shepard (1954) and Folk (1951). A revised boundary of 3% detrital clay, equivalent in the present experiments to 7% matrix sediment with a grain size <0.030 mm, appears more appropriate for the onset of bed stabilisation by cohesive forces in the seabed.
- EPS might have a similar, yet stronger, effect on the size and planform of current ripples than clay.
- Clay type had a small influence on current ripple development in mixed sand–clay–EPS, because these bedforms were generated mostly at bed clay contents for which the rheological differences between clay mineral types are small.
- Differences in cumulative flow forcing need to be accounted for in comparisons of the dynamics of current ripples in mixed sand–clay between field and laboratory.
- Pore water salinity is also expected to control bedform development in mixed sand–clay–EPS, but quantification of this influence requires further study.

- These findings have important implications for predicting current ripple properties from hydrodynamic data in hydraulic engineering and for reconstructing flow properties from current ripple size and shape in sedimentary geology.

ACKNOWLEDGEMENTS

This work was supported by the UK Natural Environment Research Council (NERC) under grant NE/I027223/1 (COHBED). We are grateful to the NOC Ocean Technology and Engineering group, and field technicians from the University of Plymouth and Bangor University for instrument set up and deployment. DMP received funding from the Marine Alliance for Science and Technology for Scotland (MASTS), funded by the Scottish Funding Council (grant reference HR09011) and contributing institutions. AM's contribution to this manuscript was partly funded by HR Wallingford Company Research project 'FineScale - Dynamics of Fine-grained Cohesive Sediments at Varying Spatial and Temporal Scales' (DDY0523). JM and DP were partially funded by a European Research Council Consolidator Award (725955). Renata Rocha kindly assisted with the field data collection as part of the Science Without Borders student exchange programme between Brazil and the UK.

REFERENCES

- Amos, C.L., Droppo, I.G., Gomez, E.A. and Murphy, T.P. (2003)** The stability of a remediated bed in Hamilton Harbour, Lake Ontario, Canada. *Sedimentology*, **50**, 149–168.
- Baas, J.H. (1993)** Dimensional analysis of current ripples in recent and ancient depositional environments. *Geologica Ultraiectina*, **106**, 199 pp.
- Baas, J.H. (1994)** A flume study on the development and equilibrium morphology of small-scale bedforms in very fine sand. *Sedimentology*, **41**, 185-209.
- Baas, J.H. (1999)** An empirical model for the development and equilibrium morphology of current ripples in fine sand. *Sedimentology*, **46**, 123–138.
- Baas, J. H., van Dam, R.L. and Storms, J.E.A. (2000)** Duration of deposition from decelerating high-density turbidity currents. *Sedimentary Geology*, **136**, 71-88.
- Baas, J.H., Davies, A.G. and Malarkey, J. (2013)**, Bedform development in mixed sand–mud: The contrasting role of cohesive forces in flow and bed. *Geomorphology*, **182**, 19-39.

- Baas, J.H., Best, J.L. and Peakall, J.** (2016) Predicting bedforms and primary current stratification in cohesive mixtures of mud and sand. *Journal of the Geological Society*, **173**, 12-45.
- Baker, M., Baas, J.H., Malarkey, J., Silva Jacinto, R., Craig, M., Kane, I. and Barker, S.** (2017) The effect of clay type on the properties of cohesive sediment gravity flows and their deposits. *Journal of Sedimentary Research*, **87**, 1176-1195.
- Best, J., Kirkbride, A. and Peakall, J.** (2001) Mean flow and turbulence structure of sediment-laden gravity currents: new insights using ultrasonic Doppler velocity profiling. In: McCaffrey, W.D., Kneller, B.C. and Peakall, J., Eds., *Particulate Gravity Currents*, International Association of Sedimentologists Special Publication, **31**, 159-172.
- Blois, G., Best, J.L., Sambrook Smith, G.H. and Hardy, R.J.** (2014) Effect of bed permeability and hyporheic flow on turbulent flow over bed forms. *Geophysical Research Letters*, **41**, 6435-6442.
- Brown, J.M. and Wolf, J.** (2009) Coupled wave and surge modelling for the eastern Irish Sea and implications for model wind-stress. *Continental Shelf Research*, **29**, 1329-1342.
- Chen, X.D., Zhang, C.K., Paterson, D.M., Thompson, C.E.L., Townend, I.H., Gong, Z., Zhou, Z. and Feng, Q.** (2017) Hindered erosion: The biological mediation of noncohesive sediment behaviour. *Water Resources Research*, **53**, 4787-4801.
- Dalrymple, R.W. and Choi, K.** (2007) Morphologic and facies trends through the fluvial-marine transition in tide-dominated depositional systems: A schematic framework for environmental and sequence-stratigraphic interpretation. *Earth-Science Reviews*, **81**, 135-174.
- Debnath, K., Nikora, V., Aberle, J., Westrich, B. and Muste, M.** (2007) Erosion of cohesive sediments: Resuspension, bed load, and erosion patterns from field experiments. *Journal of Hydraulic Engineering*, **133**, 508–520.
- Decho, A.W. and Gutierrez, T.** (2017) Microbial extracellular polymeric substances (EPSs) in ocean systems. *Frontiers in Microbiology*, **8:922**, DOI: 10.3389/fmicb.2017.00922.
- Dott Jr., R.L.** (1964) Wacke, greywacke and matrix – What approach to immature sandstone classification? *Journal of Sedimentary Petrology*, **34**, 625-632.
- Folk, R. L.** (1951) Stages of textural maturity in sedimentary rocks. *Journal of Sedimentary Petrology*, **21**, 127-130.
- Grabowski, R.C., Droppo, I.G. and Wharton, G.** (2011) Erodibility of cohesive sediment: The importance of sediment properties. *Earth-Science Reviews*, **105**, 101-120.

Hazen, A. (1911) Discussion of 'Dams on sand formations', by A.C. Koenig. *Transactions of the American Society of Civil Engineers*, **73**, 199-203.

Hillel, D. (2004) *Introduction to Environmental Soil Physics*. Elsevier Academic Press, Amsterdam, 494 pp.

Hope, J.A. (2016) The biological mediation of cohesive and non-cohesive sediment dynamics. PhD, University of St Andrews, 427 pp.

Kandiah, A. (1974) Fundamental aspects of surface erosion of cohesive soils. *PhD Thesis, University of California, Davis*, 236 pp.

Le Hir, P., Monbet, Y. and Orvain, F. (2007) Sediment erodability in sediment transport modelling: Can we account for biota effects? *Continental Shelf Research*, **27**, 1116-1142.

Lichtman, I.D., Baas, J.H., Amoudry, L.O., Thorne, P.D., Malarkey, J., Hope, J.A., Peakall, J., Paterson, D.M., Bass, S.J., Cooke, R.D., Manning, A.J., Parsons, D.R. and Ye, L. (2018) Bedform migration in a mixed sand-mud intertidal environment and implications for bed material transport predictions. *Geomorphology*, **315**, 17-32.

Malarkey, J., Baas, J.H., Hope, J.A., Aspden, R.J., Parsons, D.R., Peakall, J., Paterson, D.M., Schindler, R., Ye, L., Lichtman, I.D., Bass, S.J., Davies, A.G., Manning, A.J. and Thorne, P.D. (2015) The pervasive role of biological cohesion in bedform development. *Nature Communications*, 6:6257, doi: 10.1038/ncomms7257.

Manning, A.J., Friend, P.L., Prowse, N. and Amos, C.L. (2007) Estuarine mud flocculation properties determined using an annular mini-flume and the LabSFLOC system. *Continental Shelf Research*, **27**, 1080-1095.

McCarron C.J., Van Landeghem, K.J.J., Baas, J.H., Amoudry, L.O. and Malarkey, J. (2019, in press) The hiding-exposure effect revisited: a method to calculate the mobility of bimodal sediment mixtures. *Marine Geology*, DOI: 10.1016/j.margeo.2018.12.001.

McCrone, A.W. (1962) Clarification of the "winnowing" concept in geology. *GSA Bulletin*, **73**, 517-518.

Mehta, A.J. (2013) *An Introduction To Hydraulics Of Fine Sediment Transport*. World Scientific Publishing Company, Advanced Series on Ocean Engineering, 38, 1060 pp.

Mitchener, H and Torfs, H. (1996) Erosion of mud/sand mixtures. *Coastal Engineering*, **29**, 1-25.

Moore, D.M. and Reynolds Jr., R.C. (1997) *X-Ray Diffraction and the Identification and Analysis of Clay Minerals, 2nd Edition*. Oxford University Press, New York, 400 pp.

- Moore, R.D., Wolf, J. Souza, A.J. and Flint, S.S.** (2009) Morphological evolution of the Dee Estuary, Eastern Irish Sea, UK: A tidal asymmetry approach. *Geomorphology*, **103**, 588-596.
- Oost, A.P. and Baas, J.H.** (1994) The development of small scale bedforms in tidal environments: An empirical model and its applications. *Sedimentology*, **41**, 883-903.
- Parchure, T.M. and Mehta, A.J.** (1985) Erosion of soft cohesive sediment deposits. *Journal of Hydraulic Engineering*, **111**, 1308–1326.
- Packman, A.I. and Brooks, N.H.** (2001) Hyporheic exchange of solutes and colloids with moving bed forms. *Water Resources Research*, **37**, 2591-2605.
- Packman, A.I., Brooks, N.H. and Morgan, J.J.** (2000) A physicochemical model for colloid exchange between a stream and a sand streambed with bed forms. *Water Resources Research*, **36**, 2351-2361.
- Parsons, D.R., Schindler, R.J., Hope, J.A., Malarkey, J., Baas, J.H., Peakall, J., Manning, A.J., Ye, L., Simmons, S., Paterson, D.M., Aspden, R.J., Bass, S.J., Davies, A.G., Lichtman, I.D. and Thorne, P.D.** (2016) The role of bio-physical cohesion on subaqueous bedform size. *Geophysical Research Letters*, **43**, 1566-1573.
- Raudkivi, A.J.** (1997) Ripples on stream bed. *Journal of Hydraulic Engineering*, **123**, 58-64.
- Shields, A.** (1936) Anwendung der Ähnlichkeitsmechanik und der Turbulenzforschung auf die Geschiebebewegung. *Mitteilungen der Preussischen Versuchsanstalt für Wasserbau und Schiffbau*, **26**, 1-26.
- Schindler, R.J., Parsons, D.R., Ye, L., Hope, J.A., Baas, J.H., Peakall, J., Manning, A.J., Aspden, R.J., Malarkey, J., Simmons, S., Paterson, D.M., Lichtman, I.D., Davies, A.G., Thorne, P.D. and Bass, S.J.** (2015) Sticky stuff: Redefining bedform prediction in modern and ancient environments. *Geology*, **43**, 399-402.
- Shepard, F.P.** (1954) Nomenclature based on sand–silt–clay ratios. *Journal of Sedimentary Petrology*, **24**, 151-158.
- Soulsby, R.** (1997) *Dynamics of Marine Sands: A Manual for Practical Applications*. Thomas Telford Ltd., London, 272 pp.
- Soulsby, R.L. and Whitehouse, R.J.S.** (2005) Prediction of Ripple Properties in Shelf Seas; Mark 2 Predictor for Time Evolution. *Report TR 154, Release 2.0, HR Wallingford*, 99 pp.

Spears, B.M., Saunders, J.E., Davidson, I. and Paterson, D.M. (2008) Microalgal sediment biostabilisation along a salinity gradient in the Eden Estuary, Scotland: Unravelling a paradox. *Marine and Freshwater Research*, **59**, 313–321.

Stow, D.A.V. (2005) *Sedimentary Rocks in the Field: A Colour Guide*. Manson, London, 320 pp.

Taylor, I.S. and Paterson, D.M. (1998) Microspatial variation in carbohydrate concentrations with depth in the upper millimetres of intertidal cohesive sediments. *Estuarine Coastal and Shelf Science*, **46**, 359–370.

Tolhurst, T.J., Gust, G. and Paterson, D.M. (2002) The influence of an extracellular polymeric substance (EPS) on cohesive sediment stability. *Proceedings in Marine Science*, **5**, 409–425.

Torfs, H. (1995) Erosion of mud/sand mixtures. *PhD Thesis, Katholieke Univeriteit Leuven, Faculteit der Toegepaste Wetenschappen, Departement Burgelijke Bouwkunde, Laboratorium voor Hydraulica, Belgium*.

Underwood, G.J.C. and Paterson, D.M. (2003) The importance of extracellular carbohydrate production by marine epipelagic diatoms. *Advances in Botanical Research*, **40**, 183–240.

Van de Lageweg, W.I., McLelland, S.J. and Parsons, D.R. (2017) Quantifying biostabilisation effects of biofilm-secreted and synthetic extracellular polymeric substances (EPS) on sand substrate. *Earth Surface Dynamics*, **6**, 203–215.

Van den Berg, J.H. (1987) Bedform migration and bed-load transport in some rivers and tidal environments. *Sedimentology*, **34**, 681–698.

Van den Berg, J.H., Boersma, J.R. and van Gelder, A. (2007) Diagnostic sedimentary structures of the fluvial-tidal transition zone – Evidence from deposits of the Rhine and Meuse. *Netherlands Journal of Geosciences*, **86**, 287–306.

Van Ledden, M. (2001) Modelling of sand-mud mixtures. Part II: A process-based sand-mud model. *Delft Hydraulics, Report Z2840*, 75 pp.

Van Rijn, L.C. (1990) *Principles of Fluid Flow and Surface Waves in Rivers, Estuaries, Seas and Oceans*. Aqua Publications, Amsterdam, 335 pp.

Van Rijn, L.C. (1993) *Principles of Sediment Transport in Rivers, Estuaries and Coastal Seas, Part I: Edition 1993*. Aqua Publications, Amsterdam, 673 pp.

Villaret, C., Huybrechts, N., Davies, A.G. and Way, O. (2011), Effect of bed roughness prediction on morphodynamic modelling: Application to the Dee estuary (UK) and to the Gironde estuary (France). In: *Proceedings of 34th IAHR World Congress, Brisbane, Australia*, p. 1149–1156.

Wang, Y.C., and Sturm, T.W. (2016) Effects of physical properties on erosional and yield strengths of fine-grained sediments. *Journal of Hydraulic Engineering*, **142**, DOI: 10.1061/(ASCE)HY.1943-7900.0001193.

Yong, R.N., Nakano, M., and Pusch, R. (2012) *Environmental Soil Properties and Behavior*. CRC Press, London, 455 pp.

Table 1. Summary of morphological and textural field data.

Bed number	Ripple wavelength		Ripple planform index	Bed mud, mass %		Bed clay, mass %		Bedform type
	Mean (mm)	Number		subsurface	ripple *	subsurface	ripple *	
1	-	-	-	23.4	11.7	8.4	4.2	Flat bed
2	-	-	-	33.8	24.0	12.2	8.6	Flat bed
3	-	-	-	37.1	32.8	13.4	11.8	Flat bed
4	77	15	0.84	33.5	13.7	12.1	4.9	2D ripples
5	78	9	0.75	16.8	13.3	6.0	4.8	2D ripples
6	88	12	-	16.9	14.3	6.1	5.1	2D ripples
7	95	11	0.53	24.7	7.9	8.9	2.8	2D ripples
8	114	9	0.39	7.2	5.3	2.6	1.9	2D-3D ripples
9	116	8	0.41	16.3	7.8	5.9	2.8	2D-3D ripples
10	120	9	0.48	6.7	6.8	2.4	2.4	2D-3D ripples
11	134	7	0.41	10.9	6.5	3.9	2.3	2D-3D ripples
12	123	7	0.31	14.2	7.5	5.1	2.7	3D ripples
13	129	8	0.24	5.3	3.8	1.9	1.4	3D ripples
14	141	7	0.34	12.7	1.6	4.6	0.6	3D ripples
15	143	7	0.38	4.6	5.2	1.7	1.9	3D ripples

* Surface scrape for flat beds

APPENDIX A – CALCULATION OF RIPPLE PLANFORM INDICES

The visual descriptions of 2D, 2D-3D, and 3D ripples in the field and in the experiments of Baas *et al.* (2013) were supported quantitatively by calculating the planform index, I , of the ripples:

$$I = L_{cr}/w \quad (A1)$$

where L_{cr} is the shortest distance between the start and end of the crestline of a ripple and w is a reference width, both perpendicular to the flow direction. Here, $w = 0.3$ m, so that a direct comparison could be made between the data from the Dee Estuary and from Baas *et al.* (2013), who used a flume with a width of 0.3 m. The methodology is summarised in Fig. A1. Planform index values for the field and laboratory ripples are shown in Table 1 and Table A1, respectively. These data show that $I > 0.50$ for 2D ripples, $0.39 \leq I \leq 0.5$ for 2D-3D ripples and $I < 0.39$ for 3D ripples.

Table A1. Summary of morphological and textural data for selected experiments of Baas *et al.* (2013), including ripple planform indices.

Run number	Ripple length mean (mm)	Ripple planform index	Bed clay %		Bedform type
			initial	ripple	
08	77.7	0.57	13.8	1.2	2D ripples
06	86.7	0.44	11.8	0	2D-3D ripples
04	98.4	0.33	5.4	0.3	3D ripples
02	95.5	0.31	1.8	0	3D ripples
01	116.3	0.28	0	0	3D ripples

APPENDIX B - TIME FOR COMPLETE TRAPPING OF A COLLOID

Following the approach of Malarkey *et al.* (2015), the time for complete trapping of a colloid, such as clay or EPS, as a result of Darcy's Law when ripples are stationary, can be estimated from figure 5a of Packman *et al.* (2000). The threshold of motion according to their equations 6-9 for an effective mean water depth of 1.73 m (figure 5a) is $U = 0.361$ m s⁻¹. This value of U is approximately equivalent to 0.2 m s⁻¹ for Packman *et al.*'s (2000) water depth and ripple size. According to their figure 5a for $U = 0.2$ m s⁻¹, the time taken for the colloid concentration in the bed to reach 90% of its value in flow, t_{90} , requires that $ku_m t_{90}/P = 32.5$. Here k is ripple wave number ($= 2\pi/L$), L is the ripple wavelength, u_m is the maximum induced pore water velocity ($= kKh_m$), P is the porosity ($= 0.4$), K is the hydraulic conductivity (Hazen, 1911) and h_m is the half amplitude dynamic head (Packman *et al.*, 2000), given by:

$$K \sim 10D_{10}^2, \quad h_m = 0.14 \frac{U^2}{g} \left(\frac{H}{0.34d} \right)^{3/8}, \quad (B1a,b)$$

where K is in mm s^{-1} , D_{10} is in mm and d is the water depth. In this case, $D_{10} = 0.149$ mm, so that $K \approx 0.22 \text{ mm s}^{-1}$, $d = 1.73$ m, $U = 0.2 \text{ m s}^{-1}$, $H \approx 0.1L$ and L is in the range $77 \leq L \leq 143$ mm. For these values, $19.7 \leq t_{90} \leq 61.7$ h (0.8 to 2.6 days) or a minimum of four times the inundation period (4.7 h). The corresponding maximum pore water velocities are $0.08 \leq u_m \leq 0.13 \text{ mm min}^{-1}$, which is two orders of magnitude smaller than the 6 mm min^{-1} bedform migration rate measured in the same area on the 26th May (Lichtman *et al.*, 2018). This demonstrates that winnowing via overturning dominated over pumping for this inundation period (Packman & Brooks, 2001). The timescale for the turn-over of the ripples based on the migration rate and the range of ripple wavelengths is 0.2 to 0.4 h.

APPENDIX C – DESCRIPTION OF CRITICAL SHEAR STRESS EXPERIMENTS

The ‘vertical shear flume’ (Hydrodynamics Laboratory, Bangor University; McCarron *et al.*, 2019) was used to estimate the critical shear stress for sediment motion from beds comprising mixed sand–clay–EPS (Fig. 7). This flume consists of a straight, rectangular, channel, 1.2 m long and 0.18 m wide, with a water depth of 0.12 m, through which fresh water was recirculated by a variable-speed pump. A block-shaped sediment box was slotted onto the bottom of the channel, ensuring that the surface level of the sediment in the box was flush with the floor of the channel. The sediment column within the box was 0.14 m long, 0.1 m wide, and 0.1 m deep. Downstream of this box, a bedload trap collected sediment particles moving along the bottom of the flume in flows that exceeded the critical shear stress for sediment motion.

The experiments were conducted with well-sorted, fine sand ($D_{50} = 0.142$ mm) mixed with 0–1% xanthan gum (EPS proxy), 0–30% kaolin clay, or a mixture of up to 30% kaolin and up to 1% xanthan gum. In each experiment, steady, uniform flow was started well below the critical shear stress for the pure sand and then increased in small steps every 10 minutes. Streamwise flow velocities were measured with a vertical stack of five ultrasonic Doppler velocimetry (UVP) probes (Best *et al.*, 2001). Bed shear stresses were calculated from the UVP data by squaring the friction velocity determined from standard logarithmic law of the wall techniques (*e.g.*, van Rijn, 1990). Water samples were taken periodically to determine suspended sediment concentrations, using the standard weighing and drying method, such that an optical backscatter (OBS) instrument, which was continuously recording, could be calibrated.

For each sand–clay–EPS mixture, the critical bed shear stress was determined by a combination of three methods: (1) visual observation of the erosion of sand grains from the sediment surface; (2) initiation of infilling of the bedload trap; and (3) the first significant increase in the concentration of

suspended sediment. All critical bed shear stresses, relative to the critical bed shear stress for the pure sand, are shown in Fig. 7.

FIGURE CAPTIONS

Table 1. Summary of morphological and textural field data.

Table A1. Summary of morphological and textural data for the experiments of Baas et al. (2013), including ripple planform indices.

Fig. 1. (A) Schematic map of the Dee Estuary around Hilbre Island, with the main tidal channel in white and the study area located on the grey-coloured intertidal flat to the northwest of Little Eye. The four islands are defined by the area above the mean high water mark and by any area of bedrock exposed at low water immediately below this mark. (B) Overview of study site. The ripples in the foreground have a wavelength of approximately 150 mm.

Fig. 2. Mean ripple wavelength against bed mud content for the current ripples at the field site. The black continuous and dashed lines are the linear least-square best fits for the subsurface and the ripple samples, respectively. The colours indicate incipient ripples (yellow), 2D sinuous and straight-crested ripples (red), 2D-3D linguoid ripples (blue), and 3D linguoid ripples (green). The boundaries between these ripple types are from Baas (1999). The horizontal stippled lines denote the initial and equilibrium wavelengths of current ripples in 0.238 mm sand (*cf.*, Baas, 1999).

Fig. 3. Field examples of flat beds (A,B), 2D ripples (C,D), 2D-3D ripples (E,F), and 3D ripples (G,H). Bed mud content decreases from (A,B) to (G,H). The percentages in (A-H) denote subsurface mud contents, as shown in Table 1. The scale bar is 200 mm long.

Fig. 4. Final clay content (A) and dimensionless current ripple wavelength (B) against initial bed clay content for the ripples at the field site, with $D_{50} = 0.228$ mm, based on the subsurface samples (solid circles), and in the experiments of Baas *et al.* (2013) with $D_{50} = 0.143$ mm (open circles). In A, the black continuous lines represent different winnowing efficiencies, the dashed line corresponds to 2.8% final clay content, which is the upper limit of occurrence for 2D-3D and 3D ripples for the field data and the laboratory data, and the coloured numbers are ripple planform indices (Appendix A), for the closest experimental case for which plan-view images were available. In (B), the black continuous and dashed lines are the linear least-square best fits for the field data and the experimental data, respectively. The colours indicate the bed configurations, as in (A).

Fig. 5. Summary of the field data, collected on May 26th, 2013. (A) water depth, (B) depth-averaged flow velocity, (C) grain-related mobility parameter, (D) cumulative ripple wavelength development stage, (E) predicted temporal development of ripple wavelength. The dashed lines in (B) and (C) denote linear extrapolations. $\theta_c = 0.0508$ is critical grain-related mobility parameter; $L_E = 142$ mm is

equilibrium ripple wavelength; $L_0 = 60$ mm is initial ripple wavelength. S_L is cumulative ripple wavelength development stage.

Fig. 6. Dimensionless current ripple wavelength against initial bed clay content for the ripples in the field. All lines relate to the experimental data. ‘Laboratory’ is the least-square best fit from Fig. 4b; ‘Equation 10’ is the adjustment of the ripple development stage to that of the field data, $S_L = 1.26$. ‘Equation 11’ also includes the initiation time for ripples, based on the mixed sand-EPS experiments of Malarkey et al. (2015). The symbols and their colours represent the field data, as in Fig. 4.

Fig. 7. Relative change in critical Shields parameter (including standard error) for sand with kaolin clay, sand with EPS-proxy xanthan gum, and sand with mixtures of kaolin and xanthan gum. The term $\theta_{c,0}$ on the vertical axis denotes the critical Shields parameter for pure sand with a median grain diameter of 0.148 mm. The blue coloured surface denotes the laboratory data. The red coloured insert denotes the approximate range of the field data.

Fig. A1. Summary of the methodology used to calculate ripple planform indices. A 0.3-m wide window was placed in the centre of the picture (in green), before tracing the ripple crestlines within this window (in red). Thereafter, the shortest distance, L_{cr} , between the start and end of each crestline was measured (in blue) and divided by the width of the window to derive the ripple planform index as measure of ripple three-dimensionality. Finally, the average value of I was calculated for each bed.

Table 1. Summary of morphological and textural field data.

Bed number	Ripple length		Ripple planform		Bed mud %		Bed clay %		Bedform type
	mean (mm)	number	index		subsurface	ripple *	subsurface	ripple *	
1	-	-	-		23.4	11.7	8.4	4.2	Flat bed
2	-	-	-		33.8	24.0	12.2	8.6	Flat bed
3	-	-	-		37.1	32.8	13.4	11.8	Flat bed
4	77	15	0.84		33.5	13.7	12.1	4.9	2D ripples
5	78	9	0.75		16.8	13.3	6.0	4.8	2D ripples
6	88	12	-		16.9	14.3	6.1	5.1	2D ripples
7	95	11	0.53		24.7	7.9	8.9	2.8	2D ripples
8	114	9	0.39		7.2	5.3	2.6	1.9	2D-3D ripples
9	116	8	0.41		16.3	7.8	5.9	2.8	2D-3D ripples
10	120	9	0.48		6.7	6.8	2.4	2.4	2D-3D ripples
11	134	7	0.41		10.9	6.5	3.9	2.3	2D-3D ripples
12	123	7	0.31		14.2	7.5	5.1	2.7	3D ripples
13	129	8	0.24		5.3	3.8	1.9	1.4	3D ripples
14	141	7	0.34		12.7	1.6	4.6	0.6	3D ripples
15	143	7	0.38		4.6	5.2	1.7	1.9	3D ripples

* Surface scrape for flat beds




Fig. 1. (A) Schematic map of the Dee Estuary around Hilbre Island, with the main tidal channel in white and the study area located on the grey-coloured intertidal flat to the northwest of Little Eye. The four islands are defined by the area above the mean high water mark and by any area of bedrock exposed at low water immediately below this mark. (B) Overview of study site. The ripples in the foreground have a wavelength of approximately 150 mm.

183x262mm (300 x 300 DPI)




Fig. 3. Field examples of flat beds (A,B), 2D ripples (C,D), 2D-3D ripples (E,F), and 3D ripples (G,H). Bed mud content decreases from (A,B) to (G,H). The percentages in (A-H) denote subsurface mud contents, as shown in Table 1. The scale bar is 200 mm long.

204x130mm (150 x 150 DPI)




Fig. 4. Final clay content (A) and dimensionless current ripple wavelength (B) against initial bed clay content for the ripples at the field site, with $D_{50} = 0.228$ mm, based on the subsurface samples (solid circles), and in the experiments of Baas et al. (2013) with $D_{50} = 0.143$ mm (open circles). In A, the black continuous lines represent different winnowing efficiencies, the dashed line corresponds to 2.8% final clay content, which is the upper limit of occurrence for 2D-3D and 3D ripples for the field data and the laboratory data, and the coloured numbers are ripple planform indices (Appendix A), for the closest experimental case for which plan-view images were available. In (B), the black continuous and dashed lines are the linear least-square best fits for the field data and the experimental data, respectively. The colours indicate the bed configurations, as in (A).

190x231mm (150 x 150 DPI)




Fig. 4. Final clay content (A) and dimensionless current ripple wavelength (B) against initial bed clay content for the ripples at the field site, with $D_{50} = 0.228$ mm, based on the subsurface samples (solid circles), and in the experiments of Baas et al. (2013) with $D_{50} = 0.143$ mm (open circles). In A, the black continuous lines represent different winnowing efficiencies, the dashed line corresponds to 2.8% final clay content, which is the upper limit of occurrence for 2D-3D and 3D ripples for the field data and the laboratory data, and the coloured numbers are ripple planform indices (Appendix A), for the closest experimental case for which plan-view images were available. In (B), the black continuous and dashed lines are the linear least-square best fits for the field data and the experimental data, respectively. The colours indicate the bed configurations, as in (A).

179x165mm (600 x 600 DPI)




Fig. 5. Summary of the field data, collected on May 26th, 2013. (A) water depth, (B) depth-averaged flow velocity, (C) grain-related mobility parameter, (D) cumulative ripple wavelength development stage, (E) predicted temporal development of ripple wavelength. The dashed lines in (B) and (C) denote linear extrapolations. $\theta_c = 0.0508$ is critical grain-related mobility parameter; $L_E = 142$ mm is equilibrium ripple wavelength; $L_0 = 60$ mm is initial ripple wavelength. S_L is cumulative ripple wavelength development stage.

180x273mm (150 x 150 DPI)




Fig. 6. Dimensionless current ripple wavelength against initial bed clay content for the ripples in the field. All lines relate to the experimental data. 'Laboratory' is the least-square best fit from Fig. 4b; 'Equation 10' is the adjustment of the ripple development stage to that of the field data, $SL = 1.26$. 'Equation 11' also includes the initiation time for ripples, based on the mixed sand-EPS experiments of Malarkey et al. (2015). The symbols and their colours represent the field data, as in Fig. 4.

170x89mm (600 x 600 DPI)




Fig. 7. Relative change in critical Shields parameter (including standard error) for sand with kaolin clay, sand with EPS-proxy xanthan gum, and sand with mixtures of kaolin and xanthan gum. The term $\theta_{c,0}$ on the vertical axis denotes the critical Shields parameter for pure sand with a median grain diameter of 0.148 mm. The blue coloured surface denotes the laboratory data. The red coloured insert denotes the approximate range of the field data.

188x128mm (150 x 150 DPI)

Table A1. Summary of morphological and textural data for selected experiments of Baas *et al.* (2013), including ripple planform indices.

Run number	Ripple length mean (mm)	Ripple planform index	Bed clay %		Bedform type
			initial	ripple	
08	77.7	0.57	13.8	1.2	2D ripples
06	86.7	0.44	11.8	0	2D-3D ripples
04	98.4	0.33	5.4	0.3	3D ripples
02	95.5	0.31	1.8	0	3D ripples
01	116.3	0.28	0	0	3D ripples

Bed 7: 2D ripples, 24.7% mud




Fig. A1. Summary of the methodology used to calculate ripple planform indices. A 0.3-m wide window was placed in the centre of the picture (in green), before tracing the ripple crestlines within this window (in red). Thereafter, the shortest distance, L_{cr} , between the start and end of each crestline was measured (in blue) and divided by the width of the window to derive the ripple planform index as measure of ripple three-dimensionality. Finally, the average value of I was calculated for each bed.

97x66mm (300 x 300 DPI)