

Upward resetting of the vascular sympathetic baroreflex in middle-aged male runners

Wakeham, Denis; Lord, Rachel; Talbot, Jack ; Lodge, Freya; Dawkins, Tony; Curry, Bryony ; Simpson, Lydia; Pugh, Christopher ; Shave, Rob; Moore, Jonathan

American Journal of Physiology - Heart and Circulatory Physiology

DOI:
[10.1152/ajpheart.00106.2019](https://doi.org/10.1152/ajpheart.00106.2019)

Published: 01/07/2019

Peer reviewed version

[Cyswllt i'r cyhoeddiad / Link to publication](#)

Dyfyniad o'r fersiwn a gyhoeddwyd / Citation for published version (APA):
Wakeham, D., Lord, R., Talbot, J., Lodge, F., Dawkins, T., Curry, B., Simpson, L., Pugh, C., Shave, R., & Moore, J. (2019). Upward resetting of the vascular sympathetic baroreflex in middle-aged male runners. *American Journal of Physiology - Heart and Circulatory Physiology*, 317(1), H181-H189. <https://doi.org/10.1152/ajpheart.00106.2019>

Hawliau Cyffredinol / General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

1 **Upward resetting of the vascular sympathetic baroreflex in middle-aged male runners**

2 Denis J Wakeham¹, Rachel N Lord¹, Jack S Talbot¹, Freya M Lodge², Bryony A Curry¹, Tony
3 G Dawkins¹, Lydia L Simpson³, Rob E Shave^{1,4}, Christopher JA Pugh^{*1}, Jonathan P Moore^{*3}

4 *CJAP and JPM share senior authorship

5 ¹Cardiff Centre for Exercise and Health, School of Sport and Health Sciences, Cardiff
6 Metropolitan University, United Kingdom;

7 ²Cardiff and Vale University Health Board, University Hospital of Wales, Cardiff, United
8 Kingdom;

9 ³Physical Activity for health and Well-being (PAWB) Centre, School of Sport, Health and
10 Exercise Sciences, Bangor University, United Kingdom;

11 ⁴Centre for Heart, Lung, and Vascular Health, University of British Columbia Okanagan,
12 Kelowna, Canada

13 **Corresponding Author:**

14 Dr Jonathan P Moore, Physical Activity for health and Well-being (PAWB) Centre, School of
15 Sport, Health and Exercise Sciences, Bangor University, LL57 2PZ, United Kingdom.
16 j.p.moore@bangor.ac.uk Tel: + 44 1248 383645

17 **Running head:** Vascular sympathetic baroreflex in middle-aged runners

18 **Subject terms:** Ageing, baroreflex, blood pressure, exercise physiology, sympathetic
19 nervous system

20

21 **ABSTRACT**

22 This study focussed on the influence of habitual endurance exercise training (i.e. committed
23 runner or non-runner) on the regulation of muscle sympathetic nerve activity (MSNA) and
24 arterial pressure in middle-aged (50 to 63 years, n= 23) and younger (19 to 30 years; n=23)
25 normotensive men. Haemodynamic and neurophysiological assessments were performed at
26 rest. Indices of vascular sympathetic baroreflex function were determined from the
27 relationship between spontaneous changes in diastolic blood pressure (DBP) and MSNA.
28 Large vessel arterial stiffness and left ventricular stroke volume also were measured. Paired
29 comparisons were performed within each age-category. Mean arterial pressure and basal
30 MSNA bursts·min⁻¹ were not different between age-matched runners and non-runners.
31 However, MSNA bursts·100 heartbeats⁻¹, an index of baroreflex regulation of MSNA
32 (vascular sympathetic baroreflex operating point) was higher for middle-aged runners
33 ($P=0.006$), whereas this was not different between young runners and non-runners. The
34 slope of the DBP-MSNA relationship (vascular sympathetic baroreflex gain) was not different
35 between groups in either age-category. Aortic pulse wave velocity was lower for runners of
36 both age-categories ($P<0.03$), although carotid β stiffness was lower only for middle-aged
37 runners ($P=0.04$). For runners of both age-categories, stroke volume was larger, while heart
38 rate was lower (both $P<0.01$). In conclusion, we suggest that neural remodelling and upward
39 setting of the vascular sympathetic baroreflex compensates for cardiovascular adaptations
40 after many years committed to endurance exercise training, presumably to maintain arterial
41 blood pressure stability.

42 **NEW AND NOTEWORTHY**

43 Exercise training reduces muscle sympathetic burst activity in disease; this is often
44 extrapolated to infer a similar effect in health. We demonstrate that burst frequency of
45 middle-aged and younger men committed to endurance training is not different compared
46 with age-matched casual exercisers. Notably, well-trained middle-aged runners display
47 similar arterial pressure but higher sympathetic burst occurrence than untrained peers. We
48 suggest homeostatic plasticity and upward setting of the vascular sympathetic baroreflex
49 maintains arterial pressure stability following years of training.

50 INTRODUCTION

51 Human ageing exerts a marked influence on blood pressure, which is the primary regulated
52 variable of the cardiovascular system. Two hallmarks of cardiovascular aging are large-
53 vessel arterial stiffening (30), and chronic elevation of muscle sympathetic nerve activity,
54 (MSNA) (27). The conventional wisdom is that these factors, amongst others, contribute to
55 the age-related increase in arterial blood pressure observed in western society beyond 50
56 years of age (13).

57 Arterial baroreflex control of MSNA (i.e. vascular sympathetic baroreflex) is the
58 primary mechanism through which the autonomic nervous system regulates vasomotor tone,
59 and thus plays a pivotal role in blood pressure homeostasis. The age-related increase in
60 MSNA is underpinned by resetting of the vascular sympathetic baroreflex (31), whereby the
61 'operating point' (i.e. mean resting diastolic blood pressure [DBP] and corresponding MSNA
62 bursts per 100 heartbeats, a measure of the probability of a burst occurrence) resets upward
63 and rightward. Vascular sympathetic baroreflex 'resetting' with age occurs in the absence of
64 a change of reflex 'gain' (i.e. responsiveness to acute changes in blood pressure) (11, 25,
65 26). Notably, however, it appears that a rise in arterial pressure does not necessarily follow
66 progressive elevation in resting vascular sympathetic activity with advancing age (49). In
67 contrast, baroreflex-mediated cardiac parasympathetic control (i.e. cardiovagal baroreflex
68 gain) is progressively impaired with advancing age (11, 32). Alterations to mechanosensory
69 transduction and neural control (44) may explain these changes to the vascular sympathetic
70 and cardiovagal limbs of the arterial baroreflex with human ageing.

71 Long-term aerobic exercise training mitigates against some of the hallmarks of
72 cardiovascular aging. For example, lifelong endurance exercise training offsets age-related
73 stiffening of the aorta (51) and carotid artery (47). However, the interaction of committed
74 exercise training and age-related changes to vascular sympathetic activity is unclear. To
75 date, relatively little consensus exists among previous microneurographic studies, which
76 have found basal MSNA burst frequency for middle-aged and older endurance-trained men

77 is either higher (37), not different (38) or lower (44), compared to untrained peers.
78 Furthermore, quantification of the number of burst occurrences relative to the number of
79 opportunities for a burst (i.e. burst incidence) does not provide clarity. However, the method
80 of burst quantification provides different neurophysiological insight into regulation of vascular
81 sympathetic activity (5). Burst frequency is reflective of the amount of sympathetic activity
82 (or neurotransmitter release) that the vasculature is exposed to in a given time period (53).
83 In contrast, burst incidence indicates the probability of a sympathetic burst occurring at a
84 given arterial pressure (29). Furthermore, baroreceptor signals over a wide pressure range
85 influence both the timing and the probability of sympathetic bursts. We contend, therefore,
86 that burst incidence is an index of the baroreflex 'gating' sympathetic bursts (20, 21), rather
87 than sympathetic outflow *per se*. In the only study to consider the influence of aging and
88 chronic exercise training on vascular sympathetic baroreflex control, the training status of
89 healthy older males had no effect on MSNA burst incidence (vascular sympathetic baroreflex
90 operating point), or the MSNA responsiveness (gain) to a modified Oxford baroreceptor test
91 (44). In contrast, cross-sectional evidence from middle-aged and older men indicate that
92 vigorous long-term endurance training attenuates the ageing-related decline in cardiovagal
93 baroreflex responsiveness (34).

94 Taking the various aforementioned uncertainties into account, the primary aim of this
95 cross-sectional study was to investigate the effect that habitual endurance exercise training
96 has on regulation of vascular sympathetic burst activity and resting blood pressure in healthy
97 middle age. Because of marked sex differences in sympathetic regulation (18) and
98 autonomic support of blood pressure (6), men only were studied to experimentally isolate the
99 influence of long-term endurance training as much as possible. Furthermore, in order to
100 examine the effect of exercise training independently of ageing, a secondary aim was to
101 compare the sympathetic control of blood pressure between young runners and young non-
102 runners. To address these aims, we performed comprehensive haemodynamic and
103 neurophysiological assessment, and measured central artery stiffness and left ventricular

104 stroke volume, in four groups of healthy normotensive men: middle-aged committed runners,
105 middle-aged non-runners, younger runners and younger non-runners. Based upon limited
106 data, we hypothesised that the vascular sympathetic baroreflex control would not be different
107 between well-trained runners and non-runners.

108 **METHODS**

109 **Ethical Approval**

110 This study conformed to the most recent Declaration of Helsinki, except for registration in a
111 database. The Research Ethics Committee at the Cardiff School of Sport and Health
112 approved all study procedures (16/7/02R) and participants provided written informed consent
113 prior to entering the study.

114 **Participants**

115 Between August 2016 and August 2017, the eligibility to participate was assessed for
116 seventy men. Forty-six participants completed the study. Each participant was categorised
117 according to his age (i.e. middle-aged or young) and training status (i.e. committed runner or
118 non-runner) (Table 1). Among middle-aged men, runners performed ≥ 25 miles of moderate
119 to intense training per week for ≥ 10 years ($n=13$), whereas non-runners were casually
120 recreationally active i.e. ≤ 3 hours of structured physical activity per week for ≥ 10 years
121 ($n=10$). In the case of the young men, runners performed ≥ 50 miles of training per week
122 ($n=13$) and non-runners performed ≤ 3 hours of structured physical activity per week, for ≥ 2
123 years ($n=10$). All participants were free of known cardiovascular, metabolic or other chronic
124 diseases, normotensive ($<140/90$ mmHg when supine), non-smokers, and non-obese ($BMI <$
125 $30 \text{ kg}\cdot\text{m}^2$) as assessed by a medical history, manual sphygmomanometry (Welch Allyn, UK)
126 and measurement of height and body mass. Middle-aged men were further evaluated by
127 resting and maximal exercise electrocardiogram.

128 **Experimental overview**

129 Participants completed one screening visit and 2 days of physiological testing, with a
130 minimum of one week between the tests. All screening and physiological tests were
131 performed at the Cardiff School of Sport and Health Sciences in a quiet, temperature
132 controlled (22-24°C) environment. We requested that participants abstain from caffeine,
133 alcohol and strenuous exercise for twenty-four hours prior to arrival at the laboratory on each
134 visit; none took medication at the time of testing. On one testing day, assessment of body
135 composition (Bioelectrical impedance analysis; Bodystat 1500, Bodystat Ltd, Douglas, Isle of
136 Man) and measurement of arterial stiffness were followed by a maximal incremental exercise
137 test. On the other testing day, having fasted for six hours, participants underwent
138 cardiovascular and sympathetic neural assessments.

139 **Assessment of arterial stiffness**

140 Sequential ECG-gated arterial pressure waveforms were recorded in accordance with
141 current guidelines (50) from the carotid and femoral arteries, at the site of maximal arterial
142 pulsation, enabling the calculation of aortic pulse wave velocity (aPWV; SphygmoCor,
143 Cardie X Ltd, Australia). Furthermore, the β stiffness index of the right common carotid
144 artery was determined via high-resolution ultrasonography, using a 12-MHz linear array
145 transducer (Vivid Q, GE Medical, Norway), as previously described (19). Central blood
146 pressure was estimated to calculate β stiffness index, by applying a generalized transfer
147 function (41) to radial arterial waveforms, collected via a high fidelity micromanometer tipped
148 probe (SphygmoCor, Cardie X Ltd, Australia). Carotid artery β stiffness index is reported in
149 44 individuals (9 young non-runners, 12 young runners, 10 middle-aged non-runners, 13
150 middle-aged runners).

151 **Cardiopulmonary exercise test**

152 All participants completed an incremental exercise test to exhaustion on a cycle ergometer
153 (Lode Corival, Groningen, The Netherlands) to assess $\dot{V}O_2$ peak. Cycling was chosen for

154 reasons of safety and assessment of the exercise electrocardiogram. Each increment
155 corresponded to an increase 20 watts per minute (Middle-aged runners started at 90W and
156 young runners started at 120W; middle-aged and young non-runners started at 30W and
157 50W, respectively). During the maximal exercise test oxygen consumption was measured
158 continuously via a breath-by-breath analyser (Oxycon Pro, Jaeger, Hoechberg, Germany).
159 Heart rate was measured throughout the exercise test via either a chest strap in the young
160 groups (Polar Electro, RS400, Finland) or 12-lead electrocardiography in middle-aged men
161 (Oxycon Pro, Jaeger, Hoechberg, Germany).

162 **Hemodynamics and sympathetic neural activity**

163 Heart rate and blood pressure were monitored continuously via three-lead
164 electrocardiography and finger photoplethysmography (FinometerPro, FMS, Groningen,
165 Netherlands), with participants supine. The arterial pressure waveform was calibrated at
166 regular intervals to the average resting systolic and diastolic pressures measured via manual
167 sphygmomanometry. Echocardiograms were acquired using a commercially available
168 ultrasound system (Vivid E9, GE Medical, Norway) with a 1.5 to 4 MHz array probe. Images
169 were obtained from apical 4 and 2 chamber views by a single experienced sonographer
170 (RNL) and saved for offline analysis with commercially available software (EchoPAC, BT12,
171 GE Medical, Norway).

172 Multiunit muscle sympathetic nerve activity was obtained by microneurography using
173 a recording system (Nerve Traffic Analyser, Model 663 C, University of Iowa, Iowa City, IA)
174 and following a recognized technique (45). In brief, a unipolar tungsten microelectrode (FHC,
175 Bowdoin, ME), with shaft diameter of 0.1 mm (impedance 1-5 MW), was placed across the
176 skin at the popliteal fossa and inserted into the peroneal nerve by an experienced
177 microneurographer (JPM). A reference electrode was placed subcutaneously approximately
178 2-3 cm above from the site of the recording electrode. The recorded neurogram was
179 amplified (70, 000 to 160, 000 fold), band-pass filtered (700 to 2000 Hz), full-wave rectified
180 and integrated with a resistance-capacitance circuit (time constant 0.1 sec). Satisfactory

181 recordings of MSNA were identified, dependent on the following criteria (54), (i) pulse-
182 synchronous “bursts” of activity, (ii) increased “burst” occurrence in response to voluntary
183 apnoea, (iii) unaffected “burst” pattern during stroking of the skin, and (iv) 3:1 signal to noise
184 ratio. At least 10 minutes after an acceptable MSNA recording site was found,
185 echocardiograms and other baseline data were acquired. Hemodynamic and neural data
186 were sampled at 1000Hz using a commercial data acquisition system and stored for offline
187 data analysis (Chart Version 8, Lab Chart Pro, AD Instruments, UK).

188 **Assessment of arterial baroreflex function**

189 Hemodynamic and neural recordings were acquired for six minutes in order to characterize
190 the arterial baroreflex regulation of MSNA and interbeat RR interval. Respiratory rate was
191 monitored via a nasal cannula (Capnocheck® Sleep Capnograph, Smiths Medical, UK), to
192 ensure that the participants had a regular breathing pattern, due to the influence of breath-
193 hold on MSNA (9). Examples of the dynamic relationship between beat-by beat arterial
194 pressure and bursts of MSNA are shown in Figure 1.

195 **Data Analyses**

196 Stroke volume was estimated using the Simpson’s-biplane method (24), thus permitting
197 determination of cardiac output (\dot{Q} ; heart rate x stroke volume) and the total peripheral
198 resistance (TPR; \dot{Q} /mean arterial pressure). Satisfactory images for the quantification of
199 stroke volume were not recorded in one individual (one middle-aged runner); accordingly,
200 stroke volume, \dot{Q} and TPR data are reported for forty-five individuals.

201 Multi-unit bursts of MSNA were verified by two investigators (DJW/JPM) via visual
202 inspection following adjustment for baroreflex latency (54) (time between R wave and peak
203 burst height), which aligned each burst with the appropriate R wave of the ECG. MSNA was
204 quantified as burst frequency (bursts per minute [bursts·min⁻¹]) and burst incidence (bursts
205 per 100 heartbeats [bursts·100hb⁻¹]).

206 The slope of the stimulus-response relationship between DBP and MSNA burst
207 probability was calculated to represent vascular sympathetic baroreflex gain (21, 45). Briefly,
208 DBP was averaged into two mmHg bins, to minimize the influence of respiration on MSNA
209 and to maximize the number of data points for inclusion in the linear regression model. The
210 percentage of cardiac cycles associated with a burst of MSNA (ranging from zero to 100%),
211 per bin of DBP, was used to calculate burst probability. Data were included for further
212 analysis if, (i) at least five data points for each linear regression were available and (ii) a
213 correlation coefficient of ≥ -0.5 was present (14). Mean values and tests of statistical
214 significance are presented for 20 middle-aged (11 runners) and 20 younger men (11
215 runners). Statistical weighting was adopted for this analyses to minimize the influence of
216 differences in the number of cardiac cycles within each DBP bin (21). The operating point of
217 the vascular sympathetic baroreflex was determined from mean diastolic pressure and
218 corresponding average burst incidence.

219 Cardiovagal baroreflex gain was assessed by the sequence method using
220 customized computer software (Cardioseries version 2.4, Ribeirao Preto, São Paulo, Brazil).
221 If R-R interval was ≥ 800 milliseconds a delay of 1 beat was applied so that the SBP was
222 regressed against the following R-R interval (12). Data were included for further analysis
223 upon condition of (i) a minimum of three data points for a linear regression were available
224 and (ii) a correlation coefficient of ≥ 0.8 was present (40). The operating point of the
225 cardiovagal baroreflex was determined from mean prevailing SBP and corresponding
226 average RR interval. Data, including positive and negative ramp gains, and the number of
227 sequences, are presented for 20 middle-aged (11 runners) and 21 younger men (11
228 runners).

229 **Statistical Analyses**

230 In line with our primary (i.e. middle-aged runner *versus* age matched non-runner) and
231 secondary (i.e. younger runner *versus* age matched non-runner) aims, and after checking
232 compliance with basic parametric assumptions, we assessed between-group differences for

233 middle-aged runners and non-runners, and for young runners and non-runners, via
234 independent t-tests. Alpha was set a-priori as $P<0.05$. All statistical analyses were
235 completed using Statistics Package for Social Sciences for Windows, (Version 23, Chicago,
236 IL) and data are reported as mean (95% Confidence Intervals).

237 **RESULTS**

238 **Participant demographics**

239 By design, training and cardiorespiratory fitness ($\dot{V}O_{2peak}$) were greater for runners compared
240 to age-matched non-runners (middle-aged and young, $P<0.001$; Table 1). Runners had
241 lower body mass (middle-aged, $P=0.001$; young, $P=0.003$) and body mass index (middle-
242 aged and young, $P<0.001$), and less body fat percentage (middle-aged and young,
243 $P<0.001$), than age-matched non-runners. Systolic BP ($P=0.041$) and Diastolic BP ($P=0.027$)
244 were lower for young runners compared to age-matched non-runners. Screening blood
245 pressures were not different among middle-aged runners and untrained peers.

246 **Resting hemodynamics and vascular sympathetic neural activity**

247 Stroke volume was higher (middle-aged, $P=0.03$; young, $P<0.01$) and heart rate was lower
248 (middle-aged, $P<0.001$; young, $P<0.001$) between both groups of runners compared to age-
249 matched non-runners (Table 2). There were no other differences in resting haemodynamic
250 parameters between runners and non-runners for either age-category. Resting MSNA burst
251 frequencies were not different among middle-aged runners and non-runners, or among
252 young runners and age-matched non-runners. Burst incidence data is considered in the
253 following section.

254 **Arterial baroreflex function**

255 Among middle-aged men, there was no difference between runners and non-runners for the
256 diastolic operating pressure of the vascular sympathetic baroreflex ($P=0.57$); however, the
257 corresponding operating MSNA (i.e. bursts $\cdot 100hb^{-1}$) was higher in the runners ($P<0.01$;

258 Figure 2A). Among young men, there was no significant difference in vascular sympathetic
259 operating point between runners and non-runners (DBP, $P=0.23$; corresponding MSNA
260 bursts $\cdot 100\text{hb}^{-1}$, $P=0.24$). The vascular sympathetic baroreflex gain (i.e. slope of the DBP-
261 MSNA relationship) was not influenced by the training status of either middle-aged (-6.07 [-
262 8.80 to -3.55] vs -7.30 [-10.49 to -4.12] $\% \cdot \text{mHg}^{-1}$, $P=0.55$) or younger men (-6.68 [-13.1 to -
263 2.33] vs. -5.82 [-7.15 to -4.49] $\% \cdot \text{mHg}^{-1}$, $P=0.58$).

264 Among middle-aged runners and non-runners, there was no difference in the
265 prevailing systolic pressure for the cardiovagal baroreflex ($P=0.58$), but the corresponding
266 RR interval was higher for runners ($P<0.01$; Figure 2B). Among young men, the prevailing
267 systolic pressure was lower ($P=0.02$) and the corresponding RR interval was higher for
268 runners ($P<0.01$). The cardiovagal baroreflex gain was not different between runners and
269 non-runners of both age groups; data for positive and negative pressure ramps and the
270 number of sequences *per* ramp are presented in Table 3.

271 **Arterial stiffness**

272 Runners had lower aPWV (middle-aged, $P=0.026$; young, $P=0.027$) compared to age-
273 matched non-runners (Table 2). In contrast, the β stiffness index of the carotid artery was
274 lower only for the middle-aged runners compared to age-matched non-runners ($P=0.041$).

275 **DISCUSSION**

276 The principal findings are as follows: 1) for middle-aged men, many years of moderate to
277 vigorous endurance exercise training sets the operating point of the vascular sympathetic
278 baroreflex at a burst occurrence that is higher than for peers that have not trained; 2) higher
279 burst occurrence does not influence overall reflex gain, basal burst frequency, or resting
280 arterial pressure; 3) for younger men, endurance training has a limited effect on the
281 operating point and there are no differences in vascular sympathetic baroreflex reflex gain or
282 basal burst frequency compared with untrained peers. Taken together, these findings
283 indicate that some form of remodelling in middle-aged men following many years of

284 committed endurance exercise training plays a critical role in the baroreflex control of
285 vascular sympathetic bursts and resting blood pressure.

286 **The effect of training on vascular sympathetic baroreflex control**

287 Regardless of the training status, we observed similar frequencies of sympathetic
288 bursts in microneurographic recordings taken from middle-aged males during supine rest. An
289 intriguing finding, however, is that the well-trained men exhibit a greater MSNA burst
290 occurrence, and by some margin (40 to 50 % approximately); this occurs without any
291 obvious difference in the corresponding diastolic pressure stimulus. Together, we interpret
292 these data for MSNA burst frequency and occurrence as evidence that many years of
293 training alters the gating of sympathetic bursts (i.e. baroreflex control) without influencing the
294 frequency of sympathetic bursts *per minute* (i.e. rate of neurotransmitter release). Although
295 this might seem contradictory, burst frequency and occurrence provide slightly different
296 neurophysiological information (5, 29, 53). Furthermore, reciprocal interplay between
297 exercise bradycardia (i.e. fewer opportunities for a burst) and the higher MSNA operating
298 point (i.e. greater burst occurrence) explain why the burst frequency for trained runners and
299 non-runners is similar.

300 Our data indicate that an exercise training-induced upward setting for the MSNA
301 operating point in middle age occurs without any change in the ability to increase or
302 decrease vasoconstrictor outflow during fluctuations of resting arterial pressure. In other
303 words, vascular sympathetic baroreflex overall gain is unaffected by training. Stüding and
304 colleagues (44), using the modified Oxford baroreceptor test, also observed that overall gain
305 was similar among older trained and untrained men. Unlike the present study, however, no
306 difference was observed for sympathetic burst occurrence, and resting burst frequency was
307 marginally lower for endurance-trained versus untrained middle-aged males.

308 Other studies of trained and untrained middle-aged and older people have recorded
309 resting MSNA without specifically addressing vascular sympathetic baroreflex function.

310 Notarius and colleagues (38) observed that burst occurrence was higher, while basal
311 sympathetic burst frequency was similar, for endurance trained middle-aged men compared
312 with sedentary peers. In contrast, Ng and co-workers reported higher sympathetic burst
313 occurrence and burst frequency for older-endurance trained athletes; however, these
314 findings may reflect an older cohort, or inclusion of endurance trained females, for whom
315 burst frequency was markedly higher compared with untrained peers (37). Whilst we cannot
316 explain this lack of consensus, it may reflect the differences in the endurance phenotype
317 across the studies. Factors that influence basal vascular sympathetic outflow with human
318 ageing, such as abdominal adiposity (15), distensibility of the barosensory vessel walls (48),
319 and blood volume (2), all are influenced by the dose of endurance exercise training.

320 To isolate the effect of endurance exercise training from human ageing, we also
321 studied younger males. As with older men, we found no difference for basal burst frequency
322 between well-trained runners and non-runners. The burst occurrence was marginally higher
323 for the runners, but this difference was modest in comparison to that between the older
324 groups. These findings in young men are similar to previous cross-sectional studies (7, 43,
325 46). Furthermore, vascular sympathetic baroreflex gain is similar for trained runners and
326 non-runners. Thus, our data suggest that the endurance phenotype traits of young men does
327 not include a higher operating point for the vascular sympathetic baroreflex.

328 Differences for aortic compliance and resting heart rate between young runners and
329 non-runners are comparable with those for the middle-aged men. However, one noteworthy
330 distinction relates to the difference in resting stroke volume. For young, well-trained men,
331 stroke volume during supine rest was 50% greater than that of age-matched non-runners.
332 For older men, resting stroke volume was only 12% greater for runners compared with age-
333 matched non-runners. This lesser difference in stroke volume may explain why endurance
334 training effects the operating point for the vascular sympathetic baroreflex only for committed
335 middle-aged runners. That is, older runners rely more on vascular sympathetic neural activity
336 than cardiac output to support arterial pressure. However, further investigation of potential

337 interaction of left ventricular stroke volume and the vascular sympathetic baroreflex is
338 required.

339 Our interpretation for young men in this study is consistent with a previous report that
340 endurance training does not influence autonomic support of blood pressure in the young
341 (17). However, our findings do contrast with those of a study by Alvarez and colleagues (1).
342 As in the present study, burst occurrence was marginally higher in trained men, while basal
343 MSNA burst frequency was similar. However, when adiposity is taken into account, burst
344 occurrence and burst frequency both were greater for endurance trained versus untrained
345 men (1). Furthermore, in contrast to the present study, sympathetic baroreflex gain was
346 lower for endurance-trained compared with untrained young men, an effect regardless of
347 percentage body fat. This suggests that body composition may be important, at least in
348 younger men.

349 **The effect of training on cardiovagal baroreflex control**

350 It is well known that endurance athletes display exercise-induced bradycardia,
351 although considerable debate exists surrounding the mechanism(s) involved (3, 4).
352 Furthermore, arterial baroreceptor control of blood pressure is mediated predominantly via
353 sympathetic vascular regulation, rather than by reflex changes in heart period (8).
354 Nonetheless, we determined how habitual endurance exercise influenced the
355 responsiveness of the cardiovagal baroreflex in middle age. For well-trained middle-aged
356 men, as expected, the cardiovagal baroreflex operated around a considerably longer RR
357 interval at rest; however, the baroreflex gain was similar among runners and non-runners.
358 Previous work has shown that middle-aged endurance trained men display greater
359 cardiovagal baroreflex gain than sedentary controls, but not moderately-active, age-matched
360 peers (34). In the case of the younger trained men in this study, the cardiovagal baroreflex
361 also operated around a longer heart period, without any difference in baroreflex gain
362 compared with age-matched non-runners; this finding for gain is in agreement with previous
363 studies in younger men (1, 7, 34).

364 **Remodelling of the vascular sympathetic baroreflex**

365 Mechanosensory transduction, central mediation, and efferent neurotransmission are
366 integrated into the baroreflex regulation of vasomotor tone and arterial pressure.
367 Furthermore, it is proposed that human aging may have opposing influences on mechanical
368 and neural events (44). However, we can only speculate upon potential sites where
369 additional remodelling might have occurred in committed middle-aged runners to explain our
370 findings. Many years of training may influence the strength and/or timing of mechanosensory
371 signals controlling efferent sympathetic burst occurrence; this could arise from altered
372 vascular mechanics and/or a change to the threshold for baroreceptor activation.
373 Specifically, well-trained middle-aged men have less stiff barosensory regions; furthermore,
374 more complete elastic recoil during a longer diastolic period could lead to a longer interval of
375 'silence' in the afferent baroreceptor signal (21). However, the apparent lack of a similar
376 upward setting of the MSNA operating point for younger trained men, who also possess
377 lesser vascular stiffness and display bradycardia, argues against this. However, endurance-
378 training induced cardiovascular remodelling may only lead to upward vascular sympathetic
379 baroreflex resetting in middle-aged men due to increased autonomic support of blood
380 pressure with age (16).

381 Animal studies indicate that chronic exercise training potentially influences
382 baroreceptor control of sympathetic bursts at brain structures including, the nucleus tractus
383 solitarius, the paraventricular nucleus of the hypothalamus, and the rostral ventrolateral
384 medulla (36). Brain imaging studies have identified some of the same sites as regions of
385 baroreflex control in humans (22) (23). It is possible, therefore, that neural plasticity and
386 exercise-induced central remodelling previously observed in animals underpins the higher
387 sympathetic burst occurrence in middle-aged trained males.

388 Changes to efferent neurotransmission may also mediate upward vascular
389 sympathetic baroreflex setting. Short-term exercise training reduces alpha-adrenergic
390 vasoconstrictor responsiveness in (35), and a reduction of sympathetic vascular transduction

391 has been proposed to contribute to orthostatic intolerance observed in some highly-trained
392 individuals (52). Vasoconstrictor responsiveness to noradrenaline declines with advancing
393 age (10), which may counteract the effects of elevated MSNA burst frequency (16).
394 Furthermore, Notarius and colleagues (38) observed that sympathetic vascular transduction
395 during baroreflex-mediated sympathoexcitation may be altered further in trained middle-aged
396 men. Another possibility is that vascular sympathetic baroreflex resetting may be a
397 compensatory mechanism to offset training-induced vascular changes (42) (33). All of these
398 aforementioned possibilities require investigation. Notably, irrespective of the location(s),
399 exercise-induced remodelling does not alter vascular sympathetic baroreflex gain, at least
400 not the integrated gain.

401 **Experimental Considerations**

402 Vascular sympathetic baroreflex gain was calculated by associating spontaneous
403 fluctuations in DBP to the occurrence of bursts of MSNA. We did not take strength
404 (amplitude) of sympathetic bursts into account, because baroreceptor signals modulate burst
405 occurrence, whereas less is known of the mechanisms that govern amplitude (21, 29).
406 Furthermore, we did not assess vascular sympathetic baroreflex gain to rising and falling
407 pressures independently and we acknowledge that this does not take baroreflex hysteresis
408 into account (14).

409 It is reported that dietary salt and nitrate can influence sympathetic burst activity (28,
410 39). However, we did not control for diet in our study, therefore we cannot exclude some
411 influence on our data. Every effort was made to accurately record the number of years over
412 which an individual had exercised at their current level. In addition, we recorded lifetime
413 physical activity and exercise and observed a clear difference in maximal aerobic capacity
414 between the trained and untrained groups. However, group allocation, determined by
415 habitual endurance training, may limit the conclusions based on other components of
416 exercise training. These components include mode, intensity, duration, all of which may
417 have an impact on cardiac, vascular and neural remodelling. Because sex of the participants

418 was controlled for in this study, future studies are required to properly address potential sex
419 differences. Although our participants were non-obese, we did not specifically control for
420 adiposity, which is known to influence sympathetic burst activity. However, post hoc analysis
421 suggest that percentage body fat was not a significant covariate for any indices of
422 sympathetic activity in this study. Finally, the *a priori* intention of our study was to investigate
423 the effect that committed endurance exercise training has on elevated sympathetic neural
424 activity and vascular sympathetic baroreflex control of resting blood pressure in healthy
425 middle-aged men. However, we also studied young men in order to investigate the effect of
426 endurance training independently of cardiovascular ageing. The use of independent samples
427 t-tests reflects these *a priori* questions. To limit the chance of a type 1 error, we did not
428 perform statistical comparisons between middle-aged runners and young runners, or middle-
429 aged non-runners and young non-runners.

430 **CONCLUSION**

431 This study demonstrates upward setting of arterial baroreflex regulation of vascular
432 sympathetic bursts following committed endurance training in middle-aged men. Importantly,
433 vascular sympathetic baroreflex resetting coupled to exercise-induced bradycardia, results in
434 a similar basal burst frequency compared with untrained peers. Furthermore, the study
435 demonstrates that training status does not influence the MSNA operating point for younger
436 well-trained men, who also display similar sympathetic burst frequency compared with
437 untrained peers. In our view, remodelling within the vascular sympathetic baroreflex arc,
438 culminating in a higher MSNA operating point, is another example of phenotypic adaptation
439 to lifelong (> 25 years) training. This occurs, presumably, to maintain resting vasomotor tone
440 and blood pressure stability and to complement cardiac and vascular adaptations to many
441 years of endurance exercise training.

442 **FUNDING**

443 DJW supported by a PhD studentship from the School of Sport and Health, Cardiff
444 Metropolitan University. LLS supported by a PhD studentship from the School of Sport
445 Health and Exercise Sciences, Bangor University.

446 **CONFLICT OF INTEREST**

447 None of the authors has any conflicts of interest, financial or other.

448 **AUTHOR CONTRIBUTIONS**

449 DJW, RS, CJP and JPM conception and design; DJW, RNL, JST, FML, BAC, TGD, LLS,
450 CJAP, and JPM performed experiments. DJW, JST, RNL, and JPM data analysis. DJW and
451 JPM data interpretation. DJW and JPM drafted manuscript. DJW, RS, CJP and JPM revised
452 manuscript. All authors approved the manuscript.

453 **ACKNOWLEDGEMENTS**

454 We thank all the participants for taking part in this study.

455 **References**

- 456 1. **Alvarez GE, Halliwill JR, Ballard TP, Beske SD, and Davy KP.** Sympathetic neural regulation in
457 endurance-trained humans: fitness vs. fatness. *J Appl Physiol (1985)* 98: 498-502, 2005.
- 458 2. **Best SA, Okada Y, Galbreath MM, Jarvis SS, Bivens TB, Adams-Huet B, and Fu Q.** Age and
459 sex differences in muscle sympathetic nerve activity in relation to haemodynamics, blood volume
460 and left ventricular size. *Exp Physiol* 99: 839-848, 2014.
- 461 3. **Billman GE.** Counterpoint: Exercise training-induced bradycardia: the case for enhanced
462 parasympathetic regulation. *J Appl Physiol (1985)* 123: 686-688, 2017.
- 463 4. **Boyett MR, Wang Y, Nakao S, Ariyaratnam J, Hart G, Monfredi O, and D'Souza A.** Point:
464 Exercise training-induced bradycardia is caused by changes in intrinsic sinus node function. *J Appl*
465 *Physiol (1985)* 123: 684-685, 2017.
- 466 5. **Charkoudian N and Wallin BG.** Sympathetic neural activity to the cardiovascular system:
467 integrator of systemic physiology and interindividual characteristics. *Compr Physiol* 4: 825-850, 2014.
- 468 6. **Christou DD, Jones PP, Jordan J, Diedrich A, Robertson D, and Seals DR.** Women have lower
469 tonic autonomic support of arterial blood pressure and less effective baroreflex buffering than men.
470 *Circulation* 111: 494-498, 2005.
- 471 7. **Christou DD, Jones PP, and Seals DR.** Baroreflex buffering in sedentary and endurance
472 exercise-trained healthy men. *Hypertension* 41: 1219-1222, 2003.
- 473 8. **Dampney RAL.** Resetting of the Baroreflex Control of Sympathetic Vasomotor Activity during
474 Natural Behaviors: Description and Conceptual Model of Central Mechanisms. *Front Neurosci* 11:
475 461, 2017.
- 476 9. **Delius W, Hagbarth KE, Hongell A, and Wallin BG.** Manoeuvres affecting sympathetic
477 outflow in human muscle nerves. *Acta Physiol Scand* 84: 82-94, 1972.
- 478 10. **Dinunno FA, Dietz NM, and Joyner MJ.** Aging and forearm postjunctional alpha-adrenergic
479 vasoconstriction in healthy men. *Circulation* 106: 1349-1354, 2002.
- 480 11. **Ebert TJ, Morgan BJ, Barney JA, Denahan T, and Smith JJ.** Effects of aging on baroreflex
481 regulation of sympathetic activity in humans. *Am J Physiol* 263: H798-803, 1992.
- 482 12. **Eckberg DL and Eckberg MJ.** Human sinus node responses to repetitive, ramped carotid
483 baroreceptor stimuli. *Am J Physiol* 242: H638-644, 1982.
- 484 13. **Franklin SS, Gustin W, Wong ND, Larson MG, Weber MA, Kannel WB, and Levy D.**
485 Hemodynamic Patterns of Age-Related Changes in Blood Pressure: The Framingham Heart Study.
486 *Circulation* 96: 308-315, 1997.
- 487 14. **Hart EC, Wallin BG, Curry TB, Joyner MJ, Karlsson T, and Charkoudian N.** Hysteresis in the
488 sympathetic baroreflex: role of baseline nerve activity. *J Physiol* 589: 3395-3404, 2011.
- 489 15. **Jones PP, Davy KP, Alexander S, and Seals DR.** Age-related increase in muscle sympathetic
490 nerve activity is associated with abdominal adiposity. *Am J Physiol* 272: E976-980, 1997.
- 491 16. **Jones PP, Shapiro LF, Keisling GA, Jordan J, Shannon JR, Quaife RA, and Seals DR.** Altered
492 autonomic support of arterial blood pressure with age in healthy men. *Circulation* 104: 2424-2429,
493 2001.
- 494 17. **Jones PP, Shapiro LF, Keisling GA, Quaife RA, and Seals DR.** Is autonomic support of arterial
495 blood pressure related to habitual exercise status in healthy men? *J Physiol* 540: 701-706, 2002.
- 496 18. **Joyner MJ, Barnes JN, Hart EC, Wallin BG, and Charkoudian N.** Neural control of the
497 circulation: how sex and age differences interact in humans. *Compr Physiol* 5: 193-215, 2015.
- 498 19. **Kawasaki T, Sasayama S, Yagi S, Asakawa T, and Hirai T.** Non-invasive assessment of the age
499 related changes in stiffness of major branches of the human arteries. *Cardiovasc Res* 21: 678-687,
500 1987.
- 501 20. **Keller DM, Cui J, Davis SL, Low DA, and Crandall CG.** Heat stress enhances arterial
502 baroreflex control of muscle sympathetic nerve activity via increased sensitivity of burst gating, not
503 burst area, in humans. *J Physiol* 573: 445-451, 2006.
- 504 21. **Kienbaum P, Karlsson T, Sverrisdottir YB, Elam M, and Wallin BG.** Two sites for
505 modulation of human sympathetic activity by arterial baroreceptors? *J Physiol* 531: 861-869, 2001.

- 506 22. **Kimmerly DS, O'Leary DD, Menon RS, Gati JS, and Shoemaker JK.** Cortical regions
507 associated with autonomic cardiovascular regulation during lower body negative pressure in
508 humans. *J Physiol* 569: 331-345, 2005.
- 509 23. **Kramer HH, Ament SJ, Breimhorst M, Klega A, Schmiege K, Endres C, Buchholz HG, Elam M,
510 Schreckenberger M, and Birklein F.** Central correlation of muscle sympathetic nerve activation
511 during baroreflex unloading - a microneurography-positron emission tomography study. *Eur J*
512 *Neurosci* 39: 623-629, 2014.
- 513 24. **Lang RM, Badano LP, Mor-Avi V, Afilalo J, Armstrong A, Ernande L, Flachskampf FA, Foster
514 E, Goldstein SA, Kuznetsova T, Lancellotti P, Muraru D, Picard MH, Rietzschel ER, Rudski L, Spencer
515 KT, Tsang W, and Voigt JU.** Recommendations for cardiac chamber quantification by
516 echocardiography in adults: an update from the American Society of Echocardiography and the
517 European Association of Cardiovascular Imaging. *J Am Soc Echocardiogr* 28: 1-39 e14, 2015.
- 518 25. **Matsukawa T, Sugiyama Y, Iwase S, and Mano T.** Effects of aging on the arterial baroreflex
519 control of muscle sympathetic nerve activity in healthy subjects. *Environ Med* 38: 81-84, 1994.
- 520 26. **Matsukawa T, Sugiyama Y, and Mano T.** Age-related changes in baroreflex control of heart
521 rate and sympathetic nerve activity in healthy humans. *J Auton Nerv Syst* 60: 209-212, 1996.
- 522 27. **Matsukawa T, Sugiyama Y, Watanabe T, Kobayashi F, and Mano T.** Gender difference in
523 age-related changes in muscle sympathetic nerve activity in healthy subjects. *Am J Physiol* 275:
524 R1600-1604, 1998.
- 525 28. **Matthews EL, Brian MS, Edwards DG, Stocker SD, Wenner MM, and Farquhar WB.** Blood
526 pressure responses to dietary sodium: Association with autonomic cardiovascular function in
527 normotensive adults. *Auton Neurosci* 208: 51-56, 2017.
- 528 29. **McAllen RM and Malpas SC.** Sympathetic burst activity: characteristics and significance. *Clin*
529 *Exp Pharmacol Physiol* 24: 791-799, 1997.
- 530 30. **McEniery CM, Yasmin, Hall IR, Qasem A, Wilkinson IB, Cockcroft JR, and Investigators A.**
531 Normal vascular aging: differential effects on wave reflection and aortic pulse wave velocity: the
532 Anglo-Cardiff Collaborative Trial (ACCT). *J Am Coll Cardiol* 46: 1753-1760, 2005.
- 533 31. **Monahan KD.** Effect of aging on baroreflex function in humans. *Am J Physiol Regul Integr*
534 *Comp Physiol* 293: R3-R12, 2007.
- 535 32. **Monahan KD, Dinunno FA, Seals DR, Clevenger CM, Desouza CA, and Tanaka H.** Age-
536 associated changes in cardiovagal baroreflex sensitivity are related to central arterial compliance.
537 *Am J Physiol Heart Circ Physiol* 281: H284-289, 2001.
- 538 33. **Monahan KD, Dinunno FA, Seals DR, and Halliwill JR.** Smaller age-associated reductions in
539 leg venous compliance in endurance exercise-trained men. *Am J Physiol Heart Circ Physiol* 281:
540 H1267-1273, 2001.
- 541 34. **Monahan KD, Dinunno FA, Tanaka H, Clevenger CM, DeSouza CA, and Seals DR.** Regular
542 aerobic exercise modulates age-associated declines in cardiovagal baroreflex sensitivity in healthy
543 men. *J Physiol* 529 Pt 1: 263-271, 2000.
- 544 35. **Mortensen SP, Nyberg M, Gliemann L, Thaning P, Saltin B, and Hellsten Y.** Exercise training
545 modulates functional sympatholysis and alpha-adrenergic vasoconstrictor responsiveness in
546 hypertensive and normotensive individuals. *J Physiol* 592: 3063-3073, 2014.
- 547 36. **Mueller PJ, Clifford PS, Crandall CG, Smith SA, and Fadel PJ.** Integration of Central and
548 Peripheral Regulation of the Circulation during Exercise: Acute and Chronic Adaptations. *Compr*
549 *Physiol* 8: 103-151, 2017.
- 550 37. **Ng AV, Callister R, Johnson DG, and Seals DR.** Endurance exercise training is associated with
551 elevated basal sympathetic nerve activity in healthy older humans. *J Appl Physiol (1985)* 77: 1366-
552 1374, 1994.
- 553 38. **Notarius CF, Murai H, Morris BL, and Floras JS.** Effect of fitness on reflex sympathetic
554 neurovascular transduction in middle-age men. *Med Sci Sports Exerc* 44: 232-237, 2012.

- 555 39. **Notay K, Incognito AV, and Millar PJ.** Acute beetroot juice supplementation on sympathetic
556 nerve activity: a randomized, double-blind, placebo-controlled proof-of-concept study. *Am J Physiol*
557 *Heart Circ Physiol* 313: H59-H65, 2017.
- 558 40. **Parati G, Di Rienzo M, and Mancia G.** How to measure baroreflex sensitivity: from the
559 cardiovascular laboratory to daily life. *J Hypertens* 18: 7-19, 2000.
- 560 41. **Pauca AL, O'Rourke MF, and Kon ND.** Prospective evaluation of a method for estimating
561 ascending aortic pressure from the radial artery pressure waveform. *Hypertension* 38: 932-937,
562 2001.
- 563 42. **Prior BM, Lloyd PG, Yang HT, and Terjung RL.** Exercise-induced vascular remodeling. *Exerc*
564 *Sport Sci Rev* 31: 26-33, 2003.
- 565 43. **Seals DR.** Sympathetic neural adjustments to stress in physically trained and untrained
566 humans. *Hypertension* 17: 36-43, 1991.
- 567 44. **Studinger P, Goldstein R, and Taylor JA.** Age- and fitness-related alterations in vascular
568 sympathetic control. *J Physiol* 587: 2049-2057, 2009.
- 569 45. **Sundlof G and Wallin BG.** Human muscle nerve sympathetic activity at rest. Relationship to
570 blood pressure and age. *J Physiol* 274: 621-637, 1978.
- 571 46. **Svedenhag J, Wallin BG, Sundlof G, and Henriksson J.** Skeletal muscle sympathetic activity
572 at rest in trained and untrained subjects. *Acta Physiol Scand* 120: 499-504, 1984.
- 573 47. **Tanaka H, Dinunno FA, Monahan KD, Clevenger CM, DeSouza CA, and Seals DR.** Aging,
574 habitual exercise, and dynamic arterial compliance. *Circulation* 102: 1270-1275, 2000.
- 575 48. **Tanaka H, Dinunno FA, and Seals DR.** Reductions in central arterial compliance with age are
576 related to sympathetic vasoconstrictor nerve activity in healthy men. *Hypertens Res* 40: 493-495,
577 2017.
- 578 49. **Taylor JA and Tan CO.** BP regulation VI: elevated sympathetic outflow with human aging:
579 hypertensive or homeostatic? *Eur J Appl Physiol* 114: 511-519, 2014.
- 580 50. **Townsend RR, Wilkinson IB, Schiffrin EL, Avolio AP, Chirinos JA, Cockcroft JR, Heffernan KS,**
581 **Lakatta EG, McEniery CM, Mitchell GF, Najjar SS, Nichols WW, Urbina EM, Weber T, and American**
582 **Heart Association Council on H.** Recommendations for Improving and Standardizing Vascular
583 Research on Arterial Stiffness: A Scientific Statement From the American Heart Association.
584 *Hypertension* 66: 698-722, 2015.
- 585 51. **Vaitkevicius PV, Fleg JL, Engel JH, O'Connor FC, Wright JG, Lakatta LE, Yin FC, and Lakatta**
586 **EG.** Effects of age and aerobic capacity on arterial stiffness in healthy adults. *Circulation* 88: 1456-
587 1462, 1993.
- 588 52. **van Lieshout JJ.** Exercise training and orthostatic intolerance: a paradox? *J Physiol* 551: 401,
589 2003.
- 590 53. **Wallin BG, Esler M, Dorward P, Eisenhofer G, Ferrier C, Westerman R, and Jennings G.**
591 Simultaneous measurements of cardiac noradrenaline spillover and sympathetic outflow to skeletal
592 muscle in humans. *J Physiol* 453: 45-58, 1992.
- 593 54. **White DW, Shoemaker JK, and Raven PB.** Methods and considerations for the analysis and
594 standardization of assessing muscle sympathetic nerve activity in humans. *Auton Neurosci* 193: 12-
595 21, 2015.
- 596

Table 1 - Participant Characteristics

	Young non-runners (n=10)	Young runners (n=13)	Middle-aged non- runners (n=10)	Middle-aged runners (n=13)
<i>Demographics</i>				
Age, years	23 (21-25)	22 (21-24)	53 (52-55)	57 (54-59)
Stature, cm	178.1 (174.0-182.3)	179.9 (176.9-183.0)	175.6 (170.5-180.6)	174.7 (170.9-178.5)
Body Mass, kg	80.4 (68.8-92.0)	67.0 (63.9-70.0) *	80.9 (73.8-88.0)	66.1 (61.3-70.9) †
BMI, kg·m ²	25.4 (22.1-28.8)	20.8 (19.9-21.6) *	26.2 (24.0-28.5)	21.6 (20.7-22.6) †
Body fat (%)	19.7 (15.2-24.1)	10.7 (7.8-13.6) *	26.8 (20.4-33.3)	17.5 (15.6-19.3) †
<i>Blood Pressure</i>				
SBP, mmHg	119 (109-128)	111 (108-114) *	119 (113-124)	118 (113-123)
DBP, mmHg	71 (67-76)	66 (62-69) *	76 (73-80)	74 (70-78)
<i>Cardiorespiratory Fitness</i>				
$\dot{V}O_2$ Peak, mL·kg ⁻¹ ·min ⁻¹	36.5 (31.9-41.0)	60.6 (55.0-66.2) *	32.6 (26.6-38.6)	50.7 (47.0-54.4) †
$\dot{V}O_2$ Peak, % Predicted	86 (82-103)	116 (116-141) *	106 (87-129)	143 (129-155) †
<i>Training History</i>				
Exercise per week, miles		65 (56-73)		34 (28-39)
Training history, years		8 (5-11)		29 (28-40)

Data are presented as mean (95% Confidence Intervals). Symbols represent significant between-group differences ($P < 0.05$), * = Young runner vs. Young non-runner; † = Middle-aged runner vs. middle-aged non-runner.

601

602

603

Table 2 - Resting Haemodynamics and Basal Sympathetic Nervous System Activity

604

	Young		Middle-aged	
	non-runners (n = 10)	runners (n = 13)	non-runners (n = 10)	runners (n = 13)
<i>Central Artery Stiffness</i>				
aPWV, m·s ⁻¹	5.8 (5.2-6.3)	5.1 (4.8-5.3) *	7.5 (6.9-8.1)	6.8 (6.2-7.3) †
β stiffness index	2.96 (2.52-3.40)	2.38 (2.03-2.74)	5.06 (3.94-6.19)	4.06 (3.25-4.88) †
<i>Haemodynamics</i>				
Heart rate, beats·min ⁻¹	64 (57-70)	45 (41-48) *	56 (49-62)	43 (38-47) †
Stroke volume, ml	61 (57-64)	92 (87-97) *	62 (56-68)	70 (63-77) †
Cardiac output, L·min ⁻¹	3.8 (3.5-4.2)	4.1 (3.8-4.4)	3.4 (3.0-3.8)	3.0 (2.6-3.3)
TPR, mmHg·L·min ⁻¹	24.3 (21.2-27.4)	21.3 (19.2-23.3)	29.1 (26.8-31.3)	31.6 (28.4-34.7)
MAP, mmHg	90 (83-97)	84 (81-88)	95 (89-101)	93 (90-96)
Respiration rate, breaths·min ⁻¹	13 (10-15)	15 (14-16)	11 (9-13)	12 (10-14)
<i>Muscle Sympathetic Nerve Activity</i>				
Burst Frequency, bursts·min ⁻¹	18 (12-23)	16 (10-21)	28 (19-38)	31 (27-34)
Burst Incidence, bursts·100hb ⁻¹	27 (19-36)	36 (23-50)	50 (33-66)	72 (63-81) †

Data are presented as mean (95% Confidence Intervals). Symbols represent significant between-group differences ($P < 0.05$), * = Young runner vs. Young non-runner; † = Middle-aged runner vs. middle-aged non-runner.

Note: We were unable to quantify β stiffness index in one young non-runner and one young runner; accordingly, data are reported for forty-four individuals. Furthermore, stroke volume was unobtainable for one middle-aged runner. Accordingly, stroke volume, Q and TPR data are reported in forty-five individuals.

605

606

Table 3 – Cardiovagal baroreflex gain and the number of sequences for positive and negative pressure ramps

	Young		Middle-aged	
	non-runners (n = 10)	runners (n = 11)	non-runners (n = 9)	runners (n = 13)
'Up' Gain ($ms \cdot mmHg^{-1}$)	31 (23-39)	41 (27-55)	28 (18-37)	34 (25-44)
# sequences	20 (12-29)	8 (5-11)	17 (10-24)	11 (8-16)
'Down' Gain ($ms \cdot mmHg^{-1}$)	24 (17-32)	33 (22-45)	23 (14-32)	33 (22-45)
# sequences	30 (20-39)	9 (6-12)	20 (13-26)	13 (8-19)

Data are presented as mean (95% Confidence Intervals).

607

608

609 **Figure Legends**

610

611 **Figure 1 Example recordings of muscle sympathetic nerve activity and blood pressure during supine rest.** 20 seconds of resting muscle sympathetic
612 nerve activity (MSNA) and blood pressure (BP) data are shown from one representative participant per group: (A) Middle-aged runner; (B) Middle-aged non-
613 runner; (C) Young runner; (D) Young non-runner.

614

615 **Figure 2 Sympathetic and cardiac baroreflex function. (A)** Group mean regressions between diastolic blood pressure (DBP) and muscle sympathetic
616 nerve activity (MSNA) are presented with the sympathetic operating points superimposed on the regression lines. Middle-age runners had similar operating
617 DBP compared to middle-aged non-runners but the corresponding level of MSNA was higher (by 22 bursts·100hb⁻¹; red arrow), despite similar sympathetic
618 baroreflex gain. However, in young men training status had no influence on the operating DBP, corresponding level of MSNA or sympathetic baroreflex gain.
619 **(B)** Group mean regressions between systolic blood pressure (SBP) and R-R interval (sequence method) are shown with the operating points of the cardiac
620 baroreflex overlaid on the regression lines. Middle-aged runners had similar operating SBP and cardiovagal baroreflex gain (33.6 [24.5-42.8] vs 25.5 [16.2-
621 34.7], *P*=0.16) compared to middle-aged non-runners, but the corresponding R-R interval was longer (by 352 msec; red arrow). In contrast, when compared
622 to young non-runners, the operating SBP was set leftward (by 9 mmHg; green dashed arrow) in young runners with a longer corresponding R-R interval (by
623 418 msec; green solid arrow), despite similar cardiac baroreflex gain (37.2 [28.1-46.3] vs 26.4 [19.1-33.8], *P*=0.06). Abbreviations: M, Middle-aged non-
624 runners; MR, Middle-aged runner; Y, Young non-runner; YR, Young runner. NB: Baroreflex responsiveness data are presented from: 10 young non-runners,
625 11 young runners, 9 middle-aged non-runners, 11 middle-aged runners.

626



