

How do hedgerows influence soil organic carbon stock in livestock-grazed pasture?

Ford, Hilary; Healey, John R.; Webb, Bid; Pagella, Tim F.; Smith, Andrew R.

Soil Use and Management

DOI: 10.1111/sum.12517

Published: 31/12/2019

Peer reviewed version

Cyswllt i'r cyhoeddiad / Link to publication

Dyfyniad o'r fersiwn a gyhoeddwyd / Citation for published version (APA): Ford, H., Healey, J. R., Webb, B., Pagella, T. F., & Smith, A. R. (2019). How do hedgerows influence soil organic carbon stock in livestock-grazed pasture? *Soil Use and Management*, *35*(4), 576-584. https://doi.org/10.1111/sum.12517

Hawliau Cyffredinol / General rights Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

· Users may download and print one copy of any publication from the public portal for the purpose of private study or research.

- You may not further distribute the material or use it for any profit-making activity or commercial gain
 You may freely distribute the URL identifying the publication in the public portal ?

Take down policy If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

How do hedgerows influence soil organic carbon stock in livestock-grazed pasture?
 Hilary Ford*, John R. Healey, Bid Webb, Tim F. Pagella, Andrew. R. Smith
 School of Natural Sciences, Bangor University, Bangor, LL57 2DG, UK.

4 *Corresponding author. *Email address*: <u>hilary.ford@bangor.ac.uk</u> (H. Ford).

5 Running head title: Hedgerows and soil organic carbon

6 Abstract

7 Hedgerows have the potential to influence ecosystem function in livestock-grazed pasture. 8 Despite this, they are often ignored when quantifying farmland ecosystem service delivery. In 9 this study, we assess the contribution of hedgerows to the ecosystem function of carbon (C) 10 storage, with a particular emphasis on soil organic carbon (SOC). We measured SOC stock (kg C m⁻²), on an equivalent soil mass basis, at 0-0.15 m depth in pasture adjacent to 38 11 hedgerows (biotic) and 16 stone walls or fences (abiotic controls) across ten farms in the 12 county of Conwy, Wales, UK. Pasture SOC stock (~7 kg C m⁻²) was similar adjacent to biotic 13 and abiotic field boundaries, positively associated with soil moisture and negatively with soil 14 bulk density (BD). For biotic boundaries two further variables were significantly associated 15 16 with SOC stock, distance from hedgerow (decrease in SOC stock) and slope orientation 17 (upslope SOC stock greater than downslope). For pasture adjacent to hedgerows a model 18 combining the aforementioned variables (BD, soil moisture, distance from hedgerow, slope orientation) explained 78% of variation in SOC stock. This study demonstrates that, whilst 19 hedgerows do have subtle positive effects on SOC stock in adjacent pasture, SOC storage 20 21 adjacent to field boundaries is influenced more by soil moisture content and BD than field 22 boundary type.

Keywords: Agriculture; Agroforestry; Ecosystem function; Grassland; Soil carbon; Woody
 linear feature

25 Introduction

26 The importance of hedgerows for provision of regulatory ecosystem services, including water 27 quality, flood risk reduction, soil erosion prevention, shelter provision (livestock) and climate change mitigation (via carbon (C) storage), has been increasingly recognised over the past 28 decade (Wolton et al., 2014; Scholefield et al., 2016). Despite this, the contribution of 29 30 hedgerows, lines of trees and shrubs typically managed by regular cutting (Baudry, Bunce & Burel, 2000) and other woody linear features (e.g. lines of trees or riparian strips), are 31 32 generally not properly accounted for when quantifying ecosystem services (Scholefield et al., 2016; Cardinael *et al.*, 2018) due to a paucity of data on extent and condition. 33

The store of soil organic carbon (SOC) is usually dominated by soil organic matter (SOM), a 34 complex combination of plant- or animal-derived organic residues (e.g. leaf litter, root 35 biomass and exudates, microbial biomass, animal faeces) in various states of decomposition 36 37 (Baah-Acheamfour et al., 2015). During the process of decomposition, C within SOM is either incorporated into the soil matrix as SOC, released to the atmosphere as carbon dioxide or 38 39 methane, or transferred to ground water through leaching (Benham *et al.*, 2012). Dissolved organic C inputs, from exudation and plant-derived decomposition products of high molecular 40 weight compounds, are thought to be the main source of SOC with a long residence time in 41 terrestrial ecosystems (Sokol & Bradford, 2019). Labile C inputs from root and hyphal turnover 42 directly influence microbial resource use efficiency (Lange et al., 2015), with microbial 43 necromass forming a major part of the slow-cycling SOM/SOC pool in deeper soils. 44

Grassland SOC stocks are greatest in temperate moist-cool climates (Abdalla et al., 2018), 45 where seasonally water-logged soils reduce the availability of oxygen, limiting the breakdown 46 of SOM. Soil type is also important, with SOC positively correlated with clay content (Jobbagy 47 & Jackson, 2000), as SOM is physically protected from microbial decomposition by adsorption 48 49 onto clay minerals within the soil. The effects of livestock grazing on SOC are dependent on 50 several factors including the physical properties of soil, precipitation levels, responsiveness of 51 the plant community to grazing and depth of soil sampling (Bardgett & Wardle, 2010). 52 Livestock grazing is often associated with increased allocation of plant resources belowground, with enhanced below-ground biomass and root turnover leading to SOC 53 accumulation (Kemp & Michalk, 2007). Negative impacts on SOC are largely seen where over-54 grazing leads to sparse vegetation cover and soil erosion (Golluscio et al., 2009). There is 55 general consensus that livestock-grazed pastures in temperate zones, particularly where 56 57 rainfall is plentiful, are a net C sink (Ostle et al., 2009; Soussana et al., 2010) with SOC stock 58 broadly comparable between semi-natural grassland and woodland (Bullock et al., 2011).

59 Detailed study of spatial variation in SOC (content and stock) has shown it to be greater close to hedgerows in both grassland and arable systems (Holden *et al.*, 2019), the effect decreasing 60 with distance from the hedge boundary, for up to 4 m into neighbouring fields (Follain et al., 61 2007; D'Acunto et al., 2014; Van Vooren et al., 2017). Hedgerow woody plant root 62 architecture and depth can influence SOC storage, with deeper-rooted species associated 63 64 with greater SOC (Crossland, 2015). Position relative to hedgerows is also important, with SOC 65 greater upslope of contour-planted hedgerows, where the hedgerow acts as a physical barrier reducing soil erosion and increasing A-horizon depth (Follain *et al.*, 2007). Regular hedgerow 66 67 management (via cutting with a tractor-mounted flail) increases the amount of surface litter adjacent to the hedgerow (Axe et al., 2017) potentially increasing inputs to soil. Regular 68

cutting also decreases the shoot-to-root ratio, which can influence fine root turnover and either increase (Peter & Lehmann, 2000) or decrease (Crossland, 2015) SOC storage. Even though hedgerow age (years since planting) is highly likely to influence SOC dynamics, due to change in the quality and rate of C inputs via root and hyphal turnover, exudation, and leaf litter over time as hedgerow plants grow, to our knowledge this aspect has not been studied previously.

In this study, we focused on the contribution of hedgerows to the ecosystem function of C storage in livestock-grazed pasture, with a particular emphasis on SOC. To do this we measured SOC stock (kg C m⁻² of land area), expressed on an equivalent soil mass (ESM) basis (Cardinael et al., 2018), in pasture adjacent to 38 hedgerows (biotic) and 16 stone walls or fences (abiotic controls). We hypothesised that SOC stock would: i) increase with hedgerow age (years since planting); ii) decrease with distance from hedgerow and iii) be greater upslope of hedgerows than downslope.

82 Materials and methods

83 Study area and sampling design

84 The study area (centred on 53.04° N, 3.71° W) encompassed ten tenant farms located within 85 the county of Conwy, Wales, UK close to the village of Ysbyty Ifan, all within the same River 86 Conwy catchment. All the farms were managed as mixed livestock, primarily Welsh mountain 87 sheep, with some beef cattle. All fields were permanent pasture with no arable crops grown. Pasture type was predominantly grassland that had been semi-improved with a mixture of 88 productive grass species, in most cases mixed with clover (Trifolium spp., N-fixing), forbs and 89 90 mosses. Certain fields were cut each year for silage. Soils were classified as either stagnogleys 91 (slowly permeable, seasonally wet, acid loam or clay soil) or brown earth (freely draining,

slightly acid loam soil) (Hallett *et al.*, 2017). The study area was categorised as poor (grade 4
or 5) agricultural land (Agricultural Land Classification of England and Wales, 2018), with
elevation 175-335 m asl. Mean annual precipitation is approximately 2,500 mm, with mean
monthly maximum and minimum temperatures of 12 and 6 °C.

Before study-site selection, all field boundaries across the ten farms were verified by the 96 97 tenant farmer as being in the place indicated on a satellite-derived image. Each hedgerow was marked on the image and its age since planting, reported by the farmer, recorded in 98 99 years. All other field boundaries were assumed to be a fence or stone wall. The study design allowed more than one side of the same field (typically 3-5 straight-sided polygons) to be 100 101 considered as an independent boundary. For example, a single field could have two abiotic 102 boundaries (stone wall and fence) and two hedgerows of different ages, giving four 103 independent boundaries. In total, 82 hedgerows were recorded and stratified into each of five age-since-planting categories (1-5, 6-10, 11-15, 16-20 or \geq 40 years). Eight replicates of 104 105 each age-since-planting category were randomly selected for data collection. Abiotic 106 boundaries were also randomly selected and subject to site-visit verification to ensure that 107 eight stone walls and eight fences were sampled. This approach ensured that all ten farms, 108 typically each located in one landscape unit characterised by altitude, soil type and 109 management style, were represented in the sampling structure. All hedgerows, apart from the \geq 40 years category, were planted under the regulations of previous government agri-110 111 environment schemes: double-fencing with 2-3 m total width to exclude livestock and 112 planting with a tree/shrub species mix of pre-dominantly Crataegus monogyna, Prunus *spinosa* and *Corylus avellana*. Some of the selected \geq 40 years hedgerows were, unavoidably, 113 114 single-fenced with a road on the opposite side to the pasture. The age of the stone walls was

uncertain although, from local knowledge, it is assumed they have been in position for > 100years.

117 For each of the 56 selected boundaries, the exact position of a sampling transect (one per boundary) was decided by ground conditions [specifically, avoidance of ditches, drains or 118 119 tracks, and avoidance of stone or rush (Juncus spp.) covered ground]. The position of each 120 sampling transect was located at least 5 m away from the end of each linear boundary, with the exact position judged visually to be representative of the whole boundary. Each sampling 121 122 transect was set up perpendicular to the boundary line starting at the boundary edge (for 123 hedgerows this was defined as the fence line, ca. 1.5 m away from the centre of the hedge) and ending 4.8 m into the adjacent grazed pasture (Figure 1). Sampling transects were only 124 set up in one direction for each boundary. As ground conditions were not assessed until each 125 126 farm visit, it was difficult to ensure an equal representation of slope orientations of transects 127 in advance. The following information was recorded for each transect: i) latitude, longitude 128 and elevation above sea level logged using a Garmin etrex 20x GPS; ii) slope orientation relative to field boundary (downslope, flat or variable, upslope, across slope; from now on 129 referred to as "slope orientation"); iii) presence or absence of tree standards along the 130 131 boundary within 5 m of transect origin; iv) soil texture, classified from a sample of fresh soil 132 ~0.05 m deep using the hand texturing method (Natural England, 2008). For hedgerow boundaries only, two additional variables were recorded: v) dominant species of woody plant 133 134 in the adjacent section of hedgerow (C. monogyna, P. spinosa or C. avellana); vi) management 135 of hedgerow [simplified into three categories a) regularly cut (typically annually with a tractormounted flail); b) un-managed (no active management for 5 years, often with tall and thin 136 137 shoots and/or long horizontal shoots); c) too young for management to be appropriate]. All field work was carried out October-December 2017. 138



Figure 1 Pasture soil sampling transect schematic for biotic (hedgerow) and abiotic (stone wall
and fence) boundaries.

143

144 Soil samples

Soil samples were taken at six distances along each transect (0.15, 0.3, 0.6, 1.2, 2.4 and 4.8 m from the origin), with the 4.8 m sample assumed to be indicative of the wider pasture away from the boundary line (Figure 1). Two soil samples were taken at each distance, one at ~0.05 m depth for pH analysis using 10 g fresh mass of soil with 25 ml of distilled water in a 1:2.5 dilution method (Rowell, 1994) to determine pH (Hanna Instruments pH meter 209, Leighton

Buzzard, England). The second was an intact core (0.15 m deep, 0.05 m diameter), for 150 151 assessment of BD, field soil moisture content, soil organic matter content, and SOC stock. Where stones prevented a full-length soil core being taken, the soil sample volume was 152 adjusted. Stones and woody roots (coarse roots > 2 mm diameter) were removed from each 153 154 soil sample and their volume recorded. Fine roots were not removed. The soil sample was then dried at 105 °C for 72 hours for measurement of field soil moisture content (%) and BD 155 (g cm⁻³). Each dried sample was then ground (pestle and mortar) and sieved through 2 mm to 156 157 remove any fine roots or small stones (not previously removed). SOM (% of dry soil) was measured from a sub-sample of ~10 g using the loss-on-ignition method (375 °C for 16 hours; 158 Ball, 1964). SOC concentration (g kg⁻¹ of dry soil mass) was calculated using the conversion 159 factor of 0.55 of SOM mass (Emmett et al., 2010) with SOC stock (kg C m⁻²) of 0-0.15 m depth 160 re-calculated on an ESM basis, a layer of 0-1000 t ha⁻¹ as in Lee *et al*. (2009). Most methods 161 162 of calculating SOC stock involve multiplying SOC concentration by BD to a fixed soil depth, 163 however this can lead to misleading results as soil compaction enhances SOC stock. The ESM method used here allows SOC stock to be compared uncoupled from the influence of 164 livestock-compaction. Data for two hedgerows (one in the 6-10 and one in the 11-15 years'-165 old category) were excluded due to high SOM content (>25%) indicating peat soil (Natural 166 England, 2008). Peat soils were excluded as they are associated with very low BD values, likely 167 168 to skew results. In addition to calculating SOM content in the standard way (% of dry soil mass after stones removed), SOM content was also calculated with stones included (% of dry soil 169 with stones included) to provide an indication of the SOM content in the field conditions. 170

All statistical analyses were carried out in R v3.4.3 (R Core Team, 2018). Linear mixed effects 173 174 models were used to compare soil characteristics (at six distances 0.15, 0.3, 0.6, 1.2, 2.4 and 4.8 m along each transect) of the livestock-grazed pasture [SOC stock, SOM content (stones 175 176 removed), SOM content (stones included), BD, moisture content, stone content, woody root 177 content, pH] adjacent to three field boundary types: i) hedgerows (n = 38); ii) stone walls (n = 8) and iii) fences (n=8). Linear mixed-effects models were also used to predict SOC stock 178 adjacent to biotic and abiotic field boundaries from the sampled and measured variables. For 179 180 hedgerows the following potential explanatory variables were tested: i) age-since-planting category of hedgerow; ii) absolute hedgerow age since planting in years; iii) perpendicular 181 distance from hedgerow boundary; iv) soil type as identified by soil maps (Hallett et al., 2017); 182 183 v) pasture routinely (once a year) cut for silage (*Boolean*); vi) slope orientation (simplified into three categories: 'upslope', 'flat or variable' and 'downslope' with 'across slope' excluded due 184 185 to lack of replication); vii) management (regularly cut/not cut/too young); viii) dominant 186 hedgerow woody species; ix) standard trees present or absent (Boolean); x) soil pH; xi) soil woody root content (% by volume of intact core, before stones removed); xii) soil moisture 187 188 (% of dry soil mass); xiii) soil BD. Field soil texture was not used as all soils were assessed as 189 broadly silty-clay loam. Predictive models of SOC stock, for fence and stone-wall boundaries, tested the same explanatory variables but with i), ii) and vii) excluded. 190

Linear mixed-effects models, with 'transect' nested within 'farm' (indicative of one landscape unit characterised by altitude, soil type and management style) identified as the true level of replication [e.g. lme (SOC ~ BD + Moisture + Distance, random = ~1|Farm/Transect)] were used. Best fit models were selected on the basis of lowest Akaike Information Criterion (AIC)

value (Zuur et al., 2009), with variables excluded if non-significant (one exception was made
where a non-significant variable decreased the AIC value relative to the model including only
significant variables). Likelihood-ratio-based pseudo-R-squared values were calculated for
each model (Grömping, 2006) with results presented using the ANOVA output of the mixed
effects models.

Linear mixed effects models were also used to compare soil characteristics [SOC stock, SOM content (stones removed), BD, moisture content, stone content, woody root content, pH] adjacent to and more distant from field boundaries. Adjacent samples were located at 0.15, 0.3, 0.6 and 1.2 m distance from the boundary edge (based on estimated zone of influence of hedgerow on SOC stock as ~2 m perpendicular to the boundary edge, Figure 2c). Distant samples were located at 2.4 and 4.8 m from the boundary edge and were assumed to be more indicative of the wider pasture.

207 Results

208 Boundary type and pasture soil characteristics

There were no significant differences in SOC stock or SOM content with stones removed (0-0.15 m depth) in adjacent pasture between hedges, stone walls and fences (Table 1). However, when SOM content was adjusted to include stone content of field soil it was significantly greater adjacent to fences (~11%) than stone walls (~10%). Woody root content was greater adjacent to hedgerows than either abiotic field boundary.

214 SOC stock adjacent to field boundaries

In pasture soil adjacent to biotic (hedgerow) boundaries SOC stock was significantly: i) positively associated with soil moisture content; ii) negatively associated with BD; iii)

negatively associated with distance from hedgerow; and iv) greater upslope than downslope of the boundary (Figure 2). In pasture soil adjacent to abiotic boundaries SOC stock was positively associated with soil moisture content for both walls and fences (Table 2) and negatively associated with BD adjacent to fences.

In pasture soil adjacent to hedgerow boundaries, SOC stock was not associated with: i) age

since planting (category or years) ($P \ge 0.65$); ii) hedgerow management ($P \ge 0.32$); iii) mapped

soil type, stagnogleys versus brown earth ($P \ge 0.54$); or iv) the hedgerow's dominant woody

plant species (*C. monogyna*, *P. spinosa* or *C. avellana*) ($P \ge 0.46$).

225 Boundary effects on soil characteristics

For biotic (hedgerow) boundaries, SOM content, SOC stock, stone content and woody root volume were significantly greater, but soil BD and pH significantly lower, adjacent to the boundary than distant from it in the pasture (Table 3). Soil moisture content was not significantly different between the two zones. In contrast, for abiotic boundaries (stone walls and fences combined) stone content was significantly greater adjacent to the boundary than distant from it, whereas all other variables did not differ significantly between the two zones (Table 3).

- **Table 1** Soil characteristics of livestock-grazed pasture, adjacent to biotic (hedgerow) and
- abiotic (stone wall and fence) field boundaries. Means and standard errors of the mean are
- 236 presented alongside the significance of differences between boundary types.
- 237

Variable	Hedge (n = 38) ^d	Wall (n = 8) ^d	Fence (n = 8) ^d	Sig. ^c
SOC stock (kg C m ⁻²)	6.62 ± 0.09	7.24 ± 0.25	7.02 ± 0.28	ns
SOM content (% dry soil mass; stones removed)	12.0 ± 0.16	13.2 ± 0.45	12.8 ± 0.51	ns
SOM content (% dry soil mass; stones included)	10.7 ± 0.19 ^{ab}	10.2 ± 0.49^{b}	11.2 ± 0.59 ^a	***
Stone content (% by volume of core)	10.3 ± 1.19 ^b	21.4 ± 3.26^{a}	11.7 ± 2.55 ^b	***
BD (g cm ⁻³)	0.79 ± 0.01 ^{ab}	0.84 ± 0.02^{a}	0.77 ± 0.02 ^b	***
Moisture content (% of dry soil mass)	40.4 ± 0.39 ^{ab}	37.9 ± 0.80^{b}	44.6 ± 0.98 ^a	***
Woody root content (% by volume of core)	1.17 ± 0.18^{b}	0.17 ± 0.09^{a}	0.00 ± 0.00^{a}	**
рН	5.37 ± 0.03 ^b	5.20 ± 0.07^{b}	5.75 ± 0.05ª	***

^{ab}Superscript letters (a, b) denote significant differences between groups

239 ^cSignificance: ** *P* < 0.01; *** *P* < 0.001; *ns* not significant

^dSamples located at 0.15, 0.3, 0.6, 1.2, 2.4, 4.8 m from boundary edge

241

243 **Table 2** Best fit models of soil organic carbon stock (kg C m⁻²), expressed on an equivalent soil

244 mass (ESM) basis, for livestock-grazed pasture adjacent to abiotic (fence and stone wall) field

245 boundaries.

Explanatory variable	F-value	d.f. ^d	Significance ^c	Effect on response variable			
ESM soil organic carbon stock model: Stone walls (AIC = 158.6 , $r^2 = 0.59$) ^{ab}							
Soil moisture	31.5	39	***	Positive association			
ESM soil organic carbon stock: Fences (AIC = 150.1, r ² = 0.74) ^{ab}							
Soil moisture	27.1	38	***	Positive association			
Bulk density	53.8	38	***	Negative association			

^aANOVA output of linear mixed effects models presented with explanatory variable information

247 ^bSamples located at 0.15, 0.3, 0.6, 1.2, 2.4, 4.8 m from boundary edge

248 ^cSignificance of differences are indicated by *** *P* < 0.001

249 ^dDegrees of freedom for ANOVA term

- **Table 3** Soil of livestock-grazed pasture, adjacent to and more distant from field boundaries.
- 252 Means and standard errors of the mean are presented alongside the significance of
- 253 differences between boundary types. Biotic (hedgerow) and abiotic (fence and stone wall)
- boundaries were analysed separately.

Variable	Hedgerows (n = 38)			Fences and stone walls (n = 16)		
Proximity to boundary	Adjacent ^a	Distant ^b	Sig. ^{cde}	Adjacent ^a	Distant ^b	Sig. ^{cde}
SOM content (% dry soil	12.4 ± 0.20	11.3 ± 0.23	^ ***	12.9 ± 0.46	13.0 ± 0.46	ns
mass; stones removed)						
SOC stock (kg C m ⁻²)	6.82 ± 0.11	6.21 ± 0.13	^ ***	7.12 ± 0.25	7.15 ± 0.25	ns
BD (g cm⁻³)	0.78 ± 0.01	0.81 ± 0.01	\downarrow **	0.81 ± 0.02	0.78 ± 0.02	ns
Moisture content (% of	40.5 ± 0.78	40.3 ± 0.57	ns	41.0 ± 0.93	41.9 ± 1.09	ns
dry soil mass)						
Stone content (% by	12.5 ± 1.6	6.03 ± 1.50	^ **	20.31 ± 2.81	8.88 ± 2.54	个 **
volume of core)						
Woody root content (% by	1.55 ± 0.25	0.41 ± 0.17	^ **	0.13 ± 0.00	0.06 ± 0.00	ns
volume of core)						
рН	5.33 ± 0.03	5.43 ± 0.06	\downarrow *	5.48 ± 0.06	5.46 ± 0.09	ns

^aAdjacent samples located at 0.15, 0.3, 0.6 and 1.2 m from boundary edge (based on zone of influence of

hedgerow, figure 2)

257 ^bDistant samples located at 2.4 and 4.8 m from boundary edge, assumed to be representative of the wider

258 pasture away from the zone of influence of the boundary

259 ^cSignificance: * *P* < 0.05; ** *P* < 0.01; *** *P* < 0.001; *ns* not significant)

 d indicates that the variable is significantly higher in the adjacent zone than the distant zone

 $261 ~~^{\rm e} \downarrow$ indicates that the variable is significantly lower in the adjacent zone



Figure 2 Best fit multi-variable model (r² = 0.74) of soil organic carbon stock (kg C m⁻²; top
0.15 m of soil), expressed on an equivalent soil mass basis, for livestock-grazed pasture
adjacent to biotic (hedgerow) field boundaries consisting of four significant explanatory
variables: a) gravimetric soil moisture content; b) soil bulk density; c) distance from
hedgerow, d) slope orientation (in relation to hedgerow). ANOVA output of linear mixedeffects models (selected on the basis of lowest AIC value) are presented with F statistics and
P values (n = 228).

271 Discussion

272 Hedgerow age

273 We found no evidence of a relationship between SOC stock and either hedgerow age category 274 or exact time since planting (from 1 year to \geq 40 years). One explanation for this is that the majority of hedgerows identified by farmers as being 1-5 years since planting contained either 275 standard trees or other remnants of a previously-removed hedgerow. The management 276 practice (incentivised by agri-environment payment schemes) had been for farmers to replant 277 278 hedgerows in the position where hedgerows had previously existed (where the soil may have 279 contained a legacy of past hedgerow effects). Another legacy of the previous hedgerows was 280 high abundance of bracken (*Pteridium aquilinum*) in the land enclosed by the boundary fences of recently (re-)planted hedgerows. Bracken is a perennial fern with an extensively 281 branched rhizome system and is associated with higher levels of SOM and SOC content than 282 283 adjacent grassland (Marrs et al., 2007). Therefore, analysis of the effects of hedgerow age on 284 adjacent soil needs to take account of the longer-term land use history. One manifestation of 285 this is that adjacent soil may be affected not only by the woody plants forming the hedgerow 286 itself, but also the other vegetation that develops in the hedgerow zone, thus forming a more 287 integrated 'biotic' linear habitat feature.

288 Hedgerow zone of influence

In this study SOC stock of adjacent pasture was broadly comparable (~7 kg C m⁻²) between biotic and abiotic field boundaries. However, a much greater effect of proximity to boundary was shown for hedgerows than for abiotic boundaries. SOC stock was greatest close to hedgerow boundaries, decreasing with perpendicular distance from the hedgerow, in line with findings from other grassland and arable systems (Follain *et al.*, 2007; D'Acunto *et al.*, 2014). SOC stock reduced markedly between 1.2 and 2.4 m from the fenced hedgerow 205 boundary (equivalent to ca. 2.2 and 3.4 m from the base of the hedgerow itself, indicating a 206 slightly narrower range of influence of the hedgerow on SOC storage dynamics than the ~4 m 207 previously identified (D'Acunto *et al.*, 2014; Van Vooren *et al.*, 2017). In contrast, there was 208 no evidence of greater SOC stock adjacent to abiotic boundaries than in more distant pasture.

299 In this study soil BD was lower adjacent to fences than stone walls, with hedgerows 300 intermediate between the two. This can be attributed to two contrasting mechanisms. The increase in organic matter inputs associated with hedgerows may reduce soil BD, by 301 increasing soil porosity and aggregate structures, or diluting the mineral soil component 302 (Holland, 2004). However, where landscape features including hedgerows or walls (but not 303 fences) provide shelter and encourage livestock to congregate during adverse weather 304 305 conditions this is expected to increase soil BD (Abdalla et al., 2018), albeit asymmetrically with 306 greater compaction on the leeward side.

307 *Hedgerows and topography*

308 The effect of contour hedgerows on reducing soil erosion is well established, leading to accumulation of eroded soil upslope of hedgerows, whereas the land immediately downslope 309 of hedgerows is often a zone of net erosive soil loss (Follain *et al.*, 2007). As this eroded soil 310 is generally topsoil it can have a high SOC stock, and this may enhance the SOC content of soil 311 left in situ, measured at 0-0.15 m depth in the present study, relative to soil downslope of the 312 hedgerow. It is also possible that the roots of hedgerow woody plants grow preferentially 313 314 upslope (Caubel-Forget et al., 2001) to increase plant stability, access soil enriched by run-off and sediment, and avoid water-logged or compacted soil (Jackson et al., 2000), which would 315

lead to higher rates of fine-root turnover and associated SOC storage upslope than downslopeof contour-planted hedgerows.

318 *Relationship between SOC and multiple variables*

319 Two environmental variables, soil moisture and soil BD, were clearly associated with SOC stock for pasture adjacent to both biotic and abiotic boundaries, in combination explaining 320 more than half of the variation in SOC stock, with these relationships likely to extend across 321 the whole pasture. A positive association between SOC and soil moisture is well recognised, 322 particularly in clay-rich grasslands (Jobbagy & Jackson, 2000) and a negative relationship 323 324 between SOC content (or SOC stock in this study as expressed on an equivalent soil mass 325 basis) and BD is well established and recently confirmed in a silvopastoral setting (Upson et al., 2016). In our study, three quarters of the variation in SOC stock in pasture adjacent to 326 hedgerows was explained by a model that combined four variables [distance from hedgerow, 327 slope orientation (relative to hedgerow), soil moisture and BD], with no evidence of a 328 329 relationship between SOC stock and other measured variables (i.e. hedgerow age, 330 management, dominant woody plant species or mapped soil type). One possible explanation, for the quarter of variation in SOC stock unaccounted for, is the effect of the functional 331 composition of hedgerow woody plants and associated herbaceous plants (especially P. 332 *aquilinum*), and pasture grasses and forbs, not measured directly in this study. 333

334 Conclusions

This study aimed to quantify the influence of hedgerows on SOC storage in adjacent pasture, and is of particular relevance to upland farming systems in Wales. SOC stock in adjacent pasture was comparable (~7 kg C m⁻²) between hedgerow and abiotic field boundaries. Our first hypothesis, that SOC stock in pasture adjacent to hedgerows would increase with age

339 (years since planting) was rejected, largely due to the effect of management legacy on 340 boundary location. Our second hypothesis, that SOC stock would decrease with distance from 341 hedgerow was accepted with pasture SOC stock (to 0.15 m depth) 15% greater within 2 m of the hedgerow boundary than further into the pasture, with no such effect for abiotic 342 343 boundaries. Our third hypothesis, that SOC stock would be greater upslope of hedgerows than downslope was also accepted. However, this multi-farm study demonstrated that the 344 influence of hedgerows on the ecosystem function of SOC storage in livestock-grazed pasture 345 346 is still small in comparison with the dominant influence of BD and soil moisture.

347 Acknowledgements

All authors acknowledge the financial support provided by the Welsh Government and Higher Education Funding Council for Wales through the Sêr Cymru National Research Network for Low Carbon, Energy and Environment. Thanks are also due to the National Trust and the Fferm Ifan group that allowed access to their land.

352 **References**

- Abdalla, M., Hastings, A., Chadwick, D.R., Jones, D.L., Evans, C.D., Jones, M.B., Rees, R.M. & Smith, P. 2018. Critical review of the impacts of grazing intensity on soil organic carbon storage and other soil quality indicators in extensively managed grasslands. *Agriculture, Ecosystems and Environment*, **253**, 62-81.
- 357 Agricultural Land Classification of England and Wales. 2018. Available at: 358 https://beta.gov.wales/sites/default/files/publications/2018-02/agricultural-land-
- 359 <u>classification-frequently-asked-questions.pdf</u>.
- Axe, M.S., Grange, I.D. & Conway, J.S. 2017. Carbon storage in hedge biomass A case study
- of actively managed hedges in England. *Agriculture, Ecosystems and Environment*, **250**, 8188.
- Baah-Acheamfour, M., Chang, S.X., Carlyle, C.N. & Bork, E.W. 2015. Carbon pool size and stability are affected by trees and grassland cover types within agroforestry systems of western Canada. *Agriculture, Ecosystems and Environment*, **213**, 105-113.
- Ball, D.F. 1964. Loss-on-ignition as an estimate of organic matter and organic carbon in noncalcareous soils. *Journal of Soil Science*, **15**, 84-92.
- 368 Bardgett, R.D., Wardle, D.A. 2010. Aboveground-Belowground Linkages. Biotic Interactions,
- 369 *Ecosystem Processes and Global Change*. Oxford University Press, Oxford, UK.
- Baudrey, J., Bunce, R.G.H. & Burel, F. 2000. Hedgerows: An international perspective on their
- origin, function and management. *Journal of Environmental Management*, **60**, 7-22.

- Benham, S.E., Vanguelova, E.I. & Pitman, R.M. 2012. Short and long term changes in carbon,
 nitrogen and acidity in the forest soils under oak at the Alice Holt Environmental Change
 Network site. *Science of the Total Environment*, **421-422**, 82-93.
- Bullock, J.M., Jefferson, R.G., Blackstock, T.H., Pakeman, R.J., Emmett, B.A., Pywell, R.J.,
- 376 Grime, J.P. & Silvertown, J. 2011. Semi-natural Grasslands. In: The UK National Ecosystem
- 377 Assessment Technical Report. UK National Ecosystem Assessment, UNEP-WCMC, Cambridge.
- 378 Cardinael, R., Umulisa, V., Toudert, A., Olivier, A., Bockel, L. & Bernoux, M. 2018. Revisiting
- 379 IPCC Tier 1 coefficients for soil organic biomass carbon storage in agroforestry systems.
- 380 Environmental Research Letters, **13**, 124020.
- 381 Caubel-Forget, V., Grimaldi, C. & Rouault, F. 2001. Contrasted dynamics of nitrate and chloride
- in groundwater submitted to the influence of a hedge. *Comptes rendus de l'Académie des Sciences de Paris*, **332**, 107-113.
- Crossland, M. 2015. The carbon sequestration potential of hedges managed for woodfuel.
 The Organic Research Centre. Available at:
 http://www.organicresearchcentre.com/manage/authincludes/article_uploads/project_out
 puts/TWECOM%20ORC%20Carbon%20report%20v1.0.pdf.
- D'Acunto, L., Semmartin, M. & Ghersa, C.M. 2014. Uncropped field margins to mitigate soil
 carbon losses in agricultural landscapes. *Agriculture, Ecosystems and Environment*, **183**, 6068.
- 391 Don, A., Schumacher, J. & Friebauer, A. 2011. Impact of tropical land-use change on soil
 392 organic carbon stocks a meta-analysis. *Global Change Biology*, **17**, 1658-1670.

Emmett, B.A., Reynolds, B., Chamberlain, P.M., Rowe, E., Spurgeon, D., Brittain, S.A., ...
Woods, C. 2010. *Countryside Survey: Soils Report from 2007*. NERC/Centre for Ecology and
Hydrology 192pp. (CS Technical Report no. 9/07, CEH project number: C03259).

Follain, S., Walter, C., Legout, A., Lemercier, B. & Dutin, G. 2007. Induced effects of hedgerow

networks on soil organic carbon storage within an agricultural landscape. *Geoderma*, **142**, 8095.

Golluscio, R.A., Austin, A.T., García Martínez, G.C., Gonzalez-Polo, M., Sala, O.E. & Jackson,
R.B. 2009. Sheep grazing decreases organic carbon and nitrogen pools in the Patagonia
Steppe: combination of direct and indirect effects. *Ecosystems*, **12**, 686-697.

402 Grömping, U. 2006. Relative Importance for Linear Regression in R: The Package relaimpo.
403 *Journal of Statistical Software*, **17**, 1–27.

Hallett, S.H., Sakrabani, R., Keay, C.A. & Hannam, J.A. 2017. Developments in land information
systems: examples demonstrating land resource management capabilities and options. *Soil Use and Management*, **33**, 514-529.

Holden, J., Grayson, R.P., Berdeni, D., Bird, S., Chapman, P.J., Edmondson, J.L., Firbank, L.G.,
Helgason, T., Hodson, M.E., Hunt, S.F.P., Jones, D.T., Lappage, M.G., Marshall-Harries, E.,
Nelson, M., Prendergast-Miller, M., Shaw, H., Wade, R.N. & Leake, J.R. 2019. The role of
hedgerows in soil functioning within agricultural landscapes. *Agriculture, Ecosystems and Environment*, **273**, 1-12.

Holland, J.M. 2004. The environmental consequences of adopting conservation tillage in
Europe: reviewing the evidence. *Agriculture, Ecosystems and Environment*, **103**, 1-25.

- Jackson, R.B., Sperry, J.S. & Dawson, T.E. 2000. Root water uptake and transport: using
 physiological processes in global predictions. *Trends in Plant Science*, 5, 1360-1385.
- Jobbagy, E.G. & Jackson, R.B. 2000. The vertical distribution of soil organic carbon and its
 relation to climate and vegetation. *Ecological Applications*, **10**, 423-436.
- Kemp, D.R. & Michalk, D.L. 2007. Towards sustainable grassland and livestock management. *Journal of Agricultural Science*, **145**, 543-564.
- Lange, M., Eisenhauer, N., Sierra, C.A., Bessler, H., Engels, C., Griffiths, R.I., ... Gleixner, G.
 2015. Plant diversity increases soil microbial activity and soil carbon storage. *Nature*
- 422 *Communications*, 6:6707.
- Lee, J., Hopmans, J.W., Rolston, D.E., Baer, S.G. & Six, J. 2009. Determining soil carbon stock
 changes: Simple bulk density corrections fail. *Agriculture, Ecosystems and Environment*, **134**,
 251-256.
- Marrs, R.H., Galtress, K., Tong, C., Cox, E.S., Blackbird, S.J., Heyes, T.J., Pakeman, R.J. & Le Duc,
 M.G. 2007. Competing conservation goals, biodiversity or ecosystem services: Element losses
 and species recruitment in a managed moorland-bracken model system. *Journal of Environmental Management*, **85**, 1034-1047.
- 430 Natural England, 2008. Soil Texture. Natural England Technical Information Note TIN037.
- 431 Available at: http://publications.naturalengland.org.uk/publication/32016.
- 432 Ostle, N.J., Levy, P.E., Evans, C.D. & Smith, P. 2009. UK land use and soil carbon sequestration.
 433 Land Use Policy, 26S, S274-S283.
- 434 Peter, I. & Lehmann, J. 2000. Pruning effects on root distribution and nutrient dynamics in an
- 435 acacia hedgerow planting in northern Kenya. *Agroforestry Systems*, **50**, 59-75.

R Core Team, 2018. R: A language and environment for statistical computing. R Foundation
for Statistical Computing, Vienna, Austria. Available at: https://www.R-project.org/.

Rowell, D. 1994. Soil Science: Methods and Applications. Longman UK Ltd. Harlow, Essex, UK.
Available at: http://doi.org/10.1002/jsfa.2740660423.

Scholefield, P., Morton, D., Rowland, C., Henrys, P., Howard, D. & Norton, L. 2016. A model of
the extent and distribution of woody linear features in rural Great Britain. *Ecology and Evolution*, 6, 8893-8902.

443 Sokol, N.W. & Bradford, M.A. 2019. Microbial formation of stable soil carbon is more efficient

from belowground than aboveground input. *Nature Geoscience*, **12**, 46-53.

Soussana, J.F., Tallec, T. & Blanfort, V. 2010. Mitigating the greenhouse gas balance of
ruminant production systems through carbon sequestration in grasslands. *Animal*, 4, 334-350.

447 Upson, M.A., Burgess, P.J. & Morison, J.I.L. 2016. Soil carbon changes after establishing
448 woodland and agroforestry trees in a grazed pasture. *Geoderma*, **283**, 10-20.

449 Van Vooren, L., Reubens, B., Broekx, S., De Frenne, P., Nelissen, V., Pardon, P. & Verheyen, K.

450 2017. Ecosystem service delivery of agri-environment measures: A synthesis for hedgerows

451 and grass strips on arable land. *Agriculture, Ecosystems and Environment*, **244**, 32-51.

452 Wolton, R., Pollard, K., Goodwin, A. & Norton, L. 2014. *Regulatory services delivered by* 453 *hedges: The evidence base*. Report of Defra project LM0106. 99pp.

Zuur, A. F., Ieno, E. N., Walker, N. J., Saveliev, A. A., & Smith, G. H. 2009. *Mixed Effects Models and Extensions in Ecology with R*. Springer-Verlag, New York.