

Occurrence of legacy and emerging organic pollutants in whitemouth croakers from Southeastern Brazil

Pizzochero, Ana Carolina; de la Torre, Adrian; Sanz, Paloma; Navarro, Irene; Michel, Loic; Lepoint, Giles; Das, Krishna ; Schnitzler, Joseph; Chenery, Simon ; McCarthy, Ian; Malm, Olaf; Dorneles, Paulo; Martinez, Maria Angeles

Science of the Total Environment

DOI:

[10.1016/j.scitotenv.2019.05.213](https://doi.org/10.1016/j.scitotenv.2019.05.213)

Published: 10/09/2019

Peer reviewed version

[Cyswllt i'r cyhoeddiad / Link to publication](#)

Dyfyniad o'r fersiwn a gyhoeddwyd / Citation for published version (APA):

Pizzochero, A. C., de la Torre, A., Sanz, P., Navarro, I., Michel, L., Lepoint, G., Das, K., Schnitzler, J., Chenery, S., McCarthy, I., Malm, O., Dorneles, P., & Martinez, M. A. (2019). Occurrence of legacy and emerging organic pollutants in whitemouth croakers from Southeastern Brazil. *Science of the Total Environment*, 682, 719-728. <https://doi.org/10.1016/j.scitotenv.2019.05.213>

Hawliau Cyffredinol / General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Title: Occurrence of legacy and emerging organic pollutants in whitemouth croakers from
Southeastern Brazil

Authors: Ana Carolina Pizzochero^{1*}, Adrián de la Torre², Paloma Sanz², Irene Navarro², Loïc
N. Michel^{3**}, Gilles Lepoint³, Krishna Das³, Joseph G. Schnitzler⁴, Simon R. Chenery⁵, Ian D.
McCarthy⁶, Olaf Malm¹, Paulo R. Dorneles¹, María Ángeles Martínez ²

¹ Radioisotope Laboratory, Biophysics Institute, Federal University of Rio de Janeiro (UFRJ),
Brazil

² Persistent Organic Pollutants Group, Environmental Department, CIEMAT, Madrid, Spain

³ Freshwater and Oceanic Sciences Unit of reSearch (FOCUS), Laboratory of Oceanology,
University of Liege, Belgium

⁴ Institute for Terrestrial and Aquatic Wildlife Research (ITAW) University of Veterinary
Medicine Hannover.

⁵ British Geological Survey (BGS), Kingsley Dunham Centre, Keyworth, Nottingham, United
Kingdom

⁶ School of Ocean Sciences, Bangor University (BU), Wales, United Kingdom

*Corresponding author:

Ana Carolina Pizzochero

Universidade Federal do Rio de Janeiro (UFRJ)

Centro de Ciências da Saúde (CCS)

Instituto de Biofísica Carlos Chagas Filho (IBCCF)

Laboratório de Radioisótopos Eduardo Penna Franca (LREPF)

Avenida Carlos Chagas Filho, 373 (Edifício do CCS), sala G0-62

Cidade Universitária, 21941-902, Rio de Janeiro, RJ, Brazil

Tel.: +55 21 25615339, Fax: +55 21 22808193

E-mail: pizzochero@biof.ufrj.br; pizzocheroac@gmail.com

** Current address: Ifremer, Centre de Bretagne, REM/EEP, Laboratoire Environnement
Profond, F-29280 Plouzané, France.

Abstract


The whitemouth croaker (*Micropogonias furnieri*) is one of the most commercially important species along the Atlantic coast of South America. Moreover, some of its biological traits (long life span, inshore feeding, high trophic position) make this species a suitable sentinel of coastal pollution. Here, we investigated contamination by multiple legacy and emerging organic pollutants, such as brominated and chlorinated flame retardants, polychlorinated dibenzo-*p*-dioxins (PCDDs) and polychlorinated dibenzofurans (PCDFs), in whitemouth croakers from two estuaries (Guanabara and Sepetiba Bays) located in industrialized and urbanized areas in Rio de Janeiro State, Southeastern Brazil. Furthermore, we assessed how biological and ecological features could explain the observed contamination patterns. Regarding brominated flame retardants, concentrations of polybrominated diphenyl ethers (PBDEs) varied from 7.6 to 879.7 pg g⁻¹ wet weight (w.w.), with high contribution of tetra-, penta-, hexa- and deca-BDEs. The sum of chlorinated flame retardants (dechlorane-related compounds, Σ DRC) ranged from <LOD to 41.1 pg g⁻¹ w.w., mostly represented by Dechlorane 603 and Dechlorane Plus (DP). Concentrations of PCDDs and PCDFs varied from <LOD to 1.7 pg g⁻¹ w.w., while the Toxic Equivalent (TEQ-PCDD/Fs) levels ranged from 0.1 to 0.2 pg g⁻¹ w.w. Positive correlations between $\delta^{15}\text{N}$ and concentrations of tri-, tetra- and penta-BDEs, as well as Σ DRC, DP and *anti*-DP isomers suggested that ecological factors (namely biomagnification along the food web) influence contamination of whitemouth croakers in the estuaries studied. Moreover, the sum of PBDEs (Σ PBDE), tri- and tetra-BDEs concentrations were negatively correlated with fish size, suggesting that depuration by fishes and/or habitat shift throughout the whitemouth croaker's life cycle might also influence concentrations. Overall, our study emphasized the need for further investigations to help understand the complex patterns of bioaccumulation and biomagnification that seem to exist in Southeastern Brazil.

Keywords: *Micropogonias furnieri*, Brazil; PBDEs; Dechloranes; PCDD/Fs; Stable Isotopes.

Highlights

- Analysis suggested the presence of PBDE commercial mixtures in whitemouth croakers
- First assessment of emerging flame retardants in fish from Southwest Atlantic Ocean
- Dec 603 and DP were the predominant DRCs, followed by Mirex, Dec 602 and CP

Graphical Abstract



1. Introduction

Persistent Organic pollutants (POPs) comprise a wide range of chemicals that have received considerable attention due to their persistence in the environment, long-range transport and toxic properties (Jones and De Voogt, 1999; Walker *et al.*, 2012). Restricted or banned POPs – known as legacy POPs – are regulated by the Stockholm Convention, and include substances classified as unintentional products [*i.e.* polychlorinated dibenzo-*p*-dioxins (PCDDs) and polychlorinated dibenzofurans (PCDFs)], pesticides, and industrial chemicals [*i.e.* flame retardants (FRs), such as polybrominated diphenyl ethers (PBDEs) commercial mixtures] (UNEP, 2017). On the other hand, the term emerging pollutants refers to replacement substances for the legacy chemicals, which have been recently observed in the environment. Among the emerging pollutants, the category of FRs stand out, including brominated [*i.e.* pentabromoethylbenzene (PBEB) and 1,2-bis(2,4,6-tribromophenoxy)ethane (BTBPE)] and chlorinated flame retardants, such as the dechlorane-related compounds (DRCs) [*i.e.* Dechlorane 602 (Dec 602; CAS# 31107–44–5), 603 (Dec 603, CAS# 13560–92–4), 604 (Dec 604; CAS# 34571–16–9) and Dechlorane Plus (Dec 605 or DP, CAS# 13560–89–9) that are used as substitutes to the banned Mirex. Legacy and emerging pollutants are prone to accumulate in organisms and biomagnify throughout food webs due to their persistent and hydrophobic properties (Kelly *et al.*, 2007; Walters *et al.*, 2016; Navarro *et al.*, 2016, 2017 and 2018). Therefore, marine organisms provide opportunities to act as monitors of their environment, as levels and profiles of these contaminants can serve as intrinsic markers, reflecting the ecosystem conditions under which biota live and feed (Alonso *et al.*, 2012; Chouvelon *et al.*, 2014; 2017).

In Brazil, the whitemouth croaker, *Micropogonias furnieri* (Desmarest, 1823) (Perciformes, Sciaenidae), has been recommended as a good indicator of environmental contamination (Dorneles *et al.*, 2016) due to its distribution along the coastal waters of the western Atlantic

Ocean, to its longevity (~35 years), and high trophic position in estuarine ecosystems (Bisi *et al.*, 2012; Pizzochero *et al.*, 2018; Vazzoler, 1991). Additionally, this species constitutes a commercially-important resource in coastal demersal fisheries along the Atlantic coast of South America (FAO, 2018; Haimovici *et al.*, 2016). In this context, investigations on legacy and emerging pollutants in this species would not only provide information on the contaminants that are spreading through marine food webs in Brazilian coastal waters, but would also be relevant in public and human health assessment as ingestion of seafood constitutes the principal source of human exposure to POPs (Cruz *et al.*, 2015; Sidhu, 2003).

In the present study, concentrations of legacy and emerging POPs were measured in white muscle samples of whitemouth croakers from Rio de Janeiro state, Southeastern Brazil. We aimed to investigate whitemouth croaker exposure to POPs and its relation with stable isotopes ratios of carbon ($\delta^{13}\text{C}$), nitrogen ($\delta^{15}\text{N}$) and sulfur ($\delta^{34}\text{S}$), in order to provide a more comprehensive view of their potential use as sentinels. Combining pollutant determination with measurements of ecological tracers such as stable isotopes has been shown to be useful for better understanding sources, pathways, and the trophic flow of toxicants (Bisi *et al.*, 2012; Chouvelon *et al.*, 2014; 2017). To the authors' knowledge, this is the first study to determine emerging pollutants [brominated (HBB – hexabromobenzene, BB-153 – 2,2',4,4',5,5'-hexabromobiphenyl, PBEB and BTBPE) and chlorinated (Dec 602, Dec 603, Dec 604, DP and Chlordene Plus) flame retardants] in fish from the southwest Atlantic Ocean.

2. Materials and Methods

2.1 Study area and sample collection

Guanabara Bay and Sepetiba Bay are two important fishing areas in Rio de Janeiro state (RJ), in Southeast Brazil (Fig. 1). Located in the metropolitan area of the Rio de Janeiro city, Guanabara Bay (22°24' and 22°57' S / 43°33' and 43°00' W, 328 km²) is the most

anthropogenically-disturbed area along the Brazilian coastline (Dorneles *et al.*, 2008a, 2008b, 2013). This estuary is under the direct influence of approximately 11 million people living in its surroundings (IBGE, 2016), receiving sewage, industrial waste and consequently many contaminants that are transported along its drainage basin, which contains more than 12,000 industries (Baptista-Neto *et al.*, 2016; Kjerfve *et al.*, 1997). Sepetiba Bay (22°55' and 23° 05'S/ 43°40' and 44°40'W, 450 km²) has also been severely impacted by anthropogenic activities over the past 40 years. Its drainage basin is surrounded by a population of about 2 million people and over 400 industries, including metallurgical, petrochemical and pyrometallurgical smelters (IBGE, 2016; Molisani *et al.*, 2004). Twenty whitemouth croaker (*Micropogonias furnieri*) specimens were obtained from commercial fishery landings in Guanabara ($n = 14$) and Sepetiba ($n = 6$) Bays in the 2014 austral winter (dry season). Each fish was weighed, measured and dissected. Dorsal white muscle samples were wrapped in individual aluminium foil and kept frozen (-20 °C) until being oven-dried at 60°C to constant weight (> 72h) prior to analysis. Biological parameters [size, mass and lipid content (%)] of the specimens analyzed in the present study are presented in Table 1.

2.2 Chemicals and reagents

Complete details on the standards used are presented in Table S1 (Supplementary data). Dec 602 (95% purity), Dec 603 (98%), and Dec 604 (98%) were purchased from Toronto Research Chemical Inc. (Toronto, ON, Canada). Chlordene Plus (CP; CAS# 13560-91-3) and DP (*syn*-DP and *anti*-DP standards) were obtained from Wellington Laboratories Inc. (Guelph, ON, Canada). Mirex (CAS# 2385-85-5) was purchased from Cambridge Isotope Laboratories Inc. (Andover, MA). For brominated flame retardant (BFR) determinations, BFR-LCS (containing 14 ¹³C₁₂-PBDEs, ¹³C₆-HBB, ¹³C₁₂-BB-153 and ¹³C₆-BTBPE), BFR-ISS (containing 4 ¹³C₁₂-PBDEs) and BFR-CVS (five individual calibration solutions containing among others

35 $^{12}\text{C}_{12}$ -PBDEs, 20 $^{13}\text{C}_{12}$ -PBDEs, $^{13}\text{C}_6$ and ^{12}C -HBB, $^{13}\text{C}_{12}$ — and ^{12}C -BB-153, $^{13}\text{C}_6$ — and
 ^{12}C -BTBPE) were obtained from Wellington laboratories Inc. (Guelph, ON, Canada). For
PCDD and PCDF determinations, EPA-1613LCS (containing 15 $^{13}\text{C}_{12}$ -PCDD/Fs), EPA-
1613ISS (containing 2 $^{13}\text{C}_{12}$ -PCDDs) and EPA-1613CVS (five individual calibration solutions
containing among others 17 $^{12}\text{C}_{12}$ -PCDD/Fs and 17 $^{13}\text{C}_{12}$ -PCDD/Fs) were obtained from
Wellington laboratories Inc. (Guelph, ON, Canada). The other chemicals used, *i.e.* anhydrous
sodium sulphate, silica, sulphuric acid (95–97%) and solvents (hexane, dichloromethane, ethyl
acetate and toluene) for organic trace analysis, were all obtained from Merck (Darmstadt,
Germany).

2.3 Sample preparation and chemical analysis

The analytical methods used are described in detail elsewhere (de la Torre *et al.*, 2011, 2012).
Extractions were performed with an Accelerated Solvent Extraction system (ASE 100, Dionex,
Sunnyvale, CA, USA) using a mixture of hexane:dichloromethane (1:1 v/v) as solvent, at 100
°C, 1500 psi, 90% flush volume and three static cycles (10 min time each; 70 mL total volume).
Prior to the extraction step, the samples were spiked with $^{13}\text{C}_{12}$ labeled surrogate standards (see
Table S1). The oven-dried dorsal white muscle samples, ranging from 1 to 8 g, were
homogenized with 15 g of anhydrous sodium sulphate and introduced into a 30 mL cell
previously loaded by inserting two cellulose filters followed by 2 g of anhydrous sodium
sulphate. The resulting extract of each sample was evaporated to constant weight for
gravimetric lipid determination and then re-dissolved in hexane. Sample purification consisted
of two steps: a liquid extraction with 100 mL of hexane and 50 mL of concentrated sulfuric
acid to remove organic matter from the extracts, followed by the transfer of the organic phase
to an open glass column with 15 g acid silica modified with 44% sulphuric acid, covered with
1 g anhydrous sodium sulphate and eluted with 150 mL of hexane. The cleaned extracts were

concentrated to approximately 1 mL. The fractionation step was performed in an automated purification Power Prep™ System (FMS, Inc., USA) including acidic silica gel, basic alumina and carbon columns. Two fractions were obtained: Fraction A containing PCDD/Fs and Fraction B containing BFRs and DRCs. Both fractions were concentrated to approximately 1 mL under a flow of nitrogen using a Turbo Vap II evaporator (Vertex, Technics, Madrid, Spain), and spiked with the internal standard spiking solutions (see Table S1).

The instrumental analysis was conducted using high resolution gas chromatography coupled with high resolution mass spectrometry (HRGC–HRMS; Agilent GC 6890N connected to a Waters Micromass AutoSpec Ultima NT) at 10,000 resolving power (10% valley) and working in selected ion monitoring (SIM) mode. The GC column used for PCDD/F determination was 60 m x 0.25 mm x 0.25 µm film thickness (DB-5MS, J&W Scientific), while a short and narrow column [15 m x 0.25 mm x 0.10 µm film thickness; DB-5MS (J&W Scientific)] was used for BFR and DRC determinations. Instrument operating conditions were as described in de la Torre *et al.* (2011, 2012).

Quantification was carried out using the isotopic dilution method (US EPA, 1994). Three criteria were used to ensure the correct identification and quantification of analytes: i) ± 2 s retention time between the analyte and the standard, ii) the ratio of quantifier and qualifier ions had to be within $\pm 15\%$ of the theoretical values and iii) a signal to noise ratio greater than three. Recoveries for DP ($^{13}\text{C}_{10}\text{-syn-DP}$ and $^{13}\text{C}_{10}\text{-anti-DP}$), $^{13}\text{C}_{12}\text{-PCDD/Fs}$, and $^{13}\text{C}_{12}\text{-PBDEs}$ in this study were $81 \pm 10\%$, $79 \pm 14\%$, $82 \pm 9\%$ (mean \pm SD), respectively. The limits of detection (LODs) and quantification (LOQs) of the method were calculated as the concentration corresponding to a signal-to-noise ratio of 3 and 10 respectively (see Table S2). Procedural blanks were processed and analyzed with every batch of samples under the same conditions. In addition, instrumental blanks consisting of nonane were run before each sample injection to check for memory effects and contamination from the gas chromatograph system.

Concentrations in instrumental and procedural blanks were below LOD. For statistical descriptive calculations, samples with concentrations below LODs were considered as zero. However, for PCDD/F World Health Organization 2005 Toxic Equivalent (TEQ; Van den Berg *et al.*, 2006) calculations, not detected values were replaced by LODs.

2.4 Stable isotope measurements

The data on stable isotope ratios of carbon ($\delta^{13}\text{C}$), nitrogen ($\delta^{15}\text{N}$) and sulfur ($\delta^{34}\text{S}$) in muscle samples of whitemouth croakers from Guanabara Bay were extracted from Pizzochero *et al.* (2018). This dataset was supplemented with specimens from Sepetiba Bay. Oven-dried dorsal white muscle samples were ground into powder using mortar and pestle. Approximately 4 mg of dry powdered material were analysed. Measurements of stable isotope ratios were performed via continuous flow - elemental analysis - isotope ratio mass spectrometry (CF-EA-IRMS) at the Laboratory for Oceanology, University of Liege (Belgium), using a vario MICRO cube C-N-S elemental analyzer (Elementar Analysensysteme GMBH, Hanau, Germany) coupled to an IsoPrime100 isotope ratio mass spectrometer (Isoprime, Chesham, United Kingdom). Isotopic ratios were expressed using the widespread δ notation (Coplen, 2011), in ‰ and relative to the international references [Vienna Pee Dee Belemnite (for carbon), Atmospheric Air (for nitrogen) and Vienna Canyon Diablo Troilite (for sulfur)]. IAEA (International Atomic Energy Agency, Vienna, Austria) certified reference materials sucrose (IAEA-C-6; $\delta^{13}\text{C} = -10.8 \pm 0.5\text{‰}$; mean \pm SD), ammonium sulfate (IAEA-N-2; $\delta^{15}\text{N} = 20.3 \pm 0.2\text{‰}$) and silver sulfide (IAEA-S-1; $\delta^{34}\text{S} = -0.3\text{‰}$) were used as primary analytical standards. Sulfanilic acid (Sigma-Aldrich; $\delta^{13}\text{C} = -25.6 \pm 0.4\text{‰}$; $\delta^{15}\text{N} = -0.13 \pm 0.4\text{‰}$; $\delta^{34}\text{S} = 5.9 \pm 0.5\text{‰}$) was used as secondary analytical standard. Standard deviations on multi-batch replicate measurements of secondary and internal lab standards (animal muscle tissue) analyzed interspersed with samples (one

replicate of each standard every 15 analyses) were 0.2‰ for both $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ and 0.4‰ for $\delta^{34}\text{S}$ respectively.

2.5 Data analysis

Each dataset (i.e., each pollutant concentration, stable isotope ratio or biological parameter) was tested for normality using the Shapiro–Wilk’s W test, and non-parametric tests were applied since most datasets did not follow a Gaussian distribution. All data are presented as mean \pm standard deviation.

To test whether fishes from Sepetiba and Guanabara bays presented differences in their contamination pattern, we used one-way ANOSIM (ANalysis Of SIMilarity) to compare pollutant concentrations in fishes from the two sites. ANOSIM is a non-parametric, multivariate procedure that uses ranked dissimilarities between samples (here, fishes) to investigate the presence of significant differences between several groups. ANOSIM is permutation-based and assumption-free, which makes it a generally applicable way to test the hypothesis that one response variable (here, the sampling site) is linked with significant differences in a multivariate dataset (here, the pollutant concentrations; Clarke and Warwick, 2001). All compounds found in at least one individual fish were used as input variables. The resemblance matrix was built using Bray-Curtis similarity coefficients, and the number of permutations was set to 9999. The ANOSIM analyses were conducted using PAST 3.20 (Hammer *et al.*, 2001).

To highlight potential relationships (or the absence thereof) between pollutant concentrations and stable isotope ratios and biological parameters [length, mass and lipid content (%)], we performed correlation analyses. 10 pollutants or pollutant classes were retained for correlation analysis: tri-, tetra, penta, hexa and hepta-BDEs (summed concentrations of all PBDE congeners with 3, 4, 5, 6 and 7 bromine atoms, respectively), Σ PBDE (summed concentrations of all polybrominated compounds), Σ DRC (summed concentrations of all dechlorane-related

compounds), Dechlorane 603, anti-Dechlorane Plus, and total Dechlorane Plus (sum of *anti*- and *syn*-Dechlorane Plus concentrations). Correlation analyses between all these pollutants or pollutant categories (that could be quantified in more than 50% of the fishes) and each stable isotope ratio ($\delta^{13}\text{C}$, $\delta^{15}\text{N}$ and $\delta^{34}\text{S}$) and biological parameter (mass, size and relative lipid content) were performed. Since data did not follow a Gaussian distribution, Spearman's rank correlation coefficients (r_s) were used. Since the ANOSIM test did not reveal significant inter-site difference in fish contamination patterns (see below), correlations were performed grouping all fishes from both estuaries. Analyses were conducted using Prism 6.07 (GraphPad Software, La Jolla, U.S.A.), and the level of significance (α) was set to 0.05. To make visualization of these numerous correlations easier, results were synthesized in a correlation matrix (Fig. 3). This correlation matrix was generated using R 3.5.1 (R Core Team, 2018) and the corrplot package v. 0.84 (Wei & Simko, 2017).

3. Results

3.1 Organic pollutant levels

The sums of PCDD/Fs ($\Sigma\text{PCDD/F}$), PBDEs (ΣPBDE) and DRCs (ΣDRC) in each individual sample are listed in Table 1. Additionally, detailed concentrations in wet weight (w.w.) and lipid weight (l.w.) for all target analytes are reported in the Supplementary data (Tables S3-S9). Research budget only allowed PCDD/F investigations in 10 specimens from Guanabana Bay (Tables S3 and S4). Compounds 2,3,7,8-Tetra-CDD, 1,2,3,7,8-Penta-CDD and 1,2,3,4,7,8-Hexa-CDD could not be detected in any sample. Concentrations of PCDDs were greater than those of PCDFs in 90% of the individuals, being OCDD predominant PCDD/F in 80% of the samples, with concentrations ranging from <LOD to $1.25 \text{ pg g}^{-1} \text{ w.w}$ (Table S3). Calculated TEQ values ranged from 0.1 to $0.2 \text{ pg TEQ g}^{-1} \text{ w.w}$. (Tables 1 and S5). For PBDEs, from 35 congeners evaluated only ten presented quantification frequencies >50%: BDE-47 and BDE-

100 (100% of samples; Tables S6 and S7); BDE-154, BDE-49 & 71 and BDE-153 (95%); BDE-66 (85%); BDE-99 (75%); BDE-28 (65%); and BDE-183 (60%). The PBDE profiles observed in whitemouth croaker (Fig. 2A) indicated a high contribution of tetra-BDE (51 ± 20 %), penta-BDE (15 ± 6 %), and hexa-BDE (12 ± 7 %). The most common compounds represented were the tetra congener BDE-47 (predominant congener in 15 samples), the penta congeners BDE-100 and BDE-99 and the hexa congeners BDE-153 and BDE-154. Although BDE-209 (decaBDE) was only quantified in 40% of the fish samples (Tables S6 and S7), its contribution to total PBDE content achieved levels up to 78% (Fig. 2A). Quantification frequencies decreased for emerging brominated pollutants. PBEB, BB-153, and BTBPE levels were below LOD in all samples, while HBB was only found in one sample from Guanabara Bay (Gb#3) with a value of $0.1 \text{ pg g}^{-1} \text{ w.w.}$ (Table S6).

Amongst the dechlorane-related compounds (DRCs), Dec 604 could not be detected in any sample (Table S8). Detection frequencies of each DRC ranked as follows: Dec 603 (95% of samples), DP [65% (anti-DP: 65% and syn-DP: 45%)], Dec 602 (55%), mirex (40%) and CP (10%). Percentage contributions from individual dechlorane compounds to ΣDRC followed the same order as seen in their frequency of detection, ranking as follows: Dec 603 (61 ± 26 %), DP (25 ± 29 %), Dec 602 (7 ± 8 %), mirex (6 ± 11 %) and CP (0.2 ± 0.6 %) (Fig. 2B). Considering the samples in which any DP isomer was quantified ($n = 13$), *anti*-DP was predominant in 85% of them. The relative concentrations of the DP isomers in whitemouth croaker was explored using the values of the *anti*-DP fractions (f_{anti}), calculated as the concentration of the *anti*-DP divided by the sum of *syn*- and *anti*-DP concentrations. The f_{anti} values obtained ranged from 0.4 to 1 (0.7 ± 0.2 ; mean \pm SD) (Table 1).

3.2 Relationships between pollutant levels, stable isotopes and biological parameters

The stable isotope ratios of carbon ($\delta^{13}\text{C}$), nitrogen ($\delta^{15}\text{N}$) and sulfur ($\delta^{34}\text{S}$) measured in the whitemouth croaker muscle samples in the present study are summarized in Table 1. Considering that the ANOSIM test did not reveal significant inter-site difference in fish contamination patterns ($p = 0.17$, $R = 0.11$), correlations were performed after grouping individuals from both estuaries. Regarding correlations between pollutant levels (or grouped pollutants) and stable isotope ratios, ΣPBDE values were not correlated to any stable isotope ratios; however, using the PBDE congener groups according to the number of bromine atoms, tri-BDE was positively correlated with $\delta^{13}\text{C}$ ($r_s = 0.48$, $p = 0.03$) and $\delta^{15}\text{N}$ ($r_s = 0.45$, $p = 0.04$), while negatively correlated with $\delta^{34}\text{S}$ ($r_s = -0.48$, $p = 0.03$) values (Fig. 3; Table S10). Tetra- and penta-BDE levels increased with higher $\delta^{15}\text{N}$ values ($r_s = 0.50$, $p = 0.02$ and $r_s = 0.45$, $p = 0.045$, respectively). On the other hand, ΣDRC was positively correlated with $\delta^{15}\text{N}$ ($r_s = 0.46$, $p = 0.04$); DP and *anti*-DP were both positively correlated with $\delta^{13}\text{C}$ ($r_s = 0.47$, $p = 0.04$ and $r_s = 0.48$, $p = 0.03$, respectively) and $\delta^{15}\text{N}$ values ($r_s = 0.59$, $p = 0.006$ and $r_s = 0.62$, $p = 0.003$, respectively), while a positive correlation was found between Dec 603 and with $\delta^{34}\text{S}$ ($r_s = 0.47$, $p = 0.04$) values. Regarding biological parameters, a positive correlation was found between Dec 603 and lipid content ($r_s = 0.45$, $p = 0.048$), while size was negatively correlated with ΣPBDE , tri- and tetra-BDEs ($r_s = -0.48$, $p = 0.03$; $r_s = -0.45$, $p = 0.046$; and $r_s = -0.54$, $p = 0.01$, respectively). No significant correlations ($p > 0.05$) were found between $\Sigma\text{PCDD/F}$ or TEQ values and biological parameters or stable isotope ratios ($\delta^{13}\text{C}$, $\delta^{15}\text{N}$ and $\delta^{34}\text{S}$) (Table S10).

4 Discussion

4.1 Pollutant exposure of whitemouth croakers from Southeastern Brazil

The PCDD/Fs levels in whitemouth croakers from Guanabara Bay (from $< \text{LOD}$ to $512.7 \text{ pg g}^{-1} \text{ l.w.}$; Table S4) were apparently higher compared to those found in skipjack tuna (*Katsuwonus pelamis*) from Brazilian offshore waters (mean: $4.2 \text{ pg g}^{-1} \text{ l.w.}$) (Ueno *et al.*, 2005). This is not

313 surprising, given that Guanabara Bay is the most anthropogenically-disturbed area along the
314 Brazilian coast (Dorneles *et al.*, 2008a, b; 2013). Additionally, the predominance of OCDD in
315 whitemouth croaker samples agrees with the results observed in blubber samples of Guiana
316 dolphin (Dorneles *et al.*, 2013), as well as in sewage sludge samples from wastewater treatment
317 facilities in the Rio de Janeiro metropolitan area (Pereira *et al.*, 2005). Such predominance
318 suggests that combustion processes, *i.e.* mass combustion, unleaded gasoline and diesel fuel
319 combustion, and urban wastewater treatment plants, might be important sources of PCDD/F
320 contamination in Guanabara Bay (Guerzoni *et al.*, 2007). Regarding TEQ values, the levels
321 reported in this study were lower than the European action level (Recommendation
322 2006/88/EC) and maximum permissible level (Regulation 1881/2006) for PCDD/Fs in fish
323 muscle meat and products (set at 2.3 and 3.2 pg TEQ g⁻¹ w.w., respectively) (EFSA, 2010).
324 However, these results should be taken with caution, since previous research has demonstrated
325 that PCDD/Fs accounted for less than 1.2% of the total TEQ in all Guiana dolphins, indicating
326 that polychlorinated biphenyls (PCBs) are the main cause for environmental concern in Rio de
327 Janeiro state when compared to PCDD/Fs (Dorneles *et al.*, 2013).

328 PBDE levels in marine biota along the Brazilian coast are usually related to sampling areas
329 close to industrial and urbanized regions (Alonso *et al.*, 2012, 2017; Dorneles *et al.*, 2010;
330 Magalhães *et al.*, 2017; Quinete *et al.*, 2011; Rosenfelder *et al.*, 2012). For example, Lavandier
331 *et al.* (2013) have found PBDE values below LOD in muscle of whitemouth croakers from Ilha
332 Grande Bay, a less impacted estuary classified as a biodiversity hotspot in the south of Rio de
333 Janeiro state (Creed *et al.*, 2007). Conversely, fish from the Paraíba do Sul river (north of Rio
334 de Janeiro state) have shown apparently higher PBDE muscle concentrations than the ones
335 reported here (*i.e.*, with a mean of 2.1 ng g⁻¹ w.w.; Quinete *et al.*, 2011). While Guanabara and
336 Sepetiba bays are among the most impacted areas along the Brazilian coastline (Baptista-Neto
337 *et al.*, 2016; Dorneles *et al.*, 2008a, b; 2013; Kjerfve *et al.*, 1997; Molisani *et al.*, 2004), the

338 presence of urban (Rio de Janeiro and São Paulo cities) and industrial centres (chemicals,
339 textiles, sugar-alcohol) along the course of Paraíba do Sul river might play a role in the presence
340 of POPs contamination in its estuary (Linde-Arias *et al.*, 2008).

341 The high detection frequency and abundance of tetra- (BDE-47) and penta- (BDE-99 and -100)
342 PBDE congeners in whitemouth croakers (Fig. 2A; Table S6), could reflect the use of
343 commercial pentaBDE (C-pentaBDE) mixtures in Southeastern Brazil (la Guardia *et al.*, 2006).
344 Additionally, the high contribution of BDE-47 in whitemouth croakers reflects a worldwide
345 trend observed in aquatic biota (Barón *et al.*, 2015; Houde *et al.*, 2014; Mizukawa *et al.*, 2009,
346 Shao *et al.*, 2016), including Brazilian environments (Alonso *et al.*, 2012; Dorneles *et al.*, 2010;
347 Magalhães *et al.*, 2017; Quinete *et al.*, 2011). However, these results probably originate from
348 the combination of several factors, such as (1) higher release of BDE-47 and, consequently,
349 higher bioavailability for uptake by biota, (2) higher assimilation efficiency and resistance to
350 metabolism, and (3) metabolic transformation via debromination from higher to lower
351 brominated congeners (Munschy *et al.*, 2011; Roberts *et al.*, 2011; Stapleton *et al.*, 2006).

352 Unlike BDE-47, BDE-209 has been less reported in biota. This could be linked to its
353 physicochemical properties that cause low availability for, and low uptake by, biota, and by
354 debromination into lower brominated congeners (Tomy *et al.*, 2004; Stapleton *et al.*, 2006;
355 Munschy *et al.*, 2011; Roberts *et al.*, 2011). To the authors' knowledge, this is the first study
356 reporting BDE-209 in fish from the southwest Atlantic Ocean, as previous studies have not
357 targeted this congener (Lavandier *et al.*, 2013; Magalhães *et al.*, 2017; Quinete *et al.*, 2011).

358 The presence of BDE-209 in muscle of whitemouth croaker (Tables S6 and S7), as well as in
359 the blubber of Guiana dolphin (*Sotalia guianensis*) from Guanabara Bay (Vidal, 2015),
360 indicates the use of the commercial decaBDE (C-decaBDE) mixture (> 92 % of BDE-209; la
361 Guardia *et al.*, 2006) in Southeastern Brazil. Additionally, the presence of BDE-183 in
362 whitemouth croaker samples also suggests the recent use of commercial octaBDE (C-octaBDE)

363 mixtures in Southeastern Brazil, since this congener has not been quantified in previous studies
364 using mussels (sampled in 1996; Zhu and Hites, 2003), Guiana dolphin (from 1994 to 2006;
365 Dorneles *et al.*, 2010) and rays (Rosenfelder *et al.*, 2012) from Guanabara Bay.

366 Amongst emerging BFRs evaluated in the present study, only HBB was quantified in
367 whitemouth croaker samples. To the best of our knowledge, only two studies have previously
368 reported levels of emerging BFRs in aquatic biota from the southwest Atlantic Ocean (Alonso
369 *et al.*, 2012; de la Torre *et al.*, 2012). According to Alonso *et al.* (2012), HBB was detected in
370 13 (25% of individuals sampled) Franciscana dolphins (*Pontoporia blainvillei*) from the
371 Southeastern and Southern coasts of Brazil, while PBEB was detected in four individuals (8%
372 of the total). The low detection frequency of these compounds in aquatic biota could indicate
373 their low use in Brazil. While HBB can be used directly as flame retardant in manufactured
374 products (Covaci *et al.*, 2011; de Wit *et al.*, 2011), its presence in the environment can also
375 result from thermal degradation of commercial mixtures of PBDEs, and volatilization of
376 polymeric brominated flame retardants, such as pentabromobenzyl acrylate oligomer (de Wit
377 *et al.*, 2011; Gouteux *et al.*, 2008). These aspects, combined with the low frequency of HBB
378 detection in our samples, reinforce the hypothesis of low use of this compound in Brazil.

379 The dechlorane-related contamination pattern observed in whitemouth croakers (Dec 603 > DP
380 > mirex \approx Dec 602 > CP) in the present study suggest, for the first time, Dec 603 as the most
381 abundant DRC in environmental samples. This pattern is notable since Dec 603 is usually
382 reported as low or non-detected values in biota (Houde *et al.*, 2014; Mekni *et al.*, 2019; Rjabova
383 *et al.*, 2016). Patented by Hooker Chemicals (now Occidental Chemical Company, OxyChem,
384 United States), Dec 603 is identified as a flame retardant, and also as an impurity in technical
385 products of aldrin and dieldrin (legacy pesticides) (Shen *et al.*, 2011). Brazil allowed the
386 production of aldrin until 1990 and its use as wood preservative until 2000, while dieldrin has
387 no register of use in Brazil but its production for export occurred until 1998 (MMA, 2015;

Almeida *et al.*, 2007). However, since aldrin and dieldrin have been reported in crabs (Souza *et al.*, 2008) and in mussels (Galvão *et al.*, 2015) from Southeastern Brazil, the occurrence of the emerging pollutant Dec 603 in whitemouth croaker in this region could be associated to the production and use of these pesticides.

DP showed the second highest contribution to Σ DRC in whitemouth croaker, as well as being reported in Franciscana dolphins (*Pontoporia blainvillei*) (Mirex > DP > Dec 603 > Dec 602 > CP) from the Southeastern and Southern Brazilian coasts (de la Torre *et al.*, 2012). Since the major applications of DP are industrial polymers used for coating electrical wires and cables, connectors used in computers, and plastic roofing material (Hoh *et al.*, 2006), this suggests that DP in Southeastern Brazil could be linked to high anthropogenic influence and industrial activity. The predominance of *anti*-DP isomer in whitemouth croakers ($f_{anti} = 0.7 \pm 0.2$; Table 1) is consistent with commercial DP products (0.6 - 0.8; from Sverko *et al.* 2011 and Wang *et al.*, 2010).

Mirex and Dec 602 showed similar contributions to Σ DRC in whitemouth croaker, but this is not in agreement with previous studies in aquatic biota worldwide that report higher levels of Mirex compared to Dec 602 (de la Torre *et al.*, 2012; Peng *et al.*, 2014; Rjabova *et al.*, 2016). Mirex was widely used as a pesticide for ant control in Brazil (MMA, 2015) and, although banned in the 1990s (MMA, 2015), Mirex persistence remains an important factor to consider for understanding its detection in representatives of the Brazilian marine biota that have been recently sampled (Alonso *et al.*, 2017; de la Torre *et al.*, 2012; Santos-Neto *et al.*, 2014). In contrast to Mirex, there is no information on Dec 602 use in Brazil, however, its presence in the environment appears to be related to its use as flame retardant in manufactured products (Sverko *et al.*, 2011). The presence of Dec 602 in whitemouth croaker, as well as in marine mammals off the coast of Brazil (de la Torre *et al.*, 2012; Alonso *et al.*, 2017) reinforces the need for further research to investigate its ecotoxicological relevance.

Although the pollutant exposure of whitemouth croakers in Southeastern Brazil can be inferred through the POPs concentrations found in this study, to the best of our knowledge, no studies have focused on the mechanisms for accumulation and depuration, or on risk assessment for this species. Overall, the most likely routes of POPs uptake in fishes are dietary and respiratory via the gills and body surface area, whereas elimination is primarily via the respiratory surface, kidneys, and feces, and often involves metabolic transformation (Arnot and Gobas, 2004; Munschy *et al.*, 2011; Tierney *et al.*, 2013). Regarding risk assessment for POPs concentrations in fishes, studies on experimental exposure have demonstrated alterations in the immune and endocrine systems, as well as in their life-history traits such as reproductive success, growth and survival (Horri, *et al.*, 2018; Johnson *et al.*, 2013). For example, McCarthy *et al.* (2003) have shown that parental exposure to a commercial PCB mixture (Aroclor 1254) through the diet, during gonadal recrudescence, affected growth and survival skills of Atlantic croaker (*Micropogonias undulatus*) larvae, reducing their growth rates and impairing their startle responses. In this context, POPs exposure could not only affect the physiological responses, but also recruitment and population dynamics and, to some extent, the effects could also affect fisheries productivity for commercial species.

4.2 Linking pollutant exposure of whitemouth croakers to their ecological habits and biological features

Isotopic ratios of carbon and sulfur are usually used to establish the sources of organic matter that support food webs (Connolly *et al.* 2004; McCutchan *et al.* 2003). Nitrogen stable isotope ratios can also be used to trace organic matter sources, but are more commonly applied to provide information on the position occupied by a species in a trophic web, as nitrogen isotopes show predictable stepwise increases in values from prey to consumer (DeNiro and Epstein, 1981). In this context, our three-isotope approach showed that $\delta^{15}\text{N}$ was the isotopic ratio that

was the most commonly correlated with pollutant concentrations in whitemouth croaker. Specimens with high $\delta^{15}\text{N}$ showed higher levels of tri-, tetra- and penta-BDEs, as well as higher concentrations of DP, *anti*-DP isomer and ΣDRC . These findings suggest the occurrence of bioaccumulation and, to some extent, the biomagnification of some target pollutants through the coastal food web.

Bioaccumulation and biomagnification of organic pollutants can be influenced by many factors, such as their molecular size and octanol-water partition coefficients (K_{OW}) (Kelly *et al.*, 2007; Walters *et al.*, 2016). In aquatic food webs, chemicals with K_{OW} values between 10^5 and 10^8 would have higher bioaccumulation and biomagnification potentials, while the opposite would occur for chemicals with $K_{\text{OW}} > 10^8$ (Kelly *et al.*, 2007; Stapleton *et al.*, 2006; Walters *et al.*, 2016). Therefore, low brominated PBDEs, such as tri-, tetra- and penta-PBDEs (K_{OW} between $\sim 10^5$ and $\sim 10^7$) are prone to bioaccumulate and biomagnify in aquatic food webs, as reported previously (Barón *et al.*, 2015; Mizukawa *et al.*, 2009; Shao *et al.*, 2016), while bioaccumulation and biomagnification potentials of DP would be reduced ($K_{\text{OW}} \sim 10^9$) (Hoh *et al.*, 2006; Peng *et al.*, 2014). Due to its high hydrophobicity, DP is mainly adsorbed to organic materials, and exhibits persistence in sediment (Sverko *et al.*, 2011; Shen *et al.*, 2010). From this perspective, the use of benthic species or benthivorous demersal species, such as the whitemouth croaker, might help to demonstrate bioaccumulation and biomagnification of DP through aquatic food webs (Carlsson *et al.*, 2018; Na *et al.*, 2017; Sühring *et al.*, 2016). However, for DP isomers, aspects of stereoselective bioaccumulation potential and trophic transfer remain unclear. For instance, higher *anti*-DP concentrations upon organisms with ^{15}N -enriched values were found in aquatic biota from Lake Winnipeg (Canada) (Tomy *et al.*, 2007), as well as in the marine food webs of the Fildes Peninsula (Antarctica) (Na *et al.*, 2017); while an opposite behaviour was found in the freshwater food web from Longtang Town (China) (Wu *et al.*, 2010).

Negative correlations between fish size and Σ PBDE, tri- and tetra-BDEs were also found in the present study. This could be caused by depuration in fish, as reported in previous studies (Munschy *et al.*, 2011; Tomy *et al.*, 2004). However, it could also be linked with ontogenic habitat shifts, as older whitemouth croaker move into continental shelf waters outside of the bays, *i.e.* out of estuaries that are hotspots for contaminant exposure. This hypothesis is in accordance with the higher $\delta^{34}\text{S}$ values found in larger whitemouth croaker from Guanabara Bay (Pizzochero *et al.*, 2018), as these large fish probably mostly feed in continental shelf waters, which are ^{34}S -enriched compared to coastal zones (Connolly *et al.* 2004; Thode 1991).

5. Conclusion

This study provides new data on the contamination of Brazilian marine coastal environments by selected organic pollutants. It reveals the presence of non-PBDE brominated flame retardants and DRCs (Dec 602, 603, DP and CP) in fish from southwest Atlantic Ocean for the first time, albeit at low levels. PBDEs were detected in all samples analyzed, with the predominance of BDE -47, -99, -100, -153, 154, -183 and -209, which might reflect the use of C-pentaBDE, C-octaBDE and C-decaBDE commercial mixtures in the coastal regions of Rio de Janeiro state. Dec 603 and DP were the predominant DRCs in whitemouth croakers and their presence in fish raises concern and strengthens the need for further research not only on their toxicity and bioaccumulation potentials, but also on their occurrence and distribution in the environment. TEQ total levels for dioxins and furans ranged from 0.1 to 0.2 pg g⁻¹ w.w., which is lower than the European action and maximum permissible levels for PCDDs and PCDFs for fish muscle meat and products (set at 2.3 and 3.2 pg TEQ g⁻¹ w.w., respectively). Concentrations of tri-, tetra- and penta-BDEs, as well as Σ DRC, DP and *anti*-DP isomer were positively correlated with $\delta^{15}\text{N}$, suggesting biomagnification along the food web resulting in the contamination levels reported for whitemouth croaker. On the other hand, Σ PBDE, tri- and

tetra-BDEs were negatively correlated with fish size, which could be linked with depuration by fishes and/or habitat shift throughout the whitemouth croaker life cycle. Overall, our study confirms that whitemouth croaker might be a suitable sentinel species of coastal pollution. Moreover, it emphasizes the need for further investigations focusing on multiple species, as well as in water and sediment samples, to help understand the complex patterns of bioaccumulation and biomagnification. These processes seem to occur in Southeastern Brazil, and they could impact not only the marine biota, but also the human population dependent on this biota for food.

Acknowledgements

We thank the referees for their very helpful comments that have helped to improve the manuscript. This work was supported by the CIEMAT (public body for research in the fields of energy, environment and technology), depending on the Spanish Ministry of Science, Innovation and Universities. The authors would like to thank CIEMAT for allowing AC Pizzochero stay in the Laboratory of POPs (Madrid, Spain). This work was also supported by the Brazilian National Council for Scientific and Technological Development (CNPq) through a Universal Call CNPq-Project from PRD (proc. 432518/2016-9), as well as through a scientific cooperation established between the Brazilian Foundation for the Coordination and Improvement of Higher Level or Education Personnel (CAPES - process numbers 88881.154725/2017-01 88887.154724/2017-00) and *Wallonie Bruxelles International* (WBI, from Belgium), coordinated by PRD and KD. This study was also supported by scientific cooperation established between the Rio de Janeiro State Government Research Agency (FAPERJ, proc. E-26/170.018/2015) and the Research Councils UK (RCUK) in the context of the Newton Fund. IDM was supported by the Newton Fund Programme by a grant awarded by NERC (NE/N000889/1) on behalf of the Research Councils UK. GL is a F.R.S.-FNRS research

513 associate, and KD is a Senior F.R.S.-FNRS research associate. OM and PRD have research
514 grants from CNPq (PQ-1A proc. 306703/2014-9 and PQ-2 proc. 306847/2016-7, respectively).

515

516 **Supplementary data** related to this article can be found at XXXXX.

517

References

- Almeida, F. V., Centeno, A.J., Bisinoti, M.C., Jardim, W.F., 2007. Substâncias tóxicas persistentes (STP) no Brasil. *Quim. Nova.* 30 (8), 1976–1985.
- Alonso, M.B., Eljarrat, E., Gorga, M., Secchi, E.R., Bassoi, M., Barbosa, L., Bertozzi, C.P., Marigo, J., Cremer, M., Domit, C., Azevedo, A.F., Dorneles, P.R., Torres, J.P.M., Lailson-Brito, J., Malm, O., Barceló, D., 2012. Natural and anthropogenically-produced brominated compounds in endemic dolphins from Western South Atlantic: Another risk to a vulnerable species. *Environ. Pollut.* 170, 152–160. doi:10.1016/j.envpol.2012.06.001
- Alonso, M.B., Maruya, K.A., Dodder, N.G., Lailson-Brito, J., Azevedo, A., Santos-Neto, E., Torres, J.P.M., Malm, O., Hoh, E., 2017. Nontargeted Screening of Halogenated Organic Compounds in Bottlenose Dolphins (*Tursiops truncatus*) from Rio de Janeiro, Brazil. *Environ. Sci. Technol.*, 51 (3), 1176–1185. 10.1021/acs.est.6b04186
- Arnot, J. A., Gobas, F. A. P. C., 2004. A food web bioaccumulation model for organic chemicals in aquatic ecosystems. *Environ Toxicol Chem.* 23, 2343–2355.
- Baptista-Neto, J.A., Barreto, C.F., Vilela, C.G., Fonseca, E.M.F., Melo, G.V., Barth, O.M. 2016. Environmental change in Guanabara Bay, SE Brazil, based in microfaunal, pollen and geochemical proxies in sedimentary cores. *Ocean Coast Manag.* 143, 4–15. doi: dx.doi.org/10.1016/j.ocecoaman.2016.04.010
- Barón, E., Giménez, J., Verborgh, P., Gauffier, P., De Stephanis, R., Eljarrat, E., Barceló, D., Bioaccumulation and biomagnification of classical flame retardants, related halogenated natural compounds and alternative flame retardants in three delphinids from Southern European waters. *Environ. Pollut.* 203, 107–115. doi: 10.1016/j.envpol.2015.03.041.
- Bisi, T. L.; Lepoint, G.; Azevedo, A. d. F.; Dorneles, P. R.; Flach, L.; Das, K.; Malm, O.; Lailson-Brito, J. 2012. Trophic relationships and mercury biomagnification in Brazilian tropical coastal food webs. *Ecol. Indic.* 18, (0), 291–302. <https://doi.org/10.1016/j.ecolind.2011.11.015>
- Carlsson, P., Vrana, B., Sobotka, J., Borgå, K., Nizzetto, P.B., Varpe, Ø., 2018. New brominated flame retardants and dechlorane plus in the Arctic: Local sources and bioaccumulation potential in marine benthos. *Chemosphere.* 211, 1193–1202.
- Chouvelon, T., Caurant, F., Cherel, Y., Simon-Bouhet, B., Spitz, J., Bustamante, P., 2014. Species- and size-related patterns in stable isotopes and mercury concentrations in fish help refine marine ecosystem indicators and provide evidence for distinct management units for hake in the Northeast Atlantic. *ICES J. Mar. Sci.* 71, 1073–1087. doi:10.1093/icesjms/fst199
- Chouvelon, T., Brach-Papa, C., Auger, D., Bodin, N., Bruzac, S., Crochet, S., Degroote, M., Hollanda, S.J., Hubert, C., Knoery, J., Munsch, C., Puech, A., Rozuel, E., Thomas, B., West, W., Bourjea, J., Nikolic, N., 2017. Chemical contaminants (trace metals, persistent organic pollutants) in albacore tuna from western Indian and south-eastern Atlantic Oceans: Trophic influence and potential as tracers of populations. *Sci. Total Environ.* 15 (596–597), 481–495. doi: 10.1016/j.scitotenv.2017.04.048.
- Clarke, K.R., Warwick, R.M., 2001. Change in marine communities: an approach to statistical analysis and interpretation (2nd edition). PRIMER-E, Plymouth, 172 pp.
- Connolly, R.M., Guest, M.A., Melville, A.J., Oakes, J.M., 2004. Sulfur stable isotopes separate

561 producers in marine food-web analysis. *Oecologia*. 138, 161–167. doi:10.1007/s00442-
562 003-1415-0

563 Coplen, T.B., 2011. Guidelines and recommended terms for expression of stable-isotope-ratio
564 and gas-ratio measurement results. *Rapid Commun. Mass Spectrom.* 25, 2538–2560.
565 doi:10.1002/rcm.5129.

566 Covaci A., Harrad, S., Abdallah, M.A.-E., Ali, N., Law, R.J., Herzke, D., De wit, C.A., 2011.
567 Novel brominated flame retardants: A review of their analysis, environmental fate and
568 behaviour. *Environ. Int.* 37, (2), 532-556. <https://doi.org/10.1016/j.envint.2010.11.007>

569 Creed, J.C., Oliveira, A.E.S., Pires, D.O., Figueiredo, M.A.O., Ferreira, C.E.L., Ventura,
570 C.R.R., Brasil, A.C.S., Young, P.S., Absalão, R.S., Paiva, P.C., Castro, C.B., Serejo, C.S.,
571 2007. RAP Ilha Grande – um levantamento da biodiversidade: histórico e conhecimento
572 da biota. In: Creed, J.C., Pires, D.O., Figueiredo, M.A.O. (Eds.), *Biodiversidade Marinha*
573 *da Baía da Ilha Grande*. MMA/SBF, Brasília, pp. 43–63

574 Cruz, R., Cunha, S.C., Casal, S., 2015. Brominated flame retardants and seafood safety: A
575 review. *Environ. Int.* 77, 116–131.

576 De la Torre, A., Alonso, E., Concejero, M.A., Sanz, P., Martínez, M.A., 2011. Sources and
577 behaviour of polybrominated diphenyl ethers (PBDEs), polychlorinated dibenzo-p-dioxins
578 and dibenzofurans (PCDD/Fs) in Spanish sewage sludge. *Waste Manag.* 31, 1277–1284.
579 doi:10.1016/j.wasman.2011.01.021

580 De la Torre, A., Alonso, M.B., Martínez, A.M., Sanz, P., Shen, L., Reiner, E.J., Lailson-Brito,
581 J., Torres, J.P., Bertozzi, C., Marigo, J., Barbosa, L., Cremer, M., Secchi, E., Malm, O.,
582 Eljarrat, E., Barceló, D., 2012. Dechlorane-related compounds in franciscana dolphin
583 (*Pontoporia blainvillei*) from southeastern and southern coast of Brazil. *Environ. Sci.*
584 *Technol.* 46, 12364–12372.

585 De Wit, C.A., Kierkegaard, A., Ricklund, N., Sellstro, U. Emerging Brominated Flame
586 Retardants in the Environment. In: E. Eljarrat and D. Barceló (eds.), *Brominated Flame*
587 *Retardants*, *Hdb Env Chem.* 2011. 16: 241–286, DOI 10.1007/698_2010_73.

588 DeNiro, M.J., Epstein, S., 1981. Influence of diet on the distribution of nitrogen isotopes in
589 animal. *Geochim. Cosmochim. Acta* 45, 341–351. doi:10.1016/0016-7037(81)90244-1

590 Dorneles, P.R., Lailson-Brito, J., Azevedo, A.F., Meyer, J., Vidal, L.G., Fragoso, A.B., Torres,
591 J.P.M., Malm, O., Blust, R., Das, K., 2008a. High Accumulation of Perfluorooctane
592 Sulfonate (PFOS) in Marine Tucuxi Dolphins (*Sotalia guianensis*) from the Brazilian
593 Coast. *Environ. Sci. Technol.*, 42 (14), 5368–5373. doi: 10.1021/es800702k

594 Dorneles, P.R., Lailson-Brito, J., Fernandez, M.A.S., Vidal, L.G., Barbosa, L.A., Azevedo,
595 A.F., Fragoso, A.B.L., Torres, J.P.M., Malm, O., 2008b. Evaluation of cetacean exposure
596 to organotin compounds in Brazilian waters through hepatic total tin concentrations.
597 *Environ. Pollut.* 153 (3), 1268–1276. doi.org/10.1016/j.envpol.2008.03.007

598 Dorneles, P.R., Lailson-Brito, J., Dirtu, A.C., Weijs, L., Azevedo, A.F., Torres, J.P.M., Malm,
599 O., Neels, H., Blust, R., Das, K., Covaci, A., 2010. Anthropogenic and naturally-produced
600 organobrominated compounds in marine mammals from Brazil. *Environ. Int.* 36, 60–67.
601 doi:10.1016/j.envint.2009.10.001

602 Dorneles, P.R., Sanz, P., Eppe, G., Azevedo, A.F., Bertozzi, C.P., Martínez, M.A., Secchi, E.R.,
603 Barbosa, L.A., Cremer, M., Alonso, M.B., Torres, J.P.M., Lailson-Brito, J., Malm, O.,

Eljarrat, E., Barceló, D., Das, K., 2013. High accumulation of PCDD, PCDF, and PCB congeners in marine mammals from Brazil: A serious PCB problem. *Sci. Total Environ.* 463–464, 309–318. doi:10.1016/j.scitotenv.2013.06.015

Dorneles, P.R., Lailson-Brito, J., Bisi, T.L., Domit, C., Barbosa, L.A., Meirelles, A.C.O., Carvalho, V.L., Malm, O., Azevedo, A.F., Brose, F., Das, K., Scippo, M.L. 2016. Guiana Dolphins (*Sotalia guianensis*) and DR-CALUX for Screening Coastal Brazilian Environments for Dioxins and Related Compounds. *Arch. Environ. Contam Toxicol.* 71(3), 336-346. doi: 10.1007/s00244-016-0299-z.

EFSA, 2010. European Food Safety Authority Journal. Results of the monitoring of dioxin levels in food and feed. 8(3):1385. 36 pp.

FAO - Food and Agriculture Organization. Fisheries and aquaculture software. FishStatJ - software for fishery statistical time series. 2018. In: FAO Fisheries and Aquaculture Department [online]. Rome. Updated 21 July 2016. Available from <http://www.fao.org/fishery/>. Accessed October 2018.

Galvão, P., Henkelmann, B., Longo, R., Torres, J.P.M., Malm, O., Schramm, K.W., 2015. The brown mussel *Perna perna* (L., 1758) as a sentinel species for chlorinated pesticide and dioxin-like compounds. *Environ. Sci. Pollut. Res. Int.* 22(17), 3522-33. doi: 10.1007/s11356-015-4607-3.

Gouteux, B., Alae, M., Mabury, S.A., Pacepavicius, G., Muir, D.C., 2008. Polymeric brominated flame retardants: are they a relevant source of emerging brominated aromatic compounds in the environment? *Environ. Sci. Technol.* 42 (24), 9039-44. doi: 10.1021/es8022336

Guerzoni, S., Rossini, P., Saretta, A., Raccanelli, S., Ferrari, G., Molinaroli, E., 2007. POPs in the lagoon of Venice: budgets and pathways. *Chemosphere.* 67, 1776–1785.

Haimovici, M., Cardoso, L.G., Unpierre, R.G., 2016. Stocks and management units of *Micropogonias furnieri* (Desmarest, 1823) in southwestern Atlantic. *Lat. Am. J. Aquat. Res.* 44, (5), 1080-1095.

Hammer, O., Harper, D.A.T., Ryan, P.D., 2001. PAST: Paleontological Statistics software package for education and data analysis. *Palaeontol. Electronica.* 4(1), 1-9.

Hoh, E., Zhu, L., Hites, R.A., 2006. Dechlorane Plus, a Chlorinated Flame Retardant, in the Great Lakes. *Environ. Sci. Technol.* 40, 1184-1189.

Horri, K., Alfonso, S., Cousin, X., Munsch, C., Loizeau, V., Aroua, S., Bégout, M.-L., Ernande, B., 2018. Fish life-history traits are affected after chronic dietary exposure to an environmentally realistic marine mixture of PCBs and PBDEs. *Sci. Total Environ.* 610–611, 531-545. doi: 10.1016/j.scitotenv.2017.08.083.

Houde, M., Berryman, D., de Lafontaine, Y., Verreault, J., 2014. Novel brominated flame retardants and dechloranes in three fish species from the St. Lawrence River, Canada. *Sci. Total Environ.* 479–480, 48–56. doi:10.1016/j.scitotenv.2014.01.105.

IBGE - Instituto Brasileiro de Geografia e Estatística. Base Cartográfica Contínua, ao milionésimo - BCIM 2016: 5ª versão digital. Rio de Janeiro, 2016. Available at: http://downloads.ibge.gov.br/downloads_geociencias.htm. Accessed October 2018. [In Portuguese].

Johnson, L.L., Anulacion, B.F., Arkoosh, M.R., Burrows, D.G., Da Silva, D.A.M., Dietrich,

647 J.P., Myers, M.S., Spromberg, J.A., Ylitalo, G.M., 2013. Effects of legacy persistent
648 organic pollutants (POPs) in fish – current and future challenges. In: Tierney, K.B., Farrell,
649 A.P., Brauner, C.J. (Eds.), *Fish Physiology: Organic Chemical Toxicology of Fishes*. vol.
650 33. Academic Press, San Diego, California, pp. 53–140.

651 Jones, K., De Voogt, P., 1999. Persistent organic pollutants (POPs): state of the science.
652 *Environ. Pollut.* 100, 209–221. doi:10.1016/S0269-7491(99)00098-6

653 Kelly, B.C., Ikononou, M.G., Blair, J.D., Morin, A.E., Gobas, F.A.P.C., 2007. Food web-
654 specific biomagnification of persistent organic pollutants. *Science*. 317, 236–239.
655 doi:10.1126/science.1138275

656 Kjerfve, B., Ribeiro, C.H.A., Dias, G.T.M., Filippo, A.M., and Quaresma, V. da S., 1997.
657 Oceanographic characteristics of an impacted coastal bay: Baía de Guanabara, Rio de
658 Janeiro, Brazil. *Cont. Shelf Res.* 17(13), 1609–1643. doi:10.1016/S0278-4343(97)00028-
659 9.

660 La Guardia, M.J., Hale, R.C., Harvey, E., 2006. Detailed polybrominated diphenyl ether
661 (PBDE) congener composition of the widely used penta-, octa-, and deca-PBDE technical
662 flame-retardant mixtures. *Environ Sci Technol.* 40, 6247–6254.

663 Lavandier, R., Quinete, N., Hauser-Davis, R.A., Dias, P.S., Taniguchi, S., Montone, R.,
664 Moreira, I., 2013. Polychlorinated biphenyls (PCBs) and polybrominated diphenyl ethers
665 (PBDEs) in three fish species from an estuary in the southeastern coast of Brazil.
666 *Chemosphere*. 90, 2435–2443. doi:10.1016/j.chemosphere.2012.11.001.

667 Linde-Arias, A.R., Inacio, A.F., Novo, L.A., de Albuquerque, C., Moreira, J.C., 2008.
668 Multibiomarker approach in fish to assess the impact of pollution in a large Brazilian river,
669 Paraíba do Sul. *Environ. Pollut.* 156, 974–979

670 Magalhães, C.A., Taniguchi, S., Lourenço, R.A., Montone, R.C., 2017. Organochlorine
671 pesticides, PCBs, and PBDEs in liver and muscle tissues of *Paralichthys brasiliensis*,
672 *Trichiurus lepturus* and *Cathorops spixii* in Santos Bay and surrounding area, São Paulo,
673 Brazil. *Reg. Stud. Mar. Sci.* 16, 42–48. <https://doi.org/10.1016/j.rsma.2017.08.010>

674 McCarthy, I.D., Fuiman, L.A., Alvarez, M.C., 2003. Aroclor 1254 affects growth and survival
675 skills of Atlantic croaker *Micropogonias undulatus* larvae. *Mar. Ecol. Prog. Ser.* 252, 295–
676 301.

677 McCutchan, J.H., Lewis, W.M., Kendall, C., McGrath, C.C., 2003. Variation in trophic shift
678 for stable isotope ratios of carbon, nitrogen, and sulfur. *Oikos*. 102, 378–390.
679 doi:10.1034/j.1600-0706.2003.12098.x

680 Mekni, M., Barhoumi, B., Aznar-Alemany, O., Touil, S., Driss, M.R., Barceló, D., Eljarrat, E.,
681 2019. Occurrence of halogenated flame retardants in sediments and sea urchins
682 (*Paracentrotus lividus*) from a North African Mediterranean coastal lagoon (Bizerte,
683 Tunisia). *Sci. Total Environ.* 654, 1316–1325.

684 Mizukawa, K., Takada, H., Takeuchi, I., Ikemoto, I., Omori, K., Tsuchiya, K., 2009.
685 Bioconcentration and biomagnification of polybrominated diphenyl ethers (PBDEs)
686 through lower-trophic-level coastal marine food web. *Mar. Pollut. Bull.* 58, 1217–1224.

687 MMA - Ministry of the Environment. 2015. National Implementation Plan Brazil: Stockholm
688 Convention. Brasília: 182 p.

689 Molisani, M.M., Marins, R.V., Machado, W., Paraquetti, H.H., Bidone, E.D., Lacerda, L.D.,

2004. Environmental Changes in Sepetiba Bay, SE Brazil. *Reg. Environ. Change*, 4
(1),17–27.

Munsch, C., Héas-Moisan, K., Tixier, C., Olivier, N., Gastineau, O., Le Bayon, N., Buchet, B.,
2011. Dietary exposure of juvenile common sole (*Solea solea* L.) to polybrominated
diphenyl ethers (PBDEs): Part 1. Bioaccumulation and elimination kinetics of individual
congeners and their debrominated metabolites *Environ. Pollut.* 159, 229–237.
<https://doi.org/10.1016/j.envpol.2010.09.001>

Na, G., Yao, Y., Gao, H., Li, R., Ge, L., Titaley, I.A., Santiago-Delgado, L., Massey Simonich,
S.L., 2017. Trophic magnification of Dechlorane Plus in the marine food webs of Fildes
Peninsula in Antarctica. *Mar. Pollut. Bull.* 117, 456–461.
[doi:10.1016/j.marpolbul.2017.01.049](https://doi.org/10.1016/j.marpolbul.2017.01.049)

Navarro, I.; de la Torre, A.; Sanz, P.; Pro, J.; Carbonell, G.; Martínez, M. de los Á., 2016.
Bioaccumulation of emerging organic compounds (perfluoroalkyl substances and
halogenated flame retardants) by earthworm in biosolid amended soils. *Environ. Res.* 149,
32–39. [doi: 10.1016/j.envres.2016.05.004](https://doi.org/10.1016/j.envres.2016.05.004)

Navarro, I.; Torre, A. De; Sanz, P.; Porcel, M. Á.; Pro, J.; Carbonell, G.; Martínez, M. D. L. Á.,
2017. Uptake of perfluoroalkyl substances and halogenated flame retardants by crop plants
grown in biosolids-amended soils. *Environ. Res.* 152, 199–206. [doi:](https://doi.org/10.1016/j.envres.2016.10.018)
[10.1016/j.envres.2016.10.018](https://doi.org/10.1016/j.envres.2016.10.018)

Navarro, I.; De La Torre, A.; Sanz, P.; Porcel, M. Á.; Carbonell, G.; Martínez, M. D. L. Á.,
2018. Transfer of perfluorooctanesulfonate (PFOS), decabrominated diphenyl ether
(BDE-209) and Dechlorane Plus (DP) from biosolid-amended soils to leachate and runoff
water. *Environ. Chem.* 15 (4), 195–204.

Peng, H., Wan, Y., Zhang, K., Sun, J., Hu, J., 2014. Trophic transfer of dechloranes in the
marine food web of Liaodong Bay, north China. *Environ. Sci. Technol.* 48(10), 5458–66.
[doi: 10.1021/es500229y](https://doi.org/10.1021/es500229y).

Pereira, M.S., Kuch, B., 2005. Heavy metals, PCDD/F and PCB in sewage sludge samples from
two wastewater treatment facilities in Rio de Janeiro State, Brazil. *Chemosphere.* 60, 844–
853.

Pizzochero, A.C., Michel, L.N., Chenery, S.R., McCarthy, I.D., Vianna, M., Malm, O., Lepoint,
G., Das, K., Dorneles, P.R., 2018. Use of multi-element stable isotope ratios to investigate
ontogenetic movements of *Micropogonias furnieri* in a tropical Brazilian estuary. *Can. J.*
Fish. Aquat. Sci. 75, (6), 977–986. [Doi: https://doi.org/10.1139/cjfas-2017-0148](https://doi.org/10.1139/cjfas-2017-0148)

Quinete, N., Lavandier, R., Dias, P., Taniguchi, S., Montone, R., Moreira, I., 2011. Specific
profiles of polybrominated diphenyl ethers (PBDEs) and polychlorinated biphenyls
(PCBs) in fish and tucuxi dolphins from the estuary of Paraíba do Sul River, Southeastern
Brazil. *Mar. Pollut. Bull.* 62, 440–446. [doi:10.1016/j.marpolbul.2010.11.021](https://doi.org/10.1016/j.marpolbul.2010.11.021)

R Core Team, 2018. R: A language and environment for statistical computing. R Foundation
for Statistical Computing, Vienna, Austria. Available from <https://www.R-project.org/>.

Rjabova, J., Bartkevics, V., Zacs, D., 2016. The occurrence of Dechlorane Plus and related
norbornene-based flame retardants in Baltic wild salmon (*Salmo salar*). *Chemosphere.*
147, 210–217. [doi:10.1016/j.chemosphere.2015.12.122](https://doi.org/10.1016/j.chemosphere.2015.12.122)

Roberts, S.C., Noyes, P.D., Gallagher, E.P., Stapleton, H.M., 2011. Species-specific differences

733 and structure-activity relationships in the debromination of PBDE congeners in three fish
734 species. *Environ. Sci. Technol.* 45, 1999–2005. doi:10.1021/es103934x.

735 Rosenfelder, N., Lehnert, K., Kaffarnik, S., Torres, J.P.M., Vianna, M., Vetter, W., 2012.
736 Thorough analysis of polyhalogenated compounds in ray liver samples off the coast of Rio
737 de Janeiro, Brazil. *Environ. Sci. Pollut. Res.* 19, 379–389.

738 Santos-Neto, E.B., Azevedo-Silva, C.E., Bisi, T.L., Santos, J., Meirelles, A.C., Carvalho, V.L.,
739 Azevedo, A.F., Guimarães, J.E., Lailson-Brito, J., 2014. Organochlorine concentrations
740 (PCBs, DDTs, HCHs, HCB and MIREX) in delphinids stranded at the northeastern Brazil.
741 *Sci. Total Environ.* 472, 194–203. doi: 10.1016/j.scitotenv.2013.10.117.

742 Shao, M., Tao, P., Wang, M., Jia, H., Li, Y.F., 2016. Trophic magnification of polybrominated
743 diphenyl ethers in the marine food web from coastal area of Bohai Bay, North China.
744 *Environ. Pollut.* 213, 379–385. doi: 10.1016/j.envpol.2016.02.034

745 Shen, L., Reiner, E.J., MacPherson, K.A., Kolic, Sverko, E., Helm, P.A., Bhavsar, S.P., Brindle,
746 I.D., Marvin, C.H., 2010. Identification and screening analysis of halogenated norbornene
747 flame retardants in the Laurentian Great Lakes: Dechloranes 602, 603, and 604. *Environ.*
748 *Sci. Technol.* 44 (2), 760–766. doi: 10.1021/es902482b

749 Shen, L., Reiner, E.J., MacPherson, K.A., Kolic, T.M., Helm, P.A., Richman, L.A., Marvin,
750 C.H., Burniston, D.A., Hill, B., Brindle, I.D., McCrindle, R., Chittim, B.G., 2011.
751 Dechloranes 602, 603, 604, dechlorane plus, and chlordene plus, a newly detected
752 analogue, in tributary sediments of the Laurentian great lakes. *Environ. Sci. Technol.* 45,
753 693–699. doi:10.1021/es1027844

754 Sidhu, K.S., 2003. Health benefits and potential risks related to consumption of fish or fish oil.
755 *Regul. Toxicol. Pharmacol.* 38, 336–344. doi:10.1016/j.yrtph.2003.07.002

756 Souza, A.S., Torres, J.P.M., Meire, R.O., Neves, R.C., Couri, M.S., Serejo, C.S., 2008.
757 Organochlorine pesticides (OCs) and polychlorinated biphenyls (PCBs) in sediments and
758 crabs (*Chasmagnathus granulata*, Dana, 1851) from mangroves of Guanabara Bay, Rio
759 de Janeiro State, Brazil. *Chemosphere.* 73, S186–S192.

760 Stapleton H.M., Brazil, B., Holbrook, R.D., Mitchelmore, C.L., Benedict, R., Konstantinov, A.,
761 Potter, D., 2006. In vivo and in vitro debromination of decabromodiphenyl ether (BDE
762 209) by juvenile rainbow trout and common carp. *Environ Sci Technol.* 40, 4653–8.

763 Sühling, R., Busch, F., Fricke, N., Kötker, D., Wolschke, H., Ebinghaus, R., 2016. Distribution
764 of brominated flame retardants and dechloranes between sediments and benthic fish - A
765 comparison of a freshwater and marine habitat. *Sci. Total Environ.* 542, 578–585.

766 Sverko, E., Tomy, G.T., Reiner, E.J., Li, Y.F., McCarry, B.E., Arnot, J.A., Law, R.J., Hites,
767 R.A., 2011. Dechlorane plus and related compounds in the environment: A review.
768 *Environ. Sci. Technol.* 45, 5088–5098. doi:10.1021/es2003028

769 Thode, H.G., 1991. Sulphur isotopes in nature and the environment: an overview. In *Stable*
770 *isotopes: natural and anthropogenic sulphur in the environment.* Ed H.R. Krouse and V.A.
771 Grinenko. Chap. 1. pp. 1–26. John Wiley and Sons, Chichester, UK

772 Tierney, K.B., Kennedy, C.J., Gobas, F., Gledhill, M., Sekela, M., 2013. Organic contaminants
773 and fish. In: Tierney, K.B., Farrell, A.P., Brauner, C.J. (Eds.), *Fish Physiology: Organic*
774 *Chemical Toxicology of Fishes.* vol. 33. Academic Press, San Diego, California, pp. 1–
775 52.

776 Tomy, G.T., Palace, V.P., Halldorson, T., Braekevelt, E., Danell, R., Wautier, K., Evans, B.,
777 Brinkworth, L., Fisk, A.T., 2004. Bioaccumulation, biotransformation, and biochemical
778 effects of brominated diphenyl ethers in juvenile lake trout (*Salvelinus namaycush*).
779 Environ. Sci. Technol. 38 (5), 1496-1504. doi: 10.1021/es035070v

780 Tomy, G.T., Pleskach, K., Ismail, N., Whittle, D.M., Helm, P.A., Sverko, E.D., Zaruk, D.,
781 Marvin, C.H., 2007. Isomers of dechlorane plus in Lake Winnipeg and Lake Ontario food
782 webs. Environ. Sci. Technol. 41, 2249–2254. doi:10.1021/es062781v

783 Ueno, D., Watanabe, M., Subramanian, A., Tanaka, H., Fillmann, G., Lam, P.K., Zheng, G.J.,
784 Muchtar, M., Razak, H., Prudente, M., Chung, K.H., Tanabe, S., 2005. Global pollution
785 monitoring of polychlorinated dibenzo-p-dioxins (PCDDs), furans (PCDFs) and coplanar
786 polychlorinated biphenyls (coplanar PCBs) using skipjack tuna as bioindicator. Environ.
787 Pollut. 136(2), 303-13.

788 UNEP - United Nations Environmental Programme. Stockholm Convention on Persistent
789 Organic Pollutants (POPs). 2017. The 16 new POPs. An introduction to the chemicals
790 added to the Stockholm Convention as Persistent Organic Pollutants by the Conference of
791 the Parties. Available from: [http://chm.pops.int/Portals/0/download.aspx?d=UNEP-](http://chm.pops.int/Portals/0/download.aspx?d=UNEP-POPS-PUB-Brochure-16NewPOPs-201706.English.pdf)
792 [POPS-PUB-Brochure-16NewPOPs-201706.English.pdf](http://chm.pops.int/Portals/0/download.aspx?d=UNEP-POPS-PUB-Brochure-16NewPOPs-201706.English.pdf) (Accessed October 2018).

793 US EPA, 1994. United States Environmental Protection Agency. Tetra- through Octa-
794 chlorinated Dioxins and Furans by Isotope Dilution HRGC/HRMS: Method 1613,
795 Revision B. Available at:
796 [https://19january2017snapshot.epa.gov/sites/production/files/2015-](https://19january2017snapshot.epa.gov/sites/production/files/2015-08/documents/method_1613b_1994.pdf)
797 [08/documents/method_1613b_1994.pdf](https://19january2017snapshot.epa.gov/sites/production/files/2015-08/documents/method_1613b_1994.pdf). Accessed June 2018.

798 Van den Berg, M., Birnbaum, L.S., Denison, M., De Vito, M., Farland, W., Feeley, M., Fiedler,
799 H., Hakansson, H., Hanberg, A., Haws, L., Rose, M., Safe, S., Schrenk, D., Tohyama, C.,
800 Tritscher, A., Tuomisto, J., Tysklind, M., Walker, N., Peterson, R.E., 2006. The 2005
801 World Health Organization reevaluation of human and mammalian toxic equivalency
802 factors for dioxins and dioxin-like compounds. Toxicol. Sci. 93, 223–241.
803 doi:10.1093/toxsci/kfl055

804 Vazzoler, A.E.A.M., 1991. Síntese sobre a ecologia da corvina no Brasil. Atlantica, 13(1), 55–
805 74.

806 Vidal, L.G., 2015. Organic contaminants in boto-cinza (*Sotalia guianensis*, Cetartiodactyla:
807 Delphinidae) from the coast of Brazil. PhD Thesis. 143p. Federal University of Rio de
808 Janeiro. Brazil. [Abstract in English].

809 Walker, C.H., Hopkin, S.P., Sibly, R.M., Peakall, D.B., Principles of Ecotoxicology. Fourth
810 Edition London: Taylor & Francis, 321p. 2012.

811 Walters, D.M., Jardine, T., Cade, B., Kidd, K.A., Muir, D.C.G., 2016. Trophic magnification
812 of organic chemicals - a global synthesis. Environ. Sci. Technol. 50 (9), 4650–4658

813 Wang, D.G., Yang, M., Qi, H., Sverko, E., Ma, W.L., Li, Y.F., Alaei, M., Reiner, E.J., Shen,
814 L., 2010. An Asia-Specific source of Dechlorane Plus: concentration, isomer profiles, and
815 other related compounds. Environ. Sci. Technol. 44, 6608-6613.
816 <http://dx.doi.org/10.1021/es101224y>

817 Wei, T., Simko, V., 2017. R package "corrplot": Visualization of a Correlation Matrix (Version
818 0.84). Available from <https://github.com/taiyun/corrplot>

- 819 Wu, J.P., Zhang, Y., Luo, X.J., Wang, J., Chen, S.J., Guan, Y.T., Mai, B.X., 2010. Isomer-
820 specific bioaccumulation and trophic transfer of dechlorane plus in the freshwater food
821 web from a highly contaminated site, South China. *Environ. Sci. Technol.* 44, 606–611.
822 doi:10.1021/es902744b
- 823 Zhu L, Hites R., 2003. Determination of polybrominated diphenyl ethers in environmental
824 standard reference materials. *Anal. Chem.* 75:6696–700.

825 Table 1: Biological parameters, organic pollutants concentrations (pg g⁻¹ wet weight), f_{anti} values, TEQ values, and stable isotopes ratios of carbon ($\delta^{13}\text{C}$),
826 nitrogen ($\delta^{15}\text{N}$) and sulfur ($\delta^{34}\text{S}$) in whitemouth croaker (*Micropogonias furnieri*) muscle samples from Southeastern Brazil.

Sampling site	Code	Size (cm)	Lipid content (%)	Mass (kg)	ΣPBDE^a (pg g ⁻¹)	ΣDRC^b (pg g ⁻¹)	f_{anti}^c	$\Sigma\text{PCDD/F}^d$ (pg g ⁻¹)	TEQ ^e (pg g ⁻¹)	$\delta^{13}\text{C}$ (‰)	$\delta^{15}\text{N}$ (‰)	$\delta^{34}\text{S}$ (‰)
Sepetiba Bay	Sb#1	49	0.7	1.5	879.7	33.8	0.6	n.a.	-	-14	15.2	11.5
	Sb#2	48	4.9	1.4	101.6	1.1	n.c.	n.a.	-	-14.5	15.3	12.3
	Sb#3	47	3.4	1.4	111.1	5.1	0.5	n.a.	-	-14.7	15.5	12.3
	Sb#4	46	2.5	1.45	160.8	2.8	0.6	n.a.	-	-14.5	15.2	10.8
	Sb#5	49	4.3	1.4	40.7	3.8	0.7	n.a.	-	-15.4	14.4	12.2
	Sb#6	45	4.2	1.05	84.4	4.5	1	n.a.	-	-14.4	14.9	12.7
Mean \pm SD		47.3 \pm 1.6	3.3 \pm 1.6	1.4 \pm 0.2	229.7 \pm 320.8	8.5 \pm 12.5	0.7 \pm 0.2	-	-	-14.6 \pm 0.5	15.1 \pm 0.4	12 \pm 0.7
Min - Max		45 - 49	0.7 - 4.9	1.05 - 1.5	40.7 - 879.7	1.1 - 33.8	0.5 - 1	-	-	-15.4 - -14	14.4 - 15.5	10.8 - 12.7
Median		47.5	3.8	1.4	106.4	4.2	0.6	-	-	-14.5	15.2	12.3
Guanabara Bay	Gb#1	46	1.6	0.9	221.4	n.d.	n.c.	n.a.	-	-18.5	13.3	14.7
	Gb#2	47	1.6	1.1	7.6	0.7	n.c.	n.a.	-	-18.9	13.6	12.5
	Gb#3	49	2.7	1.09	326.2	3.0	0.4	n.a.	-	-15.6	12.8	15.0
	Gb#4	50	4.9	1.3	67.3	7.0	0.4	n.a.	-	-16.6	14.4	15.0
	Gb#5	51	2.2	1.4	59.7	9.7	1.0	0.4	0.1	-16.4	15.2	16.5
	Gb#6	52	4.9	1.4	43.1	5.1	0.7	0.6	0.1	-18.3	13.5	12.3
	Gb#7	53	3.8	1.9	53.0	2.7	n.c.	0.5	0.1	-16.7	14.0	17.3
	Gb#8	56	2.8	1.8	21.0	4.1	n.c.	n.d.	0.1	-15.8	14.2	15.3
	Gb#9	61	6.5	2.2	23.4	2.1	n.c.	0.3	0.1	-16.5	13.9	15.2
	Gb#10	61	15.9	1.8	315.2	41.7	0.6	0.2	0.1	-18.1	14.7	16.6
	Gb#11	62	3.4	2.1	30.6	6.6	0.6	1.4	0.1	-16.6	14.0	17.3

Gb#12	65	3.4	2.7	43.6	1.0	n.c.	0.01	0.1	-16.1	12.9	15.8
Gb#13	66	2.7	3	47.9	3.2	1	1.3	0.1	-15.8	14.1	16.4
Gb#14	75	1.6	4.4	14.2	1.1	1	1.7	0.2	-15.4	14.4	14.9
Mean ± SD	56.7 ± 8.5	4.1 ± 3.7	1.9 ± 0.9	91 ± 110.2	6.8 ± 10.8	0.7 ± 0.3	0.7 ± 0.6	0.1 ± 0.03	-16.8 ± 1.2	13.9 ± 0.7	15.3 ± 1.5
Min - Max	46 - 75	1.6 – 15.9	0.9 – 4.4	7.6 – 326.2	n.d. – 41.7	0.4 – 1	n.d. – 1.7	0.1 – 0.2	-18.9 – -15.4	12.8 – 15.2	12.3 – 17.3
Median	54.5	3.1	1.8	45.8	3.2	0.6	0.5	0.1	-16.6	14	15.3
TOTAL											
Mean ± SD	53 ± 8.3	3.9 ± 3.2	1.8 ± 0.8	132.6 ± 199.1	7.3 ± 11.1	0.7 ± 0.2	-	-	-16.1 ± 1.4	14.3 ± 0.8	14.3 ± 2
Min - Max	45 - 75	0.7 – 15.9	0.9 – 4.4	7.6 – 879.7	n.d. – 41.1	0.4 – 1	-	-	-18.9 – -14	12.8 – 15.5	10.8 – 17.3
Median	50.5	3.4	1.5	56.4	3.8	0.6	-	-	-16	14.3	15

n.d. = not detected.

n.c. = not calculated due *syn*-DP value below LOD.

n.a. = not analysed.

^aΣPBDE: sum of PBDEs (IUPAC congener numbers: 17, 28, 47, 49 & 71, 66, 77, 85, 99, 100, 119, 126, 139, 140, 153, 154, 156 & 169, 183, 184, 206, 207, 208 and 209).

^bΣDRC: sum of Mirex, Dechlorane 602, Dechlorane 603, Dechlorane Plus and Chlordene Plus.

^c*f_{anti}*: *anti*-DP divided by the sum of *syn*-DP and *anti*-DP.

^dΣPCDD/F: sum of 1,2,3,6,7,8– HexaCDD; 1,2,3,7,8,9– HexaCDD; 1,2,3,4,6,7,8– HeptaCDD; OctaCDD (OCDD); 1,2,7,8-TCDF; 2,3,7,8-TCDF; 1,2,3,7,8-PCDF; 2,3,4,7,8-PCDF; 1,2,3,4,7,8-HCDF; 1,2,3,6,7,8-HCDF; 1,2,3,7,8,9-HCDF; 2,3,4,6,7,8-HCDF; 1,2,3,4,6,7,8-HpCDF; 1,2,3,4,7,8,9-HpCDF and OCDF.

^eTEQ: sum of TEQ of PCDDs and PCDFs.


Figure captions

Fig. 1: Map of South America showing Brazil and Rio de Janeiro state. The insert shows the locations of Sepetiba Bay (A) and Guanabara Bay (B) within Rio de Janeiro state.


Fig. 2: (A) Relative contribution of PBDEs grouped by the number of bromine atoms in the molecule to Σ PBDE, and (B) relative contribution of individual dechlorane-related compounds to Σ DRC in muscle samples of whitemouth croakers from Southeastern Brazil. The figure presents the individual code of each fish, which includes the sampling area (Sep: Sepetiba Bay; Gb: Guanabara Bay) and the specimen number (#1, #2...).

Fig. 3: Spearman rank correlation matrix between organic pollutants and biological parameters [length, mass and lipid content (%)] and stable isotope ratios of carbon ($\delta^{13}\text{C}$), nitrogen ($\delta^{15}\text{N}$) and sulfur ($\delta^{34}\text{S}$) in muscle samples of whitemouth croakers from Southeastern Brazil. Statistically-significant spearman rank correlations (r_s , $p < 0.05$) are shown in blue (positive correlation) and red (negative correlation) color scale (color intensity related to r_s value), while non-significant correlations are left blank.


855 Fig.1



856
857



861 Fig. 3



862

863

864 Supplementary data

865 Occurrence of legacy and emerging organic pollutants in whitemouth croakers from Southeastern
866 Brazil

867 Authors: Ana Carolina Pizzochero^{1*}, Adrián de la Torre², Paloma Sanz², Irene Navarro², Loïc N.
868 Michel^{3**}, Gilles Lepoint³, Krishna Das³, Joseph G. Schnitzler⁴, Simon R. Chenery⁵, Ian D. McCarthy⁶, Olaf
869 Malm¹, Paulo R. Dorneles¹, María Ángeles Martínez ²

870

871 ¹ Radioisotope Laboratory, Biophysics Institute, Federal University of Rio de Janeiro (UFRJ), Brazil

872 ² Persistent Organic Pollutants Group, Environmental Department, CIEMAT, Madrid, Spain

873 ³ Freshwater and Oceanic Sciences Unit of reSearch (FOCUS), Laboratory of Oceanology,

874 University of Liege, Belgium

875 ⁴ Institute for Terrestrial and Aquatic Wildlife Research (ITAW) University of Veterinary Medicine
876 Hannover.

877 ⁵ British Geological Survey (BGS), Kingsley Dunham Centre, Keyworth, Nottingham, United Kingdom

878 ⁶ School of Ocean Sciences, Bangor University (BU), Wales, United Kingdom

879

880 *Corresponding author:

881 Ana Carolina Pizzochero

882 Universidade Federal do Rio de Janeiro (UFRJ)

883 Centro de Ciências da Saúde (CCS)

884 Instituto de Biofísica Carlos Chagas Filho (IBCCF)

885 Laboratório de Radioisótopos Eduardo Penna Franca (LREPF)

886 Avenida Carlos Chagas Filho, 373 (Edifício do CCS), sala G0-62

887 Cidade Universitária, 21941-902, Rio de Janeiro, RJ, Brazil

888 Tel.: +55 21 25615339, Fax: +55 21 22808193

889 E-mail: pizzochero@biof.ufri.br; pizzocheroac@gmail.com

890

891 **Current address: Ifremer, Centre de Bretagne, REM/EEP, Laboratoire Environnement Profond, F-
892 29280 Plouzané, France.

893

894 **List of Tables**

Table S1.	Details of standards used for dechlorane-related compounds (DRCs), brominated flame retardants (BFRs) and polychlorinated dibenzo- <i>p</i> -dioxins and polychlorinated dibenzofurans (PCDD/Fs) determinations.	3
Table S2.	Limits of detection (LODs; pg g ⁻¹ wet weight) and quantification (LOQs; pg g ⁻¹ wet weight) in whitemouth croaker.	4
Table S3.	Concentrations (pg g ⁻¹ wet weight) of polychlorinated dibenzo- <i>p</i> -dioxins and polychlorinated dibenzofurans (PCDD/Fs) in whitemouth croakers from Guanabara Bay, Southeastern Brazil.	5
Table S4.	Concentrations (pg g ⁻¹ lipid weight) of polychlorinated dibenzo- <i>p</i> -dioxins and polychlorinated dibenzofurans (PCDD/Fs) in whitemouth croakers from Guanabara Bay, Southeastern Brazil.	6
Table S5.	Toxic equivalent (TEQ) concentrations of dibenzo- <i>p</i> -dioxins and polychlorinated dibenzofurans (PCDD/Fs), expressed in pg g ⁻¹ wet weight, using WHO-TEQ 2005 values, in whitemouth croakers from Guanabara Bay, Southeastern Brazil.	7
Table S6.	Concentrations (pg g ⁻¹ wet weight) of brominated flame retardants (BFRs) in whitemouth croakers from Guanabara and Sepetiba bays, Southeastern Brazil.	8
Table S7.	Concentrations (pg g ⁻¹ lipid weight) of brominated flame retardants (BFRs) in whitemouth croakers from Guanabara and Sepetiba bays, Southeastern Brazil.	9
Table S8.	Concentrations (pg g ⁻¹ wet weight) of dechlorane-related compounds (DRCs) in whitemouth croakers from Guanabara and Sepetiba bays, Southeastern Brazil.	10
Table S9.	Concentrations (pg g ⁻¹ lipid weight) of dechlorane-related compounds (DRCs) in whitemouth croakers from Guanabara and Sepetiba bays, Southeastern Brazil.	10
Table S10.	Spearman's correlations coefficients between pollutant levels (or grouped pollutants) and biological parameters or stable isotope ratios	11

895

Table S1. Details of standards used for dechlorane-related compounds (DRCs), brominated flame retardants (BFRs) and polychlorinated dibenzo-*p*-dioxins and polychlorinated dibenzofurans (PCDD/Fs) determinations.

Type	Analytes	Standard Solutions
Surrogate standard solutions (LCS)	DRCs	¹³ C ₁₀ -syn-DP ^a and ¹³ C ₁₀ -anti-DP ^a
	BFRs	BFR-LCS ^b containing: ¹³ C ₁₂ -BDE-28, -47, -77, -99, -100, -126, -153, -154, -169, -183, -197, -205, -207, -209, ¹³ C ₆ -HBB, ¹³ C ₁₂ -BB-153, ¹³ C ₆ -BTBPE.
	PCDD/Fs	EPA-1613LCS ^b containing ¹³ C ₁₂ labeled 2,3,7,8-TCDD, 1,2,3,7,8-PCDD, 1,2,3,4,7,8-HxCDD, 1,2,3, 6,7,8- HxCDD, 1,2,3,4,6,7,8-HpCDD; OCDD, 2,3,7,8-TCDF, 1,2,3,7,8-PCDF, 2,3,4,7,8- PCDF, 1,2,3,4,7,8-HxCDF, 1,2,3,6,7,8-HxCDF, 1,2,3,7,8, 9-HxCDF, 2,3,4,6,7,8-HxCDF, 1,2,3,4,6,7,8-HpCDF, and 1,2,3,4,7,8,9-HpCDF.
Internal Standard Spiking Solution (ISS)	DRCs	BFR-ISS ^b containing: ¹³ C ₁₂ BDE-79, -139, -180, -206
	BFRs	
	PCDD/Fs	EPA1613-ISS ^b containing: ¹³ C ₁₂ -1,2,3,4-TCDD and ¹³ C ₁₂ -1,2,3,7,8,9- HxCDD
Calibration Solutions	DRCs	Five individual calibration solutions prepared from natives (Dec 602 ^c , Dec 603 ^c , Dec 604 ^c , CP ^b , Mirex ^a , <i>syn</i> -DP ^b and <i>anti</i> -DP ^b) and labeled (¹³ C ₁₂ -syn-DP ^a and ¹³ C ₁₂ -anti-DP ^a).
	BFRs	BFR-CVS ^b five individual calibration solutions containing natives (¹² C ₁₂ -BDE-17, -28, -30 -47, -49, -66, -71,-77, -85, -99, -100, -119,-126, -138, -139, -140, -153, -154, -156, -169, -171, -180, -183, -184, -191, -196, -197, -201, -203, -204, -205, -206, -207, -208, -209, PBEB, HBB, BB-153, BTBPE.) and labeled (¹³ C ₁₂ -BDE-28, -47, -77, -79, -99, -100, -126, -139, -153, -154, -169, -180 -183, -197, -205, -206, -207, -209, ¹³ C ₆ -HBB, ¹³ C ₁₂ -BB-153, ¹³ C ₆ -BTBPE)
	PCDD/Fs	EPA 1613CVS ^b five individual calibration solutions containing natives (2,3,7,8-TCDD, 1,2,3,7,8-PCDD, 1,2,3,4,7,8- HxCDD, 1,2,3, 6,7,8- HxCDD, 1,2,3,7,8,9-HxCDD, 1,2,3,4,6,7,8-HpCDD; OCDD, 2,3,7,8-TCDF, 1,2,3,7,8-PCDF, 2,3,4,7,8- PCDF, 1,2,3,4,7,8-HxCDF, 1,2,3,6,7,8-HxCDF, 1,2,3,7,8, 9-HxCDF, 2,3,4,6,7,8-HxCDF, 1,2,3,4,6,7,8-HpCDF, 1,2,3,4,7,8,9-HpCDF; OCDF) and labeled (¹³ C ₁₂ -2,3,7,8-TCDD, -1,2,3,4-TCDD, -1,2,3,7,8-PCDD, -1,2,3,4,7,8- HxCDD, -1,2,3, 6,7,8- HxCDD, -1,2,3,7,8,9- HxCDD, -1,2,3,4,6,7,8-HpCDD, OCDD, -2,3,7,8-TCDF, -1,2,3,7,8-PCDF, -2,3,4,7,8- PCDF, -1,2,3,4,7,8-HxCDF, -1,2,3,6,7,8-HxCDF, -1,2,3,7,8, 9-HxCDF, -2,3,4,6,7,8-HxCDF, -1,2,3,4,6,7,8-HpCDF, and -1,2,3,4,7,8,9-HpCDF

^a Cambridge Isotope Labs (USA) trading house ^b Wellington Labs (Canada) ^c Toronto Research Chemical Inc. (Toronto, ON, Canada).

903 **Table S2.** Limits of detection (LODs; pg g⁻¹ wet weight) and quantification (LOQs; pg g⁻¹ wet weight) in
904 whitemouth croaker.

		LODs	LOQs
Polychlordibenzo-<i>p</i>-dioxin and dibenzofuranss	PCDD/Fs	(pg g ⁻¹)	(pg g ⁻¹)
2,3,7,8-Tetrachlordibenzo- <i>p</i> -dioxin	2,3,7,8-TCDD	0.004	0.013
1,2,3,7,8- Pentachlordibenzo- <i>p</i> -dioxin	1,2,3,7,8-PCDD	0.005	0.017
1,2,3,4,7,8- Hexachlordibenzo- <i>p</i> -dioxin	1,2,3,4,7,8- HxCDD	0.006	0.021
1,2,3, 6,7,8- Hexachlordibenzo- <i>p</i> -dioxin	1,2,3, 6,7,8- HxCDD	0.003	0.010
1,2,3,7,8,9- Hexachlordibenzo- <i>p</i> -dioxin	1,2,3,7,8,9-HxCDD	0.007	0.022
1,2,3,4,6,7,8- Heptachlordibenzo- <i>p</i> -dioxin	1,2,3,4,6,7,8-HpCDD	0.007	0.022
Octachlordibenzo- <i>p</i> -dioxin	OCDD	0.012	0.041
2,3,7,8-Tetrachlordibenzofuran	2,3,7,8-TCDF	0.004	0.013
1,2,3,7,8-Pentachlordibenzofuran	1,2,3,7,8-PCDF	0.003	0.010
2,3,4,7,8- Petachlordibenzofuran	2,3,4,7,8- PCDF	0.006	0.021
1,2,3,4,7,8-Hexachlordibenzofuran	1,2,3,4,7,8-HxCDF	0.005	0.018
1,2,3,6,7,8-Hexachlordibenzofuran	1,2,3,6,7,8-HxCDF	0.005	0.016
1,2,3,7,8, 9-Hexachlordibenzofuran	1,2,3,7,8, 9-HxCDF	0.005	0.018
2,3,4,6,7,8-Hexachlordibenzofuran	2,3,4,6,7,8-HxCDF	0.008	0.027
1,2,3,4,6,7,8-Heptachlordibenzofuran	1,2,3,4,6,7,8-HpCDF	0.004	0.015
1,2,3,4,7,8,9-Heptachlordibenzofuran	1,2,3,4,7,8,9-HpCDF	0.006	0.021
Octachlordibenzofuran	OCDF	0.009	0.029
Brominated flame retardants	BFRs		
2,4,6-Tribromodiphenyl ether	BDE-30	0.023	0.075
2,4',4-Tribromodiphenyl ether	BDE-17	0.015	0.048
2,4,4'-Tribromodiphenyl ether	BDE-28	0.014	0.045
Pentabromoethylbenzene	PBEB	0.006	0.021
Hexabromobenzene	HBB	0.013	0.044
2,2',4,5'&2,3',4',6-Tetrabromodiphenyl ether	BDE-49&71	0.011	0.035
2,2',4,4'-Tetrabromodiphenyl ether	BDE-47	0.004	0.015
2,3',4,4'-Tetrabromodiphenyl ether	BDE-66	0.008	0.026
3,3',4,4'-Tetrabromodiphenyl ether	BDE-77	0.004	0.015
2,2',4,4',6-Pentabromodiphenyl ether	BDE-100	0.019	0.062
2,3',4,4',6-Pentabromodiphenyl ether	BDE-119	0.029	0.097
2,2',4,4',5-Pentabromodiphenyl ether	BDE-99	0.026	0.085
2,2',3,4,4'-Pentabromodiphenyl ether	BDE-85	0.029	0.095
3,3',4,4',5-Pentabromodiphenyl ether	BDE-126	0.025	0.083

2,2',4,4',5,6'-Hexabromodiphenyl ether	BDE-154	0.026	0.087
2,2',4,4',5,5'-Hexabromobiphenyl	BB-153	0.016	0.052
2,2',4,4',5,5'-Hexabromodiphenyl ether	BDE-153	0.027	0.090
2,2',3,4,4',6-Hexabromodiphenyl ether	BDE-139	0.033	0.109
2,2',3,4,4',6'-Hexabromodiphenyl ether	BDE-140	0.041	0.135
2,2',3,4,4',5'-Hexabromodiphenyl ether	BDE-138	0.033	0.110
2,3,3',4,4',5&3,3',4,4',5,5'-Hexabromodiphenyl ether	BDE-156&169	0.046	0.150
2,2',3,4,4',6,6'-Heptabromodiphenyl ether	BDE-184	0.013	0.043
2,2',3,4,4',5',6-Heptabromodiphenyl ether	BDE-183	0.014	0.045
2,3,3',4,4',5',6-Heptabromodiphenyl ether	BDE-191	0.029	0.096
1,2-Bis(2,4,6-tribromophenoxy)ethane	BTBPE	0.375	1.238
2,2',3,4,4',5,5'-Heptabromodiphenyl ether	BDE-180	0.055	0.180
2,2',3,3',4,4',6-Heptabromodiphenyl ether	BDE-171	0.055	0.182
2,2',3,3',4,5',6,6'-Octabromodiphenyl ether	BDE-201	0.130	0.430
2,2',3,4,4',5,6,6'-Octabromodiphenyl ether	BDE-204	0.067	0.220
2,2',3,3',4,4',6,6'-Octabromodiphenyl ether	BDE-197	0.054	0.177
2,2',3,4,4',5,5',6-Octabromodiphenyl ether	BDE-203	0.079	0.261
2,2',3,3',4,4',5,6'-Octabromodiphenyl ether	BDE-196	0.094	0.309
2,3,3',4,4',5,5',6-Octabromodiphenyl ether	BDE-205	0.088	0.291
2,2',3,3',4,5,5',6,6'-Nonabromodiphenyl ether	BDE-208	0.108	0.357
2,2',3,3',4,4',5,6,6'-Nonabromodiphenyl ether	BDE-207	0.074	0.244
2,2',3,3',4,4',5,5',6-Nonabromodiphenyl ether	BDE-206	0.064	0.211
Decabromodiphenyl ether	BDE-209	0.270	0.892

Dechloranes and related compounds

DRCs

Dechlorane 602	Dec 602	0.010	0.034
Dechlorane 603	Dec 603	0.036	0.120
Dechloranes 604	Dec 604	0.169	0.559
syn-Dechlorane 605 or syn-Dechlorane Plus	<i>syn</i> -DP	0.032	0.107
anti-Dechlorane 605 or anti-Dechlorane Plus	<i>anti</i> -DP	0.029	0.094
Chlordene Plus	CP	0.024	0.080
Dechlorane or Hexachlorocyclopentadiene dimer	Mirex	0.007	0.022

905

906

907

908
909

Table S3. Concentrations (pg g⁻¹ wet weight) of polychlorinated dibenzo-*p*-dioxins and polychlorinated dibenzofurans (PCDD/Fs) in whitemouth croakers from Guanabara Bay, Southeastern Brazil. n.d.: not detected; ^a ΣPCDD/F: sum of all the PCDD/Fs

PCDD/Fs	Gb#5	Gb#6	Gb#7	Gb#8	Gb#9	Gb#10	Gb#11	Gb#12	Gb#13	Gb#14
2,3,7,8-TCDD	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
1,2,3,7,8-PeCDD	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
1,2,3,4,7,8-HxCDD	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
1,2,3,6,7,8-HxCDD	0.08	0.05	n.d.	n.d.	n.d.	n.d.	0.12	0.01	n.d.	n.d.
1,2,3,7,8,9-HxCDD	0.09	0.06	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	0.20
1,2,3,4,6,7,8-HpCDD	n.d.	0.12	0.08	n.d.	n.d.	n.d.	0.13	n.d.	n.d.	n.d.
OCDD	0.16	0.27	0.19	n.d.	0.30	0.15	0.26	n.d.	1.25	0.75
2,3,7,8-TCDF	n.d.	n.d.	0.02	n.d.	n.d.	n.d.	0.04	n.d.	0.03	n.d.
1,2,3,7,8-PeCDF	n.d.	n.d.	0.04	n.d.	0.02	n.d.	0.06	n.d.	n.d.	n.d.
2,3,4,7,8-PeCDF	n.d.	0.04	0.05	n.d.	n.d.	n.d.	0.10	n.d.	n.d.	n.d.
1,2,3,4,7,8-HxCDF	n.d.	0.06	n.d.	n.d.	n.d.	n.d.	0.09	n.d.	n.d.	n.d.
1,2,3,6,7,8-HxCDF	n.d.	n.d.	0.03	n.d.	n.d.	n.d.	0.07	n.d.	n.d.	n.d.
2,3,4,6,7,8-HxCDF	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	0.10	n.d.	n.d.	n.d.
1,2,3,7,8,9-HxCDF	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	0.14	n.d.	n.d.	0.18
1,2,3,4,6,7,8-HpCDF	n.d.	n.d.	n.d.	n.d.	0.02	n.d.	0.10	n.d.	0.02	n.d.
1,2,3,4,7,8,9-HpCDF	0.07	0.03	n.d.	n.d.	n.d.	n.d.	0.08	n.d.	n.d.	n.d.
OCDF	0.04	n.d.	0.08	n.d.	n.d.	n.d.	0.12	n.d.	n.d.	0.60
ΣPCDD/F ^a	0.4	0.6	0.5	0.0	0.3	0.2	1.4	0.01	1.3	1.7

910

911 **Table S4.** Concentrations (pg g⁻¹ lipid weight) of polychlorinated dibenzo-*p*-dioxins and polychlorinated dibenzofurans (PCDD/Fs) in whitemouth
912 croakers from Guanabara Bay, Southeastern Brazil. n.d.: not detected; a ΣPCDD/F: sum of all the PCDD/Fs

PCDD/Fs	Gb#5	Gb#6	Gb#7	Gb#8	Gb#9	Gb#10	Gb#11	Gb#12	Gb#13	Gb#14
2,3,7,8-TCDD	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
1,2,3,7,8-PeCDD	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
1,2,3,4,7,8-HxCDD	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
1,2,3,6,7,8-HxCDD	17.2	5.2	n.d.	n.d.	n.d.	n.d.	18.6	1.6	n.d.	n.d.
1,2,3,7,8,9-HxCDD	21.3	6	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	58.9
1,2,3,4,6,7,8-HpCDD	n.d.	12.9	10.2	n.d.	n.d.	n.d.	19.2	n.d.	n.d.	n.d.
OCDD	37.6	27.8	23.5	n.d.	15.8	4.1	39.6	n.d.	173.7	222.2
2,3,7,8-TCDF	n.d.	n.d.	2.3	n.d.	n.d.	n.d.	5.7	n.d.	4.7	n.d.
1,2,3,7,8-PeCDF	n.d.	n.d.	4.4	n.d.	1.0	n.d.	9.3	n.d.	n.d.	n.d.
2,3,4,7,8-PeCDF	n.d.	4	6.6	n.d.	n.d.	n.d.	15.1	n.d.	n.d.	n.d.
1,2,3,4,7,8-HxCDF	n.d.	6.6	n.d.	n.d.	n.d.	n.d.	13.6	n.d.	n.d.	n.d.
1,2,3,6,7,8-HxCDF	n.d.	n.d.	3.8	n.d.	n.d.	n.d.	10.6	n.d.	n.d.	n.d.
2,3,4,6,7,8-HxCDF	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	15.5	n.d.	n.d.	n.d.
1,2,3,7,8,9-HxCDF	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	22.1	n.d.	n.d.	54.1
1,2,3,4,6,7,8-HpCDF	n.d.	n.d.	n.d.	n.d.	1.4	n.d.	17.0	n.d.	2.2	n.d.
1,2,3,4,7,8,9-HpCDF	16	3.6	n.d.	n.d.	n.d.	n.d.	12.6	n.d.	n.d.	n.d.
OCDF	9	n.d.	9.4	n.d.	n.d.	n.d.	18.8	n.d.	n.d.	177.5
ΣPCDD/F^a	101.1	66.1	60.2	0.0	18.2	4.1	217.7	1.6	180.6	512.7

914 **Table S5.** Toxic equivalent (TEQ) concentrations of dibenzo-*p*-dioxins and
915 polychlorinated dibenzofurans (PCDD/Fs), expressed in pg g⁻¹ wet weight, using
916 WHO-TEQ 2005 values, in whitemouth croakers from Guanabara Bay, Southeastern
917 Brazil.

Congener	WHO-TEQ 2005									
	Gb#5	Gb#6	Gb#7	Gb#8	Gb#9	Gb#10	Gb#11	Gb#12	Gb#13	Gb#14
2,3,7,8-TCDD	0.0211 2	0.0266 3	0.0229 1	0.0274 6	0.0294 3	0.0211 3	0.0139 4	0.0144 6	0.0220 1	0.0248
1,2,3,7,8-PeCDD	0.0537 6	0.0441 7	0.0355 0	0.0521 0	0.0333 8	0.0363 8	0.0226 9	0.0332 7	0.0445 8	0.0833
1,2,3,4,7,8-HxCDD	0.0037 1	0.0043 0	0.0070 4	0.0051 3	0.0030 4	0.0024 4	0.0063 8	0.0017 0	0.0022 4	0.0050
1,2,3,6,7,8-HxCDD	0.0075 1	0.0050 6	0.0050 7	0.0051 3	0.0032 1	0.0026 1	0.0121 3	0.0011 0	0.0022 5	0.0041
1,2,3,7,8,9-HxCDD	0.0093 2	0.0058 4	0.0061 3	0.0053 9	0.0032 8	0.0026 6	0.0058 9	0.0018 0	0.0023 6	0.0199
1,2,3,4,6,7,8-HpCDD	0.0002 9	0.0012 5	0.0008 5	0.0003 7	0.0003 5	0.0002 7	0.0012 5	0.0001 6	0.0004 5	0.0008
OCDD	0.0000 8	0.0000 8	0.0000 6	0.0000 4	0.0000 9	0.0000 5	0.0000 8	0.0000 1	0.0003 8	0.0002
2,3,7,8-TCDF	0.0022 4	0.0029 5	0.0019 0	0.0032 2	0.0021 8	0.0028 2	0.0037 4	0.0014 6	0.0033 9	0.0041
1,2,3,7,8-PeCDF	0.0011 5	0.0012 6	0.0011 0	0.0011 4	0.0005 6	0.0008 2	0.0018 2	0.0005 4	0.0006 2	0.0010
2,3,4,7,8-PeCDF	0.0101 7	0.0116 7	0.0163 8	0.0101 4	0.0069 3	0.0080 3	0.0296 3	0.0052 2	0.0060 0	0.0101
1,2,3,4,7,8-HxCDF	0.0026 7	0.0063 6	0.0036 3	0.0033 1	0.0033 7	0.0024 2	0.0088 7	0.0016 1	0.0019 5	0.0044
1,2,3,6,7,8-HxCDF	0.0025 7	0.0031 7	0.0031 3	0.0032 2	0.0033 5	0.0023 3	0.0069 4	0.0015 0	0.0019 3	0.0041
2,3,4,6,7,8-HxCDF	0.0027 0	0.0034 1	0.0076 1	0.0035 5	0.0035 2	0.0025 2	0.0101 1	0.0017 3	0.0020 8	0.0045
1,2,3,7,8,9-HxCDF	0.0037 6	0.0046 1	0.0063 8	0.0045 0	0.0045 8	0.0032 6	0.0144 2	0.0020 6	0.0026 9	0.0183
1,2,3,4,6,7,8-HpCDF	0.0002 4	0.0003 2	0.0002 7	0.0002 8	0.0002 6	0.0001 6	0.0011 1	0.0001 2	0.0001 6	0.0003

1,2,3,4,7,8, 9-HpCDF	0.0007 0	0.0003 5	0.0003 2	0.0004 5	0.0004 0	0.0002 7	0.0008 2	0.0002 2	0.0002 2	0.0005
OCDF	0.0000 1	0.0000 1	0.0000 2	0.0000 2	0.0000 1	0.0000 1	0.0000 4	0.0000 1	0.0000 1	0.0002
ΣTEQ- PCDD/F	0.1220 1	0.1214 3	0.1183 2	0.1254 5	0.0979 5	0.0881 9	0.1398 5	0.0669 3	0.0933 2	0.1856 2
ΣTEQ- PCDD/F	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.2

Brominated flame retardants	Sepetiba Bay						Guanabara Bay													
	Sb#1	Sb#2	Sb#3	Sb#4	Sb#5	Sb#6	Gb#1	Gb#2	Gb#3	Gb#4	Gb#5	Gb#6	Gb#7	Gb#8	Gb#9	Gb#10	Gb#11	Gb#12	Gb#13	Gb#14
BDE-30	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
BDE-17	n.d.	n.d.	n.d.	0.8	n.d.	1.4	n.d.	n.d.	0.3	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
BDE-28	4.6	1.3	n.d.	3.5	1.5	3.5	n.d.	0.1	0.8	0.8	0.2	0.8	n.d.	0.3	n.d.	1.5	n.d.	n.d.	0.1	n.d.
PBEB	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
HBB	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	0.1	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
BDE-49&71	7.0	6.8	10.7	9.8	6.1	9.8	n.d.	0.9	9.4	4.6	4.5	3.6	4.3	1.4	1.0	18.2	2.9	2.3	3.0	1.1
BDE-47	45.3	64.2	59.1	107.7	19.6	48.2	67.5	3.6	60.6	24.3	26.0	19.4	27.3	11.5	7.1	79.6	14.3	6.9	25.3	7.2
BDE-66	2.0	2.6	3.7	3.8	2.9	2.0	0.3	0.3	n.d.	0.9	1.2	0.7	0.4	n.d.	n.d.	3.9	0.3	0.6	0.7	0.2
BDE-77	n.d.	n.d.	n.d.	n.d.	n.d.	0.5	n.d.	n.d.	0.2	0.2	n.d.	0.1	0.2	n.d.	0.1	0.6	0.1	n.d.	n.d.	0.0
BDE-100	8.4	9.6	10.5	10.2	4.4	7.3	10.0	1.1	16.4	6.7	12.8	5.9	8.8	3.9	2.8	23.4	5.5	3.1	9.8	3.2
BDE-119	n.d.	n.d.	1.7	1.0	0.4	n.d.	n.d.	n.d.	n.d.	1.8	n.d.	1.0	1.0	0.4	n.d.	5.1	n.d.	0.5	1.0	n.d.
BDE-99	17.1	9.2	0.5	13.1	1.0	3.7	12.0	n.d.	2.6	1.2	0.2	1.1	n.d.	n.d.	n.d.	4.6	0.6	0.1	0.7	n.d.
BDE-85	n.d.	0.4	0.3	0.2	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	0.5	n.d.	0.1	n.d.	n.d.	0.9	n.d.	n.d.	0.3	n.d.
BDE-126	n.d.	n.d.	0.3	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	0.3	n.d.
BDE-154	1.6	2.3	6.3	4.5	3.1	5.7	n.d.	0.9	6.8	7.4	9.1	4.3	4.0	2.3	1.8	18.7	4.0	1.3	4.1	1.2
BB-153	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
BDE-153	2.3	2.8	2.6	2.2	1.2	2.3	n.d.	0.7	4.6	3.5	4.5	2.3	2.3	1.2	1.1	11.2	2.1	1.6	2.3	0.9
BDE-139	n.d.	n.d.	n.d.	3.4	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	3.2	3.9	n.d.	n.d.	n.d.	0.4	n.d.	n.d.	n.d.
BDE-140	n.d.	n.d.	0.3	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	0.7	0.3	0.3	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	0.2

BDE-138	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
BDE-156&169	n.d.	1.3	3.0	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
BDE-184	n.d.	0.1	n.d.	0.3	0.2	n.d.	n.d.	n.d.	0.2	n.d.	n.d.	0.2	0.2	n.d.	0.4	n.d.	0.2	n.d.	0.1	0.1
BDE-183	3.6	1.0	n.d.	0.3	0.3	n.d.	n.d.	n.d.	0.5	1.7	n.d.	0.2	0.2	n.d.	0.7	n.d.	0.2	n.d.	0.2	0.1
BDE-191	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
BTBPE	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
BDE-180	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
BDE-171	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
BDE-201	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
BDE-204	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
BDE-197	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
BDE-203	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
BDE-196	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
BDE-205	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
BDE-208	26.1	n.d.	1.0	n.d.	n.d.	n.d.	n.d.	n.d.	2.3	0.6	n.d.	n.d.	n.d.	n.d.	0.9	1.3	n.d.	1.4	n.d.	n.d.
BDE-207	35.4	n.d.	1.2	n.d.	n.d.	n.d.	n.d.	n.d.	3.6	0.3	n.d.	n.d.	n.d.	n.d.	0.9	3.9	n.d.	1.7	n.d.	n.d.
BDE-206	36.5	n.d.	0.8	n.d.	n.d.	n.d.	n.d.	n.d.	4.9	0.2	n.d.	n.d.	n.d.	n.d.	n.d.	3.8	n.d.	1.3	n.d.	n.d.
BDE-209	689.8	n.d.	9.1	n.d.	n.d.	n.d.	131.6	n.d.	213.0	13.1	n.d.	n.d.	n.d.	n.d.	6.6	138.5	n.d.	22.8	n.d.	n.d.
ΣPBDE^a	879.7	101.6	111.1	160.8	40.7	84.4	221.4	7.6	326.2	67.3	59.7	43.1	53.0	21.0	23.4	315.2	30.6	43.6	47.9	14.2
ΣBFR^b	879.7	101.6	111.1	160.8	40.7	84.4	221.4	7.6	326.3	67.3	59.7	43.1	53.0	21.0	23.4	315.2	30.6	43.6	47.9	14.2

920 n.d.: not detected; ^a ΣPBDE: sum of 36 PBDE congeners; ^b ΣBFR: sum of PBEB, HBB, BTBPE, BB-153 and PBDEs congeners.

[illegible]

BDE-156&169	n.d.	124	440	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
BDE-184	n.d.	12	n.d.	57	23	n.d.	n.d.	n.d.	50	n.d.	n.d.	17	18	n.d.	21	n.d.	23	n.d.	18	18
BDE-183	2992	91	n.d.	58	42	n.d.	n.d.	n.d.	98	155	n.d.	18	21	n.d.	34	n.d.	27	n.d.	23	28
BDE-191	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
BTBPE	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
BDE-180	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
BDE-171	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
BDE-201	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
BDE-204	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
BDE-197	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
BDE-203	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
BDE-196	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
BDE-205	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
BDE-208	21441	n.d.	143	n.d.	n.d.	n.d.	n.d.	n.d.	487	43	n.d.	n.d.	n.d.	n.d.	47	35	n.d.	201	n.d.	n.d.
BDE-207	29097	n.d.	180	n.d.	n.d.	n.d.	n.d.	n.d.	758	32	n.d.	n.d.	n.d.	n.d.	47	104	n.d.	248	n.d.	n.d.
BDE-206	29971	n.d.	115	n.d.	n.d.	n.d.	n.d.	n.d.	1034	23	n.d.	n.d.	n.d.	n.d.	n.d.	100	n.d.	194	n.d.	n.d.
BDE-209	566710	n.d.	1332	n.d.	n.d.	n.d.	37702	n.d.	45177	1219	n.d.	n.d.	n.d.	n.d.	347	3666	n.d.	3346	n.d.	n.d.
ΣPBDE ^a	722,693	9,515	16,238	31,047	5,500	9,853	63,423	2,642	69,191	6,255	13,650	4,458	6,402	3,867	1,235	8,345	4,692	6,407	6,637	4,180
ΣBFR ^b	722,693	9,515	16,238	31,047	5,500	9,853	63,423	2,642	69,203	6,255	13,650	4,458	6,402	3,867	1,235	8,345	4,692	6,407	6,637	4,180

n.d.: not detected; ^a ΣPBDE: sum of 36 PBDE congeners; ^b ΣBFR: sum of PBEB, HBB, BTBPE, BB-153 and PBDEs congeners.

922

923

924 **Table S8.** Concentrations (pg g⁻¹ wet weight) of dechlorane-related compounds (DRCs) in whitemouth croakers from Guanabara and Sepetiba bays, Southeastern Brazil.

Dechlorane-related compounds	Sepetiba Bay						Guanabara Bay													
	Sb#1	Sb#2	Sb#3	Sb#4	Sb#5	Sb#6	Gb#1	Gb#2	Gb#3	Gb#4	Gb#5	Gb#6	Gb#7	Gb#8	Gb#9	Gb#10	Gb#11	Gb#12	Gb#13	Gb#14
Mirex	n.d.	n.d.	0.10	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	0.18	0.82	1.74	n.d.	1.16	n.d.	4.22	0.43	n.d.	0.90	n.d.
Dec 602	0.06	n.d.	0.49	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	1.41	2.12	0.37	0.24	0.63	n.d.	7.03	1.2	0.21	0.15	n.d.
Dec 603	1.67	1.07	2.29	1.0	1.11	2.19	n.d.	0.70	2.48	4.46	5.50	2.59	2.47	2.32	2.05	25.97	3.27	0.83	1.67	0.61
Dec 604	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
CP	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	0.10	n.d.	n.d.	n.d.	n.d.	n.d.	0.9	n.d.	n.d.	n.d.	n.d.
<i>syn</i> -DP	12.92	n.d.	1.11	0.77	0.68	n.d.	n.d.	n.d.	0.30	0.46	n.d.	0.15	n.d.	n.d.	n.d.	1.54	0.66	n.d.	n.d.	n.d.
<i>anti</i> -DP	19.18	n.d.	1.15	1.07	2.02	2.28	n.d.	n.d.	0.21	0.37	1.24	0.29	n.d.	n.d.	n.d.	2.05	1.01	n.d.	0.52	0.48
ΣDRC^a	33.8	1.1	5.1	2.8	3.8	4.5	-	0.7	3.0	7.0	9.7	5.1	2.7	4.1	2.1	41.7	6.6	1.0	3.2	1.1
DP ^b	32.1	-	2.3	1.8	2.7	2.3	-	-	0.5	0.8	1.2	0.4	-	-	-	3.6	1.7	-	0.5	0.5
<i>f_{anti}</i> ^c	0.6	-	0.5	0.6	0.7	1.0	-	-	0.4	0.4	1.0	0.7	-	-	-	0.6	0.6	-	1.0	1.0

925 n.d.: not detected; ^a ΣDRC: sum of Mirex, Dec 602, Dec 603, Dec 604, CP, *syn*-DP and *anti*-DP; ^b DP: sum of *syn*- and *anti*-DP; ^c *f_{anti}*: *anti*-DP divided by DP.

926

927

928

929

930 **Table S9.** Concentrations (pg g⁻¹ lipid weight) of dechlorane-related compounds (DRCs) in whitemouth croakers from Guanabara and Sepetiba bays, Southeastern Brazil.

Dechlorane-related compounds	Sepetiba Bay						Guanabara Bay													
	Sb#1	Sb#2	Sb#3	Sb#4	Sb#5	Sb#6	Gb#1	Gb#2	Gb#3	Gb#4	Gb#5	Gb#6	Gb#7	Gb#8	Gb#9	Gb#10	Gb#11	Gb#12	Gb#13	Gb#14
Mirex	n.d.	n.d.	15	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	17	187	181	n.d.	215	n.d.	112	66	n.d.	123	n.d.
Dec 602	46	n.d.	71	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	131	485	38	29	115	n.d.	185	186	23	22	n.d.
Dec 603	1370	101	334	192	149	256	n.d.	242	526	414	1257	268	299	429	108	688	502	126	231	180
Dec 604	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
CP	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	10	n.d.	n.d.	n.d.	n.d.	n.d.	24	n.d.	n.d.	n.d.	n.d.
<i>syn</i> -DP	10613	n.d.	163	149	92	n.d.	n.d.	n.d.	63	43	n.d.	16	n.d.	n.d.	n.d.	41	102	n.d.	n.d.	n.d.
<i>anti</i> -DP	15761	n.d.	168	206	276	266	n.d.	n.d.	45	34	284	30	n.d.	n.d.	n.d.	54	154	n.d.	72	143
ΣDRC ^a	27,790	101	751	547	518	522	-	242	634	649	2,213	533	328	759	108	1,104	1,010	149	448	323
DP ^b	26,374	-	331	355	368	266	-	-	108	77	284	46	-	-	-	95	256	-	72	143
<i>f_{anti}</i> ^c	0.6	-	0.5	0.6	0.7	1.0	-	-	0.4	0.4	1.0	0.7	-	-	-	0.6	0.6	-	1.0	1.0

931 n.d.: not detected; ^a ΣDRC: sum of Mirex, Dec 602, Dec 603, Dec 604, CP, *syn*-DP and *anti*-DP; ^b DP: sum of *syn*- and *anti*-DP; ^c *f_{anti}*: *anti*-DP divided by DP.

932

933

934

935 **Table S10.** Spearman's correlations coefficients between pollutant levels (or grouped pollutants) and biological parameters or stable isotope ratios.

936

	Σ PBDE	Tri-BDE	Tetra-BDE	Penta-BDE	Hexa-BDE	Hepta-BDE	Dec 603	anti-DP	DP	Σ DRC	Σ PCDD/F [#]	Σ TEQ-PCDD/F [#]
Lipid content (%)	0.01	0.14	0.13	-0.05	0.37	0.19	0.45*	-0.07	-0.002	0.20	-0.36	-0.43
Mass	-0.41	-0.38	-0.40	-0.31	-0.20	0.07	-0.05	-0.08	-0.11	-0.02	0.37	-0.06
Size	-0.48*	-0.45*	-0.54*	-0.40	-0.09	-0.02	0.21	-0.15	-0.18	0.10	0.34	-0.06
$\delta^{13}\text{C}$	0.33	0.48*	0.36	0.29	0.08	0.30	-0.22	0.48*	0.47*	0.12	0.06	0.18
$\delta^{15}\text{N}$	0.34	0.45*	0.50*	0.45*	0.38	0.10	0.12	0.62*	0.59*	0.46*	0.06	0.29
$\delta^{34}\text{S}$	-0.22	-0.48*	-0.24	-0.17	0.18	-0.32	0.47*	-0.21	-0.24	0.12	0	-0.10

937 * Significant correlation ($p < 0.05$)

938 [#] n = 10 specimens

939