Mangrove area and propagule number planting targets produce sub-optimal rehabilitation and afforestation outcomes

Wodehouse, Dominic C.J.; Rayment, Mark B.

Estuarine, Coastal and Shelf Science

DOI:

[10.1016/j.ecss.2019.04.003](https://doi.org/10.1016/j.ecss.2019.04.003)

Published: 30/06/2019

Peer reviewed version

[Cyswllt i'r cyhoeddiad / Link to publication](https://doi.org/10.1016/j.ecss.2019.04.003)

Dyfyniad o'r fersiwn a gyhoeddwyd / Citation for published version (APA):

Wodehouse, D. C. J., & Rayment, M. B. (2019). Mangrove area and propagule number planting targets produce sub-optimal rehabilitation and afforestation outcomes. *Estuarine, Coastal and Shelf Science*, 222, 91-102. <https://doi.org/10.1016/j.ecss.2019.04.003>

Hawliau Cyffredinol / General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Mangrove Area and Propagule Number Planting Targets Produce Sub-Optimal Rehabilitation and Afforestation Outcomes

Dominic C.J. Wodehouse^{1,2}, Mark B. Rayment¹

¹ School of Natural Sciences, Thoday Building, Bangor University, Deiniol Rd,
Bangor, Wales, LL57 2UW, UK.

² Mangrove Action Project. PO Box 1854. Port Angeles. WA. 98362-0279. USA.

Corresponding author: d.wodehouse@bangor.ac.uk +44 1248 382 281

Academic Editors: Dr Tim Jennerjahn and Dr Martin Zimmer

ABSTRACT

Mangrove rehabilitation projects often fail completely or fail to meet their objectives. This study examines village-level rehabilitation planting carried out in 13 villages (119 rehabilitation attempts at 74 sites) across two countries in southeast Asia, to assess village-level rehabilitation effectiveness, and to identify what factors influenced outcomes. Mean propagule survival across all rehabilitation attempts was 20% with a median of 10%. Sixty six percent of attempts had a survival rate of less than 20%. Mid mangrove zone projects were more successful (mean 30%) than rehabilitation projects at other elevations. Planting on mudflats, representing 32% of rehabilitation / afforestation attempts, achieved only a 1.4% propagule survival rate. The overall low success rate was due to several inter-related factors. Poor site/species matching on high and low elevation sites was common; for example,

Rhizophora spp. was used alone or in combination at least 65% of the time, including on mudflats where this genus is ecologically unlikely to establish. Site selection was often driven by the desire to achieve centrally defined area or propagule *planting* targets, rather than *survivorship* targets, and thus required large, uncontested project areas. Conversely, the presence of natural regeneration, even if in small amounts, was associated with higher than average success. Therefore, it was estimated that only 16% of planting attempts were actually necessary.

Highlights

- Mean survival of all mangrove rehabilitation planting attempts was 20%, median 10%
- Only 16% of planting attempts deemed necessary vs. natural recruitment potential
- Better hydrological connection significantly improved survivorship (mid/high zones)
- Area or propagule planting targets should be changed to survivorship targets

Keywords: mangrove restoration, mangroves, failure, survivorship, area target, planting target, Thailand, Philippines

1. Introduction

1.1 Mangroves are Particularly Beneficial to the Poorest Coastal Villagers

Mangroves form highly productive ecosystems, (Alongi, 2009) which provide many direct and indirect benefits and services (Moberg and Rönnbäck, 2003; Saenger, 2002; van Oudenhoven et al., 2015). These services are particularly valuable and relevant to the poorest members of coastal villages (Glaser and da Silva Oliveira, 2004; Kairo et al., 2001; Springate-Baginski and Than, 2011; Stevenson et al., 1999; Sunderlin et al., 2005). They include a nursery function for fish and shrimp (Saenger et al., 2013; Salmo III et al., 2018) and provision of wood for construction and fuel for cooking (Moberg and Rönnbäck, 2003).

1.2 Mangrove Losses Declining, but Measurement is Challenging

In previous decades, management of mangrove loss has proved challenging. This was as a result of low-resolution remote sensing imagery and of classifying mangrove as opposed to other types of vegetation. Furthermore, there has been a lack of a definitive methodology for identifying and classifying 'mangrove' at different levels of degradation and canopy cover (Giri et al., 2011; Hamilton and Casey, 2016). Within the 21st century, global losses have been reported to be 0.16% - 0.39% per year, indicating a slowing of the rate of loss from the 1980s (0.99%) and 1990s (0.7%) (Hamilton and Casey, 2016).

Southeast Asia has historically seen some of the greatest losses. For example, mangrove cover in the Philippines has fallen from 450,000 ha in 1900 to 120,000 ha in 1995 but the rate of loss has slowed to 0.1% per year since 2000 (Long et al., 2014; Primavera and Esteban, 2008). In Thailand, using 1961 as a baseline, less

than half of all the mangroves remain (Aksornkoae, 2004). From 2000 - 2012, Thai losses were 0.69% per year (Hamilton and Casey, 2016).

1.3 Mangrove Rehabilitation Initiated for Many Reasons and by a Variety of Actors

Following natural disasters such as the Indian Ocean tsunami of 2004 and cyclone Haiyan / Yolanda in the Philippines, and to recover some of the goods and services provided by mangroves, there have been significant efforts to rehabilitate areas of converted or degraded mangroves (Aung et al., 2011; Primavera et al., 2011; Primavera and Esteban, 2008). This effort has included attempts to afforest new areas that were previously not inhabited by mangroves. Most often, governments have commissioned mangrove rehabilitation programs through forestry and environment departments, which in turn have sometimes engaged local villages to assist. Occasionally, communities and villages have initiated their own rehabilitation projects. Examples include Pred Nai in Thailand (Fisher, 2000; Senyk, 2005) and Myanmar (Springate-Baginski and Than, 2011). There are also reports of rehabilitation by individuals within coastal villages, in the Philippines (Walters, 2004, 2000, 1997; Walters et al., 2005) and Thailand (pers. obs.).

1.4 ‘Restoration’ or ‘Rehabilitation’?

The scientific literature often uses ‘restoration’ and ‘rehabilitation’ interchangeably (van Oudenhoven et al., 2015) or uses other words including repair, reclamation, reforestation, conservation, afforestation or eco-development (Duke, 1996). Restoration might be defined as recovering an area back to an assumed original ‘pristine’ ecosystem, implicitly including the restoration of mangrove functionality (Kairo et al., 2001; Stevenson et al., 1999; Walters et al., 2008). However, the word

is often used more broadly (McDonald et al., 2016). Rehabilitation is an attempt to recover *some* of the ecosystem functions or to find another stable use for the land (McDonald et al., 2016; Stevenson et al., 1999; Walters, 2008; but see Field, 1999a). The debate concerning these terms continues (see Dale et al., 2014 for a review). While acknowledging this debate, and the need for clarity of definition for legal purposes and for setting expectations (Dale et al., 2014), this paper will use the term rehabilitation. Similarly, care is needed when describing areas as ‘degraded’, because the perception of whether an area is partially degraded or not is affected by cultural expectation and land management intensity (Hobbs, 2016). Furthermore, changes to an ecosystem’s state may be adjustments beyond those caused by normal forest growth and development processes, leading to a new equilibrium as a result of climate change or long-term variation of weather patterns (Hobbs, 2016; Mansourian et al., 2017).

1.5 What is ‘Successful’ Rehabilitation?

In principle project outcomes should be assessed in relation to stated project objectives, and this is crucial in the planning of any mangrove rehabilitation work (Field, 1999b; Lewis, 2000; Saenger, 2002). When viewing rehabilitation outcomes from a narrow standpoint, ‘success’ may be claimed after five to seven years, because this indicates probable long-term survivorship (i.e. to reproductive maturity) and eventual (re)establishment of a mangrove stand (Bosire et al., 2008; Kodikara et al., 2017). Salmo III et al., (2013) focused on vegetation and soil parameters in a study of monospecific plantations. Their study suggested that mangrove ecosystem stability might be reached by 11 years, and that ecological characteristics resembled natural mangroves after 25 years. Other indicators of success have focused on the whole ecosystem and consider that success can be claimed when the hydrological

normality of a mangrove has returned (Asaeda et al., 2016). Alternatively rehabilitation assessment might compare project sites to natural mangroves (McDonald et al., 2016) but not in terms of succession (Ellison, 2000). Despite sometimes being an appropriate long-term measure, comparing project sites to old-growth mangroves is particularly difficult in countries such as Thailand and the Philippines. This is because much of the natural mangroves have been cut-over for charcoal or fuelwood and replanted with a less diverse range of species (Alongi, 2002). Resources permitting, a more comprehensive approach is an *ecological rehabilitation* perspective (the literature often uses *restoration* in this case) (Asaeda et al., 2016; Ellison, 2000; Lewis, 2005; Walters et al., 2008). This approach looks for the return of full ecosystem function, including outflow of organic material to, and habitat connectivity with, linked seagrass and coral systems. This can be relatively complete within five years (Saenger et al., 2013).

While social factors are pertinent, here we focus on a strictly biological (or silvicultural) definition of success – whether planted seeds / propagules survive to establishment.

1.6 Rehabilitation and Afforestation Successes and Failures

There have been positive mangrove afforestation survivorship outcomes in Bangladesh (Saenger and Siddiqi, 1993, but see Moberg and Rönnbäck, 2003) and successful mangrove rehabilitation in Florida (Brockmeyer et al., 1996; Lewis, 2005; Lewis and Gilmore, 2007), Philippines (Asaeda et al., 2016; Primavera et al., 2012; Walters, 2004) Indonesia, (Lewis and Brown, 2014) and Myanmar (pers. obs.). However, many rehabilitation projects fail completely or do not achieve their objectives (Elliott et al., 2016; Erftemeijer and Lewis, 1999; Field, 1996; IUCN, 2017;

Lewis, 2005; UNEP, 2007) or at best produce limited positive results (Alongi, 2002; Aung et al., 2011; Barbier, 2006; Ellison, 2000; Memon and Chandio, 2011; Moberg and Rönnbäck, 2003). Mangrove rehabilitation projects that have become established often resemble even-age class, mono-specific plantations rather than natural mangrove (Bosire et al., 2006; Ellison, 2000; Field, 1996; Lewis, 2005), bearing little or no similarity to the original mangrove (Alongi, 2002). In addition they exhibit only limited species zoning and biodiversity (Bosire et al., 2008). However, of greater immediate concern are the often extremely low propagule survival rates of these rehabilitation programs.

Sanyal (1998) reported that in West Bengal more than 9,000 ha were planted with only 1.5% probable survival. In the Philippines, despite significant efforts and financial inputs over the last twenty years, survival of planted mangroves remains low at 10-20% (Primavera, 2015; Primavera and Esteban, 2008; Samson and Rollon, 2008; Walters, 1997). Similar conclusions have been drawn from Sri Lankan rehabilitation programs (Samarakoon, 2012).

1.7 Technical Reasons for Previous Failures

Why do so many rehabilitation projects fail? Here we consider the suggestion that the most common technical reason for planting failure is poor site/species matching, i.e. choosing an unsuitable species to plant for a given site (Aung et al., 2011; Bosire et al., 2006; Lewis et al., 2016; Primavera and Esteban, 2008; Walters et al., 2008). What is meant by unsuitable? Individual species have differing tolerances to specific biogeochemical factors and gradients present across the intertidal area (Saenger, 2002). These include salinity, soil type, soil anoxia, sulphate levels, nutrient levels, pH, wave energy, temperature, light levels, inundation regimes (Alongi, 2009;

Tomlinson, 2016; van Loon et al., 2016), tides and wind distribution of propagules and seeds (van der Stocken et al., 2012), and species-selective predation by herbivores (Elster, 2000; Sousa et al., 2003). Species therefore exhibit differing 'preferences' for elevation and location within the intertidal zone (Duke, 2006; Snedaker, 1982; Tomlinson, 2016).

Closely related to site/species matching is poor site choice. Insufficient regard is often given to understanding local hydrology, topography relative to sea level, and the effects these have on soil conditions. These features greatly affect planting outcomes (Aung et al., 2011; Elster, 2000; Hashim et al., 2010; Kairo et al., 2001; Lewis, 2005). Duration of inundation is particularly important (van Loon et al., 2016). In some cases rehabilitation can be achieved simply by reconnecting or improving site hydrology, e.g. by installing culverts under a road, reconnecting former aquaculture ponds or reopening lagoons (Brown et al., 2014; Elster, 2000; Ferreira and Lacerda, 2016; Lewis, 2014; Twilley et al., 1999). In contrast, sites that have an elevation below that of a natural front mangrove fringe are likely to have permanently saturated soil with poor drainage, leading to anoxic and potentially acidic soil (Holguin et al., 2001; Kristensen and Alongi, 2006). These factors have a significant negative impact on the outcomes of projects attempting to afforest mudflats and seagrass beds (Asaeda et al., 2016; Samson and Rollon, 2008; Stevenson et al., 1999).

Many rehabilitation projects start planting first without fully understanding the original cause of mangrove loss or why there is no natural regeneration on site (Asaeda et al., 2016; Lewis, 2005, 2000; Walters et al., 2008). Both of these factors might be mitigated by reducing and removing mangrove stressors specific to a site, such as obstructed hydrology or unsustainable anthropogenic activities (e.g. harvesting of

187 mangrove wood) (Lewis et al., 2016). Other reported reasons for failure include
188 herbivore grazing and footfall damage, poor planting method, lack of aftercare (e.g.
189 weeding) and monitoring (Kodikara et al., 2017), barnacle infestation and high wave
190 energy (i.e. inappropriate site choice).

191 In this paper we describe a study of community-level planting projects to assess
192 survival rates and to identify factors that determine success or failure at the project
193 level.

2. Materials and Methods

2.1 Location of Study Sites

Although specific regional and local contexts are very important and highly variable, multiple villages within two countries were studied in an attempt to produce some general conclusions. Thailand and the Philippines share the same Indo-Malesia bio-geo-climatic zone within the Indo-West Pacific (Duke, 2006; Tomlinson, 2016). They have extensive mangrove areas, on which a substantial proportion of the coastal inhabitants depend for their livelihoods and food (Balmford et al., 2002). Since 1945, both countries have experienced significant mangrove conversion to aquaculture and degradation for charcoal and fuelwood production, among other causes (Richards and Friess, 2016).

Table 1 lists the Thai and the Philippine villages studied in this large-scale investigation. This study combined ecological and social research to examine mangrove rehabilitation in the context of biophysical, silvicultural and social factors. Villages were chosen because they were located either within or near to an extensive riverine mangrove delta or had a significant area of mangroves nearby. In all cases mangroves were considered an important village resource and were used in some ways by a substantial part of the village population. Finally, village members had attempted mangrove rehabilitation or afforestation in the past. The exception to these selection criteria was village P3A in the Philippines (Table 1) which was included as it had conducted a record-setting ‘1 million propagules in an hour’ planting project (Escandor, 2012). Except for this final record-setting planting, all Philippine planting discussed here was funded by the National Greening Program (Department of Environment and Natural Resources, 2016), a large-scale bio-shield

establishment scheme initiated after typhoon Yolanda / Haiyan in 2013. Some rehabilitation or afforestation sites had been attempted more than once and each attempt was assessed. A site was defined as an individual plot or area villagers had attempted to rehabilitate or afforest as a discrete project. In total 119 attempts at rehabilitating 74 sites were assessed.

Table 1 Site information on mangrove rehabilitation projects evaluated in Thailand and Philippines

Thailand							
Village Number	1	2	3	4	5	6	
Village Code	T1A	T1B	T2A	T2B	T3A	T3B	
Approx. Lat Long	6.89° N 99.79° E	6.89° N 99.79° E	7.89° N, 99.16° E	7.89° N, 99.16° E	8.44° N, 99.96° E	8.44° N, 99.96° E	
Province	Satun	Satun	Krabi	Krabi	Nakorn Sri Thammarat	Nakorn Sri Thammarat	
Number of Rehabilitation Sites Assessed	13	11	9	6	5	3	
Village Mangrove Area (Ha) ¹	407	592	319	176	3,894	257	
Approximate Research Dates	Sept – Nov 2013	Dec 2013 – Feb 2014	Feb – May 2014	June – Aug 2014	Oct 2014 – Feb 2015	Feb – May 2015	

Philippines							
Village Number	1	2	3	4	5	6	7
Village Code	P1A	P1B	P2A	P2B	P2C	P2D	P3A
Approx. Lat Long	10.81° N, 119.5° E	10.81° N, 119.5° E	14.06° N 123.3° E	14.3° N 123.3° E	13.9° N, 123.2° E	14.0° N, 123.2° W	13.8° N, 122.8° E
Province	Northern Palawan	Northern Palawan	San Miguel Bay, Luzon	San Miguel Bay, Luzon	San Miguel Bay, Luzon	San Miguel Bay, Luzon	Camarines Sur, Luzon
Number of Rehabilitation Sites Assessed	10	8	3	3	1	1	1
Village Mangrove Area (Ha) ¹	126	856	Unknown	Unknown	Unknown	Unknown	NA
Approximate Research Dates	Sept – Dec 2015	Jan – Apr 2016	May 2016	May 2016	May 2016	May 2016	May 2016

1. Source: Local Dept. for Marine and Coastal Resources field offices (Thailand) and Dept. of Environment and Natural Resources field office (Philippines).

Many mangrove rehabilitation projects were attempted by Thailand's Department for Marine and Coastal Resources (DMCR) and the Philippine Department of Environment and Natural Resources (DENR) in the 1980s and 1990s. However, this study looked only at more recent planting from approximately 2007 onwards, which involved participation by local villages. Assessment of village P1A's (Philippines) rehabilitation ability was expanded because this village was commissioned by the government to plant not only within its own territory, but also in neighbouring villages.

232 All accessible P1A-rehabilitated sites were assessed because the planting team and
233 the techniques used were the same. Some rehabilitation was carried out in both
234 Thailand and the Philippines while the lead author was present in the village, (T1B
235 Jan 2014, T3A Dec 2014, Oct 2015) providing an opportunity to act as an observer
236 and witness techniques.

2.2 Assessment Method

2.2.1 Initial Visits with Village Mangrove Expert

During initial scoping interviews with villagers and village leaders, opinions were sought to ascertain which resident was most knowledgeable about their mangroves. In all villages consensus about a mangrove expert readily emerged, thus negating the need to perform a village expert ranking exercise (Davis and Wagner, 2003; Chalmers and Fabricius, 2007). All rehabilitation sites were then visited initially with the village mangrove expert to record site history, reasons for the previous degradation, history of the rehabilitation effort(s), planting dates, details of site preparation, silvicultural practice and species choice. Site details recorded included presence / absence of trees, presence / absence of natural regeneration (indicating whether a site might naturally regenerate on its own) and hydrological connectivity. Also recorded were site elevation (section 2.2.3), soil type (sand, silt or clay), presence of standing water and post-hoc interventions such as the use of fencing. In addition other factors likely to affect rehabilitation and plant establishment were noted, such as evidence or presence of grazing livestock or trampling damage. Soil salinity was measured either from available soil pore water or groundwater sourced from minor excavations up to 15cm deep (Bellingham and Stanley handheld refractometer). However, it should be noted that it was not always possible to measure salinity in some of the high mangrove zone assessments because of a lack of available soil water. This might have skewed the resulting analysis. The direction of this potential skew is uncertain. The locations of the boundaries of all the rehabilitation sites were recorded via a handheld GPS (Garmin 62stc). Subsequently these GPS waypoints were employed to calculate the area of each site using Google Earth Pro. All site features were photographed.

Because several sites were planted more than once, a distinction has been made between 'attempt' (n = 119) and 'site' (n = 74). Wherever possible, all previous attempts on the same site were evaluated (38% of assessments) as well as the final (or only) attempt on a site (62%). Seventy-five of the attempts were in Thailand, 44 in the Philippines. The majority of the rehabilitation sites were <0.3 ha, ranging from 0.001 – 50 ha (SD $\pm$ 7.73 ha). The cumulative total area assessed was 164 ha.

2.2.2 Mangrove Establishment: Counts, Extrapolations and Area Calculations

Although there are a range of factors that could be measured when assessing rehabilitation (see Dale et al., 2014 for a review), propagule or seedling survival is an unequivocal measure of whether the plants had managed to establish and survive or not. Presence or absence of natural regeneration was noted at the time of assessment - distinguishable from planted material by not being in lines, unevenly spaced, without canes and often of a pioneer species - but which did not contribute to survival scores. Plant health and vigour was also noted at the time of assessment. Three techniques were used to assess survival depending on different planting ages and types of sites.

Preferentially, we used a 'full count' method for more recent planting events as both Thai and Philippine villages usually used canes which indicated where planting material had been inserted. Planting was frequently conducted in straight lines and even spacing. Therefore, for more recent planting projects (i.e. less than 1-2 years old) in less exposed sites, missing or absent plants were immediately obvious due to the resulting gap left in the lines of plants. Where possible every plant was counted for each generation of planting (if applicable). However, ten large-scale planting attempts were too extensive to allow each surviving plant to be counted. These

286 extensive sites were stratified by elevation, exposure and channel edge / interior.
287 Sub-plots were assessed to incorporate all significant variation of a site in order to
288 achieve a minimum sample of at least 10% of the surviving plants.

289 Where full count inventory was not possible, we extrapolated counts from surviving
290 patches of planting to the whole site. Some rehabilitation sites were too small or too
291 fragmented to justify planting in lines. If present, surviving patches indicated how
292 densely the site had been planted originally. In combination with a site history and
293 the opinion of the village expert, total numbers planted were estimated and
294 contrasted with survivors present, to produce a survivorship percentage for each
295 generation of planting (if applicable).

296 In some cases, particularly on mudflat sites, there was often little trace of planting
297 activity, or insufficient survivors to assess survivorship either via the 'full count'
298 method or by extrapolation of surviving patches. Therefore we asked the village
299 expert and participants of the planting to indicate as accurately as possible the
300 boundary of the planted area, which was then marked by GPS. We then counted
301 every surviving plant within this defined planting area. Subsequently Google Earth
302 Pro was used to determine the area of the planting site. The stated planting
303 spacing, normally 2x2m, was then used to calculate numbers originally planted. By
304 dividing the number of surviving plants by the estimated number that had originally
305 been planted, a survivorship percentage could be produced. For example, a
306 planting area described by the village expert, marked by GPS, drawn as a polygon
307 on Google Earth Pro might be revealed to cover 10 ha. If the stated planting density
308 was 2x2 m spacing, this area would have originally had 25,000 plants. If the
309 survivors counted within this defined planting area numbered 500, the survivorship
310 was 2%.

311 These three different methods of survivorship assessment engendered different
312 levels of confidence. To reconcile possible differences between methods, a post-hoc
313 resampling was conducted using the extensive field photography to reassess all 74
314 sites. On the assumption that the 'full count' method produced the most accurate,
315 bias-free estimate of survival, we recalibrated 'extrapolation from surviving patches'
316 against the 'full count'. This post-hoc reassessment suggested that there was only a
317 minor under-estimation of survival at low levels of survival, and a corresponding
318 minor over-estimation of survival at higher levels of survival. Therefore, once
319 reconciled, all three data sets were subsequently treated in the same manner. 'Full
320 count' method was used for 38% assessments, 'extrapolation from surviving
321 patches' 52% and 'counts within a defined area' 10% of attempts.

322 **2.2.3 Mangrove Zones and the Quality of the Hydrological Connection**

323 On any of the sites examined, several biotic and abiotic gradients were potentially
324 affecting where mangroves lived, resulting in distinct bands of species. Most
325 influential among these factors was the frequency and duration of a site's inundation
326 due to its elevation relative to sea level (van Loon et al., 2016). Following Duke
327 (2006) and Tomlinson (2016), bands of mangrove species were classified into three
328 zones. The 'low' zone, which started at approximately mean sea level, received
329 inundation at high tides >45 times a month and was characterised by species such
330 as *Sonneratia alba* and *Avicennia alba*. 'Mid' zones were inundated by normal high
331 tides 20 to 45 times a month and were the home of *Rhizophora* spp. and *Ceriops*
332 *tagal*. 'High' or 'back' zones received inundation <20 times a month at high tide and
333 included back mangrove species such as *Heritiera littoralis*, *Lumnitzera* spp.,
334 *Scyphiphora hydrophyllacea* and *Acrostichum* spp. Mudflats, which normally

occurred between lowest water and mean sea level, were inundated by every high tide.

Table 2 illustrates the distribution of rehabilitation attempts within these mangrove zones. Most rehabilitation or afforestation attempts were on mid mangrove areas (52%), mudflats (30%), together with high zone areas (13%).

Table 2. Frequency of rehabilitation attempt by mangrove zone, by country

		Position of Rehabilitation Attempt Relative to Tidal Inundation Regime				Total
		Mudflat	Low Zone	Mid Zone	High Zone	
Country	Thailand	7	1	52	15	75
	Philippines	29	4	10	1	44
Total		36	5	62	16	119

Mid and high zone rehabilitation sites varied greatly in hydrological connection to tidal flushing, thus elevation per se was not necessarily a good indicator. Instead better hydrological connection was judged by the following indicators:

- greater number of days a month the site was inundated, according to the village expert
- many seeds / propagules present on the ground not directly under a potential seeding tree (indicating that inundations were able to transport them onto the site)
- greater presence of established mangrove natural regeneration (indicating that soil drainage was adequate for plant growth)
- wet soil and other evidence of the area having been recently inundated (e.g. visible tide line)

- a lack of extensive areas of standing water (indicating better drainage and suggesting a better quality of soil, as saturated soils are less well suited for mangrove establishment)
- a lack of visible salt crystals on the soil surface (indicating that sufficient inundation was avoiding a build-up of salt – a stressor for all mangroves)
- limited plant / tree stress indicators (e.g. canopy die-back, stunted plants, abnormally small leaves, a proliferation of prop roots on *Rhizophora* sp.)
- fewer dead leaves on the ground (indicating that they had been washed away)
- no significant debris within the channels, e.g. from cutting for charcoal production (debris would slow water flows, inhibit the distribution of seeds and propagules and increase the chance of sedimentation within the channels)

A qualitative decision was made by weighting all the above criteria equally. We classified each site's hydrological connectivity as either 'good' or 'partial / poor' based on the preponderance of indicators of good connection compared to indicators of poor connection. Mudflats and lower mangrove elevations by definition have good connectivity to the local hydrology and therefore were not assessed for the quality of their hydrological connection.

2.2.4 Time Since Planting

One hundred and nineteen attempts at mangrove rehabilitation or afforestation were evaluated over 74 sites. Of these attempts 36 were assessed less than 12 months after planting. We attempted to achieve a balance between including the maximum amount of data possible yet avoiding false-positives by excluding planting that had not yet had sufficient time to either establish or fail to establish. The cut-off was set at one year. The exception to this cut-off period was planting attempts where

survivorship was $\leq 5\%$ (14 planting attempts) as the planting within these attempts had already clearly failed. Therefore 97 attempts were retained for analysis.

Table 3. Time between planting and assessment, by frequency of rehabilitation attempt

Time between Planting Attempt and Assessment	All Attempts Assessed	Planting Attempts Retained for Further Analysis
< 1 Year	36	14
13 – 24 Months	29	29
> 2 Years	54	54
Total	119	97

2.2.5 Criteria for Judging Whether Planting was Required

An assessment was made as to whether each mangrove rehabilitation site might have regenerated naturally, whether rehabilitation should never have been attempted at that site, or whether planting was necessary and appropriate. Whether planting was necessary and appropriate or not was assessed by the following indicators:

- an absence of natural regeneration and / or a lack of successful establishment of natural regeneration
- appropriate site elevation for mangrove establishment relative to sea level, and resulting inundation regime, (i.e. within either low, mid or high mangrove zones, with duration and frequency of flooding, as described in section 2.2.3.). Mudflats, extending from approximately mean sea level down to lowest water, hydrological channels and areas of standing water were deemed inappropriate places for planting and ecologically unsuitable for mangrove establishment (Lewis, 2005)
- low expected wave energy (the assumption was that young plants that were subject to significant wave energy will be damaged or uprooted and washed

399 away.) Assessment included evidence of erosion, whether the site was
400 directly open to the sea and the opinion of a local mangrove expert

- 401 ● soil mechanically firm enough to anchor a propagule or seedling
- 402 appropriately, not so soft as to allow the researcher to sink into the mud up to
- 403 the knees
- 404 ● minimal levels of significantly-sized debris on site (which might be lifted by the
- 405 tide and damage vulnerable plants)
- 406 ● no inhibiting social factors such as uncontrolled animal grazing, boat impact,
- 407 damage from footfall or destruction from the use of damaging fishing gear
- 408 which scrapes along the sediment surface uprooting natural regeneration

409 A qualitative decision was made by weighting all the above criteria equally, and
410 classifying each site as either requiring planting, able to naturally regenerate on its
411 own, or an inappropriate site that will never become mangrove, based on the
412 preponderance of indicators described above.

3. Results

3.1 Site Descriptors and Demographics for All 119 Attempts

Hydrological connection, by definition, was complete for mudflats and low zone mangroves. However, for mid and high mangrove zones, some only had partial / poor connection and drainage (section 2.2.3) with a limited exchange of water at each tidal flushing, and areas of standing water. Of all the attempts within mid and high mangrove zones (n = 80) only a quarter (26%) had a good hydrological connection. There was no evidence that any measures had been taken to improve hydrological connection in those sites with partial / poor connection. Occasionally, hydrology was made worse (e.g. village T2A), by skimming the grass off a site with a bulldozer during site preparation, thereby filling the drainage channels in the process. On other sites, previous tree felling for charcoal had left brush and debris in the channels (e.g. village P1A), slowing the flow of water and increasing sedimentation in the channels.

High and mid zone mangrove soil salinities (both 27ppt, SD ± 2 and SD ± 8 respectively) were slightly less than low zone salinity (33ppt, SD ± 2.3), which in turn was less than sea water (normally approximately 35ppt). Partial / poor hydrology appeared not to affect average mangrove soil salinity as much as the presence of fresh water input from rivers flowing into mangrove deltas.

A majority of rehabilitation attempts (65%) 'direct planted' propagules into the soil. Thirteen percent of attempts (all in the Philippines) used 'wildlings', young plants with 2-5 leaf pairs, pulled out of their original location and transplanted as bare-root stock. Eleven percent of rehabilitation attempts used polybagged seedlings. Rehabilitation was left to natural regeneration in only two instances, which have been included in

the analysis because using this form of rehabilitation was a conscious decision on the part of the village (T2A) conservation group.

Fig. 1 illustrates in which zone the different types of planting material were used. This broad distribution suggests there was little relationship between planting material used and site elevation. Direct planting of propagules was the most common across all species except *Nypa fruticans*.

Fig. 1. Frequency of mangrove zone planted, by planting material used. (Natural regeneration has been excluded for clarity.)

By species, *Rhizophora* spp. was used in the majority of planting; alone in 52% of attempts and in conjunction with other mid mangrove species (e.g. *Ceriops tagal*, *Bruguiera* spp.) another 13% of the time.

Other mid mangrove species such as *Ceriops tagal* and *Bruguiera* spp. were planted 19% of the time (Fig. 2) . 'Mix' (n = 6) denotes when a selection of (rarely more than three) species was used from more than one mangrove zone. These often but not always included *Rhizophora* spp., along with *C. tagal*, *Bruguiera* spp. and very

occasionally mangrove associate *Pandanus tectorius* (Kitamura et al., 1998; Tomlinson, 2016).

Fig. 2. Proportions of mangrove species used by mangrove zone.

3.2. Rehabilitation Successes and Failures

All analyses from this point onwards excludes the 22 attempts assessed as ‘too recent to judge’ (section 2.2.4) unless otherwise stated. For this reduced subset of rehabilitation attempts (n = 97), the mean survival rate was 20% (SD $\pm$ 23.4) with a median of 10%, Fig. 3, the median or middle score being less affected by a non-normal or skewed distribution of data and extreme scores (Field, 2018).

Fig. 3. Percentage survival rates by frequency.

The difference between 13-24 month and > 2-year planting survivorship was not significantly different (p = 0.54), indicating that most propagule death occurred within the first year after planting, Table 4.

Table 4. Mean survivorship by time between planting and assessment

Mean Survivorship by Time Between	< 1 Year	13-24 months	> 2 Years	Total
-----------------------------------	----------	--------------	-----------	-------

Planting and Assessment				
Mean Survival %	1.6	23.9	22.8	
SD	2	27.4	22	
N	14	29	54	97

475

476 Mean planting survival varied significantly ($p < 0.001$) between Thailand (26%, SD
477 ± 24.3 , $n = 58$) and the Philippines (11%, SD ± 18.8 , $n = 39$). This reflected a
478 tendency to attempt to afforest mudflats in the Philippines. The mean survival for
479 mudflats was low (1.4%, SD ± 3.6 , $n = 31$) compared to mid mangrove zones (30.1%,
480 SD ± 22.5 , $n = 48$) or high mangrove zones (25%, SD ± 28.3 , $n = 13$). Salinity
481 exhibited a significant inverse relationship with planting survivorship ($p < 0.001$).

482 Fig. 4 shows the survival rates by mangrove species. The mean survival rate of
483 *Rhizophora* spp. was 11% (SD ± 20.4 , $n = 50$), despite being the most popular choice
484 for planting. The establishment of *Nypa fruticans* was similarly poor (9%, SD ± 10.4 ,
485 $n = 5$). Other mid-mangrove species fared better with a mean survival of 29% (SD
486 ± 26.8 , $n = 20$), as did 'Mix' (i.e. a range of species from more than one mangrove
487 zone, 46.2%, SD ± 22.8 , $n = 6$).

488

489
490
491
492
493
494
495
496
497
498
499
500
501

Fig. 4. Survival rates by species planted.

While there was no significant difference in survival between directly planted propagules and bagged plants, ($p = 0.32$), there was a significant difference between propagules and wildlings ($p = 0.024$), and between bagged plants and wildlings, ($p = 0.022$), Fig 5.

Bagged plants (mean survival 27.6%, SD ± 26.6 , $n = 9$) were either *Nypa fruticans*, *Ceriops tagal* or very occasionally mangrove associate *Pandanus tectorius*.

Otherwise, planting was ‘direct planting’ of propagules into the substrate (mean survival 20.2%, SD ± 23.9 , $n = 62$). Transplanted ‘wildlings’ (bare root stock, always *Rhizophora* spp., mean survival 5.4%, SD ± 8.6 , $n = 14$) were only used in the Philippines.

502
503
504
505

Fig. 5. Planting survival by type of planting material used.

Within the mid and high zone mangrove areas, mean survival of planted material was significantly lower ($p = 0.038$) in sites with partial / poor hydrological connection

at 24.8% (SD $\pm$ 21.9, n = 48), compared to areas with good hydrology where the survival rate was 39.9% (SD $\pm$ 26, n = 15).

3.3 Interactions Between Factors

Table 2 shows that mudflats and mid mangrove elevations were frequently chosen as sites for afforestation / rehabilitation planting. Mudflats proved resistant to planting whatever form of planting material was employed. However, mid and high zone sites had better survival of all planted materials, and bagged plants (n = 9) in particular, Fig. 6.

Fig. 6. Planting survival by mangrove zone, by planting material.
(Planting material category 'Don't Know' (n = 10) and 'Mixed' (n = 2) have been omitted for clarity.)

Fig. 2 (section 3.1) indicated the proportions of different species used for planting, at differing elevations. Fig. 7 illustrates that on mudflats and in back mangrove zones, *Rhizophora* sp. (n = 52) was not an appropriate genus to use. However, when

524 *Rhizophora* sp. was used in zones suitable for this genus (low and mid zones), its
525 survival rate improved but was no more successful than ‘Other Mid-Mangrove
526 Species’ (n = 21) which was only used in mid-to-back elevations. *Nypa fruticans* (n
527 = 5) also performed poorly on mudflats.

Fig. 7. Survival of species planted by mangrove zone.
(Mixed species (n = 6), natural regeneration (n = 2) have been omitted for clarity.)

3.4 Was Planting Necessary?

Planting survivorship was significantly higher ($p < 0.001$) when natural regeneration was present on a rehabilitation site (mean survival 26.3%, SD ± 20.6 , $n = 51$) compared to when there was no natural regeneration present (mean survival 13%, SD ± 24.4 , $n = 46$).

Using the criteria described previously (section 2.2.5) natural regeneration would have been sufficient, and planting unnecessary in 37% of attempts, largely within mid and back mangrove zones. Another 47% of attempts 'will never be mangrove' because of inappropriate hydrology or being located at an unsuitable inter-tidal elevation. For clarity of depiction, these two categories have been combined in Fig. 8 to contrast against the 16% of planting attempts that were considered to have been necessary, by mangrove zone.

Fig. 8. Frequency of unnecessary/inappropriate planting against necessary planting, by mangrove zone.

4. Discussion

4.1 Research Method Limitations

More sites were assessed in Thailand than in the Philippines. Where possible, information from the village expert was cross-checked against interviews with government mangrove agency field office staff, villagers, and by direct observation, to reduce recall error.

Plant health, vigour and biomass characteristics were not included in survivorship assessment, but were taken into account for the quality of hydrological connection, the appropriateness of the site and species choice, and whether or not a site required planting. This inevitably involved an element of judgement and site interpretation, and consideration of factors such as the frequent seasonal floods in southern Thailand and watershed-scale hydrological disturbance in villages T3A and T3B (Osbeck et al., 2010; Prabnarong and Kaewrat, 2006), or the reduction of precipitation due to the 'El Nino Southern Oscillation' event that occurred during the research period (L'Heureux et al., 2017).

Separating and discarding 22 planting attempts which were 'too early to judge' (section 2.2.4) from those which had had 'enough time' to establish or fail, may have negatively affected survivorship results, but may also have removed potential real positives as well as false positives.

Although benchmarking against other mangroves (McDonald et al., 2016) might have been suitable in countries where there is pristine mangrove nearby, Thailand and the Philippines have very little mangrove which has not been replanted after charcoal / fuelwood concessions, subjected to species-selective felling or had natural *Sonneratia* / *Avicennia* forests replaced by *Rhizophora* spp. planting, such as in

Banacon Island, Philippines, or Pak Phanang Bay, Thailand (Macintosh et al., 2002; Osbeck et al., 2010; Walters, 2005). Mangrove workers who have suggested that rehabilitation projects should aim for and be judged by ecological rehabilitation criteria (Asaeda et al., 2016; Ellison, 2000; Lewis, 2000; Walters et al., 2008) are by implication working towards the conditions which allow the return of full ecosystem function (Saenger et al., 2013). Although appropriate in theory, using such criteria presents a practical problem as a result of the extensive time needed between planting and full recovery of ecosystem function. To have a chance of returning an area to functioning mangrove forest, the initial planting must first survive any transplant shock and establish itself. The data presented here only describe this initial establishment. We acknowledge that planting which might become established and grow into a new stand and might therefore be deemed successful, could still potentially fail to deliver the full suite of ecosystem benefits. Examples of this later failure include mangroves used to stabilise the walls of aquaculture ponds but which have little hydrological connection, or when mangroves are planted in drainage channels which block the local hydrological connection, leading to eventual ecosystem failure.

4.2 Discussion of the Results

The majority of sites that would probably have recovered through natural regeneration alone were mid and back mangrove, Fig. 8. Within these zones, some areas viewed by the villagers as 'degraded' were simply mangroves with natural gaps and desirable forest complexity. However, because they were seen as degraded they were re-planted, which sometimes included clearing biodiverse natural regeneration and 'crown lifting' of existing non-*Rhizophora* sp. trees (Walters, 2004). Only a few sites that had previously been mangrove before being cleared or

degraded, normally for charcoal, were not regenerating (16%). Typically, this was due to poor hydrology, hard smooth soil making the retention of 'volunteer' propagules / seeds difficult or for other reasons such as a lack of fencing to keep out grazing animals (Field, 1996). In these cases planting was necessary and might ultimately facilitate quicker mangrove succession (Ferreira and Lacerda, 2016; Lewis et al., 2016), but with no guarantee of success because the other site-specific mangrove stressors, discussed here and in section 2.2.5, were often not resolved.

If planted sites would have regenerated on their own, without planting intervention as demonstrated by Lewis (2005) and Brown et al. (2014), this could avoid planting costs and liberate financial and labour resources for other management tasks.

Generally, the presence of natural regeneration is a good indicator that a potential site is suitable for rehabilitation. However, natural regeneration can also start to establish in hydrological channels following the failure of the local hydrology.

Similarly, although some of the mudflat afforestation attempts were situated near the fringe of existing mangrove, and hence were interspersed with a limited amount of pioneer species natural regeneration, this did not mean that these sites were potential mangrove areas. In short, open mudflats and mangrove drainage channels (47% of 97 attempts) were not ecologically appropriate sites, and rehabilitation / afforestation should not have been attempted in these locations.

The mangrove zone within the inter-tidal range runs from above mean sea level (Alongi, 2002; Kairo et al., 2001; Lewis, 2005) or upper third (Saenger, 2002) to highest high water. Knowledge of mangrove species zoning is essential for successful rehabilitation (Kairo et al., 2001). So-called site / species matching has been offered as a key reason for planting failure – i.e. planting inappropriate species for a given site and its inherent conditions (Alongi, 2002; Aung et al., 2011; Bosire et

al., 2006; IUCN, 2017; Kodikara et al., 2017; Primavera and Esteban, 2008; Saenger, 2002; Walters et al., 2008). The failure to improve planting performance despite increased financial spend by NGO-led projects in the Philippines (Samson and Rollon, 2008; Walters, 2008) or the 1bn Peso (£14m) 'National Greening Programme' in the Philippines (Ranada, 2015) is in part due to the frequent planting of mid zone *Rhizophora* sp. in all zones (Fig. 2). This is possibly because its propagules are easy to collect and handle and do not require growing-on in a nursery (Lewis, 2014; Primavera, 2015; Primavera et al., 2011; Primavera and Esteban, 2008). The research described here demonstrates the improved success rates associated with planting the correct species for the specific mangrove zone (Fig.7).

While acknowledging the challenges of hydrological assessment (van Loon et al., 2016), an understanding of site hydrology, topography and drainage, and the effects these have on salinity and the species chosen, is vital for successful mangrove rehabilitation (Aung et al., 2011; Elliott et al., 2016; Elster, 2000; Hashim et al., 2010; Kairo et al., 2001; Lewis, 2005; Oh et al., 2017). Some sites can be restored simply by hydrological reconnection or improvement if propagules are available from nearby stands via hydrochory (Prach and del Moral, 2015; Stevenson et al., 1999). Unlike Elster's Colombian experience (2000) and Brown et al., (2014) in Indonesia, hydrology was rarely considered at our study sites, having been discussed only once at one Thai site (village T3A). Occasionally site hydrology was made worse by inappropriate site preparation. This study has documented the significant difference improved / adequate hydrology makes to rehabilitation success. This therefore suggests that many of the mid and back mangrove sites would have benefited from improved hydrological connectivity and drainage. However, guidelines for hydrologic

648 rehabilitation are sparse and communication between researchers and mangrove
649 managers appears to have been insufficient to change rehabilitation activities.

650 Although there was no significant difference in the survival rates of directly planted
651 propagules and bagged seedlings (Fig. 5), extrapolation of these results should be
652 done with caution. Bagged seedlings tended to be *Ceriops tagal* and *Nypa fruticans*
653 not *Rhizophora* sp. and were likely to be used in a more appropriate zone (Fig. 6).
654 However, planting of bagged *N. fruticans* on mudflats resulted in total mortality
655 (village T3B). Bagged material was only deployed if it was provided by the
656 government, rather than for ecological or silvicultural reasons and used much less
657 often than direct planting of propagules (section 3.1). Excluding special cases, the
658 resulting small sample sizes were too small to make further analysis appropriate.

659 Clump planting propagules close together (i.e. < 30cm apart) to allow planted
660 material to benefit from a mutually improved rhizosphere (Chan and Baba, 2010;
661 Lavieren et al., 2012) was never attempted. Root-balled 'wildlings' were not
662 attempted by any group. Bare-root wildlings were only used in the Philippines,
663 where villagers and government staff believed they were more reliable than
664 propagules. Contrary to this local belief, bare-root wildlings were significantly less
665 likely to establish than other planting material (Fig. 5). However, because these
666 wildlings were most frequently deployed on mudflats, their very low survival (mean =
667 5.4%) also found by Primavera et al. (2011), cannot definitively be ascribed to bare-
668 root wildlings being an intrinsically poor silvicultural method. Furthermore, poor
669 handling, for example allowing exposed roots to dry out in direct sunlight before
670 being re-planted, cannot be ruled out. Poor survival of directly planted propagules
671 might also have resulted from propagules being collected from trees before maturity,

672 and planters having only a partial understanding of the effects of pests such as
673 *Poecilips fallax* beetle on propagules.

674 Protection from storms and strong winds is often a key motivator for mangrove
675 planting and afforestation, particularly in the Philippines where village planting was
676 funded by the 'National Greening Programme' (Department of Environment and
677 Natural Resources, 2016). In the medium-term, the rehabilitation projects described
678 here will produce densely stocked, even-aged plantations with limited structural
679 complexity. This lack of complexity should be a cause for concern as research has
680 shown that older plantation stands of *Rhizophora* spp. are more vulnerable to strong
681 winds than other species. Furthermore, they have a poor ability to recover from
682 storm damage because they lack latent buds and cannot re-grow from the base
683 when the stem is damaged (Bosire et al., 2008; Salmo III et al., 2014; Villamayor et
684 al., 2016). In addition the smooth canopy of an even-aged class stand slows wind
685 less than a mixed-aged stand of uneven height (Villamayor et al., 2016). Structural
686 complexity is characterised by a number of forest attributes such as basal area, tree
687 height, tree species, tree density, biomass, foliage arrangement, canopy cover and
688 understory (McElhinny et al., 2005). This complexity develops over time but could be
689 accelerated through planting a diversity of species at a variety of spatial densities.

690 In order to implement the 'National Greening Programme', the Department of
691 Environment and Natural Resources of the Philippines passes down extensive
692 planting area quotas to the department's field offices. To fulfil these quotas, mudflats
693 are frequently selected as they offer the necessary spatial extent (Primavera, 2015).
694 Although mudflats in both countries might have been considered silviculturally
695 inappropriate, these areas typically have uncontested land tenure (for a description
696 of the land tenure issues, see Primavera et al., 2015, 2011; Samson and Rollon,

2011). They are therefore easily available, as other researchers have reported (Lewis and Brown, 2014; Primavera, 2015; Primavera et al., 2011; Samson and Rollon, 2008; Walters et al., 2008). Thus despite evidence in the published scientific literature, rehabilitation manuals and national media (Primavera, 2015; Primavera et al., 2011; Ranada, 2015), planting continues on mudflats, and sometimes even seagrass beds, even though mean survival rates were shown here to be <2%. Mudflats are particularly valuable for feeding shorebirds, producing income for local gleaners and food security (Primavera et al., 2011). Therefore on the rare occasions that mudflat planting survived, normally due to rapid accretion or deposition of sediment (pers. obs.), the value of substituting one ecosystem for another has been questioned (Erftemeijer and Lewis, 1999; Lewis, 2005).

Similarly in Thailand much of the mangrove management activity was driven by national propagule planting targets delegated to the mangrove agency field offices. These targets originated from successive four-year National Economic and Social Development Plans (for example, Office of the National Economic and Social Development Board, 2011, 2001). Field offices also received additional directives such as planting 840,000 propagules to celebrate a national event (National News Bureau of Thailand, 2016). Furthermore there was often a general desire by villages to carry out communal planting activity on national holidays. However, some field offices are starting to negotiate the return of aquaculture ponds which had been illegally established within the mangroves and other encroached former mangrove areas. Consequently, more planting was carried out in mid and high mangrove zones (section 3.1). Although the overall success rate was higher, the question remains as to how much of the planting was actually necessary.

This paper and others (Dale et al., 2014; Lewis, 2005; Primavera and Esteban, 2008; Salmo III et al., 2007; Samson and Rollon, 2008;) have suggested that, despite being largely unnecessary, planting has tended to dominate mangrove management activity. This is typically endorsed at the national level. Area planting targets set by the Philippines' National Greening Programme have produced sub-optimal outcomes, and planting has also arguably received too much emphasis in Thailand. Although such target-driven planting provides quantifiable measures (Mansourian et al., 2017), this is unlikely to be aligned with silvicultural best practices. Propagule survivorship would be a more appropriate measure, perhaps combined with an emphasis on recovering abandoned aquaculture ponds. The area of abandoned ponds in Thailand and the Philippines is not known, but in Indonesia alone there is estimated to be around 250,000 ha (Gusmawati et al., 2017). Aquaculture ponds are frequently located in mid and high zone mangrove areas which this study and others have shown to be a more appropriate elevation for mangrove rehabilitation. Restoring hydrological connectivity to these abandoned ponds to rehabilitate them back to functioning mangrove ecosystems (Primavera et al., 2011; Villamayor et al., 2016) would arguably be a more appropriate management task, particularly over the coming decades as sea level rise requires mangroves to retreat landward and upward (Gilman et al., 2008; Primavera et al., 2011).

5. Conclusion

This research suggests that attention to a few key factors can enhance rehabilitation outcomes. First, mangrove workers should ensure that the appropriate species are planted in the mangrove zone for which they are best suited. Second, appropriate

745 hydrological connectivity with good tidal flushing and drainage improves project
746 outcomes. Third, it is suggested that much mangrove rehabilitation is either
747 unnecessary or conducted on sites which are inappropriate. Fourth, attempted
748 afforestation of mudflat sites usually fails and is not recommended. Finally,
749 rehabilitation projects should focus on survivorship rather than meeting area or
750 propagule number targets which typically produce sub-optimal outcomes.

751

6. Acknowledgements

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

The authors would like to pass on their most heart-felt thanks to the Editors of ECSS and the three anonymous reviewers whose inputs greatly improved this article.

Sincere thanks go to many people who assisted this research: Assoc. Prof. Boosya Bunnag and Dr Pavinee 'Jin' Patanachan of King Mongkut University for acting as Thai supervisors and helping with village selection; the National Research Council of Thailand; the directors and staff of the Royal Thai Government's Department Marine and Coastal Resources mangrove field offices for their generosity with their time and knowledge; Dr Antonio Dano and Dr Portia Lapitan from the Philippine Ecosystems Research and Development Bureau and John Pontillas of the Palawan Council for Sustainable Development and the members of the Malampaya Sound Protected Area Management Board for permission to conduct this research.

Thank you also to Khun Phonsawan 'Gift' Jinowat, Khun Kamondham 'Tuan' Truatmakha and Khun Petchrung 'Aey' Sukpong in Thailand; Valiente Milan and Jevy Simion in Palawan and Agnes 'Ging' Faustino and Jerry Qutorio for their insightful interpretation; Khun Dear, Yosuke Fukushima san and Khun Namtan 'Namy' Kunja for helping with the transects; Jim Enright and Khun Jaruan 'Ning' Kaewmahanin of NGO Mangrove Action Project for the use of the office, research advice and long-term support; Assoc. Prof. Dan Friess of the National University of Singapore for his GIS expertise; Dr Jean Yong of the Singapore University of Technology and Design for his mangrove species identification PDF; Prof. Rene Rollon of University of Philippines and Assit. Prof. Sev Salmo III of Ateneo University

776 for their research advice. Finally thank you to the excellent people in the mangrove
777 villages studied, for their generosity and willingness to share their lives.

778 **7. Declaration of Interests**

779 The lead author is a volunteer for NGO Mangrove Action Project (MAP) and
780 periodically carries out paid ecological mangrove rehabilitation teaching on its behalf.
781 MAP did not contribute in any way to the funding, design, analysis or writing of this
782 paper. MAP has not run training for either DMCR or DENR. The second author
783 declares no conflict of interest.

784 **8. Authors' Contributions**

785 Dominic Wodehouse carried out the fieldwork, analysis and the writing of the paper.
786 Mark Rayment developed the research questions, designed the research matrix,
787 contributed several data capture techniques, added to the statistical analysis and
788 edited the paper.

789

9. References

- Aksornkoae, S., 2004. Sustainable Use and Conservation of Mangrove Forest Resources with Emphasis on Policy and Management Practices in Thailand, in: Vanucci, M. (Ed.), Mangrove Management and Conservation : Present and Future. United Nations Press, New York, USA., pp. 149–160.
- Alongi, D.M., 2009. Paradigm shifts in mangrove biology, in: Perillo, G.M.E., Wolanski, E., Cahoon, D.R., Brinson, M.M. (Eds.), Coastal Wetlands An Integrated Ecosystem Approach, Coastal Wetlands: An Integrated Ecosystems Approach. Elsevier, Amsterdam, pp. 615–640. <https://doi.org/10.1016/B978-0-444-53103-2.00022-3>
- Alongi, D.M., 2002. Present state and future of the world's mangrove forests. Environ. Conserv. 29, 331–349. <https://doi.org/10.1017/S0376892902000231>
- Asaeda, T., Barnuevo, A., Sanjaya, K., Fortes, M.D., Kanesaka, Y., Wolanski, E., 2016. Mangrove plantation over a limestone reef - Good for the ecology? Estuar. Coast. Shelf Sci. 173, 57–64. <https://doi.org/10.1016/j.ecss.2016.02.017>
- Aung, T.T., Than, M.M., Katsuhiko, O., Yukira, M., 2011. Assessing the status of three mangrove species restored by the local community in the cyclone-affected area of the Ayeyarwady Delta, Myanmar. Wetl. Ecol. Manag. 19, 195–208. <https://doi.org/10.1007/s11273-011-9211-9>
- Balmford, A., Bruner, A., Cooper, P., Costanza, R., Farber, S., Green, R.E., Jenkins, M., Jefferiss, P., Jessamy, V., Madden, J., Munro, K., Myers, N., Naeem, S., Paavola, J., Rayment, M., Rosendo, S., Roughgarden, J., Trumper, K., Kerry Turner, R., 2002. Economic Reasons for Conserving Wild Nature. Science (80). 297, 950–953.

812 <https://doi.org/10.1126/science.1073947>

813 Barbier, E.B., 2006. Natural barriers to natural disasters: Replanting mangroves after the
814 tsunami. *Front. Ecol. Environ.* 4, 124–131. [https://doi.org/10.1890/1540-](https://doi.org/10.1890/1540-9295(2006)004[0124:NBTNDR]2.0.CO;2)
815 9295(2006)004[0124:NBTNDR]2.0.CO;2

816 Bosire, J.O., Dahdouh-Guebas, F., Kairo, J.G., Wartel, S., Kazungu, J., Koedam, N., 2006.
817 Success rates of recruited tree species and their contribution to the structural
818 development of reforested mangrove stands. *Mar. Ecol. Prog. Ser.* 325, 85–91.
819 <https://doi.org/10.3354/meps325085>

820 Bosire, J.O., Dahdouh-Guebas, F., Walton, M., Crona, B.I., Lewis, R.R., Field, C., Kairo, J.G.,
821 Koedam, N., 2008. Functionality of restored mangroves: A review. *Aquat. Bot.* 89, 251–
822 259. <https://doi.org/10.1016/j.aquabot.2008.03.010>

823 Brockmeyer, R.E., Rey, J.R., Virnstein, R.W., Gilmore, R.G., Earnest, L., 1996. Rehabilitation of
824 impounded estuarine wetlands by hydrologic reconnection to the Indian River Lagoon,
825 Florida (USA). *Wetl. Ecol. Manag.* 4, 93–109. <https://doi.org/10.1007/BF01876231>

826 Brown, B., Fadillah, R., Nurdin, Y., Soulsby, I., Ahmad, R., 2014. Community Based Ecological
827 Mangrove Rehabilitation (CBEMR) in Indonesia. *S.a.p.i.e.n.s. Surv. Perspect. Integr.*
828 *Environ. Soc.* 7, 1–13.

829 Chalmers, N., Fabricius, C., 2007. Expert and Generalist Local Knowledge about Land-cover
830 Change on South Africa’s Wild Coast: Can Local Ecological Knowledge Add Value to
831 Science ? *Ecol. Soc.* 12, 10. <https://doi.org/10.1042/BST0391533>

832 Chan, H.T., Baba, S., 2010. Manual on Guidelines for Rehabilitation of Coastal Forests

833 Damaged by Natural Hazards in the Asia-Pacific Region. International Society for
 834 Mangrove Ecosystems and International Tropical Timber Organization, Okinawa, Japan.

835 Dale, P.E.R., Knight, J.M., Dwyer, P.G., 2014. Mangrove rehabilitation: a review focusing on
 836 ecological and institutional issues. *Wetl. Ecol. Manag.* 22, 587–604.
 837 <https://doi.org/10.1007/s11273-014-9383-1>

838 Davis, A., Wagner, J.R., 2003. Who knows? On the importance of identifying “experts” when
 839 researching local ecological knowledge. *Hum. Ecol.* 31, 463–489.
 840 <https://doi.org/10.1023/A:1025075923297>

841 Department of Environment and Natural Resources, 2016. Mangrove, beach forest
 842 rehabilitation in Yolanda-hit areas completed. [WWW Document]. URL
 843 [http://oldsite.denr.gov.ph/news-and-features/latest-news/2465-mangrove-beach-](http://oldsite.denr.gov.ph/news-and-features/latest-news/2465-mangrove-beach-forest-rehab-in-yolanda-hit-areas-completed.html)
 844 [forest-rehab-in-yolanda-hit-areas-completed.html](http://oldsite.denr.gov.ph/news-and-features/latest-news/2465-mangrove-beach-forest-rehab-in-yolanda-hit-areas-completed.html) (accessed 2.3.18).

845 Duke, N., 1996. Mangrove reforestation in Panama: an evaluation of planting in areas
 846 deforested by a large oil spill, in: Field, C.D. (Ed.), *Restoration of Mangrove Ecosystems*,
 847 *Restoration of Mangrove Ecosystems*. International Society for Mangrove Ecosystems,
 848 Okinawa, Japan., pp. 209–232.

849 Duke, N.C., 2006. *Australia’s Mangroves: The Authoritative Guide to Australia’s Mangrove*
 850 *Plants*. University of Queensland, Brisbane, Australia.

851 Elliott, M., Mander, L., Mazik, K., Simenstad, C., Valesini, F., Whitfield, A., Wolanski, E., 2016.
 852 *Ecoengineering with Ecohydrology: Successes and failures in estuarine restoration*.
 853 *Estuar. Coast. Shelf Sci.* 176, 12–35. <https://doi.org/10.1016/j.ecss.2016.04.003>

854 Ellison, A.M., 2000. Mangrove restoration: Do we know enough? *Restor. Ecol.* 8, 219–229.
855 <https://doi.org/10.1046/j.1526-100X.2000.80033.x>

856 Elster, C., 2000. Reasons for reforestation success and failure with three mangrove species
857 in Colombia. *For. Ecol. Manage.* 131, 201–214. [https://doi.org/10.1016/S0378-](https://doi.org/10.1016/S0378-1127(99)00214-5)
858 [1127\(99\)00214-5](https://doi.org/10.1016/S0378-1127(99)00214-5)

859 Erftemeijer, P., Lewis, R.R., 1999. Planting Mangroves on Intertidal Mudflat: Habitat
860 Rehabilitation or Habitat Conversion?, in: ECOTONE-VIII Seminar “Enhancing Coastal
861 Ecosystem Restoration for the 21st Century.” FAO, Ranong, Thailand, p. 9.
862 [https://doi.org/ISBN: 978-92-807-3173-6](https://doi.org/ISBN:978-92-807-3173-6)

863 Escandor, J.J., 2012. In Camarines Sur, it took an hour to plant 1M mangroves [WWW
864 Document]. *Inquirer.Net*. URL [http://newsinfo.inquirer.net/159243/in-camarines-sur-](http://newsinfo.inquirer.net/159243/in-camarines-sur-it-took-an-hour-to-plant-1m-mangroves)
865 [it-took-an-hour-to-plant-1m-mangroves](http://newsinfo.inquirer.net/159243/in-camarines-sur-it-took-an-hour-to-plant-1m-mangroves) (accessed 9.24.17).

866 Ferreira, A.C., Lacerda, L.D., 2016. Degradation and conservation of Brazilian mangroves,
867 status and perspectives. *Ocean Coast. Manag.* 125, 38–46.
868 <https://doi.org/10.1016/j.ocecoaman.2016.03.011>

869 Field, A., 2018. *Discovering statistics using IBM SPSS Statistics*, 5th ed. SAGE Publications
870 Ltd., London, UK.

871 Field, C.D., 1999a. Mangrove rehabilitation: Choice and necessity. *Hydrobiologia* 413, 47–52.
872 <https://doi.org/10.1023/A:1003863415354>

873 Field, C.D., 1999b. Rehabilitation of mangrove ecosystems: An overview. *Mar. Pollut. Bull.*
874 37, 383–392. [https://doi.org/10.1016/S0025-326X\(99\)00106-X](https://doi.org/10.1016/S0025-326X(99)00106-X)

875 Field, C.D. (Ed.), 1996. Restoration of mangrove ecosystems, 1st ed. International Tropical
876 Timber Organization, Okinawa, Japan.

877 Fisher, R.J., 2000. Poverty Alleviation and Forests: Experiences from Asia, in: In Workshop
878 'Forest Ecospaces, Biodiversity and Environmental Security, Vol. 5. RECOFT, Bangkok,
879 Thailand.

880 Gilman, E.L., Ellison, J., Duke, N.C., Field, C., 2008. Threats to mangroves from climate
881 change and adaptation options: A review. *Aquat. Bot.* 89, 237–250.
882 <https://doi.org/10.1016/j.aquabot.2007.12.009>

883 Giri, C., Ochieng, E., Tieszen, L.L., Zhu, Z., Singh, A., Loveland, T., Masek, J., Duke, N.C., 2011.
884 Status and distribution of mangrove forests of the world using earth observation
885 satellite data. *Glob. Ecol. Biogeogr.* 20, 154–159. [https://doi.org/10.1111/j.1466-](https://doi.org/10.1111/j.1466-8238.2010.00584.x)
886 [8238.2010.00584.x](https://doi.org/10.1111/j.1466-8238.2010.00584.x)

887 Glaser, M., da Silva Oliveira, R., 2004. Prospects for the co-management of mangrove
888 ecosystems on the North Brazilian coast: Whose rights, whose duties and whose
889 priorities?, in: *Natural Resources Forum*. Wiley Online Library, pp. 224–233.
890 <https://doi.org/10.1111/j.1477-8947.2004.00092.x>

891 Gusmawati, N., Selmaoui-Folcher, N., Proisy, C., Mustafa, A., Gendre, R. Le, Laugier, T.,
892 Lemonnier, H., Caledonia, N., Caledonia, N., 2017. Surveying shrimp aquaculture pond
893 activity using multitemporal VHR satellite images - case study from the Perancak
894 Estuary, Bali, Indonesia. *Mar. Pollut. Bull.* 4, 1–36.
895 <https://doi.org/10.1016/j.marpolbul.2017.03.059>

896 Hamilton, S.E., Casey, D., 2016. Creation of a high spatio-temporal resolution global

897 database of continuous mangrove forest cover for the 21st century (CGMFC-21). Glob.
 898 Ecol. Biogeogr. 25, 729–738. <https://doi.org/10.1111/geb.12449>

899 Hashim, R., Kamali, B., Tamin, N.M., Zakaria, R., 2010. An integrated approach to coastal
 900 rehabilitation: Mangrove restoration in Sungai Haji Dorani, Malaysia. Estuar. Coast.
 901 Shelf Sci. 86, 118–124. <https://doi.org/10.1016/j.ecss.2009.10.021>

902 Hobbs, R.J., 2016. Degraded or just different? Perceptions and value judgements in
 903 restoration decisions. Restor. Ecol. 24, 153–158. <https://doi.org/10.1111/rec.12336>

904 Holguin, G., Vazquez, P., Bashan, Y., 2001. The role of sediment microorganisms in the
 905 productivity, conservation, and rehabilitation of mangrove ecosystems: An overview.
 906 Biol. Fertil. Soils 33, 265–278. <https://doi.org/10.1007/s003740000319>

907 IUCN (International Union for Conservation of Nature), 2017. Mass mangrove restoration:
 908 driven by good intentions but offering limited results. [WWW Document]. URL
 909 [https://www.iucn.org/news/forests/201702/mass-mangrove-restoration-driven-good-](https://www.iucn.org/news/forests/201702/mass-mangrove-restoration-driven-good-intentions-offering-limited-results)
 910 [intentions-offering-limited-results](https://www.iucn.org/news/forests/201702/mass-mangrove-restoration-driven-good-intentions-offering-limited-results) (accessed 2.3.18).

911 Kairo, J.G., Dahdouh-Guebas, F., Bosire, J., Koedam, N., 2001. Restoration and management
 912 of mangrove systems - a lesson for and from the East African region. South African J.
 913 Bot. 67, 383–389. [https://doi.org/https://doi.org/10.1016/S0254-6299\(15\)31153-4](https://doi.org/https://doi.org/10.1016/S0254-6299(15)31153-4)

914 Kitamura, S., Anwar, C., Chaniago, A., Baba, S., 1998. Handbook of Mangroves in Indonesia.
 915 ISME, Okinawa, Japan.

916 Kodikara, K.A.S., Mukherjee, N., Jayatissa, L.P., Dahdouh-Guebas, F., Koedam, N., 2017. Have
 917 mangrove restoration projects worked? An in-depth study in Sri Lanka. Restor. Ecol. 25,

918 1–12. <https://doi.org/10.1111/rec.12492>

919 Kristensen, E., Alongi, D.M., 2006. Control by fiddler crabs (*Uca vocans*) and plant roots
 920 (*Avicennia marina*) on carbon, iron, and sulfur biogeochemistry in mangrove sediment.
 921 *Limnol. Oceanogr.* 51, 1557–1571. <https://doi.org/10.4319/lo.2006.51.4.1557>

922 L’Heureux, M.L., Takahashi, K., Watkins, A.B., Barnston, A.G., Becker, E.J., Di Liberto, T.E.,
 923 Gamble, F., Gottschalck, J., Halpert, M.S., Huang, B., Mosquera-Vásquez, K.,
 924 Wittenberg, A.T., 2017. Observing and predicting the 2015/16 El Niño. *Bull. Am.*
 925 *Meteorol. Soc.* 98, 1363–1382. <https://doi.org/10.1175/BAMS-D-16-0009.1>

926 Lavieren, H. Van, Spalding, M., Alongi, D.M., Kainuma, M., Clusener-Godt, M., Adeel, Z.,
 927 2012. Securing the future of mangrove. A Policy Brief. UNU-INWEH, UNESCO-MAB with
 928 ITTO, FAO, UNEP-WCMC TNC. 53. <https://doi.org/http://www.inweh.unu.edu>

929 Lewis, R.R., 2014. Habitat-specific approaches, in: Bozzano, M., Jalonen, R., Thomas, E.,
 930 Boshier, D., Gallo, L., Cavers, S., Bordács, S., Smith, P., Loo, J. (Eds.), *Genetic*
 931 *Considerations in Ecosystem Restoration Using Native Tree Species*. FAO, Rome, pp.
 932 195–200. <https://doi.org/https://doi.org/10.1016/j.foreco.2014.07.015>

933 Lewis, R.R., 2005. Ecological engineering for successful management and restoration of
 934 mangrove forests. *Ecol. Eng.* 24, 403–418.
 935 <https://doi.org/10.1016/j.ecoleng.2004.10.003>

936 Lewis, R.R., 2000. Ecologically based goal setting in mangrove forest and tidal marsh
 937 restoration. *Ecol. Eng.* 15, 191–198. [https://doi.org/10.1016/S0925-8574\(00\)00070-7](https://doi.org/10.1016/S0925-8574(00)00070-7)

938 Lewis, R.R., Brown, B.M., 2014. *Ecological Mangrove Rehabilitation A field manual for*

- 939 practitioners, 3rd ed. Oxfam, Canadian International Development Agency, USAID.
- 940 Lewis, R.R., Gilmore, R.G., 2007. Important considerations to achieve successful mangrove
941 forest restoration with optimum fish habitat. *Bull. Mar. Sci.* 80, 823–837.
- 942 Lewis, R.R., Milbrandt, E.C., Brown, B., Krauss, K.W., Rovai, A.S., Beever, J.W., Flynn, L.L.,
943 2016. Stress in mangrove forests: Early detection and preemptive rehabilitation are
944 essential for future successful worldwide mangrove forest management. *Mar. Pollut.*
945 *Bull.* 109, 764–771. <https://doi.org/10.1016/j.marpolbul.2016.03.006>
- 946 Long, J., Napton, D., Giri, C., Graesser, J., 2014. A Mapping and Monitoring Assessment of
947 the Philippines' Mangrove Forests from 1990 to 2010. *J. Coast. Res.* 294, 260–271.
948 <https://doi.org/10.2112/JCOASTRES-D-13-00057.1>
- 949 Macintosh, D.J., Ashton, E.C., Havanon, S., 2002. Mangrove rehabilitation and intertidal
950 biodiversity: A study in the Ranong mangrove ecosystem, Thailand. *Estuar. Coast. Shelf*
951 *Sci.* 55, 331–345. <https://doi.org/10.1006/ecss.2001.0896>
- 952 Mansourian, S., Stanturf, J.A., Derkyi, M.A.A., Engel, V.L., 2017. Forest Landscape
953 Restoration: increasing the positive impacts of forest restoration or simply the area
954 under tree cover? *Restor. Ecol.* 25, 178–183. <https://doi.org/10.1111/rec.12489>
- 955 McDonald, T., Gann, G., Jonson, J., Dixon, K.W., 2016. International standards for the
956 practice of ecological restoration—including principles and key concepts. Society for
957 Ecological Restoration, Washington, D. C.
958 <https://doi.org/10.1177/15257401070280021001>
- 959 McElhinny, C., Gibbons, P., Brack, C., Bauhus, J., 2005. Forest and woodland stand structural

960 complexity: Its definition and measurement. *For. Ecol. Manage.* 218, 1–24.

961 <https://doi.org/10.1016/j.foreco.2005.08.034>

962 Memon, J.A., Chandio, A.A., 2011. A critical appreciation of Restoration and conservation of

963 Degraded Mangroves in Thailand. *Int. J. Environ. Rural Dev.* 2, 108–113.

964 Moberg, F., Rönnbäck, P., 2003. Ecosystem services of the tropical seascape: Interactions,

965 substitutions and restoration. *Ocean Coast. Manag.* 46, 27–46.

966 [https://doi.org/10.1016/S0964-5691\(02\)00119-9](https://doi.org/10.1016/S0964-5691(02)00119-9)

967 National Economic and Social Development Board, 2011. Eleventh National Economic and

968 Social Development Plan (2012-2016). Government of Thailand, Bangkok, Thailand.

969 National News Bureau of Thailand, 2016. The First Division Army Area Commander launched

970 the conservation and restoration of 840,000 mangrove seedling in Honor of Her

971 Majesty Queen Sirikit in Chonburi area [WWW Document]. *Natl. News Bur. Thai.* URL

972 [http://nwnt.prd.go.th/centerweb/News/NewsDetail?NT01_NewsID=WNEVN59040100](http://nwnt.prd.go.th/centerweb/News/NewsDetail?NT01_NewsID=WNEVN5904010010013)

973 10013 (accessed 2.21.17).

974 Office of the National Economic and Social Development Board, 2001. The Ninth National

975 Economic and Social Development Plan (2002-2006).

976 Oh, R.R.Y., Friess, D.A., Brown, B.M., 2017. The role of surface elevation in the rehabilitation

977 of abandoned aquaculture ponds to mangrove forests, Sulawesi, Indonesia. *Ecol. Eng.*

978 100, 325–334. <https://doi.org/10.1016/j.ecoleng.2016.12.021>

979 Osbeck, M., Powell, N., Boromthananarat, S., 2010. Linking policy processes and stakeholder

980 agency to coastal change: A case study from Nakhon Si Thammarat, Thailand, in:

981 Fujikura, R., Kawanishi, M. (Eds.), *Climate Change Adaptation and International*
 982 *Development*. Routledge, London, UK, pp. 173–191.
 983 <https://doi.org/10.4324/9781849775267>

984 Prabnarong, P., Kaewrat, J., 2006. The Uthokawiphatprasit Watergate: A Man-Made Change
 985 in Pak Phanang River Basin. *Walailak J. Sci. Technol.* 3, 131–143.

986 Prach, K., del Moral, R., 2015. Passive restoration is often quite effective: Response to
 987 Zahawi et al. (2014). *Restor. Ecol.* 23, 344–346. <https://doi.org/10.1111/rec.12224>

988 Primavera, J.H., 2015. Stop seafront planting of bakhaw propagules [WWW Document].
 989 INQUIRER.net. URL [http://business.inquirer.net/187523/stop-seafront-planting-of-](http://business.inquirer.net/187523/stop-seafront-planting-of-bakhaw-propagules)
 990 [bakhaw-propagules](http://business.inquirer.net/187523/stop-seafront-planting-of-bakhaw-propagules) (accessed 9.25.17).

991 Primavera, J.H., Esteban, J.M.A., 2008. A review of mangrove rehabilitation in the
 992 Philippines: successes, failures and future prospects. *Wetl. Ecol. Manag.* 16, 345–358.
 993 <https://doi.org/10.1007/s11273-008-9101-y>

994 Primavera, J.H., Rollon, R.N., Samson, M.S., 2011. Mangrove rehabilitation in the Philippines:
 995 The challenges of coastal protection and pond reversion, in: Wolanski, E., McKlusky, D.
 996 (Eds.), *Treatise on Estuarine and Coastal Science*. Waltham Academic Press, pp. 217–
 997 244. <https://doi.org/10.1016/B978-0-12-374711-2.01010-X>

998 Primavera, J.H., Savaris, J.P., Bajoyo, B.E., Coching, J.D., Curnick, D.J., Golbeque, R.L.,
 999 Guzman, A.T., Henderin, J.Q., Joven, R. V, Loma, R.A., 2012. *Manual for community-*
 1000 *based mangrove rehabilitation*, 1st ed, London, Zoological Society of London,
 1001 *Mangrove Manual Series*. ZSL, London, UK.

- 1002 Primavera, J.H., Savaris, J.P., Loma, R.J.A., Coching, J.D., Montilijao, C.L., 2015. Manual for
1003 trainers: mangrove and beach forest rehabilitation and conservation., 3rd ed. ZSL,
1004 London, UK.
- 1005 Ranada, P., 2015. “Wrong” mangrove rehab in Yolanda areas wasting funds [WWW
1006 Document]. Rappler News Website Philipp. URL [http://www.rappler.com/science-](http://www.rappler.com/science-nature/environment/89163-unscientific-mangrove-rehabilitation-yolanda)
1007 [nature/environment/89163-unscientific-mangrove-rehabilitation-yolanda](http://www.rappler.com/science-nature/environment/89163-unscientific-mangrove-rehabilitation-yolanda) (accessed
1008 10.25.17).
- 1009 Richards, D.R., Friess, D.A., 2016. Rates and drivers of mangrove deforestation in Southeast
1010 Asia, 2000–2012. *Proc. Natl. Acad. Sci.* 113, 344–349.
1011 <https://doi.org/10.1073/pnas.1510272113>
- 1012 Saenger, P., 2002. Mangrove ecology, silviculture, and conservation. Kluwer Academic
1013 Publishers, Dordrecht; Boston.
- 1014 Saenger, P., Gartside, D., Funge-Smith, S., 2013. A review of mangrove and seagrass
1015 ecosystems and their linkage to fisheries and fisheries management, Food and
1016 Agriculture Organization of the United Nations. FAO Regional Office for Asia and the
1017 Pacific, Bangkok, Thailand. [https://doi.org/ISBN: 9789251077733](https://doi.org/ISBN:9789251077733)
- 1018 Saenger, P., Siddiqi, N.A., 1993. Land from the sea: The mangrove afforestation program of
1019 Bangladesh. *Ocean Coast. Manag.* 20, 23–39.
1020 [https://doi.org/http://dx.doi.org/10.1016/0964-5691\(93\)90011-M](https://doi.org/http://dx.doi.org/10.1016/0964-5691(93)90011-M)
- 1021 Salmo III, S.G., Lovelock, C., Duke, N.C., 2013. Vegetation and soil characteristics as
1022 indicators of restoration trajectories in restored mangroves. *Hydrobiologia* 720, 1–18.
1023 <https://doi.org/10.1007/s10750-013-1617-3>

- 1024 Salmo III, S.G., Lovelock, C.E., Duke, N.C., 2014. Assessment of vegetation and soil conditions
1025 in restored mangroves interrupted by severe tropical typhoon “Chan-hom” in the
1026 Philippines. *Hydrobiologia* 733, 85–102. <https://doi.org/10.1007/s10750-013-1766-4>
- 1027 Salmo III, S.G., Tibbetts, I.R., Duke, N.C., 2018. Nekton communities as indicators of habitat
1028 functionality in Philippine mangrove plantations. *Mar. Freshw. Res.* 69, 477–485.
1029 <https://doi.org/10.1071/MF17116>
- 1030 Salmo III, S.G., Torio, D., Esteban, J., 2007. Evaluation of rehabilitation strategies and
1031 management schemes for the improvement of mangrove management programs in
1032 Lingayen Gulf. *Sci. Diliman* 19, 24–34.
- 1033 Samarakoon, J.I., 2012. Mangrove restoration and planting in micro-tidal barrier-built
1034 estuaries and lagoons in Asia - ideology versus sustainable ecosystem science?, in:
1035 Macintosh, D.J., Mahindapala, R., Markopoulos, M. (Eds.), *Mangroves for the Future*
1036 *Regional Colloquium: Sharing Lessons on Mangrove Restoration*. MFF Secretariat,
1037 IUCN, Bangkok, Thailand, Chennai, India, pp. 141–157.
- 1038 Samson, M.S., Rollon, R.N., 2011. Mangrove Revegetation Potentials of Brackish-Water Pond
1039 Areas in the Philippines, *Aquaculture and the environment - A shared destiny*. INTECH
1040 Open Access Publisher.
- 1041 Samson, M.S., Rollon, R.N., 2008. Growth performance of planted mangroves in the
1042 Philippines: Revisiting forest management strategies. *Ambio* 37, 234–240.
1043 [https://doi.org/10.1579/0044-7447\(2008\)37\[234:GPOPMI\]2.0.CO;2](https://doi.org/10.1579/0044-7447(2008)37[234:GPOPMI]2.0.CO;2)
- 1044 Sanyal, P., 1998. Rehabilitation of degraded mangrove forests of the Sunderbans of India.,
1045 in: Anon (Ed.), *Program of the International Workshop on the Rehabilitation of*

- 1046 Degraded Coastal Systems. Phuket Marine Biological Centre, Phuket, Thailand, pp. 19–
1047 24.
- 1048 Senyk, J.P.J., 2005. Lessons from the Equator Initiative : Community-based management by
1049 Pred Nai Community Forestry Group in the mangroves of Southeastern Thailand.
1050 Winnipeg, Canada.
- 1051 Snedaker, S.C., 1982. Mangrove species zonation: why?, in: Sen, D.N., Rajpurohit, K.. (Eds.),
1052 Contributions to the Ecology of Halophytes. Tasks for Vegetation Science, Contributions
1053 to the Ecology of Halophytes. Springer, pp. 111–125.
1054 https://doi.org/https://doi.org/10.1007/978-94-009-8037-2_8
- 1055 Sousa, W.P., Kennedy, P.G., Mitchell, B.J., 2003. Propagule size and predispersal damage by
1056 insects affect establishment and early growth of mangrove seedlings. *Oecologia* 135,
1057 564–575. <https://doi.org/10.1007/s00442-003-1237-0>
- 1058 Springate-Baginski, O., Than, M.M., 2011. Community Forestry in Myanmar : Some field
1059 realities. School of International Development, University of East Anglia, Pyoe Pin
1060 Programme and Ecosystem Conservation and Community Development Initiative,
1061 Yangon, Myanmar.
- 1062 Stevenson, N.J., Lewis, R.R., Burbridge, P.R., 1999. Disused shrimp ponds and mangrove
1063 rehabilitation, in: Streever, W. (Ed.), An International Perspective on Wetland
1064 Rehabilitation, An International Perspective on Wetland Rehabilitation. Kluwer
1065 Academic Publishers, The Netherlands, pp. 277–297. [https://doi.org/10.1007/978-94-](https://doi.org/10.1007/978-94-011-4683-8_28)
1066 [011-4683-8_28](https://doi.org/10.1007/978-94-011-4683-8_28)
- 1067 Sunderlin, W.D., Angelsen, A., Belcher, B., Burgers, P., Nasi, R., Santoso, L., Wunder, S., 2005.

1068 Livelihoods, forests, and conservation in developing countries: An Overview. World
 1069 Dev. 33, 1383–1402. <https://doi.org/10.1016/j.worlddev.2004.10.004>
 1070 Tomlinson, P.B., 2016. The botany of mangroves, 2nd ed. Cambridge University Press.
 1071 <https://doi.org/https://doi.org/10.1017/CBO9781139946575>
 1072 Twilley, R.R., Rivera-Monroy, V.H., Chen, R., Botero, L., 1999. Adapting an Ecological
 1073 Mangrove Model to Simulate Trajectories in Restoration Ecology. Mar. Pollut. Bull. 37,
 1074 404–419. [https://doi.org/10.1016/S0025-326X\(99\)00137-X](https://doi.org/10.1016/S0025-326X(99)00137-X)
 1075 UNEP (United Nations Environment Program), 2007. After the Tsunami - Coastal Ecosystem
 1076 Restoration. Lessons Learnt [WWW Document]. URL
 1077 [https://wedocs.unep.org/bitstream/handle/20.500.11822/17467/after_tsunami.pdf?s](https://wedocs.unep.org/bitstream/handle/20.500.11822/17467/after_tsunami.pdf?sequence=1&isAllowed=y)
 1078 [equence=1&isAllowed=y](https://wedocs.unep.org/bitstream/handle/20.500.11822/17467/after_tsunami.pdf?sequence=1&isAllowed=y) (accessed 10.10.17).
 1079 van der Stocken, T., Ryck, D.D., Balke, T., Bouma, T.J., Nitto, D. Di, Kairo, J.G., Dahdouh-
 1080 Guebas, F., Koedam, N., 2012. Biological and environmental drivers in mangrove
 1081 propagule dispersal and recruitment: a field and modelling approach, in: Dahdouh-
 1082 Guebas, F., Satyanarayana, B. (Eds.), Proceedings of the International Conference:
 1083 Meeting on Mangrove Ecology, Functioning and Management. 2-6 July, 2012. Galle, Sri
 1084 Lanka. ULB/VUB, Brussels, Belgium, p. 181.
 1085 van Loon, A.F., Brake, B. Te, Huijgevoort, M.H.J. Van, Dijksma, R., 2016. Hydrological
 1086 classification, a practical tool for mangrove restoration. PLoS One 11.
 1087 <https://doi.org/https://doi.org/10.1371/journal.pone.0150302>
 1088 van Oudenhoven, A.P.E., Siahainenina, A.J., Sualia, I., Tonneijck, F.H., van der Ploeg, S., de
 1089 Groot, R.S., Alkemade, R., Leemans, R., 2015. Effects of different management regimes

1090 on mangrove ecosystem services in Java, Indonesia. *Ocean Coast. Manag.* 116, 353–
 1091 367. <https://doi.org/10.1016/j.ocecoaman.2015.08.003>

1092 Villamayor, B.M.R., Rollon, R.N., Samson, M.S., Albano, G.M.G., Primavera, J.H., 2016.
 1093 Impact of Haiyan on Philippine mangroves: Implications to the fate of the widespread
 1094 monospecific *Rhizophora* plantations against strong typhoons. *Ocean Coast. Manag.*
 1095 132, 1–14. <https://doi.org/10.1016/j.ocecoaman.2016.07.011>

1096 Walters, B., 2008. Mangrove forests and human security. *CAB Rev. Perspect. Agric. Vet. Sci.*
 1097 *Nutr. Nat. Resour.* 3, 1–9. <https://doi.org/10.1079/PAVSNNR20083064>

1098 Walters, B.B., 2005. Ecological effects of small-scale cutting of Philippine mangrove forests.
 1099 *For. Ecol. Manage.* 206, 331–348. <https://doi.org/10.1016/j.foreco.2004.11.015>

1100 Walters, B.B., 2004. Local Management of Mangrove Forests in the Philippines: Successful
 1101 Conservation or Efficient Resource Exploitation? *Hum. Ecol.* 32, 177–195.
 1102 <https://doi.org/10.1023/B:HUEC.0000019762.36361.48>

1103 Walters, B.B., 2000. Local Mangrove Planting in the Philippines: Are Fisherfolk and Fishpond
 1104 Owners Effective Restorationists? *Restor. Ecol.* 8, 237–246.
 1105 <https://doi.org/10.1046/j.1526-100x.2000.80035.x>

1106 Walters, B.B., 1997. Human ecological questions for tropical restoration: experiences from
 1107 planting native upland trees and mangroves in the Philippines. *For. Ecol. Manage.* 99,
 1108 275–290. [https://doi.org/10.1016/S0378-1127\(97\)00211-9](https://doi.org/10.1016/S0378-1127(97)00211-9)

1109 Walters, B.B., Rönnbäck, P., Kovacs, J.M., Crona, B.I., Hussain, S.A., Badola, R., Primavera,
 1110 J.H., Barbier, E.B., Dahdouh-Guebas, F., 2008. Ethnobiology, socio-economics and

1111 management of mangrove forests: A review. *Aquat. Bot.* 89, 220–236.
1112 <https://doi.org/10.1016/j.aquabot.2008.02.009>
1113 Walters, B.B., Sabogal, C., Snook, L.K., de Almeida, E., 2005. Constraints and opportunities
1114 for better silvicultural practice in tropical forestry: an interdisciplinary approach. *For.*
1115 *Ecol. Manage.* 209, 3–18. <https://doi.org/10.1016/j.foreco.2005.01.003>
1116
1117