Hotspots and hot moments of amino acid N in soil: Real-time insights using continuous microdialysis sampling

Hill, Elliot; Jones, Davey L.; Paterson, E.; Hill, Paul

Soil Biology and Biochemistry

DOI:

[10.1016/j.soilbio.2018.12.026](https://doi.org/10.1016/j.soilbio.2018.12.026)

Published: 01/04/2019

Peer reviewed version

[Cyswllt i'r cyhoeddiad / Link to publication](#)

Dyfyniad o'r fersiwn a gyhoeddwyd / Citation for published version (APA):

Hill, E., Jones, D. L., Paterson, E., & Hill, P. (2019). Hotspots and hot moments of amino acid N in soil: Real-time insights using continuous microdialysis sampling. *Soil Biology and Biochemistry*, 131, 40-43. <https://doi.org/10.1016/j.soilbio.2018.12.026>

Hawliau Cyffredinol / General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Hotspots and hot moments of amino acid N in soil: real-time insights using continuous microdialysis sampling

Elliot J. Hill^{a,*}, Davey L. Jones^{a,b}, Eric Paterson^c, Paul W. Hill^a

^a*School of Natural Sciences, Bangor University, Gwynedd, LL57 2UW, UK*

^b*UWA School of Agriculture and Environment, University of Western Australia, Crawley, WA 6009, Australia*

^c*The James Hutton Institute, Craigiebuckler, Aberdeen, AB15 8QH, UK*

Corresponding author: Elliot J. Hill*

Corresponding author address: School of Natural Sciences, Bangor University, Bangor, Gwynedd, LL57 2UW, UK

Corresponding author Tel: +447980398914

Corresponding author E-mail: elliott.hill@bangor.ac.uk

Declarations of Conflict of Interest: None

Article type: Short Communication

ABSTRACT Protein hotspots in soil, such as those associated with decaying soil fauna or plant litter, may produce ephemeral patches of disproportionately high soil nutrients. These hotspots may occur at the macro and microscale in close proximity to plant roots, however, the likely concentration of soluble products produced in these hotspots remains poorly understood. To address this, we buried two contrasting biomass residues in soil, namely earthworm (*Lumbricus terrestris*) and clover (*Trifolium repens*). Their transformation to amino acids, NH_4^+ and NO_3^- was monitored continually over 6 days using microdialysis. All treatments showed greater soluble nitrogen (N) concentrations compared to the unamended controls. The highest concentrations of both amino acids (12.9 mM after 12 h) and NH_4^+ (45.3 mM after 6 h) were generated in the vicinity of decomposing earthworm. In comparison, dried clover residues yielded 2.7 mM of amino acids at 6 h. After 12 h, amino acid and NH_4^+ concentrations in both earthworm and dried clover treatments showed a steep decline, returning close to background levels ($<20 \mu\text{M}$). Through the use of microdialysis we are able to show that soil nutrient hotspots may provide nearby roots with concentrations of amino acids and NH_4^+ several orders of magnitude higher than found in the bulk soil solution.

Keywords: Dissolved organic nitrogen; DON; Mineralization; Plant-microbial competition; Proteases

Amino acids (AAs) and oligopeptides are the most abundant and first quantitatively-significant protein breakdown products to be directly available as nitrogen (N) sources for plants and soil microbes (Sauheitl et al., 2009; Farrell et al., 2011; Warren, 2013; Moran-Zuloaga et al., 2015). Consequently, transformation of proteins to AAs is a major factor limiting N availability in soil and competition between plant roots and soil microbes for AAs is fierce (Jones and Kielland, 2002; Bardgett et al., 2003; Jan et al., 2009; Hill et al., 2011; Hill and Jones, 2018). The concentration of AAs in soil appears to be a key determinant in

the outcome of competition, with higher concentrations probably favouring plant AA-N capture (Jones et al., 2005). Therefore establishing true concentrations of AAs in soil is crucial to our understanding of plant N acquisition, and N cycling in soil. Typically, total free AA concentrations in contrasting ecosystems have been reported to remain fairly constant at $23 \pm 5 \mu\text{M}$, with the concentration of individual AAs typically ranging from 0.1-5.0 μM (Jones et al., 2009). These measurements in the bulk soil reflect the balance between slow rates of AA production and, rapid microbial AA consumption. However, biogeochemical ‘hotspots’, ephemeral patches which yield disproportionately high nutrient levels relative to the surrounding soil matrix, may supply high quantities of AAs to nearby roots (McClain et al., 2003; Schimel and Bennett, 2004; Jones et al., 2005; Kuzyakov and Blagodatskaya, 2015).

Using plant and soil fauna residues and microdialysis, we measured concentrations of protein breakdown products which may realistically occur close to roots. Microdialysis is a membrane-based sampling technique which offers non-invasive measurement of the soil solution phase, allowing probes to be positioned in close spatial proximity to samples and therefore yielding a high spatial resolution. We hypothesised that AAs generated from soil hotspots of protein breakdown transiently occur at concentrations greatly exceeding those found in measurements of bulk soil solution.

Microdialysis probes were single use, 100 kDa cut-off, 10 mm membrane length, 0.5 mm membrane diameter, 2.6 μL membrane internal volume and inlet internal volume of 1.4 μL (CMA Microdialysis, Torshamnsgatan, Sweden). Probes with a 100 kDa membrane pore size have been shown by other researchers to recover a great fraction of amino acids, compared to 20 kDa probes (Buckley et al., 2017). Probes were calibrated for relative recovery using amino acid (L-alanine), NH_4Cl and KNO_3 standard solutions of 0.1, 0.25, 0.5, 0.75, 1 and 25 mM (Inselsbacher et al., 2011; Lange, 2012). Probes ($n = 8$) were positioned

in standard solutions at room temperature, and perfused with ultra-pure deionised-water (5 $\mu\text{L min}^{-1}$). Aliquots of dialysate were collected at 30 min intervals during a 120 min perfusion period. Total amino acids in dialysate were measured fluorometrically according to Jones et al. (2002) while NH_4^+ and NO_3^- were determined colorimetrically according to Mulvaney (1996) and Miranda et al. (2001), respectively. Relative recovery was found to be $14.5 \pm 5.5\%$ of the supplied standard solution, irrespective of N-form, standard solution concentration or time. The relative recovery was later used to estimate actual concentrations of target analytes in the soil; although we cannot completely exclude some difference between standard solution calibration and measurements in soil solution. Collection vials were weighed before and after perfusion to ensure that there was no net loss or gain of water due to transmembrane flux or leakage.

Soil (0-15 cm) was collected from the Ah horizon of an agricultural grassland at Henfaes Agricultural Research Station, N. Wales, UK (53°23'N, 4°01'W, 19 m.a.s.l). The sandy clay loam textured soil is classified as a Eutric Cambisol (FAO), derived from post-glacial alluvial deposits. The main properties of the soil and added protein sources are shown in Table S1.

Soil was sieved to 2 mm, homogenised and added to 1.5 ml microcentrifuge tubes, each tube receiving *ca.* 1.5 g of field-moist soil. Three protein sources were added to the soil in a 1:10 w/w protein source to soil ratio as follows: a) fresh foliage of *Trifolium repens* L. (clover), b) dried foliage of *T. repens*, and c) fresh necromass of the earthworm, *Lumbricus terrestris* L., with soil controls receiving no protein addition. All treatments were replicated three times. These protein sources, added by weight, represent different levels of N input, however our hypothesis is predicated on a greater understanding of realistically occurring hotspots in the soil, rather than a direct comparison between treatments. Protein sources were added to the centre of each tube. Each replicate was sealed using gas-permeable film

(Parafilm M[®], Bemis Inc., USA) to restrict water loss by evaporation. Probes were positioned to a depth of 10 mm in the centre of the tube, using a syringe introducer and perfused with ultra-pure deionised water, at a continuous rate of 5 $\mu\text{L min}^{-1}$, using a multi-channel syringe pump (NE-1200 Multi-Phaser, New Era Pump Systems Inc., NY, USA) over a sampling period of 144 h. Samples were taken at hourly intervals for the first 6 h, then every 6 h thereafter (samples at 30 h were contaminated during storage and therefore omitted from analysis). Samples were analysed for amino acids and inorganic N as described above and for pH and electrical conductivity (EC) with standard microelectrodes. Outliers were identified using Grubbs' test (9 of 1008 measurements were removed). Data were analysed using a mixed two-way ANOVA with a Tukey HSD post-hoc test (SPSS v22; IBM, New York, USA), with a $p \leq 0.05$ cut-off for statistical significance.

Amino acid concentrations in controls did not exceed a mean concentration of 0.018 mM at any point during the experiment. Initially, levels of AA's were significantly higher in all treatments within the first 4 h following protein addition (Fig. 1). We attribute this to immediate loss of soluble components e.g. from damaged cells. All treatments had greater ($p \leq 0.05$) AA and NH_4^+ concentrations than controls during the majority of sampling intervals. Amino acids in dried clover and earthworm treatments showed a sharp rise in concentration after ~4 h, with those in dried clover rising to 2.7 mM after 6 h and to 12.9 mM in the earthworm treatment after 12 h, before sharply decreasing and levelling off after ~72 h. In contrast, fresh clover showed initially low AA concentrations (lowest mean concentration of 0.006 mM at 12 h), with only slightly higher concentrations than the control at some time points until ~54 h, thereafter reaching a peak of 0.099 mM at ~72 h, which was conversely the time at which the other treatments returned to levels comparable with controls. In comparison to the air-dried clover which was dead at the point of addition to the soil, it is likely that the fresh clover leaves underwent slow autolysis and death after excision. During

this autolysis, leaf protein and amino acids are catabolised internally for energy production (with plant cell walls probably remaining largely intact) and are less likely to be lost to the soil (Marella et al., 2017). The single highest concentration of AAs was observed in the earthworm treatment at 6 h, with one replicate of 29.3 mM of AAs, though this was identified as an outlier.

Concentrations of NH_4^+ in the control treatment did not exceed 0.21 mM during the entirety of the experiment. In contrast, in the earthworm treatment, concentrations peaked at 45.3 mM at 12 h within the same replicate as the greatest AA concentration. Generally, however, ammonium concentrations in other treatments were low in comparison to the earthworm, reaching peak mean concentrations of 2.67 mM and 1.71 mM for the dried and fresh clover respectively, with significantly ($p \leq 0.05$) greater concentrations than the control observed at every time point for dried clover, and from 24 h onwards for fresh clover (Fig. 2). Interestingly, mean nitrate concentrations of all treatments remained low throughout the sampling period (< 1 mM) with the highest concentration of 0.71 mM seen in the earthworm treatment at 6 h. We observed no clear relationship between pH and treatment and/or time (Fig. S1), however, EC was significantly higher in the earthworm and dried clover treatments than the control during the majority of sampling times (12-144 h) (Fig. S2).

The temporal pattern of our results suggests that the rise in levels of free AA's is matched by increased microbial consumption and growth that rapidly lowers the concentration. This suggests that a distinction should be made between concentration hotspots (where breakdown rate is high and exceeds microbial demand leading to accumulation in solution) and flux hotspots (where breakdown rate is high but is matched by microbial demand and where little change in solution concentration occurs). Likewise, microbial demand may have become saturated earlier on in the sampling period due in part to adding N at a higher concentrations (despite being the same weight) in earthworm and dried

clover treatments than that of fresh clover. However, hotspots occurring naturally within the soil would reflect this, yielding a range of concentrations from the macroscale, down to very fine microscales.

Results indicate that transformation of proteins to amino acids in ephemeral hotspots of protein addition may provide orders of magnitude greater levels of amino acids than previously measured in bulk soil solution (Jones et al., 2002; Jones et al., 2005; Hill et al., 2011). Further, although our experiment used relatively large fragments, it seems likely that hotspots occur at a much finer scale, for instance in the vicinity of dead microbial cells or damaged root cells in the detritusphere, as hypothesised by other researchers (Marschner et al., 2012; Kuzyakov and Blagodatskaya, 2015). As a consequence of this, it is therefore probable that the form and quantity of N acquired by plants varies considerably on a fine temporal and spatial scale.

Acknowledgments

This work was supported by the Biotechnology and Biological Sciences Research Council; and the Natural Environment Research Council [Grant number NE/M009106/1], by a Soils Training and Research Studentships (STARS) grant to Elliot J. Hill. STARS is a consortium consisting of Bangor University, British Geological Survey, Centre for Ecology and Hydrology, Cranfield University, James Hutton Institute, Lancaster University, Rothamsted Research and the University of Nottingham.

References

Bardgett, R., Streeter, T., Bol, R., 2003. Soil microbes compete effectively with plants for organic-nitrogen inputs to temperate grasslands. *Ecology* 84, 1277-1287.

- Buckley, S., Brackin, R., Näsholm, T., Schmidt, S., & Jämtgård, S., 2017. Improving in situ recovery of soil nitrogen using the microdialysis technique. *Soil Biology and Biochemistry*, 114, 93-103.
- de Lange, E.C.M., 2012. Recovery and calibration techniques: Toward quantitative microdialysis. In: *Microdialysis in Drug Development*, (M. Müller, ed.), AAPS Advances, Pharmaceutical Sciences Series, pp. 13-33. Springer Nature, Switzerland.
- Farrell, M., Hill, P., Wanniarachchi, S., Farrar, J., Bardgett, R., Jones, D., 2011. Rapid peptide metabolism: A major component of soil nitrogen cycling? *Global Biogeochemical Cycles* 25, GB3014.
- Hill, P.W., Jones, D.L., 2018. Plant-microbe competition: does injection of isotopes of C and N into the rhizosphere effectively characterise plant use of soil N? *New Phytologist*, In Press.
- Hill, P., Quilliam, R., DeLuca, T., Farrar, J., Farrell, M., Roberts, P., Newsham, K., Hopkins, D., Bardgett, R., Jones, D., 2011. Acquisition and assimilation of nitrogen as peptide-bound and D-enantiomers of amino acids by Wheat. *PLoS ONE* 6,e19220.
- Inselsbacher, E., Öhlund, J., Jämtgård, S., Huss-Danell, K. and Näsholm, T., 2011. The potential of microdialysis to monitor organic and inorganic nitrogen compounds in soil. *Soil Biology and Biochemistry* 43, 1321-1332.
- Jan, M., Roberts, P., Tonheim, S., Jones, D.L., 2009. Protein breakdown represents a major bottleneck in nitrogen cycling in grassland soils. *Soil Biology and Biochemistry* 41, 2272-2282.
- Jones, D.L., Owen, A., Farrar, J., 2002. Simple method to enable the high resolution determination of total free amino acids in soil solutions and soil extracts. *Soil Biology and Biochemistry* 34, 1893-1902.

191 Jones, D.L., Kielland, K., 2002. Soil amino acid turnover dominates the nitrogen flux in
 192 permafrost-dominated taiga forest soils. *Soil Biology and Biochemistry* 34, 209 -
 193 219.

194 Jones, D.L., Kielland, K., Sinclair, F.L., Dahlgren, R.A., Newsham, K.K., Farrar, J.F.,
 195 Murphy, D.V., 2009. Soil organic nitrogen mineralization across a global latitudinal
 196 gradient. *Global Biogeochemical Cycles* 23, G81016.

197 Jones, D.L., Owen, A., Farrar, J., 2002. Simple method to enable the high resolution
 198 determination of total free amino acids in soil solutions and soil extracts. *Soil Biology*
 199 *and Biochemistry* 34, 1893-1902.

200 Jones, D.L., Shannon, D., Junvee-Fortune, T., Farrar, J., 2005. Plant capture of free amino
 201 acids is maximized under high soil amino acid concentrations. *Soil Biology and*
 202 *Biochemistry* 37, 179-181.

203 Kuzyakov, Y., Blagodatskaya, E., 2015. Microbial hotspots and hot moments in soil: Concept
 204 and Review. *Soil Biology and Biochemistry* 83, 184-199.

205 Marella, V.S., Roberts, P., Hill, P., Jones, D.L., 2017. Different ways in which CO₂ can be
 206 released during the turnover of roots in soil. *Biology and Fertility of Soils* 53, 369-
 207 374.

208 Marschner, P., Marhan, S., Kandeler, E., 2012. Microscale distribution and function of soil
 209 microorganisms in the interface between rhizosphere and detritusphere. *Soil Biology*
 210 *and Biochemistry* 49, 174-183.

211 McClain, M., Boyer, E., Dent, C., Gergel, S., Grimm, N., Groffman, P., Hart, S., Harvey, J.,
 212 Johnston, C., Mayorga, E., McDowell, W., Pinay, G., 2003. Biogeochemical hot spots
 213 and hot moments at the interface of terrestrial and aquatic ecosystems. *Ecosystems*
 214 6, 301-312.

215 Miranda, K.M., Espey, M.G., Wink, D.A., 2001. A rapid simple spectrophotometric method
 216 for simultaneous detection of nitrate and nitrite. *Biology and Chemistry* 5, 62-71.

217 Moran-Zuloaga, D., Dippold, M., Glaser, B., Kuzyakov, Y., 2015. Organic nitrogen uptake
 218 by plants: reevaluation by position-specific labeling of amino acids. *Biogeochemistry*
 219 125, 359-374.

220 Mulvaney, R., 1996. Nitrogen—inorganic forms. In: Sparks, D.L. (Ed.). *Methods of Soil*
 221 *Analysis. Part 3. Chemical Methods*. Soil Science Society of America, Madison, WI,
 222 USA, pp. 1123–1184.

223 Rousk, J., Jones, D., 2010. Loss of low molecular weight dissolved organic carbon (DOC)
 224 and nitrogen (DON) in H₂O and 0.5 M K₂SO₄ soil extracts. *Soil Biology and*
 225 *Biochemistry* 42, 2331-2335.

226 Sauheitl, L., Glaser, B., Weigelt, A., 2009. Uptake of intact amino acids by plants depends on
 227 soil amino acid concentrations. *Environmental and Experimental Botany* 66, 145-
 228 152.

229 Schimel, J., Bennett, J., 2004. Nitrogen mineralization: Challenges of a changing paradigm.
 230 *Ecology* 85, 591-602.

231 Warren, C., 2013. Organic N molecules in the soil solution: what is known, what is unknown
 232 and the path forwards. *Plant and Soil* 375, 1-19.

233

234

Figure legends

Figure 1. Temporal dynamics of soil solution amino acid concentrations in response to the addition of protein-rich hotspots (dead earthworm (A), dried (B) or fresh clover leaves (C)) to soil and measurement by microdialysis. Values represent means + SEM ($n=4$).

Figure 2. Temporal dynamics of soil solution ammonium (A) and nitrate concentrations (B) in response to the addition of protein-rich hotspots (dead earthworm, dried or fresh clover leaves) to soil and measurement by microdialysis. Values represent means + SEM ($n=4$).

247

248 **Table 1.** Physiochemical properties of the A-horizon of the Eutric Cambisol (Means + SEM).

Soil parameters	Values
Soil moisture (%)	31.47 ± 0.22
pH	5.33 ± 0.04
EC ($\mu\text{S cm}^{-1}$)	45.62 ± 3.25
Total carbon (%)	2.55 ± 0.18
Total nitrogen (%)	0.28 ± 0.02
Nitrate (N mg kg ⁻¹ oven dry soil)	0.872 ± 0.10
Ammonium (N mg kg ⁻¹ oven dry soil)	3.94 ± 1.02

249

250

251 **Table 2.** Total C & N ratios of protein additions (Means + SEM).

	Carbon (%)	Nitrogen (%)
Clover	48.6 ± 0.2	4.81 ± 0.06
Earthworm	43.73 ± 3.78	9.72 ± 1.03

252