The effect of ocean acidification on the intertidal hermit crab *Pagurus criniticornis* is not modulated by cheliped amputation and sex

turra, alexander; Ragagnin, Marilia; McCarthy, Ian; Fernandez, Wellington

Marine Environmental Research

DOI:

[10.1016/j.marenvres.2019.104794](https://doi.org/10.1016/j.marenvres.2019.104794)

Published: 01/01/2020

Peer reviewed version

[Cyswllt i'r cyhoeddiad / Link to publication](#)

Dyfyniad o'r fersiwn a gyhoeddwyd / Citation for published version (APA):

turra, A., Ragagnin, M., McCarthy, I., & Fernandez, W. (2020). The effect of ocean acidification on the intertidal hermit crab *Pagurus criniticornis* is not modulated by cheliped amputation and sex. *Marine Environmental Research*, 153, [104794].
<https://doi.org/10.1016/j.marenvres.2019.104794>

Hawliau Cyffredinol / General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

The effect of ocean acidification on the intertidal hermit crab *Pagurus criniticornis* is not modulated by cheliped amputation and sex

Alexander Turra^{a*}; Marilia N. Ragagnin; Ian D. McCarthy^b; Wellington S. Fernandez^a;

^a Oceanographic Institute; University of São Paulo, São Paulo, 05508-120, SP, Brazil.

^b School of Ocean Sciences, Bangor University, Menai Bridge, Anglesey, LL59 5AB, United Kingdom.

* Corresponding author: Laboratório de Manejo, Ecologia e Conservação Marinha, Departamento de Oceanografia Biológica, Instituto Oceanográfico, Universidade de São Paulo (USP), Praça do Oceanográfico, 191, 05508-120, São Paulo, SP, Brazil.

E-mail address: turra@usp.br (A. Turra)

Highlights

- First study to assess long-term combined effects of OA, autotomy and sex on crustaceans.
- Crabs exposed to OA exhibited reduced survivorship, molting frequency and lipid content.
- Males showed increased molting frequency and higher regeneration rate than females.
- Interactive effects of pH, autotomy and sex were evidenced only for calcium content.
- There are no evident synergy of autotomy and sex on the effects of OA on hermit crabs.

ABSTRACT

Impacts of the interactive effects of ocean acidification (OA) with other anthropogenic environmental stressors on marine biodiversity are receiving increasing attention in recent years. However, little is known about how organismal responses to OA may be influenced by common phenomena such as autotomy and sexual dimorphism. This study evaluated the long-term (120 days) combined effects of OA (pH 7.7), experimental cheliped amputation and sex on physiological stress (mortality, growth, number of molts, cheliped regeneration and startle response) and energy budget (lipid and calcium contents) in the intertidal sexually-dimorphic hermit crab *Pagurus criniticornis*. Crabs exposed to OA reduced survivorship (46%), molting frequency (36%) and lipid content (42%). Autotomised crabs and males molted more frequently (39% and 32%,

respectively). Males presented higher regeneration (33%) and lower lipid content (24%). The few synergistic effects recorded did not indicate any clear pattern among treatments however, (1) a stronger reduction in lipid content was recorded in non-autotomised crabs exposed to low pH; (2) calcium content was higher in males than females only for autotomised crabs under control pH; and (3) autotomised females showed a proportionally slower activity recovery than autotomised males. Although our results suggest an effect of long-term exposure to low pH on the physiological stress and energy budget of *Pagurus criniticornis*, the physiological repertoire and plasticity associated with limb regeneration and the maintenance of dimorphism in secondary sexual characters may provide resilience to long-term exposure to OA.

Keywords: global change; environmental impact; water chemistry; seawater pH; physiological stress; energy budget; limb loss; sexual dimorphism.

1. Introduction

Carbon dioxide concentrations in the atmosphere are increasing, driven by anthropogenic activities, however, approximately one-third of the CO₂ released in the Industrial Age has been absorbed by the oceans modifying seawater chemistry in a process that has been termed ocean acidification (OA) (Sabine et al., 2004; Zeebe, 2012). To date, average seawater pH has declined by 0.1 units compared to pre-industrial values (*i.e.*, from *ca.* 8.2 to *ca.* 8.1) and future ‘business-as-usual’ IPCC scenario predictions estimate a further reduction of up to 0.4 units by 2100 (IPCC, 2014).

The impacts of OA on marine taxa and biodiversity have been widely discussed in recent reviews (Hofmann et al., 2010; Kelley and Lunden, 2017; Kroeker et al., 2010; Wittmann and Pörtner, 2013). OA can synergistically interact with other human-induced environmental changes (Mostofa et al., 2015), especially ocean warming (Byrne and Przeslawski, 2013; Przeslawski et al., 2015) and it is becoming clear that future OA research should focus on multi-stressor impacts as these may vary among taxa/species and are dependent on habitat, metabolic characteristics, activity patterns and life-cycle (Hofmann and Todgham, 2010; Pörtner et al., 2004; Widdicombe and Spicer, 2008).

Investigations on the effects of OA have generally focused on vulnerable calcifying groups (*e.g.*, corals, Gómez et al., 2015; molluscs, Parker et al., 2013;

echinoderms, Dupont et al., 2010), however, the impacts of OA on crustaceans, have been recognised (Whiteley, 2011) and research on this animal group is gaining momentum (*e.g.*, Borges et al., 2018; Coffey et al., 2017; Lim and Harley, 2018; Ragagnin et al., 2018; Whiteley et al., 2018). Although crustaceans are likely to be more tolerant to changes in seawater pH (Long et al., 2017; Taylor et al., 2015), mainly due to their acid-base regulation ability (Small et al., 2010; Wheatly and Henry, 1992), there is increasing evidence that important ecological aspects may be affected by OA, such as behavioral patterns (Dodd et al., 2015; Roggatz et al., 2016) and reproductive success (Borges et al., 2018). Furthermore, crustacean vulnerability to reduced seawater pH may increase when concurrently exposed to other anthropogenic stressors (Dissanayake and Ishimatsu, 2011; Ragagnin et al., 2018), although this is an issue that still is poorly understood.

Several invertebrate taxa are able to autotomise body parts as a defense response to increase chances of survival and decapod crustaceans represent the most widely-studied group exhibiting this phenomenon (Fleming et al., 2007). However, despite the prevalence of autotomy among crustacean species, the interactive multi-stressor effects of OA and autotomy in crustaceans have not been studied, although data are available for polychaetes (Pires et al., 2015), starfish (McCarthy et al., under review; Schram et al., 2011), sea urchins (Emerson et al., 2017) and brittle stars (Christensen et al., 2017; Hu et al., 2014; Wood et al., 2008, 2010).

Crustaceans may undergo limb loss (self-amputation or autotomy), usually chelipeds, during intra- and interspecific interactions and especially as a strategy to escape from predators (Mace and Curran, 2011; Maginnis, 2006). Although advantageous for survival, energy budgets can be affected with body resources being reallocated to regeneration, potentially at the expense of growth, mating success and ability to compete for resources (Juanes and Smith, 1995; Maginnis, 2006; Mariappan et al., 2000). Thus, the stress/cost of limb regeneration may increase the vulnerability of the organism to other stressors or threats (Mariappan et al., 2000) such as OA.

In addition, the synergistic effects of OA and autotomy may also be sex-dependent, a factor that has been largely neglected in the literature (Ellis et al., 2017). The few studies conducted to date on crustaceans are equivocal with either some (Kurihara et al., 2008), or no (Donohue et al., 2012) evidence of any sex-related effects of OA. The potential for an interactive effect of sex and OA may be especially relevant in sexually-dimorphic taxa where differences in gonad size, body size and secondary sexual characters are observed (Ellis et al., 2017; Shine, 1989). Crustaceans, especially

decapods, exhibit sexual dimorphism in relation to body size (males normally larger than females; Subramoniam, 2017) and in some species cheliped asymmetry is also observed (Salmon, 1987). Cheliped asymmetry is observed in some hermit crabs, *e.g.* in *Pagurus* spp. the right cheliped is larger (Nucci and Melo, 2011) and in *Calcinus* spp. the left cheliped is larger (Nucci and Melo, 2015). In the genus *Pagurus*, males may possess a relatively larger right cheliped than females (Matsuo et al., 2015). Besides the fundamental role in defense, feeding and mating behaviors (Yasuda et al., 2011, 2014), the major cheliped has a particular function for hermit crabs as a weapon during contests for gastropod shells (Arnott and Elwood, 2007; Elwood et al., 2006) and, in the case of males, for mates (Turra, 2005). Therefore, any impairment of claw integrity or regeneration due to reduced pH may potentially impact on fitness at the individual or population level.

Hermit crabs are considered appropriate biological models to both understand the consequences of limb autotomy and the importance of the regeneration process for individual fitness and population maintenance (Yasuda et al., 2014) and to evaluate behavioral effects of ocean acidification, since they demonstrate clear impaired responses under environmental changes and stimuli (Briffa et al., 2012). Studies on the effects of OA on hermit crabs have shown decreases in growth (Ragagnin et al., 2018), “antennular flicking” (de la Haye et al., 2011; 2012; Kim et al., 2016), shell exchange (de la Haye et al., 2011), locomotory activity (de la Haye et al., 2012), the ability to approach food (Newman and Dubuque, 2013) and to identify signals of shell availability (Ragagnin et al., 2018), as well as increased mortality (Ragagnin et al., 2018) and an increase in the time needed to select a new shell (de la Haye et al., 2011). Some of these effects are related to changes in the olfactory function and represent significant impacts on information gathering in hermit crabs (Kim et al., 2016). However, the studies evaluating chemosensory capacity and behavioral responses of hermit crabs have been under short-term exposure to reduced pH as a single stressor (de la Haye et al., 2011; 2012; Newman and Dubuque, 2013), and little is known about the potential effects of long-term exposure to OA. Therefore, considering the potential impacts on hermit crabs caused by changes in seawater chemistry, especially when it occurs simultaneously with other stressors (Ragagnin et al., 2018), we predict that the long-term effects of reduced pH may be amplified by the physiological repertoire that has evolved in hermit crabs regarding limb regeneration and the development of secondary sexual characters.

In this context, this study evaluated the potential long-term (120 day exposure) synergistic effects of ocean acidification expected by the end of the 21st century (pH 7.7; RCP8.5 *p*CO₂ levels predicted for 2100; IPCC, 2014) with experimental right cheliped amputation and sexual dimorphism on physiological stress (mortality, growth, number of molts, cheliped regeneration, and startle response) and energy budget (lipid and calcium contents) of the intertidal and shallow subtidal tropical hermit crab *Pagurus criniticornis* (Dana, 1852) (Decapoda, Anomura).

2. Materials and methods

2.1. Sampling and experimental design

Individuals of *Pagurus criniticornis* (Dana, 1852) were collected by hand in August 2016 at Araçá Bay, located on the northern coast of São Paulo State (23°48'47''S, 45°24'30''W). Immediately after collection, individuals were placed in thermal boxes containing seawater aerated using battery-powered air pumps and with empty gastropod shells to minimize agonistic interactions, and transported to the Oceanographic Institute at the University of São Paulo. In the laboratory, crabs were acclimatized for two weeks in seawater aquaria (salinity 31; Light:dark photoperiod of 12:12 hours; temperature ca. 23 °C, pH 8.1) with constant aeration and biological and mechanical filtration and fed *ad libitum* with pelleted food for crustaceans (JBL, NovoPrawn, Germany).

After the acclimation period, the hermit crabs were removed from their gastropod shells to identify sex and, after allowing each crab to return to its shell, 160 individuals (80 males and 80 females; shield length: 4.45 ± 0.51 mm and 4.35 ± 0.58 mm, respectively; although males are on average larger than females, crab size was controlled in the experiment.) were individually allocated to plastic containers with drilled walls to ensure good water circulation. Eight combinations of treatments (n=20 crabs each) were set up in an orthogonal design and maintained for 120 days (25th August to 22nd December, 2016). The experimental treatments consisted of two levels of pH [pH 8.25 (Control) and pH 7.7 (reduced pH)], cheliped amputation (autotomised and non-autotomised individuals), and sex (male and female).

For the treatments with autotomised crabs (i.e., experimentally amputated), 80 individuals (40 males and 40 females) were individually anesthetized with 7.5% magnesium chloride (MgCl₂) and their right cheliped (*i.e.* larger) was sectioned with a pair of scissors between the ischium and the merus.

Two independent recirculating water systems were used, one for each pH treatment, using artificial seawater (HW Marinemix Reefer, HW Wiegandt, Germany). Each system (*ca.* 440 L of total seawater volume) consisted of a seawater reservoir (310 L), 10 tanks used to house the hermit crabs (9.5 L each) and a tank for biological filtration (35 L). The control pH treatment was initially planned at pH 8.1 to represent an average actual pH in the ocean. However, the system stabilised at a value of 8.25 approximately 3 weeks after the experiment had started due to the biological filtering system. We decided to not correct the control pH by modifying the biological filtering system (the same in the low pH treatment) or introducing other substances in the aquaria to avoid additional influences and oscillations in the system. Since *Pagurus criniticornis* is an intertidal species commonly occupying tide pools (Turra and Denadai, 2003; Nucci and Melo, 2011), where pH tend to be more alkaline (Truchot and Duhamel-Jouve, 1980; Wolfe et al., 2013), this relatively high “control” pH represents a natural condition for this species and can be used to be contrasted to the experimental (low) pH. In this way, the experimental treatments are robust to allow unequivocal considerations about the effect of reduced pH on the tested species of hermit crab. In addition, we note that the mortality rates for *P. criniticornis* exposed to control pH values of 8.25 for 120 days in the current study (see results), were comparable to those recorded for control *P. criniticornis* reared at pH 8.1 for 98 days (Ragagnin et al., 2018).

The reduced pH treatment was based on the expected reduction of 0.3-0.5 pH units by the end of the 21st century (the ‘business-as-usual’ IPCC scenario; IPCC, 2014). Crabs were introduced into this system and the pH from the experimental system was slowly reduced over 48 hours to reach 7.7 units in order to avoid any effect on survivorship related to a sudden pH decrease. The reduced pH recirculating system was maintained at pH 7.7 by bubbling CO₂ in the seawater reservoir through a solenoid valve linked to a pH controller system (accuracy of ± 0.01 unit; Aqua Medic, Germany).

Eight separately-housed individuals were allocated per tank, two individuals of each sex and amputation treatments. All crabs were fed *ad libitum* with pelleted food for crustaceans (JBL, NovoPrawn, Germany) during the experimental period and three empty gastropod shells were provided for each separately-housed individual to avoid growth limitation. Salinity, pH and temperature were measured daily. Dissolved inorganic carbon (DIC) and total alkalinity (TA) samples were taken monthly, and analyzed in triplicate by infrared detection (LICOR-AIRICA, Marianda, Belgium) and potentiometric titration

(Titirino, Metrohm, Brazil), respectively (Dickson et al., 2003). Data were corrected using certified reference materials (Scripps Institution of Oceanography, USA).

2.2. Physiological stress

2.2.1. Mortality

Any dead individuals were recorded daily during the 120-day experimental period and immediately removed from the tanks, in order to minimize changes in water quality. Mortality patterns were evaluated from the cumulative mortality (*i.e.*, cumulative number of deaths) per 30-day experimental period (*i.e.*, after 30, 60, 90, and 120 days respectively) and compared among treatments after 120 days.

2.2.2. Growth and number of molts

Shield length (± 0.001 mm) was measured from the first molt and from the live crabs at the end of the experiment. These measurements were taken from digital images of the molts (or crabs) obtained using a stereomicroscope coupled with a camera and analyzed using the ImageJ 1.51d software (Abramoff et al., 2004) through the “Measure” function after scale setting (“Set Scale” function). Molts with damaged shields were excluded from the analysis and only individuals that survived until the end of the experiment were included in the statistical analysis. For these individuals, the total number of molts was recorded at the end of the experiment.

2.2.3. Cheliped regeneration

The cheliped regenerative capacity was evaluated by comparison to the length of the right cheliped amputated at the beginning of the experiment. These measurements (to the nearest mm, using digital images collected as described above) were taken at the end of the 120-day experiment for all surviving experimentally-autotomised males and females.

2.2.4. Startle Response

To verify changes in behavioral responses due to the experimental conditions, the startle response (Briffa et al., 2008; White and Briffa, 2016) of all surviving hermit crabs was analysed at the end of the 120 days of experiment. Each individual was removed from the experimental system, placed in a nine-liter aquarium with the same physico-

chemical parameters as their treatment water and allowed to acclimate for 10 minutes. Then, the hermit crab shells were turned, using forceps, so the shell aperture was uppermost, and the time taken (seconds) by each individual to turn the shell with the locomotory appendages back to the substrate was recorded (Briffa et al., 2008; White and Briffa, 2016) using a stopwatch (accuracy ± 0.01 s).

2.3. Energy budget

2.3.1. Lipid content

The total lipid content was measured in all individuals that survived to the end of the experiment using the extraction method of Folch et al. (1957) and quantified following Frings et al (1972). For this analysis, the whole body of each crab was used, except the left cheliped, which was used for calcium content analysis (see below). Each individual was weighed (g) and homogenized in a 2:1 solution of chloroform and methanol (according to Folch et al. 1957) using an ultrasonic processor. Then, 0.5 ml distilled water was added, the solution was centrifuged for 5 minutes at 1,000 rpm and dried in liquid nitrogen for 45 minutes. Following this procedure, the total lipid content was determined according to Frings et al. (1972). Briefly, 100 μ l of the sample was pipetted into 50 ml falcon tubes in triplicate, 1 ml of chloroform was added, the solution mixed, samples were evaporated for 15 min in an oven at 60 °C. Then, 200 μ l of concentrated sulfuric acid was added and the solution was heated for 10 minutes in boiling water on a heating plate at 100 °C. The solution was then cooled in an ice bath for 5 minutes, 5 ml of phosphovaniline was added, and the samples were reheated on heating plate for 15 minutes at 37 °C. Finally, absorbance at 540 nm was measured by spectrophotometry and related to a calibration curve prepared with cod liver oil (Sigma – Cod liver oil fatty acid methyl esters – C5650 10g) as a standard.

2.3.2. Calcium content

The calcium content in the left cheliped was measured in all surviving individuals. The samples were dried in an oven at 60 °C for 48 hours, weighed (g), digested in solution of nitric acid (HNO₃) and hydrogen peroxide (H₂O₂), maintained in boiling water in a heating plate at 100 °C for one hour, and 10 ml of deionized water was added. Calcium concentration was then analyzed by flame atomic absorption spectrometry [AAS vario 6 model with automatic sampler AS52 (AnalytikJenaAG, Jena, Germany) and hollow cathode lamp (Narva, Germany)].

2.4 Statistical analyses

All experimental data were first tested for normality (Shapiro-Wilk test) and homogeneity of variance (Levene's test) and met the assumptions for parametric statistical analysis (Zar, 2010) except for the startle response data that was transformed as $\text{Log}_{10}(\text{SR}+1)$ prior to analysis. Measures of physiological stress (mortality, growth, number of molts and startle response) and energy budget (lipid and calcium content) after 120 days were each analysed using a three-way ANOVA (pH, sex and autotomy plus two-way and three-way interactions) followed by Tukey HSD *post-hoc* tests (Zar, 2010). Crab growth was evaluated by the difference between the initial and final shield length (mm). Differences in cheliped regeneration between treatments after 120 days were analyzed using a two-way ANOVA (pH and sex), followed by Tukey HSD *post-hoc* tests (Zar, 2010).

3. Results

During the 120-day experimental period, the physico-chemical parameters of the seawater (pH, salinity, temperature, dissolved inorganic carbon, and total alkalinity) exhibited little variation over time (Table 1). The exception being the pH of the control treatment that increased from 8.1 to 8.25 during the first 3 weeks of the study but then stabilised for the remainder of the experiment.

Cumulative mortality increased over time in all treatments (Fig. 1A) with a total of 63 crabs (39%) dying by the end of the experiment. The accumulated mortality over the 120 days showed that autotomised males reared in reduced pH tended to present a higher number of deaths from the beginning of the experiment compared to all other treatments (15 crabs *cf.* 3 to 11 crabs in the other treatments; Fig. 1A). However, ANOVA analysis on total mortality after 120 days revealed that only pH as a single factor affected crab survival with no interactions among treatments (Table 2). Crabs maintained at low pH represented 73% ($n = 46$) of the total mortality, 2.7 times higher than the mortalities of crabs maintained at control pH ($n = 17$) (Table 2; Fig. 1B). Alternatively, survival rate was 46% lower in crabs maintained at low pH (34 out of 80; 43%) in comparison to control pH (63 out of 80; 79%).

Crab growth during the 120-day experiment, expressed as the difference between the final and initial shield length, was not influenced by pH, amputation or sex, with no interactions observed between any of these three treatments (Table 2; Fig. 2). A total of

21 crabs did not molt during the experimental period: 3 males and 7 females from the reduced pH condition and 3 males and 8 females from the control pH. Hermit crabs performed a maximum of five molts with the average number of molts during the 120-day experiment of 1.5 ± 1.1 . Amputated males reared in control pH performed the highest average number of molts (2.3 ± 1.8) with non-amputated females maintained in reduced pH performing the lowest average of number of molts (0.6 ± 0.7). Although crab growth was not affected by treatments, the average number of molts was influenced by pH, amputation and sex, with no interactive effects observed between treatments (Fig. 3, Table 2). In general, males showed a higher average number of molts (32% higher) than females (1.7 ± 0.6 and 1.2 ± 0.5 , respectively; Fig. 3A). In addition, the average number of molts of autotomised crabs was 39% higher than non-autotomised (1.8 ± 0.5 and 1.1 ± 0.6 , respectively; Fig. 3B), while it was 36% lower for individuals reared in reduced pH compared to those maintained in control pH (1.8 ± 0.5 and 1.1 ± 0.5 , respectively; Fig. 3C).

Cheliped regeneration was influenced by crab sex but not pH (Table 3), with cheliped length on average 33% greater in males than in females after 120 days (18.40 mm and 13.79 mm, respectively; Fig. 4). No significant interaction between sex and pH was recorded (Table 3).

The startle response on day 120 did not show any clear patterns among treatments (Table 2; Fig. 5), with a significant interaction between amputation and sex. Autotomised females showed a proportionally slower response time than autotomised males. Autotomised males showed the highest SR values, *i.e.* responded faster compared to autotomised females (average response times: 76.5 seconds and 21.5 seconds, respectively) (Fig. 5).

Lipid content after 120 days was affected by pH, sex and an interaction between pH and amputation (Table 4) with a greater reduction in lipid content recorded in non-autotomised crabs exposed to low pH (Fig. 6A). In general, OA reduced average lipid content by 42% in comparison to that recorded in crabs reared at the control pH (0.85 mg/g and 1.45 mg/g, respectively; Fig. 6A). In addition, average lipid content in males was 24% lower than the lipid content in female crabs (0.99 mg/g and 1.30 mg/g, respectively; Fig. 6A). The calcium content of the left cheliped was influenced by the interaction between amputation and sex (Fig. 6B; Table 4), with higher values in males than females only for autotomised crabs under control pH conditions. In general, calcium content was 6 % higher in males (176.75 mg/g) than females (166.13 mg/g).

4. Discussion

The present study has highlighted some potential effects of long-term exposure to the reduced surface seawater pH predicted for the end of this century (pH 7.7), combined with autotomy events (as represented by experimental cheliped amputation) and sexual dimorphism, on the performance of the intertidal, sexually-dimorphic hermit crab *Pagurus criniticornis*. No synergistic effects of pH, amputation and sex were demonstrated on mortality, growth, and cheliped regeneration, although single treatment effects of pH (mortality) and sex (cheliped regeneration) were observed. The few interactive effects recorded were: (1) between sex and amputation for startle response, with amputated females responding more slowly than amputated males; (2) between pH and amputation for lipid content, with a significant reduction in lipid content in non-amputated crabs exposed to reduced pH; and (3) among pH, sex and autotomy for calcium content - the only interactive effect between the three conditions - with higher calcium concentrations in males than females, but only for amputated crabs under control pH condition. Thus, no clear patterns of synergistic effects were recorded in this study.

Ocean acidification has been reported to significantly increase mortality in marine crustaceans (Dissanayake et al., 2010; Dissanayake and Ishimatsu, 2011; Findlay et al., 2010; Kurihara et al., 2008; Long et al., 2013; Ragagnin et al., 2018). Our results indicated a similar pattern of mortality, with long-term exposure to reduced pH resulting in higher mortality rates than control pH conditions. One aim of our study was to determine any interaction between sex and pH on mortality, however, no sex-related effects were observed. Since *Pagurus criniticornis* is a sexually dimorphic species, with larger cheliped asymmetry in males than in females, we expected that amputation of the right (larger) cheliped would amplify the potential cumulative role of sex on the effects of OA. However, this effect was not supported by the data gathered. Few studies have investigated how sex may modulate the effects of OA on crustaceans, with the available data indicating that sex-related effects may vary among species. For example, in the Pacific grass shrimp *Palaemon pacificus*, males exhibited higher survival rates (but not growth) than females when exposed to reduced seawater pH values of 7.89 or 7.64 compared to control seawater at pH 8.16 for 30 weeks (Kurihara et al., 2008). In contrast, no evidence of sex-related effects were seen in the burrowing shrimp *Upogebia deltaura* exposed to reduced seawater pH values of 7.64, 7.35 or 6.71 compared to control seawater at pH 7.99 for 35 days (Donohue et al., 2012). Since sensitivities to OA can vary among

373 taxa, species and populations (Kroeker et al., 2013; Przeslawski et al., 2015; Whiteley,
374 2011; Wittman and Pörtner, 2013), more research is needed to determine whether there
375 are any consistent effect of sex on mortality rates in crustaceans.

376 It was expected that amputation would increase the effect of reduced pH on
377 mortality, however, no differences were observed between autotomised and non-
378 autotomised individuals. Autotomy is a natural defense mechanism and despite the
379 immediate survival benefit, the regrowth of an appendage may be physiologically
380 expensive, especially in cases of the loss of the major cheliped (Maginnis et al., 2014).
381 Although autotomy itself may not increase mortality risk *per se*, crabs may become more
382 vulnerable as a result of the limb loss, since it may impair feeding ability (Flynn et al.,
383 2015) and/or increase susceptibility to predators and competitors (Darnell et al., 2018;
384 Maginnis et al., 2014). However, mortality associated with limb loss in crustaceans may
385 depend on its function in the behavior and ecology of a species. For example, stone crabs
386 *Menippe* spp. display increased mortality in the natural environment following forced
387 claw removal during fishing activities since amputated crabs have limited ability to crush
388 their bivalve prey (Duermit et al., 2015, 2017). In addition, autotomised hermit crabs
389 change shells less frequently (Matsuo et al., 2014), which may prevent crabs from finding
390 and using more adequate shells within which they can withdraw and protect themselves
391 from predators. Crabs in low adequacy shells (*i.e.*, shells relatively smaller than crabs)
392 are more susceptible to predation than crabs in adequate shells (Vance, 1972). Since
393 chelipeds are also relevant to hermit crabs for feeding (Turra and Denadai, 2003),
394 burrowing (Rebach, 1974) and mating (Yasuda et al., 2011, 2014; Turra, 2005), autotomy
395 may have chronic effects at both the individual and population level. These effects may
396 be intensified under OA, which was demonstrated here to reduce survivorship and lipid
397 content (*e.g.*, energy for reproduction). Nevertheless, further studies on the ecological
398 responses under these natural situations in the context of OA would be necessary to
399 understand species-specific vulnerability.

400 In the present study, neither sex, pH or autotomy affected the overall growth of *P.*
401 *criniticornis* but individually these factors all influenced the number of molts. Although
402 no interaction was observed, molting rate was reduced at low pH (*i.e.* molting frequency
403 was 36% lower than crabs exposed to control pH), as has already been demonstrated for
404 other crustaceans (Findlay et al., 2010; Kurihara et al., 2008; Long et al., 2013; Zheng et
405 al., 2015). Previously, we have shown that juvenile *P. criniticornis* exhibited reduced
406 growth at pH 7.6 (Ragagnin et al., 2018), indicating that different responses may be

observed among life stages (Byrne, 2011; Byrne and Przeslawski, 2013). Reduced growth will be associated with energetic trade-offs, where the costs to physiological maintenance may impair processes related to growth and reproduction (Kurihara et al., 2013; Pörtner et al., 2004; Wood et al., 2008). Thus, it is likely that hermit crabs exposed to reduced pH presented fewer molt cycles due to the higher energetic costs incurred due to OA (Whiteley, 2011).

In addition, the higher number of molts in autotomised compared to non-autotomised individuals, and in males compared to females (39% and 32%, respectively) may be associated with the important role the major cheliped plays, especially in males crabs, in mating behavior and male-male contests (Yasuda et al., 2014). Previous work has shown that autotomy accelerates molting cycles, possibly as an adaptive response to recover the major claw considering its functional role (Darnell et al., 2018). Males may regenerate their major cheliped at the first molt in some species (Yasuda et al., 2014). Indeed, our results showed that males had a significantly higher regeneration rate of the right cheliped compared to females (*ca.* 33%), with no influence of reduced pH exposure. These results highlight the importance of energy allocation to a fast growth of the major claw in males of *Pagurus criniticornis* due to the potential ecological costs related to this condition (*e.g.*, disadvantage in male-male contests for females and shells and less protection against predators), as demonstrated for other *Pagurus* species (Yasuda et al., 2011, 2014).

Regarding behavioral responses, autotomised females demonstrated a clearly slower startle response (*i.e.*, activity recovery) than autotomised males, irrespective of pH treatment. Sex-dependent differences in startle response between males and females have been rarely explored in behavioral studies. To the best of our knowledge, the only studies considering the effect of sex on startle response are by Briffa et al. (2008) and White and Briffa (2016), who showed no influence of sex in *Pagurus bernhardus* in the field and in the laboratory in the absence/presence of predator cues (Briffa et al., 2008) or exposed to high concentrations of copper White and Briffa (2016). Sex also had no effect on shell abandoning response by the hermit crab *Pagurus criniticornis* when exposed to experimental entrapment of the shell (Gorman et al., 2015). In the present study, the faster response of males may be related to increased aggressiveness expected in autotomised males compared to autotomised females, considering the importance of aggression in the frequent male-male contests observed in hermit crabs (Suzuki et al., 2012; Yasuda and Koga, 2016). Unexpectedly, reduced pH did not affect behavioral activity of *P.*

441 *criniticornis*, contradicting some studies reporting changes in startle response in *Pagurus*
442 *spp.* exposed to other environmental stressors (temperature; Briffa et al. 2013; copper,
443 White and Briffa, 2016) as well as reduction in displacement behavior in *P. criniticornis*
444 exposed to combined effect of AO and shadow in response to gastropod odor (Ragagnin
445 et al., 2018).

446 In our study, pH-dependent differences in lipid concentrations were observed,
447 with lower mean lipid concentrations recorded in individuals maintained in the acidified
448 treatment, thus providing evidence of higher energy expenditure under OA (Carter et al.,
449 2013). There is also evidence that lipids reserves were significantly reduced under
450 increased temperature (25.2 ± 0.6 °C) and $p\text{CO}_2$ (763.0 ± 104.6 ppm) on the whelk
451 *Dicathais orbita* (Valles-Regino et al., 2015). Further, lipid reserves decreased in
452 crustaceans exposed to other stressful environmental conditions, for example cadmium
453 (freshwater crab *Sinopotamon henanense*; Yang et al., 2013) and crude oil (blue crab,
454 *Callinectes sapidus*; Wang and Stickle, 1988), showing that lipid content of marine
455 invertebrates can be affected by adverse environmental conditions. In addition, it is still
456 unclear how these effects may induce consequences on the community scale, but there is
457 evidence that impacts on fatty acids composition of lower trophic levels organisms may
458 impair growth and reproduction of consumers (Rossol et al., 2012).

459 In the present study, an interaction among the three treatments (pH, autotomy and
460 sex) was observed for calcium content. In general, males presented a higher calcification
461 than females, feature that was maintained in autotomised males compared to autotomised
462 females reared in control pH, but not under low pH. Although it is well known that several
463 species of crustaceans may demonstrate both maintenance or increased rates of
464 calcification when specimens are subjected to long-term exposure to ocean acidification
465 conditions (Long et al., 2013; McDonald et al., 2009; Ries et al., 2009; Ragagnin et al.,
466 2018; Small et al., 2010; Taylor et al., 2014), in the present study, exposure to low pH
467 eliminated the difference in cheliped calcification between autotomised males and
468 females. Based on the arguments presented above on the importance of chelipeds for
469 males, pH may have an additional effect on male fitness through cheliped weakening.

470 In conclusion, this is the first study to evaluate the potential interactive effects of
471 ocean acidification with intrinsic physiological characteristics and behaviors, *i.e.*
472 autotomy combined with startle responses, and to consider the influence of sexual
473 dimorphism on such effects in crustaceans. In addition, few studies have investigated the
474 synergistic effects of OA with other environmental stressors under a long-term exposure

perspective. In general, our results highlight the negative effects of long-term exposure to reduced pH, through reducing survival, molt frequency and lipid content. Although our results did not provide evidence for clear patterns of synergistic impacts of the tested factors, the few interactive effects recorded highlight the different responses of these organisms to a scenario of reduced pH exposure and cheliped loss. The results indicate that males may invest more energy into faster regeneration, resulting in lower concentrations of lipids and greater number of molts, processes not governed by low pH itself *per se*. Such a response may be a consequence of evolutionary processes and osmoregulatory adaptations of intertidal species, as *Pagurus criniticornis*, as already suggested by Whiteley et al. (2018) for sea urchins. Additional evidence for the synergistic effect of amputation and sexual dimorphism with low pH might be provided by studies on subtidal hermit crabs species that exhibit cheliped asymmetry. In addition, the already existing adaptations regarding limb regeneration and the maintenance of dimorphism on the secondary sexual characters overcame the effect of OA, potentially alleviating some additional effects due to alterations in behavioral interactions. However, other impacts of exposure to low pH (reduced survival, number of molts and lipid content) may reveal the potential chronic effects that can be up-scaled to higher levels of organization.

Acknowledgments

We thank Camila Andrade from the Oceanographic Institute, University of São Paulo, for support, maintenance and logistics during the experimental procedures.

Funding: This work was supported by the São Paulo Research Foundation (FAPESP) [grant numbers 2013/19826-6 and 2014/12879-0, WSF; 2013/50197-5, AT/IDM; 2015/02727-0, MNR] and the National Council for Scientific and Technological Development (CNPq) [grant number 309697/2015-8, AT].

References

- Abramoff, M.D., Magalhaes, P.J., Ram, S.J., 2004. Image Processing with ImageJ. *Biophotonics International* 11, 36-42.
- Arnott, G., Elwood, R.W., 2007. Fighting for shells: How private information about resource value changes hermit crab pre-fight displays and escalated fight behaviour. *Proceedings of the Royal Society B: Biological Sciences* 274, 3011–3017.

Borges, F.O., Figueiredo, C., Sampaio, E., Rosa, R., Grilo, T.F., 2018. Transgenerational deleterious effects of ocean acidification on the reproductive success of a keystone crustacean (*Gammarus locusta*). *Marine Environmental Research* 138, 55–64.

Briffa, M., Bridger, D., Biro, P. A., 2013. How does temperature affect behaviour? Multilevel analysis of plasticity, personality and predictability in hermit crabs. *Animal Behaviour* 86, 47-54.

Briffa, M., de la Haye, K., Munday, P.L., 2012. High CO₂ and marine animal behaviour: Potential mechanisms and ecological consequences. *Marine Pollution Bulletin* 64, 1519–1528.

Briffa, M., Rundle, S.D., Fryer, A., 2008. Comparing the strength of behavioural plasticity and consistency across situations: animal personalities in the hermit crab *Pagurus bernhardus*. *Proceedings of the Royal Society B: Biological Sciences* 275, 1305–1311.

Byrne, M., 2011. Impact of ocean warming and ocean acidification on marine invertebrate life history stages: Vulnerabilities and potential for persistence in a changing ocean. *Oceanography and Marine Biology: An Annual Review* 49: 1-42.

Byrne, M., Przeslawski, R., 2013. Multistressor impacts of warming and acidification of the ocean on marine invertebrates' life histories. *Integrative and Comparative Biology* 53, 582–596.

Carter, H.A., Caballos-Osuna, L., Miller, N.A., Stillman, J.H., 2013. Impact of ocean acidification on metabolism and energetics during early life stages of the intertidal porcelain crab *Petrolisthes cinctipes*. *The Journal of Experimental Biology* 216: 1412-1422.

Christensen, A.B., Radivojevic, K.O., Pyne, M.I., 2017. Effects of CO₂, pH and temperature on respiration and regeneration in the burrowing brittle stars *Hemipholis cordifera* and *Microphiopholis gracillima*. *Journal of Experimental Marine Biology and Ecology* 495, 13-23.

Coffey, W.D., Nardone, J.A., Yarram, A., Long, W.C., Swiney, K.M., Foy, R.J., Dickinson, G.H., 2017. Ocean acidification leads to altered micromechanical properties of the mineralized cuticle in juvenile red and blue king crabs. *Journal of Experimental Marine Biology and Ecology* 495, 1–12.

de la Haye, K. L., Spicer, J. I., Widdicombe, S., Briffa, M., 2011. Reduced sea water pH disrupts resource assessment and decision making in the hermit crab *Pagurus bernhardus*. *Animal Behaviour* 82, 495-501.

- de la Haye, K. L., Spicer, J. I., Widdicombe, S., Briffa, M., 2012. Reduced pH sea water disrupts chemo-responsive behavior in an intertidal crustacean. *Journal of Experimental Marine Biology and Ecology* 412, 134-140.
- Darnell, M.Z., Rittschof, C.C., Rittschof, J., Beach, C., Rittschof, D., 2018. Autotomy of the major claw stimulates molting and suppresses feeding in fiddler crabs. *Journal of Experimental Marine Biology and Ecology* 509, 66-70.
- Dickson, A.G., Afghan, J.D., Anderson, G.C., (2003). Reference materials for oceanic CO₂ analysis: a method for the certification of total alkalinity. *Marine Chemistry* 80, 185-197
- Dissanayake, A., Ishimatsu, A., 2011. Synergistic effects of elevated CO₂ and temperature on the metabolic scope and activity in a shallow-water coastal decapod (*Metapenaeus joyneri*; Crustacea: Penaeidae). *ICES Journal of Marine Science* 68, 1147–1154.
- Dissanayake, A., Clough, R., Spicer, J.I., Jones, M.B., 2010. Effects of hypercapnia on acid–base balance and osmo-iono-regulation in prawns (Decapoda: Palaemonidae). *Aquatic Biology* 11, 27–36.
- Dodd, L.F., Grabowski, J.H., Piehler, M.F., Westfield, I., Ries, J.B., 2015. Ocean acidification impairs crab foraging behavior. *Proceedings of the Royal Society B: Biological Sciences* 282, 1–9.
- Donohue, P.J.C., Calosi, P., Bates, A.H., Laverock, B., Rastrick, S., Mark, F.C., Strobel, A., Widdicombe, S., 2012. Impact of exposure to elevated *p*CO₂ on the physiology and behaviour of an important ecosystem engineer, the burrowing shrimp *Upogebia deltaura*. *Aquatic Biology* 15, 73-86.
- Duermit, E., Kingsley-Smith, P.R., Wilber, D.H. 2015. The consequences of claw removal on stone crabs *Menippe* spp. and the ecological and fishery implications. *North American Journal of Fisheries Management* 35, 895-905.
- Duermit, E., Shervitte, V., Whitaker, J.D., Kingsley-Smith, P.R., Wilber, D.H. 2017. A field assessment of claw removal impacts on the movement and survival of stone crabs *Menippe* spp. *Fisheries Research* 193, 43-50.
- Dupont, S., Ortega-Martinez, O., Thorndyke, M., 2010. Impact of near-future ocean acidification on echinoderms. *Ecotoxicology* 19, 449-462.
- Ellis, R.P., Davison, W., Queirós, A.M., Kroeker, K.J., Calosi, P., Dupont, S., Spicer, J.I., Wilson, R.W., Widdicombe, S., Urbina, M.A., 2017. Does sex really matter?

576 Explaining intraspecies variation in ocean acidification responses. *Biology Letters*
577 13, 20160761.

578 Elwood, R.W., Pothanikat, R.M.E., Briffa, M., 2006. Honest and dishonest displays,
579 motivational state and subsequent decisions in hermit crab shell fights. *Animal*
580 *Behaviour* 72, 853–859.

581 Emerson, C.E., Reinardy, H.C., Bates, N.R., Bodnar, A.G., 2017. Ocean acidification
582 impacts spine integrity but not regenerative capacity of spines and tube feet in adult
583 sea urchins. *Royal Society Open Science* 4, 170140.

584 Findlay, H.S., Kendall, M.A., Spicer, J.I., Widdicombe, S. 2010. Relative influences of
585 ocean acidification and temperature on intertidal barnacle post-larvae at the northern
586 edge of their geographic distribution. *Estuarine, Coastal and Shelf Science* 86, 675–
587 682.

588 Fleming, P.A., Muller, D., Bateman, P.W., 2007. Leave it all behind: A taxonomic
589 perspective of autotomy in invertebrates. *Biological Reviews* 82, 481–510.

590 Flynn, P.S.T., Mellish, C.L., Pickering, R.T., Quijón, P.A., 2015. Effects of claw
591 autotomy on green crab (*Carcinus maenas*) feeding rates. *Journal of Sea Research*
592 103, 113–119.

593 Folch, J., Lees, M., Sloane Stanley, G.H., 1957. A simple method for the isolation and
594 purification of total lipids from animal tissues. *The Journal of Biological Chemistry*
595 226, 497–509.

596 Frings, C.S., Queen, C.A., Dunn, R.T., Fendley, T.W., 1972. Improved determination of
597 total serum lipids by sulfo-phospho-vanillin reaction. *Clinical Chemistry* 18, 673–
598 674.

599 Gómez, C.E., Paul, V.J., Ritson-Williams, R., Muehllehner, N., Langdon, C., Sánchez,
600 J., 2015. Responses of the tropical gorgonian coral *Eunicea fusca* to ocean
601 acidification conditions. *Coral Reefs* 34, 451–460.

602 Gorman, D., Barros, F., Turra, A., 2015. What motivates hermit crabs to abandon trapped
603 shells? Assessing the influence of shell value, olfactory attractants, and previous
604 experience. *Hydrobiologia* 743: 285–297.

605 Hofmann, G.E., Barry, J.P., Edmunds, P.J., Gates, R.D., Hutchins, D.A., Klinger, T.,
606 Sewell, M.A., 2010. The effect of ocean acidification on calcifying organisms in
607 marine ecosystems: an organism-to-ecosystem perspective. *Annual Review of*
608 *Ecology, Evolution, and Systematics* 41, 127–147.

- Hofmann, G.E., Todgham, A.E., 2010. Living in the now: physiological mechanisms to tolerate a rapidly changing environment. *Annual Review of Physical Chemistry* 72, 127-145.
- Hu, M.Y., Casties, I., Stumpp, M., Ortega-Martinez, O., Dupont, S., 2014. Energy metabolism and regeneration are impaired by seawater acidification in the infaunal brittlestar *Amphiura filiformis*. *Journal of Experimental Biology* 217, 2411–21.
- IPCC, 2014. Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change [Core Writing Team, R.K. Pachauri and L.A. Meyer (eds.)]. IPCC, Geneva, Switzerland, 151 pp.
- Juanes, F., Smith, L.D., 1995. The ecological consequences of limb damage and loss in decapod crustaceans: a review and prospectus. *Journal of Experimental Marine Biology and Ecology* 193, 197–223.
- Kelley, A.L., Lunden, J.J., 2017. Meta-analysis identifies metabolic sensitivities to ocean acidification. *AIMS Environmental Science* 4, 709–729.
- Kim, T.W., Taylor, J., Lovera, C., Barry, J.P., 2016. CO₂-driven decrease in pH disrupts olfactory behaviour and increases individual variation in deep-sea hermit crabs. *ICES Journal of Marine Science* 73, 613-619.
- Kroeker, K.J., Kordas, R.L., Crim, R.N., Singh, G.G., 2010. Meta-analysis reveals negative yet variable effects of ocean acidification on marine organisms. *Ecology Letters* 13, 1419–1434.
- Kurihara, H., Matsui, M., Furukawa, H., Hayashi, M., Ishimatsu, A., 2008. Long-term effects of predicted future seawater CO₂ conditions on the survival and growth of the marine shrimp *Palaemon pacificus*. *Journal of Experimental Marine Biology and Ecology* 367, 41-46.
- Lim, E.G., Harley, C.D.G., 2018. Caprellid amphipods (*Caprella spp.*) are vulnerable to both physiological and habitat-mediated effects of ocean acidification. *PeerJ* 6, e5327.
- Long W.C., Swiney, K.M., Harris, C., Page, H.N., Foy, R.J., 2013. Effects of ocean acidification on juvenile red king crab (*Paralithodes camtschaticus*) and tanner crab (*Chionoecetes bairdi*) growth, condition, calcification, and survival. *PLoS ONE* 8, e60959.

- Long, W.C., Van Sant, S.B., Swiney, K.M., Foy, R.J., 2017. Survival, growth, and morphology of blue king crabs: Effect of ocean acidification decreases with exposure time. *ICES Journal of Marine Science* 74, 1033–1041.
- Mace, M.M., Curran, M.C., 2011. Differences in the use of cheliped autotomy by the mud fiddler crab *Uca pugnax* (Smith, 1870) (Decapoda, Ocypodidae) when escaping predation by the blue crab *Callinectes sapidus* (Rathbun, 1896). *Crustaceana* 84, 1281–1293.
- Maginnis, T.L., 2006. The costs of autotomy and regeneration in animals: a review and framework for future research. *Behavioral Ecology* 17, 857–872.
- Mariappan, P., Balasundaram, C., Schmitz, B., 2000. Decapod crustacean chelipeds: An overview. *Journal of Biosciences* 25, 301–313.
- Matsuo, K., Yasuda, C.I., Wada, S., 2014. Effect of cheliped loss due to autotomy on shell selection behavior in the hermit crab *Pagurus middendorffii*. *Crustacean Research* 43: 41–46.
- Matsuo, K., Tanikawa, D., Yasuda, C.I., Wada, S., 2015. Sex-related differences in size, function and regeneration of the major cheliped in the hermit crab *Pagurus filholi*. *Marine Ecology* 36, 1391–1399.
- McCarthy, I.D., Whiteley, N.M., Fernandez, W.S., Ragagnin, M.N., Cornwell, T.O., Suckling, C., Turra, A., Under review. Elevated pCO₂ does not impair performance in autotomised individuals of the intertidal predatory starfish *Asterias rubens* (Linnaeus, 1758). *Marine Environmental Research*.
- McDonald, M.R.; Mcclintock, J.B.; Amsler, C.D.; Rittschof, D.; Angus, R.A.; Orihuela, B.; Lutostanski, K., 2009. Effects of ocean acidification over the life history of the barnacle *Amphibalanus amphitrite*. *Marine Ecology Progress Series* 385, 179–187.
- Mostofa, K.M.G., Liu, C.Q., Zhai, W.D., Minella, M., Vione, D., Gao, K., Minakata, D., Arakaki, T., Yoshioka, T., Hayakawa, K., Konohira, E., Tanoue, E., Akhand, A., Chanda, A., Wang, B., Sakugawa, H., 2015. Reviews and syntheses: ocean acidification and its potential impacts on marine ecosystems. *Biogeosciences* 13, 1767–1786.
- Newman, O., Dubuque, C., 2013. The effects of ocean acidification on the food location behavior and locomotion of *Pagurus longicarpus*. *Journal of Emerging Investigators* 30, 1–6.
- Nucci, P.R., Melo, G.A., 2011. Hermit crabs from Brazil: family Paguridae (Crustacea: Decapoda: Paguroidea), except *Pagurus*. *Zootaxa* 3104, 26–41.

- Nucci, P.R., Melo, G.A., 2015. Hermit crabs from Brazil: family Diogenidae (Crustacea: Decapoda: Paguroidea), except *Paguristes*. Zootaxa 3947, 327-346.
- Parker, L., Ross, P., O'Connor, W., Pörtner, H., Scanes, E., Wright, J., 2013. Predicting the response of molluscs to the impact of ocean acidification. Biology (Basel) 2, 651–692.
- Pires, A., Figueira, E., Moreira, A., Soares, A.M.V.M., Freitas, R. 2015. The effects of water acidification, temperature and salinity on the regenerative capacity of the polychaete *Diopatra neapolitana*. Marine Environmental Research 106, 30-41.
- Pörtner, H.O., Langebuch, M., Reipschläger, A., 2004. Biological impact of elevated ocean CO₂ concentrations: Lessons from animal physiology and Earth history. Journal of Oceanography 60, 705-718.
- Przeslawski, R., Byrne, M., Mellin, C., 2015. A review and meta-analysis of the effects of multiple abiotic stressors on marine embryos and larvae. Global Change Biology 21, 2122–2140.
- Ragagnin, M.N., McCarthy, I.D., Fernandez, W.S., Tschiptschin, A.P., Turra, A., 2018. Vulnerability of juvenile hermit crabs to reduced seawater pH and shading. Marine Environmental Research 142, 130–140.
- Rebach, S., 1974. Burying behavior in relation to substrate and temperature in the hermit crab *Pagurus longicarpus*. Ecology 55: 195-198.
- Ries, J.B., Cohen, A.L., McCorkle, D.C., 2009. A nonlinear calcification response to CO₂-induced ocean acidification by the coral *Oculina arbuscula*. Coral Reefs 29, 661–674.
- Roggatz, C.C., Lorch, M., Hardege, J.D., Benoit, D.M., 2016. Ocean acidification affects marine chemical communication by changing structure and function of peptide signalling molecules. Global Change Biology 22, 3914–3926.
- Rossol, D., Bermúdez, R., Hauss, H., Schulz, K. G., Riebesell, U., Sommer, U., Winder, M., 2012. Ocean acidification-induced food quality deterioration constrains trophic transfer. PLoS ONE 7, e34737.
- Sabine, C.L., Feely, R.A., Gruber, N., Key, R.M., Lee, K., Bullister, J.L., Wanninkhof, R., Wong, C.S., Wallace, D.W.R., Tilbrook, B., Millero, F.J., Peng, T.-H., Kozyr, A., Ono, T., Rios, A.F., 2004. The oceanic sink for anthropogenic CO₂. Science 305, 367–371.

- Salmon, M., 1987. On the reproductive behavior of the fiddler crab *Uca thayeri*, with comparisons to *U. pugilator* and *U. vocans*: evidence for behavioral convergence. *Journal of Crustacean Biology* 7, 25-44.
- Schram, J.B., Mcclintock, J.B., Angus, R.A., Lawrence, J.M., 2011. Regenerative capacity and biochemical composition of the sea star *Luidia clathrata* (Say) (Echinodermata:Asteroidea) under conditions of near-future ocean acidification. *Journal of Experimental Marine Biology and Ecology* 407, 266–274.
- Shine, R., 1989. Ecological causes for the evolution of sexual dimorphism: a review of the evidence. *The Quarterly Review of Biology* 64, 419-461.
- Small, D., Calosi, P., White, D., Spicer, J.I., Widdicombe, S., 2010. Impact of medium-term exposure to CO₂-enriched seawater on the physiological functions of the velvet swimming crab *Necora puber*. *Aquatic Biology* 10, 11–21.
- Subramoniam, T., 2017. Sexual biology and reproduction in Crustaceans, first ed. Academic Press/Elsevier, Cambridge.
- Suzuki, Y., Yasuda, C., Takeshita, F., Wada, S., 2012. Male mate choice and male–male competition in the hermit crab *Pagurus nigrofascia*: importance of female quality. *Marine Biology* 159, 1991-1996.
- Taylor, J.R.A., Gilleard, J.M., Allen, M.C., Deheyn, D.D., 2015. Effects of CO₂-induced pH reduction on the exoskeleton structure and biophotonic properties of the shrimp *Lysemata californica*. *Scientific Reports* 5, 10608.
- Taylor, J.R., Lovera, C., Whaling, P.J., Buck, K.R., Pane, E.F., Barry, J.P., 2014. Physiological effects of environmental acidification in the deep-sea urchin *Strongylocentrotus fragilis*. *Biogeosciences* 11, 1413–1423.
- Truchot, J.P., Duhamel-Jouve, A., 1980. Oxygen and carbon dioxide in the marine intertidal environment: diurnal and tidal changes in rockpools. *Respiration Physiology* 39: 241-254.
- Turra, A., 2005. Reproductive behavior of intertidal hermit crabs (Decapoda, Anomura) in southeastern Brazil. *Revista Brasileira de Zoologia* 22, 313-319.
- Turra, A., Denadai, M.R., 2003. Daily activity of four tropical intertidal hermit crabs from southeastern Brazil. *Brazilian Journal of Biology* 63: 537-544.
- Valles-Regino, R., Tate, R., Kelaher, B., Savins, D., Dowell, A., Benkendorff, K., 2015. Ocean warming and CO₂-induced acidification impact the lipid content of a marine predatory gastropod. *Marine Drugs* 13, 6019-6037.

- Vance, R.R., 1972. The role of shell adequacy in behavioral interactions involving hermit crabs. *Ecology* 53: 1075-1083.
- Wang, S.Y., Stickle, W.B., 1988. Biochemical composition of the blue crab *Callinectes sapidus* exposed to the water-soluble fraction of crude oil. *Marine Biology* 98, 23-30.
- White, S.J., Briffa, M., 2016. How do anthropogenic contaminants (ACs) affect behaviour? Multi-level analysis of the effects of copper on boldness in hermit crabs. *Oecologia* 183, 391-400.
- Wheatly, M.G., Henry, R.P., 1992. Extracellular and intracellular acid-base regulation in crustaceans. *Journal of Experimental Zoology* 263, 127-142.
- Whiteley, N.M., 2011. Physiological and ecological responses of crustaceans to ocean acidification. *Marine Ecology Progress Series* 430, 257–271.
- Whiteley, N.M., Suckling, C.C., Ciotti, B.J., Brown, J., McCarthy, I.D., Gimenez, L., Hauton, C., 2018. Physiological compensation for elevated $p\text{CO}_2$ in marine crabs depends on susceptibilities to salinity change. *Scientific Reports* 8, 15639.
- Widdicombe, S., Spicer, J.I., 2008. Predicting the impact of ocean acidification on benthic biodiversity: What can animal physiology tell us? *Journal of Experimental Marine Biology and Ecology* 366, 187–197.
- Wittmann, A.C., Pörtner, H.O., 2013. Sensitivities of extant animal taxa to ocean acidification. *Nature Climate Change* 3, 995–1001.
- Wolfe, K., Dworjanyn, S.A., Byrne, M., 2013. Thermal and pH/ $p\text{CO}_2$ fluctuations in the intertidal habitat of *Helicoidaris erythrogramma*: effects on post-metamorphic juveniles. *Cahiers du Biologie Marine* 54: 657-666.
- Wood, H.L., Spicer, J.I., Lowe, D.M., Widdicombe, S., 2010. Interaction of ocean acidification and temperature; the high cost of survival in the brittlestar *Ophiura ophiura*. *Marine Biology* 157, 2001–2013.
- Wood, H.L., Spicer, J.I., Widdicombe, S., 2008. Ocean acidification may increase calcification rates, but at a cost. *Proceedings of the Royal Society B: Biological Sciences* 275, 1767–1773.
- Yang, J., Liu, D., Jing, W., Dahms, H-U., Wang, L., 2013. Effects of cadmium on lipid storage and metabolism in the freshwater crab *Sinopotamon henanense*. *PLoS ONE* 8, e77569.

- 772 Yasuda, C., Suzuki, Y., Wada, S., 2011. Function of the major cheliped in male-male
773 competition in the hermit crab *Pagurus nigrofascia*. *Marine Biology* 158, 2327–
774 2334.
- 775 Yasuda, C.I., Matsuo, K., Wada, S., 2014. Rapid regeneration of the major cheliped in
776 relation to its function in male-male contests in the hermit crab *Pagurus*
777 *middendorffii*. *Plankton & Benthos Research* 9, 122–131.
- 778 Yasuda, C.I., Koga, T., 2016. Importance of weapon size in all stages of male-male
779 contests in the hermit crab *Pagurus minutus*. *Behavioral Ecology and Sociobiology*
780 70, 2175–2183.
- 781 Zar, J.H., 2010. *Biostatistical Analysis*, fifth ed. Prentice Hall, Upper Saddle River.
- 782 Zeebe, R.E., 2012. History of seawater carbonate chemistry, atmospheric CO₂, and ocean
783 acidification. *Annual Review of Earth and Planetary Sciences* 40, 141–165.
- 784 Zheng, C., Jeswin, J., Shen, K., Lablache, M., Wang, K., Liu, H., 2015. Detrimental effect
785 of CO₂-driven seawater acidification on a crustacean brine shrimp, *Artemia sinica*.
786 *Fish and Shellfish Immunology* 43, 181-190.

787

788 **Figure captions**

789

790 Figure 1. Mortality of males and females of the hermit crab *Pagurus criniticornis*
791 subjected to experimental amputation of the right cheliped (autotomised and non-
792 autotomised) and maintained at different pH treatments (control: pH 8.25; reduced: pH
793 7.70), during a 120-day experimental period. (A) Accumulated mortality, represented by
794 the accumulated number of dead individuals across the time intervals (30, 60, 90 and 120
795 days); and (B) Average mortality [number of dead individuals per experimental tank
796 (n=10) per treatment; Mean $\pm$ Standard deviation] at the end of the experiment (T120).

Figure 2. Growth (Mean $\pm$ Standard deviation), expressed by the difference between the final and initial shield length (mm), of males and females of the hermit crab *Pagurus criniticornis* subjected to experimental amputation of the right cheliped (autotomised and non-autotomised) and maintained at different pH treatments (control: pH 8.25; reduced: pH 7.70) during a 120-day experimental period.

Figure 3. Number of molts (Mean $\pm$ Standard deviation) of males and females (A) of the hermit crab *Pagurus criniticornis* subjected to experimental amputation of the right cheliped (autotomised and non-autotomised) (B) and maintained at different pH treatments (control: pH 8.25; reduced: pH 7.70) (C) during a 120-day experimental period.

Figure 4. Cheliped regeneration, represented by the right cheliped length (Mean $\pm$ Standard deviation; mm) at the end of the experiment, of males and females of the hermit crab *Pagurus criniticornis* subjected to experimental amputation and maintained at different pH treatments (control: pH 8.25; reduced: pH 7.70) during a 120-day experimental period.

819

820 Figure 5. Startle response (Mean $\pm$ Standard deviation; s^{-1}) of males and females of the
821 hermit crab *Pagurus criniticornis* subjected to experimental amputation of the right
822 cheliped (autotomised and non-autotomised) and maintained at different pH treatments
823 (control: pH 8.25; reduced: pH 7.70) during a 120-day experimental period.

824

826

827 Figure 6. Energy budget, represented by (A) Lipid content (Mean $\pm$ Standard deviation;
828 mg/g) and (B) Calcium content of the left cheliped (Mean $\pm$ Standard deviation; mg/g),
829 of males and females of the hermit crab *Pagurus criniticornis* subjected to experimental
830 amputation of the right cheliped (autotomised and non-autotomised) and maintained at
831 different pH treatments (control: pH 8.25; reduced: pH 7.70) during a 120-day
832 experimental period. * indicate significant difference in calcium content between males
833 and females within the combination of amputation and pH treatments.

834

Table caption

Table 1. Mean ($\pm$ standard deviation; SD) of abiotic seawater parameters measured daily (n=120; pH, salinity, temperature) and monthly (n=4; dissolved inorganic carbon – DIC; total alkalinity – TA) in the control and experimental treatments.

Parameters	Mean ($\pm$ SD)	
	Control	Experimental
pH	8.25 (± 0.11)	7.70 (± 0.11)
Salinity	31.2 (± 0.1)	31.1 (± 0.3)
Temperature ($^{\circ}\text{C}$)	23.2 (± 0.5)	23.2 (± 0.4)
DIC ($\mu\text{mol/kgSW}$)	2,573 (± 192)	2,872 (± 70)
TA ($\mu\text{mol/kgSW}$)	2,366 (± 19.3)	2,287 (± 149)

849

850 Table 2. Three-way ANOVA of the physiological stress indicators of males and females of the hermit crab *Pagurus criniticornis* subjected to
 851 experimental amputation of the right cheliped (autotomised and non-autotomised) and maintained at different pH treatments (control: pH 8.25;
 852 reduced: pH 7.70) during a 120-day experimental period. A) Mortality (average number of dead individuals per experimental tank; n=10); B)
 853 Growth (average difference in the shield length between final and initial conditions; mm); C) Number of molts (average number of molts per
 854 individual) and D) Startle response (s^{-1} ; data transformed as $\text{Log}_{10}[\text{SR}+1]$ prior to ANOVA).

855

Effect	A) Mortality				B) Growth				C) Number of molts				D) Startle Response			
	$\frac{D}{F}$	MS	F	p	$\frac{D}{F}$	MS	F	p	$\frac{D}{F}$	MS	F	p	DF	MS	F	p
Intercept	1	49.613	133.787	<0.001	1	6.991	59.418	<0.001	1	183.898	171.618	<0.001	1	203.744	751.403	<0.001
pH	1	10.513	28.348	<0.001	1	0.017	0.141	0.709	1	8.605	8.031	0.006	1	0.009	0.033	0.855
Sex	1	0.113	0.303	0.583	1	0.165	1.400	0.244	1	6.615	6.173	0.015	1	1.822	6.942	0.010
Autotomy	1	0.113	0.303	0.583	1	0.093	0.794	0.378	1	10.645	9.934	0.002	1	0.0004	0.001	0.973
pH*Sex	1	1.01	2.73	0.102	1	0.011	0.096	0.758	1	0.028	0.026	0.872	1	0.145	0.536	0.466
pH*Autotomy	1	0.313	0.842	0.361	1	0.043	0.365	0.549	1	0.720	0.672	0.414	1	0.139	0.514	0.475
Sex*Autotomy	1	0.613	1.651	0.202	1	0.011	0.091	0.765	1	0.149	0.139	0.710	1	1.529	5.640	0.020
pH*Sex*Autotomy	1	0.013	0.033	0.854	1	0.038	0.321	0.574	1	1.185	1.105	0.296	1	0.115	0.423	0.518
Error	72	0.37			40	0.118			92	1.072			81	0.271		

856

857

858

Table 3. Two-way ANOVA of the cheliped regeneration, represented by the right cheliped length (mm) at the end of the experiment, of males and females of the hermit crab *Pagurus criniticornis* subjected to experimental amputation and maintained at different pH treatments (control: pH 8.25; reduced: pH 7.70) during a 120-day experimental period.

Effect	<i>SS</i>	<i>DF</i>	<i>MS</i>	<i>F</i>	<i>p</i>
Intercept	10583.53	1	10583.53	2210.618	<0.001
Sex	210.59	1	210.59	43.987	<0.001
pH	8.77	1	8.77	1.831	0.183
Sex*pH	5.12	1	5.12	1.07	0.306
Error	201.08	42	4.79		

875 Table 4. Three-way ANOVA of the energy budget (lipid and calcium contents; mg/g) of males and females of the hermit crab *Pagurus criniticornis*
876 subjected to experimental amputation of the right cheliped (autotomised and non-autotomised) and maintained at different pH treatments (control:
877 pH 8.25; reduced: pH 7.70) during a 120-day experimental period.

878

Effect	A) Total lipid content				B) Calcium content			
	<i>DF</i>	<i>MF</i>	<i>F</i>	<i>p</i>	<i>DF</i>	<i>MF</i>	<i>F</i>	<i>p</i>
Intercept	1	122.546	946.564	<0.001	1	4129006.422	9270.433	<0.001
pH	1	8.9	68.741	<0.001	1	261.128	0.586	0.445
Sex	1	2.302	17.777	<0.001	1	3958.665	8.888	0.003
Autotomy	1	0.021	0.164	0.687	1	357.14	0.802	0.372
pH*Sex	1	0.002	0.017	0.898	1	78.11	0.175	0.676
pH*Autotomy	1	0.66	5.095	0.027	1	1576.124	3.539	0.062
Sex*Autotomy	1	0.012	0.092	0.762	1	22.667	0.051	0.822
pH*Sex*Autotomy	1	0.111	0.856	0.357	1	3917.977	8.797	0.004
Error	86	0.13			133	445.395		

879

880