



The effect of unilateral hand contractions on psychophysiological activity during motor performance: Evidence of verbal-analytical engagement

Hoskens, Merel; Bellomo, Eduardo; Uiga, Liis; Cooke, Andrew; Masters, Rich

Psychology of Sport and Exercise

DOI:

[10.1016/j.psychsport.2020.101668](https://doi.org/10.1016/j.psychsport.2020.101668)

Published: 30/05/2020

Peer reviewed version

[Cyswllt i'r cyhoeddiad / Link to publication](#)

Dyfyniad o'r fersiwn a gyhoeddwyd / Citation for published version (APA):

Hoskens, M., Bellomo, E., Uiga, L., Cooke, A., & Masters, R. (2020). The effect of unilateral hand contractions on psychophysiological activity during motor performance: Evidence of verbal-analytical engagement. *Psychology of Sport and Exercise*, 48, Article 101668. <https://doi.org/10.1016/j.psychsport.2020.101668>

Hawliau Cyffredinol / General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?


Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Journal Pre-proof

The effect of unilateral hand contractions on psychophysiological activity during motor performance

Merel C.J. Hoskens, Eduardo Bellomo, Liis Uiga, Andrew Cooke, Rich S.W. Masters



PII: S1469-0292(19)30778-2

DOI: <https://doi.org/10.1016/j.psychsport.2020.101668>

Reference: PSYSPO 101668

To appear in: *Psychology of Sport & Exercise*

Received Date: 12 November 2019

Revised Date: 23 January 2020

Accepted Date: 9 February 2020

Please cite this article as: Hoskens, M.C.J., Bellomo, E., Uiga, L., Cooke, A., Masters, R.S.W., The effect of unilateral hand contractions on psychophysiological activity during motor performance, *Psychology of Sport & Exercise* (2020), doi: <https://doi.org/10.1016/j.psychsport.2020.101668>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 Published by Elsevier Ltd.

Merel Hoskens: Conceptualization, methodology, formal analysis, investigation, writing - original draft, visualization,

Eduardo Bellomo: Methodology, formal analysis, investigation, writing – review & editing

Liis Uiga: Conceptualization, methodology, writing – review & editing and supervision

Andrew Cooke: Conceptualization, methodology, formal analysis, writing – review & editing and supervision

Rich Masters: Conceptualization, methodology, formal analysis, writing – review & editing and supervision

Journal Pre-proof

The effect of unilateral hand contractions on psychophysiological activity
during motor performance

Merel C.J. Hoskens^{1}, Eduardo Bellomo², Liis Uiga¹, Andrew Cooke² & Rich S.W. Masters¹*

1: Te Huataki Waiora School of Health, University of Waikato, NZ

2: School of Sport, Health & Exercise Sciences, Bangor University, UK

Corresponding author*:

Merel Hoskens

School of Health

University of Waikato

Private Bag 3105, Hamilton 3240, New Zealand

E: mcjhh1@students.waikato.ac.nz

Abstract

Objectives: Conscious engagement in movement control can influence motor performance. In most cases, the left hemisphere of the brain plays an important role in verbal-analytical processing and reasoning, so changes in the balance of hemispheric activation may influence conscious engagement in movement. Evidence suggests that unilateral hand contractions influence hemispheric activation, but no study has investigated whether there is an associated effect of hand contractions on verbal-analytical processing and psychophysiological activity during motor performance. This study was designed to examine whether pre-performance unilateral hand contraction protocols change verbal-analytical involvement and psychophysiological activity during motor performance. **Design:** A repeated measures crossover design was employed. **Methods:** Twenty-eight participants completed three hand contraction protocols in a randomised order: left, right and no-hand contractions. Electroencephalography (EEG) measures of hemispheric asymmetry were computed during hand contractions. A golf putting task was conducted after each protocol. EEG connectivity between sites overlying the left verbal-analytical temporal region (T7) and the motor planning region (Fz) was computed for the 3-sec prior to movement initiation. Additionally, electrocardiography (ECG) and electromyography (EMG) signals were analysed 6-sec prior to movement initiation until 6-sec after. Golf putting performance was obtained by distance from the target and putter swing kinematics. **Results:** Contralateral hemisphere activity was revealed for the left and right-hand contraction conditions. During motor planning, the left-hand contraction protocol led to significantly lower T7-Fz connectivity, and the right-hand contraction protocol led to significantly higher T7-Fz connectivity than the other conditions. EMG, ECG and kinematic measures did not differ as a function of condition. Importantly, T7-Fz connectivity mediated the relationship between hand squeezing and motor performance (distance from the target). **Conclusion:** The EEG results suggest that pre-

performance unilateral hand contractions influence the extent of verbal-analytical engagement in motor planning, which in turn influences motor performance. However, the hand contractions did not influence cardiac activity, muscle activity or kinematics.

Key words: hand contraction protocol; hemisphere-specific priming; EEG; heart rate; movement kinematics

Journal Pre-proof

Journal Pre-proof

Introduction

A link between conscious processes and motor performance is found in studies using electroencephalography (EEG) to examine communication (synchronization) between different regions of the brain (Babiloni et al., 2011; Deeny, Hillman, Janelle, & Hatfield, 2003; Gallicchio, Cooke, & Ring, 2016; Zhu, Poolton, Wilson, Maxwell, & Masters, 2011). Evidence from these studies suggests that high conscious engagement in motor performance is associated with more synchronous neuronal activity, indexing greater functional communication between the left temporal T7 region of the brain (involved in verbal-analytical processing), and the frontal midline Fz region of the brain (involved in motor planning) (Babiloni et al., 2011; Deeny et al., 2003; Gallicchio et al., 2016; Zhu et al., 2011).

Compelling evidence for the link between conscious control of movements and verbal-analytical processes has been reported by Zhu et al. (2011, Experiment 1). They measured propensity to consciously control motor skills using the Movement Specific Reinvestment Scale (MSRS, Masters, Eves, & Maxwell, 2005). Participants with a lower propensity to consciously control movements displayed lower T7-Fz communication (e.g., coherence) than participants with a higher propensity for conscious control, during the 4-sec preceding golf putts (Zhu et al., 2011). Co-activation between the left temporal and frontal regions is also associated with motor performance. For example, Gallicchio et al. (2016) reported that T7-Fz connectivity was lower in the final seconds preceding successful golf putts compared to unsuccessful golf putts, suggesting that reduced or suppressed verbal-analytical processing is a feature of effective motor performance. In sum, reduced left temporal-frontal synchronicity may be associated with less verbal, more procedural, processing of movements.

Attempts to reduce verbal-analytical engagement during motor performance have used neuro-stimulation to suppress activity in the left hemisphere (Landers et al., 1991;

26 Snyder et al., 2003; Zhu et al., 2015). For instance, Zhu et al. (2015) found that cathodal (i.e.,
27 inhibitory) transcranial Direct Current Stimulation (tDCS) over the left dorsolateral prefrontal
28 cortex promoted lower verbal-analytical engagement when practicing a golf putting task,
29 compared to sham stimulation (i.e., placebo). However, tDCS is not a practical or accessible
30 training method for the majority of performers, and ethical concerns about such extreme
31 training methods have been raised (Davis, 2013).

32 Using a slightly less shocking method, Beckmann, Gröpel, and Ehrlenspiel (2013) and
33 Gröpel and Beckmann (2017) asked semi-professional athletes (gymnastics, soccer,
34 badminton and taekwondo) to squeeze a stress ball in either the left hand or the right hand for
35 45-sec before performing under competitive pressure. They reasoned that due to the
36 contralateral coupling between our hands and our brain (i.e., the brain area controlling the
37 right hand resides in left hemisphere, and vice-versa), squeezing the right hand should prime
38 the left (verbal-analytic) hemisphere and squeezing the left hand should prime the right
39 (visual-spatial) hemisphere. Results showed that left-hand contractions resulted in more
40 stable performance under pressure than right-hand contractions. The authors argued that left-
41 hand contractions prevented breakdown under pressure by activating the right hemisphere
42 and deactivating the left hemisphere, which reduced disruptive verbal-analytical control of
43 the movements (Beckmann et al., 2013; Gröpel & Beckmann, 2017). Beckmann et al. (2013,
44 Experiment 3) additionally found that right-hand contractions magnified the effect of
45 pressure, with participants performing worse when they carried out right-hand contractions
46 prior to performing. They suggested that since right-hand contractions activated the left
47 hemisphere, they potentially increased the likelihood that pressure would cause disruptive
48 verbal-analytical involvement in performance. However, it is important to note that this
49 interpretation cannot be confirmed since Beckmann and colleagues did not directly measure
50 cortical activity in their studies.

51 Studies that did record cortical activity during unilateral hand contractions have
52 revealed inconsistent results. For example, some studies revealed that unilateral hand
53 contractions result in lower alpha power (i.e., increased brain activity) in the contralateral
54 hemisphere (Gable, Poole, & Cook, 2013; Harmon-Jones, 2006; Peterson, Shackman, &
55 Harmon-Jones, 2008; Schiff, Guirguis, Kenwood, & Herman, 1998). However, Cross-
56 Villasana, Gropel, Doppelmayr, and Beckmann (2015) revealed that unilateral hand
57 contractions produced lower alpha power over both hemispheres. Furthermore, they revealed
58 that immediately after left-hand contractions ceased, whole scalp alpha power increased,
59 indicating widespread deactivation (Cross-Villasana et al., 2015). This latter finding
60 challenges Beckmann and colleagues suggestion that left-hand contractions are beneficial
61 because they activate the right hemisphere. However, it does support the argument that left-
62 hand contractions can deactivate the left hemisphere, perhaps suppressing verbal-analytical
63 engagement in motor planning. Taken together, these findings indicate that hemispheric
64 activity can be altered by hand contraction protocols. However, their effects on verbal-
65 analytical processes have yet to be established. Specifically, no study has examined the effect
66 of unilateral hand contractions on T7-Fz connectivity during the final moments of motor
67 preparation. These final moments are important for establishing the level of conscious
68 monitoring and control of the movement (e.g., Deeny et al., 2003; Gallicchio et al., 2016; Zhu
69 et al., 2011). Therefore, measurement of cortical activity, especially T7-Fz connectivity, is
70 required to more rigorously examine the proposed relations between left-hand contractions,
71 verbal-analytical engagement and motor performance.

72 Finally, no studies have investigated the effects of hand contraction protocols on
73 physiological and kinematic measures that may also relate to verbal-analytical engagement
74 and motor performance outcomes (Cooke, Kavussanu, McIntyre, & Ring, 2010). Although
75 Cooke et al. (2014) did not examine hand contractions, they did report greater heart rate

76 deceleration during the 6-sec preceding motor performance in skilled versus low skilled
77 golfers. Therefore, heart rate deceleration could offer another corroborative physiological
78 measure that is sensitive to the amount of verbal-analytical engagement during motor
79 planning (Cooke et al., 2014; Neumann & Thomas, 2009; Neumann & Thomas, 2011; Radlo,
80 Steinberg, Singer, Barba, & Melnikov, 2002). Similarly, more automatic motor control is also
81 associated with lower muscle activity (Lohse, Sherwood, & Healy, 2010; Vance, Wulf,
82 Tollner, McNevin, & Mercer, 2004; Zachry, Wulf, Mercer, & Bezodis, 2005). For example,
83 Lohse et al. (2010) revealed lower muscle activity when participants adopted an external
84 focus of attention while throwing darts, compared to when they consciously monitored their
85 technique. Finally, movement kinematics can also be linked to verbal-analytical engagement
86 in motor planning (Cooke et al., 2014; Malhotra, Poolton, Wilson, Omuro, & Masters, 2015;
87 Masters, Poolton, Maxwell, & Raab, 2008; Maxwell, Masters, & Eves, 2003). For example,
88 Maxwell et al. (2003) revealed that verbal-analytic engagement in motor planning was
89 associated with a less fluid technique. The assessment of such measures alongside T7-Fz
90 connectivity may therefore provide new insight into the mechanisms underpinning the effects
91 of unilateral hand contraction protocols on performance.

92 The present study is the first to investigate the effect of unilateral hand contraction
93 protocols on psychophysiological and behavioural markers of golf putting performance. The
94 aim was to gain a better understanding of whether pre-performance unilateral hand
95 contractions have an effect on verbal-analytical processes involved in motor performance.
96 Three hand contraction protocols (left, right and no-hand) were performed in a repeated
97 measures crossover design, before performance of a golf putting task. Measures of alpha
98 power (8-12 Hz) between homologous electrode pairs were first computed during the hand
99 contraction protocols to verify that left-hand contractions activated the right hemisphere, and
100 that right-hand contractions activated the left hemisphere. Cortical activity was then

101 examined further by measuring the high-alpha power (10-12 Hz) connectivity level between
102 the verbal-analytical left temporal (T7) region and the motor planning (Fz) region during
103 preparation for each golf putt. Cardiac activity (electrocardiography), muscle activity
104 (electromyography), kinematics, and golf performance were tested as supporting measures of
105 verbal-analytical engagement in motor planning. Mediation analyses were employed to
106 examine whether our EEG and psychophysiological indices of verbal-analytic engagement
107 are the mechanisms underpinning any effect of hand contractions on performance.

108 Based on the behavioural findings of Beckmann et al. (2013) and Gröpel and
109 Beckmann (2017), we predicted that unilateral hand contractions would influence verbal-
110 analytical involvement (i.e., inferred by changes in T7-Fz connectivity) during movement
111 planning. Specifically, we predicted that the left-hand contractions would lower verbal-
112 analytical involvement during motor planning compared to right-hand and no-hand
113 contractions, and that right-hand contractions would raise verbal-analytical involvement in
114 motor planning compared to left-hand and no-hand contractions. Consequently, lower verbal-
115 analytical engagement during the left-hand contraction protocol was expected to promote
116 greater heart rate deceleration, lower muscular activity, smoother kinematics when initiating
117 the golf putt and better outcome performance compared to the right-hand and no-hand
118 contraction protocols (Cooke et al., 2014; Lohse et al., 2010; Neumann & Thomas, 2009;
119 Radlo et al., 2002; Zachry et al., 2005). The opposite effects were predicted for the right-hand
120 contraction protocol. Finally, we predicted that the effects of hand contractions on T7-Fz
121 connectivity and our ECG, EMG and kinematic measures would mediate the relationship
122 between hand contraction protocols and performance.

123

Methods**124 Participants and design**

125 Twenty-eight people were recruited to participate in the experiment. Three
126 participants who had major artefacts in their EEG signal were excluded from further analysis,
127 resulting in a final sample of twenty-five participants (mean age = 26.52, SD = 5.08, female =
128 15). To control for handedness, only right-handed participants were included (> 70 ,
129 Edinburgh Handedness Inventory, Oldfield, 1971). All participants had normal/corrected
130 vision. The participants were instructed not to consume alcohol or drugs 24-hours prior to
131 testing or caffeine 3-hours prior to testing, and to obtain at least 6-hours of sleep the night
132 before testing. A repeated measures crossover design was adopted, with participants
133 performing three different protocols (right, left and no-hand contractions). The order of
134 protocols was counterbalanced within participants. This study was approved by the
135 University (Human) Research ethics committee.

136 Task

137 The experiment consisted of a pre-performance hand contraction protocol followed by
138 a golf putting task. The hand contraction protocol required participants to firmly contract a
139 stress ball at a self-paced rate for 45-sec either with their left hand or right hand, or to place
140 their hands on their lap and hold them still for 45-sec (no-hand contraction condition). The
141 researcher instructed the participants to sit quietly and to not talk or make large movements
142 during these protocols, in order to control for muscle activity artefacts.

143 After each protocol, participants performed 25 golf putts on an artificial grass surface,
144 using a standard length (90-cm) golf putter and a regular-size (diameter 4.7-cm) golf ball.
145 The target was a 1-cm diameter white sticker on the putting surface positioned 2.4-m from
146 the initial starting point. Mean radial error (mean distance in any direction from the target)
147 was assessed.

148 **Measures**149 **Psychophysiological measures.**

150 EEG data was used to assess cortical activity during the pre-performance hand
151 contraction protocols (e.g., Gable et al., 2013) and during preparation of the golf putts (e.g.,
152 Zhu et al., 2011). EEG was recorded from thirty-two (32) active electrodes positioned using
153 the 10-20 system (Jaspers, 1958): Fp1, Fp2, AF3, AF4, F7, F3, Fz, F4, F8, FC5, FC1, FC2,
154 FC6, T7, C3, Cz, C4, T8, CP5, CP1, CP2, CP6, P7, P3, Pz, P4, P8, PO3, PO4, O1, Oz, and
155 O2. Additionally, active electrodes were positioned on each mastoid, at the outer canthus and
156 below each eye to record vertical and horizontal electrooculogram (EOG). Monopolar
157 recorded signals were sampled at 1024 Hz, without an online filter, using an ActiveTwo
158 amplifier (Biosemi, The Netherlands).

159 During the pre-performance protocols, we were primarily interested in cortical
160 asymmetry (i.e., right hemisphere minus left hemisphere) in the broad alpha band frequency
161 (i.e., 8-12 Hz), as previous studies have demonstrated the effects of unilateral hand
162 contractions on broad-band alpha (Cross-Villasana et al., 2015; Gable et al., 2013; Harmon-
163 Jones, 2006; Peterson et al., 2008). During preparation of the golf putt, we were interested in
164 connectivity in the high-alpha frequency band (i.e., 10-12 Hz), as this portion of the alpha
165 frequency is thought to be specifically related to task specific attentional processes and
166 cortico-communication (Smith, McEvoy, & Gevins, 1999; for a review see Klimesch, 1999).

167 Electrocardiography (ECG) was used during golf putting performance, to assess
168 cardiac activity (Cooke et al., 2014; Cooke, Kavussanu, McIntyre, Boardley, & Ring, 2011).
169 Silver/silver chloride spot electrodes (BlueSensor SP, Ambu, Cambridgeshire, UK) were
170 placed on each clavicle and on the lowest left rib. The ECG signal was amplified (Bagnoli-4,
171 Delsys, Boston, MA), filtered (1-100 Hz) and digitized at 2500 Hz with 16-bits resolution

172 (CED Power 1401, Cambridge Electronic Design, Cambridge, UK) using Spike2 software
173 (version 5, Cambridge Electronic Design).

174 Electromyography (EMG) was used to obtain muscle activity during golf putting for
175 the extensor carpi radialis and flexor carpi ulnaris muscles in the left arm (Cooke et al., 2014;
176 Cooke et al., 2011). Differential surface electrodes (DE 2.1, Delsys) were placed on the belly
177 of the muscles and a ground electrode (BleuSensor SP, Ambu, Cambridgeshire, UK) was
178 placed on the left collarbone. The EMG signal was amplified (Bagnoli-4, Delsys), filtered
179 (20-45 Hz), and digitized at 2500 Hz with 16-bit resolution (Power 1401) using Spike2
180 software.

181 **Golf putting performance measures.**

182 The golf putting performance was determined by the mean radial error (cm),
183 representing the mean distance between the final position of the ball and the centre of the
184 target. This measure was computed with *ScorePutting* software (written in National
185 Instruments LabVIEW), which uses the photographs from a camera system directly placed
186 above the targets to control for angle differences (Neumann & Thomas, 2008).

187 **Golf kinematics.**

188 A triaxial accelerometer (LIS3L06AL, ST Microelectronics, Geneva, Switzerland)
189 and amplifier (frequency response of DC to 15 Hz) were attached to the rear of the putter
190 head in order to measure movement kinematics (Cooke et al., 2014; Cooke et al., 2011).
191 Acceleration of the golf putter from downswing until ball contact was calculated for the x, y
192 and z-axes (representing the lateral, vertical and back-and-forth movement of the club head),
193 to determine club head orientation, swing height and impact force (Spike2, version 5,
194 Cambridge Electronic Design).

195 Procedure

196 Participants were informed about the context of the study and signed an informed
197 consent form prior to the start of the experimental procedure. The EEG, ECG and EMG
198 equipment were set up and a 2-min EEG resting state measurement was performed (1-min
199 open eyes and 1-min closed eyes).

200 Participants first completed 130 putts as part of a separate investigation of the
201 psychophysiological corollaries of practice (data not reported here). The putts served to
202 familiarise participants with the task. This was followed by performing one of the three pre-
203 performance hand-contraction protocols (left, right or no-hand contractions) while seated.
204 Immediately after each protocol, participants were instructed to stand-up and perform 25 self-
205 paced golf putts, aiming for the target as accurately as possible. The time lag between the end
206 of the squeezing protocol and the start of the putting task was approximately 10-sec. A
207 photograph of the final position of the golf ball was taken after each trial. The researcher then
208 collected the golf ball and positioned it for the next trial, thereby standardising the inter-trial
209 interval, and reducing the need for participants to move in-between putts. This procedure was
210 repeated for all conditions (three times in total) and took on average 5-min and 53-sec per
211 condition.

212 Analysis**213 Pre-performance hand contraction protocols.**

214 EEG signals captured during the hand contraction protocols were processed offline
215 with EEGLAB software (Delorme & Makeig, 2004) running on MATLAB (Mathwork, Inc.,
216 USA version 2018b) to compute the power asymmetry. The signals were first resampled to
217 250 Hz, re-referenced to the average of all electrodes, and filtered (.01-30 Hz bandpass filter).
218 The IAF toolbox was used to adjust the alpha frequency band for each participant based on

219 their individual alpha frequency peak, determined from the baseline measure (Corcoran,
220 Alday, Schlesewsky, & Bornkessel-Schlesewsky, 2018).

221 The signals were then subjected to a threshold-based artefact removal procedure,
222 where any 250-ms window containing signal fluctuations exceeding $\pm 150 \mu\text{V}$ was rejected
223 (ERPLAB Toolbox, Lopez-Calderon & Luck, 2014). Independent Component Analyses were
224 then performed via the RunICA infomax algorithm (Makeig, Bell, Jung, & Sejnowski, 1996)
225 to identify and remove any remaining artefacts and non-neural activity (e.g., eye-blinks) from
226 the signal. An average of 5.76 components were rejected. The clean signal was then subjected
227 to a time frequency analysis, to obtain the estimate of instantaneous alpha power for the 38-
228 sec of the hand contraction protocols. The total of 45-sec was reduced by 7-sec, due to some
229 participants showing increased artefacts at the end. This analysis was performed by
230 convolving the Fast-Fourier Transform (FFT) power spectrum of the signal with a family of
231 complex Morlet wavelets and eventually taking the inverse FFT (Cohen, 2014). All power
232 values were then log transformed to control for skewness and inter-individual differences.
233 Finally, the transformed values were used to compute the asymmetry scores of the
234 homologous electrode pairs close to the cortical regions involved in hand movements (e.g.,
235 Grefkes, Eickhoff, Nowak, Dafotakis, & Fink, 2008): T8-T7, P4-P3, P8-P7, F4-F3, F8-F7,
236 C4-C3, FC2-FC1, FC6-FC5, CP2-CP1, CP6-CP5 (right – left). This is a common way of
237 calculating alpha asymmetry to identify the effects of a state manipulation (e.g., unilateral
238 hand contractions) on the relative activation of the right hemisphere versus left hemisphere of
239 the brain (e.g., Harmon-Jones, 2006). A higher asymmetry score signifies more activity in the
240 left hemisphere (inverse of alpha activity) compared to the right hemisphere (Harmon-Jones,
241 2006; Wolf et al., 2015).

242 **Golf putting task.**

243 An optical sensor and microphone were used to mark movement initiation and ball
 244 contact in the continuous data (Spike2 and ActiView software, Biosemi), in order to analyse
 245 the psychophysiological measures prior to and during the golf putts. The optical sensor (S51-
 246 PA-2-C10PK, Datasensor, Monte San Pietro, Italy) was used to identify swing-onset by
 247 detecting when the infrared beam was broken by movement of the putter head. The
 248 microphone (NT1, Rode, Silverwater, Australia) was linked to a mixing desk (Club 2000,
 249 Studiomaster, Leighton Buzzard, UK) to detect putter-to-ball contact.

250 Connectivity prior to movement initiation was computed offline by processing the
 251 EEG signals (EEGLAB software) computed during the golf putt preparation. The signals
 252 were cut into epochs of 5-sec (4-sec prior to and 1-sec after movement initiation). Thereafter,
 253 the signals were filtered and cleaned with the same methods as for the *hand contraction*
 254 *protocols*. The signals were then baseline corrected (-.2 to 0-sec, where 0 = movement
 255 initiation; Ring et al., 2015) and time-frequency analysis was performed (see *hand*
 256 *contraction protocols*) to obtain the phase angles. These phase angles were then used to
 257 compute connectivity between the left temporal (T7) and frontal (Fz) regions for the 3-sec
 258 prior to movement initiation, by calculating inter-site phase clustering (ISPC, Cohen, 2014).¹
 259 We calculated $ISPC_{time}$ measuring phase angle differences across the electrodes over time:²

$$ISPC_{xy}(f) = \left| n^{-1} \sum_{t=1}^n e^{i(\theta_x(tf) - \theta_y(tf))} \right|$$

¹ Two different methods have been used to measure synchronization in the sport science literature. Earlier work (e.g., Deeny et al., 2003) measured magnitude squared *coherence*; however, more recent research has measured inter-site phase *connectivity* (ISPC). ISPC is based on phase information only, which makes it independent of fluctuations in absolute power (Gallicchio et al., 2016).

² Cohen (2014) suggests that the ISPC *time* measure is appropriate when having relatively long epochs, with 3-sec considered as long.

260 N is the number of data points; i is the imaginary operator; θ_x and θ_y are the phase angles of
 261 the recorded signal at two different scalp locations; t is the time point and f is the frequency
 262 bin. The $e^{i(\theta_x(tf) - \theta_y(tf))}$ represents the complex vector with magnitude 1 and angle $\theta_x - \theta_y$;
 263 $n^{-1} \sum_{t=1}^n (\cdot)$ denotes averaging over time points, and $|\cdot|$ is the module of the averaged vector
 264 (Cohen, 2014; Lachaux, Rodriguez, Martinerie, & Varela, 1999). ISPC is given as a value
 265 between 0 (no functional connection) and 1 (perfect functional connection). Finally, values
 266 were Z-transformed (inverse hyperbolic tangent) to ensure normal distribution (Gallicchio et
 267 al., 2016).

268 The EMG and ECG signals 6-sec prior to until 6-sec after movement initiation were
 269 analysed offline in epochs of 1-sec (Cooke et al., 2014; Moore, Vine, Cooke, Ring, & Wilson,
 270 2012; Neumann & Thomas, 2011). Heart rate was corrected for artefacts and R-wave peaks
 271 were identified. The intervals between the successive R-waves peaks were calculated and
 272 instantaneous heart rate (beats per minute, BPM) was calculated as $6000/(\text{R-R interval})$.
 273 Muscle activity was assessed by rectifying the EMG signal and averaging over 0.5-sec
 274 windows, such that the mean activity between 6.25 and 5.75-sec prior to movement was used
 275 to calculate muscle activity 6-sec before movement, and so on (Cooke et al., 2014).

276 The acceleration of each putt was determined from the initiation of the downswing
 277 phase until the point of contact (Cooke et al., 2014; Cooke et al., 2010; Moore et al., 2012).
 278 Average acceleration was calculated for the x, y, and z-axes. Besides impact velocity, Root
 279 Mean Square (RMS) jerk and smoothness on the z-axis were computed, as the z-axis is the
 280 main axis involved in the putting swing (Cooke et al., 2011; Maxwell et al., 2003).

281 **Statistical analysis.**

282 The cortical activity manipulation check was subjected to a 3 x 10 repeated measures
 283 analysis of variance (ANOVA): Condition (Left, Right, No-hand) x Homologous electrode
 284 pairs (T8-T7, P4-P3, P8-P7, F4-F3, F8-F7, C4-C3, FC2-FC1, FC6-FC5, CP2-CP1, CP6-

285 CP5). The T7-Fz connectivity measure during preparation of the golf putt was subjected to a
286 one-way ANOVA of Condition (Left, Right, No-hand). Cardiac and muscle activity were
287 subjected to a 3 x 13 repeated measures ANOVA: Condition (Left, Right, No-hand) x Time
288 Bin (-6, -5, -4, -3, -2, -1, 0, +1, +2, +3, +4, +5, +6). Golf putting kinematics and golf putting
289 performance were both subjected to a one-way ANOVA of Condition (Left, Right, No-hand).


290 Sphericity was checked and corrected using the Huynh-Feldt correction when
291 necessary. Separate ANOVAs with Bonferroni corrections or polynomial trend analysis were
292 performed when main effects or interactions were found. Effect sizes are reported as partial η
293 squared (η_p^2). The statistical tests were performed using SPSS (IBM, version 25.0) computer
294 software. Significance was set at $p = .05$ for all statistical tests.

295 MEMORE for SPSS (MEdiation and MOderation analysis for REpeated measure
296 designs, Montoya & Hayes, 2017) was used to test within-subject mediation effects on golf
297 putting performance associated with left-hand and right-hand contractions. Mediators were
298 individually tested and included EEG, EMG, ECG and kinematics (i.e., club head orientation,
299 swing height and impact force). The mediation effect (B), standard error (BootSE) and 95%
300 CI (low and high) were reported (Montoya & Hayes, 2017).

301 **Results**

302 **Manipulation check**

303 The results revealed a main effect of Condition, $F(2,42) = 3.95$, $p = .027$, $\eta_p^2 = .16$,
304 with post-hoc analysis revealing a significantly lower asymmetry score for left-hand
305 contractions compared with right-hand contractions ($p = .015$, see Fig. 1). No significant
306 effects were revealed for left-hand contractions compared with no-hand contractions (p
307 $= .180$) or right-hand contractions compared with no-hand contractions ($p = 1.00$). No main
308 effect was found for Homologous electrode pairs, $F(3.20,67.15) = 0.93$, $p = .438$, $\eta_p^2 = .04$.



309

310 **Fig. 1.** Alpha power asymmetry score per condition. Asymmetry score was calculated by: right hemisphere –
 311 left hemisphere (positive values represent higher right-hemisphere power and negative values represent higher
 312 left-hemisphere power). Error bars represent standard error of the mean. (* $p < .05$).

313 **Cortical activity preceding golf putts**

314 The results revealed a main effect of Condition, $F(2,48) = 122.5$, $p < .001$, $\eta_p^2 = .84$.


315 Post-hoc tests revealed that left-hand contractions led to significantly lower T7-Fz

316 connectivity, than right-hand contractions ($p < .001$) or no-hand contractions ($p < .001$, see

317 Fig. 2). Right-hand contractions revealed the opposite effect with significantly higher T7-Fz

318 connectivity compared to left-hand contractions ($p < .001$) and no-hand contractions (p

319 $< .001$, see Fig. 2).




320

321 **Fig. 2.** T7-Fz ISPCtime connectivity during each condition and time bin. Error bars represent standard error of
 322 the mean. (** $p < .001$).

323 Muscle activity


324 No Condition x Time Bin interactions were evident for the extensor carpi radialis,
 325 ($F(24,432) = 1.15, p = .290, \eta_p^2 = .06$, or the flexor carpi ulnaris, $F(24,480) = 0.82, p = .715,$
 326 $\eta_p^2 = .04$. A main effect of Time Bin was evident for the extensor carpi radialis, $F(3.73,67.11)$
 327 $= 9.99, p < .001, \eta_p^2 = .36$, and the flexor carpi ulnaris, $F(4.18,83.61) = 13.51, p < .001, \eta_p^2$
 328 $= .40$. Post-hoc analysis revealed that for the extensor carpi radialis the variance for Time Bin
 329 was best described by a quadratic trend ($p < .001, \eta_p^2 = .53$), with a gradual increase of
 330 activity until peak in activity during movement initiation (time zero), which quickly drops
 331 back to baseline (see Fig. 3). For the flexor carpi ulnaris, variance for Time Bin was also best
 332 described by a quadratic trend ($p < .001, \eta_p^2 = .68$), with similar trends to the extensor carpi
 333 radialis (see Fig. 4). Main effects of Condition were not evident for the extensor carpi radialis,
 334 $F(2,36) = 1.74, p = .191, \eta_p^2 = .09$, or the flexor carpi ulnaris, $F(2,40) = 0.69, p = .510, \eta_p^2$
 335 $= .03$.



336

337 **Fig. 3.** Activity of the extensor carpi radialis in each condition over time. Error bars represent standard error of

338 the mean.




339

340 **Fig. 4.** Activity for of the flexor carpi ulnaris in each condition over time. Error bars represent standard error of

341 the mean.

342 **Cardiac activity**343 The ECG analysis did not reveal a Condition x Time Bin interaction, $F(24,567) = 0.95$,344 $p = .532$, $\eta_p^2 = .04$, or a main effect of Condition, $F(2,48) = 0.62$, $p = .542$, $\eta_p^2 = .03$. A main345 effect of Time Bin was evident, $F(1.57,37.61) = 17.26$, $p < .001$, $\eta_p^2 = .42$. Post-hoc analysis346 revealed that heart rate differences over time was best described by a cubic trend ($p < .001$,347 $\eta_p^2 = .56$). Heart rate decreased during approximately 2-sec preceding movement initiation

348 and then gradually returned to baseline in the 6-sec after movement initiation (see Fig. 5).



349

350 **Fig. 5.** Heart rate in each condition over time (6-sec before until 6-sec after movement initiation). Error bars
 351 represent standard error of the mean.

352 Golf kinematics

353 No differences were evident between conditions for any of the kinematic measures:
 354 acceleration on the x-axis, $F(2,48) = 2.60, p = .085, \eta_p^2 = .10$; acceleration on the y-axis,
 355 $F(1.59,38.26) = 0.65, p = .493, \eta_p^2 = .03$; acceleration on the z-axis, $F(2,44) = 0.55, p = .581,$
 356 $\eta_p^2 = .02$; impact speed, $F(1.52,36.39) = 0.25, p = .718, \eta_p^2 = .01$; RMS jerk, $F(2,46) = 0.31, p$
 357 $= .738, \eta_p^2 = .01$; smoothness, $F(1.59,38.03) = 0.46, p = .592, \eta_p^2 = .02$.

358 Golf putting performance

359 No differences were evident between conditions for mean radial error, $F(2,48) = 1.75,$
 360 $p = .184, \eta_p^2 = .07$.

361 Mediation analysis

362 Mediation analyses were used to examine whether EEG, EMG, ECG or kinematics
 363 mediated the relationship between hand contractions and golf putting performance (mean
 364 radial error). Although there was no significant difference in performance between the
 365 different hand contraction conditions, there was a significant indirect effect of hand
 366 squeezing on performance via T7-Fz connectivity. Within-subject changes in performance
 367 following left-hand versus right-hand contractions were mediated by the changes in EEG T7-

368 Fz connectivity induced by these protocols, $B = -12.41$, $\text{BootSE} = 4.12$, 95% CI [-21.07, -
369 4.94]. The other mediators did not reveal significant indirect effects on performance.

370 **Discussion**

371 The present study was conducted to examine whether pre-performance unilateral hand
372 contraction protocols influence verbal-analytical engagement in motor performance. A
373 repeated measures crossover design was adopted, measuring psychophysiological markers
374 (neural, cardiovascular and muscular) and performance (distance from the target and
375 movement kinematics) of a golf putting task that was completed immediately after
376 performing a hand contraction protocol (left, right and no-hand). During the hand contraction
377 protocols, measures of alpha power spectra between homologous electrode pairs were
378 computed as a manipulation check to determine whether hand contractions caused different
379 hemispheric activation.

380 The manipulation check revealed a significant difference in hemispheric asymmetry
381 between left-hand and right-hand contraction protocols, with the left-hand contraction
382 protocol resulting in more right-hemisphere activity and the right-hand contraction protocol
383 resulting in higher left-hemisphere activity (see Fig. 1). These findings are consistent with
384 previous studies (Gable et al., 2013; Harmon-Jones, 2006; Peterson et al., 2008).

385 Our study is the first to include a no-hand contractions, which makes it possible to
386 compare the effect of left-hand and right-hand contractions relative to no contractions.
387 Asymmetry during the no-hand contraction protocol was not significantly different from
388 either contraction condition, which suggests that hand contractions did not create different
389 asymmetry compared to no-hand contractions. However, hand contractions did achieve
390 different asymmetry compared to each other. The slight rightward bias evident during the no-
391 hand condition is in line with previous studies revealing that right-handedness is related to a

392 bias to rightward hemisphere asymmetry (greater left-hemisphere activity) for resting state
393 alpha power (e.g., Ocklenburg et al., 2019).

394 As hypothesized, a lower level of T7-Fz connectivity during preparation for putts was
395 revealed after left-hand contractions, compared to right-hand and no-hand contractions. The
396 opposite effect was found for right-hand contractions, revealing higher T7-Fz connectivity
397 compared to left-hand and no-hand contractions. Previous studies have suggested that lower
398 T7-Fz connectivity reflects less verbal-analytical engagement in movements (e.g., Deeny et
399 al., 2003; Gallicchio et al., 2016; Zhu et al., 2011). Left-hand contractions in the present
400 study may therefore have lowered T7-Fz connectivity and reduced verbal-analytical
401 engagement in the putting task, compared to right-hand and no-hand contractions.

402 Although there was no significant effect of hand contractions on golf putting
403 performance,³ mediation analysis suggested that hand contractions influenced T7-Fz
404 connectivity, which in turn influenced performance. Beckmann et al. (2013) and Gröpel and
405 Beckmann (2017) speculated that top-down verbal-analytical control processes are the
406 mechanism by which hand contractions influence performance under pressure. Many
407 explanations of skill failure, such as the theory of reinvestment (Masters, 1992; see Masters
408 & Maxwell, 2008 for a review), suggest that attempts to consciously control movements
409 (characterised by verbal-analytical processing), can disrupt normally efficient motor
410 behaviours. Given the hypothesised link between T7-Fz connectivity and conscious verbal
411 engagement of movement, our mediation findings provide some support for their speculation.

412 Although the hand contraction protocols clearly influenced neurophysiological
413 activity, their effects did not extend to the cardiac, muscular or kinematic measures. There

³ It is acceptable to conduct mediation analysis when there is no significant effect of the independent variable (hand contractions) on the dependent variable (golf putting performance) (see e.g., Kenny, Kashy, & Bolger, 1998).

414 were no condition effects for these variables and there were no mediational effects to
415 implicate any of these variables in the relationship between hand contractions and
416 performance. From a theoretical perspective it makes sense that neural measures should be
417 more sensitive to the effects of hand contraction protocols than peripheral measures such as
418 heart rate, because verbal-analytic processes originate from the brain, and any effects they
419 might have on the heart and muscles would be always be secondary. Any effects of
420 psychological processes on cardiac and muscular activity could also have been masked by
421 any physical strain on these variables caused by the golf putting task (e.g., standing posture,
422 swinging arms, etc.).

423 Despite the indirect effect of hand contractions on performance through T7-Fz
424 connectivity, there were no significant performance differences between the different hand
425 contraction protocols. Our participants only performed 130 trials prior to the first hand
426 contraction condition, so they remained relatively inexperienced novices with high inter and
427 intra person performance variability that may have camouflaged any subtle (direct) hand
428 contraction effects. A more cognitively challenging task may reveal performance differences.
429 Zhu et al (2015) also manipulated T7-Fz coherence, using real versus sham tDCS, and also
430 failed to find an effect on golf putting performance alone. However, Zhu et al. (2015) did
431 report a differential effect on golf putting performance under dual-task load (e.g., backwards
432 counting). Alternatively, replicating the experiment with more experienced performers could
433 also increase the likelihood of performance differences. For example, the theory of
434 reinvestment (Masters & Maxwell, 2008) argues that verbal-analytic engagement (e.g., right-
435 hand contractions) would be more detrimental to the performance of autonomous experts than
436 cognitive novices. Effects of condition on the cardiac, muscular and kinematic measures
437 would also be more likely with experienced performers for the same reasons.

438 A limitation of this study is that we did not control force of grip used by participants
439 during the hand contraction protocol. Consequently, differences in hemisphere asymmetry
440 might have been a function of effort or strength. For example, Hirao and Masaki (2018)
441 showed that force and duration of left-hand contractions had differential effects on
442 hemisphere activity. Additionally, a requirement to achieve a specific force during
443 contractions may require more cognitive resources (e.g., Derosière et al., 2014; Hirao &
444 Masaki, 2018). One solution might simply be to measure grip force and include it as a
445 covariate in analysis of hemisphere asymmetry. This issue should be addressed in further
446 studies.

447 Another limitation is that we were unable to determine the longevity of the hand
448 contractions with respect to their effect on cortical activity. Studies suggest that the effects of
449 hand contraction protocols last at least 15-min (e.g., Baumer, Munchau, Weiller, and Liepert
450 (2002). Participants in our study completed 25 trials over approximately a 6-min duration, so
451 it is likely that the effects remained. However, there is little doubt that further research is
452 needed to gain greater understanding of the timecourse of hand contraction effects.

453 To our knowledge this is the first study reporting neural evidence that left-hand
454 contractions lower verbal-analytical engagement in motor planning of a golf putting task. The
455 additional markers (ECG, EMG, kinematics and performance) did not, however, provide
456 supporting evidence of this effect. These secondary markers may have been insufficiently
457 sensitive to reveal the brain's influence over the body. Nevertheless, it appears that the body
458 (the hands) influenced the brain!

459

460

References

- 461 Babiloni, C., Infarinato, F., Marzano, N., Iacoboni, M., Dassu, F., Soricelli, A., . . . Del
462 Percio, C. (2011). Intra-hemispheric functional coupling of alpha rhythms is related to
463 golfer's performance: a coherence EEG study. *International Journal of*
464 *Psychophysiology*, 82(3), 260-268. <https://doi.org/10.1016/j.ijpsycho.2011.09.008>
- 465 Baumer, T., Munchau, A., Weiller, C., & Liepert, J. (2002). Fatigue suppresses ipsilateral
466 intracortical facilitation. *Experimental Brain Research*, 146, 467–473.
467 <https://doi.org/10.1007/s00221-002-1202-x>
- 468 Beckmann, J., Gröpel, P., & Ehrlenspiel, F. (2013). Preventing motor skill failure through
469 hemisphere-specific priming: cases from choking under pressure. *Journal of*
470 *Experimental Psychology General*, 142(3), 679-691.
471 <https://doi.org/10.1037/a0029852>
- 472 Cohen, M. X. (2014). *Analyzing neural time series data: Theory and practice*. Cambridge,
473 MA: MIT Press.
- 474 Cooke, A., Kavussanu, M., Gallicchio, G., Willoughby, A., McIntyre, D., & Ring, C. (2014).
475 Preparation for action: Psychophysiological activity preceding a motor skill as a
476 function of expertise, performance outcome, and psychological pressure.
477 *Psychophysiology*, 51, 374-384. <https://doi.org/10.1111/psyp.12182>
- 478 Cooke, A., Kavussanu, M., McIntyre, D., Boardley, I. D., & Ring, C. (2011). Effects of
479 competitive pressure on expert performance: Underlying psychological, physiological,
480 and kinematic mechanisms. *Psychophysiology*, 48, 1146-1156.
481 <https://doi.org/10.1111/j.1469-8986.2011.01175.x>
- 482 Cooke, A., Kavussanu, M., McIntyre, D., & Ring, C. (2010). Psychological, muscular and
483 kinematic factors mediate performance under pressure. *Psychophysiology*, 47(6),
484 1109-1118. <https://doi.org/10.1111/j.1469-8986.2010.01021.x>

- 485 Corcoran, A. W., Alday, P. M., Schlesewsky, M., & Bornkessel-Schlesewsky, I. (2018).
486 Toward a reliable, automated method of individual alpha frequency (IAF)
487 quantification. *Psychophysiology*, *55*(7), 1-21. <https://doi.org/10.1111/psyp.13064>
- 488 Cross-Villasana, F., Gropel, P., Doppelmayr, M., & Beckmann, J. (2015). Unilateral left-
489 hand contractions produce widespread depression of cortical activity after their
490 execution. *PLoS One*, *10*(12), e0145867.
491 <https://doi.org/10.1371/journal.pone.0145867>
- 492 Davis, N. J. (2013). Neurodoping: brain stimulation as a performance-enhancing measure.
493 *Sports Medicine*, *43*(8), 649-653. <https://doi.org/10.1007/s40279-013-0027-z>
- 494 Deeny, S. P., Hillman, C. H., Janelle, C. M., & Hatfield, B. D. (2003). Cortico-cortical
495 communication and superior performance in skilled marksmen: An EEG coherence
496 analysis. *Journal of Sport and Exercise Psychology*, *25*, 188-204.
497 <https://doi.org/10.1123/jsep.25.2.188>
- 498 Delorme, A., & Makeig, S. (2004). EEGLAB: an open source toolbox for analysis of single-
499 trial EEG dynamics including independent component analysis. *Journal of*
500 *Neuroscience Methods*, *134*, 9-21.
- 501 Derosière, G., Alexandre, F., Bourdillon, N., Mandrick, K., Ward, T. E., & Perrey, S. (2014).
502 Similar scaling of contralateral and ipsilateral cortical responses during graded
503 unimanual force generation. *Neuroimage*, *85*, 471-477.
504 <https://doi.org/10.1016/j.neuroimage.2013.02.006>
- 505 Gable, P. A., Poole, B. D., & Cook, M. S. (2013). Asymmetrical hemisphere activation
506 enhances global-local processing. *Brain and Cognition*, *83*(3), 337-341.
507 <https://doi.org/10.1016/j.bandc.2013.09.012>
- 508 Gallicchio, G., Cooke, A., & Ring, C. (2016). Lower left temporal-frontal connectivity
509 characterizes expert and accurate performance: High-alpha T7-Fz connectivity as a

- 510 marker of conscious processing during movement. *Sport, Exercise, and Performance*
511 *Psychology*, 5(1), 14-24. <https://doi.org/http://dx.doi.org/10.1037/spy0000055>
- 512 Grefkes, C., Eickhoff, S. B., Nowak, D. A., Dafotakis, M., & Fink, G. R. (2008). Dynamic
513 intra- and interhemispheric interactions during unilateral and bilateral hand
514 movements assessed with fMRI and DCM. *Neuroimage*, 41(4), 1382-1394.
515 <https://doi.org/10.1016/j.neuroimage.2008.03.048>
- 516 Gröpel, P., & Beckmann, J. (2017). A pre-performance routine to optimize competition
517 performance in artistic gymnastics. *The Sport Psychologist*, 31(2), 199-207.
518 <https://doi.org/10.1123/tsp.2016-0054>
- 519 Harmon-Jones, E. (2006). Unilateral right-hand contractions cause contralateral alpha power
520 suppression and approach motivational affective experience. *Psychophysiology*, 43(6),
521 598-603. <https://doi.org/10.1111/j.1469-8986.2006.00465.x>
- 522 Hirao, T., & Masaki, H. (2018). Effects of unilateral hand contraction on the persistence of
523 hemispheric asymmetry of cortical activity. *Journal of Psychophysiology*, 1-8.
524 <https://doi.org/10.1027/0269-8803/a000215>
- 525 Jaspers, H. H. (1958). The ten twenty electrode system of the International Federation.
526 *Electroencephalography and Clinical Neurophysiology*, 10, 371-375.
- 527 Kenny, D. A., Kashy, D. A., & Bolger, N. (1998). Data analysis in social psychology.
528 *Handbook of social psychology*, 1, 233-265.
- 529 Lachaux, J. P., Rodriguez, E., Martinerie, J., & Varela, F. J. (1999). Measuring phase
530 synchrony in brain signals. *Human Brain Mapping*, 8(4), 194-208.
531 [https://doi.org/10.1002/\(SICI\)1097-0193\(1999\)8:4<194::AID-HBM4>3.0.CO;2-C](https://doi.org/10.1002/(SICI)1097-0193(1999)8:4<194::AID-HBM4>3.0.CO;2-C)
- 532 Landers, D. M., Petruzzello, S. J., Salazar, W., Crews, D. J., Kubitz, K. A., Gannon, T. L., &
533 Han, M. (1991). The influence of electrocortical biofeedback on performance in pre-

- 534 elite archers. *Medicine & Science in Sports & Exercise*.
- 535 <https://doi.org/http://dx.doi.org/10.1249/00005768-199101000-00018>
- 536 Lohse, K. R., Sherwood, D. E., & Healy, A. F. (2010). How changing the focus of attention
537 affects performance, kinematics, and electromyography in dart throwing. *Human*
538 *Movement Science*, 29(4), 542-555. <https://doi.org/10.1016/j.humov.2010.05.001>
- 539 Lopez-Calderon, J., & Luck, S. J. (2014). ERPLAB: an open-source toolbox for the analysis
540 of event-related potentials. *Frontiers in Human Neuroscience*, 8.
541 <https://doi.org/10.3389/fnhum.2014.00213>
- 542 Makeig, S., Bell, A. J., Jung, T. P., & Sejnowski, T. J. (1996). Independent component
543 analysis of electroencephalographic data. In D. Touretzky, M. Mozer, & M. Hasselmo
544 (Eds.), *Advances in neural information processing systems* (Vol. 8, pp. 145-151).
545 Cambridge, MA: MIT Press.
- 546 Malhotra, N., Poolton, J. M., Wilson, M. R., Omuro, S., & Masters, R. S. W. (2015).
547 Dimensions of movement specific reinvestment in practice of a golf putting task.
548 *Psychology of Sport and Exercise*, 18, 1-8.
549 <https://doi.org/10.1016/j.psychsport.2014.11.008>
- 550 Masters, R. S. W. (1992). Knowledge, knerves and know-how: The role of explicit versus
551 implicit knowledge in the breakdown of a complex motor skill under pressure. *British*
552 *Journal of Psychology*, 83(3), 343-358. [https://doi.org/10.1111/j.2044-](https://doi.org/10.1111/j.2044-8295.1992.tb02446.x)
553 [8295.1992.tb02446.x](https://doi.org/10.1111/j.2044-8295.1992.tb02446.x)
- 554 Masters, R. S. W., Eves, F. F., & Maxwell, J. (2005, 15-19 August 2005). *Development of a*
555 *Movement Specific Reinvestment Scale*. Paper presented at the ISSP 11th World
556 Congress of Sport Psychology, Sydney, Australia.

- 557 Masters, R. S. W., & Maxwell, J. P. (2008). The theory of reinvestment. *International Review*
558 *of Sport and Exercise Psychology*, 1(2), 160-183.
559 <https://doi.org/10.1080/17509840802287218>
- 560 Masters, R. S. W., Poolton, J. M., Maxwell, J. P., & Raab, M. (2008). Implicit motor learning
561 and complex decision making in time-constraints environments. *Journal of Motor*
562 *Behavior*, 40(1), 71-79. <https://doi.org/10.3200/JMBR.40.1.71-80>
- 563 Maxwell, J. P., Masters, R. S. W., & Eves, F. F. (2003). The role of working memory in
564 motor learning and performance. *Consciousness and Cognition*, 12(3), 376-402.
565 [https://doi.org/10.1016/s1053-8100\(03\)00005-9](https://doi.org/10.1016/s1053-8100(03)00005-9)
- 566 Montoya, A. K., & Hayes, A. F. (2017). Two-condition within-participant statistical
567 mediation analysis: A path-analytic framework. *Psychological Methods*, 22(1), 6–27.
568 <https://doi.org/10.1037/met0000086>
- 569 Moore, L. J., Vine, S. J., Cooke, A., Ring, C., & Wilson, M. R. (2012). Quiet eye training
570 expedites motor learning and aids performance under heightened anxiety: the roles of
571 response programming and external attention. *Psychophysiology*, 49(7), 1005-1015.
572 <https://doi.org/10.1111/j.1469-8986.2012.01379.x>
- 573 Neumann, D. L., & Thomas, P. R. (2008). A camera-based scoring system for evaluating
574 performance accuracy during a golf putting task. *Behavior Research Methods*, 40(3),
575 892-897. <https://doi.org/10.3758/brm.40.3.892>
- 576 Neumann, D. L., & Thomas, P. R. (2009). The relationship between skill level and patterns in
577 cardiac and respiratory activity during golf putting. *International Journal of*
578 *Psychophysiology*, 72(3), 276-282. <https://doi.org/10.1016/j.ijpsycho.2009.01.001>
- 579 Neumann, D. L., & Thomas, P. R. (2011). Cardiac and respiratory activity and golf putting
580 performance under attentional focus instructions. *Psychology of Sport and Exercise*,
581 12(4), 451-459. <https://doi.org/10.1016/j.psychsport.2011.02.002>

- 582 Ocklenburg, S., Friedrich, P., Schmitz, J., Schluter, C., Genc, E., Gunturkun, O., . . .
583 Grimshaw, G. (2019). Beyond frontal alpha: investigating hemispheric asymmetries
584 over the EEG frequency spectrum as a function of sex and handedness. *Laterality*,
585 24(5), 505-524. <https://doi.org/10.1080/1357650X.2018.1543314>
- 586 Oldfield, R. C. (1971). The assessment and analysis of handedness: the Edinburgh inventory.
587 *Neuropsychologia*, 9(1), 97-113. [https://doi.org/10.1016/0028-3932\(71\)90067-4](https://doi.org/10.1016/0028-3932(71)90067-4)
- 588 Peterson, C. K., Shackman, A. J., & Harmon-Jones, E. (2008). The role of asymmetrical
589 frontal cortical activity in aggression. *Psychophysiology*, 45(1), 86-92.
590 <https://doi.org/10.1111/j.1469-8986.2007.00597.x>
- 591 Radlo, S. J., Steinberg, G. M., Singer, R. N., Barba, D. A., & Melnikov, A. (2002). The
592 influence of an attentional focus strategy on alpha brain wave activity, heart rate and
593 dart-throwing performance. *International Journal of Sport Psychology*.
- 594 Schiff, B. B., Guirguis, M., Kenwood, C., & Herman, C. P. (1998). Asymmetrical
595 hemispheric activation and behavioral persistence: effects of unilateral muscle
596 contractions. *Neuropsychology*, 12(4), 526-532. <https://doi.org/10.1037/0894-4105.12.4.526>
- 597
598 Smith, M. E., McEvoy, L. K., & Gevins, A. (1999). Neurophysiological indices of strategy
599 development and skill acquisition. *Cognitive Brain Research*, 7, 389-404.
600 [https://doi.org/10.1016/S0926-6410\(98\)00043-3](https://doi.org/10.1016/S0926-6410(98)00043-3)
- 601 Snyder, A. W., Mulcahy, E., Taylor, J. L., Mitchell, D. J., Sachdev, P., & Gandevia, S. C.
602 (2003). Savant-like skills exposed in normal people by suppressing the left fronto-
603 temporal lobe. *Journal of Integrative Neuroscience*, 2(2), 149-158.
604 <https://doi.org/10.1142/S0219635203000287>

- 605 Vance, J., Wulf, G., Tollner, T., McNevin, N., & Mercer, J. (2004). EMG activity as a
606 function of the performer's focus of attention. *Journal of Motor Behavior*, 36(4), 450-
607 459. <https://doi.org/10.3200/JMBR.36.4.450-459>
- 608 Wolf, S., Brölz, E., Keune, P. M., Wesa, B., Hautzinger, M., Birbaumer, N., & Strehl, U.
609 (2015). Motor skill failure or flow-experience? Functional brain asymmetry and brain
610 connectivity in elite and amateur table tennis players. *Biological Psychology*, 105, 95-
611 105. <https://doi.org/http://dx.doi.org/10.1016/j.biopsycho.2015.01.007>
- 612 Zachry, T., Wulf, G., Mercer, J., & Bezodis, N. (2005). Increased movement accuracy and
613 reduced EMG activity as the result of adopting an external focus of attention. *Brain*
614 *Research Bulletin*, 67(4), 304-309. <https://doi.org/10.1016/j.brainresbull.2005.06.035>
- 615 Zhu, F. F., Poolton, J. M., Wilson, M. R., Maxwell, J. P., & Masters, R. S. W. (2011). Neural
616 co-activation as a yardstick of implicit motor learning and the propensity for
617 conscious control of movement. *Biological Psychology*, 87(1), 66-73.
618 <https://doi.org/10.1016/j.biopsycho.2011.02.004>
- 619 Zhu, F. F., Yeung, A. Y., Poolton, J. M., Lee, T. M. C., Leung, G. K. K., & Masters, R. S. W.
620 (2015). Cathodal Transcranial Direct Current Stimulation over left dorsolateral
621 prefrontal cortex area promotes implicit motor learning in a golf putting task. *Brain*
622 *Stimulation*, 8(4), 784-786. <https://doi.org/10.1016/j.brs.2015.02.005>

623

Highlights:

- The effects of unilateral hand contractions during motor performance was investigated
- Unilateral hand contractions influenced the verbal-analytical engagement in motor planning
- Left-hand contractions caused lower verbal-analytical engagement in motor planning
- It appears that the body (the hands) influenced the brain!

Journal Pre-proof

Declaration of interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests:

Journal Pre-proof