



## Genome Sequence of the Acidophilic Sulfate-Reducing Peptococcaceae Strain CEB3

Petzsch, Patrick; Poehlein, Anja; Johnson, D Barrie; Daniel, Rolf; Schlömann, Michael; Mühling, Martin

### Genome Announcements

DOI:

[10.1128/genomeA.00886-15](https://doi.org/10.1128/genomeA.00886-15)

Published: 06/08/2015

Publisher's PDF, also known as Version of record

[Cyswllt i'r cyhoeddiad / Link to publication](#)

*Dyfyniad o'r fersiwn a gyhoeddwyd / Citation for published version (APA):*

Petzsch, P., Poehlein, A., Johnson, D. B., Daniel, R., Schlömann, M., & Mühling, M. (2015). Genome Sequence of the Acidophilic Sulfate-Reducing Peptococcaceae Strain CEB3. *Genome Announcements*, 3(4). <https://doi.org/10.1128/genomeA.00886-15>

#### Hawliau Cyffredinol / General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

#### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

# Genome Sequence of the Acidophilic Sulfate-Reducing *Peptococcaceae* Strain CEB3

Patrick Petzsch,<sup>a</sup> Anja Poehlein,<sup>b</sup> D. Barrie Johnson,<sup>c</sup> Rolf Daniel,<sup>b</sup> Michael Schlömann,<sup>a</sup> Martin Mühling<sup>a</sup>

Institute of Biological Sciences, TU Bergakademie Freiberg, Freiberg, Germany<sup>a</sup>; Genomic and Applied Microbiology & Göttingen Genomics Laboratory, Georg-August-Universität Göttingen, Göttingen, Germany<sup>b</sup>; College of Natural Sciences, Bangor University, Bangor, United Kingdom<sup>c</sup>

P.P. and A.P. contributed equally to this work.

**We report the draft genome of the *Peptococcaceae* strain CEB3 that originated from an acidic (pH 2.5) stream draining an abandoned copper mine. Strain CEB3 is one of the very few reported acidophilic sulfate-reducing isolates. The 5.04-Mb draft genome harbors 5,069 predicted protein-encoding and 66 RNA genes.**

Received 27 June 2015 Accepted 30 June 2015 Published 6 August 2015

**Citation** Petzsch P, Poehlein A, Johnson DB, Daniel R, Schlömann M, Mühling M. 2015. Genome sequence of the acidophilic sulfate-reducing *Peptococcaceae* strain CEB3. *Genome Announc* 3(4):e00886-15. doi:10.1128/genomeA.00886-15.

**Copyright** © 2015 Petzsch et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](https://creativecommons.org/licenses/by/3.0/).

Address correspondence to Martin Mühling, [martin.muehling@ioez.tu-freiberg.de](mailto:martin.muehling@ioez.tu-freiberg.de).

Acidophilic sulfate-reducing bacteria are of particular interest for the treatment of sulfate- and metal-enriched acid mine drainage (AMD) waters since sulfidogenesis is a proton-consuming process (thereby promoting amelioration of mine water acidity) and the hydrogen sulfide produced can be used to selectively precipitate transition metals present in AMD (1). Strain CEB3, which, based on its 16S rRNA gene sequence, is likely to be related to a new genus within the *Peptococcaceae* (*Firmicutes*), is a spore-forming, motile, obligately anaerobic sulfidogen that was isolated as one of the dominating microbial strains present in a sulfidogenic bioreactor operated at pH 2.4 ( $\pm 0.1$ ) for the selective recovery of metals (1). The inoculum for this bioreactor originated from an anaerobic streamer mat growing in an acidic (pH 2.5) stream draining an abandoned copper mine in southwest Spain (1).

Chromosomal DNA of *Peptococcaceae* strain CEB3 was isolated with the MasterPure complete DNA purification kit (Epicentre) and used to generate 454-shotgun and Illumina-shotgun libraries according to the manufacturers' protocols. The libraries were sequenced using a 454 GS-FLX system (Titanium Chemistry, Roche Life Science) and the Genome Analyzer IIx (2 $\times$ 112 bp paired-end sequencing; Illumina). Sequencing resulted in coverages of 8.03 and 78.26, respectively. The assembly of the genome sequence was performed with the Roche Newbler assembler v2.9 and Mira v3.4 (2) and resulted in 64 contigs.

The draft genome of strain CEB3 comprises a single chromosome of 5.04 Mb with an overall G+C content of 52.58%. Prodigal (3) was used for automated gene prediction, and identification of rRNA and tRNA genes was carried out using RNAmmer (4) and tRNAscan (5), respectively. The IMG-ER (Integrated Microbial Genomes-Expert Review) system (6) was used for automated annotation, which was subsequently manually curated using the Swiss-Prot, TrEMBL, and InterPro databases (7). The draft genome harbored 9 rRNA genes, 57 tRNA genes, and 3,898 predicted protein-encoding genes with function prediction and 1,171 genes coding for hypothetical proteins.

Analyses of the protein-coding genes revealed that strain CEB3 appears to be a metabolically versatile acidophile. For instance, strain CEB3 uses both phosphate and phosphonate (via C-P lyase activity) as a P source. Similarly, it can utilize several nitrogen sources for biomass production (nitrate, ammonium, and dinitrogen gas). The assimilated nitrogen is predicted to be stored in the form of cyanophycin, which is known to play a role in heavy-metal resistance (8).

Strain CEB3 appears to be able to utilize sulfate as a source of sulfur and as an electron acceptor for dissimilatory reduction. Utilization of trithionate and tetrathionate or thiosulfate may provide alternative avenues for sulfur acquisition and respiration, though only one subunit of the thiosulfate reductase has been detected in the genome sequence. Strain CEB3 may assimilate carbon either through CO<sub>2</sub> fixation via the reductive acetyl-CoA pathway or take up organic carbon molecules via monosaccharide and disaccharide transporters.

**Nucleotide sequence accession numbers.** The results from this genome sequencing project have been deposited at GenBank under the accession number [LDXJ000000000](https://www.ncbi.nlm.nih.gov/nuccore/LDXJ000000000). The version described in this paper is version [LDXJ000000000.1](https://www.ncbi.nlm.nih.gov/nuccore/LDXJ000000000).

## ACKNOWLEDGMENTS

The sequence analysis is part of GETGEOWEB (project 100101363), which was funded by the European Social Fund (ESF).

We thank Frauke-Dorothee Meyer, Kathleen Gollnow, and Sarah Hofmann for technical support.

## REFERENCES

1. Nancucheo I, Johnson DB. 2012. Selective removal of transition metals from acidic mine waters by novel consortia of acidophilic sulfidogenic bacteria. *Microb Biotechnol* 5:34–44. <http://dx.doi.org/10.1111/j.1751-7915.2011.00285.x>.
2. Chevreaux B, Wetter T, Suhai S. 1999. Genome sequence assembly using trace signals and additional sequence information, p 45–56. In Wingender E (ed), *Computer science and biology: proceedings of the German Conference on Bioinformatics (GCB) 1999 Hannover, Germany*. GBF-Braunschweig, Department of Bioinformatics, Braunschweig, Germany.

3. Hyatt D, Chen G, Locascio PF, Land ML, Larimer FW, Hauser LJ. 2010. Prodigal: prokaryotic gene recognition and translation initiation site identification. *BMC Bioinformatics* 11:119. <http://dx.doi.org/10.1186/1471-2105-11-119>.
4. Lagesen K, Hallin P, Rødland EA, Stærfeldt H-H, Rognes T, Ussery DW. 2007. RNAmmer: consistent and rapid annotation of ribosomal RNA genes. *Nucleic Acids Res* 35:3100–3108. <http://dx.doi.org/10.1093/nar/gkm160>.
5. Lowe TM, Eddy SR. 1997. tRNAscan-SE: a program for improved detection of transfer RNA genes in genomic sequence. *Nucleic Acids Res* 25: 955–964. <http://dx.doi.org/10.1093/nar/25.5.0955>.
6. Markowitz VM, Chen I-MA, Palaniappan K, Chu K, Szeto E, Pillay M, Ratner A, Huang J, Woyke T, Huntemann M, Anderson I, Billis K, Varghese N, Mavromatis K, Pati A, Ivanova NN, Kyrpides NC. 2014. IMG 4 version of the integrated microbial genomes comparative analysis system. *Nucleic Acids Res* 42:D560–D567. <http://dx.doi.org/10.1093/nar/gkt963>.
7. Zdobnov EM, Apweiler R. 2001. InterProScan—an integration platform for the signature-recognition methods in InterPro. *Bioinformatics* 17: 847–848. <http://dx.doi.org/10.1093/bioinformatics/17.9.847>.
8. Wood JM. 1983. Selected biochemical reactions of environmental significance. *Chem Scr* 21:155–160.