

**Genetic variation in a small bivalve along a retreating glacier fjord, King George Island, Antarctica.**

Muñoz-Ramirez, Carlos; Beltrán-Concha, Maribel ; Pérez-Araneda, Karla; Sands, Chester; Barnes, David; Roman-Gonzalez, Alejandro; De Lecea, Ander; Retallick, Katherine; Van Landeghem, Katrien; Sheen, Katy; Gonnelli, Kelly; Scourse, James D.; Bascur, Miguel; Brante, Antonio

Revista de Biología Marina y Oceanografía

DOI:

<https://doi.org/10.22370/rbmo.2021.56.2.3059>

Published: 31/12/2021

Peer reviewed version

[Cyswllt i'r cyhoeddiad / Link to publication](#)

Dyfyniad o'r fersiwn a gyhoeddwyd / Citation for published version (APA):

Muñoz-Ramirez, C., Beltrán-Concha, M., Pérez-Araneda, K., Sands, C., Barnes, D., Roman-Gonzalez, A., De Lecea, A., Retallick, K., Van Landeghem, K., Sheen, K., Gonnelli, K., Scourse, J. D., Bascur, M., & Brante, A. (2021). Genetic variation in a small bivalve along a retreating glacier fjord, King George Island, Antarctica. *Revista de Biología Marina y Oceanografía*, 56(2), 151-156. <https://doi.org/10.22370/rbmo.2021.56.2.3059>

Hawliau Cyffredinol / General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Research Note:

Genetic variation in a small bivalve along a retreating glacier fjord, King George Island,

Antarctica

Variación genética en un pequeño bivalvo a lo largo de un fiordo con retroceso glaciar, Isla Rey Jorge,

Antártica

Short title: Genetic diversity along an Antarctic fjord

Carlos P. Muñoz-Ramírez^{1,2,3,4}, Maribel Beltrán-Concha², Karla Pérez-Araneda², Chester Sands⁴, David K A Barnes⁴, Alejandro Roman-Gonzalez⁵, Ander De Lecea⁶, Katherine Retallick⁷, Katrien Van Landeghem⁷, Katy Sheen⁵, Kelly Gonnelli², James Scourse⁵, Miguel Bascur², Antonio Brante^{2,3,4}

¹ Instituto de Entomología, Universidad Metropolitana de Ciencias de la Educación, Santiago, Chile.

² Facultad de Ciencias, Universidad Católica de la Santísima Concepción, Concepción, Chile.

³ Centro de Investigación en Biodiversidad y Ambientes Sustentables (CIBAS), Universidad Católica de la Santísima Concepción, Concepción, Chile.

⁴ British Antarctic Survey, NERC, Madingley Road, Cambridge, CB3 0ET, UK

⁵ Falmouth Campus, University of Exeter, Penryn, UK

⁶ South Atlantic Environmental Research Institute, Stanley, Falkland Islands

⁷ School of Ocean Sciences, Bangor University, Menai Bridge, United Kingdom

⁸ Corresponding author: carlos.munoz@umce.cl

Genetic diversity along an Antarctic fjord

Abstract.- Climate change is strongly influencing regions of Antarctica but the consequences on microevolutionary processes have been little studied. Patterns of population genetic diversity were analysed in the Antarctic bivalve *Nuculana inaequisculpta* (Protobranchia: Nuculanidae) from a fjord with 70 years of documented climate-forced glacier retreat. Thirty-nine individuals from five sites at different distances from the glacier terminus were collected, and the COI gene was sequenced from each individual. No statistically significant genetic differentiation was found between sites nor a significant correlation between the proximity of glaciers and genetic diversity, suggesting a high dispersal capability and therefore, a planktonic larval stage for this species.

Key words: COI, climate change, dispersal capability, Western Antarctic Peninsula.

Introduction

West Antarctica is a hotspot for physical Climate Change, and ~90% of all glaciers along the West Antarctic Peninsula (WAP) are rapidly retreating (Cook *et al.* 2016). Marine species, especially those in the shallows, are subject to drastic environmental changes, including those which may negatively impact species survival (*e.g.*, increased ice scour) or favor colonization of new available habitats exposed by retreating glaciers. Several studies have documented ecological impacts of glacial retreat and melting on marine communities (*e.g.*, Moon *et al.* 2015, Sahade *et al.* 2015, Barnes *et al.* 2020, Bascur *et al.* 2020). However, it is less well known how such perturbations may drive microevolutionary responses of marine species at population genetic scales. There is a lack of studies examining this question in Antarctic ecosystems. Studies testing patterns of genetic diversity in relation to climate-change forced distributional shifts in other areas have found mixed responses. For instance, simulation studies have found that the edge of a shifting range, under climate change, has low levels of neutral genetic diversity (Cobben *et al.* 2011, Arenas *et al.* 2012). In contrast, some empirical data, for example of *Larix decidua*, have shown high genetic diversity at the leading edge of an expanding population following a retreating glacier in the Swiss Alps (Pluess 2011).

Genetic diversity along an Antarctic fjord

Genetic population responses of a species after perturbation is also related to aspects of its biology, such as its dispersal capability. For instance, a species with low dispersal capability might be expected to present a stronger pattern of genetic diversity (*e.g.*, decreasing genetic diversity along an expansion transect) than a species with high dispersal capability. This is because migration between demes is not fast enough to dilute the pattern (Klopfstein *et al.* 2006). In contrast, high dispersal capability would be likely to drive high gene flow between populations and therefore, rapid homogenization of any incipient genetic differentiation (Slatkin 1987).

In Antarctica, it has been argued that climatic conditions have filtered for life-history strategies that reduce or lack larval stages in benthic organisms, especially those which are planktotrophic (Pearse *et al.* 2009). However, planktonic larvae have been described for many Antarctic benthic species (*e.g.*, Stanwell-Smith *et al.* 1997, 1999), and their actual number may have been greatly underestimated due to the lack of detailed morphological and molecular studies on planktonic larvae (Thatje 2012). Studies around Antarctica have shown that developmental type influences patterns of genetic differentiation across barriers. For example, Poulin *et al.* (2014) found that species lacking planktonic larvae were more divergent across the Antarctic Circumpolar Current than species with planktonic larvae. Similarly, Muñoz-Ramirez *et al.* (2020) found that *Aequiyoldia eightsii*, a species with a lecitotrophic larvae, showed genetic patterns that were congruent with ocean current dispersal. Dispersal capability of species likely influences colonization rates and genetic patterns on ecological time scales. However, no studies have examined the genetic consequences at the temporal scale of current climate change in Antarctica.

Nuculana inaequisculpta (Lamy, 1906) (Fig. 1D) is a small bivalve species that belongs to the Protobranchia subclass, a taxonomic group that contains the most ancient bivalves in the world (Villarroel & Stuardo 1998, Yonge 1939). It is an abundant infaunal species, patchy distributed in the South Shetland Islands and along the Antarctic Peninsula (Cattaneo-Vietti *et al.* 2000, Lovell & Trego 2003, Gordillo *et al.* 2017). This species lives in benthic, muddy marine habitats and feeds mainly on organic deposits in sediments (Zardus 2002). It occurs as deep as ca. 800 m, and range in size from 2.5-16 mm in shell length

Genetic diversity along an Antarctic fjord

(Cattaneo-Vietti *et al.* 2000, Gordillo *et al.* 2017). In general, protobranchia species have two separate sexes, with a similar (1:1) sex ratio, high fecundity, and lecithotrophic larval development with a pericalima larva (Zardus 2002). Recent bioenergetic analyses have found they are sensitive to food availability and exhibit lower fitness in recently colonized areas, perhaps due to lower food supply and higher perturbations near the glacier fronts (Bascur *et al.* 2020).

In this study, sequences of the Cytochrome C Oxidase Subunit I (COI) gene obtained from *Nuculana inaequisculpta* were analysed to investigate patterns of genetic diversity along a fjord with a retreating glacier. A strong pattern of decreasing genetic diversity towards the glacier front could be considered indicative of low dispersal capability as expected under a direct development life-history strategy or benthic larvae, and would suggest impacts of current climate change on population diversity. In contrast, a lack of correlation would indicate high dispersal capability, as expected for planktonic larval stages, and highlight the importance of high dispersion as a mechanism of resilience against climate-change induced perturbations.

Materials and methods

Sampling sites and molecular data collection

Thirty-nine individuals of *N. inaequisculpta* were collected from five sites (6-8 individuals per site) at Marian Cove, King George Island, Antarctica, a fjord with a significant glacier retreat documented since 1950 (Park *et al.* 1998, Cook *et al.* 2016, Barnes *et al.* 2020) (Fig. 1, Table 1). Genomic DNA was extracted with the Qiagen DNeasy Blood & Tissue kit, following the manufacturer's protocol. A 688 bp fragment of the cytochrome c oxidase subunit I (COI) gene was amplified using universal primers from Folmer *et al.* (1994) with final concentrations for PCR components per 25 μ L reaction as follows: 25 ng template DNA, 0.25 μ M of each primer, 0.625 units of GoTaq DNA polymerase (Promega, Madison, WI, USA), 0.1 mM of each dNTP, 2.5 μ L of 10 reaction buffer and 2.5 mM MgCl₂. Amplification parameters were as follows: 95 °C for 2 min followed by 35 cycles of 95 °C for 30 s, 48 °C for 30 s,

Genetic diversity along an Antarctic fjord

72 °C for 90 s, and 72 °C for 7 min. Purification and sequencing were conducted at MACROGEN Inc. (South Korea). Chromatograms were edited in CodonCode Aligner 8.0.2 (Dedham, MA, USA). Sequences were imported to BioEdit 7.0.5.2 (Hall *et al.* 2011), aligned using the Clustal W algorithm, available within BioEdit, and checked by eye. All sequences were deposited in GenBank (accession numbers MT510924-MT510962).

Genetic differentiation and diversity

To estimate levels of population differentiation, F_{st} values between sites were calculated using the R-packages *ADEGENET* (Jombart 2008), and *HIERFSTAT* (Goudet 2005). Genetic diversity indexes, haplotype richness (S), haplotype diversity (Hd) and nucleotide diversity (π) for each site and for the entire fjord were estimated using *PEGAS* (Paradis 2010). To visually display relationships between haplotypes, a haplotype network was estimated using the median-joining network algorithm of Bandelt *et al.* (1999) implemented in the program POPART (Leigh & Bryant 2015).

Based on results of low genetic diversity for the closest site to the glacier terminus, a test was conducted to estimate the probability of obtaining the lowest π value for that site under a model of panmixia and chance. For this, a panmictic population was assumed and samples were randomly sampled to build a frequency distribution of 1,000 simulated π values, selecting six sample at each iteration to calculate π . Subsequently, the empirical π value was compared to the frequency distribution of random π values. Support for a significantly low value is given if the empirical π value is lower than the 95% of the simulated π values.

Results and discussion

A total of eighteen haplotypes were present among the thirty-nine individuals of *N. inaequisculpta* at Marian Cove. The average number of nucleotide differences between samples was 0.402. Genetic structure between populations was low. Genetic differentiation was greater between sites MC2 and MC5 ($F_{st}= 0.1057$), whilst the lowest differentiation values were between MC3 versus MC1 and MC3 versus

Genetic diversity along an Antarctic fjord

MC4 (both $F_{st}= 0.0279$). However, none of the F_{st} values between sites were statistically significant at the 0.05 level. A similar lack of structure was observed in the haplotype network (Fig. 2A). A central, high-frequency haplotype was present at all sites at relatively high frequency. Most of the remaining haplotypes were derived from that, with low frequency and separated by only few mutational steps. This star-like topology is typical of populations that have undergone a recent bottleneck and subsequent population expansion (Slatkin & Hudson 1991). Living space on the Antarctic shelf has undergone cyclic expansions and contractions due to the advance and retreat of the Antarctic Ice Shield during glacial cycles (Augustin *et al.* 2004). These major historical events can explain the signal of demographic expansion, which could have initiated after the Last Glacial Maximum. However, this could also be explained, at least in part, due to recent colonization of the fjord. We lack the data to distinguish these hypotheses, but larger geographic sampling, including populations from the entire distribution of the species could help resolve this question in a future study.

Haplotype diversity (Hd) across sites varied from 0.643 (in MC1) to 1 (in MC2) (Fig. 2B). There was no clear tendency for Hd to increase or decrease with glacier proximity, and the lowest values were found at the opposite sites of the transect ($Hd_{MC1}= 0.643$; $Hd_{MC5}= 0.8$). Nucleotide diversity tended to increase with proximity to the glacier from sites MC1 ($\pi_{MC1}= 0.0032$) to MC4 ($\pi_{MC4}= 0.0053$), but it strongly decreased at site MC5, presenting the lowest value among all sites ($\pi_{MC5}= 0.0021$) (Fig. 2C). No significant correlation was found between geographic proximity to the glacier terminus and genetic diversity for both Hd ($r= -0.307$; P -value= 0.6157) and π ($r= 0.06$; P -value = 0.9216), suggesting high dispersal capability for *N. inaequisculpta*, and a planktonic larval stage. However, the lowest π value found at MC5 (the closest site to the glacier terminus) suggest that strong perturbations (e.g. ice scouring, glacier meltwater discharge, high sedimentation) near the glacier terminus may impact genetic diversity. To test if this low value of genetic diversity could imply a deterministic force (e.g., impact of perturbations on genetic diversity), the probability of having the lowest π value at MC5 was estimated under population panmixia and chance. The probability of obtaining a π value equal or lower than the

Genetic diversity along an Antarctic fjord

observed value ($\pi=0.00201$) only by chance was $P=0.1897$, and therefore, not significant. This indicates that there is not enough evidence that the lowest π value at MC5 is due to deterministic forces (e.g. recent colonization of the area or perturbations). However, evidence of low fitness in *Nuculana* specimens from near the glacier terminus has been recently provided by Bascur *et al.* (2020), who found, by conducting bioenergetic analyses, that individuals from MC5 had the lowest energy content in their tissues. These congruence between bioenergetic and genetic results strongly suggest an impact of glacier-derived perturbations on the species's performance.

The pattern of low genetic diversity towards MC1 (the farthest to the glacier terminus) is puzzling, and may suggest a more complex scenario including low competitive abilities. If *N. inaequisculpta* is a species that exploits its dispersal capability to reach recently emptied habitats, but is a generally weak competitor, it is possible that populations that are present in longer-established communities (e.g., those which are further away from the glacier terminus) may be at disadvantage, and therefore maintain lower population sizes. This could explain low genetic diversity at MC1 and the progressively higher diversity towards MC4 (although not significant). Indeed, an additional site (MC6) furthest from the glacier terminus (not shown on the map) was also sampled, but it yielded no individuals of *N. inaequisculpta*, so it was not included in this study. This MC6 site was rich and diverse in other species though (data not shown), suggesting that the community was old and long established. This seems to support the idea that *N. inaequisculpta* could be a weak competitor given its absence in this site. However, it is recognized that this is speculative, and as a tentative hypothesis it will require further study. Therefore, it is proposed that the comparative genetic study of multiple species with different life history strategies (e.g., with and without planktonic larval stage) and competitive abilities will allow testing this hypothesis in future studies.

There are clear limitations to the inferential power of the marker used here (one single mitochondrial gene). Nevertheless, the results provided in this study are a first step towards a more comprehensive

Genetic diversity along an Antarctic fjord

understanding of the patterns and processes governing the genetic diversity of this bivalve species, which can be largely improved in future studies by the adoption of more powerful genomic markers such as RADSeq-derived single-nucleotide polymorphisms.

Acknowledgements

We thank INACH and the research grants CONICYT-NERC project ICEBERGs (PII20150078) and CONICYT-FONDECYT Postdoctorado (3180331) and especially the crew of the RRS James Clark Ross for their great work and effort in the pursue of a successful cruise. We also appreciate the helpful feedback provided by two anonymous reviewers that greatly improved the manuscript.

Literature cited

- Augustin L, C Barbante, PR Barnes, JM Barnola, M Bigler, E Castellano, O Cattani, J Chappellaz, D Dahl-Jensen, B Delmonte & G Dreyfus, G Durand, S Falourd, H Fischer, J Flückiger, ME Hansson, P Huybrechts, G Jugie, SJ Johnsen, J Jouzel, P Kaufmann, J Kipfstuhl, F Lambert, V Y Lipenkov, GC Littot, A Longinelli, R Lorrain, V Maggi, V Masson-Delmotte, H Miller, R Mulvaney, J Oerlemans, H Oerter, G Orombelli, F Parrenin, DA Peel, J-R Petit, D Raynaud, C Ritz, U Ruth, J Schwander, U Siegenthaler, R Souchez, B Stauffer, JP Steffensen, B Stenni, TF Stocker, IE Tabacco, R Udisti, RSW Van De Wal, M Van Den Broeke, J Weiss, F Wilhelms, J-G Winther, EW Wolff, M Zucchelli & EPICA community members. 2004.** Eight glacial cycles from an Antarctic ice core. *Nature* 429: 623-628.
- Arenas M, N Ray, M Currat & L Excoffier. 2012.** Consequences of range contractions and range shifts on molecular diversity. *Molecular Biology and Evolution* 29: 207-218.
- Bandelt HJ, P Forster & A Röhl. 1999.** Median-joining networks for inferring intraspecific phylogenies. *Molecular Biology and Evolution* 16: 37-48.

- Barnes DKA, CJ Sands, A Cook, F Howard, A Roman-Gonzalez, CP Muñoz-Ramirez, K Retallick, J Scourse, K Van Landegh & N Zserschke. 2020.** Blue carbon gains from glacial retreat along Antarctic fjords: What should we expect? *Global Change Biology* 26: 2750-2755.
- Bascur M, CP Muñoz-Ramírez, A Román-González, K Sheen, DKA Barnes, CJ Sands, A Brante & A Urzúa. 2020.** The influence of glacial melt and retreat on the nutritional condition of the bivalve *Nuculana inaequisculpta* (Protobranchia: Nuculanidae) in the West Antarctic Peninsula. *PLoS ONE* 15(5): e0233513. <<https://doi.org/10.1371/journal.pone.0233513>>
- Cattaneo-Vietti R, M Chiantore, S Schiaparelli & G Albertelli. 2000.** Shallow- and deep-water mollusc distribution at Terra Nova Bay (Ross Sea, Antarctica). *Polar Biology* 23: 173-182.
- Cobben MMP, J Verboom J, PFM Opdam, RF Hoekstra, R Jochem, P Arens & MJM Smulders. 2011.** Projected climate change causes loss and redistribution of genetic diversity in a model metapopulation of a medium good disperser. *Ecography* 34: 920-932.
- Cook AJ, PR Holland, MP Meredith, T Murray, A Luckman & DG Vaughan. 2016.** Ocean forcing of glacier retreat in the western Antarctic Peninsula. *Science* 353: 283-286.
- Folmer O, M Black, W Hoeh & R Vrijenhoek. 1994.** DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Molecular Marine Biology and Biotechnology* 3: 294-299.
- Goudet J. 2005.** hierfstat, a package for r to compute and test hierarchical F-statistics. *Molecular Ecology Notes* 5:184-186.
- Gordillo S, M Malvé & G Moran. 2017.** Benthic mollusc assemblages in West Antarctica: taxa composition and ecological insights. *Marine and Freshwater Research* 68: 2095-2105.
- Hall T. 2011.** BioEdit: An important software for molecular biology. *GERF Bulletin of Biosciences* 2: 60-61.
- Jombart T. 2008.** adegenet: a r package for the multivariate analysis of genetic markers. *Bioinformatics* 24: 1403-1405.

Genetic diversity along an Antarctic fjord

- Klopfstein S, M Currat & L Excoffier. 2006.** The fate of mutations surfing on the wave of a range expansion. *Molecular Biology and Evolution* 23: 482–490.
- Lovell LL & KD Trego. 2003.** The epibenthic megafaunal and benthic infaunal invertebrates of Port Foster, Deception Island (South Shetland Islands, Antarctica). *Deep-Sea Research Part II* 50: 1799-1819.
- Leigh JW & D Bryant. 2015.** POPART: Full-feature software for haplotype network construction. *Methods in Ecology and Evolution* 6: 1110-1116.
- Moon HW, WMR Wan Hussin, HC Kim & IY Ahn. 2015.** The impacts of climate change on Antarctic nearshore mega-epifaunal benthic assemblages in a glacial fjord on King George Island: Responses and implications. *Ecological Indicators* 57: 280-292.
- Muñoz-Ramírez CP, DKA Barnes, L Cárdenas, M Meredith, S Morley, A Roman-Gonzalez, C Sands, J Scourse & A Brante. 2020.** Gene flow in the Antarctic bivalve *Aequiyoldia eightsii* (Jay, 1839) suggests a role for the Antarctic Peninsula Coastal Current in larval dispersal. *Royal Society Open Science* 7: 200603. <<https://doi.org/10.1098/rsos.200603>>
- Paradis E. 2010.** Pegas: an R package for population genetics with an integrated-modular approach. *Bioinformatics* 26: 419-420.
- Park BK, SK Chang, HI Yoon & H Chung. 1998.** Recent retreat of ice cliffs, King George Island, South Shetland Islands, Antarctic Peninsula. *Annals of Glaciology* 27: 633-635.
- Pearse JS, R Mooi, SJ Lockhart & A Brandt. 2009.** Brooding and species diversity in the Southern Ocean: selection for brooder or speciation within brooding clades? In: Krupnik I, MA Lang, SE Miller, editors. *Smithsonian Poles Symposium*. Washington (DC): Smithsonian Institute p. 181–96.
- Pluess AR. 2011.** Pursuing glacier retreat: genetic structure of a rapidly expanding *Larix decidua* population. *Molecular Ecology* 20: 473-485.

- Poulin E, C González-Wevar, A Díaz, K Gérard & M Hüne. 2014.** Divergence between Antarctic and South American marine invertebrates: What molecular biology tells us about Scotia Arc geodynamics and the intensification of the Antarctic Circumpolar Current. *Global and Planetary Change* 123: 392-399.
- Sahade R, C Lager, L Torre, F Momo, P Monien, I Schloss, DK Barnes, N Servetto, S Tarantelli, M Tatián & N Zamboni. 2015.** Climate change and glacier retreat drive shifts in an Antarctic benthic ecosystem. *Science Advances* 1(10), e1500050. <DOI: 10.1126/sciadv.1500050>
- Slatkin M. 1987.** Gene flow and the geographic structure of natural populations. *Science* 236: 787-792.
- Slatkin M & RR Hudson. 1991.** Pairwise comparisons of mitochondrial DNA sequences in stable and exponentially growing populations. *Genetics* 129: 555-562.
- Stanwell-Smith D, A Hood & L Peck. 1997.** A field guide to the pelagic invertebrate larvae of the maritime Antarctic, 152 pp. British Antarctic Survey, Cambridge.
- Stanwell-Smith D, LS Peck, A Clarke, AWA Murray & CD Todd. 1999.** The distribution, abundance, and seasonality of pelagic marine invertebrate larvae in the maritime Antarctic. *Philosophical Transactions of the Royal Society of London B, Biological Sciences* 354: 471-484.
- Thatje S. 2012.** Effects of capability for dispersal on the evolution of diversity in Antarctic benthos. *Integrative and Comparative Biology* 52: 470-482.
- Villarroel M & J Stuardo. 1998.** Protobranchia (mollusca: Bivalvia) of recent distribution in Chile and some fossils. *Malacologia* 40: 113-229.
- Yonge CM. 1939.** The Protobranchiate Mollusca: A functional interpretation of their structure and evolution. *Philosophical Transactions of the Royal Society A* 230: 79-148.
- Zardus JD. 2002.** Protobranch bivalves. *Advances in Marine Biology* 42: 1-65.

Table 1. Sampling details, genetic diversity and genetic differentiation (F_{st}) between sampling sites based on COI sequences for the Antarctic bivalve *Nuculana inaequisculpta*. F_{st} values were not

Genetic diversity along an Antarctic fjord

statistically significant based on a 999-permutation test. N: Number of individuals. Hd: Haplotype diversity. π : Nucleotide diversity / Detalles de muestreo, diversidad genética y diferenciación genética (F_{st}) entre sitios de muestreo basado en secuencias del gen COI para el bivalvo antártico *Nuculana inaequisculpta*. Los valores de F_{st} no fueron significativos basado en un test de 999 permutaciones. N: Número de individuos. Hd: Diversidad Haplotípica. π : Diversidad nucleotídica

Sites	Latitude	Longitude	N	Hd	π
MC1	-62.2176	-58.7292	8	0.6429	0.0032
MC2	-62.21273	-58.7689	8	1.0000	0.0036
MC3	-62.20815	-58.7479	8	0.9286	0.0040
MC4	-62.20588	-58.7399	9	0.9722	0.0053
MC5	-62.20373	-58.7314	6	0.8000	0.0021
F_{st} between populations	MC1	MC2	MC3	MC4	MC5
MC1	0.0000	0.0819	0.0279	0.0425	0.0848
MC2	0.0819	0.0000	0.0630	0.0388	0.1057
MC3	0.0279	0.0630	0.0000	0.0279	0.0583
MC4	0.0425	0.0388	0.0279	0.0000	0.0758
MC5	0.0848	0.1057	0.0583	0.0758	0.0000

N: Number of individuals. Hd: Haplotype diversity. π : Nucleotide diversity

Genetic diversity along an Antarctic fjord




Figure 1: Sample sites for *Nuculana inaequisculpta* at Marian Cove, King George Island, South Shetlands Islands, Antarctica. A) Antarctic Peninsula. B) King George Island. C) Marian Cove and sampling sites. D) Specimen of *Nuculana inaequisculpta*. AP: Antarctic Peninsula. Dashed lines across the fjord represent the location of the glacier front in different years / Sitios de muestreo para *Nuculana inaequisculpta* en Marian Cove, Isla Rey Jorge, Shetlands del Sur, Antártica. A) Península Antártica. B) Isla Rey Jorge. C) Marian Cove con los sitios de muestreo. D) individuo de *N. inaequisculpta*. AP: Península Antártica. Líneas segmentadas indican la posición del frente glaciar en diferentes años




Figure 2. A) Haplotype network, B) haplotype diversity (H_d), and C) nucleotide diversity (π) for *Nuculana inaequisculpta* based on COI sequences. Black dots and dashes in the haplotype network represent nodes and mutational steps, respectively. / A) Red de haplotipos, B) diversidad haplotípica (H_d) y C) diversidad nucleotídica (π) para *Nuculana inaequisculpta* basada en secuencias COI. Los círculos negros y las líneas cortas perpendiculares sobre las líneas que conectan haplotipos corresponden a nodos y pasos mutacionales, respectivamente.