

Investigating awareness, fear and control associated with norovirus and other pathogens and pollutants using best-worst scaling

Farkas, Kata; Green, Emma; Rigby, Dan; Cross, Paul; Tyrrel, Sean; Malham, Shelagh; Jones, Davey L.

Scientific Reports

DOI: 10.1038/s41598-021-90704-7

Published: 27/05/2021

Publisher's PDF, also known as Version of record

Cyswllt i'r cyhoeddiad / Link to publication

Dyfyniad o'r fersiwn a gyhoeddwyd / Citation for published version (APA): Farkas, K., Green, E., Rigby, D., Cross, P., Tyrrel, S., Malham, S., & Jones, D. L. (2021). Investigating awareness, fear and control associated with norovirus and other pathogens and pollutants using best–worst scaling. *Scientific Reports*, *11*(1), Article 11194. https://doi.org/10.1038/s41598-021-90704-7

Hawliau Cyffredinol / General rights Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

• Users may download and print one copy of any publication from the public portal for the purpose of private study or research.

- You may not further distribute the material or use it for any profit-making activity or commercial gain
 You may freely distribute the URL identifying the publication in the public portal ?

Take down policy If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

scientific reports

OPEN



Investigating awareness, fear and control associated with norovirus and other pathogens and pollutants using best–worst scaling

Kata Farkas^{1,2,6,7^{\infty}}, Emma Green^{1,7}, Dan Rigby³, Paul Cross¹, Sean Tyrrel⁴, Shelagh K. Malham² & David L. Jones^{1,5}

Pollutants found in the water and air environment represent an ever-growing threat to human health. Contact with some air-, water- and foodborne pathogens (e.g. norovirus) results in gastrointestinal diseases and outbreaks. For future risk mitigation, we aimed to measure people's awareness of waterborne and foodborne norovirus relative to other environment-associated pollutants (e.g. pesticides, bioaerosols, antibiotic resistant bacteria) and well-known risks (e.g. diabetes, dementia, terrorist attack). We used an online survey, which included a best-worst scaling component to elicit personal levels of control and fear prompted by norovirus relative to 15 other risks. There was a negative correlation between levels of fear vs. control for all 16 measured risks. Perceived infection control levels were higher amongst women compared to men and correlated with age and the level of qualification in both groups. Participants who had sought advice regarding the symptoms caused by norovirus appeared to have more control over the risks. Norovirus is associated with high levels of fear, however, the levels of control over it is low compared to other foodborne illnesses, e.g. *Salmonella*. Addressing this deficit in the public's understanding of how to control exposure to the pathogen in an important health need.

Increased agricultural and industrial activities alongside progressive urbanization have caused the emergence and spread of a wide range of pollutants in the environment, leading to global health threats. For example, air pollutants (including bioaerosols) and pesticides are leading to an increasing number of skin and respiratory syndromes¹⁻⁴. The overuse of antibiotics in healthcare and agriculture has led to a high prevalence of antibiotic resistant bacteria ('superbugs') resulting in an increasing number of untreatable bacterial infections⁵.

The number of infections and outbreaks associated with gastrointestinal symptoms are also on the rise⁶⁻⁹. Among these pathogens, norovirus (often known as 'winter vomiting bug') is the most common cause of nonbacterial gastroenteritis globally^{10,11}. Norovirus is highly infectious, spreads rapidly and hence is responsible for outbreaks leading to the temporary closure of public places (e.g. schools, restaurants and hospitals). The symptoms, including diarrhea, vomiting, nausea, fever and abdominal pain, are usually mild and resolve in 2–5 days, however, in some cases, the infection can be life threatening¹⁰. The estimated annual number of cases is 685 million with approximately 200,000 deaths worldwide^{10,12} with a total of US\$4.2 billion in direct health system costs and US\$60.3 billion in societal costs¹³.

The main route of transmission is direct (person-to-person) contact, however, the number of cases associated with consumption of contaminated food and water is increasing^{14,15}. Norovirus can be found at high concentrations in the stools of infected individuals even weeks after the symptoms are resolved¹⁶. As the virus is extremely resistant to traditional wastewater treatment procedures¹⁷⁻¹⁹, significant loads are discharged into the

¹School of Natural Sciences, Bangor University, Deiniol Road, Bangor, Gwynedd LL57 2UW, UK. ²School of Ocean Sciences, Bangor University, Menai Bridge, Anglesey LL53 5AB, UK. ³Department of Economics, University of Manchester, Oxford Road, Manchester M13 9PL, UK. ⁴School of Water, Energy and Environment, Cranfield University, Cranfield MK43 0AL, UK. ⁵UWA Oceans Institute, The University of Western Australia, Perth, WA 6009, Australia. ⁶Marine Centre Wales, Menai Bridge, Anglesey LL59 5AB, UK. ⁷These authors contributed equally: Kata Farkas and Emma Green. [⊠]email: fkata211@gmail.com

environment contaminating recreational and drinking water bodies and irrigation systems^{20–22}. Furthermore, norovirus is transported in environmental waters over long distances, accumulating in sediment²³ and taken up by filter feeder aquatic animals, such as shellfish, harvested for human consumption^{24,25}. As little is known about the transport patterns and behavior of norovirus, the contamination and health risks associated with water and shellfish are hard to predict²⁶.

As the virus is highly contagious and no vaccination is available, prophylactic measures are critical to the control of disease spread^{27,28}. For instance, in some developed countries, information on the risks associated with norovirus and other foodborne agents are disseminated in public places (e.g. healthcare units, pharmacies, restaurant menus) and on governmental websites. Research has been done to understand the persistence and associated risks of norovirus and other pollutants in the environment^{26,29-33}. Several studies have attempted to evaluate the perception of food safety and microbial contamination in general³⁴⁻³⁷. However, little is known about people's awareness and perception of norovirus and its transmission³⁸⁻⁴⁰ despite the importance of such knowledge in shaping the behaviors which may reduce the spread of the disease.

This study investigated the public's awareness and perceptions of norovirus and other environment-associated health risks. This was undertaken via analysis of data from a UK national survey. Within this survey, a choice-based approach (Best Worst Scaling; BWS)³⁵ was used to determine the levels of fear and control people associated with norovirus in comparison to a set of other risks. Specifically, we investigated:

- I. the level of public awareness of norovirus
- II. the level of fear associated with norovirus compared to other hazards (both environment-related and more general) with established perception
- III. the level of control people believed they had over norovirus compared to other hazards (both environment-related and more general) with established perception
- IV. how these perceptions of control and fear varied over socio-demographic characteristics and the implications of these variations for efforts to increase awareness of norovirus and reduce the disease burden.

Results

The survey questions and results from respondents are available at the Environmental Information Data Centre (EIDC, www.eidc.ceh.uk). https://doi.org/10.5285/0869d961-99ca-4946-9192-f35afccdda38.

Respondents' characteristics and experience with norovirus sources. The online survey was completed by 1006 adults between the 15th and 20th February 2018. In order to assess the awareness of people most vulnerable to water- and foodborne illnesses, individuals who had consumed bivalve shellfish and had been in contact with environmental waters within the previous year were asked to participate. After the removal of incomplete responses, those who had completed the survey so quickly as to suggest inadequate consideration of their responses (n = 194), the sample comprised 806 responses (80% of total responses). Of those 806 responders, 47% were male, 96% white, 7% were 18–24 years old, 30% were aged 24–44 years old, 38% were 45–64 years old and 25% were over 65 (Table S1). Of the respondents, 72% were parents or legal guardians with 62% having one child and 19% having two children. 47% of the children were older than 7 years (25% of the children were 7–12 and 22% were 13–18 years old).

The respondents consumed shellfish cooked in a restaurant or food outlet (89%, n = 718), at home (79%, n = 636), as ready meals (68%, n = 548) or as pickles (54%, n = 437) in the past year. Approximately one half of the sample consumed shellfish raw in restaurants (48%, n = 387) and a third consumed them raw at home (32%, n = 255). Of all respondents, 16% (n = 125) believed that they had contracted gastroenteritis (defined as "upset stomach", see "Methods" section for details) due to shellfish consumed in a restaurant in the UK or abroad (9%, n = 70 and 5%, n = 37, respectively), takeaway in the UK or abroad (5%, n = 38 and 3%, n = 22, respectively) or as a home-cooked meal in the UK or abroad (4%, n = 34 and 0.6%, n = 5, respectively). Among the 125 respondents who believed their illness associated with consuming shellfish, most respondents started to make sure the shellfish they were eating were well cooked (75%, n = 94), told others about the experience (61%, n = 76), had a break from eating shellfish (61%, n = 76) or raw shellfish (43%, n = 54), permanently stopped eating raw shellfish at home (27%, n = 34) or in a restaurant (28%, n = 35).

The majority of the respondents (87%, n = 701) had contact with UK recreational waters 4–5 times a year and 29% (n = 234) of the respondents had experienced gastroenteritis-related illness after the activity. Due to the illness, 60% (n = 140) of those people started to use hand sanitizers after recreational activity, 52% (n = 121) told others about the experience, 51% (n = 119) took a break from using recreational waters, 38% (n = 90) changed the places they visited and 24% (n = 57) changed or stopped their recreational activities. Of all 806 respondents, 22% (n = 178) had contracted gastroenteritis after nursing someone with gastroenteritis in the UK and 14% (n = 113) became ill after taking care of a sick person abroad.

Respondents' awareness of norovirus. Over the previous year, 40% (n=324) of the respondents had looked for advice or guidance regarding gastroenteritis (Table 1). Of those who visited one or more of the governmental websites (National Health Services choices website, Public Health England and Wales, Health Protection Scotland, Food Standards Agency, Food Standard Scotland), 69% (n=97) found the information useful and all would use these resources in the future. Of those who did not use governmental websites as resources, 74% (n=136) were not aware of the resources and indicated that they would use these resources in the future. Of those respondents who did not look for advice on governmental websites (Table 1), 27% (n=216) would use governmental websites. Of those who would not use governmental websites, 77% (n=454) were not aware of

	Sought information on gastroenteritis (n = 324), n(%)	Would consider seeking information on gastroenteritis (n = 482), n(%)		
Accident and Emergency Department	43 (13%)	42 (9%)		
General practitioner/family doctor	121 (37%)	238 (49%)		
Medical center	47 (15%)	127 (26%)		
NHS 111	52 (16%)	215 (45%)		
NHS Choices website	105 (32%)	188 (39%)		
PHE/PHW/HPS website	45 (14%)	57 (12%)		
FSA/FSS website	24 (7%)	33 (7%)		
Newspapers	15 (5%)	7 (1%)		
Online newspapers	22 (7%)	16 (3%)		
Friends, family social media	77 (24%)	117 (24%)		
Internet	112 (35%)	184 (38%)		
Pharmacy	7 (2%)	43 (9%)		
Other	4 (1%)	0 (0%)		

Table 1. Summary on the use of information sources on gastroenteritis. NHS: National Health Service; PHE:Public Health England; PHW: Public Health Wales; HPS: Health Protection Scotland; FSA: Food StandardsAgency; FSS: Food Standards Scotland.



Figure 1. Respondents' knowledge on pathogens causing gastroenteritis. *Methicillin-resistant *Staphylococcus aureus*.

the resources, 12% (n = 71) believed those resources would not be useful and 11% (n = 65) were not interested or would rather ask a health care professional in person.

The respondents were also asked about their knowledge of pathogens associated with gastroenteritis, including a fictive pathogen (*Perginella*) as a control. *Salmonella* was the pathogen which most people indicated they knew about followed by *E. coli* (Fig. 1). Norovirus was ranked 3rd (with MRSA) with 73% claiming to know a little or a lot about it. The equivalent figure for 'winter vomiting bug' (synonym for norovirus) was 67%. *Listeria* was ranked 6th and *Campylobacter* ranked 9th (21% said they knew a little about it, 6% claimed to know a lot) along with rotavirus, *Shigella* and the fictive pathogen, *Perginella* was reported as the least-known pathogen, with 3% of respondents claiming to know a lot about it.

Respondents' perceptions of fear. The respondents' ratio-scaled fear and control scores were determined based on the BWS experiment. The BWS part included 16 risks, such as norovirus, other air, food and waterborne illnesses and other hazards (Table 2). Ratio-scaled fear scores from estimation of the mixed logit model are reported in Table 3 and displayed in Fig. 2A. The most feared risk was heart attack, followed by getting dementia and becoming ill with skin cancer. The most feared environmental risk, ranked 4th, was getting lung

No	Variable name	Risk as defined in the BWS tasks
1	Terrorist attack	Being a victim of a terrorist attack
2	Dementia	Getting dementia in your lifetime
3	Car accident	Being injured in a car accident
4	Lung disease	Getting lung disease as a result of air pollution
5	Antibiotic resistance	Getting ill from bugs that are not killed by antibiotics
6	Fire	Fire at home
7	Diabetes	Getting diabetes in your lifetime
8	Heart attack	Suffering a heart attack in your lifetime
9	Pesticide residues	Becoming ill from eating substances that control pests or weeds that would remain on or in food
10	Skin cancer	Getting skin cancer in your lifetime
11	Common cold	Getting the common cold
12	Dog bite	Being bitten by a dog
13	Bioaerosols	Having breathing difficulties from inhaling spores from mold
14	Salmonella	Getting food poisoning from Salmonella
15	Lightning	Being struck by lightning
16	Norovirus	Catching a bug which causes nausea, vomiting, diarrhea and abdominal cramping. Sometimes called the winter vomiting bug

Table 2. Risks used in the best worst scaling (BWS).

	Fear			Control		Control (anchored) ^a			
Risk	RSS	95% Lower	95% Upper	RSS	95% Lower	95% Upper	RSSA	95% Lower	95% Upper
Terrorist attack	8.439	7.980	8.898	1.706	1.450	1.962	- 29.595	- 32.328	-26.863
Dementia	12.643	12.304	12.982	1.692	1.472	1.911	-24.027	-26.637	-21.416
Car accident	7.750	7.398	8.103	6.367	5.997	6.737	10.793	8.297	13.289
Lung disease	9.217	8.920	9.513	3.321	3.122	3.521	-3.831	- 6.042	- 1.621
Antibiotic resistance	6.196	5.864	6.529	4.510	4.257	4.762	-0.063	-2.320	2.195
Fire	7.869	7.466	8.273	12.210	11.860	12.559	37.436	34.953	39.919
Diabetes	4.794	4.520	5.067	9.519	9.192	9.846	21.964	19.409	24.519
Heart attack	13.496	13.256	13.735	5.249	4.920	5.579	6.809	4.314	9.303
Pesticide residues	2.882	2.668	3.096	6.616	6.312	6.921	5.571	3.137	8.006
Skin cancer	10.992	10.683	11.302	7.529	7.140	7.918	15.286	12.606	17.967
Common cold	0.548	0.409	0.687	6.926	6.495	7.356	10.262	7.321	13.202
Dog bite	1.341	1.142	1.539	9.794	9.424	10.163	18.412	15.827	20.998
Bioaerosols	3.527	3.329	3.725	5.736	5.534	5.938	3.129	0.911	5.347
Salmonella	3.857	3.617	4.097	9.348	9.075	9.622	20.007	17.456	22.557
Lightning	1.848	1.607	2.090	3.944	3.578	4.310	- 12.750	-15.711	- 9.789
Norovirus	4.601	4.288	4.914	5.533	5.239	5.827	5.285	2.827	7.743

Table 3. Ratio-scaled fear and control scores (RSS, sum to 100) with 95% confidence intervals. The control scores are presented as unanchored (RSS) and anchored (RSSA) scores. ^aZero = threshold for taking action to reduce the risk.

disease from air pollution. Norovirus was ranked 10th in terms of fear, with a level of fear equal to that from diabetes. The fear associated with norovirus was 18% greater than associated with *Salmonella*, 30% greater than that from bioaerosols, and 60% greater than the fear of illness from pesticide residues. We have found no significant differences in the fear levels of people with different age, gender, education, employment status, family background, income or ethnicity.

Respondents' perceptions of control. Respondents believed they had the greatest levels of control over fire at home, dog bites, developing diabetes and suffering food poisoning from *Salmonella* (Fig. 2B). Norovirus was ranked 10th with the perceived level of control very similar to that for bioaerosols (i.e. breathing difficulties from inhaling spores from mold). Developing lung disease from air pollution ranked 14th as the least controlled environmental risk. The level of control people felt they had over norovirus was 70% lower than the perceived level of control over *Salmonella* food poisoning and 20% lower that the level of control over becoming ill from



Figure 2. The ranking of ratio-scaled scores (RSS) relating to (**A**) fear and (**B**) control for the 16 risk items (n = 806). Error bars represent 95% confidence intervals. Red striped bar shows risk associated with norovirus and orange, crosshatched bars show other environment-associated risks.



Figure 3. Anchored ratio-scaled control scores (RSSA). Zero = threshold for taking action to reduce the risk. The error bars represent 95% confidence intervals. Red striped bar shows risk associated with norovirus and orange, crosshatched bars show other environment-associated risks.

pesticide residues. People believed they had far greater (72%) levels of control over acquiring diabetes than they did over norovirus.

The consideration thus far has been relative—the degree of fear and control has been considered in relation to other risks included in the survey. The mixed logit model was re-estimated incorporating the information on which risks respondents took action to lower their risk (i.e. they exerted some control) with threshold normalized to zero (Fig. 3; Table 3). The anchored results indicate that norovirus is one of the risks people typically think they are able to control, as they do with illness from pesticide residues and bioaerosols. The marked difference in perceived control between norovirus and both diabetes and *Salmonella* is present in the anchored (RSSA) results. Getting dementia or developing lung disease are the two illness risks for which people did not take actions to lower their risk and hence scored below the zero threshold—along with 'events', such as lightning strike and terrorist attack.

The results reported thus far used the mean fear and control scores for each risk, generated by estimation of the mixed logit model. The model also yielded individual level estimates, conditional on their control BWS choice data and the population parameters. These estimates allowed us to consider sample level heterogeneity and systematic differences in perceptions of control, over observable characteristics. This revealed differences between age groups, genders, education levels and between those who had, and had not, sought advice on gastroenteritis (Figure S2). The analysis showed no trends or significant differences in the perception of individuals in different ethnic groups, employment status, having children or previously experiencing water- or shellfishborne gastroenteritis.





Females perceived themselves to have higher levels of control than males for risks including norovirus, *Salmo-nella*, getting diabetes, illness from pesticide residues, skin cancer and illness from bioaerosols. People aged 18–24 perceived themselves to have lower control over suffering a heart attack in their lifetime than those aged 45–64.

All risks were considered more controllable among the higher educated population. The difference was significant for most risk items, except fire at home, getting the common cold, being bitten by a dog and being struck by lightning. Respondents who had sought advice on gastroenteritis believed they had more control on all the risks than those who did not. The differences were significant except for fire at home, getting diabetes, suffering a heart attack, getting skin cancer and getting breathing difficulties due to bioaerosols.

Fear vs. control. The results from the fear and (unanchored) control models are combined in Fig. 4 which shows the ratio-scaled fear and control scores (from Table 3). The re-scaling sums the scores to 100 and the axes represent mean levels of fear (and control) which, with 16 hazards included, is equal to score 6.25 (Fig. 4). Hence, risks 'north' of the x-axis indicate above average levels of control, and those 'east' of the y-axis indicate above average levels of control, and those 'east' of the y-axis indicate above average levels of fear. There was a negative correlation between the risk dimensions with higher levels of fear associated with lower levels of control. Risks, such as dementia, terrorist attack, having a heart attack or developing lung disease from air pollution, were associated with relatively high levels of fear and relatively low levels of control.

Salmonella was characterized by relatively low fear levels and high levels of control. Illness from bioaerosols and pesticide residues were associated with similar levels of fear as *Salmonella*, but significantly lower levels of control. Norovirus generated higher levels of fear and less control than these three hazards.

Discussion

This study evaluated the public's level of awareness of norovirus infection in relation to both environmental and general risks using an online survey. In order to identify the most vulnerable populations, only people regularly exposed to environmental risks (i.e. recreational water users and shellfish eaters) were included. It is estimated that 51% of the UK population consumes shellfish⁴¹, suggesting that the survey covered a great proportion of the population. However, there is no data available on people's recreational activities.

Our study revealed that 28% of the respondents had gastroenteritis-related symptoms after using recreational water. Jones et al.⁴² have also found that 20% of people had diarrhea, 12% had nausea and 4% vomited up to three weeks after exposure to recreational water at Swansea, UK. In an UK-wide survey, Fleisher et al.⁴³ also found that 14.9% of participants had gastroenteritis after bathing in natural waters. In both cases, the rates of illnesses were significantly higher among people exposed to water than in the control groups suggesting strong association between bathing and developing symptoms.

Our study revealed that participants had gastroenteritis after being in direct contact with a sick person (22%) and eating shellfish (16%), suggesting that these routes may also be significant routes of gastroenteritis transmission. Becoming ill from shellfish or recreational activities caused significant changes in people's behavior; most people started to take precautions after being sick, such as increased use of hand sanitizers or no longer eating raw or lightly cooked shellfish.

The survey revealed the resources people used, or would use, to learn more about gastrointestinal diseases. We found, that although detailed information on the transmission and prevention of norovirus, and other environment-associated risks, is available on governmental websites, 74% of respondents were not aware of

those resources. Understanding how and why people (fail to) become aware of the availability of such resources, could improve reduction and management of the risks. Improved understanding of those processes would be particularly valuable in relation to those groups most vulnerable to the risks.

Many respondents claimed to have good knowledge of some of the most common food- and waterborne pathogens, e.g. Salmonella and norovirus (Fig. 1). Their claimed levels of knowledge regarding norovirus and its synonym, 'winter vomiting bug', were very similar, suggesting that both terms are familiar to people. Norovirus is responsible for approx. 600,000 to one million cases each year in the UK⁴⁴, hence the recognition of the pathogen was expected. Participants claimed to have far less knowledge on Listeria, a food pathogen with the most severe typical impact on health (a typical loss per case of 4.03 Quality Adjusted Life Years (QALYs) compared to 0.67 for norovirus⁴⁵. Participants also claimed little knowledge of Campylobacter, even though this pathogen is the most common foodborne pathogen in the UK with over 300,000 cases and the highest annual burden on the economy (£424.4 M), followed by norovirus (£248.5 M) and Salmonella (£143.9 M)⁴⁵. Furthermore, rotavirus was not recognized by 48% of respondents. Rotavirus had been the most common cause of gastroenteritis among children under five years old until the vaccination introduced in 2013, which lowered the number of cases requiring hospitalization by 83%⁴⁶. A similar decline in cases of Salmonella was observed between 2000 and 2017 due to improved legislation, food safety advice, and an industry-led flock vaccination program⁴⁷, however, the number of lab-confirmed cases is still approx. 9000/year⁴⁸. As all these foodborne pathogens cause mild gastroenteritis in the majority of cases, people may not be familiar with the different causative agents, and may believe their symptoms are caused by a pathogen they are familiar with (e.g. Salmonella, norovirus). Therefore, more information should be disseminated for people on the different kinds of pathogens. With that knowledge, people would be better able to assess their symptoms and more likely to seek medical help when necessary.

The main aim of this study was to investigate perceptions of norovirus compared to both other emerging risks associated with contaminated air and water and more established risks (e.g. heart attack, hit by lightning). The results of the BWS revealed low levels of fear and control regarding norovirus and other emerging environmental health risks compared to more general hazards. A possible reason may be that some of the risks, such as norovirus or *Salmonella* infection, mostly cause mild illness that resolve without treatment within days^{10,49}.

Norovirus fear levels were equal to those from diabetes. This is remarkable given that over three million people in the UK have diabetes with this number estimated to rise to 4.6 million people by 2030⁵⁰. Norovirus prompted significantly higher levels of fear than *Salmonella*, pesticide residues and bioaerosols. This correlates with the number of cases and economic burden of the two foodborne pathogens⁴⁵. The levels of fear from lung disease and antibiotic resistance were higher than the level of fear prompted by norovirus. In contrast, reverse association was observed in terms of control (Fig. 2; Table 3). This suggests that along with the severity of symptoms, active management (i.e. control) of a risk also influence the overall perception of the risk. With more knowledge on the precautions one can take to avoid certain hazards, the fear prompted by the hazard may be reduced. Therefore, taking into account the high number of norovirus cases each year, the lack of concern may be due to misinformation or the lack of information on the transmittance and prevention of the disease.

The results show significant gender differences in the perceived levels of control when anchored analysis was used. In general, women believed they had more control over the risks. However, many previous studies showed that women have similar or lower levels of control than men on a wide range of risks, such as being run over/burgled, stressed environment (including technological, behavioral, and land use hazards) and earthquakes^{51–54}. We found the highest differences between men and women in the control of risks associated with food preparation (getting diabetes, becoming ill due to pesticides or *Salmonella*/norovirus infection) and personal hygiene and health (developing skin cancer, common cold and antibiotic resistance pathogens). In these areas, women may believe they are more informed on the risks and take measures to reduce them by for instance thoroughly washing vegetables, adequately cooking meat and cleaning and washing hands regularly^{55,56}.

Our results also showed that the level of risk perception in terms of control increases with age and educational qualifications. These findings suggest that people are more aware of the risks with more experience and knowledge. We found no significant differences in the control levels of people with different employment status, family background, income or ethnicity. Many studies have explored the effect of socio-demographic factors on behavior, e.g. disaster preparedness or consumer behavior^{57–59}, however, little information is available on the effect of these factors on risk perception. More targeted analysis is therefore necessary to explore the effect of socio-demographic factors on risk perception.

The BWS results suggest that those who sought advice on gastroenteritis perceived themselves to have higher levels of control over risks (related and unrelated) to the illness. That also indicates that people who sought advice are generally more aware and cautious of risks, and hence take measures to better understand and control them. As frequent exposure to messages on health and well-being have been shown to positively influence health-related behavior⁶⁰, the targeted dissemination of information on environment-associated health risks may increase control over risk and hence reduce the number of illnesses. The information to be disseminated may include detailed information on the transmission of norovirus and options to mitigate risk of infection, e.g. hand washing, cooking shellfish and washing vegetables before consumption, good kitchen hygiene⁶¹. According to our analysis, men and people with low education levels believe they have the least control on environmental risks (Figure S2), hence distribution of information targeting these socio-demographic groups may be required. Our survey revealed that the information available on public health and food safety-related websites was regarded as adequate, however, many participants were not aware of these resources. Therefore, more information (including QR code and other links to official web resources) should be circulated at places where people would look for such advice, e.g. doctors' offices, hospitals, medical centers and the health services' websites. Albeit controversial, more information on food safety could also be placed on shellfish food packaging.

Conclusions

In this study, we assessed the public's perception of norovirus infection in relation to other environment-associated and general health risks. Our results suggest that the level of norovirus awareness is relatively high compared to other foodborne pathogens (e.g. *Listeria, Cryptosporidium*, rotavirus or *Campylobacter*) and slightly lower than the claimed knowledge on *Salmonella, E. coli* and MRSA. Furthermore, our BWS results suggest that the levels of fear associated with norovirus are comparable to those from diabetes. Despite the considerable levels of awareness and fear, the level of control people felt they had over norovirus was low, significantly below that for *Salmonella* or diabetes.

Given the scale of the public health burden associated with norovirus, addressing this deficit in the public's understanding of how to control exposure to the pathogen is an important public health need. However, our study revealed some association between perceived control on norovirus and age, gender and education. Further research is therefore needed to understand the effect of sociodemographic factors and behavior on the perception of norovirus and other food- and waterborne illnesses. Our current findings suggest that different approaches are needed to disseminate information regarding transmission of the virus to increase awareness and reduce risk of infection.

Methods

Survey design. The online survey we designed comprised of two sections. The first section elicited demographic information (age, gender, income, qualification, ethnicity etc.). The respondents were also asked about behaviors relevant to exposure to norovirus, their knowledge of the pathogen and their use of information sources related to it. Specifically:

- Type and frequency of shellfish consumption.
- How often, and where, they come into contact with recreational waters (including rivers, lakes and seawater in the UK).
- Whether they had had contact with anyone with gastroenteritis or food/shellfish-related gastroenteritis over the past year and if so, where the shellfish was obtained.
- Whether they had sought information about gastroenteritis over the past year and if so, whether they found the information useful.
- The reason for not seeking information and where they would consider looking.
- How much they had heard about certain enteric pathogens (*Salmonella*, methicillin-resistant *Staphylococcus aureus*—MRSA, *Cryptosporidium*, norovirus, *Escherichia coli*, *Clostridium difficile*, *Listeria*, winter vomiting bug, *Shigella*, *Campylobacter*, rotavirus) and a fictive one (*Perginella*), which acted as a control. We used the terms 'norovirus' and its synonym, 'winter vomiting bug', separately to assess which term is better known.

The survey referred to gastroenteritis as "upset stomach" defined as onset of stomach pains that can strike quickly with force and make a person feel very sick, which typically resolves within 2–3 days with symptoms of nausea, vomiting, diarrhea and abdominal cramping. The full list of questions is detailed in the Supplementary Material.

The second part of the survey comprised the Best Worst Scaling (BWS) exercise. BWS is a choice-based survey method designed to elicit the relative importance of multiple items enabling accurate ranking. Rather than asking respondents to rank large numbers of items, they are asked to choose the 'most' and 'least' from repeated subsets of items³⁵. In our study, the items were the risks considered in the study (see Table 1) and respondents were asked to choose they had most and least control over (Figure S1). The task was repeated with varying combinations of four items.

The BWS experiment contained 16 risks, including norovirus, other air, food and waterborne illnesses and other hazards (Table 1). Some of the risks were well-established in the risk perception literature (e.g. heart attack, car accident), whereas others were more novel (e.g. antibiotic resistance, bioaerosols). The allocation of risks into sets was determined by an experimental design which varied the combinations of risk presented, the frequency with which they occur and co-occur and their position in the sets (top, middle, bottom). Respondents completed 12 tasks using each criteria (fear, control) with the risks randomly allocated.

The approach employed here resembles that of Erdem and Rigby⁵³, who elicited perceptions of a set of risks in terms of fear and control—as these had been identified in the psychometric risk perception literature as dimensions on which risks could be meaningfully located. We augmented their approach by the introduction of an absolute threshold into the analysis of control levels. The BWS method, as typically employed, is a relative assessment of the items under consideration. In this case, as well as eliciting the relative degree of fear that the hazards induced, and the degree of control people perceived they had over them, we also asked, for each risk, whether the respondent ever amended their behavior to reduce the risk. We then incorporated that threshold information into the analysis of control. Further details on survey and experiment design and model specification can be found in the Supplementary Material.

Data collection and analysis. The survey was distributed by an online marketing company (Research Now, UK). Only adults (over 18 years old) who had consumed bivalve shellfish and had been in contact with environmental waters within the previous year were included in the study. The survey was piloted on the 13th February 2018 with 100 participants. When the required number of answers was reached, the survey was paused, and survey responses were checked to ensure that sensible answers were received. Based on the responses, the skip logic was working, people saw the right responses based on their answers and the ratio of replies were what we predicted and people who did not meet the criteria were disqualified. The survey was then administered

between the 15th and 20th February 2018 with 1006 participants. After the removal of incomplete responses, those who had completed the survey so quickly as to suggest inadequate consideration of their responses (n = 194), the sample comprised 806 responses.

The BWS data were analyzed via estimation of random utility models⁶² and mixed logit models^{63,64}, as detailed in the Supplementary Material. Best–worst raw scores were calculated for each risk and then transformed into standardized ratio-scaled (0–100) scores (RSS) and anchored ratio-scaled scores (RSSA) incorporating threshold data using Lighthouse Studio⁶⁵.

Ethical statement. The ethical approval for this study was obtained from the Ethics Committee of Bangor University. The online survey was handled by Research Now, UK. All procedures performed in studies involving human participants were in accordance with the 1964 Helsinki declaration and its later amendments. This manuscript does not contain any individual person's data in any form. The introductory text of the survey informed the participants that the survey was part of a scientific project with its aims and methods explained, and that participation was optional. Informed consent for participation was obtained by Research now, UK.

Received: 28 November 2019; Accepted: 10 May 2021 Published online: 27 May 2021

References

- Cohen, A. J. *et al.* The global burden of disease due to outdoor air pollution. *J. Toxicol. Environ. Heal. Part A* 68, 1301–1307 (2005).
 Lin, W. E., Carpentieri, M. & Savory, E. Transmission of human respiratory disease by indoor bioaerosols. in *Physmod 2017–Inter-*
- Lin, W. E., Carpentieri, M. & Savory, E. Transmission of numan respiratory disease by indoor bioaerosois. in *Physmol 2017—Inter*national Workshop on Physical Modelling of Flow and Dispersion Phenomena, Dynamics of Urban and Coastal Atmosphere—LHEEA -École Centrale de Nantes, France 23–25 August 2017 (2017).
- 3. Kallawicha, K. *et al.* Exposure to ambient bioaerosols is associated with allergic skin diseases in Greater Taipei residents. *Environ. Pollut.* **216**, 845–850 (2016).
- Trout, D., Bernstein, J., Martinez, K., Biagini, R. & Wallingford, K. Bioaerosol lung damage in a worker with repeated exposure to fungi in a water-damaged building. *Environ. Health Perspect.* 109, 641–644 (2001).
- O'Neill, J. Antimicrobial Resistance: Tackling a Crisis for the Health and Wealth of Nations, 2014. (2014). https://doi.org/10.1038/ 510015a.
- Efstratiou, A., Ongerth, J. E. & Karanis, P. Waterborne transmission of protozoan parasites: Review of worldwide outbreaks—An update 2011–2016. Water Res. 114, 14–22 (2017).
- Thongprachum, A., Khamrin, P., Maneekarn, N., Hayakawa, S. & Ushijima, H. Epidemiology of gastroenteritis viruses in Japan: Prevalence, seasonality, and outbreak. J. Med. Virol. 88, 551–570 (2016).
- Chatziprodromidou, I. P., Bellou, M., Vantarakis, G. & Vantarakis, A. Viral outbreaks linked to fresh produce consumption: A systematic review. J. Appl. Microbiol. 124, 932–942 (2018).
- 9. Ahmed, S. M. *et al.* Global prevalence of norovirus in cases of gastroenteritis: A systematic review and meta-analysis. *Lancet Infect. Dis.* 14, 725–730 (2014).
- Katayama, H. & Vinje, J. Norovirus and other Caliciviruses. in Global Water Pathogen Project (eds. Meschke, J. S. & Gironés, R.) (Michigan State University, E. Lansing, MI, UNESCO, 2017).
- 11. Barardi, C. R. et al. Monitoring viruses in environmental samples. Int. J. Environ. Sci. Eng. Res. 3, 62-79 (2012).
- 12. CDC. Norovirus worldwide. (2016). (Accessed 29 May 2019). https://www.cdc.gov/norovirus/worldwide.html.
- Bartsch, S. M., Lopman, B. A., Ozawa, S., Hall, A. J. & Lee, B. Y. Global economic burden of norovirus gastroenteritis. *PLoS ONE* 11, 1–16 (2016).
- 14. Bellou, M., Kokkinos, P. & Vantarakis, A. Shellfish-borne viral outbreaks: A systematic review. Food Environ. Virol. 5, 13-23 (2013).
- 15. Radin, D. New trends in food- and waterborne viral outbreaks. Arch. Biol. Sci. 66, 1–9 (2014).
- Aoki, Y. et al. Duration of norovirus excretion and the longitudinal course of viral load in norovirus-infected elderly patients. J. Hosp. Infect. 75, 42–46 (2010).
- 17. Sidhu, J. P. S., Sena, K., Hodgers, L., Palmer, A. & Toze, S. Comparative enteric viruses and coliphage removal during wastewater treatment processes in a sub-tropical environment. *Sci. Total Environ.* **616**, 669–677 (2017).
- Qiu, Y. *et al.* Assessment of human virus removal during municipal wastewater treatment in Edmonton, Canada. J. Appl. Microbiol. 119, 1729–1739 (2015).
- Kitajima, M., Iker, B. C., Pepper, I. L. & Gerba, C. P. Relative abundance and treatment reduction of viruses during wastewater treatment processes—Identification of potential viral indicators. *Sci. Total Environ.* 488, 290–296 (2014).
- Iaconelli, M. et al. One-year surveillance of human enteric viruses in raw and treated wastewaters, downstream river waters, and drinking waters. Food Environ. Virol. 9, 79–88 (2017).
- Farkas, K. et al. Seasonal and spatial dynamics of enteric viruses in wastewater and in riverine and estuarine receiving waters. Sci. Total Environ. 634, 1174–1183 (2018).
- Silverman, A. I., Nelson, K. L., Akrong, M. O., Amoah, P. & Drechsel, P. Quantification of human norovirus GII, human adenovirus, and fecal indicator organisms in wastewater used for irrigation in Accra, Ghana. J. Water Health 11, 473–488 (2013).
- 23. Hassard, F. *et al.* Abundance and distribution of enteric bacteria and viruses in coastal and estuarine sediments—A review. *Front. Microbiol.* 7, 1692 (2016).
- Landry, E. F., Vaughn, J. M., Vicale, T. J. & Mann, R. Accumulation of sediment-associated viruses in shellfish. Appl. Environ. Microbiol. 45, 238–247 (1983).
- Lowther, J. A., Gustar, N. E., Powell, A. L., Hartnell, R. E. & Lees, D. N. Two-year systematic study to assess norovirus contamination in oysters from commercial harvesting areas in the United Kingdom. *Appl. Environ. Microbiol.* 78, 5812–5817 (2012).
- Wang, J. & Deng, Z. Modeling and prediction of oyster norovirus outbreaks along Gulf of Mexico Coast. *Environ. Health Perspect.* 124, 627–633 (2016).
- 27. MacCannell, T. *et al.* Guideline for the prevention and control of norovirus gastroenteritis outbreaks in healthcare settings. *Infect. Control Hosp. Epidemiol.* **32**, 939–969 (2011).
- Centers for Disease Control and Prevention. Updated norovirus outbreak management and disease prevention guidelines. MMWR Recomm. Reports 60, 1–18 (2011).
- Baert, L. et al. Review: Norovirus prevalence in Belgian, Canadian and French fresh produce: A threat to human health?. Int. J. Food Microbiol. 151, 261–269 (2011).

- Viau, E. J., Lee, D. & Boehm, A. B. Swimmer risk of gastrointestinal illness from exposure to tropical coastal waters impacted by terrestrial dry-weather runoff. *Environ. Sci. Technol.* 45, 7158–7165 (2011).
- Mara, D. & Sleigh, A. Estimation of norovirus infection risks to consumers of wastewater-irrigated food crops eaten raw. J. Water Health 8, 39–43 (2010).
- 32. Sano, D. et al. Risk management of viral infectious diseases in wastewater reclamation and reuse: Review. Environ. Int. 91, 220–229 (2016).
- 33. Liverani, M. et al. Understanding and managing zoonotic risk in the new livestock industries. Environ. Health Perspect. 121, 873–877 (2013).
- Anacleto, P., Barrento, S., Nunes, M. L., Rosa, R. & Marques, A. Portuguese consumers' attitudes and perceptions of bivalve molluscs. Food Control 41, 168–177 (2014).
- Finn, A. & Louviere, J. J. Determining the appropriate response to evidence of public concern: The case of food safety. J. Public Policy Mark. 11, 12–25 (1992).
- Lin, C.-T.J. & Milon, J. W. Attributes and safety perception in a double-hurdle model of shellfish consumption. Am. J. Agric. Econ. 75, 724–729 (1993).
- 37. Le Bihan, V., Pardo, S. & Guillotreau, P. Risk perception and risk management strategies of oyster farmers. *Mar. Resour. Econ.* 28, 285–304 (2013).
- Borchardt, M. A., Spencer, S. K., Kieke, B. A., Lambertini, E. & Loge, F. J. Viruses in nondisinfected drinking water from municipal wells and community incidence of acute gastrointestinal illness. *Environ. Health Perspect.* 120, 1272–1279 (2012).
- Edelstein, M., Wallensten, A., Zetterqvist, I. & Hulth, A. Detecting the norovirus season in Sweden using search engine data— Meeting the needs of hospital infection control teams. *PLoS ONE* 9, 1–8 (2014).
- Knight, A. J., Worosz, M. R. & Todd, E. C. D. Serving food safety: Consumer perceptions of food safety at restaurants. Int. J. Contemp. Hosp. Manag. 19, 476–484 (2007).
- 41. Mintel. Executive summary: Fish and Shellfish. (2017).
- Jones, F., Kay, D., Stanwell-Smith, R. & Wyer, M. Results of the first pilot-scale controlled cohort epidemiological investigation into the possible health effects of bathing in seawater at Langland Bay, Swansea. Water Environ. J. 5, 91–98 (1991).
- Fleisher, J. M., Kay, D., Wyer, M. D. & Godfree, A. F. Estimates of the severity of illnesses associated with bathing in marine recreational waters contaminated with domestic sewage. *Int. J. Epidemiol.* 27, 722–726 (1998).
- 44. HPA. Norovirus Fact Sheet. (2011).
- FSA. Estimating Quality Adjusted Life Years and Willingness to Pay Values for Microbiological Foodborne Disease (Phase 2). (2017).
 Hungerford, D. et al. Early impact of rotavirus vaccination in a large paediatric hospital in the UK. J. Hosp. Infect. 93, 117–120
- Hungertord, D. et al. Early impact of rotavirus vaccination in a large paediatric hospital in the UK. J. Hosp. Infect. 93, 117–120 (2016).
- 47. O'Brien, S. J. The, "decline and fall" of nontyphoidal salmonella in the United Kingdom. Clin. Infect. Dis. 56, 705-710 (2013).
- 48. FSA. Consolidated annual report and accounts 2017/18. (Food Standards Agency, 2018).
- 49. D'Aoust, J.-Y. Pathogenicity of foodborne Salmonella. Int. J. Food Microbiol. 12, 17-40 (1991).
- Holman, N., Forouĥi, N. G., Goyder, E. & Wild, S. H. Epidemiology The Association of Public Health Observatories (APHO) Diabetes Prevalence Model: Estimates of total diabetes prevalence for England, 2010–2030. *Diabet. Med.* 28, 575–582 (2011).
- Greenberg, M. R. & Schneider, D. F. Gender differences in risk perception: Effects differ in stressed vs. non-stressed environments. *Risk Anal.* 15, 503–511 (1995).
- 52. Ming-Chou, H., Daigee, S., Shuyeu, L. & Yao-Chu, C. How do disaster characteristics influence risk perception?. *Risk Anal.* 28, 635–643 (2008).
- Erdem, S. & Rigby, D. Investigating heterogeneity in the characterization of risks using best worst scaling. *Risk Anal.* 33, 1728–1748 (2013).
- Yi-Wen, K. & Sue-Huei, C. Perception of earthquake risk in Taiwan: Effects of gender and past earthquake experience. *Risk Anal.* 32, 1535–1546 (2012).
- 55. Jay, L. S., Comar, D. & Govenlock, L. D. A National Australian food safety telephone survey. J. Food Prot. 62, 921-928 (2016).
- 56. Lake, A. A. *et al.* Food shopping and preparation among the 30-somethings: Whose job is it? (The ASH30 study). *Br. Food J.* **108**, 475–486 (2006).
- Miceli, R., Sotgiu, I. & Settanni, M. Disaster preparedness and perception of flood risk: A study in an alpine valley in Italy. J. Environ. Psychol. 28, 164–173 (2008).
- Verbeke, W. Consumer acceptance of functional foods: Socio-demographic, cognitive and attitudinal determinants. Food Qual. Prefer. 16, 45–57 (2005).
- Nunes, F., Madureira, T., Oliveira, J. V. & Madureira, H. The consumer trail: Applying best-worst scaling to classical wine attributes. Wine Econ. Policy 5, 78–86 (2016).
- 60. Wakefield, M. A., Loken, B. & Hornik, R. C. Use of mass media campaigns to change health behaviour. *Lancet* 376, 1261–1271 (2010).
- Guix, S., Pintó, R. & Bosch, A. Final consumer options to control and prevent foodborne norovirus infections. Viruses 11, 333 (2019).
- 62. McFadden, D. Conditional logit analysis of qualitative choice behavior. In *Frontiers in Econometrics* (ed. Zarembka, P.) 105–142 (Academic Press, 1973).
- 63. Train, K. Discrete Choice Methods with Simulation (Cambridge University Press, 2003).
- 64. McFadden, D. & Train, K. Mixed MNL models of discrete response. J. Appl. Econom. 15, 447-470 (2000).
- 65. Sawtooth. Sawtooth Software. (2020).

Acknowledgements

This work was funded by the Natural Environment Research Council (NERC) under the Environmental Microbiology and Human Health (EMHH) Programme (NE/M010996/1) and by the Shellfish Centre RD&I operation, part-funded by the EU's West Wales and the Valleys European Regional Development Fund (ERDF) Operational Programme through the Welsh Government. The authors wish to thank David Lees, Andrew Turner, Andrew Younger and David Walker (Centre for Environment, Fisheries and Aquaculture Science, UK) for their help constructing the survey.

Author contributions

Conceptualization, K.F., E.G., D.R., P.C., S.M., D.J., S.T.; Methodology, K.F., E.G., D.R., P.C., S.M., D.J., S.T.; Data analysis, K.F., E.G., D.R.; Writing—original draft preparation, K.F., E.G., D.R., P.C.; Writing—review & editing, K.F., E.G., D.R., P.C., S.M., D.J., S.T.; Funding acquisition, D.J., S.T.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary Information The online version contains supplementary material available at https://doi.org/ 10.1038/s41598-021-90704-7.

Correspondence and requests for materials should be addressed to K.F.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/.

© The Author(s) 2021