

Precision and cost-effectiveness of bioindicators to estimate nutrient regimes on coral reefs

Vaughan, Eleanor J.; Wynn, Peter M. ; Wilson, Shaun K.; Williams, Gareth J.; Barker, Philip A. ; Graham, Nicholas A.J

Marine Pollution Bulletin

DOI:

<https://doi.org/10.1016/j.marpolbul.2021.112606>

Published: 01/09/2021

Peer reviewed version

[Cyswllt i'r cyhoeddiad / Link to publication](#)

Dyfyniad o'r fersiwn a gyhoeddwyd / Citation for published version (APA):

Vaughan, E. J., Wynn, P. M., Wilson, S. K., Williams, G. J., Barker, P. A., & Graham, N. A. J. (2021). Precision and cost-effectiveness of bioindicators to estimate nutrient regimes on coral reefs. *Marine Pollution Bulletin*, 170, Article 112606.
<https://doi.org/10.1016/j.marpolbul.2021.112606>

Hawliau Cyffredinol / General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Precision and cost-effectiveness of bioindicators to estimate nutrient regimes on coral reefs

Eleanor J. Vaughan^{1*}, Peter M. Wynn¹, Shaun K. Wilson^{2,4}, Gareth J. Williams³, Philip A. Barker¹, and Nicholas A.J. Graham¹

¹Lancaster Environment Centre, Lancaster University, Lancaster LA1 4YQ, UK ; ²Department of Biodiversity, Conservation and Attractions, Kensington, Perth, Western Australia 6151, Australia. ³School of Ocean Sciences, Bangor University, Menai Bridge, Anglesey LL59 5AB, UK; ⁴Oceans Institute, University of Western Australia, Crawley, WA 6009, Australia

* Corresponding author: e.vaughan@lancaster.ac.uk.

Highlights

- Multiple bioindicators and stable isotopes provide comprehensive, spatio-temporal assessments of nutrient regimes on coral reefs.
- N- and C-based nutrient signatures were assessed across eight bioindicators both within and among reefs as well as between degraded reef states, the most precise being brown macroalgae, green macroalgae, and zoanthids.
- There was low congruency between signatures of these three indicators due to differences in internal nutrient processing.
- Turf algae and sediment were more widespread, but their signatures were variable and did not reflect their local environment.

Abstract

Bioindicators are useful for determining nutrient regimes in marine environments, but their ability to evaluate corals reefs in different ecological states is poorly understood. The precision, availability and congruency of eight potential bioindicators (brown macroalgae, green macroalgae, turf algae, cyanobacteria, soft corals, zoanthids, sponges, and sediment) and their stable isotopic and elemental signatures ($\delta^{15}\text{N}$, $\delta^{13}\text{C}$, %N, %C, and C:N Ratio) were assessed across 21 reefs in the Inner Seychelles. The coefficient of variation (CoV) for $\delta^{15}\text{N}$ showed that green and brown macroalgae were highly precise (2.47 ± 0.95 , $n=11$; 4.68 ± 1.33 , $n=16$, respectively), though were less common on recently-bleached reefs relative to macroalgal-dominated ones. Zoanthids were also highly precise for $\delta^{15}\text{N}$ (2.98 ± 1.20), but were more readily available regardless of reef state ($n=18$). Congruency was low among these indicators, suggesting that different physiological mechanisms for nutrient processing have a stronger influence on a bioindicator's effectiveness than reef state.

Keywords: *Pollution; stable isotopes; macroalgae; environmental monitoring; regime shifts*

1. Introduction

Coral reefs are facing global declines in live coral cover due to climate change (Hughes et al., 2018), and local-scale degradation from overfishing and pollution (Burkepile & Hay, 2006; Littler et al., 2006; Zaneveld et al., 2016; MacNeil et al., 2019). Increased anthropogenic nutrient loads and reduced herbivory can cause the proliferation of opportunistic species such as fleshy macroalgae, which may lead to a regime shift from a coral-dominated to an algal-dominated reef (Littler et al., 2006; Hughes et al., 2007; Fulton et al., 2019). Monitoring the state of coral reefs relative to anthropogenic stressors provides insights into causes of decline in reef condition, potentially instigating management actions. Two particularly widespread

local stressors are fishing and eutrophication (Fabricius et al., 2005; Burkepile & Hay, 2006; Littler et al., 2006; Zaneveld et al., 2016). While there has been significant progress in understanding the effects of fishing (e.g. Cinner et al. 2018), it has been more difficult to detect and quantify nutrient loads that cause eutrophication in the marine environment, due to high spatio-temporal variability in the water column (Fabricius et al., 2005; Wyatt et al., 2013; D'Angelo & Wiedenmann, 2014; Briand et al., 2015; Lowe & Falter, 2015; Clausing & Fong, 2016; MacNeil et al., 2019). It is therefore critical to identify more cost-effective methods of capturing nutrient enrichment to improve assessments of coral reef health over different spatial scales as part of routine environmental monitoring strategies (Fabricius et al., 2012; Bal et al., 2020).

Bioindicators are used widely to capture nutrient regimes in tropical marine systems, as they provide an ecologically relevant response to bioavailable nutrients in the surrounding water column (Fichez et al., 2005; Cooper et al., 2009; Fabricius et al., 2012). As such, bioindicators are cost-effective alternatives to direct measures of seawater nutrients, which can be highly variable and require frequent sampling that do not always capture fine-scale temporal variation or wider ecological impacts (Fabricius et al., 2012). Suitable bioindicators are defined in Cooper et al. (2009) as those with biological responses that are a) specific towards a driver of change or stressor, b) reflective of the magnitude of any changes, c) consistent across different scales, d) cost-effective, and e) ecologically relevant. Non-biological indicators, conversely, are those which can still reflect drivers of change, but not through biological responses (i.e. nutrients stored in reef sediments) (Linton & Warner, 2003; Fichez et al., 2005).

Previous studies have measured the presence: absence ratio of selected bioindicators to investigate water quality (Fichez et al., 2005; Cooper et al., 2009), however, using this type of methodology alone does not take into account other biophysical factors that may influence their abundance (Linton & Warner, 2003). Therefore, measuring stable isotope signatures ($\delta^{15}\text{N}$ and $\delta^{13}\text{C}$) and concentration levels (%N, %C and C:N ratio) in the tissues of a selected bioindicator allows scientists and environmental managers to assess both the source(s) and concentration of nutrient regimes, respectively, better determine the spatio-temporal variability of nutrient regimes and detect and map the spatial ecological impacts (Costanzo et al., 2001). Fleshy macroalgae are widely used for such a purpose, because they respond rapidly to high nutrient concentrations by assimilating bioavailable nutrients from their local environment into their tissues over their active growth periods, thereby capturing temporal variation in nutrients (Costanzo et al., 2001). They are also easy to collect and survey in the field, especially in nutrient-rich coastal areas (Fichez et al., 2005; García-Seoane et al., 2018a&b; Zubia et al., 2018).

One of the main limitations of using only a single species of macroalgae, even with stable isotopic analyses, are the spatio-temporal gaps in their distribution, which are driven by a number of abiotic factors such as wave exposure, irradiance, temperature, rainfall and seasonality (Linton & Warner, 2003; Williams et al., 2013; Clausing & Fong, 2016; Duran et al., 2016; Fulton et al. 2019), and biotic factors such as herbivory and competition (Burkepile & Hay, 2006; Duran et al., 2016). These limiting factors may also affect the ability of macroalgae to proliferate on some reefs that have experienced significant disturbances (Littler et al., 1991; Graham et al., 2015). These distributional gaps can also lead to inconclusive or even misleading findings in any studies or monitoring programs, particularly if they are quantifying the abundance of a particular species across a range of target sites

(Linton & Warner, 2003). As such, the utility of alternative bioindicators to capture nutrient regimes is of importance to monitoring programmes.

A range of other marine organisms have been used as bioindicators in water quality or nutrient enrichment studies, such as scleractinian corals (Hoegh-Guldberg et al., 2004), soft corals (Fleury et al., 2000; Risk, 2014), and sponges (Ward-Paige et al., 2005). In addition, multiple candidate bioindicators have been used to assess water quality depending upon their response time to a change in their local nutrient environment (Cooper et al., 2009), or on the extent of their abundance and distribution, which also allows the spatial extent of nutrient runoff to be assessed (Fabricius et al., 2012). Some bioindicators may take longer to find or process than others, particularly in areas where they are relatively uncommon or rare. Selection of bioindicators should therefore also consider the cost-effectiveness of the collection and subsequent processing of samples (Risk et al., 2001; Drummond & Connell, 2008; Bal et al., 2020). This will be especially important for researchers and managers tasked with monitoring water quality over large spatial and temporal scales, such as entire reef systems (De'ath & Fabricius, 2010; Graham et al., 2015).

Few studies have tested whether patterns in nutrient signatures of different bioindicators are congruent (i.e. they are able to show the same relative trends in isotopic values between indicators) across different spatio-temporal scales or gradients (Tucker et al., 1999; Gartner et al., 2002; Pitt et al., 2009), and this multi-taxa approach is even less common in coral reef studies,(Connolly et al., 2013; Kürten et al., 2014; Graham et al., 2018). Untested variability in isotopic composition within and between different reefs , bioindicators, and even studies could therefore reduce the reproducibility, or else the comparability of large-scale and long-term monitoring assessments (Pitt et al., 2009; Connolly et al., 2013).

If multiple bioindicators can demonstrate similarly precise and congruent spatial patterns of nutrients over a large-scale gradient, then other taxa, particularly as those from multiple trophic positions, may become useful proxies in areas where macroalgae are scarce, such as on reefs that are dominated by reef-building corals or turf algae (den Haan et al., 2014; Fulton et al, 2019). However, some of these bioindicators may not be directly comparable with others due to the way they take up and process nutrients internally or how other biophysical drivers could potentially influence their signatures (Raimonet et al., 2013; Viana & Bode, 2013; Clausen & Fong, 2016). In addition, species at different trophic levels have different $\delta^{15}\text{N}$ signatures due to isotopic fractionation (Boecklen et al., 2011). This may therefore impact the overall effectiveness of a suite of bioindicators, so additional measures are needed to directly compare their compatibility before they can be used for monitoring programs.

In this study, we investigated the precision and cost-effectiveness of a suite of eight potential bioindicators collected from coral reefs across the Inner Seychelles Islands for measuring nutrient regimes. The specific objectives of the study were to (1) quantify the precision of different bioindicators for measuring stable isotopic and elemental signatures of nitrogen and carbon, (2) determine how much variation exists within bioindicators across different coral reef sites which vary in ecological condition, (3) consider whether there is congruency between selected precise bioindicators based on their nitrogen (N)- and carbon (C)-based measurements, and (4) assess cost-effectiveness of using different bioindicators and the tasks involved.

2. Methods

2.1 Study Sites and Sample Collections

The inner Seychelles islands (43°S, 55°30'E) are comprised of high granitic islands with well-developed carbonate fringing reefs (Littler et al., 1991; Dajka et al., 2019). Bioindicator samples were collected from 21 coral reef sites around the populated islands of Mahé and Praslin, between 11th – 22nd April 2017. These sites have been used as part of a 23-year long-term coral reef monitoring survey, of the reefs of the Inner Seychelles Islands (Suppl. Table 1; Graham et al., 2015; Wilson et al., 2019). The 21 reefs in this study were formed on habitats of either granite, contiguous carbonate or patches that are surrounded by sand or rubble. Twelve of these reefs were defined as “recovering” live coral from a mass bleaching event in 1998, and nine as “regime-shifted” where macroalgae had proliferated (Wilson et al., 2019). However, another mass bleaching event in 2016 caused mass coral mortality on the recovering reefs (Wilson et al., 2019), and so here we define them as “coral-mortality” reefs. Using nitrogen content of brown macroalgae collected from these sites, Graham et al. (2015) also found that nutrient regimes are one of the key determinants of whether a reef can recover or experience a regime shift after a major disturbance like bleaching.

To assess the availability of potential bioindicators, eight replicate 7-m radius point counts were surveyed along the reef slope at each site, and within each point count area, the percent cover of benthic groups such as hard coral, soft coral, macroalgae, sand, rubble, and rock was quantified using eight replicate 10m line-intercept transects (Wilson et al., 2019). Along each transect, the distance of tape occupied by different benthic organisms and substrates was recorded, including live hard coral, soft coral, macroalgae, sponge, cyanobacteria, zoanthids, sand, rubble and rock. For the purpose of this study, the percent cover of dead hard coral and rubble was pooled for an estimate of turf algae per site. Up to ten replicate samples of eight

different bioindicators (i.e. each replicate was a separate individual or sample) were collected haphazardly using SCUBA from within the same area used for the benthic surveys on each reef. However, there were not always ten available replicate samples at all sites, and some reefs had none of some types at all. Bioindicators were selected based on their presence in long-term benthic composition data and their use in previous nutrient enrichment and bioindicator studies (Risk et al., 2001; Fichez et al., 2005; Cooper et al., 2009; Fabricius et al., 2012). Bioindicators included whole fronds of mature foliose brown macroalgae with the apical tips (*Sargassum* sp., Littler et al., 1991; Schaffelke, 1999), filamentous green macroalgae (*Chlorodesmis* sp., Schaffelke, 1999), cyanobacteria (Ford et al., 2018), soft corals (*Sarcophyton* sp., Fleury et al., 2000), turf algal matrix (Graham et al., 2018), sponges (Demospongiae: Ward-Paige et al., 2005; Lamb et al., 2012), and zoanthids (*Palythoa* sp., Leal et al., 2017). For turf algae, branches of dead *Acropora* spp. coral densely covered in turf algal assemblages were broken off and scraped with a scalpel to collect enough material to make up ten replicate samples. Marine sediment (< 4 cm depth; Fichez et al., 2005; Umezawa et al., 2008) which was considered as a non-biological indicator in this study, was also collected to determine nutrient signatures as an important store of nutrients on coral reefs. All samples were frozen at -20°C for up to one month.

2.2 Stable Isotopic and Elemental Analyses

Sample processing and preparation for isotopic analyses were conducted between the Seychelles Fishing Authority laboratory, Victoria, Mahé, Seychelles and Lancaster Environment Centre, Lancaster University, UK. All frozen samples were defrosted, rinsed thoroughly with distilled water and replicate samples were placed in a drying oven for ~48 hr at 60°C.. Once dried, samples were each ground into a fine powder using a ball mill and stored in individual airtight containers at SFA. All dried samples were weighed, alongside the

relevant standards (IAEA 600, cornflour, wheatflour and LEC flour), for stable isotopic analyses at LEC. For bioindicators which contained inorganic carbon material (i.e. calcifying organisms such as soft corals, sponges, and zoanthids), additional acidification was required to remove the inorganic carbonate which can affect carbon-based signatures (Schlacher & Connolly, 2014). ~10g of material was digested in 10% v/v hydrochloric acid (HCl) at room temperature until all constituent carbonate had been removed. Samples were then centrifuged, repeatedly washed until all traces of acidity had been removed, and left to dry prior to analysis for carbon stable isotope composition. The carbon stable isotopic and elemental signatures could not be measured in sediments in this study, because the samples were almost entirely composed of inorganic carbon material, so almost all of the test sediment material dissolved during initial runs of the acidification process. In addition, a subset of all calcified samples were not acidified so that they could be used for nitrogen-based stable isotopic signatures, as acidification can alter $\delta^{15}\text{N}$ signatures in some organisms (Schlacher & Connolly, 2014).

Stable isotopic and elemental analyses for nitrogen stable isotopes ($\delta^{15}\text{N}$), carbon stable isotopes ($\delta^{13}\text{C}$), nitrogen content (%N), carbon content (%C), and C:N Ratio (calculated from dividing the values of %C over %N) were undertaken within the Lancaster Environment Centre stable isotope facility, using an Isoprime100 Isotope Ratio Mass Spectrometer (IRMS) linked to an Elementar VARIO MICROcube Elemental Analyser. Combustion of samples within tin capsules at 950°C yielded N_2 and CO_2 for determination of $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ respectively. Analyses were standardised to AIR (for $\delta^{15}\text{N}$) and VPDB (for $\delta^{13}\text{C}$) using internal reference materials calibrated to international standards. Within-run replication (1 σ) was <0.3 ‰ for $\delta^{15}\text{N}$ and <0.1 ‰ for $\delta^{13}\text{C}$ for both standards and samples.

2.3 Cost-Effectiveness Analyses

To evaluate the cost-effectiveness of each of the techniques used to quantify the nutrient signatures in the eight different bioindicators, the time taken for collection, processing and analysis was calculated as follows. Collection time involved the time taken to search for and retrieve samples from up to 21 sites, where the average time recorded for each dive was ~1 h. Processing time included sample drying, crushing, weighing, and/or acidifying. Drying time represented the time taken to completely dry each sample in the drying oven, while crushing time was the time taken to crush each dried sample into a fine power. For weighing, the average time weighing standards for each mass spectrometric analysis was added to the time taken to weigh each individual sample, and stable isotope analysis time represented the time per analysis. The time taken to acidify each sample of the four calcified bioindicators was also included, though these samples had to be run twice to obtain results for both N and C signatures, with the first subset of samples unacidified, and the second subset acidified. All recorded and calculated times were then standardised to hours (h). The time taken per unit sample was used as a measure of “cost” instead of monetary value in this study, because the methods used to collect, process and analyse them were the same, except for the carbonate-containing samples which needed to be weighed and analysed twice.

2.4 Statistical Analyses

Availability of the bioindicators was assessed in two ways. Firstly, the abundance of the selected groups from the benthic composition data across the 21 sites was averaged and pooled for the two different types of reef state. Secondly, the number of sites that the different bioindicator types were collected from were totalled and categorised according to reef state (i.e. coral-mortality or regime-shifted). The percentage of sites from which each bioindicator was collected, relative to each reef state (i.e. out of 12 for coral-mortality reefs,

and out of 9 for regime-shifted reefs), was calculated, as there were different numbers in each category. The mean and standard deviation of the five nutrient signatures ($\delta^{15}\text{N}$, $\delta^{13}\text{C}$, %N, %C and C:N Ratio) from samples of each bioindicator, collected from up to 21 sites, were then analysed in R (R-Core-Team 2018).

The spatial variation for nutrient signatures of each bioindicator was assessed across all available sites using generalized linear models (GLM). All model fits were inspected for normality using visual plots, and GLMs were used on those with non-normal distributions. A GLM was used to determine the impact of the bioindicator, reef state and individual site on the five nutrient signatures (i.e. the response variables), using the following model for each individual signature:

Model 1: Nutrient Signature ~ Bioindicator + Reef State + Site

Where the nutrient signature was either $\delta^{15}\text{N}$, $\delta^{13}\text{C}$, %N, %C and C:N Ratio, and bioindicator (eight levels), reef state (two levels) and site (up to 21 levels) as fixed factors for each of the five response variables, (C-based signatures in sediment were omitted, as there was no data available). A total of 37 models were therefore run for the overall analysis ($\alpha = 0.05$).

The coefficient of variation (CoV) was used to calculate the overall precision of each bioindicator across all available sites. CoV is the ratio of the sample standard deviation to the same mean, for a given set number of data points, and was used in this study because it is a unitless measure of variation, which is useful when testing the statistical effectiveness (i.e. precision) of the signatures across the different bioindicators. High precision is defined in this study as a small standard deviation compared to the mean, which increases the ability to detect statistical significance, both between the replicate samples of each bioindicator collected at each site, and over all the sites from which each bioindicator was collected. Low

precision, conversely, is a large standard deviation compared to the mean (Conquest, 1983). Though there is not one set standard in the literature, it is generally assumed that values of $\text{CoV} < 10$ can be regarded as “precise”. CoV was calculated from the raw measurements detected in the replicate samples of each bioindicator collected from individual sites. Following this, the CoV of the N- and C-based signatures were compared across all the sites from which each bioindicator was collected with five linear models (Model 2), which were run separately for each nutrient signature:

Model 2: $\text{CoV} \sim \text{Bioindicator} + \text{Reef State} + \text{Site}$

Where CoV was the CoV value for $\delta^{15}\text{N}$, $\delta^{13}\text{C}$, %N, %C and C:N Ratio, and Bioindicator (eight levels), reef state (two levels) and site (up to 21 levels) were the fixed factors. The overall mean and standard deviation for the CoV each bioindicator were also summarised in box-plots.

A principal components analysis (PCA) (PRIMER-E Ltd, V.6.1.5, Plymouth, UK) based on a Bray–Curtis similarity matrix was used to visualise the similarities between averaged values of the five different nutrient measurements and the different bioindicators as a way of assessing the level of congruency of the bioindicators (Clarke & Warwick, 2001). The selection of a subset of bioindicators for this analysis (brown macroalgae, green macroalgae and zoanthids) was based on their level of precision, and the number of sites used, out of 21, depended upon the availability of each of these three indicators. Therefore nine sites were selected, as they had sufficient replicates of all three bioindicators to compare across sites ($n=4$), and the nutrient measurements were averaged at site level to compensate for the varying numbers of replicate samples available at each site. However, for C-based signatures, zoanthid samples from one site could not be acidified due to limited material so for these, eight sites were used. A correlation matrix was also constructed to assess the different

correlation values between the three selected indicators, where a p-value < 0.05 was considered significant.

To statistically assess the cost-effectiveness of each bioindicator, another GLM was used (as the data was not normally distributed) to compare the average times taken (per sample per bioindicator) for (a) collecting from the field, (b) drying and crushing of samples, (c) weighing and preparing samples (i.e. acidification) for isotopic analyses, and (d) running isotopic analyses. In this model, “Time” was the response variable, and “Bioindicator” and “Task” were the fixed factors (eight and two levels in each factor, respectively):

Model 3: Time ~ Bioindicator * Task

The interaction between these two fixed factors in Model 3 was also analysed to determine whether the “Bioindicator” (eight levels), “Task” (4-5 levels, depending on whether or not the bioindicator was acidified), or the interaction between them affects the time per unit sample. Reef State was also used as a fixed factor (with two levels) during initial statistical analyses, but was not included in this study as it showed no significant effect.

3. Results

3.1 Sample Collection and Benthic Cover

Across the 21 sites, a total of 150 samples of brown macroalgae (*Sargassum* sp.), 91 green macroalgae (*Chlorodesmis* sp.), 103 cyanobacteria, 59 soft corals, 112 sponges, 134 zoanthids (*Palythoa* sp.), 171 turf algal assemblages, and 204 sediment samples were collected. Availability of bioindicator varied between regime-shifted and coral-mortality reefs, as did the percentage of sites within these two categories where they were present (Table 1). Average cover of *Sargassum* sp. was significantly higher at the regime-shifted sites where it was an order of magnitude greater than on the coral-mortality sites. As such, there were specimens available at 100% of the regime-shifted sites, whereas they were only found at 58% of regime-shifted reefs. There was a similar percent cover of sediment across sites (along the line-intersect transect) regardless of reef state, and sediment samples were collected from all 21 sites. Percent cover of turf algae on coral-mortality reefs was 32.8 ± 23.8 %, compared to 12.2 ± 8.11 % on regime-shifted reefs, but still had 100% availability in both reef states. Cyanobacteria, soft coral and sponge all had higher percent cover and were also present on a higher percentage of coral-mortality sites than on regime-shifted ones.

Table 1. Summary table for percent cover (% cover) of candidate bioindicators (BM = brown macroalgae; CYB = Cyanobacteria; GM = Green Macroalgae; SED = Sediment; SC = Soft Coral; TA = Turf Algae; ZO = Zoanthid) from the line-intercept transect surveys at 21 coral reefs around the Inner Seychelles Islands. Percentage of Sites represents the percentage of sites relative to the total number in each reef state (out of n=12 for “coral-mortality” reefs versus n=9 “regime-shifted” reefs). Mean \pm S.D for percent cover.

| Bioindicator | Regime-Shifted Reefs (n=9) | | Coral-Mortality Reefs (n=12) | |
|--------------------------|-------------------------------|-------------------------|---------------------------------|-------------------------|
| | Mean \pm S.D. (%) | Percentage of Sites (%) | Mean \pm S.D. (%) | Percentage of Sites (%) |
| <i>Sargassum</i> (BM) | 36.9 \pm 20.3 | 100 | 2.7 \pm 8.47 | 58 |
| Cyanobacteria (CYB) | 1.2 \pm 2.8 | 44 | 2.5 \pm 5.0 | 75 |
| <i>Chlorodesmis</i> (GM) | 0.2 \pm 0.3 | 89 | 0.3 \pm 0.4 | 25 |
| Soft Coral (SC) | 0.1 \pm 0.8 | 11 | 1.2 \pm 2.5 | 67 |
| Sediment (SED) | 6.7 \pm 3.4 | 100 | 9.52 \pm 11.5 | 100 |
| Sponge (SP) | 0.00* | 56 | 1.4 \pm 2.1 | 75 |
| Turf Algae (TA) | 12.2 \pm 8.1 | 100 | 32.8 \pm 23.8 | 100 |
| <i>Palythoa</i> (ZO) | 0.2 \pm 0.4 | 67 | 1.3 \pm 1.0 | 100 |

3.2 Spatial Variation of Nutrient Signatures in Bioindicators

The type of bioindicator had variable effects on each of the five nutrient signatures. Overall, brown and green macroalgae (BM and GM, respectively) not only had lower average $\delta^{15}\text{N}$ signatures than the other indicators, but they also had the smallest variations in signatures across all of their sites (5.58 ± 0.82 and $5.33 \pm 0.45\text{‰}$, respectively. *Fig. 1a*). Bioindicators representing higher trophic levels, such as sponges (SP), soft corals (SC), and zoanthids (ZO) (7.51 ± 0.67 ; 7.61 ± 1.27 , and $9.08 \pm 0.88\text{‰}$, respectively) had more enriched average $\delta^{15}\text{N}$ signatures, as did sediment (SED) ($9.61 \pm 1.41 \text{‰}$). After acidification, the four bioindicators that contained inorganic carbon (soft corals, sponges, and turf algae (TA)) showed similar signatures of $\delta^{13}\text{C}$ on average (-16.3 ± 1.29 ; -17.4 ± 0.38 ; and $-18.5 \pm 3.16 \text{‰}$, respectively), though it was less negative in zoanthids ($-13.7 \pm 0.88 \text{‰}$). The two types of macroalgae also differed (BM: -16.2 ± 1.58 , and GM: $-21.3 \pm 0.96 \text{‰}$) whereas cyanobacteria (CYB) ($-21.3 \pm 3.36 \text{‰}$) was similar to green macroalgae (*Fig. 1b*).

Turf algae had a similar average signature for %N ($1.53 \pm 0.45\text{‰}$) relative to brown macroalgae ($1.10 \pm 0.18 \text{‰}$) but green macroalgae had a much higher value ($4.32 \pm 0.48 \text{‰}$), which was even higher than cyanobacteria ($3.31 \pm 1.25 \text{‰}$). The N content of brown macroalgae was also most similar to zoanthids ($1.06 \pm 0.22 \text{‰}$). N content was also much lower in sediment ($0.05 \pm 0.11 \text{‰}$) (*Fig. 1c*). There was much higher C content in green macroalgae than in the other bioindicators ($42.2 \pm 2.40 \text{‰}$), followed by brown macroalgae ($31.0 \pm 1.41 \text{‰}$), and cyanobacteria ($28.7 \pm 5.52 \text{‰}$). Zoanthids had the lowest %C (11.2 ± 2.74) (*Fig. 1d*). Brown macroalgae had higher C:N Ratio signatures with a large range due to high %C content and low %N content (28.8 ± 4.99). The other five groups were quite similar to one another, with the exception of sponge (0.85 ± 0.11) (*Fig. 1e*).

The GLMs showed that the type of bioindicator had a strong influence on the variability of nutrient signatures, with significance evident across almost all signatures. However, both types of macroalgae were statistically similar for $\delta^{15}\text{N}$, as were brown macroalgae, turf algae and zoanthid for %N (Suppl. Table 2). However, the effect of reef state varied among both bioindicators and nutrient signatures. For instance, differences in $\delta^{15}\text{N}$ signatures in BM ($p=0.0002$), CYB ($p=0.002$), GM ($p<0.0001$), SED ($p=0.01$), TA ($p=0.02$) and ZO ($p<0.0001$) were significant, whereas the difference in %N for GM between reef states was not ($p=0.93$). Reef state was also significantly different for $\delta^{13}\text{C}$ in cyanobacteria ($p=0.002$), green macroalgae ($p < 0.0001$), sediment ($p=0.01$), turf algae ($p=0.02$) and zoanthids ($p<0.0001$). For %N, reef state also significantly differed in BM ($p < 0.0001$), CYB ($p<0.0001$) and ZO ($p=0.04$). For %C, reef state differed significantly for CYB ($p<0.0001$) and ZO ($p=0.01$), and for C:N Ratio, only BM ($p=0.04$) and TA ($p=0.0002$) differed significantly.




Figure 1. Box (median and 50% quantile) and whisker (95% quantile) plots of the variation of the average values of nutrient signatures measured in the eight bioindicators for (a) $\delta^{15}\text{N}$, (b) $\delta^{13}\text{C}$, (c) %N, (d) %C and (e) C:N Ratio from up to 21 reefs. Each black dot represents the average value from an individual site that each bioindicator was collected from to also show the spread of variation within each bioindicator (BM = Brown Macroalgae; CYB = Cyanobacteria; GM = Green Macroalgae; SED = Sediment; SC = Soft Coral; SP = Sponge; TA = Turf Algae, and ZO = Zoanthid).

3.3 Precision of Bioindicators

The precision of the bioindicators was assessed using CoV, as this standardised the nutrient signatures between bioindicators (including the non-biological indicator sediment) and controlled for differences in isotopic fractionation in measurements, particularly between trophic levels. Green macroalgae had the lowest and most consistent CoV within and across reefs, and therefore the highest precision for all N-based nutrient measurements ($\delta^{15}\text{N}$: 2.47 ± 0.95 ; %N: 7.53 ± 4.29 ; C:N Ratio: 5.76 ± 5.39), however this pattern was not as distinct for C-only signatures ($\delta^{13}\text{C}$: -1.87 ± 1.06 and %C: 3.60 ± 1.67) (*Fig. 2*). This was closely followed by brown macroalgae ($\delta^{15}\text{N}$: 4.68 ± 1.33 ‰; $\delta^{13}\text{C}$: -6.03 ± 3.12 ; %N: 11.3 ± 4.07 ; %C: 4.07 ± 1.12 , and C:N Ratio: 9.92 ± 3.75). Turf algal assemblages had much more variable average signatures for all five measures, especially those that were N-based ($\delta^{15}\text{N}$: 8.30 ± 4.90 ; $\delta^{13}\text{C}$: $-5.14 \pm$; %N: 20.5 ± 20.1 ; %C: 9.54 ± 10.6 , and C:N Ratio: 10.6 ± 10.3).

Zoanthids had lower average CoV values for N-based signatures than higher trophic organisms and were more similar to the two macroalgal types ($\delta^{15}\text{N}$: 2.98 ± 1.20 , and %N: 14.3 ± 5.52), as well as for $\delta^{13}\text{C}$ (-5.14 ± 2.43), though the CoV values for both %C and C:N Ratio were much higher than for any of the other bioindicators (11.8 ± 8.57 and 20.0 ± 24.1 , respectively). The other higher trophic level organisms, such as soft corals ($\delta^{15}\text{N}$: 6.26 ± 4.87 ; $\delta^{13}\text{C}$: -6.20 ± 1.86 ; %N: 30.4 ± 17.6 ; %C: 17.4 ± 12.2 , and C:N Ratio: 11.6 ± 8.68) and sponges ($\delta^{15}\text{N}$: 6.82 ± 5.24 ; $\delta^{13}\text{C}$: -1.44 ± 1.08 ; %N: 20.0 ± 10.3 ; %C: 7.24 ± 3.94 , and C:N Ratio: 7.58 ± 12.1) showed inconsistent levels of precision across the five signatures. Though sediment had similar precision for $\delta^{15}\text{N}$ to the other candidates (7.97 ± 3.90), it had the highest range of CoV values for %N (17.4 ± 40.2) (*Fig. 2a*).

Overall, the CoV analyses showed that both brown and green macroalgae had low average CoV values for N-based signatures, as well as small variations in CoV across the sites. In addition, while the C-based signatures were more variable for zoanthids, the N-based results were more precise compared to the other higher-trophic bioindicators. There was also no overall significant effect of reef state or site-level variation on CoV for any of the five nutrient signatures, suggesting that precision did not vary over different spatial scales or between the coral-mortality and regime-shifted reefs. The statistical models showed variable patterns for each nutrient signature type across the eight bioindicators, however for %C and C:N Ratio, zoanthids were the only bioindicator that significantly differed from brown macroalgae due to its high variation (Suppl. Table 3).






Figure 2. Box (median and 50% quantile) and whisker (95% quantile) plots of the spread of the coefficient of variation (CoV) of the eight bioindicators for (a) $\delta^{15}\text{N}$, (b) $\delta^{13}\text{C}$, (c) %N, (d) %C and (e) C:N Ratio up to 21 reefs (mean \pm S.D.). Each black dot represents the average CoV from the individual sites from which each bioindicator was collected to also show the spread of variation within- and among sites (BM = Brown Macroalgae; CYB = Cyanobacteria; GM = Green Macroalgae; SED = Sediment; SC = Soft Coral; SP = Sponge; TA = Turf Algae, and ZO = Zoanthid). CoV for each nutrient measurement in each bioindicator collected from each site was calculated by the ratio of standard deviation to the mean of a given number of replicate data points (i.e. up to 5 samples per indicator per site).

3.4. Congruency of Bioindicators

A principal components analysis (PCA) was used to assess congruency between the three selected bioindicators. Brown and green macroalgae had low correlation, especially for signatures of N, while zoanthids had no significant relationships with either macroalgae. There were weak positive relationships between N-based signatures of green and brown macroalgae (Table 2), but these explain <40% of the variance and are not significant at $\alpha = 0.05$ (Fig. 3). This was also shown by Pearson's correlation analyses between the different combinations of bioindicators (Table 2). The two types that showed the highest similarity for N-based signatures were between brown and green macroalgae for C:N Ratio measurements ($r^2 = 0.61$), closely followed for those of %N ($r^2 = 0.60$) and $\delta^{15}\text{N}$ ($r^2 = 0.55$) signatures, though none of these were significantly correlated. However, the highest similarity for C-only signatures was between %C of brown and green macroalgae ($r^2 = 0.81$), but was very low for $\delta^{13}\text{C}$ ($r^2 = 0.041$) (Table 2).

Table 2. Pearson's correlation analyses between the three selected bioindicators (brown macroalgae versus green macroalgae; brown macroalgae versus zoanthids; green macroalgae versus zoanthids) to determine amount of correlation between them (correlation coefficient) The significance level for the p-values is $\alpha = 0.05$.

| Bioindicator | $\delta^{15}\text{N}$ | $\delta^{13}\text{C}$ | %N | %C | C:N Ratio |
|--------------|-----------------------|-----------------------|----------|----------|-----------|
| BM vs. GM | 0.55 | 0.041 | 0.60 | 0.81 | 0.61 |
| | (p=0.12) | (p=0.92) | (p=0.09) | (p=0.02) | (p=0.08) |
| BM vs. ZO | 0.10 | 0.11 | 0.18 | -0.005 | 0.07 |
| | (p=0.79) | (p=0.80) | (p=0.64) | (p=0.99) | (p=0.68) |
| GM vs. ZO | 0.28 | 0.64 | 0.23 | -0.23 | -0.36 |
| | (p=0.47) | (p=0.09) | (p=0.55) | (p=0.58) | (p=0.34) |



525






Figure 3. Principal Components Analyses (PCA) quantifying congruency between a selection of bioindicators ($n=3$) (BM = Brown Macroalgae; GM = Green Macroalgae; ZO = Zoanthids) all present at a subset number of sites ($n=9$) for measurements of (a) $\delta^{15}\text{N}$, (b) $\delta^{13}\text{C}$, (c) %N, (d) %C and (e) C:N Ratio.

3.5 Cost-Effectiveness of Bioindicators

The time taken for the whole process, from collection to stable isotopic analyses, per unit sample, differed among the eight bioindicators (Table 3; Suppl. Table 4). The GLMs suggested that both bioindicator and task can have a significant effect on the time taken, per sample, to use each bioindicator for capturing measure nutrient regimes, but reef state does not. Overall, it took a similar amount of time to collect the two macroalgae and cyanobacteria, whereas soft corals, sponges, turf algae and zoanthids took significantly longer to find. Sediment, in contrast, took the least time overall to find and collect (Table 3). Each task differed significantly as well, with “Drying and Crushing” taking the most time to complete and “Field Collection” took the least time, but significance varied between the bioindicators. The time taken to process the four calcified bioindicators was much greater, because each sample of these indicators required the additional step of “Acidification”.

Table 3. Summary of the mean time taken (per unit sample, per hour) for each task undertaken to process each bioindicator for the cost-effectiveness. *Acidifying only includes the four bioindicators that were acidified, and thus weighed and analysed in the mass spectrometer. Significance Level is $p < 0.05$. Normality inspected using visual plots. Mean \pm S.D.

| BIOINDICATOR | FIELD COLLECTION | DRYING & CRUSHING | ACIDIFICATION | WEIGHING | STABLE ISOTOPIC ANALYSES |
|------------------------------|---------------------------------------|------------------------------------|--------------------------------------|----------------------------------|---|
| Brown Macroalgae (BM) | 0.038 \pm 0.04 ($p < 0.0001$) | 24.8 \pm 0.5 ($p < 0.0001$) | - | 1.5 \pm 0.01 (N/A) | 0.18 \pm 0.03 ($p < 0.0001$) |
| Cyanobacteria (CYB) | 0.35 \pm 0.4 ($p = 0.52$) | 23.2 \pm 1.4 ($p < 0.0001$) | - | 1.5 \pm 0.03 (N/A) | 0.21 \pm 0.1 ($p = 0.93$) |
| Green Macroalgae (GM) | 0.078 \pm 0.08 ($p = 0.86$) | 24.1 \pm 0.005 ($p = 0.26$) | - | 1.5 \pm 0.01 (N/A) | 0.17 \pm 0.05 ($p = 0.98$) |
| Soft Coral (SC) | 0.25 \pm 0.3 ($p = 0.001$) | 22.2 \pm 1.2 ($p < 0.0001$) | 0.17 \pm 0.001 ($p < 0.0001$) | 3.1 \pm 0.06 ($p = 0.95$) | 0.48 \pm 0.2 ($p = 0.004$) |
| Sediment (SED) | 0.015 \pm 0.003 ($p = 0.0002$) | 22.7 \pm 1.2 ($p = 0.05$) | - | 0.14 \pm 0.02 (N/A) | 0.25 \pm 0.1 ($p < 0.0001$) |
| Sponge (SP) | 0.24 \pm 0.3 ($p = 0.0006$) | 22.6 \pm 1.3 ($p < 0.0001$) | 0.17 \pm 0.0 ($p < 0.0001$) | 3.1 \pm 0.00 ($p = 0.89$) | 0.37 \pm 0.07 ($p = 0.002$) |
| Turf Algae (TA) | 0.03 \pm 0.04 ($p = 0.0002$) | 24.6 \pm 0.5 ($p = 0.0003$) | 0.17 \pm 0.002 ($p < 0.0001$) | 3.0 \pm 0.02 ($p = 0.98$) | 0.34 \pm 0.08 ($p = 0.0005$) |
| Zoanthids | 0.18 \pm 0.2 ($p < 0.0001$) | 23.0 \pm 1.5 ($p < 0.0001$) | 0.17 \pm 0.0 ($p < 0.0001$) | 3.0 \pm 0.00 (N/A) | 0.41 \pm 0.03 ($p = 0.0004$) |

Although the time taken per sample to collect each bioindicator from the field did not differ between reef states, the availability of samples on the different reef did (*Table 1*). There was a strong negative correlation between average time taken per sample to collect and the percentage of sites from which each indicator was available on regime-shifted reefs (relative to the total number of sites, i.e. $n=9$) ($r^2 = 0.94$), whereas there was a very weak negative relationship between average time taken and sample availability on coral-mortality sites ($r^2 = 0.15$; $n=12$) (*Fig. 4*). This suggests that although the time taken varied more among bioindicators on regime-shifted reefs (i.e. it took over an hour, on average, to find one sample of soft coral), it is a better predictor for finding specific bioindicator(s) on sites dominated by macroalgae. For coral-mortality reefs, in contrast, the times among bioindicators were more similar, but sample availability was more variable. Brown macroalgae had similar collections times between reef states (regime-shifted: 0.01 ± 0.01 ; coral-mortality: 0.07 ± 0.05 h), but

there was 100% availability on regime-shifted sites relative to 58% on coral-mortality sites. Turf algae and sediment, in contrast, not only had 100% availability on both reef states, but they took the least time to collect .




Figure 4. The relationships between the average time taken, per unit sample (h) and the availability of samples on both reef states. Each individual point in red represent the total average time, per sample, for the eight bioindicators collected from regime-shifted sites versus the percentage of sites they were available to collect at (n=12), and the individual point in blue represented each indicator from coral-mortality sites. $r^2 = 0.94$ on regime-shifted reefs, and $r^2 = 0.15$ on coral-mortality reefs. BM = Brown Macroalgae; CYB = Cyanobacteria; GM = Green Macroalgae; SED = Sediment; SC = Soft Coral; SP = Sponge; TA = Turf Algae, and ZO = Zoanthid.

4. Discussion

The principal aims of this study were to identify precise, cost-effective, and congruent bioindicators for capturing nutrient regimes on coral reefs, particularly over those in different ecological conditions. Overall, nutrient signatures of brown macroalgae, green macroalgae and zoanthids were considered to meet these criteria, relative to the other candidates. While the macroalgae were more consistent indicators for reefs that have undergone a regime shift, zoanthids were more common on both types of reef state. Turf algae and sediment took the least time to collect and were also the most abundant and available samples across the 21 reefs studied, regardless of reef state, but their utility as indicators is limited by their highly variable CoV values. There was low congruency between the three most precise indicators (brown macroalgae, green macroalgae and zoanthids), which suggested that physiological processing of nutrients within each bioindicator has a greater influence on N- and C-based signatures than its local environment. Congruency between multiple taxa could be improved by either choosing a suite of indicators from the same functional group, such as macroalgae with comparable nutrient uptake mechanisms, or by tracing the accumulation of nutrient signatures across different trophic levels from the same food chain.

4.1 Spatial Variation, Precision and Congruency of Nutrient Signatures in Bioindicators

The N- and C-based nutrient signatures of the bioindicators in the current study appear typical of measurements reported in the literature (Atkinson & Smith, 1983; Smit, 2001). For instance, the range of absolute values of $\delta^{15}\text{N}$ signatures in all of the bioindicators are quite consistent (5 – 10 ‰), though they are slightly high relative to other marine systems (Sigman & Casciotti, 2001). In addition, the $\delta^{13}\text{C}$ signatures reflect that of a carbonate-dominated system, which for instance lies within the range of -10 to -30‰ for most marine macrophytes (Smit, 2001; Raven et al., 2002). The N-based signatures also follow trophic status whereby

those organisms at higher trophic levels are relatively more enriched than those of primary producer status (Boecklen et al., 2011; Lamb et al., 2012)..

Spatial variation of the different nutrient signatures, both within- and among-reefs, varied widely across the inner Seychelles. The N-based signatures also showed a significant difference between coral-mortality and regime-shifted reefs for a number of the bioindicators, including $\delta^{15}\text{N}$ in the two macroalgae and zoanthids, whereas signatures tended to be more similar across sites for the C-based signatures. Being able to capture variability in nutrient regimes, especially across different spatial scales or even different reef states, is another important aspect of a good bioindicator (Cooper et al., 2009), so this study provides supporting evidence that $\delta^{15}\text{N}$ and %N are particularly effective proxies of nutrient regimes (Lin & Fong, 2008). For instance, Littler et al., (1991) found that nutrient concentrations in a number of algal species were generally higher on reefs around the high granitic, populated islands like Mahe and Praslin, relative to the low, remote carbonate atolls in the wider Seychelles Archipelago. In a related study in Vaughan et al. (2021), the use of macroalgal $\delta^{15}\text{N}$ helped to determine that the dead coral tissue released into the water column after the 2016 coral bleaching event in the Seychelles may have been subsequently taken up and retained by *Sargassum* on the coral-mortality reefs. However, the high variability shown across nutrient signatures in the current study, particularly in $\delta^{15}\text{N}$, may not be solely due to differences in local sources of nutrients. Other studies, for example, have found that differences in signatures are not always consistent with distinct sources of nutrient loads (i.e. in areas with known anthropogenic run-off), which implied that external inputs are not always the cause of variations in nutrient regimes captured in bioindicators (Raimonet et al., 2013; Viana & Bode, 2013).

There were discrepancies found in some of the signatures even between different primary producers in this study, such as between brown (*Sargassum* sp.) and green macroalgae (*Chlorodesmis* sp.). For instance, although they had similar $\delta^{15}\text{N}$ values across the sites, the other four signatures varied on average between these two bioindicators, particularly for %N, which was much higher in green macroalgae, although it was similar between reef states (Fig. 2a&c). This could be because nitrogen content in *Chlorodesmis* is affected by both biological nutrient uptake mechanisms and environmental factors (Fong et al., 2001; Raimonet et al., 2013; Viana & Bode, 2013; Clausing & Fong, 2016), and therefore do not reflect either inorganic concentrations or the $\delta^{15}\text{N}$ of their surrounding environment (Viana & Bode, 2013). Slower-growing algal species like *Chlorodesmis* have a greater capacity for internal nutrient storage so are not as nutrient-limited, and therefore are less responsive to fluctuations in nutrients as other, more opportunistic species like *Sargassum* (Schaffelke., 1999; García-Seoane et al., 2018a&b).

Zoanthids are positioned at a higher trophic level than benthic algae so their nutrient signatures tend to fractionate and become more enriched (Fig. 1a; Zanden & Rasmussen, 2001; Fox et al., 2018). There has been little research into zoanthids as potential indicators of nutrient runoff (Leal et al., 2017), but Costa Jr. et al. (2008) found that phosphorus and silica water concentrations had positive effects on both algal and zoanthid growth, and negative effects on coral cover. However, unlike primary producers, zoanthids have to balance auto- and heterotrophic processes for acquiring sources of C and N (Smit, 2001; Leal et al., 2017) because, like scleractinian corals, they have photosynthetic symbionts in their tissues (Hoegh-Guldberg et al., 2004; Fox et al., 2018). This could explain the large variations in %C and C:N Ratio, both within- and among-reefs in this study (Fig. 2d &e; Suppl. Table 2), as they represent the combined signatures from both host and symbiont (Leal et al., 2017).

671

672 Even though the three most precise bioindicators (brown macroalgae, green macroalgae and
673 zoanthids) all showed significant differences in $\delta^{15}\text{N}$ between the two reef states for the
674 spatial variation analyses, their CoV values did not. This suggests that these bioindicators are
675 not only consistently precise among reefs and reef states, but are also able to detect
676 differences in nutrient regimes across the same areas, which is why $\delta^{15}\text{N}$ is such a versatile
677 tool for monitoring water quality (Costanzo et al., 2001; Lin & Fong, 2008). However, when
678 compared directly, the congruency among these three bioindicators was relatively low. This
679 could be due to the differences in nutrient processing between the different bioindicators.
680 Congruency is important, as a single-species approach may result in an underestimation of
681 spatial patterns in nutrient regimes (Linton & Warner, 2003), and it has been shown across
682 multiple taxa in previous studies (Connolly et al., 2013), but these studies were also
683 conducted along strong nutrient gradients (i.e. with increasing distance from a sewage outfall)
684 (Fernandes et al., 2012). This suggests that in the current study, the biological mechanisms of
685 individual species may have outweighed the effect of environmental factors on their isotopic
686 and elemental signatures.

687

688 The other (bio)indicators included in this study were found to have variable and inconsistent
689 nutrient signatures across sites and the two reef states, which was why they were not included
690 in the congruency analyses. Like macroalgae, turf algal assemblages and cyanobacteria are
691 primary producers that not only take up and utilise bioavailable nutrients but are becoming
692 more prevalent on reefs across a range of reef states, particularly following a disturbance (den
693 Haan et al., 2016; Zaneveld et al., 2016, Ford et al., 2018). However, this study showed that
694 both bioindicators had variable precision among the five nutrient signatures with no clear
695 spatial patterns between reefs, which implied they were also more influenced by biological

factors (i.e. multiple species within the turf assemblage) than their local environment (Steneck & Dethier, 1994; Raimonet et al., 2013). Similarly to zoanthids, soft corals can also harbour symbionts (Fleury et al., 2000; Risk, 2014; Williams et al., 2018), and while sponges are not photosynthetic, they do have symbiotic relationships with cyanobacteria, which is reflected in their $\delta^{13}\text{C}$ signatures (Smit, 2001; Lamb et al., 2012). Sediments can also capture a range of nutrients within a reef, which can be resuspended within local biogeochemical cycles through various biophysical factors and thus provide an additional source (Fabricius, 2005; Umezawa et al., 2008). However, some studies have found sediments to be an overall poor indicator (Fichez et al., 2005). In the current study, for instance, very little N was detected in the subsamples of sediment analysed even before acidification, so the low precision calculated for it was more likely due to random error than environmental factors, and so was not comparable for either N- or C-based signatures.

4.2 Cost-Effectiveness of Bioindicators

Cost-effectiveness is often mentioned as an important criteria in previous bioindicator studies (Fichez et al. 2005; Cooper et al., 2009; Risk et al., 2001). However, analyses are rarely conducted to quantify these in ecological studies (Drummond & Connell, 2008; Bal et al., 2020) even though the “cost” of any particular indicator can be affected by various different factors. For instance, the average time taken to collect an individual sample from a study site depended upon its availability and/ or abundance, which is why there was a significant difference in collection time with reef state. While it only took ~1 to 2 minutes on average to collect samples of turf algae and sediments from each site, regardless of ecological condition, it took significantly less time to collect brown macroalgae from regime-shifted reefs than it did on coral-mortality reefs. Differences in availability on those reefs could be influenced by

nutrient loads, abundance of herbivores, depth, structural complexity, and juvenile coral cover (Graham et al., 2015; Dajka et al., 2019). The findings of both the sample collection and the line-intercept survey of benthic cover at the 21 sites illustrated the importance of considering the local abundance of a bioindicator when assessing nutrient regimes (Cooper et al., 2009; Fabricius et al., 2012). For instance, turf algae and sediments were ubiquitous at all sites, so could be considered as more “cost-effective” in terms of sampling availability and abundance. However, as turf algae are composed of an assemblage of varying functional groups, and there was very little N detected in sediment, it is difficult to interpret results for nutrient signatures from either bioindicator, and therefore to rely on them for capturing nutrient regimes precisely, despite their widespread abundance.

4.3 Future Directions in Bioindicator Research

This study investigated novel ways of assessing potential bioindicators for monitoring programs across coral reefs under different ecological states. However precision and effectiveness of bioindicators used in this study could be improved, even if these improvements will increase costs. For instance, to reduce the CoV of turf algal assemblages, cyanobacteria, and symbiotic organisms, future studies could isolate and individually measure the different functional groups within assemblages (Steneck & Dethier, 1985), individual strains of cyanobacteria (Thacker & Paul, 2001), or the host and symbiont fractions in zoanthids and soft corals (Hoegh-Guldberg et al., 2004; Leal et al., 2017) so that the nutrient signatures of each group can be measured and interpreted separately. Conversely, such techniques will increase the time taken to process and analyse samples, and thus will increase their “costs” as a bioindicator.

It was also difficult to determine the accuracy of the bioindicator nutrient signatures, as there is little reference data for nutrient levels around the inner Seychelles Islands, even from seawater samples, and especially at the spatio-temporal scales required for this study. Further research should therefore also investigate the accuracy of cost-effective bioindicators such as macroalgae for capturing either natural or anthropogenic sources by additionally measuring stable isotopic signatures of potential point sources (Costanzo et al., 2001; Dailer et al., 2010; Fernandes et al., 2012; den Haan et al., 2012).. Another approach could entail building up a suite of relatively similar bioindicators by focusing on specific functional group(s), appropriately matched to the scale of the ecological process being investigated (Fong & Fong, 2014). If this option is not possible, for instance, when a group of congruent bioindicators (i.e. fleshy macroalgae) is only found on reefs in a certain ecological state, then nutrient signatures could be compared across a suite of bioindicators to see the accumulation of this energy source across different trophic levels within the same food chain (Smit, 2001; Pitt et al., 2009; Connolly et al., 2013; Kürten et al., 2014; Graham et al., 2018).

5. Conclusion

In conclusion, the stable isotopic and elemental signatures of fleshy macroalgae were found to be precise and cost-effective bioindicators across coral reefs in the inner Seychelles, as primary producers with widespread distribution and consistent measurements within their tissues. If the precision of bioindicators can be increased, it would provide additional opportunities to determine differences in bioavailable nutrient regimes between reefs. This could be particularly useful in remote coastal areas where environmental monitoring efforts to assess the local anthropogenic impacts of coastal run-off and excessive nutrient loads on coral reefs are currently limited, but would be highly beneficial to assessing overall ecosystem health. If remote reefs have been subjected to any large disturbance, such as a

mass bleaching event, having precise and cost-effective bioindicators to detect whether any areas have excessive nutrient loads, could enable better-informed efforts to improve water quality and mediate coral recovery potential.

6. Author Contributions

Eleanor Vaughan: Conceptualisation, Data curation, Formal analysis, Investigation, Methodology, Visualisation, Writing – original draft, Writing – review and editing. **Nicholas Graham:** Conceptualisation, Funding acquisition, Project administration, Resources, Supervision, Writing – review and editing. **Shaun Wilson:** Funding acquisition, Investigation, Writing – review & editing. **Peter Wynn:** Methodology, Resources, Supervision. **Gareth Williams:** Supervision, Writing – review and editing. **Phillip Barker:** Supervision, Writing – review & editing.

7. Declaration of Interest Statement

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

8. Funding

The project was funded via grants from the Royal Society (UF140691 and RG150710).

9. Acknowledgements

We thank Seychelles Fishing Authority, Seychelles Marine Parks Authority and Jan Dajka for field and laboratory assistance in the Seychelles, and David Hughes and Jon Crosse from Lancaster Environment Centre for extensive help with preparations for fieldwork and with the subsequent stable isotopic and elemental analyses. James Robinson and Jeneen Hadj-Hammou gave advice on statistical analyses.

7. References

- Atkinson, M.J. and Smith, S.V. (1983) C:N:P ratios of benthic marine plants, *Limnology and Oceanography*, 28(3): 568-574
- Bal, P., Rhodes, J.R., Carwardine, J., Legge, S., Tulloch, A., Game, E., Martin, T.G., Possingham, H.P., & McDonald-Madden, E. (2020) How to choose a cost-effective indicator to trigger conservation decisions? *Methods in Ecology and Evolution*, 12: 520-529
- Boecklen, W.J., Yarnes, C.T., Cook, B.A., and James, A.C. (2011) On the use of stable isotopes in trophic ecology. *Annu. Rev. Ecol. Evol. Syst.*, 42: 411-440
- Briand, M.J., Bonnet, X., Goiran, C., Guillou, G. and Letourneur, Y. (2015) Major sources of organic matter in a complex coral reef lagoon: identification from isotopic signatures ($\delta^{13}\text{C}$ and $\delta^{15}\text{N}$). *PLoS ONE*, 10(7): e0131555, doi:10.1371/journal.pone.0131555
- Burkepile, D.E. and Hay, M.E. (2006) Herbivore vs. nutrient control of marine primary producers: context-dependent effects, *Ecology*, 87(12): 3128-3139

815 Cinner, J.E., Maire, E., Huchery, C., MacNeil, M.A., Graham, N.A.J., Mora, C. ... and Mouillot, D.
816 (2018) Gravity of human impacts mediates coral reef conservation gains, *PNAS*, 115(27):
817 E6116-E6125

818 Clarke K.R. and Warwick R.M. (2001) *Change in Marine Communities: An Approach to Statistical*
819 *Analysis and Interpretation*, 2nd ed. PRIMER-E: Plymouth, UK, 38pp.

820 Clausing, R.J., and Fong, P. (2016) Environmental variability drives rapid and dramatic changes in
821 nutrient limitation of tropical macroalgae with different ecological strategies. *Coral Reefs*, 35:
822 669-680

823 Connolly, R.M., Gorman, D., Hindell, J.S., Kildea, T.N. and Schlacher, T.A. (2013) High congruence
824 of isotope sewage signals in multiple marine taxa. *Mar. Pollut. Bull.*, 71: 152-158

825 Conquest, L.L. (1983) Assessing the statistical effectiveness of ecological experiments: utility of the
826 coefficient of variation, *Intern. J. Environmental Studies*, 20: 209-221

827 Cooper, T.F., Gilmour, J.P. and Fabricius, K.E. (2009) Bioindicators of changes in water quality on
828 coral reefs: review and recommendations for monitoring programmes. *Coral Reefs*, 28: 589-
829 606

830 Costa Jr., O.S., Nimmo, M. and Attrill, M.J. (2008) Coastal nutrification in Brazil: A review of the
831 role of nutrient excess on coral reef demise. *Journal of South American Earth Sciences*, 25:
832 257-270

833 Costanzo, S.D., O'Donohue, M.J., Dennison, W.C., Loneragan, N.R., and Thomas, M. (2001) A new
834 approach for detecting and mapping sewage impacts. *Mar. Pollut. Bull.*, 42(2): 149-156

835 Dajka, J-C, Wilson, S.K., Robinson, J.P.W., Chong-Seng, K.M., Harris, A. and Graham, N.A.J.
836 (2019) Uncovering drivers of juvenile coral density following mass bleaching. *Coral Reefs*,
837 38: 637-649

838 Dailer, M.L., Knox, R.S., Smith, J.E., Napier, M. and Smith, C.M. (2010) Using $\delta^{15}\text{N}$ values in algal
839 tissue to map locations and potential sources of anthropogenic nutrient inputs on the island of
840 Maui, Hawai'i, USA. *Mar. Pollut. Bull.*, 60(5): 655-671

841 D'Angelo, C. and Wiedenmann, J. (2014) Impacts of nutrient enrichment on coral reefs: new
842 perspectives and implications for coastal management and reef survival. *Current Opinion in*
843 *Environmental Sustainability*, 7: 82-93

844 De'ath, G. and Fabricius, K. (2010) Water quality as a regional driver of coral biodiversity and
845 macroalgae on the Great Barrier Reef. *Ecol. Appl.*, 20(3): 840-850

846 den Haan, J., Visser, P.M., Ganase, A.E., Gooren, A.E., Stal, L.J., van Duyl, F.C., Vermeij, M.J.A,
847 and Huisman, J. (2014) Nitrogen fixation rates in algal turf communities of a degraded versus
848 less degraded coral reef. *Coral Reefs*, 33(4): 1003-1015

849 Drummond, S.P. and Connell, S.D. (2008) Quantifying percent cover of subtidal organisms on rocky
850 coasts: a comparison of the costs and benefits of standard methods. *Marine and Freshwater*
851 *Research*, 56: 865-876

852 Duran, A., Collado-Videz, L., and Burkepile, D.E. (2016), Seasonal regulation of herbivory and
853 nutrient effects on macroalgal recruitment and succession in a Florida coral reef. *PeerJ*
854 4:e2643; DOI 10.7717/peerj.2643

855 Fabricius, K.E. (2005) Effects of terrestrial runoff on the ecology of corals and coral reefs: Review
856 and synthesis. *Marine Pollution Bulletin*, 50: 125-146

857 Fabricius, K., De'ath, G., McCook, L., Turak, E. and Williams, D. McB. (2005) Changes in algal,
858 coral and fish assemblages along water quality gradients on the inshore Great Barrier Reef.
859 *Mar. Pollut. Bull.*, 51 : 384-398

860 Fabricius, K.E., Cooper, T.F., Humphrey, C., Uthicke, S., De'ath, G., Davidson, J., LeGrand, H.,
861 Thompson, A., and Schaffelke, B. (2012) A bioindicator system for water quality on inshore
862 coral reefs of the Great Barrier Reef. *Mar. Pollut. Bull.*, 65: 320-332

863 Fernandes, M., Bengert, S., Sharma, S.K., Gaylard, S., Kildea, T., Hoare, S., Braley, M., and Irving,
864 A.D. (2012) The use of $\delta^{15}\text{N}$ signatures of translocated macroalgae to map coastal plumes:
865 improving species selection and spatial analysis of metropolitan datasets. *J. Environ. Monit.*,
866 14: 2399-2410

867 Fichez, R., Adjeroud, M., Bozec, Y-M., Breau, L., Chancerelle, Y., Chevillon, C. ... and Thebault, J.
868 (2005) A review of selected indicators of particle, nutrient and metal inputs in coral reef
869 lagoon systems. *Aquatic Living Resources*, 18: 125-147

870 Fleury, B.G., Coll, J.C., Tentori, E., Duquesne, S., and Figueiredo, L. (2000) Effect of nutrient
871 enrichment on the complementary (secondary) metabolite composition of the soft coral
872 *Sarcophyton ehrenbergi* (Cnidaria: Octocorallia: Alcyonaceae) of the Great Barrier Reef.
873 *Marine Biology*, 136: 63-68

874 Fong, P., Kamer, K., Boyer, K.E., and Boyle, K.A. (2001) Nutrient content of macroalgae with
875 differing morphologies may indicate sources of nutrients for tropical marine systems. *Mar.*
876 *Ecol. Prog. Ser.*, 220: 137-152

877 Fong, C.F. and Fong, P. (2014) Why species matter: an experimental assessment of assumptions and
878 predictive ability of two functional-group models. *Ecology*, 95(8): 2055-2061

879 Ford, A.K., Bejarano, S., Nugues, M.M., Visser, P.M., Albert, S., and Ferse, S.C.A. (2018) Reefs
880 under siege – the rise, putative drivers, and consequences of benthic cyanobacterial mats.
881 *Front. Mar. Sci.*, 5(18): 1-15

882 Fox, M.D., Williams, G.J., Johnson, M.D., Radice, V.Z., Zgliczynski, B.J., Kelly, E.L.A. ... and
883 Smith, J.E. (2018) Gradients in primary production predict trophic strategies of mixotrophic
884 corals across spatial scales. *Current Biology*, 28(21): 3355-3363

885 Fulton, C.J., Abesamis, R.A., Berkstrom, C., Depczynski, M., Graham, N.A.J., Holmes, T.H. ... and
886 Wilson, S.K. (2019) Form and function of tropical macroalgal reefs in the Anthropocene.
887 *Functional Ecology*, 1-11

888 García-Seoane, R., Aboal, J.R., Boquete, M.T., and Fernández, J.A. (2018) Biomonitoring coastal
889 environments with transplanted macroalgae: A methodological review. *Mar. Pollut. Bull.*,
890 135: 988-999

891 García-Seoane, R., Fernández, J.A., Villares, R., and Aboal, J.R. (2018b) Use of macroalgae to
892 biomonitor pollutants in coastal waters: Optimization of the methodology, *Ecological*
893 *Indicators*, 84: 710-726

894 Gartner, A., Lavery, P., and Smit, A.J. (2002) Use of $\delta^{15}\text{N}$ signatures of different functional forms of
895 macroalgae and filter-feeders to reveal temporal and spatial patterns in sewage dispersal.
896 *Marine Ecology Progress Series*, 235: 63-73

897 Graham, N.A.J., Jennings, S., MacNeil, M.A., Mouillot, D. and Wilson, S.K. (2015) Predicting
898 climate-driven regime shifts versus rebound potential in coral reefs. *Nature*, 518(7537): 94-97

899 Graham, N.A.J., Wilson, S.K., Carr, P., Hoey, A.S., Jennings, S., and Macneil, M.A. (2018) Seabirds
900 enhance coral reef productivity and functioning in the absence of invasive rats. *Nature*,
901 559(7713): 250-253

902 Hoegh-Guldberg, O., Muscatine, L., Goiran, C., Siggard, D., and Marion, G. (2004) Nutrient-induced
903 perturbations to $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ in symbiotic dinoflagellates and their coral hosts. *Mar. Ecol.*
904 *Prog. Ser.*, 280: 105-114

905 Hughes, T.P., Rodrigues, M.J., Bellwood, D.R., Ceccarelli, D., Hoegh-Guldberg, O., McCook, L. ...
906 and Willis, B. (2007) Phase shifts, herbivory, and the resilience of coral reefs to climate
907 change. *Curr. Biol.*, 17: 360-365

908 Hughes, T., Anderson, K.D., Connolly, S.R., Heron, S.F., Kerry, J.T., Lough, J.M. ... and Wilson, S.
909 (2018) Spatial and temporal patterns of mass bleaching of corals in the Anthropocene.
910 *Science*, 359: 80-83

911 Kürten, B., A-Aideroos, A.M., Struck, U., Sulaiman, K., Gharbawi, W.Y. and Sommer, U. (2014)
912 Influence of environmental gradients on C and N stable isotope ratios in coral reef biota of the
913 Red Sea, Saudi Arabia. *Journal of Sea Research*, 85: 379-394

914 Lamb, K., Swart, P.K. and Altabet, M.A. (2012) Nitrogen and carbon isotopic systematics of the
915 Florida Reef Tract. *Bulletin of Marine Science*, 88(1): 1-28

916 Leal, M.C., Rocha, R.J.M., Anaya-Rojas, J.M., Cruz, I.C.S., and Ferrier-Pages, C. (2017) Trophic and
917 stoichiometric consequences of nitrification for the intertidal tropical zoanthid *Zoanthus*
918 *sociatus*. *Mar. Pollut. Bull.*, 119(1): 169-175

919 Lin, D.T. and Fong, P. (2008) Macroalgal bioindicators (growth, tissue N, $\delta^{15}\text{N}$) detect nutrient
920 enrichment from shrimp farm effluent entering Opunohu Bay, Moorea, French Polynesia.
921 *Marine Pollution Bulletin.*, 56: 245-249

922 Linton, D.M. and Warner, G.F. (2003) Biological indicators in the Caribbean coastal zone and their
 923 role in integrated coastal management. *Ocean and Coastal Management*, 46: 261-276
 924 Littler, M.M., Littler, D.S., and Titlyanov, E.A. (1991) Comparisons of N- and P-limited productivity
 925 between high granitic islands versus low carbonate atolls in the Seychelles Archipelago: a test
 926 of the relative-dominance paradigm. *Coral Reefs*, 10: 199-209
 927 Littler, M.M., Littler, D.S., and Brooks, B.L. (2006) Harmful algae on tropical coral reefs: Bottom-up
 928 eutrophication and top-down herbivory. *Harmful Algae*, 5: 565-585
 929 Lowe, R.J. and Falter, J.L. (2015) Oceanic forcing of coral reefs. *Annu. Rev. Mar.Sci.*, 7: 43-66
 930 MacNeil, M.A., Mellin, C., Matthews, S., Wolff, N.H., McClanahan, T.R., Devlin, M. ... and
 931 Graham, N.A.J. (2019) Water quality mediates resilience on the Great Barrier Reef. *Nat Ecol.*
 932 *Evol.*, 3: 620-627
 933 Pitt, K.A., Connolly, R.M., Meziane, T. (2008) *Stable isotope and fatty acid tracers in energy and*
 934 *nutrient studies in jellyfish: a review*. In: Pitt, K.A. and Purcell, J.E. (eds) Jellyfish Blooms:
 935 Causes, Consequences, and Recent Advances. Developments in Hydrobiology, vol., 206.
 936 Springer, Dordrecht.
 937 Raimonet, M., Guillou, G., Mornet, F., and Richard, P. (2013) Macroalgae $\delta^{15}\text{N}$ values in well-mixed
 938 estuaries: Indicator of anthropogenic nitrogen input or macroalgae metabolism? *Estuarine,*
 939 *Coastal and Shelf Science*, 119: 126-138
 940 Raven, J.A., Johnston, A.M., Kübler, J.E., Korb, R., McInroy, S.G., Handley, L.L. ... and Dunton,
 941 K.H. (2002) Mechanistic interpretation of carbon isotope determination by marine
 942 macroalgae and seagrasses. *Funct. Plant. Biol.*, 29: 355-378
 943 R Core Team (2018). R: A language and environment for statistical computing. R Foundation for
 944 Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.
 945 Risk, M.J., Heikoop, J.M., Edinger, E.N., and Erdmann, M.V. (2001) The assessment “toolbox”:
 946 community-based reef evaluation methods coupled with geochemical techniques to identify
 947 sources of stress. *Bull. Mar. Sci.* 62(2): 443-458
 948 Risk, M.J. (2014) Assessing the effects of sediments and nutrients on coral reefs. *Current Opinion in*
 949 *Environmental Sustainability*, 7: 108-117

950 Schaffelke, B. and Klumpp, D.W. (1998) Nutrient-limited growth of the coral reef macroalga
951 *Sargassum baccularia* and experimental growth enhancement by nutrient addition in
952 continuous flow culture. *Mar. Ecol. Prog. Ser.*, 164: 199-211

953 Schaffelke, B. (1999) Short-term nutrient pulses as tools to assess responses of coral reef macroalgae
954 to enhance nutrient availability. *Mar. Ecol. Prog. Ser.*, 182: 305-310

955 Schlacher, T.A. and Connolly, R.M. (2014) Effects of acid treatment on carbon and nitrogen stable
956 isotope ratios in ecological samples: a review and synthesis. *Method of Ecology and*
957 *Evolution*, 5: 541-550

958 Sigman, D.M. and Casciotti, K.L. (2001) Nitrogen isotopes in the ocean. *Encyclopedia of Ocean*
959 *Sciences*, 1884-1894

960 Smit, A.J. (2001) 'Source Identification in Marine Ecosystems: Food Web Studies using $\delta^{13}C$ and
961 $\delta^{15}N$ ', in: Unkovich, M., Pate, J., McNeill, A., and Gibbs, D.J. (ed.) Stable Isotope
962 Techniques in the Study of Biological Processes and Functioning of Ecosystems, Kluwer
963 Academic Publishers, pp. 219-246

964 Steneck, R., & Dethier, M. (1994) A functional group approach to the structure of algal-dominated
965 communities. *Oikos*, 69(3), 476-498

966 Thacker, R.W. and Paul, V.J. (2001) Are benthic cyanobacterial indicators of nutrient enrichment?
967 Relationships between cyanobacterial abundance and environmental factors on the reef flats
968 of Guam. *Bulletin of Marine Science*, 69(2): 497-501

969 Tucker, J., Sheats, N., Giblin, A.E., Hopkinson, C.S., and Montoya, J.P. (1999) Using stable isotopes
970 to trace sewage-derived material through Boston Harbor and Massachusetts Bay. *Marine*
971 *Environmental Research*, 48: 4-5: 353-375

972 Umezawa, Y., Miyajima, T. and Koike, I. (2008) Stable nitrogen isotope composition in sedimentary
973 organic matter as a potential proxy of nitrogen sources for primary producers at a fringing
974 coral reef. *Journal of Oceanography*, 64: 899-909

975 Vaughan, E.J., Wilson, S.K., Howlett, S.J, Parravicini, V., Williams, G.J., and Graham, N.A.J. (2021)
976 Nitrogen enrichment in macroalgae following mass coral mortality. *Coral Reefs*, 40: 767–776

- Viana, I.G. and Bode, A. (2013) Stable nitrogen isotopes in coastal macroalgae: Geographic and anthropogenic variability. *Science of the Total Environment*, 443: 887-895
- Ward-Paige, C.A., Risk, M.J., Sherwood, O.A., and Jaap, W.C. (2005) Clinoid sponge surveys on the Florida Reef Track suggest land-based nutrient inputs, *Mar. Pollut. Bull.*, 51: 570-579
- Williams, G.J., Smith, J.E., Conklin, E.J., Gove, J.M., Sala, E., and Sandin, S.A. (2013), Benthic communities at two remote Pacific coral reefs: effects of reef habitat, depth, and wave energy gradients on spatial patterns. *PeerJ*, 1: e81; DOI 10.7717/peerj.81909
- Williams, G.J., Sandin, S.A., Zgliczynski, B.J., Fox, M.D., Gove, J.M., Rogers, J.S. ... and Smith, J.E. (2018) Biophysical drivers of coral trophic depth zonation. *Marine Biology*, 165: 60-75
- Wilson, S.K., Robinson, J.P.W., Chong-Seng, K, Robinson, J. and Graham, N.A.J. (2019) Boom and bust of keystone structure on coral reefs. *Coral Reefs*, 38: 625-635
- Wyatt, A.S.J., Lowe, R.J., Humphries, S. and Waite, A.M. (2013) Particulate nutrient fluxes over a fringing coral reef: Source-sink dynamics inferred from carbon to nitrogen ratios and stable isotopes, *Limnol. Oceanogr.*, 58(1): 409-427
- Zanden, M.J.V. and Rasmussen, J.B. (2001) Variation in $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ trophic fractionation: Implications for aquatic food web studies. *Limnol. Oceanogr.*, 46(8): 2016-2066
- Zaneveld, J.R., Burkepile, D.E., Shantz, A.A., Pritchard, C.E., McMinds, R., Payet, J.P. ... and Thurber, R.V. (2016) Overfishing and nutrient pollution interact with temperature to disrupt coral reefs down to microbial scales. *Nature Communications*, 7(11833): 1-12
- Zubia, M., Depetris, M., Flores, O., Turquet, J., and Cuet, P. (2018) Macroalgae as a tool for assessing the ecological status of coral reefs under the Water Framework Directive: A case study on the reef flats of La Reunion (Indian Ocean), *Marine Pollution Bulletin.*, 137: 339-351

Supplementary Figures

Supplementary Table 1. Summary of the 21 coral reefs surveyed around the inner Seychelles islands, including latitude, longitude, habitat type and reef state as categorised in 2017 (CM= coral-mortality reef; RS – regime-shifted reef). * denotes the sites added to the 2017 survey in place of the three sites around Cousin Island that were not surveyed that year (Graham et al., 2015).

| Site | Lat | Long | Habitat Type | Reef State |
|------------------------------------|-----------|----------|--------------|------------|
| Mahe West patch reef | -4.684675 | 55.43472 | Patch | CM |
| Mahe West carbonate | -4.669121 | 55.40025 | Carbonate | CM |
| Mahe West granitic reef | -4.659828 | 55.36099 | Granitic | CM |
| Mahe North West carbonate | -4.634994 | 55.37612 | Carbonate | CM |
| Mahe North West patch reef | -4.614482 | 55.41627 | Patch | CM |
| Mahe North West granitic | -4.562673 | 55.43691 | Granitic | CM |
| Ste. Anne granitic reef | -4.605095 | 55.51353 | Granitic | CM |
| Ste. Anne patch reef | -4.618086 | 55.5094 | Patch | CM |
| Ste Anne carbonate | -4.609864 | 55.49636 | Carbonate | RS |
| Mahe East granitic reef | -4.734961 | 55.52896 | Granitic | RS |
| Mahe East carbonate | -4.710589 | 55.52704 | Carbonate | RS |
| Mahe East patch reef | -4.703574 | 55.5282 | Patch | CM |
| Praslin North East patch reef | -4.303653 | 55.74655 | Patch | CM |
| Praslin North East carbonate | -4.315847 | 55.75669 | Carbonate | RS |
| Praslin NE granitic reef | -4.290079 | 55.7075 | Granitic | CM |
| Praslin SW granitic reef | -4.313662 | 55.67872 | Granitic | CM |
| Praslin SW patch reef | -4.333943 | 55.69204 | Patch | RS |
| Praslin SW carbonate | -4.350873 | 55.70152 | Carbonate | RS |
| Curieuse South West carbonate* | -4.28007 | 55.71199 | Carbonate | RS |
| Curieuse North East granitic reef* | -4.27987 | 55.74425 | Granitic | RS |
| Baie Ste Anne patch reef* | -4.34278 | 55.76919 | Patch | RS |

Supp. Table 2. Model (1) for each nutrient measurement and for each bioindicator: Nutrient Signature ~ Bioindicator + Reef State + Site. Model type was selected for each individual model based on normality of distribution. Sediment (SED) values were not available and so were not included for C-based signatures. Significance is noted as: '****' p < 0.001; '***' p < 0.01; '**' p < 0.05; and ',' p < 0.1.

| Bioindicator | Model Type (Family) | Intercept | Lower C.I. (5%) | Upper C.I. (95%) | p-value |
|---|---------------------|-----------|-----------------|------------------|----------------------|
| $\delta^{15}\text{N}$ | | | | | |
| Brown Macroalgae | | | | | |
| BSAP_Coral Mortality (Intercept) | GLM | 0.15 | 0.14 | 0.16 | < 0.0001 **** |
| Reef State: Regime Shift | GLM | 0.020 | 0.010 | 0.029 | 0.0002 *** |
| CNEG | GLM | 0.020 | 0.0095 | 0.031 | 0.0005 *** |
| CSWC | GLM | 0.013 | 0.0029 | 0.024 | 0.16 * |
| MEC | GLM | 0.0090 | -0.0014 | 0.019 | 0.095. |
| MEG | GLM | 0.046 | 0.034 | 0.057 | < 0.0001 **** |
| MEP | GLM | 0.0028 | -0.0063 | 0.012 | 0.55 |
| MNWP | GLM | -0.012 | -0.021 | -0.0032 | 0.010 * |
| PNEC | GLM | 0.064 | 0.052 | 0.076 | < 0.0001 **** |
| PNEG | GLM | 0.028 | 0.018 | 0.037 | < 0.0001 **** |
| PNEP | GLM | 0.033 | 0.023 | 0.043 | < 0.0001 **** |
| PSWC | GLM | 0.039 | 0.028 | 0.050 | < 0.0001 **** |
| PSWG | GLM | 0.033 | 0.023 | 0.044 | < 0.0001 **** |
| PSWP | GLM | 0.047 | 0.035 | 0.059 | < 0.0001 **** |
| SAC | GLM | 0.017 | 0.0062 | 0.028 | 0.003 ** |
| SAG | GLM | 0.0091 | -0.00012 | 0.018 | 0.058. |
| SAP | GLM | N/A | N/A | N/A | N/A |
| Cyanobacteria | | | | | |
| Intercept (BSAP_Coral Mortality) | GLM | 0.12 | 0.10 | 0.13 | < 0.0001 **** |
| Reef State: Regime Shift | GLM | 0.039 | 0.017 | 0.062 | 0.002 ** |
| MEC | GLM | -0.013 | -0.038 | 0.013 | 0.35 |
| MEG | GLM | 0.0055 | -0.019 | 0.030 | 0.66 |
| MEP | GLM | 0.019 | 0.00051 | 0.038 | 0.054. |
| MNWC | GLM | 0.015 | -0.0023 | 0.033 | 0.10. |
| MNWG | GLM | 0.020 | 0.0029 | 0.039 | 0.03 * |
| MWC | GLM | 0.037 | 0.018 | 0.056 | 0.0006 **** |
| MWG | GLM | 0.031 | 0.0097 | 0.053 | 0.009 ** |
| PNEC | GLM | 0.017 | -0.022 | 0.060 | 0.42 |
| PSWG | GLM | 0.033 | 0.015 | 0.052 | 0.0014 ** |
| SAG | GLM | 0.026 | -0.0046 | 0.061 | 0.12 |
| SAP | GLM | N/A | N/A | N/A | N/A |

Green Macroalgae

| | | | | | |
|----------------------------------|-----|--------|--------|---------|----------------------|
| Intercept (BSAP_Coral Mortality) | GLM | 0.17 | 0.16 | 0.17 | <0.0001*** |
| Reef State: Regime Shift | GLM | 0.030 | 0.023 | 0.036 | <0.0001*** |
| CNEG | GLM | -0.029 | -0.035 | -0.023 | <0.0001*** |
| CSWC | GLM | -0.014 | -0.021 | -0.0078 | <0.0001*** |
| MEC | GLM | -0.021 | -0.027 | -0.015 | <0.0001*** |
| PNEG | GLM | -0.010 | -0.017 | -0.0037 | 0.004** |
| PNEG | GLM | 0.013 | 0.0072 | 0.020 | 0.0001*** |
| PSWC | GLM | 0.15 | 0.0067 | 0.023 | 0.0009*** |
| PSWG | GLM | 0.020 | 0.013 | 0.026 | <0.0001*** |
| PSWP | GLM | 0.026 | 0.019 | 0.033 | <0.0001*** |
| SAC | GLM | -0.014 | -0.021 | -0.0077 | <0.0001*** |
| SAG | GLM | N/A | N/A | N/A | N/A |

Soft Coral

| | | | | | |
|---------------------------------|-----|---------|---------|--------|----------------------|
| Intercept (MEC_Coral Mortality) | GLM | 0.12 | 0.11 | 0.13 | <0.0001*** |
| Reef State: Regime Shift | GLM | 0.016 | -0.004 | 0.038 | 0.14 |
| MEP | GLM | 0.026 | 0.012 | 0.041 | 0.001** |
| MNWC | GLM | 0.012 | 0.00097 | 0.024 | 0.042* |
| MNWP | GLM | 0.10 | 0.084 | 0.12 | <0.0001*** |
| MWC | GLM | 0.0060 | -0.0052 | 0.017 | 0.31 |
| MWG | GLM | 0.015 | 0.0035 | 0.027 | 0.017* |
| MWP | GLM | 0.0058 | -0.0054 | 0.017 | 0.32 |
| PNEG | GLM | -0.0039 | -0.015 | 0.0068 | 0.48 |
| PSWG | GLM | N/A | N/A | N/A | N/A |

Sediment

| | | | | | |
|----------------------------------|----|-------|-------|--------|----------------------|
| Intercept (BSAP_Coral Mortality) | LM | 9.4 | 8.7 | 10.2 | <0.0001*** |
| Reef State: Regime Shift | LM | 1.4 | 0.35 | 2.5 | 0.010* |
| CNEG | LM | 0.076 | -0.99 | 1.1 | 0.89 |
| CSWC | LM | 0.90 | -0.17 | 2.0 | 0.10 |
| MEC | LM | -1.3 | -2.3 | -0.18 | 0.022* |
| MEG | LM | -1.1 | -2.3 | 0.15 | 0.085. |
| MEP | LM | -1.0 | -2.1 | 0.051 | 0.062. |
| MNWC | LM | -1.1 | -2.1 | 0.0018 | 0.050. |
| MNWG | LM | 0.52 | -0.55 | 1.6 | 0.34 |
| MNWP | LM | -0.95 | -2.0 | 0.12 | 0.081. |
| MWC | LM | -1.2 | -2.3 | -0.13 | 0.029* |
| MWG | LM | 1.2 | 0.021 | 2.3 | 0.046* |
| MWP | LM | 1.5 | 0.43 | 2.6 | 0.0064** |
| PNEG | LM | -2.6 | -3.7 | -1.5 | <0.0001*** |
| PNEG | LM | 1.2 | 0.09 | 2.2 | 0.034* |
| PNEP | LM | -0.12 | -1.2 | 0.95 | 0.83 |

| | | | | | | |
|-------------------|----------------------------------|-----|----------|---------|---------|----------------------|
| | PSWC | LM | -1.2 | -2.2 | -0.080 | 0.035* |
| | PSWG | LM | 1.3 | 0.25 | 2.4 | 0.012* |
| | PSWP | LM | -3.7 | -4.8 | -2.6 | <0.0001*** |
| | SAC | LM | -2.1 | -3.2 | -1.0 | 0.0002*** |
| | SAG | LM | 0.95 | -0.12 | 2.0 | 0.082. |
| | SAP | LM | N/A | N/A | N/A | N/A |
| Sponge | | | | | | |
| | Intercept (CNEG_Coral Mortality) | GLM | 0.13 | 0.12 | 0.14 | <0.0001*** |
| | Reef State: Regime Shift | GLM | 0.0013 | -0.016 | 0.019 | 0.89 |
| | CSWC | GLM | 0.0026 | 0.0062 | 0.046 | 0.013* |
| | MEG | GLM | -0.0034 | -0.029 | 0.023 | 0.80 |
| | MEP | GLM | 0.0043 | -0.010 | 0.020 | 0.56 |
| | MNWG | GLM | -0.0024 | -0.016 | 0.011 | 0.72 |
| | MWC | GLM | 0.0098 | -0.0041 | 0.024 | 0.17 |
| | MWG | GLM | 0.01 | -0.0038 | 0.024 | 0.16 |
| | MWP | GLM | 0.0011 | -0.012 | 0.015 | 0.87 |
| | PNEC | GLM | 0.0067 | -0.012 | 0.025 | 0.47 |
| | PNEG | GLM | 0.0064 | -0.0073 | 0.020 | 0.37 |
| | PSWC | GLM | 0.0097 | -0.017 | 0.038 | 0.49 |
| | PSWG | GLM | 0.0062 | -0.0075 | 0.020 | 0.38 |
| | PSWP | GLM | 0.0017 | -0.017 | 0.020 | 0.86 |
| | SAG | GLM | -0.0036 | -0.017 | 0.0095 | 0.59 |
| | SAP | GLM | N/A | N/A | N/A | N/A |
| Turf Algae | | | | | | |
| | Intercept (BSAP_Coral Mortality) | GLM | 0.13 | 0.12 | 0.15 | <0.0001*** |
| | Reef State: Regime Shift | GLM | -0.029 | -0.052 | -0.0044 | 0.020* |
| | CNEG | GLM | 0.0066 | -0.017 | 0.028 | 0.57 |
| | CSWC | GLM | 0.041 | 0.016 | 0.065 | 0.001** |
| | MEC | GLM | 0.026 | 0.0012 | 0.048 | 0.036* |
| | MNWC | GLM | -0.0044 | -0.021 | 0.012 | 0.60 |
| | MNWG | GLM | 0.0055 | -0.011 | 0.022 | 0.53 |
| | MNWP | GLM | 0.0037 | -0.014 | 0.022 | 0.69 |
| | MWC | GLM | 0.018 | -0.0027 | 0.039 | 0.10 |
| | MWG | GLM | 0.0015 | -0.015 | 0.018 | 0.86 |
| | PNEC | GLM | 0.30 | 0.26 | 0.34 | <0.0001*** |
| | PNEG | GLM | -0.0064 | -0.023 | 0.0098 | 0.44 |
| | PNEP | GLM | -0.00036 | -0.017 | 0.016 | 0.97 |
| | PSWC | GLM | 0.054 | 0.018 | 0.093 | 0.006** |
| | PSWG | GLM | -0.0043 | -0.021 | 0.012 | 0.61 |
| | PSWP | GLM | 0.093 | 0.062 | 0.12 | <0.0001*** |
| | SAC | GLM | 0.028 | 0.0029 | 0.05 | 0.026* |

| | | | | | | |
|------------------|----------------------------------|-----|---------|---------|-----------|------------|
| | SAG | GLM | 0.019 | -0.0050 | 0.043 | 0.14 |
| | SAP | GLM | N/A | N/A | N/A | N/A |
| Zoanthids | | | | | | |
| | Intercept (BSAP_Coral Mortality) | GLM | 0.12 | 0.11 | 0.12 | <0.0001*** |
| | Reef State: Regime Shift | GLM | 0.017 | 0.012 | 0.022 | <0.0001*** |
| | CNEG | GLM | 0.017 | -0.024 | -0.014 | <0.0001*** |
| | MEG | GLM | -0.019 | -0.031 | -0.16 | <0.0001*** |
| | MEP | GLM | -0.024 | 0.0069 | 0.020 | <0.0001*** |
| | MNWC | GLM | 0.013 | -0.011 | -0.0018 | 0.0001*** |
| | MNWG | GLM | -0.011 | -0.016 | -0.0071 | 0.008** |
| | MNWP | GLM | -0.018 | -0.022 | -0.014 | <0.0001*** |
| | MWC | GLM | -0.013 | -0.018 | -0.0091 | <0.0001*** |
| | MWG | GLM | -0.012 | -0.017 | -0.0082 | <0.0001*** |
| | MWP | GLM | -0.022 | -0.026 | -0.018 | <0.0001*** |
| | PNEC | GLM | -0.011 | -0.016 | -0.0058 | <0.0001*** |
| | PNEG | GLM | -0.0044 | -0.0088 | -0.000024 | 0.056. |
| | PNEP | GLM | -0.0014 | -0.0058 | 0.0031 | 0.55 |
| | PSWC | GLM | -0.0037 | -0.0087 | 0.0014 | 0.16 |
| | PSWG | GLM | -0.0039 | -0.0083 | 0.00053 | 0.089. |
| | PSWP | GLM | -0.025 | -0.030 | -0.020 | <0.0001*** |
| | SAC | GLM | -0.033 | -0.038 | -0.028 | <0.0001*** |
| | SAG | GLM | -0.0043 | -0.0090 | 0.00031 | 0.072. |
| | SAP | GLM | N/A | N/A | N/A | N/A |

$\delta^{13}\text{C}$

| | | | | | | |
|--|----------------------------------|-----|-------|--------|-------|------------|
| | Brown Macroalgae | GLM | | | | |
| | Intercept (BSAP_Coral Mortality) | GLM | -17.0 | -17.9 | -16.1 | <0.0001*** |
| | Reef State: Regime Shift | GLM | -0.25 | -1.6 | 1.1 | 0.72 |
| | CNEG | GLM | -0.42 | -1.7 | 0.91 | 0.54 |
| | CSWC | GLM | 1.2 | -0.079 | 2.6 | 0.070. |
| | MEC | GLM | 0.44 | -0.89 | 1.8 | 0.52 |
| | MEG | GLM | 0.16 | -1.2 | 1.5 | 0.82 |
| | MEP | GLM | 0.20 | -1.1 | 1.5 | 0.77 |
| | MNWP | GLM | 4.1 | 2.8 | 5.4 | <0.0001*** |
| | PNEC | GLM | 1.9 | 0.54 | 3.2 | 0.008** |
| | PNEG | GLM | 2.5 | 1.2 | 3.8 | 0.0005*** |
| | PNEP | GLM | 1.6 | 0.31 | 3.0 | 0.02* |
| | PSWC | GLM | -0.18 | -1.5 | 1.1 | 0.79 |
| | PSWG | GLM | -0.60 | -2.0 | 0.81 | 0.41 |
| | PSWP | GLM | 1.2 | -0.17 | 2.5 | 0.093. |
| | SAC | GLM | 2.0 | 0.70 | 3.4 | 0.004** |
| | SAG | GLM | 0.36 | -0.96 | 1.7 | 0.59 |

| | | | | | |
|----------------------------------|-----|-------|-------|-------|------------|
| SAP | GLM | N/A | N/A | N/A | N/A |
| Cyanobacteria | | | | | |
| Intercept (BSAP_Coral Mortality) | GLM | -14.6 | -16.5 | -12.7 | <0.0001*** |
| Reef State: Regime Shift | GLM | -9.3 | -12.2 | -6.4 | <0.0001*** |
| MEC | GLM | -2.4 | -3.3 | 2.8 | 0.88 |
| MEG | GLM | 0.40 | -2.4 | 3.2 | 0.78 |
| MEP | GLM | -8.8 | -11.5 | -6.1 | <0.0001*** |
| MNWC | GLM | -2.4 | -5.0 | 0.11 | 0.069. |
| MNWG | GLM | -7.9 | -10.4 | -5.3 | <0.0001*** |
| MNWP | GLM | -7.2 | -9.7 | -4.6 | <0.0001*** |
| MWC | GLM | -5.0 | -7.5 | -2.5 | 0.0004*** |
| MWG | GLM | -9.2 | -12.1 | -6.4 | <0.0001*** |
| PNEC | GLM | 3.1 | -1.2 | 7.5 | 0.17 |
| PSWG | GLM | -7.8 | -10.4 | -5.3 | <0.0001*** |
| SAG | GLM | -8.5 | -12.7 | -4.3 | 0.0004*** |
| SAP | GLM | N/A | N/A | N/A | N/A |
| Green Macroalgae | | | | | |
| Intercept (BSAP_Coral Mortality) | GLM | -21.0 | -21.4 | -20.5 | <0.0001*** |
| Reef State: Regime Shift | GLM | 1.2 | 0.64 | 1.8 | 0.0002*** |
| CNEG | GLM | -2.6 | -3.2 | -2.0 | <0.0001*** |
| CSWC | GLM | -1.1 | -1.6 | -0.52 | 0.0005*** |
| MEC | GLM | -1.4 | -2.0 | -0.83 | <0.0001*** |
| PNEC | GLM | -2.7 | -3.2 | -2.1 | <0.0001*** |
| PNEG | GLM | 0.89 | 0.29 | 1.5 | 0.006** |
| PSWC | GLM | -1.8 | -2.5 | -1.2 | <0.0001*** |
| PSWG | GLM | -0.98 | -1.6 | -0.39 | 0.002** |
| PSWP | GLM | -1.6 | -2.2 | -1.1 | <0.0001*** |
| SAC | GLM | -2.3 | -2.9 | -1.8 | <0.0001*** |
| SAG | GLM | N/A | N/A | N/A | N/A |
| Soft Coral | | | | | |
| Intercept (MEC_Coral Mortality) | GLM | -17.7 | -18.6 | 16.7 | <0.0001*** |
| Reef State: Regime Shift | GLM | 0.93 | -1.3 | 3.2 | 0.42 |
| MEP | GLM | 3.0 | 1.6 | 4.5 | 0.0003*** |
| MNWC | GLM | 2.3 | 1.0 | 3.6 | 0.001** |
| MNWG | GLM | 1.1 | -0.19 | 2.4 | -0.10 |
| MNWP | GLM | 1.3 | 0.028 | 2.6 | 0.053. |
| MWC | GLM | 2.2 | 0.89 | 3.5 | 0.002** |
| MWG | GLM | 0.41 | -0.88 | 1.7 | 0.54 |
| MWP | GLM | 2.1 | 0.76 | 3.3 | 0.004** |
| PNEG | GLM | 0.81 | -0.48 | 2.1 | 0.23 |
| PSWG | GLM | N/A | N/A | N/A | N/A |

| | | | | | |
|-----------------------------------|-----|--------|-------|--------|----------------------|
| Sponge | | | | | |
| Intercept (CNEG_Coral Mortality?) | GLM | -17.2 | -17.5 | -16.9 | <0.0001*** |
| Reef State: Regime Shift | GLM | -0.034 | -0.53 | 0.47 | 0.89 |
| CSWC | GLM | -0.023 | -0.52 | 0.48 | 0.93 |
| MEG | GLM | 0.13 | -0.60 | 0.86 | 0.73 |
| MEP | GLM | 0.15 | -0.22 | 0.53 | 0.43 |
| MNWG | GLM | -0.53 | -0.91 | -0.15 | 0.008** |
| MWC | GLM | -0.12 | -0.49 | 0.26 | 0.55 |
| MWG | GLM | -0.60 | -0.98 | -0.22 | 0.003** |
| MWP | GLM | -0.44 | -0.81 | -0.059 | 0.028* |
| PNEC | GLM | 0.31 | -0.19 | 0.81 | 0.24 |
| PNEG | GLM | -0.30 | -0.68 | 0.079 | 0.13 |
| PSWC | GLM | -0.42 | -1.1 | 0.31 | 0.27 |
| PSWG | GLM | -0.44 | -0.82 | -0.064 | 0.027* |
| SAG | GLM | -0.21 | -0.58 | 0.17 | 0.29 |
| SAP | GLM | N/A | N/A | N/A | N/A |
| Turf Algae | | | | | |
| Intercept (BSAP_Coral Mortality) | GLM | -14.3 | -15.4 | -13.2 | <0.0001*** |
| Reef State: Regime Shift | GLM | -1.6 | -4.3 | 0.97 | 0.22 |
| CNEG | GLM | -1.8 | -4.4 | 0.85 | 0.19 |
| CSWC | GLM | 0.74 | -1.9 | 3.4 | 0.58 |
| MEC | GLM | -0.98 | -3.6 | 1.6 | 0.46 |
| MNWC | GLM | -6.6 | -8.1 | -5.1 | <0.0001*** |
| MNWG | GLM | -3.0 | -4.5 | -1.5 | 0.0002*** |
| MNWP | GLM | -6.4 | -8.0 | -4.8 | <0.0001*** |
| MWC | GLM | -8.2 | -10.0 | -6.5 | <0.0001*** |
| MWG | GLM | -10.1 | -11.6 | -8.6 | <0.0001*** |
| PNEC | GLM | 2.7 | 0.078 | 5.3 | 0.048* |
| PNEG | GLM | -5.1 | -6.6 | -3.5 | <0.0001*** |
| PNEP | GLM | -4.5 | -6.0 | -3.0 | <0.0001*** |
| PSWC | GLM | -0.60 | -3.3 | 2.1 | 0.65 |
| PSWG | GLM | -5.2 | -6.7 | -3.7 | <0.0001*** |
| PSWP | GLM | -6.9 | -9.5 | -4.2 | <0.0001*** |
| SAC | GLM | -2.4 | -5.0 | 0.24 | 0.080. |
| SAG | GLM | -4.9 | -6.9 | -2.9 | <0.0001*** |
| SAP | GLM | N/A | N/A | N/A | N/A |
| Zoanthids | | | | | |
| Intercept (BSAP_Coral Mortality) | GLM | -13.8 | -14.6 | 13.1 | <0.0001*** |
| Reef State: Regime Shift | GLM | 1.1 | 0.14 | 2.2 | 0.031* |
| CNEG | GLM | -2.2 | -3.4 | -1.1 | 0.0002*** |
| MEG | GLM | -1.5 | -3.2 | 0.14 | 0.079. |

| | | | | | |
|------|-----|---------|-------|--------|---------------|
| MEP | GLM | 0.33 | -0.97 | 1.6 | 0.62 |
| MNWC | GLM | 0.60 | -0.47 | 1.7 | 0.27 |
| MNWG | GLM | -0.12 | -1.1 | 0.90 | 0.82 |
| MNWP | GLM | 0.10 | -0.91 | 1.1 | 0.84 |
| MWC | GLM | 0.56 | -0.45 | 1.6 | 0.29 |
| MWG | GLM | -0.23 | -1.2 | 0.78 | 0.66 |
| MWP | GLM | 0.57 | -0.44 | 1.6 | 0.28 |
| PNEC | GLM | -0.84 | -1.9 | 0.17 | 0.11 |
| PNEG | GLM | 0.044 | -1.6 | 1.7 | 0.96 |
| PNEP | GLM | -0.55 | -1.9 | 0.75 | 0.41 |
| PSWC | GLM | -1.3 | -2.3 | -2.5 | 0.018* |
| PSWG | GLM | -0.0030 | -1.3 | 1.3 | 0.99 |
| PSWP | GLM | -1.0 | -2.0 | 0.0065 | 0.057. |
| SAG | GLM | -0.97 | -2.0 | 0.039 | 0.066. |
| SAP | GLM | N/A | N/A | N/A | N/A |

%N

| | | | | | |
|----------------------------------|-----|---------|---------|--------|----------------------|
| Brown Macroalgae | | | | | |
| Intercept (BSAP_Coral Mortality) | LM | 1.07 | 0.96 | 1.2 | <0.0001*** |
| Reef State: Regime Shift | LM | 0.18 | 0.053 | 0.31 | 0.040* |
| CNEG | LM | -0.41 | -0.57 | -0.25 | <0.0001*** |
| CSWC | LM | -0.16 | -0.32 | 0.0052 | 0.058. |
| MEC | LM | 0.13 | -0.033 | 0.29 | 0.12 |
| MEG | LM | -0.072 | -0.23 | 0.091 | 0.38 |
| MEP | LM | 0.19 | 0.025 | 0.35 | 0.024* |
| MNWP | LM | 0.053 | -0.11 | 0.22 | 0.52 |
| PNEC | LM | -0.20 | -0.36 | -0.034 | 0.019* |
| PNEG | LM | 0.16 | -0.0028 | 0.32 | 0.054. |
| PNEP | LM | -0.088 | -0.25 | 0.075 | 0.28 |
| PSWC | LM | 0.037 | -0.13 | 0.20 | 0.65 |
| PSWG | LM | -0.0072 | -0.18 | 0.17 | 0.93 |
| PSWP | LM | -0.26 | -0.42 | -0.096 | 0.002** |
| SAC | LM | -0.11 | -0.27 | 0.053 | 0.18 |
| SAG | LM | 0.037 | -0.13 | 0.2 | 0.65 |
| SAP | LM | N/A | N/A | N/A | N/A |
| Cyanobacteria | | | | | |
| Intercept (BSAP_Coral Mortality) | GLM | 0.65 | 0.50 | 0.83 | <0.0001*** |
| Reef State: Regime Shift | GLM | -0.41 | -0.60 | -0.24 | <0.0001*** |
| MEC | GLM | -0.024 | -0.12 | 0.07 | 0.62 |
| MEG | GLM | -0.023 | -0.11 | 0.060 | 0.60 |
| MEP | GLM | -0.44 | -0.62 | -0.28 | <0.0001*** |
| MNWC | GLM | -0.063 | -0.28 | 0.14 | 0.56 |

| | | | | | |
|----------------------------------|-----|---------|--------|---------|----------------------|
| MNWG | GLM | -0.32 | -0.51 | -0.16 | 0.0007*** |
| MWC | GLM | -0.22 | -0.42 | -0.035 | 0.030* |
| MWG | GLM | -0.41 | -0.60 | -0.24 | <0.0001*** |
| PNEC | GLM | -0.017 | -0.14 | 0.13 | 0.80 |
| PSWG | GLM | -0.37 | -0.56 | -0.21 | 0.0002*** |
| SAG | GLM | -0.41 | -0.62 | -0.20 | 0.0004*** |
| SAP | GLM | N/A | N/A | N/A | N/A |
| Green Macroalgae | | | | | |
| Intercept (BSAP_Coral Mortality) | GLM | 0.23 | 0.21 | 0.25 | <0.0001*** |
| Reef State: Regime Shift | GLM | -0.0012 | -0.028 | 0.026 | 0.93 |
| CNEG | GLM | 0.030 | 0.0033 | 0.057 | 0.034* |
| CSWC | GLM | 0.0023 | -0.023 | 0.028 | 0.86 |
| MEC | GLM | -0.026 | -0.050 | -0.0019 | 0.041* |
| PNEC | GLM | 0.015 | -0.011 | 0.041 | 0.28 |
| PNEG | GLM | -0.012 | -0.039 | 0.014 | 0.37 |
| PSWC | GLM | -0.026 | -0.053 | 0.0014 | 0.069. |
| PSWG | GLM | 0.029 | 0.0012 | 0.058 | 0.047* |
| PSWP | GLM | -0.013 | -0.038 | 0.011 | 0.30 |
| SAC | GLM | 0.014 | -0.013 | 0.040 | 0.32 |
| SAG | GLM | N/A | N/A | N/A | N/A |
| Soft Coral | | | | | |
| Intercept (MEC_Coral Mortality) | GLM | 0.43 | 0.32 | 0.57 | <0.0001*** |
| Reef State: Regime Shift | GLM | -0.14 | -0.35 | 0.12 | 0.24 |
| MEP | GLM | 0.023 | -0.18 | 0.25 | 0.83 |
| MNWC | GLM | -0.20 | -0.35 | -0.064 | 0.0098** |
| MNWP | GLM | -0.29 | -0.43 | -0.17 | 0.0002*** |
| MWC | GLM | 0.0010 | -0.18 | -0.18 | 0.99 |
| MWG | GLM | -0.16 | -0.32 | -0.020 | 0.41* |
| MWP | GLM | -0.12 | -0.28 | 0.033 | 0.14 |
| PNEG | GLM | -0.12 | -0.28 | 0.031 | 0.14 |
| PSWG | GLM | N/A | N/A | N/A | N/A |
| Sediment | | | | | |
| Intercept (BSAP_Coral Mortality) | GLM | 27.5 | 18.3 | 39.4 | <0.0001*** |
| Reef State: Regime Shift | GLM | -5.1 | -19.3 | 8.3 | 0.46 |
| CNEG | GLM | 1.3 | -11.3 | 14.1 | 0.84 |
| CSWC | GLM | 1.9 | -10.8 | 15.0 | 0.76 |
| MEC | GLM | 10.3 | -4.4 | 26.4 | 0.19 |
| MEG | GLM | 10.4 | -6.4 | 30.9 | 0.27 |
| MEP | GLM | 1.1 | -14.3 | 16.6 | 0.89 |
| MNWC | GLM | 8.6 | -8.48 | 26.8 | 0.34 |
| MNWG | GLM | 5.8 | -10.6 | 23.1 | 0.49 |

| | | | | | |
|----------------------------------|-----|---------|---------|-------|----------------------|
| MNWP | GLM | 0.81 | -14.5 | 16.3 | 0.92 |
| MWC | GLM | 5.1 | -11.2 | 22.1 | 0.54 |
| MWG | GLM | 0.10 | -15.7 | 16.6 | 0.99 |
| MWP | GLM | 0.84 | -14.5 | 16.3 | 0.91 |
| PNEC | GLM | 7.2 | -6.7 | 22.2 | 0.32 |
| PNEG | GLM | -3.7 | -18.1 | 10.2 | 0.61 |
| PNEP | GLM | 2.3 | -13.4 | 18.2 | 0.78 |
| PSWC | GLM | 9.9 | -4.7 | 25.8 | 0.20 |
| PSWG | GLM | -23.7 | -35.7 | -14.3 | <0.0001*** |
| PSWP | GLM | -2.6 | -14.4 | 8.8 | 0.65 |
| SAC | GLM | 5.9 | -7.7 | 20.3 | 0.41 |
| SAG | GLM | -4.6 | -18.9 | 9.0 | 0.51 |
| SAP | GLM | N/A | N/A | N/A | N/A |
| Sponge | | | | | |
| Intercept (CNEG_Coral Mortality) | GLM | 0.53 | 0.44 | 0.64 | <0.0001*** |
| Reef State: Regime Shift | GLM | -0.055 | -0.22 | 0.12 | 0.53 |
| CSWC | GLM | 0.15 | -0.044 | 0.34 | 0.13 |
| MEG | GLM | 0.41 | 0.058 | 0.86 | 0.046* |
| MEP | GLM | 0.25 | 0.068 | 0.45 | 0.012* |
| MNWG | GLM | -0.061 | -0.19 | 0.07 | 0.37 |
| MWC | GLM | 0.064 | -0.08 | 0.21 | 0.40 |
| MWG | GLM | 0.029 | -0.11 | 0.17 | 0.69 |
| MWP | GLM | 0.24 | 0.076 | 0.43 | 0.008** |
| PNEC | GLM | -0.0010 | -0.18 | 0.16 | 0.99 |
| PNEG | GLM | 0.16 | -0.0034 | 0.32 | 0.064. |
| PSWC | GLM | -0.027 | -0.25 | 0.22 | 0.82 |
| PSWG | GLM | -0.079 | -0.21 | 0.050 | 0.24 |
| PSWP | GLM | 0.052 | 0.133 | 0.23 | 0.57 |
| SAG | GLM | 0.16 | -0.018 | 0.33 | 0.056. |
| SAP | GLM | N/A | N/A | N/A | N/A |
| Turf Algae | | | | | |
| Intercept (BSAP_Coral Mortality) | GLM | 0.64 | 0.41 | 0.96 | <0.0001*** |
| Reef State: Regime Shift | GLM | -0.058 | -0.39 | 0.21 | 0.70 |
| CSWC | GLM | 0.031 | -0.13 | 0.20 | 0.71 |
| MNWC | GLM | -0.16 | -0.49 | 0.10 | 0.30 |
| MNWG | GLM | -0.0069 | -0.35 | 0.28 | 0.97 |
| MWG | GLM | -0.063 | -0.42 | 0.24 | 0.71 |
| PNEC | GLM | 0.42 | 0.20 | 0.65 | 0.0007*** |
| PNEG | GLM | -0.059 | -0.39 | 0.21 | 0.70 |
| PNEP | GLM | -0.074 | -0.27 | 0.36 | 0.64 |
| PSWC | GLM | 0.052 | -0.11 | 0.22 | 0.54 |

| | | | | | |
|----------------------------------|-----|----------|----------|---------|----------------------|
| PSWG | GLM | -0.053 | -0.40 | 0.24 | 0.74 |
| SAC | GLM | 0.11 | -0.065 | 0.28 | 0.23 |
| SAG | GLM | 1.4 | 0.74 | 2.1 | 0.0003*** |
| SAP | GLM | N/A | N/A | N/A | N/A |
| Zoanthids | | | | | |
| Intercept (BSAP_Coral Mortality) | GLM | 0.94 | 0.82 | 1.1 | <0.0001*** |
| Reef State: Regime Shift | GLM | 0.22 | 0.021 | 0.42 | 0.036* |
| CNEG | GLM | -0.30 | -0.49 | -0.11 | 0.0037** |
| MEG | GLM | -0.29 | -0.58 | 0.027 | 0.62 |
| MEP | GLM | 0.33 | 0.044 | 0.64 | 0.034* |
| MNWC | GLM | 0.12 | -0.079 | 0.33 | 0.25 |
| MNWG | GLM | -0.010 | -0.19 | 0.17 | 0.91 |
| MNWP | GLM | 0.064 | -0.12 | 0.25 | 0.50 |
| MWC | GLM | -0.085 | -0.26 | 0.085 | 0.33 |
| MWG | GLM | -0.040 | -0.21 | 0.13 | 0.66 |
| MWP | GLM | 0.025 | -0.15 | 0.21 | 0.79 |
| PNEC | GLM | 0.14 | -0.10 | 0.40 | 0.26 |
| PNEG | GLM | -0.07 | -0.24 | 0.10 | 0.42 |
| PNEP | GLM | 0.14 | -0.050 | 0.33 | 0.15 |
| PSWC | GLM | -0.42 | -0.61 | -0.24 | <0.0001*** |
| PSWG | GLM | -0.11 | -0.28 | 0.058 | 0.21 |
| PSWP | GLM | -0.16 | -0.38 | -0.52 | 0.14 |
| SAC | GLM | -0.14 | -0.37 | 0.10 | 0.26 |
| SAG | GLM | 0.26 | -0.43 | -0.90 | 0.004** |
| SAP | GLM | N/A | N/A | N/A | N/A |
| %C | | | | | |
| Brown Macroalgae | | | | | |
| Intercept (BSAP_Coral Mortality) | GLM | 0.031 | 0.030 | 0.032 | <0.0001*** |
| Reef State: Regime Shift | GLM | 0.0010 | -0.00060 | 0.0027 | 0.22 |
| CNEG | GLM | -0.00012 | -0.0018 | 0.0015 | 0.89 |
| CSWC | GLM | 0.00033 | -0.0013 | 0.0020 | 0.70 |
| MEC | GLM | -0.00074 | -0.0024 | 0.00091 | 0.38 |
| MEG | GLM | 0.0018 | 0.00013 | 0.0036 | 0.039* |
| MEP | GLM | 0.0025 | 0.00080 | 0.0042 | 0.0052** |
| MNWP | GLM | 0.0011 | -0.00056 | 0.0028 | 0.20 |
| PNEC | GLM | -0.00038 | -0.0020 | 0.0013 | 0.65 |
| PNEG | GLM | 0.0012 | -0.00042 | 0.0029 | 0.15 |
| PNEP | GLM | 0.00041 | -0.0012 | 0.0020 | 0.62 |
| PSWC | GLM | 0.00019 | -0.0015 | 0.0019 | 0.82 |
| PSWG | GLM | 0.0022 | 0.00044 | 0.0040 | 0.017* |
| PSWP | GLM | 0.00087 | -0.00082 | 0.0026 | 0.32 |

| | | | | | | |
|-------------------------|----------------------------------|-----|----------|----------|---------|----------------------|
| | SAC | GLM | -0.00075 | -0.0024 | 0.00090 | 0.38 |
| | SAG | GLM | 0.0022 | 0.00051 | 0.0039 | 0.013* |
| | SAP | GLM | N/A | N/A | N/A | N/A |
| Cyanobacteria | | | | | | |
| | Intercept (BSAP_Coral Mortality) | GLM | 0.053 | 0.047 | 0.059 | <0.0001*** |
| | Reef State: Regime Shift | GLM | -0.024 | -0.031 | -0.017 | <0.0001*** |
| | MEC | GLM | -0.00033 | -0.0055 | 0.0049 | 0.90 |
| | MEG | GLM | 0.0042 | -0.00078 | 0.0091 | 0.10 |
| | MEP | GLM | -0.023 | -0.030 | -0.017 | <0.0001*** |
| | MNWC | GLM | -0.0071 | -0.015 | 0.00015 | 0.065. |
| | MNWG | GLM | -0.021 | -0.027 | -0.014 | <0.0001*** |
| | MNWP | GLM | -0.017 | -0.024 | -0.011 | <0.0001*** |
| | MWC | GLM | -0.011 | -0.019 | -0.0045 | 0.0030** |
| | MWG | GLM | -0.020 | -0.027 | -0.013 | <0.0001*** |
| | PNEC | GLM | 0.00027 | -0.0069 | 0.0080 | 0.94 |
| | PSWG | GLM | -0.020 | -0.027 | -0.013 | <0.0001*** |
| | SAG | GLM | -0.022 | -0.031 | -0.013 | <0.0001*** |
| | SAP | GLM | N/A | N/A | N/A | N/A |
| Green Macroalgae | | | | | | |
| | Intercept (BSAP_Coral Mortality) | GLM | 0.024 | 0.023 | 0.025 | <0.0001*** |
| | Reef State: Regime Shift | GLM | 0.00045 | -0.0017 | 0.00082 | 0.49 |
| | CNEG | GLM | 0.00073 | -0.0019 | 0.00043 | 0.23 |
| | CSWC | GLM | 0.0010 | -0.0021 | 0.00016 | 0.098. |
| | MEC | GLM | 0.0019 | 0.0006 | 0.0031 | 0.005** |
| | PNEC | GLM | 0.0010 | -0.0023 | 0.00021 | 0.11 |
| | PNEG | GLM | 0.00095 | -0.0022 | 0.00030 | 0.14 |
| | PSWC | GLM | 0.000086 | -0.0013 | 0.0015 | 0.90 |
| | PSWG | GLM | 0.00043 | -0.00084 | 0.0017 | 0.50 |
| | PSWP | GLM | 0.00068 | -0.0018 | 0.00048 | 0.26 |
| | SAC | GLM | 0.0023 | 0.0011 | 0.0038 | 0.0006*** |
| | SAG | GLM | N.A | N/A | N/A | N/A |
| Soft Coral | | | | | | |
| | Intercept (MEC_Coral Mortality) | GLM | 51.5 | 49.8 | 53.1 | <0.0001*** |
| | Reef State: Regime Shift | GLM | 2.5 | -1.6 | 6.6 | 0.25 |
| | MEP | GLM | -0.13 | -2.9 | 2.6 | 0.93 |
| | MNWC | GLM | 1.6 | -0.81 | 3.9 | 0.20 |
| | MNWG | GLM | -0.87 | -3.2 | 1.5 | 0.48 |
| | MNWP | GLM | 2.1 | -0.21 | 4.5 | 0.084. |
| | MWC | GLM | 2.2 | -0.21 | 4.5 | 0.084. |
| | MWG | GLM | 0.75 | -1.6 | 3.1 | 0.54 |
| | MWP | GLM | 0.43 | -1.9 | 2.8 | 0.72 |

| | | | | | | | |
|----------------------------------|----------------------------------|----------|-----------|-----------|------------|------------|--|
| Sponge | PNEG | GLM | 0.56 | -1.8 | 2.9 | 0.65 | |
| | PSWG | GLM | N/A | N/A | N/A | N/A | |
| | Intercept (CNEG_Coral Mortality) | GLM | 0.096 | 0.092 | 0.10 | <0.0001*** | |
| | Reef State: Regime Shift | GLM | 0.0011 | -0.0066 | 0.0090 | 0.79 | |
| | CSWC | GLM | -0.00032 | -0.0082 | 0.0074 | 0.94 | |
| | MEG | GLM | -0.0039 | -0.015 | 0.0074 | 0.50 | |
| | MEP | GLM | 0.0032 | -0.0028 | 0.0091 | 0.30 | |
| | MNWG | GLM | 0.011 | 0.0050 | 0.017 | 0.00096*** | |
| | MWC | GLM | 0.0032 | -0.0027 | 0.0092 | 0.29 | |
| | MWG | GLM | 0.0043 | -0.0017 | 0.010 | 0.17 | |
| | MWP | GLM | 0.0016 | -0.0043 | 0.0075 | 0.60 | |
| | PNEC | GLM | 0.0028 | -0.0052 | 0.011 | 0.49 | |
| | PNEG | GLM | 0.0038 | -0.0021 | 0.0098 | 0.21 | |
| | PSWC | GLM | 0.0083 | -0.0036 | 0.021 | 0.19 | |
| | PSWG | GLM | -0.0020 | -0.0078 | 0.0038 | 0.51 | |
| | SAG | GLM | 0.0086 | 0.0025 | 0.015 | 0.0084** | |
| | SAP | GLM | N/A | N/A | N/A | N/A | |
| | Turf Algae | | | | | | |
| | Intercept (BSAP_Coral Mortality) | GLM | 0.024 | 0.023 | 0.024 | <0.0001*** | |
| Reef State: Regime Shift | GLM | -0.0013 | -0.0026 | 0.00012 | 0.075. | | |
| CNEG | GLM | 0.0017 | 0.00029 | 0.0031 | 0.020* | | |
| CSWC | GLM | 0.00049 | -0.00089 | 0.0018 | 0.49 | | |
| MEC | GLM | 0.00037 | -0.0010 | 0.0071 | 0.60 | | |
| MNWC | GLM | -0.0010 | -0.0018 | -0.000021 | 0.016* | | |
| MNWG | GLM | -0.0012 | -0.0020 | -0.00042 | 0.004*** | | |
| MNWP | GLM | 0.0029 | 0.0020 | 0.0039 | <0.0001*** | | |
| MWC | GLM | 0.00025 | -0.00071 | 0.0012 | 0.61 | | |
| MWG | GLM | 0.000068 | -0.00076 | 0.00089 | 0.87 | | |
| PNEC | GLM | 0.00067 | -0.000070 | 0.0020 | 0.34 | | |
| PNEG | GLM | -0.00026 | -0.0011 | 0.00056 | 0.54 | | |
| PNEP | GLM | 0.00067 | 0.0016 | -0.000030 | 0.047* | | |
| PSWC | GLM | -0.00084 | -0.0011 | 0.0017 | 0.67 | | |
| PSWG | GLM | -0.00078 | -0.0016 | 0.000024 | 0.062. | | |
| SAC | GLM | 0.0012 | -0.00020 | 0.0025 | 0.094. | | |
| SAG | GLM | 0.0033 | 0.0021 | 0.0044 | <0.0001*** | | |
| | | | N/A | N/A | N/A | N/A | |
| Zoanthids | | | | | | | |
| Intercept (BSAP_Coral Mortality) | GLM | 0.038 | 0.026 | 0.053 | <0.0001*** | | |
| Reef State: Regime Shift | GLM | 0.056 | 0.026 | 0.091 | 0.001** | | |
| CNEG | GLM | 0.040 | -0.018 | 0.11 | 0.21 | | |

| | | | | | | |
|-------------------------|----------------------------------|-----|----------|---------|---------|----------------------|
| | MEG | GLM | -0.071 | -0.11 | -0.038 | 0.0001*** |
| | MEP | GLM | 0.0052 | -0.018 | 0.033 | 0.69 |
| | MNWC | GLM | 0.14 | 0.081 | 0.21 | <0.0001*** |
| | MNWG | GLM | 0.10 | 0.059 | 0.15 | <0.0001*** |
| | MNWP | GLM | 0.11 | 0.063 | 0.16 | <0.0001*** |
| | MWC | GLM | 0.0076 | -0.012 | 0.027 | 0.44 |
| | MWG | GLM | 0.025 | 0.0022 | 0.050 | 0.042* |
| | MWP | GLM | 0.0048 | -0.014 | 0.024 | 0.62 |
| | PNEC | GLM | -0.055 | -0.089 | -0.024 | 0.002** |
| | PNEG | GLM | 0.11 | 0.026 | 0.23 | 0.047* |
| | PNEP | GLM | -0.010 | -0.029 | 0.0097 | 0.30 |
| | PSWC | GLM | -0.060 | -0.10 | -0.034 | 0.0003*** |
| | PSWG | GLM | 0.060 | 0.017 | 0.12 | 0.023* |
| | PSWP | GLM | -0.040 | -0.077 | -0.0067 | 0.027* |
| | SAG | GLM | 0.036 | 0.011 | 0.064 | 0.010* |
| | SAP | GLM | N/A | N/A | N/A | N/A |
| C:N Ratio | | | | | | |
| Brown Macroalgae | | | | | | |
| | Intercept (BSAP_Coral Mortality) | GLM | 0.033 | 0.030 | 0.036 | <0.0001*** |
| | Reef State: Regime Shift | GLM | 0.0052 | 0.00049 | 0.0099 | 0.035* |
| | CENG | GLM | -0.0134 | -0.018 | -0.0092 | <0.0001*** |
| | CSWC | GLM | -0.0046 | -0.0093 | 0.00013 | 0.062. |
| | MEC | GLM | 0.0029 | -0.0023 | 0.0081 | 0.28 |
| | MEG | GLM | -0.00027 | -0.0053 | 0.0047 | 0.92 |
| | MEP | GLM | 0.0088 | 0.0039 | 0.014 | 0.0009*** |
| | MNWP | GLM | 0.0028 | -0.0017 | 0.0073 | 0.23 |
| | PNEC | GLM | -0.0072 | -0.012 | -0.0026 | 0.003** |
| | PNEG | GLM | 0.0063 | 0.0016 | 0.011 | 0.012* |
| | PNEP | GLM | -0.0031 | -0.0073 | 0.0010 | 0.14 |
| | PSWC | GLM | 0.0013 | -0.0038 | 0.0064 | 0.61 |
| | PSWG | GLM | 0.0017 | -0.0030 | 0.0065 | 0.47 |
| | PSWP | GLM | -0.0073 | -0.012 | -0.0028 | 0.002** |
| | SAC | GLM | -0.0043 | -0.0090 | 0.00047 | 0.083. |
| | SAG | GLM | 0.0034 | -0.0012 | 0.0079 | 0.15 |
| | SAP | GLM | N/A | N/A | N/A | N/A |
| Cyanobacteria | | | | | | |
| | Intercept (BSAP_Coral Mortality) | GLM | 0.12 | 0.099 | 0.14 | <0.0001*** |
| | Reef State: Regime Shift | GLM | 0.027 | -0.0033 | 0.057 | 0.087. |
| | MEC | GLM | -0.016 | -0.047 | 0.016 | 0.34 |
| | MEG | GLM | N/A | N/A | N/A | N/A |
| | MEP | GLM | 0.016 | -0.015 | 0.046 | 0.32 |

| | | | | | | |
|-------------------------|----------------------------------|-----|----------|---------|----------|----------------------|
| | MNWC | GLM | -0.019 | -0.046 | 0.0065 | 0.16 |
| | MNWG | GLM | -0.020 | -0.046 | 0.0057 | 0.15 |
| | MWC | GLM | -0.029 | -0.055 | -0.0046 | 0.031* |
| | MWG | GLM | 0.0094 | -0.021 | 0.039 | 0.54 |
| | PNEC | GLM | 0.010 | -0.033 | 0.060 | 0.66 |
| | PSWG | GLM | -0.0014 | 0.031 | 0.027 | 0.92 |
| | SAP | GLM | -0.048 | -0.073 | -0.024 | 0.0006*** |
| Green Macroalgae | | | | | | |
| | Intercept (BSAP_Coral Mortality) | GLM | 0.10 | 0.096 | 0.11 | <0.0001*** |
| | Reef State: Regime Shift | GLM | -0.0015 | -0.012 | 0.0093 | 0.78 |
| | CNEG | GLM | -0.016 | -0.025 | -0.0063 | 0.002** |
| | CSWC | GLM | -0.0052 | -0.015 | 0.0047 | 0.31 |
| | MEC | GLM | 0.022 | 0.11 | 0.033 | 0.0004*** |
| | PNEC | GLM | -0.012 | -0.022 | -0.0015 | 0.031* |
| | PNEG | GLM | 0.00082 | -0.010 | 0.012 | 0.88 |
| | PSWC | GLM | 0.014 | 0.0010 | 0.026 | 0.042* |
| | PSWG | GLM | -0.010 | -0.021 | -0.00012 | 0.056. |
| | PSWP | GLM | 0.00067 | -0.0095 | 0.011 | 0.90 |
| | SAC | GLM | 0.0034 | -0.0069 | 0.014 | 0.52 |
| | SAG | GLM | N/A | N/A | N/A | N/A |
| Soft Coral | | | | | | |
| | Intercept (MEC_Coral Mortality) | GLM | 0.018 | 0.16 | 0.20 | <0.0001*** |
| | Reef State: Regime Shift | GLM | -0.00021 | -0.053 | 0.060 | 0.99 |
| | MEP | GLM | 0.021 | -0.018 | 0.062 | 0.32 |
| | MNWC | GLM | 0.0088 | -0.024 | 0.042 | 0.61 |
| | MNWG | GLM | 0.027 | -0.0074 | 0.063 | 0.13 |
| | MNWP | GLM | -0.0046 | -0.037 | 0.027 | 0.78 |
| | MWC | GLM | -0.027 | -0.057 | 0.0031 | 0.090. |
| | MWG | GLM | 0.021 | -0.014 | 0.055 | 0.25 |
| | MWP | GLM | 0.016 | -0.018 | 0.050 | 0.37 |
| | PNEG | GLM | 0.00012 | -0.032 | 0.033 | 0.99 |
| | PSWG | GLM | N/A | N/A | N/A | N/A |
| Sponge | | | | | | |
| | Intercept (CNEG_Coral Mortality) | GLM | 1.2 | 1.0 | 1.3 | <0.0001*** |
| | Reef State: Regime Shift | GLM | -0.041 | -0.31 | 0.24 | 0.77 |
| | CSWC | GLM | 0.030 | -0.25 | 0.30 | 0.83 |
| | MEG | GLM | -0.041 | -0.42 | 0.37 | 0.84 |
| | MEP | GLM | -0.026 | -0.24 | 0.19 | 0.81 |
| | MNWG | GLM | 0.022 | -0.19 | 0.24 | 0.84 |
| | MWC | GLM | 0.011 | -0.20 | 0.23 | 0.92 |
| | MWG | GLM | -0.030 | -0.24 | 0.18 | 0.78 |

| | | | | | | |
|-------------------|----------------------------------|-----|----------|--------|---------|----------------------|
| | MWP | GLM | -0.024 | -0.24 | 0.19 | 0.83 |
| | PNEC | GLM | 0.23 | -0.09 | 0.51 | 0.13 |
| | PNEG | GLM | 0.0041 | -0.27 | 0.30 | 0.98 |
| | PSWC | GLM | 0.12 | -0.29 | 0.57 | 0.59 |
| | PSWG | GLM | -0.054 | -0.26 | 0.15 | 0.61 |
| | SAG | GLM | -0.011 | -0.22 | 0.20 | 0.92 |
| | SAP | GLM | N/A | N/A | N/A | N/A |
| Turf Algae | | | | | | |
| | Intercept (BSAP_Coral Mortality) | GLM | 0.068 | 0.059 | 0.078 | <0.0001*** |
| | Reef State: Regime Shift | GLM | 0.11 | 0.062 | 0.18 | 0.0002*** |
| | CNEG | GLM | -0.060 | -0.12 | -0.0068 | 0.048* |
| | CSWC | GLM | -0.97 | -0.16 | -0.045 | 0.0014** |
| | MEC | GLM | -0.78 | -0.14 | -0.026 | 0.0098** |
| | MNWC | GLM | 0.056 | 0.037 | 0.077 | <0.0001*** |
| | MNWG | GLM | 0.053 | 0.034 | 0.072 | <0.0001*** |
| | MNWP | GLM | 0.057 | 0.037 | 0.079 | <0.0001*** |
| | MWC | GLM | 0.044 | 0.023 | 0.067 | 0.0002*** |
| | MWG | GLM | 0.039 | 0.021 | 0.056 | <0.0001*** |
| | PNEC | GLM | -0.084 | -0.15 | -0.031 | 0.0059** |
| | PNEG | GLM | 0.064 | 0.044 | 0.085 | <0.0001*** |
| | PNEP | GLM | 0.068 | 0.046 | 0.092 | <0.0001*** |
| | PSWC | GLM | -0.13 | -0.20 | -0.084 | <0.0001*** |
| | PSWG | GLM | 0.049 | 0.030 | 0.070 | <0.0001*** |
| | PSWP | GLM | -0.030 | -0.095 | 0.026 | 0.33 |
| | SAC | GLM | -0.057 | -0.12 | -0.0039 | 0.059. |
| | SAG | GLM | 0.067 | 0.038 | 0.099 | <0.0001*** |
| | SAP | GLM | N/A | N/A | N/A | N/A |
| Zoanthids | | | | | | |
| | Intercept (BSAP_Coral Mortality) | GLM | 0.25 | 0.19 | 0.32 | <0.0001*** |
| | Reef State: Regime Shift | GLM | -0.011 | -0.095 | 0.071 | 0.80 |
| | CNEG | GLM | 0.014 | -0.074 | 0.11 | 0.76 |
| | MEG | GLM | 0.0069 | -0.11 | 0.16 | 0.92 |
| | MEP | GLM | -0.00023 | -0.10 | 0.11 | 0.99 |
| | MNWC | GLM | -0.17 | -0.24 | -0.11 | <0.0001*** |
| | MNWG | GLM | -0.17 | -0.24 | -0.11 | <0.0001*** |
| | MNWP | GLM | -0.17 | -0.24 | -0.11 | <0.0001*** |
| | MWC | GLM | -0.072 | -0.15 | 0.00014 | 0.065. |
| | MWG | GLM | -0.050 | -0.14 | 0.048 | 0.30 |
| | MWP | GLM | 0.21 | 0.093 | 0.33 | 0.0014** |
| | PNEC | GLM | 0.012 | -0.070 | 0.096 | 0.78 |
| | PNEG | GLM | -0.033 | -0.15 | 0.11 | 0.61 |

| | | | | | |
|------|-----|---------|--------|--------|-----------------|
| PNEP | GLM | -0.0067 | -0.11 | 0.11 | 0.90 |
| PSWC | GLM | -0.0044 | -0.084 | 0.077 | 0.91 |
| PSWG | GLM | -0.13 | -0.21 | -0.059 | 0.0012** |
| PSWP | GLM | -0.0077 | -0.086 | 0.073 | 0.85 |
| SAG | GLM | -0.095 | -0.17 | -0.026 | 0.012* |
| SAP | GLM | N/A | N/A | N/A | N/A |

1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065

Supp. Table 3. Model (2) for the CoV of each nutrient measurement in each bioindicator across all sites: $\text{CoV} \sim \text{Bioindicator} + \text{Reef State} + \text{Site}$. Model type selected for each individual model based on normality of distribution. Sediment (SED) values were not available and so were not included for C-based signatures. Significance is noted as: '****' $p < 0.001$; '***' $p < 0.01$; '**' $p < 0.05$; and '.' $p < 0.1$.

| Bioindicator | Model Type | Estimate | Lower C.I. (%) | Upper C.I. (%) | p-value |
|-------------------------------------|------------|----------|----------------|----------------|-----------------------|
| $\Delta^{15}\text{N}$ | | | | | |
| Intercept (BM_BSAP_Coral Mortality) | GLM | 0.26 | 0.17 | 0.37 | <0.0001**** |
| CYB | GLM | -0.083 | -0.16 | -0.012 | 0.027* |
| GM | GLM | 0.17 | 0.042 | 0.33 | 0.019* |
| SC | GLM | -0.059 | -0.14 | 0.028 | 0.17 |
| SED | GLM | -0.086 | -0.15 | -0.027 | 0.008** |
| SP | GLM | -0.071 | -0.14 | -0.0011 | 0.053. |
| TA | GLM | -0.099 | -0.17 | -0.038 | 0.003** |
| ZO | GLM | 0.11 | 0.0077 | 0.21 | 0.042* |
| Reef State: Regime Shift | GLM | -0.056 | -0.18 | 0.066 | 0.36 |
| CNEG | GLM | 0.048 | -0.077 | 0.18 | 0.45 |
| CSWC | GLM | -0.045 | -0.15 | 0.042 | 0.35 |
| MEC | GLM | 0.15 | -0.0048 | 0.34 | 0.079. |
| MEG | GLM | -0.064 | -0.17 | 0.018 | 0.16 |
| MEP | GLM | -0.082 | -0.19 | 0.015 | 0.12 |
| MNWC | GLM | -0.051 | -0.17 | 0.058 | 0.37 |
| MNWG | GLM | -0.0048 | -0.13 | 0.12 | 0.94 |
| MNWP | GLM | -0.069 | -0.18 | 0.037 | 0.21 |
| MWC | GLM | -0.053 | -0.17 | 0.048 | 0.32 |
| MWG | GLM | 0.16 | -0.0079 | 0.36 | 0.083. |
| MWP | GLM | -0.047 | -0.17 | 0.079 | 0.45 |
| PNEC | GLM | -0.034 | -0.14 | 0.057 | 0.49 |
| PNEG | GLM | -0.051 | -0.17 | 0.054 | 0.36 |
| PNEP | GLM | -0.050 | -0.17 | 0.074 | 0.42 |
| PSWC | GLM | 0.12 | -0.056 | 0.34 | 0.24 |
| PSWG | GLM | -0.0037 | -0.13 | 0.11 | 0.95 |
| PSWP | GLM | 0.024 | -0.095 | 0.14 | 0.68 |
| SAC | GLM | 0.017 | -0.10 | 0.14 | 0.78 |
| SAG | GLM | -0.019 | -0.14 | 0.11 | 0.76 |
| SAP | GLM | N/A | N/A | N/A | N/A |
| $\Delta^{13}\text{C}$ | | | | | |
| Intercept (BM_BSAP_Coral Mortality) | LM | -6.5 | -10.0 | -3.0 | 0.0004**** |
| CYB | LM | -0.93 | -3.8 | 1.9 | 0.52 |
| GM | LM | 4.2 | 1.4 | 7.0 | 0.004** |
| SC | LM | 0.61 | -2.6 | 3.8 | 0.70 |
| SP | LM | 4.9 | 2.1 | 7.7 | 0.0008**** |
| TA | LM | 1.4 | -1.1 | 3.9 | 0.26 |
| ZO | LM | 1.2 | -1.4 | 3.7 | 0.37 |
| Reef State: Regime Shift | LM | 0.72 | -4.0 | 5.4 | 0.76 |
| CNEG | LM | -2.1 | -6.8 | 2.5 | 0.36 |
| CSWC | LM | -0.21 | -5.2 | 4.7 | 0.93 |
| MEC | LM | 1.8 | -3.1 | 6.7 | 0.46 |
| MEG | LM | 2.9 | -3.2 | 8.9 | 0.35 |
| MEP | LM | -0.48 | -4.9 | 3.9 | 0.83 |
| MNWC | LM | -3.3 | -8.0 | 1.4 | 0.16 |
| MNWG | LM | -1.1 | -5.5 | 3.3 | 0.62 |
| MNWP | LM | 1.2 | -3.2 | 5.6 | 0.58 |
| MWC | LM | -1.0 | 5.4 | 3.4 | 0.64 |

| | | | | | |
|-------------------------------------|-----|--------|--------|--------|----------------------|
| MWG | LM | 1.3 | -3.1 | 5.7 | 0.56 |
| MWP | LM | -0.61 | -5.8 | 4.5 | 0.81 |
| PNEC | LM | -0.73 | -3.2 | 3.9 | 0.75 |
| PNEG | LM | -0.41 | -4.8 | 4.0 | 0.85 |
| PNEP | LM | -.17 | -6.8 | 3.4 | 0.50 |
| PSWC | LM | -1.3 | -6.2 | 3.6 | 0.59 |
| PSWG | LM | 1.6 | -2.5 | 5.6 | 0.44 |
| PSWP | LM | 0.62 | -4.3 | 5.5 | 0.80 |
| SAC | LM | -3.6 | -8.9 | 1.7 | 0.18 |
| SAG | LM | 3.3 | -1.1 | 7.7 | 0.14 |
| SAP | LM | N/A | N/A | N/A | N/A |
| %N | | | | | |
| Intercept (BM_BSAP_Coral Mortality) | GLM | 14.8 | -6.4 | 36.0 | 0.17 |
| CYB | GLM | 8.8 | -8.8 | 26.4 | 0.33 |
| GM | GLM | -3.7 | -20.4 | 13.1 | 0.67 |
| SC | GLM | 14.6 | -5.0 | 34.1 | 0.15 |
| SED | GLM | 5.2 | -8.9 | 19.4 | 0.47 |
| SP | GLM | 5.0 | -11.3 | 21.4 | 0.55 |
| TA | GLM | 7.2 | -9.4 | 23.8 | 0.40 |
| ZO | GLM | 1.0 | -13.8 | 15.8 | 0.89 |
| Reef State: Regime Shift | GLM | -11.0 | -37.7 | 15.8 | 0.42 |
| CNEG | GLM | 12.8 | -12.9 | 38.5 | 0.33 |
| CSWC | GLM | 1.4 | -25.6 | 28.2 | 0.92 |
| MEC | GLM | 0.050 | -28.1 | 28.2 | 0.99 |
| MEG | GLM | 3.1 | -27.8 | 34.0 | 0.85 |
| MEP | GLM | 0.18 | -25.3 | 25.7 | 0.99 |
| MNWC | GLM | -8.1 | -35.1 | 18.9 | 0.56 |
| MNWG | GLM | -2.0 | -27.6 | 23.6 | 0.88 |
| MNWP | GLM | -8.4 | -37.0 | 20.2 | 0.57 |
| MWC | GLM | -0.33 | -27.1 | 26.4 | 0.98 |
| MWG | GLM | 7.3 | -18.4 | 33.0 | 0.58 |
| MWP | GLM | 3.9 | -24.7 | 32.5 | 0.79 |
| PNEC | GLM | 8.5 | 17.1 | 34.2 | 0.52 |
| PNEG | GLM | -3.1 | -28.1 | 21.8 | 0.81 |
| PNEP | GLM | -1.1 | -29.6 | 27.4 | 0.94 |
| PSWC | GLM | 5.8 | -21.0 | 32.6 | 0.67 |
| PSWG | GLM | 16.5 | -7.6 | 40.7 | 0.18 |
| PSWP | GLM | 8.2 | -18.6 | 34.9 | 0.55 |
| SAC | GLM | 5.9 | -20.9 | 32.7 | 0.67 |
| SAG | GLM | -7.9 | -33.7 | 17.8 | 0.55 |
| SAP | GLM | N/A | N/A | N/A | N/A |
| %C | | | | | |
| Intercept (BM_BSAP_Coral Mortality) | GLM | 0.34 | 0.053 | 0.84 | 0.075 |
| CYB | GLM | -0.13 | -0.22 | -0.047 | 0.004** |
| GM | GLM | 0.023 | -0.10 | 0.16 | 0.72 |
| SC | GLM | 0.081 | -0.059 | 0.24 | 0.29 |
| SP | GLM | -0.017 | -0.12 | 0.093 | 0.76 |
| TA | GLM | 0.21 | 0.060 | 0.37 | 0.011* |
| ZO | GLM | -0.19 | -0.27 | -0.12 | <0.0001*** |
| Reef State: Regime Shift | GLM | -0.046 | -0.53 | 0.22 | 0.80 |
| CNEG | GLM | -0.083 | -0.22 | 0.0058 | 0.13 |
| CSWC | GLM | 0.038 | -0.19 | 0.32 | 0.77 |
| MEC | GLM | 0.065 | -0.14 | 0.33 | 0.58 |
| MEG | GLM | 0.055 | -0.16 | 0.37 | 0.67 |
| MEP | GLM | -0.13 | -0.63 | 0.15 | 0.48 |
| MNWC | GLM | -0.11 | -0.61 | 0.18 | 0.56 |
| MNWG | GLM | -0.12 | -0.61 | 0.18 | 0.57 |
| MNWP | GLM | -0.13 | -0.62 | 0.17 | 0.54 |

| | | | | | | |
|--------------------------|------|------|--------|-------|---------|--------|
| | MWC | GLM | -0.13 | -0.63 | 0.15 | 0.48 |
| | MWG | GLM | -0.12 | -0.62 | 0.17 | 0.54 |
| | MWP | GLM | -0.088 | -0.59 | 0.21 | 0.65 |
| | PNEC | GLM | -0.091 | -0.22 | -0.0019 | 0.097. |
| | PNEG | GLM | -0.065 | -0.58 | 0.28 | 0.75 |
| | PNEP | GLM | -0.11 | -0.61 | 0.18 | 0.56 |
| | PSWC | GLM | -0.080 | -0.21 | 0.011 | 0.15 |
| | PSWG | GLM | -0.096 | -0.59 | 0.20 | 0.62 |
| | PSWP | GLM | -0.097 | -0.23 | -0.018 | 0.072. |
| | SAC | GLM | -0.091 | -0.29 | 0.15 | 0.39 |
| | SAG | GLM | -0.075 | -0.58 | 0.23 | 0.70 |
| | SAP | GLM | -0.13 | -0.63 | 0.16 | 0.49 |
| C:N Ratio | | | | | | |
| Intercept (BM_BSAP_Coral | GLM | 2.6 | -20.0 | 25.2 | 0.82 | |
| Mortality) | | | | | | |
| | CYB | GLM | 1.9 | -9.7 | 13.5 | 0.75 |
| | GM | GLM | -4.9 | -15.5 | 5.8 | 0.37 |
| | SC | GLM | 0.14 | -11.9 | 12.2 | 0.98 |
| | SP | GLM | -5.9 | -16.6 | 4.7 | 0.28 |
| | TA | GLM | -0.87 | -10.2 | 8.5 | 0.86 |
| | ZO | GLM | 9.2 | -0.46 | 18.9 | 0.066. |
| Reef State: Regime Shift | GLM | 0.65 | -21.2 | 22.5 | 0.95 | |
| | CNEG | GLM | 13.2 | -5.2 | 31.6 | 0.17 |
| | CSWC | GLM | 5.6 | -13.9 | 25.0 | 0.58 |
| | MEC | GLM | 4.1 | -15.3 | 23.5 | 0.68 |
| | MEG | GLM | 8.6 | -15.0 | 32.2 | 0.48 |
| | MEP | GLM | 1.6 | -22.0 | 26.1 | 0.90 |
| | MNWC | GLM | 4.0 | -21.1 | 29.2 | 0.75 |
| | MNWG | GLM | 5.2 | -19.3 | 29.7 | 0.68 |
| | MNWP | GLM | 1.2 | -24.4 | 26.7 | 0.93 |
| | MWC | GLM | 18.0 | -6.6 | 42.5 | 0.16 |
| | MWG | GLM | 17.8 | -6.77 | 42.3 | 0.16 |
| | MWP | GLM | 8.0 | -18.9 | 34.9 | 0.56 |
| | PNEC | GLM | 14.7 | -3.0 | 32.5 | 0.11 |
| | PNEG | GLM | 9.9 | -14.7 | 34.5 | 0.43 |
| | PNEP | GLM | 4.0 | -22.7 | 30.7 | 0.77 |
| | PSWC | GLM | 3.1 | -16.2 | 22.4 | 0.75 |
| | PSWG | GLM | 14.4 | -9.1 | 37.9 | 0.23 |
| | PSWP | GLM | 1.7 | -17.6 | 21.0 | 0.86 |
| | SAC | GLM | 8.6 | -12.3 | 29.5 | 0.42 |
| | SAG | GLM | 8.5 | 16.1 | 33.1 | 0.50 |
| | SAP | GLM | 12.0 | -12.5 | 36.4 | 0.34 |

1072

1073

1074

1075

1076

1077

1078

1079

1080

1081

Supp. Table 4. Generalised Linear Model (3) for the cost-effectiveness analyses to determine the effect of Bioindicator, Task, Reef State and the interaction between them on the time per unit sample (per hour): Time ~ Bioindicator * Task * Reef State. Normality inspected using visual plots. Significance is noted as: ‘***’ p < 0.001; ‘**’ p < 0.01; ‘*’ p < 0.05; and ‘,’ p < 0.1.

| | Intercept | Lower C.I. (2.5%) | Upper C.I. (97.5%) | p-value |
|---------------------------|-----------|----------------------|-----------------------|------------------------|
| BM (Intercept) | -1.33 | -1.89 | -0.767 | <0.0001*** |
| CYB | 0.0109 | -0.498 | 0.520 | 0.966 |
| GM | 0.00803 | -0.689 | 0.704 | 0.982 |
| SC | 1.50 | 0.831 | 2.16 | <0.0001*** |
| SED | -1.37 | -1.85 | -0.891 | <0.0001*** |
| SP | 1.49 | 0.838 | 2.15 | <0.0001*** |
| TA | 1.45 | 0.864 | 2.13 | <0.0001*** |
| ZO | 1.49 | 1.01 | 1.97 | <0.0001*** |
| DRY-CRUSH | 26.0 | 25.3 | 26.7 | <0.0001*** |
| FIELD | 1.40 | 0.719 | 2.08 | <0.0001*** |
| SIA | 1.51 | 0.830 | 2.19 | <0.0001*** |
| WEIGH | 2.84 | 2.43 | 3.25 | <0.0001*** |
| REEF STATE- REGIME | 0.00376 | -0.873 | 0.881 | 0.993 |
| SHIFT | | | | |
| CYB-DRY | -1.91 | -2.63 | -1.19 | <0.0001*** |
| GM-DRY | -0.562 | -1.55 | 0.424 | 0.264 |
| SC-DRY | -3.94 | -4.79 | -3.094 | <0.0001*** |
| SED-DRY | -0.681 | -1.36 | -0.00183 | 0.0500* |
| SP-DRY | -3.83 | -4.66 | -3.00 | <0.0001*** |
| TA-DRY | -1.48 | -2.27 | -0.684 | 0.000293*** |
| ZO-DRY | -3.10 | -3.78 | -2.42 | <0.0001*** |
| CYB-FIELD | 0.237 | -0.483 | 0.957 | 0.519 |
| GM-FIELD | 0.0864 | -0.899 | 1.072 | 0.864 |
| SC-FIELD | -1.42 | -2.27 | -0.573 | 0.00109** |
| SED-FIELD | 1.32 | 0.636 | 2.00 | 0.000166*** |
| SP-FIELD | -1.46 | -2.29 | -0.629 | <0.000620*** |
| TA-FIELD | -1.55 | -2.34 | -0.753 | <0.000152*** |
| ZO-FIELD | -1.45 | -2.13 | -0.767 | <0.0001*** |
| CYB-SIA | 0.0336 | -0.686 | 0.753 | 0.927 |
| GM-SIA | 0.0158 | -0.970 | 1.00 | 0.975 |
| SC-SIA | -1.26 | -2.12 | -0.410 | 0.00380** |
| SED-SIA | 1.40 | 0.720 | 2.08 | <0.0001*** |
| SP-SIA | -1.29 | -2.12 | -0.459 | 0.00246** |
| TA-SIA | -1.43 | -2.22 | -0.635 | 0.000461*** |
| ZO-SIA | -1.25 | -1.92 | -0.566 | 0.000361*** |
| CYB-WEIGH | NA | NA | NA | NA |
| GM-WEIGH | NA | NA | NA | NA |
| SC-WEIGH | 0.0198 | -0.632 | 0.672 | 0.953 |
| SED-WEIGH | NA | NA | NA | NA |
| SP-WEIGH | 0.0476 | -0.585 | 0.675 | 0.889 |
| TA-WEIGH | 0.00659 | -0.576 | 0.590 | 0.982 |
| ZO-WEIGH | NA | NA | NA | NA |
| CYB-REGIME | -0.00667 | -0.799 | 0.785 | 0.987 |
| GM-REGIME | -0.0157 | -0.868 | 0.837 | 0.971 |
| SC-REGIME | -0.00710 | -1.39 | 1.38 | 0.992 |
| SED-REGIME | 0.0110 | -0.665 | 0.687 | 0.975 |
| SP-REGIME | -0.00376 | -1.046 | 1.038 | 0.994 |
| TA-REGIME | -0.00710 | -0.991 | 0.976 | 0.989 |
| ZO-REGIME | -0.00376 | -0.721 | 0.713 | 0.992 |
| DRY-REGIME | 0.203 | -0.811 | 1.22 | 0.695 |
| FIELD-REGIME | -0.0627 | -1.08 | 0.951 | 0.904 |
| SIA-REGIME | -0.00429 | -1.02 | 1.01 | 0.993 |
| WEIGH-REGIME | <0.0001 | -0.714 | 0.714 | 1.00 |
| CYB-DRY-REGIME | 1.13 | 0.0128 | 2.25 | 0.0480* |
| GM-DRY-REGIME | -0.194 | -1.40 | 1.01 | 0.752 |
| SC-DRY-REGIME | -0.313 | -2.14 | 1.50 | 0.737 |
| SED-DRY-REGIME | 0.0877 | -0.869 | 1.04 | 0.857 |

| | | | | |
|-------------------------|-----------|--------|-------|-------|
| SP-DRY-REGIME | 0.518 | -0.771 | 1.81 | 0.431 |
| TA-DRY-REGIME | -0.343 | -1.54 | 0.850 | 0.573 |
| ZO-DRY-REGIME | -0.176 | -1.90 | 0.838 | 0.734 |
| CYB-FIELD-REGIME | 0.149 | -0.972 | 1.27 | 0.795 |
| GM-FIELD-REGIME | -0.0465 | -1.25 | 1.16 | 0.940 |
| SC-FIELD-REGIME | 0.969 | -0.854 | 2.79 | 0.298 |
| SED-FIELD-REGIME | 0.0462 | -0.910 | 1.00 | 0.925 |
| SP-FIELD-REGIME | 0.443 | -0.846 | 1.73 | 0.501 |
| TA-FIELD-REGIME | 0.0873 | -1.11 | 1.28 | 0.886 |
| ZO-FIELD-REGIME | 0.259 | -0.755 | 1.27 | 0.617 |
| CYB-SIA-REGIME | -0.0562 | -1.18 | 1.06 | 0.922 |
| GM-SIA-REGIME | -0.0279 | -1.23 | 1.18 | 0.964 |
| SC-SIA-REGIME | 0.520 | -1.30 | 2.34 | 0.577 |
| SED-SIA-REGIME | 0.0742 | -0.882 | 1.03 | 0.879 |
| SP-SIA-REGIME | -0.0365 | -1.33 | 1.25 | 0.956 |
| TA-SIA-REGIME | 0.233 | -0.970 | 1.42 | 0.714 |
| ZO-SIA-REGIME | 0.0749 | -1.09 | 0.939 | 0.885 |
| CYB-WEIGH-REGIME | NA | NA | NA | NA |
| GM-WEIGH-REGIME | NA | NA | NA | NA |
| SC-WEIGH-REGIME | 0.164 | -1.51 | 1.84 | 0.848 |
| SED-WEIGH-REGIME | NA | NA | NA | NA |
| SP-WEIGH-REGIME | - <0.0001 | -1.07 | 1.07 | 1.00 |
| TA-WEIGH-REGIME | 0.0133 | -0.939 | 0.965 | 0.978 |
| ZO-WEIGH-REGIME | NA | NA | NA | NA |

1087