

**Bangor University**

## **DOCTOR OF PHILOSOPHY**

### **Turnover of S-containing amino acids (cysteine and methionine) in grassland soils and their availability to plants**

Wang, Deying

*Award date:*  
2021

*Awarding institution:*  
Bangor University

[Link to publication](#)

#### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

#### **Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

# **Turnover of S-containing amino acids (cysteine and methionine) in grassland soils and their availability to plants**

Deying Wang

2021

A thesis submitted to Bangor University  
in candidature for the degree Philosophiae Doctor

School of Natural Sciences,  
Bangor University,  
Bangor,  
Gwynedd,  
LL57 2UW



## **Declaration and consent**

---

### *Details of the Work*

I hereby agree to deposit the following item in the digital repository maintained by Bangor University and/or in any other repository authorized for use by Bangor University.

Author Name: Deying Wang

Title: Miss

Supervisor/Department: School of Natural Resources

Funding body (if any): Bangor University and China Scholarship Council (CSC)

Qualification/Degree obtained: Philosophiae Doctor

This item is a product of my own research endeavours and is covered by the agreement below in which the item is referred to as “the Work”. It is identical in content to that deposited in the Library, subject to point 4 below.

### *Non-exclusive Rights*

Rights granted to the digital repository through this agreement are entirely non-exclusive. I am free to publish the Work in its present version or future versions elsewhere.

I agree that Bangor University may electronically store, copy or translate the Work to any approved medium or format for the purpose of future preservation and accessibility. Bangor University is not under any obligation to reproduce or display the Work in the same formats or resolutions in which it was originally deposited.

*Bangor University Digital Repository*

I understand that work deposited in the digital repository will be accessible to a wide variety of people and institutions, including automated agents and search engines via the World Wide Web.

I understand that once the Work is deposited, the item and its metadata may be incorporated into public access catalogues or services, national databases of electronic theses and dissertations such as the British Library's EThOS or any service provided by the National Library of Wales.

I understand that the Work may be made available via the National Library of Wales Online Electronic Theses Service under the declared terms and conditions of use (<http://www.llgc.org.uk/index.php?id=4676>). I agree that as part of this service the National Library of Wales may electronically store, copy or convert the Work to any approved medium or format for the purpose of future preservation and accessibility. The National Library of Wales is not under any obligation to reproduce or display the Work in the same formats or resolutions in which it was originally deposited.

*Statement 1:*

This work has not previously been accepted in substance for any degree and is not being concurrently submitted in candidature for any degree unless as agreed by the University for approved dual awards.

Signed ..... (candidate)

Date .....



*Statement 2:*

This thesis is the result of my own investigations, except where otherwise stated. Where correction services have been used, the extent and nature of the correction is clearly marked in a footnote(s).

All other sources are acknowledged by footnotes and/or a bibliography.

Signed ..... (candidate)

Date .....

*Statement 3:*

I hereby give consent for my thesis, if accepted, to be available for photocopying, for interlibrary loan and for electronic storage (subject to any constraints as defined in statement 4), and for the title and summary to be made available to outside organisations.

Signed ..... (candidate)

Date .....

*Statement 4:*

I agree to deposit an electronic copy of my thesis (the Work) in the Bangor University (BU) Institutional Digital Repository, the British Library ETHOS system, and/or in any other repository authorized for use by Bangor University and where necessary have gained the required permissions for the use of third party material.

In addition to the above I also agree to the following:

1. That I am the author or have the authority of the author(s) to make this agreement and do hereby give Bangor University the right to make available the Work in the way described above.

2. That the electronic copy of the Work deposited in the digital repository and covered by this agreement, is identical in content to the paper copy of the Work deposited in the Bangor University Library, subject to point 4 below.

3. That I have exercised reasonable care to ensure that the Work is original and, to the best of my knowledge, does not breach any laws – including those relating to defamation, libel and copyright.

4. That I have, in instances where the intellectual property of other authors or copyright holders is included in the Work, and where appropriate, gained explicit permission for the inclusion of that material in the Work, and in the electronic form of the Work as accessed through the open access digital repository, or that I have identified and removed that material for which adequate and appropriate permission has not been obtained and which will be inaccessible via the digital repository.

5. That Bangor University does not hold any obligation to take legal action on behalf of the Depositor, or other rights holders, in the event of a breach of intellectual property rights, or any other right, in the material deposited.

6. That I will indemnify and keep indemnified Bangor University and the National Library of Wales from and against any loss, liability, claim or damage, including without limitation any related legal fees and court costs (on a full indemnity bases), related to any breach by myself of any term of this agreement.

Signature..... Date.....

## Thesis Summary

---

Sulphur (S) is one of the most important elements in nature with large quantities cycling through the biosphere, atmosphere and geosphere annually. In recent decades an increased frequency of S deficiency has been observed in soils from many regions of the world. Consequently, in agricultural systems with low S inputs from fertilizers and atmospheric deposition, plants must rely heavily on the release of S from soil organic matter. Typically, less than 10% of the soil S pool occurs as inorganic sulphate with most soil S held in an organic form. Sulphate, together with simple organic S compounds (low molecular weight organic S compounds such as cysteine (Cys) and methionine (Met)), are regarded as potentially being plant-available. However, little information is available on the mineralization and availability of soil organic S, and this has severely limited our capacity to understand the factors that regulate its persistence, bioavailability, and movement in soil. Therefore, the overall aims of this thesis were to explore the bioavailability and biodegradation of Cys and Met in grassland soils. This was achieved by first investigating the intrinsic dissolved organic sulphur (DOS) and inorganic sulphate concentrations in grassland soils (Chapter 3). The results revealed that DOS is the dominant fraction of dissolved S in grassland soils. These results enabled us to optimise the design of the following experimental chapters. The following experiment (Chapter 4) quantified microbial uptake and mineralization of Cys and Met using  $^{14}\text{C}$  labelling. My results revealed that  $^{14}\text{C}$ -labelled Cys and Met were directly and rapidly assimilated by soil microbes, with assimilation rates being an order of magnitude (or more) faster than microbial mineralisation rates measured via  $^{14}\text{CO}_2$  evolution. The considerable delay between uptake and mineralization indicates that the turnover of Cys and Met in soil solution was largely biotically mediated. In a subsequent experiment (Chapter 7), I investigated the concurrent mineralization of S, C and N from Cys and Met, as well as the influence of available C, N and S on this process. My results revealed that while a large proportion of added Cys-C and Met-C were used for microbial respiration and microbial biomass incorporation, N and S were excreted as by-products into soil ( $\text{NH}_4^+$  and  $\text{SO}_4^{2-}$ ). We assume this was due to the low C/N ratio of these amino acids (C/N ratio = 3 for Cys, and 5 for Met). This assumption was supported by the results that glucose addition promoted a more complete utilization of both amino acids, whereas nutrient addition had less effect. In Chapter 6, I quantified the gross S mineralization rates and the size of the total labile S pools in a closed incubation experiment (70 days) using a  $^{35}\text{S}$  isotope-pool dilution-based method. Another critical question is whether plants can acquire S containing amino acids from soil solution, and if so, how does this compare with the uptake rate of inorganic sulphate? To answer this question, I measured the root uptake of three individual S compounds, namely Cys, Met, and sulphate labelled with  $^{14}\text{C}$  or  $^{35}\text{S}$ , over short time periods (24 h) at an ecologically relevant concentration (100  $\mu\text{M}$ ; Chapter 8). Our results revealed that sulphate is preferred to S-containing amino acids by maize plants. A large proportion of the exogenously applied Cys (66%) and Met (73%) could have been taken up rapidly intact by the roots under sterile hydroponic conditions, indicating that they may supply a significant proportion of S to plants when microbial populations are low. I then studied the microbial-plant competition for S-containing amino acids in a rhizosphere context (Chapter 5). My results revealed that < 10% of the added amino acid- $^{14}\text{C}$  was captured by the plant, while the rhizosphere microbial community assimilated > 75%. The addition of inorganic N and S, not C, reduced the uptake of Cys and Met from soil by the maize plants, indicating that amino acid utilization may be regulated by inorganic N and S availability. In conclusion, this research provides a more detailed understanding of the turnover of Cys and Met in soils and their availability to maize plants. In addition, it demonstrated the importance of DOS, its rapid turnover in soils, and the intense microbial and plant competition which exists for this resource. The results will also help develop more accurate models of S cycling in soils.

# Thesis Content

<b>Declaration and consent .....</b>	<b>ii</b>
<b>Thesis Summary.....</b>	<b>vi</b>
<b>Thesis Content.....</b>	<b>vii</b>
<b>Acknowledgements .....</b>	<b>xiv</b>
<b>Abbreviations .....</b>	<b>xv</b>
<b>List of tables.....</b>	<b>xvi</b>
<b>List of figures.....</b>	<b>xxii</b>
<b>Chapter 1 .....</b>	<b>1</b>
<b>Introduction.....</b>	<b>1</b>
<b>1.1 Introduction.....</b>	<b>2</b>
<b>1.2 Thesis aims and objectives .....</b>	<b>4</b>
<b>1.2.1 Thesis aims.....</b>	<b>4</b>
<b>1.2.2 Thesis objectives.....</b>	<b>4</b>
1.2.2.1 Objective 1: Determine the size of the inorganic and organic S pools in grassland soils .....	4
1.2.2.2 Objective 2: Investigate the effects of nutrient amendment on the mineralization of cysteine and methionine in soil.....	5
1.2.2.3 Objective 3: Compare the uptake of organic and inorganic S by soil microorganisms and plants .....	5
<b>1.2.3 Experimental chapters.....</b>	<b>5</b>
<b>1.2.4 References.....</b>	<b>8</b>
<b>Chapter 2 .....</b>	<b>11</b>
<b>Literature Review .....</b>	<b>11</b>
<b>2. Literature review .....</b>	<b>12</b>
<b>2.1. The global sulphur cycle.....</b>	<b>12</b>
<b>2.1.1. Sulphur cycling in the atmosphere .....</b>	<b>12</b>
2.1.1.1. Sources of atmospheric sulphur .....	12
2.1.1.2 Deposition of atmospheric sulphur .....	13
<b>2.1.2 Sulphur cycling in freshwater ecosystems .....</b>	<b>14</b>
2.1.2.1 Sources of sulphur input into freshwater ecosystems .....	14
2.1.2.2 Content of sulphur in freshwater sediments.....	14
2.1.2.3 Transformations of sulphur in freshwater sediments .....	15
<b>2.1.3 Sulphur cycling in terrestrial ecosystems .....</b>	<b>16</b>

2.1.3.1 Input and output of sulphur in soil .....	16
2.1.3.2 Sulphur forms in soil .....	18
2.1.3.3 Sulphur transformations in soil .....	21
<b>2.1.4 Enzymes involved in sulphur cycle .....</b>	<b>27</b>
2.1.4.1 Enzymes involved in the soil sulphur cycle .....	27
2.1.4.2 Enzymes involved in plant sulphur cycling .....	29
<b>2.2 Nutrient cycling in the rhizosphere .....</b>	<b>30</b>
2.2.1 Plant-microbe interactions in the rhizosphere .....	31
2.2.2 Carbon input by plants into the rhizosphere .....	32
2.2.3 Secondary productivity by soil microorganisms in the rhizosphere .....	33
2.2.4 DOC flows in rhizosphere .....	34
2.2.5 DON flows in the rhizosphere .....	34
2.2.6 DOS flows in the rhizosphere .....	35
<b>2.3 The role of sulphur for plant nutrition .....</b>	<b>36</b>
2.3.1 Sulphur nutrition and sulphur fertilizers .....	36
2.3.2 Uptake and allocation of sulphate by plants .....	38
2.3.3 Sulphur metabolites in plant metabolism .....	40
<b>2.4 Sulphur analysis methods, including isotopic methods .....</b>	<b>41</b>
2.4.1 Extraction of sulphate and organic sulphur in soils .....	41
2.4.2 Determination of volatile sulphur compounds .....	42
2.4.3 Determination of microbial biomass sulphur .....	43
2.4.4 Determination of transformation rates of sulphur in soils .....	44
2.4.5 Quantification of S uptake by plant roots .....	45
<b>2.5 Summary .....</b>	<b>46</b>
<b>2.6 References .....</b>	<b>47</b>
<b>Chapter 3 .....</b>	<b>85</b>
<b>Dissolved organic and inorganic sulphate in soils along a grassland productivity gradient .....</b>	<b>85</b>
<b>3.1 Introduction .....</b>	<b>87</b>
<b>3.2 Material and Methods .....</b>	<b>88</b>
3.2.1 Site description and soil sampling .....	88
3.2.2 Soil chemical characteristics .....	89
3.2.3 Sorption of <sup>35</sup> S-Cys, Met and Na <sub>2</sub> SO <sub>4</sub> in sterile grassland soil .....	90
3.2.4 Statistics and data analysis .....	91
<b>3.3 Results and Discussion .....</b>	<b>91</b>
3.3.1 DOS and SO <sub>4</sub> <sup>2-</sup> concentration in soils along the grassland productivity gradient .....	91

3.3.2 Factors influencing DOS concentration in grassland soils .....	94
3.3.3 Sorption of <sup>35</sup> S-Cys, Met and Na <sub>2</sub> SO <sub>4</sub> in sterile grassland soil .....	95
3.4 Conclusions .....	98
3.5 Acknowledgements .....	98
3.6 References .....	98
Chapter 4 .....	103
Rapid microbial uptake and mineralization of cysteine and methionine along a grassland productivity gradient .....	103
4.1 Introduction .....	105
4.2 Materials and methods .....	107
4.2.1 Soil sampling and characteristics .....	107
4.2.1 S-amino acid depletion from soil .....	110
4.2.2 S-amino acid mineralization .....	111
4.2.3 Statistical and data analysis .....	111
4.3 Results .....	113
4.3.1 S-amino acid depletion from soil solution .....	113
4.3.1.1 Substrate depletion from soil solution affected by soil conditions .....	113
4.3.1.2 Substrate depletion from soil solution as affected by substrate concentration.....	116
4.3.2 Microbial C use efficiency for individual substrates.....	118
4.3.2.1 S-amino acid mineralization from autoclaved soils .....	118
4.3.2.2 S-amino acid mineralization affected by soil conditions .....	119
4.3.2.3 S-amino acid mineralization affected by substrate concentrations .....	121
4.3.3 Carbon turnover rates in soil .....	128
4.4 Discussion.....	129
4.4.1 Kinetics of Cys and Met uptake by soil microbial community .....	129
4.4.2 Dynamics of Cys and Met mineralization by soil microbial community .....	131
4.5 Conclusions .....	132
4.6 Acknowledgements .....	132
4.7 References .....	132
Chapter 5 .....	141
Competition for S-containing amino acids (cysteine and methionine) between soil microorganisms and maize roots in the rhizosphere .....	141
5.1 Introduction.....	143
5.2 Materials and methods .....	146
5.2.1 Soil sampling and chemical characterisation .....	146

5.2.2 Rhizosphere microbial uptake and mineralization of $^{14}\text{C}$ -Cys and Met .....	149
5.2.3 Rhizosphere microbial oxidation of Cys and Met in soil using $^{35}\text{S}$ labelling .....	150
5.2.4 Amino acid uptake by plants under sterile hydroponic conditions ( $^{14}\text{C}$ and $^{35}\text{S}$ ).....	151
5.2.5 Competition for Cys and Met between plant and soil microorganisms in the rhizosphere .....	152
5.2.5.1 Competition for $^{14}\text{C}$ - Cys and Met in the rhizosphere .....	152
5.2.5.2 Competition for $^{35}\text{S}$ -Cys and Met in the rhizosphere .....	153
5.2.6 Statistics and data analysis.....	155
5.3 Results .....	156
5.3.1 Use of isotopes for tracing Cys and Met utilization by microorganisms in the rhizosphere .....	156
5.3.1.1 $^{14}\text{C}$ -amino acid depletion from soil solution .....	157
5.3.1.2 $^{14}\text{C}$ -Amino acid mineralization in soil .....	162
5.3.1.3 Incorporation of $^{14}\text{C}$ -amino acids into microbial biomass .....	165
5.3.2 Amino acid- $^{35}\text{S}$ utilization by soil microorganisms in the rhizosphere .....	165
5.3.3 Amino acid uptake by maize plants under hydroponic conditions .....	168
5.3.4 Competition for $^{14}\text{C}$ from Cys and Met between plants and soil microorganisms in the rhizosphere as affected by nutrient amendment .....	169
5.3.5 Competition for $^{35}\text{S}$ from Cys and Met between plants and soil microorganisms as affected by sulphate amendment .....	172
5.4 Discussion.....	173
5.5 Conclusions .....	176
5.6 Acknowledgements .....	177
5.7 References.....	177
Chapter 7 .....	188
Tracing the mineralization rates of C, N and S from cysteine and methionine in a grassland soil: A $^{14}\text{C}$ and $^{35}\text{S}$ dual-labelling study .....	188
7.1 Introduction.....	190
7.2 Materials and methods .....	195
7.2.1 Study sites and soil sampling.....	195
7.2.2 Incubation experiments .....	196
7.2.3 Partitioning of the $^{14}\text{C}$ into different compartments.....	197
7.2.3.1 Microbial mineralization of $^{14}\text{C}$ -labelled Cys and Met.....	198
7.2.3.2 Determination of microbial biomass $^{14}\text{C}$ .....	198
7.2.4 Partitioning of the $^{35}\text{S}$ into different compartments .....	199
7.2.4.1 Production of volatile sulphur compounds from Cys or Met mineralization.....	199

7.2.4.2 Determination of microbial biomass $^{35}\text{S}$ .....	199
<b>7.2.5 Microbial biomass analysis .....</b>	<b>200</b>
<b>7.2.6 Statistical and data analysis .....</b>	<b>201</b>
7.2.6.1 Half-time of amino acid- $^{14}\text{C}$ mineralization.....	201
7.2.6.2 Half-time of amino acid- $^{14}\text{C}$ depletion in soil.....	201
7.2.6.3 Half-time of amino acid- $^{35}\text{S}$ depletion in soil .....	202
7.2.6.4 Half-time of microbial biomass turnover in soil .....	202
<b>7.3 Results .....</b>	<b>203</b>
<b>7.3.1 Amino acid-<math>^{14}\text{C}</math> mineralization in soil .....</b>	<b>203</b>
<b>7.3.2 Amino acid depletion from soil .....</b>	<b>206</b>
7.3.2.1 Amino acid- $^{14}\text{C}$ depletion from soil .....	206
7.3.2.2 Amino acid- $^{35}\text{S}$ depletion from soil .....	207
<b>7.3.3 Soil microbial biomass C and N.....</b>	<b>209</b>
<b>7.3.4 Turnover time of microbial biomass <math>^{14}\text{C}</math> and <math>^{35}\text{S}</math>.....</b>	<b>210</b>
7.3.4.1 $^{14}\text{C}$ tracer incorporation into microbial biomass and turnover .....	210
7.3.4.2 $^{35}\text{S}$ tracer incorporation into microbial biomass and turnover .....	211
<b>7.3.5 Inorganic nitrogen release from cysteine and methionine decomposition.....</b>	<b>213</b>
<b>7.4 Discussion.....</b>	<b>214</b>
<b>7.5 Conclusions and outlook.....</b>	<b>216</b>
<b>7.6 Acknowledgements .....</b>	<b>218</b>
<b>7.7 References.....</b>	<b>218</b>
<b>Chapter 8 .....</b>	<b>227</b>
<b>Short-term uptake and efflux of sulphur containing amino acids and inorganic sulphate by maize roots under sterile hydroponic conditions .....</b>	<b>227</b>
<b>8.1 Introduction.....</b>	<b>229</b>
<b>8.2 Materials and methods .....</b>	<b>233</b>
<b>8.2.1 Plant material and nutrient solution .....</b>	<b>233</b>
<b>8.2.2 Experiment 1: Plant uptake and partitioning of <math>^{14}\text{C}</math>-Cys and Met under hydroponic conditions .....</b>	<b>235</b>
<b>8.2.3 Experiment 2: Plant uptake and distribution of <math>^{35}\text{S}</math>-Cys, Met and sulphate under hydroponic conditions .....</b>	<b>236</b>
<b>8.2.4 Experiment 3: Efflux of <math>^{35}\text{S}</math>-labelled Cys, Met and <math>\text{SO}_4^{2-}</math> from maize roots .....</b>	<b>240</b>
<b>8.2.5 Experiment 4: A split root system-cycling of amino acid S and other nutrients between shoots and roots in young maize plants.....</b>	<b>243</b>
<b>8.2.6 Statistics and data analysis.....</b>	<b>245</b>
<b>8.3 Results.....</b>	<b>245</b>



8.3.1 Plant uptake and distribution of <sup>14</sup> C-labelled Cys and Met under sterile hydroponic conditions .....	245
8.3.2 Uptake and distribution of <sup>35</sup> S-labelled Met, Cys and sulphate by maize plant roots under sterile hydroponic conditions .....	247
8.3.3 Intact uptake of cysteine and methionine .....	249
8.3.4. Efflux of cysteine, methionine and inorganic sulphate from maize roots .....	250
8.3.5 Cycling of sulphur compounds between maize shoot and root via split root systems	253
8.4 Discussion.....	255
8.5 Conclusions and outlook.....	256
8.6 Acknowledgements .....	258
8.7 References.....	258
Chapter 9 .....	271
Summary, discussion and recommendations for future research .....	271
9.1 Introduction.....	272
9.2 Synthesis of findings .....	272
9.3 Methodological strengths and limitations .....	276
9.4 Future research .....	280
9.4.1 Determination the dynamics of inorganic and organic S compounds in soil solution	281
9.4.2 What specific members of microbial community play a dominant role in soil S cycling?.....	281
9.4.3 Predicting S availability from C/S ratio of plant residues.....	282
9.4.4 Is organic S a significant S source for plants <i>in situ</i> ?.....	283
9.4.5 Seasonal and spatial responses in microbial biomass sulphur .....	284
9.5 Concluding remarks .....	285
9.6 References.....	285
Appendix 1 .....	291
Chapter 6 Estimation of sulphur transformation rates in grassland soils using <sup>35</sup> S isotope pool dilution (This chapter is not complete due to author being unable to read <sup>32</sup> S data for the final isotope dilution model) .....	291
6.1 Introduction.....	293
6.2 Materials and methods .....	297
6.2.1 Field sites and soil sampling .....	297
6.2.2 Extraction of total labile S.....	298
6.2.3 Incubation experiment using <sup>35</sup> S isotope pool dilution .....	299
6.2.4 Compartment model .....	301
6.3 Results .....	302

<b>6.3.1 Labile sulphur pool size.....</b>	<b>302</b>
<b>6.3.2 Total labile S in relation to S concentration extracted by different extractants .....</b>	<b>302</b>
<b>6.3.3 Gross S transformation rates estimated with an established model.....</b>	<b>302</b>
<b>6.3.4 Effects of glucose or SO<sub>4</sub><sup>2-</sup> addition on S transformations.....</b>	<b>302</b>
<b>6.4. Discussion and conclusions.....</b>	<b>304</b>
<b>6.5 Acknowledgements .....</b>	<b>304</b>
<b>6.6 References.....</b>	<b>304</b>

---

## Acknowledgements

---

First of all, I would like to express sincere appreciation to my supervisor team: Davey Jones, Dave Chadwick, Paul Hill and Tida Ge for all of their expert guidance, assistance and immense encouragement throughout my four-year PhD. I am deeply grateful for the support and quick feedback you have given me over the past few years. Thank you for challenging me to think further and work harder. Working under your supervision has been truly a privilege, I have learned so much from you both professionally and personally that I will carry for the rest of my career. I would also like to thank Bangor University and China Scholarship Council for providing funding for my study.

I am grateful to many people without whose assistance this work could not have been completed. I would like to thank my committee members: Dr Paula Roberts and Dr Prysor Williams for their insightful advice and continuous help. I would also like to thank Joseph Cotton and Alan Jones for their assistance in my field work. Many thanks to Sarah Chesworth, Jonathan Roberts who offered me technical support in the laboratory.

I would also like to thank my fellow colleagues in ECW for their understanding and support throughout my study. I felt like an awkward alien when I first arrived in the group, but various social activities organised by ECW helped me settle quickly into my new study life. Thank you all for sharing your cultures with me, it has been such a pleasure working with an international group of individuals who are always kind and passionate. Particular thanks to Timothy David Peters, Francesca Brailsford, Lucy Greenfield, Paul Bryn Llewellyn George, Fiona M. Seaton, Rob Brown, Jinyang Wang, Karina Marsden and Emily Cooledge for their invaluable help.

Finally, I would like to express gratitude to my family, I am forever in debt to my parents for their unconditional trust, support and love. A special acknowledgement goes to my partner, who has provided endless encouragement and support, who is always willing to share my joys and sorrows of doing a PhD patiently. Thanks also to my cat Tommy, who didn't physically help me with my thesis, but has always been a spirit of courage, exploring and independence. I would never have done this without the help and support from you all.

## Abbreviations

---

**%** – Percentage

**°C** – Degree Celsius

**ANOVA** – Analysis of variance

**C** – Carbon

**CO<sub>2</sub>** – Carbon dioxide

**cm** – Centimetre (s)

**Cys** – Cysteine

**d** – Day (s)

**DOC** – Dissolved organic carbon

**DON** – Dissolved organic nitrogen

**DOS** – Dissolved organic sulphur

**DW** – Dry weight

**EC** – Electrical conductivity

**h** – Hour (s)

**h<sup>-1</sup>** – Per hour

**IC** – Ion chromatography

**ICP** – Inductively coupled plasma

**kBq** – Kilobecquerel

**KCl** – Potassium chloride

**kg** – Kilogram

**K<sub>2</sub>SO<sub>4</sub>** – Potassium sulphate

**L** – Litre

**LMW** – Low molecular weight

**Met** – Methionine

**mg** – Milligram (s)

**min** – minute (s)

**min<sup>-1</sup>** – Per minute (s)

**mL** – Millilitre

**mM** – Millimolar

**MW** – Molecular weight

**NaOH** – Sodium hydroxide

**Na<sub>2</sub>SO<sub>4</sub>** – Sodium sulphate

**N** – Nitrogen

**NH<sub>4</sub><sup>+</sup>** – Ammonium

**NO<sub>3</sub><sup>-</sup>** – Nitrate

**PCA** – Principal component analysis

**S** – Sulphur

**SEM** – Standard error of mean

**UK** – United Kingdom

**μM** – Micromolar

**w/v** – Weight per volume

## List of tables

---

<b>Table 3.1</b> General site and soil properties (samples collected in late May 2019). Values represent means $\pm$ SEM ( $n = 3$ ) for either topsoil (top) or subsoil (sub). Data is expressed on a dry soil weight basis.....	92
<b>Table 3.2</b> Major soil cation concentrations measured by ICP-OES (samples collected in late May 2019). Values represent means $\pm$ SEM ( $n = 3$ ) for either topsoil (top) or subsoil (sub). Data is expressed on a dry soil weight basis. NA indicates not applicable.....	93
<b>Table 3.3</b> Stepwise multiple linear regressions results.....	95
<b>Table 3.4</b> Correlation matrixes between DOS concentration and cations, ammonium, and MBC. ** and * represents significance at level $p < 0.01$ and $p < 0.05$ , respectively.....	97
<b>Table 4.1</b> Characteristics of the 5 grassland soils used in the experiment. Values represent means $\pm$ SEM, $n = 3$ . Superscript letters represent significant differences between soils at the $P < 0.05$ level. Data are presented on a dry weight basis.....	109
<b>Table 4.2</b> Coefficients of single first order curves describing the depletion of Cys and Met from soil solution over time in 5 contrasting grassland soils along an altitudinal productivity gradient. $A_1$ and $A_2$ are estimated pool sizes for fast and slow substrate uptake routes, and $K_1$ and $K_2$ are the rate constants for the fast and slow uptake routes, respectively. Half time values for pools $A_1$ and $A_2$ are represented by $t_{1/2}$ values and these are determined from $K_1$ and $K_2$ . Values represent means $\pm$ SEM ( $n = 3$ ).....	123
<b>Table 4.3</b> Influence of amino acid concentration on the coefficients of single (0.01 mM and 0.05 mM) and double (0.1-100 mM) first order curve fits describing the depletion of $^{14}\text{C}$ from	

soil solution over time.  $A_1$  and  $A_2$  are estimated pool sizes for fast and slow substrate uptake routes, and  $K_1$  and  $K_2$  are the rate constants for the fast and slow uptake routes, respectively. Half time values for pools  $A_1$  and  $A_2$  are represented by  $t_{1/2}$  values and these are determined from  $K_1$  and  $K_2$ , respectively. Values represent means  $\pm$  SEM ( $n = 3$ ).....124

**Table 4.4** Kinetic coefficients of single and double exponential models describing the depletion of  $^{14}\text{C}$ -Cys and Met from soil and loss in microbial respiration over time in 5 contrasting grassland soils along an altitudinal productivity gradient.  $AA_{\text{catabolic}}$  and  $AA_{\text{anabolic}}$  are estimated pool sizes for fast and slow substrate uptake routes, and  $K_3$  and  $K_4$  are the rate constants for the fast and slow uptake routes, respectively. Half-life values for pools  $AA_{\text{catabolic}}$  and  $AA_{\text{anabolic}}$  are determined from  $K_3$  and  $K_4$ , respectively. Values represent means  $\pm$  SEM ( $n = 3$ ).....125

**Table 4.5** Influence of amino acid concentration on the kinetic coefficients of double exponential models describing the depletion of  $^{14}\text{C}$ -Cys and Met from soil and loss in microbial respiration over time over a 48 h-incubation. Values represent means  $\pm$  SEM ( $n = 3$ ).....126

**Table 4.6** Influence of amino acid concentration on the uptake and mineralization of either Cys or Met in the first 30 minutes after substrate addition, and the percent of amino acid taken up which was respired as  $^{14}\text{CO}_2$  during the same time period. Values represent means  $\pm$  SEM ( $n = 3$ ).....127

**Table 4.7** Influence of amino acid concentration on microbial C use efficiency ( $\text{CUE}_{\text{mic}}$ ). Values represent means  $\pm$  SEM ( $n = 3$ ).....128

**Table 5.1** Summary of the main properties of the soil used in the experiments. Values represent means  $\pm$  SEM ( $n = 3$ ). Data are presented on a dry soil weight basis.....147

**Table 5.2** Kinetic coefficients of single (0.1 mM and 1 mM) and double (10 mM) first order curves describing the depletion of  $^{14}\text{C}$ -Cys from soil solution over a 24 h-incubation period in the presence of an excess of C (+ Glucose), inorganic N (+ N) or sulphate (+ S). Values

represent means  $\pm$  SEM ( $n = 3$ ).  $a_1$  and  $a_2$  are estimated pool sizes for fast and slow substrate uptake routes,  $b_1$  and  $b_2$  are the rate constants for the fast and slow uptake routes, respectively. Half-time values for each pool are calculated according to Eqn. 2.....160

**Table 5.3** Kinetic coefficients of single (0.1 mM and 1 mM) and double (10 mM) first order curves describing the depletion of  $^{14}\text{C}$  labelled Met from soil solution over a 24 h incubation period in the presence of an excess of C (+ Glucose), inorganic N (+ N) or sulphate (+ S). Values represent means  $\pm$  SEM ( $n = 3$ ).  $a_1$  and  $a_2$  are estimated pool sizes for fast and slow substrate uptake routes,  $b_1$  and  $b_2$  are the rate constants for the fast and slow uptake routes, respectively. Half-time values for each pool are calculated according to equation 2. Values represent means  $\pm$  standard deviation,  $n = 3$ .....161

**Table 5.4** Coefficients of double (0.1 and 1 mM) and single (10 mM) first order curve fits to depletion of  $^{14}\text{C}$ -cysteine from soil by microbial mineralization over a 24 h-incubation period in the presence of an excess of C (+ Glucose), inorganic N (+ N) or sulphate (+ S).  $c_1$  and  $c_2$  are estimated pool sizes for fast and slow phases of mineralization,  $d_1$  and  $d_2$  are the rate constants for the fast and slow phases of mineralization, respectively. Half-time values for each pool are calculated according to equation 2. Values represent means  $\pm$  standard deviation,  $n = 3$ .....163

**Table 5.5** Coefficients of double (0.1 and 1 mM) and single (10 mM) first order curve fits to depletion of  $^{14}\text{C}$ -methionine from soil by microbial mineralization over a 24 h incubation period in the presence of an excess of C (+ Glucose), inorganic N (+ N) or sulphate (+ S).  $c_1$  and  $c_2$  are estimated pool sizes for fast and slow phases of mineralization,  $d_1$  and  $d_2$  are the rate constants for the fast and slow phases of mineralization, respectively. Half-time values for each pool are calculated according to equation 2. Values represent means  $\pm$  standard deviation,  $n = 3$ .....164

<b>Table 5.6</b> Carbon and S yields for rhizosphere microorganisms and axenic maize plants after growth on two S-containing amino acids (Cys and Met, 0.1 mM) over a 24 h-period separately. Values represent means $\pm$ SEM ( $n = 3$ ). NA indicates not applicable.....	168
<b>Table 5.7</b> Percentage partitioning of $^{14}\text{C}$ label after the introduction of 0.1 mM $^{14}\text{C}$ -Cys into the maize rhizosphere over a 24 h incubation period in the presence of an excess of C (+ Glucose), inorganic N (+ N) or sulphate (+ S). Values represent means $\pm$ SEM ( $n = 3$ ). *Different letters of each row indicate significant differences ( $p < 0.05$ ) between treatments.....	170
<b>Table 5.8</b> Percentage partitioning of $^{14}\text{C}$ label after the introduction of 0.1 mM $^{14}\text{C}$ -Met into maize rhizosphere over a 24 h incubation period in the presence of an excess of C (+ Glucose), inorganic N (+ N) or sulphate (+ S). Values represent means $\pm$ SEM ( $n = 3$ ). *Different letters of each row indicate significant differences ( $p < 0.05$ ) between treatments.....	170
<b>Table 5.9</b> Partitioning of $^{14}\text{C}$ label after the introduction of $^{14}\text{C}$ -labelled cysteine or methionine into the maize rhizosphere for 24 h at three different concentrations. Values represent means $\pm$ SEM ( $n = 3$ ). Average plant shoot weight is 86.5 mg ( $n = 18$ ), and average plant root weight is 56.9 mg ( $n = 18$ ).....	171
<b>Table 5.10</b> Percentage partitioning of $^{35}\text{S}$ -Cys, Met or $\text{Na}_2^{35}\text{SO}_4$ (0.1 mM) between maize tissues and soil microorganisms in the rhizosphere over a 24 h-incubation period. Values represent means $\pm$ SEM ( $n = 3$ ). *Different letters of each row indicate significant differences ( $p < 0.05$ ) among treatments.....	171
<b>Table 6.1</b> Summary of the main pools used in the isotope dilution analysis.....	301
<b>Table 7.1</b> Physiochemical properties of the soil used in this study. Data is presented on a soil dry weight basis. Values represent means $\pm$ SEM ( $n = 3$ ).....	196



**Table 7.2** Kinetic coefficients of double first order exponential decay models describing the depletion of  $^{14}\text{C}$ -Cys or Met from soil by mineralization over a 168 h-incubation period in the absence (control) or presence of either glucose (+G) or nutrients (+NPS). The corresponding mineralization of  $^{14}\text{C}$ -glucose in the presence of Cys and Met is also presented.  $a_1$  and  $a_2$  are estimated pool sizes for the fast and slow phases of mineralization, and  $b_1$  and  $b_2$  are the rate constants for the fast and slow phases of mineralization, respectively.  $t_{1/2}$  values are the half-times for pool  $a_1$  and  $a_2$  determined from  $b_1$  and  $b_2$ . Values represent means  $\pm$  SEM ( $n = 3$ ). The  $R^2$  represents the goodness of fit to the experimental data.....204

**Table 7.3** Turnover of soil microbial biomass  $^{14}\text{C}$ -labelled Cys or Met in a grassland soil after either glucose-C (+G) or nutrient (+NPS) input. The corresponding mineralization of  $^{14}\text{C}$ -glucose in the presence of Cys and Met is also presented. A first order exponential decay equation Eqn. 5 was fitted to the experimental data. Values represent means  $\pm$  SEM ( $n = 3$ ). The  $R^2$  represents the goodness of fit to the experimental data.....212

**Table 7.4** Turnover of soil microbial biomass  $^{35}\text{S}$ -Cys or Met in grassland soil after glucose-C (+G) or nutrient (+NPS) input. A first order exponential decay equation Eqn. 5 was fitted to the experimental data. Values represent means  $\pm$  SEM ( $n = 3$ ). The  $R^2$  represents the goodness of fit to the experimental data..... 212

**Table 8.1** Parameters of amino acids (Cys and Met) and sulphate release from intact maize roots. Efflux data was fitted to Eqn. 2. The parameters  $a$  and  $c$  represent the  $^{35}\text{S}$  held in the cytoplasm and vacuole respectively, while  $b$  and  $d$  are the efflux rate constants for these two pools, respectively. Values represent means  $\pm$  SEM ( $n = 3$ ).....252

**Table 8.2** Translocation and utilization of three sulphur compounds by maize plants over a 24-h period. Both root compartments received the same amount of nutrient solution. Three

independent measurement from replicate plants were made for each treatment. Values  
represent means  $\pm$  SEM ( $n = 3$ ).....254

## List of figures

---

<b>Fig. 2.1</b> Sulphur cycle in agricultural soils (Chalk, Inácio and Chen, 2017).....	17
<b>Fig. 2.2</b> Simplified schematic representation of main processes involved in sulphur cycle within plant-soil-atmosphere systems. Modified from (Brown, 1982).....	22
<b>Fig. 3.1</b> Sorption of Cys, Met and Na <sub>2</sub> SO <sub>4</sub> to the soil's solid phase, panel A shows sorption onto autoclaved soils, panel B shows sorption onto oven heated soils. The legend is the same for both panels. Values represent means $\pm$ SEM ( $n = 6$ ).....	96
<b>Fig. 4.1</b> Percent deviation from perfect mixing after addition of <sup>14</sup> C-Cys or Met solution (0.01 mM) to autoclaved soils collected from 5 contrasting grassland sites. A 200 $\mu$ l of <sup>14</sup> C-amino acid (Cys or Met) was added to 1g of soil containing non <sup>14</sup> C labelled water. Percentages are based on the theoretical amount of <sup>14</sup> C available in solution assuming no biotic loss. The dashed line at 0% represents the point at which perfect mixing has occurred throughout the soil. Values above 0% (dotted line) indicates incomplete mixing of added solution and soil native soil solution (values above 0%), values below 0% indicates abiotic sorption. Lines represent first order exponential equations fit to the experimental data. Data points represent means $\pm$ SEM ( $n = 3$ ).....	114
<b>Fig. 4.2</b> Amount of <sup>14</sup> C-label remaining in soil solution after addition of a <sup>14</sup> C labelled Cys and Met (0.01 mM) to soil collected from 5 grassland sites of increasing elevation and decreasing productivity within 60 minutes, corrected from autoclaved soil data as described in material and methods section. Percentages are based on theoretical amount of <sup>14</sup> C available in solution assuming perfect mixing of the substrate with soil native water (Eqn. 1). Values represent	

means  $\pm$  SEM ( $n = 3$ ), lines represent fits of single first order exponential equations to the experimental data.....115

**Fig. 4.3** Deviation from perfect mixing when a  $^{14}\text{C}$ -Cys and Met solution (0.01 to 100 mM) is added to autoclaved soil containing non  $^{14}\text{C}$  labelled water. Percentages are based on the theoretical amount of  $^{14}\text{C}$  available in solution assuming perfect mixing of the substrate with the native soil solution. Deviations from 0% (dotted line) indicates incomplete mixing (values above 0) or abiotic sorption (values below 0%). Lines represent fits of first order kinetic equations (Eqn. 2) to the experimental data. Values represent means  $\pm$  SEM ( $n = 3$ ).....117

**Fig. 4.4** Amount of  $^{14}\text{C}$  label in soil solution after the addition of a  $^{14}\text{C}$  labelled Cys and Met to non-sterile grassland soil (site 1). Lines represent fits of single (0.01 to 0.1 mM) and double (1 to 100 mM) first order kinetic equations to the experimental data. Values represent means  $\pm$  SEM ( $n = 3$ ).....118

**Fig. 4.5** Kinetics of the  $^{14}\text{C}$  released from cumulative microbial respiration after addition of  $^{14}\text{C}$  labelled Cys or Met (0.01 mM) to autoclaved grassland soil (site 1) within 48h. Values represent means  $\pm$  SEM ( $n = 3$ ).....118

**Fig. 4.6** Amount of  $^{14}\text{C}$ -label remaining in soil solution and microbial biomass after the injection of  $^{14}\text{C}$  labelled Cys and Met (0.01 mM) to soil collected from five contrasting grassland sites. Values represent means  $\pm$  SEM ( $n = 3$ ). Lines represent fits a double first order kinetic equations to the experimental data.....120

**Fig. 4.7** Amount of  $^{14}\text{C}$ - Cys and Met remaining in soil solution and soil microbial biomass within 48 hours added at 6 concentrations ranging from 0.01 to 100 mM (soil 1). Data points represent means  $\pm$  SEM ( $n = 3$ ). Lines represent fits of single or double first order exponential equations to the experimental data.....122

**Fig. 5.1** Detailed photographs of the area of our study.....148

**Fig. 5.2** Schematic representation of the experimental apparatus showing three-leaf stage maize plant growing down a rhizotube containing grassland soil in which  $^{14}\text{C}$  or  $^{35}\text{S}$  labelled amino acid is injected through pre-drilled holes. The uptake of amino acid into either plant biomass, respiration, or soil microbial biomass and respiration was determined over a 24 h chase period.....154

**Fig. 5.3** Depletion of  $^{14}\text{C}$ -labelled cysteine or methionine and subsequent incorporation of the  $^{14}\text{C}$  amino acid-C into microbial biomass or respiration ( $^{14}\text{CO}_2$ ) as affected by the addition of excess C, N and S over a 24 h-incubation period at three amino acid concentrations (0.1 mM, 1 mM, 10 mM). Values represent means  $\pm$  SEM,  $n = 3$ .....159

**Fig. 5.4** Depletion of  $^{35}\text{S}$ -labelled cysteine or methionine in soil solution, the formation of extractable  $^{35}\text{S}$ -sulphate in soil and the subsequent incorporation of the  $^{35}\text{S}$  amino acid-S into microbial biomass over a 24 h-incubation period at three added amino acid concentrations (0.1 mM, 1 mM, 10 mM). Values represent means  $\pm$  SEM,  $n = 3$ .....167

**Fig. 6.1** Schematic representation of the two ongoing processes in our inorganic sulphate incubation experiment. On one hand, there is quick immobilization of  $^{35}\text{S}$  in newly formed microbial biomass, which will decay and again be released as inorganic sulphate. On the other hand, inorganic sulphate is also slowly turned into labile organic sulphur, which could be mineralized to inorganic sulphate again, this process is mediated by microorganisms. In our study, however, we only took immobilization and mineralization between inorganic sulphate and labile organic sulphur into account.....294

**Fig. 6.2** Changes with time in the recovery rate of extractable specific activity of  $^{35}\text{S}$  for soils 1, 5, 6 and 7 with or without substrate amendments. Values represent means  $\pm$  SEM ( $n = 3$ ).....303

<b>Fig. 7.1</b> Schematic representation of the microbial degradation of -S-containing amino acids in soil (biological mineralization process). In this study, microbial respiration and amino acid consumption were determined, as well as gross rates of inorganic N and S fluxes, namely, N mineralization, ammonium consumption, nitrification, nitrate consumption, S mineralization and sulphate consumption.....	194
<b>Fig. 7.2</b> Cumulative $^{14}\text{CO}_2$ evolution from a grassland soil after the addition of $^{14}\text{C}$ -labelled Cys or Met (1 mM). Lines represent fits of a double exponential first kinetic equation to the experimental data. Values represent means $\pm$ SEM ( $n = 3$ ).....	205
<b>Fig. 7.3</b> Amount of $^{14}\text{C}$ -label remaining in soil solution after addition of $^{14}\text{C}$ -Cys or Met to a grassland soil relative to the amount added at time zero (100%). Values represent means $\pm$ standard deviation ( $n = 3$ ). Lines represent fits of a double first order exponential decay equation to the experimental data ( $r^2 > 0.80$ in all cases; Eq. (3)).....	207
<b>Fig. 7.4</b> Amount of $^{35}\text{S}$ -label remaining in soil solution after addition of $^{35}\text{S}$ -labelled Cys or Met to grassland soil relative to the amount added at time zero (100%). Values represent means $\pm$ standard deviation ( $n = 3$ ). Lines represents fist of a double first order exponential decay equation to the experimental data ( $r^2 > 0.80$ in all cases; Eq. (4)).....	208
<b>Fig. 7.5</b> Dissolved organic carbon (TOC; mg C kg $^{-1}$ DW soil) and N (TDN; mg N kg $^{-1}$ DW soil) concentrations in grassland soil solution subjected to treatments of C or NPS addition. Values represent means $\pm$ standard deviation ( $n = 3$ ).....	209
<b>Fig. 7.6</b> The fraction of $^{14}\text{C}$ -C $_{\text{mic}}$ in soil after following the addition of amino acids to soil in the presence or absence of glucose-C or inorganic nutrients (+NPS). Values represent means $\pm$ SEM ( $n = 3$ ). Lines represent $^{14}\text{C}_{\text{mic}}$ obtained by the fumigation extraction method, adjusted with the extraction factor of 0.35.....	211

<b>Fig. 7.7</b> The fraction of $^{35}\text{S}$ labelled $\text{S}_{\text{mic}}$ after substrate addition. Values represent means $\pm$ SEM ( $n = 3$ ). Lines represent $^{35}\text{S}_{\text{mic}}$ obtained by fumigation extraction method, adjusted with the extraction factor of 0.35.....	212
<b>Fig. 7.8</b> Generation of ammonium and nitrate from added Cys and Met (1 mM) in soil solution over a 7-day period. Data expressed on a dry soil weight basis. Values represent means $\pm$ SEM ( $n = 3$ ).....	213
<b>Fig. 7.9</b> Schematic presentation of carbon, nitrogen and sulphur mineralization from the S-containing amino acid-cysteine in grassland soils. a: labelled amino acid uptake into soil microbial biomass; b: subsequent microbial mineralization.....	217
<b>Fig. 8.1</b> Potential routes for root uptake of carbon, nitrogen and sulphur from cysteine added to the soil solution.....	232
<b>Fig. 8.2</b> Growth stages of maize plants ( <i>Zea mays</i> L.) from seedling to the three-leave stage.....	234
<b>Fig. 8.3</b> Schematic representation of the maize microcosm into which $^{14}\text{C}$ -labelled amino acid (Cys or Met) was injected. Each polypropylene tube was filled with 20 ml of 10%-strength Long Ashton nutrient solution. The final concentration of the two sulphur-containing amino acids in nutrient solution were brought to 100 $\mu\text{M}$ separately. The $^{14}\text{C}$ -radioisotope was then injected into the 10%-strength Long Ashton nutrient solution in the polypropylene tubes. The $^{14}\text{C}$ -amino acid movement and incorporation into different root and shoot sections was measured after 24 h after the injection. $^{14}\text{CO}_2$ evolved from shoot compartment was trapped in 10 ml of 1 M NaOH.....	238
<b>Fig. 8.4</b> Schematic representation of the maize microcosm into which $^{35}\text{S}$ -labelled Cys, Met or inorganic sulphate was injected. The same experimental set up as experiment 1 was adopted here, except that NaOH traps were not placed inside Lock & Lock plastic containers. The final	

concentration of Cys, Met or inorganic sulphate in nutrient solution were brought to 100  $\mu\text{M}$  separately. The  $^{35}\text{S}$ -radioisotope was then injected into the 10%-strength Long Ashton nutrient solution in the polypropylene tubes. The  $^{35}\text{S}$  depletion from nutrient solution as well as the incorporation into plant root and shoot sections were measured after 24 h after the injection.....239

**Fig. 8.5** Schematic illustrating how  $^{35}\text{S}$  labelling were used to obtain an estimate of sulphur efflux by maize plants (*Zea mays* L.).....242

**Fig. 8.6** Schematic representation of the experimental apparatus showing the maize plant growing in a split root system with only ‘donor root’ exposed to radioisotope. This system was allowed to develop over a 24 h-period.....244

**Fig. 8.7** Distribution of  $^{14}\text{C}$ -labelled Met and Cys in maize plants after 24 h of a)  $^{14}\text{C}$ -Cys recovered in the shoots; b)  $^{14}\text{C}$ -Cys recovered in the roots; c)  $^{14}\text{C}$ -Cys respiration from the maize plant; d)  $^{14}\text{C}$ -Met recovered in the shoots; e)  $^{14}\text{C}$ - Met recovered in the roots; f)  $^{14}\text{C}$ -Met respiration from the maize plant. Bars and lines represent mean  $\pm$  SEM ( $n = 3$ ). Common lowercase letters denote a lack of significant difference between treatments (One way ANOVA: LSD test,  $p < 0.05$ ).....246

**Fig. 8.8** Partitioning of  $^{35}\text{S}$  label after the introduction of radio isotope into hydroponic maize plants for 24 h. a)  $^{35}\text{S}$ -Cys recovered in the shoots; b)  $^{35}\text{S}$ -Met recovered in the shoots; c)  $^{35}\text{S}$ - $\text{SO}_4^{2-}$  recovered in the shoots; d)  $^{35}\text{S}$ -Cys recovered in the roots; e)  $^{35}\text{S}$ -Met recovered in the roots; f)  $^{35}\text{S}$ - $\text{SO}_4^{2-}$  recovered in the roots. Values represent means  $\pm$  SEM ( $n = 3$ ).....248

**Fig. 8.9** Relationship between the total accumulation of  $^{14}\text{C}$  and  $^{35}\text{S}$  radiotracer ( $\text{nmol g}^{-1}$  root DW  $\text{h}^{-1}$ ) in maize roots and shoots after 24 h of exposure to the labelled solution containing 100  $\mu\text{M}$  cysteine or methionine ( $n = 9$ ).....250

**Fig. 8.10** Transmembrane sulphate movement.....251



**Fig. 8.11** Cumulative efflux of added  $^{35}\text{S}$ -Cys, Met or sulphate per unit dry mass of maize roots. Prior to measuring efflux, plants were pre-treated with radioisotopes for 60 minutes. Efflux was determined by measuring the increase of radioisotope in the root bathing medium solutions.  $\text{BaCl}_2$  was applied to separate organic and inorganic S in solution. Data represent means  $\pm$  SEM ( $n = 3$ ).....252

# **Chapter 1**

---

## **Introduction**

## 1.1 Introduction

Sulphur (S) is increasingly being recognized as the fourth major plant nutrient after nitrogen (N), phosphorus (P) and potassium (K). It is a key element ranking in importance with N and P in the formation of proteins (Scherer, 2001). To be more specific, plants require S for the synthesis of essential amino acids (cysteine and methionine) (Wirtz and Droux, 2005), peptides (e.g. glutathione (GSH) and phytochelatins (PCs)) (Ravilious and Jez, 2012), iron-sulphur clusters (Seeber, 2002), membrane sulfolipids (Shimajima, 2011), cell wall components, vitamins and cofactors, and other secondary metabolites (e.g. glucosinolates and alliins). S is mainly taken up as the inorganic sulphate ion ( $\text{SO}_4^{2-}$ ) against the root's electrochemical potential gradient through the activity of plasma membrane sulphate-specific transporters (De Kok *et al.*, 2017; Hawkesford, 2017). In addition, atmospheric sulphur dioxide can enter the leaf via stomata and be used as a source of S (Spedding, 1969; Van Der Kooij *et al.*, 1997). Therefore, S availability is of great importance for plant growth. However, many European countries are currently facing the problem of S deficiency in their soils as a result of reduced S deposition (Engardt *et al.*, 2017), a consequence of cleaner coal fired power stations, higher yields and consequent greater S removal, and decreased use of S containing fertilisers (McGrath, Zhao and Blake-Kalff, 2003).

Owing to its vital and indispensable role in various metabolic and enzymatic processes of plants, it is important to understand its typical and complex behaviour in soils to manage and use S sources more efficiently. Since soil is the primary source of nutrient supply for plants, it is, therefore, of great importance to understand the complete S cycle. Soil S occurs in organic matter and in mineral forms, among which more than 90% is organic bonded (Jamal, Moon and Abdin, 2010). Although not readily plant available, the large organic S fraction is an important source of S supply to plants due to its rapid mineralization to inorganic  $\text{SO}_4^{2-}$  (Ghosh, De and Maiti, 2018), which can thereafter be taken up by plants. Therefore, apart from

anthropogenic inputs of  $\text{SO}_4^{2-}$ , the mineralization of organic S to  $\text{SO}_4^{2-}$  may represent an equally important source of S for plants, especially in areas where there is a deficiency of available inorganic sulphate and organic S is the only S available. There is a great diversity of organic S compounds in soil, among which S-containing amino acids may constitute an important proportion (Scott, Bick and Anderson, 1981). Cysteine and methionine are the only two S-containing amino acids from twenty essential amino acids that make up proteins (Colovic *et al.*, 2018). Since the mineralization of cysteine and methionine to sulphate in soil represents a critical process in S availability to plants, they will be the main focus of the research described in this thesis.

More recently, a number of studies have confirmed that various plant species from diverse ecosystems such as boreal forests (Nordin, Högberg and Näsholm, 2001; Persson and Näsholm, 2001b; Lupi *et al.*, 2013), agricultural plants (Jämtgård, Näsholm and Huss-Danell, 2008; Gioseffi, de Neergaard and Schjørring, 2012), arctic plants (Kielland, 1994) as well as bromeliads plants (Endres and Mercier, 2003) have the capacity to take up amino acids and oligopeptides as a source of N, bypassing the common mineralization process and the need to take up ammonium ( $\text{NH}_4^+$ ) and nitrate ( $\text{NO}_3^-$ ). There is also evidence that cysteine and methionine can be actively transported into cultured tobacco cells (Wright, 1962; Harrington and Smith, 1977; Persson and Näsholm, 2001b), indicating plants may have active mechanisms for enhancing their access to amino acid-S in soil. While the root uptake of amino acids demonstrates the potential for plants to acquire organic-S, understanding the quantitative significance of this process in soil S cycling remains unknown. In addition, the concentration of amino acids in soil solution is known to vary both temporarily and spatially, suggesting that plants may acquire both organic and inorganic S depending on the local conditions. Further, whether plants can compete with soil microorganisms for amino acid-N and S sources in natural environments is still unclear.

## **1.2 Thesis aims and objectives**

### **1.2.1 Thesis aims**

Based on the knowledge gaps highlighted above, this PhD thesis mainly focuses on the potential importance of two S containing amino acids (cysteine and methionine; Cys and Met) as a S source in the soil-plant system. Radioisotope tracers ( $^{14}\text{C}$  and  $^{35}\text{S}$ ) were applied to quantify the turnover of these labile organic S compounds in a series of experimental chapters. This thesis quantified the mineralization of cysteine and methionine to inorganic sulphate in soil solution and measured the importance of abiotic and biotic factors affecting this process. In addition, this thesis also explored the direct uptake of Cys and Met by maize roots under sterile conditions, as well as the rhizosphere competition between plant roots and soil microorganisms for these amino acids in soil solution. The overall hypotheses were that 1) microbial mineralization of Cys and Met from soil solution is extremely rapid, releasing plant available inorganic N and S, and 2) plant roots have the capacity to compete with soil microorganisms for Cys and Met in soil solution to enhance their access to low molecular weight organic-S.

### **1.2.2 Thesis objectives**

#### *1.2.2.1 Objective 1: Determine the size of the inorganic and organic S pools in grassland soils*

Dissolved S in soil solution was fractionated into dissolved organic S (DOS) and inorganic sulphate. In chapter 3, the size of the DOS and sulphate pools in grassland soils were compared, providing evidence of the spatial variation of these S fractions. In chapter 4, DOS (cysteine and methionine) biodegradation rates were investigated using an isotopic labelling approach ( $^{14}\text{C}$ ), highlighting their capacity to supply S-containing amino acids to plants due to their high rates of replenishment in soil solution. In chapter 6, the size of the labile organic S (LOS) pool in grassland soils was estimated using a  $^{35}\text{S}$  isotope dilution technique.

#### *1.2.2.2 Objective 2: Investigate the effects of nutrient amendment on the mineralization of cysteine and methionine in soil*

Soil microorganisms play an important role in mineralizing organic S to inorganic forms that are potentially more plant available. In chapter 6, the mineralization and immobilization rates of S from grassland soils was estimated in a closed laboratory incubation system using an isotope-based method. In chapter 7, the concurrent mineralization of C, N and S from Cys and Met was investigated, and the effects of C and nutrient addition on this mineralization process was also studied.

#### *1.2.2.3 Objective 3: Compare the uptake of organic and inorganic S by soil microorganisms and plants*

Plant S nutrition is thought to mainly depend on the uptake of inorganic sulphate. However, research has demonstrated that S-containing amino acids (Cys and Met) are also plant-available. Chapter 8 addressed the question: to what extent can plants acquire S-containing amino acids, and how does this compare with that of (inorganic) sulphate under sterile conditions? In chapter 5, the competition between maize roots and soil microorganisms for three S compounds ( $\text{SO}_4^{2-}$ , Cys and Met) was quantified in an incubation experiment.

### **1.2.3 Experimental chapters**

A brief summary of each experimental chapter is provided below:

Chapter 3 investigated dissolved organic S (DOS) and inorganic sulphate in soils from across a grassland altitudinal gradient (0-400 m a.s.l.). Sulphate concentration was measured by ion chromatography (IC), total dissolved S (TDS) and other cations were analysed using ICP-OES. DOS concentration was obtained by the difference between TDS and inorganic sulphate. The hypotheses were that: (1) DOS is the dominant fraction of TDS in grassland soils; (2) there is a strong interrelationship between DOS and dissolved organic carbon (DOC) and

total dissolved N (TDN) in soil solution. The data demonstrated that DOS constituted 24 to 95% of TDS in all soils with the amount present strongly correlated with TDS, Al, B, Ba, Fe, K, Mn,  $\text{NH}_4^+$ , DON, DOC concentrations and pH in soil extracts.

Chapter 4 quantified the rapid microbial uptake and mineralization of  $^{14}\text{C}$ -labelled Cys and Met in a series of five contrasting grassland soils from an elevation gradient (32-402 m a.s.l.) with decreasing primary productivity. Substrate depletion from soil solution was measured using a centrifugal-drainage procedure, and the subsequent production of  $^{14}\text{CO}_2$  from microbial mineralization was measured by 1 M NaOH traps. The hypotheses were that: (1) turnover rates of Cys and Met are faster in soil from more productive and lower altitude grassland than in soil from less intensive, higher altitude grasslands; (2) DOS turnover rates estimated by the centrifugal-drainage procedure are higher than when determined by measuring mineralization rates. These results indicate that alongside  $\text{SO}_4^{2-}$ , Cys and Met could also be important sources of S for plants.

Chapter 5 evaluated the uptake and rhizosphere competition for Cys and Met between soil microorganisms and maize roots.  $^{14}\text{C}$  and  $^{35}\text{S}$  labelling was used to quantify the incorporation of amino acids into plant tissues or soil microorganisms. The hypotheses were that: (1) soil microorganisms and maize plants both possess a high capacity to utilize Cys and Met; (2) when allowed to compete, maize plants possess a lower capacity to capture free Cys and Met from soil solution compared to soil microorganisms. Results from the incubation experiment showed that soil microorganisms captured amino acids at around  $400 \text{ nmol } ^{14}\text{C} (\text{kg soil})^{-1} \text{ h}^{-1}$ , while plant roots incorporated amino acids at around  $200 \text{ nmol } ^{14}\text{C} (\text{g root DW})^{-1} \text{ h}^{-1}$ . When allowed to compete together in microcosms, maize plants captured around 10% of the  $^{14}\text{C}$ -labelled Cys and Met, while the rhizosphere microbial community captured more than 70%.

Chapter 6 explored the intrinsic rates of S turnover in seven contrasting soils along a natural grassland gradient using  $^{35}\text{S}$  isotope labelling and a pool-dilution model. In a closed incubation experiment (70 d), we quantified the temporal changes in extractable  $^{35}\text{SO}_4^{2-}$  and stable  $^{35}\text{SO}_4^{2-}$  concentrations. The hypotheses were that: (1) the size of the labile S pool derived from the model are compatible with measured extractable concentrations; (2) the addition of non-labelled sulphate would retard the incorporation of  $^{35}\text{S}$ -sulphate into the organic-S pool, while the addition of glucose would enhance this incorporation process.

Chapter 7 describes the importance of C and nutrient availability in regulating microbial decomposition of Cys and Met in a grassland soil using dual-isotope labelling with  $^{14}\text{C}$  and  $^{35}\text{S}$ . This allowed the concurrent mineralization of S, C and N during the incubation to be investigated. The hypotheses were that: (1) Cys and Met mineralization is a biological mediated process, with amino acid-C used for respiration and retained in the microbial biomass, and inorganic N and S released back into soil solution; (2) C and nutrient amendments regulate microbial decomposition of both Cys and Met due to altered microbial elemental stoichiometry. The results showed that microbial communities have an innate capacity to utilize Cys and Met, with inorganic N ( $\text{NH}_4^+$  and  $\text{NO}_3^-$ ) and  $\text{SO}_4^{2-}$  being generated as a result of microbial C acquisition, through which process both soil microorganisms and plants could potentially benefit.

Chapter 8 determined the short-term uptake and efflux of organic and inorganic S (Cys, Met,  $\text{SO}_4^{2-}$ ) by maize plants in sterile conditions. Plants have the ability to take up intact amino acids as an N source, but this has not been thoroughly studied for S-containing amino acids. The hypotheses were that: (1) Cys and Met can be taken up intact by plant roots under sterile hydroponic conditions, yet inorganic sulphate is preferred over organic S by maize plants; (2) root efflux of Cys and Met occurs through passive diffusion. Our results showed that a considerable proportion of Cys (66%) and Met (73%) could have been rapidly taken up intact



under sterile hydroponic conditions, when inorganic sulphate was available, indicating that S-containing amino acids could be important nutrient sources for plants under certain growing conditions.

#### 1.2.4 References

- Colovic, M. B. *et al.* (2018) ‘Sulphur-containing amino acids: protective role against free radicals and heavy metals’, *Current medicinal chemistry*. Bentham Science Publishers, 25(3), pp. 324–335.
- Endres, L. and Mercier, H. (2003) ‘Amino acid uptake and profile in bromeliads with different habits cultivated in vitro’, *Plant Physiology and Biochemistry*, 41(2), pp. 181–187. doi: 10.1016/S0981-9428(02)00025-6.
- Engardt, M. *et al.* (2017) ‘Deposition of sulphur and nitrogen in Europe 1900–2050. Model calculations and comparison to historical observations’, *Tellus B: Chemical and Physical Meteorology*. Taylor & Francis, 69(1), p. 1328945.
- Ghosh, P. K., De, T. K. and Maiti, T. K. (2018) *Role of ACC deaminase as a stress ameliorating enzyme of plant growth-promoting rhizobacteria useful in stress agriculture: A review, Role of Rhizospheric Microbes in Soil: Stress Management and Agricultural Sustainability*. doi: 10.1007/978-981-10-8402-7\_3.
- Gioseffi, E., de Neergaard, A. and Schjørring, J. K. (2012) ‘Interactions between uptake of amino acids and inorganic nitrogen in wheat plants’, *Biogeosciences*. Copernicus GmbH, 9(4), p. 1509.
- Harrington, H. M. and Smith, I. K. (1977) ‘Cysteine Transport into Cultured Tobacco Cells’, *Plant Physiology*, 60(6), pp. 807–811. doi: 10.1104/pp.60.6.807.
- Hawkesford, M. J. (2017) ‘Sulfate Transport in Plants: A Personal Perspective’, in *Sulfur*

- Metabolism in Higher Plants-Fundamental, Environmental and Agricultural Aspects*. Springer, pp. 3–12.
- Jamal, A., Moon, Y. S. and Abdin, M. Z. (2010) ‘Sulphur -a general overview and interaction with nitrogen’, *Australian Journal of Crop Science*, 4(7), pp. 523–529.
- Jämtgård, S., Näsholm, T. and Huss-Danell, K. (2008) ‘Characteristics of amino acid uptake in barley’, *Plant and Soil*, 302(1–2), pp. 221–231. doi: 10.1007/s11104-007-9473-4.
- Kielland, K. (1994) ‘Amino acid absorption by arctic plants: implications for plant nutrition and nitrogen cycling’, *Ecology*. Wiley Online Library, 75(8), pp. 2373–2383.
- De Kok, L. J. *et al.* (2017) *Sulfur Metabolism in Higher Plants-Fundamental, Environmental and Agricultural Aspects*. Springer.
- Van Der Kooij, T. A. W. *et al.* (1997) ‘Uptake and metabolism of sulphur dioxide by *Arabidopsis thaliana*’, *New Phytologist*, 135(1), pp. 101–107. doi: 10.1046/j.1469-8137.1997.00619.x.
- Lupi, C. *et al.* (2013) ‘Role of soil nitrogen for the conifers of the boreal forest: a critical review’, *International Journal of Plant & Soil Science*, 2(2), pp. 155–189.
- McGrath, S. P., Zhao, F. J. and Blake-Kalff, M. M. (2003) ‘History and outlook for sulphur fertilizers in Europe’, *Fertilizers Fertilization*, 2(15), pp. 5–27.
- Nordin, A., Högberg, P. and Näsholm, T. (2001) ‘Soil nitrogen form and plant nitrogen uptake along a boreal forest productivity gradient’, *Oecologia*, 129(1), pp. 125–132. doi: 10.1007/s004420100698.
- Persson, J. and Näsholm, T. (2001) ‘Amino acid uptake: A widespread ability among boreal forest plants’, *Ecology Letters*, 4(5), pp. 434–438. doi: 10.1046/j.1461-0248.2001.00260.x.

- Ravilious, G. E. and Jez, J. M. (2012) 'Structural biology of plant sulfur metabolism: From assimilation to biosynthesis', *Natural Product Reports*, 29(10), pp. 1138–1152. doi: 10.1039/c2np20009k.
- Scherer, H. W. (2001) 'Sulphur in crop production - Invited paper', *European Journal of Agronomy*, 14(2), pp. 81–111. doi: 10.1016/S1161-0301(00)00082-4.
- Scott, N. M., Bick, W. and Anderson, H. A. (1981) 'The measurement of sulphur-containing amino acids in some scottish soils', *Journal of the Science of Food and Agriculture*, 32(1), pp. 21–24. doi: 10.1002/jsfa.2740320105.
- Seeber, F. (2002) 'Biogenesis of iron-sulphur clusters in amitochondriate and apicomplexan protists', *International Journal for Parasitology*, 32(10), pp. 1207–1217. doi: 10.1016/S0020-7519(02)00022-X.
- Shimajima, M. (2011) 'Biosynthesis and functions of the plant sulfolipid', *Progress in Lipid Research*. Elsevier Ltd, 50(3), pp. 234–239. doi: 10.1016/j.plipres.2011.02.003.
- Spedding, D. J. (1969) 'Uptake of sulphur dioxide by barley leaves at low sulphur dioxide concentrations', *Nature*. Springer, 224(5225), pp. 1229–1231.
- Wirtz, M. and Droux, M. (2005) 'Synthesis of the sulfur amino acids: Cysteine and methionine', *Photosynthesis Research*, 86(3), pp. 345–362. doi: 10.1007/s11120-005-8810-9.
- Wright, D. E. (1962) 'Amino acid uptake by plant roots', *Archives of Biochemistry and Biophysics*, 97(1), pp. 174–180. doi: 10.1016/0003-9861(62)90061-9.

## **Chapter 2**

---

### **Literature Review**

## **2. Literature review**

### **2.1. The global sulphur cycle**

Sulphur (S) is an essential element for all living systems (Mitchell, 1996) and is closely associated with numerous fundamental biotic and abiotic geochemical reactions and cycling processes within the earth's atmosphere, hydrosphere, biosphere, and geosphere (Granat, Rodhe and Hallberg, 1976). This key role is a consequence of its large variety of different oxidation states (Charlson, Anderson and McDuff, 1992; Lomans *et al.*, 2002), ranging from completely reduced (oxidation state, -2) to completely oxidized (oxidation state, +6). Among these oxidation states, only three are abundant in nature, -2 in the form of sulfhydryl and sulphide, 0 in the form of elemental S, and +6 in the form of sulphate. These compounds are continuously converted to each other by a combination of biological (Rennenberg *et al.*, 1990; Rennenberg and Polle, 1994; Sors, Ellis and Salt, 2005; Davidian and Kopriva, 2010), biochemical (Kabdasli, Tünay and Orhon, 1995), and geochemical processes (Maher and Geoscience, 2013; Cao *et al.*, 2018), forming the global S cycle. Therefore, it is useful to consider both the amounts of S in these individual S states and the fluxes between these S pools.

#### **2.1.1. Sulphur cycling in the atmosphere**

##### *2.1.1.1. Sources of atmospheric sulphur*

S compounds are released into the atmosphere by both anthropogenic and natural processes (Shinn and Lynn, 1979; Galloway and Whelpdale, 1980; Last, 1982). Natural atmospheric sources of S include the following: ocean-derived sea spray containing sulphate, organic compounds produced from the microbial decomposition of organic matter, reduction of sulphate in oxygen depleted waters and soils, volcanoes, and forest fires (Bates *et al.*, 1992). Anthropogenic-derived sources of atmospheric S include the combustion of fossil fuels and the

smelting of non-ferrous ores (Cullis and Hirschler, 1980). Overall, the global emission of all S-containing gases is 251 million tonnes year<sup>-1</sup>, with about 41% being of anthropogenic origin (Castro, Williams and Ogram, 2000). Due to the ubiquitous presence of S in fossil fuels and metal sulphide deposits, man-made atmospheric S emissions have increased in step with industrialization (Rodhe, 1999; Smith *et al.*, 2011). Atmospheric S occurs in a range of forms, including SO<sub>2</sub> emitted during volcanic eruptions, microbial oxidation and from coal-fired power plants and metal smelters. In contrast, H<sub>2</sub>S is emitted by anaerobic bacteria in soils and sediments, from volcanoes, and deep-sea vents. Therefore the majority of SO<sub>2</sub> emitted to the atmosphere is related to human activities, while most H<sub>2</sub>S emissions are natural (Rasmussen and Kabel, 1974). In addition, dimethyl-sulphide (DMS) and other volatile S compounds produced by marine algae (Liss, Malin and Turner, 1993) and livestock (Zicari *et al.*, 2013) may also contribute to atmospheric S loading.

#### *2.1.1.2 Deposition of atmospheric sulphur*

After being transported and mixed by winds and turbulence, atmospheric S compounds are re-deposited on the land and water surface as either wet or dry deposition. Briefly, SO<sub>2</sub> becomes oxidized to sulphate where it electrostatically interacts with cations (e.g. NH<sub>4</sub><sup>+</sup>, Ca<sup>2+</sup>, H<sup>+</sup>) in small particulates or moisture droplets (Park, 2013; Fowler, 1980; Garland, 1978). Both wet and dry deposition processes are efficient in scavenging gaseous and particulate S from the atmosphere, with about half of the SO<sub>2</sub> emitted to the atmosphere removed by dry deposition, and the remainder oxidised to sulphate and removed by precipitation (Garland, 1978; Wałaszek, Kryza and Dore, 2013).

In the 1980s, the amount of S deposited annually in rainfall in SE England ranged from 10 to 15 kg S ha<sup>-1</sup> from coal-fired power stations, while the absorption of atmospheric-derived S by crops and soil was estimated to be 20 and 50 kg S ha<sup>-1</sup> y<sup>-1</sup> (Martin, 1980). However, due

to the move away from coal-fired power stations and S-containing fertilizers, very little S is now deposited directly onto UK agricultural fields ( $2\text{--}4 \text{ kg S ha}^{-1} \text{ y}^{-1}$ ; Battarbee *et al.*, 2014). Consequently, atmospheric S deposition can no longer meet crop S demand, resulting in potential crop quality and yields losses. Whilst uncertainties still exist in components of these S budgets, several studies have attempted to provide a holistic circulation of S within terrestrial, freshwater and marine systems (Möller, 1984; Hamilton, 1985; Liss *et al.*, 1997; Granat, 2013).

### **2.1.2 Sulphur cycling in freshwater ecosystems**

#### *2.1.2.1 Sources of sulphur input into freshwater ecosystems*

The weathering of S-containing rocks in catchments and the oxidation of organic S from terrestrial sources represent the main sources by which S enters freshwaters (Zak *et al.*, 2021). In addition, S may reach water bodies from atmospheric deposition, the application of S-based products to soil and their subsequent leaching and runoff (e.g. wastes, fertilisers, pesticides and fungicides. Hinckley *et al.*, 2020; Onwona-Kwakye *et al.*, 2020), and in irrigation water (Schuler *et al.*, 2019).

#### *2.1.2.2 Content of sulphur in freshwater sediments*

Previous work has established that the sediment and its interface with overlying water are major sites for S accumulation and transformation (Jørgensen, 1990; Holmer and Storkholm, 2001; Zhang *et al.*, 2018). In freshwater sediments, there are many different S compounds including inorganic and organic forms. Inorganic S in freshwater sediments occurs in a variety of forms such as pyrite, other iron sulphide compounds due to redox gradients and elemental S. The organic S fraction can be subdivided into three components: protein, sulphate ester, and non-protein carbon-bonded S (sulfolipids, vitamins, and a large diversity of S containing amino acids; King and Klug, 1982).

### 2.1.2.3 Transformations of sulphur in freshwater sediments

Freshwater sediments contain a high density and diversity of microorganisms which are involved in a variety of biogeochemical processes including both reductive and oxidative processes. The co-operative action of these gives rise to the S cycle in freshwater sediments. Although based on a mass balance budget, sediment S is predominantly present in an organic form (Mitchell *et al.*, 1984; Losher and Kelts, 1989; Couture *et al.*, 2016). The net mineralization of this organic S released  $\text{SO}_4^{2-}$  into the water column. In addition, despite low  $\text{SO}_4^{2-}$  concentrations (Holmer and Storkholm, 2001), the S cycle in freshwater sediments is dominated by  $\text{SO}_4^{2-}$  reduction in terms of the quantity of S transformed (King and Klug, 1982).

Sulphate reducing bacteria (SRB) are a complex physiological bacteria group (Castro, Williams and Ogram, 2000), including *Desulfovibrio*, *Desulfomicrobium*, *Desulfobulbus*, *Desulfobacter*, *Desulfococcus*, *Desulfosarcina*, *Desulfobacterium*, *Desulfonema*, *Desulfotomaculum* and *Thermodesulfobacterium* (Warren *et al.*, 2005). The groups are characterized by their use of sulphate as a terminal electron acceptor during anaerobic respiration. The first described and best known genus of S-reducing bacteria in sediments and other natural environments is *Desulfovibrio* (Postgate and Campbell, 1966; Odom and Peck Jr, 1981; Fauque *et al.*, 1988; Blumenberg *et al.*, 2006). The concentration of sulphate is a key factor controlling the rate of sulphate reduction in freshwaters. Further, the addition of sulphate to sediments enhances sulphate reduction and promotes the cycling of C, N, P and Fe (Schindler, 1981; Koschorreck, 2008; Chen *et al.*, 2016). Atmospheric precipitation supplies most of the sulphate to natural waters (Fisher *et al.*, 1968) with the concentration of sulphate in freshwater typically ranging from 10 to 500  $\mu\text{M}$ , which is much lower than in seawater (Holmer and Storkholm, 2001).

Sulphides produced during sulphate reduction may undergo re-oxidation by several processes: chemical oxidation with oxygen, bacterial oxidation under oxidized conditions,




phototrophic oxidation, anoxic chemical oxidation and bacterial oxidation under anoxic conditions (Holmer and Storkholm, 2001). Previous studies have shown that microorganisms from the genera *Sulfuricella* (*Betaproteo bacteria*) (Watanabe, Kojima and Fukui, 2014), *Thiobacillus* (*Betaproteobacteria*) (Haaijer *et al.*, 2006) and *Sulfuricurvum* (*Epsilonproteobacteria*) (Haaijer *et al.*, 2008) can undertake S oxidation. Further, aerobic S oxidizing bacteria (SOB) can grow autotrophically using inorganic S as an electron donor and play an important role in the conversion of inorganic S compounds in freshwater ecosystems. Previous research has shown that up to 90% of reduced S was re-oxidized to sulphate (Holmer and Storkholm, 2001), and this high rate of re-oxidation of reduced S compounds in freshwater sediments may revert the sediments from a sulphate sink to a source to the overlying water in some cases.

### **2.1.3 Sulphur cycling in terrestrial ecosystems**

#### *2.1.3.1 Input and output of sulphur in soil*

The terrestrial S cycle involves significant interactions between the pedosphere, the hydrosphere, the biosphere, and the atmosphere. In comparison to other elements (e.g. C, N, P), the study of S in soils has received relatively little attention since adequate amounts were supplied from rock weathering, S fertilizers, irrigation water and from atmospheric deposition (volcanoes, atmospheric pollution, and sea spray). Consequently, until recently, there has been little economic drive to fund research on S cycling. Vegetation can also provide an important source of S in soil ecosystems because living roots release organic and inorganic S into soils (i.e., root exudation), and on plant death, decay of plant tissues releases S back to the soil. In addition, S is added to soil due to the widespread use of S-containing fungicides and insecticides (Griffith, Woodrow and Seiber, 2015). A simplified illustration of the sulphur cycle in agricultural soils is depicted in Fig. 2.1.



**Fig. 2.1** Sulphur cycle in agricultural soils (Chalk, Inácio and Chen, 2017).

S is removed from soil systems by leaching, erosion and surface runoff (Barrows and Kilmer, 1963), S volatilisation (Cullis and Hirschler, 1980; Peñuelas *et al.*, 2014), and uptake by plants (Plante, 2006). Of these, most S is removed from agroecosystems by crop harvesting, although a large amount of S may ultimately be returned to the soil via their residues or in livestock waste. S losses can also occur via leaching of sulphate alongside base cations (Ercoli *et al.*, 2012). Other forms of S (elemental S and organic S) are less mobile in soil (Riley, Zhao and McGrath, 2002), and must be converted to sulphate before significant leaching can occur. The amount of sulphate leaching is dependent on a range of factors including S status of the soil, texture, weather patterns, Fe content, O<sub>2</sub> status and soil structure. In addition, volatile forms of S are emitted from soils (e.g. H<sub>2</sub>S from waterlogged soils supplemented with sulphate; Fitzgerald, 1976). Other forms of S are generated and emitted from soils under both aerobic and anaerobic conditions, e.g. CH<sub>3</sub>-SH, CH<sub>3</sub>-CH<sub>2</sub>-CH<sub>2</sub>-SH has been shown to be produced following addition of cysteine to soils, and CH<sub>3</sub>-SH, CH<sub>3</sub>SCH<sub>3</sub> and CH<sub>3</sub>-S-S-CH<sub>3</sub> during methionine degradation in soils (W. L. Banwart and Bremner, 1975).

#### 2.1.3.2 Sulphur forms in soil

S is present in soils as both organically-bonded S and inorganic sulphate. Organically bonded S provides the major S reservoir and accounts for about 90% of total S in most soils (Eriksen *et al.*, 1998; Scherer, 2009; Freney *et al.*, 1969; Houle and Carignan, 1992; Kertesz and Mirleau, 2004; Yang *et al.*, 2007), from which plant-available sulphate is released upon microbial oxidation of organic-S. In general, inorganic S species can be operationally fractionated into water-soluble sulphate, adsorbed sulphate, dilute-HCl soluble S, volatile S, and pyritic S (Johnson, Henderson and Todd, 1981; Scherer, 2009). In contrast to inorganic S, the organic S pool contains a diverse mixture of compounds contained within organic matter (SOM) and the microbial biomass (Jamal, Moon and Abdin, 2010). Organic S can be further

divided into two distinct fractions: ester S as well as carbon-bonded S such as S-containing amino acids (Freney, Melville and Williams, 1970; Landerst, David and Mitchell, 1983; Scherer, 2009).

**(1). Ester sulphates:** Organic forms of S are a heterogeneous mixture of humus and plant residues, and can be subdivided into two main fractions according to the susceptibility to reduction by hydroiodic acid (HI), namely, (i) carbon-bonded S (directly C-S bonded), and (ii) ester organic S (S is bonded to oxygen or N). Ester organic S is easily hydrolysed to inorganic sulphate by mild physical or chemical treatments as well as extracellular enzymes termed arylsulphatases, therefore, they can be considered to be more biologically active (labile) in soil (Scherer, 2001). As the measurement of this form involves reduction with hydroiodic acid (HI), it is often referred to as HI-reducible-S. The organic sulphate fraction includes compounds containing C-O-S linkage (ester sulphate), C-N-S linkage (sulfamates) and N-O-S linkage (sulphated thioglycosides) (Fitzgerald, 1976), and constitutes 30-75% of the total organic S in soil. Ester sulphates include compounds such as choline sulphate, sulphated polysaccharides and phenolic sulphates (Edwards, 1998).

**(2). Carbon-bonded S:** C-bonded S constitutes up to 30% of the organic S in soil. C-bonded S includes S-amino acids (e.g. methionine, taurine, cystathionine, cysteine and cystine), polypeptides and proteins (Fe-S proteins called ferredoxins), and other S-containing compounds such as biotin, thiamine, coenzyme A as well as lipoic acid (Brosnan and Brosnan, 2006; Cronan, 2014; Colovic *et al.*, 2018). S-containing amino acids constitute an important proportion of the soil organic S (Norman M. Scott and Anderson, 1976), however, estimates of cysteine and methionine in soil hydrolysates are generally low because (i) S-containing amino acids are rapidly degraded and metabolized to sulphate in soil; and (ii) amino acids are mostly present as protein or peptides rather than as monomers, making their estimation difficult.

In addition to anthropogenic inputs of sulphate, the mineralization of organic S also represents an important source of sulphate in soils (McGill and Cole, 1981). Both ester sulphates and C-bonded S can be mineralized to plant-available sulphate, and it is generally accepted that soil microorganisms (including bacteria, archaea, and fungi) play a major role in this mineralization process (Kertesz and Mirleau, 2004a; McLaren and Swift, 1977). Previous studies have shown that S present in cysteine can be converted to sulphate by a mixed population of soil microorganisms, with cysteinesulphinic acid, cysteic acid, sulphite and  $\beta$ -hydroxypyruvic as intermediate forms (Freney, 1960). In addition, various fungi and bacteria can convert methionine to sulphate, although in terms of a source of soil sulphate, methionine is thought to be less readily mineralized than cysteine (Fitzgerald and Watwood, 1988). Therefore, for soils lacking significant S or  $S^{2-}$  oxidative capacity, mineralization of S-containing amino acids may represent an important mechanism for sulphate generation.

**(3). Microbial biomass-S:** The soil microbial biomass (i.e. bacteria, fungi and protozoa) generally accounts for ca. 1.5-3% of the total soil organic S (Saggar, Bettany and Stewart, 1981; Chapman, 1987; Smith and Paul, 1990). It plays a vital role in the S cycle as both a source and sink of plant-available S and is frequently considered to be relatively labile and a highly active pool of S turnover in soil (Banerjee and Chapman, 1996; Stevenson and Cole, 1999). The rate at which microbial biomass-S turns over in soil is an important parameter for soil S modelling. One might expect microbial biomass-S to turnover on an annual timeframe since microbial cells assimilate C, N and S in a relatively fixed ratio (Spohn, 2016; Spohn *et al.*, 2016). However, there is evidence that the microbial biomass C:S ratio is not fixed and can vary somewhat (Khan and Joergensen, 2019), therefore microbial biomass-S could turnover at a different rate from that of microbial biomass C.

The major S forms in microbial cells are proteins, amino acids, and other minor metabolites and essential vitamins (Moat, Foster and Spector, 2003). In addition, the microbial


biomass is a major driving force regulating the nutrient transformations in soil (Wu and Syers, 1997). By lysing microbial cells with chloroform, microbial biomass-S can be measured in a similar way to the measurement of biomass C and N (Voroney, Brookes and Beyaert, 2007). Plant residues are a major source of organic inputs to soil, exerting a primary role in maintaining SOM, microbial biomass and activity, and the size of the soil nutrient pool. Following addition of plant residues to soil, the microbial community grows rapidly in response to the addition of labile C, often inducing a rapid immobilization of available soil nutrients, particularly N, P and S by the microbial biomass (Wu, O'Donnell and Syers, 1993). In addition, mineralization of N, P and S may occur. Ultimately, this depends on the stoichiometric C:N:P:S ratio of the residues.

**(4). Inorganic sulphate:** Inorganic forms of S are usually much less abundant than organic S and can be divided into four major categories: (i) highly mobile sulphate in soil water; (ii) adsorbed sulphate, which can be desorbed; (iii) sulphate minerals, which may precipitate and re-dissolve (Barber, 1995); and (iv) sulphate, which is irreversibly retained (Wang *et al.*, 2011). The content and availability of sulphate in soil is continually undergoing changes due to: 1) rapid mineralization and immobilization by soil microbes; 2) leaching; 3) additions from the atmosphere in gaseous form or from wet deposition, and 4) additions from application of fertilisers and livestock waste (e.g. urine, faeces, manure, slurry).

#### *2.1.3.3 Sulphur transformations in soil*

Generally, nutrients are first released into soil due to mineral weathering (dissolution and desorption; Comerford, 2005), as well as from SOM mineralization (Nannipieri, Kandeler and Ruggiero, 2002). Soluble S can then move through the soil solution to the root surface by means of mass flow and diffusion where it can be taken up. Plants mainly acquire S primarily through the assimilation of sulphate, although small amounts of SO<sub>2</sub> can be acquired from the

atmosphere via stomatal uptake (Garsed and Read, 1977). When plant residues and livestock remains are returned to the soil, they are decomposed by microorganisms with the liberation of inorganic S. These inorganic S compounds may be oxidised to sulphates or reduced to hydrogen sulphide.



**Fig. 2.2** Simplified schematic representation of main processes involved in sulphur cycle within plant-soil-atmosphere systems. Modified from (Brown, 1982).

A simplified schematic representation of the main processes involved in sulphur cycle within plant-soil-atmosphere systems is shown in Fig. 2.2. S compounds undergo many transformations in soils as a result of biological activities including plants, animals and microorganisms, among which the microbial community plays a crucial role in the

transformation of various elements such as S, N, C and Fe due to the combined action of mineralization and immobilization. The key reactions include: 1) mineralization, the decomposition of organic S into inorganic sulphate; 2) immobilization, the assimilation of these inorganic and organic S compounds into microbial tissues; 3) oxidation, the conversion of reduced, inorganic forms (sulphides, thiosulphates, polythionates and elemental S) to sulphate; 4) reduction, the conversion of sulphate and other oxidised forms of S into sulphides; 5) sulphate adsorption-precipitation phenomena; 6) leaching. The importance of these processes is dependent on the form of S present, the physico-chemical environment, and the size, diversity and physiological state of the soil microbial community. Microbial activity is affected by many soil physical and chemical properties, especially pH, temperature, moisture availability, plant type, agricultural management regime etc. (White, 1959; Freney and Spencer, 1960).

**(1). Mineralization of S:** In well-drained soil surface layers, the amount of inorganic sulphate is often too small to provide adequate S for sustained rapid plant growth (Scherer, 2001), so plants may be largely dependent upon the conversion of soil organic S to sulphate for satisfactory S nutrition. Soil organic S must be broken down into smaller molecules by extracellular enzymes (e.g. proteases) before further mineralization can occur. S mineralization is the transformation of organic S into  $\text{SO}_4^{2-}$  and  $\text{CO}_2$  (McGill and Cole, 1981). The biological process is considered to be driven by the microbial consumption of organic C in response to microbial energy demand, thus S which is directly bonded to C in soil organic compounds (carbon-bonded S, e.g. S containing amino acids) is released as  $\text{SO}_4^{2-}$  (Blum et al., 2013). In contrast, sulphate can be released due to the cleavage of ester sulphates by extracellular or periplasmic microbial sulphatases (T. Sherene, 2017). This biochemical process is controlled by the supply of S rather than the need for energy, thus when soil sulphate is too low to meet microbial demand, sulphate esters are hydrolysed by sulphatases (McGill and Cole, 1981).



Mineralization of soil organic S has been shown to contribute substantially to plant S uptake and leaching (Pirela and Tabatabai, 1988; Sakadevan, Mackay and Hedley, 1993; Gahan and Schmalenberger, 2014; Coyne, Mikkelsen and Mineralization, 2015). An open system of incubation showed that soil microbes could mineralise C-bonded S to satisfy their S or possibly N requirements, and C-bonded S was identified as a major contributor to mineralised  $\text{SO}_4^{2-}$  (Ghani, 1989). Therefore, under circumstances where the storage of sulphate is too low to provide plants with adequate S for plant growth, the conversion of organic S to sulphate by the action of soil microbial becomes critical.

**(2). Immobilization of S:** S applied to soil is subject to physical/electrostatic (sorbed onto soil particles) or biological immobilization. Biological S immobilization is the process by which mineral S is incorporated into the microbial biomass (Freney, 1967; Shahsavani and Gholami, 2009). Since C availability greatly controls microbial activity, the extent to which sulphate-S could be immobilized is determined by both the amount of available-S and the availability of labile C. Addition of labile low-molecular-weight (MW) organic compounds such as glucose can result in a rapid increase in microbial biomass and thus S immobilization (Vong et al., 2008, 2010; Kertesz and Mirleau, 2004a; Scherer, 2009). Incubation studies (Freney et al., 1971; Freney et al., 1975) have also shown that  $^{35}\text{S}$ -labelled sulphate is incorporated into two major organic S fractions in soil, namely hydriodic acid reducible S, in which S is not directly bonded to carbon and is reduced to hydrogen sulphide by hydriodic acid, and soluble C-bonded S (e.g. Cys and Met). This is in line with other studies (Williams and Donald, 1957) showing that applied sulphate can be readily recovered in SOM over longer timescales.

**(3). Oxidation of inorganic sulphur:** Microbial oxidation of S is an important process where reduced forms of S are present, and involves the microbial oxidation of reduced inorganic S forms (e.g. sulphides, elemental S, thiosulfates) to higher oxidation states (e.g. sulphate). Microbial S oxidation is generally beneficial to soil in different ways: (1) solubilized inorganic

salts contain plant nutrients and thereby increase the level of soluble phosphate, potassium, calcium and magnesium for plant nutrition; (2) the acidity produced by oxidation of S can also be used to improve alkaline soils especially under warm and wet conditions (Germida and Janzen, 1993).

Although chemical oxidation of S occurs in soils, this process is mainly microbially mediated and largely involves three groups of microorganisms: (1) chemoautotrophic S bacteria (*Thiobacteriaceae*, *Beggiatoaceae* and *Achromatiaceae*); (2) photosynthetic S bacteria (*Thiorhodaceae* and *Chlorobacteriaceae*); and (3) certain heterotrophic microorganisms including some actinomycetes, bacteria and fungi (e.g. *Pseudomonas* spp., *Arthrobacter* spp., *Flavobacterium* spp. and *Bacillus* spp; Kuenen and Beudeker, 1982; Vidyalakshmi et al., 2009). Of these, *Thiobacilli* are chemoautotrophic bacteria who gain energy from oxidising reduced forms of S (Vidyalakshmi, Paranthaman and Bhakayaraj, 2009). Five species of the genus *Thiobacillus* are regarded as important in S oxidation in soils, viz. *T. denitrificans*, *T. ferrooxidans*, *T. thioparus* and *T. thiooxidans*.

**(4). Reduction of sulphate:** Sulphate can be directly reduced to sulphide by sulphate reducing bacteria (SRB) via either assimilatory or dissimilatory pathways (Kushkevych *et al.*, 2020). Both assimilatory and dissimilatory reduction of sulphate begin with the activation of sulphate by adenosine triphosphate (ATP). Adenosine phosphosulphate (APS), formed by ATP and sulphate, is then catalysed by the enzyme ATP sulphurylase (Abdulina *et al.*, 2020). In dissimilatory reduction, the sulphate moiety of APS is directly reduced to sulphite ( $\text{SO}_3^{2-}$ ) by the enzyme APS reductase, while in assimilatory reduction, another P atom is added to APS to form phosphoadenosine phosphosulphate (PAPS), PAPS is then reduced to sulphite (Romero *et al.*, 2014). Once sulphite is formed, it is oxidized to sulphide by the enzyme sulphite reductase. The assimilatory pathway generates reduced S compounds for biosynthesis of amino acids and proteins, and therefore does not lead to direct excretion of sulphide. In contrast, in

dissimilatory reduction, sulphate is reduced to inorganic sulphide by obligatory anaerobic sulphate reducing bacteria (Koschorreck, 2008).

Sulphate can be reduced to hydrogen sulphide ( $\text{H}_2\text{S}$ ) by SRB, reducing the availability of S for plant nutrition. The conversion of sulphate to  $\text{H}_2\text{S}$  may seem undesirable from a soil fertility perspective, but it could be of great significance in soils under alkaline and anaerobic conditions, as the acidity produced through this process can be used to lower soil pH. The predominant sulphate reducing bacteria genera in soils are *Desulfovibrio*, *Desulfotomaculum*, and *Desulfomonas* (Luptakova, 2007), all of which excrete the enzymes desulphurases or bisulphate reductase, which are responsible for reduction of S (Meena and Improvement, 2018).

**(5). Sulphate adsorption-precipitation phenomena:** The mobility and concentration of sulphate in soil solution can be affected by adsorption and desorption reactions with the solid phase (De Vries and Breeuwsma, 1987). Sulphate can be sorbed to mineral surfaces (e.g. Fe/Al hydroxides) and organic matter in acidic soils (Schoenau and Malhi, 2008), leading to greater retention and reduced leaching losses (Curtin and Syers, 1990; Eriksson, 1988). Sulphate sorption varies widely between soils and is strongly affected by soil pH (Muktamar, 1993), and sulphate concentration (Haque and Walmsley, 1973). Sulphate adsorption and desorption is also important in buffering soil systems against extreme climatic events (Moldan *et al.*, 2012). Sulphate sorption-desorption curves are typically used to predict the concentration-dependent availability of sulphate in soil and thus plant availability (Ghosh and Dash, 2012).

**(6). Leaching:** Unlike phosphate, sulphate  $\text{SO}_4^{2-}$  is weakly retained on soil surfaces and is therefore more susceptible to leaching than P in most soils (Barrow, 1978). The amount of S leaching depends on management practices and various soil characteristics, such as the surface charge characteristics of the soil, which can be manipulated by the addition of soil amendments (e.g., P and lime) (Bolan *et al.*, 1988). Addition of lime (with associated pH increase) and P

fertiliser decreases sulphate sorption by increasing the net negative charge of the soil as well as blocking sorption sites. They can also promote S mineralization, thus increasing sulphate movement in soil (Chao, Harward and Fang, 1962). Sulphate is susceptible to leaching and may be lost to the subsoil (Churka Blum et al., 2013). In comparison to topsoils, our knowledge of S cycling in subsoils is very poor. Sulphate leaching down the soil profile mainly occurs in the autumn and winter in response to an increase in seasonal rainfall, however, cover crops can be used to reduce these losses in arable systems (Lehmann and Schroth, 2003).

#### **2.1.4 Enzymes involved in sulphur cycle**

A wide range of enzymes are excreted by plant roots and soil microorganisms into soil where they are involved in the cycling of S. The following section summarises the role of these enzymes.

##### *2.1.4.1 Enzymes involved in the soil sulphur cycle*

**(1). Carbon-S lyases:** The release of sulphate from organic S may proceed by two mechanisms: oxidation of the C skeleton to yield energy with sulphate being released as a by-product; or directly after hydrolysis of ester sulphate linkages comprising the organic S. There are two enzymes that belong to the family of lyases, specifically the class of carbon-S lyases. Methionine gamma-lyase (MGL) catalyses the formation of methanethiol,  $\alpha$ -ketobutyrate and ammonia from L-methionine (Esaki *et al.*, 1979; Johnston *et al.*, 1981). Cysteine desulphydrase is an enzyme which catalyses the degradation of L-cysteine to pyruvate, ammonia and hydrogen sulphide (Kumagai *et al.*, 1974). These enzymes have been found in a range of bacteria belonging to the genera *Escherichia*, *Aerobacter*, *Serratia*, *Proteus*, *Alcaligenes*, *Agrobaaterium*, *Miarococcus* and *Sarcina*.

**(2). Sulphatases:** Sulphatases release sulphate from sulphate esters and have been the focus for much soil S research (Gahan and Schmalenberger, 2014). In soils, different types of

sulfatases occur: arylsulfatases (ARS), alkylsulfatases, steroidsulfatases, glucosesulfatases, chondrosulfatases, and myrosulfatases (Schinner *et al.*, 2012). Of these, ARS is the key enzyme involved in soil organic S mineralization by hydrolysing ester S to sulphate (Tabatabai and Bremner, 1970; Cooper, 1972). It has been detected in plants, animals and soil microorganisms (Stressler *et al.*, 2016). In soils, enzymes are located either in living cells (intracellular enzymes) or outside cells (extracellular enzymes). By use of the chloroform fumigation extraction methods, studies have shown that ca. 57% of the total arylsulphatase was intracellular with the remainder being extracellular (Klose and Tabatabai, 1999). However, increased microbial arylsulfatase activity may not lead to increased amounts of sulphate accumulation, indicating that sulfatase enzyme activity is not the rate-limiting step in the biochemical release of sulphate-S from sulphate esters (Ganeshamurthy and Nielsen, 1990).

The activity of microbial and soil-extracted arylsulfatase increases with increasing sulphate concentration, while barley root arylsulfatase activity can be repressed by sulphate, indicating that arylsulfatase produced by soil microorganisms and plants are controlled differently (Ganeshamurthy and Nielsen, 1990). Current evidence suggests that the activities of ARS are influenced by a range of soil chemical and biological properties. For example, a study in boreal forest soils revealed that maximum sulfatase activity occurred below the humus layer, possibly due to the pH optimum of sulfatase being more typical of subsoils than highly acidic topsoils (Wittmann *et al.*, 2004). Arylsulphatase activity was also found to be positively correlated with the total soil C, total organic S and reducible organic S in soil (Cooper, 1972).

**(3). Rhodanese:** Rhodanese (Thiosulphate cyanide sulphurtransferase) is another enzyme involved in the S cycle which has been detected and characterized in soil and which catalyses the formation of thiocyanate from  $\text{S}_2\text{O}_3^{2-}$  and cyanide according to the following reaction:  $\text{S}_2\text{O}_3^{2-} + \text{CN}^- = \text{SCN}^- + \text{SO}_3^{2-}$  (Dick and Deng, 1991).

#### 2.1.4.2 Enzymes involved in plant sulphur cycling

In higher plants the enzymes of assimilatory sulphate reduction are localized mainly in the chloroplasts of green leaves, where the reduction is strongly dependent on light intensity (Chen *et al.*, 2018). S assimilation starts with the formation of adenosine phosphosulphate (APS), a reaction catalysed by the enzyme ATP sulphurylase, which has been found in many higher plant species (Osslund, Chandler and Segel, 1982; Lunn *et al.*, 1990; Hatzfeld *et al.*, 2000; Herrmann *et al.*, 2014). APS is then phosphorylated by another ATP molecule to form adenosine 3'-phospho-5'-phosphosulphate (PAPS), which is catalysed by the enzyme APS kinase (Coughlan, 1977). Another inorganic S compound produced by the assimilatory reduction of sulphate by photosynthetic organisms is sulfide, which can be incorporated into an amino acid skeleton to form cysteine (Ngo and Shargool, 1974). Cysteine biosynthesis is driven by the sequential reaction of two enzymes: 1) serine acetyl transferase (SAT), which synthesizes the intermediary product O-acetylserine (OAS) from acetyl-CoA and serine; and 2) OAS (thiol) lyase (OASTL), which incorporates sulfide into OAS, producing cysteine, requiring pyridoxal-5'-phosphate as cofactor (Wirtz and Droux, 2005). Depending on their location in the cytosol, plastid, or mitochondrion, a variety of isoforms of SAT and OASTL exist in plants (Romero *et al.*, 2014).

Atmospheric S can enter plants via their stomata where it readily reacts with water to form sulphite, including  $\text{HSO}_3^-$  and  $\text{SO}_3^{2-}$  (Rennenberg, 1984). At low concentrations (e.g. < 0.25 ppm), plants are able to utilize  $\text{SO}_2$ , however, above a certain threshold (e.g. > 0.25 ppm), which differs between plant species,  $\text{SO}_2$  toxicity often leads to visible negative effects such as chlorosis, necrosis and long-term yield reduction (Swain and Padhi, 2015). Therefore, plants have developed mechanisms to control sulphite levels by the action of various enzymes, including sulphite oxidase (SO), a key enzyme for protecting plants against  $\text{SO}_2$ , catalysing the reaction  $\text{SO}_3^{2-} + \text{H}_2\text{O} \rightarrow 2\text{H}^+ + \text{SO}_4^{2-} + 2\text{e}^-$  (Brychkova *et al.*, 2007). Sulphite oxidase

belongs to the class of Mo cofactor (Moco) containing enzymes that catalyse electron redox reactions (Hille, 1996), and is essential for detoxifying excessive amounts of sulphite in the cell (Lang *et al.*, 2007).

## **2.2 Nutrient cycling in the rhizosphere**

The rhizosphere is a realm where complex biological and ecological processes occur, encompassing the few millimetres of soil which surrounds the root (Curl and Truelove, 2012). An abundant population of archaea, bacteria, protists, fungi and animals live here alongside plant roots, the activities of each influencing those of the others across spatial and temporal scales (Cardon and Whitbeck, 2011). Rhizosphere soil effectively forms an interface layer between roots and the surrounding soil. As roots and soil act as both sources and sinks for a diverse range of compounds, this interface layer of soil mediates large fluxes of soluble and volatile compounds (Belnap, Hawkes and Firestone, 2003).

Rhizodeposition was first defined as all material lost from plant roots, including water soluble exudates, secretions of insoluble materials, lysates, dead fine roots, and gases such as CO<sub>2</sub> and ethylene (Lynch and Whipps, 1990). In a broad sense, the processes by which C enters the soil e.g. root cap and border cell loss, death and lysis of root cells (cortex, root hairs etc.), flow of C to root-associated symbionts living in the soil, gaseous losses, leakage of solutes from living cells (root exudates), and insoluble polymer secretion from living cells are collectively known as rhizodeposition (D. L. Jones, Nguyen and Finlay, 2009). Rhizodeposition is central to a diverse range of functions in plant nutrition and soil ecology (Cardon and Whitbeck, 2011). Some of these compounds are able to improve nutrient availability, e.g. phytosiderophores for Fe, organic acids for P and phenolics for micronutrients (Carvalhais *et al.*, 2011), while organic acids can relieve Al<sup>3+</sup> rhizotoxicity (Heim *et al.*, 2001). Furthermore, apart from serving as an important C and energy source for rhizosphere soil

microorganisms, root exudates also act as allelochemicals, or as signalling substances for the establishment of symbiotic relationships between plant roots and microorganisms (Paterson, 2003). Yet despite the involvement in many soil ecological functions, due to the complex pathways and chain of reactions, our knowledge of the amount, composition and turnover of these root-derived compounds, especially those that influence S cycling is still very limited.

Although the quantities of organic compounds exuded from roots is not large (Kuzyakov and Domanski, 2000), they exert a very strong influence on soil microorganisms and may be significant in affecting plant nutrient availability (Rovira, 1969a; Bertin, Yang and Weston, 2003). As soon as a plant germinates, the root starts releasing root exudates (e.g. sugars, amino acids) into the rhizosphere, with the exact composition and quantity excreted dependent upon plant species, age, temperature, light intensity, plant nutritional status, microorganisms, soil bulk density, soil moisture, and root damage (Curl and Truelove, 2012). Root exudates are often divided into two classes: (1) low MW compounds such as amino acids, organic acids, sugars, phenolics, and other secondary metabolites, and (2) high MW compounds, such as mucilage (polysaccharides), peptides and proteins (Bais *et al.*, 2006).

### **2.2.1 Plant-microbe interactions in the rhizosphere**

The rhizosphere microbial community contributes strongly to nutrient cycling, plant growth and root health as they (1) decompose SOM and release inorganic nutrients to the plant; (2) store nutrients in, and release nutrients from the microbial biomass; (3) affect nutrient availability by undertaking solubilisation, chelation, oxidation and reduction reactions; (4) affect plant growth by the release of stimulating or inhibiting substances. Interactions involving plants roots in the rhizosphere include root-root, root-insect and root-microbe interactions, which may be classified as positive, negative associations and neutral associations. Positive interactions include symbiotic associations with mycorrhizal fungi, and root colonization by



bacterial biocontrol agents and plant growth-promoting bacteria (PGPB). Negative interactions include competition or parasitism among plants, pathogenesis by bacteria or fungi, and invertebrate herbivory (Bais *et al.*, 2006). The factors that determine whether the chemical signature of a plant's root exudates will be perceived as a "negative" or a "positive" signal is only beginning to be understood. Due to the release of organic C and N from the root, microorganisms are generally more numerous than in the bulk soil, such that there is fierce competition for nutrients. Soil microorganisms in the rhizosphere can be roughly divided into two categories, those that compete with the plant root for N, P, S, and other nutrients; and others that provide plant inorganic nutrients by the decomposition of SOM.

### **2.2.2 Carbon input by plants into the rhizosphere**

Attempts have been made to achieve total C balances for the soil-plant system (Kuzyakov and Domanski, 2000; Kuzyakov, 2006). Typically, labelling of photo-assimilates with  $^{13}\text{CO}_2$  or  $^{14}\text{CO}_2$  is used to trace C flow through the plant into the soil and further transformations in the plant-microbial-soil system monitored. From this, it has been estimated that the amount of C lost in root exudation ranges from 1 to 3% of a plant's net fixed C (Pinton, Varanini and Nannipieri, 2007). Previous estimates suggest that ca. 50% of rhizosphere respiration is due to the turnover of rhizodeposits and 50% to direct root (and mycorrhizal) respiration (Kuzyakov, 2002). Root and microbial respiration can alter the pH of the soil solution and can lead to the creation of anoxic and hypoxic hotspots in soil. This C flow can also flow beyond the rhizosphere by 2 main mechanisms: (1) investment of C in the production of intra- and extra-radical mycorrhizal structures which can extend many cm from the root (mycorrhizosphere; Leake, 2004); (2) release of volatile C that can travel very long distances in soil.

Root exudates can be classified into four groups depending on their mode of delivery: 1) water soluble exudates (e.g. sugars, amino acids, hormones, vitamins) which are lost by passive diffusion and whose loss rate strongly depends upon the concentration gradient between the root cytoplasm and soil solution and the permeability of the plasma membrane (Neumann and Römheld, 2007; Vranová *et al.*, 2010); 2) secretions, such as polymeric carbohydrates and enzymes which are actively excreted (Maseko and Dakora, 2013); 3) lysates, released when cells autolyse (Oburger and Jones, 2018); 4) gases such as ethylene and CO<sub>2</sub> (Jones, Hodge and Kuzyakov, 2004). These groups cover all stages of plant growth and development with the balance of these various processes changing with age of the plant, although water soluble exudates are thought to dominate rhizosphere C flow (Wen *et al.*, 2006).

### **2.2.3 Secondary productivity by soil microorganisms in the rhizosphere**

Microbial growth in the rhizosphere is stimulated by the continual input of readily assimilable organic substrates from roots. In ecological terms, the substrate flowing from the roots is the product of photosynthesis and is thus primary productivity, and the utilization of these organic substrates by rhizosphere microorganisms results in secondary productivity. As the biggest reservoir of terrestrial C, SOM contains more organic C than global vegetation and the atmosphere combined (Lehmann and Kleber, 2015). Plant residues, including roots and rhizodeposits, quantitatively represent the largest inputs of C to soil (Schmidt *et al.*, 2011). In return, SOM retains nutrients, which improves plant growth and protects water quality. To date, SOM research has mainly focussed on the topsoil (0-30 cm) (Rumpel, Eusterhues and Kögel-Knabner, 2004; Song *et al.*, 2005; Hati *et al.*, 2007), yet an emphasis should also be given to deeper soil C to gain a better understanding of the key factors controlling soil C storage. Key processes regulating C dynamics are controlled by microorganisms at the microscale, but the related processes are often studied at larger scales.

#### **2.2.4 DOC flows in rhizosphere**

Dissolved organic matter (DOM) is a continuum of organic molecules of different sizes, charges and structures that pass through a 0.45 µm-pore size filter (Bolan *et al.*, 2011). DOM plays an important role in soil formation, mineral weathering and transport of pollutants, and the transport of nutrients from soil to surface waters. Dissolved organic C (DOC) is commonly used as a proxy for DOM (Wang *et al.*, 2012a). DOC is an important fraction of the C pool in soils, and an understanding of DOC dynamics requires a knowledge of its origin, function, and fate. Generally, DOC contains a wide range of molecules, ranging from simple amino acids through to complex high MW materials (Neff and Asner, 2001), which are released from vegetation and SOM. DOC concentrations can vary considerably both spatially and temporally with rates of production dependent on inputs from vegetation and SOM, consumption by soil organisms, sorption by mineral particles and water movement through the soil. As it is relatively mobile, DOC plays an important role in the short- and long-distance transport of N, P and S.

#### **2.2.5 DON flows in the rhizosphere**

Dissolved organic N (DON) refers to a diverse array of compounds which can be operationally grouped into high (>1 kDa) and low (<1 kDa) MW components. It is now clear that when soil N mineralization is slow and concentrations of inorganic N are low, plants can take up DON directly from soil (e.g. amino acids, peptides) and that this may contribute significantly to the plant's N demand (Kielland, 1994; Persson and Näsholm, 2001b, 2001a). Therefore, an understanding of DON fluxes between rhizosphere microorganisms and plant roots is critical to quantifying the significance of DON in terrestrial ecosystems (Christou *et al.*, 2005).

DON flow at the soil-root interface is bidirectional with N being lost from roots by root exudation at the same time that DON is also taken up from the soil (Ford *et al.*, 2007; D. L. Jones, Nguyen and Finlay, 2009). Clearly, it is the net balance between these two processes that determines the amount of organic N captured by the root from the soil. The magnitude of N flow into the rhizosphere is dependent upon a wide range of biotic and abiotic factors including plant species, plant ecotype/cultivar, age and environmental conditions (D. L. Jones, Nguyen and Finlay, 2009). As free amino acids and proteins represent only a minor component of root exudates (typically 1–2%), it has been assumed that they contribute little to plant N rhizodeposition (D. L. Jones, Nguyen and Finlay, 2009). Although the size of the DON pool is important, it is the flux rate (pool replenishment rate), however, that is more critical in understanding the importance of DON to plant nutrition (Christou, Avramides and Jones, 2006; Rousk and Jones, 2010). Typically, low MW DON turns over rapidly in soil, persisting from minutes to hours. This has led to the hypothesis that the bottleneck in soil organic N cycling lies in the breakdown of higher MM DON to smaller units that can be directly assimilated by the microbial biomass (Lucy M. Greenfield *et al.*, 2020). Once released into the soil DON undergoes a number of fates including movement away from the root due to diffusion and mass flow, capture by soil microorganisms, and sorption to the soil solid (Sacchi *et al.*, 2000).

#### **2.2.6 DOS flows in the rhizosphere**

Dissolved organic S (DOS) makes up a significant part of the S cycle in both terrestrial and aquatic environments (Wang *et al.*, 2012a). Whilst numerous studies have focused on DOC, DON and DOP in soil and freshwater, our knowledge of DOS remains poor, particularly in a rhizosphere context (Marschner, 1992; Wang *et al.*, 2012). Mineralization of DOS in the rhizosphere can be divided into biological and biochemical processes (McGill and Cole, 1981). C-bonded S is thought to be mineralized by biological processes with sulphate released as a

by-product during the oxidation of the C skeleton to CO<sub>2</sub> (Kertesz and Mirleau, 2004b). Although the evidence base is weak, more ester bonded S was found in non-rhizosphere (bulk) soil than in the rhizosphere (Hu *et al.*, 2002), while no significant differences existed between C-bonded S in the rhizosphere and in the bulk soil, indicating that C-bonded S may be more important for plant S nutrition (Hu *et al.*, 2003).

## **2.3 The role of sulphur for plant nutrition**

### **2.3.1 Sulphur nutrition and sulphur fertilizers**

Although required in smaller quantities than other macro elements, S is hugely important in plant nutrition, yet it has received comparatively less attention. This is due to the cheap availability of S fertilizers and atmospheric S inputs which have replenished soil S reserves. During the 25 years, however, S deficiency has become more widespread in the UK and other European countries (Murphy and Boggan, 1988; McGrath, Zhao and Blake-Kalff, 2003; Kühn-institut, 2019b) as a result of strict controls on industrial emissions (Campbell and Smith, 1996). In addition, changes away from S fertilisers (e.g. ammonium sulphate) to low-S or even S-free fertilisers has also led to decreased inputs of S to soils (Ceccotti, Morris and Messick, 1998).

S is a constituent of three essential amino acids namely cysteine, cystine and methionine (Schnug, Haneklaus and Murphy, 1993) which are fundamental to protein/enzyme synthesis. Insufficient S supply therefore affects both the quality and the quantity of plants (Schnug, 1990; Filipek-Mazur *et al.*, 2019) as it prevents protein synthesis and the production of many secondary metabolites (Ahmad *et al.*, 2007). For example, S deficiency reduces nutritional quality due to a re-direction of protein synthesis to low S vicilin proteins at the expense of legumin, which is rich in S-containing amino acids, cysteine and methionine (Byers and Bolton, 1979; Millerd, Thomson and Randall, 1979). The baking quality of wheat is thus

impaired by S deficiency because the S-containing amino acids in the gluten fraction of flour are responsible for the elasticity of the dough and the bread volume (Wrigley *et al.*, 1984). S deficiency can cause phenotypic symptoms that become first visible at chlorosis in leaves, because S is required in the photosynthetic processes. The increasing recognition of the importance of S for plant growth and crop yield, as well as nutritional importance of S for human and animal diets, has led to an increased emphasis on the understanding of S metabolism in plants.

Studies have shown that S fertilization increases N incorporation into protein synthesis, thereby enhancing photosynthetic rates (Ahmad and Abdin, 2000), indicating that S can directly affect the use efficiency of other nutrients. This could be explained by the law of minimum, that balanced supply of all nutrients is the prerequisite of correct crop nutrition (Jungk, 2009). In addition, S can improve the availability of Ca, Mg, P, Cu, Mn and Zn, as acidity produced on oxidation of reduced inorganic S in soil can promote their solubility (Karimizarchi *et al.*, 2014). Leaves can also take up volatile S from the atmosphere (e.g. SO<sub>2</sub> and H<sub>2</sub>S) via the stomata and incorporate this S into S organic compounds (Rennenberg and Polle, 1994). In addition, leaves can also emit S in the form of H<sub>2</sub>S when leaves or roots are exposed to relatively high concentrations of sulphate (Rennenberg *et al.*, 1990).

S enters soils in both inorganic and organic fertiliser. Inorganic S fertilizers mainly comprise sulphates and elemental S. Sulphate fertilizers (e.g. gypsum, ammonium sulphate) are highly soluble and provide an immediate source of S to the plants, whereas elemental S contains very high concentrations of S (70-100%) and is poorly soluble. Elemental S fertilisers have the potential advantage of offering a continual release of S during the growing season with minimal leaching losses (Boswell and Friesen, 1993; Riley, Zhao and McGrath, 2002).

In ecosystems with low mineral fertilizer inputs, organic wastes (e.g. manures, compost, slurry) can be used to supply S to the growing crops. These often contain a more

balanced supply of essential nutrients. However, these S-containing fertilizers often have a low efficiency as S can be released slowly and immobilised in the microbial biomass (Eriksen, 2001). Lloyd (1994) showed that the average annual grass dry matter yield from a cattle slurry application was equivalent to 55% of that when the equivalent quantity of S was applied as gypsum. In less economically developed countries, S is largely delivered to crops in manure collected from cattle and pigs rather than mineral fertilisers (Erikseil and Mortensen, 1999).

Previous studies have shown large differences in S availability between crop residues. For instance, cruciferous crops, such as rape, can provide a large and rapid release of available S, while cereal residues release only minimal amounts of available S (Scherer, 2001). In addition, the release and the subsequent immobilization of S largely depends on the C-to-S ratio of the added organic material (S. J. Chapman, 1997). Previous research has shown that the incorporation of cattle manure compost with a low C-to-S ratio can provide a large amount of plant-available S, while sawdust compost with a high C-to-S ratio results in severe S deficiency, indicating the release and subsequent immobilization of S (Chowdhury *et al.*, 2000).

### **2.3.2 Uptake and allocation of sulphate by plants**

Growing plants have a high demand for S thus inadequate acquisition of S can limit plant growth (Kopriva *et al.*, 2016). The increasing recognition of the importance of S for plant growth (Zhao *et al.*, 1997; Zhao, Hawkesford and McGrath, 1999; Scherer, 2001; Aulakh, 2003; Walker and Booth, 2003; Kühn-institut, 2019a), as well as the nutritional importance of S for human and animal diets (Virtanen, 1962; Donoso-sin, 1963; Kandylis, 1984; Grimble, 2006; Nimni, Han and Cordoba, 2007), has led to an increased emphasis on research on sulphate uptake, transport and assimilation in plants (Coughlan, 1977; Smith, Rae and Hawkesford, 2000; Takahashi *et al.*, 2000; Hawkesford *et al.*, 2003; Buchner, Takahashi and

Hawkesford, 2004). The S requirement of a crop for optimal growth varies between 0.1 and 0.5% of the dry weight of plants. In general, to produce 1 tonne of grain under S-sufficient conditions, winter wheat needs 2-3 kg S (Harwood and Nicholls, 1979), but it should be noted that this is plant cultivar dependent (Vong, Nguyen and Guckert, 2007).

Sulphate uptake across the root plasma membrane is the first step involved in the synthesis of organic-S compounds in plants. This uptake by the root epidermis and cortical cells, as well as sulphate movements to other tissues and organs, requires high affinity transporters. Studies have shown that sulphate uptake is mediated by a combination of saturable high-affinity transporters and non-saturable low-affinity transporters (Epstein, 1966; Yoshimoto *et al.*, 2002). High affinity sulphate transporters are highly regulated by changes in S status of plants, encoded by a group 1 sulphate-transporter clade, while low affinity transporters play an important role in internal translocation of sulphate in plants, and all belong to a group 2 clade (Takahashi *et al.*, 1996, 2000; Gent *et al.*, 1997; Smith *et al.*, 1997). After root uptake, subcellular and long-distance transport can occur (i.e., root to shoot). Sulphate is subsequently delivered to plastids, where assimilatory reductive pathway of sulphate takes place, as well as into storage vacuoles, where the internal nutritional reservoir plays a crucial role in maintaining cytosolic and plastidic sulphate homeostasis (Leustek, 2002).

Sulphate is activated, reduced to sulphide and incorporated into carbohydrate skeletons by assimilatory sulphate reduction before it can be used in protein synthesis in plants (Brunold, 1990). The final product of assimilatory sulphate reduction in plants is cysteine. From this amino acid, all other reduced S compounds including methionine (Rennenberg *et al.*, 1990), glutathione (Strohm *et al.*, 1995), and phytochelatins (Rauser, 1995) are synthesized in a connected set of metabolic pathways. In some cases, these metabolic transformations occur in leaves with newly synthesized organic-S compounds returned to the root via the phloem. This is of vital importance in plant nutrition, as some plant species have a very low capacity for



reducing N and S in their roots, and thus depend on the translocation of nutrients for synthesis of protein.

The traditional view is that only mineral S (i.e.  $\text{SO}_4^{2-}$ ) can be utilized by plants (Haynes, 1986; Rennenberg *et al.*, 1990). However, observations that plants can take up organic N in the form of amino acids and peptides suggests that this could also be a potential route of organic-S uptake (Kielland, 1994; Smith and Steenkamp, 1992; Falkengren-Grerup *et al.*, 2000; Näsholm *et al.*, 1998; Nordin *et al.*, 2001; Owen and Jones, 2001; Raab *et al.*, 1996; Schimel and Chapin, 1996). By analogy to N, this uptake of organic S by roots may be particularly important when inorganic S becomes limiting in soil.

### **2.3.3 Sulphur metabolites in plant metabolism**

Cysteine and methionine are the major end-products of sulphate assimilation in plants, comprising up to 90% of the total S in most plants, predominantly bound in protein moieties (99% or more) (Giovanelli, Mudd and Datko, 1980; Rennenberg *et al.*, 1990). S, however, can be found in many co-enzymes and prosthetic groups and various other secondary metabolites. These secondary metabolites are important for plant fitness and their ability to cope with stress. One key example is glutathione which is a tripeptide of glycine, glutamate, cysteine and a major thiol in plants. It functions as a transient storage pool of reduced S, and is a powerful antioxidant being involved in defence against oxidative stress, detoxification of heavy metals or xenobiotics and also in biotic interactions (Mullineaux and Rausch, 2005; Rausch and Wachter, 2005). Another important secondary metabolite group are phytochelatins which consist of repetitive glutamyl-cysteine units (between 2 and more than 10) with a terminal glycine (Inouhe, 2005). They bind heavy metal cations (e.g.  $\text{Zn}^{2+}$ ) via thiol coordination making them non-toxic (Salt and Rauser, 1995; Yadav, 2010). Thioredoxins are low-MW proteins of about 12 kDa with two well-conserved cysteine residues which form a redox-active,

intermolecular disulphide bridge, constituting another important family of thiols in higher plants (Attiwill *et al.*, 1993; Vieira Dos Santos and Rey, 2006). S also plays a vital role in chlorophyll formation (Rahul *et al.*, 2018). When S is deficient, plants are often characterized by yellowing of leaves and generally show symptoms that resemble those of N starvation. In addition, alliin and glucosinolates are also two important secondary metabolites. Alliin (S-allyl-L-cysteine sulfoxide) are abundant in the *Allium* family of plants and are related with defence response reactions. Glucosinolates, the best known example of preformed defence compounds, are stored in vacuoles and their hydrolysis is catalysed by the cytosolic enzyme myrosinase, and are important for plant interactions with herbivores and pathogens (Wittstock and Halkier, 2002; Martínez-Ballesta, Moreno and Carvajal, 2013; Variyar *et al.*, 2014).

## **2.4 Sulphur analysis methods, including isotopic methods**

### **2.4.1 Extraction of sulphate and organic sulphur in soils**

Historically, one of the main reasons why so little work has been done on organic-S, relative to other plant nutrients, is the difficulty of measuring it in plants and soils. In contrast to organic-S, the development of ion chromatography (IC) (Brown and Morra, 1991) and inductively coupled plasma atomic emission spectroscopy (ICP-AES) (Munter and Grande, 1981) has spurred much research on inorganic S. IC with a conductivity detector is suitable technique for rapid determination of ions including sulphate (Raue, Brauch and Frimmel, 1991; Zhao and McGrath, 1994; Turrión, Gallardo and González, 1999; Meneses, Maniasso and Zagatto, 2005). ICP-AES has proved to be an accurate and rapid method for measurement of total dissolved S (TDS) for soil and plant materials (Novozamsky *et al.*, 1986; Perrott *et al.*, 1991), allowing a simultaneous measurement of  $\text{SO}_4^{2-}$ -S and dissolved organic S by difference with TDS. However, ICP-AES methods fail to identify the exact species of S, therefore near edge X-ray absorption fine structure spectroscopy (NEXAFS), also known as X-ray absorption

near-edge spectroscopy (XANES), the practice of determining oxidation states involves using a set of conventional rules, has been used to identify and quantify multiple organic S composition in coal (Spiro *et al.*, 1984; Kasrai *et al.*, 1990, 1996), minerals (Marusak and Tongson, 1979; Sugiura, 1981; Kubono *et al.*, 2017), biological samples (Frank *et al.*, 1987, 1994; Shadle *et al.*, 1993; Pickering *et al.*, 1998), and humic substances isolated from marine sediments (Vairavamurthy *et al.*, 1994, 1997; Xia *et al.*, 1998), and soils (Jokic *et al.*, 2003; Solomon, Lehmann and Martínez, 2003; Solomon *et al.*, 2005; Zhao *et al.*, 2006; Prietzel *et al.*, 2013).

#### **2.4.2 Determination of volatile sulphur compounds**

Research on S cycling in soil was hindered by the lack of sensitive methods for quantifying the flux of volatile organosulfur compounds (VOS) produced by soil microorganisms. However, the advancement of GC-MS and VOS capture technologies now allows us to identify and trace VOS compounds. VOS compounds play important roles in the S cycle, owing to their transfer from solution to the gas phase and *vice versa*. In particular, low MW VOS compounds, such as dimethylsulfide (DMS), dimethyldisulfide (DMDS), methanethiol (MT), carbon disulphide (CS<sub>2</sub>) and carbonyl sulfide (COS) are recognised as important source of S to the atmosphere (Kiene, 1996). The release of VOS compounds primarily occurs from soils treated with plant materials (Bending and Lincoln, 1999) or from the breakdown of S-containing amino acids such as methionine and cysteine (W. L. Banwart and Bremner, 1976). Waterlogging appears to be one of the most important factors affecting VOS emissions from soils, with increased VOS emissions following a decrease in oxygen availability (Chapman *et al.*, 1996). Temperature (Staubes *et al.*, 1989) as well as the soil organic S content (Bending and Lincoln, 1999) has also been shown to be important in regulating VOS emissions from soils.

In a water saturated soil, aerobic microorganisms quickly consume all the available oxygen, and anaerobic microorganisms begin to use  $\text{Fe}^{3+}$ ,  $\text{Mn}^{4+}$  and  $\text{SO}_4^{2-}$  etc. as terminal electron acceptors to obtain energy for growth (Lovley, Holmes and Nevin, 1991). In wetland soils and freshwater environments with appreciable  $\text{SO}_4^{2-}$ , microbial reduction of sulphate to  $\text{S}^-$  or  $\text{H}_2\text{S}$  occurs (Pester *et al.*, 2012). Several species of bacteria, such as sulphate reducers of the genera *Desulphovibrio*, *Desulfobacter*, *Desulfococcus*, are predominantly responsible for the volatilization of reduced S gases in waterlogged soils (Brychkova *et al.*, 2007). These bacteria use  $\text{SO}_4^{2-}$  as an oxidizing agent and produce large quantities of  $\text{H}_2\text{S}$  as one of their main metabolic products. Sulphate can also be reduced to  $\text{H}_2\text{S}$  under well-aerated conditions by the aerobic *Bacillus megaterium* in partially sterilized soil (Brychkova *et al.*, 2007).

#### **2.4.3 Determination of microbial biomass sulphur**

The soil microbial biomass acts as the driving force behind mineralization–immobilization, and oxidation–reduction transformations and is therefore central to S cycling (Banerjee and Chapman, 1996). Methods have been developed for estimating microbial biomass-S (Strick and Nakas, 1984; Wu *et al.*, 1994; Banerjee and Chapman, 1996), which accounts for < 3% of organic S in soil (Stevenson and Cole, 1999). The most commonly used method to measure microbial biomass-S is the  $\text{CHCl}_3$  fumigation-extraction method, in a way analogous to the measurement of microbial biomass-C.

This method involves lysing the microbial cells with  $\text{CHCl}_3$ , evacuating the  $\text{CHCl}_3$ , and then directly extracting the S from both fumigated and non-fumigated soils with 0.01 M  $\text{CaCl}_2$ . Microbial biomass-S is calculated as  $F/K_s$ , where  $F$  is the total amount of S released from fumigated soil minus that released from non-fumigated soil, and  $K_s=0.35$  represents the efficiency of extraction of microbial biomass-S (Voroney, R.P.; Brookes, P.C.; Beyaert, 2008).

#### 2.4.4 Determination of transformation rates of sulphur in soils

The rates at which nutrients are released to, and removed from, the mineral nutrient pool is an important determinant of nutrients supply to plants. These transformation rates therefore need to be considered when developing nutrient management strategies for sustainable production. Typically, net S mineralisation has traditionally been studied using a closed incubation system which measure net changes in sulphate concentration with time. However, such determinations have contributed very little to the understanding of processes involved in S transformations within the soil system. More recently, by measuring sulphatase activity (A. N. Ganeshamurthy and Nielsen, 1990), gross S mineralization rates can be estimated, yet this only accounts for some of the enzymes responsible for S mineralization. Alternatively, periodic leaching of soils allows the determination of S mineralization rates (D. G. Maynard, Stewart and Bettany, 1983; Ghani, McLaren and Swift, 1991, 1992; Valeur and Nilsson, 1993; Zhou *et al.*, 1999; J., K. and R., 2003), but this method alters soil conditions leading to an overestimation of gross mineralization.

The isotopic dilution technique is also used for determining gross nutrient transformation rates in soil. Isotopic dilution was first described as a method for analysing the lead content of rocks (Lockwood, 1954). Yet it was not until eight years later that this technique was shown to have a great analytical utility and applied for the analysis of complex mixtures of organic compounds (Rittenberg and Foster, 1940). These authors recognized the potential of this method for determining the relative D or L-isomeric content of isolated amino acids. This technique involves labelling a soil mineral nutrient pool, e.g.  $\text{NH}_4^+$ ,  $\text{NO}_3^-$ , (NISHIO, 1991; Delaune *et al.*, 1998),  $\text{PO}_4^{3-}$  (Walbridge and Vitousek, 1987; Wanek *et al.*, 2019), or  $\text{SO}_4^{2-}$  (Di, Cameron and McLaren, 2000; Nziguheba, Smolders and Merckx, 2005), and monitoring the changes of the size of the labelled nutrient pool and the abundance of the rarer isotopes (atom%, if stable isotope is used) or specific activity (if radioisotope is added).

The element S has four naturally occurring stable isotopes.  $^{32}\text{S}$ ,  $^{33}\text{S}$ ,  $^{34}\text{S}$  and  $^{35}\text{S}$ , with approximate abundances of 95.02%, 0.75%, 4.21% and 0.02%, respectively (Day and Moynier, 2014). The only radioactive isotope of S which is suitable for biochemical investigations is  $^{35}\text{S}$ . Therefore,  $^{35}\text{S}$  has been used to trace S processes in soil, including microbial mineralization, immobilisation, reduction, sorption, leaching and plant uptake (Di, Cameron and McLaren, 2000). By tracing the flow of added  $^{35}\text{S}$  from one S fraction to the other, the dynamics of S mineralization and immobilization within soil can be examined. When fertilizer  $^{35}\text{S}\text{-Na}_2\text{SO}_4$  was added to soil, studies showed that it is readily converted to organic forms of both hydriodic acid reducible and carbon-bonded S (Freney, Melville and Williams, 1971; Schindler *et al.*, 1986). Double labelled fertilizer such as  $(^{15}\text{NH}_4)_2^{35}\text{SO}_4$  as well as triple-labelled ( $^{13}\text{C}$ ,  $^{15}\text{N}$ ,  $^{34}\text{S}$ ) manure (Andriuzzi and Schmidt, 2014) have been synthesized, yet their experimental application have not been reported.

#### **2.4.5 Quantification of S uptake by plant roots**

Current techniques to quantify the S uptake by the roots, include depletion methods (in which uptake rates are calculated from changes in solution concentrations over time), and tracer methods (in which the uptake of tracer into root or shoot tissue is measured) (Lucash *et al.*, 2007). These techniques provide valuable information about the uptake rates, but they are also subject to several methodological problems. First, uptake rates can be determined by analysing tracer accumulation in the roots using the tracer method (Larsson *et al.*, 1991). The tracer method is cost-effective and sensitive and it also offers the advantage of tracking the radioisotope in the shoots by autoradiography, thereby enabling determination of translocation rates of tracer through the plant (Rubio, Sorgona and Lynch, 2004). For the tracer method, both excised and intact roots are used. Of these, the use of excised roots has been used extensively since the 1960s to measure inorganic (Sheat, Fletcher and Street, 1959; Huang *et al.*, 1992) and

organic (McFarlane and Wickliff, 1985; Su and Zhu, 2007) nutrient uptake by roots. However, some scientists have argued that damage to root tissues caused by excision prior to experiments, may often artificially increase the nutrient loss from roots and thereby reduce net uptake, providing less realistic estimates of root uptake than intact root systems (Bloom and Caldwell, 1988).

Second, the depletion method of measuring the concentration of nutrients in solution before and after the experiment, can be used to determine the amount of nutrient taken up by the roots. The depletion method offers a possible improvement over the excised root method, as that the roots are still attached to the plant and can continue nutrient transportation throughout the experiment. Another advantage of this method is that isotopes are not required, so the approach is more affordable, and fewer handling restrictions are involved. The main limitation of the depletion method is that the difference must be detectable and, thus, requires either low starting amino acid concentrations or long uptake periods.

## **2.5 Summary**

The five key findings of this literature review are:

1. Most S held in soils is present in organic matter and the microbial biomass.
2. Many groups of soil microorganisms are involved in the conversion of organic S to  $\text{SO}_4^{2-}$ .
3. Our understanding of whether plants can take up organic-S from soils is still very limited.
4. The rates of conversion of low MW organic-S to  $\text{SO}_4^{2-}$  involves many enzymatic pathways but remains poorly understood.
5. While we know what the major pools of S are in the agricultural S cycle, we still know little about rates of flux between these pools and their co-dependence on C and N cycling.

Hence, the main aims of this thesis are to i) study the release and subsequent availability of C, N and S from cysteine and methionine following addition into soil and the factors that control this mineralization process (e.g., nutrient addition, substrate concentrations and soil properties); and ii) compare the uptake and competition of organic and inorganic S nutrients by plant and soil microbes in a rhizosphere context.

## 2.6 References

- Abdulina, D. *et al.* (2020) ‘ATP sulfurylase activity of sulfate-reducing bacteria from various ecotopes’, *3 Biotech.* Springer International Publishing, 10(2), pp. 1–9. doi: 10.1007/s13205-019-2041-9.
- Ahmad, A. and Abdin, M. Z. (2000) ‘Photosynthesis and its related physiological variables in the leaves of Brassica genotypes as influenced by sulphur fertilization’, *Physiologia Plantarum*, 110(1), pp. 144–149. doi: 10.1034/j.1399-3054.2000.110119.x.
- Ahmad, G. *et al.* (2007) ‘Influence of nitrogen and sulfur fertilization on quality of canola (*Brassica napus* L.) under rainfed conditions’, *Journal of Zhejiang University Science B.* Springer, 8(10), pp. 731–737.
- Andriuzzi, W. S. and Schmidt, O. (2014) ‘Production of rabbit dung triple-labelled with C, N and S stable isotopes’, *Pedobiologia*, 57(3), pp. 155–160. doi: 10.1016/j.pedobi.2014.02.001.
- Attiwill, P. M. *et al.* (1993) ‘Published by : Wiley on behalf of the New Phytologist Trust  
Linked references are available on JSTOR for this article : Tansley Review No . 50  
Nutrient cycling in forests’, 124(50), pp. 561–582.
- Aulakh, M. S. (2003) ‘Crop Responses to Sulphur Nutrition’, in *Sulphur in Plants*. Springer, pp. 341–358. doi: 10.1007/978-94-017-0289-8\_19.



- Bais, H. P. *et al.* (2006) 'THE ROLE OF ROOT EXUDATES IN RHIZOSPHERE INTERACTIONS WITH PLANTS AND OTHER ORGANISMS', *Annual Review of Plant Biology*, 57(1), pp. 233–266. doi: 10.1146/annurev.arplant.57.032905.105159.
- Banerjee, M. R. and Chapman, S. J. (1996) 'The significance of microbial biomass sulphur in soil', *Biology and Fertility of Soils*, 22(1), pp. 116–125. doi: 10.1007/s003740050085.
- Banwart, W. L. and Bremner, J. M. (1975) 'Formation of volatile sulfur compounds by microbial decomposition of sulfur-containing amino acids in soils', *Soil Biology and Biochemistry*, pp. 359–364. doi: 10.1016/0038-0717(75)90050-4.
- Banwart, W. L. and Bremner, J. M. (1976) 'Evolution of volatile sulfur compounds from soils treated with sulfur-containing organic materials', *Soil Biology and Biochemistry*, 8(5), pp. 439–443. doi: 10.1016/0038-0717(76)90047-X.
- Barber, S. A. (1995) *Soil nutrient bioavailability: a mechanistic approach*. John Wiley & Sons.
- Barrow, N. J. (1978) 'Inorganic reactions of phosphorus, sulphur and molybdenum in soil', *Mineral nutrition of legumes in tropical and subtropical soils*. CSIRO, Aust.
- Barrows, H. L. and Kilmer, V. J. (1963) 'Plant Nutrient Losses from Soils by Water Erosion', *Advances in Agronomy*. Elsevier, 15(C), pp. 303–316. doi: 10.1016/S0065-2113(08)60401-0.
- Bates, T. S. *et al.* (1992) 'Sulfur emissions to the atmosphere from natural sources', *Journal of Atmospheric Chemistry*, 14(1–4), pp. 315–337. doi: 10.1007/BF00115242.
- Battarbee, R. W. *et al.* (2014) 'Recovery of acidified surface waters from acidification in the United Kingdom after twenty years of chemical and biological monitoring (1988–2008)', *Ecological Indicators*. Elsevier Ltd, pp. 267–273. doi: 10.1016/j.ecolind.2013.10.011.

- Belnap, J., Hawkes, C. V. and Firestone, M. K. (2003) 'Boundaries in miniature: Two examples from soil', *BioScience*, 53(8), pp. 739–749. doi: 10.1641/0006-3568(2003)053[0739:BIMTEF]2.0.CO;2.
- Bending, G. D. and Lincoln, S. D. (1999) 'Characterisation of volatile sulphur-containing compounds produced during decomposition of Brassica juncea tissues in soil', *Soil Biology and Biochemistry*, 31(5), pp. 695–703. doi: 10.1016/S0038-0717(98)00163-1.
- Bertin, C., Yang, X. and Weston, L. A. (2003) 'The role of root exudates and allelochemicals in the rhizosphere', *Plant and Soil*, 256(1), pp. 67–83. doi: 10.1023/A:1026290508166.
- Bloom, A. J. and Caldwell, R. M. (1988) 'Root excision decreases nutrient absorption and gas fluxes', *Plant Physiology. Am Soc Plant Biol*, 87(4), pp. 794–796.
- Blumenberg, M. *et al.* (2006) 'Biosynthesis of hopanoids by sulfate-reducing bacteria (genus *Desulfovibrio*)', *Environmental Microbiology*, 8(7), pp. 1220–1227. doi: 10.1111/j.1462-2920.2006.01014.x.
- Bolan, N. S. *et al.* (1988) 'Effect of liming and phosphate additions on sulphate leaching in soils', *Journal of soil science. Wiley Online Library*, 39(4), pp. 493–504.
- Bolan, N. S. *et al.* (2011) *Dissolved Organic Matter. Biogeochemistry, Dynamics, and Environmental Significance in Soils*. 1st edn, *Advances in Agronomy*. 1st edn. Elsevier Inc. doi: 10.1016/B978-0-12-385531-2.00001-3.
- Boswell, C. C. and Friesen, D. K. (1993) 'Elemental sulfur fertilizers and their use on crops and pastures', *Fertilizer Research*, 35(1–2), pp. 127–149. doi: 10.1007/BF00750226.
- Brosnan, J. T. and Brosnan, M. E. (2006) 'The sulfur-containing amino acids: an overview', *The Journal of nutrition*. Oxford University Press, 136(6), pp. 1636S-1640S.
- Brown, K. A. (1982) 'Sulphur in the environment: A review', *Environmental Pollution. Series*

- B, Chemical and Physical*, 3(1), pp. 47–80. doi: 10.1016/0143-148X(82)90042-8.
- Brown, P. D. and Morra, M. J. (1991) ‘Ion Chromatographic Determination of SCN in Soils’, *Journal of Agricultural and Food Chemistry*. Wiley Online Library, 39(7), pp. 1226–1228. doi: 10.1021/jf00007a006.
- Brunold, C. (1990) ‘Reduction of sulphate to sulphide. Sulphur Nutrition and Sulphur Assimilation in Higher Plants’. SPB Academic Publishing, The Hague.
- Brychkova, G. *et al.* (2007) ‘Sulfite oxidase protects plants against sulfur dioxide toxicity’, *Plant Journal*, 50(4), pp. 696–709. doi: 10.1111/j.1365-313X.2007.03080.x.
- Buchner, P., Takahashi, H. and Hawkesford, M. J. (2004) ‘Plant sulphate transporters: Co-ordination of uptake, intracellular and long-distance transport’, *Journal of Experimental Botany*, 55(404), pp. 1765–1773. doi: 10.1093/jxb/erh206.
- Byers, M. and Bolton, J. (1979) ‘Effects of nitrogen and sulphur fertilisers on the yield, N and S content, and amino acid composition of the grain of spring wheat’, *Journal of the Science of Food and Agriculture*. Wiley Online Library, 30(3), pp. 251–263.
- Campbell, G. W. and Smith, R. I. (1996) ‘Spatial and temporal trends in atmospheric sulphur deposition to agricultural surfaces in the United Kingdom’, *Proceedings-Fertiliser Society (United Kingdom)*. Fertiliser Society.
- Cao, X. *et al.* (2018) ‘Tracing the origin and geochemical processes of dissolved sulphate in a karst-dominated wetland catchment using stable isotope indicators’, *Journal of Hydrology*. Elsevier, 562(May), pp. 210–222. doi: 10.1016/j.jhydrol.2018.04.072.
- Cardon, Z. G. and Whitbeck, J. L. (2011) *The rhizosphere: an ecological perspective*. Elsevier.
- Carvalhais, L. C. *et al.* (2011) ‘Root exudation of sugars, amino acids, and organic acids by maize as affected by nitrogen, phosphorus, potassium, and iron deficiency’, *Journal of*

- Plant Nutrition and Soil Science*, 174(1), pp. 3–11. doi: 10.1002/jpln.201000085.
- Castro, H. F., Williams, N. H. and Ogram, A. (2000) ‘Phylogeny of sulfate-reducing bacteria 1’, 31.
- Ceccotti, S. P., Morris, R. J. and Messick, D. L. (1998) ‘A global overview of the sulphur situation: industry’s background, market trends, and commercial aspects of sulphur fertilizers’, in *Sulphur in agroecosystems*. Springer, pp. 175–202.
- Chalk, P. M., Inácio, C. T. and Chen, D. (2017) ‘Tracing S dynamics in agro-ecosystems using <sup>34</sup>S’, *Soil Biology and Biochemistry*, 114, pp. 295–308. doi: 10.1016/j.soilbio.2017.07.001.
- Chao, T. T., Harward, M. E. and Fang, S. C. (1962) ‘Movement of S<sup>35</sup> tagged sulfate through soil columns’, *Soil Science Society of America Journal*. Wiley Online Library, 26(1), pp. 27–32.
- Chapman, S. J. (1987) ‘Microbial sulphur in some scottish soils’, *Soil Biology and Biochemistry*, 19(3), pp. 301–305. doi: 10.1016/0038-0717(87)90013-7.
- Chapman, S. J. *et al.* (1996) ‘Influence of temperature and oxygen availability on the flux of volatile sulphur compounds from wetlands: A comparison of peat and paddy soils’, *Soil Science and Plant Nutrition*, 42(2), pp. 279–288. doi: 10.1080/00380768.1996.10415098.
- Chapman, S. J. (1997) ‘Barley straw decomposition and S immobilization’, *Soil Biology and Biochemistry*, 29(2), pp. 109–114. doi: 10.1016/S0038-0717(97)00001-1.
- Charlson, R. J., Anderson, T. L. and McDuff, R. E. (1992) ‘The sulfur cycle’, *Global biogeochemical cycles*. Academic Press, London, pp. 285–300.
- Chen, M. *et al.* (2016) ‘Increasing sulfate concentrations result in higher sulfide production

- and phosphorous mobilization in a shallow eutrophic freshwater lake’, *Water Research*. Elsevier Ltd, 96, pp. 94–104. doi: 10.1016/j.watres.2016.03.030.
- Chen, Y. *et al.* (2018) ‘Formation and change of chloroplast-located plant metabolites in response to light conditions’, *International Journal of Molecular Sciences*, 19(3). doi: 10.3390/ijms19030654.
- Chowdhury, M. A. H. *et al.* (2000) ‘Microbial biomass, S mineralization and S uptake by African millet from soil amended with various composts’, *Soil Biology and Biochemistry*, 32(6), pp. 845–852. doi: 10.1016/S0038-0717(99)00214-X.
- Christou, M. *et al.* (2005) ‘Dissolved organic nitrogen in contrasting agricultural ecosystems’, *Soil Biology and Biochemistry*, 37(8), pp. 1560–1563. doi: 10.1016/j.soilbio.2005.01.025.
- Christou, M., Avramides, E. J. and Jones, D. L. (2006) ‘Dissolved organic nitrogen dynamics in a Mediterranean vineyard soil’, *Soil Biology and Biochemistry*, 38(8), pp. 2265–2277. doi: 10.1016/j.soilbio.2006.01.025.
- Churka Blum, S. *et al.* (2013) ‘Sulfur forms in organic substrates affecting S mineralization in soil’, *Geoderma*, 200–201, pp. 156–164. doi: 10.1016/j.geoderma.2013.02.003.
- Churka Blum, S., Caires, E. F. and Alleoni, L. R. (2013) ‘Lime and phosphogypsum application and sulfate retention in subtropical soils under no-till system’, *Journal of soil science and plant nutrition*. Chilean Society of Soil Science, 13(2), pp. 279–300.
- Colovic, M. B. *et al.* (2018) ‘Sulphur-containing amino acids: protective role against free radicals and heavy metals’, *Current medicinal chemistry*. Bentham Science Publishers, 25(3), pp. 324–335.
- Comerford, N. B. (2005) ‘Soil factors affecting nutrient bioavailability’, in *Nutrient acquisition*

- by plants. Springer, pp. 1–14.
- Cooper, P. J. M. (1972) 'Aryl sulphatase activity in northern Nigerian soils', *Soil Biology and Biochemistry*. Elsevier, 4(3), pp. 333–337.
- Coughlan, S. (1977) 'Sulphate uptake in *Fucus serratus*', *Journal of Experimental Botany*, 28(5), pp. 1207–1215. doi: 10.1093/jxb/28.5.1207.
- Couture, R. M. *et al.* (2016) 'Non-steady state diagenesis of organic and inorganic sulfur in lake sediments', *Geochimica et Cosmochimica Acta*. Elsevier Ltd, 194, pp. 15–33. doi: 10.1016/j.gca.2016.08.029.
- Coyne, M. S., Mikkelsen, R. and Mineralization, S. (2015) 'Soil microorganisms contribute to plant nutrition and root health', *Better Crops*, 99(1), pp. 18–20.
- Cronan, J. E. (2014) 'Biotin and lipoic acid: synthesis, attachment and regulation', *EcoSal Plus*. NIH Public Access, 6(1).
- Cullis, C. F. and Hirschler, M. M. (1980) 'Atmospheric sulphur: Natural and man-made sources', *Atmospheric Environment (1967)*, 14(11), pp. 1263–1278. doi: 10.1016/0004-6981(80)90228-0.
- Curl, E. A. and Truelove, B. (2012) *The rhizosphere*. Springer Science & Business Media.
- Curtin, D. and Syers, J. K. (1990) 'Extractability and adsorption of sulphate in soils', *Journal of soil science*. Wiley Online Library, 41(2), pp. 305–312.
- Davidian, J. C. and Kopriva, S. (2010) 'Regulation of sulfate uptake and assimilation - The same or not the same?', *Molecular Plant*. The Authors. All rights reserved., 3(2), pp. 314–325. doi: 10.1093/mp/ssq001.
- Day, J. M. D. and Moynier, F. (2014) 'Evaporative fractionation of volatile stable isotopes and

- their bearing on the origin of the Moon’, *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 372(2024). doi: 10.1098/rsta.2013.0259.
- Delaune, R. D. *et al.* (1998) ‘Nitrification and denitrification estimates in a Louisiana swamp forest soil as assessed by <sup>15</sup>N isotope dilution and direct gaseous measurements’, *Water, Air, and Soil Pollution*, 106(1–2), pp. 149–161. doi: 10.1023/A:1004953626415.
- Di, H. J., Cameron, K. C. and McLaren, R. G. (2000) ‘Isotopic dilution methods to determine the gross transformation rates of nitrogen, phosphorus, and sulfur in soil: a review of the theory, methodologies, and limitations’, *Soil Research*, 38(1), pp. 213–230. doi: 10.1071/SR99005.
- Dick, R. P. and Deng, S. (1991) ‘Multivariate factor analysis of sulfur oxidation and rhodanese activity in soils’, *Biogeochemistry*, 12(2), pp. 87–101. doi: 10.1007/BF00001808.
- Donoso-sin, M. G. (1963) ‘RELATIONSHIP BETWEEN THE SULPHUR / NITROGEN’, 14, pp. 345–349.
- Edwards, P. J. (1998) *Sulfur cycling, retention, and mobility in soils: a review*. US Department of Agriculture, Forest Service, Northeastern Research Station.
- Ercoli, L. *et al.* (2012) ‘Management of sulphur fertiliser to improve durum wheat production and minimise S leaching’, *European Journal of Agronomy*. Elsevier B.V., 38(1), pp. 74–82. doi: 10.1016/j.eja.2011.12.004.
- Erikseil, J. and Mortensen, J. V. (1999) ‘Soil sulphur status following long-term annual application of animal manure and mineral fertilizers’, *Biology and Fertility of Soils*, 28(4), pp. 416–421. doi: 10.1007/s003740050514.

- Eriksen, J. (2001) 'Organic manures as sources of fertiliser sulphur', *International Fertiliser Society Proceedings*, 505, pp. 1–19.
- Eriksson, E. (1988) 'Retention and release of sulphate in soils', *Critical loads for sulphur and nitrogen*. Nordic Council of Ministers Copenhagen, pp. 151–184.
- Esaki, N. *et al.* (1979) 'Catalytic Action of L-Methionine  $\gamma$ -Lyase on Selenomethionine and Selenols†', *Biochemistry*, 18(3), pp. 407–410. doi: 10.1021/bi00570a003.
- Fauque, G. *et al.* (1988) 'The three classes of hydrogenases from sulfate-reducing bacteria of the genus *Desulfovibrio*', *FEMS Microbiology Letters*, 54(4), pp. 299–344. doi: 10.1016/0378-1097(88)90248-0.
- Filipek-Mazur, B. *et al.* (2019) 'Effect of sulfur-containing fertilizers on the quantity and quality of spring oilseed rape and winter wheat yield', *Journal of Elementology*. -, 24(4).
- Fisher, D. W. *et al.* (1968) 'Atmospheric contributions to water quality of streams in the Hubbard Brook Experimental Forest, New Hampshire', *Water Resources Research*. Wiley Online Library, 4(5), pp. 1115–1126.
- Fitzgerald, J. W. (1976) 'Sulfate ester formation and hydrolysis: a potentially important yet often ignored aspect of the sulfur cycle of aerobic soils', *Bacteriological Reviews*, pp. 698–721. doi: 10.1128/mmbr.40.3.698-721.1976.
- Fitzgerald, J. W. and Watwood, M. E. (1988) 'Amino-acid metabolism in forest soil-Isolation and turnover of organic matter covalently labelled with  $^{35}\text{S}$ -methionine', *Soil Biology and Biochemistry*, 20(6), pp. 833–838. doi: 10.1016/0038-0717(88)90089-2.
- Ford, C. R. *et al.* (2007) 'Soil DIC uptake and fixation in *Pinus taeda* seedlings and its C contribution to plant tissues and ectomycorrhizal fungi.', *Tree physiology*, 27(3), pp.



375–383. doi: 10.1093/treephys/27.3.375.

Fowler, D. (1980) ‘Wet and dry deposition of sulphur and nitrogen compounds from the atmosphere’, in *Effects of acid precipitation on terrestrial ecosystems*. Springer, pp. 9–27.

Frank, P. *et al.* (1987) ‘A Large Reservoir of Sulfate and Sulfonate Resides within Plasma Cells from *Ascidia ceratodes*, Revealed by X-ray Absorption Near-Edge Structure Spectroscopy’, *Biochemistry*, 26(16), pp. 4975–4979. doi: 10.1021/bi00390a014.

Frank, P. *et al.* (1994) ‘Interaction of Vanadium and Sulfate in Blood Cells from the Tunicate *Ascidia ceratodes*: Observations Using X-ray Absorption Edge Structure and EPR Spectroscopies’, *Inorganic Chemistry*, 33(17), pp. 3794–3803. doi: 10.1021/ic00095a024.

Freney, J. (1960) ‘The Oxidation of Cysteine to Sulphate in Soil’, *Australian Journal of Biological Sciences*, 13(3), p. 387. doi: 10.1071/bi9600387.

Freney, J. R. (1967) ‘Sulfur containing organics’. New York, Marcel Dekker.

Freney, J. R., Melville, G. E. and Williams, C. H. (1970) ‘The determination of carbon bonded sulfur in soil’, *Soil Science. LWW*, 109(5), pp. 310–318. doi: 10.1097/00010694-197005000-00008.

Freney, J. R., Melville, G. E. and Williams, C. H. (1971) ‘Organic sulphur fractions labelled by addition of <sup>35</sup>S-sulphate to soil’, *Soil Biology and Biochemistry*, 3(2), pp. 133–141. doi: 10.1016/0038-0717(71)90006-X.

Freney, J. R., Melville, G. E. and Williams, C. H. (1975) ‘Soil organic matter fractions as sources of plant-available sulphur’, *Soil Biology and Biochemistry*, 7(3), pp. 217–221. doi: 10.1016/0038-0717(75)90041-3.

- Freney, J. R. and Spencer, K. (1960) 'Soil sulphate changes in the presence and absence of growing plants', *Australian Journal of Agricultural Research*, 11(3), pp. 339–345. Available at: <https://doi.org/10.1071/AR9600339>.
- Gahan, J. and Schmalenberger, A. (2014) 'The role of bacteria and mycorrhiza in plant sulfur supply', *Frontiers in Plant Science*. Frontiers, 5(DEC), p. 723. doi: 10.3389/fpls.2014.00723.
- Galloway, J. N. and Whelpdale, D. M. (1980) 'An atmospheric sulfur budget for eastern North America', *Atmospheric Environment* (1967), 14(4), pp. 409–417. doi: 10.1016/0004-6981(80)90205-X.
- Ganeshamurthy, A N and Nielsen, N. E. (1990) 'Arylsulfatase and the biochemical mineralization of soil organic sulphur.', *Soil biology & biochemistry*, 22(8), pp. 1163–1165.
- Ganeshamurthy, A. N. and Nielsen, N. E. (1990) 'Arylsulphatase and the biochemical mineralization of soil organic sulphur', *Soil Biology and Biochemistry*, 22(8), pp. 1163–1165. doi: 10.1016/0038-0717(90)90045-2.
- Garland, J. A. (1978) 'Dry and wet removal of sulphur from the atmosphere', in *Sulfur in the Atmosphere*. Elsevier, pp. 349–362.
- Garsed, S. G. and Read, D. J. (1977) 'The uptake and metabolism of  $^{35}\text{SO}_2$  in plants of differing sensitivity to sulphur dioxide', *Environmental Pollution* (1970). Elsevier, 13(3), pp. 173–186.
- Gent, U. *et al.* (1997) 'Regulation of sulfur assimilation in higher plants : A sulfate transporter induced in sulfate-starved roots plays a central role in *Arabidopsis thaliana*', 94(September), pp. 11102–11107.

- Germida, J. J. and Janzen, H. H. (1993) 'Factors affecting the oxidation of elemental sulfur in soils', *Fertilizer Research*, 35(1–2), pp. 101–114. doi: 10.1007/BF00750224.
- Ghani, A. (1989) 'Role of organic sulphur in supplying sulphate for plant growth'. Lincoln College, University of Canterbury.
- Ghani, A., McLaren, R. G. and Swift, R. S. (1991) 'Sulphur mineralisation in some New Zealand soils', *Biology and Fertility of Soils*, 11(1), pp. 68–74. doi: 10.1007/BF00335838.
- Ghani, A., McLaren, R. G. and Swift, R. S. (1992) 'Sulphur mineralisation and transformations in soils as influenced by additions of carbon, nitrogen and sulphur', *Soil Biology and Biochemistry*, 24(4), pp. 331–341. doi: 10.1016/0038-0717(92)90193-2.
- Ghosh, G. K. and Dash, N. R. (2012) 'Sulphate sorption–desorption characteristics of lateritic soils of west Bengal, India', *International Journal of Plant, Animal and Environmental Sciences*, 2(1), pp. 167–176.
- Giovanelli, J. S., Mudd, S. H. and Datko, A. H. (1980) 'Sulfur Amino Acids in Plants', *Amino Acids and Derivatives*. New York: Academic Press. doi: org/10.1016/B978-0-12-675405-6.50018-8.
- Granat, L. (2013) 'Oikos Editorial Office The Global Sulphur Cycle Author ( s ): L . Granat , H . Rodhe and R . O . Hallberg Published by : Oikos Editorial Office', 7(22), pp. 89–134.
- Granat, L., Rodhe, H. and Hallberg, R. O. (1976) 'The Global Sulphur Cycle', in *Nitrogen, Phosphorus and Sulphur - Global Cycles. SCOPE Report 7*. JSTOR, pp. 89–134. Available at: <http://www.jstor.org/stable/20112523>.
- Greenfield, L. M. *et al.* (2020) 'Do plants use root-derived proteases to promote the uptake of

- soil organic nitrogen?', *Plant and Soil*. Plant and Soil, 456(1–2), pp. 355–367. doi: 10.1007/s11104-020-04719-6.
- Griffith, C. M., Woodrow, J. E. and Seiber, J. N. (2015) 'Environmental behavior and analysis of agricultural sulfur', *Pest Management Science*, pp. 1486–1496. doi: 10.1002/ps.4067.
- Grimble, R. F. (2006) 'The Effects of Sulfur Amino Acid Intake on Immune Function in Humans', *The Journal of Nutrition*, 136(6), pp. 1660S-1665S. doi: 10.1093/jn/136.6.1660s.
- Haaijer, S. C. M. *et al.* (2006) 'Evidence for the involvement of betaproteobacterial Thiobacilli in the nitrate-dependent oxidation of iron sulfide minerals', *FEMS Microbiology Ecology*, 58(3), pp. 439–448. doi: 10.1111/j.1574-6941.2006.00178.x.
- Haaijer, S. C. M. *et al.* (2008) 'Bacteria associated with iron seeps in a sulfur-rich, neutral pH, freshwater ecosystem', *ISME Journal*, 2(12), pp. 1231–1242. doi: 10.1038/ismej.2008.75.
- Hamilton, E. I. (1985) 'The global biogeochemical sulphur cycle', *Science of The Total Environment*, 41(2), pp. 195–196. doi: 10.1016/0048-9697(85)90190-1.
- Haque, I. and Walmsley, D. (1973) 'Adsorption and desorption of sulphate in some soils of the West Indies', *Geoderma*. Elsevier, 9(4), pp. 269–278.
- Harwood, J. L. and Nicholls, R. G. (1979) 'The plant sulpholipid-a major component of the sulphur cycle', *Biochem Soc Trans*, 7(2), pp. 440–447. doi: 10.1042/bst0070440.
- Hati, K. M. *et al.* (2007) 'Changes in soil physical properties and organic carbon status at the topsoil horizon of a vertisol of central India after 28 years of continuous cropping, fertilization and manuring', *Agriculture, Ecosystems and Environment*, 119(1–2), pp.

127–134. doi: 10.1016/j.agee.2006.06.017.

Hatzfeld, Y. *et al.* (2000) 'Functional characterization of a gene encoding a fourth ATP sulfurylase isoform from *Arabidopsis thaliana*', *Gene*. Elsevier, 248(1–2), pp. 51–58.

Hawkesford, M. J. *et al.* (2003) 'Sulphate Uptake and Transport', in *Sulphur in Plants*. Springer, pp. 71–86. doi: 10.1007/978-94-017-0289-8\_4.

Haynes, R. J. (1986) 'Uptake and assimilation of mineral nitrogen by plants', *Mineral nitrogen in the plant-soil system*. Academic Press San Diego, pp. 303–378.

Heim, A. *et al.* (2001) 'Root exudation, organic acids, and element distribution in roots of Norway spruce seedlings treated with aluminum in hydroponics', *Journal of Plant Nutrition and Soil Science*. Wiley Online Library, 164(5), pp. 519–526.

Herrmann, J. *et al.* (2014) 'Structure and mechanism of soybean ATP sulfurylase and the committed step in plant sulfur assimilation', *Journal of Biological Chemistry*. Elsevier, 289(15), pp. 10919–10929.

Hille, R. (1996) 'The mononuclear molybdenum enzymes', *Chemical Reviews*, 96(7), pp. 2757–2816. doi: 10.1021/cr950061t.

Hinckley, E. L. S. *et al.* (2020) 'A shift in sulfur-cycle manipulation from atmospheric emissions to agricultural additions', *Nature Geoscience*. Springer US, 13(9), pp. 597–604. doi: 10.1038/s41561-020-0620-3.

Holmer, M. and Storkholm, P. (2001) 'Sulphate reduction and sulphur cycling in lake sediments: A review', *Freshwater Biology*, 46(4), pp. 431–451. doi: 10.1046/j.1365-2427.2001.00687.x.

Hu, Z. *et al.* (2002) 'Effect of crop growth on the distribution and mineralization of soil sulfur fractions in the rhizosphere', *Journal of Plant Nutrition and Soil Science*, 165(3), pp.

249–254. doi: 10.1002/1522-2624(200206)165:3<249::AID-JPLN1111249>3.0.CO;2-N.

Hu, Z. *et al.* (2003) ‘Comparison of mineralization and distribution of soil sulfur fractions in the rhizosphere of oilseed rape and rice’, *Communications in Soil Science and Plant Analysis*, 34(15–16), pp. 2243–2257. doi: 10.1081/CSS-120024061.

Huang, Z.-Z. *et al.* (1992) ‘Short-term experiments on ion transport by seedlings and excised roots: technique and validity’, *Plant physiology*. Am Soc Plant Biol, 100(4), pp. 1914–1920.

Inouhe, M. (2005) ‘Phytochelatin’, *Brazilian Journal of Plant Physiology*. SciELO Brasil, 17(1), pp. 65–78.

J., P., K., G. and R., M. (2003) ‘Comparison of three different methods of determining soil sulphur mineralization in relation to plant sulphur availability in soils’, *Biology and Fertility of Soils*, 34(3), pp. 131–139. doi: 10.1007/s003740100378.

Jamal, A., Moon, Y. S. and Abdin, M. Z. (2010) ‘Sulphur -a general overview and interaction with nitrogen’, *Australian Journal of Crop Science*, 4(7), pp. 523–529.

Johnson, D. W., Henderson, G. S. and Todd, D. E. (1981) ‘Evidence of modern accumulations of adsorbed sulfate in an east Tennessee forested Ultisol’, *Soil Science*, 132(6), pp. 422–426. doi: 10.1097/00010694-198112000-00006.

Johnston, M. *et al.* (1981) ‘Mechanistic Studies on Reactions of Bacterial Methionine  $\gamma$ -Lyase with Olefinic Amino Acids’, *Biochemistry*, 20(15), pp. 4325–4333. doi: 10.1021/bi00518a014.

Jokic, A. *et al.* (2003) ‘Organic carbon and sulphur compounds in wetland soils: Insights on structure and transformation processes using K-edge XANES and NMR spectroscopy’,

- Geochimica et Cosmochimica Acta*, 67(14), pp. 2585–2597. doi: 10.1016/S0016-7037(03)00101-7.
- Jones, D. L., Hodge, A. and Kuzyakov, Y. (2004) ‘Plant and mycorrhizal regulation of rhizodeposition’, *New Phytologist*, 163(3), pp. 459–480. doi: 10.1111/j.1469-8137.2004.01130.x.
- Jones, D. L., Nguyen, C. and Finlay, R. D. (2009) ‘Carbon flow in the rhizosphere: Carbon trading at the soil-root interface’, *Plant and Soil*, pp. 5–33. doi: 10.1007/s11104-009-9925-0.
- Jørgensen, B. B. (1990) ‘The sulfur cycle of freshwater sediments: Role of thiosulfate’, *Limnology and Oceanography*, 35(6), pp. 1329–1342. doi: 10.4319/lo.1990.35.6.1329.
- Jungk, A. (2009) ‘Carl sprengel-the founder of agricultural chemistry a re-appraisal commemorating the 150th anniversary of his death’, *Journal of Plant Nutrition and Soil Science*, 172(5), pp. 633–636. doi: 10.1002/jpln.200900065.
- Kabdasli, I., Tünay, O. and Orhon, D. (1995) ‘Sulfate removal from indigo dyeing textile wastewaters’, *Water Science and Technology*. IWA Publishing, 32(12), p. 21.
- Kandyliis, K. (1984) ‘The role of sulphur in ruminant nutrition. A review’, *Livestock Production Science*, 11(6), pp. 611–624. doi: 10.1016/0301-6226(84)90075-7.
- Karimizarchi, M. *et al.* (2014) ‘Elemental sulphur application effects on nutrient availability and sweet maize (*Zea mays* L.) response in a high pH soil of Malaysia’, *Malaysian Journal of Soil Science*, 18, pp. 75–86.
- Kasrai, M. *et al.* (1990) ‘Characterization of sulphur in coal from sulphur L-edge XANES spectra’, *Fuel*. Elsevier, 69(4), pp. 411–414.
- Kasrai, M. *et al.* (1996) ‘Sulphur characterization in coal from X-ray absorption near edge

- spectroscopy', *International Journal of Coal Geology*, 32(1–4), pp. 107–135. doi: 10.1016/S0166-5162(96)00033-X.
- Kertesz, M. A. and Mirleau, P. (2004a) 'The role of soil microbes in plant sulphur nutrition', *Journal of Experimental Botany*, 55(404), pp. 1939–1945. doi: 10.1093/jxb/erh176.
- Kertesz, M. A. and Mirleau, P. (2004b) 'The role of soil microbes in plant sulphur nutrition', *Journal of Experimental Botany*, 55(404), pp. 1939–1945. doi: 10.1093/jxb/erh176.
- Kielland, K. (1994) 'Amino acid absorption by arctic plants: implications for plant nutrition and nitrogen cycling', *Ecology*. Wiley Online Library, 75(8), pp. 2373–2383.
- Kiene, R. P. (1996) 'Microbial cycling of organosulfur gases in marine and freshwater environments', *Internationale Vereinigung für Theoretische und Angewandte Limnologie: Mitteilungen*. Taylor & Francis, 25(1), pp. 137–151.
- King, G. M. and Klug, M. J. (1982) 'Comparative aspects of sulfur mineralization on sediments of a eutrophic lake basin', *Applied and Environmental Microbiology*, 43(6), pp. 1406–1412. doi: 10.1128/AEM.43.6.1406-1412.1982.
- Klose, S. and Tabatabai, M. A. (1999) 'Arylsulfatase Activity of Microbial Biomass in Soils', *Soil Science Society of America Journal*, 63(3), pp. 569–574. doi: 10.2136/sssaj1999.03615995006300030020x.
- Kopriva, S. *et al.* (2016) 'Editorial: Frontiers of sulfur metabolism in plant growth, development, and stress response', *Frontiers in Plant Science*, 6(JAN2016), pp. 2015–2017. doi: 10.3389/fpls.2015.01220.
- Koschorreck, M. (2008) 'Microbial sulphate reduction at a low pH', *FEMS Microbiology Ecology*, 64(3), pp. 329–342. doi: 10.1111/j.1574-6941.2008.00482.x.
- Kubono, I. *et al.* (2017) 'Mössbauer spectra of iron (III) sulfide particles', *Hyperfine*



- Interactions*, 238(1), pp. 1–5. doi: 10.1007/s10751-017-1465-z.
- Kuenen, J. G. and Beudeker, R. F. (1982) ‘Microbiology of thiobacilli and other sulphur-oxidizing autotrophs, mixotrophs and heterotrophs.’, *Philosophical transactions of the Royal Society of London. Series B, Biological sciences*, 298(1093), pp. 473–497. doi: 10.1098/rstb.1982.0093.
- Kühn-institut, J. (2019a) ‘Aus dem Institut für Pflanzenernährung und Bodenkunde Schnug E Sulphur nutrition and quality of vegetables Manuskript , zu finden in [www.fal.de](http://www.fal.de) Braunschweig Bundesforschungsanstalt für Landwirtschaft ( FAL ) Erscheinungsjahr 1990 Also available at : [http](http://www.fal.de)’, (January 1990).
- Kühn-institut, J. (2019b) ‘Aus dem Institut für Pflanzenernährung und Bodenkunde Schnug E Sulphur nutritional status of European crops and consequences for agriculture Manuskript , zu finden in [www.fal.de](http://www.fal.de) Braunschweig Bundesforschungsanstalt für Landwirtschaft ( FAL ) Erscheinungsj’, (May 2016).
- Kumagai, H. *et al.* (1974) ‘Synthesis of S-alkyl-L-cysteine from pyruvate, ammonia and alkyl-mercaptan by cysteine desulphydrase of *Aerobacter aerogenes*’, *Biochemical and Biophysical Research Communications*, 59(2), pp. 789–795. doi: 10.1016/S0006-291X(74)80049-5.
- Kushkevych, I. *et al.* (2020) ‘Recent Advances in Metabolic Pathways of Sulfate Reduction in Intestinal Bacteria’, *Cells*, 9(3), p. 698. doi: 10.3390/cells9030698.
- Kuzyakov, Y. (2002) ‘Separating microbial respiration of exudates from root respiration in non-sterile soils: A comparison of four methods’, *Soil Biology and Biochemistry*, 34(11), pp. 1621–1631. doi: 10.1016/S0038-0717(02)00146-3.
- Kuzyakov, Y. (2006) ‘Sources of CO<sub>2</sub> efflux from soil and review of partitioning methods’,

- Soil biology and biochemistry*. Elsevier, 38(3), pp. 425–448.
- Kuzyakov, Y. and Domanski, G. (2000) ‘Carbon input by plants into the soil. Review’, *Journal of Plant Nutrition and Soil Science*, 163(4), pp. 421–431. doi: 10.1002/1522-2624(200008)163:4<421::AID-JPLN421>3.0.CO;2-R.
- Landerst, D. H., David, M. B. and Mitchell, M. J. (1983) ‘Analysis of Organic and Inorganic Sulfur Constituents in Sediments, Soils and Water’, *International Journal of Environmental Analytical Chemistry*. Taylor & Francis, 14(4), pp. 245–256. doi: 10.1080/03067318308071623.
- Lang, C. *et al.* (2007) ‘Sulphite oxidase as key enzyme for protecting plants against sulphur dioxide’, *Plant, Cell and Environment*, 30(4), pp. 447–455. doi: 10.1111/j.1365-3040.2006.01632.x.
- Larsson, C. *et al.* (1991) ‘Translocation and cycling through roots of recently absorbed nitrogen and sulphur in wheat (*Triticum aestivum*) during vegetative and generative growth’, *Physiologia Plantarum*. Wiley Online Library, 82(3), pp. 345–352.
- Last, F. T. (1982) ‘Effects of atmospheric sulphur compounds on natural and man-made terrestrial and aquatic ecosystems’, *Agriculture and Environment*, 7(3–4), pp. 299–387. doi: 10.1016/0304-1131(82)90022-4.
- Leake, J. R. (2004) ‘Myco-heterotroph/epiparasitic plant interactions with ectomycorrhizal and arbuscular mycorrhizal fungi’, *Current Opinion in Plant Biology*, pp. 422–428. doi: 10.1016/j.pbi.2004.04.004.
- Lehmann, J. and Kleber, M. (2015) ‘The contentious nature of soil organic matter’, *Nature*, 528(7580), pp. 60–68. doi: 10.1038/nature16069.
- Lehmann, J. and Schroth, G. (2003) ‘Nutrient leaching’, *Trees, Crops and Soil Fertility*, CABI

- Publishing, Wallingford*, pp. 151–166.
- Leustek, T. (2002) ‘Sulfate metabolism’, *The Arabidopsis Book/American Society of Plant Biologists*. American Society of Plant Biologists, 1.
- Liss, P. S. *et al.* (1997) ‘Marine sulphur emissions’, *Philosophical Transactions of the Royal Society B: Biological Sciences*, 352(1350), pp. 159–169. doi: 10.1098/rstb.1997.0011.
- Liss, P. S., Malin, G. and Turner, S. M. (1993) *Production of DMS by marine phytoplankton, Dimethylsulphide: oceans, atmosphere and climate. Proc. international symposium, Belgirate, 1992*. Kluwer Dordrecht. doi: 10.1007/978-94-017-1261-3\_1.
- Lockwood, H. C. (1954) ‘Determination of lead in foodstuffs’, in *The Analyst*, pp. 143–146. doi: 10.1039/an9547900143.
- Lomans, B. P. *et al.* (2002) ‘Cellular and Molecular Life Sciences Microbial cycling of volatile organic sulfur compounds’, *Cellular and molecular life sciences : CMLS*, 59, pp. 575–588. doi: 10.1007/s00018-002-8450-6.
- Losher, A. J. and Kelts, K. R. (1989) ‘Organic sulphur fixation in freshwater lake sediments and the implication for US ratios’, *Terra Nova*. Wiley Online Library, 1(3), pp. 253–261.
- Lovley, D. R., Holmes, D. E. and Nevin, K. P. (1991) ‘Dissimilatory Fe ( ) and Mn (IV) reduction’, *Microbiological Reviews*, 55(2), pp. 259–287.
- Lucash, M. S. *et al.* (2007) ‘Estimating nutrient uptake by mature tree roots under field conditions: Challenges and opportunities’, *Trees - Structure and Function*, pp. 593–603. doi: 10.1007/s00468-007-0160-0.
- Lunn, J. E. *et al.* (1990) ‘Localization of ATP sulfurylase and O-acetylserine (thiol) lyase in spinach leaves’, *Plant Physiology*. Am Soc Plant Biol, 94(3), pp. 1345–1352.

- Luptakova, A. (2007) 'Importance of Sulphate Reducing Bacteria in Environment', *Nova Biotechnology*, 7(1), pp. 17–22.
- Lynch, J. M. and Whipps, J. M. (1990) 'Substrate flow in the rhizosphere', *Plant and Soil*, 129(1), pp. 1–10. doi: 10.1007/BF00011685.
- Maher, C. A. and Geoscience, S. C. (2013) 'Examining geochemical processes in acid sulphate soils using stable sulphur isotopes Examining geochemical processes in acid sulfate soils using stable sulfur isotopes'.
- Martin, A. (1980) 'Sulphur in air and deposited from air and rain over Great Britain and Ireland', *Environmental Pollution. Series B, Chemical and Physical*, 1(3), pp. 177–193. doi: 10.1016/0143-148X(80)90010-5.
- Martínez-Ballesta, M. del C., Moreno, D. A. and Carvajal, M. (2013) 'The physiological importance of glucosinolates on plant response to abiotic stress in Brassica', *International Journal of Molecular Sciences*, 14(6), pp. 11607–11625. doi: 10.3390/ijms140611607.
- Marusak, L. A. and Tongson, L. L. (1979) 'Soft x-ray emission and Auger electron spectroscopic study of FeS, Fe<sub>0.9</sub>S, Fe<sub>0.875</sub>S, and Fe<sub>0.5</sub>S', *Journal of Applied Physics*, 50(6), pp. 4350–4355. doi: 10.1063/1.326420.
- Maseko, S. T. and Dakora, F. D. (2013) 'Plant enzymes, root exudates, cluster roots and mycorrhizal symbiosis are the drivers of P nutrition in native legumes growing in P deficient soil of the Cape fynbos in South Africa', *Journal of Agricultural Science and Technology*. A. David Publishing Company, Inc., 3(5A), p. 331.
- Maynard, D. G., Stewart, J. W. B. and Bettany, J. R. (1983) 'Sulfur and nitrogen mineralization in soils compared using two incubation techniques', *Soil Biology and Biochemistry*,

- 15(3), pp. 251–256. doi: 10.1016/0038-0717(83)90067-6.
- McFarlane, C. and Wickliff, C. (1985) ‘Excised Barley root uptake of several  $^{14}\text{C}$  labeled organic compounds’, *Environmental monitoring and assessment*. Springer, 5(4), pp. 385–391.
- McGill, W. B. and Cole, C. V. (1981) ‘Comparative aspects of cycling of organic C, N, S and P through soil organic matter’, *Geoderma*. Elsevier, 26(4), pp. 267–286. doi: 10.1016/0016-7061(81)90024-0.
- McGrath, S. P., Zhao, F. J. and Blake-Kalff, M. M. (2003) ‘History and outlook for sulphur fertilizers in Europe’, *Fertilizers Fertilization*, 2(15), pp. 5–27.
- McLaren, R. G. and Swift, R. S. (1977) ‘Changes in soil organic sulphur fractions due to the long term cultivation of soils’, *Journal of Soil Science*. Wiley Online Library, 28(3), pp. 445–453. doi: 10.1111/j.1365-2389.1977.tb02252.x.
- Meena, V. S. and Improvement, C. (2018) *Role of Rhizospheric Microbes in Soil, Role of Rhizospheric Microbes in Soil*. doi: 10.1007/978-981-13-0044-8.
- Meneses, S. R. P., Maniasso, N. and Zagatto, E. A. G. (2005) ‘Spectrophotometric flow-injection determination of sulphate in soil solutions’, *Talanta*, 65(5), pp. 1313–1317. doi: 10.1016/j.talanta.2004.09.010.
- Millerd, A., Thomson, J. A. and Randall, P. J. (1979) ‘Heterogeneity of sulphur content in the storage proteins of pea cotyledons’, *Planta*. Springer, 146(4), pp. 463–466.
- Mitchell, M. J. *et al.* (1984) ‘Organic and inorganic sulfur constituents of the Sediments in three New York lakes: Effect of site, sediment depth and season’, *Water, Air, and Soil Pollution*, 21(1–4), pp. 231–245. doi: 10.1007/BF00163627.
- Mitchell, S. (1996) *Biology of sulfur*. CRC Press: New York.

- Moat, A. G., Foster, J. W. and Spector, M. P. (2003) *Microbial physiology*. John Wiley & Sons.
- Moldan, F. *et al.* (2012) 'Experimental simulation of the effects of extreme climatic events on major ions, acidity and dissolved organic carbon leaching from a forested catchment, Gårdsjön, Sweden', *Biogeochemistry*. Springer, 107(1–3), pp. 455–469.
- Möller, D. (1984) 'On the global natural sulphur emission', *Atmospheric Environment (1967)*, 18(1), pp. 29–39. doi: 10.1016/0004-6981(84)90226-9.
- Muktamar, Z. (1993) 'Sorption and desorption of sulfate from Kentucky soils.'
- Mullineaux, P. M. and Rausch, T. (2005) 'Glutathione, photosynthesis and the redox regulation of stress-responsive gene expression', *Photosynthesis Research*, 86(3), pp. 459–474. doi: 10.1007/s11120-005-8811-8.
- Munter, R. C. and Grande, R. E. (1981) 'Plant tissue and soil extract analysis by ICP-AES', in *Developments in atomic plasma spectrochemical analysis*, pp. 653–673.
- Murphy, A. M. D. and Boggan, J. M. (1988) 'Sulphur Deficiency in Herbage in Ireland : 1 . Causes and Extent', 27(1), pp. 83–90. Available at: <https://www.jstor.org/stable/25556208>.
- Nannipieri, P., Kandeler, E. and Ruggiero, P. (2002) 'Enzyme Activities and Microbiological and Biochemcial Processes in Soil', in *Enzymes in the Environment*. Marcel Dekker, New York, pp. 1–33. doi: 10.1201/9780203904039.ch1.
- Neff, J. C. and Asner, G. P. (2001) 'Dissolved organic carbon in terrestrial ecosystems: Synthesis and a model', *Ecosystems*, 4(1), pp. 29–48. doi: 10.1007/s100210000058.
- Neumann, G. and Römheld, V. (2007) 'The release of root exudates as affected by the plant physiological status', *The Rhizosphere: Biochemistry and organic substances at the soil-plant interface*. CRC Press Boca Raton, FL, 2, pp. 23–72.

- Ngo, T. T. and Shargool, P. D. (1974) 'The enzymatic synthesis of L-cysteine in higher plant tissues', *Canadian journal of biochemistry*. NRC Research Press Ottawa, Canada, 52(6), pp. 435–440.
- Nimni, M. E., Han, B. and Cordoba, F. (2007) 'Are we getting enough sulfur in our diet?', *Nutrition and Metabolism*, pp. 1–12. doi: 10.1186/1743-7075-4-24.
- NISHIO, T. (1991) '<sup>15</sup>N-NH<sub>4</sub><sup>+</sup> Isotope Dilution Method for Analyzing Nitrogen Transformation in Upland Soils', *JARQ. Japan Agricultural Research Quarterly*, 25(2), pp. 88–92.
- Novozamsky, I. *et al.* (1986) 'Determination of total sulphur and extractable sulphate in plant materials by inductively-coupled plasma atomic emission spectrometry', *Communications in Soil Science and Plant Analysis*. Taylor & Francis, 17(11), pp. 1147–1157. doi: 10.1080/00103628609367780.
- Nziguheba, G., Smolders, E. and Merckx, R. (2005) 'Sulphur immobilization and availability in soils assessed using isotope dilution', *Soil Biology and Biochemistry*, 37(4), pp. 635–644. doi: 10.1016/j.soilbio.2004.09.007.
- Oburger, E. and Jones, D. L. (2018) 'Sampling root exudates – Mission impossible?', *Rhizosphere*. Elsevier, 6(June), pp. 116–133. doi: 10.1016/j.rhisph.2018.06.004.
- Odom, J. M. and Peck Jr, H. D. (1981) 'Hydrogen cycling as a general mechanism for energy coupling in the sulfate-reducing bacteria, *Desulfovibrio* sp', *FEMS Microbiol. Lett*, 12(1).
- Onwona-Kwakye, M., Hogarth, J. N. and Van den Brink, P. J. (2020) 'Environmental risk assessment of pesticides currently applied in Ghana', *Chemosphere*. Elsevier Ltd, 254, p. 126845. doi: 10.1016/j.chemosphere.2020.126845.

- Osslund, T., Chandler, C. and Segel, I. H. (1982) 'ATP sulfurylase from higher plants: purification and preliminary kinetics studies on the cabbage leaf enzyme', *Plant physiology*. Am Soc Plant Biol, 70(1), pp. 39–45.
- Park, C. C. (2013) *Acid Rain (routledge revivals): rhetoric and reality*. Routledge.
- Paterson, E. (2003) 'Importance of rhizodeposition in the coupling of plant and microbial productivity', *European Journal of Soil Science*, 54(4), pp. 741–750. doi: 10.1046/j.1351-0754.2003.0557.x.
- Peñuelas, J. *et al.* (2014) 'Biogenic volatile emissions from the soil', *Plant, Cell and Environment*, pp. 1866–1891. doi: 10.1111/pce.12340.
- Perrott, K. W. *et al.* (1991) 'Determination of total sulphur in soil using inductively coupled plasma-atomic emission spectrometry', *Communications in Soil Science and Plant Analysis*. Taylor & Francis, 22(13–14), pp. 1477–1487. doi: 10.1080/00103629109368507.
- Persson, J. and Näsholm, T. (2001a) 'A GC-MS method for determination of amino acid uptake by plants', *Physiologia Plantarum*, 113(3), pp. 352–358. doi: 10.1034/j.1399-3054.2001.1130308.x.
- Persson, J. and Näsholm, T. (2001b) 'Amino acid uptake: A widespread ability among boreal forest plants', *Ecology Letters*, 4(5), pp. 434–438. doi: 10.1046/j.1461-0248.2001.00260.x.
- Pester, M. *et al.* (2012) 'Sulfate-reducing microorganisms in wetlands - fameless actors in carbon cycling and climate change', *Frontiers in Microbiology*, 3(FEB), pp. 1–19. doi: 10.3389/fmicb.2012.00072.
- Pickering, I. J. *et al.* (1998) 'Sulfur K-edge X-ray absorption spectroscopy for determining the



- chemical speciation of sulfur in biological systems', *FEBS Letters*, 441(1), pp. 11–14.  
doi: 10.1016/S0014-5793(98)01402-1.
- Pinton, R., Varanini, Z. and Nannipieri, P. (2007) *The rhizosphere: biochemistry and organic substances at the soil-plant interface*. CRC press.
- Pirela, H. J. and Tabatabai, M. A. (1988) 'Sulfur mineralization rates and potentials of soils', *Biology and Fertility of Soils*, 6(1), pp. 26–32. doi: 10.1007/BF00257916.
- Plante, A. F. (2006) 'Soil biogeochemical cycling of inorganic nutrients and metals', *Soil Microbiology, Ecology and Biochemistry: Third Edition*, pp. 389–432.
- Postgate, J. R. and Campbell, L. L. (1966) 'Classification of Desulfovibrio species, the nonsporulating sulfate-reducing bacteria.', *Bacteriological reviews*, 30(4), pp. 732–738. doi: 10.1128/membr.30.4.732-738.1966.
- Prietz, J. *et al.* (2013) 'Soil sulphur speciation in two glacier forefield soil chronosequences assessed by S K-edge XANES spectroscopy', *European Journal of Soil Science*, 64(2), pp. 260–272. doi: 10.1111/ejss.12032.
- Rahul, S. *et al.* (2018) 'Sulphur nutrition in maize—a critical review', *Int J Pure Appl Biosci*, 5, pp. 1582–1596.
- Rasmussen, K. H. and Kabel, R. L. (1974) 'POLLUTANTS', 4(1975), pp. 33–64.
- Raue, B., Brauch, H. J. and Frimmel, F. H. (1991) 'Determination of sulphate in natural waters by ICP/OES - comparative studies with ion chromatography', *Fresenius' Journal of Analytical Chemistry*, 340(6), pp. 395–398. doi: 10.1007/BF00321590.
- Rausch, T. and Wachter, A. (2005) 'Sulfur metabolism: A versatile platform for launching defence operations', *Trends in Plant Science*, 10(10), pp. 503–509. doi: 10.1016/j.tplants.2005.08.006.

- Rauser, W. E. (1995) 'Phytochelatins and related peptides. Structure, biosynthesis, and function', *Plant Physiology*, 109(4), pp. 1141–1149. doi: 10.1104/pp.109.4.1141.
- Rennenberg, E. H. *et al.* (1990) *Higher Plants*. Available at: [https://www.researchgate.net/profile/Luit\\_De\\_Kok/publication/257820230\\_Sulfur\\_Nutrition\\_and\\_Sulfur\\_Assimilation\\_in\\_Higher\\_Plants/links/55a7acb008aea2222c74875a/Sulfur-Nutrition-and-Sulfur-Assimilation-in-Higher-Plants.pdf#page=61](https://www.researchgate.net/profile/Luit_De_Kok/publication/257820230_Sulfur_Nutrition_and_Sulfur_Assimilation_in_Higher_Plants/links/55a7acb008aea2222c74875a/Sulfur-Nutrition-and-Sulfur-Assimilation-in-Higher-Plants.pdf#page=61).
- Rennenberg, H. (1984) 'The Fate of Excess Sulfur in Higher Plants', *Annual Review of Plant Physiology*, 35(1), pp. 121–153. doi: 10.1146/annurev.pp.35.060184.001005.
- Rennenberg, H. and Polle, A. (1994) 'Metabolic consequences of atmospheric sulphur influx into plants', in *Plant responses to the gaseous environment*. Springer, pp. 165–180.
- Riley, N. G., Zhao, F. J. and McGrath, S. P. (2002) 'Sulphur Fertilizers : a ® Eld Lysimeter Study', pp. 120–126. doi: 10.1079/SUM2002115.
- Riley, N. G., Zhao, F. J. and McGrath, S. P. (2002) 'Leaching losses of sulphur from different forms of sulphur fertilizers: A field lysimeter study', *Soil Use and Management*, 18(2), pp. 120–126. doi: 10.1111/j.1475-2743.2002.tb00229.x.
- Rittenberg, D. and Foster, G. L. (1940) 'A new procedure for quantitative analysis by isotope dilution, with application to the determination of amino acids and fatty acids', *Journal of Biological Chemistry*, 133(3), pp. 737–744. doi: 10.1016/s0021-9258(18)73304-8.
- Rodhe, H. (1999) 'Human impact on the atmospheric sulfur balance', *Tellus B*. Wiley Online Library, 51(1), pp. 110–122.
- Romero, L. C. *et al.* (2014) 'Cysteine and cysteine-related signaling pathways in arabidopsis thaliana', *Molecular Plant*, 7(2), pp. 264–276. doi: 10.1093/mp/sst168.
- Rousk, J. and Jones, D. L. (2010) 'Loss of low molecular weight dissolved organic carbon

- (DOC) and nitrogen (DON) in H<sub>2</sub>O and 0.5M K<sub>2</sub>SO<sub>4</sub> soil extracts', *Soil Biology and Biochemistry*. Elsevier Ltd, 42(12), pp. 2331–2335. doi: 10.1016/j.soilbio.2010.08.017.
- Rovira, A. D. (1969) 'Plant root exudates', *The Botanical Review*, 35(1), pp. 35–57. doi: 10.1007/BF02859887.
- Rubio, G., Sorgona, A. and Lynch, J. P. (2004) 'Spatial mapping of phosphorus influx in bean root systems using digital autoradiography', *Journal of experimental botany*. Oxford University Press, 55(406), pp. 2269–2280.
- Rumpel, C., Eusterhues, K. and Kögel-Knabner, I. (2004) 'Location and chemical composition of stabilized organic carbon in topsoil and subsoil horizons of two acid forest soils', *Soil Biology and Biochemistry*, 36(1), pp. 177–190. doi: 10.1016/j.soilbio.2003.09.005.
- Sacchi, G. a *et al.* (2000) 'Efflux and active re-absorption of glucose in roots of cotton plants grown under saline conditions', *Plant and Soil*, 220, pp. 1-2 ST-Efflux and active re-absorption of gluco. doi: 10.1023/A:1004701912815.
- Saggar, S., Bettany, J. R. and Stewart, J. W. B. (1981) 'Measurement of microbial sulfur in soil', *Soil Biology and Biochemistry*, 13(6), pp. 493–498. doi: 10.1016/0038-0717(81)90040-7.
- Saifullah Khan, K. and Georg Joergensen, R. (2019) 'Stoichiometry of the soil microbial biomass in response to amendments with varying C/N/P/S ratios', *Biology and Fertility of Soils*. Biology and Fertility of Soils, 55(3), pp. 265–274. doi: 10.1007/s00374-019-01346-x.
- Sakadevan, K., Mackay, A. D. and Hedley, M. J. (1993) 'Sulphur cycling in New Zealand hill country pastures. II. The fate of fertilizer sulphur', *Journal of Soil Science*. Wiley Online Library, 44(4), pp. 615–624.

- Salt, D. E. and Rauser, W. E. (1995) 'MgATP-dependent transport of phytochelatins across the tonoplast of oat roots', *Plant Physiology*, 107(4), pp. 1293–1301. doi: 10.1104/pp.107.4.1293.
- Scherer, H. W. (2001) 'Sulphur in crop production - Invited paper', *European Journal of Agronomy*, 14(2), pp. 81–111. doi: 10.1016/S1161-0301(00)00082-4.
- Scherer, H. W. (2009) 'Sulfur in soils', *Journal of Plant Nutrition and Soil Science*, 172(3), pp. 326–335. doi: 10.1002/jpln.200900037.
- Schindler, D. W. (1981) 'Interrelationships between the cycles of elements in freshwater ecosystems.', *Some Perspectives on the Major Biogeochemical Cycles*, pp. 113–123.
- Schindler, S. C. *et al.* (1986) 'Incorporation of <sup>35</sup>S- sulfate into inorganic and organic constituents of two forest soils.', *Soil Science Society of America Journal*, 50(2).
- Schinner, F. *et al.* (2012) *Methods in soil biology*. Springer Science & Business Media.
- Schmidt, M. W. I. *et al.* (2011) 'Persistence of soil organic matter as an ecosystem property', *Nature*, 478(7367), pp. 49–56. doi: 10.1038/nature10386.
- Schnug, E. (1990) 'Sulphur nutrition and quality of vegetables', *Sulphur Agric*, 14, pp. 3–7.
- Schnug, E., Haneklaus, S. and Murphy, D. (1993) 'Impact of sulphur fertilization on fertilizer nitrogen efficiency', *Sulphur in Agriculture*, 17(May 2016), pp. 8–12.
- Schoenau, J. J. and Malhi, S. S. (2008) 'Sulfur forms and cycling processes in soil and their relationship to sulfur fertility', *Sulfur: A missing link between soils, crops, and nutrition*. Wiley Online Library, 50, pp. 1–10.
- Schuler, M. S. *et al.* (2019) 'Regulations are needed to protect freshwater ecosystems from salinization', *Philosophical Transactions of the Royal Society B: Biological Sciences*,

374(1764). doi: 10.1098/rstb.2018.0019.

Scott, N. M. and Anderson, G. (1976) 'Organic sulphur fractions in scottish soils', *Journal of the Science of Food and Agriculture*, 27(4), pp. 358–366. doi: 10.1002/jsfa.2740270410.

Shadle, S. E. *et al.* (1993) 'X-ray Absorption Spectroscopic Studies of the Blue Copper Site: Metal and Ligand K-Edge Studies To Probe the Origin of the EPR Hyperfine Splitting in Plastocyanin', *Journal of the American Chemical Society*, 115(2), pp. 767–776. doi: 10.1021/ja00055a057.

Shahsavani, S. and Gholami, A. (2009) 'Study on the effect of sulphur, glucose, nitrogen and plant residues on the immobilization of sulphate-s in soil', *World Academy of Science, Engineering and Technology*, 37, pp. 114–118.

Sheat, D. E. G., Fletcher, B. H. and Street, H. E. (1959) 'Studies on the growth of excised roots: VIII. The growth of excised tomato roots supplied with various inorganic sources of nitrogen', *New Phytologist*. Wiley Online Library, 58(2), pp. 128–141.

Shinn, J. H. and Lynn, S. (1979) 'Do man-made sources affect the sulfur cycle of northeastern states?', *Environmental Science & Technology*, 13(9), pp. 1062–1067. doi: 10.1021/es60157a003.

Smith, F. W. *et al.* (1997) 'Regulation of expression of a cDNA from barley roots encoding a high affinity sulphate transporter', *Plant Journal*, 12(4), pp. 875–884. doi: 10.1046/j.1365-313X.1997.12040875.x.

Smith, F. W., Rae, A. L. and Hawkesford, M. J. (2000) 'Molecular mechanisms of phosphate and sulphate transport in plants', *Biochimica et Biophysica Acta - Biomembranes*, pp. 236–245. doi: 10.1016/S0005-2736(00)00141-3.

- Smith, J. L. and Paul, E. A. (1990) 'The significance of soil microbial biomass estimations', *Soil biochemistry. Vol. 6*. Marcel Dekker New York, pp. 357–396.
- Smith, S. J. *et al.* (2011) 'Anthropogenic sulfur dioxide emissions: 1850-2005', *Atmospheric Chemistry and Physics*, 11(3), pp. 1101–1116. doi: 10.5194/acp-11-1101-2011.
- Solomon, D. *et al.* (2005) 'Sulphur speciation and biogeochemical cycling in long-term arable cropping of subtropical soils: Evidence from wet-chemical reduction and S K-edge XANES spectroscopy', *European Journal of Soil Science*, 56(5), pp. 621–634. doi: 10.1111/j.1365-2389.2005.00702.x.
- Solomon, D., Lehmann, J. and Martínez, C. E. (2003) 'Sulfur K-edge XANES spectroscopy as a tool for understanding sulfur dynamics in soil organic matter', *Soil Science Society of America Journal*, 67(6), p. 1721. doi: 10.2136/sssaj2003.1721.
- Song, G. *et al.* (2005) 'Topsoil organic carbon storage of China and its loss by cultivation', *Biogeochemistry*, 74(1), pp. 47–62. doi: 10.1007/s10533-004-2222-3.
- Sors, T. G., Ellis, D. R. and Salt, D. E. (2005) 'Selenium uptake, translocation, assimilation and metabolic fate in plants', *Photosynthesis Research*, 86(3), pp. 373–389. doi: 10.1007/s11120-005-5222-9.
- Spiro, C. L. *et al.* (1984) 'X-ray absorption spectroscopic investigation of sulfur sites in coal: Organic sulfur identification', *Science*. American Association for the Advancement of Science, 226(4670), pp. 48–50.
- Spohn, M. (2016) 'Element cycling as driven by stoichiometric homeostasis of soil microorganisms', *Basic and Applied Ecology*. Elsevier GmbH, pp. 471–478. doi: 10.1016/j.baae.2016.05.003.
- Spohn, M. *et al.* (2016) 'Soil microbial carbon use efficiency and biomass turnover in a long-

- term fertilization experiment in a temperate grassland', *Soil Biology and Biochemistry*, 97, pp. 168–175. doi: 10.1016/j.soilbio.2016.03.008.
- STAUBES, R., GEORGII, H. -W and OCKELMANN, G. (1989) 'Flux of COS, DMS and CS<sub>2</sub> from various soils in Germany', *Tellus B*, 41 B(3), pp. 305–313. doi: 10.1111/j.1600-0889.1989.tb00309.x.
- Stevenson, F. J. and Cole, M. A. (1999) *Cycles of soils: carbon, nitrogen, phosphorus, sulfur, micronutrients*. John Wiley & Sons.
- Stressler, T. *et al.* (2016) 'Detection, production, and application of microbial arylsulfatases', *Applied Microbiology and Biotechnology*. Applied Microbiology and Biotechnology, 100(21), pp. 9053–9067. doi: 10.1007/s00253-016-7838-4.
- Strick, J. E. and Nakas, J. P. (1984) 'Calibration of a microbial sulfur technique for use in forest soils', *Soil Biology and Biochemistry*, 16(3), pp. 289–291. doi: 10.1016/0038-0717(84)90020-8.
- Strohm, M. *et al.* (1995) 'Regulation of glutathione synthesis in leaves of transgenic poplar (*Populus tremula* x *P. alba*) overexpressing glutathione synthetase', *The Plant Journal*. Wiley Online Library, 7(1), pp. 141–145.
- Su, Y.-H. and Zhu, Y.-G. (2007) 'Transport mechanisms for the uptake of organic compounds by rice (*Oryza sativa*) roots', *Environmental Pollution*. Elsevier, 148(1), pp. 94–100.
- Sugiura, C. (1981) 'Sulfur K x-ray absorption spectra of FeS, FeS<sub>2</sub>, and Fe<sub>2</sub>S<sub>3</sub>', *The Journal of Chemical Physics*. American Institute of Physics, 74(1), pp. 215–217.
- Swain, S. C. and Padhi, S. K. (2015) 'Effect of Sulphur dioxide on growth, Chlorophyll and sulphur contents of Pomegranate', *Tropical Agricultural Research and Extension*, 16(1), p. 21. doi: 10.4038/tare.v16i1.5269.

- T. Sherene (2017) 'Role of Soil Enzymes in Nutrient Transformation: A Review', *Bio Bulletin*, 3 (1)(1), pp. 109–131.
- Tabatabai, M. A. and Bremner, J. M. (1970) 'Arylsulfatase Activity of Soils', *Soil Science Society of America Journal*, 34(2), pp. 225–229. doi: 10.2136/sssaj1970.03615995003400020016x.
- Takahashi, H. *et al.* (1996) 'Isolation and characterization of a cDNA encoding a sulfate transporter from *Arabidopsis thaliana*', *FEBS Letters*, 392(2), pp. 95–99. doi: 10.1016/0014-5793(96)00787-9.
- Takahashi, H. *et al.* (2000) 'The roles of three functional sulphate transporters involved in uptake and translocation of sulphate in *Arabidopsis thaliana*', *Plant Journal*, 23(2), pp. 171–182. doi: 10.1046/j.1365-3113X.2000.00768.x.
- Turrión, M. B., Gallardo, J. F. and González, M. I. (1999) 'Extraction of soil-available phosphate, nitrate, and sulphate ions using ion exchange membranes and determination by ion exchange chromatography', *Communications in Soil Science and Plant Analysis*, 30(7–8), pp. 1137–1152. doi: 10.1080/00103629909370274.
- Vairavamurthy, A. *et al.* (1994) 'Sulfonates: A novel class of organic sulfur compounds in marine sediments', *Geochimica et Cosmochimica Acta*, 58(21), pp. 4681–4687. doi: 10.1016/0016-7037(94)90200-3.
- Vairavamurthy, M. A. *et al.* (1997) 'Characterization of sulfur-containing functional groups in sedimentary humic substances by X-ray absorption near-edge structure spectroscopy', *Energy and Fuels*, 11(3), pp. 546–553. doi: 10.1021/ef960212a.
- Valeur, I. and Nilsson, I. (1993) 'Effects of lime and two incubation techniques on sulfur mineralization in a forest soil', *Soil Biology and Biochemistry*, 25(10), pp. 1343–1350.



doi: 10.1016/0038-0717(93)90049-H.

Variyar, P. S. *et al.* (2014) 'Role of glucosinolates in plant stress tolerance', in *Emerging technologies and management of crop stress tolerance*. Elsevier, pp. 271–291.

Vidyalakshmi, R., Paranthaman, R. and Bhakayaraj, R. (2009) 'Sulphur Oxidizing Bacteria and Pulse Nutrition - A Review', 5(3), pp. 270–278.

Vieira Dos Santos, C. and Rey, P. (2006) 'Plant thioredoxins are key actors in the oxidative stress response', *Trends in Plant Science*, 11(7), pp. 329–334. doi: 10.1016/j.tplants.2006.05.005.

Virtanen, A. I. (1962) 'Some organic sulfur compounds in vegetables and fodder plants and their significance in human nutrition', *Angewandte Chemie International Edition in English*. Wiley Online Library, 1(6), pp. 299–306.

Vong, P. C., Nguyen, C. and Guckert, A. (2007) 'Fertilizer sulphur uptake and transformations in soil as affected by plant species and soil type', *European Journal of Agronomy*, 27(1), pp. 35–43. doi: 10.1016/j.eja.2007.01.011.

Voroney, R.P.; Brookes, P.C.; Beyaert, R. P. (2008) *Chapter 49: Soil Microbial Biomass C,N,P, and S, Soil Sampling and Methods of Analysis, Second Edition*. doi: 10.1201/9781420005271.ch18.

Voroney, R. P., Brookes, P. C. and Beyaert, R. . (2007) 'Soil Microbial Biomass C,N,P and S', in *Soil Sampling and Methods of Analysis, Second Edition*. CRC Press Boca Raton, FL, pp. 637–652. doi: 10.1201/9781420005271.ch49.

Vranová, V. *et al.* (2010) 'Dominant amino acids, organic acids and sugars in water-soluble root exudates of C4 plants: A mini-review', *Acta Universitatis Agriculturae et Silviculturae Mendelianae Brunensis*, 58(5), pp. 441–446. doi:

10.11118/actaun201058050441.

De Vries, W. and Breeuwsma, A. (1987) 'The relation between soil acidification and element cycling', *Water, Air, and Soil Pollution*. Springer, 35(3–4), pp. 293–310.

Wałaszek, K., Kryza, M. and Dore, A. J. (2013) 'The impact of precipitation on wet deposition of sulphur and nitrogen compounds', *Ecological Chemistry and Engineering S*, 20(4), pp. 733–745. doi: 10.2478/eces-2013-0051.

Walbridge, M. R. and Vitousek, P. M. (1987) 'Phosphorus mineralization potentials in acid organic soils: Processes affecting  $^{32}\text{PO}_4^{3-}$  isotope dilution measurements', *Soil Biology and Biochemistry*, 19(6), pp. 709–717. doi: 10.1016/0038-0717(87)90053-8.

Walker, K. C. and Booth, E. J. (2003) 'Sulphur Nutrition and Oilseed Quality', in *Sulphur in Plants*. Springer, pp. 323–339. doi: 10.1007/978-94-017-0289-8\_18.

Wanek, W. *et al.* (2019) 'A novel isotope pool dilution approach to quantify gross rates of key abiotic and biological processes in the soil phosphorus cycle', *Biogeosciences*, 16(15), pp. 3047–3068. doi: 10.5194/bg-16-3047-2019.

Wang, Zhanyi *et al.* (2011) 'Accumulation of different sulfur fractions in Chinese forest soil under acid deposition', *Journal of Environmental Monitoring*, 13(9), pp. 2463–2470. doi: 10.1039/c1em10313j.

Wang, Zhanyi *et al.* (2012) 'Dissolved organic sulfur in streams draining forested catchments in southern China', *Journal of Environmental Sciences*, 24(4), pp. 704–710. doi: 10.1016/S1001-0742(11)60781-X.

Warren, Y. A. *et al.* (2005) 'Biochemical differentiation and comparison of *Desulfovibrio* species and other phenotypically similar genera', *Journal of Clinical Microbiology*, 43(8), pp. 4041–4045. doi: 10.1128/JCM.43.8.4041-4045.2005.

- Watanabe, T., Kojima, H. and Fukui, M. (2014) 'Complete genomes of freshwater sulfur oxidizers *Sulfuricella denitrificans* skB26 and *Sulfuritalea hydrogenivorans* sk43H: Genetic insights into the sulfur oxidation pathway of betaproteobacteria', *Systematic and Applied Microbiology*. Elsevier GmbH., 37(6), pp. 387–395. doi: 10.1016/j.syapm.2014.05.010.
- Wen, F. *et al.* (2006) 'Extracellular Proteins in Pea Root Tip and Border Cell Exudates', *PLANT PHYSIOLOGY*, 143(2), pp. 773–783. doi: 10.1104/pp.106.091637.
- White, J. G. (1959) 'Mineralisation of nitrogen and sulphur in sulphur-deficient soils', *New Zealand Journal of Agricultural Research*, 2(2), pp. 255–258. doi: 10.1080/00288233.1959.10420311.
- Williams, C. H. and Donald, C. M. (1957) 'Changes in organic matter and pH in a podzolic soil as influenced by subterranean clover and superphosphate', *Australian Journal of Agricultural Research*, 8(2), pp. 179–189. Available at: <https://doi.org/10.1071/AR9570179>.
- Wirtz, M. and Droux, M. (2005) 'Synthesis of the sulfur amino acids: Cysteine and methionine', *Photosynthesis Research*, 86(3), pp. 345–362. doi: 10.1007/s11120-005-8810-9.
- Wittmann, C. *et al.* (2004) 'Areal activities and stratification of hydrolytic enzymes involved in the biochemical cycles of carbon, nitrogen, sulphur and phosphorus in podsolized boreal forest soils', *Soil Biology and Biochemistry*, 36(3), pp. 425–433. doi: 10.1016/j.soilbio.2003.10.019.
- Wittstock, U. and Halkier, B. A. (2002) 'Glucosinolate research in the Arabidopsis era', *Trends in Plant Science*, 7(6), pp. 263–270. doi: 10.1016/S1360-1385(02)02273-2.

- Wrigley, C. W. *et al.* (1984) 'Changes in polypeptide composition and grain quality due to sulfur deficiency in wheat', *Journal of cereal science*. Elsevier, 2(1), pp. 15–24.
- Wu, J. *et al.* (1994) 'Fumigation-extraction method for the measurement of soil microbial biomass-S', *Soil Biology and Biochemistry*, 26(1), pp. 117–125. doi: 10.1016/0038-0717(94)90203-8.
- Wu, J., O'Donnell, A. G. and Syers, J. K. (1993) 'Microbial growth and sulphur immobilization following the incorporation of plant residues into soil', *Soil Biology and Biochemistry*, 25(11), pp. 1567–1573. doi: 10.1016/0038-0717(93)90012-Z.
- Wu, Z. L. H. J. and Syers, A. G. O. D. J. K. (1997) 'Seasonal responses in microbial biomass carbon , phosphorus and sulphur in soils under pasture', pp. 421–428.
- Xia, K. *et al.* (1998) 'XANES Studies of Oxidation States of Sulfur in Aquatic and Soil Humic Substances', *Soil Science Society of America Journal*, 62(5), p. 1240. doi: 10.2136/sssaj1998.03615995006200050014x.
- Yadav, S. K. (2010) 'Heavy metals toxicity in plants: An overview on the role of glutathione and phytochelatins in heavy metal stress tolerance of plants', *South African Journal of Botany*. Elsevier B.V., 76(2), pp. 167–179. doi: 10.1016/j.sajb.2009.10.007.
- Zak, D. *et al.* (2021) 'Sulphate in freshwater ecosystems: A review of sources, biogeochemical cycles, ecotoxicological effects and bioremediation', *Earth-Science Reviews*. doi: 10.1016/j.earscirev.2020.103446.
- Zhang, M. *et al.* (2018) 'The Coupling of Carbon, Nitrogen and Sulphur Transformational Processes in River Sediments Based on Correlationship among the Functional Genes', *Biogeosciences Discussions*. Copernicus GmbH, pp. 1–39. doi: 10.5194/bg-2018-369.
- Zhao, F. J. *et al.* (1997) 'Sulphur nutrition: An important factor for the quality of wheat and

- rapeseed', *Soil Science and Plant Nutrition*, 43(SPEC. ISS.), pp. 1137–1142. doi: 10.1080/00380768.1997.11863731.
- Zhao, F. J. *et al.* (2006) 'Sulphur speciation and turnover in soils: evidence from sulphur K-edge XANES spectroscopy and isotope dilution studies', *Soil Biology and Biochemistry*, 38(5), pp. 1000–1007. doi: 10.1016/j.soilbio.2005.08.013.
- Zhao, F. J., Hawkesford, M. J. and McGrath, S. P. (1999) 'Sulphur assimilation and effects on yield and quality of wheat', *Journal of Cereal Science*, 30(1), pp. 1–17. doi: 10.1006/jcrs.1998.0241.
- Zhao, F. and McGrath, S. P. (1994) 'Extractable sulphate and organic sulphur in soils and their availability to plants', *Plant and Soil*, 164(2), pp. 243–250. doi: 10.1007/BF00010076.
- Zhou, W. *et al.* (1999) 'Mineralization of organic sulfur and its importance as a reservoir of plant-available sulfur in upland soils of north China', *Biology and Fertility of Soils*, 30(3), pp. 245–250. doi: 10.1007/s003740050615.
- Zicari, G. *et al.* (2013) 'Livestock farming and atmospheric emissions.', *Igiene e sanità pubblica*, 69(4), pp. 445–457.

## **Chapter 3**

---

### **Dissolved organic and inorganic sulphate in soils along a grassland productivity gradient**

## Abstract

Nitrogen (N), phosphorus (P) and sulphur (S) predominate in most soils in solid organic forms. However, prior to plant and microbial uptake these must first be converted to dissolved organic or inorganic forms. Although the role of some dissolved organic nutrients in plant nutrition is well established (e.g. N), the role of dissolved organic S (DOS) in plant nutrition remains unknown. However, protein breakdown is predicted to lead to the release of large quantities of DOS compounds into soil solution (e.g. S-containing amino acids and oligopeptides). We hypothesize that, similar to N, the soil solution pool will contain significant quantities of DOS and that the DOS-to-sulphate ratio will be inversely correlated to ecosystem productivity. The aim of this study was therefore to compare the relative concentrations of DOS with inorganic sulphate ( $\text{SO}_4^{2-}$ ) in a contrasting range of grassland soils. The top- (0 – 10 cm) and sub- (10 – 20 cm) soil were collected from 7 grassland pastures along a natural primary productivity gradient. The relative amounts of DOS and  $\text{SO}_4^{2-}$  present in water extracts from each soil were then evaluated. Our results showed that the concentration of DOS in all soils ranged from 10.6 to 270.3 mg kg<sup>-1</sup> for topsoils, while it ranged from 41.2 to 309.3 mg kg<sup>-1</sup> for subsoils. Higher concentrations of DOS, along with higher concentrations of  $\text{SO}_4^{2-}$ , were found in high altitude soils. The DOS pool constituted 23.8 to 94.8% of total dissolved sulphur (DOS plus  $\text{SO}_4^{2-}$ , TDS) across the range of soils. Statistical analysis showed that DOS concentration co-correlated with the concentrations of Al, B, Fe, K, Mn, ammonium in the soil extracts as well as DOC. This study shows that DOS is an important contributor of dissolved S in grassland soils, and therefore is important in S cycling and transport in soils. Further investigation is needed to evaluate the relative importance of DOS to plant S acquisition in comparison to inorganic forms of S.

**Keywords:** Dissolved organic sulphur; Sulphate; Grassland; Productivity gradient; Spatial Variation.

### 3.1 Introduction

Dissolved organic nutrients are increasingly being recognised as a major nutrient loss pathway in some ecosystems as well as being a direct source of nutrients to plants and microorganisms. This pool is often collectively known as dissolved organic matter (DOM) which is operationally defined as the continuum of organic molecules of different sizes, charges and structures that can pass through a 0.45 µm-pore filter (He *et al.*, 2016). DOM also plays a major role in freshwater ecosystems with most of the DOM entering freshwaters originating from soil. DOM is therefore receiving increasing attention due to its key role in nutrient cycling and its ability to transfer from soils to freshwater and marine ecosystems (Stedmon, Markager and Bro, 2003; Zsolnay, 2003; Amon, Benner and Saunders, 2013). Most of the studies to date on soluble organic nutrients have focused on N (Jones *et al.*, 2004; Bronk, Gilbert and Ward, 2007; Neff, III and Vitousek, 2007), with a smaller number on phosphorus (P) (Rowland and Haygarth, 1997; McDowell and Koopmans, 2006). Although sulphur (S) is the second most abundant macronutrient in DOM, information on dissolved organic sulphur (DOS) is lacking. This is surprising considering that, like N and P, organic forms of S represent the dominant form of S in soils. We therefore hypothesize that DOS must play an important role in the soil S cycle.

Sulphur occurs as reduced (e.g. sulfide and thiol) or as oxidized species (e.g. sulfonate and sulfate) in DOM. Incorporation into organic matter is an important pathway to retain S derived from inorganic S fertiliser application or atmospheric deposition. A few studies indicate that DOS constitutes a significant part of the total flux of S from terrestrial to aquatic ecosystems (Wieder and Lang, 1988; Kaiser and Guggenberger, 2005; Goller *et al.*, 2006). However, the DOS content of soil varies considerably with season and space (Peuravuori *et al.*, 2005; Wang *et al.*, 2012a). Therefore, a fundamental mechanistic understanding of the factors governing the spatial and temporal variation of DOS in the landscape will greatly



enhance our ability to develop more sustainable agricultural ecosystems as well as better understanding the different S loss pathways.

In the context of the discussion above, the first aim of this study was to quantify the spatial variations in DOS and  $\text{SO}_4^{2-}$  concentration in a range of grassland soils. The second aim was to identify parameters governing any patterns of spatial variation in soil S concentration. We approached this by analysing the chemical composition of DOS as well as  $\text{SO}_4^{2-}$  extracted from both topsoil and subsoil across a grassland productivity gradient.

## **3.2 Material and Methods**

### **3.2.1 Site description and soil sampling**

Soil was collected from seven sites located along a gradient of grassland productivity, altitude, and soil DON availability in Abergwyngregyn, Gwynedd, North Wales (Table 3.1; 53°14'N, 4°01'W; May 2019). The seven selected grassland sites represent a gradient from high productivity, inorganic N dominated grassland to low productivity, high organic matter and organic N dominated grassland. Most of these soils are assumed to be low in absorbed  $\text{SO}_4^{2-}$  and receive relatively little atmospheric sulphur (S) deposition. Although our study is based on a single elevation gradient (0-400 m a.s.l.), it was selected to include a broad range of vegetation, soil, and climatic conditions. Soil characteristics for all seven sites are presented in Table 3.1.

At each site, three independent samples from randomly positioned replicate plots ( $5 \times 5 \text{ m}^2$ ; representing 3 replicates) were collected separately from the topsoil (0 - 10 cm) and subsoil (10 - 20 cm). Soil samples were placed in gas-permeable plastic bags and transported immediately to the laboratory, where the soil was gently sieved to  $< 2 \text{ mm}$  for homogenization and to remove roots and big stones. The soil samples were then stored at  $4^\circ\text{C}$  prior to further analysis.

### 3.2.2 Soil chemical characteristics

Soil bulk density (100 cm<sup>3</sup> cores) and moisture content were determined by oven drying soil at 80 °C. Soil pH and electrical conductivity analysis was performed on a 1:2.5 w/v soil: distilled water suspension using standard electrodes. Soil microbial biomass C and N were determined using the chloroform fumigation-extraction (FE) method (Voroney, Brookes and Beyaert, 2007). For each soil sample, 5 g of field-moist soil was extracted using distilled water (ratio 1:5 w/v). A further 5 g of soil was fumigated in a CHCl<sub>3</sub> atmosphere for 48 h and then extracted in the same way. The soil and distilled water mixtures were shaken on an end-over-end shaker for 30 mins at 200 rpm (20 ± 1°C), centrifuged at 3000 rpm for 10 mins and then filtered through a Whatman No. 42 filter paper. The soil solution recovered was analysed for dissolved organic C (DOC) and total dissolved N (TDN) using a TOC/TN analyser (Shimadzu Corp., Kyoto, Japan). Microbial biomass C and N were calculated from the concentration difference from between directly extracted soil and fumigated soil samples, by using K<sub>ec</sub> and K<sub>en</sub> factors of 0.35 (Wu *et al.*, 1990) and 0.5, respectively. Field-moist soil (5 g) was shaken for 30 min with 0.5 M K<sub>2</sub>SO<sub>4</sub> (1:5 w/v ratio) for analysis of NH<sub>4</sub><sup>+</sup> and NO<sub>3</sub><sup>-</sup> with subsequent analysis of the extracts using a SynergyMX microtitre plate reader (Mulvaney, 1996; Miranda, Espey and Wink, 2001). Dissolved organic nitrogen (DON) was calculated as the difference between TDN and dissolved inorganic N (NH<sub>4</sub><sup>+</sup> plus NO<sub>3</sub><sup>-</sup>).

Additionally, SO<sub>4</sub><sup>2-</sup> and other major anions were determined in soil: distilled water extracts (1:5 w/v; filtered through 0.45 µm syringe filters) by ion chromatography (IC; Bradfield and Cooke, 1985). Distilled water was adopted as extract in our study because other strong extracts such as KCL, NaHCO<sub>3</sub> etc, may cause large background anion concentration, leading to inaccurate readings of sulphate concentrations by IC. Total dissolved sulphur (DOS plus SO<sub>4</sub><sup>2-</sup>) and other major cations were determined in 1:5 (w/v) soil: 0.01 M Ca(H<sub>2</sub>PO<sub>4</sub>)<sub>2</sub> extracts using inductively coupled plasma-optical emission spectroscopy (ICP-OES; Anderson

et al., 1992). Since ICP-AES measures total dissolved S, while IC measures only  $\text{SO}_4\text{-S}$ , the difference between the two methods can be taken as extractable organic S (David and Mitchell, 1985; Mitchell *et al.*, 1986). All chemical analyses were made in triplicate and the results expressed on an oven-dry soil weight basis.

### **3.2.3 Sorption of $^{35}\text{S}$ -Cys, Met and $\text{Na}_2\text{SO}_4$ in sterile grassland soil**

To determine the sorption of Cys, Met and  $\text{Na}_2\text{SO}_4$  to grassland soil's solid phase, soil samples were heat sterilised to prevent microbial activity. Briefly, 1 g of soil (top soil from site 1) was placed in pre labelled 15 ml polypropylene tubes and these tubes were capped to prevent soil water loss (six replicates per treatment). Immediately before the sorption experiment, to kill soil micro-organisms, centrifuge tubes containing soils were either heated for 80 °C for 60 minutes, or autoclaved at 121 °C for 60 minutes. Once soil is heat sterilised, 5 ml of either uniformly  $^{35}\text{S}$  labelled Cys (specific activity = 0.1 kBq  $\text{ml}^{-1}$ ), Met (specific activity = 0.3 kBq  $\text{ml}^{-1}$ ) or  $\text{Na}_2\text{SO}_4$  (specific activity = 0.4 kBq  $\text{ml}^{-1}$ ) solution were added to the soil to give five separate concentrations ranging from 0.01 to 100 mM (0.01 mM, 0.1 mM, 1 mM, 10 mM and 100 mM).

The soil samples containing sorption solutions were then shaken at 200 rpm on a flat-shaking bed for 5 minutes at room temperature ( $21 \pm 1^\circ\text{C}$ ). Immediately after shaking, 1.5 ml supernatant solution was transferred to a 1.7 ml microfuge tube and centrifuged for 5 minutes at 14000 g. 1 ml supernatant was then removed from the microfuge tubes, and the amount of  $^{35}\text{S}$  in the supernatant solution was determined by liquid scintillation counting using a Wallac 1404 scintillation counter with automated quench correction (Wallac EG&G, Milton Keynes, UK). The quantity of  $^{35}\text{S}$  adsorbed was calculated as the difference between  $^{35}\text{S}$  added initially and equilibrium  $^{35}\text{S}$  concentration in the solution.

### 3.2.4 Statistics and data analysis

All data analyses were carried out using SPSS 25.0 (IBM UK Ltd, Portsmouth, UK). Simple correlation analysis and multiple linear regression were used to empirically identify parameters influencing the DOS concentrations from soil solution. These parameters included pH, DOC, DON,  $\text{NH}_4^+$ , major cations (Mn, Fe, Zn, Na, Al etc.) in soil water. The statistical methods employed were simple Bivariate correlation analysis, stepwise multiple linear regression. All values are presented as mean  $\pm$  the standard error of the mean (SEM) ( $n = 3$ ).

## 3.3 Results and Discussion

### 3.3.1 DOS and $\text{SO}_4^{2-}$ concentration in soils along the grassland productivity gradient

The field site locations and soil property descriptions are reported in table 3.1 and 3.2. All soil samples contained water-extractable sulphate which ranged in amount from 9.7 to 139.5 mg  $\text{kg}^{-1}$  DW soil (Table 3.1). Higher concentrations of  $\text{SO}_4^{2-}$  in soil were found at high altitude, low productivity sites (site 4, 5, 6 and 7). We ascribe this to lower rates of S removal by plant uptake at higher altitudes. At all seven sites, the concentrations of  $\text{SO}_4^{2-}$  in topsoil were higher than that from subsoils (topsoil vs. subsoil;  $p < 0.05$  for sites 3, 4, 5 and 6). However, subsoils may contain large amounts of adsorbed sulphate (Schoenau and Malhi, 2008; Anderson, 2020), which may not have been extracted in our study, therefore the contribution of subsoils to plant S uptake can't be ignored.

Sulphate concentrations were consistently smaller than the total extractable S as measured by ICP-AES. The concentrations of total dissolved sulphur (DOS plus sulphate) in most soil samples were found to be less than 400 mg S  $\text{kg}^{-1}$  soil DW. In general, higher concentrations of TDS were detected from subsoils. DOS concentration accounted for 23.9 to 74.8% of TDS in topsoils, while accounting for between 58.5 to 94.8% in subsoils, this is in line with other studies on soil sulphur fractions (Tabatabai, 2015).

**Table 3.1** General site and soil properties (samples collected in late May 2019). Values represent means  $\pm$  SEM ( $n = 3$ ) for either topsoil (top) or subsoil (sub). Data is expressed on a dry soil weight basis.

	Site 1 - top	Site 2 - top	Site 3 - top	Site 4 - top	Site 5 - top	Site 6- top	Site 7 - top
Altitude (m)	23	87	134	209	277	333	404
NO <sub>3</sub> <sup>-</sup> -N (mg kg <sup>-1</sup> soil DW)	8.1 $\pm$ 0.2	7.6 $\pm$ 1.5	11.5 $\pm$ 0.2	16.6 $\pm$ 1.3	23.4 $\pm$ 0.5	33.9 $\pm$ 4.7	21.6 $\pm$ 2.3
NH <sub>4</sub> <sup>+</sup> -N (mg kg <sup>-1</sup> soil DW)	2.7 $\pm$ 0.6	15.3 $\pm$ 2.1	5.0 $\pm$ 0.3	12.0 $\pm$ 2.4	10.6 $\pm$ 1.0	20.8 $\pm$ 3.5	64.5 $\pm$ 0.7
Organic matter (mg kg <sup>-1</sup> soil DW)	7.3 $\pm$ 0.1	12.4 $\pm$ 7.4	18.7 $\pm$ 2.5	22.4 $\pm$ 0.4	20.4 $\pm$ 0.2	23.8 $\pm$ 0.5	85.9 $\pm$ 0.8
pH	6.0 $\pm$ 0.04	5.1 $\pm$ 0.02	4.6 $\pm$ 0.03	4.8 $\pm$ 0.04	5.1 $\pm$ 0.04	4.5 $\pm$ 0.12	4.1 $\pm$ 0.05
Water content (%)	19.7 $\pm$ 0.2	24.3 $\pm$ 12.5	25.3 $\pm$ 0.6	34.6 $\pm$ 0.2	38.5 $\pm$ 0.2	26.3 $\pm$ 0.6	65.2 $\pm$ 0.3
Sulphate-S (mg S kg <sup>-1</sup> soil DW)	32.8 $\pm$ 2.7	65.8 $\pm$ 30.7	60.3 $\pm$ 13.2	46.9 $\pm$ 16.5	83.4 $\pm$ 15.7	84.9 $\pm$ 5.7	137.9 $\pm$ 16.5
DOS (mg kg <sup>-1</sup> soil DW)	10.6 $\pm$ 3.7	186.0 $\pm$ 21.8	232.8 $\pm$ 66.7	196.3 $\pm$ 17.8	65.5 $\pm$ 22.7	79.3 $\pm$ 15.8	270.3 $\pm$ 17.4
DOC (mg kg <sup>-1</sup> soil DW)	108.0 $\pm$ 1.2	192.3 $\pm$ 30.6	174.1 $\pm$ 8.7	390.1 $\pm$ 8.0	226.8 $\pm$ 6.2	329.3 $\pm$ 1.2	1537.2 $\pm$ 18.6
DON (mg kg <sup>-1</sup> soil DW)	20.7 $\pm$ 0.5	38.6 $\pm$ 6.9	29.4 $\pm$ 1.0	74.4 $\pm$ 1.5	50.7 $\pm$ 1.1	65.1 $\pm$ 1.2	202.3 $\pm$ 3.8
MBC (mg kg <sup>-1</sup> soil DW)	81.8 $\pm$ 1.0	158.7 $\pm$ 25.5	143.5 $\pm$ 8.8	349.6 $\pm$ 7.8	190.0 $\pm$ 4.6	286.1 $\pm$ 2.1	1348.1 $\pm$ 20.7
MBN (mg kg <sup>-1</sup> soil DW)	10.1 $\pm$ 0.1	20.8 $\pm$ 3.7	16.7 $\pm$ 0.9	51.5 $\pm$ 0.9	22.8 $\pm$ 1.5	40.1 $\pm$ 0.4	148.7 $\pm$ 4.7

	Site 1 - sub	Site 2 - sub	Site 3 - sub	Site 4 - sub	Site 5 - sub	Site 6- sub	Site 7 - sub
NO <sub>3</sub> <sup>-</sup> -N (mg kg <sup>-1</sup> soil DW)	6.2 $\pm$ 0.1	7.7 $\pm$ 0.2	10.2 $\pm$ 0.8	10.5 $\pm$ 0.2	12.5 $\pm$ 0.9	19.0 $\pm$ 0.5	44.5 $\pm$ 2.9
NH <sub>4</sub> <sup>+</sup> -N (mg kg <sup>-1</sup> soil DW)	2.3 $\pm$ 0.6	7.6 $\pm$ 0.3	4.4 $\pm$ 0.4	4.7 $\pm$ 0.3	7.9 $\pm$ 1.6	6.5 $\pm$ 0.3	23.5 $\pm$ 2.2
Organic matter (mg kg <sup>-1</sup> soil DW)	6.9 $\pm$ 0.2	10.8 $\pm$ 0.8	12.8 $\pm$ 0.3	14.7 $\pm$ 0.4	14.5 $\pm$ 0.5	14.9 $\pm$ 0.2	66.2 $\pm$ 0.5
pH	6.4 $\pm$ 0.03	5.2 $\pm$ 0.04	4.6 $\pm$ 0.02	4.9 $\pm$ 0.02	5.2 $\pm$ 0.02	4.6 $\pm$ 0.02	3.9 $\pm$ 0.02
Water content (%)	17.8 $\pm$ 0.1	25.8 $\pm$ 1.2	30.4 $\pm$ 0.5	32.2 $\pm$ 0.3	32.6 $\pm$ 0.6	29.9 $\pm$ 0.2	65.3 $\pm$ 0.1
Sulphate-S (mg S kg <sup>-1</sup> soil DW)	9.7 $\pm$ 2.1	18.2 $\pm$ 9.6	17.0 $\pm$ 2.9	12.1 $\pm$ 3.3	48.5 $\pm$ 7.8	47.2 $\pm$ 4.4	139.5 $\pm$ 25.3
DOS (mg kg <sup>-1</sup> soil DW)	41.2 $\pm$ 8.1	120.4 $\pm$ 16.9	317.5 $\pm$ 24.3	350.0 $\pm$ 241.0	87.1 $\pm$ 5.6	97.9 $\pm$ 6.9	209.5 $\pm$ 85.9
DOC (mg kg <sup>-1</sup> soil DW)	96.5 $\pm$ 4.4	140.1 $\pm$ 1.9	127.4 $\pm$ 2.2	189.5 $\pm$ 3.5	160.1 $\pm$ 0.7	177.7 $\pm$ 0.4	1241.7 $\pm$ 23.5
DON (mg kg <sup>-1</sup> soil DW)	15.7 $\pm$ 0.4	25.6 $\pm$ 0.4	20.8 $\pm$ 0.1	32.3 $\pm$ 0.4	28.9 $\pm$ 1.0	34.6 $\pm$ 1.1	149.0 $\pm$ 3.5
MBC (mg kg <sup>-1</sup> soil DW)	69.8 $\pm$ 4.2	111.8 $\pm$ 2.7	98.5 $\pm$ 0.6	158.8 $\pm$ 3.3	128.0 $\pm$ 0.8	147.3 $\pm$ 1.5	1073.7 $\pm$ 27.3
MBN (mg kg <sup>-1</sup> soil DW)	9.4 $\pm$ 0.2	14.9 $\pm$ 0.5	12.4 $\pm$ 0.5	20.7 $\pm$ 0.3	15.0 $\pm$ 1.0	18.3 $\pm$ 0.9	99.4 $\pm$ 4.5

**Table 3.2** Major soil cation concentrations measured by ICP-OES (samples collected in late May 2019). Values represent means  $\pm$  SEM ( $n = 3$ ) for either topsoil (top) or subsoil (sub). Data is expressed on a dry soil weight basis. NA indicates not applicable.

	Site 1 - top	Site 2 - top	Site 3 - top	Site 4 - top	Site 5 - top	Site 6- top	Site 7 - top
Al (mg kg <sup>-1</sup> soil DW)	8.6 $\pm$ 1.2	67.6 $\pm$ 29.4	120.9 $\pm$ 28.5	114.7 $\pm$ 24.8	56.3 $\pm$ 12.9	111.8 $\pm$ 2.2	518.4 $\pm$ 3.1
B (mg kg <sup>-1</sup> soil DW)	2.6 $\pm$ 0.3	3.6 $\pm$ 1.3	2.8 $\pm$ 0.8	7.3 $\pm$ 0.7	6.6 $\pm$ 3.5	3.3 $\pm$ 0.5	10.2 $\pm$ 9.1
Ba (mg kg <sup>-1</sup> soil DW)	4.1 $\pm$ 0.2	1.5 $\pm$ 0.7	0.9 $\pm$ 0.3	3.6 $\pm$ 0.9	0.9 $\pm$ 0.2	3.6 $\pm$ 0.02	28.5 $\pm$ 0.4
Co (mg kg <sup>-1</sup> soil DW)	0.1 $\pm$ 0.01	0.4 $\pm$ 0.06	0.5 $\pm$ 0.03	0.2 $\pm$ 0.04	0.1 $\pm$ 0.04	0.3 $\pm$ 0.02	0.7 $\pm$ 0.04
Cr (mg kg <sup>-1</sup> soil DW)	0.6 $\pm$ 0.03	0.6 $\pm$ 0.11	0.7 $\pm$ 0.02	0.8 $\pm$ 0.01	1.0 $\pm$ 0.02	0.7 $\pm$ 0.02	1.6 $\pm$ 0.01
Fe (mg kg <sup>-1</sup> soil DW)	2.3 $\pm$ 0.7	20.4 $\pm$ 6.7	14.9 $\pm$ 1.3	37.7 $\pm$ 5.8	21.2 $\pm$ 7.0	33.0 $\pm$ 0.2	68.2 $\pm$ 1.8
K (g kg <sup>-1</sup> soil DW)	1.0 $\pm$ 0.03	1.2 $\pm$ 0.19	0.2 $\pm$ 0.01	1.4 $\pm$ 0.03	0.8 $\pm$ 0.02	1.5 $\pm$ 0.03	6.9 $\pm$ 0.1
Mg (g kg <sup>-1</sup> soil DW)	0.6 $\pm$ 0.01	0.9 $\pm$ 0.14	0.3 $\pm$ 0.01	1.3 $\pm$ 0.01	1.1 $\pm$ 0.01	1.1 $\pm$ 0.01	2.5 $\pm$ 0.02
Mn (mg kg <sup>-1</sup> soil DW)	47.5 $\pm$ 1.0	217.9 $\pm$ 34.3	34.1 $\pm$ 0.9	253.5 $\pm$ 6.0	110.8 $\pm$ 3.7	350.6 $\pm$ 8.6	691.3 $\pm$ 3.5
Na (g kg <sup>-1</sup> soil DW)	0.2 $\pm$ 0.01	0.9 $\pm$ 0.14	0.3 $\pm$ 0.01	0.4 $\pm$ 0.01	0.5 $\pm$ 0.01	0.5 $\pm$ 0.01	1.5 $\pm$ 0.02
Si (mg kg <sup>-1</sup> soil DW)	43.2 $\pm$ 0.3	37.8 $\pm$ 4.5	46.6 $\pm$ 1.0	58.8 $\pm$ 4.5	57.3 $\pm$ 6.2	55.3 $\pm$ 2.1	111.5 $\pm$ 1.2
Sr (mg kg <sup>-1</sup> soil DW)	38.4 $\pm$ 0.2	17.4 $\pm$ 2.9	10.3 $\pm$ 0.1	17.9 $\pm$ 0.4	17.5 $\pm$ 0.1	16.0 $\pm$ 0.1	48.9 $\pm$ 0.3
Zn (mg kg <sup>-1</sup> soil DW)	1.0 $\pm$ 0.1	10.1 $\pm$ 2.6	2.7 $\pm$ 0.1	17.4 $\pm$ 1.5	8.9 $\pm$ 0.4	17.8 $\pm$ 0.5	46.3 $\pm$ 0.3
	Site 1 - sub	Site 2 - sub	Site 3 - sub	Site 4 - sub	Site 5 - sub	Site 6- sub	Site 7 - sub
Al (mg kg <sup>-1</sup> soil DW)	6.8 $\pm$ 1.9	40.7 $\pm$ 7.5	134.5 $\pm$ 16.2	66.4 $\pm$ 43.7	40.2 $\pm$ 2.7	134.0 $\pm$ 0.6	441.7 $\pm$ 7.0
B (mg kg <sup>-1</sup> soil DW)	3.3 $\pm$ 1.3	5.3 $\pm$ 2.5	2.6 $\pm$ 0.1	5.5 $\pm$ 1.9	4.0 $\pm$ 2.1	3.3 $\pm$ 1.6	4.1 $\pm$ 1.1
Ba (mg kg <sup>-1</sup> soil DW)	4.0 $\pm$ 0.3	0.7 $\pm$ 0.2	1.6 $\pm$ 0.9	1.1 $\pm$ 0.9	0.3 $\pm$ 0.03	2.3 $\pm$ 0.01	33.5 $\pm$ 0.6
Co (mg kg <sup>-1</sup> soil DW)	NA	0.2 $\pm$ 0.01	0.5 $\pm$ 0.03	0.2 $\pm$ 0.02	0.1 $\pm$ 0.01	0.3 $\pm$ 0.02	0.7 $\pm$ 0.06
Cr (mg kg <sup>-1</sup> soil DW)	0.6 $\pm$ 0.01	0.6 $\pm$ 0.02	0.7 $\pm$ 0.04	0.7 $\pm$ 0.07	0.9 $\pm$ 0.03	0.7 $\pm$ 0.03	1.6 $\pm$ 0.03
Fe (mg kg <sup>-1</sup> soil DW)	1.8 $\pm$ 0.9	7.5 $\pm$ 1.4	23.1 $\pm$ 6.4	13.9 $\pm$ 9.8	8.6 $\pm$ 0.4	20.3 $\pm$ 1.8	69.3 $\pm$ 2.4
K (g kg <sup>-1</sup> soil DW)	1.1 $\pm$ 0.02	0.5 $\pm$ 0.01	0.5 $\pm$ 0.40	0.6 $\pm$ 0.03	0.4 $\pm$ 0.01	0.8 $\pm$ 0.01	5.5 $\pm$ 0.1
Mg (g kg <sup>-1</sup> soil DW)	0.5 $\pm$ 0.01	0.7 $\pm$ 0.01	0.4 $\pm$ 0.12	0.7 $\pm$ 0.01	0.7 $\pm$ 0.01	0.6 $\pm$ 0.01	2.3 $\pm$ 0.01
Mn (mg kg <sup>-1</sup> soil DW)	18.8 $\pm$ 0.5	76.7 $\pm$ 2.3	64.6 $\pm$ 35.8	87.1 $\pm$ 1.0	53.9 $\pm$ 1.2	97.5 $\pm$ 0.9	1123.0 $\pm$ 3.4
Na (g kg <sup>-1</sup> soil DW)	0.2 $\pm$ 0.01	0.4 $\pm$ 0.01	0.4 $\pm$ 0.12	0.4 $\pm$ 0.01	0.5 $\pm$ 0.01	0.3 $\pm$ 0.03	1.3 $\pm$ 0.02
Si (mg kg <sup>-1</sup> soil DW)	49.5 $\pm$ 0.9	32.6 $\pm$ 1.2	51.0 $\pm$ 4.0	51.6 $\pm$ 2.1	37.2 $\pm$ 1.8	39.9 $\pm$ 3.1	78.6 $\pm$ 2.4
Sr (mg kg <sup>-1</sup> soil DW)	35.9 $\pm$ 0.3	15.4 $\pm$ 0.4	12.3 $\pm$ 2.0	11.8 $\pm$ 0.3	12.7 $\pm$ 0.1	11.9 $\pm$ 0.1	50.9 $\pm$ 0.1
Zn (mg kg <sup>-1</sup> soil DW)	0.6 $\pm$ 0.1	5.4 $\pm$ 0.1	4.5 $\pm$ 1.9	5.8 $\pm$ 0.5	4.3 $\pm$ 0.4	8.4 $\pm$ 0.2	53.4 $\pm$ 0.8

### 3.3.2 Factors influencing DOS concentration in grassland soils

Concentrations of the major cations in soil solution were summarized in Table 3.2. Stepwise linear regressions were employed to identify parameters which may help explain the spatial variation in DOS concentrations across the seven selected grassland sites. In this method, DOS was selected as the dependent variable, and TDS, Al, B, Ba, Fe, K, Mn, Na, Sr, Zn,  $\text{NH}_4^+$  content from soil water, and soil organic matter, pH, DOC, DON, were selected as possible independent variables. According to our linear regression analysis, TDS, DON along with pH were found to be the significant factors in the model, together they are able to explain over 95% of the spatial variation in DOS in soil solution (Table 3.3).

The data were then analysed using Bivariate correlation, this analysis results confirm that TDS, DON concentration and pH in soil plays a significant role in governing the DOS concentration. The analysis results were summarized in table 3.4, results showed that Al, B, Ba, Fe, K, Mn,  $\text{NH}_4^+$ , DOC and soil organic matter also co-correlate with DOS concentration in soil solution, while Na, Sr and Zn correlated less well (Table 3.4). Al and Fe concentrations are positively correlated to DOS in the multiple regression, this may be due to that they are inherently and positively correlated to  $\text{SO}_4^{2-}$ . In contrast, soil pH is negatively correlated to DOS content. Soil pH affects the microbial population and their activities, therefore, a pH increase would lead to higher DOS mineralization (Neina, 2019).

Previous studies indicated that the major organic sulphur constituents in soil solution are carbon-bonded sulphur, including some amino acids, and ester sulphates such as choline sulphates and sulphated thioglycides (Zhao *et al.*, 2006). Hence, there is often a close correspondence between organic C and organic S in soils since carbon-bonded sulphur is generally the dominant organic sulphur (Homann *et al.*, 1990; Houle *et al.*, 2011). In our study, a strong relationship between DOS and DOC was also found, indicating that soil organic

sulphur is conceptually an explanatory parameter for DOS as both parameters are inherently related to the amount of DOM.

**Table 3.3** Stepwise multiple linear regressions results.


Model	Predictor in model	B	SE	R	Adjusted R <sup>2</sup>	R <sup>2</sup> change
1	(Constant)	-13.41	11.55	0.92	0.85	0.85
	TDS	0.78	0.05			
2	(Constant)	-20.46	7.34	0.97	0.94	0.94
	TDS	1.02	0.05			
	DON	-0.71	0.09			
3	(Constant)	-176.31	45.28	0.98	0.96	0.95
	TDS	1.13	0.05			
	DON	-.69	0.08			
	pH	26.64	7.66			

Predictor in model: variables selected into the model; B: partial regression coefficient; SE: standard errors.

### 3.3.3 Sorption of <sup>35</sup>S-Cys, Met and Na<sub>2</sub>SO<sub>4</sub> in sterile grassland soil

Previous studies indicated that most UK soils have limited capacity to adsorb SO<sub>4</sub>-S (Curtin and Syers, 1990; Zhao and McGrath, 1994), this is also found in our study, according to our results (Fig. 3.1), less than 4% of <sup>35</sup>S-Na<sub>2</sub>SO<sub>4</sub> was adsorbed by our selected grassland soil, this in line with previous studies. Although the SO<sub>4</sub>-S sorption capacity of soils could vary widely depending on factors such as the pH, Fe, Ca and Al content, clay minerals, soil organic matter content, temperature, water content etc. (Tabatabai, 1987; Sokolova and Alekseeva, 2008). In general, the soil sorption capacities of <sup>35</sup>S-Cys and Met were slightly higher than that of inorganic sulphate, with 5-10 % of amino acid-<sup>35</sup>S lost during our experiment. Due to the large solution to soil ratio and rapid experimentation, we assume that microorganisms played little, if any, part in our sorption experiment.





**Fig. 3.1** Sorption of Cys, Met and  $\text{Na}_2\text{SO}_4$  to the soil's solid phase, panel A shows sorption onto autoclaved soils, panel B shows sorption onto oven heated soils. The legend is the same for both panels. Values represent means  $\pm$  SEM ( $n = 6$ ).

**Table 3.4** Correlation matrixes between DOS concentration and cations, ammonium, and MBC. \*\* and \* represents significance at level  $p < 0.01$  and  $p < 0.05$ , respectively.

	DOS	TDS	Al	B	Ba	Fe	K	Mn	Na	Sr	Zn	NH <sub>4</sub> <sup>+</sup>	Organic matter	pH	DOC	DON
DOS	1	0.92 **	0.68 **	-0.49 **	0.35 *	0.57 **	-0.56 **	0.41 **	0.31 *	-0.36 *	0.32 *	0.41 **	0.47 **	-0.63 **	0.44 **	0.41 **
TDS		1	0.86 **	-0.66 **	0.59 **	0.74 **	-0.60 **	0.66 **	0.43 **	-0.3	0.59 **	0.65 **	0.73 **	-0.78 **	0.70 **	0.68 **
Al			1	-0.65 **	0.78 **	0.90 **	-0.48 **	0.82 **	0.50 **	-0.2	0.74 **	0.74 **	0.87 **	-0.86 **	0.82 **	0.78 **
B				1	-0.46 **	-0.48 **	0.72 **	-0.52 **	-0.3	0.3	-0.45 **	-0.60 **	-0.62 **	0.58 **	-0.64 **	-0.66 **
Ba					1	0.69 **	-0.02	0.95 **	0.45 **	0.44 **	0.82 **	0.77 **	0.92 **	-0.49 **	0.95 **	0.90 **
Fe						1	-0.36 *	0.77 **	0.69 **	-0.2	0.76 **	0.66 **	0.79 **	-0.84 **	0.72 **	0.68 **
K							1	-0.2	-0.1	0.75 **	-0.2	-0.23	-0.29 **	0.68 **	-0.19	-0.02
Mn								1	0.53 **	0.2	0.89 **	0.68 **	0.91 **	-0.64 **	0.92 **	0.87 **
Na									1	-0.1	0.59 **	0.53 **	0.53 **	-0.46 **	0.55 **	0.54 **
Sr										1	0.05	0.12	0.15	0.53 **	0.27	0.22
Zn											1	0.76 **	0.87 **	-0.72 **	0.85 **	0.85 **
NH <sub>4</sub> <sup>+</sup>												1	0.90 **	-0.55 **	0.89 **	0.93 **
Organic Matter													1	-0.68 **	0.96 **	0.95 **
pH														1	-0.58 **	-0.58 **
DOC															1	0.98 **
DON																1

### 3.4 Conclusions

DOS and  $\text{SO}_4^{2-}$  concentration in seven grassland soils in North Wales were studied. We found that the DOS concentration in all soils ranged from 10.6 to 270.3  $\text{mg kg}^{-1}$  DW soil from topsoils, while it ranged from 41.2 to 309.3  $\text{mg kg}^{-1}$  DW soil in subsoils; concentrations of  $\text{SO}_4^{2-}$  in all seven sites ranged from 46.9 to 137.9  $\text{mg kg}^{-1}$  DW soil in topsoil, while ranged from 9.7 to 139.5  $\text{mg kg}^{-1}$  DW soil in subsoil. The concentrations of total dissolved sulphur (DOS plus  $\text{SO}_4^{2-}$ ) in most of grassland soils were found to be  $< 350 \text{ mg kg}^{-1}$  DW soil (Table 3.1). DOS concentrations were found to be higher in highland grassland soil compared to lowland soils. The DOS constituted 23.8 to 93.8% of total dissolved sulphur (DOS plus  $\text{SO}_4^{2-}$ ) in all soils; this may be because our sites currently receive little atmospheric S deposition and are low in extractable  $\text{SO}_4^{2-}$ , therefore soil S stored is primarily in the organic form (Table 3.1). Soluble organic S seems to be intimately related with soluble carbon, suggesting the production of dissolved organic matter and mineralisation of  $\text{SO}_4^{2-}$  may be regulated by the same factors.

### 3.5 Acknowledgements

I would like to thank Joseph Cotton and Mark Hughes for the help with soil sample collection, Sarah Chesworth and Jonathan Roberts for their technical assistance in the laboratory. This research was funded by Bangor University and the China Scholarship Council (201606510012).

### 3.6 References

- Amon, R. M. W., Benner, R. and Saunders, D. O. M. (2013) 'Bacterial utilization of different size classes of dissolved matter organic', *Limnology and Oceanography*, 41(1), pp. 41–51.
- Anderson, G. C. (2020) 'Sulfate sorption measured by a buffering index over a range of

- properties of soils from south Western Australia', *Soil Research*, 58(7), pp. 651–661.  
doi: 10.1071/SR20005.
- Anderson, G. L. *et al.* (1992) 'Soil sulphur testing', *Sulphur in agriculture*, pp. 6–14.
- Bronk, D. A., Gilbert, P. M. and Ward, B. B. (2007) 'Nitrogen Uptake , Dissolved Organic Nitrogen', 265(5180), pp. 1843–1846.
- Curtin, D. and Syers, J. K. (1990) 'Extractability and adsorption of sulphate in soils', *Journal of soil science*. Wiley Online Library, 41(2), pp. 305–312.
- David, M. B. and Mitchell, M. J. (1985) 'Sulfur constituents and cycling in waters, seston, and sediments of an oligotrophic lake', *Limnology and Oceanography*, 30(6), pp. 1196–1207. doi: 10.4319/lo.1985.30.6.1196.
- Edgell, K. W., Longbottom, J. E. and Pfaff, J. D. (1994) 'Determination of inorganic anions in water by ion chromatography: a collaborative study.', *Journal of AOAC International*. Royal Society of Chemistry, 77(5), pp. 1253–1263. doi: 10.1093/jaoac/77.5.1253.
- Goller, R. *et al.* (2006) 'Dissolved nitrogen, phosphorus, and sulfur forms in the ecosystem fluxes of a montane forest in Ecuador', *Biogeochemistry*, 77(1), pp. 57–89. doi: 10.1007/s10533-005-1061-1.
- He, W. *et al.* (2016) 'Dynamic exchanges between DOM and POM pools in coastal and inland aquatic ecosystems: A review', *Science of the Total Environment*. Elsevier, pp. 415–428. doi: 10.1016/j.scitotenv.2016.02.031.
- Homann, P. S. *et al.* (1990) 'Organic sulfur in throughfall, stem flow, and soil solutions from temperate forests', *Canadian Journal of Forest Research*. NRC Research Press, 20(9), pp. 1535–1539. doi: 10.1139/x90-204.
- Houle, D. *et al.* (2011) 'in southwestern Quebec lakes : carbon and sulfur Dissolved organic

- and lake properties with catchment Relationships’, *Limnology*, 40(4), pp. 710–717.
- Jones, D. L. *et al.* (2004) ‘Role of dissolved organic nitrogen (DON) in soil N cycling in grassland soils’, *Soil Biology and Biochemistry*, 36(5), pp. 749–756. doi: 10.1016/j.soilbio.2004.01.003.
- Kaiser, K. and Guggenberger, G. (2005) ‘Dissolved organic sulphur in soil water under *Pinus sylvestris* L. and *Fagus sylvatica* L. stands in northeastern Bavaria, Germany - Variations with seasons and soil depth’, *Biogeochemistry*, 72(3), pp. 337–364. doi: 10.1007/s10533-004-0155-5.
- McDowell, R. W. and Koopmans, G. F. (2006) ‘Assessing the bioavailability of dissolved organic phosphorus in pasture and cultivated soils treated with different rates of nitrogen fertiliser’, *Soil Biology and Biochemistry*, 38(1), pp. 61–70. doi: 10.1016/j.soilbio.2005.03.026.
- Miranda, K. M., Espey, M. G. and Wink, D. A. (2001) ‘A rapid, simple spectrophotometric method for simultaneous detection of nitrate and nitrite’, *Nitric Oxide - Biology and Chemistry*, 5(1), pp. 62–71. doi: 10.1006/niox.2000.0319.
- Mitchell, M. J. *et al.* (1986) ‘Sulfur constituents in soils and streams of a watershed in the Rocky Mountains of Alberta’, *Canadian Journal of Forest Research*. NRC Research Press, 16(2), pp. 315–320. doi: 10.1139/x86-053.
- Mulvaney, R. L. (1996) ‘Nitrogen—inorganic forms’, *Methods of soil analysis: Part 3 Chemical methods*. Wiley Online Library, 5, pp. 1123–1184. doi: 10.2136/sssabookser5.3.c38.
- Neff, J. C., III, F. S. C. and Vitousek, P. M. (2007) ‘Breaks in the Cycle: Dissolved Organic Nitrogen in Terrestrial Ecosystems’, *Frontiers in Ecology and the Environment*, 1(4),

p. 205. doi: 10.2307/3868065.

Neina, D. (2019) 'The Role of Soil pH in Plant Nutrition and Soil Remediation', *Applied and Environmental Soil Science*, 2019(3). doi: 10.1155/2019/5794869.

Peuravuori, J. *et al.* (2005) 'Comparative study for separation of aquatic humic-type organic constituents by DAX-8, PVP and DEAE sorbing solids and tangential ultrafiltration: Elemental composition, size-exclusion chromatography, UV-vis and FT-IR', *Talanta*, 65(2 SPEC. ISS.), pp. 408–422. doi: 10.1016/j.talanta.2004.06.042.

Rowland, A. P. and Haygarth, P. M. (1997) 'Determination of Total Dissolved Phosphorus in Soil Solutions', *Journal of Environmental Quality*. Madison, WI: American Society of Agronomy, Crop Science Society of America, and Soil Science Society of America, 26, pp. 410–415. doi: 10.2134/jeq1997.00472425002600020011x.

Schoenau, J. J. and Malhi, S. S. (2008) 'Sulfur forms and cycling processes in soil and their relationship to sulfur fertility', *Sulfur: A missing link between soils, crops, and nutrition*. Wiley Online Library, 50, pp. 1–10.

Sokolova, T. A. and Alekseeva, S. A. (2008) 'Adsorption of sulfate ions by soils (a review)', *Eurasian Soil Science*, 41(2), pp. 140–148. doi: 10.1134/s106422930802004x.

Stedmon, C. A., Markager, S. and Bro, R. (2003) 'Tracing dissolved organic matter in aquatic environments using a new approach to fluorescence spectroscopy', *Marine Chemistry*, 82(3–4), pp. 239–254. doi: 10.1016/S0304-4203(03)00072-0.

Tabatabai, M. A. (1987) 'Physicochemical fate of sulfate in soils', *Journal of the Air Pollution Control Association*, 37(1), pp. 34–38. doi: 10.1080/08940630.1987.10466197.

Tabatabai, M. A. (2015) 'Sulfur in Agriculture', *Sulfur in Agriculture*, (1), pp. 1–652. doi: 10.2134/agronmonogr27.

- Voroney, R. P., Brookes, P. C. and Beyaert, R. . (2007) 'Soil Microbial Biomass C,N,P and S', in *Soil Sampling and Methods of Analysis, Second Edition*. CRC Press Boca Raton, FL, pp. 637–652. doi: 10.1201/9781420005271.ch49.
- Wang, Zhanyi *et al.* (2012) 'Dissolved organic sulfur in streams draining forested catchments in southern China', *Journal of Environmental Sciences*, 24(4), pp. 704–710. doi: 10.1016/S1001-0742(11)60781-X.
- Wieder, R. K. and Lang, G. E. (1988) 'Cycling of inorganic and organic sulfur in peat from Big Run Bog, West Virginia', *Biogeochemistry*, 5(2), pp. 221–242. doi: 10.1007/BF02180229.
- Wu, J. *et al.* (1990) 'Measurement of soil microbial biomass C by fumigation-extraction-an automated procedure.', *Soil Biology & Biochemistry*, 22(8), pp. 1167–1169. doi: 10.1016/0038-0717(90)90046-3.
- Zhao, F. J. *et al.* (2006) 'Sulphur speciation and turnover in soils: evidence from sulphur K-edge XANES spectroscopy and isotope dilution studies', *Soil Biology and Biochemistry*, 38(5), pp. 1000–1007. doi: 10.1016/j.soilbio.2005.08.013.
- Zhao, F. and McGrath, S. P. (1994) 'Extractable sulphate and organic sulphur in soils and their availability to plants', *Plant and Soil*, 164(2), pp. 243–250. doi: 10.1007/BF00010076.
- Zsolnay, Á. (2003) 'Dissolved organic matter: Artefacts, definitions, and functions', *Geoderma*, 113(3–4), pp. 187–209. doi: 10.1016/S0016-7061(02)00361-0.

## **Chapter 4**

---

### **Rapid microbial uptake and mineralization of cysteine and methionine along a grassland productivity gradient**



## Abstract

Cysteine (Cys) and methionine (Met) constitute important sources of directly available carbon (C), nitrogen (N), and sulphur (S) for both plant nutrition and microbial growth and are central to terrestrial S cycling. The factors controlling turnover rates of these S-amino acids in soil, however, remains understudied. Due to their multi-nutrient composition (C, N and S) we hypothesized that intense competition would exist in the microbial community for these substrates. Further, we hypothesized that this competition would be inversely related to plant productivity and thus C inputs to soil. To address this, we studied the short-term, concentration-dependent microbial uptake and mineralization of  $^{14}\text{C}$ -labelled Cys and Met in five grassland soils taken from an altitude-driven primary productivity gradient. Substrate depletion from soil solution was measured over a 60-minute period using a centrifugal drainage procedure and the subsequent evolution of  $^{14}\text{CO}_2$  measured over a 48 h-period. Our results revealed that  $^{14}\text{C}$ -Cys and Met are directly and rapidly assimilated by soil microbes, with half-times ranging from 0.34 - 2.14 mins. This is an order of magnitude (or more) faster than when determined from measurement of  $^{14}\text{CO}_2$  evolution. This considerable delay between microbial  $^{14}\text{C}$  removal from soil solution and subsequent  $^{14}\text{CO}_2$  evolution indicates that the degradation of Cys and Met in soil solution was mainly through biological processes. The use of different substrate concentrations suggested that when lower concentrations of  $^{14}\text{C}$ -Cys and Met were added to soil (0.01 - 0.1 mM), soil microbial uptake occurred via a single carrier system, while a second transport system/mechanism became important when the substrates were added at higher concentrations (1 - 100 mM). Rates of substrate mineralization by soil microbes declined in less productive, nutrient limited grassland soils with lower levels of microbial biomass, suggesting that the turnover of organic N and S, and subsequent availability for plant uptake is likely to be controlled by the size and activity of the soil microbial community. We conclude that despite the relatively low concentration of Cys and Met in soil solution, these DON/DOS compounds could be of great significance in terrestrial nitrogen and sulphur cycling due to their rapid microbial turnover.

**Keywords:** Cysteine, methionine, biodegradation, mineralization, grassland productivity gradient, free amino acid

## 4.1 Introduction

Recent decades has seen a major reduction in global anthropogenic SO<sub>2</sub> emissions due to a decrease in fossil fuel burning and greater regulatory controls on sulphur (S) release to the atmosphere (Fowler et al., 2007; Hinckley et al., 2020). In combination with the adoption of low S-containing fertilizers and greater crop yields (Ceccotti and Messick, 1997), this has resulted in a negative S balance in many agricultural systems (McGrath and Zhao, 1995; Hioang et al, 2020; Pariasca-Tanaka et al., 2020). As a consequence, these agroecosystems are becoming more reliant on the supply of S from native soil organic matter (SOM). In comparison to other macronutrients (e.g. N, P), however, relatively few studies have focused on the mineralization of organic S to SO<sub>4</sub><sup>2-</sup> and its capacity to alleviate S deficiency in plants (Fitzgerald et al., 1988; Fitzgerald and Andrew, 1984; Chalk and Inacio, 2017).

Typically, more than 95% of the S present in soil is held in a solid organic form. Most of this S enters soil from plant residues and therefore includes an abundance of free and combined S-containing amino acids (i.e. cysteine and methionine; Cys and Met) in addition to a range of other S compounds such as thiols (e.g. glutathione), sulfolipids and secondary S compounds (e.g. allicins, glucosinolates, phytochelatins; Falloon and Smith, 2000). Unsurprisingly therefore, studies have shown the presence of large quantities of S-containing amino acids in hydrolysates of SOM (Freney, Stevenson and Beavers, 1972; Scott, Bick and Anderson, 1981).

In contrast to SOM, several studies have reported that the concentration of free S-containing amino acids in soil solution is very low (ca. 0.1 to 10 µM; Broughton et al., 2015), particularly in relation to the levels of SO<sub>4</sub><sup>2-</sup> (ca. 50 - 1000 µM; Cambier et al., 2014; Kölling and Prietzel, 1995). This may suggest that inorganic S dominates plant and microbial uptake. However, studies in freshwaters suggest that this assumption may not always be valid, especially when both pool size and flux rate are considered together (Brailsford *et al.*, 2020). Their low concentration in soil solution could be explained by (i) their low rate of production,

(ii) their rapid removal by roots and microorganisms, (iii) their external breakdown by free sulfatases, or (iv) sorption to mineral surfaces (Broughton et al., 2015; Ma et al., 2020; Sanchez-Arenillas et al., 2017). The relative importance and complementarity of these mechanisms in regulating free Cys and Met levels in soil, however, remains unknown. Cys and Met can be considered multi-nutritional as they contain C, N and S. It might be expected therefore that C- or N-limited microbial communities would prefer these substrates to  $\text{SO}_4^{2-}$ . From a thermodynamic perspective, it is also energetically more efficient to take up amino acids which can be directly incorporated into common metabolic pathways than to take up and assimilate  $\text{SO}_4^{2-}$  (Kertesz, 2000). We hypothesize that S mineralization from C-bonded organic forms (e.g. Cys and Met), occurs mainly through biological processes (McGill and Cole, 1981; Jones and Shannon, 1999), driven by microbial demand for energy and C skeletons for maintenance and growth, with excess  $\text{SO}_4^{2-}$  and  $\text{NH}_4^+$  being released as by-products (Ruiz-Herrera and Starkey, 1969a; Sohn and Ho, 1995; Zhao, Wu and McGrath, 1996). These processes may be regulated by many physical, chemical and biological factors which would ultimately determine the quality and timing of inorganic N and S becoming available for plant growth (Zehr, Axler and Goldman, 1985; Grant, Juma and McGill, 1993).

The purpose of this study was therefore to evaluate the uptake and subsequent mineralization of Cys and Met in a contrasting range of grassland soils. We chose Cys and Met as they are likely to represent the dominant forms of organic S entering grassland soils (Scott, Bick and Anderson, 1981). The specific aims of this study were therefore to: (1) determine the effects of substrate concentration on the mineralization of Cys and Met; (2) compare the turnover rates of Cys and Met under differing pasture productivity conditions; (3) ascertain whether the microbial uptake and mineralization of Cys and Met were temporally uncoupled in soil, and (4) determine if Met and Cys had different microbial C use efficiencies.

## 4.2 Materials and methods

### 4.2.1 Soil sampling and characteristics

Five soils were sampled along an altitudinal gradient on a north facing slope at the Henfaes Research Station, Abergwyngregyn, Gwynedd, North Wales, UK (53°14'N; 4°10'W; Table 4.1). They were not chosen to be intrinsically linked, rather just to provide a diverse array of soils with inherent differences in fertility and grassland productivity. This elevation gradient is characterised by a reduction in soil fertility and grassland productivity with increased elevation (Farrell, Hill, Farrar, *et al.*, 2011). Further characteristics of the altitudinal catena sequence are described in Wilkinson *et al.* (2014a), Greenfield *et al.* (2020) and Withers *et al.* (2020). The mean annual temperature ranged from 11 °C at low altitude (Site 1) to 8 °C at the top of the gradient (Site 5), with annual rainfall ranging from 960 mm at Site 1 to 1800 mm at Site 5. Five sites with contrasting soil characteristics were selected along the gradient. More soil characteristics are provided in Table 4.1. At each site, 2 kg of topsoil (0 - 10 cm) was collected from three independent randomly positioned replicate plots (5 × 5 m<sup>2</sup>; Withers *et al.*, 2020). The soil was placed in gas-permeable plastic bags and transported immediately back to the laboratory where it was sieved (2 mm) and stored at 4 °C (< 48 h) until required for further experimentation.

Moisture content was determined by oven drying (80 °C, 24 h). Soil pH and electrical conductivity (EC) were determined in a 1:5 (w/v) soil: distilled water suspensions with standard electrodes. Soil bulk density was determined by taking intact cores (100 cm<sup>3</sup>) from the field followed by oven drying and stone correction (Rowell, 2014). Available NH<sub>4</sub><sup>+</sup> and NO<sub>3</sub><sup>-</sup> were determined by extracting 5 g of field-moist soil with 25 ml 0.5 M K<sub>2</sub>SO<sub>4</sub> (200 rev min<sup>-1</sup>; 20°C, 30 min), filtering the suspensions (Whatman No. 42 filter paper) before colorimetric analysis using a Synergy-MX microplate reader (BioTek Ltd, Swindon, UK). Nitrate was determined according to the vanadate procedure of Miranda *et al.* (2001) while ammonium was determined

using the salicylic acid procedure of Mulvaney (1996). Extractable phosphorus (P) was extracted using a 0.5 M acetic acid (1:5 w/v) shaken for 1 hour (200 rev min<sup>-1</sup>; 20°C), then centrifuged for 10 min at 3220 g before passing through a Whatman 42 filter (Quevauviller, 2007). P was then analysed by the colorimetric method of Murphy and Riley (1962). Total free amino acids were determined in the 0.5 M K<sub>2</sub>SO<sub>4</sub> extracts according to Jones et al. (2002). Microbial biomass C (MBC) and N (MBN) were calculated from the differences between C and N concentrations in CHCl<sub>3</sub>-fumigated and unfumigated samples after extracting the soil with 0.5 M K<sub>2</sub>SO<sub>4</sub> as described above and analysis of extractable C and N using a TOC-V-TN analyser (Shimadzu Corp., Kyoto, Japan;(Brookes *et al.*, 1985)). Total soil C and N were determined on oven dry soil using a TruSpec CN analyser (Leco Corp., St Joseph, MI). Extraction efficiency conversion factors (Kec and Ken) of 0.35 and 0.50 were used to calculate MBC and MBN, respectively (Voroney, Brookes and Beyaert, 2007). Sulphate and other major anions were analysed in 1:5 w/v soil: distilled water extracts (200 rev min, 20°C, 30 min; (Zhao and McGrath, 1994)) using an ICS-2011 ion chromatograph (Dionex Inc., Sunnyvale, CA). Total dissolved S (dissolved organic and inorganic S) and other major cations were analysed by inductively coupled plasma-optical emission spectroscopy after extraction of the soil (200 rev min, 20°C, 30 min) in 1:5 w/v soil: 0.01 M Ca(H<sub>2</sub>PO<sub>4</sub>)<sub>2</sub> extracts using a Varian 710ES ICP-OES (Agilent Technologies, Santa Clara, CA). Dissolved organic sulphur (DOS) content was calculated from the difference between total dissolved S and inorganic SO<sub>4</sub><sup>2-</sup> (David and Mitchell, 1985).

Differences in soil properties along the altitudinal gradient were assessed using ANOVA with Tukey's LSD post hoc test using SPSS version 25.0 (IBM UK Ltd., Portsmouth, UK). *P* < 0.05 was used as the cut-off for statistical significance.

**Table 4.1** Characteristics of the 5 grassland soils used in the experiment. Values represent means  $\pm$  SEM,  $n = 3$ . Superscript letters represent significant differences between soils at the  $P < 0.05$  level. Data are presented on a dry weight basis.

	Site1	Site2	Site3	Site4	Site5
Altitude (m)	32	53	222	342	400
Bulk density <sup>a</sup> (g cm <sup>-3</sup> )	0.96 $\pm$ 0.04 <sup>a</sup>	0.72 $\pm$ 0.03 <sup>b</sup>	0.47 $\pm$ 0.06 <sup>cd</sup>	0.51 $\pm$ 0.05 <sup>c</sup>	0.55 $\pm$ 0.08 <sup>c</sup>
Water content <sup>b</sup> (%)	17.1 $\pm$ 0.2 <sup>e</sup>	53.5 $\pm$ 2.1 <sup>d</sup>	81.9 $\pm$ 1.3 <sup>b</sup>	77.1 $\pm$ 1.5 <sup>c</sup>	84.6 $\pm$ 0.5 <sup>a</sup>
pH <sup>c</sup>	5.92 $\pm$ 0.02 <sup>a</sup>	5.74 $\pm$ 0.06 <sup>b</sup>	5.67 $\pm$ 0.01 <sup>c</sup>	5.43 $\pm$ 0.01 <sup>d</sup>	4.20 $\pm$ 0.01 <sup>e</sup>
Electrical conductivity <sup>c</sup> ( $\mu$ S cm <sup>-1</sup> )	73.5 $\pm$ 3.6 <sup>a</sup>	56.8 $\pm$ 1.0 <sup>d</sup>	90.1 $\pm$ 0.8 <sup>b</sup>	42.1 $\pm$ 1.1 <sup>e</sup>	153.0 $\pm$ 1.4 <sup>a</sup>
NH <sub>4</sub> <sup>+</sup> -N <sup>d</sup> (mg kg <sup>-1</sup> )	6.6 $\pm$ 0.6 <sup>c</sup>	10.2 $\pm$ 6.1 <sup>c</sup>	9.7 $\pm$ 1.1 <sup>c</sup>	19.9 $\pm$ 9.2 <sup>b</sup>	61.5 $\pm$ 4.2 <sup>a</sup>
NO <sub>3</sub> <sup>-</sup> -N <sup>d</sup> (mg kg <sup>-1</sup> )	0.89 $\pm$ 0.06 <sup>e</sup>	12.6 $\pm$ 0.1 <sup>c</sup>	21.4 $\pm$ 2.1 <sup>b</sup>	4.0 $\pm$ 0.3 <sup>d</sup>	49.2 $\pm$ 1.2 <sup>a</sup>
Available P <sup>d</sup> (mg kg <sup>-1</sup> )	0.47 $\pm$ 0.05 <sup>ab</sup>	0.42 $\pm$ 0.01 <sup>b</sup>	0.51 $\pm$ 0.02 <sup>a</sup>	0.52 $\pm$ 0.09 <sup>a</sup>	0.58 $\pm$ 0.03 <sup>a</sup>
Dissolved organic C <sup>e</sup> (mg kg <sup>-1</sup> )	35.1 $\pm$ 8.7 <sup>a</sup>	40.5 $\pm$ 23.0 <sup>a</sup>	41.3 $\pm$ 3.7 <sup>a</sup>	40.6 $\pm$ 1.2 <sup>a</sup>	45.8 $\pm$ 7.0 <sup>a</sup>
Microbial biomass-C <sup>e</sup> (g kg <sup>-1</sup> )	0.67 $\pm$ 0.02 <sup>e</sup>	2.26 $\pm$ 0.16 <sup>d</sup>	2.94 $\pm$ 0.20 <sup>c</sup>	3.23 $\pm$ 0.38 <sup>b</sup>	94 $\pm$ 0.32 <sup>a</sup>
Total dissolved S <sup>f</sup> (mg S kg <sup>-1</sup> )	31.6 $\pm$ 2.1 <sup>d</sup>	34.4 $\pm$ 3.7 <sup>c</sup>	42.8 $\pm$ 0.6 <sup>b</sup>	42.9 $\pm$ 2.4 <sup>b</sup>	62.9 $\pm$ 6.6 <sup>a</sup>
Sulphate-S <sup>g</sup> (mg S kg <sup>-1</sup> )	4.37 $\pm$ 0.79 <sup>c</sup>	4.65 $\pm$ 0.33 <sup>c</sup>	5.53 $\pm$ 0.54 <sup>bc</sup>	6.71 $\pm$ 0.48 <sup>b</sup>	11.89 $\pm$ 1.77 <sup>a</sup>

<sup>a</sup>Dry bulk density measured by taking intact 100 cm<sup>3</sup> cores in the field and then oven drying at 105 °C overnight.

<sup>b</sup>Measured by drying at 105 °C overnight and expressed on a wet weight basis.

<sup>c</sup>Measured in a 1:5 (v/v) soil-to-distilled water extract.

<sup>d</sup>Measured in a 1:5 (w/v) 0.5 M K<sub>2</sub>SO<sub>4</sub> extract followed by colorimetry analysis via a Synergy-MX microplate reader.

<sup>e</sup>Measured in a 1:5 (w/v) 0.5 M K<sub>2</sub>SO<sub>4</sub> extract using a TOC-V-TN analyser.

<sup>f</sup>Measured in 1:5 w/v soil: 0.01 M Ca(H<sub>2</sub>PO<sub>4</sub>)<sub>2</sub> extracts using a Varian 710ES ICP-OES.

<sup>g</sup>Measured in 1:5 w/v soil: distilled water extracts using an ICS-2011 ion chromatograph.

#### 4.2.1 S-amino acid depletion from soil

A centrifugal-drainage (snap elution) approach was used to measure the rate of Met and Cys depletion from soil solution as described by Hill et al. (2008) and Wilkinson et al. (2014). Briefly, 1 g of field-moist soil was placed in a 1.5 cm<sup>3</sup> polypropylene micro-centrifuge tube in which a small hole (0.55 mm dia.) had been drilled in the base. This tube was then placed into another intact 1.5 cm<sup>3</sup> micro-centrifuge tube. A uniformly <sup>14</sup>C-labelled Cys or Met solution (200 µl, 0.5 kBq ml<sup>-1</sup>; PerkinElmer Inc., Waltham, MA) was added to the soil surface. Infiltration of the Cys and Met solution into the soil took less than 5 s. The range of concentrations used (0.01, 0.05, 0.1, 1, 10 or 100 mM) were chosen to reflect those likely to be naturally found in soil solution (0.01 - 0.05 mM), those present in the rhizosphere due to root exudation (0.1 - 1 mM), those present in soil after a root cell bursts (1 - 10 mM), and those in amino acid-based fertilisers (10 - 100 mM) (Jones and Darrah, 1994; Moe, 2013). After incubating <sup>14</sup>C-Cys and <sup>14</sup>C-Met in the soil for varying times (1, 3, 5, 10, 20, 30, 40, 60 mins), soil solution was recovered in the lower receptacle by centrifuging the soils for 1 min at 14000 g at 20°C. 100 µl of the recovered soil solution was mixed with Optiphase Hisafe 3 scintillation fluid (PerkinElmer Inc.) and <sup>14</sup>C determined by liquid scintillation counting using a Wallac 1404 scintillation counter with automated quench correction (Wallac EG&G, Milton Keynes, UK). All treatments were carried out in triplicate.

To account for any abiotic effects (e.g. imperfect mixing, dilution with the intrinsic non-labelled soil solution and sorption; Hill et al., 2008), the same experiment was carried out on field-moist soil that had been autoclaved (121°C, 30 min) and allowed to cool down immediately prior to use. This treatment was used to eliminate the microbial activity in the samples.

#### 4.2.2 S-amino acid mineralization

Cys and Met mineralization was measured by placing 5 g of field-moist soil into sterile polypropylene tubes (50 cm<sup>3</sup>). Subsequently, 1 ml of uniformly labelled <sup>14</sup>C-Cys or <sup>14</sup>C-Met (0.5 kBq ml<sup>-1</sup>; 0.01 to 100 mM; PerkinElmer Inc.) was added to soil surface. A 6 cm<sup>3</sup> polypropylene vial containing 1 ml of 1 M NaOH solution was then placed above the soil to trap any <sup>14</sup>CO<sub>2</sub> emitted. All tubes were then sealed and maintained at 20°C in the dark. The NaOH traps were changed after 0.25, 0.5, 1, 3, 6, 24 and 48 h. The <sup>14</sup>CO<sub>2</sub> in the NaOH traps was determined by liquid scintillation counting as described earlier. All treatments were carried out in triplicate. The same experiment was also carried out on field-moist soil that had been autoclaved as described above.

#### 4.2.3 Statistical and data analysis

After addition of the <sup>14</sup>C-labelled amino acid solution to the soil, the <sup>14</sup>C-Cys or Met solution gradually equilibrates with the intrinsic soil solution. The theoretical final concentration of <sup>14</sup>C-amino acid in the soil solution (<sup>14</sup>C<sub>max</sub>) after equilibration is therefore:

$$^{14}\text{C}_{\text{max}} = ^{14}\text{C}_{\text{added}} \times V_{\text{added}} / (V_{\text{added}} + V_{\text{native}}) \quad (\text{Eqn. 1})$$

where <sup>14</sup>C<sub>added</sub> is the activity of the <sup>14</sup>C solution added (kBq ml<sup>-1</sup>), V<sub>added</sub> is the volume of <sup>14</sup>C solution added and V<sub>native</sub> the volume of water already present in the soil. To investigate the time required to achieve perfect isotopic mixing, the dynamics of isotopic pool dilution was described in the autoclaved soil (i.e. no microbial removal) by a single first order exponential decay function:

$$^{14}\text{C}_{\text{sol}} = A + B \times \exp^{(-k \times t)} \quad (\text{Eqn.2})$$

Where <sup>14</sup>C<sub>sol</sub> is the amount of <sup>14</sup>C recovered in autoclaved soil solution over time, A represents the point when perfect mixing is achieved, t is the time after <sup>14</sup>C solution addition, and B describes the size of the added amino acid pool and, k is the rate constant describing the time



taken to reach equilibrium. It should be noted that an equilibrium was not reached in some soils. Consequently, the amount of  $^{14}\text{C}$  in solution recovered from sterile soils by centrifugation was higher than the theoretical maximum in some cases.

The half-time required to reach equilibrium ( $t_{1/2}$ ) was calculated from:

$$t_{1/2} = \ln(2)/k \quad (\text{Eqn.3})$$

In the non-sterile soils, the depletion of  $^{14}\text{C}$ -Cys or  $^{14}\text{C}$ -Met from soil solution was described by a range of first order kinetic equations. For the low substrate concentration (0.01 - 0.1 mM) a single first order exponential decay model, was fitted to the experimental data as follows:

$$^{14}\text{C}_{\text{sol}} = A_1 \times \exp(-k_1 \times t) \quad (\text{Eqn. 4})$$

Where  $C_{\text{sol}}$  is the  $^{14}\text{C}$  remaining in soil solution,  $k_1$  is the rate constant,  $t$  is time and  $A_1$  is the initial amino acid concentration at time,  $t_0$ .

At higher  $^{14}\text{C}$ -Cys or  $^{14}\text{C}$ -Met concentrations (1 - 100 mM) the depletion from soil solution as biphasic. Consequently, a double first order exponential decay model was fitted to the experimental data as follows:

$$^{14}\text{C}_{\text{sol}} = A_1 \times \exp(-k_1 \times t) + A_2 \times \exp(-k_2 \times t) \quad (\text{Eqn. 5})$$

Where  $A_1$  and  $A_2$  represent the size of the fast and slow depletion pools, respectively, and  $k_1$  and  $k_2$  represent the rate constants for these two pools, respectively.

It is well documented that the mineralization of low molecular weight substrates in soils is biphasic (Scow, Simkins and Alexander, 1986; Glanville *et al.*, 2012). Therefore, a double first order exponential decay equation was fitted to the experimental mineralization data as follows:

$$^{14}\text{C}_{\text{total}} = AA_{\text{catabolic}} \times \exp(-k_3 \times t) + AA_{\text{anabolic}} \times \exp(-k_4 \times t) \quad (\text{Eqn. 6})$$

where  $^{14}\text{C}_{\text{total}}$  is the total amount of  $^{14}\text{C}$  remaining in soil,  $AA_{\text{catabolic}}$  is the amount of amino acid-derived  $^{14}\text{C}$  partitioned into microbial catabolic processes and  $AA_{\text{anabolic}}$  is the amount of amino acid-derived  $^{14}\text{C}$  partitioned into microbial anabolic processes. The rate constants  $k_3$  and

$k_4$  describe the turnover rate of these two pools, respectively. For a full description of the model see Glanville et al. (2016). Half-lives of the two C pools was calculated according to Eqn. 3.

Microbial C use efficiency ( $CUE_{mic}$ ) for each substrate was calculated according to Jones et al. (2018) as follows:

$$CUE_{mic} = AA_{anabolic} / (AA_{catabolic} + AA_{anabolic}) \quad (\text{Eqn. 7})$$

All experiments were carried out in triplicate. After checking for normality, data were subjected to one-way ANOVA with LSD post hoc test using SPSS version 25.0 (IBM UK Ltd., Portsmouth, UK).  $P < 0.05$  was used as the cut-off for statistical significance.


## 4.3 Results

### 4.3.1 S-amino acid depletion from soil solution

#### 4.3.1.1 Substrate depletion from soil solution affected by soil conditions

The change in soil solution concentration of  $^{14}\text{C}$ -Cys and Met in the autoclaved soils (where biotic removal is prevented) allowed us to calculate the dynamics of isotope pool dilution (i.e. the mixing of the introduced isotope with the native soil solution). After 1 h, 11.6%, 33.0% and 21.9% of added  $^{14}\text{C}$ -Cys was removed from the autoclaved soils 1, 2 and 3 respectively due to abiotic factors (such as adsorption, or occlusion within micropores that are not drained by centrifugation), and 32.6% and 27.0% of added  $^{14}\text{C}$ -Met was removed in this way from soil 1 and 2 separately (Fig. 4.1). More  $^{14}\text{C}$ -Cys was recovered from soil 4 and 5 (107.7%, 120.6% respectively), and more  $^{14}\text{C}$ -Met from soil 3, 4 and 5 (145.5%, 108.1% and 140.2% separately) after 1 h than would be predicted following perfect mixing of substrate and soil, indicating that the added substrate had not fully mixed with soil prior to starting the centrifugal drainage procedure. We found that recovery of added  $^{14}\text{C}$  from autoclaved soil followed first order exponential decay equation (Eqn. 2;  $R^2 > 0.95$ ). From this, we calculated half-time required for the two solution pools to mix, the majority of  $^{14}\text{C}$  Cys and Met mixing


and abiotic depletion occurred rapidly within the first 40 minutes (half times ranged from 3.1 minutes for Cys in soil 3 to 25.4 minutes for Cys in soil 4), this excludes values for soil 1 and 2 which undergo a slower exponential decay.



**Fig. 4.1** Percent deviation from perfect mixing after addition of  $^{14}\text{C}$ -Cys or Met solution (0.01 mM) to autoclaved soils collected from 5 contrasting grassland sites. A 200  $\mu\text{l}$  of  $^{14}\text{C}$ -amino acid (Cys or Met) was added to 1g of soil containing non  $^{14}\text{C}$  labelled water. Percentages are based on the theoretical amount of  $^{14}\text{C}$  available in solution assuming no biotic loss. The dashed line at 0% represents the point at which perfect mixing has occurred throughout the soil. Values above 0% (dotted line) indicates incomplete mixing of added solution and soil native soil solution (values above 0%), values below 0% indicates abiotic sorption. Lines represent first order exponential equations fit to the experimental data. Data points represent means  $\pm$  SEM ( $n = 3$ ).

Based on this isotopic pool dilution dynamics in autoclaved soils, depletion of  $^{14}\text{C}$ -amino acid from soil solution was corrected for mixing and abiotic processes, the data presented hereafter refer to  $^{14}\text{C}$  directly available for microbial uptake. As shown in Fig. 4.2,  $^{14}\text{C}$ -Cys removal from soil solution by the microbial community was extremely rapid, with  $87.4 \pm 1.2\%$  of available  $^{14}\text{C}$ -Cys taken up by microbes from soil 1 after only 60 s. The

proportion of  $^{14}\text{C}$ -Cys removed from soil solution by microbes fell significantly ( $p < 0.05$ ) in soil from high elevation, low productivity sites to  $33.4 \pm 7.3\%$  (soil 2),  $42.5 \pm 12.7\%$  (soil 3),  $45.8 \pm 8.9\%$  (soil 4) and  $43.3 \pm 14.7\%$  (soil 5) within one minute. Similarly, available  $^{14}\text{C}$  uptake by soil microbes from soil solution also occurred very quickly after Met addition, especially in soil 1 and 2 where  $59.4 \pm 2.2\%$  and  $57.3 \pm 5.7\%$  of the available  $^{14}\text{C}$ -Met was taken up by the microbial biomass within 1 min. The amount of  $^{14}\text{C}$ -Met removed from soil solution fell significantly ( $p < 0.05$ ) in soils from high elevation, low productivity sites, to  $42.0 \pm 3.9\%$  (soil 3),  $39.5 \pm 10.1\%$  (soil 4) and  $20.2 \pm 8.0\%$  (soil 5) when compared with soil 1 and 2.



**Fig. 4.2** Amount of  $^{14}\text{C}$ -label remaining in soil solution after addition of a  $^{14}\text{C}$  labelled Cys and Met (0.01 mM) to soil collected from 5 grassland sites of increasing elevation and decreasing productivity within 60 minutes, corrected from autoclaved soil data as described in material and methods section. Percentages are based on theoretical amount of  $^{14}\text{C}$  available in solution assuming perfect mixing of the substrate with soil native water (Eqn. 1). Values represent means  $\pm$  SEM ( $n = 3$ ), lines represent fits of single first order exponential equations to the experimental data.

The depletion of  $^{14}\text{C}$  from soil solution at sites 1 and 2 was best described by a single exponential decay equation (Eqn. 4) for additions of Cys ( $r^2 > 0.96$ ) or Met ( $r^2 > 0.89$ ). The half time ( $t_{1/2}$ ) for Cys depletion from soil solution was significantly shorter than that for Met ( $p <$


0.05) in soil 1, but there were no significant differences between the half times in soil 2. Furthermore,  $^{14}\text{C}$  was removed significantly faster ( $p < 0.05$ ) from soil 1 than soil 2 following both Cys and Met addition. By contrast, soils 3, 4 and 5 behaved differently from soil 1 and 2, and  $^{14}\text{C}$  depletion from soil solution was best described by a double exponential decay equation (Eq (4);  $r^2 > 0.98$ ). This implies that in grassland soils from higher elevation, lower productivity soils,  $^{14}\text{C}$  depletion occurred in two phases. Roughly two thirds of added  $^{14}\text{C}$ -Cys was depleted in a rapid initial phase (Table 4.2;  $t_{1/2} < 1\text{min}$ ), while the remaining third was depleted more slowly, with half times up to 27 mins (no significant difference between these three soils). Furthermore, compared to Met, more Cys was taken up by soil microbes after one minute in soil 1 and 2 ( $p < 0.05$ ), there were no significant differences between soil 3, 4 and 5.

#### *4.3.1.2 Substrate depletion from soil solution as affected by substrate concentration*

As described earlier, by monitoring the recovery of  $^{14}\text{C}$ -amino acid in soil solution extracted from autoclaved soil over time, we obtained deviation from perfect mixing of added  $^{14}\text{C}$  labelled solution with native soil solution due to abiotic process including adsorption and occlusion within micro-pores that are not drained by our centrifugal drainage procedure. After one hour, 11.8%, 25.3%, 14.3%, 7.7%, 22.7%, 11.0% of the added  $^{14}\text{C}$ -Cys was removed from the autoclaved soil at concentration of 0.01, 0.05, 0.1 mM, and 1, 10, 100 mM respectively due to abiotic factors, and 33.6%, 30.8%, 26.3%, 27.3%, 25.2% and 23.4% of added Met were removed in this way, respectively (Fig. 4.3).  $^{14}\text{C}$ -amino acid recovery from sterile soil followed a first order exponential decay (Eqn. 2;  $R^2 > 0.86$ ), with half-lives ranging from 1.8 min at 1 mM (Cys) to 75.3 min at 0.01 mM (Met). Perfect mixing of native soil solution and added  $^{14}\text{C}$ -label was achieved within approximately 40 mins after substrate addition.


In the non-sterile soil, the microbial community removed  $87.4 \pm 1.2\%$  and  $59.4 \pm 2.2\%$  of the added  $^{14}\text{C}$ -Cys and Met from the soil solution within 1 minute when added at 0.01 mM,

respectively. This fell to  $30.2 \pm 1.9\%$  and  $29.1 \pm 3.9\%$  when added at the highest concentration (100 mM). In absolute terms, however, the actual amount of Cys and Met removed from soil solution by soil microbes increased with increasing amino acid concentration, at the highest added amino acid concentration (100 mM), uptake rates were equivalent to  $7.1 \text{ mmol } ^{14}\text{C-Cys (kg}^{-1} \text{ DW soil) min}^{-1}$  and  $6.8 \text{ mmol } ^{14}\text{C-Met (kg}^{-1} \text{ DW soil) min}^{-1}$ , respectively after only one minute.



**Fig. 4.3** Deviation from perfect mixing when a  $^{14}\text{C}$ -Cys and Met solution (0.01 to 100 mM) is added to autoclaved soil containing non  $^{14}\text{C}$  labelled water. Percentages are based on the theoretical amount of  $^{14}\text{C}$  available in solution assuming perfect mixing of the substrate with the native soil solution. Deviations from 0% (dotted line) indicates incomplete mixing (values above 0) or abiotic sorption (values below 0%). Lines represent fits of first order kinetic equations (Eqn. 2) to the experimental data. Values represent means  $\pm$  SEM ( $n = 3$ ).


The dynamics of both Cys and Met depletion from soil solution were dependent upon the amount of substrate added to the soil (Fig. 4.4). The depletion of  $^{14}\text{C}$ -Cys and Met from soil solution was best described by a single exponential decay equation ( $R^2 > 0.92$  in all cases) when added at 0.01, 0.05 and 0.1 mM, but better described by a double exponential kinetic decay model ( $R^2 > 0.98$  in all cases) when added at higher concentrations (1 mM, 10 mM, 100 mM; Table 4.3). The exponential coefficients and half-lives ( $t_{1/2}$ ) describing the loss of Cys and Met from the soil solution are presented in Table 4.3.



**Fig. 4.4** Amount of  $^{14}\text{C}$  label in soil solution after the addition of a  $^{14}\text{C}$  labelled Cys and Met to non-sterile grassland soil (site 1). Lines represent fits of single (0.01 to 0.1 mM) and double (1 to 100 mM) first order kinetic equations to the experimental data. Values represent means  $\pm$  SEM ( $n = 3$ ).

#### 4.3.2 Microbial C use efficiency for individual substrates

##### 4.3.2.1 S-amino acid mineralization from autoclaved soils



**Fig. 4.5** Kinetics of the  $^{14}\text{C}$  released from cumulative microbial respiration after addition of  $^{14}\text{C}$  labelled Cys or Met (0.01 mM) to autoclaved grassland soil (site 1) within 48h. Values represent means  $\pm$  SEM ( $n = 3$ ).

The supply of  $^{14}\text{C}$ -Cys or Met to autoclaved grassland soil led to little  $^{14}\text{CO}_2$  respiration within 48 hours, the amount of  $^{14}\text{C}$ -Cys or Met released as  $^{14}\text{CO}_2$  corresponded to less than 1% of total  $^{14}\text{C}$ -substrate input (Fig. 4.5). The low recovery of  $^{14}\text{C}$  from microbial respiration indicates that autoclaving process was successful at killing the active soil microorganisms in short term, these results concur with the findings of similar studies on autoclaved soils (Carter, Yellowlees and Tibbett, 2007).


#### 4.3.2.2 *S-amino acid mineralization affected by soil conditions*

The  $^{14}\text{C}$ -amino acid mineralization was divided into two main phases: the first phase, rapid respired  $^{14}\text{CO}_2$  may be attributable to the immediate use of the added substrate for catabolic process, in the second phase,  $^{14}\text{CO}_2$  is released much more slowly following microbial turnover through catabolic processes (Eqn. 5;  $K_3$  and  $K_4$ ; Table 4.4). This is in line with what has previously been reported (Vinolas, Vallejo and Jones, 2001; Wilkinson, Paul W. Hill, *et al.*, 2014). The first order double exponential decay equation gave a good fit to these biphasic experimental data (Fig. 4.6;  $R^2 > 0.99$ ). The exponential coefficients and half-lives of substrate mineralization are presented in Table 4.4. As little  $^{14}\text{CO}_2$  evolution from soil solution was observed in the autoclaved soil treatments (Fig. 4.5), we assume that  $^{14}\text{C}$ -Cys and Met mineralization from soil solution occurred almost entirely due to microbial activity.

In soil 2, 3 and 4, more  $^{14}\text{C}$ -Cys was allocated to the initial rapid mineralization pool than in the secondary mineralization pool ( $p < 0.01$ ), yet there were no significant differences ( $p < 0.01$ ) in two Cys pools in soil 1 and 5, where marginally more  $^{14}\text{C}$ -Cys was allocated to the fast turnover pool. Half-lives for the  $^{14}\text{C}$ -Cys fast mineralization pool  $c_1$  ranged from 0.6 h (site 3) to approximately 1.8 h (site 5), while the slower turnover pool has a much longer half-life, ranging from ca. 10.1 d (site 5) to 67.8 d (site 4). On the other hand, between 7.9% (soil 5)



and 38.0% (soil 2) of  $^{14}\text{C}$ -Met was allocated to the rapid mineralization pool in soils, with the slow turnover pool accounting for 66.4 - 92.4% of added  $^{14}\text{C}$ -Met.



**Fig. 4.6** Amount of  $^{14}\text{C}$ -label remaining in soil solution and microbial biomass after the injection of  $^{14}\text{C}$  labelled Cys and Met (0.01 mM) to soil collected from five contrasting grassland sites. Values represent means  $\pm$  SEM ( $n = 3$ ). Lines represent fits a double first order kinetic equations to the experimental data.


More  $^{14}\text{C}$ -Met was retained in soil solution and soil microbial biomass one hour after incubation from all five sites compared to Cys (0.01 mM; Fig. 4.6;  $p < 0.001$ ), due to faster mineralization of Cys than that of Met. One hour after substrate addition, the cumulative  $^{14}\text{CO}_2$  evolved from added  $^{14}\text{C}$ -Cys was  $16.7 \pm 0.5\%$  (site 5)  $31.4 \pm 1.1\%$  (site 4),  $42.7 \pm 0.5\%$  (site 3),  $31.0 \pm 0.9\%$  (site 2), and  $14.5 \pm 0.5\%$  (site 1), whereas significantly ( $p < 0.05$ ) less  $^{14}\text{C}$ -Met was mineralized in all five soils,  $3.0 \pm 0.3\%$  (site 5),  $6.1 \pm 0.6\%$  (site 4),  $11.8 \pm 0.4\%$  (site 3),  $7.9 \pm 0.6\%$  (site 2), and  $4.6 \pm 0.1\%$  (site 1). Furthermore, rates of amino acid mineralization in soil 5 were slower, with the amount of  $^{14}\text{C}$  remaining in soil and soil microbial biomass being significantly greater than that in other four soils ( $p < 0.05$ ).

#### 4.3.2.3 *S*-amino acid mineralization affected by substrate concentrations

The double exponential decay equation gave a good fit to the depletion of both  $^{14}\text{C}$ -Cys and Met from soil by microbial mineralization over time ( $R^2 > 0.98$  in all cases). The exponential coefficients and half-lives of double exponential decay equations are presented in Table 4.5.

The proportion of Cys released through microbial respiration in the initial rapid phase of  $^{14}\text{CO}_2$  evolution range from  $3.5 \pm 0.6\%$  (100 mM) to  $60.4 \pm 0.1\%$  (0.05 mM), with half-lives ranging from  $1.5 \pm 0.1$  h to  $14.5 \pm 0.7$  h, while the proportion of Cys respired in the secondary slower phase range from  $41.4 \pm 1.0\%$  (1 mM) to  $96.5 \pm 0.6$  (100 mM), with half-lives ranging from  $6.6 \pm 0.04$  to  $43.8 \pm 3.7$  d. More Cys was allocated in the first fast-turnover pool than in the secondary slow-turnover pool when added at 0.01 mM, 0.05 mM, 0.1 mM and 1 mM, yet more Cys was allocated in the second slow-turnover pool when added at 10 mM and 100 mM.

On the other hand, more Met was retained in soil solution and microbial biomass compared with Cys at all concentrations ( $p < 0.001$ ) at all our sampling time points (Fig. 4.7). Less than 50% of added Met was allocated to the first initial rapid mineralization pool at concentrations of 0.01 mM, 0.05 mM, 0.1 mM and 100 mM with half-lives for the fast turnover of  $c_1$  pool ranging from  $4.0 \pm 0.1$  to  $781.7 \pm 12.8$  h, while the slower pool ( $c_2$ ) had a much slower half-life, ranging from  $17.2 \pm 0.5$  to  $32.5 \pm 0.5$  d following substrate addition. More Met was allocated to the first initial rapid phase when added at 1mM and 10 mM, with half-life of  $17.5 \pm 0.1$ ,  $114.8 \pm 19.2$  h for the turnover of fast pool,  $57.3 \pm 7.7$ ,  $4.8 \pm 0.8$  d for the turnover of slow pool, respectively.



**Fig. 4.7** Amount of <sup>14</sup>C- Cys and Met remaining in soil solution and soil microbial biomass within 48 hours added at 6 concentrations ranging from 0.01 to 100 mM (soil 1). Data points represent means  $\pm$  SEM ( $n = 3$ ). Lines represent fits of single or double first order exponential equations to the experimental data.

In the non-sterile soil, within 30 minutes after substrate application, following addition of the <sup>14</sup>C-labelled Cys or Met (0.01 mM) to the soil, <sup>14</sup>C released from cumulative microbial respiration accounted for  $5.3 \pm 0.3\%$  and  $1.9 \pm 0.01\%$  of the added <sup>14</sup>C-Cys and Met, this fell to  $0.5 \pm 0.1\%$  and  $0.2 \pm 0.03\%$  when added at the highest concentration (100 mM). In absolute terms, however, the actual amount of Cys and Met removed from soil solution by soil microbial respiration increased with increasing amino acid concentration, at the highest added amino acid concentration (100 mM), mineralization rates were equivalent to  $3.7 \mu\text{mol } ^{14}\text{C-Cys (kg}^{-1} \text{ DW soil) min}^{-1}$  and  $1.3 \mu\text{mol } ^{14}\text{C-Met (kg}^{-1} \text{ DW soil) min}^{-1}$ , respectively after only one minute.

According to our results,  $5.63 \pm 0.33\%$  and  $1.94 \pm 0.01\%$  of the added <sup>14</sup>C-Cys and Met taken up by the microbial community from the soil solution within 30 minutes were subsequently used for microbial respiration when added at 0.01 mM, respectively (Table 4.6). This fell to  $0.98 \pm 0.13\%$  and  $0.40 \pm 0.05\%$  when added at the highest concentration (100 mM).

**Table 4.2** Coefficients of single first order curves describing the depletion of Cys and Met from soil solution over time in 5 contrasting grassland soils along an altitudinal productivity gradient.  $A_1$  and  $A_2$  are estimated pool sizes for fast and slow substrate uptake routes, and  $K_1$  and  $K_2$  are the rate constants for the fast and slow uptake routes, respectively. Half time values for pools  $A_1$  and  $A_2$  are represented by  $t_{1/2}$  values and these are determined from  $K_1$  and  $K_2$ . Values represent means  $\pm$  SEM ( $n = 3$ ).

Substrate	Site	$A_1$	$K_1$	$A_2$	$K_2$	$A_1 t_{1/2}$ (min)	$A_2 t_{1/2}$ (min)
Cys	Site 1	99.9 $\pm$ 5.9	2.03 $\pm$ 0.46			0.34 $\pm$ 0.02	
	Site 2	100.3 $\pm$ 8.0	0.44 $\pm$ 0.08			1.58 $\pm$ 0.15	
	Site 3	61.3 $\pm$ 6.6	0.93 $\pm$ 0.25	37.3 $\pm$ 5.1	0.03 $\pm$ 0.01	0.83 $\pm$ 0.55	25.6 $\pm$ 4.5
	Site 4	59.7 $\pm$ 2.8	1.38 $\pm$ 0.18	40.4 $\pm$ 2.0	0.03 $\pm$ 0.00	0.48 $\pm$ 0.18	21.8 $\pm$ 3.6
	Site 5	68.9 $\pm$ 3.2	0.96 $\pm$ 0.11	31.1 $\pm$ 2.4	0.03 $\pm$ 0.00	0.76 $\pm$ 0.37	27.9 $\pm$ 3.6
Met	Site 1	100.1 $\pm$ 1.7	0.91 $\pm$ 0.04			0.76 $\pm$ 0.03	
	Site 2	91.3 $\pm$ 9.8	0.42 $\pm$ 0.11			1.64 $\pm$ 0.32	
	Site 3	50.0 $\pm$ 3.5	1.55 $\pm$ 0.28	50.0 $\pm$ 2.7	0.05 $\pm$ 0.01	0.40 $\pm$ 0.24	12.9 $\pm$ 3.3
	Site 4	35.8 $\pm$ 7.2	2.92 $\pm$ 3.11	64.2 $\pm$ 6.1	0.09 $\pm$ 0.01	0.22 $\pm$ 0.38	8.0 $\pm$ 1.2
	Site 5	82.4 $\pm$ 15.1	0.24 $\pm$ 0.05	17.1 $\pm$ 15.5	0.04 $\pm$ 0.03	2.14 $\pm$ 1.65	9.0 $\pm$ 7.6

**Table 4.3** Influence of amino acid concentration on the coefficients of single (0.01 mM and 0.05 mM) and double (0.1-100 mM) first order curve fits describing the depletion of  $^{14}\text{C}$  from soil solution over time.  $A_1$  and  $A_2$  are estimated pool sizes for fast and slow substrate uptake routes, and  $K_1$  and  $K_2$  are the rate constants for the fast and slow uptake routes, respectively. Half time values for pools  $A_1$  and  $A_2$  are represented by  $t_{1/2}$  values and these are determined from  $K_1$  and  $K_2$ , respectively. Values represent means  $\pm$  SEM ( $n = 3$ ).

Substrate	Concentration	$A_1$	$K_1$	$A_2$	$K_2$	$A_1 t_{1/2}$ (min)	$A_2 t_{1/2}$ (min)	$R^2$
Cys	0.01 mM	99.9 $\pm$ 5.9	2.03 $\pm$ 0.46			0.34 $\pm$ 0.02		0.98
	0.05 mM	92.1 $\pm$ 7.2	0.41 $\pm$ 0.08			1.69 $\pm$ 0.12		0.97
	0.1 mM	85.0 $\pm$ 11.3	0.28 $\pm$ 0.09			2.45 $\pm$ 0.15		0.92
	1 mM	48.5 $\pm$ 4.5	3.42 $\pm$ 2.78	51.4 $\pm$ 2.6	0.01 $\pm$ 0.002	0.17 $\pm$ 0.15	53.2 $\pm$ 15.9	0.99
	10 mM	40.6 $\pm$ 5.2	1.38 $\pm$ 0.49	59.1 $\pm$ 3.0	0.01 $\pm$ 0.001	0.52 $\pm$ 0.15	109.7 $\pm$ 23.8	0.98
	100 mM	38.8 $\pm$ 3.0	1.43 $\pm$ 0.32	61.1 $\pm$ 1.7	0.004 $\pm$ 0.001	0.48 $\pm$ 0.14	155.0 $\pm$ 41.9	0.99
Met	0.01 mM	100.0 $\pm$ 1.7	0.91 $\pm$ 0.04			0.76 $\pm$ 0.03		0.99
	0.05 mM	87.6 $\pm$ 9.8	0.30 $\pm$ 0.08			2.29 $\pm$ 0.04		0.94
	0.1 mM	88.9 $\pm$ 7.0	0.24 $\pm$ 0.04			2.85 $\pm$ 0.13		0.97
	1 mM	44.2 $\pm$ 1.9	2.80 $\pm$ 0.69	55.7 $\pm$ 1.0	0.01 $\pm$ 0.001	0.25 $\pm$ 0.04	59.4 $\pm$ 3.0	0.99
	10 mM	41.9 $\pm$ 3.5	1.45 $\pm$ 0.35	57.9 $\pm$ 1.9	0.004 $\pm$ 0.001	0.48 $\pm$ 0.08	185.9 $\pm$ 87.6	0.99
	100 mM	35.2 $\pm$ 1.7	1.70 $\pm$ 0.26	64.7 $\pm$ 0.9	0.002 $\pm$ 0.001	0.35 $\pm$ 0.31	307.5 $\pm$ 142.0	0.99

**Table 4.4** Kinetic coefficients of single and double exponential models describing the depletion of  $^{14}\text{C}$ -Cys and Met from soil and loss in microbial respiration over time in 5 contrasting grassland soils along an altitudinal productivity gradient.  $AA_{\text{catabolic}}$  and  $AA_{\text{anabolic}}$  are estimated pool sizes for fast and slow substrate uptake routes, and  $K_3$  and  $K_4$  are the rate constants for the fast and slow uptake routes, respectively. Half-life values for pools  $AA_{\text{catabolic}}$  and  $AA_{\text{anabolic}}$  are determined from  $K_3$  and  $K_4$ , respectively. Values represent means  $\pm$  SEM ( $n = 3$ ).

Substrate	Site	$AA_{\text{catabolic}}$	$K_3$	$AA_{\text{anabolic}}$	$K_4$	Catabolic $t_{1/2}$ (h)	Anabolic $t_{1/2}$ (d)	$R^2$
Cys	Site 1	$51.4 \pm 3.4$	$0.4 \pm 0.1$	$51.0 \pm 3.5$	$0.001 \pm 0.002$	$1.6 \pm 0.1$	$31.2 \pm 6.3$	0.99
	Site 2	$64.7 \pm 3.2$	$0.7 \pm 0.1$	$37.9 \pm 3.0$	$0.003 \pm 0.002$	$1.1 \pm 0.0$	$11.5 \pm 5.0$	0.99
	Site 3	$62.8 \pm 4.0$	$1.0 \pm 0.2$	$38.9 \pm 3.1$	$0.001 \pm 0.002$	$0.6 \pm 0.0$	$10.4 \pm 2.5$	0.99
	Site 4	$61.2 \pm 5.1$	$0.7 \pm 0.1$	$37.2 \pm 4.7$	$0.002 \pm 0.004$	$0.9 \pm 0.2$	$67.8 \pm 50.1$	0.99
	Site 5	$53.3 \pm 0.8$	$0.4 \pm 0.0$	$46.9 \pm 0.9$	$0.002 \pm 0.001$	$1.8 \pm 0.4$	$10.1 \pm 7.8$	0.99
Met	Site 1	$26.3 \pm 2.3$	$0.2 \pm 0.0$	$74.1 \pm 2.3$	$0.002 \pm 0.001$	$3.2 \pm 0.1$	$11.2 \pm 0.6$	0.99
	Site 2	$38.0 \pm 3.0$	$0.2 \pm 0.0$	$61.2 \pm 3.1$	$0.001 \pm 0.001$	$3.7 \pm 0.9$	$20.7 \pm 7.6$	0.99
	Site 3	$34.0 \pm 1.8$	$0.4 \pm 0.0$	$66.4 \pm 1.9$	$0.007 \pm 0.001$	$1.9 \pm 0.1$	$4.3 \pm 0.3$	0.99
	Site 4	$29.3 \pm 1.7$	$0.2 \pm 0.0$	$71.4 \pm 1.8$	$0.006 \pm 0.001$	$2.9 \pm 0.4$	$4.2 \pm 0.5$	0.99
	Site 5	$7.9 \pm 0.6$	$0.4 \pm 0.1$	$92.4 \pm 0.7$	$0.009 \pm 0.000$	$1.9 \pm 0.9$	$3.0 \pm 0.2$	0.99

**Table 4.5** Influence of amino acid concentration on the kinetic coefficients of double exponential models describing the depletion of  $^{14}\text{C}$ -Cys and Met from soil and loss in microbial respiration over time over a 48 h-incubation. Values represent means  $\pm$  SEM ( $n = 3$ ).

Substrate	Concentration	$AA_{\text{catabolic}}$	$K_3$	$AA_{\text{anabolic}}$	$K_4$	Catabolic $t_{1/2}$ (h)	Anabolic $t_{1/2}$ (d)	$R^2$
Cys	0.01 mM	$52.1 \pm 2.0$	$0.4 \pm 0.03$	$50.5 \pm 2.0$	$0.001 \pm 0.0001$	$1.5 \pm 0.1$	$43.8 \pm 3.7$	0.99
	0.05 mM	$60.4 \pm 0.1$	$0.2 \pm 0.001$	$41.8 \pm 0.1$	$0.001 \pm 0.0001$	$3.2 \pm 0.02$	$33.8 \pm 1.9$	0.99
	0.1 mM	$60.1 \pm 0.9$	$0.2 \pm 0.003$	$41.7 \pm 0.8$	$0.001 \pm 0.0001$	$3.8 \pm 0.1$	$23.3 \pm 2.6$	0.99
	1 mM	$59.2 \pm 1.1$	$0.08 \pm 0.001$	$41.4 \pm 1.0$	$0.001 \pm 0.00003$	$8.4 \pm 0.1$	$28.2 \pm 0.9$	0.99
	10 mM	$41.2 \pm 1.9$	$0.04 \pm 0.002$	$58.9 \pm 1.8$	$0.003 \pm 0.0001$	$14.5 \pm 0.7$	$11.7 \pm 0.3$	0.99
	100 mM	$3.5 \pm 0.6$	$0.2 \pm 0.02$	$96.5 \pm 0.6$	$0.004 \pm 0.00003$	$4.4 \pm 0.4$	$6.6 \pm 0.04$	0.99
Met	0.01 mM	$30.5 \pm 0.6$	$0.2 \pm 0.01$	$69.6 \pm 0.6$	$0.001 \pm 0.00003$	$4.0 \pm 0.1$	$25.9 \pm 0.8$	0.99
	0.05 mM	$40.2 \pm 0.2$	$0.1 \pm 0.01$	$60.2 \pm 0.3$	$0.002 \pm 0.00003$	$5.9 \pm 0.5$	$18.8 \pm 0.4$	0.99
	0.1 mM	$46.1 \pm 0.5$	$0.1 \pm 0.002$	$54.5 \pm 0.6$	$0.002 \pm 0.00005$	$7.6 \pm 0.2$	$17.2 \pm 0.5$	0.99
	1 mM	$57.0 \pm 0.9$	$0.04 \pm 0.0002$	$44.4 \pm 0.9$	$0.001 \pm 0.00007$	$17.5 \pm 0.1$	$57.3 \pm 7.7$	0.99
	10 mM	$50.1 \pm 0.3$	$0.01 \pm 0.001$	$51.1 \pm 0.1$	$0.006 \pm 0.001$	$114.8 \pm 19.2$	$4.8 \pm 0.8$	0.98
	100 mM	$44.1 \pm 0.1$	$0.001 \pm 0.00002$	$55.6 \pm 0.1$	$0.001 \pm 0.00002$	$781.7 \pm 12.8$	$32.5 \pm 0.5$	0.99

**Table 4.6** Influence of amino acid concentration on the uptake and mineralization of either Cys or Met in the first 30 minutes after substrate addition, and the percent of amino acid taken up which was respired as  $^{14}\text{CO}_2$  during the same time period. Values represent means  $\pm$  SEM ( $n = 3$ ).

		Rate of amino acid uptake (nmol amino acid kg <sup>-1</sup> DW soil min <sup>-1</sup> )	Rate of amino acid mineralization (nmol amino acid kg <sup>-1</sup> DW soil min <sup>-1</sup> )	Percentage of $^{14}\text{C}$ taken up that is subsequently respired (%)
Cys	0.01 mM	63 $\pm$ 1	3.5 $\pm$ 0.1	5.63 $\pm$ 0.33
	0.05 mM	329 $\pm$ 1	9.63 $\pm$ 0.4	2.93 $\pm$ 0.14
	0.1 mM	637 $\pm$ 1	15.0 $\pm$ 0.5	2.37 $\pm$ 0.09
	1 mM	4558 $\pm$ 239	92.8 $\pm$ 2.5	2.04 $\pm$ 0.16
	10 mM	35460 $\pm$ 2950	666.3 $\pm$ 95.4	1.88 $\pm$ 0.24
	100 mM	326084 $\pm$ 25950	3196.3 $\pm$ 610.4	0.98 $\pm$ 0.13
Met	0.01 mM	65 $\pm$ 1	1.2 $\pm$ 0.0	1.94 $\pm$ 0.01
	0.05 mM	319 $\pm$ 1	4.7 $\pm$ 0.0	1.50 $\pm$ 0.02
	0.1 mM	649 $\pm$ 2	7.3 $\pm$ 0.2	1.12 $\pm$ 0.04
	1 mM	4112 $\pm$ 139	21.8 $\pm$ 2.6	0.53 $\pm$ 0.05
	10 mM	33071 $\pm$ 3350	148.1 $\pm$ 8.1	0.45 $\pm$ 0.04
	100 mM	288617 $\pm$ 25564	1141.9 $\pm$ 201.2	0.40 $\pm$ 0.05



### 4.3.3 Carbon turnover rates in soil

Microbial mineralization of amino acid- $^{14}\text{C}$  to  $^{14}\text{CO}_2$  was extremely fast, but slower than amino acid- $^{14}\text{C}$  uptake from the soil solution. A maximum of 59.9% of added amino acid was respired with half times of 1.6 - 4.5 min when taken up at amino acid concentration close to those found naturally in the soil solution (0.01 – 0.05 mM; Table 4.5; Table 4.7). The remainder was cycled through the microbial biomass before mineralization. At steady state, the rates of amino acid removal from the soil solution should be matched with new inputs from microbial/plant root turnover, exudation and hydrolysis of soil organic matter. Therefore assuming 0.01 – 0.05 mM are the closest concentrations to those in natural soil solution, we estimate that the rate of S-containing amino acids inputs to grassland soil was between 6.35 - 329.0 nmol amino acid  $\text{kg}^{-1}$  DW soil  $\text{min}^{-1}$ .

**Table 4.7** Influence of amino acid concentration on microbial C use efficiency ( $\text{CUE}_{\text{mic}}$ ). Values represent means  $\pm$  SEM ( $n = 3$ ).

Substrate	Soil solution concentration	$\text{CUE}_{\text{mic}}$
Cys	0.01 mM	$0.49 \pm 0.02$
	0.05 mM	$0.41 \pm 0.001$
	0.1 mM	$0.41 \pm 0.01$
	1 mM	$0.41 \pm 0.01$
	10 mM	$0.59 \pm 0.02$
	100 mM	$0.96 \pm 0.01$
Met	0.01 mM	$0.69 \pm 0.01$
	0.05 mM	$0.59 \pm 0.002$
	0.1 mM	$0.54 \pm 0.005$
	1 mM	$0.43 \pm 0.008$
	10 mM	$0.51 \pm 0.001$
	100 mM	$0.56 \pm 0.001$

## 4.4 Discussion

### 4.4.1 Kinetics of Cys and Met uptake by soil microbial community

Across all five sites, DON was the dominant form of soluble N within the soil solution, contributing between 22.5 to 82.3% of total DON in five sites. Similarly, DOS was the dominant form of soluble S within the soil solution, contributing between 81.1 to 87.1% of total dissolved S across the five sites (Table 4.1). This is in line with our findings in Chapter 3 (Table 3.1). The concentration of DOS was of the same order magnitude in all five soils ranging from 27.5 to 51.0 mg kg<sup>-1</sup> soil DW, while the sulphate concentrations were one order of magnitude lower, varying from 4.4 to 11.9 mg kg<sup>-1</sup> soil DW. Therefore, in our selected grasslands, DON and DOS represent important reservoirs of soluble N and S in the soil. This is also in agreement with previous studies (Farrell, Hill, Farrar, *et al.*, 2011; Meena and Improvement, 2018).

The results presented here are direct evidence to support earlier work that the uptake of amino acids by soil microbial community from soil solution can be exceedingly rapid (Jones and Kielland, 2012; Farrell, Prendergast-Miller, *et al.*, 2014; Wilkinson, Paul W. Hill, *et al.*, 2014; Carswell *et al.*, 2016). Within only 10 minutes after substrate addition, over 80% of added <sup>14</sup>C-Cys or Met had been removed from soil solution, with the exact amount varying depending on soil conditions and substrate concentrations. By the end of our 60-minute incubation experiment, less than 5% of the added amino acid-<sup>14</sup>C were retained in soil solution. Moreover, given that we removed living roots in this study (and associated symbionts), it is possible that substrate depletion could occur even more rapidly *in situ* than those measured here.

Calculated half-lives of both Cys and Met depletion indicated that both the substrate itself and soil properties are strong factors influencing the rates of amino acid turnover in soil solution. On the one hand, Cys removal from soils collected from lower altitude, more

productive sites (e.g. site 1) was twice as fast as that from high altitude, less productive sites (e.g. site 5), with the same true for Met. This is supported by previous work showing that amino acids and peptide turnover rates were much faster in higher productivity grassland than in low productivity grasslands (Farrell, Macdonald, *et al.*, 2014; Wilkinson, Paul W. Hill, *et al.*, 2014). On the other hand, the substrate itself is also an important factor affecting amino acid turnover rate. For all five soils, the half-life of Met depletion was greater than that of Cys depending on soil conditions. This is consistent with previous studies indicating that, Met, a “resistant” amino acid, was degraded slower than Cys in soil (Greenwood and Lees, 1960).

The range of Cys and Met concentrations employed here was chosen to reflect those likely to be found in soils. Lower concentrations may be expected to occur after dilution of soil native water with rainfall (0.01 - 0.1 mM; (Werdin-Pfisterer, Kielland and Boone, 2009)), while higher concentrations may occur upon lysis of root cell in the rhizosphere or the addition of amino acid-based fertilisers (1 - 100 mM). Under all concentrations, the depletion rate of Cys and Met from soil solution was extremely high. Our results suggested that under low Cys and Met concentrations (0.01, 0.05 and 0.1 mM), soil microbial uptake of amino acids occurred via a single carrier system, while a second uptake pathway became important when substrate was added at higher concentrations (1 mM, 10 mM, 100 mM). The concentration of low molecular weight S-containing amino acid at which the secondary transport system becomes important here (1 mM) is higher than that found in a previous study (Paul W. Hill, Farrar and Jones, 2008), in which 0.01 and 0.1 mM was found to be the point where the second transport system became important in the microbial uptake of glucose. It is not clear whether these differences resulted from different experimental conditions or soil conditions.

#### 4.4.2 Dynamics of Cys and Met mineralization by soil microbial community

As Cys or Met are weakly adsorbed in the grassland soil used here (data shown in Fig. 3.2, chapter 3), and very little  $^{14}\text{CO}_2$  evolution was observed in the autoclaved treatment within 48 h during our experiment time-period (Fig. 4.5), we assume that amino acid mineralization from soil solution occurred entirely due to microbial activity. It is likely that mineralization of biological sulphur to inorganic sulphate under the action of microbial community is the rate determining step in the soil sulphur cycle (Meena and Improvement, 2018; Udayana *et al.*, 2021). It is striking that within only one hour of substrate addition (0.01 mM) to soil, soil microbes mineralized >10% of added  $^{14}\text{C}$ -Cys and > 2% of added  $^{14}\text{C}$ -Met, suggesting that soil microbes are severely carbon limited. Overall, the mineralization rates of Cys and Met by the soil microbial community were extremely rapid, but much slower than substrate uptake from soil solution, indicating that the uptake and mineralization of cysteine and methionine were decoupled in time. We ascribe this lag in mineralization to the need to strip the N and S from the amino acids prior to incorporation of the carbon skeletons into respiratory cycles. Across most sites,  $^{14}\text{CO}_2$  evolution showed a biphasic pattern, which coincides with those reported previously for other low MW substrates (Paul W. Hill, Farrar and Jones, 2008; Wilkinson, Paul W. Hill, *et al.*, 2014). However, with a similar pool size in general,  $^{14}\text{C}$  allocated to the rapid mineralization phase was respired much faster than that of the secondary slower pool (by two orders of magnitude). This supports the model that the second pool is due to the slow turnover of  $^{14}\text{C}$  incorporated into the microbial biomass (e.g. from meosfaunal grazing).

Across all five grassland sites, when added at 0.01 mM the proportion of added  $^{14}\text{C}$ -Cys recovered as  $^{14}\text{CO}_2$  was higher than 50%, while the reverse was true for  $^{14}\text{C}$ -Met. These results suggested that Cys is a more readily available substrate for soil microbes (i.e. anabolic processes), hence more Cys was broken down in the initial phase, while more Met was allocated to anabolic pools. When added at 100 mM, the proportion of added  $^{14}\text{C}$ -Cys released

as  $^{14}\text{CO}_2$  following rapid substrate mineralization was low (3.5%, Table 4.5), the remainder was cycled through microbial biomass before mineralization, emphasizing the importance of microbial biomass cycle in the process of microbial substrate mineralization.

#### **4.5 Conclusions**

In all five grassland soils, the microbial depletion of added  $^{14}\text{C}$ -Cys and Met (added at a concentration of 0.01 mM) from soil solution was extremely rapid. However, although our experiments were carried out on soil shortly after collection from the field (less than 48h), due to the fact that we had removed plant roots, which have been proved to contribute to nutrient cycling (Cheng and Kuzyakov, 2005; Azcón-Aguilar and Barea, 2015), the actual turnover rates of Cys and Met in situ could be even faster. Therefore, although Cys and Met may represent only a minor component of DON and DOS pools in soil, the importance of Cys and Met for soil microbes and plant nutrition may have been underestimated due to its very fast turnover and replenishment rates in soil. We assume it's the cleavage of proteins into LMW DOS compounds rather than the uptake of amino acids from soil that limits the soil organic S cycle.

#### **4.6 Acknowledgements**

I would like to thank Joseph Cotton and Mark Hughes for soil sample collection, and Sarah Chesworth and Jonathan Roberts for their technical assistance in the laboratory. This research was funded by the China Scholarship Council (201606510012) and Bangor University.

#### **4.7 References**

Azcón-Aguilar, C. and Barea, J. M. (2015) 'Nutrient cycling in the mycorrhizosphere', *Journal*

*of Soil Science and Plant Nutrition*, 15(2), pp. 372–396. doi: 10.4067/s0718-95162015005000035.

Brailsford, F. L. *et al.* (2020) ‘Rapid depletion of dissolved organic sulphur (DOS) in freshwaters’, *Biogeochemistry*. Springer, 149(1), pp. 105–113. doi: 10.1007/s10533-020-00669-4.

Brookes, P. C. *et al.* (1985) ‘Chloroform fumigation and the release of soil nitrogen: a rapid direct extraction method to measure microbial biomass nitrogen in soil’, *Soil biology and biochemistry*. Elsevier, 17(6), pp. 837–842. doi: 10.1016/0038-0717(85)90144-0.

Broughton, R. C.I. *et al.* (2015) ‘Differential acquisition of amino acid and peptide enantiomers within the soil microbial community and its implications for carbon and nitrogen cycling in soil’, *Soil Biology and Biochemistry*. Elsevier Ltd, 88, pp. 83–89. doi: 10.1016/j.soilbio.2015.05.003.

Broughton, R C I *et al.* (2015) ‘Differential acquisition of amino acid and peptide enantiomers within the soil microbial community and its implications for carbon and nitrogen cycling in soil’, *Soil Biology and Biochemistry*. Elsevier, 88, pp. 83–89. doi: 10.1016/j.soilbio.2015.05.003.

Brydges, T. G. and Wilson, R. B. (1990) ‘Acid rain since 1985 – times are changing’, *Proceedings of the Royal Society of Edinburgh. Section B. Biological Sciences*. 2011/12/05. Royal Society of Edinburgh Scotland Foundation, 97, pp. 1–16. doi: 10.1017/S0269727000005261.

Cambier, P. *et al.* (2014) ‘Impact of long-term organic residue recycling in agriculture on soil solution composition and trace metal leaching in soils’, *Science of the total environment*. Elsevier, 499, pp. 560–573. doi: 10.1016/j.scitotenv.2014.06.105.

- Carswell, A. M. *et al.* (2016) ‘Short-term biotic removal of dissolved organic nitrogen (DON) compounds from soil solution and subsequent mineralisation in contrasting grassland soils’, *Soil Biology and Biochemistry*. Elsevier Ltd, 96, pp. 82–85. doi: 10.1016/j.soilbio.2016.01.017.
- Carter, D. O., Yellowlees, D. and Tibbett, M. (2007) ‘Autoclaving kills soil microbes yet soil enzymes remain active’, *Pedobiologia*, 51(4), pp. 295–299. doi: 10.1016/j.pedobi.2007.05.002.
- Ceccotti, S. P. and Messick, D. L. (1997) ‘global review of crop requirements, supply, and environmental impact on nutrient sulphur balance’, *Sulphur metabolism in higher plants: molecular, ecophysiological and nutritional aspects*. Leiden: Backhuys, c1997., pp. 155–163.
- Cheng, W. and Kuzyakov, Y. (2005) ‘Root effects on soil organic matter decomposition’, *Roots and soil management: interactions between roots and the soil. Agronomy monograph No. 48*, (48), pp. 119–144. doi: 10.2134/agronmonogr48.c7.
- David, M. B. and Mitchell, M. J. (1985) ‘Sulfur constituents and cycling in waters, seston, and sediments of an oligotrophic lake’, *Limnology and Oceanography*, 30(6), pp. 1196–1207. doi: 10.4319/lo.1985.30.6.1196.
- Falloon, P. D. and Smith, P. (2000) *Modelling refractory soil organic matter*. Third Edit, *Biology and Fertility of Soils*. Third Edit. Elsevier Inc. doi: 10.1016/B978-0-08-047514-1.50016-0.
- Farrell, M. *et al.* (2011) ‘Seasonal variation in soluble soil carbon and nitrogen across a grassland productivity gradient’, *Soil Biology and Biochemistry*. Elsevier Ltd, 43(4), pp. 835–844. doi: 10.1016/j.soilbio.2010.12.022.

- Farrell, M., Macdonald, L. M., *et al.* (2014) 'Amino acid dynamics across a grassland altitudinal gradient', *Soil Biology and Biochemistry*. Elsevier Ltd, 76(December 2008), pp. 179–182. doi: 10.1016/j.soilbio.2014.05.015.
- Farrell, M., Prendergast-Miller, M., *et al.* (2014) 'Soil microbial organic nitrogen uptake is regulated by carbon availability', *Soil Biology and Biochemistry*. Elsevier Ltd, 77, pp. 261–267. doi: 10.1016/j.soilbio.2014.07.003.
- Fitzgerald, J. W. and Andrew, T. L. (1984) 'Mineralization of methionine sulphur in soils and forest floor layers', *Soil Biology and Biochemistry*, 16(6), pp. 565–570. doi: 10.1016/0038-0717(84)90073-7.
- Fitzgerald, J. W., Hale, D. D. and Swank, W. T. (1988) 'Sulphur-containing amino acid metabolism in surface horizons of a hardwood forest', *Soil Biology and Biochemistry*, 20(6), pp. 825–831. doi: 10.1016/0038-0717(88)90088-0.
- Fowler, D. *et al.* (2007) 'Long Term Trends in Sulphur and Nitrogen Deposition in Europe and the Cause of Non-linearities BT - Acid Rain - Deposition to Recovery', in Brimblecombe, P. *et al.* (eds). Dordrecht: Springer Netherlands, pp. 41–47. doi: 10.1007/978-1-4020-5885-1\_5.
- Freney, J. R., Spencer, K. and Jones, M. B. (1978) 'The diagnosis of sulphur deficiency in wheat', *Australian Journal of Agricultural Research*. CSIRO, 29(4), pp. 727–738. doi: 10.1071/AR9780727.
- Freney, J. R., Stevenson, F. J. and Beavers, A. H. (1972) 'Sulfur-containing amino acids in soil hydrolysates', *Soil Science*. LWW, 114(6), pp. 468–476. doi: 10.1016/j.soilbio.2015.11.016.
- Glanville, H. *et al.* (2012) 'Soil Biology & Biochemistry Mineralization of low molecular



- weight carbon substrates in soil solution under laboratory and field conditions', *Soil Biology and Biochemistry*. Elsevier Ltd, 48, pp. 88–95. doi: 10.1016/j.soilbio.2012.01.015.
- Glanville, H. C. *et al.* (2016) 'Combined use of empirical data and mathematical modelling to better estimate the microbial turnover of isotopically labelled carbon substrates in soil', *Soil Biology and Biochemistry*. Elsevier, 94, pp. 154–168. doi: 10.1016/j.soilbio.2015.11.016.
- Grant, R. F., Juma, N. G. and McGill, W. B. (1993) 'Simulation of carbon and nitrogen transformations in soil: Microbial biomass and metabolic products', *Soil Biology and Biochemistry*, 25(10), pp. 1331–1338. doi: 10.1016/0038-0717(93)90047-F.
- Greenfield, L. M. *et al.* (2020) 'Is soluble protein mineralisation and protease activity in soil regulated by supply or demand?', *Soil Biology and Biochemistry*. Elsevier, 150, p. 108007. doi: 10.1016/j.soilbio.2020.108007.
- Greenwood, D. J. and Lees, H. (1960) 'Studies on the decomposition of amino acids in soils', *Plant and Soil*, 12(1), pp. 69–80. doi: 10.1007/bf01377762.
- Hill, P. W., Farrar, J. F. and Jones, D. L. (2008) 'Decoupling of microbial glucose uptake and mineralization in soil', *Soil Biology and Biochemistry*, 40(3), pp. 616–624. doi: 10.1016/j.soilbio.2007.09.008.
- Jones, D. L. *et al.* (2018) 'Moisture activation and carbon use efficiency of soil microbial communities along an aridity gradient in the Atacama Desert', *Soil Biology and Biochemistry*. Elsevier, 117, pp. 68–71. doi: 10.1016/j.soilbio.2017.10.026.
- Jones, D. L. and Kielland, K. (2012) 'Amino acid, peptide and protein mineralization dynamics in a taiga forest soil', *Soil Biology and Biochemistry*. Elsevier Ltd, 55(3), pp. 60–69.

doi: 10.1016/j.soilbio.2012.06.005.

Jones, D. L., Owen, A. G. and Farrar, J. F. (2002) 'Simple method to enable the high resolution determination of total free amino acids in soil solutions and soil extracts', *Soil Biology and Biochemistry*. Elsevier, 34(12), pp. 1893–1902. doi: 10.1016/S0038-0717(02)00203-1.

Jones, D. L. and Shannon, D. (1999) 'Mineralization of Amino Acids Applied to Soils Impact of Soil Sieving, Storage, and Inorganic Nitrogen Additions', *Soil Science Society of America Journal*, 63(5), pp. 1199–1206. doi: 10.2136/sssaj1999.6351199x.

Kertesz, M. A. (2000) 'Riding the sulfur cycle—metabolism of sulfonates and sulfate esters in Gram-negative bacteria', *FEMS Microbiology Reviews*. Blackwell Publishing Ltd Oxford, UK, 24(2), pp. 135–175. doi: 10.1016/S0168-6445(99)00033-9.

Kölling, C. and Prietzel, J. (1995) 'Correlations between nitrate and sulfate in the soil solution of disturbed forest ecosystems', *Biogeochemistry*. Springer, 31(3), pp. 121–138. doi: 10.1007/bf00004045.

Ma, Q. *et al.* (2020) 'Soil carbon, nitrogen, and sulphur status affects the metabolism of organic S but not its uptake by microorganisms', *Soil Biology and Biochemistry*. Elsevier, 149, p. 107943. doi: 10.1016/j.soilbio.2020.107943.

McGill, W. B. and Cole, C. V. (1981) 'Comparative aspects of cycling of organic C, N, S and P through soil organic matter', *Geoderma*. Elsevier, 26(4), pp. 267–286. doi: 10.1016/0016-7061(81)90024-0.

McGrath, S. P. and Zhao, F. J. (1995) 'A risk assessment of sulphur deficiency in cereals using soil and atmospheric deposition data', *Soil Use and Management*, 11(3), pp. 110–114. doi: 10.1111/j.1475-2743.1995.tb00507.x.

- Meena, V. S. and Improvement, C. (2018) *Role of Rhizospheric Microbes in Soil, Role of Rhizospheric Microbes in Soil*. doi: 10.1007/978-981-13-0044-8.
- Miranda, K. M., Espey, M. G. and Wink, D. A. (2001) 'A rapid, simple spectrophotometric method for simultaneous detection of nitrate and nitrite', *Nitric Oxide - Biology and Chemistry*, 5(1), pp. 62–71. doi: 10.1006/niox.2000.0319.
- Moe, L. A. (2013) 'Amino acids in the rhizosphere: From plants to microbes', *American Journal of Botany*, 100(9), pp. 1692–1705. doi: 10.3732/ajb.1300033.
- Mulvaney, R. L. (1996) 'Nitrogen—inorganic forms', *Methods of soil analysis: Part 3 Chemical methods*. Wiley Online Library, 5, pp. 1123–1184. doi: 10.2136/sssabookser5.3.c38.
- Murphy, J. and Riley, J. P. (1962) 'A modified single solution method for the determination of phosphate in natural waters', *Analytica chimica acta*. Elsevier, 27, pp. 31–36. doi: 10.1016/S0003-2670(00)88444-5.
- Quevauviller, P. (2007) *Method performance studies for speciation analysis*. Royal Society of Chemistry. doi: 10.1039/9781847551405.
- Rowell, D. L. (2014) *Soil science: Methods & applications*. Routledge.
- Ruiz-Herrera, J. and Starkey, R. L. (1969) 'Dissimilation of methionine by a demethiolase of *Aspergillus* species.', *Journal of Bacteriology*, 99(3), pp. 764–770. doi: 10.1128/jb.99.3.764-770.1969.
- Sanchez-Arenillas, M., Galvez-Martinez, S. and Mateo-Marti, E. (2017) 'Sulfur amino acids and alanine on pyrite (100) by X-ray photoemission spectroscopy: Surface or molecular role?', *Applied Surface Science*. Elsevier, 414, pp. 303–312. doi: 10.1016/j.apsusc.2017.04.049.

- Scott, N. M., Bick, W. and Anderson, H. A. (1981) 'The measurement of sulphur-containing amino acids in some scottish soils', *Journal of the Science of Food and Agriculture*, 32(1), pp. 21–24. doi: 10.1002/jsfa.2740320105.
- Scow, K. M., Simkins, S. and Alexander, M. (1986) 'Kinetics of mineralization of organic compounds at low concentrations in soil', *Applied and Environmental Microbiology*, 51(5), pp. 1028–1035. doi: 10.1128/aem.51.5.1028-1035.1986.
- Sohn, M. and Ho, C. T. (1995) 'Ammonia Generation during Thermal Degradation of Amino Acids', *Journal of Agricultural and Food Chemistry*, 43(12), pp. 3001–3003. doi: 10.1021/jf00060a001.
- Udayana, K. S. *et al.* (2021) ' Sulphur: A boon in agriculture', *The Pharma Innovation Journal*, 10(7), pp. 912–921. Available at: <http://www.thepharmajournal.com>.
- Vinolas, L. C., Vallejo, V. R. and Jones, D. L. (2001) 'Control of amino acid mineralization and microbial metabolism by temperature', *Soil Biology and Biochemistry*, 33(7–8), pp. 1137–1140. doi: 10.1016/S0038-0717(00)00243-1.
- Voroney, R. P., Brookes, P. C. and Beyaert, R. . (2007) 'Soil Microbial Biomass C,N,P and S', in *Soil Sampling and Methods of Analysis, Second Edition*. CRC Press Boca Raton, FL, pp. 637–652. doi: 10.1201/9781420005271.ch49.
- Werdin-Pfisterer, N. R., Kielland, K. and Boone, R. D. (2009) 'Soil amino acid composition across a boreal forest successional sequence', *Soil Biology and Biochemistry*. Elsevier Ltd, 41(6), pp. 1210–1220. doi: 10.1016/j.soilbio.2009.03.001.
- Wilkinson, A., Hill, Paul W, *et al.* (2014) 'Rapid microbial uptake and mineralization of amino acids and peptides along a grassland productivity gradient', *Soil Biology and Biochemistry*. Elsevier, 72, pp. 75–83. doi: 10.1016/j.soilbio.2014.01.026.

- Wilkinson, A., Hill, Paul W., *et al.* (2014) 'Rapid microbial uptake and mineralization of amino acids and peptides along a grassland productivity gradient', *Soil Biology and Biochemistry*. Elsevier Ltd, 72, pp. 75–83. doi: 10.1016/j.soilbio.2014.01.026.
- Withers, E. *et al.* (2020) 'Use of untargeted metabolomics for assessing soil quality and microbial function', *Soil Biology and Biochemistry*. Elsevier, 143, p. 107758. doi: 10.1016/j.soilbio.2020.107758.
- Zehr, J. P., Axler, R. P. and Goldman, C. R. (1985) 'Heterotrophic mineralization of amino acid nitrogen in subalpine Castle Lake, California', *Marine Chemistry*, 16(4), pp. 343–350. doi: 10.1016/0304-4203(85)90055-6.
- Zhao, F. J., Wu, J. and McGrath, S. P. (1996) 'Soil Organic Sulphur and its Turnover', *Humic Substances in Terrestrial Ecosystems*, pp. 467–506. doi: 10.1016/b978-044481516-3/50013-x.
- Zhao, F. and McGrath, S. P. (1994) 'Extractable sulphate and organic sulphur in soils and their availability to plants', *Plant and Soil*, 164(2), pp. 243–250. doi: 10.1007/BF00010076.

## **Chapter 5**

---

### **Competition for S-containing amino acids (cysteine and methionine) between soil microorganisms and maize roots in the rhizosphere**

## Abstract

The direct uptake of amino acids from soil solution has been proposed as a potential route by which plants can acquire nitrogen (N), particularly in nutrient-limiting ecosystems. The factors regulating the potential acquisition of sulphur-containing amino acids by roots, however, remains poorly understood. Using a  $^{14}\text{C}$ -labelling approach, we independently investigated the short-term (24 h) uptake of two S-containing amino acids, cysteine and methionine, by the soil microbial community and maize plants (*Zea mays* L.). We then studied their relative ability to capture amino acids when in competition with each other. Our results show that individually, both soil microbes and maize plants have a high ability to utilize cysteine and methionine (added at a concentration of 0.1 mM). Soil microbes captured cysteine and methionine at a rate of  $413 \pm 2$  and  $408 \pm 2$  nmol (kg soil) $^{-1}$  h $^{-1}$  respectively, while uptake rates of individual amino acids by maize roots were  $203 \pm 36$  nmol (g root DW) $^{-1}$  h $^{-1}$  for cysteine, and  $192 \pm 31$  nmol (g root DW) $^{-1}$  h $^{-1}$  for methionine. When in competition with each other, the capture of free cysteine and methionine by the maize plants was very low compared to soil microbes, with only  $8.8 \pm 1.6$  of the added amino acid captured by the plant ( $8.8 \pm 1.9\%$  or the total), compared to the rhizosphere microbial community ( $76.9 \pm 8.9\%$ ). In addition, to examine the significance of cysteine and methionine for plant sulphur nutrition in the rhizosphere, we also examined how maize plants and soil microbes vary in their ability to take up and compete for organic ( $^{35}\text{S}$ -cysteine,  $^{35}\text{S}$ -methionine) and inorganic sulphur ( $\text{Na}_2^{35}\text{SO}_4$ ) over a short time scale (24 h) using  $^{35}\text{S}$  tracer techniques. Our results showed that soil microbes could capture 16.9% of added inorganic sulphur, with 30.7% captured by plant roots; while maize plants could take up less than 10% of added amino acid- $^{35}\text{S}$ , over 50% was recovered in the microbial biomass. These results, however, indicate that all three experimentally added sulphur forms contributed available sulphur for plant uptake, although rhizosphere microbial community showed higher capacity for capture of free cysteine and methionine in the

rhizosphere than maize plants. We suggest that this is a result of relatively high availability of inorganic nutrient in soil solution and the lack of transmembrane for amino acids on plant root cells, as well as the rapid turnover of amino acids by soil microbes in the rhizosphere. Our results also show that the addition of inorganic N and S, not C, significantly reduced the uptake of cysteine and methionine by maize plants in the rhizosphere, indicating that amino acid utilization by maize plants in our study could be regulated by inorganic N and S availability. Overall, our results imply that (1) cysteine and methionine are sources of available N and S for maize plants, contributing to total plant N and S demand, (2) amino acid mineralization rates underestimate N and S supply rates to plants, (3) plants may compete better with soil microbes when concentrations of free cysteine and methionine in soil are high.

**Keywords:** Cysteine; Methionine; Dissolved organic sulphur (DOS); Plant-microbial competition; Rhizosphere.

## 5.1 Introduction

Sulphur (S) is an essential nutrient for plants, being required for the biosynthesis of essential amino acids (i.e. cysteine and methionine; Wirtz and Droux, 2005), oligopeptides (e.g. glutathione and phytochelatins; Na and Salt, 2011), vitamins (e.g. biotin, thiamine), enzyme co-factors (e.g. Fe-S sulphur clusters), and a variety of secondary metabolites (e.g. glucosinolates and alliins; Ravilious and Jez, 2012). An adequate supply of S is therefore essential to ensure optimal crop growth, however, there is increasing evidence that S deficiency is becoming more widespread in agricultural systems (Aula *et al.*, 2019). Although S fertilisers are produced relatively cheaply as a by-product from petrochemical processing, S use efficiency in agriculture is typically very low (<25%) (Eriksen, 2009b). There is therefore a need to better understand plant-soil-microbial S cycling to enable us to devise more sustainable agricultural systems. It is generally assumed that plants can only acquire S in an inorganic form



(SO<sub>4</sub><sup>2-</sup>; Prasad and Shivay, 2018). However, there is growing evidence that plants can also take up a range of nutrients (e.g. N, P, Fe) held in low molecular weight (MW) organic forms, whether this applies to S remains largely unknown.

Generally, more than 90% of S held in soil is present in an insoluble organic form (Johnson, 1984). Of this, a large proportion is present in amino acids (e.g. cysteine, methionine) which are contained in proteinaceous material which makes up ca. 20-30% of soil organic matter (SOM; Stevenson, 1982). Soil microorganisms (including bacteria, archaea and fungi) play a major role in releasing plant-available sulphate through the oxidation of organic S (McLaren and Swift, 1977; Michael A. Kertesz and Mirleau, 2004a). Specifically, the release of microbially-derived sulfatases can release SO<sub>4</sub><sup>2-</sup> from outward facing S groups in SOM. In addition, the microbial community can release exo- and endo-peptidases to release S-containing amino acids and oligopeptides into soil solution. In contrast, S embedded within SOM particles or bound to mineral surfaces may not be readily available for plants due to chemical or physical protection (Scherer, 2001).

Observations in Arctic and sub-Antarctic ecosystems have shown that the supply of inorganic N is often insufficient to meet the annual N requirement of many plant species (Smith and Steenkamp, 1992; Kielland, 1994). This has raised awareness that free amino acids and oligopeptides in soil solution, may represent a readily available source of N and C for plants as well as soil microorganisms (Geisseler *et al.*, 2010). Quantitative estimates of the contribution of this type of organic S to plant nutrition, however, remain unknown. There is very good evidence to show that plant roots can take up exogenous amino acids and oligopeptides intact (Abuzinadah *et al.*, 1986; Yamagata and Ae, 1999; Nishizawa and Mori, 2001; Weigelt *et al.*, 2005; Tegeder and Rentsch, 2010; Hill *et al.*, 2011). Traditionally, however, plant roots have been considered poor competitors with soil microbes for organic nutrients in soil (Kuzyakov and Jones, 2006; Løerke *et al.*, 2008). Despite this, roots possess a wide range of H<sup>+</sup>-ATPase

fuelled amino acid and sugar co-transporters which appear to be constitutively expressed (Rentsch, Schmidt and Tegeder, 2007; D L Jones, Nguyen and Finlay, 2009). These have the capacity to take up a selected range of low MW solutes (e.g. amino acids, sugars, peptides, polyamines, urea) but not others (e.g. amino sugars, tricarboxylic acids; Jones et al., 2009; Warren, 2016). It is assumed that the selective uptake and incorporation of N, and presumably S, from amino acids requires less energy in comparison to the assimilation of  $\text{NO}_3^-$ ,  $\text{NH}_4^+$  or  $\text{SO}_4^{2-}$  (Yamagata and Ae, 1999). Others have hypothesized that uptake of amino acids from the rhizosphere is an important way to recapture C lost in root exudation to prevent excessive microbial growth in the rhizosphere and simultaneously reduce nutrient loss to the soil (Jones et al., 2009). Alternatively, these root transporters may be used in root signal transduction cascades and used to regulate developmental processes (Walch-Liu *et al.*, 2006).

Here we focus on the uptake of the two main S-containing amino acids in plants, cysteine and methionine (Noji and Saito, 2003). These compounds represent 70% of the S contained in plant tissues, although they are often only present in low abundance relative to other amino acids (Kumar *et al.*, 2019). The potential importance to crop plants of taking up S-containing amino acids from soil is likely to depend on the level of competition between the root and the rhizosphere microbial community (Cheng and Bledsoe, 2004; Xu *et al.*, 2008). Based on studies of other organic-N containing solutes, we expect that this competition will be particularly strong when concentrations of organic S are low in the soil solution (A. G. Owen and Jones, 2001). This competition may also be influenced by a range of abiotic factors, including the relative availability of inorganic N and S. Our primary aim was therefore to investigate the capacity of maize (*Zea mays* L.) to take up cysteine (Cys) and methionine (Met) and to directly compare these rates to those of the soil microbial community. As S uptake and metabolism has been shown to be influenced by intracellular levels of S-containing amino acids in plants (García et al., 2015; Miranda et al., 2001a; Smith et al., 1997), it also suggests that

there could be a close interplay between inorganic and organic S availability in the rhizosphere and root transport processes. Our secondary aim was therefore to manipulate the availability of C, N and S in the rhizosphere to determine its impact on Met and Cys capture. We hypothesized that in the short term, elevated C availability would promote increased plant capture of Cys and Met due to the provision of an alternative C supply to the microbial community. Further, we predicted that N and S addition would have no effect on Cys and Met uptake if the root amino acid transporters are constitutively expressed.

## **5.2 Materials and methods**

### **5.2.1 Soil sampling and chemical characterisation**

An agricultural soil (Eutric Cambisol) was collected from the Ah horizon (0-10 cm depth) of a temperate *Lolium perenne* L. dominated grassland site located at Abergwyngregyn, Gwynedd, North Wales (53°14'N, 4°01'W). This lowland grassland site is unfertilized, free draining, and lightly grazed by sheep (Fig. 5.1). Properties of the soil are listed in Table 5.1. Approximately 1 kg of soil was collected from three randomly positioned replicate plots, located 2 m apart, within our study site (5 × 5 m<sup>2</sup>). In terms of N availability, the site is characterised as being spatially homogenous (Shaw et al., 2016). The soil was then placed in gas permeable plastic bags and transferred immediately to the laboratory. These samples represented the three independent replicates for all experiments. On return to the laboratory, the crumb structured, sandy clay loam textured soil was sieved to pass 2 mm before being stored at 4 °C until required for experimentation.

Soil moisture content was assessed by oven drying soil at 80 °C. Soil pH and electrical conductivity were analysed in a 1:5 (w/v) soil: deionised water suspension using standard electrodes. To determine the levels of available N, fresh soil (5 g) was shaken (200 rev min<sup>-1</sup>, 15 min) with 0.5 M K<sub>2</sub>SO<sub>4</sub> (25 ml), the suspension centrifuged (10,000 g, 15 min) and the

supernatant retained for analysis. The concentration of  $\text{NH}_4^+$  in the extracts was determined colorimetrically on a Synergy MX micro-plate reader using the salicylic acid method of Mulvaney (1996), while  $\text{NO}_3^-$  was determined colorimetrically using the vanadate procedure of Miranda et al. (2001b). Total free amino acids in the extracts were determined fluorometrically using the *o*-phthaldialdehyde/ $\beta$ -mercaptoethanol procedure of Jones et al. (2002), while total protein was determined colorimetrically according to Jones and Kielland (2002). To determine the levels of available S, soil was shaken ( $200 \text{ rev min}^{-1}$ , 15 min) with distilled water (1:5 w/v). All extracts were centrifuged ( $8000 \text{ g}$ , 10 min),  $0.45 \mu\text{m}$  syringe filtered and frozen at  $-20^\circ\text{C}$  prior to analysis. The concentration of sulphate and other major anions in the extract was determined by ion chromatography (IC; Dionex corporation, ICS 2100, USA; Zhao and McGrath, 1994) according to ISO 10304-1:2009. Total dissolved S (DOS plus inorganic sulphate) and other major cations were determined by inductively coupled plasma-optical emission spectroscopy (ICP-OES; Varian 710ES, Agilent Technologies, USA) according to ISO 11885:2016. Dissolved organic S (DOS) concentrations were calculated from the difference between TDS and inorganic sulphate (David and Mitchell, 1985).

**Table 5.1** Summary of the main properties of the soil used in the experiments. Values represent means  $\pm$  SEM ( $n = 3$ ). Data are presented on a dry soil weight basis.

Parameter	
Altitude (m)	32
Water content <sup>a</sup> (%)	$17.11 \pm 0.17$
pH <sup>b</sup>	$5.92 \pm 0.02$
Electrical conductivity <sup>b</sup> ( $\mu\text{S cm}^{-1}$ )	$73.50 \pm 3.55$
$\text{NH}_4^+\text{-N}^c$ ( $\text{mg kg}^{-1}$ )	$1.14 \pm 0.14$
$\text{NO}_3^-\text{-N}^c$ ( $\text{mg kg}^{-1}$ )	$17.05 \pm 0.33$
Amino acids- $\text{N}^c$ ( $\text{mg N kg}^{-1}$ )	$0.61 \pm 0.15$
Protein- $\text{N}^c$ ( $\text{mg N kg}^{-1}$ )	$2.95 \pm 1.37$
Sulphate- $\text{S}^d$ ( $\text{mg S kg}^{-1}$ )	$4.37 \pm 0.79$
Total dissolved S <sup>e</sup> ( $\text{mg S kg}^{-1}$ )	$31.62 \pm 2.14$

<sup>a</sup>Measured by drying at  $105^\circ\text{C}$  overnight and expressed on a wet weight basis.

<sup>b</sup>Measured in a 1:5 (v/v) soil-to-distilled water extract.

<sup>c</sup>Measured in a 1:5 (w/v) 0.5 M  $\text{K}_2\text{SO}_4$  extract followed by colorimetry analysis via a Synergy-MX microplate reader.

<sup>d</sup>Measured in 1:5 w/v soil: distilled water extracts using an ICS-2011 ion chromatograph.

<sup>e</sup>Measured in 1:5 w/v soil: 0.01 M  $\text{Ca}(\text{H}_2\text{PO}_4)_2$  extracts using a Varian 710ES ICP-OES.



**Fig. 5.1** Detailed photographs of the area of our study.

### 5.2.2 Rhizosphere microbial uptake and mineralization of $^{14}\text{C}$ -Cys and Met

Maize plants were grown in soil-filled microcosms (rhizotubes) as described in Ström *et al.* (2002). The rhizotubes were constructed from nylon tube (250 mm long, 9 mm dia.) which expanded over a 0.5 cm span to a 50 mm long, 20 mm diameter section which was used to hold the seed. Holes (0.5 mm diameter) were pierced at 10 mm intervals down the length of the main rhizotube to ensure aeration. Before the addition of the pre-germinated plant, the microcosms were filled  $15 \pm 0.5$  g of soil to a bulk density of  $1.16 \text{ g cm}^{-3}$  to reflect that in the field. After 7 d of plant growth, all of the soil volume contains primary and secondary roots and was effectively classed as rhizosphere soil. The microcosms were then harvested, the roots removed and the rhizosphere soil retained for further experimentation.

To determine the microbial turnover of Cys and Met in the rhizosphere soil, 0.5 ml of uniformly radiolabelled  $^{14}\text{C}$ -Cys or Met (0.1, 1 or 10 mM; PerkinElmer Inc, Waltham, MA) was added to 5 g of soil contained in a  $50 \text{ cm}^3$  sterile polypropylene tube. This range was chosen to reflect the free amino acid concentrations naturally present in rhizosphere soil solution or which arise from the lysis of a root or microbial cell (Dietz *et al.*, 1990; Jones, Shannon, *et al.*, 2005). After accounting for dilution in the native soil solution, this resulted in an actual final concentration of 0.03, 0.3 or 3 mM in soil solution.

To assess whether the breakdown of these two amino acids by the microbial biomass was affected by the availability of C, inorganic N or inorganic S, a very high concentration of either glucose ( $360 \text{ mg C kg}^{-1}$  soil),  $\text{NH}_4\text{Cl}$  ( $30 \text{ mg N kg}^{-1}$  soil) or  $\text{Na}_2\text{SO}_4$  ( $30 \text{ mg S kg}^{-1}$  soil) was added to the soil surface together with the  $^{14}\text{C}$ -labelled amino acid solution. These levels were chosen based on a ca. 10-fold excess of glucose relative to Met and Cys at their highest addition rate, a C use efficiency of 50% and a microbial C:N:S ratio of 6:1. A control treatment was included where labelled Cys or Met was applied in distilled water alone.

After addition of either  $^{14}\text{C}$ -labelled Cys or Met to the soil surface, 1 M NaOH traps (1 ml) were suspended above the soil to catch any respired  $^{14}\text{CO}_2$ . The tubes were then hermetically sealed and incubated at room temperature ( $20 \pm 1^\circ\text{C}$ ) in the dark. There were three replicates of each treatment. The NaOH traps were replaced after 0.5, 1, 2, 3, 12 and 24 h. The efficiency of the NaOH traps was  $> 98\%$  (as determined by collecting  $^{14}\text{CO}_2$  generated from adding excess 0.1 M HCl to 0.001 M  $\text{NaH}^{14}\text{CO}_3$ ). Alongside the collection of  $^{14}\text{CO}_2$ , the amount of  $^{14}\text{C}$  label remaining in soil solution was determined by destructive harvesting of replicate tubes and extraction with 25 ml of 0.5 M  $\text{K}_2\text{SO}_4$  (200 rev  $\text{min}^{-1}$ , 15 min; Rousk and Jones, 2010). After shaking, the  $\text{K}_2\text{SO}_4$  extracts were centrifuged (14,000 g, 5 min) and the supernatant recovered for analysis. The amount of  $^{14}\text{C}$  in the  $\text{K}_2\text{SO}_4$  extracts and NaOH traps was measured with HiSafe 3<sup>®</sup> scintillation cocktail (PerkinElmer, Waltham, MA) and a Wallac 1404 liquid scintillation counter (Wallac EG&G, UK) with automated quench correction. The amount of  $^{14}\text{C}$  immobilised in the soil microbial biomass was calculated by the difference between  $^{14}\text{C}$  microbial respiration plus that recovered in soil solution and the total amount of  $^{14}\text{C}$  added to the soil (Glanville et al., 2016).

### **5.2.3 Rhizosphere microbial oxidation of Cys and Met in soil using $^{35}\text{S}$ labelling**

A  $^{35}\text{S}$  labelling approach was used to study the oxidation of Cys and Met to sulphate by the rhizosphere microbial community. As described above, 5 g of rhizosphere soil was placed in a 50  $\text{cm}^3$  polypropylene tube, except that 0.5 ml of either a  $^{35}\text{S}$ -labelled Cys or Met solution was added uniformly to the soil surface at three concentrations (0.1, 1 or 10 mM; PerkinElmer Inc.). The samples were then incubated in the dark at room temperature.

After periods of 0.5, 1, 2, 3, 12 and 24 h, the soil was extracted with 25 ml of 0.01 M  $\text{CaCl}_2$ . After centrifugation (10,000 g, 15 min), the supernatant was divided into two portions. On one half, the total  $^{35}\text{S}$  content was measured by liquid scintillation counting as described



above. This fraction contains any  $^{35}\text{S}$ -Met or -Cys remaining in solution, any  $^{35}\text{S}$ -intermediates and  $^{35}\text{SO}_4^{2-}$ . The remaining half was shaken with an equal volume of 0.1 M  $\text{BaCl}_2$  (200 rev  $\text{min}^{-1}$ , 10 min) and centrifuged (14,000 g, 15 min) to remove any inorganic S (as a  $\text{Ba}^{35}\text{SO}_4$  precipitate). The  $^{35}\text{S}$  content of the supernatant was then determined (i.e. this fraction contains any  $^{35}\text{S}$ -Met or -Cys remaining, any  $^{35}\text{S}$ -intermediate products minus  $^{35}\text{SO}_4^{2-}$ ). It was assumed that any  $^{35}\text{S}$  not recovered from the soil was retained in the soil microbial biomass. All  $^{35}\text{S}$  counts were decay-corrected to the start of the incubation (half-life 87.4 d). This approach relies on the low amount of  $\text{SO}_4^{2-}$  sorbed to the solid phase in this soil (Data shown in Fig. 3.1, Chapter 3).

#### **5.2.4 Amino acid uptake by plants under sterile hydroponic conditions ( $^{14}\text{C}$ and $^{35}\text{S}$ )**

Maize seedlings were grown under aseptic conditions in hydroponic culture using a 10% strength Long Ashton nutrient solution (Hewitt, 1952; Smith, Johnston and Cornforth, 1983). After 7 d, the maize plants were transferred to sealed plastic containers containing either  $^{14}\text{C}$ -labelled or  $^{35}\text{S}$  labelled S-containing amino acid (Cys or Met; 0.1 mM; (Jones and Darrah, 1992)).

For  $^{14}\text{C}$  treatments,  $^{14}\text{CO}_2$  respired from the plant was trapped using a 1 M NaOH trap (5 ml) while the amount of  $^{14}\text{C}$  amino acid remaining in the nutrient solution was measured by removing small aliquots over time. After being exposed to the  $^{14}\text{C}$ -amino acid for 24 h, the maize roots were rinsed with unlabelled nutrient solution and oven-dried (80 °C for 24 h). The  $^{14}\text{C}$  content of the maize shoot and root were then measured with an OX-400 Biological Sample Oxidizer (RJ Harvey Instrument Corp., Hillsdale, NJ).  $^{14}\text{C}$  in all the samples was measured by liquid scintillation counting as described above.

For the parallel  $^{35}\text{S}$  treatments, after 24 h, the maize tissues were rinsed with unlabelled nutrient solution and oven-dried (80 °C for 24 h). The amount of  $^{35}\text{S}$  in the plant tissues was



then determined by dissolving the plant tissues in Soluene-350 (40 °C, 4 h; PerkinElmer Inc.) followed by liquid scintillation counting as described above.

### **5.2.5 Competition for Cys and Met between plant and soil microorganisms in the rhizosphere**

Maize seeds were soaked for 12 h in ultrapure water and then distributed evenly on moistened filter paper for 48 h to allow them to germinate at room temperature. Just after germination, uniform seedlings ( $n = 54$ ) were transplanted into individual mesocosms (rhizotubes) as described above. When the maize roots were ca. 12 cm long (7 d after transplantation), a solution containing a  $^{14}\text{C}$ - or  $^{35}\text{S}$ -labelled solution of Cys or Met (0.1 mM; 0.25 ml) was injected at four different points along the rhizotubes through the aeration holes. This enabled a uniform labelling of the soil along the root length.

To determine the influence of nutrient addition on plant capture of free cysteine and methionine from soil solution, an excess of C, N or S (as glucose,  $\text{NH}_4\text{Cl}$  or  $\text{Na}_2\text{SO}_4$ ) was added simultaneously with the  $^{14}\text{C}$ -Met or Cys as described above. Again, the control treatment consisted of Cys or Met applied alone in distilled water. In addition, two higher amino acid concentrations (1 and 10 mM) were chosen to study the influence of free amino acid concentration on plant-microbial competition.

#### *5.2.5.1 Competition for $^{14}\text{C}$ - Cys and Met in the rhizosphere*


Individual mesocosms were then transferred to transparent gas-tight polypropylene containers (11×8 cm base and 27 cm high; Lock & Lock Ltd., Seoul, Republic of Korea; Fig. 5.2) with removable caps. Each container cap had a hole drilled in it (ca. 0.5 cm diameter), to allow physical separation of the plant shoot and root compartments. The shoot was sealed in the hole using non-phytotoxic silicon paste (Swinnen, 1994; Kuzyakov and Siniakina, 2001).

To recover  $^{14}\text{CO}_2$  from the root and shoot compartments, 1 M NaOH traps (5 ml) were placed in each compartment.

24 h after injection of the amino acids, all the rhizotubes were destructively harvested by splitting the rhizo-tubes vertically with a razor blade, allowing the separation of the root, shoot, and soil components. The roots were rinsed with 0.01 M  $\text{CaCl}_2$  for 30 s before being further washed with distilled water to visually remove all adhering soil. The  $^{14}\text{C}$  or  $^{35}\text{S}$  content of the washings was also determined by liquid scintillation counting and added to the soil solution fraction. After harvesting, the rhizosphere soil was divided into two parts: one part was shaken with 0.5 M  $\text{K}_2\text{SO}_4$  (20 min, 200 rev  $\text{min}^{-1}$ ), centrifuged (3800 g, 5 min) and the  $^{14}\text{C}$  content of the supernatant measured as above. The remaining portion was used to determine the  $^{14}\text{C}$  contained in the soil microbial biomass using the  $\text{CHCl}_3$  fumigation- $\text{K}_2\text{SO}_4$  extraction method and an extraction efficiency correction factor of 0.35 (Voroney et al., 2008). Plant shoots and roots were oven-dried (80 °C for 24 h), after which their  $^{14}\text{C}$  content was measured as described above.

#### *5.2.5.2 Competition for $^{35}\text{S}$ -Cys and Met in the rhizosphere*

For the  $^{35}\text{S}$  labelling experiments, the amount of  $^{35}\text{S}$  in the oven-dried plant tissues was determined by dissolving the plant tissues in Soluene-350 (40 °C, 4 h; PerkinElmer Inc.) followed by liquid scintillation counting as described above.  $^{35}\text{S}$  was extracted from the fumigated and unfumigated soil using 0.01 M  $\text{CaCl}_2$  (1:5 w/v) followed by liquid scintillation counting. Differences between the unfumigated and fumigated fraction were assumed to be retained in the soil microbial biomass (after correction using an extraction efficiency value of 0.35).



**Fig. 5.2** Schematic representation of the experimental apparatus showing three-leaf stage maize plant growing down a rhizotube containing grassland soil in which  $^{14}\text{C}$  or  $^{35}\text{S}$  labelled amino acid is injected through pre-drilled holes. The uptake of amino acid into either plant biomass, respiration, or soil microbial biomass and respiration was determined over a 24 h chase period.

### 5.2.6 Statistics and data analysis

To determine the half time ( $t_{1/2}$ ) of cysteine and methionine depletion in soil solution, a single first order exponential decay model was fitted to the experimental data for the 0.1 mM and 1 mM treatments according to:

$$S = y_0 + a_1 \times \exp^{(-b_1 \times t)} \quad (\text{Eqn. 1})$$

Where  $S$  is the  $^{14}\text{C}$  remaining in soil solution,  $b_1$  is the exponential coefficient describing the rate of depletion of  $^{14}\text{C}$  from soil solution by the soil microorganisms,  $a_1$  describes the pool size and  $t$  refers to time. The half time  $t_{1/2}$  of cysteine/methionine soil solution pools can be defined as:

$$t_{1/2} = \ln(2)/b_1 \quad (\text{Eqn. 2})$$

For the 10 mM treatment, the depletion of  $^{14}\text{C}$ -Cys or Met from the soil solution was better described by a double first order exponential decay equation (3).

$$S = a_1 \times \exp^{(-b_1 \times t)} + a_2 \times \exp^{(-b_2 \times t)} \quad (\text{Eqn. 3})$$

The two exponential parts of the equation were assumed to represent independent transport systems (high and low affinity). The half times of each pools were calculated following Equation 2. A double first order kinetic decay equation was also fitted to the  $^{14}\text{CO}_2$  evolution data.

$$C = c_1 \times \exp^{(-d_1 \times t)} + c_2 \times \exp^{(-d_2 \times t)} \quad (\text{Eqn. 4})$$

Where  $c_1$  describes the amount of  $^{14}\text{C}$  allocated to the first mineralization pool and  $d_1$  is the rate constant for  $a_1$ ,  $c_2$  is the proportion partitioned into the second slower pool described by rate constant  $d_2$ . Half time of two mineralization pools  $c_1$  and  $c_2$  can be calculated as described above. Further details of the modelling approach are provided in (H. C. Glanville *et al.*, 2016).

All statistical analyses were carried out in IBM SPSS statistics v25.0 (IBM UK Ltd., Portsmouth, UK). Graphs and curve fitting were produced using SigmaPlot 13.0 (Systat

software Inc., London). To assess the effects of carbon and nutrient amendment on Cys or Met mineralization and depletion rates from soil solution, Pearson's correlations were carried out against the components of the first order exponential decay equation, and half times were derived from these. One-way analysis of variance (ANOVA) followed by Tukey's post-hoc test was used to identify group differences, with the significance level of the P-value being set at  $p \leq 0.05$ . All results are presented in figures and tables as mean  $\pm$  the standard error.

## **5.3 Results**

### **5.3.1 Use of isotopes for tracing Cys and Met utilization by microorganisms in the rhizosphere**

Although mineralization of N from low molecular weight compounds such as amino acids has been studied extensively (Barracough, 1997; O'Dowd, Barracough and Hopkins, 1999; Jones and Kielland, 2012; Zhang *et al.*, 2015; Fujii *et al.*, 2018), few attempts have been made to study S mineralization from amino acids in soils. Therefore, we investigated the biodegradation and release of C and S moieties from cysteine and methionine by soil microorganisms using uniformly  $^{14}\text{C}$ - or  $^{35}\text{S}$ -labelled amino acids. We chose cysteine and methionine as model substrates for this study as the majority of the organic S in plants and microorganisms occur in S-containing amino acids (Giovanelli, Mudd and Datko, 1980; Scott, Bick and Anderson, 1981; Prasad and Shivay, 2018). In addition, cysteine and methionine are likely to be important bioavailable S sources due to their high turnover rates in soils and freshwaters (Andrew and Island, 1984; Fitzgerald, Hale and Swank, 1988; Brailsford *et al.*, 2020). Here, a  $^{35}\text{S}$  labelling technique was applied in this study to track the fate of the  $^{35}\text{S}$  moiety from amino acids in rhizosphere soil, enabling us to determine the rate of formation of  $^{35}\text{S}$ -sulphate and incorporation of  $^{35}\text{S}$  into microbial biomass, while the  $^{14}\text{C}$  labelling technique

allowed us to determine substrate uptake into soil microbial biomass and its subsequent mineralization (CO<sub>2</sub>).

#### 5.3.1.1 <sup>14</sup>C-amino acid depletion from soil solution

<sup>14</sup>C-amino acid removal rates under all three concentrations from soil solution were extremely rapid (Fig. 5.3). For example, the microbial community removed  $93.6 \pm 0.7\%$  and  $92.9 \pm 0.7\%$  of added Cys and Met from soil solution separately within 30 mins when added at 0.1 mM. This fell to  $74.9 \pm 4.4$  and  $74.8 \pm 1.9\%$  at an intermediate amino acid concentration (1 mM), and to  $48.2 \pm 2.1\%$  and  $35.4 \pm 3.6\%$  at the highest amino acid concentration (10 mM). The amount of <sup>14</sup>C remaining in soil solution after 30 mins indicated that cysteine-<sup>14</sup>C was taken up by the microbial community at a similar rate to methionine-<sup>14</sup>C.


However, on a C basis, methionine-C was taken up at a faster rate than cysteine-C due to the greater number of C atoms present. In absolute terms (i.e.  $\mu\text{mol } ^{14}\text{C kg}^{-1} \text{ soil}$ ), the actual amount of cysteine or methionine removed from soil solution increased with amino acid concentration: at the lowest amino acid concentration (0.1 mM), cysteine and methionine-<sup>14</sup>C uptake rates were  $28.1 \pm 0.2$  and  $46.5 \pm 0.3 \mu\text{mol C (kg}^{-1} \text{ soil)}$  respectively within 30 mins of substrate addition. This increased to  $224.9 \pm 13.4$ ,  $373.9 \pm 9.5 \mu\text{mol C (kg}^{-1} \text{ soil)}$ , respectively, when added at 1 mM, and to  $1446.8 \pm 62.2$ ,  $1771.7 \pm 183.1 \mu\text{mol C (kg}^{-1} \text{ soil)}$  when added at 10 mM.

The temporal dynamics of amino acid-<sup>14</sup>C depletion from the soil solution were dependent upon the concentration of amino acid supplied. This is also in agreement with what we found in Chapter 4 (4.3.1.2). The depletion of <sup>14</sup>C-amino acids from soil solution at 0.1 mM and 1 mM was best described by a single first order exponential decay model ( $R^2 > 0.98$  in all cases; Eqn. 1, Table 5.2 and 5.3). We estimated the time necessary for the initial amount of <sup>14</sup>C-amino acids to decrease by 50% (half time) using the exponential decay model. Using this

approach, we estimated the half-life to be  $6.6 \pm 0.2$  min for Cys and  $5.9 \pm 0.3$  min for Met when added at 0.1 mM. This increased to  $12.6 \pm 1.7$  min for Cys and  $10.9 \pm 0.6$  min for Met when added at 1 mM. When added at the highest concentration (10 mM), the depletion of  $^{14}\text{C}$ - amino acids from the soil solution was better described by a double first order exponential decay model ( $R^2 > 0.99$  in all cases; Eqn. 3, Table 5.2 and 5.3), implying that  $^{14}\text{C}$ -Cys and Met were depleted from soil solution in two phases under higher concentrations. Approximately 40% of added Met- $^{14}\text{C}$  (10 mM) was depleted in an initial rapid phase ( $t_{1/2} < 10$  mins), and the remaining 60% was depleted more slowly, with half times up to around 3 d.  $^{14}\text{C}$ -Cys depletion from soil solution (10 mM) also occurred in two phases, with 57% of Cys turning over rapidly ( $t_{1/2} = 13$  minutess), and the remaining formed a pool that turned over much slower ( $t_{1/2} = 24.7$  h).

The concentration of S-containing amino acid at which the secondary transport system becomes important in this chapter (10 mM) is higher than that found in Chapter 4 (4.3.1.2), in which 1mM was found to be the point where the second transport system became important in the microbial uptake of Cys and Met.

At all three amino acid concentrations, exogenous addition of N as  $\text{NH}_4\text{Cl}$ , S as  $\text{Na}_2\text{SO}_4$  or C as glucose exerted a significant influence on the depletion rate of methionine ( $p < 0.05$ ). When added at higher concentration (10 mM), addition of C and S, not N, increased the half-time for both the rapid and slow uptake phases ( $p < 0.05$ ), indicating microbial uptake of cysteine is driven by the microbial demand for organic C to provide energy, as well as for organic sulphur to maintain microbial growth.



**Fig. 5.3** Depletion of  $^{14}\text{C}$ -labelled cysteine or methionine and subsequent incorporation of the  $^{14}\text{C}$  amino acid-C into microbial biomass or respiration ( $^{14}\text{CO}_2$ ) as affected by the addition of excess C, N and S over a 24 h-incubation period at three amino acid concentrations (0.1 mM, 1 mM, 10 mM). Values represent means  $\pm$  SEM,  $n = 3$ .



**Table 5.2** Kinetic coefficients of single (0.1 mM and 1 mM) and double (10 mM) first order curves describing the depletion of  $^{14}\text{C}$ -Cys from soil solution over a 24 h-incubation period in the presence of an excess of C (+ Glucose), inorganic N (+ N) or sulphate (+ S). Values represent means  $\pm$  SEM ( $n = 3$ ).  $a_1$  and  $a_2$  are estimated pool sizes for fast and slow substrate uptake routes,  $b_1$  and  $b_2$  are the rate constants for the fast and slow uptake routes, respectively. Half-time values for each pool are calculated according to Eqn. 2.

Substrate		Cysteine						
		$y_0$	$c_1$	$d_1$	$c_2$	$d_2$	$c_1\ t_{1/2}\ (\text{min})$	$c_2\ t_{1/2}\ (\text{h})$
0.1 mM	None	$2.2 \pm 0.5$	$97.7 \pm 1.1$	$6.3 \pm 0.5$			$6.6 \pm 0.2\text{b}$	
	+Glucose	$2.9 \pm 0.6$	$97.0 \pm 1.6$	$7.6 \pm 1.5$			$5.4 \pm 0.2\text{c}$	
	+N	$3.1 \pm 0.8$	$96.8 \pm 1.9$	$5.8 \pm 0.7$			$7.1 \pm 0.2\text{ab}$	
	+S	$4.4 \pm 1.0$	$95.5 \pm 2.4$	$5.8 \pm 0.9$			$7.1 \pm 0.4\text{a}$	
1 mM	None	$9.1 \pm 2.1$	$90.7 \pm 4.7$	$3.3 \pm 0.5$			$12.6 \pm 1.7$	
	+Glucose	$10.6 \pm 1.7$	$89.3 \pm 4.0$	$3.5 \pm 0.5$			$11.9 \pm 1.6$	
	+N	$14.3 \pm 3.2$	$85.5 \pm 7.6$	$3.5 \pm 1.0$			$11.8 \pm 1.2$	
	+S	$12.7 \pm 2.6$	$87.1 \pm 6.2$	$3.5 \pm 0.8$			$11.6 \pm 1.2$	
10 mM	None		$57.6 \pm 4.4$	$3.2 \pm 0.7$	$42.2 \pm 2.8$	$0.03 \pm 0.01$	$12.9 \pm 1.7\text{b}$	$24.7 \pm 1.7\text{c}$
	+Glucose		$59.9 \pm 3.3$	$2.5 \pm 0.3$	$32.8 \pm 2.2$	$0.02 \pm 0.01$	$16.3 \pm 1.5\text{a}$	$27.7 \pm 1.9\text{c}$
	+N		$52.4 \pm 2.9$	$2.7 \pm 0.3$	$47.4 \pm 1.8$	$0.01 \pm 0.01$	$15.1 \pm 1.3\text{ab}$	$46.3 \pm 5.9\text{a}$
	+S		$55.5 \pm 2.3$	$2.6 \pm 0.3$	$44.3 \pm 1.5$	$0.02 \pm 0.01$	$15.9 \pm 2.6\text{a}$	$36.8 \pm 5.4\text{b}$

**Table 5.3** Kinetic coefficients of single (0.1 mM and 1 mM) and double (10 mM) first order curves describing the depletion of  $^{14}\text{C}$  labelled Met from soil solution over a 24 h incubation period in the presence of an excess of C (+ Glucose), inorganic N (+ N) or sulphate (+ S). Values represent means  $\pm$  SEM ( $n = 3$ ).  $a_1$  and  $a_2$  are estimated pool sizes for fast and slow substrate uptake routes,  $b_1$  and  $b_2$  are the rate constants for the fast and slow uptake routes, respectively. Half-time values for each pool are calculated according to equation 2. Values represent means  $\pm$  standard deviation,  $n = 3$ .

Substrate		Methionine						
		$y_0$	$c_1$	$d_1$	$c_2$	$d_2$	$c_1 t_{1/2}$ (min)	$c_2 t_{1/2}$ (h)
0.1 mM	None	$4.1 \pm 0.6$	$95.8 \pm 1.6$	$6.9 \pm 1.2$			$5.9 \pm 0.3$	
	+ G	$4.5 \pm 0.8$	$95.4 \pm 2.1$	$6.5 \pm 1.2$			$6.3 \pm 0.4$	
	+ N	$4.7 \pm 0.6$	$95.1 \pm 1.5$	$6.4 \pm 0.7$			$6.4 \pm 0.4$	
	+ S	$3.1 \pm 0.8$	$95.6 \pm 1.9$	$5.8 \pm 0.7$			$5.4 \pm 0.3$	
1 mM	None	$13.3 \pm 2.4$	$86.3 \pm 7.4$	$3.8 \pm 0.9$			$10.9 \pm 0.6$	
	+ G	$15.4 \pm 3.2$	$84.3 \pm 7.4$	$3.4 \pm 0.9$			$12.3 \pm 1.2$	
	+ N	$15.4 \pm 3.4$	$84.3 \pm 7.9$	$3.6 \pm 1.1$			$11.5 \pm 0.2$	
	+ S	$13.4 \pm 2.4$	$86.5 \pm 5.7$	$3.9 \pm 0.9$			$10.5 \pm 0.3$	
10 mM	None		$37.7 \pm 0.9$	$5.2 \pm 0.7$	$62.6 \pm 0.6$	$0.01 \pm 0.01$	$7.2 \pm 4.8$	$64.7 \pm 10.8$
	+ G		$32.8 \pm 1.1$	$4.1 \pm 0.4$	$61.7 \pm 0.6$	$0.01 \pm 0.01$	$8.6 \pm 6.8$	$68.2 \pm 11.8$
	+ N		$39.8 \pm 1.5$	$3.2 \pm 0.3$	$60.2 \pm 0.9$	$0.01 \pm 0.01$	$12.3 \pm 3.4$	$53.3 \pm 6.7$
	+ S		$39.7 \pm 2.0$	$4.9 \pm 1.1$	$60.2 \pm 1.1$	$0.01 \pm 0.01$	$8.3 \pm 3.5$	$52.0 \pm 7.7$

### 5.3.1.2 $^{14}\text{C}$ -Amino acid mineralization in soil

The mineralization of  $^{14}\text{C}$ -labeled Cys and Met to  $^{14}\text{CO}_2$  is shown in Fig. 5.1. Following addition of  $^{14}\text{C}$ -labeled amino acids to the soil, there was an initial rapid phase of  $^{14}\text{CO}_2$  evolution followed by a secondary slower phase of evolution (Fig. 5.3).  $^{14}\text{C}$ - $\text{CO}_2$  evolution data fitted well to a first order double exponential model when added at all three concentrations (0.1 mM, 1 mM and 10 mM;  $R^2 > 0.99$ ; Eqn. 4; Fig.5.3). The biphasic exponential dynamics of  $^{14}\text{CO}_2$  evolution from cysteine and methionine were similar to those previously reported for other amino acids (Wilkinson, Paul W. Hill, *et al.*, 2014). The exponential coefficients and half-times ( $t_{1/2}$ ) of amino acid mineralization are presented in Table 5.4 and 5.5.

Less than 40% of the added amino acid- $^{14}\text{C}$  was allocated to the initial rapid mineralization pool ( $c_1$ ) among all treatments, with the half-times for the turnover of this pool ranging from ca. 1.1 to 38.7 h. In contrast, the slow turnover pool ( $c_2$ ) accounted for 51.4 – 91.1% of the added  $^{14}\text{C}$  and had a much slower half time, ranging from ca. 0.5 to 1.8 d following amino acid addition. For all three amino acid concentrations used in this study, the addition of excess C as glucose significantly enhanced the partitioning of amino acid-C into catabolic and anabolic processes ( $p < 0.05$ ).

**Table 5.4** Coefficients of double (0.1 and 1 mM) and single (10 mM) first order curve fits to depletion of  $^{14}\text{C}$ -cysteine from soil by microbial mineralization over a 24 h-incubation period in the presence of an excess of C (+ Glucose), inorganic N (+ N) or sulphate (+ S).  $c_1$  and  $c_2$  are estimated pool sizes for fast and slow phases of mineralization,  $d_1$  and  $d_2$  are the rate constants for the fast and slow phases of mineralization, respectively. Half-time values for each pool are calculated according to equation 2. Values represent means  $\pm$  standard deviation,  $n = 3$ .

Substrate		$c_1$	$d_1$	$c_2$	$d_2$	$c_1$ $t_{1/2}$ (h)	$c_2$ $t_{1/2}$ (d)
0.1 mM Cysteine	None	$19.8 \pm 1.2$	$0.5 \pm 0.1$	$81.3 \pm 1.4$	$0.04 \pm 0.003$	$1.3 \pm 0.1$	$0.8 \pm 0.1$
	+ G	$27.2 \pm 5.7$	$0.4 \pm 0.1$	$74.7 \pm 5.4$	$0.06 \pm 0.01$	$1.7 \pm 0.3$	$0.5 \pm 0.1$
	+ N	$13.6 \pm 4.4$	$0.7 \pm 0.2$	$87.3 \pm 4.6$	$0.03 \pm 0.004$	$1.1 \pm 0.4$	$1.0 \pm 0.1$
	+ S	$15.6 \pm 6.5$	$0.7 \pm 0.2$	$85.0 \pm 6.8$	$0.02 \pm 0.003$	$1.1 \pm 0.3$	$1.2 \pm 0.2$
1 mM Cysteine	None	$18.3 \pm 1.5$	$0.6 \pm 0.1$	$82.5 \pm 1.5$	$0.03 \pm 0.003$	$1.3 \pm 0.1$	$1.0 \pm 0.1$
	+ G	$40.0 \pm 10.0$	$0.3 \pm 0.1$	$61.3 \pm 10.1$	$0.02 \pm 0.003$	$2.3 \pm 0.5$	$1.3 \pm 0.2$
	+ N	$9.0 \pm 2.9$	$0.7 \pm 0.3$	$91.1 \pm 3.1$	$0.02 \pm 0.003$	$1.0 \pm 0.4$	$1.4 \pm 0.2$
	+ S	$12.0 \pm 5.1$	$0.7 \pm 0.3$	$88.6 \pm 4.8$	$0.02 \pm 0.004$	$1.2 \pm 0.5$	$1.5 \pm 0.3$
10 mM Cysteine	None	$3.5 \pm 1.2$	$0.7 \pm 0.2$	$96.8 \pm 1.1$	$0.02 \pm 0.001$	$1.1 \pm 0.4$	$1.3 \pm 0.1$
	+ G	$14.4 \pm 23.0$	$0.6 \pm 0.4$	$85.8 \pm 23.4$	$0.03 \pm 0.001$	$6.8 \pm 12.0$	$1.1 \pm 0.05$
	+ N	$36.4 \pm 22.7$	$0.2 \pm 0.4$	$63.2 \pm 23.0$	$0.02 \pm 0.001$	$32.4 \pm 21.1$	$1.8 \pm 0.1$
	+ S	$47.9 \pm 0.2$	$0.02 \pm 0.001$	$51.4 \pm 0.1$	$0.02 \pm 0.001$	$38.7 \pm 0.9$	$1.6 \pm 0.04$

**Table 5.5** Coefficients of double (0.1 and 1 mM) and single (10 mM) first order curve fits to depletion of  $^{14}\text{C}$ -methionine from soil by microbial mineralization over a 24 h incubation period in the presence of an excess of C (+ Glucose), inorganic N (+ N) or sulphate (+ S).  $c_1$  and  $c_2$  are estimated pool sizes for fast and slow phases of mineralization,  $d_1$  and  $d_2$  are the rate constants for the fast and slow phases of mineralization, respectively. Half-time values for each pool are calculated according to equation 2. Values represent means  $\pm$  standard deviation,  $n = 3$ .

Substrate		$c_1$	$d_1$	$c_2$	$d_2$	$c_1 t_{1/2}$ (h)	$c_2 t_{1/2}$ (d)
0.1 mM Methionine	None	$10.4 \pm 0.7$	$0.8 \pm 0.1$	$90.6 \pm 0.7$	$0.02 \pm 0.003$	$0.9 \pm 0.1$	$1.4 \pm 0.2$
	+ G	$13.1 \pm 0.8$	$0.8 \pm 0.1$	$88.3 \pm 0.7$	$0.03 \pm 0.003$	$0.9 \pm 0.1$	$0.9 \pm 0.1$
	+ N	$9.6 \pm 1.7$	$0.6 \pm 0.1$	$91.4 \pm 1.9$	$0.02 \pm 0.002$	$1.1 \pm 0.2$	$1.8 \pm 0.2$
	+ S	$9.9 \pm 2.2$	$0.5 \pm 0.1$	$91.0 \pm 2.1$	$0.02 \pm 0.001$	$1.3 \pm 0.2$	$1.9 \pm 0.1$
1 mM Methionine	None	$8.0 \pm 0.8$	$0.6 \pm 0.1$	$92.9 \pm 0.6$	$0.01 \pm 0.001$	$1.3 \pm 0.2$	$2.3 \pm 0.1$
	+ G	$13.8 \pm 2.8$	$0.2 \pm 0.04$	$86.8 \pm 2.8$	$0.02 \pm 0.002$	$4.2 \pm 1.1$	$1.9 \pm 0.2$
	+ N	$7.9 \pm 4.6$	$0.4 \pm 0.2$	$92.9 \pm 4.4$	$0.01 \pm 0.003$	$1.6 \pm 0.6$	$3.2 \pm 1.3$
	+ S	$19.0 \pm 7.7$	$0.2 \pm 0.04$	$82.2 \pm 7.4$	$0.004 \pm 0.005$	$2.9 \pm 0.6$	$7.8 \pm 6.5$
10 mM Methionine	None	$47.3 \pm 0.3$	$0.008 \pm 0.0005$	$53.2 \pm 0.2$	$0.008 \pm 0.0005$	$82.9 \pm 2.2$	$3.5 \pm 0.1$
	+ G	$48.4 \pm 0.4$	$0.01 \pm 0.0008$	$52.6 \pm 0.2$	$0.01 \pm 0.0008$	$58.7 \pm 3.6$	$2.4 \pm 0.1$
	+ N	$46.8 \pm 0.4$	$0.006 \pm 0.0005$	$53.6 \pm 0.3$	$0.006 \pm 0.0005$	$102.5 \pm 6.3$	$4.3 \pm 0.3$
	+ S	$46.5 \pm 0.3$	$0.007 \pm 0.0005$	$53.8 \pm 0.2$	$0.007 \pm 0.0005$	$98.4 \pm 5.7$	$4.1 \pm 0.2$

### 5.3.1.3 Incorporation of $^{14}\text{C}$ -amino acids into microbial biomass

Within three hours of substrate addition, the majority of C from cysteine and methionine was taken up into the microbial biomass among all amendments, indicating that soil microorganisms can rapidly assimilate amino acids, even in the presence of readily available C, N and S (Fig. 5.3). The amount of amino acid-C taken up by the soil microbial biomass declined rapidly as the substrate was depleted from soil solution. By the end of our incubation,  $9.5 \pm 0.9$ ,  $25.1 \pm 1.9 \mu\text{mol } ^{14}\text{C kg}^{-1}$  soil (0.1 mM),  $111 \pm 12$ ,  $312 \pm 12 \mu\text{mol } ^{14}\text{C kg}^{-1}$  soil (1 mM),  $1014 \pm 32$ ,  $1665 \pm 89 \mu\text{mol } ^{14}\text{C kg}^{-1}$  soil (10 mM) of Cys and Met remained in the soil microbial biomass.

When added at lower concentrations (0.1 mM, 1mM), more Met was recovered in the microbial biomass than Cys by the end of incubation period ( $p < 0.05$ ). This difference proved non-significant when added at a higher concentration (10 mM). It is also worth noting that for all three concentrations we used in the study, the addition of excess C as glucose generally reduced the amount of amino acid-C retained by the soil microbial biomass ( $p < 0.05$ ), while the addition of N as  $\text{NH}_4\text{Cl}$  or S as  $\text{Na}_2\text{SO}_4$  generally increased C allocation to this pool ( $p < 0.05$ ) (except for 1 mM Cys in the presence of C and for 10 mM Cys in the presence of N or S as these proved non-significant,  $p > 0.05$ ).


### 5.3.2 Amino acid- $^{35}\text{S}$ utilization by soil microorganisms in the rhizosphere

Microbial biomass sulphur is an important pool in soil (Heinze *et al.*, 2021). Fig. 5.4 shows rapid accumulation of  $^{35}\text{S}$  into the microbial biomass and mineralization of  $^{35}\text{S}$  from amino acids due to the action of the microbial community. When added at a concentration of 0.1 mM, 73.2%, 85.9% of the  $^{35}\text{S}$  derived from Cys or Met was incorporated into the microbial biomass by the end of incubation. This decreased to 66.8%, 75.8% when added at 1 mM, and to 44.6%, 40.1% when added at 10 mM. It could be highly likely that cysteine and methionine

were taken up intact by the microbial biomass as a closer amount of  $^{14}\text{C}$  and  $^{35}\text{S}$  was recovered in the microbial biomass after 30 mins. These results indicate that mineralization of cysteine and methionine was biotically mediated, which is controlled by the microbial requirement for energy and C skeletons for the maintenance and growth of microorganisms, with sulphate released as a by-product of S-amino acid turnover inside the cell.

Once taken up into the microbial biomass, excess S is rapidly released from the internal transformation of cysteine and methionine as  $\text{SO}_4^{2-}$ . When added at a concentration of 0.1 mM, ca. 12.4%, 7.2% of the total added  $^{35}\text{S}$ -amino acid (Cys, Met) was released as  $^{35}\text{SO}_4^{2-}$  after only 30 mins. Under all three concentrations, even higher amount of  $^{35}\text{S}$  were retained in the microbial biomass from methionine. The amount of  $\text{SO}_4^{2-}$  released from methionine was lower than for cysteine, indicating that cysteine serves as a more important source of sulphate than methionine in the short term. This is in line with previous studies that cysteine showed a higher mineralization potential in soil in comparison to methionine (J. W. Fitzgerald, Hale and Swank, 1988).

In addition, some of the newly released sulphate was converted into soil organic sulphur through subsequent immobilization by soil microbial biomass. When added at 0.1 mM, around 4.8%, 1.7% of the generated  $^{35}\text{SO}_4^{2-}$  from Cys or Met was incorporated into the microbial biomass or organic sulphur separately by the end of the incubation. A similar observation was made when amino acids were added at 1 mM, where the incorporation of released sulphate corresponded to approximate 14.5% and 13.4% of the total  $^{35}\text{S}$ -Cys and Met added, respectively. However, when added at a higher concentration (10 mM), an accumulation of generated  $^{35}\text{SO}_4^{2-}$  in soil solution was found with incubation time, with around 26.5% and 22.3% of total  $^{35}\text{S}$ -Cys and Met recovered as  $^{35}\text{SO}_4^{2-}$  at the end of the incubation period.



**Fig. 5.4** Depletion of  $^{35}\text{S}$ -labelled cysteine or methionine in soil solution, the formation of extractable  $^{35}\text{S}$ -sulphate in soil and the subsequent incorporation of the  $^{35}\text{S}$  amino acid-S into microbial biomass over a 24 h-incubation period at three added amino acid concentrations (0.1 mM, 1 mM, 10 mM). Values represent means  $\pm$  SEM,  $n = 3$ .



### 5.3.3 Amino acid uptake by maize plants under hydroponic conditions

Results of amino acid uptake by maize plants in isolation under hydroponic conditions indicated that sterile maize roots were also capable of taking up both amino acids (Table 5.6). In total,  $12.6 \pm 0.8\%$  and  $14.5 \pm 0.4\%$  of the added  $^{14}\text{C}$ -Cys and Met were taken up by the maize roots after 24 h, after which  $^{14}\text{C}$  was incorporated into both new cell biomass and utilised for respiration. The addition of inorganic sulphate ( $\text{Na}_2\text{SO}_4$ ) did not affect this partitioning process significantly ( $p > 0.05$ ). Overall, the two S-containing amino acids behaved differently with regards to their internal partitioning in the plant. Based on the results presented in Table 5.4,  $46.4 \pm 3.4\%$  and  $25.5 \pm 1.5\%$  of the added  $^{14}\text{C}$ -Cys and Met taken up were respired by the maize plants, with the rest immobilised in the plant biomass. In addition, a similar amount of  $^{35}\text{S}$ -cysteine and methionine was recovered in the plant by the end of the 24 h incubation, suggesting that amino acid could have been taken up intact by the maize roots.

In total, the amino acid uptake rate by the maize plant roots over the 24 h period was  $0.87 \text{ nmol-}^{14}\text{C} (\text{cm}^{-1} \text{ root}) \text{ h}^{-1}$  for Cys, and for  $1.00 \text{ nmol-}^{14}\text{C} (\text{cm}^{-1} \text{ root}) \text{ h}^{-1}$  Met when added at 0.1 mM. This compares to uptake rates into soil microorganisms for Cys of  $0.41 \text{ nmol (g soil)}^{-1} \text{ h}^{-1}$  and for Met of  $0.40 \text{ nmol (g soil)}^{-1} \text{ h}^{-1}$  when added at the same concentration. Overall, this indicates a comparable capacity for taking up externally applied S-containing amino acids from the environment (more details are shown in chapter 8).

**Table 5.6** Carbon and S yields for rhizosphere microorganisms and axenic maize plants after growth on two S-containing amino acids (Cys and Met, 0.1 mM) over a 24 h-period separately. Values represent means  $\pm$  SEM ( $n = 3$ ). NA indicates not applicable.

Partitioning (nmol-amino acid $\text{cm}^{-1}$ root)		$^{14}\text{C}$ -Cys	$^{35}\text{S}$ -Cys	$^{14}\text{C}$ -Met	$^{35}\text{S}$ -Met
Plant	Respiration	$9.8 \pm 1.3$	NA	$6.2 \pm 0.3$	NA
	Plant biomass	$11.2 \pm 0.4$	$36.5 \pm 3.6$	$18.0 \pm 0.8$	$39.1 \pm 4.4$
Soil microorganisms	Microbial respiration	$6.7 \pm 0.3$	NA	$4.8 \pm 0.4$	NA
	Microbial biomass	$3.2 \pm 0.3$	$7.3 \pm 0.02$	$5.0 \pm 0.3$	$8.6 \pm 0.03$

### **5.3.4 Competition for $^{14}\text{C}$ from Cys and Met between plants and soil microorganisms in the rhizosphere as affected by nutrient amendment**

In accordance with results from the soil-only experiments, a rapid use of Cys and Met was seen in the microcosms containing both soil microorganisms and plant roots, with <10% of the  $^{14}\text{C}$ -amino acid recovered from soil solution after 24 h (Table 5.7 and 5.8). The competition results indicated that plant roots were poor competitors for free Cys and Met in soil solution, with only  $2.1 \pm 0.2\%$  and  $8.3 \pm 0.6\%$  of  $^{14}\text{C}$ -Cys and Met recovered in the plant biomass. A further  $4.9 \pm 0.3\%$  and  $2.1 \pm 0.3\%$  was also released through shoot respiration. Overall, the total plant capture of  $^{14}\text{C}$ -Cys and Met amounted to  $12.6 \pm 0.8\%$  and  $14.5 \pm 0.4\%$  for Cys and Met, respectively. In contrast, a large proportion of  $^{14}\text{C}$ -Cys and Met was recovered in the soil microbial biomass ( $36.7 \pm 5.1\%$ ,  $32.8 \pm 5.9\%$ , respectively), along with  $46.8 \pm 0.4$ ,  $37.4 \pm 0.5\%$  evolved in microbial respiration. The total uptake of Cys and Met by the microbial community was therefore estimated to be  $83.5 \pm 5.1\%$  and  $70.2 \pm 6.3\%$ , respectively, of the total amino acids added to the microcosm. A caveat to this is that we could not account for root respiration, however, we predict this to only contribute a small amount to soil  $^{14}\text{CO}_2$  evolution based on the sterile, plant-only  $^{14}\text{C}$  partitioning results.

The addition of N or S did not significantly influence the accumulation of  $^{14}\text{C}$ -Cys and Met into maize plant root tissues, but significantly reduced translocation of  $^{14}\text{C}$  to the plant shoot and thus the proportion of  $^{14}\text{CO}_2$  respired by the shoot ( $p < 0.05$ ). In total, the addition of N and S led to a significant reduction of total capture of  $^{14}\text{C}$ -Cys and Met by the maize plants, while the addition of glucose-C to the soil did not seem to have a significant influence on plant capture of either amino acid. On the other hand, addition of glucose-C resulted in a significant increase in the recovery of  $^{14}\text{C}$ -Cys and Met as  $^{14}\text{C}$ - $\text{CO}_2$  from the soil ( $p < 0.05$ ), whilst addition of N and S gave caused a significant reduction in the amount of  $^{14}\text{C}$ -Cys and Met respired as  $\text{CO}_2$  from the soil ( $p < 0.05$ ).

**Table 5.7** Percentage partitioning of  $^{14}\text{C}$  label after the introduction of 0.1 mM  $^{14}\text{C}$ -Cys into the maize rhizosphere over a 24 h incubation period in the presence of an excess of C (+ Glucose), inorganic N (+ N) or sulphate (+ S). Values represent means  $\pm$  SEM ( $n = 3$ ). \*Different letters of each row indicate significant differences ( $p < 0.05$ ) between treatments.

		$^{14}\text{C}$ -Cys	$^{14}\text{C}$ -Cys + N	$^{14}\text{C}$ -Cys + S	$^{14}\text{C}$ -Cys + G
Plant	Shoot respiration	$4.9 \pm 0.3^a$	$2.0 \pm 0.4^b$	$2.6 \pm 0.7^b$	$5.7 \pm 0.7^a$
	Shoot	$1.6 \pm 0.1^a$	$1.2 \pm 0.02^b$	$1.0 \pm 0.1^b$	$1.1 \pm 0.2^b$
	Root	$0.5 \pm 0.2^{ab}$	$0.4 \pm 0.1^b$	$0.4 \pm 0.02^b$	$0.6 \pm 0.1^a$
	Total	$7.4 \pm 0.6^a$	$3.6 \pm 0.5^b$	$3.9 \pm 0.8^b$	$7.3 \pm 0.8^a$
Soil	Microbial biomass	$36.8 \pm 5.1^b$	$33.1 \pm 1.8^{bc}$	$49.3 \pm 7.8^a$	$26.8 \pm 1.3^c$
	Microbial respiration	$46.8 \pm 0.4^b$	$40.6 \pm 1.1^c$	$38.9 \pm 0.7^c$	$57.6 \pm 2.1^a$
	Remaining in solution	$2.3 \pm 0.9$	$5.5 \pm 3.7$	$6.1 \pm 2.9$	$2.2 \pm 0.5$
	Total	$83.6 \pm 5.1^c$	$73.7 \pm 1.3^b$	$88.2 \pm 7.3^{ac}$	$84.4 \pm 2.9^c$
Total recovery		$90.6 \pm 5.6$	$77.3 \pm 3.6$	$92.2 \pm 8.1$	$91.7 \pm 3.4$

**Table 5.8** Percentage partitioning of  $^{14}\text{C}$  label after the introduction of 0.1 mM  $^{14}\text{C}$ -Met into maize rhizosphere over a 24 h incubation period in the presence of an excess of C (+ Glucose), inorganic N (+ N) or sulphate (+ S). Values represent means  $\pm$  SEM ( $n = 3$ ). \*Different letters of each row indicate significant differences ( $p < 0.05$ ) between treatments.

		$^{14}\text{C}$ -Met	$^{14}\text{C}$ -Met + N	$^{14}\text{C}$ -Met + S	$^{14}\text{C}$ -Met + G
Plant	Shoot respiration	$2.1 \pm 0.3a$	$1.4 \pm 0.1b$	$1.4 \pm 0.2b$	$1.9 \pm 0.1a$
	Shoot	$5.4 \pm 0.3a$	$3.4 \pm 0.5b$	$3.4 \pm 0.4b$	$5.77 \pm 1.5a$
	Root	$3.0 \pm 0.4$	$2.7 \pm 0.2$	$3.0 \pm 0.5$	$2.5 \pm 0.5$
	Total	$10.4 \pm 0.4a$	$7.5 \pm 0.6b$	$7.8 \pm 0.5b$	$10.2 \pm 1.1a$
Soil	Microbial biomass	$32.8 \pm 5.9$	$42.4 \pm 9.0$	$31.7 \pm 3.6$	$29.83 \pm 8.4$
	Microbial respiration	$37.4 \pm 0.5b$	$32.6 \pm 0.4c$	$31.1 \pm 0.6c$	$41.7 \pm 2.1a$
	Remaining in solution	$6.4 \pm 0.8$	$6.9 \pm 1.1$	$9.4 \pm 1.9$	$9.3 \pm 2.6$
	Total	$76.6 \pm 7.1$	$81.9 \pm 9.0$	$72.2 \pm 4.3$	$80.80 \pm 10.1$
Total recovery		$80.7 \pm 6.3$	$82.4 \pm 8.9$	$70.5 \pm 3.7$	$81.7 \pm 7.3$

**Table 5.9** Partitioning of  $^{14}\text{C}$  label after the introduction of  $^{14}\text{C}$ -labelled cysteine or methionine into the maize rhizosphere for 24 h at three different concentrations. Values represent means  $\pm$  SEM ( $n = 3$ ). Average plant shoot weight is 86.5 mg ( $n = 18$ ), and average plant root weight is 56.9 mg ( $n = 18$ ).

Percentage (%)	$^{14}\text{C}$ -Cys			$^{14}\text{C}$ -Met		
	0.1 mM	1 mM	10 mM	0.1 mM	1 mM	10 mM
Plant shoot respiration	4.9 $\pm$ 0.3	4.2 $\pm$ 0.5	2.2 $\pm$ 0.2	2.1 $\pm$ 0.3	1.6 $\pm$ 0.1	0.4 $\pm$ 0.1
Microbial respiration	46.8 $\pm$ 0.3	38.1 $\pm$ 3.1	20.1 $\pm$ 2.8	37.4 $\pm$ 0.5	32.1 $\pm$ 4.8	15.4 $\pm$ 1.4
Plant shoot uptake	1.6 $\pm$ 0.1	1.1 $\pm$ 0.3	0.7 $\pm$ 0.1	5.4 $\pm$ 0.3	4.5 $\pm$ 0.6	1.4 $\pm$ 0.1
Plant root uptake	0.5 $\pm$ 0.2	0.3 $\pm$ 0.04	0.1 $\pm$ 0.02	3.0 $\pm$ 0.4	0.7 $\pm$ 0.1	0.5 $\pm$ 0.1
Remaining in soil solution	2.3 $\pm$ 0.9	9.5 $\pm$ 2.9	29.2 $\pm$ 3.5	6.4 $\pm$ 0.8	19.2 $\pm$ 2.9	38.4 $\pm$ 4.9
Microbial storage	36.7 $\pm$ 5.1	31.2 $\pm$ 3.9	19.2 $\pm$ 2.4	32.8 $\pm$ 5.9	31.9 $\pm$ 1.1	24.2 $\pm$ 3.9
Plant capture	7.4 $\pm$ 0.5	5.6 $\pm$ 0.4	3.1 $\pm$ 0.3	10.4 $\pm$ 0.4	6.8 $\pm$ 0.7	2.3 $\pm$ 0.2
Microbial capture	83.6 $\pm$ 5.1	69.3 $\pm$ 3.7	39.5 $\pm$ 3.2	76.6 $\pm$ 7.1	64.0 $\pm$ 6.1	39.6 $\pm$ 3.7
Total recovered	90.6 $\pm$ 5.6	84.4 $\pm$ 6.1	71.5 $\pm$ 3.3	80.7 $\pm$ 6.3	90.2 $\pm$ 2.8	90.3 $\pm$ 5.1

**Table 5.10** Percentage partitioning of  $^{35}\text{S}$ -Cys, Met or  $\text{Na}_2^{35}\text{SO}_4$  (0.1 mM) between maize tissues and soil microorganisms in the rhizosphere over a 24 h-incubation period. Values represent means  $\pm$  SEM ( $n = 3$ ). \*Different letters of each row indicate significant differences ( $p < 0.05$ ) among treatments.

		$^{35}\text{S}$ -Cys	$^{35}\text{S}$ -Cys + $\text{Na}_2\text{SO}_4$	$^{35}\text{S}$ -Met	$^{35}\text{S}$ -Met + $\text{Na}_2\text{SO}_4$	$^{35}\text{S}$ - $\text{Na}_2\text{SO}_4$
Plant	Shoot	6.8 $\pm$ 0.9b	7.2 $\pm$ 1.1b	3.5 $\pm$ 0.8b	2.8 $\pm$ 0.4b	29.3 $\pm$ 7.6a
	Root	2.8 $\pm$ 0.6b	2.6 $\pm$ 0.4b	2.4 $\pm$ 0.5b	1.7 $\pm$ 0.2b	8.7 $\pm$ 1.2a
	Total	9.6 $\pm$ 1.1b	9.8 $\pm$ 1.5b	5.9 $\pm$ 1.2b	4.6 $\pm$ 0.5b	30.7 $\pm$ 7.5a
	Microbial biomass	51.7 $\pm$ 1.3ab	41.2 $\pm$ 14.8b	61.5 $\pm$ 2.5a	52.6 $\pm$ 7.3ab	16.9 $\pm$ 7.6c
Soil	Remaining in solution	18.0 $\pm$ 0.5a	16.7 $\pm$ 1.4a	5.9 $\pm$ 1.0c	5.1 $\pm$ 0.6c	13.7 $\pm$ 1.0b
	Total	69.8 $\pm$ 1.3a	57.8 $\pm$ 13.7a	67.5 $\pm$ 3.6a	57.7 $\pm$ 7.2a	30.7 $\pm$ 7.5b
Total recovery		79.3 $\pm$ 0.2	67.7 $\pm$ 15.1	73.4 $\pm$ 3.9	62.3 $\pm$ 7.7	68.7 $\pm$ 6.7

Our observations are consistent with other studies on plant uptake of amino acids (Persson and Näsholm, 2001; Kielland, McFarland and Olson, 2006; Hill and Jones, 2019), suggesting that cysteine and methionine may constitute a potentially important source of N and S for plant growth. Accordingly, it may be suggested that amino acid uptake rates are related to soil amino acids concentration, as microbial utilization of amino acids may influence the availability of amino acids to plant roots in soil solution (Lipson *et al.*, 1999) and thus the degree of competition between plant roots and the soil microbial community (Sauheitl, Glaser and Weigelt, 2009). Here we hypothesized that plant-microbial competition for Cys and Met would be diminished at high concentrations of free amino acid in soil solution. To test this hypothesis, the effect of three amino acid N (S) concentrations (0.1 mM, 1 mM, 10 mM) on the competition for Cys and Met between plants and soil microorganisms was investigated using  $^{14}\text{C}$  labelling. Our results revealed that total plant  $^{14}\text{C}$ -amino acid acquisition was significantly affected by amino acid concentration (Table 5.9). Overall, higher amino acid concentrations increased the availability of the amino acids allowing greater capture and reduced competition with soil microorganisms.

### **5.3.5 Competition for $^{35}\text{S}$ from Cys and Met between plants and soil microorganisms as affected by sulphate amendment**

The  $^{35}\text{S}$  results revealed that among all three S sources (added at 0.1 mM), inorganic sulphate is the preferred S source for maize plants, with around 30% of the added sulphate incorporated into plant biomass within 24 h (Table 5.10). This was higher than that of both S-containing amino acids (3 times higher than that of cysteine, 5 times higher than that of methionine). In contrast, organic amino acid-S was the preferred S source by soil microorganisms in comparison to inorganic sulphate, with more than 40% of the added  $^{35}\text{S}$  derived from amino acids retained in soil microbial biomass, while only 16.9% of the  $^{35}\text{S}$ -

sulphate was recovered in the soil microbial biomass. For all three S sources, more  $^{35}\text{S}$  was transported and retained in the plant shoot, indicating that S taken up by the plant is readily mobile and distributed through the whole plant to optimize performance.

## 5.4 Discussion

It is widely accepted from pure culture studies that soil microbial growth can rely solely on amino acids, with inorganic N released via deamination or transamination, and C skeletons released as C sources for glucogenic or ketogenic catabolic pathways (Moe, 2013). While C and N mineralization rates of amino acid in soils have been quantified and shown to be very rapid (Jones, 1999; Jones and Kielland, 2002; Wilkinson *et al.*, 2014; Almulla, Jones and Roberts, 2018; Jones *et al.*, 2018), mineralization of organic S in soils is not well understood. This is partly due to the complexity caused by multiple pathways involved in the mineralization of soil organic S. For example, S mineralization can occur through a range of biochemical processes such as the hydrolyzation of organic S by intra- and extra-cellular sulfatases, driven by microbial S demand, while biological process is driven by microbial energy needs, with S releases as a by-product (McGill and Cole, 1981; Zhao, Wu and McGrath, 1996). Specifically, the transformation of cysteine and methionine are first dominated with deamination and the release of ammonium (Greenwood and Lees, 1956; Ruiz-Herrera and Starkey, 1969a; Sohn and Ho, 1995), followed by desulphination of substituted pyruvic acid to sulphate (Freney, 1960; Andrew and Island, 1984; Hale and Fitzgerald, 1990). Further work could explore the metabolic profile of the soil after S-amino acid addition to investigate the relative importance of these different catabolic and anabolic pathways.

From our  $^{14}\text{C}$ -radiotracer results we concluded that the soil cysteine and methionine pool (0.1 mM) would turnover in the rhizosphere 553-685 times per day, with utilization rates and internal partitioning of Cys and Met affected by the inorganic N and S status of the soil.

However, as the concentrations of Cys and Met found *in situ* are typically lower (Autry and Fitzgerald, 1990; Werdin-Pfisterer, Kielland and Boone, 2009), actual C mineralization from cysteine and methionine in soil solution under field conditions may be even faster. These results support our hypothesis that cysteine and methionine are ubiquitously rapidly utilized by soil microorganisms and therefore there is likely to be strong competition both within the soil microbial community and with plants for this low molecular weight S resource pool. It is likely that strong inter-plant competition may also occur.

Previous studies on the relationships between S, C and N mineralization indicated that S mineralization did not occur in parallel with other elements (Homann and Harrison, 1992; Haque *et al.*, 2007). Rapid  $^{35}\text{S}$ -cysteine and methionine incorporation into the microbial biomass indicated that the mineralization of the two S-containing amino acids in soil was biotically mediated, rather than abiotic or occurring by extracellular enzymes residual in the soil. This decoupling of elemental cycles was also confirmed by faster C mineralization ( $^{14}\text{C}$ -amino acid to  $^{14}\text{CO}_2$ ) relative to S mineralization ( $^{35}\text{S}$ -amino acid to  $\text{SO}_4^{2-}$ ). We ascribe this to the microbial requirement to strip N and S from the amino acids to produce C skeletons that can enter respiratory cycles for energy production. It would therefore be expected that C and N excretion would occur in advance of  $^{14}\text{CO}_2$  production. It is also interesting to note that Cys was mineralized to a slightly greater extent than Met in all treatments. From this, we assume that Cys may serve as a more important S source for plants and soil microorganisms compared to Met due to its more rapid turnover as well as higher utilization efficiency by soil microorganisms. This probably also reflects the greater proportion of Cys relative to Met in proteins which enter the soil in plant and microbial residues.

It is a commonly held view that plants are capable of taking up amino acids and utilizing them as a N source for growth under sterile hydroponic conditions (Schmidt and Stewart, 1999; Jämtgård, Näsholm and Huss-Danell, 2008). Amino acids are transported into plant root cells

through active, proton-coupled symporters (Bush, 1993; Tegeder and Rentsch, 2010). Following uptake into the root cytoplasm, amino acids are used for the production of new cell biomass and to produce energy. When it comes to S, the observation that sulphate uptake by root cells is inhibited by organic S compounds such as cysteine and methionine implies that these compounds may also be bioavailable to root cells (Hart and Filner, 1969). Previous evidence suggests that cysteine and methionine can be rapidly taken up by tobacco cells (Parthier, Malaviya and Mothes, 1964; Harrington and Smith, 1977) and algae (Deane and O'Brien, 1981). However, little is known about potential contribution of S-containing amino acids to higher plant S nutrition such as agricultural crops. Here our results under hydroponic conditions revealed that comparable amounts of  $^{14}\text{C}$ -Cys and Met were taken up by maize roots, followed by translocation to the shoot where it was transformed and used for respiration. When cysteine and methionine were  $^{35}\text{S}$  labelled, incorporation of  $^{35}\text{S}$  into the plant confirmed that maize plants possess the capacity to access amino acids from soil solution should they become available.

In short-term rhizosphere competition studies for amino acids, plants are generally considered poor competitors against soil microorganisms (Hill and Jones, 2019). Our data indicated, however, that maize plants competed quite well for Cys and Met against the microbial community, especially at high exogenous amino acid concentrations. This indicates that organic N and S mineralization to  $\text{NH}_4^+$  and  $\text{SO}_4^{2-}$  is not a prerequisite for plant uptake of these nutrients. A significant change in the rate of plant Cys and Met uptake occurred when inorganic N and S was added to the soil (but not glucose-C), indicating that S-containing amino acids utilization by maize plants could be regulated by inorganic N and S availability. Further research is needed to reveal the distribution of organic S in specific root tissues, cells and compartments to better understand S transport and metabolism in response to the surrounding



environment. The top-down control in response to shoot S and N status on the root uptake of Cys and Met is also required (i.e. source-sink relationships).

One interesting question addressed here was whether plant-microbial competition in the rhizosphere is affected by different S forms (i.e. inorganic vs. organic). Our results showed that maize plants recovered more  $^{35}\text{S}$  than soil microorganisms in the inorganic S treatment, while soil microorganisms recovered more  $^{35}\text{S}$  than maize plants in the organic S ( $^{35}\text{S}$ -Cys and Met) treatments. We ascribe this to the fact that heterotrophic soil microorganisms demand C and N as well as S for their growth, whereas autotrophic plants only require N and S from soil.

## 5.5 Conclusions

In conclusion, experimental evidence was presented here that both maize plant roots and soil microorganisms have a similar capacity to take up free Cys and Met from the external solution when studied in isolation. However, when they are allowed to compete together, maize roots showed a lower ability to utilize both S-containing amino acids compared to the rhizosphere microbial community. Our results suggest that plants compete better for Cys and Met when their levels in soil are high. This suggests that plants may be better than microorganisms at exploiting ephemeral hotspots of organic N and S in soil. The present study greatly expands our understanding of terrestrial S cycling, as previously it was assumed that low molecular weight S compounds such as Cys and Met must be extracellularly cleaved to inorganic forms before plant uptake. Our study also points out the clear need for studies of plant acquisition of Cys and Met *in situ* where amino acid concentrations are low, in order to learn more about the contribution of low molecular weight S containing compounds for plant N and S nutrition. It would also be useful to study plants which have a greater S demand such as oilseed rape and other Brassicas (Basumatary *et al.*, 2021). In addition, DOS in soil does not just exist as free Cys and Met. It would therefore be desirable to investigate the plant and

microbial uptake of a wider variety of DOS compounds such as oligopeptides and S-containing secondary metabolites which may be released into soil. Lastly, the contribution of inorganic and organic S forms to the production of volatile organic S (VOS) would also be worthwhile as very little is known about the factors that control the production, consumption and transport of VOS in soil.

## 5.6 Acknowledgements

I thank Jonathan Roberts and Sarah Chesworth for their technical support. This work was supported by Bangor University (Grant number: B37-915) and the China Scholarship Council (Grant number: 201606510012).

## 5.7 References

- Almulla, A. A., Jones, D. and Roberts, P. (2018) 'Substrate influences temperature sensitivity of dissolved organic carbon (DOC) and nitrogen (DON) mineralization in arid agricultural soils', *Soil Systems*, 2(2), pp. 1–20. doi: 10.3390/soilsystems2020028.
- Andrew, T. L. and Island, S. (1984) 'Mineralization of methionine sulphur in soils and forest floor layers', 16(6), pp. 565–570. doi: org/10.1016/0038-0717(84)90073-7.
- Aula, L. *et al.* (2019) 'World sulfur use efficiency for cereal crops', *Agronomy Journal*, 111(5), pp. 2485–2492. doi: 10.2134/agronj2019.02.0095.
- Autry, A. R. and Fitzgerald, J. W. (1990) 'Sulfonate S: A major form of forest soil organic sulfur', *Biology and Fertility of Soils*, 10(1), pp. 50–56. doi: 10.1007/BF00336124.
- Barraclough, D. (1997) 'The direct or mit route for nitrogen immobilization: A <sup>15</sup>N mirror image study with leucine and glycine', *Soil Biology and Biochemistry*, 29(1), pp. 101–108. doi: 10.1016/S0038-0717(96)00241-6.

- Basumatary, A. *et al.* (2021) 'Impact of sulfur and boron fertilization on yield, quality of crop and nutrient use efficiencies in rapeseed in subtropical acidic soil of Assam, India', *Journal of Plant Nutrition*. Taylor & Francis, 44(12), pp. 1779–1793. doi: 10.1080/01904167.2021.1884708.
- Brailsford, F. L. *et al.* (2020) 'Rapid depletion of dissolved organic sulphur (DOS) in freshwaters', *Biogeochemistry*. Springer, 149(1), pp. 105–113. doi: 10.1007/s10533-020-00669-4.
- Bush, D. R. (1993) 'Proton-coupled sugar and amino acid transporters in plants', *Annual Review of Plant Physiology and Plant Molecular Biology*, 44(1), pp. 513–542. doi: 10.1146/annurev.pp.44.060193.002501.
- Cheng, X. and Bledsoe, C. S. (2004) 'Competition for inorganic and organic N by blue oak (*Quercus douglasii*) seedlings, an annual grass, and soil microorganisms in a pot study', *Soil Biology and Biochemistry*, 36(1), pp. 135–144. doi: 10.1016/j.soilbio.2003.08.024.
- David, M. B. and Mitchell, M. J. (1985) 'Sulfur constituents and cycling in waters, seston, and sediments of an oligotrophic lake', *Limnology and Oceanography*, 30(6), pp. 1196–1207. doi: 10.4319/lo.1985.30.6.1196.
- Deane, E. M. and O'Brien, R. W. (1981) 'Uptake of sulphate, taurine, cysteine and methionine by symbiotic and free-living dinoflagellates', *Archives of Microbiology*, 128(3), pp. 311–319. doi: 10.1007/BF00422537.
- Dietz, K. J. *et al.* (1990) 'Amino acid transport across the tonoplast of vacuoles isolated from barley mesophyll protoplasts: Uptake of alanine, leucine, and glutamine', *Plant Physiology*, 92(1), pp. 123–129. doi: 10.1104/pp.92.1.123.
- Eriksen, J. (2009) 'Soil sulfur cycling in temperate agricultural systems', *Advances in*

- Agronomy*. Elsevier, 102, pp. 55–89. doi: org/10.1016/S0065-2113(09)01002-5.
- Fitzgerald, J. W., Hale, D. D. and Swank, W. T. (1988) ‘Sulphur-containing amino acid metabolism in surface horizons of a hardwood forest’, *Soil Biology and Biochemistry*. Pergamon, 20(6), pp. 825–831. doi: 10.1016/0038-0717(88)90088-0.
- Freney, J. (1960) ‘The Oxidation of Cysteine to Sulphate in Soil’, *Australian Journal of Biological Sciences*, 13(3), p. 387. doi: 10.1071/bi9600387.
- Fujii, K. *et al.* (2018) ‘Another bottleneck for nitrogen mineralization in temperate forest soils: Arginine metabolism in microorganisms’, *Soil Biology and Biochemistry*. Elsevier, 126(August), pp. 22–30. doi: 10.1016/j.soilbio.2018.08.005.
- García, I., Romero, L. C. and Gotor, C. (2015) *Cysteine homeostasis, Amino Acids in Higher Plants*. CABI Publishing.
- Geisseler, D. *et al.* (2010) ‘Pathways of nitrogen utilization by soil microorganisms - A review’, *Soil Biology and Biochemistry*. Elsevier Ltd, 42(12), pp. 2058–2067. doi: 10.1016/j.soilbio.2010.08.021.
- Giovanelli, J. S., Mudd, S. H. and Datko, A. H. (1980) ‘Sulfur Amino Acids in Plants’, *Amino Acids and Derivatives*. New York: Academic Press. doi: org/10.1016/B978-0-12-675405-6.50018-8.
- Glanville, H. C. *et al.* (2016) ‘Combined use of empirical data and mathematical modelling to better estimate the microbial turnover of isotopically labelled carbon substrates in soil’, *Soil Biology and Biochemistry*. Elsevier Ltd, 94, pp. 154–168. doi: 10.1016/j.soilbio.2015.11.016.
- Greenwood, D. J. and Lees, H. (1956) ‘Studies on the decomposition of amino acids in soils - I. A preliminary survey of techniques’, *Plant and Soil*, 7(3), pp. 253–268. doi:

10.1007/BF01666122.

- Hale, D. D. and Fitzgerald, J. W. (1990) 'Generation of sulphate from cysteine in forest soil and litter.', *Soil Biology and Biochemistry*, 22(3), pp. 427–429. doi: 10.1016/0038-0717(90)90124-I.
- Haque, M. A. *et al.* (2007) 'Organic sulphur and nitrogen in pasture-cereal mixed cropping soils and related factors –a review', 1(July), pp. 39–54.
- Harrington, H. M. and Smith, I. K. (1977) 'Cysteine Transport into Cultured Tobacco Cells', *Plant Physiology*, 60(6), pp. 807–811. doi: 10.1104/pp.60.6.807.
- Hart, J. W. and Filner, P. (1969) 'Regulation of Sulfate Uptake by Amino Acids in Cultured Tobacco Cells', *Plant Physiology*, 44(9), pp. 1253–1259. doi: 10.1104/pp.44.9.1253.
- Heinze, S. *et al.* (2021) 'Microbial biomass sulphur—an important yet understudied pool in soil', *Agronomy*, 11(8), pp. 1–13. doi: 10.3390/agronomy11081606.
- Hewitt, E. J. (1952) *Sand and water culture methods used in the study of plant nutrition.*, *Sand and water culture methods used in the study of plant nutrition.*
- Hill, P. W. and Jones, D. L. (2019) 'Plant–microbe competition: does injection of isotopes of C and N into the rhizosphere effectively characterise plant use of soil N?', *New Phytologist*, 221(2), pp. 796–806. doi: 10.1111/nph.15433.
- Homann, P. S. and Harrison, R. B. (1992) 'Relationships among N, P, and S in temperate forest ecosystems', in *Atmospheric Deposition and Forest Nutrient Cycling*. Springer, pp. 214–232.
- Jämtgård, S., Näsholm, T. and Huss-Danell, K. (2008) 'Characteristics of amino acid uptake in barley', *Plant and Soil*, 302(1–2), pp. 221–231. doi: 10.1007/s11104-007-9473-4.

- Johnson, D. W. (1984) 'Sulfur cycling in forests', *Biogeochemistry*, 1(1), pp. 29–43. doi: 10.1007/BF02181119.
- Jones, D. L. (1999) 'Amino acid biodegradation and its potential effects on organic nitrogen capture by plants', *Soil Biology and Biochemistry*, 31(4), pp. 613–622. doi: 10.1016/S0038-0717(98)00167-9.
- Jones, D. L. *et al.* (2005) 'Plant capture of free amino acids is maximized under high soil amino acid concentrations', *Soil Biology and Biochemistry*, 37(1), pp. 179–181. doi: 10.1016/j.soilbio.2004.07.021.
- Jones, D. L. *et al.* (2018) 'Microbial competition for nitrogen and carbon is as intense in the subsoil as in the topsoil', *Soil Biology and Biochemistry*, 117(November 2017), pp. 72–82. doi: 10.1016/j.soilbio.2017.10.024.
- Jones, D. L. and Kielland, K. (2002) 'Soil amino acid turnover dominates the nitrogen flux in permafrost-dominated taiga forest soils', *Soil Biology and Biochemistry*, 34(2), pp. 209–219. doi: 10.1016/S0038-0717(01)00175-4.
- Jones, D. L. and Kielland, K. (2012) 'Amino acid, peptide and protein mineralization dynamics in a taiga forest soil', *Soil Biology and Biochemistry*. Elsevier Ltd, 55(3), pp. 60–69. doi: 10.1016/j.soilbio.2012.06.005.
- Jones, D. L., Nguyen, C. and Finlay, R. D. (2009) 'Carbon flow in the rhizosphere: Carbon trading at the soil-root interface', *Plant and Soil*, pp. 5–33. doi: 10.1007/s11104-009-9925-0.
- Jones, D. L., Owen, A. G. and Farrar, J. F. (2002) 'Simple method to enable the high resolution determination of total free amino acids in soil solutions and soil extracts', *Soil Biology and Biochemistry*, 34(12), pp. 1893–1902. doi: 10.1016/S0038-0717(02)00203-1.

- Kertesz, M. A. and Mirleau, P. (2004) 'The role of soil microbes in plant sulphur nutrition', *Journal of Experimental Botany*, 55(404), pp. 1939–1945. doi: 10.1093/jxb/erh176.
- Kielland, K., McFarland, J. and Olson, K. (2006) 'Amino acid uptake in deciduous and coniferous taiga ecosystems', *Plant and Soil*, 288(1–2), pp. 297–307. doi: 10.1007/s11104-006-9117-0.
- Kumar, V. *et al.* (2019) 'Amino acids distribution in economical important plants: a review', *Biotechnology Research and Innovation*, 3(2), pp. 197–207. doi: 10.1016/j.biori.2019.06.004.
- Kuzyakov, Y. and Jones, D. L. (2006) 'Glucose uptake by maize roots and its transformation in the rhizosphere', *Soil Biology and Biochemistry*, 38(5), pp. 851–860. doi: 10.1016/j.soilbio.2005.07.012.
- Kuzyakov, Y. and Siniakina, S. V (2001) 'A novel method for separating root-derived organic compounds from root respiration in non-sterilized soils', 164, pp. 511–517. doi: org/10.1002/1522-2624(200110)164:5<511::AID-JPLN511>3.0.CO;2-T.
- Lipson, D. A. *et al.* (1999) 'Variation in competitive abilities of plants and microbes for specific amino acids', *Biology and Fertility of Soils*, 29(3), pp. 257–261. doi: 10.1007/s003740050550.
- Lærkedal, P. *et al.* (2008) 'The Regents of the University of Colorado , a body corporate , contracting on behalf of the University of Colorado at Boulder for the benefit of INSTAAR Plant and Microbial Uptake and Allocation of Organic and Inorganic Nitrogen Related to Plant Growth Fo', *Arctic, Antarctic, and Alpine Research*, 8(2), pp. 139–150. doi: 10.1657/1523-0430(06-114).
- McGill, W. B. and Cole, C. V. (1981) 'Comparative aspects of cycling of organic C, N, S and

- P through soil organic matter', *Geoderma*. Elsevier, 26(4), pp. 267–286. doi: 10.1016/0016-7061(81)90024-0.
- McLaren, R. G. and Swift, R. S. (1977) 'Changes in soil organic sulphur fractions due to the long term cultivation of soils', *Journal of Soil Science*. Wiley Online Library, 28(3), pp. 445–453. doi: 10.1111/j.1365-2389.1977.tb02252.x.
- Miranda, K. M., Espey, M. G. and Wink, D. A. (2001) 'A rapid, simple spectrophotometric method for simultaneous detection of nitrate and nitrite', *Nitric Oxide - Biology and Chemistry*, 5(1), pp. 62–71. doi: 10.1006/niox.2000.0319.
- Miranda, M. *et al.* (2001) 'Amino acid permeases in developing seeds of *Vicia faba* L.: Expression precedes storage protein synthesis and is regulated by amino acid supply', *Plant Journal*, 28(1), pp. 61–71. doi: 10.1046/j.1365-313X.2001.01129.x.
- Moe, L. A. (2013) 'Amino acids in the rhizosphere: From plants to microbes', *American Journal of Botany*, 100(9), pp. 1692–1705. doi: 10.3732/ajb.1300033.
- Mulvaney, R. L. (1996) 'Nitrogen—inorganic forms', *Methods of soil analysis: Part 3 Chemical methods*. Wiley Online Library, 5, pp. 1123–1184. doi: 10.2136/sssabookser5.3.c38.
- Na, G. N. and Salt, D. E. (2011) 'The role of sulfur assimilation and sulfur-containing compounds in trace element homeostasis in plants', *Environmental and Experimental Botany*. Elsevier B.V., 72(1), pp. 18–25. doi: 10.1016/j.envexpbot.2010.04.004.
- Noji, M. and Saito, K. (2003) *Sulphur amino acids: biosynthesis of cysteine and methionine, Sulphur in Plants*. Springer.
- O'Dowd, R. W., Barraclough, D. and Hopkins, D. W. (1999) 'Nitrogen and carbon mineralization in soil amended with D- and L-leucine', *Soil Biology and Biochemistry*,



- 31(11), pp. 1573–1578. doi: 10.1016/S0038-0717(99)00083-8.
- Owen, A. G. and Jones, D. L. (2001) ‘Competition for amino acids between wheat roots and rhizosphere microorganisms and the role of amino acids in plant n acquisition’, *Soil Biology and Biochemistry*, 33(4–5), pp. 651–657. doi: 10.1016/S0038-0717(00)00209-1.
- Parthier, B., Malaviya, B. and Mothes, K. (1964) ‘Effects of Chloramphenicol and Kinetin on Uptake and Incorporation of Amino Acids By Tobacco Leaf Disks’, *Plant and Cell Physiology*, 5. doi: 10.1093/oxfordjournals.pcp.a079060.
- Persson, J. and Näsholm, T. (2001) ‘Amino acid uptake: A widespread ability among boreal forest plants’, *Ecology Letters*, 4(5), pp. 434–438. doi: 10.1046/j.1461-0248.2001.00260.x.
- Prasad, R. and Shivay, Y. S. (2018) ‘Sulphur in Soil, Plant and Human Nutrition’, *Proceedings of the National Academy of Sciences India Section B - Biological Sciences*. Springer India, 88(2), pp. 429–434. doi: 10.1007/s40011-016-0769-0.
- Ravilious, G. E. and Jez, J. M. (2012) ‘Structural biology of plant sulfur metabolism: From assimilation to biosynthesis’, *Natural Product Reports*, 29(10), pp. 1138–1152. doi: 10.1039/c2np20009k.
- Rentsch, D., Schmidt, S. and Tegeder, M. (2007) ‘Transporters for uptake and allocation of organic nitrogen compounds in plants’, *FEBS Letters*, 581(12), pp. 2281–2289. doi: 10.1016/j.febslet.2007.04.013.
- Rousk, J. and Jones, D. L. (2010) ‘Loss of low molecular weight dissolved organic carbon (DOC) and nitrogen (DON) in H<sub>2</sub>O and 0.5M K<sub>2</sub>SO<sub>4</sub> soil extracts’, *Soil Biology and Biochemistry*. Elsevier Ltd, 42(12), pp. 2331–2335. doi: 10.1016/j.soilbio.2010.08.017.

- Ruiz-Herrera, J. and Starkey, R. L. (1969) 'Dissimilation of methionine by a demethiolase of *Aspergillus* species.', *Journal of Bacteriology*, 99(3), pp. 764–770. doi: 10.1128/jb.99.3.764-770.1969.
- Sauheitl, L., Glaser, B. and Weigelt, A. (2009) 'Uptake of intact amino acids by plants depends on soil amino acid concentrations', *Environmental and Experimental Botany*, 66(2), pp. 145–152. doi: 10.1016/j.envexpbot.2009.03.009.
- Scherer, H. W. (2001) 'Sulphur in crop production - Invited paper', *European Journal of Agronomy*, 14(2), pp. 81–111. doi: 10.1016/S1161-0301(00)00082-4.
- Schmidt, S. and Stewart, G. R. (1999) 'Glycine metabolism by plant roots and its occurrence in Australian plant communities', *Functional Plant Biology*. CSIRO, 26(3), pp. 253–264. doi: org/10.1071/PP98116.
- Scott, N. M., Bick, W. and Anderson, H. A. (1981) 'The measurement of sulphur-containing amino acids in some scottish soils', *Journal of the Science of Food and Agriculture*, 32(1), pp. 21–24. doi: 10.1002/jsfa.2740320105.
- Smith, F. W. *et al.* (1997) 'Regulation of expression of a cDNA from barley roots encoding a high affinity sulphate transporter', *Plant Journal*, 12(4), pp. 875–884. doi: 10.1046/j.1365-313X.1997.12040875.x.
- Smith, G. S., Johnston, C. M. and Cornforth, I. S. (1983) 'Comparison of nutrient solutions for growth of plants in sand culture', *New phytologist*. Wiley Online Library, 94(4), pp. 537–548.
- Sohn, M. and Ho, C. T. (1995) 'Ammonia Generation during Thermal Degradation of Amino Acids', *Journal of Agricultural and Food Chemistry*, 43(12), pp. 3001–3003. doi: 10.1021/jf00060a001.

- Stevenson, F. J. (1982) 'Organic forms of soil nitrogen', *Nitrogen in agricultural soils*. Wiley Online Library, 22, pp. 67–122. doi: [org/10.2134/agronmonogr22.c3](https://doi.org/10.2134/agronmonogr22.c3).
- Ström, L. *et al.* (2002) 'Organic acid mediated P mobilization in the rhizosphere and uptake by maize roots', *Soil Biology and Biochemistry*, 34(5), pp. 703–710. doi: [10.1016/S0038-0717\(01\)00235-8](https://doi.org/10.1016/S0038-0717(01)00235-8).
- Swinnen, J. (1994) 'Evaluation of the use of a model rhizodeposition technique to separate root and microbial respiration in soil', *Plant and Soil*, 165(1), pp. 89–101. doi: [10.1007/BF00009966](https://doi.org/10.1007/BF00009966).
- Tegeder, M. and Rentsch, D. (2010) 'Uptake and partitioning of amino acids and peptides', *Molecular Plant*. The Authors 2010, 3(6), pp. 997–1011. doi: [10.1093/mp/ssq047](https://doi.org/10.1093/mp/ssq047).
- Voroney, R.P.; Brookes, P.C.; Beyaert, R. P. (2008) *Chapter 49: Soil Microbial Biomass C,N,P, and S, Soil Sampling and Methods of Analysis, Second Edition*. doi: [10.1201/9781420005271.ch18](https://doi.org/10.1201/9781420005271.ch18).
- Walch-Liu, P. *et al.* (2006) 'Evidence that L-glutamate can act as an exogenous signal to modulate root growth and branching in *Arabidopsis thaliana*', *Plant and Cell Physiology*, 47(8), pp. 1045–1057. doi: [10.1093/pcp/pcj075](https://doi.org/10.1093/pcp/pcj075).
- Warren, C. R. (2016) 'Simultaneous efflux and uptake of metabolites by roots of wheat', *Plant and Soil*. Plant and Soil, 406(1–2), pp. 359–374. doi: [10.1007/s11104-016-2892-3](https://doi.org/10.1007/s11104-016-2892-3).
- Werdin-Pfisterer, N. R., Kielland, K. and Boone, R. D. (2009) 'Soil amino acid composition across a boreal forest successional sequence', *Soil Biology and Biochemistry*. Elsevier Ltd, 41(6), pp. 1210–1220. doi: [10.1016/j.soilbio.2009.03.001](https://doi.org/10.1016/j.soilbio.2009.03.001).
- Wilkinson, A. *et al.* (2014) 'Rapid microbial uptake and mineralization of amino acids and peptides along a grassland productivity gradient', *Soil Biology and Biochemistry*.

- Elsevier Ltd, 72, pp. 75–83. doi: 10.1016/j.soilbio.2014.01.026.
- Wirtz, M. and Droux, M. (2005) ‘Synthesis of the sulfur amino acids: Cysteine and methionine’, *Photosynthesis Research*, 86(3), pp. 345–362. doi: 10.1007/s11120-005-8810-9.
- Xu, X. *et al.* (2008) ‘Light affects competition for inorganic and organic nitrogen between maize and rhizosphere microorganisms’, *Plant and Soil*, 304(1–2), pp. 59–72. doi: 10.1007/s11104-007-9519-7.
- Yamagata, M. and Ae, N. (1999) ‘Direct acquisition of organic nitrogen by crops’, *Japan Agricultural Research Quarterly*, 33(1), pp. 15–21.
- Zhang, W. *et al.* (2015) ‘Differentiating the mineralization dynamics of the originally present and newly synthesized amino acids in soil amended with available carbon and nitrogen substrates’, *Soil Biology and Biochemistry*, 85, pp. 162–169. doi: 10.1016/j.soilbio.2015.03.004.
- Zhao, F. J., Wu, J. and McGrath, S. P. (1996) ‘Soil Organic Sulphur and its Turnover’, *Humic Substances in Terrestrial Ecosystems*, pp. 467–506. doi: 10.1016/b978-044481516-3/50013-x.
- Zhao, F. and McGrath, S. P. (1994) ‘Extractable sulphate and organic sulphur in soils and their availability to plants’, *Plant and Soil*, 164(2), pp. 243–250. doi: 10.1007/BF00010076.

## Chapter 7

---

**Tracing the mineralization rates of C, N and S from cysteine and methionine in a grassland soil: A  $^{14}\text{C}$  and  $^{35}\text{S}$  dual-labelling study**

## Abstract

Cysteine (Cys) and Methionine (Met) represent the two main sulphur (S)-containing amino-acids found in soil solution. Although general measures of S cycling (e.g., sulfatase activity) provide valuable information concerning the cycling of labile organic S in soil, detailed information regarding the microbial transformation pathways of Cys and Met at a molecular level remain poorly characterised. Therefore, in this study a  $^{14}\text{C}$  and  $^{35}\text{S}$  dual-isotopic labelling approach was used to trace the fate of C and S derived from Cys and Met in an agricultural grassland soil over a 7-day incubation period. Microbial biomass C, N, and S were analysed by the  $\text{CHCl}_3$  fumigation-extraction method, and  $\text{CO}_2$  evolution along with inorganic nutrient release ( $\text{NH}_4^+$ ,  $\text{NO}_3^-$  and  $\text{SO}_4^{2-}$ ) were also measured. We then imposed an excess of C as glucose, or excess NPS (as  $\text{NH}_4\text{NO}_3$ ,  $\text{KH}_2\text{PO}_4$  and  $\text{K}_2\text{SO}_4$ ) to investigate whether the Cys and Met mineralization process was affected by manipulating C, N and S availability in the soil solution. Our results showed that after 168 h, 2.7 – 19.5% of the  $^{14}\text{C}$  derived from Cys and Met had been immobilised in the microbial biomass, 67.2 – 89.2% had been respired as  $^{14}\text{CO}_2$  while the recovery of  $^{35}\text{S}$  label in the soil microbial biomass ranged from 11.9 – 41.8%. Overall, our results indicated that microbial communities have a high capacity to utilize Cys and Met but that they enter multiple metabolic pathways once inside the cell. While some of the amino acids may be directly used in protein synthesis, other metabolic pathways lead to  $\text{NH}_4^+$  and  $\text{SO}_4^{2-}$  being released back into the soil with the  $\text{NH}_4^+$  then rapidly converted to  $\text{NO}_3^-$  by the nitrifier community. The resulting C-skeletons were dissimilated and used to produce energy, leading to the release of  $^{14}\text{CO}_2$ . The significant differences in C, N and S mineralization processing demonstrate a decoupling of the S and C cycles at the molecular level. Further, glucose-C addition shifted the allocation of C inside the cell to more catabolic processes and greater mineralization, while in comparison inorganic N, P and S availability had much less effect on resource partitioning.

**Keywords:** Biodegradation; Dissolved organic sulphur; Nutrient availability; Radioisotope tracers;  $^{14}\text{C}$  tracer;  $^{35}\text{S}$  tracer; Grassland soil; Turnover.

## 7.1 Introduction

Agricultural mineral soils have a quasi-constant mean carbon (C): nitrogen (N): phosphorous (P): sulphur (S) ratio of 10000: 833: 200: 143 on a mass basis (Kirkby *et al.*, 2011). Consequently, changes in the abundance of one element are often balanced by increased or decreases in another. Over the next few decades, it is predicted that global C and N inputs to soil will increase due to the need to improve plant productivity and food security (Liang *et al.*, 2011; Sillen & Dieleman, 2012). Thus, terrestrial ecosystems are expected to develop a higher demand for other nutrients, including S and other micronutrients stripped from the soil system at harvest. (Jones *et al.*, 2013). Typically, N, P, and S are seen as the major nutrients that are most likely to limit plant and microbial growth (Fermoso *et al.*, 2019), with their availability in soil a key constraint for the productivity of terrestrial ecosystems (Marschner & Rengel, 2012; Vitousek & Howarth, 1991). In most agroecosystems, plant nutrition relies in part on the decomposition and mineralization of organically bound N and S mediated by the soil microbial community (Sahrawat, 1981; Michael A. Kertesz and Mirleau, 2004a). Many of the labile organic N- and S-containing compounds in soil are present in a low molecular weight (MW) soluble form within the dissolved organic matter (DOM) pool. This pool represents a key component of C, N and S cycling (Brailsford, 2019), and the breakdown of DOM in soil is a critical step in supplying available nutrients for plant growth (Marschner & Kalbitz, 2003).

More than 90% of the soil's S reserves are bound within soil organic matter (SOM; Eriksen, 2009) with most of this held within protein moieties. Soil microorganisms secrete large quantities of proteases into soil which facilitate the breakdown of proteins and peptides into free amino acids (AA) (Jan *et al.*, 2009; Kandeler *et al.*, 1999). Therefore, despite the fact that plants mainly absorb S as inorganic sulphate, organic S (carbon-bonded S and ester S) may be also be an important source of S to plants, especially at the height of the growing season,

or when there is a shortfall in the supply of inorganic sulphate (Freney et al., 1981). Acid hydrolysis has revealed that amino acids are the largest pool of organic in soil (Stevenson, 1982), and that they also represent an important energy and N source for soil organisms. Within this amino acid pool, S-containing cysteine (Cys) and methionine (Met), constitute the most important reserve of combined C, N and S in soil (Scott et al., 1981) and they also constitute the majority of organic S in plants (Allaway and Thompson, 1966). Therefore, understanding the bioavailability and fate of these S-containing amino acids in the plant-soil-microbial system is important step in understanding and modelling terrestrial N and S cycling.

Turnover rates of low MW dissolved organic C in soil have been measured by adding isotopically labelled (e.g.  $^{13}\text{C}$ ,  $^{14}\text{C}$ ) substrates to soil in laboratory incubations and measuring the evolved  $^{13}\text{CO}_2/^{14}\text{CO}_2$  (Hill et al., 2008a; O'Dowd & Hopkins, 1998). The isotopic approach also provides information on the internal partitioning of the added  $^{14}\text{C}$  in the microbial biomass. After uptake, the amino acid-C taken up by the soil microbial community is partitioned into two pools associated with either catabolic or anabolic processes (Jones et al., 2009). The catabolic production of  $^{14}\text{CO}_2$  is normally characterized by an initial rapid phase (< 3 h) followed by a slower phase that continues for several weeks. The first rapid phase of  $^{14}\text{CO}_2$  production is attributable to the immediate use of the substrate in catabolic processes (i.e. respiration) and typically accounts for approximately 7-16% of the added amino acid- $^{14}\text{C}$  (Boddy et al., 2007; Wilkinson et al., 2014a). This is followed by a secondary slower phase of  $^{14}\text{CO}_2$  production which is attributable to the subsequent turnover of (i.e. the  $^{14}\text{C}$  incorporated into the anabolic pool (i.e. maintenance respiration) and ultimately accounts for 48-87% of the added amino acid- $^{14}\text{C}$  (Boddy *et al.*, 2007; Farrell, Hill, Wanniarachchi, *et al.*, 2011).

The mineralization of amino acid-N occurs via the direct route in which organic N compounds are taken up intact by soil microorganisms, assimilated and then any excess N is released as  $\text{NH}_4^+$  back into the soil (Barracough et al., 2015; Finzi & Berthrong, 2005).



Alternatively, amino acid-N held in exposed amino groups can be released extracellularly through the action of deaminases, after which the  $\text{NH}_4^+$  can be taken up and assimilated (Jansson and Persson, 2015). Similarly, based on the stoichiometric relationship between C, N and S, it is suggested that both biological and biochemical mineralisation processes are involved in the mineralisation of organic S (David et al., 1983; McGill & Cole, 1981). However, it should be noted that sulfatases cannot act on Cys and Met. Biochemical mineralization is mainly controlled by the S supply to soil microorganisms rather than their need for energy, which is responsible for the release of sulphates from ester sulphates by the arylsulphatase enzyme (Tabatabai and Bremner, 1970), while biological mineralization is driven by the energy demand of actively growing microorganisms (Wu et al., 1993) (shown in Fig. 7.1), releasing inorganic sulphate as a by-product, which is responsible for the release of sulphate from carbon-bonded S (Ghani et al., 1992). Some of the newly released inorganic N and sulphate is then converted to soil organic S through subsequent immobilization by the microbial biomass (Barraclough, 1997; Vermeiren et al., 2018).


Due to the lack of sensitive and specific methods for detection, identification and estimation of individual S compounds, the biodegradation processes of S from DOS in soils remain uncertain. Previous studies indicate that Cys is decomposed rapidly in soil to inorganic sulphate (Freney, 1960), following the sequence: cysteine  $\rightarrow$  cystine  $\rightarrow$  cystine disulphoxide  $\rightarrow$  cysteine sulphinic  $\rightarrow$  inorganic sulphate (Levine, 1982). In contrast, the most frequently reported end-product for Met decomposition is methanethiol (Ruiz-Herrera and Starkey, 1969b; Segal and Starkey, 1969), with other products reported as dimethyl disulphide (Ruiz-Herrera and Starkey, 1969c; W L Banwart and Bremner, 1976), dimethyl sulfide (W. L. Banwart and Bremner, 1975), hydrogen sulfide (Nader and Walker, 1970; Zinder and Brock, 1978), and inorganic sulphate (Fitzgerald and Andrew, 1984). Based on the multiple steps required to break down these amino acids extracellularly, it seems more energetically efficient

to take them inside the cell and assimilate them using as required based on the prevailing conditions.

From the current knowledge of S decomposition processes in soils, it is possible that microbial demand for C is the dominant driver for both biochemical and biological mineralisation of organic S (Ghani, McLaren and Swift, 1992). Therefore, the supply of a readily available C source to the soil may have a considerable effect on organic S mineralization. If available C concentrations in the soil are too low to meet microbial demand, mineralisation of organic S would take place, and during this process greater S excretion is likely. Conversely, in the presence of high amounts available C in soil, immobilisation of inorganic S is likely to occur. Similarly, since sulphatase activity is related to concentrations of its end-product (inorganic sulphate), when inorganic sulphate concentrations are too low to meet microbial demand, sulphatase enzymes are used to hydrolyse sulphate esters. Conversely, addition of sulphate to the soil may hinder the mineralisation of soil native organic S. By experimentally manipulating the amounts and the availability of C, N and S in soils, we aim to gain a better understanding of what regulates the fate of Cys and Met in soil.

Net nutrient mineralization represents the sum of gross mineralization and immobilization (Schimel and Bennett, 2004; Scherer, 2009). The mineralization and immobilization processes in soil are tightly linked to microbial activity. Soil microbial activity in soil is regulated by the supply of energy and nutrients. Generally, it is reported that the cut-off point of the C/N ratio with respect to mineralization/immobilization is around 25 (Kumar and Goh, 1999). Materials with a narrow C/N ratio (below 25) are expected to result in net mineralization, whereas materials with wider C/N ratio favour immobilization due to the abundance of C relative to N (Kumar and Goh, 2003). Therefore, experimental manipulation of C-to-nutrient ratios in soil could help differentiate between the effects of nutrient availability on the biodegradation and microbial use of the organic N and S pools in soil. It should be noted

that the C:N:S ratio for Cys (3:1:1) and Met (5:1:1) will always favour excretion of excess  $\text{SO}_4^{2-}$  and  $\text{NH}_4^+$  when supplied as a sole C source to microbial cells (bacterial C:N:S ratio of 8:1:0.25) (Fagerbakke et al., 1996).



**Fig. 7.1** Schematic representation of the microbial degradation of -S-containing amino acids in soil (biological mineralization process). In this study, microbial respiration and amino acid consumption were determined, as well as gross rates of inorganic N and S fluxes, namely, N mineralization, ammonium consumption, nitrification, nitrate consumption, S mineralization and sulphate consumption.

In this study, a non-fertilized grassland soil was chosen to avoid the effects of long-term fertilization on soil microbial community structure and functioning. After adding <sup>14</sup>C- or <sup>35</sup>S-labelled Cys and Met to soil, we tracked their fate in terms of immobilization in the microbial biomass, C mineralisation and inorganic nutrient release (Fig. 7.1). The specific objectives of

this study were to (1) compare short-term concurrent mineralization of C, N and S derived Cys and Met over 7 d; and (2) assess the effect of inorganic nutrient availability on microbial mineralization and utilization of Cys and Met. Such studies would provide valuable information for future modelling of S transformations and nutrient coupling in soil. Further, the information will improve our fundamental mechanistic understanding of S cycling with the aim of designing better fertiliser management regimes to address S deficiency.

## **7.2 Materials and methods**

### **7.2.1 Study sites and soil sampling**

Replicate batches of soil samples ( $n = 3$ ) were collected in May 2018 from the Ah horizon (0 -10 cm) of an unfertilized grassland plot at Abergwyngregyn, Gwynedd, UK (53°14'N 4°11'W). On return to the laboratory, the soil was sieved (< 2 mm) to remove stones, fine roots, and other plant debris, where it was stored at < 4 °C prior to further analysis. Soil pH and electrical conductivity (EC) were determined in 1:5 (w/v) soil-to-distilled water extracts using standard electrodes. Microbial biomass C and N (MBC/N) were determined by the chloroform-fumigation extraction procedure of Voroney et al. (2008). Briefly, the amount of DOC and DON was determined before and after  $\text{CHCl}_3$  fumigation (48 h) with 0.5 M  $\text{K}_2\text{SO}_4$  extracts (30 min, 200 rev  $\text{min}^{-1}$ ) using  $K_{\text{EC}}$  and  $K_{\text{EN}}$  extraction factors of 0.35 and 0.5, respectively (Voroney et al., 2008). For  $\text{NO}_3^-$  and  $\text{NH}_4^+$  analysis, the soil was extracted with 0.5 M  $\text{K}_2\text{SO}_4$  (1:5 w/v) and  $\text{NO}_3^-$  in the extracts determined colorimetrically on a Synergy MX microplate reader using the vanadate procedure of Miranda et al. (2001), while  $\text{NH}_4^+$  was determined colorimetrically using the salicylate procedure of Mulvaney (1996). Free amino acids and hydrolysable protein was determined by the *o*-phthaldialdehyde fluorescence method of Jones et al. (2002) using a Cary Eclipse fluorimeter. All values are reported on a dry soil weight basis (Table 7.1).

**Table 7.1** Physiochemical properties of the soil used in this study. Data is presented on a soil dry weight basis. Values represent means  $\pm$  SEM ( $n = 3$ ).

Parameter	Values
Altitude (m)	32
Water content (%)	$17.1 \pm 0.2$
pH	$5.92 \pm 0.02$
Electrical conductivity ( $\mu\text{S cm}^{-1}$ )	$74 \pm 4$
Microbial biomass C ( $\text{g kg}^{-1}$ )	$0.67 \pm 0.02$
Microbial biomass N ( $\text{mg kg}^{-1}$ )	$16.8 \pm 21.6$
Total dissolved N ( $\text{mg kg}^{-1}$ )	$42.6 \pm 0.7$
Total dissolved C ( $\text{mg kg}^{-1}$ )	$35.1 \pm 8.7$
Total dissolved C:N of bulk soil	$5.34 \pm 0.19$
Ammonium-N ( $\text{mg kg}^{-1}$ )	$1.14 \pm 0.14$
Nitrate-N ( $\text{mg kg}^{-1}$ )	$17.1 \pm 0.3$
Amino acids ( $\text{mg N kg}^{-1}$ )	$0.61 \pm 0.15$
Protein ( $\text{mg N kg}^{-1}$ )	$2.95 \pm 1.37$
Sulphate-S ( $\text{mg S kg}^{-1}$ )	$4.37 \pm 0.79$
Total dissolved SS ( $\text{mg S kg}^{-1}$ )	$31.6 \pm 2.1$

### 7.2.2 Incubation experiments

The incubation experiments lasted for 7 d, as previous studies have shown that the response of the microbial biomass was relatively fast (several hours to several days) when labile substrates are added to soil (Blagodatskaya & Kuzyakov, 2013; Blagodatskaya et al., 2007, 2009). The incubations were carried out at room temperature ( $21 \pm 1$  °C) in the dark, in the absence of plants. All treatments were conducted in triplicate. Samples for microbial biomass C, N and S determination (MBC, MBN, MBS) were taken before application of the experimental treatments. Each experimental unit comprised 5 g of field-moist soil held in individual 50 cm<sup>3</sup> polypropylene centrifuge tubes with a screw cap. All radiolabelled substrates were made up in 1 ml of deionized water, before being uniformly applied to the soil surface dropwise using a pipette. All centrifuge tubes were set up at the same time under the same experimental conditions. The incubation experiment consisted of two parallel experiments, one involving <sup>14</sup>C and the other using <sup>35</sup>S.

Briefly, soil in each unit was spiked with 1 ml of <sup>14</sup>C-labelled or <sup>35</sup>S-labelled Cys or Met (1 mM; 0.3 kBq ml<sup>-1</sup>; PerkinElmer Inc, Waltham, MA) either in the presence or absence

of glucose-C or inorganic nutrients (N, P and S). Glucose (+G treatment) was applied to soil at a rate of 360 mg C kg<sup>-1</sup> soil, whereas inorganic N, P and S (+NPS treatments) were added as 100 mg N (NH<sub>4</sub>NO<sub>3</sub>), 30 mg P (KH<sub>2</sub>PO<sub>4</sub>), 30 mg S (K<sub>2</sub>SO<sub>4</sub>) per kg soil. Glucose addition was equivalent to ca. 50% of the native microbial biomass C content. NPS application rates were chosen to represent common UK field fertiliser application practices (DEFRA, 2016). To measure the speed of glucose turnover, an additional treatment was used in which the <sup>14</sup>C-amino acids were replaced with <sup>14</sup>C-glucose (0.3 kBq ml<sup>-1</sup>; Sigma-Aldrich Ltd., Poole, UK). We chose Cys and Met as model substrates as C-bonded S as they represent the dominant low molecular weight dissolved organic S compounds entering soil (Fitzgerald et al., 1988; Yazzie et al., 1994; Yeoh & Watson, 1982). The concentration of Cys and Met (1 mM) was chosen to reflect those likely to occur in the rhizosphere upon lysis of root cells.

### 7.2.3 Partitioning of the <sup>14</sup>C into different compartments

After application of the <sup>14</sup>C-labelled amino acid solution to the soil surface, <sup>14</sup>C either remained in solution, or was taken up by the microbial community. After uptake, the amino acids can be used either for microbial growth (synthesis of new cells; <sup>14</sup>C-biomass) or for production of energy (<sup>14</sup>CO<sub>2</sub>). Therefore, we determined their partitioning of the added <sup>14</sup>C isotope into a range of pools as follows:

- <sup>14</sup>C<sub>O</sub>: <sup>14</sup>C-substrate added at time zero, determined by counting the <sup>14</sup>C-substrate solution (kBq);
- <sup>14</sup>C<sub>uf</sub>: <sup>14</sup>C in the 0.5 M K<sub>2</sub>SO<sub>4</sub> extract of non-fumigated soils (kBq);
- CO<sub>2</sub> partitioning: (<sup>14</sup>CO<sub>2</sub> trapped in NaOH/<sup>14</sup>C<sub>O</sub>) × 100 (%);
- <sup>14</sup>C<sub>FE</sub>: (<sup>14</sup>C in the 0.5 M K<sub>2</sub>SO<sub>4</sub> extract of fumigated soils - <sup>14</sup>C<sub>uf</sub>) × 100/<sup>14</sup>C<sub>O</sub> (%);
- <sup>14</sup>C<sub>MB</sub>: Microbial biomass <sup>14</sup>C = (<sup>14</sup>C<sub>O</sub> - CO<sub>2</sub> - <sup>14</sup>C<sub>uf</sub>) × 100/<sup>14</sup>C<sub>O</sub> (%);
- <sup>14</sup>C<sub>FNON-FE</sub>: <sup>14</sup>C<sub>MB</sub> - <sup>14</sup>C<sub>FE</sub> (%);

- $^{14}\text{C}_{\text{efficiency}} = ^{14}\text{C}_{\text{MB}}/^{14}\text{C}_{\text{O}}$ ;
- $K_{\text{ec}}: (^{14}\text{C}_{\text{FE}}/^{14}\text{C}_{\text{MB}})$ .

### 7.2.3.1 Microbial mineralization of $^{14}\text{C}$ -labelled Cys and Met

After addition of  $^{14}\text{C}$  labeled amino acid solution to the soil surface, a 6 ml polypropylene tube containing 1 M NaOH (1 ml) was placed inside the incubation vessels above the soil to trap respired  $^{14}\text{CO}_2$ . These NaOH  $\text{CO}_2$  traps were changed hourly up to 24 h and then after 48, 72, 96, 120, 144 and 168 h after substrate addition to quantify microbial respiration rates. After removal, the amount of  $^{14}\text{CO}_2$  in the 1 M NaOH traps was determined by liquid scintillation counting using a Wallac 1404 scintillation counter (Wallac EG&G, Milton Keynes, UK) and Optiphase HiSafe 3 alkali-compatible scintillation fluid (Perkin Elmer Inc., Waltham, MA).

### 7.2.3.2 Determination of microbial biomass $^{14}\text{C}$

Microbial biomass  $^{14}\text{C}$  was measured by the  $\text{CHCl}_3$  fumigation incubation method (Voroney et al., 2008). Briefly, at nine different time points (6, 12, 24, 48, 72, 96, 120, 144 and 168 h) after substrate addition, 5 g of soil was exposed to chloroform vapour for 24 h. After removal of the fumigant, the soil was extracted with 25 ml of 0.5 M  $\text{K}_2\text{SO}_4$  (30 min, 200 rev  $\text{min}^{-1}$ ) to recover any  $^{14}\text{C}$ -label remaining in the soil solution or held on soil exchange surfaces. The extracts were centrifuged (4000 rev  $\text{min}^{-1}$ , 5 min), and  $^{14}\text{C}$  in the supernatant determined by liquid scintillation counting as described previously. A non-fumigated control was extracted alongside the fumigated counterparts. The  $^{14}\text{C}$  contained in the microbial biomass was calculated as the differences of  $^{14}\text{C}$  recovered in the fumigated and non-fumigated samples, adjusted with the extraction factor of 0.35.

#### 7.2.4 Partitioning of the $^{35}\text{S}$ into different compartments

Similar to the approach taken above for  $^{14}\text{C}$ , the partitioning of  $^{35}\text{S}$  was determined as follows:

- $^{35}\text{S}_\text{O}$ :  $^{35}\text{S}$  -substrate added at time zero, determined by counting the  $^{35}\text{S}$ -substrate solution (kBq);
- $^{35}\text{S}_\text{uf}$ :  $^{35}\text{S}$  in the 0.01 M  $\text{CaCl}_2$  extract of non-fumigated soils (kBq);
- $^{35}\text{S}_\text{FE}$ : ( $^{35}\text{S}$  in the 0.01 M  $\text{CaCl}_2$  extract of fumigated soils -  $^{35}\text{S}_\text{uf}$ )  $\times 100 / ^{35}\text{S}_\text{O}$  (%);
- $^{35}\text{S}_\text{MB}$ : Microbial biomass  $^{35}\text{S} = (^{35}\text{S}_\text{O} - ^{35}\text{S}_\text{uf}) \times 100 / ^{35}\text{S}_\text{O}$  (%);
- $^{35}\text{S}_\text{FNON-FE}$ :  $^{35}\text{S}_\text{MB} - ^{35}\text{S}_\text{FE}$  (%);
- $K_\text{ec}$ : ( $^{35}\text{S}_\text{FE} / ^{35}\text{S}_\text{MB}$ ).

##### 7.2.4.1 Production of volatile sulphur compounds from Cys or Met mineralization

To determine the rate of volatile sulphur compounds produced after the addition of the  $^{35}\text{S}$ -labelled Cys or Met to the soil, 5 g of soil was placed in a 50  $\text{cm}^3$  centrifuge tube, and 1 ml of 1 mM  $^{35}\text{S}$ -labelled Cys or Met added to the soil surface as described above. A 6 ml polypropylene tubes filled with 1 ml of ethanol (> 98%; Sigma-Aldrich Ltd.) was placed inside the closed incubation vessel to trap any volatile sulphur compounds evolved. The ethanol traps were collected and changed after 6, 12, 24, 48, 72, 96, 122, 144 and 168 hours, and  $^{35}\text{S}$  was determined by liquid scintillation counting as described previously. It should be noted that this is not a proven technology for trapping volatile S and in retrospect is unlikely to capture all forms of volatile S.

##### 7.2.4.2 Determination of microbial biomass $^{35}\text{S}$

At each sampling time,  $^{35}\text{S}$  remaining free in the soil was extracted by adding 25 ml of 0.01 M  $\text{CaCl}_2$  solution to the soil. The soils were then extracted on an end-over-end shaker (30



min, 200 rev min<sup>-1</sup>), and subsequently centrifuged (5 min, 4000 rev min<sup>-1</sup>). The supernatant was then analysed for total <sup>35</sup>S activity by liquid scintillation counting as described previously. To separate organic-S and sulphate-S in the soil extract, a BaCl<sub>2</sub> turbidimetric/precipitation approach was used (Combs et al., 1998). Briefly, 10 ml of the centrifuged extract was mixed with 10 ml of 1 M BaCl<sub>2</sub> on an end-over-end shaker for 10 min at 200 rev min<sup>-1</sup>. Subsequently, the mixture was centrifuged (5 min, 4000 rev min<sup>-1</sup>) and the supernatant analysed for organic <sup>35</sup>S (the inorganic S having precipitated as BaSO<sub>4</sub>). The difference between <sup>35</sup>S activity in soil solution before and after BaSO<sub>4</sub> precipitation was considered as inorganic sulphate generated from mineralisation of the added amino acids.

To estimate microbial biomass-S, chloroform fumigation-extraction was used as described above except that 0.01 M CaCl<sub>2</sub> was used as the extractant. <sup>35</sup>S retained in the microbial biomass was calculated as the differences in CaCl<sub>2</sub> extractable <sup>35</sup>S between fumigated and non-fumigated samples, adjusted with an extraction factor of 0.35 (Voroney et al., 2008). All <sup>35</sup>S data were corrected for radioactive decay relative to the start of the incubation (half time of 87.4 d; Holtzhauer, 2006; Zoon, 1987).

### **7.2.5 Microbial biomass analysis**

The chloroform fumigation-extraction procedure used above was used to determine microbial biomass (C<sub>mic</sub>, N<sub>mic</sub>, S<sub>mic</sub>). In this case, total dissolved organic carbon (DOC) and total dissolved nitrogen (TDN) in the 0.5 M K<sub>2</sub>SO<sub>4</sub> extracts from both the fumigated and unfumigated samples were analysed with a Multi N/C 2100 analyser (AnalytikJena, Jena, Germany). C<sub>mic</sub> and N<sub>mic</sub> were determined as the difference of TOC or TN concentrations between the fumigated and non-fumigated samples, corrected by the extraction factor 0.35 (K<sub>EC</sub>) and 0.5 (K<sub>EN</sub>), respectively (Joergensen et al., 1998; Vance et al., 1987; Wu et al., 1990).

For  $S_{mic}$ , the total dissolved S in the fumigated and unfumigated extracts was determined by ICP-OES.

## 7.2.6 Statistical and data analysis

### 7.2.6.1 Half-time of amino acid- $^{14}C$ mineralization

By difference, the amount of respired  $^{14}CO_2$  was used to calculate the  $^{14}C$  remaining in soil and the microbial biomass pools (% of total added available to soil microbes). Previous studies have demonstrated that low MW substrate mineralization occurs in two distinct phases (Hill et al., 2008b; Wilkinson et al., 2014b). In this study, the mineralization of Cys and Met to  $^{14}CO_2$  was described by a double first order exponential decay equation:

$$y_I = (a_1 \times \exp^{-b_1 t}) + (a_2 \times \exp^{-b_2 t}) \quad (\text{Eqn. 1})$$

Where  $y_I$  the amount of  $^{14}C$  remaining in the soil and microbial biomass,  $t$  is time,  $b_1$  is the exponential coefficient describing the primary mineralization phase,  $b_2$  is the exponential coefficient describing the secondary mineralization of the microbial biomass, while  $a_1$  and  $a_2$  represent the size of two pools. Half-times of the soil/microbial biomass pools can therefore be defined as:

$$t_{1/2} = \ln(2)/b_1 \text{ or } t_{1/2} = \ln(2)/b_2 \quad (\text{Eqn. 2})$$

### 7.2.6.2 Half-time of amino acid- $^{14}C$ depletion in soil

To determine the half-time ( $t_{1/2}$ ) of amino acid- $^{14}C$  depletion in the soil solution, a double first order exponential decay was fitted to the experimental data according to:

$$y_2 = (a_3 \times \exp^{-b_3 t}) + (a_4 \times \exp^{-b_4 t}) \quad (\text{Eqn. 3})$$

Where  $y_2$  is the  $^{14}C$  remaining in soil solution,  $b_3$  and  $b_4$  the exponential coefficient describing the rate of amino acid- $^{14}C$  depletion by soil microbial community,  $a_3$  and  $a_4$  describes the sizes of pools and  $t$  is time. The half-time of the soil solution pools  $a_3$  and  $a_4$  can therefore be

calculated according to Eqn. 2. We assumed that the two exponential parts of the equation represented uptake of substrate by two independent carrier systems.

#### 7.2.6.3 Half-time of amino acid-<sup>35</sup>S depletion in soil

Similar to the approach taken for <sup>14</sup>C above, a double first order exponential decay equation was also used to calculate the <sup>35</sup>S-amino acid depletion from soil solution as follows:

$$y_3 = (a_5 \times \exp^{-b_5 t}) + (a_6 \times \exp^{-b_6 t}) \quad (\text{Eqn. 4})$$

Where  $y_3$  is the <sup>35</sup>S remaining in the soil solution,  $b_5$  and  $b_6$  the exponential coefficient describing depletion by soil microbial community,  $a_5$  and  $a_6$  describes the sizes of pools and  $t$  is time. The half-time of the soil solution pools  $a_5$  and  $a_6$  can therefore be calculated according to Eqn. 2. We assumed that the two exponential parts of the equation represented uptake of substrate by two independent carrier systems.

#### 7.2.6.4 Half-time of microbial biomass turnover in soil

The turnover time of  $C_{\text{mic}}$  and  $S_{\text{mic}}$  was estimated from the decrease of <sup>14</sup>C and <sup>35</sup>S in microbial biomass with time when a clear exponential decay pattern was detectable. Here, we focused solely on the production of C that was incorporated into microbial biomass, the respired C was excluded. Generally, a first order exponential decay equation fitted well to the data (based on the amount of <sup>14</sup>C or <sup>35</sup>S-labelled  $C_{\text{mic}}$  and  $S_{\text{mic}}$  at each sampling time) where:

$$A_t = A_0 \times \exp^{-kt} \quad (\text{Eqn. 5})$$

Where  $A_0$  and  $A_t$  are the content of <sup>14</sup>C<sub>mic</sub> or <sup>35</sup>S<sub>mic</sub> at times  $t = 0$  and  $t$ , respectively. The initial time point of  $t$  was set at 24 h after substrate addition, as this was the time at which an approximate equilibrium was attained between exchangeable soil C and <sup>14</sup>C, soil S and <sup>35</sup>S. In addition, the <sup>14</sup>C<sub>mic</sub> and <sup>35</sup>S<sub>mic</sub> reached the peak and started to decline at this time.  $k$  is the decay rate, therefore the turnover time is calculated by Eqn. 6:

$$t_{1/2} = \ln(2)/k \quad (\text{Eqn. 6})$$

All treatments were replicated three times. All statistical analyses were carried out using SPSS v25.0 (SPSS Inc., Chicago, IL, USA), with  $p < 0.05$  used as the upper limit for statistical significance. The exponential decay curves for  $^{14}\text{C}_{\text{mic}}$  and  $^{35}\text{S}_{\text{mic}}$  were fitted to the experimental results using SigmaPlot v13.0 (SPSS Inc., Chicago, IL), and the adjusted  $R^2$  for the curves as well as the significance of each parameter were calculated.

## 7.3 Results


### 7.3.1 Amino acid- $^{14}\text{C}$ mineralization in soil

Following addition of  $^{14}\text{C}$ -labeled Cys and Met to soil, there was an initial rapid phase of  $^{14}\text{CO}_2$  evolution followed by a secondary slower phase (Fig. 7.2). This pattern was similar in all treatments. The kinetics of  $^{14}\text{CO}_2$  evolution was biphasic for all treatments and the rate constants ( $b_1$  and  $b_2$ ) for these two phases are presented in Table 7.2. The loss of  $^{14}\text{CO}_2$  from the soil was best described by a double first order exponential equation (Eq (1); Fig. 7.2;  $r^2 > 0.99$ ), which is in line with our findings in Chapter 4, 5 and previous studies (Boddy et al., 2008; Glanville et al., 2012b; Mariano et al., 2016; Scow et al., 1986). The recovery of  $^{14}\text{CO}_2$  after the addition of  $^{14}\text{C}$ -glucose to the soil was significantly smaller than that recovered after the addition of  $^{14}\text{C}$ -amino acids ( $p < 0.05$ ).

Overall, the mineralization of added amino acid- $^{14}\text{C}$  by the microbial community was extremely rapid, suggesting that soil microbes are severely C-limited. Generally, the addition of glucose-C or nutrients did not affect the two-phase microbial respiration pattern of Cys or Met, yet noticeable differences in the rate constants for the two pools were found. Overall, addition of glucose-C increased the allocation of  $^{14}\text{C}$  to the fast pool, whereas the addition of NPS decreased allocation to this pool.

**Table 7.2** Kinetic coefficients of double first order exponential decay models describing the depletion of  $^{14}\text{C}$ -Cys or Met from soil by mineralization over a 168 h-incubation period in the absence (control) or presence of either glucose (+G) or nutrients (+NPS). The corresponding mineralization of  $^{14}\text{C}$ -glucose in the presence of Cys and Met is also presented.  $a_1$  and  $a_2$  are estimated pool sizes for the fast and slow phases of mineralization, and  $b_1$  and  $b_2$  are the rate constants for the fast and slow phases of mineralization, respectively.  $t_{1/2}$  values are the half-times for pool  $a_1$  and  $a_2$  determined from  $b_1$  and  $b_2$ . Values represent means  $\pm$  SEM ( $n = 3$ ). The  $R^2$  represents the goodness of fit to the experimental data.

Substrate	Treatments	$a_1$	$b_1$	$a_2$	$b_2$	$a_1 t_{1/2}$ (h)	$a_2 t_{1/2}$ (d)	$R^2$
Cys	+G	$65.1 \pm 2.7$	$0.07 \pm 0.003$	$30.4 \pm 2.9$	$0.004 \pm 0.0010$	$12.8 \pm 5.3$	$7.3 \pm 3.3$	0.99
	Control	$62.7 \pm 0.9$	$0.08 \pm 0.002$	$34.8 \pm 1.1$	$0.001 \pm 0.0003$	$9.2 \pm 8.4$	$23.1 \pm 5.2$	0.99
	+NPS	$44.7 \pm 2.0$	$0.08 \pm 0.006$	$51.7 \pm 2.2$	$0.003 \pm 0.0003$	$9.5 \pm 1.4$	$10.7 \pm 1.6$	0.99
Met	+G	$43.0 \pm 0.6$	$0.04 \pm 0.004$	$42.3 \pm 0.9$	$0.04 \pm 0.004$	$17.3 \pm 1.7$	$0.7 \pm 0.1$	0.99
	Control	$72.9 \pm 1.5$	$0.04 \pm 0.001$	$28.4 \pm 1.6$	$0.003 \pm 0.0005$	$18.6 \pm 5.9$	$29.6 \pm 7.3$	0.99
	+NPS	$49.3 \pm 2.0$	$0.07 \pm 0.005$	$53.4 \pm 2.2$	$0.003 \pm 0.0001$	$9.3 \pm 1.0$	$10.7 \pm 3.9$	0.99
Glucose	+Cys	$33.7 \pm 1.4$	$0.06 \pm 0.004$	$67.5 \pm 1.5$	$0.003 \pm 0.0002$	$11.9 \pm 2.4$	$72.4 \pm 3.7$	0.99
	+Met	$36.2 \pm 1.6$	$0.06 \pm 0.004$	$65.4 \pm 1.7$	$0.007 \pm 0.0001$	$12.9 \pm 2.4$	$40.4 \pm 6.4$	0.99



**Fig. 7.2** Cumulative  $^{14}\text{CO}_2$  evolution from a grassland soil after the addition of  $^{14}\text{C}$ -labelled Cys or Met (1 mM). Lines represent fits of a double exponential first kinetic equation to the experimental data. Values represent means  $\pm$  SEM ( $n = 3$ ).

There was a close similarity between the dynamics of  $^{14}\text{CO}_2$  evolution from Cys and Met throughout the 7-d period. Within the first 48 h, the proportion of Cys- $^{14}\text{C}$  mineralized was higher than that of Met. However, due to Met being more enriched in C on a molar basis, the actual rate of mineralization of Met- $^{14}\text{C}$  started to exceed that of Cys- $^{14}\text{C}$  10 h after addition. According to our calculation, the mineralization rate for Met for the first 10 h after addition remained relatively constant, ranging from  $292 \pm 31$  to  $309 \pm 12 \mu\text{g } ^{14}\text{C-Cys DW soil kg}^{-1} \text{ h}^{-1}$ , which is lower than that for Cys, ranging from  $297 \pm 71$  to  $699 \pm 79 \mu\text{g } ^{14}\text{C-Met DW soil kg}^{-1} \text{ h}^{-1}$ . From 10 h onwards, the mineralization rates of Cys decreased dramatically to slightly below that of Met.

The half-times of the  $^{14}\text{C}$  substrates in the soil solution, calculated using the double exponential kinetic model, are presented in Table 7.2. The results showed that the turnover of both glucose and amino acids in soil solution was extremely rapid, with an average first phase half-time of  $17.7 \pm 4.4$  h. ANOVA indicated that neither substrate type nor nutrient amendment had a significant effect on the rate of loss of these compounds from soil solution ( $p > 0.05$ ). In contrast to the half-time for the first phase of mineralisation ( $12.4 \pm 2.2$  h for glucose- $^{14}\text{C}$  and

12.8 ± 4.9 h for amino acid-<sup>14</sup>C), the rate constant for the second phase was significantly longer, with an average value of 56.4 ± 18.2 d for glucose-<sup>14</sup>C and 16.3 ± 9.3 d for the amino acid-<sup>14</sup>C.

Studies on the uptake and use of amino acids in pure microbial cultures have shown that they are transported into the cell by amino acid group (e.g. neutral, basic, acidic) and isomer-specific membrane transporters (Popova, Dietz and Golldack, 2003; Tilsner *et al.*, 2005), and are differentially metabolized (Bender, 2012). It is likely that internal metabolic control of microbial amino acid transport and synthesis pathways significantly influence amino acid mineralization rates. Previous studies have shown that bacteria, actinomycetes, and filamentous fungi are all capable of decomposing Met, with the release of ammonia, volatile S and ketobutyric acid (Segal and Starkey, 1969), while a mixed population of soil microorganisms oxidized Cys to sulphate (Freney, 1967). Our results suggested that although Cys and Met are both weakly sorbed in soil (Data shown in Fig. 3.1, Chapter 3), they exhibited very different respiration rates and microbial assimilation efficiencies (Ma) ( $\text{Ma} = \text{biomass-}^{14}\text{C} / (\text{biomass-}^{14}\text{C} + ^{14}\text{CO}_2)$ ).


### **7.3.2 Amino acid depletion from soil**

#### *7.3.2.1 Amino acid-<sup>14</sup>C depletion from soil*

Overall, the depletion of both <sup>14</sup>C-labelled Cys and Met from soil solution was extremely rapid (Fig. 7.3). For example, after 3 h, 81.2 ± 0.6% of the added Cys had been taken up by the microbial community, whereas 74.4 ± 0.5% of the Met was taken up. At the end of the 24 h incubation period, marginally more <sup>14</sup>C-Met remained in the soil than <sup>14</sup>C-Cys ( $p < 0.01$ ) in all three treatments. In addition, after 24 h, the content of 0.5 M K<sub>2</sub>SO<sub>4</sub>-extractable <sup>14</sup>C in the soil had declined to < 10% of the applied amino acid-<sup>14</sup>C, indicating that > 90% of the added amino acid-<sup>14</sup>C was consumed by the microbial community. This rapid disappearance

of substrate- $^{14}\text{C}$  is consistent with other studies (Vinolas, Vallejo and Jones, 2001; Jones *et al.*, 2018).

No significant difference ( $p > 0.05$ ) between the amount of  $\text{K}_2\text{SO}_4$ -extractable  $^{14}\text{C}$  was observed in the three different treatments. In contrast, the extractable  $^{14}\text{C}$  content derived from glucose in soil for the first 12 h was higher by a factor of two than that derived from both amino acids, and smaller by a factor of three afterwards. This indicates different utilization mechanisms for glucose and the two amino acids by the microbial community.



**Fig. 7.3** Amount of  $^{14}\text{C}$ -label remaining in soil solution after addition of  $^{14}\text{C}$ -Cys or Met to a grassland soil relative to the amount added at time zero (100%). Values represent means  $\pm$  standard deviation ( $n = 3$ ). Lines represent fits of a double first order exponential decay equation to the experimental data ( $r^2 > 0.80$  in all cases; Eq. (3)).


#### 7.3.2.2 Amino acid- $^{35}\text{S}$ depletion from soil

In general,  $^{35}\text{S}$ -Cys and  $^{35}\text{S}$ -Met depletion from the soil solution was also extremely rapid, with a much higher proportion of  $^{35}\text{S}$ -Cys removed from the soil solution than  $^{35}\text{S}$ -Met. After only 6 h, the total  $\text{CaCl}_2$  extractable  $^{35}\text{S}$  ( $\text{SO}_4^{2-}\text{-S} + \text{org-S}$ ) derived from Cys and Met control treatments was  $52.1 \pm 1.2$  and  $25.5 \pm 4.1\%$ , respectively, suggesting around 47.9% and 74.5% of the Cys- $^{35}\text{S}$  and Met- $^{35}\text{S}$  was removed from soil solution by microorganisms. Three



days after substrate addition, the total extractable  $^{35}\text{S}$  content significantly decreased, by more than half, and then remained consistently low ( $< 15\%$  of added) until the end of the experiment. The addition of glucose-C or inorganic nutrients did not affect this trend significantly ( $p > 0.05$ ). The depletion of  $^{35}\text{S}$ -Cys and  $^{35}\text{S}$ -Met from soil solution was best described by a double first order exponential decay equation ( $r^2 > 0.80$  in all cases; Eq, (3); Fig. 7.4).


As sources of soil sulphate, our results showed that Cys is more readily mineralized than Met. By applying the precipitation method, we found that the majority of  $^{35}\text{S}$  radioactivity detected from soil solution was recovered as inorganic sulphate, indicating that immediately after addition of  $^{35}\text{S}$ -Cys and Met,  $^{35}\text{S}$  was mineralised and released back to the soil as sulphate. This sulphate pool was then gradually incorporated back into the microbial biomass or was present as non-extractable organic sulphur compounds. While there was clear evidence of the oxidation of Cys and Met to sulphate, the mechanisms by which they were decomposed, and the intermediates formed during this process remain unknown.



**Fig. 7.4** Amount of  $^{35}\text{S}$ -label remaining in soil solution after addition of  $^{35}\text{S}$ -labelled Cys or Met to grassland soil relative to the amount added at time zero (100%). Values represent means  $\pm$  standard deviation ( $n = 3$ ). Lines represent fit of a double first order exponential decay equation to the experimental data ( $r^2 > 0.80$  in all cases; Eq, (4)).

### 7.3.3 Soil microbial biomass C and N

Within 9 h of substrate addition,  $C_{mic}$  increased from  $782 \pm 52$  mg C kg<sup>-1</sup> DW soil (Control soil) to  $1006 \pm 20$  mg C kg<sup>-1</sup> soil and  $917 \pm 67$  mg C kg<sup>-1</sup> soil for Cys and Met, respectively (Fig. 7.5). Thereafter, it decreased gradually to the level seen before amino acid addition. An increase in  $C_{mic}$  concentrations due to glucose addition alone were significant ( $p < 0.01$ ) for the whole incubation period; the extent of the increase varied from 47.6 to 76.9% with time, whereas the addition of S combined with NPS did not affect  $C_{mic}$  to a significant extent ( $p > 0.05$ ).



**Fig. 7.5** Dissolved organic carbon (TOC; mg C kg<sup>-1</sup> DW soil) and N (TDN; mg N kg<sup>-1</sup> DW soil) concentrations in grassland soil solution subjected to treatments of C or NPS addition. Values represent means  $\pm$  standard deviation ( $n = 3$ ).


Similarly,  $N_{mic}$  decreased gradually after substrate addition. NPS addition did not greatly influence this trend, whereas following glucose-C addition,  $N_{mic}$  increased from  $63.3 \pm 13.2$  (Control soil) to  $69.4 \pm 7.9$  mg  $N_{mic}$  kg<sup>-1</sup> DW soil and  $66.5 \pm 1.1$  mg  $N_{mic}$  kg<sup>-1</sup> soil within 9 h for Cys and Met individually. After 9 h,  $N_{mic}$  content decreased to a level similar to seen before substrate addition and then remained relatively stable for the rest of the incubation period.

### 7.3.4 Turnover time of microbial biomass <sup>14</sup>C and <sup>35</sup>S

The turnover time of <sup>14</sup>C and <sup>35</sup>S in the microbial community was estimated from the decline of radioisotope in the biomass over time. In this study, between 24-168 h was chosen to calculate this because isotope incorporation into the microbial biomass between this time-period followed a clear first order exponential decay (Table 7.3 and Table 7.4).

#### 7.3.4.1 <sup>14</sup>C tracer incorporation into microbial biomass and turnover

During the incubation period, <sup>14</sup>C recovery in the microbial biomass increased immediately after substrate addition, then decreased afterwards. Overall, the <sup>14</sup>C recovery rate from Met in the microbial biomass was slightly higher than that from <sup>14</sup>C-Cys, indicating different amino acid-C uptake mechanisms (Fig. 7.6). After 24 h, as much as 30% of the <sup>14</sup>C-Cys was recovered in the microbial biomass. The <sup>14</sup>C recovery in the microbial biomass was close to the glucose and NPS amended soils, showing that C and nutrient addition did not significantly change the microbial carbon uptake of Cys or Met-C. The <sup>14</sup>C decline rate in MBC was best fitted with an exponential decay equation. From this, the calculated half-time for <sup>14</sup>C-Met in the biomass was 6.3 d, this is quite similar to that of <sup>14</sup>C-Cys (6.2 d). There were no significant differences among the three treatments. We estimated that the degradation of MBC was the main source of second phase microbial mineralization for both Cys and Met, which could be regulated by a range of factors.




**Fig. 7.6** The fraction of  $^{14}\text{C}$ - $\text{C}_{\text{mic}}$  in soil after following the addition of amino acids to soil in the presence or absence of glucose-C or inorganic nutrients (+NPS). Values represent means  $\pm$  SEM ( $n = 3$ ). Lines represent  $^{14}\text{C}_{\text{mic}}$  obtained by the fumigation extraction method, adjusted with the extraction factor of 0.35.

#### 7.3.4.2 $^{35}\text{S}$ tracer incorporation into microbial biomass and turnover

After amino acid addition, there was a rapid incorporation of  $^{35}\text{S}$  into the microbial biomass for the first 12 h, after which a gradual decrease followed. A higher proportion of  $^{35}\text{S}$  derived from Met was measured in microbial biomass than from Cys. Three hours after substrate addition, 38.2% of  $^{35}\text{S}$ -Cys was recovered in the microbial biomass, while 67.6% was recovered from the  $^{35}\text{S}$ -Met. This result is consistent with  $^{35}\text{S}$  depletion from soil solution, where an opposite trend was found for the two amino acids.

The soil microbial biomass S fraction in soil is often considered to be relatively labile and the most active S pool. Clearly,  $^{35}\text{S}$  was turning over rapidly in microbial the biomass. Specifically, the turnover time of  $^{35}\text{S}_{\text{mic}}$  was longest in soil amended with NPS (10.4 and 15.6 d for Cys and Met, respectively), followed by control soil (7.4 and 10.1 d), and soil amended with glucose (3.5 and 7.8 d). Overall, the turnover of  $^{35}\text{S}$ -Cys was faster than  $^{35}\text{S}$ -Met (Fig. 7.7, Table 7.4).



**Fig. 7.7** The fraction of  $^{35}\text{S}$  labelled  $\text{S}_{\text{mic}}$  after substrate addition. Values represent means  $\pm$  SEM ( $n = 3$ ). Lines represent  $^{35}\text{S}_{\text{mic}}$  obtained by fumigation extraction method, adjusted with the extraction factor of 0.35.

**Table 7.3** Turnover of soil microbial biomass  $^{14}\text{C}$ -labelled Cays or Met in a grassland soil after either glucose-C (+G) or nutrient (+NPS) input. The corresponding mineralization of  $^{14}\text{C}$ -glucose in the presence of Cys and Met is also presented. A first order exponential decay equation Eqn. 5 was fitted to the experimental data. Values represent means  $\pm$  SEM ( $n = 3$ ). The  $R^2$  represents the goodness of fit to the experimental data.


Substrate	Treatments	$A_0\text{-}^{14}\text{C}$	$K\text{-}^{14}\text{C}$	$t_{1/2}$ (d)	$R^2$
Cys	+G	$21.3 \pm 2.8$	$0.013 \pm 0.002$	$3.7 \pm 0.8$	0.92
	Control	$35.8 \pm 7.5$	$0.008 \pm 0.003$	$6.2 \pm 0.5$	0.69
	+NPS	$39.6 \pm 3.9$	$0.008 \pm 0.001$	$6.5 \pm 2.2$	0.90
Met	+G	$42.4 \pm 5.8$	$0.013 \pm 0.002$	$4.3 \pm 0.8$	0.91
	Control	$49.8 \pm 7.4$	$0.009 \pm 0.002$	$6.3 \pm 1.4$	0.83
	+NPS	$56.3 \pm 5.9$	$0.009 \pm 0.001$	$6.7 \pm 1.6$	0.91
Glucose	+Cys	$35.3 \pm 3.7$	$0.008 \pm 0.001$	$6.5 \pm 1.3$	0.89
	+Met	$47.9 \pm 5.9$	$0.008 \pm 0.002$	$4.8 \pm 0.8$	0.89

**Table 7.4** Turnover of soil microbial biomass  $^{35}\text{S}$ -Cys or Met in grassland soil after glucose-C (+G) or nutrient (+NPS) input. A first order exponential decay equation Eqn. 5 was fitted to the experimental data. Values represent means  $\pm$  SEM ( $n = 3$ ). The  $R^2$  represents the goodness of fit to the experimental data.

Substrate	Treatments	$A_0\text{-}^{35}\text{S}$	$K\text{-}^{35}\text{S}$	$t_{1/2}$ (d)	$R^2$
Cys	+G	$32.2 \pm 6.2$	$0.013 \pm 0.003$	$3.6 \pm 1.5$	0.83
	Control	$43.3 \pm 5.5$	$0.006 \pm 0.002$	$7.4 \pm 2.0$	0.74
	+NPS	$63.5 \pm 6.7$	$0.004 \pm 0.001$	$10.4 \pm 2.6$	0.73
Met	+G	$50.7 \pm 3.5$	$0.005 \pm 0.001$	$7.8 \pm 0.6$	0.80
	Control	$80.8 \pm 9.0$	$0.002 \pm 0.001$	$10.1 \pm 3.2$	0.67
	+NPS	$82.4 \pm 5.9$	$0.003 \pm 0.001$	$15.7 \pm 1.4$	0.71

### 7.3.5 Inorganic nitrogen release from cysteine and methionine decomposition

It is now well established that the breakdown of organic N compounds in soil involves a cascade of N transformations (e.g. ammonification, nitrification) (Verma, Chaudhary and Goyal, 2018; Fujii *et al.*, 2020; Ma *et al.*, 2021). In this study,  $\text{NH}_4^+$  and  $\text{NO}_3^-$  concentrations over the 7-day incubation period was measured. Control treatments were included where substrate solution was replaced with distilled water.



**Fig. 7.8** Generation of ammonium and nitrate from added Cys and Met (1 mM) in soil solution over a 7-day period. Data expressed on a dry soil weight basis. Values represent means  $\pm$  SEM ( $n = 3$ ).

After a preliminary lag phase of about 24 h, the production of ammonium reached a maximum for all treatments (Fig. 7.8). During this ammonification period, there was a sharp rise in the rate of  $\text{CO}_2$  loss from the soil, indicating an extensive breakdown of the amino acids.

After ammonium-N generation by the breakdown of amino acids, nitrification occurred, with ca. 70% of the ammonium-N being converted to nitrate-N. The course of ammonification followed by nitrification is shown in Fig. 7.8.

## 7.4 Discussion

Immediately after substrate addition, amino acid- $^{14}\text{C}$  was rapidly removed from soil solution by soil microbes, with half-lives calculated to be  $< 3$  h, suggesting that the free amino acid pool in soil turns over many times daily. This supports previous studies that have also revealed short amino acid half-lives in soil (Geisseler & Horwath, 2014; Ma et al., 2017). The results from all three treatments all conformed to a biphasic exponential dynamics of  $^{14}\text{CO}_2$  evolution, similar to those reported previously (Chotte et al., 1997; Glanville et al., 2012a; Saggar et al., 1996). Overall, the mineralization of Cys by the soil microbial community was extremely rapid, with the first rapid phase of  $\text{CO}_2$  production attributable to the immediate use of the substrate in catabolic process (i.e. respiration), and the remaining substrate being taken up and immobilized in the microbial biomass (i.e. formation of new biomass or storage polymers). The slower second phase of  $^{14}\text{CO}_2$  production is then attributable to the subsequent turnover of the soil microbial community or storage polymers leading to the production of  $^{14}\text{CO}_2$ . Glucose addition promoted greater mineralization of both Cys and Met, while it had less of an effect on the proportion of  $^{14}\text{C}$  incorporated in microbial biomass, resulting in a more complete utilization of both amino acids. In contrast, nutrient (NPS) addition had less effect on Cys and Met mineralization. We ascribe the lack effect of NPS addition to the low C/N ratio of these amino acids (C/N ratio = 3 for Cys and 5 for Met) which results in excess microbial  $\text{NH}_4^+$  being excreted into the soil.

It has been demonstrated that, in the case of simple and generally weakly sorbed amino acids, soil microorganisms tend to take up intact amino acids into the cells using specific

transport proteins (Padan, 2009), instead of mineralizing the amino acids outside the cell prior to the assimilation of the remaining C skeleton. To determine the mineralization rate of amino acid-N in soil, the excretion of amino acid derived  $\text{NH}_4^+$  and  $\text{NO}_3^-$  into the soil was measured over the 7-day incubation period. We provide clear evidence of successive stages of amino acid breakdown. A rapid loss of amino acid was associated with an increase of free ammonium in the soil solution, as well as a sharp rise in the  $^{14}\text{CO}_2$  release indicating an extensive breakdown of the added amino acid. Then the ammonium-N produced by the breakdown of amino acids in soil becomes available to the nitrifier community leading to  $\text{NO}_3^-$  production.

Sulphur mineralization from organic-S together with immobilization of new released inorganic sulphate regulates S availability to plants (Li et al., 2001). In this study, S derived from both Cys and Met was rapidly converted to sulphate. Results indicated that conversion of Cys and Met to sulphate was almost complete 24 h after amino acid addition. This is in line with previous studies (Meena and Improvement, 2018). A similar trend for the distribution of amino acid-S between sulphate release and microbial biomass S incorporation was found for Cys and Met. Amount of inorganic sulphate decreased gradually throughout the incubation period, indicating a re-capture of sulphate by microorganisms. The proportion of amino acid-S incorporated into the microbial biomass showed a similar trend through the incubation, although lagged behind inorganic sulphate production:  $^{14}\text{C}_{\text{mic}}$  reached a maximum before our first sampling (3 h after substrate addition), while  $^{35}\text{S}_{\text{mic}}$  reached maximum with a slight lag ( $^{35}\text{S}_{\text{mic}}$  from Cys peaked around 9 h after substrate addition, while  $^{35}\text{S}_{\text{mic}}$  from Met reached peak after 6 h).

It is also interesting to note that by the end of the incubation there was incomplete recovery of the total  $^{35}\text{S}$  added in both amino acids (total recovery < 100%). It is possible that this may be associated with analytical errors, including incomplete mixing of soil sampled. But some of this ‘missing’ S could also be due to volatile S emissions from soil (Banwart and




Bremner, 1975; Brown *et al.*, 2021), or incorporation into organic matter or absorbed by mineral colloids (Gustafsson, Akram and Tiberg, 2015; Heinze *et al.*, 2021) and therefore not extractable by 0.01 M CaCl<sub>2</sub>. In this study, we set up ethanol traps to capture any volatile sulphur compounds released, yet these all proved to have minimal <sup>35</sup>S activity. This does not necessarily mean that volatile sulphur compounds were not produced by microbial decomposition of Cys and Met, just that we used the wrong approach to capture them. Previous studies have shown that some soils have a substantial capacity for sorption of volatile sulphur suggesting that a heating step may be needed to induce their desorption from soil prior to capture (Banwart and Bremner, 1975; Ko and Chu, 2005; Ko, Chu and Tseng, 2006). Further work is required to investigate the microbial use of other dissolved organic sulphur (DOS) compounds and also compare microbial preference for S forms when they are exposed to multiple sources of DOS compounds, as would occur under field conditions.

## **7.5 Conclusions and outlook**

In conclusion, our results indicate that decomposition of S-containing amino acids (Cys and Met) to CO<sub>2</sub>, NH<sub>4</sub><sup>+</sup>, NO<sub>3</sub><sup>-</sup>, and SO<sub>4</sub><sup>2-</sup> in soil mainly occurs as a result of biological oxidation (Fig. 7.9). This is also in agreement with previous studies on the breakdown of low molecular weight organic acids in soil (Roy and White, 2013; Tanikawa *et al.*, 2013; Creamer *et al.*, 2014). Due to the low C/N ratio of Cys and Met, short-term addition of inorganic NPS to soil had no detectable effect on amino acid mineralization rates, while the addition of a readily available C source, such as glucose, led to a high and rapid rate of sulphate incorporation into soil organic fractions.

Understanding S transformations in soils is necessary for predicting the S supply from the soil organic matter to crops. The evidence is conclusive that there is rapid interconversion of organic and inorganic S forms in the soil, such as mineralization (the transformation of

organic S into inorganic sulphate) and immobilization (incorporation of sulphate into soil organic compounds or soil microbial biomass). This cycling is microbiologically mediated. Yet little is still known about the specific microbial species or genera that play important roles in the soil S transformation cycle.



**Fig. 7.9** Schematic presentation of carbon, nitrogen and sulphur mineralization from the S-containing amino acid-cysteine in grassland soils. a: labelled amino acid uptake into soil microbial biomass; b: subsequent microbial mineralization.

## 7.6 Acknowledgements

I would like to thank Joseph Cotton and Mark Hughes for soil sample collection, and Jonathan Roberts for assistance with isotope analysis in the laboratory. This work was performed during a studentship funded by Bangor University and the China Scholarship Council (201606510012).

## 7.7 References

- Allaway, W. H. and Thompson, J. F. (1966) 'Sulfur in the nutrition of plants and animals', *Soil Science*, 101(4), pp. 240–247. doi: 10.1097/00010694-196604000-00003.
- Banwart, W. L. and Bremner, J. M. (1975) 'Formation of volatile sulfur compounds by microbial decomposition of sulfur-containing amino acids in soils', *Soil Biology and Biochemistry*, pp. 359–364. doi: 10.1016/0038-0717(75)90050-4.
- Banwart, W L and Bremner, J. M. (1975) 'Formation of volatile sulfur compounds by microbial decomposition of sulfur-containing amino acids in soils', *Soil Biology and Biochemistry*, pp. 359–364. doi: 10.1016/0038-0717(75)90050-4.
- Banwart, W. L. and Bremner, J. M. (1976) 'Evolution of volatile sulfur compounds from soils treated with sulfur-containing organic materials', *Soil Biology and Biochemistry*, 8(5), pp. 439–443. doi: 10.1016/0038-0717(76)90047-X.
- Bender, D. A. (2012) *Amino acid metabolism*. John Wiley & Sons.
- Blagodatskaya, E. and Kuzyakov, Y. (2013) 'Active microorganisms in soil: Critical review of estimation criteria and approaches', *Soil Biology and Biochemistry*. Elsevier, pp. 192–211. doi: 10.1016/j.soilbio.2013.08.024.
- Blagodatskaya, E. V *et al.* (2007) 'Priming effects in Chernozem induced by glucose and N in relation to microbial growth strategies', *Applied Soil Ecology*. Elsevier, 37(1–2), pp.

95–105. doi: 10.1016/j.apsoil.2007.05.002.

Blagodatskaya, E. V *et al.* (2009) ‘Contrasting effects of glucose, living roots and maize straw on microbial growth kinetics and substrate availability in soil’, *European Journal of Soil Science*, 60(2), pp. 186–197. doi: 10.1111/j.1365-2389.2008.01103.x.

Boddy, E. *et al.* (2007) ‘Fast turnover of low molecular weight components of the dissolved organic carbon pool of temperate grassland field soils’, *Soil Biology and Biochemistry*, 39(4), pp. 827–835. doi: 10.1016/j.soilbio.2006.09.030.

Brailsford, F. (2019) ‘Dissolved organic matter (DOM) in freshwater ecosystems’. Bangor University.

Brown, R. W. *et al.* (2021) ‘Dependence of thermal desorption method for profiling volatile organic compound (VOC) emissions from soil’, *Soil Biology and Biochemistry*. Elsevier Ltd, 160(March), p. 108313. doi: 10.1016/j.soilbio.2021.108313.

Creamer, C. A. *et al.* (2014) ‘Stoichiometric controls upon low molecular weight carbon decomposition’, *Soil Biology and Biochemistry*. Elsevier Ltd, 79, pp. 50–56. doi: 10.1016/j.soilbio.2014.08.019.

David, M. B. *et al.* (1983) ‘Importance of organic and inorganic sulfur to mineralization processes in a forest soil’, *Soil Biology and Biochemistry*, 15(6), pp. 671–677. doi: 10.1016/0038-0717(83)90031-7.

DEFRA (2016) *British Survey of Fertiliser Practise*.

Eriksen, J. (2009) ‘Chapter 2 Soil Sulfur Cycling in Temperate Agricultural Systems’, *Advances in Agronomy*, 102(09), pp. 55–89. doi: org/10.1016/S0065-2113(09)01002-5.

Farrell, M. *et al.* (2011) ‘Rapid peptide metabolism: A major component of soil nitrogen

- cycling?', *Global Biogeochemical Cycles*, 25(3), pp. 1–11. doi: 10.1029/2010GB003999.
- Fermoso, F. G. *et al.* (2019) *Trace Elements in Anaerobic Biotechnologies, Trace Elements in Anaerobic Biotechnologies*. doi: 10.2166/9781789060225.
- Fitzgerald, J. W. and Andrew, T. L. (1984) 'Mineralization of methionine sulphur in soils and forest floor layers', *Soil Biology and Biochemistry*, 16(6), pp. 565–570. doi: 10.1016/0038-0717(84)90073-7.
- Freney, J. (1960) 'The Oxidation of Cysteine to Sulphate in Soil', *Australian Journal of Biological Sciences*, 13(3), p. 387. doi: 10.1071/bi9600387.
- Freney, J. R. (1967) 'Oxidation of sulphur in soils', *Mineralium Deposita*, 2(3), pp. 181–187. doi: 10.1007/BF00201914.
- Freney, J. R., Melville, G. E. and Williams, C. H. (1975) 'Soil organic matter fractions as sources of plant-available sulphur', *Soil Biology and Biochemistry*, 7(3), pp. 217–221. doi: 10.1016/0038-0717(75)90041-3.
- Fujii, K. *et al.* (2020) 'Decoupling of protein depolymerization and ammonification in nitrogen mineralization of acidic forest soils', *Applied Soil Ecology*. Elsevier, 153(January), p. 103572. doi: 10.1016/j.apsoil.2020.103572.
- Geisseler, D. and Horwath, W. R. (2014) 'Investigating amino acid utilization by soil microorganisms using compound specific stable isotope analysis', *Soil Biology and Biochemistry*. Elsevier Ltd, 74, pp. 100–105. doi: 10.1016/j.soilbio.2014.02.024.
- Ghani, A., McLaren, R. G. and Swift, R. S. (1992) 'Sulphur mineralisation and transformations in soils as influenced by additions of carbon, nitrogen and sulphur', *Soil Biology and Biochemistry*, 24(4), pp. 331–341. doi: 10.1016/0038-0717(92)90193-2.

- Gustafsson, J. P., Akram, M. and Tiberg, C. (2015) 'Predicting sulphate adsorption/desorption in forest soils: Evaluation of an extended Freundlich equation', *Chemosphere*. Elsevier Ltd, 119, pp. 83–89. doi: 10.1016/j.chemosphere.2014.05.067.
- Heinze, S. *et al.* (2021) 'Microbial biomass sulphur—an important yet understudied pool in soil', *Agronomy*, 11(8), pp. 1–13. doi: 10.3390/agronomy11081606.
- Hill, P. W., Farrar, J. F. and Jones, D. L. (2008) 'Decoupling of microbial glucose uptake and mineralization in soil', *Soil Biology and Biochemistry*, 40(3), pp. 616–624. doi: 10.1016/j.soilbio.2007.09.008.
- Holtzhauer, M. (2006) 'Radioactive labeling'. Springer.
- Jan, M. T. *et al.* (2009) 'Protein breakdown represents a major bottleneck in nitrogen cycling in grassland soils', *Soil Biology and Biochemistry*. Elsevier Ltd, 41(11), pp. 2272–2282. doi: 10.1016/j.soilbio.2009.08.013.
- Jansson, S. L. and Persson, J. (2015) 'Mineralization and immobilization of soil nitrogen', in *Nitrogen in Agricultural Soils*. Wiley Online Library, pp. 229–252. doi: 10.2134/agronmonogr22.c6.
- Jones, D. L. *et al.* (2009) 'Soil organic nitrogen mineralization across a global latitudinal gradient', *Global Biogeochemical Cycles*, 23(1), pp. 1–5. doi: 10.1029/2008GB003250.
- Jones, D. L. *et al.* (2018) 'Role of substrate supply on microbial carbon use efficiency and its role in interpreting soil microbial community-level physiological profiles (CLPP)', *Soil Biology and Biochemistry*, 123(May), pp. 1–6. doi: 10.1016/j.soilbio.2018.04.014.
- Kandeler, E. *et al.* (1999) 'Xylanase, invertase and protease at the soil-litter interface of a loamy sand', *Soil Biology and Biochemistry*, 31(8), pp. 1171–1179. doi: 10.1016/S0038-

0717(99)00035-8.

- Kertesz, M. A. and Mirleau, P. (2004) 'The role of soil microbes in plant sulphur nutrition', *Journal of Experimental Botany*, 55(404), pp. 1939–1945. doi: 10.1093/jxb/erh176.
- Kirkby, C. A. *et al.* (2011) 'Stable soil organic matter: A comparison of C:N:P:S ratios in Australian and other world soils', *Geoderma*. Elsevier B.V., 163(3–4), pp. 197–208. doi: 10.1016/j.geoderma.2011.04.010.
- Ko, T. H. and Chu, H. (2005) 'Spectroscopic study on sorption of hydrogen sulfide by means of red soil', *Spectrochimica Acta - Part A: Molecular and Biomolecular Spectroscopy*, 61(9), pp. 2253–2259. doi: 10.1016/j.saa.2004.09.016.
- Ko, T. H., Chu, H. and Tseng, J. J. (2006) 'Feasibility study on high-temperature sorption of hydrogen sulfide by natural soils', *Chemosphere*, 64(6), pp. 881–891. doi: 10.1016/j.chemosphere.2006.01.056.
- Kumar, K. and Goh, K. M. (1999) 'Crop residues and management practices: effects on soil quality, soil nitrogen dynamics, crop yield, and nitrogen recovery', *Advances in agronomy*. Elsevier, 68, pp. 197–319.
- Kumar, K. and Goh, K. M. (2003) 'Nitrogen release from crop residues and organic amendments as affected by biochemical composition', *Communications in Soil Science and Plant Analysis*, 34(17–18), pp. 2441–2460. doi: 10.1081/CSS-120024778.
- Levine, N. D. (1982) 'The genus *atoplasma* (Protozoa, Apicomplexa).', *The Journal of parasitology*, 68(4), pp. 719–723. doi: 10.2307/3280933.
- Li, S., Lin, B. and Zhou, W. (2001) 'Soil organic sulfur mineralization in the presence of growing plants under aerobic or waterlogged conditions', *Soil Biology and Biochemistry*, 33(6), pp. 721–727. doi: 10.1016/S0038-0717(00)00205-4.

- Liang, Z. *et al.* (2011) ‘Effects of elevated CO<sub>2</sub> concentration and nitrogen deposition on the biomass accumulation and allocation in south subtropical main native tree species and their mixed communities.’, *Yingyong Shengtai Xuebao*, 22(8).
- Ma, H. *et al.* (2017) ‘Mineralization of amino acids and its signs in nitrogen cycling of forest soil’, *Shengtai Xuebao/ Acta Ecologica Sinica*. Ecological Society of China, 37(1), pp. 60–63. doi: 10.1016/j.chnaes.2016.09.001.
- Ma, H. *et al.* (2021) ‘Contrasting Effects of Alanine and Methionine on Nitrogen Ammonification and Nitrification, and Nitrous Oxide Emissions in Subtropical Forest Soil’, *Journal of Soil Science and Plant Nutrition*. Springer International Publishing, pp. 2967–2979. doi: 10.1007/s42729-021-00582-1.
- Marschner, B. and Kalbitz, K. (2003) ‘Controls of bioavailability and biodegradability of dissolved organic matter in soils’, *Geoderma*. Elsevier, 113(3–4), pp. 211–235.
- Marschner, P. and Rengel, Z. (2012) *Soil Biology, Nutrient Cycling in Terrestrial Ecosystems*, Springer. doi: 10.1017/CBO9781107415324.004.
- McGill, W. B. and Cole, C. V. (1981) ‘Comparative aspects of cycling of organic C, N, S and P through soil organic matter’, *Geoderma*. Elsevier, 26(4), pp. 267–286. doi: 10.1016/0016-7061(81)90024-0.
- Meena, V. S. and Improvement, C. (2018) *Role of Rhizospheric Microbes in Soil, Role of Rhizospheric Microbes in Soil*. doi: 10.1007/978-981-13-0044-8.
- Nader, C. J. and Walker, D. J. (1970) ‘Metabolic fate of cysteine and methionine in rumen digesta.’, *Applied microbiology*, 20(5), pp. 677–681. doi: 10.1128/AEM.20.5.677-681.1970.
- Padan, E. (2009) ‘Bacterial membrane transport: superfamilies of transport proteins’, *eLS*.



Wiley Online Library.

- Popova, O. V., Dietz, K. J. and Golldack, D. (2003) 'Salt-dependent expression of a nitrate transporter and two amino acid transporter genes in *Mesembryanthemum crystallinum*', *Plant Molecular Biology*, 52(3), pp. 569–578. doi: 10.1023/A:1024802101057.
- Roy, E. D. and White, J. R. (2013) 'Measurements of nitrogen mineralization potential in wetland soils', *Methods in Biogeochemistry of Wetlands*. Soil Science Society of America Madison, WI, pp. 465–471.
- Ruiz-Herrera, J. and Starkey, R. L. (1969a) 'Dissimilation of methionine by a demethiolase of *Aspergillus* species.', *Journal of bacteriology*, 99(3), pp. 764–770. doi: 10.1128/jb.99.3.764-770.1969.
- Ruiz-Herrera, J. and Starkey, R. L. (1969b) 'Dissimilation of methionine by fungi.', *Journal of Bacteriology*, 99(2), pp. 544–551. doi: 10.1128/jb.99.2.544-551.1969.
- S.Combs, J.Denning, K. D. F. (1998) 'Sulfate-Sulfur', *Recommended Chemical Soil Test Procedures for the North Central Region Agricultural*, 221(221), pp. 35–41.
- Sahrawat, K. L. (1981) 'Mineralization of biuret nitrogen in soil', *Plant and Soil*, 62(3), pp. 469–471. doi: 10.1007/BF02374144.
- Scherer, H. W. (2009) 'Sulfur in soils', *Journal of Plant Nutrition and Soil Science*, 172(3), pp. 326–335. doi: 10.1002/jpln.200900037.
- Schimel, J. P. and Bennett, J. (2004) 'Nitrogen mineralization: challenges of a changing paradigm', *Ecology*. Wiley Online Library, 85(3), pp. 591–602.
- Scott, N. M., Bick, W. and Anderson, H. A. (1981) 'The measurement of sulphur-containing amino acids in some scottish soils', *Journal of the Science of Food and Agriculture*, 32(1), pp. 21–24. doi: 10.1002/jsfa.2740320105.

- Segal, W. and Starkey, R. L. (1969) 'Microbial decomposition of methionine and identity of the resulting sulfur products.', *Journal of Bacteriology*, 98(3), pp. 908–913.
- Sillen, W. M. A. and Dieleman, W. I. J. (2012) 'Effects of elevated CO<sub>2</sub> and N fertilization on plant and soil carbon pools of managed grasslands: A meta-analysis', *Biogeosciences*, 9(6), pp. 2247–2258. doi: 10.5194/bg-9-2247-2012.
- Stevenson, F. J. (1982) 'Organic forms of soil nitrogen', *Nitrogen in agricultural soils*. Wiley Online Library, 22, pp. 67–122. doi: org/10.2134/agronmonogr22.c3.
- Tabatabai, M. A. and Bremner, J. M. (1970) 'Arylsulfatase Activity of Soils', *Soil Science Society of America Journal*, 34(2), pp. 225–229. doi: 10.2136/sssaj1970.03615995003400020016x.
- Tanikawa, T. *et al.* (2013) 'Sequential transformation rates of soil organic sulfur fractions in two-step mineralization process', *Biology and Fertility of Soils*, 50(2), pp. 225–237. doi: 10.1007/s00374-013-0849-8.
- Tilsner, J. *et al.* (2005) 'Amino acid contents and transport in oilseed rape (*Brassica napus* L.) under different nitrogen conditions', *Planta*, 221(3), pp. 328–338. doi: 10.1007/s00425-004-1446-8.
- Verma, N., Chaudhary, S. and Goyal, S. (2018) 'Long Term Effects of Inorganic Fertilizers and Organic Amendments on Ammonification and Nitrification Activity of Soils under Cotton-Wheat Cropping System', *International Journal of Current Microbiology and Applied Sciences*, 7(04), pp. 718–724. doi: 10.20546/ijcmas.2018.704.080.
- Vinolas, L. C., Vallejo, V. R. and Jones, D. L. (2001) 'Control of amino acid mineralization and microbial metabolism by temperature', *Soil Biology and Biochemistry*, 33(7–8), pp. 1137–1140. doi: 10.1016/S0038-0717(00)00243-1.

- Vitousek, P. M. and Howarth, R. W. (1991) 'Nitrogen limitation on land and in the sea: How can it occur?', *Biogeochemistry*, 13(2), pp. 87–115. doi: 10.1007/BF00002772.
- Voroney, R.P.; Brookes, P.C.; Beyaert, R. P. (2008) *Chapter 49: Soil Microbial Biomass C,N,P, and S, Soil Sampling and Methods of Analysis, Second Edition*. doi: 10.1201/9781420005271.ch18.
- Wilkinson, A. *et al.* (2014) 'Rapid microbial uptake and mineralization of amino acids and peptides along a grassland productivity gradient', *Soil Biology and Biochemistry*. Elsevier Ltd, 72, pp. 75–83. doi: 10.1016/j.soilbio.2014.01.026.
- Zinder, S. H. and Brock, T. D. (1978) 'Methane, carbon dioxide, and hydrogen sulfide production from the terminal methiol group of methionine by anaerobic lake sediments', *Applied and Environmental Microbiology*, 35(2), pp. 344–352. doi: 10.1128/aem.35.2.344-352.1978.
- Zoon, R. A. (1987) 'Safety with <sup>32</sup>P- and <sup>35</sup>S-labeled compounds', *Methods in enzymology*, 152.

## Chapter 8

---

**Short-term uptake and efflux of sulphur containing amino acids and inorganic sulphate by maize roots under sterile hydroponic conditions**

## Abstract

A variety of plants possess the ability to utilize organic nitrogen (N) compounds such as amino acids, however, this has not been thoroughly studied for sulphur (S)-containing amino acids. A critical question is whether plants can acquire dissolved organic S (DOS) without assistance from soil microorganisms, and if so, how does this compare with that of inorganic sulphate uptake? To address this, we measured the uptake of three S compounds ((cysteine, methionine and sulphate) by maize (*Zea mays* L.). Plants were labelled in hydroponic culture with either  $^{14}\text{C}$  or  $^{35}\text{S}$  at ecologically relevant concentrations (100  $\mu\text{M}$ ) over short time periods (24 h). Efflux of  $^{35}\text{S}$ -compounds was also estimated by monitoring the increase of  $^{35}\text{S}$  content in the root bathing solution after pre-feeding the plants with each compound. The uptake results showed that sulphate was the preferred S source by maize, with a two-fold greater S accumulation compared to that of Cys or Met. However, sulphate uptake was inhibited by the presence of the organic S sources. In addition, we estimated that at least 66% of Cys and 73% of Met was rapidly taken up intact by the roots, even when inorganic sulphate was available. This indicates that low molecular weight S-containing amino acids could theoretically constitute a significant proportion of the plant's S supply under certain conditions. The uptake of Cys and Met was, however, sensitive to the presence of each other, indicating their uptake occurred via a common membrane transport system. In addition, we present evidence to show that once taken up by the roots, S transitorily accumulates in the root cell vacuoles where it is more susceptible to root efflux through passive diffusion. However, a large proportion of the S taken up is rapidly transformed and translocated to the shoot preventing efflux. In conclusion, we present direct evidence for the uptake and assimilation of dissolved organic S compounds and that maize plants are very effective in retaining and cycling S at the whole-plant level.

**Keywords:** Cysteine; Methionine; Sulphate; Uptake; Root exudation; Radiotracer;  $^{14}\text{C}$ ;  $^{35}\text{S}$ ; Maize.

## 8.1 Introduction

It is generally thought that sulphur (S) is predominantly taken up by plant roots in an inorganic form (i.e. sulphate; Moss, 1978). In actively growing plants, the S is then transported in the xylem via a selective distribution/redistribution system to the expanding leaves (Adiputra and Anderson, 1992, 1993), where assimilation into organic S takes place in the light (Leustek, 2002; Takahashi, 2010; Takahashi *et al.*, 2011). It should be noted, however, that in the case of N, many plants are opportunistic, being capable of taking up a range of organic or inorganic-N forms depending on the prevailing conditions in the soil (Moreau *et al.*, 2019). Further, it is now clear that plants can take up significant quantities of organic nitrogen (N), especially simple forms such as amino acids and oligopeptides and when inorganic N levels in soil are low (Falkengren-Grerup, Månsson and Olsson, 2000; Hawkins, Johansen and George, 2000; Henry and Jefferies, 2003; Weigelt, Bol and Bardgett, 2005; Gallet-Budynek *et al.*, 2009; Ge *et al.*, 2009; Czaban *et al.*, 2016; Song *et al.*, 2016), yet only a few studies have focussed on S-containing amino acids.

It has been reported that over 90% of soil S is present in an organic form, and that a large proportion of this is present as carbon (C)-bonded S (e.g. cysteine, methionine; Cys and Met) (Scott, Bick and Anderson, 1981). Therefore, direct evidence is required to determine if S-containing amino acids are taken up intact by plant roots, as well as their quantitative contribution to plant S demand compared to inorganic sulphate. In previous studies, nutrient uptake in plants is often studied in simplified systems, such as excised roots in hydroponic culture (Ferrari and Renosto, 1972; Soldal and Nissen, 1978). This technique has been used extensively and has provided valuable information about specific uptake rates of nutrients at the molecular and cellular level. Some researchers, however, have argued that excised roots may artificially increase the loss of nutrients from roots and thereby inhibit net uptake (Lucash *et al.*, 2007), resulting in an unrealistic estimation of root uptake. Excising roots also alters the

source-sink relationships within the plant. This may feedback on root membrane transport systems and repress uptake if the above-ground sink is removed. It is therefore essential to study nutrient uptake in intact plants.

Cysteine (Cys) and methionine (Met) are two essential S-containing amino acids required for primary and secondary metabolism in plants (Droux, 2004). Cys is the first reduced S product resulting from the sulphate assimilation pathway (Saito, 2000), while both Cys and Met play a major role in the growth and development of plant cells (Ravanel *et al.*, 1998; Saito, 2000; Wirtz and Droux, 2005). Evidence has been presented that shows that Cys can be actively transported into cultured tobacco cells (Harrington and Smith, 1977), where it can be rapidly metabolized, the products being pyruvate, ammonium and S-sulfocysteine (Tishel and Mazelis, 1966). In addition, the transport of Met into excised plant roots has also been studied (Wright, 1962; Persson and Näsholm, 2001), with observations suggesting that both Cys and Met can be taken up by the same membrane transport system as other free amino acids.

It is known that most nutrients taken up by roots are transported to the shoot via the xylem. These can then be returned in reduced forms to the root via the phloem (Scott and Brewer, 1980; Duarte and Larsson, 1993; Atkins and Smith, 2007). This translocation of organic compounds from leaves, and release of root exudates such as sugars, amino acids and organic acids by roots, are particularly important when plants are growing in nutrient deficient soils or when plant species have a very low capacity for reducing nutrients in their roots (Carvalhais *et al.*, 2011). Therefore, root systems of plants can not only import water and nutrients from soil solution, but also release low and high molecular-weight-compounds into the environment (Smith, 1969; Dakora and Phillips, 2002). Amino acids are generally considered to be the second most abundant class in terms of the total amount exuded by plant root systems, after sugars (Smith, 1976; Jaeger *et al.*, 1999). Depending on the cause and mechanisms, amino acid release from roots may include active transport (Badri *et al.*, 2009;

Lesuffleur and Cliquet, 2010) or passive diffusion (Brophy and Heichel, 1989; Rroço, Kosegarten and Mengel, 2002; Vives-Peris *et al.*, 2020). Passive diffusion of amino acids is caused by the large concentration gradient between the cytoplasm of root cells (e.g. 1 – 10 mM) and the outside soil solution (0.1 – 10  $\mu$ M) (Phillips *et al.*, 2004), while active transport of amino acids is mediated by proteins located in the root plasma membrane and can release amino acids against the electrochemical potential gradient (Okumoto *et al.*, 2004).

In previous studies, dual labelled ( $^{13}\text{C}$ ,  $^{15}\text{N}$ ) compounds have been used to estimate amino acid uptake into plants (Näsholm, Huss-Danell and Högberg, 2000; Streeter, Bol and Bardgett, 2000; Wei, Chen and Yu, 2015). In these experiments, intact uptake is implied if the slope of the correlation of  $^{13}\text{C}$  to  $^{15}\text{N}$  excess in the plant tissue is the same as in the parent amino acid compounds fed to the plants. In this present study, dual isotope labelling:  $^{14}\text{C}$  (Biernath, Fischer and Kuzyakov, 2008; Rasmussen *et al.*, 2010) and  $^{35}\text{S}$  were used to estimate intact uptake of amino acids.  $^{14}\text{C}$  labelling has been used in studies of hydroponic (soil free, nutrient water) systems (Pratelli, Boyd and Pilot, 2016; Oburger and Jones, 2018). A  $^{14}\text{C}$  tracer was chosen in this study to eliminate the problem of  $^{13}\text{C}$  dilution by the high  $^{12}\text{C}$  content in plant tissues, as  $^{13}\text{C}$  isotope can be strongly diluted in plant tissues making it difficult to detect  $^{13}\text{C}$  in bulk plant tissues. A  $^{14}\text{C}$  tracer approach allows estimation of the incorporation of amino acid- $^{14}\text{C}$  into plant tissues, as well as the amino acid- $^{14}\text{C}$  loss in the form of  $^{14}\text{CO}_2$  produced during deamination and breakdown of the C skeleton in the TCA cycle (Näsholm and Persson, 2001), or in processes relating to photorespiration (Bauwe, Hagemann and Fernie, 2010). In this study, the intact uptake of amino acids can be implied if the slope of the correlation of  $^{14}\text{C}$  to  $^{35}\text{S}$  excess in the plant tissue is the same as in the intact amino acid (e.g. 3  $^{14}\text{C}$  : 1  $^{35}\text{S}$  in the case of added  $^{14}\text{C}$ ,  $^{35}\text{S}$ -Cys, 5  $^{14}\text{C}$  : 1  $^{35}\text{S}$  in the case of added  $^{14}\text{C}$ ,  $^{35}\text{S}$ -Met).






Although a large number of studies have focused on inorganic sulphate metabolism, data on the regulation of S-containing amino acids uptake, transport and efflux in plants are relatively scarce. Therefore, the primary objective of this study was to quantify the importance of intact uptake of Cys and Met relative to inorganic sulphate uptake by maize under sterile hydroponic conditions using isotope-based methods; the secondary objective was to assess S translocation and efflux from roots (i.e. root exudation). We chose maize as the model plant as it is known to possess a high demand for both N and S (Rasheed, Ali and Mahmood, 2004; Sutar, 2017), and is capable of taking up exogenously applied amino acids (Salmenkallio and Sopanen, 1989; Moran-Zuloaga *et al.*, 2015).

## **8.2 Materials and methods**

### **8.2.1 Plant material and nutrient solution**

Maize (*Zea mays* L.) seeds were sterilised in 2% sodium hypochlorite (1 min) and rinsed twice with sterile water (Cuero, Smith and Lacey, 1986; Sauer and Burroughs, 1986). The seeds were then soaked for 24 h in sterile deionized water and allowed to germinate on moist filter paper at room temperature (ca. 20 °C) under sterile conditions. After 48 h, each seedling was transferred into individual microcosms. Each microcosm consisted of 25 ml polypropylene containers filled with 20 ml of full-strength S-free Long Ashton nutrient solution (Hewitt, 1952; Smith, Johnston and Cornforth, 1983). The composition of the full-strength nutrient solution was as follows (g 10 L<sup>-1</sup>) MgCl<sub>2</sub>·6H<sub>2</sub>O, 3.05; KCl, 1.49; CaCl<sub>2</sub>·2H<sub>2</sub>O, 5.88; NaH<sub>2</sub>PO<sub>4</sub>·2H<sub>2</sub>O, 2.92; Na<sub>2</sub>HPO<sub>4</sub>·12H<sub>2</sub>O, 0.47; H<sub>3</sub>BO<sub>3</sub>, 0.86; MnCl<sub>2</sub>·H<sub>2</sub>O, 0.30; ZnCl<sub>2</sub>, 0.03; CuCl<sub>2</sub>·2H<sub>2</sub>O, 0.06; Na<sub>2</sub>MoO<sub>4</sub>·2H<sub>2</sub>O, 0.005; FeEDTA, 0.33; MES buffer, 0.19; NaNO<sub>3</sub>, 3.40; NH<sub>4</sub>Cl, 2.14. After the addition of an individual seedling, the microcosms were placed in a climate-controlled cabinet with 16-h photoperiod maintained at 25 ± 0.5 °C. All sulphate nutrient salts were replaced with chloride salts in this study, so plants growing in our nutrient

solution were therefore expected to be S deficient once seed reserves are exhausted. Seven days after transplanting, plants were transferred to 10%-strength S-free Long Ashton solution for a further 3 d. All experiments were conducted on 10-d-old plants in 10%-strength hydroponic solution, with three fully expanded leaves on the main shoot.



**Fig. 8.2** Growth stages of maize plants (*Zea mays* L.) from seedling to the three-leaf stage.

At the three-leaf stage, there are three elongated leaves and the tip of the fourth leaf appears at the centre of the leaf whorl (Fig. 8.2). This stage is a pivot point of maize growth from heterotrophic growth (i.e. growth relying on seed reserves) to autotrophic growth (i.e. growth relying on photosynthesis after the exhaustion of seed reserves) (Cooper and MacDonald, 1970; Hanway, 1966). All three compounds were chosen to reflect possible organic (Cys, Met) and inorganic ( $\text{Na}_2\text{SO}_4$ ) S compounds typically released and exposed to plant roots during the breakdown of soil organic matter.

To ensure and maintain sterile conditions, all nutrient stock solutions, deionized water and containers (polypropylene vials, syringe, pipette tips etc) used in the experiments were autoclaved. In addition, roots were rinsed with sterile 10%-strength Long Ashton nutrient solution (to remove any exoenzymes and exudates) prior to transfer to the  $^{35}\text{S}/^{14}\text{C}$ -labelled solution. Light was excluded from the nutrient solution by tightly wrapping the root compartment with aluminium foil. Experiments were carried out during daylight hours with plants exposed to the same light intensity and temperatures as described above.

### **8.2.2 Experiment 1: Plant uptake and partitioning of $^{14}\text{C}$ -Cys and Met under hydroponic conditions**

Briefly, maize plants were rinsed with sterile 10%-strength Long Ashton nutrient solution before being placed into individual 25 ml polypropylene containers filled with 20 ml of 10%-strength S-free Long Ashton nutrient solution. Experiment one included six treatments:

- (a)  $^{14}\text{C}$ -Cys
- (b)  $^{14}\text{C}$ -Cys + Met
- (c)  $^{14}\text{C}$ -Cys +  $\text{Na}_2\text{SO}_4$
- (d)  $^{14}\text{C}$ -Met
- (e)  $^{14}\text{C}$ - Met + Cys
- (f)  $^{14}\text{C}$ - Met +  $\text{Na}_2\text{SO}_4$

The concentration of each S compound in the final nutrient solution was 100  $\mu\text{M}$  (i.e. in treatment b, the concentration of Cys and Met are 100  $\mu\text{M}$  separately), which was in the range of previously reported amino acid concentrations in soil solution (Jones and Darrah, 1994; Johnson and Pregitzer, 2007). After injection of labelled material(s) into the nutrient solution, the plant-solution system was sealed and the  $^{14}\text{CO}_2$  efflux from plant tissues was trapped by placing a container of 1 M NaOH solution (10 ml) inside the container (Fig. 8.3). After 24 h,

$^{14}\text{C}$  activity in each compartment was determined separately. Plant roots and shoots were destructively harvested. Plant materials were first rinsed with 0.01 M  $\text{CaCl}_2$  for 30 s to remove any isotope adhered to the plant surface, then rinsed with unlabelled 10%-strength Long Ashton nutrient solution. Plant shoots and roots were then oven dried (80 °C, 24 h), weighed and ground to a powder separately.  $^{14}\text{C}$  activity in the plant tissues (shoots and roots) was determined with an OX-400 Biological Sample Oxidizer (RJ Harvey Instrument Corp., Hillsdale, NJ). The liberated  $^{14}\text{CO}_2$  was collected in Oxosol scintillation fluid (National Diagnostics, Hesse, UK).  $^{14}\text{C}$  was then quantified by liquid scintillation counting using a Wallac 1404 scintillation counter with automated quench correction (Wallac EG&G, Milton Keynes, UK). The amount of  $^{14}\text{C}$ -amino acid remaining in the nutrient solution alongside the amount of plant respiration was also determined.

### **8.2.3 Experiment 2: Plant uptake and distribution of $^{35}\text{S}$ -Cys, Met and sulphate under hydroponic conditions**

Maize germination, transplanting and nutrient provision were the same as in Experiment 1 except that  $^{14}\text{C}$  was replaced with a  $^{35}\text{S}$  tracer and no NaOH traps was required (Fig. 8.4). The concentration of each  $^{35}\text{S}$ -labelled compound in the final nutrient solution was 100  $\mu\text{M}$  and the chase period was 24 h at which time the incorporation of  $^{35}\text{S}$  into plant tissues was determined. Experiment 2 included nine treatments:

- (a)  $^{35}\text{S}$ -Cys
- (b)  $^{35}\text{S}$ -Cys + Met
- (c)  $^{35}\text{S}$ -Cys +  $\text{Na}_2\text{SO}_4$
- (d)  $^{35}\text{S}$ -Met
- (e)  $^{35}\text{S}$ -Met + Cys
- (f)  $^{35}\text{S}$ -Met +  $\text{Na}_2\text{SO}_4$


(g)  $^{35}\text{S}\text{-Na}_2^{35}\text{SO}_4$

(h)  $^{35}\text{S}\text{-Na}_2^{35}\text{SO}_4 + \text{Cys}$


(i)  $^{35}\text{S}\text{-Na}_2^{35}\text{SO}_4 + \text{Met}$

To determine the liberation of inorganic S in nutrient solution during the course of experiment 2 (e.g., from exudation or exoenzyme activity), the nutrient solution at the end of the experiment was divided equally into two parts. Half of the nutrient solution was used for  $^{35}\text{S}$  quantification (i.e. inorganic sulphate mineralized from Cys or Met, plus organic  $^{35}\text{S}$ ) directly by liquid scintillation counting. The remaining half was shaken (200 rev min<sup>-1</sup>; 5 min) with the same volume of 0.1 M BaCl<sub>2</sub> (10 ml) and centrifuged (4000 rev min<sup>-1</sup>; 5 min) to precipitate and remove any inorganic sulphate present in the Cys or Met treatments (i.e. as Ba<sup>35</sup>SO<sub>4</sub>). In both cases the amount of  $^{35}\text{S}$  in the resultant solutions was determined by liquid scintillation counting as described above.

The amount of  $^{35}\text{S}$  incorporated into the plant materials was also determined. Plant materials were first rinsed with 0.01 M CaCl<sub>2</sub> for 30 s to remove any isotope adhered to the plant surface, then rinsed with unlabelled 10%-strength Long Ashton nutrient solution. Plant material was then divided into roots and shoots, weighed and dried at 80 °C in the oven prior to further measurements. To determine the total amount of  $^{35}\text{S}$  incorporated into plant tissues, aliquots of 40 mg of powdered dried samples were placed in glass vials and 1 ml of Soluene-350 (PerkinElmer Life Sciences, Inc) added. The vials were then capped and incubated (40°C, 4 h) until the samples were fully digested and an almost colourless. This prevents the presence of pigments, chlorophyll in our case, causing inaccurate readings on the scintillation counter (Gibson, 1980; Smith and Lang, 1987; Thomson and Temple, 2020). The amount of  $^{35}\text{S}$  was then determined by liquid scintillation counting as described above.



**Fig. 8.3** Schematic representation of the maize microcosm into which  $^{14}\text{C}$ -labelled amino acid (Cys or Met) was injected. Each polypropylene tube was filled with 20 ml of 10%-strength Long Ashton nutrient solution. The final concentration of the two sulphur-containing amino acids in nutrient solution were brought to 100  $\mu\text{M}$  separately. The  $^{14}\text{C}$ -radioisotope was then injected into the 10%-strength Long Ashton nutrient solution in the polypropylene tubes. The  $^{14}\text{C}$ -amino acid movement and incorporation into different root and shoot sections was measured after 24 h after the injection.  $^{14}\text{CO}_2$  evolved from shoot compartment was trapped in 10 ml of 1 M NaOH.



**Fig. 8.4** Schematic representation of the maize microcosm into which  $^{35}\text{S}$ -labelled Cys, Met or inorganic sulphate was injected. The same experimental set up as experiment 1 was adopted here, except that NaOH traps were not placed inside Lock & Lock plastic containers. The final concentration of Cys, Met or inorganic sulphate in nutrient solution were brought to  $100\ \mu\text{M}$  separately. The  $^{35}\text{S}$ -radioisotope was then injected into the 10%-strength Long Ashton nutrient solution in the polypropylene tubes. The  $^{35}\text{S}$  depletion from nutrient solution as well as the incorporation into plant root and shoot sections were measured after 24 h after the injection.



### 8.2.4 Experiment 3: Efflux of $^{35}\text{S}$ -labelled Cys, Met and $\text{SO}_4^{2-}$ from maize roots

Nine uniform sterile maize plants were removed from 10%-strength Long Ashton nutrient solution and transferred into the open barrel of individual 25 ml polypropylene syringes with a two-way stopcock connected at the bottom (Fig. 8.5). Each syringe was filled with 20 ml of one isotopically labelled S compound (i.e.  $^{35}\text{S}$ - $\text{Na}_2\text{SO}_4$ ,  $^{35}\text{S}$ -Cys or  $^{35}\text{S}$ -Met; 100  $\mu\text{M}$ ) in 10%-strength Long Ashton S-free nutrient solution. This simple axenic system facilitated collection of root-derived  $^{35}\text{S}$  efflux and minimizes root damage and overestimation of efflux (Ayers and Thornton, 1968).

After being transferred to the new nutrient solution, maize plants were supplied with each  $^{35}\text{S}$  compound for 1 h in the external root bathing medium. After 1 h, the labelled nutrient solution was removed by opening the valve at the bottom of each syringe and the plants rinsed with sterile deionized water to remove any isotope adhering to the roots. The syringe was then refilled with 20 ml of non- $^{35}\text{S}$ -labelled nutrient solution. This root bathing solution was collected and replaced every 10 mins over an 80 min period. The amount of  $^{35}\text{S}$  label present in the collected solutions in either an organic or inorganic form was determined using the 0.1 M  $\text{BaCl}_2$  precipitation procedure described above. This enabled the efflux of both sulphate and organic S from maize roots to be determined.  $^{35}\text{S}$  efflux rate from roots ( $S_{\text{efflux}}$ ) was expressed as  $\mu\text{mol g}^{-1} \text{ DW root h}^{-1}$  as the increase of  $^{35}\text{S}$  in the bathing solution between the start ( $T_0$ ) and the end ( $T_t$ ) of the sampling period, where  $R$  is dry root biomass and  $T$  denotes sampling time.

$$S_{\text{efflux}} = (T_t - T_0) / (R \times T) \quad (\text{Eqn. 1})$$

Many studies have investigated the efflux of low MW organic solutes and ions (e.g.  $\text{K}^+$ ,  $\text{Cl}^-$ , sugars, etc.) after pre-loading the roots. This efflux process typically involves three distinct root compartments, namely, the apoplast, cytoplasm and vacuole (Thoiron *et al.*, 1981; Saftner, Daie and Wyse, 1983). In this study, we can discount the fast-exchanging (<1 min) apoplastic compartment as we washed the roots prior to monitoring efflux. Here we fitted a mathematical

model to the experimental efflux data (Rauser, 1987) in which the leakage of S compounds to the outer bathing solution was considered as the sum of two diffusional processes from the cytoplasm and vacuole to the root bathing medium:


$$y = a \times (1 - \exp^{(-c \times t)}) + b \times (1 - \exp^{(-d \times t)}) \quad (\text{Eqn. 2})$$

Where  $y$  is the accumulated  $^{35}\text{S}$  washed out of plant roots,  $a$  and  $b$  is the size of the S storage pool in the cytoplasm and vacuole respectively,  $c$  and  $d$  are the exponential coefficient describing the rate of  $^{35}\text{S}$  release from these pools into the external root bathing solution. The half time ( $t_{1/2}$ ) of each pool can then be calculated as:

$$t_{1/2} = \ln(2)/c \text{ or } t_{1/2} = \ln(2)/d \quad (\text{Eqn. 3})$$

The initial volume of different S compounds in the two compartments A and B can be calculated by:

$$A = a - c / d \times a; B = b + c / d \times a \quad (\text{Eqn. 4})$$




**Fig. 8.5** Schematic illustrating how  $^{35}\text{S}$  labelling were used to obtain an estimate of sulphur efflux by maize plants (*Zea mays* L.).

#### **8.2.5 Experiment 4: A split root system-cycling of amino acid S and other nutrients between shoots and roots in young maize plants**

In the fourth experiment, fifteen uniform maize plants were selected and the roots of each maize plant split approximately equally between two separate containers of nutrient solution. At the start of the experiment, one of the root compartments was exposed to a radioisotope solution (i.e. either  $^{35}\text{S}$ -Cys,  $^{14}\text{C}$ -Cys,  $^{35}\text{S}$ -Met,  $^{14}\text{C}$ -Met or  $^{35}\text{S}$ - $\text{Na}_2\text{SO}_4$ ; 100  $\mu\text{M}$ ). Roots in this compartment were termed ‘donor’ roots; while the other root compartment was immersed into unlabelled nutrient solution, and roots in this compartment were termed ‘receiver’ roots. The ‘donor root’ compartment was used for the labelling and ‘receiver root’ compartment was used for determining the internal cycling and subsequent release of radioisotope.

Each experiment unit was placed inside a 2 L translucent sealable plastic container (Lock & Lock; Really Useful Products Ltd, West Yorkshire, UK; Fig. 8.6). After 24 h, the shoots were removed and the roots harvested and rinsed with 0.01 M  $\text{CaCl}_2$  for 30 s to remove any surface isotope contamination. Nutrient solution from both compartments was also collected to determine the amount of isotope depletion by the donor root and exudation from the receiver root. Radioactivity in the plants and solutions were determined by liquid scintillation counting as described above.

For  $^{14}\text{C}$  treatments, 1 M NaOH trap (10 ml) were placed inside the 2 L plastic container beside maize plant to catch any  $^{14}\text{CO}_2$  evolved from the plant. For  $^{35}\text{S}$  treatments, no NaOH traps were placed.



**Fig. 8.6** Schematic representation of the experimental apparatus showing the maize plant growing in a split root system with only 'donor root' exposed to radioisotope. This system was allowed to develop over a 24 h-period.


### 8.2.6 Statistics and data analysis

All experiments had three replicates per treatment and used plants with similar shoot heights (ca. 12 cm) and root lengths (ca. 9 cm). When calculating root S influx, it was assumed that efflux of the S compounds was minimal during the exposure period. Similarly, for calculation of S efflux, it was assumed that S uptake was minimal during the exposure period. All data analysis was carried out in IBM SPSS Statistics v25 (IBM UK Ltd., Portsmouth, UK). One way ANOVA followed by Tukey's post-hoc test was used to identify treatment differences, with the significance level set at  $p < 0.05$ . Graphs and curve fitting were produced using SigmaPlot v13.0 (Systat software Inc., London). The results are presented as means  $\pm$  SEM ( $n = 3$ ) and only significant differences are discussed ( $p < 0.05$ ).

## 8.3 Results

### 8.3.1 Plant uptake and distribution of $^{14}\text{C}$ -labelled Cys and Met under sterile hydroponic conditions

In experiment 1, the plant uptake of  $^{14}\text{C}$ -labelled Cys and Met over time was measured under sterile hydroponic conditions. The results indicated that sterile maize roots rapidly took up both Cys and Met, after which the amino acid-C was incorporated into both new cell biomass and utilised for respiration (Fig. 8.7). A similar amino acid incorporation rate was recorded for both amino acids:  $203.4 \pm 35.7 \text{ nmol } ^{14}\text{C} (\text{g root DW})^{-1} \text{ h}^{-1}$  for Cys, and  $191.9 \pm 30.7 \text{ nmol } ^{35}\text{S} (\text{g root DW})^{-1} \text{ h}^{-1}$  for Met. However, the partitioning of  $^{14}\text{C}$  among the different plant compartments varied for the two amino acids. Overall, a higher proportion of Cys was partitioned into plant respiration while a higher proportion of Met was partitioned into plant biomass ( $p < 0.05$ ). Based on our calculation,  $5.8 \pm 0.7 \%$  and  $3.7 \pm 0.2 \%$  of the added Cys and Met were respired by the maize plants, respectively, whereas  $6.7 \pm 0.3\%$  and  $10.8 \pm 0.5\%$  were incorporated into plant biomass (shoot plus root tissues), respectively.



**Fig. 8.7** Distribution of  $^{14}\text{C}$ -labelled Met and Cys in maize plants after 24 h of a)  $^{14}\text{C}$ -Cys recovered in the shoots; b)  $^{14}\text{C}$ -Cys recovered in the roots; c)  $^{14}\text{C}$ -Cys respiration from the maize plant; d)  $^{14}\text{C}$ -Met recovered in the shoots; e)  $^{14}\text{C}$ -Met recovered in the roots; f)  $^{14}\text{C}$ -Met respiration from the maize plant. Bars and lines represent mean  $\pm$  SEM ( $n = 3$ ). Common lowercase letters denote a lack of significant difference between treatments (One way ANOVA: LSD test,  $p < 0.05$ ).

Overall, the total percentage of  $^{14}\text{C}$  recovered in plant shoots, roots, solution and  $\text{CO}_2$  evolution from root-solution system exceeded 80% for all treatments. The highest amount of  $^{14}\text{C}$  recovered in the  $^{14}\text{C}$ -Cys among all three compartments (shoots, roots, and respiration) was for  $\text{CO}_2$  evolution, constituting  $5.8 \pm 0.7\%$  of  $^{14}\text{C}$ -Cys input. Only a small fraction of the  $^{14}\text{C}$  derived from Cys was retained in plant roots after uptake ( $2.6 \pm 0.1\%$ ), while a larger proportion ( $4.1 \pm 0.2\%$ ) was transported to the shoots. Total plant utilization of  $^{14}\text{C}$  derived from Met ( $14.5 \pm 0.4\%$ ) was similar to that of Cys ( $12.6 \pm 0.8\%$ ), however, the highest  $^{14}\text{C}$  content for Met was found in the shoots ( $6.3 \pm 0.3\%$ ), followed by the roots ( $4.5 \pm 0.5\%$ ) and respiration ( $3.7 \pm 0.2\%$ ).

Plant uptake of Cys and Met decreased in the presence of each other. This implies that competition between Cys and Met occurred for entry into the root cells, probably due to a


common transport carrier. Cys supply led to a decrease in Met partitioning into respiration, shoot tissue, and root tissue by  $29.3 \pm 10.3\%$ ,  $49.3 \pm 13.1\%$ , and  $57.9 \pm 10.6\%$  respectively, while Met supply led to a decrease in Cys incorporation by  $48.3 \pm 12.1\%$ ,  $42.1 \pm 10.1\%$ , and  $55.3 \pm 6.7\%$ , respectively. In contrast, plant uptake of Cys and Met was not markedly affected by the presence of inorganic sulphate ( $p < 0.05$ ) (Fig. 8.7). This implies that Cys and Met may be a more favourable source of S even under situations of high access to inorganic sulphate. It should be noted that the lower concentrations of Cys and Met in soil solutions *in situ* may limit the actual contribution of these compounds to plant nutrition.

### **8.3.2 Uptake and distribution of $^{35}\text{S}$ -labelled Met, Cys and sulphate by maize plant roots under sterile hydroponic conditions**

In experiment two, the uptake of  $^{35}\text{S}$ -labelled Cys, Met and sulphate was assessed over 24 h. The uptake period was chosen to ensure that sterility was maintained as well as ensuring that all the S in the external medium was not depleted during the incubation period (Barber and Gunn, 1974; Gaume, Mächler and Frossard, 2001). The uptake of all three S forms were similar in that the labelled S taken up was not retained in root tissues, but was rapidly transported to the shoots. However, there were striking differences in the ability of the maize plants to utilize the different S compounds. Cys and Met uptake by the whole plant was  $347 \pm 15$  and  $390 \pm 54$  nmol  $^{35}\text{S}$  g<sup>-1</sup> root DW h<sup>-1</sup>, respectively over the 24-h period. However, there was a much greater accumulation of sulphate, being 2.2- and 1.9-fold higher than Cys and Met, respectively. The higher uptake could be explained by faster  $^{35}\text{S}$ -sulphate transport from root to shoot tissue, as similar values for root retention of all three S sources were obtained. Overall,  $10.3 \pm 1.4\%$ ,  $10.1 \pm 0.8\%$ , and  $11.5 \pm 1.1\%$  of the added  $^{35}\text{S}$  was recovered in the plant roots from Cys, Met, and sulphate respectively. However, in terms of transportation to shoot tissue, sulphate was more mobile than both amino acids as only  $11.6 \pm 0.9\%$  and  $13.3 \pm 1.9\%$  of Cys and Met was



detected in shoot tissues, whereas a significantly greater proportion ( $34.1 \pm 2.2\%$ ) of the added sulphate ( $p < 0.05$ ) was found in the shoots.



**Fig. 8.8** Partitioning of  $^{35}\text{S}$  label after the introduction of radio isotope into hydroponic maize plants for 24 h. a)  $^{35}\text{S-Cys}$  recovered in the shoots; b)  $^{35}\text{S-Met}$  recovered in the shoots; c)  $^{35}\text{S-SO}_4^{2-}$  recovered in the shoots; d)  $^{35}\text{S-Cys}$  recovered in the roots; e)  $^{35}\text{S-Met}$  recovered in the roots; f)  $^{35}\text{S-SO}_4^{2-}$  recovered in the roots. Values represent means  $\pm$  SEM ( $n = 3$ ).


The presence of other S sources on the root uptake of each S compound (i.e. competition) was also examined (Fig. 8.8). The root uptake of both amino acids was unaffected by the presence of sulphate ( $p > 0.05$ ), but transportation of  $^{35}\text{S-Cys}$  and  $^{35}\text{S-Met}$  to the shoot was markedly decreased in the presence of sulphate by 29.8 and 21.3 % ( $p < 0.05$ ), respectively. This resulted in a significant inhibition of Cys uptake when sulphate was present ( $p < 0.05$ ), although it proved non-significant for Met. In contrast, root  $^{35}\text{S-sulphate}$  uptake and transportation to shoots was markedly decreased by 32.6% and 47.3% in the presence of Cys, and by 22.6 % and 33.5 % in the presence of Met respectively, indicating that there was a downregulation of inorganic S uptake by organic S compounds. In addition, Cys and Met resulted in significantly decreased levels of both root uptake and shoot transportation of each

other ( $p < 0.05$ ). Cys supply effectively decreased root uptake and shoot transportation of Met by 35.2% and 29.7%, respectively ( $p < 0.05$ ), while Met supply effectively decreased root uptake and transportation of Cys by 42.1% and 30.1%, respectively ( $p < 0.05$ ).

### 8.3.3 Intact uptake of cysteine and methionine

To determine whether Cys and Met were taken up as intact molecules or as inorganic compounds after enzymatic or microbial degradation, results of the uptake techniques (i.e. via  $^{14}\text{C}$  and  $^{35}\text{S}$  labelling) were combined (Fig. 8.9). The co-location of  $^{14}\text{C}$  and  $^{35}\text{S}$  appears a reasonable measure of intact uptake by plants roots under sterile hydroponic conditions. Our results clearly show that a large proportion of the supplied tracer may have been taken up intact. This was illustrated by plots of total  $^{14}\text{C}$  vs total  $^{35}\text{S}$  in maize roots ( $R^2 = 0.68$  for Cys,  $R^2 = 0.83$  for Met).

A higher proportion of  $^{35}\text{S}$  (experiment two) than  $^{14}\text{C}$  (experiment one) derived from both amino acids in plant material was detected in the plant tissue. The discrepancy between expected and measured ratios of  $^{14}\text{C}$  to  $^{35}\text{S}$  may be explained by several possibilities. First, the difference of measurement of  $^{14}\text{C}$  and  $^{35}\text{S}$  from biological samples ( $^{14}\text{C}$  was measured by dry combustion, while  $^{35}\text{S}$  was measured by wet digestion) could have led to different recovery rates of the two radiotracers. Second, under the action of enzymes released by plant roots, part of added amino acids could have been degraded in nutrient solution to inorganic compounds ( $^{14}\text{CO}_2$ ,  $\text{NO}_3^-$ ,  $\text{NH}_4^+$  and  $^{35}\text{SO}_4^{2-}$ ; Fig. 8.1) prior to being taken up independently (Jones, Healey, *et al.*, 2005). This rapid enzymatic degradation of amino acids may contribute to a higher  $^{35}\text{S}$  recovery in plant materials due to the fast uptake rates of inorganic sulphate by maize roots (Astolfi *et al.*, 2004). Some  $^{14}\text{CO}_2$  may also have been lost in respiration during the washing and drying of the root and shoot material. In addition, rapid post-uptake metabolism of amino acids may also explain the anomalous relationships between  $^{14}\text{C}$  and  $^{35}\text{S}$ .




**Fig. 8.9** Relationship between the total accumulation of  $^{14}\text{C}$  and  $^{35}\text{S}$  radiotracer ( $\text{nmol g}^{-1} \text{ root DW h}^{-1}$ ) in maize roots and shoots after 24 h of exposure to the labelled solution containing  $100 \mu\text{M}$  cysteine or methionine ( $n = 9$ ).

### 8.3.4. Efflux of cysteine, methionine and inorganic sulphate from maize roots

Previous studies on the loss of organic compounds from intact roots have focused on the accumulation of amino acids over long experiment periods, yet according to our results from experiment one and two, plant roots have the capacity to actively take up amino acids rapidly, which may in turn result in a lower detection of root efflux. Experiment three examined S efflux from maize roots. Rapid sampling was used to minimize the negative effect of root re-capture of the compounds lost from root efflux. It also helped to increase the concentration gradient between the root cytosol and the root bathing solution. The results showed that of the  $^{35}\text{S}$  loaded into the plant, 37.1, 27.6, and 27.5% the Cys, Met, and sulphate-S was recovered in the root bathing medium, respectively. Efflux rates exceeded influx rates for all three compounds, which is in line with previous studies performed under axenic conditions (Lesuffleur and Cliquet, 2010). The results showed that Cys and Met efflux rates were in the same range, between  $1.0 \pm 0.1$  and  $4.2 \pm 0.3 \mu\text{mol g}^{-1} \text{ root DW h}^{-1}$  for Cys, and  $0.7 \pm 0.2$  and  $3.2 \pm 0.1 \mu\text{mol g}^{-1} \text{ root DW h}^{-1}$  for Met (Fig. 8.8), indicating rapid efflux of low molecular S compounds within the short monitoring period. The release of sulphate was similar to that of the amino acids, rates ranging from  $1.2 \pm 0.1$  to  $4.2 \pm 0.3 \mu\text{mol g}^{-1} \text{ root DW h}^{-1}$ . In this study, the efflux rate of Cys and Met from attached maize roots was about 3-13 fold greater than the


rate of uptake, implying a minor effect of amino acid uptake on the overall efflux pattern of amino acids over the short monitoring period. Application of  $\text{BaCl}_2$  allowed the separation of organic and inorganic S in the root exudates. The results revealed that efflux of all three S compounds were in the form they were taken up, suggesting efflux of low molecular weight compounds in short period occurs via passive leakage. This is in line with previous studies, which suggested that amino acid efflux rates are small or insignificant, this efflux is generally regarded as not carrier-mediated but occurs by passive leakage (Jones and Darrah, 1993; Paynel, Murray and Bernard Cliquet, 2001), and could be recaptured by roots.



**Fig. 8.10** Transmembrane sulphate movement

**Table 8.1** Parameters of amino acids (Cys and Met) and sulphate release from intact maize roots. Efflux data was fitted to Eqn. 2. The parameters *a* and *c* represent the  $^{35}\text{S}$  held in the cytoplasm and vacuole respectively, while *b* and *d* are the efflux rate constants for these two pools, respectively. Values represent means  $\pm$  SEM ( $n = 3$ ).

Substrates	Pool <i>a</i>	Pool <i>c</i>	$t_{1/2}$ (fast pool) hour	$t_{1/2}$ (slow pool) day	A ( $\mu\text{mol} \cdot (\text{g root DW})^{-1}$ )	B ( $\mu\text{mol} \cdot (\text{g root DW})^{-1}$ )	R <sup>2</sup>
Cys	0.56 $\pm$ 0.07	1.09 $\pm$ 0.01	2.98 $\pm$ 0.62	1.78 $\pm$ 0.26	1.03 $\pm$ 0.18	0.66 $\pm$ 0.02	0.99
Met	0.55 $\pm$ 0.04	0.93 $\pm$ 0.45	3.14 $\pm$ 0.41	6.59 $\pm$ 2.32	1.10 $\pm$ 0.35	0.58 $\pm$ 0.03	0.99
Na <sub>2</sub> SO <sub>4</sub>	0.58 $\pm$ 0.04	1.27 $\pm$ 0.02	3.72 $\pm$ 0.33	1.39 $\pm$ 0.24	1.14 $\pm$ 0.05	0.71 $\pm$ 0.06	0.99



**Fig. 8.11** Cumulative efflux of added  $^{35}\text{S}$ -Cys, Met or sulphate per unit dry mass of maize roots. Prior to measuring efflux, plants were pre-treated with radioisotopes for 60 minutes. Efflux was determined by measuring the increase of radioisotope in the root bathing medium solutions.  $\text{BaCl}_2$  was applied to separate organic and inorganic S in solution. Data represent means  $\pm$  SEM ( $n = 3$ ).

The release of  $^{35}\text{S}$  into the bathing solution indicated two distinct compartments (Fig. 8.10), which may be interpreted as two pools: the cytoplasmic and vacuole compartments (Cooper and Clarkson, 1989; Paynel, Murray and Bernard Cliquet, 2001). The rate of  $^{35}\text{S}$  release from the roots decreased sharply over the course of the efflux period. A double first order exponential decay equation fitted well to the efflux data ( $R^2 > 0.99$ ; Fig. 8.11)). This predicted that the half-life for the slower exchanging compartment (vacuole) were 1.7, 6.6, and 1.4 d for Cys, Met, and sulphate, respectively (Table 8.1), while half-lives for the faster exchanging compartment (cytoplasm) were 2.9, 3.1 and 3.7 h for Cys, Met and sulphate, respectively. Based on calculation from Eqn. 4, the cytoplasmic S pool was estimated to range from 0.58 to 0.71  $\mu\text{mol g}^{-1}$  root DW, which was smaller than concentration in the vacuole which ranged from 1.03 to 1.14  $\mu\text{mol g}^{-1}$  root DW.

### **8.3.5 Cycling of sulphur compounds between maize shoot and root via split root systems**

In experiment four, a split root system under sterile conditions was used to investigate the cycling of S compounds between shoot and root in young maize plants. Here, both the translocation of S compounds from ‘donor root to shoot’ and ‘shoot to receiver roots’ were monitored. Isotope labels ( $^{14}\text{C}$  and  $^{35}\text{S}$ ) were used to establish the quantitative significance of S cycling in S deficient plants. Overall, the results indicated that S compounds entered the plant via donor roots and were then cycled through the shoots and to the receiver-roots.

Calculation showed that  $> 50\%$  of the  $^{14}\text{C}$  tracer loaded from the donor root was cycled through the whole plant. The fraction of isotope tracer in each compartment (donor root, shoot, receiver root) is shown in Table 8.2. By the end of this incubation experiment, less than 10% of the  $^{14}\text{C}$ -Cys from the donor root was transported and retained in the shoot within 24 h, while a much higher proportion (nearly 50%) was respired from the shoot,  $^{14}\text{C}$ -Cys partitioning in the receiver root reached a similar level as the donor root (around 20%). Similarly, a high

proportion of  $^{14}\text{C}$  derived from Met was cycled from donor root to the whole plant, although less  $^{14}\text{C}$  was respired and a higher proportion was retained in the shoot relative to Cys.

The distribution pattern for  $^{35}\text{S}$  differed from that of  $^{14}\text{C}$  in that a higher proportion of S was retained in the donor root tissue. Nearly half of the  $^{35}\text{S}$  taken up from the nutrient solution was retained in the donor root, ca. 40% was transported to the shoot, from where less than half was subsequently translocated to the receiver root. One possible explanation for the difference in  $^{35}\text{S}$  and  $^{14}\text{C}$  responses is that after being taken up by the root tissues, the amino acids are metabolized (deaminated, transaminated etc.) prior to transport to the shoots (Warren, 2012a). Overall, the actual amount of sulphate cycled from nutrient solution to the donor root was around three times higher than that of the two amino acids. By the end of the 24-h cycling period, however, negligible amounts of radioactivity were observed in the nutrient solution in the receiver root compartment. Therefore, it is likely that under S deficient conditions, maize plants are highly effective at maximising S use efficiency within the plant and minimise root exudation (or recapture any S lost to the external medium).

**Table 8.2** Translocation and utilization of three sulphur compounds by maize plants over a 24-h period. Both root compartments received the same amount of nutrient solution. Three independent measurement from replicate plants were made for each treatment. Values represent means  $\pm$  SEM ( $n = 3$ ).

	$^{14}\text{C}$ -Cys	$^{35}\text{S}$ -Cys	$^{14}\text{C}$ -Met	$^{35}\text{S}$ -Met	$^{35}\text{S}$ - $\text{Na}_2\text{SO}_4$
$^{14}\text{CO}_2$ partitioning (% $^{14}\text{C}$ taken up)	47.2 $\pm$ 9.3		32.4 $\pm$ 2.9		
Shoot tissue	6.0 $\pm$ 1.9	20.5 $\pm$ 5.2	11.4 $\pm$ 1.4	29.0 $\pm$ 2.5	21.1 $\pm$ 2.5
Donor root	22.9 $\pm$ 3.1	63.7 $\pm$ 2.7	32.0 $\pm$ 1.6	49.2 $\pm$ 3.2	62.9 $\pm$ 1.9
Receiver root	23.9 $\pm$ 5.0	15.8 $\pm$ 3.2	24.1 $\pm$ 1.8	21.8 $\pm$ 2.9	16.0 $\pm$ 3.7

## 8.4 Discussion

A complication about quantifying intact amino acid uptake is that theoretically the same correlation in isotope enrichment could be observed if the dual labelled amino acids were mineralized to inorganic compounds external to the root and then taken up (e.g. by carbon-sulphur lyases). In this short plant uptake experiment, considering the efforts to minimize bacteria growth in nutrient solution before the conduction of experiments, it is likely that the pre-mineralization of Cys and Met was negligible.

Previous studies have addressed the importance of carbon-sulphur-lyases in the degradation of amino acids to inorganic compounds, among which methionine gamma-lyase degrades Met to  $\alpha$ -keto acids, ammonia and thiols (Rébeillé *et al.*, 2006; Goyer *et al.*, 2007; Huang, Joshi and Jander, 2014), while D-cysteine desulphydrase degrades Cys to pyruvate, sulphide and ammonia (Schmidt, 1982; Ramírez and Whitaker, 1998; Riemenschneider *et al.*, 2005). Therefore, it is possible that enzymatic transformation of Cys and Met took place during our experiment, and the breakdown products of these metabolites were taken up separately by maize roots, however, to our knowledge these enzymes do not exist extracellularly.

Data from experiment two revealed that plants utilized inorganic sulphate preferentially over Cys and Met, and that organic S supply has a negative effect on sulphate uptake. This agrees with the widely accepted view that initial root uptake of inorganic sulphate is energy dependent through a proton/sulphate coupled co-transport in the plasma membrane of root cells and is well adjusted to the S status of the plant. When other sulphur sources (Cys or GSH) are provided to the plant, uptake is repressed in a negative feedback loop (Herschbach and Rennenberg, 1994; Bolchi *et al.*, 1999; Hawkesford *et al.*, 2003; Davidian and Kopriva, 2010; Noctor *et al.*, 2011), while during S starvation, uptake is enhanced by activating the expression of high affinity sulphate transporters (Maruyama-Nakashita *et al.*, 2004). However, the fact that maize plants could take up DOS does imply that it may have ecological significance under



some circumstances. The results clearly showed that a considerable proportion of the supplied amino acids was absorbed intact, this was illustrated by plots of excess  $^{14}\text{C}$  against excess  $^{35}\text{S}$  in roots (Fig. 8.3), with 66% and 73% of Cys and Met taken up intact separately. Observation that both Cys and Met inhibit uptake of each another supported our hypothesis that these amino acids enter root cells via the same transport system.

It is well documented that influx of amino acids involves proton-coupled amino acid transporters (Bush, 1993; Delrot, Atanassova and Maurousset, 2000), whereas passive efflux is driven primarily by large differences in concentrations between the inside (10 mM) and the outside (0.1 – 10  $\mu\text{M}$ ) of roots cells (Jones and Darrah, 1993; Moore *et al.*, 2003). However, the mechanisms behind S efflux from root tissue are still unclear. In this study, rapid efflux of intact S compounds was detected from maize roots in experiment three. Considering S influx across the plasma membrane is energy dependent process, it is therefore surprising that S appears to leak out again rapidly. We therefore assume that after uptake, each compound transitorily accumulates in the cellular compartments (cytoplasm, vacuole) that are sensitive to efflux. Once they enter a complex reductive metabolic pathway, they are less likely to leak out again. Even so, the total amino acid efflux in the present study may have been under-estimated since amino acids could be re-absorbed by plant roots and these re-absorbed amino acids would not have been detected in efflux. In addition, high exogenous amino acid concentrations applied may have stimulated influx and diminished efflux which is assumed to be concentration dependent.

## **8.5 Conclusions and outlook**

We have presented direct experimental evidence to support our hypothesis that plants (e.g., *Zea mays* L.) can directly take up S-containing amino acids intact under hydroponic conditions. These results indicate that dissolved organic matter in the form of free amino acids

(e.g. Cys and Met) may represent a readily available source of N, S and C for plants. We also provided evidence for the rapid redistribution of S within the plant following root uptake. S efflux from roots indicated two distinct S compartments, with the vacuole being the slower releasing compartment and the cytoplasm being the smaller storage compartment.

However, many gaps in our understanding remain. First, the correlation between  $^{14}\text{C}$  and  $^{35}\text{S}$  in plant tissues could arise if the amino acid were mineralized in the nutrient solution and the products taken up independently. However, this would require enzymes such as L-cysteine desulfhydrase which are unlikely to exist as exoenzymes in sterile hydroponic culture. Therefore, stronger evidence using compound-specific ( $^{13}\text{C}/^{34}\text{S}$ ) isotope ratio mass spectrometry (IRMS) could be used to examine this further.

Second, despite evidence that plants take up free amino acids from hydroponic solution, it is unclear to what extent Cys and Met are taken up intact by plants *in situ*, where plant roots and rhizosphere microorganisms compete intensely for organic N (A G Owen and Jones, 2001). It is also not clear whether plant roots are capable of taking up other dissolved organic S forms (e.g., peptides and proteins). These may also play an important role in the N and S dynamics where inorganic N and S are inadequate for plant growth.

It has previously been suggested that most amino acids are minor components of the xylem sap (e.g. glycine) and must be metabolized within the roots (e.g. to glutamine) before transport to the shoots, where post uptake metabolism takes place (Warren, 2012b). To our knowledge, we know little about the metabolic fate of Cys and Met following uptake (i.e. which are the dominant pathways involved in their transformation and in which root cells this takes place).

Also, to avoid disturbance in sampling solution due to microbial decomposition, the majority of studies on root efflux have been carried out under sterile conditions. However, as it is difficult to maintain sterile root systems over long periods, most efflux experiments only

last for a short time (up to several weeks). Sodium hydroxide trapping method normally involves placing plants in a closed system that is not at a steady state. This major drawback also makes it difficult to maintain steady atmosphere in the system for a longer experimental period. Therefore, there is a need to develop new methods for maintaining the root system under sterile conditions, so that exudates from various root types could be collected for longer experiment period.

## 8.6 Acknowledgements

This work was supported by Bangor University and the China Scholarship Council. I would also like to thank Jonathan Roberts and Sarah Chesworth for technical support.

## 8.7 References

- Adiputra, I. G. K. and Anderson, J. W. (1992) 'Distribution and redistribution of sulphur taken up from nutrient solution during vegetative growth in barley', *Physiologia Plantarum*, 85(3), pp. 453–460. doi: 10.1111/j.1399-3054.1992.tb05811.x.
- Adiputra, I. G. K. and Anderson, J. W. (1993) 'Effect of sulphur nutrition and leaf excision on distribution and redistribution of sulphur in barley', in *Plant Nutrition—from Genetic Engineering to Field Practice*. Springer, pp. 239–242.
- Astolfi, S. *et al.* (2004) 'Influence of iron nutrition on sulphur uptake and metabolism in maize (*Zea mays* L.) roots', *Soil Science and Plant Nutrition*, 50(7), pp. 1079–1083. doi: 10.1080/00380768.2004.10408577.
- Atkins, C. A. and Smith, P. M. C. (2007) 'Translocation in legumes: Assimilates, nutrients, and signaling molecules', *Plant Physiology*, 144(2), pp. 550–561. doi: 10.1104/pp.107.098046.

- Ayers, W. A. and Thornton, R. H. (1968) 'Exudation of amino acids by intact and damaged roots of wheat and peas', *Plant and Soil*, 28(2), pp. 193–207. doi: 10.1007/BF01880238.
- Badri, D. V. *et al.* (2009) 'An ABC transporter mutation alters root exudation of phytochemicals that provoke an overhaul of natural soil microbiota', *Plant Physiology*, 151(4), pp. 2006–2017. doi: 10.1104/pp.109.147462.
- Barber, D. A. and Gunn, K. B. (1974) 'The effect of mechanical forces on the exudation of organic substances by the roots of cereal plants grown under sterile conditions', *New Phytologist*. JSTOR, pp. 39–45.
- Bauwe, H., Hagemann, M. and Fernie, A. R. (2010) 'Photorespiration: players, partners and origin', *Trends in Plant Science*, 15(6), pp. 330–336. doi: 10.1016/j.tplants.2010.03.006.
- Biernath, C., Fischer, H. and Kuzyakov, Y. (2008) 'Root uptake of N-containing and N-free low molecular weight organic substances by maize: A <sup>14</sup>C/<sup>15</sup>N tracer study', *Soil Biology and Biochemistry*, 40(9), pp. 2237–2245. doi: 10.1016/j.soilbio.2008.04.019.
- Bolchi, A. *et al.* (1999) 'Coordinate modulation of maize sulfate permease and ATP sulfurylase mRNAs in response to variations in sulfur nutritional status: Stereospecific down-regulation by L-cysteine', *Plant Molecular Biology*, 39(3), pp. 527–537. doi: 10.1023/A:1006148815106.
- Brophy, L. S. and Heichel, G. H. (1989) 'Nitrogen release from roots of alfalfa and soybean grown in sand culture', *Plant and Soil*, 116(1), pp. 77–84. doi: 10.1007/BF02327259.
- Bush, D. R. (1993) 'Proton-coupled sugar and amino acid transporters in plants', *Annual Review of Plant Physiology and Plant Molecular Biology*, 44(1), pp. 513–542. doi:

10.1146/annurev.pp.44.060193.002501.

- Carvalhais, L. C. *et al.* (2011) 'Root exudation of sugars, amino acids, and organic acids by maize as affected by nitrogen, phosphorus, potassium, and iron deficiency', *Journal of Plant Nutrition and Soil Science*, 174(1), pp. 3–11. doi: 10.1002/jpln.201000085.
- Cooper, C. S. and MacDonald, P. W. (1970) 'Energetics of Early Seedling Growth in Corn (*Zea mays* L.) 1', *Crop Science*. Wiley Online Library, 10(2), pp. 136–139.
- Cooper, H. D. and Clarkson, D. T. (1989) 'Cycling of amino-nitrogen and other nutrients between shoots and roots in cereals—a possible mechanism integrating shoot and root in the regulation of nutrient uptake', *Journal of Experimental Botany*. Oxford University Press, 40(7), pp. 753–762.
- Cuero, R. G., Smith, J. E. and Lacey, J. (1986) 'The influence of gamma irradiation and sodium hypochlorite sterilization on maize seed microflora and germination', *Food Microbiology*, 3(2), pp. 107–113. doi: 10.1016/S0740-0020(86)80034-X.
- Czaban, W. *et al.* (2016) 'Direct acquisition of organic N by white clover even in the presence of inorganic N', *Plant and Soil*. Plant and Soil, 407(1–2), pp. 91–107. doi: 10.1007/s11104-016-2896-z.
- Dakora, F. D. and Phillips, D. A. (2002) 'Root exudates as mediators of mineral acquisition in low-nutrient environments', in *Food Security in Nutrient-Stressed Environments: Exploiting Plants' Genetic Capabilities*. Springer, pp. 201–213.
- Davidian, J. C. and Kopriva, S. (2010) 'Regulation of sulfate uptake and assimilation - The same or not the same?', *Molecular Plant*. The Authors. All rights reserved., 3(2), pp. 314–325. doi: 10.1093/mp/ssq001.
- Delrot, S., Atanassova, R. and Maurousset, L. (2000) 'Regulation of sugar, amino acid and

- peptide plant membrane transporters', *Biochimica et Biophysica Acta - Biomembranes*, 1465(1–2), pp. 281–306. doi: 10.1016/S0005-2736(00)00145-0.
- Droux, M. (2004) 'Sulfur assimilation and the role of sulfur in plant metabolism: A survey', *Photosynthesis Research*, 79(3), pp. 331–348. doi: 10.1023/B:PRES.0000017196.95499.11.
- Duarte, P. J. P. and Larsson, C.-M. (1993) 'Translocation of nutrients in N-limited, non-nodulated pea plants', *Journal of Plant Physiology*. Elsevier, 141(2), pp. 182–187.
- Falkengren-Grerup, U., Månsson, K. F. and Olsson, M. O. (2000) 'Uptake capacity of amino acids by ten grasses and forbs in relation to soil acidity and nitrogen availability', *Environmental and Experimental Botany*, 44(3), pp. 207–219. doi: 10.1016/S0098-8472(00)00068-X.
- Ferrari, G. and Renosto, F. (1972) 'Regulation of Sulfate Uptake by Excised Barley Roots in the Presence of Selenate', *Plant Physiology*, 49(2), pp. 114–116. doi: 10.1104/pp.49.2.114.
- Gallet-Budynek, A. *et al.* (2009) 'Intact amino acid uptake by northern hardwood and conifer trees', *Oecologia*, 160(1), pp. 129–138. doi: 10.1007/s00442-009-1284-2.
- Gaume, A., Mächler, F. and Frossard, E. (2001) 'Aluminum resistance in two cultivars of *Zea mays* L.: Root exudation of organic acids and influence of phosphorus nutrition', *Plant and Soil*, 234(1), pp. 73–81. doi: 10.1023/A:1010535132296.
- Ge, T. *et al.* (2009) 'Amino acids as a nitrogen source for tomato seedlings: The use of dual-labeled (<sup>13</sup>C, <sup>15</sup>N) glycine to test for direct uptake by tomato seedlings', *Environmental and Experimental Botany*, 66(3), pp. 357–361. doi: 10.1016/j.envexpbot.2009.05.004.

- Gibson, J. A. B. (1980) 'Modern techniques for measuring the quenching correction in a liquid scintillation counter: a critical review', in *Physical Aspects*. ACADEMIC PRESS, INC., pp. 153–172. doi: 10.1016/b978-0-12-549901-9.50021-1.
- Goyer, A. *et al.* (2007) 'Functional characterization of a methionine  $\gamma$ -lyase in arabidopsis and its implication in an alternative to the reverse trans-sulfuration pathway', *Plant and Cell Physiology*, 48(2), pp. 232–242. doi: 10.1093/pcp/pcl055.
- Hanway, J. (1966) 'How a corn plant develops.; Spec. Publ. 48.', *Iowa Agricultural and Home Economics Experiment Station Publications*, 48(Sept), pp. 1–18.
- Harrington, H. M. and Smith, I. K. (1977) 'Cysteine Transport into Cultured Tobacco Cells', *Plant Physiology*, 60(6), pp. 807–811. doi: 10.1104/pp.60.6.807.
- Hawkesford, M. J. *et al.* (2003) 'Sulphate Uptake and Transport', in *Sulphur in Plants*. Springer, pp. 71–86. doi: 10.1007/978-94-017-0289-8\_4.
- Hawkins, H. J., Johansen, A. and George, E. (2000) 'Uptake and transport of organic and inorganic nitrogen by arbuscular mycorrhizal fungi', *Plant and Soil*, 226(2), pp. 275–285. doi: 10.1023/A:1026500810385.
- Henry, H. A. L. and Jefferies, R. L. (2003) 'Plant amino acid uptake, soluble N turnover and microbial N capture in soils of a grazed Arctic salt marsh', *Journal of Ecology*, 91(4), pp. 627–636. doi: 10.1046/j.1365-2745.2003.00791.x.
- Herschbach, C. and Rennenberg, H. (1994) 'Influence of glutathione (GSH) on net uptake of sulfate and sulfate transport in tobacco plants', *Journal of Experimental Botany*, 45(277), pp. 1069–1076. doi: 10.1093/jxb/45.8.1069.
- Hewitt, E. J. (1952) *Sand and water culture methods used in the study of plant nutrition.*, *Sand and water culture methods used in the study of plant nutrition.*

- Huang, T., Joshi, V. and Jander, G. (2014) 'The catabolic enzyme methionine gamma-lyase limits methionine accumulation in potato tubers', *Plant Biotechnology Journal*, 12(7), pp. 883–893. doi: 10.1111/pbi.12191.
- Jaeger, C. H. *et al.* (1999) 'Mapping of sugar and amino acid availability in soil around roots with bacterial sensors of sucrose and tryptophan', *Applied and Environmental Microbiology*, 65(6), pp. 2685–2690. doi: 10.1128/aem.65.6.2685-2690.1999.
- Johnson, R. M. and Pregitzer, K. S. (2007) 'Concentration of sugars, phenolic acids, and amino acids in forest soils exposed to elevated atmospheric CO<sub>2</sub> and O<sub>3</sub>', *Soil Biology and Biochemistry*, 39(12), pp. 3159–3166. doi: 10.1016/j.soilbio.2007.07.010.
- Jones, D. L. *et al.* (2005) 'Dissolved organic nitrogen uptake by plants - An important N uptake pathway?', *Soil Biology and Biochemistry*, 37(3), pp. 413–423. doi: 10.1016/j.soilbio.2004.08.008.
- Jones, D. L. and Darrah, P. R. (1993) 'Influx and efflux of amino acids from Zea mays L. roots and their implications for N nutrition and the rhizosphere Author (s): D. L. JONES and P. R. DARRAH Source: Plant and Soil, Vol. 155 / 156, Selected papers from the PROCEEDINGS OF THE TWEL', *Plant and Soil*, 155/156(1992), pp. 87–90.
- Jones, D. L. and Darrah, P. R. (1993) 'Influx and efflux of amino acids from Zea mays L. roots and their implications for N nutrition and the rhizosphere Author (s): D. L. JONES and P. R. DARRAH Source: Plant and Soil, Vol. 155 / 156, Selected papers from the PROCEEDINGS OF THE TWEL', *Plant and Soil*, 155/156(1992), pp. 87–90.
- Jones, D. L. and Darrah, P. R. (1994) 'Amino-acid influx at the soil-root interface of Zea mays L. and its implications in the rhizosphere', *Plant and Soil*, 163(1), pp. 1–12. doi: 10.1007/BF00033935.



- Lesuffleur, F. and Cliquet, J. B. (2010) 'Characterisation of root amino acid exudation in white clover (*Trifolium repens* L.)', *Plant and Soil*, 333(1), pp. 191–201. doi: 10.1007/s11104-010-0334-1.
- Leustek, T. (2002) 'Sulfate metabolism', *The Arabidopsis Book/American Society of Plant Biologists*. American Society of Plant Biologists, 1.
- Lucash, M. S. *et al.* (2007) 'Estimating nutrient uptake by mature tree roots under field conditions: Challenges and opportunities', *Trees - Structure and Function*, pp. 593–603. doi: 10.1007/s00468-007-0160-0.
- Maruyama-Nakashita, A. *et al.* (2004) 'Regulation of high-affinity sulphate transporters in plants: Towards systematic analysis of sulphur signalling and regulation', *Journal of Experimental Botany*, 55(404), pp. 1843–1849. doi: 10.1093/jxb/erh175.
- Moore, J. C. *et al.* (2003) 'TOP-DOWN IS BOTTOM-UP : DOES PREDATION IN THE RHIZOSPHERE Special Feature', *Ecology*, 84(4), pp. 846–857.
- Moran-Zuloaga, D. *et al.* (2015) 'Organic nitrogen uptake by plants: reevaluation by position-specific labeling of amino acids: Reevaluation of organic N uptake by plants by position-specific labeling', *Biogeochemistry*. Springer International Publishing, 125(3), pp. 359–374. doi: 10.1007/s10533-015-0130-3.
- Moss, M. R. (1978) 'Sources of sulfur in the environment; the global sulfur cycle', *Sulfur in the environment*. John Wiley and Sons New York, pp. 23–50.
- Näsholm, T., Huss-Danell, K. and Högberg, P. (2000) 'Uptake of organic nitrogen in the field by four agriculturally important plant species', *Ecology*, 81(4), pp. 1155–1161. doi: 10.1890/0012-9658(2000)081[1155:UOONIT]2.0.CO;2.
- Näsholm, T. and Persson, J. (2001) 'Plant acquisition of organic nitrogen in boreal forests',

- Physiologia Plantarum*, 111(4), pp. 419–426. doi: 10.1034/j.1399-3054.2001.1110401.x.
- Noctor, G. *et al.* (2011) ‘Graham Noctor’, pp. 1–32. doi: 10.1199/tab.0142.
- Oburger, E. and Jones, D. L. (2018) ‘Sampling root exudates – Mission impossible?’, *Rhizosphere*. Elsevier, 6(June), pp. 116–133. doi: 10.1016/j.rhisph.2018.06.004.
- Okumoto, S. *et al.* (2004) ‘Root phloem-specific expression of the plasma membrane amino acid proton co-transporter AAP3’, *Journal of Experimental Botany*, 55(406), pp. 2155–2168. doi: 10.1093/jxb/erh233.
- Owen, A. G. and Jones, D. L. (2001) ‘Competition for amino acids between wheat roots and rhizosphere microorganisms and the role of amino acids in plant N acquisition’, *Soil Biology and Biochemistry*, 33(4–5), pp. 651–657. doi: 10.1016/S0038-0717(00)00209-1.
- Paynel, F., Murray, P. J. and Bernard Cliquet, J. (2001) ‘Root exudates: A pathway for short-term N transfer from clover and ryegrass’, *Plant and Soil*, 229(2), pp. 235–243. doi: 10.1023/A:1004877214831.
- Persson, J. and Näsholm, T. (2001) ‘Amino acid uptake: A widespread ability among boreal forest plants’, *Ecology Letters*, 4(5), pp. 434–438. doi: 10.1046/j.1461-0248.2001.00260.x.
- Phillips, D. A. *et al.* (2004) ‘Microbial products trigger amino acid exudation from plant roots.’, *Plant physiology*, 136(1), pp. 2887–2894. doi: 10.1104/pp.104.044222.
- Pratelli, R., Boyd, S. and Pilot, G. (2016) ‘Analysis of amino acid uptake and translocation in arabidopsis with a low-cost hydroponic system’, *Journal of Plant Nutrition and Soil Science*, 179(2), pp. 286–293. doi: 10.1002/jpln.201500464.

- Ramírez, E. C. and Whitaker, J. R. (1998) 'Cystine lyases in plants: A comprehensive review', *Journal of Food Biochemistry*, 22(5), pp. 427–440. doi: 10.1111/j.1745-4514.1998.tb00254.x.
- Rasheed, M., Ali, H. and Mahmood, T. (2004) 'Impact of Nitrogen and Sulfur Application on Growth and Yield of Maize (*Zea Mays* L.) Crop', *J. res. Sci*, 15(2), pp. 153–157.
- Rasmussen, J. *et al.* (2010) 'Plant uptake of dual-labeled organic N biased by inorganic C uptake: Results of a triple labeling study', *Soil Biology and Biochemistry*, 42(3), pp. 524–527. doi: 10.1016/j.soilbio.2009.11.032.
- Rausser, W. E. (1987) 'Compartmental Efflux Analysis and Removal of Extracellular Cadmium from Roots', *Plant Physiology*, 85(1), pp. 62–65. doi: 10.1104/pp.85.1.62.
- Ravanel, S. *et al.* (1998) 'The specific features of methionine biosynthesis and metabolism in plants', *Proceedings of the National Academy of Sciences of the United States of America*, 95(13), pp. 7805–7812. doi: 10.1073/pnas.95.13.7805.
- Rébeillé, F. *et al.* (2006) 'Methionine catabolism in Arabidopsis cells is initiated by a  $\gamma$ -cleavage process and leads to S-methylcysteine and isoleucine syntheses', *Proceedings of the National Academy of Sciences of the United States of America*, 103(42), pp. 15687–15692. doi: 10.1073/pnas.0606195103.
- Riemenschneider, A. *et al.* (2005) 'Isolation and characterization of a D-cysteine desulfhydrase protein from Arabidopsis thaliana', *FEBS Journal*, 272(5), pp. 1291–1304. doi: 10.1111/j.1742-4658.2005.04567.x.
- Rroço, E., Kosegarten, H. and Mengel, K. (2002) 'Importance of plasmalemma H<sup>+</sup>-ATPase activity for N losses from intact roots of spring wheat (*Triticum aestivum* L.)', *European Journal of Agronomy*, 16(3), pp. 187–196. doi: 10.1016/S1161-

0301(01)00123-X.

- Saftner, R. A., Daie, J. and Wyse, R. E. (1983) 'Sucrose Uptake and Compartmentation in Sugar Beet Taproot Tissue', *Plant Physiology*, 72(1), pp. 1–6. doi: 10.1104/pp.72.1.1.
- Saito, K. (2000) 'Regulation of sulfate transport and synthesis of sulfur-containing amino acids', *Current Opinion in Plant Biology*, 3(3), pp. 188–195. doi: 10.1016/s1369-5266(00)80064-3.
- Salmenkallio, M. and Sopanen, T. (1989) 'Amino Acid and Peptide Uptake in the Scutella of Germinating Grains of Barley, Wheat, Rice, and Maize', *Plant Physiology*, 89(4), pp. 1285–1291. doi: 10.1104/pp.89.4.1285.
- Sauer, D. B. and Burroughs, R. (1986) 'Disinfection of seed surfaces with sodium hypochlorite.', *Phytopathology*. Saint Paul, 76(7), pp. 745–749.
- Schmidt, A. (1982) 'A cysteine desulfhydrase from spinach leaves specific for D-cysteine', *Zeitschrift für Pflanzenphysiologie*. Elsevier, 107(4), pp. 301–312.
- Scott, H. D. and Brewer, D. W. (1980) 'Translocation of nutrients in soybeans', *Soil Science Society of America Journal*. Wiley Online Library, 44(3), pp. 566–569.
- Scott, N. M., Bick, W. and Anderson, H. A. (1981) 'The measurement of sulphur-containing amino acids in some scottish soils', *Journal of the Science of Food and Agriculture*, 32(1), pp. 21–24. doi: 10.1002/jsfa.2740320105.
- Smith, G. S., Johnston, C. M. and Cornforth, I. S. (1983) 'Comparison of nutrient solutions for growth of plants in sand culture', *New phytologist*. Wiley Online Library, 94(4), pp. 537–548.
- Smith, I. K. and Lang, A. L. (1987) 'Decoloration and solubilization of plant tissue prior to determination of  $^3\text{H}$ ,  $^{14}\text{C}$ , and  $^{35}\text{S}$  by liquid scintillation', *Analytical Biochemistry*,

164(2), pp. 531–536. doi: 10.1016/0003-2697(87)90529-X.

Smith, W. H. (1969) 'Release of organic materials from the roots of tree seedlings', *Forest Science*. Oxford University Press, 15(2), pp. 138–143.

Smith, W. H. (1976) 'Character and significance of forest tree root exudates', *Ecology*. Wiley Online Library, 57(2), pp. 324–331.

Soldal, T. and Nissen, P. E. R. (1978) 'Multiphasic uptake of amino acids by barley roots', *Physiologia Plantarum*. Wiley Online Library, 43(3), pp. 181–188.

Song, S. *et al.* (2016) 'Uptake Kinetics of Different Nitrogen Forms by Chinese Kale', *Communications in Soil Science and Plant Analysis*. Taylor & Francis, 47(11), pp. 1372–1378. doi: 10.1080/00103624.2016.1178279.

Streeter, T. C., Bol, R. and Bardgett, R. D. (2000) 'Amino acids as a nitrogen source in temperate upland grasslands: The use of dual labelled ((<sup>13</sup>C, (<sup>15</sup>N) glycine to test for direct uptake by dominant grasses', *Rapid Communications in Mass Spectrometry*, 14(15), pp. 1351–1355. doi: 10.1002/1097-0231(20000815)14:15<1351::AID-RCM23>3.0.CO;2-9.

Sutar, R. K. (2017) 'Sulphur Nutrition in Maize - A Critical Review', *International Journal of Pure & Applied Bioscience*, 5(6), pp. 1582–1596. doi: 10.18782/2320-7051.6092.

Takahashi, H. (2010) 'Regulation of sulfate transport and assimilation in plants', in *International review of cell and molecular biology*. Elsevier, pp. 129–159.

Takahashi, H. *et al.* (2011) 'Sulfur Assimilation in Photosynthetic Organisms: Molecular Functions and Regulations of Transporters and Assimilatory Enzymes', *Annual Review of Plant Biology*, 62(1), pp. 157–184. doi: 10.1146/annurev-arplant-042110-103921.

Thoirion, A. *et al.* (1981) 'The compartmental analysis of sulphate transport in cells of',

- Biochimica et Biophysica Acta (BBA) - Biomembranes*, 644(1), pp. 24–35. Available at: [https://doi.org/10.1016/0005-2736\(81\)90054-7](https://doi.org/10.1016/0005-2736(81)90054-7).
- Thomson, J. and Temple, S. (2020) ‘Sample preparation techniques for liquid scintillation analysis’, in *Handbook of Radioactivity Analysis*. Elsevier Inc., pp. 803–860. doi: 10.1016/b978-0-12-814397-1.00007-8.
- Tishel, M. and Mazelis, M. (1966) ‘Enzymatic degradation of L-cystine by cytoplasmic particles from cabbage leaves [26]’, *Nature*, 211(5050), pp. 745–746. doi: 10.1038/211745a0.
- Vives-Peris, V. *et al.* (2020) ‘Root exudates: from plant to rhizosphere and beyond’, *Plant Cell Reports*. Springer Berlin Heidelberg, 39(1), pp. 3–17. doi: 10.1007/s00299-019-02447-5.
- Warren, C. R. (2012a) ‘Post-uptake metabolism affects quantification of amino acid uptake’, *New Phytologist*, 193(2), pp. 522–531. doi: 10.1111/j.1469-8137.2011.03933.x.
- Warren, C. R. (2012b) ‘Post-uptake metabolism affects quantification of amino acid uptake’, *New Phytologist*, 193(2), pp. 522–531. doi: 10.1111/j.1469-8137.2011.03933.x.
- Wei, L., Chen, C. and Yu, S. (2015) ‘Uptake of organic nitrogen and preference for inorganic nitrogen by two Australian native Araucariaceae species’, *Plant Ecology and Diversity*. Taylor & Francis, 8(2), pp. 259–264. doi: 10.1080/17550874.2013.871656.
- Weigelt, A., Bol, R. and Bardgett, R. D. (2005) ‘Preferential uptake of soil nitrogen forms by grassland plant species’, *Oecologia*, 142(4), pp. 627–635. doi: 10.1007/s00442-004-1765-2.
- Wirtz, M. and Droux, M. (2005) ‘Synthesis of the sulfur amino acids: Cysteine and methionine’, *Photosynthesis Research*, 86(3), pp. 345–362. doi: 10.1007/s11120-005-

8810-9.

Wright, D. E. (1962) 'Amino acid uptake by plant roots', *Archives of Biochemistry and Biophysics*, 97(1), pp. 174–180. doi: 10.1016/0003-9861(62)90061-9.

## Chapter 9

---

### Summary, discussion and recommendations for future research



## 9.1 Introduction

In this section, the main findings of the experimental work (presented in Chapters 3-8) are summarised and discussed in relation to the overall initial objectives of the thesis and published literature. The broader implications of the findings are also addressed. Detailed discussions of the results from individual experiments are provided alongside an outline of the main strengths and potential limitations of the research. This is followed by an assessment of the challenges and uncertainties identified in the thesis that need to be addressed by future research.

## 9.2 Synthesis of findings

The overarching aim of this thesis was to address three fundamental issues relative to the behaviour of S-containing amino acids in the plant-soil system:

- (i) To gain further insight into the size of the organic and inorganic S pools from a range of grassland sites across an altitudinal gradient, as well as the mineralization rates of S-containing amino acids (Cys and Met);
- (ii) Determine the relative contribution of amino acid-S to plant S uptake compared to inorganic  $\text{SO}_4^{2-}$  under sterile hydroponic conditions;
- (iii) Quantify the short-term competition for two S-containing amino acids between the rhizosphere microbial community and maize plant roots.

The results of each experimental chapter, in relation to the thesis objectives, are presented below:

The literature review highlighted that soil organic S is present as a heterogeneous mixture of S compounds, many of which likely remain uncharacterised. This pool can be separated into S immobilised within the soil microbial biomass and mesofaunal community, that present in soil organic matter and that present in living plant biomass (Michael A. Kertesz

and Mirleau, 2004a). It is also clear that most organic S is present in soil in an insoluble form and that a significant proportion may be chemically and/or physically protected, and therefore not susceptible to mineralization (Eriksen, Lefroy and Blair, 1995a). In contrast, there appears to be a small pool of organic S that can be readily mineralized (here termed labile organic S, LOS; Vermeiren et al., 2018). Examples of compounds in the LOS pool include S-containing proteins, peptides and amino acids. To better understand soil S dynamics, it is therefore essential to explore the size and turnover rate of this LOS pool. Carbon-bonded S (including S-containing amino acids, organic sulphonates) is a key component of this labile S pool and the breakdown product of many higher molecular weight organic-S moieties. To gain a better understanding of the S forms and their transformation in soils, radiolabelled and stable isotope tracers ( $^{14}\text{C}$ ,  $^{35}\text{S}$ ,  $^{32}\text{S}$ ) were applied to track the turnover of Cys and Met in soils.

In chapter 3, P-containing extractants were used to extract total dissolved S (DOS plus  $\text{SO}_4^{2-}$ , TDS) from a range of different grassland soil types. Evidence showed that the concentration of DOS in all soils ranged from 11 to 309  $\text{mg kg}^{-1}$  DW soil, constituting 24 to 94% of TDS in all selected grassland soils. Statistical analysis also showed that the DOS concentration correlated well with Al, B, Fe, K, Mn and  $\text{NH}_4^+$  concentrations in soil extracts and MBC. A better understanding of the factors governing the seasonal and spatial variation in DOS in contrasting soils is therefore a prerequisite to make more accurate S budgets in soil.

Chapter 4 investigated the short-term, concentration-dependent microbial uptake and mineralization of  $^{14}\text{C}$ -labelled Cys and Met in five grassland soils collected from an altitude-driven primary productivity gradient. The results revealed that  $^{14}\text{C}$ -Cys and  $^{14}\text{C}$ -Met were directly and rapidly assimilated by soil microbes, with half times ranging from 0.34 – 2.14 min. This is an order of magnitude (or more) faster than mineralisation rates determined from measurement of  $^{14}\text{CO}_2$  evolution and indicated that the DOS pool may be turning over many thousands of times in a year. This considerable delay between microbial  $^{14}\text{C}$ -amino acid

removal from the soil solution (plant uptake) and subsequent  $^{14}\text{CO}_2$  evolution (mineralisation) indicates that the degradation of  $^{14}\text{C}$ -labelled Cys and Met in soil solution was mainly through microbial biological process rather than abiotic removal processes. This rapid turnover mirrors findings for other low molecular weight solutes in soil (e.g. sugars, organic acids; (Paul W. Hill, Farrar and Jones, 2008; Gunina *et al.*, 2017)) suggesting that the low molecular weight DOC, DOS, DOP and DON pools are all turning over very fast in concert with each other.

On the other hand, Cys and Met also constitute important sources of directly available S for both plant nutrition and microbial growth. Traditionally, gross S mineralization rates can be estimated by measuring sulphatase activity, or by periodically leaching the soils to prevent re-immobilization of newly formed sulphate. However, the legitimacy of both methods has been questioned by researchers; sulphatase enzymes only account for part of the enzymes responsible for S mineralization, while leaching sulphate changes soil conditions leading to overestimation of gross mineralization rates. In addition, enzymatic approaches for measuring transformation rates in soil have also been heavily criticised due to a lack of standardisation in methodological approaches (e.g. substrates, concentrations, pH, detection method etc; (Burns *et al.*, 2013)). In addition, difficulties have arisen due to problems in directly measuring S transformation processes and being able to design realistic laboratory studies which reflect the real world. Alternatively, the application of the S isotope dilution technique allows the determination of both gross and net S transformation rates. In Chapter 6, by quantifying 0.01 M  $\text{Ca}(\text{H}_2\text{PO}_4)_2$  extractable  $^{35}\text{SO}_4^{2-}$  and stable  $\text{SO}_4^{2-}$  concentrations every 7 d in a closed incubation experiment (for 70 d), the size of the labile organic S pool, and soil S turnover rates in grassland soils with contrasting soil properties were quantified via an established isotope-based method.

Chapter 7 investigated concurrent microbial mineralization and utilization of C, N and S derived from Cys and Met in a grassland soil over a 7-d incubation period, using  $^{14}\text{C}$  and  $^{35}\text{S}$

dual-labelling. Results indicated that microbial communities retain the capacity to utilize Cys-C and Met-C in the presence of inorganic N ( $\text{NH}_4^+$  and  $\text{NO}_3^-$ ) and S ( $\text{SO}_4^{2-}$ ) and that inorganic products derived from the amino acid are excreted into the soil, benefiting plants. In accordance with results from chapter 4, I confirmed that the decomposition of S-containing amino acids (Cys and Met) in my grassland soil mainly occurs as a result of microbial oxidation.

Results from Chapter 8 showed that a considerable proportion of Cys (66%) and methionine (73%) was also rapidly taken up intact by plant roots under sterile hydroponic conditions, even when sulphate was available, indicating that organic S may constitute a readily available source of S to plants, particularly in S-limiting ecosystems. This uptake pathway be important in providing an alternative source of S to plants and in recapturing amino acids previously lost in root exudates or when they are directly adjacent to decomposing organic matter. I also provided evidence of the rapid redistribution of S within the plant after been taken up by the root system.

Chapter 5 investigated the competition for S-containing amino acids between plant roots and microorganisms in the rhizosphere ( $^{14}\text{C}$ ,  $^{35}\text{S}$ ). Results showed that the capture of free Cys and Met by maize plants was very low compared to soil microbes, with < 10% of the added amino acid captured by the plant, compared to the rhizosphere microbial community (around 80%). I conclude that both Cys and Met are sources of available C, N and S for maize plants, both pre- and post-mineralization. Therefore, studies relying solely on soil organic matter mineralization rates may have underestimated its N and S supply potential to plants.

In this thesis, I addressed a range of fundamental questions relating to the potential significance of amino acids to the overall plant S budget, with emphasis on S supply in different agroecological contexts. The results clearly showed that i) when studied in isolation, both soil microorganisms and maize roots have the capacity to take up free amino acid-S from soil solution, and ii) when allowed to compete, soil microorganisms outcompete maize roots for

amino acid-S in soil. Taken together, these findings have important implications for ecosystems S cycling and the more accurate design of ecosystem S cycling models.

### 9.3 Methodological strengths and limitations

Soil organic matter contains an important source of plant-available S and good methods are needed to estimate its potential contribution to total plant S supply. My work brings together a wide range of experimental techniques, leading to meaningful conclusions about the potential significance of Cys and Met to graminaceous plant nutrition.

In the experimental chapters where the aim was to capture the intrinsic variability in S cycling in different soils and keep conditions closer to natural conditions, soil samples were collected, transported immediately back to laboratory, and kept in the dark at  $\leq 5^{\circ}\text{C}$  before conducting the experiments. All experiments conducted in this thesis commenced within  $\leq 48$  h of soil collection, to minimise artefacts which may arise from sample preparation and storage (Ross, 1992; Jones and Willett, 2006; Meyer, Welp and Amelung, 2019). However, it should also be considered whether the fast turnover measured in the laboratory was a result of C starvation in the microbial community following the removal of plant roots and rhizodeposition. On the other hand, the sometimes large concentrations of substrate added to the soil may not reflect steady state concentrations in the soil solution which are often at the nanomolar level (Jones, Shannon, *et al.*, 2005; Boddy *et al.*, 2007; R C I Broughton *et al.*, 2015). Therefore, there is a strong need for actual *in situ* field experiments to better estimate soil S turnover (Oburger and Jones, 2009).

A few previous studies indicate that DOS constitutes a significant part of the S flux from terrestrial to aquatic environments (Wang *et al.*, 2012b) where it can be turned over rapidly (Brailsford *et al.*, 2020). In chapter 3, spatial variation of two S fractions: DOS and inorganic sulphate were assessed based on S fractionation results from seven grassland soil

samples. Instrumental methods of S determination, such as ion chromatography (IC) and inductively coupled plasma atomic emission spectrometry (ICP-AES), gave rapid and precise estimation of  $\text{SO}_4^{2-}$ -S and organic S concentrations. However, despite previous studies showing that extractable organic S was directly related to mineralizable organic S (Watkinson, Perrott and Thorrold, 1991), it is advisable for future studies to develop a sound understanding of the labile organic S pool, which can become directly available to plants through mineralisation. Furthermore, future work should also include evaluations of the seasonal variation of DOS content in soil solution due to dilution or microbial activity caused by changes in moisture regime (i.e. rainfall, drought), seasonal and daily temperature patterns, management regime (e.g. tillage), crop growth stage, the presence of livestock etc.

High molecular weight C inputs (e.g. protein, hemicellulose, lignin, and cellulose) into soil are first broken down extracellularly to smaller units by the soil microbial community (e.g. fungi and bacteria). These low molecular weight breakdown products (such as peptides, amino acids etc) are then taken up into the cell, entering catabolic or anabolic pathways. Traditionally, the measurements of substrate turnover in soil typically has relied on the addition of isotopically labelled substrates (e.g.  $^{13}\text{C}$ ,  $^{14}\text{C}$ ) to soil in the laboratory and measurement of  $^{14}\text{CO}_2$  evolution. An important advantage of using the  $^{14}\text{C}$  tracer technique compared with traditional methods is that it is possible to calculate the distribution of C between different measured C pools accurately with high sensitivity, low cost, and ease of sample preparation in comparison to other non-isotopic approaches (e.g., LC-MS, NMR). However, while isotopic based measurements of  $^{14}\text{CO}_2$  evolution greatly facilitate our understanding of the transformations involved in amino acid turnover in soil, it may underestimate their rates of cycling in the soil due to the delay between microbial uptake and mineralization (Paul W. Hill, Farrar and Jones, 2008). There are also issues about isotopic pool dilution and incomplete extraction of C pools that need to be considered (H. C. Glanville *et al.*, 2016). In Chapter 4,

therefore, I adopted the centrifugal drainage procedure to study substrate depletion from soil solution. This method allows small quantities of soil samples to be used so that soil solution can be recovered extremely rapidly; this method also allowed us to explore the temporal decoupling of Cys and Met- $^{14}\text{C}$  uptake and subsequent mineralization.

Of all four S containing amino acids (Met, Cys, homocysteine, and taurine), only Cys and Met are biosynthetically incorporated into proteins and are therefore expected to represent the main forms of DOS entering soil. In chapter 7, with dual labelling ( $^{14}\text{C}$ ,  $^{35}\text{S}$ ), I linked S-amino acid C and S dynamics to the size and activity of the microbial biomass in grassland soils. However, previous studies have indicated S mineralization seems to be driven by the form of S in the starting materials (Churka Blum *et al.*, 2013b), where the mineralization process of the ester S pool is mostly governed by the enzymatic hydrolysis, while mineralization of C-bonded S is governed by the need of C to provide energy to the microorganisms. Therefore, my study emphasizes the importance of a more detailed analysis of uptake and mineralization of ester S in soil solution.

Due to the continual degradation and synthesis of S fractions by soil organisms, it is not easy to examine the dynamics of S mineralization and immobilization within soil as these occur concurrently. The use of  $^{35}\text{S}$  (Chapter 6) allowed the tracing of separate S pools and the determination of both gross and net transformation rates. The model I adopted in my study provided more realistic estimates of transformation rates as it takes the re-mineralisation of previously immobilized  $^{35}\text{S}$  into account. However, as my study was carried out in a closed laboratory incubation system with no plants and rhizosphere microbial activity, future research and models should also take into account leaching, plant uptake, fertilization etc., performed under field conditions where possible at various points during the growing season. It would also be useful to investigate the diversity and function of sulfatases in soil and their rates of reaction.

Although it has been suggested that plants have the physiological capacity to take up a wide range of DOM in the form of amino acids from the environment prior to microbial degradation to inorganic nutrients, this has not been thoroughly studied for S-containing amino acids. In chapter 8, radiolabelled compounds were used to measure plant uptake.  $^{14}\text{C}$  and  $^{35}\text{S}$  were combined in parallel treatments to directly trace plant uptake of the carbon or S skeleton of added amino acids, but the results did not indicate whether the amino acid was taken up intact or cleaved prior to uptake of the molecule (Hill and Jones, 2019). It would be possible to better study plant uptake at a molecular level, if  $^{15}\text{N}/^{34}\text{S}/^{13}\text{C}$  labelled S-containing amino acids were also utilised coupled with compound-specific isotope ratio mass spectrometry (Knowles *et al.*, 2010; Charteris, 2019). Future work should therefore build from the results in this thesis to increase our understanding of the molecular mechanisms behind DOS processing in soils and their subsequent incorporation in different plant species.

The availability of amino acids to plant roots may be constrained by differences between amino acids in their potential for mineralization, microbial assimilation, and sorption to soil solids, as well as the relative soil reserves of inorganic nutrients. Most studies to date have focused on the direct uptake of organic N solutes such as amino acids. However, the degree to which S-containing amino acids contribute to the overall S budget of plants, remains poorly understood. This is especially the case in the rhizosphere where microbial activity can be an order of magnitude greater than in the bulk soil, and competition for labile organic S between plant roots and soil microorganisms could be intense. The main aim of Chapter 5 was therefore to quantify the competition between rhizosphere microorganisms and plant roots for two S-containing amino acids namely, Cys and Met. In agreement with previous studies, only small amounts of S-containing amino acids were captured by maize plant roots. My study also indicates the clear need for *in situ* studies of plant acquisition of Cys and Met where amino acid concentrations are low, in order to learn more about the realistic contribution of low



molecular weight S containing compounds to plant N and S nutrition. In addition, the capacity of roots to take up alternative forms of organic S (e.g. oligopeptides, glutathione, S-lipids) remains to be evaluated.

Sulfurous air pollutants may act as both toxins and nutrients to plants (De Kok and Tausz, 2001). Atmospheric S gases can be taken up directly by the foliage, forming bisulfite ( $\text{HSO}_3^-$ ) and sulfite ( $\text{SO}_3^{2-}$ ) due to its high solubility in the apoplastic water of the mesophyll (Omasa *et al.*, 2012). Sulfite may directly enter the S reduction pathway and be reduced to sulphide, incorporated into Cys and subsequently into other S containing compounds (Romero *et al.*, 2014). Therefore, some plants may even benefit from elevated levels of atmospheric S gases, since they contribute to plant S nutrition, especially when inorganic sulphate is low in soil. In chapter 5, the total recovery rate of  $^{35}\text{S}$  in the soil-plant system was  $< 80\%$  for all five treatments. It is possible that volatile S compounds (VOS) were produced during the incubation, and this was not measured in my experiments. Future studies are needed to assess the magnitude of evolution of VOS and inorganic S compounds from both soil and plants. It is likely that VOS will be just as important in soils as it is in marine environments (Carpenter and Milyo, 2012; Tang, 2020).

#### **9.4 Future research**

The studies presented in this thesis have provided pivotal information about the significance of amino acid-S to plant roots and soil microbes. However, several research gaps have also been identified during the research. It is clear from the results presented in this study that a much deeper analysis of many areas covered in this subject are still required. Some of these are detailed below:

#### **9.4.1 Determination the dynamics of inorganic and organic S compounds in soil solution**

Determining S availability in soil based on extraction methods is very challenging since many forms of available S are found in soils, including readily soluble inorganic sulphate, adsorbed sulphate and a portion of organic S (carbon bonded S or ester S). Most of the methods of soil S testing involving extraction of soil with a weak salt solution have been criticised for poor precision and lack of accuracy due to the effects of serious chemical interferences. In addition, these extraction methods fail to estimate the pool of labile organic S which becomes available to plants through mineralization. Some loss of organic S is also expected during the extraction procedure due to microbial mineralisation (Rousk and Jones, 2010). More recently, isotope dilution methods have provided a much deeper fundamental understanding of soil N cycle processes, modelling of gross N fluxes and understanding of individual microbial metabolic pathways (Sprent, Knowles and Blackburn, 1993). In my research, I also adopted this method to quantify the size of total labile S pool and gross mineralization/immobilization rates. Further experiments and modelling, which include a wider range of soils and organic S substrates are required to improve our understanding and prediction of the dynamics of inorganic and organic soil S. It would also be useful to see if the rates of S cycling are coupled to those of other nutrients such as P and K.

#### **9.4.2 What specific members of microbial community play a dominant role in soil S cycling?**

Soil microbial communities are strongly involved in the biological and biochemical mineralization of soil S. S oxidizing bacteria, such as *Thiobacillus*, *Thiothrix*, *Chlorobiaceae*, *Beggiatoa*, *Chromatiaceae*, and *Ectothiorhodospiraceae*, are able to oxidize organic S to inorganic sulphate. In addition, the biochemical S mineralization is driven mainly by arylsulfatases, which are partly produced by plant roots but mostly produced by soil microbes.

It seems likely that particular microbial species or genera in the rhizosphere may play a greater role in S cycling than others, but to date, assays for determining the overall rate of entire metabolic processes such as respiration, or specific enzyme activities do not allow any identification of the specific microbial species directly involved in each measured process. The development of modern molecular techniques which do not rely on cultivation (e.g. gene knockouts, mutants metagenomics, transcriptomics) now allow us to better explore the composition and function of soil microbial communities. Despite this, little is known about the specific microbial species or genera that play important roles in the soil organo-S cycle. For a better understanding of the microbial role in S cycling, therefore, future studies incorporating traditional approaches (e.g. phospholipid fatty acid analysis) with modern molecular techniques (e.g. proteomics, transcriptomics, metabolomics, stable isotope probing; Nannipieri *et al.*, 2003) are needed. This may also be coupled to stable isotope imaging techniques (e.g. NanoSIMS) for imaging microbial S dynamics (Higuchi *et al.*, 2020; Stuart *et al.*, 2020). A better understanding of the mechanisms and processes regarding the effects of soil microbes on the availability of nutrients is particularly important for the design of new biofertilizers. The right combination of microbes in these products may enhance the solubility and plant availability of a wide range of different macro- and micro-nutrients (N, P, K and S etc.) making this an economically and environmentally friendly approach to improving plant nutrition (Miransari, 2013).

#### **9.4.3 Predicting S availability from C/S ratio of plant residues**

Where S inputs from fertilizer and atmospheric deposition are low, the release of S from organic forms becomes important for the supply of S to plants. Therefore, estimation of the potential contribution that the organic S pool makes to plant available S, especially following the addition of organic S materials such as composts, manures, biosolids, fresh plant residues

etc., is critical. S mineralization seems to be driven by the form of S added (Churka Blum *et al.*, 2013b), with the S mineralization process mostly governed by biological mineralization of C-bonded S-rich residues, with liberation of inorganic sulphate as a secondary product. In contrast, the application of highly oxidized S compounds is mainly governed by enzymatic hydrolysis of the ester S pool. Therefore, future studies providing valuable information for modelling of S transformations from different crop residues in more soil types are recommended.

#### **9.4.4 Is organic S a significant S source for plants *in situ*?**

Although my study did not take into account plants from widely differing habitats, results showing uptake of S-containing amino acids in maize provides an indication of the potential role of Cys and Met for plant S nutrition under certain conditions. However, it is not clear to what extent amino acid-S contributes to the total S budget of the plant when grown under field conditions, where concentrations of free amino acids and inorganic  $\text{SO}_4^{2-}$  may differ. The concentration of Cys and Met in soil solution *in situ* is typically in the 1-50  $\mu\text{M}$  range, while the concentration of  $\text{SO}_4^{2-}$  in soils is typically one or more orders of magnitude greater. My research points to the need for longer term studies on the acquisition of S-containing amino acids by various plant species in the field to explore the actual importance of these low molecular weight organic S for plant S nutrition. In addition, to date, most studies on amino acid absorption by plant roots have mainly focused on gross influx rates, excluding the possibility of simultaneous root leakage, which could affect the calculation of net uptake rates. Clearly, in the future, more experimentation is needed to expand our mechanistic understanding of this root uptake processes, with the application of new approaches and techniques (e.g. cloning and *ex situ* expression of transporters to better understand their function, signal

transduction cascades which are triggered under S deficiency, internal S recycling in plants and microbes).

#### **9.4.5 Seasonal and spatial responses in microbial biomass sulphur**

Some of the more recent studies have shown a growing interest in soil microbial biomass S, as it not only contains a labile pool of nutrients but also drives the cycling of organic matter and nutrients in soil, thereby affecting S availability to plants (Banerjee and Chapman, 1996). Previous studies have compared seasonal responses of microbial biomass C and N to changes to environmental factors such as temperature (Van Gestel *et al.*, 2011), moisture (He *et al.*, 1997), plant growth (Franzluebbers, Hons and Zuberer, 1994) and fertilizer application (Ge *et al.*, 2010), to assess the importance of the microbial biomass in regulating the plant availability of C and N in soil. However, research for other major plant nutrients, such as S, are still limited.

Microbial biomass-S in soil is considered as intermediary between soil inorganic and organic S, and therefore the retention or release of this S pool is vital to support plant growth. Although only a relatively small fraction of the soil organic S resides in microbial biomass-S, this fraction is relatively labile and is thought to be the most active S pool for soil S turnover. Microbial biomass S can form a significant proportion of the organic S pool which is involved in cycling and potentially available to plants (Chapman, 1987), and its availability depends on its turnover rate. Therefore, additional studies are particularly needed to improve our understanding of the seasonal variations of microbial biomass S under more soil types with different plant species. In addition, the partitioning of microbial S between C-bonded S and C-O-S and how these two fractions turnover seasonally is also of particular interest.

## 9.5 Concluding remarks

In this thesis, I addressed fundamental questions related to amino acid-S such as their mineralization rates, contribution to plant nutrition that could have key implications for soil nutrient management purposes. This thesis provided the following essential information: 1) the mineralization of S containing amino acids is extremely rapid, representing an important short-term source of C, N and S in grassland soils; 2) maize plant roots can acquire S-containing amino acids directly under hydroponic conditions, even when inorganic sulphate was available, indicating low molecular organic S compounds like S-containing amino acids could constitute a significant proportion of S supply to plant growth under certain conditions; and 3) when in competition, the capture of free Cys and Met-C (or S) by the maize plants was low compared to soil microbes, with < 10% of the added amino acid captured by the plant roots, compared to the rhizosphere microbial community (around 80%). Despite these advances, there is still much more to explore about S cycling in both an agricultural and natural ecosystem settings. Additional research will be important to enhance our understanding of how to optimise the formulation and use of S fertilisers in agriculture and for understanding the impacts and legacy of anthropogenic S deposition on natural ecosystems.

## 9.6 References

- Banerjee, M. R. and Chapman, S. J. (1996) 'The significance of microbial biomass sulphur in soil', *Biology and Fertility of Soils*, 22(1), pp. 116–125. doi: 10.1007/s003740050085.
- Boddy, E. *et al.* (2007) 'Fast turnover of low molecular weight components of the dissolved organic carbon pool of temperate grassland field soils', *Soil Biology and Biochemistry*, 39(4), pp. 827–835. doi: 10.1016/j.soilbio.2006.09.030.
- Brailsford, F. L. *et al.* (2020) 'Rapid depletion of dissolved organic sulphur (DOS) in freshwaters', *Biogeochemistry*. Springer, 149(1), pp. 105–113. doi: 10.1007/s10533-

- Broughton, R. C. I. *et al.* (2015) 'Differential acquisition of amino acid and peptide enantiomers within the soil microbial community and its implications for carbon and nitrogen cycling in soil', *Soil Biology and Biochemistry*. Elsevier, 88, pp. 83–89. doi: 10.1016/j.soilbio.2015.05.003.
- Burns, R. G. *et al.* (2013) 'Soil enzymes in a changing environment: current knowledge and future directions', *Soil Biology and Biochemistry*. Elsevier, 58, pp. 216–234.
- Carpenter, M. and Milyo, J. (2012) 'E.,(2013)', *Differences Between an Acid Detergent Fiber & a Neutral Detergent Fibe.*
- Chapman, S. J. (1987) 'Microbial sulphur in some scottish soils', *Soil Biology and Biochemistry*, 19(3), pp. 301–305. doi: 10.1016/0038-0717(87)90013-7.
- Charteris, A. F. (2019) 'Compound-Specific Amino Acid 15 N Stable Isotope Probing of Nitrogen Assimilation by the Soil Microbial Biomass Using Gas Chromatography-Combustion-Isotope Ratio Mass Spectrometry', in *15N Tracing of Microbial Assimilation, Partitioning and Transport of Fertilisers in Grassland Soils*. Springer, pp. 57–78.
- Churka Blum, S. *et al.* (2013) 'Sulfur forms in organic substrates affecting S mineralization in soil', *Geoderma*. Elsevier B.V., 200–201, pp. 156–164. doi: 10.1016/j.geoderma.2013.02.003.
- Eriksen, J., Lefroy, R. D. B. and Blair, G. J. (1995) 'Physical protection of soil organic S studied by extraction and fractionation of soil organic matter', *Soil Biology and Biochemistry*, 27(8), pp. 1011–1016. doi: 10.1016/0038-0717(95)00032-A.
- Franzluebbers, A. J., Hons, F. M. and Zuberer, D. A. (1994) 'Seasonal changes in soil microbial

- biomass and mineralizable c and n in wheat management systems’, *Soil Biology and Biochemistry*, 26(11), pp. 1469–1475. doi: 10.1016/0038-0717(94)90086-8.
- Ge, G. *et al.* (2010) ‘Soil biological activity and their seasonal variations in response to long-term application of organic and inorganic fertilizers’, *Plant and Soil*, 326(1), pp. 31–44. doi: 10.1007/s11104-009-0186-8.
- Van Gestel, N. C. *et al.* (2011) ‘Reductions in daily soil temperature variability increase soil microbial biomass C and decrease soil N availability in the Chihuahuan Desert: Potential implications for ecosystem C and N fluxes’, *Global Change Biology*, 17(12), pp. 3564–3576. doi: 10.1111/j.1365-2486.2011.02479.x.
- Glanville, H. C. *et al.* (2016) ‘Combined use of empirical data and mathematical modelling to better estimate the microbial turnover of isotopically labelled carbon substrates in soil’, *Soil Biology and Biochemistry*. Elsevier Ltd, 94, pp. 154–168. doi: 10.1016/j.soilbio.2015.11.016.
- Gunina, A. *et al.* (2017) ‘Microbial uptake and utilization of low molecular weight organic substrates in soil depend on carbon oxidation state’, *Biogeochemistry*. Springer, 133(1), pp. 89–100.
- He, Z. L. *et al.* (1997) ‘Seasonal responses in microbial biomass carbon, phosphorus and sulphur in soils under pasture’, *Biology and Fertility of Soils*, 24(4), pp. 421–428. doi: 10.1007/s003740050267.
- Higuchi, T. *et al.* (2020) ‘Sulfur assimilation in corals with aposymbiotic and symbiotic zooxanthellae’, *Environmental Microbiology Reports*. Wiley Online Library.
- Hill, P. W., Farrar, J. F. and Jones, D. L. (2008) ‘Decoupling of microbial glucose uptake and mineralization in soil’, *Soil Biology and Biochemistry*, 40(3), pp. 616–624. doi:



10.1016/j.soilbio.2007.09.008.

Hill, P. W. and Jones, D. L. (2019) 'Plant–microbe competition: does injection of isotopes of C and N into the rhizosphere effectively characterise plant use of soil N?', *New Phytologist*. Wiley Online Library, 221(2), pp. 796–806.

Jones, D. L. *et al.* (2005) 'Plant capture of free amino acids is maximized under high soil amino acid concentrations', *Soil Biology and Biochemistry*, 37(1), pp. 179–181. doi: 10.1016/j.soilbio.2004.07.021.

Jones, D. L. and Willett, V. B. (2006) 'Experimental evaluation of methods to quantify dissolved organic nitrogen (DON) and dissolved organic carbon (DOC) in soil', *Soil Biology and Biochemistry*. Elsevier, 38(5), pp. 991–999.

Kertesz, M. A. and Mirleau, P. (2004) 'The role of soil microbes in plant sulphur nutrition', *Journal of Experimental Botany*, 55(404), pp. 1939–1945. doi: 10.1093/jxb/erh176.

Knowles, T. D. J. *et al.* (2010) 'Tracing the rate and extent of N and C flow from <sup>13</sup>C,<sup>15</sup>N-glycine and glutamate into individual de novo synthesised soil amino acids', *Organic Geochemistry*. Elsevier Ltd, 41(12), pp. 1259–1268. doi: 10.1016/j.orggeochem.2010.09.003.

De Kok, L. J. and Tausz, M. (2001) 'The role of glutathione in plant reaction and adaptation to air pollutants', in *Significance of glutathione to plant adaptation to the environment*. Springer, pp. 185–205.

Meyer, N., Welp, G. and Amelung, W. (2019) 'Effect of sieving and sample storage on soil respiration and its temperature sensitivity (Q<sub>10</sub>) in mineral soils from Germany', *Biology and Fertility of Soils*. Springer, 55(8), pp. 825–832.

Miransari, M. (2013) 'Soil microbes and the availability of soil nutrients', *Acta Physiologiae*

- Plantarum*, pp. 3075–3084. doi: 10.1007/s11738-013-1338-2.
- Nannipieri, P. *et al.* (2003) ‘Microbial diversity and soil functions’, *European Journal of Soil Science*, 54(4), pp. 655–670. doi: 10.1046/j.1351-0754.2003.0556.x.
- Oburger, E. and Jones, D. L. (2009) ‘Substrate mineralization studies in the laboratory show different microbial C partitioning dynamics than in the field’, *Soil Biology and Biochemistry*. Elsevier, 41(9), pp. 1951–1956.
- Omasa, K. *et al.* (2012) *Air pollution and plant biotechnology: prospects for phytomonitoring and phytoremediation*. Springer Science & Business Media.
- Romero, L. C. *et al.* (2014) ‘Cysteine and cysteine-related signaling pathways in arabidopsis thaliana’, *Molecular Plant*, 7(2), pp. 264–276. doi: 10.1093/mp/sst168.
- Ross, D. J. (1992) ‘Influence of sieve mesh size on estimates of microbial carbon and nitrogen by fumigation-extraction procedures in soils under pasture’, *Soil biology and biochemistry*. Elsevier, 24(4), pp. 343–350.
- Rousk, J. and Jones, D. L. (2010) ‘Loss of low molecular weight dissolved organic carbon (DOC) and nitrogen (DON) in H<sub>2</sub>O and 0.5M K<sub>2</sub>SO<sub>4</sub> soil extracts’, *Soil Biology and Biochemistry*. Elsevier Ltd, 42(12), pp. 2331–2335. doi: 10.1016/j.soilbio.2010.08.017.
- Sprent, J. I., Knowles, R. and Blackburn, T. H. (1993) ‘Nitrogen Isotope Techniques.’, *The Journal of Applied Ecology*. Elsevier, 30(3), p. 576. doi: 10.2307/2404198.
- Stuart, R. K. *et al.* (2020) ‘Bidirectional C and N transfer and a potential role for sulfur in an epiphytic diazotrophic mutualism’, *The ISME Journal*. Nature Publishing Group, 14(12), p. 3068.
- Tang, K. (2020) ‘Chemical diversity and biochemical transformation of biogenic organic sulfur in the ocean’, *Frontiers in Marine Science*. Frontiers, 7, p. 68.

- Vermeiren, C. *et al.* (2018) 'Model-based rationalization of sulphur mineralization in soils using <sup>35</sup>S isotope dilution', *Soil Biology and Biochemistry*. Elsevier, 120(May 2017), pp. 1–11. doi: 10.1016/j.soilbio.2018.01.013.
- Wang, Zhanyi *et al.* (2012) 'Dissolved organic sulfur in streams draining forested catchments in southern China', *Journal of Environmental Sciences*, 24(4), pp. 704–710. doi: 10.1016/S1001-0742(11)60781-X.
- Watkinson, J. H., Perrott, K. W. and Thorrold, B. S. (1991) 'Relationship between the MAF pasture development index of soil and extractable organic sulphur', *Soil and Plant Testing for Nutrient Deficiencies and Toxicities*. Eds. RE White and LD Currie, pp. 66–71.

## Appendix 1

---

**Chapter 6 Estimation of sulphur transformation rates in grassland soils using  $^{35}\text{S}$  isotope pool dilution (This chapter is not complete due to author being unable to read  $^{32}\text{S}$  data for the final isotope dilution model)**

## Abstract


Due to reductions in fossil fuel-derived SO<sub>2</sub> emissions and atmospheric sulphur (S) deposition, many agroecosystems are now experiencing a greater incidence of S deficiency. This has led to a renewed interest in understanding the processes which regulate the supply of S to plants. In well drained grassland soils, most S is locked up in soil organic matter (SOM), however, a large proportion of this organic S is poorly bioavailable to soil microorganisms. The primary objective of this study was to quantify the size of labile organic S pool, and to characterize soil S turnover rates in grassland soils with contrasting soil properties. To achieve this, we used an isotope (<sup>35</sup>SO<sub>4</sub><sup>2-</sup>) pool dilution method in a closed incubation experiment over a 70-d period. After addition of <sup>35</sup>S-labelled SO<sub>4</sub><sup>2-</sup> to each soil, we measured the progressive immobilization of this <sup>35</sup>S into the SOM pool on a weekly basis. The amount of <sup>35</sup>SO<sub>4</sub><sup>2-</sup> remaining in the soil at each measurement time was estimated by extracting the soil with 0.01 M Ca(H<sub>2</sub>PO<sub>4</sub>)<sub>2</sub>. Gross S mineralization rates and the size of total labile S pools were quantified using an established isotope pool dilution model. The modelling approach provided a more accurate estimations of gross S cycling processes, as it takes into account the immobilization of SO<sub>4</sub><sup>2-</sup> into the labile organic S pool (LOS), and the re-mineralization of this newly formed LOS pool into account. In addition, it is clear that S cycling (mineralization, immobilization) in the soil is microbially mediated, and affected by nutrient conditions. For example, S mineralization in soil is attributed to either biological or biochemical processes, where biological processes are governed by the microbial search for energy from carbon-bonded S, and where the biochemical process is via enzymatic hydrolysis of sulphate esters controlled by S supply. On the other hand, S immobilization also depends on soil conditions, as carbon availability largely controls soil microbial activity, whereas SO<sub>4</sub><sup>2-</sup> availability was found to retard <sup>35</sup>S immobilization. Therefore, a secondary objective was to evaluate effects of additions of various substrates (glucose, SO<sub>4</sub><sup>2-</sup>-S or a combination of both) on the turnover of <sup>35</sup>S in soil.

**Keywords:** Isotope pool dilution method; Labile organic S; S turnover; Grassland soil.

## 6.1 Introduction

Sulphur (S) is an essential element for plant growth and development (Zhao *et al.*, 1997) as well as being involved in a range of processes involved in biotic and abiotic stress mitigation (Bloem, Haneklaus and Schnug, 2005). It is a key component of the amino acids, cysteine and methionine, as well as several co-enzymes. Historically, agricultural production in most European nations was not limited by S availability due to the emissions and subsequent atmospheric deposition of S from fossil-fuel burning. However, the almost complete removal of SO<sub>2</sub> from many power plant emissions and the adoption of low S containing fertilizers within Europe has led to an increase in S deficiency in many agricultural regions (Engardt *et al.*, 2017). The increased recognition of the importance of maintaining an adequate S supply for optimal crop growth has therefore led to a renewed interest in understanding the key factors that regulate S cycling in soil.

Typically, more than 95% of the S present in soil is bound within organic matter (SOM) (Ghani, McLaren and Swift, 1993), typically in the form of sulphate ester S and carbon-bonded S (Norman M Scott and Anderson, 1976; Lou and Warman, 1992). Although a proportion of this organic S pool is quasi-stable due to chemical and physical protection, the unprotected proportion is thought to be extremely dynamic, with S being continuously cycled between inorganic and organic forms (McLaren, Keer and Swift, 1985). This is likely to be particularly true in the rhizosphere where microbial activity is greatest and plant demand continually removes inorganic S from the soil solution. Specifically, the transformation of soil organic-S to inorganic SO<sub>4</sub><sup>2-</sup> (i.e. S mineralization as well as enzymatic cleavage), and the reverse process, immobilization (the incorporation of SO<sub>4</sub><sup>2-</sup> into soil organic-S), are two key processes governing S availability in soils (Scherer, 2001).



**Fig. 6.1** Schematic representation of the two ongoing processes in our inorganic sulphate incubation experiment. On one hand, there is quick immobilization of  $^{35}\text{S}$  in newly formed microbial biomass, which will decay and again be released as inorganic sulphate. On the other hand, inorganic sulphate is also slowly turned into labile organic sulphur, which could be mineralized to inorganic sulphate again, this process is mediated by microorganisms. In our study, however, we only took immobilization and mineralization between inorganic sulphate and labile organic sulphur into account.

In well drained surface soils, the amounts of inorganic  $\text{SO}_4^{2-}$  present is typically too small to satisfy plant demand unless it is regularly replenished. Consequently, mineralization plays a key role in resupplying this pool and maintaining plant S nutrition (Williams, 1967). Studies on the turnover of organic S in soils will, therefore, facilitate an improved prediction of S response to reduced S inputs (Haque and Walmsley, 1972; A N Ganeshamurthy and Nielsen, 1990). In grassland soils, a significant fraction of the soil organic S is either insoluble or physically protected on clay surfaces, and is therefore not directly available for microbial attack (Eriksen et al., 1995). This pool is defined as non-labile organic S. By contrast, labile soil organic S can be readily mineralized and the size of this pool can be used as an indicator of soil quality (Fig. 6.1; (Maynard, Stewart and Bettany, 1984)). To date, numerous soil extraction methods have been developed to assess the size of the labile organic S pool (e.g. by extracting with hot water or with alkaline solutions) (Dick, Kost and Chen, 2008); (Spencer and Freney, 1960; Fox, Olson and Rhoades, 1964). However, these methods are often time consuming, and strong extractants may alter the nature of organic S during the extraction procedure, while milder extractants may result in incomplete recovery from the soil (Freney, Melville and Williams, 1969). Therefore, it is difficult to gain a clear idea of the size and dynamics of the labile S pool(s) using a conventional chemical extraction approach.

Previously, low molecular weight (MW) organic-S compounds (e.g. methionine (Fitzgerald and Andrew, 1984), cysteine (Hale and Fitzgerald, 1990), sulphate esters (Houghton and Rose, 1976) and sulphoquinovose (Fitzgerald, 1983)) have been added to soil and the generation of inorganic S measured. However, most soil organic-S in soil is thought to be associated with higher MW substances (e.g., protein-humic complexes), so it may be inappropriate to extrapolate results from S mineralization studies that solely focus on simple organic compounds.



It is highly likely that the net rate of S mineralization in soil will be greatly influenced by the availability of other key nutrients (e.g., C, N and P), as well as the amount of inorganic S present. Their availability will also influence the final form of S incorporated into the microbial biomass. For example, low  $\text{SO}_4^{2-}$  concentrations in soil may stimulate the production of sulfohydrolases by plant roots or soil microorganisms (Maynard, Stewart and Bettany, 1984), leading to increased biochemical mineralization of ester sulphates. In contrast, a high soil  $\text{SO}_4^{2-}$  content may stimulate S immobilization into higher ester sulphates, since sulphate ester production is a mechanism for soil microorganisms, especially fungi, to store S when there is an adequate supply of  $\text{SO}_4^{2-}$  (Ghani, McLaren and Swift, 1992). In addition, amendment with a readily available C source (e.g. glucose) may stimulate rapid immobilization of  $^{35}\text{S}$  into newly formed microbial biomass, leading to an increased incorporation of  $\text{SO}_4^{2-}$  into organic-S (S J Chapman, 1997; Shahsavani and Gholami, 2009).

Quantifying the dynamics of soil S is difficult as mineralization and immobilization processes take place simultaneously (Michael A Kertesz and Mirleau, 2004). The mineralization rate of soil S can be investigated using either “closed” or “open” incubations techniques in the laboratory. The closed incubation technique allows mineralized sulphate to accumulate (D G Maynard, Stewart and Bettany, 1983; Valeur and Nilsson, 1993), but this technique does not take account of the loss of newly formed  $\text{SO}_4^{2-}$  (e.g. via immobilization, plant uptake or leaching). ‘Open’ incubation techniques involve periodic leaching of  $\text{SO}_4^{2-}$  (D G Maynard, Stewart and Bettany, 1983; Ghani, McLaren and Swift, 1991), or quantifying plant S uptake plus increases in soil  $\text{SO}_4^{2-}$  at regular intervals (Eriksen *et al.*, 1995; McLaren, 2001). The ‘open’ approach more closely replicates field conditions, where mineralized sulphate is either taken up by plant roots or lost by leaching. However, periodic leaching limits the potential for  $\text{SO}_4^{2-}$  immobilization (as it is removed from the system) and can therefore only be used to quantify gross mineralization. Further, repeated leaching may alter soil conditions (e.g.

removing  $\text{NO}_3^-$ , altering soil pH). Together all these factors may lead to overestimation of S mineralization with the ‘open’ approach.

An alternative approach is the use of stable or radio-isotopes (e.g.  $^{32}\text{S}$ ,  $^{34}\text{S}$ ,  $^{35}\text{S}$ ) to monitor the flow of S between different pools in soil. Previous studies using  $\text{Na}_2^{35}\text{SO}_4$  have shown the added  $^{35}\text{S}$  label can become readily incorporated into both hydriodic acid reducible S (sulphate esters) and C-bonded S (Freney, Melville and Williams, 1971), and the nature of the resulting organic S compounds largely determines the subsequent mineralization rate of S (Zhao, Wu and McGrath, 1996). The isotope pool dilution technique provides quantitative information on nutrient dynamics in the soil-plant system and when combined with mathematical modelling can be used to determine both gross and net S transformation rates in soil (Di, Cameron and McLaren, 2000; Eriksen, 2005; Vermeiren *et al.*, 2018b).

In the following study, the rates of  $^{35}\text{S}$  released to and removed from the soil inorganic pool were estimated using the isotopic pool dilution method. The objectives of this study were to (1) quantify the size of the total labile S pool (sum of labile organic S and inorganic  $\text{SO}_4^{2-}$ ) in grassland soils using  $^{35}\text{S}$  as a tracer; (2) compare the modelled results of the labile S pool with those measured by conventional chemical extractant techniques; (3) evaluate gross S transformation rates with an established model; and (4) assess the effects of C availability (glucose) or  $\text{SO}_4^{2-}$  availability on S transformation rates.

## **6.2 Materials and methods**

### **6.2.1 Field sites and soil sampling**

Topsoil (0 - 10 cm) and subsoil (10 - 20 cm) was collected from seven sites along a natural grassland productivity gradient, located in Abergwyngregyn, Gwynedd, North Wales. Approximately 2 kg of soil was collected from each of three random points located 2 m apart from each other, placed in gas-permeable plastic bags and transferred immediately to the

laboratory. These samples represent the three replicates used for all experiments. After collection, all soils were sieved to < 2 mm and stored at 4 °C. Soil bulk density and moisture content of the original top- and subsoil was assessed by oven drying (80 °C, 24 h) intact 100 cm<sup>3</sup> cores collected from the field. Soil pH and electrical conductivity (EC) measurements were performed on 1:1.25 w/v soil-distilled water suspensions using standard electrodes. Selected properties of these soils are presented in Table 3.1 and 3.2.

### **6.2.2 Extraction of total labile S**

Three established single-step extraction tests were performed to estimate total labile S on each untreated soil.

For the hot water extraction (Ghani, Dexter and Perrott, 2003; Vong *et al.*, 2010), 5 g of field-moist soil was weighed into 50 ml polypropylene centrifuge tubes, after which, deionized water (25 ml) was added, and the suspensions manually shaken. The mixture was then incubated for 18 h at 80 °C. The second extractant was 0.5 M NaHCO<sub>3</sub> (pH 8.5), mirroring the common Olsen-P method used to measure available P (Abdu, 2006). NaHCO<sub>3</sub> (20 ml) was added to 5 g of field-moist soil and the suspension placed on an end-over-end shaker for 1 h at 20 °C. The third extractant was 0.25 M KCl (Blair, Lefroy and Chinoim, 1993). This procedure involved the addition of 25 ml of 0.25 M KCl to 5 g of field-moist soil after which the suspension was manually shaken and then incubated at 40 ± 0.5 °C for 3 h.

For all three extractions, the samples were left to cool (20 °C) before being centrifuged (3500 rpm, 20 min), filtered (0.45 µm sterile PES syringe filter; Fisher Scientific), before S analysis using an inductively coupled plasma optical emission spectroscopy (ICP-OES; Varian 710ES, Agilent Technologies, USA).

### 6.2.3 Incubation experiment using $^{35}\text{S}$ isotope pool dilution

The experiment had 4 treatments to test how the availability of inorganic S and glucose-C affected  $^{35}\text{S}$  cycling in soil. These were undertaken in a factorial design as follows:

1. -S -C (control)
2. -S +C (glucose addition only)
3. +S -C (sulphate addition only)
4. +S +C (glucose and sulphate addition)

The -S treatments just contained a very small amount of S ( $1 \mu\text{g SO}_4^{2-}\text{-S kg}^{-1}$ ) to label the native  $\text{SO}_4^{2-}$  pool, but importantly without changing its intrinsic concentration. The +S additions were designed to greatly increase the native  $\text{SO}_4^{2-}$  pool and involved the addition of  $30 \text{ mg SO}_4^{2-}$  (as  $\text{Na}_2\text{SO}_4$ )  $\text{kg}^{-1}$  soil (equivalent to  $30 \text{ kg S ha}^{-1}$ ), while the C addition was  $5 \text{ g C}$  (as glucose)  $\text{kg}^{-1}$  soil, respectively. The +S treatment represents that typically used as a fertiliser dose in agricultural fields (Kulczycki, 2021).

To maintain the natural moisture content of the soil, the  $^{35}\text{SO}_4^{2-}$  tracer ( $\pm$  glucose) was added to the soil using an inert quartz sand carrier. To prepare the carrier, a solution of  $\text{Na}_2^{35}\text{SO}_4$  (American Radiolabelled Chemicals Inc., St Louis, MO) and/or glucose was mixed with pure sterile quartz sand and left to dry at room temperature. Glucose was chosen as a readily available C source, as it is highly abundant in root exudates (Rovira, 1969b). For the experiments, the  $^{35}\text{S}$ -labelled quartz sand was mixed with the soil in a ratio of 1:10 (w/w; quartz sand: soil). Briefly,  $25 \text{ g}$  of field-moist soil were placed in individual polypropylene containers ( $5 \text{ cm}$  internal diameter;  $7.3 \text{ cm}$  height; Fisher Scientific, Leicestershire, UK). Quartz sand ( $2.5 \text{ g}$ ) carrying the  $^{35}\text{S}$  ( $\pm$  glucose) was then thoroughly mixed with the soil to give a final  $^{35}\text{S}$  activity of  $0.35 \text{ kBq g}^{-1}$  soil.

The soil was then compressed to match the bulk density measured in the field. All containers were then incubated at  $20^\circ\text{C}$  in the dark for 70 d. Containers were loosely covered

with plastic lids and opened regularly to ensure adequate aeration. Subsamples of soil (2 g) were taken from each container every 7 d for analysis of extractable  $^{35}\text{S-SO}_4^{2-}$  and  $^{32}\text{S-SO}_4^{2-}$ . To measure the amount of inorganic  $^{35}\text{SO}_4^{2-}$  remaining in the soil at each sampling point, the 2 g sub-sample was extracted with 10 ml of 0.01 M  $\text{Ca}(\text{H}_2\text{PO}_4)_2$  on an end-over-end shaker (30 min, 200 rev min<sup>-1</sup>), centrifuged (4000 rpm, 10 min) and filtered (0.45  $\mu\text{m}$  syringe filter) prior to analysis. A 1 ml aliquot of the filtrate was used for analysis of  $^{35}\text{S-SO}_4^{2-}$  by liquid scintillation counting using a Wallac 1404 scintillation counter with automated quench correction (Wallac EG&G, Milton Keynes, UK) and Optiphase HiSafe 3 scintillation fluid (PerkinElmer, Waltham, MA). The  $^{32}\text{S-SO}_4^{2-}$  content of the filtrate was measured by ion chromatography using an ICS2100 ion chromatograph (Dionex Corp., Sunnyvale, CA; (Zhao and McGrath, 1994)). The  $^{35}\text{S}$  present in our 0.01 M  $\text{Ca}(\text{H}_2\text{PO}_4)_2$  extracts represented the non-immobilized  $^{35}\text{S}$ , and the difference between  $^{35}\text{S}$  initially applied and  $^{35}\text{S}$  found in the  $\text{Ca}(\text{H}_2\text{PO}_4)_2$  extracts was assumed to be  $^{35}\text{S}$  that had been immobilized. Prior to data analysis, the  $^{35}\text{S}$  measurements were all corrected for the natural decay of  $^{35}\text{S}$  (half-life 87.1 d).

$^{35}\text{S}$  has been used to successfully trace a range of S cycling processes in soil including microbial immobilisation, reduction, sorption, leaching and plant uptake (Di et al., 2000). The isotope pool dilution method is best used when the  $^{35}\text{S}$  radiotracer is combined with measurements of its stable isotope counterpart (i.e.  $^{32}\text{S}$ ). The ratio of introduced label ( $^{35}\text{S}$ -labelled) with the native pool ( $^{32}\text{S}$ -labelled) can then be used to calculate flux rates (Table 6.1). For example,  $W_x$  can be calculated according to equation:

$$W_x = W_d \times A_d / A_r, \quad (\text{Eqn. 1})$$


Where  $W_d$ ,  $A_d$  and  $A_r$  are known parameters.

**Table 6.1** Summary of the main pools used in the isotope dilution analysis.

	Total radioactivity	Weight	Ratio
Native analyte	0	$W_x$	0
Adding radioactive analyte	A	$W_d$	$A_d$
Mixture	A	$W_x + W_d$	$A_r$

#### 6.2.4 Compartment model

The fluxes of S in soil can be described as follows:



Where  $M$  is defined as the mineralization rate and  $I$  is the immobilization rate, both in  $\text{mg kg}^{-1} \text{ soil day}^{-1}$ . Some of the S immobilized during the incubation can be re-mineralized, however, this cannot be measured. Overall, this can be expressed by the following equation:

$$f = (1 - f_{\text{bio}}) \times (f_s + (1 - f_s) \times \exp^{-k_{\text{los}} \times t}) \quad (\text{Eqn. 2})$$

Where  $f_{\text{bio}}$  represents the fraction of  $^{35}\text{S}$  that is quickly incorporated into the microbial biomass, and  $f$  is the fraction of  $^{35}\text{S}$  in the total added S,  $k_{\text{los}}$  is the rate constant for exchange between the sulphate and LOS pool,  $f_s$  is the fraction of  $^{35}\text{S}$  in the total labile S pool (Total labile S = sulphate + LOS) present as  $\text{SO}_4^{2-}$  at steady state. By fitting data into this equation, the best fitted parameters for  $f_{\text{bio}}$ ,  $k_{\text{los}}$  and  $f_s$  can be obtained.

Since previous studies have reported that glucose addition can result in rapid incorporation of  $^{35}\text{S}$  into microbial biomass, with little change in total biomass S, this indicates that original non-labelled biomass-S was replaced by  $^{35}\text{S}$ . Based on parameters ( $f_{\text{bio}}$ ,  $k_{\text{los}}$  and  $f_s$ ) derived from above, LBS, LOS and  $M$  can be obtained according to following equations separately:

$$[\text{LBS}] = [\text{SO}_4^{2-}\text{-S}] \times \ln(1 / (1 - f_{\text{bio}})),$$

$$[\text{LOS}] = (1 / f_s - 1) \times [\text{SO}_4^{2-}\text{-S}];$$

$$M = I = k_{\text{los}} \times [\text{SO}_4^{2-}\text{-S}] \times [\text{LOS}] / ([\text{SO}_4^{2-}\text{-S}] + [\text{LOS}]).$$

Subsequently, the effect of glucose (5 g C kg<sup>-1</sup> soil) and sulphate (10 mg S kg<sup>-1</sup> soil) addition on sulphur transformation was also explored by repeating the same experiment as described above after substrate addition.

$$f = \exp ([LBS] / [SO_4^{2-}-S] \times \exp (-k_{bio} - 1)) \quad (\text{Eqn. 3})$$

Where [LBS] is the concentration of labile biomass S (the SO<sub>4</sub>-S immobilized in biomass when all glucose is decomposed), which can be derived by fitting data to equation (3), and where [SO<sub>4</sub><sup>2-</sup>-S] is considered to be a constant. These two parameters can be obtained by fitting equation (2) to the data with *k<sub>bio</sub>* set as a constant.

Equation (2) and (3) were added to account for both LOS and LBS, and since no analytical solution can be derived from equation 3, numerical solutions were derived using time steps of 0.25 d.

$$d_f / d_t = (M / [LOS] \times (1 - f) - I / [SO_4^{2-}-S] \times f) - (f \times [LBS] / [SO_4^{2-}-S] \times k_{bio} \times \exp (k_{bio} \times t)) \quad (\text{Eqn. 4})$$


## 6.3 Results

### 6.3.1 Labile sulphur pool size

### 6.3.2 Total labile S in relation to S concentration extracted by different extractants

### 6.3.3 Gross S transformation rates estimated with an established model

### 6.3.4 Effects of glucose or SO<sub>4</sub><sup>2-</sup> addition on S transformations



**Fig. 6.2** Changes with time in the recovery rate of extractable specific activity of  $^{35}\text{S}$  for soils 1, 5, 6 and 7 with or without substrate amendments. Values represent means  $\pm$  SEM ( $n = 3$ ).



#### 6.4. Discussion and conclusions

Results of the recovery of  $^{35}\text{S}$  in soils showed that, the –S-C treatment had the highest recovery rate, followed by +S-C, +S+C and –S+C treatments. Within a time period of 10 weeks, soil incorporated a large proportion of carrier free  $\text{Na}_2^{35}\text{SO}_4$  into the non-extractable fraction, ranging from  $27.0 \pm 4.8$  to  $92.8 \pm 1.6\%$  (Fig. 6.2), depending on specific soils and substrate amendments. The addition of carrier sulphate was found to retard the incorporation of  $^{35}\text{S}$  into the organo-sulphur pool, while the addition of glucose as a readily utilized carbon source was found to enhance this incorporation process due to an encouragement of rapid microbial growth. Our recovery test with quartz sand extracted by 0.01 M  $\text{Ca}(\text{H}_2\text{PO}_4)_2$  showed a recovery rate ca. 100% along our 70-day-incubation experiment for all sampling times (Fig. 6.2).

#### 6.5 Acknowledgements

I would like to thank Joseph Cotton and Mark Hughes for help with soil sample collection, and Sarah Chesworth and Jonathan Roberts for their technical assistance in the laboratory. This research was funded by the China Scholarship Council (201606510012) and Bangor University.

#### 6.6 References

- Abdu, N. (2006) ‘Soil-phosphorus extraction methodologies : A review’, 1(5), pp. 159–161.
- Blair, G. J., Lefroy, R. O. D. D. B. and Chinoim, N. A. N. T. H. A. N. A. (1993) ‘Sulfur soil testing’, pp. 383–386.
- Bloem, E., Haneklaus, S. and Schnug, E. (2005) ‘Significance of sulfur compounds in the protection of plants against pests and diseases’, *Journal of Plant Nutrition*, 28(5), pp. 763–784. doi: 10.1081/PLN-200055532.

- Chapman, S. J. (1997) 'Carbon substrate mineralization and sulphur limitation', *Soil Biology and Biochemistry*. Elsevier, 29(2), pp. 115–122.
- Di, H. J., Cameron, K. C. and McLaren, R. G. (2000) 'Isotopic dilution methods to determine the gross transformation rates of nitrogen, phosphorus, and sulfur in soil: a review of the theory, methodologies, and limitations', *Soil Research*, 38(1), pp. 213–230. doi: 10.1071/SR99005.
- Dick, W. A., Kost, D. and Chen, L. (2008) 'Availability of sulfur to crops from soil and other sources', *Sulfur: A missing link between soils, crops, and nutrition*. Wiley Online Library, 50, pp. 59–82.
- Engardt, M. *et al.* (2017) 'Deposition of sulphur and nitrogen in Europe 1900–2050. Model calculations and comparison to historical observations', *Tellus B: Chemical and Physical Meteorology*. Taylor & Francis, 69(1), p. 1328945.
- Eriksen, J. *et al.* (1995) 'Sulphur mineralisation in five Danish soils as measured by plant uptake in a pot experiment', *Agriculture, Ecosystems and Environment*, 56(1), pp. 43–51. doi: 10.1016/0167-8809(95)00632-X.
- Eriksen, J. (2005) 'Gross sulphur mineralisation-immobilisation turnover in soil amended with plant residues', *Soil Biology and Biochemistry*, 37(12), pp. 2216–2224. doi: 10.1016/j.soilbio.2005.04.003.
- Eriksen, J., Lefroy, R. D. B. and Blair, G. J. (1995) 'Physical protection of soil organic S studied using acetylacetone extraction at various intensities of ultrasonic dispersion', *Soil Biology and Biochemistry*, 27(8), pp. 1005–1010. doi: 10.1016/0038-0717(95)00031-9.
- Fitzgerald, J. W. (1983) 'Mineralization of sulphur in sulphooquinovose by forest soils', 15(3),

pp. 347–349. doi: [https://doi.org/10.1016/0038-0717\(83\)90082-2](https://doi.org/10.1016/0038-0717(83)90082-2).

Fitzgerald, J. W. and Andrew, T. L. (1984) ‘Mineralization of methionine sulphur in soils and forest floor layers’, *Soil Biology and Biochemistry*, 16(6), pp. 565–570. doi: 10.1016/0038-0717(84)90073-7.

Fox, R. L., Olson, R. A. and Rhoades, H. F. (1964) ‘Evaluating the sulfur status of soils by plant and soil tests’, *Soil Science Society of America Journal*. Wiley Online Library, 28(2), pp. 243–246.

Freney, B. J. R., Melville, G. E. and Williams, C. H. (1969) ‘Extraction, chemical nature, and properties of soil organic sulphur’, 20(42).

Freney, J. R., Melville, G. E. and Williams, C. H. (1971) ‘Organic sulphur fractions labelled by addition of <sup>35</sup>S-sulphate to soil’, *Soil Biology and Biochemistry*, 3(2), pp. 133–141. doi: 10.1016/0038-0717(71)90006-X.

Ganeshamurthy, A. N. and Nielsen, N. E. (1990) ‘Arylsulfatase and the biochemical mineralization of soil organic sulphur.’, *Soil biology & biochemistry*, 22(8), pp. 1163–1165.

Ghani, A., Dexter, M. and Perrott, K. W. (2003) ‘Hot-water extractable carbon in soils: A sensitive measurement for determining impacts of fertilisation, grazing and cultivation’, *Soil Biology and Biochemistry*, 35(9), pp. 1231–1243. doi: 10.1016/S0038-0717(03)00186-X.

Ghani, A., McLaren, R. G. and Swift, R. S. (1991) ‘Sulphur mineralisation in some New Zealand soils’, *Biology and Fertility of Soils*, 11(1), pp. 68–74. doi: 10.1007/BF00335838.

Ghani, A., McLaren, R. G. and Swift, R. S. (1992) ‘Sulphur mineralisation and transformations

- in soils as influenced by additions of carbon, nitrogen and sulphur', *Soil Biology and Biochemistry*, 24(4), pp. 331–341. doi: 10.1016/0038-0717(92)90193-2.
- Ghani, A., McLaren, R. G. and Swift, R. S. (1993) 'The incorporation and transformations of <sup>35</sup>S in soil: Effects of soil conditioning and glucose or sulphate additions', *Soil Biology and Biochemistry*, 25(3), pp. 327–335. doi: 10.1016/0038-0717(93)90131-T.
- Hale, D. D. and Fitzgerald, J. W. (1990) 'Generation of sulphate from cysteine in forest soil and litter.', *Soil Biology and Biochemistry*, 22(3), pp. 427–429. doi: 10.1016/0038-0717(90)90124-I.
- Haque, I. and Walmsley, D. (1972) 'Incubation studies on mineralization of organic sulphur and organic nitrogen', *Plant and Soil*. Springer, 37(2), pp. 255–264.
- Houghton, C. and Rose, F. A. (1976) 'Liberation of sulfate from sulfate esters by soils', *Applied and Environmental Microbiology*, 31(6), pp. 969–976. doi: 10.1128/aem.31.6.969-976.1976.
- Kertesz, M. A. and Mirleau, P. (2004) 'The role of soil microbes in plant sulphur nutrition', *Journal of Experimental Botany*, 55(404), pp. 1939–1945. doi: 10.1093/jxb/erh176.
- Lou, G. and Warman, P. R. (1992) 'Labile ester sulphate in organic matter extracted from podzolic soils', *Biology and Fertility of Soils*, 14(4), pp. 267–271. doi: 10.1007/BF00395462.
- Maynard, D. G., Stewart, J. W. B. and Bettany, J. R. (1983) 'Sulfur and nitrogen mineralization in soils compared using two incubation techniques', *Soil Biology and Biochemistry*, 15(3), pp. 251–256. doi: 10.1016/0038-0717(83)90067-6.
- Maynard, D. G., Stewart, J. W. B. and Bettany, J. R. (1984) 'Sulfur cycling in grassland and parkland soils', *Biogeochemistry*, 1(1), pp. 97–111. doi: 10.1007/BF02181123.

- McLaren, J. P. K. M. G. R. G. (2001) 'Comparison of three different methods of determining soil sulphur mineralization in relation to plant sulphur availability in soils', pp. 131–139. doi: 10.1007/s003740100378.
- McLaren, R. G., Keer, J. I. and Swift, R. S. (1985) 'Sulphur transformations in soils using sulphur-35 labelling', *Soil Biology and Biochemistry*, 17(1), pp. 73–79. doi: 10.1016/0038-0717(85)90092-6.
- Rovira, A. D. (1969) 'Plant root exudates', *The Botanical Review*, 35(1), pp. 35–57. doi: 10.1007/BF02859887.
- Scherer, H. W. (2001) 'Sulphur in crop production - Invited paper', *European Journal of Agronomy*, 14(2), pp. 81–111. doi: 10.1016/S1161-0301(00)00082-4.
- Scott, N. M. and Anderson, G. (1976) 'Organic Sulphur Fractions in Scottish Soils', (October 1975), pp. 358–366.
- Shahsavani, S. and Gholami, A. (2009) 'Study on the effect of sulphur, glucose, nitrogen and plant residues on the immobilization of sulphate-s in soil', *World Academy of Science, Engineering and Technology*, 37, pp. 114–118.
- Spencer, K. and Freney, J. R. (1960) 'A comparison of several procedures for estimating the sulphur status of soils', *Australian Journal of Agricultural Research*, 11(6), pp. 948–960. doi: 10.1071/AR9600948.
- Valeur, I. and Nilsson, I. (1993) 'Effects of lime and two incubation techniques on sulfur mineralization in a forest soil', *Soil Biology and Biochemistry*, 25(10), pp. 1343–1350. doi: 10.1016/0038-0717(93)90049-H.
- Vermeiren, C. *et al.* (2018) 'Model-based rationalization of sulphur mineralization in soils using  $^{35}\text{S}$  isotope dilution', *Soil Biology and Biochemistry*. Elsevier, 120(January), pp.

1–11. doi: 10.1016/j.soilbio.2018.01.013.

Vong, P. C. *et al.* (2010) ‘Effects of low-molecular-weight organic compounds on sulphur immobilization and re-mineralization and extraction of immobilized sulphur by hot-water and acid hydrolysis’, *European Journal of Soil Science*, 61(2), pp. 287–297. doi: 10.1111/j.1365-2389.2010.01228.x.

Williams, C. H. (1967) ‘Some factors affecting the mineralization of organic sulphur in soils’, *Plant and Soil*. Springer, 26(2), pp. 205–223.

Zhao, F. J. *et al.* (1997) ‘Sulphur nutrition: An important factor for the quality of wheat and rapeseed’, *Soil Science and Plant Nutrition*, 43(SPEC. ISS.), pp. 1137–1142. doi: 10.1080/00380768.1997.11863731.

Zhao, F. J., Wu, J. and McGrath, S. P. (1996) ‘Soil Organic Sulphur and its Turnover’, *Humic Substances in Terrestrial Ecosystems*, pp. 467–506. doi: 10.1016/b978-044481516-3/50013-x.

Zhao, F. and McGrath, S. P. (1994) ‘Extractable sulphate and organic sulphur in soils and their availability to plants’, *Plant and Soil*, 164(2), pp. 243–250. doi: 10.1007/BF00010076.