

Pre-competitive anxiety and autonomic responses in professional U-20 futsal players: Effect of the competition phase and game location

Paludo, Ana-Carolina ; Woodman, Tim; Owen, Julian; Rabelo, Felipe N.; Bernacikovà, Martina ; Simões, Antonio Carlos

Physiology and Behavior

DOI: https://doi.org/10.1016/j.physbeh.2022.113903

Published: 01/10/2022

Peer reviewed version

Cyswllt i'r cyhoeddiad / Link to publication

Dyfyniad o'r fersiwn a gyhoeddwyd / Citation for published version (APA): Paludo, A.-C., Woodman, T., Owen, J., Rabelo, F. N., Bernacikovà, M., & Simões, A. C. (2022). Pre-competitive anxiety and autonomic responses in professional U-20 futsal players: Effect of the competition phase and game location. Physiology and Behavior, 254, Article 113903. https://doi.org/10.1016/j.physbeh.2022.113903

Hawliau Cyffredinol / General rights Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

• Users may download and print one copy of any publication from the public portal for the purpose of private study or research.

- · You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

1	Pre-competitive anxiety and autonomic responses in professional U-20 futsal players:
2	effect of the competition phase and game location
3	
4	Ana Carolina Paludo ¹ , Tim Woodman ² , Julian A. Owen ² , Felipe N Rabelo ³ , Martina
5	Bernacikova ¹ , Antonio Carlos Simões ⁴ .
6	
7	¹ Faculty of Sport Studies Masaryk University: Masarykova Univerzita, Czechia
8	² School of Human and Behavioural Sciences, Bangor University, Wales, UK
9	³ Real Valladolid Club de Fútbol S.A.D, 47014 Valladolid, Spain
10	⁴ Department of Sport, School of Physical Education and Sport, São Paulo University, São
11	Paulo, Brazil
12	
13	
14	
15	Address for Correspondence:
16	Dr Ana Carolina Paludo,
17	Faculty of Sport Studies Masaryk University: Masarykova Univerzita Czechia.
18	Telephone: +420 725 164 109. Email: carolina.paludo@fsps.muni.cz
19	
20	Running title: Anxiety and autonomic responses in futsal athletes

Conflict of interest: The authors declare no conflict of interest.

22

ABSTRACT

23	The study examined whether pre-competition anxiety and autonomic responses in elite futsal
24	players change across the playoffs at away vs home venues. Heart rate variability, somatic and
25	cognitive anxiety and self-confidence (by CSAI-2R questionnaire) were evaluated in fourteen
26	male futsal players from a professional U-20 team, before competitive matches at the quarter-
27	final, semi-final and final stages played at home and away venues. Two-way ANOVA for
28	repeated measures was used, with factor one the game location and factor two the playoff stage,
29	with significance set at $p < 0.05$. Significant effects of the playoff stage were demonstrated in
30	the Mean of HR (F=4.643; p=0.014) and SD2 index (F= 14.83; p=< 0.001). No difference
31	was found for somatic and cognitive anxiety and self-confidence between the two factors. The
32	results suggest that play at the final stage of the playoff, regardless of the game location, may
33	cause higher physiological stress, demonstrated by elevated HR and decrease of SD2 index.
34	
35	Keywords: anxiety, cardiac autonomic, athletes, futsal, competition
36	
37	
38	
39	
40	
41	
42	

43 **1. Introduction**

Competition can be considered a stressful situation, increasing anxiety-state, triggering 44 physiological and emotional changes in athletes' responses, leading to changes in their 45 performance [1,2]. Regarding the autonomous nervous system (ANS), an increase in 46 sympathetic and a decrease in parasympathetic activation is observed when faced with a new 47 stressors [3]. In the sport settings, measurement and evaluation of the ANS via heart rate 48 49 variability is sensitive to describe the changes in the sympathovagal balance in stressful situations, such as the pre-competition [4-6]. The competitive environment includes several 50 51 contextual factors that may increase the perceived stress and lead to different behavior in athletes. 52

In team sport, contextual factors, such as the stage of competition, standard of 53 opponent, and game location could conceivably influence players' pre-competitive anxiety. 54 55 Previous studies demonstrated the effect of these contextual factors on pre-competitive anxiety and physiological responses such as salivary hormone concentrations [7,8] and autonomic 56 57 responses [4] are equivocal. Yet, it is unclear how athletes respond to these contextual factors, all together, in the field settings. In sports such as Futsal, the playoff structure consists of 58 playing home and away in each classification stage (e.g., quarterfinal, semi-final, and final). 59 Therefore, it might be expected that players' physiological and psychological responses may 60 61 be different depending on game location and the perceived importance of the competition stage. 62 Moreover, players in the transition from youth-professional to first-team have reported that they put more pressure on themselves to perform well and experience anxiety about moving to 63 senior sport [9]. 64

Therefore, we aimed to examine the effect of two contextual factors on pre-competition
anxiety and cardiac autonomic response in Futsal youth-professional players. Consequently,
this study aimed to examine pre-competition anxiety and autonomic responses during the

playoff quarter-final, semi-final, and final stages in games played away vs. home venue in a professional U-20 futsal team. It was postulated that playing in different contexts during the playoffs would promote different psychological and physiological responses in futsal athletes. We hypothesized that players would demonstrate higher scores of cognitive and somatic anxiety, lower self-confidence, together with changes in HRV suggestive of increased sympathetic tone (or vagal activity) when playing away (*versus* home) and in the final stage (*versus* quarter and semi-final).

75

76 **2.Material and methods**

77 2.1. Participants

Fourteen professional male futsal players belonging to the same team volunteered to 78 participate in the study. However, data from nine players who completed data collection from 79 80 all games wre analyzed (mean \pm sd: age, 19 \pm 0.68 years; height, 176.9 \pm 5.4 cm; body mass, 73.07± 5.87 kg). The team was participating in the Under-20 State Futsal Championship in São 81 Paulo, Brazil, and was ranked 1st in the State Championship during the period investigated. 82 Players were not taking prescribed medications or any special dietary supplement and did not 83 report any cardiovascular disease. All procedures were conducted with the approval of the 84 Ethics Committee of the local University (n° 544.410/2014), and informed consent was 85 obtained from each participant before study commencement. 86

87

88 2.2 Experimental design

Players were assessed in six games during the playoff stages (quarter-final, semi-final
and final), with each stage being played at home and away. The games occurred between 19:00
and 20:00h. Measurements were recorded approximately 35min prior to each game (before the
warm-up procedures). The first measurement was the cardiac autonomic response, assessed by

the recording of heart rate variability (HRV) for 10 min, followed by the completion of the
state-anxiety questionnaire (CSAI-2R). Data collection was performed inside the locker room
in both home and away games. Players were instructed to refrain from caffeine and alcohol in
the previous 24h.

97

98 *2.3. Autonomic response*

99 The autonomic response was measured via resting heart rate variability (HRV). Beatto-beat heart rate was recorded for 10min, using a portable heart rate monitor (Polar Team²) 100 101 Pro, Kempele, Finland) with an electrode belt firmly held around the thorax. The players remained in a seated position inside the locker room (21-22°C) and were instructed to stay 102 quiet, without speaking, breathing spontaneously, with eyes open. After collection, the data 103 104 was downloaded onto a computer using recognized software (Polar®Pro Trainer, Kempele, 105 Finland) and exported as a text file for analysis using Kubios 2.0 software (Biosignal Analysis and Medical Imaging Group, Finland). Data analysis was performed as previously described 106 [10]. The 10 min sampling time was chosen from standardized protocols for HRV data 107 collection under resting conditions [11], and the final 5min were selected to be analyzed due 108 to the most reliable method demonstrated in this sample [12]. 109

Each HRV index measures different features of ANS activity, with common methods 110 used to evaluate HRV at rest being time domain and spectral analyses. Complementary, non-111 112 linear indices were also calculated. Time-domain indices included mean heart rate (Mean HR), absolute square root of the mean of the sum of the squares of differences between adjacent RR 113 interval (RMSSD) and natural logarithm transformed (lnRMSSD). For the frequency domain 114 analysis, firstly, a Smooth Prior function was applied to the entire signal in order to remove the 115 slow trend components. Afterward, the signal was re-sampled at 4Hz using cubic splines, and 116 the Fast Fourier Transform (FFT) method was used to perform the spectral decomposition of 117

the signal. The bands within low (LF: 0.04-0.15Hz) and high (HF: 0.15-0.4 Hz) frequencies, and natural logarithm transformed (ln) of these indices, were calculated. The ratio between lnLF and lnHF was also reported. Finally, regarding the non-linear analysis, the SD1 (instantaneous beat-to-beat variability) and SD2 (long-term beat-to-beat variability) indices were calculated from the Poincaré plot. These indexes have been used before in team-sport athletes to assess the effect of competition stressor [4-6], especially the RMSSD and SD1 indexes due to the relation with parasympathetic modulation [13].

125

126 *2.4. Anxiety state*

The Revised Competitive State Anxiety Inventory-2 (CSAI-2R) [14] was administered 127 pre-competition to estimate the athletes' cognitive and somatic anxiety, as well as self-128 129 confidence levels. The CSAI-2R consists of 17 items scored on a 4-point Likert Scale (from 1 = not at all to 4 = very much so). A higher score related to cognitive and somatic anxiety 130 indicates a higher level of anxiety, and an elevated score on the self-confidence subscale 131 corresponds to a higher level of self-confidence. The Cronbach's alpha coefficients of internal 132 consistency were 0.84 for cognitive anxiety and self-confidence, and 0.88 for somatic anxiety 133 reported previously in Brazilian athletes [15]. 134

135

136 *2.5 Statistical analysis*

To analyze the effect of venue and playoffs factors, the two-way ANOVA for repeated measures was used. Factor one was the game location (home *versus* away) and factor two was the playoff stage (quarters, semi and finals). The assumptions of normality and heterogeneity of the variances were tested by Shapiro Wilk and Levene's tests, respectively. If the main effects or interaction effects were significant, Bonferroni post hoc tests were performed for multiple comparisons of different conditions. All Levene's values were higher than 0.05. The normality of the data was verified for each combination of the two factors as suggested
elsewhere [16], and significant deviations were observed for some indices of HRV and the
anxiety parameters. Considering the ANOVA is a robust test for violation of normality [17,18],
it was opted to use the test in the comparison of the two factors. The significance level was set
at p<0.05. Data were analyzed using JAMOVI statistical software package.

148

149 **3. Results**

Table 1 shows the outcomes of pre-competition HRV indices and anxiety state during 150 151 the playoff stage, whilst playing home and away, describing also the interaction of the two factors (stage * venue). Significant effects of the playoff stage were demonstrated by the Mean 152 of HR (F=4.643; p=0.014) and SD2 index (F= 14.83; p=<0.001). Based on these findings, 153 Figure 1 shows the effect of the playoff stages only on these variables, in which playing at the 154 final stage was a higher stressful condition compared to the quarter-final stage. No difference 155 was found between the game location for any the HRV indices. Somatic and cognitive anxiety 156 and self-confidence differed neither by the playoff stage nor the game location. 157

	Quarter-Finals		Semi-Finals		Finals		P-value	<i>P</i> -value	<i>P</i> -value
Variable	Home	Away	Home	Away	Home	Away	venue	payoff stage	interaction
Mean HR	70.3 ± 6.95	65.8 ±9.26	72.2 ± 6.75	80.3 ± 9.79	74.6 ± 10.3	74.7 ± 7.76	0.603	0.014*	0.095
RMSSD	46.2 ± 17.7	61.6 ± 26.5	47.6 ± 27.0	38.1 ± 19.7	40.9 ± 20.2	48.9 ±22.6	0.452	0.305	0.246
lnRMSSD	3.77 ± 0.39	3.48 ± 0.85	3.74 ± 0.50	3.31 ± 0.91	3.60 ± 0.70	3.79 ± 0.50	0.335	0.760	0.359
lnLF	7.57 ± 0.82	7.23 ± 0.62	7.34 ± 0.76	7.23 ± 0.87	7.25 ± 0.88	7.30 ± 0.58	0.529	0.852	0.743
lnHF	6.35 ± 0.869	6.77 ± 0.87	6.14 ± 1.01	6.02 ± 0.970	5.96 ± 1.08	6.36 ± 1.03	0.396	0.295	0.645
lnLF/HF	1.20 ± 0.13	1.08 ± 0.14	1.20 ± 0.17	1.21 ± 0.09	1.23 ± 0.09	1.17 ± 0.13	0.097	0.261	0.339
SD1	33.1 ± 12.6	44.1 ± 18.9	34.1 ± 19.2	27.3 ± 14.0	29.0 ± 14.3	34.6 ± 16.0	0.452	0.290	0.244
SD2	107 ± 34.4	135 ± 40.7	102 ± 35.4	95.5 ± 28.1	63.1 ± 20.4	67.2 ± 19.3	0.315	<0.001*	0.234
Anxiety-state									
Somatic	8.67 ± 2.74	9.33 ± 2.96	9.56 ± 3.43	9.78 ± 3.07	10.1 ± 3.76	10.8 ± 3.60	0.564	0.424	0.973
Cognitive	7.78 ± 2.22	7.89 ± 2.20	7.22 ± 2.22	7.56 ± 2.46	7.22 ± 2.22	7.33 ± 2.35	0.767	0.743	0.986
Self-confidence	18.4 ± 2.19	18.6 ± 1.67	18.9 ± 1.83	19.1 ± 1.69	19.3 ± 1.41	19.0 ± 1.73	1.000	0.505	0.883

1 Table 1. Pre-competition HRV and anxiety considering the game location and playoff stage (n=9).

2 *Significant difference in playoff stage (p<0.05).



Figure 1. Descriptive outcomes of the mean of heart rate (A) and SD2 index (B) during the
 playoff stage (quarter-final, semi-final and final). ^a Significant difference from quarter-final
 compared to semi-final and final. ^b Significant difference from final to quarte-final and semi-final.

8 4. Discussion

1

The purpose of the present study was to examine the effect of game location and playoff 9 10 stage on pre-competition cardiac autonomic responses and anxiety state in U-20 futsal players. It was hypothesized that players would demonstrate higher scores of cognitive and somatic 11 anxiety, lower self-confidence, together with changes in HRV suggestive of increased 12 sympathetic tone (or vagal activity) when playing away (versus home) and in the final stage 13 (versus quarter and semi-final). The main findings of the present study partially corroborate 14 15 the hypothesis formulated, indicating that playing the final stage was a challenging situation, compared to the quarter and semi-finals, showed by players' higher values of the Mean HR 16 17 and a decrease in SD2 values. Nonetheless, the hypothesis about the game location, somatic and cognitive anxiety and self-confidence were refuted in the study. No difference among the 18 19 playoff stages venue (home versus away) for pre-competition HRV and anxiety-state was verified. This result suggests that the players in the present study consider the final stage a 20 21 stressful factor independent of the game location.

1 Considering the contextual factors included in the present study, the game location was 2 chosen due to address the concept of territoriality and dominance, in which athletes seem to protect their territory against the opponent, demonstrating an increase in physiological arousal 3 [19]. However, the futsal players exhibited contradictory responses, with no difference in HRV 4 indices, it seems that players in the present study weigh the importance and demand of the 5 playoff stage more than the game location. Most studies have reported differences in HRV 6 7 indices in athletes' before competition compared to pre-training [4,5,20] but no comparison was found regarding playoff stage playing away versus at home. Recently, a study with female 8 9 soccer players evaluated the pre-competitive HRV and anxiety responses according to the demanding level of the match and found that a highly-demanding game led to a decrease in 10 RMSSD and SD1 indexes pre-competition. Also, the authors described an increase in cognitive 11 anxiety when compared with a lowly-demanding match [5]. Our results demonstrated that 12 besides the changes in Mean HR and SD2 indices regarding the playoff stage, the somatic and 13 cognitive anxiety and self-confidence did not change. A variation in somatic anxiety was 14 expected due to evidence of a positive correlation between pre-game HRV and somatic anxiety 15 state [6]. A possible explanation may be due to the physiological arousal level and individual 16 perception, in which, it is likely that individuals perceive physiological arousal (e.g., increases 17 in cardiac autonomic responses) when it was under higher levels. Maybe the increases in Mean 18 HR and decrease of SD2 index were not that higher for players who perceived the physiological 19 20 arousal changes.

A pre-competition context seems to inhibit the parasympathetic activity, leading to higher physiological arousal, usually shown with a reduction in the RMSSD and the SD1 indices [4,5,13]. Interestingly, the results of the current study presented a change during the playoff stages by the Mean HR and SD2 index only. The SD1 is directly linked to parasympathetic activity, however, the SD2 still needs a clear physiological meaning. Some reported that SD2 seems to be inversely proportional to sympathetic activity [21,22]. As described elsewhere, *if we take the inverse of SD2, we have a direct index for sympathetic activity... especially in a relationship to evaluate the sympathetic-parasympathetic balance* [23]. To improve the understanding of the SD2 index, Orellana et al [23] suggested a stress score (SS) based on the SD2 index (1000 x 1/SD2) for elite soccer players. The authors described the ratio SS: SD1 as a new index than can improve the physiological meaning of HRV by the Poincaré plot-analysis method.

8 One limitation of this study is the assessment of small sample size (n=9) from a selected 9 sport-team. The outcomes from the present study are specific samples; therefore, the current 10 results should be interpreted with caution. Besides the limitations, to our knowledge, this is the 11 first study involving pre-competitive cardiac autonomic response and anxiety state analyzed in 12 two different contexts: playoff stage and game location. Additionally, the assessment of HVR 13 performed in the field setting increases the ecological validity and the major strength of this 14 study.

As a practical application, coaches and physical trainers could monitor the players' HRV responses, especially through the playoff stage, in order to identify stressful levels before games. Interventions related to decreasing physiological arousal, such as biofeedback or neuromodulation, could be provided to manage the stress levels that the final stage of playoffs may be required.

In conclusion, the results indicated that the playoff stage, but not the game location, may be perceived as a stressful factor for professional U-20 futsal players. An increase in Mean HR and a decrease of SD2 index as shown by the players in the final stage of playoffs. These findings highlight the importance of monitoring cardiac autonomic status before the playoff games. The monitoring could be useful in identifying situations where interventions to manage pre-competition stress may be required. 1

2

Conflict of Competing Interests

or financial relationships that could be construed as a potential conflict of interest. 3

The authors declare that the research was conducted in the absence of any commercial

Acknowledgments 4

5

The authors wish to acknowledge the committed participation of all futsal players, staff, and coaches involved in this study. 6

7

References 8

9 [1] Woodman T, Mawn L, Martin C, Eklund BT. (Ed.). Models and theories of emotion-

performance. In Encyclopedia of sport and exercise psychology (2014 ed., pp. 448-452). Sage. 10

[2] Van Paridon KN, Timmis MA, Nevison CM, Bristow M. The anticipatory stress response 11

to sport competition; a systematic review with meta-analysis of cortisol reactivity. BMJ Open 12 Sport Exerc Med. 3(2017) e000261. 13

[3] Freeman R. Assessment of cardiovascular autonomic function. Clin. Neurophysiol. 14 117(2006) 716-730. 15

[4] Blásquez JCC, Font GR, Ortís LC. Heart-rate variability and precompetitive anxiety in 16 swimmers. Psicothema. 21 (2009) 531-536. 17

[5] Ayuso-Moreno RM, Fuentes-García JP, Collado-Mateo D, Villafaina S. Heart rate 18 variability and pre-competitive anxiety according to the demanding level of the match in female 19 20 soccer athletes. Physiol Behav. 222 (2020)112926.

[6] Mateo M, Blasco-Lafarga C, Martínez-Navarro I, Guzmán JF, Zabala M. Heart rate 21 variability and pre-competitive anxiety in BMX discipline. Eur J Appl Physiol. (2011) 1-11. 22

[7] Arruda AF, Aoki MS, Paludo AC, Moreira, A. Salivary steroid response and competitive 23

anxiety in elite basketball players: Effect of opponent level. Physiol Behav.177 (2017) 291-24

296. 25

- 1 [8] Paludo AC, Rabelo FN, Batista MM, Maciel IR, Tartaruga MP, Simões AC. Game location
- 2 effect on pre-competition cortisol concentration and anxiety state: A case study in a futsal team.
- 3 Rev de Psicol del Deporte. 29 (2020)105-112.
- 4 [9] Morris R, Tod D, Eubank, M. From youth team to first team: An investigation into the
- 5 transition experiences of young professional athletes in soccer. Int J Sport Exerc
 6 Psychol. 15(2017) 523-539.
- [10] Task Force. Heart rate variability: standards of measurement, physiological interpretation
 and clinical use. Task Force of the European Society of Cardiology and the North American
 Society of Pacing and Electrophysiology. Circulation 1996; 93:1043-1065.
- 10 [11] Peçanha T, Prodel E, Bartels R, Nasario-Junior O, Paula RB, Silva LP, Laterza MC, Lima
- JRP. 24-h cardiac autonomic profile after exercise in sedentary subjects. Int J Sports Med. 35
 (2014) 245-252.
- [12] Paludo AC, Peçanha T, Rabelo, FN, Bartels R, Fecchio RY, Simões AC, Nakamura FY.
 Reliability of heart rate variability in futsal players. Rev Bras Educ Fís Esp. 34 (2020) 673683.
- 16 [13] Buchheit M. Monitoring training status with HR measures: do all roads lead to17 Rome?. Front Physiol. 5 (2014) 73.
- 18 [14] Cox RH, Martens MP, Russell WD. Measuring anxiety in athletics: The revised
 19 competitive state anxiety inventory-2. J Sport Exercise Psy. 25 (2003)519-533.
- 20 [15] Fernandes MG, Nunes AS, Raposo JV, Fernandes HM, Brustad R. The CSAI-2: An
- 21 examination of the instrument's factorial validity and reliability of the intensity, direction and
- frequency dimensions with Brazilian athletes. J Appl Sport Psychol. 25 (2013) 377-391.
- 23 [16] O'Donoghue, P. (2012). Statistics for sport and exercise studies: An introduction.
 24 Routledge.

- [17] Newell, J., Aitchison, T., & Grant, S. (2010). Statistics for sport and exercise science: A
 practical approach. Prentice Hall.
- 3 [18] Blanca JM, Alar<u>cón</u> R, Arnau J, Bono R, Bendayan R. Non-normal data: is ANOVA still
- 4 a valid option? *Psicothema*. 29 (2017) 552-557.
- 5 [19] Terry PC, Walrond N, Carron AV. The influence of game location on athletes'
 6 psychological states. J. Sci. Med. Sport. 1(1998) 29-37.
- [20] Souza, R. A., Beltran, O. A., Zapata, D. M. et al. (2019). Heart rate variability, salivary
 cortisol and competitive state anxiety responses during pre-competition and pre-training
 moments. Bio Sport. 36 (2019) 39-46.
- 10 [21] Lerma C, Infante O, Pérez-Groyas H, José MV. Poincaré plot indexes of heart rate
- 11 variability capture dynamic adaptations after haemodialysis in chronic renal failure patients.
- 12 Clin Physiol Funct Imaging. 23(2003):72–80.
- 13 [22] Hoshi RA, Pastre CM, Vanderlei LC, Godoy MF. Poincaré plot indexes of heart rate
- variability: relationships with other nonlinear variables. Auton Neurosci. 177(2013) 271–274.
- 15 [23] Orellana JN, Torres BC, Cachadina ES, Hoyo M, Cobo AD. Two new indexes for the
- assessment of autonomic balance in elite soccer players. Int J Sports Physiol Perform. 10
- **17** (2015) 452-457.