

## Madagascar's extraordinary biodiversity: Threats and opportunities

Ralimanana, Hélène; Perrigo, Allison L; Smith, Rhian J; Borrell, James S; Faurby, Søren; Rajaonah, Mamy Tiana; Randriamboavonjy, Tianjanahary; Vorontsova, Maria S; Cooke, Robert S C; Phelps, Leanne N; Sayol, Ferran; Andela, Niels; Andermann, Tobias; Andriamanohera, Andotiana M; Andriambololonera, Sylvie; Bachman, Steven P; Bacon, Christine D; Baker, William J; Belluardo, Francesco; Birkinshaw, Chris; Cable, Stuart; Canales, Nataly A; Carrillo, Juan D; Clegg, Rosie; Clubbe, Colin; Crottini, Angelica; Damasco, Gabriel; Dhanda, Sonia; Edler, Daniel; Farooq, Harith; de Lima Ferreira, Paola; Fisher, Brian L; Forest, Félix; Gardiner, Lauren M; Goodman, Steven M; Grace, Olwen M; Guedes, Thaís B; Hackel, Jan; Henniges, Marie C; Hill, Rowena; Lehmann, Caroline E R; Lowry, Porter P; Marline, Lovanomenjanahary; Matos-Maraví, Pável; Moat, Justin; Neves, Beatriz; Nogueira, Matheus G C; Onstein, Renske E; Papadopoulos, Alexander S T; Perez-Escobar, Oscar A; Phillipson, Peter B; Pironon, Samuel; Przelomska, Natalia A S; Rabarimanarivo, Marina; Rabehevitra, David; Raharimampionona, Jeannie; Rajaonary, Fano; Rajaovelona, Landy R; Rakotoarinivo, Mijoro; Rakotoarisoa, Amédée A; Rakotoarisoa, Solofo E; Rakotomalala, Herizo N; Rakotonasolo, Franck; Ralaiveloarisoa, Berthe A; Ramirez-Herranz, Myriam; Randriamamonjy, Jean Emmanuel N; Randrianasolo, Vonona; Rasolohery, Andriambolantsoa; Ratsifandrihamanana, Anitry N; Ravololomanana, Noro; Razafiniary, Velosoa; Razanajatovo, Henintsoa; Razanatsoa, Estelle; Rivers, Malin; Silvestro, Daniele; Testo, Weston; Torres Jiménez, Maria F; Walker, Kim; Walker, Barnaby E; Wilkin, Paul; Williams, Jenny; Ziegler, Thomas; Zizka, Alexander; Antonelli, Alexandre

### Science

DOI:  
[10.1126/science.adf1466](https://doi.org/10.1126/science.adf1466)

Published: 02/12/2022

Peer reviewed version

[Cyswllt i'r cyhoeddiad / Link to publication](#)

*Dyfyniad o'r fersiwn a gyhoeddwyd / Citation for published version (APA):*

Ralimanana, H., Perrigo, A. L., Smith, R. J., Borrell, J. S., Faurby, S., Rajaonah, M. T., Randriamboavonjy, T., Vorontsova, M. S., Cooke, R. S. C., Phelps, L. N., Sayol, F., Andela, N., Andermann, T., Andriamanohera, A. M., Andriambololonera, S., Bachman, S. P., Bacon, C. D., Baker, W. J., Belluardo, F., ... Antonelli, A. (2022). Madagascar's extraordinary biodiversity: Threats and opportunities. *Science*, 378(6623), Article eadf1466. <https://doi.org/10.1126/science.adf1466>

## Madagascar's extraordinary biodiversity: Threats and opportunities

Papadopoulos, Alexander S. T.

### Science

DOI:

[10.1126/science.adf1466](https://doi.org/10.1126/science.adf1466)

Published: 02/12/2022

Peer reviewed version

[Cyswllt i'r cyhoeddiad / Link to publication](#)

*Dyfyniad o'r fersiwn a gyhoeddwyd / Citation for published version (APA):*

Papadopoulos, A. S. T. (2022). Madagascar's extraordinary biodiversity: Threats and opportunities. *Science*, 378(6623). <https://doi.org/10.1126/science.adf1466>

#### Hawliau Cyffredinol / General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

#### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

**Title:** Madagascar's extraordinary biodiversity: Threats and opportunities

**Authors:** Hélène Ralimanana<sup>1\*†</sup>, Allison L. Perrigo<sup>2,3†</sup>, Rhian J. Smith<sup>4,3†</sup>, James S. Borrell<sup>4</sup>, Søren Faurby<sup>2,3</sup>, Mamy Tiana Rajaonah<sup>1</sup>, Tianjanahary Randriamboavonjy<sup>1</sup>, Maria S. Vorontsova<sup>4</sup>, Robert S. C. Cooke<sup>5,2,3</sup>, Leanne N. Phelps<sup>6,7</sup>, Ferran Sayol<sup>8,3</sup>, Niels Andela<sup>9</sup>, Tobias Andermann<sup>10,11,2,3</sup>, Andotiana M. Andriamanohera<sup>1</sup>, Sylvie Andriambololonera<sup>12</sup>, Steven P. Bachman<sup>4</sup>, Christine D. Bacon<sup>2,3</sup>, William J. Baker<sup>4</sup>, Francesco Belluardo<sup>13,14,15</sup>, Chris Birkinshaw<sup>12,16</sup>, Stuart Cable<sup>4</sup>, Nataly A. Canales<sup>17</sup>, Juan D. Carrillo<sup>18,3,11,19</sup>, Rosie Clegg<sup>20,4</sup>, Colin Clubbe<sup>4</sup>, Angelica Crottini<sup>13,14,15</sup>, Gabriel Damasco<sup>21,2</sup>, Sonia Dhanda<sup>4</sup>, Daniel Edler<sup>22,2,3</sup>, Harith Farooq<sup>2,3,23</sup>, Paola de Lima Ferreira<sup>24,2,3</sup>, Brian L. Fisher<sup>25</sup>, Félix Forest<sup>4</sup>, Lauren M. Gardiner<sup>26</sup>, Steven M. Goodman<sup>27,28</sup>, Olwen M. Grace<sup>4</sup>, Thaís B. Guedes<sup>29</sup>, Jan Hackel<sup>4</sup>, Marie C. Henniges<sup>4,30</sup>, Rowena Hill<sup>4,30</sup>, Caroline E.R. Lehmann<sup>7,6</sup>, Porter P. Lowry II<sup>16,31</sup>, Lovanomenjanahary Marline<sup>1,3,27</sup>, Pável Matos-Maravi<sup>24,3</sup>, Justin Moat<sup>4</sup>, Beatriz Neves<sup>32,3</sup>, Matheus G. C. Nogueira<sup>32,33,3</sup>, Renske E. Onstein<sup>34,35</sup>, Alexander S. T. Papadopoulos<sup>36</sup>, Oscar A. Perez-Escobar<sup>4</sup>, Peter B. Phillipson<sup>16,31</sup>, Samuel Pironon<sup>4,37</sup>, Natalia A. S. Przelomska<sup>4,38</sup>, Marina Rabarimanarivo<sup>12</sup>, David Rabehevitra<sup>1</sup>, Jeannie Raharimampionona<sup>12</sup>, Fano Rajaonary<sup>12</sup>, Landy R. Rajaovelona<sup>1</sup>, Mijoro Rakotoarinivo<sup>39</sup>, Amédée A. Rakotoarisoa<sup>1</sup>, Solofo E. Rakotoarisoa<sup>1</sup>, Herizo N. Rakotomalala<sup>1</sup>, Franck Rakotonasolo<sup>1</sup>, Berthe A. Ralaiveloarisoa<sup>1</sup>, Myriam Ramirez-Herranz<sup>40,3,41</sup>, Jean Emmanuel N. Randriamamonjy<sup>1</sup>, Vonona Randrianasolo<sup>1</sup>, Andriambolantsoa Rasolohery<sup>42</sup>, Anitry N. Ratsifandrihamanana<sup>43</sup>, Noro Ravololomanana<sup>12</sup>, Veloso Razafiniary<sup>1</sup>, Henintsoa Razanajatovo<sup>1</sup>, Estelle Razanatsoa<sup>44</sup>, Malin Rivers<sup>45</sup>, Daniele Silvestro<sup>11,19,2,3</sup>, Weston

22 Testo<sup>28,2,3</sup>, Maria F. Torres Jiménez<sup>46,2,3</sup>, Kim Walker<sup>4,47</sup>, Barnaby E. Walker<sup>4</sup>, Paul Wilkin<sup>4</sup>, Jenny  
23 Williams<sup>4</sup>, Thomas Ziegler<sup>48,49</sup>, Alexander Zizka<sup>50</sup>, Alexandre Antonelli<sup>2,3,4,51\*†</sup>.

24

## 25 Affiliations:

26 <sup>1</sup> Royal Botanic Gardens, Kew, Kew Madagascar Conservation Centre, Antananarivo,  
27 Madagascar.

28 <sup>2</sup> Department of Biological and Environmental Sciences, University of Gothenburg, Gothenburg,  
29 Sweden.

30 <sup>3</sup> Gothenburg Global Biodiversity Centre, University of Gothenburg, Gothenburg, Sweden.

31 <sup>4</sup> Royal Botanic Gardens, Kew, Richmond, Surrey, UK.

32 <sup>5</sup> UK Centre for Ecology and Hydrology, Wallingford, UK.

33 <sup>6</sup> School of GeoSciences, University of Edinburgh, Edinburgh, UK.

34 <sup>7</sup> Royal Botanic Garden Edinburgh, Edinburgh, UK.

35 <sup>8</sup> Centre for Biodiversity and Environment Research, Department of Genetics, Evolution and  
36 Environment, University College London, London, UK.

37 <sup>9</sup> School of Earth and Environmental Sciences, Cardiff University, Cardiff, Wales, UK.

38 <sup>10</sup> Department of Organismal Biology, SciLifeLab, Uppsala University, Sweden.

39 <sup>11</sup> Department of Biology, University of Fribourg, Fribourg, Switzerland.

40 <sup>12</sup> Missouri Botanical Garden, Madagascar Program, Antananarivo, Madagascar.

41 <sup>13</sup> CIBIO, Centro de Investigação em Biodiversidade e Recursos Genéticos, InBIO Laboratório  
42 Associado, Universidade do Porto, Vairão, Portugal.

43 <sup>14</sup> Departamento de Biologia, Faculdade de Ciências, Universidade do Porto, Porto, Portugal

44 <sup>15</sup> BIOPOLIS Program in Genomics, Biodiversity and Land Planning, CIBIO, Campus de Vairão,  
45 Vairão, Portugal.

46 <sup>16</sup> Missouri Botanical Garden, St Louis, Missouri, USA.

47 <sup>17</sup> Natural History Museum of Denmark, University of Copenhagen, Copenhagen, Denmark.

48 <sup>18</sup> CR2P, Muséum National d'Histoire Naturelle, Paris, France.

49 <sup>19</sup> Swiss Institute of Bioinformatics, Fribourg, Switzerland.

50 <sup>20</sup> Department of Geography, University of Exeter, Exeter, Devon, UK.

51 <sup>21</sup> Departamento de Botânica e Zoologia, Universidade Federal do Rio Grande do Norte, Natal,  
52 Rio Grande do Norte, Brazil.

53 <sup>22</sup> Integrated Science Lab, Department of Physics, Umeå University, Umeå, Sweden.

54 <sup>23</sup> Faculty of Natural Sciences, Lúrio University, Pemba, Cabo Delgado Province, Mozambique.

55 <sup>24</sup> Biology Centre CAS, Institute of Entomology, České Budějovice, Czech Republic.

56 <sup>25</sup> California Academy of Sciences, San Francisco, California, USA.

57 <sup>26</sup> Cambridge University Herbarium, Department of Plant Sciences, University of Cambridge,  
58 Cambridge, UK.

59 <sup>27</sup> Association Vahatra, Antananarivo, Madagascar.

60 <sup>28</sup> Field Museum of Natural History, Chicago, Illinois, USA.

61 <sup>29</sup> Instituto de Biologia, Universidade Estadual de Campinas, Unicamp, Campinas, São Paulo,  
62 Brazil.

<sup>30</sup> School of Biological and Behavioural Sciences, Queen Mary University of London, London, UK.

<sup>31</sup> Institut de Systématique, Évolution, et Biodiversité (ISYEB), Muséum National d'Histoire Naturelle, Paris, France.

<sup>32</sup> Museu Nacional, Universidade Federal do Rio de Janeiro, Rio de Janeiro, Brazil.

<sup>33</sup> Universidade Federal do Rio de Janeiro, Rio de Janeiro, Brazil.

<sup>34</sup> Naturalis Biodiversity Center, Darwinweg 2, 2333CR Leiden, the Netherlands

<sup>35</sup> German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, Leipzig, Germany.

<sup>36</sup> School of Natural Sciences, Bangor University, Bangor, Gwynedd, Wales, UK.

<sup>37</sup> UN Environment Programme World Conservation Monitoring Center (UNEP-WCMC), Cambridge, UK.

<sup>38</sup> Department of Anthropology, Smithsonian National Museum of Natural History, Washington, D.C., USA.

<sup>39</sup> Department of Plant Biology and Ecology, University of Antananarivo, Antananarivo, Madagascar.

<sup>40</sup> Instituto de Ecología y Biodiversidad, University of La Serena, La Serena, Chile.

<sup>41</sup> Programa de Doctorado en Biología y Ecología Aplicada, Universidad Católica del Norte, Universidad de La Serena, La Serena, Chile.

<sup>42</sup> Ileiry Geospatial Services, Antananarivo, Madagascar.

<sup>43</sup> WWF, Antananarivo, Madagascar.

<sup>44</sup> Plant Conservation Unit, Department of Biological Sciences, University of Cape Town, South Africa.

<sup>45</sup> Botanic Gardens Conservation International, Kew, Richmond, Surrey, UK.

<sup>46</sup> Institute of Biosciences, Life Sciences Centre, Vilnius University, Lithuania.

<sup>47</sup> Royal Holloway, University of London, Egham, Surrey, UK.

<sup>48</sup> Cologne Zoo, Cologne, Germany.

<sup>49</sup> Institute of Zoology, University of Cologne, Cologne, Germany.

<sup>50</sup> Department of Biology, Philipps-University Marburg, Marburg, Germany.

<sup>51</sup> Department of Biology, University of Oxford, Oxford, UK.

\* Correspondence to: [H.Ralimanana@kew.org](mailto:H.Ralimanana@kew.org); [a.antonelli@kew.org](mailto:a.antonelli@kew.org)

† Authors contributed equally

## Abstract:

Madagascar's unique biota is heavily impacted by human activity and under intense threat. Here, we review the current state of knowledge on the conservation status of Madagascar's terrestrial and freshwater biodiversity by presenting data and analyses on documented and predicted species-level conservation status, the most prevalent and relevant threats, *ex situ* collections and programs, and the coverage and comprehensiveness of protected areas. The existing terrestrial protected area

network in Madagascar covers 10.4% of its land area and includes at least part of the range of the majority of described native species of vertebrates with known distributions (97.1% of freshwater fishes, amphibians, reptiles, birds and mammals combined) and plants (67.7%). The overall figures are higher for threatened species (97.7% of threatened vertebrates and 79.6% of threatened plants occurring within at least one protected area). IUCN Red List assessments and Bayesian neural network analyses for plants identify overexploitation of biological resources and unsustainable agriculture as the most prominent threats to biodiversity. We highlight five opportunities for action at multiple levels to ensure that conservation and ecological restoration objectives, programs and activities take account of complex underlying and interacting factors and produce tangible benefits for the biodiversity and people of Madagascar.

**One Sentence Summary:** Current knowledge on Madagascar's biodiversity and its decline indicates an urgent need for inclusive actions.

**Main text:**

Madagascar's biota, the result of millions of years of evolution in relative isolation, is both unique and under threat. At the same time as the scientific description of new species is accelerating (1), so is the overall rate of extinction (2), and many species may be disappearing before they are even documented. In this review, we aim to consolidate information on the conservation status of some of the main elements of Madagascar's biodiversity, evaluate the many and varied threats faced by species assessed under the criteria for the International Union for Conservation of Nature (IUCN) Red List of Threatened Species, and provide some perspectives on future opportunities to ensure the future of this hyperdiverse and unique biota.

**Threats to Madagascar's biodiversity**

Madagascar's biodiversity is in decline, with some groups more threatened than others (Fig. 1). In our review of threatened species, we follow the IUCN Red List data (3) and threat categories (4), unless otherwise specified. Threatened species are those listed as Critically Endangered (CR), Endangered (EN) or Vulnerable (VU). At one extreme, 22% (35 species) of assessed birds are threatened, while, at the other end of the scale, approximately 73% (66 species) of freshwater fishes and 75% (173 species) of magnoliid plants are threatened. Trees are particularly important in terms of their broad ecological functions and human uses, and 63% of the 3,118 assessed tree species in Madagascar are threatened (5). Humans have impacted the environment since arrival on Madagascar, not only in recent years. To avoid a shifting baseline effect, it is necessary to view changes in light of human settlement beginning hundreds or even thousands of years ago (1). For example, despite the relatively low proportion of bird species currently threatened with extinction,

Madagascar has already lost at least 14 species (7% of all species) that were present when humans first settled the island (Fig. 1). The rate of anthropogenic extinction is even higher in mammals, with 23 species (10%) extirpated since first human settlement. Vertebrate extinctions include the loss of lineages representing millions of years of evolution – e.g., the sloth-, koala- and monkey-lemurs (families Palaeopropithecidae, Megaladapidae, and Archaeolemuridae) and two species of hippopotamus (family Hippopotamidae). The extinction of four species of elephant birds (order Aepyornithiformes) represents the global loss of a functionally unique clade (6, 7). Extinctions, especially those of megafauna such as these, have broad scale implications for ecosystem functioning (6-8).

In total, 13 endemic animal species are listed as Extinct (EX), defined as extinctions after 1500 AD, and an additional 33 are listed as Extinct Prehistorically [EP], defined as anthropogenic extinctions prior to 1500 AD (see (9) for a full list of documented anthropogenic extinctions before 1500 AD). A further nine have been categorized as Critically Endangered (Possibly Extinct) – CR(PE). For plants, no species has been assessed as Extinct, and only one species (*Aloe silicicola*) is categorized as Extinct in the Wild (EW). A further 118 plant species are listed by IUCN as CR(PE) (111 spp.) or Critically Endangered (Possibly Extinct in the Wild) – CR(PEW) (7 spp.). Of those currently listed as CR(PE), five species are present in *ex situ* living collections, and their status should therefore be updated to CR(PEW) (3, 10).

Malagasy species feature prominently among animal groups that have been considered by the EDGE of Existence program (11-13), which ranks species according to their evolutionary distinctiveness and the level of threat they face (EDGE = Evolutionary Distinct and Globally Endangered). Almost one in five species of amphibians (18 spp.), reptiles (17 spp.), and mammals



(17 spp.) in the top 100 EDGE species of each group are found in Madagascar (13). Yet only one in 20 (4 spp.) of the top 100 EDGE species of birds are found on the island.

Given the narrow geographic range of many Malagasy species (e.g., (14)), numerous undetected anthropogenic extinctions are likely to have taken place (15), such as CR *Aloe* species, which may have become extinct in the wild since they were last recorded. This may be especially pronounced in groups with high levels of micro-endemism, for example freshwater fishes and amphibians (16). Ascertaining extinction events is difficult due to sampling biases, insufficient taxonomic knowledge regarding the morphological features of extant species, and the challenges of comparisons with fossil and subfossil remnants in certain groups, such as frogs (e.g., (17)).

#### *Reliability of species conservation assessments*

Conservation assessments rely on taxonomic classification, and different opinions on species limits and numbers may influence the proportion of threatened species (e.g., (18)). This proportion may also be biased by an over-assessment of well-known and widespread taxa, or, alternatively, range-restricted species that are more likely to be threatened. To investigate indications of bias, we calculated the fraction of threatened species across different plant groups based on two sets of species: taxa with full threat-status assessments in the Red List compiled by the IUCN and their partners (19); and those estimated with a Bayesian neural network approach (Fig. 1; (9, 20)), which inferred the threat status for all remaining species. Using this method, we predicted the threat status of 8,821 species with an estimated test accuracy of >65%. All taxa with a full threat-status assessment were included, although some assessments may be out of date and could underestimate threat levels.

The neural network approach combined with current IUCN assessments revealed a similar fraction of species inferred to be threatened across most taxonomic groups (Fig. 1). Large deviations from the proportion of threatened species in the current IUCN assessments occur in the ferns and lycophytes, and to a lesser extent the magnoliids. The neural network results combined with the known IUCN categories predicted a far higher proportion of threatened ferns and lycophytes (146 of 306 spp; 47.7% [95%CI: 38.5-56.7%]) than reflected in published IUCN assessments (1 of 33 spp; 3.0%), suggesting a bias towards assessing more common species. In the magnoliids, the combined results predict a lower proportion of threatened species (211 of 294 spp; 71.8% [95%CI: 68.0-75.9%]) compared to published IUCN assessments alone (173 of 225 spp; 76.9%), suggesting a bias towards assessing rare species in that group.

### *Genetic erosion*

The reduction of genetic diversity within species resulting from the extirpation of subpopulations is a crucial, yet easily overlooked, facet of biodiversity loss that is often a precursor to extinction. Genetic erosion has negative effects on the individual fitness, the health of populations, and a species' ability to adapt to changing environments, reducing their resilience to further change, and potentially incurring extinction debt (21, 22). In practice, genetic factors are not directly incorporated into IUCN assessments, which are based on measures of the probability of extinction due to population declines, restricted geographic ranges, and small population sizes (23).

The reduction in population sizes of wild plants and animals, together with their fragmentation and isolation, is generally expected to increase inbreeding and genetic load, reducing genetic diversity

and fitness over time (22, 24). The few studies of intraspecific diversity in Malagasy species to date reveal that some species have maintained high genetic diversity in spite of habitat fragmentation (e.g., (25, 26)), whereas others have relatively low diversity, possibly as a result of anthropogenic effects (e.g., (25, 27-29)). Results differ even within species, such as in the palm *Beccariophoenix madagascariensis*, in which only some populations show strong signals of inbreeding, reflected by an excess of homozygotes (30). It is important to note that under some circumstances, population decline may outstrip the speed with which genetic diversity is eroded due to inbreeding. Estimates of heterozygosity may therefore not indicate the true genetic health and long-term prospects of populations when considered in isolation (31, 32).

A more powerful, although less explored, approach is to use coalescence-based demographic modeling, which uses genome-wide data to estimate the longer-term trends in population size, providing more information than metrics of contemporary genetic diversity alone (25, 33). In *Cheirogaleus* dwarf lemurs, genomic analysis suggests that four species have experienced population size declines in the last 50,000 years, with one decline (*C. cf. medius*) starting as long as 300,000 years ago – all clearly in pre-human times and resulting in lower genetic diversity (29). In contrast, another genomic study shows that five out of ten analyzed plant species with varying extinction risk have experienced substantial population declines since human colonization of Madagascar (25). In the golden-crowned sifaka (*Propithecus tattersalli*) (26), mouse lemurs (*Microcebus* spp.) (28), *Mantella* frogs (34), and the Milne-Edwards' sportive lemur (*Lepilemur edwardsi*) (35) demographic declines also appear to have taken place after the arrival of humans on the island (although the inherent uncertainties of mutation rates in the microsatellite data used makes the timing of these declines less certain).

The risks of inbreeding and increased genetic load may represent substantial and likely underestimated longer-term threats to the survival of Malagasy species. This is especially relevant considering the high level of fragmentation of native habitats in some vegetation types, such as the humid forests, and is worthy of further investigation.

### **Predicting future extinction: direct drivers of loss**

Identifying direct threats is part of the IUCN Red List Assessment process, and even species that are not explicitly threatened (i.e., those that are Least Concern [LC], Near Threatened [NT], or Data Deficient [DD]) can still have threats listed. Here we discuss these threats and how they apply to all species. Our analysis of IUCN assessments indicates that overexploitation and agriculture are the most frequently listed threats to Malagasy fauna (excluding invertebrates) and flora (Fig. 2), mirroring global findings (36). Overexploitation is unsustainable biological resource use as defined by the IUCN (37), including hunting and collecting for subsistence use or national/international trade. Overexploitation is linked in some cases to illegal harvesting – for example, the illegal logging of rosewood for trade (*Dalbergia* spp.) – which is banned under the Convention on International Trade in Endangered Species of Wild Fauna and Flora since 2013 and under Malagasy law since 2010.

We estimated that 62.1% of vertebrates and 87.1% of plants are threatened by overexploitation and that 56.8% of vertebrates and 87.8% of plants are threatened by agriculture. These two major threats, almost equal in magnitude (Fig. 2), have different modes of impact – overexploitation is more targeted and tends to occur over relatively restricted areas compared to the broad effects of land clearance for agriculture.

Agriculture, and to a lesser extent overexploitation, are also the primary causes of deforestation in Madagascar. Approximately 44% of the land area covered by native forest in 1953 was deforested by 2014 (38). The rate of deforestation has steadily increased, reaching 99.0 kha/yr between 2010 and 2014 (38), and according to Global Forest Watch remains very high at 72.9 kha/yr (2014–2020) (39). Deforestation in Madagascar reflects global patterns (40) and is primarily driven by the small-scale but widespread practice of swidden agriculture (also known as shifting cultivation; in Madagascar referred to as *tavy* for rice cultivation in humid and subhumid areas, and *hatsake* for cassava and maize in dry and subarid areas). Additionally, cash crop production, particularly maize and peanut, has become a major driver of deforestation (41), alongside the production of products for international markets, such as forest-derived vanilla (42). The most frequent threats listed for plants and vertebrates suggest that this trend of increasing deforestation rates will continue, with forest loss and degradation a consequence of clearance of land for agriculture, potentially associated with small-scale fire activity (43) and overexploitation through selective logging and highly targeted activities such as the collection of palm hearts. Additionally, natural system modifications (threats from actions that convert or degrade habitat, e.g., anthropogenic fire in forests or changes in water management; Fig. 2), adds to deforestation and threatens 23.2% of vertebrates and is estimated to threaten 68.9% of plants. Some predictions indicate that in the absence of an effective strategy against deforestation, 38–93% of forest present in 2000 will be no longer present in 2050 (41).

For vertebrates, the greatest threat after overexploitation and agriculture is ‘invasive and problematic species and emerging infectious diseases’ (referred to as invasives/diseases in Fig. 2), which impacts 27% of all species (360 spp.; Fig. 2). This category includes non-native invasive species, as well as problematic native species and diseases of any origin. Changes in habitat due

to the spread of non-native plant species can have a large effect, and one study reports that of a total of 546 naturalized non-native plants in Madagascar, 101 have been found to display invasive characteristics (44). Many non-native plants, such as the Mexican yellow pine (*Pinus patula*) in terrestrial systems (45), and common water hyacinth (*Pontederia crassipes*) in freshwater systems (46), are aggressively invasive and transformative in semi-natural habitats, and are clearly impacting native fauna and flora. Even within reserves and protected areas, the issue can be pronounced. For example, three species of invasive/problematic plants – strawberry guava (*Psidium cattleianum*), Molucca raspberry (*Rubus moluccanus*), and wild cardamom (*Aframomum angustifolium*) – together occupy 17.6% of the Betampona Nature Reserve (47) and are also widespread in Ranomafana National Park and other protected areas.

Not all impacts are negative, however, and there is some evidence to suggest that, due to their potential for faster growth, some non-native plants are better able to combat the rapid fragmentation of native vegetation, and may be beneficial for endemic vertebrates, providing refuge, food, and vegetation corridors, while also improving human livelihoods (48). The potential for such species to become invasive or readily burn must however be fully considered before embarking on any planting initiatives (49). In addition, effects must be considered at different scales. For examples, the presence of strawberry guava has been reported to locally increase species richness in frugivores, but as they are primary dispersers of the seed this further contributes to the spread and to associated changes in floral and faunal community structure and reduction in taxonomic richness (50).


Non-native vertebrates have also had marked and diverse impacts, which we also here illustrate with some examples. Introduced rats (*Rattus rattus*; present since at least the 14th century) are now ubiquitous, even in remote areas, and there is evidence that their presence is associated with

declines in native small mammals (51). In freshwater habitats, competition and predation by exotic fish species is considered a major factor in the decline of native freshwater fish (52), which have been completely replaced by non-native species across much of the Central Highlands and western areas (53). While not yet listed in current assessments, the recent invasion of the toxic Asian common toad (*Duttaphrynus melanostictus*), along with the predicted vulnerability of most native vertebrates to its toxins (54), is expected to represent a new threat to many nocturnal carnivores. The effects of other introduced and naturalized animals on native biodiversity are not well studied; this includes widely occurring species such as dogs (*Canis familiaris*), cats (*Felis catus*), the common myna (*Acridotheres tristis*), and the marbled crayfish (*Procambarus virginalis*). The threat of emerging infectious diseases is primarily driven by the occurrence of the chytrid fungus *Batrachochytrium dendrobatidis*, widely documented across Madagascar over the last decade and a potential threat to all amphibians, although no mass mortalities associated with chytridiomycosis have been reported in the country (55). Species often face multiple threats at the same time, although the impact of each threat can vary between species (Fig. 2).

Among vertebrates, amphibians have the highest number of IUCN-identified threats per species (Fig. 2A), with a mean of 4.8 threats per species, followed by mammals (mean 2.5 threats/species), and reptiles (mean 2.2 threats/species). For plants (Fig. 2B), magnoliids have the most threats per species (mean 2.9 threats/species), followed by rosids (mean 2.8 threats/species), and other eudicots (mean 2.8 threats/species). Although there might be some variation in the perception and documentation of threats between the specialists carrying out assessments, all follow the same protocols (4).

The number and relative impact of these threats may change in coming decades. The impact of climate change on Malagasy biodiversity remains understudied and it is not currently indicated in


IUCN assessments as a major threat. However, this impact is expected to increase in the future (56-59), and could potentially result in synergistic negative effects with unsustainable agriculture associated with land clearance, invasive alien species, and inappropriate management of fire regimes that can increase future fire risk (43, 56, 57, 60). Extinctions in one group could also have effects on others that depend on them, such as in cases of strong plant–animal mutualisms (61, 62). Although coextinction is hard to quantify, with substantial knowledge and data gaps (63), models suggest that the effects of extinction can be amplified as a result of the interactions between species within and between trophic levels, with the potential to lead to secondary and even cascading extinctions (64, 65).





**Fig. 1. Madagascar's threatened and lost biodiversity.** IUCN Red List assessment categories of major groups of plants and animals from Madagascar. Assessment categories and coloration follow the standards used by the IUCN Red List. Category distributions for animal groups include ray-finned fishes, (Actinopterygii, freshwater species only, N=91), mammals (Mammalia, N=231 species), amphibians (Amphibia, N=296), mollusks (Mollusca, N=67), reptiles (Reptilia, N=340), arthropods (Arthropoda, N=374), and birds (Aves, N=209). Category distributions for plants, indicated with saturated, wider bars, include magnoliids (N=225), gymnosperms (N=6), rosids (N=1,704), monocots (N=822), asterids (N=1,105 species), other eudicots (N=81), and ferns & lycophytes (N=33). Thinner, unsaturated bars indicate the relative proportion of plant taxa in each threat category for IUCN Red List assessments combined with the taxa where the threat category was predicted in a Bayesian neural network analysis: asterids (N=2,924 species), rosids (N=2,990), other eudicots (N=312), magnoliids (N=294), monocots (N=1,965), and ferns & lycophytes (N=306). The number indicated above each bar with “+” is the number of taxa for which the threat category was predicted using the neural network analysis. IUCN Red List Assessment categories include Least Concern (LC) and Near Threatened (NT), together making up the “not threatened” category; while Vulnerable (VU); Endangered (EN); Critically Endangered (CR); Critically Endangered, Possibly Extinct (CR(PE)); Extinct in the Wild (EW); Extinct (EX; i.e., extinct after 1500 CE), and Extinct Prehistorically (EP; sensu (66), i.e., extinct before 1500 CE but with dated records within the last 130,000 years) make up the group “threatened and extinct.” Silhouettes below the bars depict taxonomic orders with EP, EX, EW, and CR(PE) species, with the number of species in each category per order. For some plant groups, additional orders with single CR(PE) species are indicated with a star. Depicted orders are, from left to right and top to bottom: Perciformes, Cyprinodontiformes, Cetartiodactyla, Carnivora, Rodentia, Primates, Afrosoricida, Venerida, Unionoida, Perciformes,

Cyprinodontiformes, Squamata, Testudines, Crocodilia, Orthoptera, Spirobolida, Araneae,  
Calanoida, Cyclopoida, Podicipediformes, Cuculiformes, Coraciiformes, Charadriiformes,  
Gruiformes, Anseriformes, Aepyornithiformes, Accipitriformes, Laurales, Magnoliales, Pinales,  
Oxalidales, Sapindales, Myrtales, Malvales, Malpighiales, Fabales, Asparagales, Poales,  
Ericales, Boraginales, Gentianales, Asterales, Saxifragales.



**Fig. 2. Threats to Malagasy biodiversity.** Alluvial plots showing threats, as defined by the IUCN, and their associations with major groups of terrestrial and freshwater (A) vertebrates (1,332 species with IUCN assessments, of which 993 species have at least one listed threat) and (B) plants (9,268 species with IUCN assessments or predictions, all of which have at least one listed threat; includes gymnosperms [6 species], which could not be visualized). Widths of the boxes/lines reflect the number of species impacted by each threat. Threats for vertebrates are further divided into sub-threats, whereas only the highest threat classification was available for assessed plants. The estimates for plants include predictions for unassessed species based on a Bayesian neural network analysis (9). The color scheme is consistent across panels. The “Other” threat class includes Pollution, Climate change, Transportation, and Human disturbance, plus Invasives/diseases for plants. Some threat classes have been renamed for brevity/clarity, including the IUCN category “biological resource use”, which is labeled “overexploitation” here and in the text, for brevity and in line with IPBES terminology (36).

## **Conservation efforts and effectiveness**

### *Protected Areas*

Protected areas (PAs) are the central political and scientific accomplishment of Madagascar's conservation strategy. The network has been continuously developed since the first PA was established in 1927 (67-71). Our data compilation shows that the network now encompasses 10.4% of the land area of Madagascar, having grown by more than a third over the last two decades (Fig. 3). This recent and extensive designation of new PAs was carried out via a multi-stakeholder consultative process, in combination with data and literature analyses, through the Durban Vision

initiative conceived in 2003. In addition to preserving diverse ecosystems and landscapes, the focus has been on species groups for which sufficient diversity and distribution data were available, primarily vertebrates (including birds, mammals, amphibians, and reptiles), and some plant groups. Despite the production of considerable new data since the Durban Vision began (e.g., many newly described species; (1), the network designed during that process remains highly taxonomically comprehensive. From a global perspective, the PA network also excels at capturing the vast majority of Madagascar's many EDGE species: 14 out of 18 amphibians, 15 out of 17 reptiles, 16 out of 17 mammals, and all four birds (13).

As of November 2020, there were 110 terrestrial PAs with permanent protected status in Madagascar, covering 61,300 km<sup>2</sup> across the country (Fig. 3) (70, 72, 73). Eleven of these are “orphan PAs” – sites abandoned by their former managers with responsibility reverting to the Ministry of Environment and Sustainable Development (70). An additional 89 sites (15,200 km<sup>2</sup>), predominantly comprising Key Biodiversity Areas (KBAs), are not under formal protection (70, 72, 74, 75).

The long-term security and effective management of Madagascar's PAs is therefore crucial to addressing the country's biodiversity challenges. Providing evidence of their effectiveness and co-benefits, such as ecosystem service provision, will be critical to securing ongoing support and management from local communities, as well as from local and national governments. However, measuring PA effectiveness is challenging (e.g., at avoiding deforestation, or providing alternative livelihoods) while accounting for numerous covariates (76), particularly in Madagascar with comparatively little long-term biodiversity monitoring data (77). Recent counterfactual analyses (78) have sought to address this question by identifying protected and non-protected sites that are similar across multiple social and environmental variables, and then comparing indicators of

conservation effectiveness, such as deforestation rate. These analyses indicate that PAs have a small, but significant, effect at reducing deforestation (9).

We show that since 1990, human impacts have measurably increased across all terrestrial PAs (Table S8 (9)), a trend documented worldwide (76). Human activity by local communities inside PAs is not necessarily detrimental to biodiversity, and land use and conservation are therefore not mutually exclusive. Nevertheless, land conversion and unsustainable exploitation remain major drivers of biodiversity loss. This suggests that protecting and realizing the potential of Madagascar's comprehensive PA network will require the application of rigorous monitoring and evaluation strategies, matched with extensive community collaboration, to understand co-benefits and minimize detrimental human impacts.

Scores for deforestation and management effectiveness – for example, from the self-reported Management Effectiveness Tracking Tool (79) – have been the main metrics used to monitor effectiveness to date. However, these are not always reliable indicators of management effectiveness (77). New and expanded capacity of variables such as remote-sensed fire and stable night lights, with increased temporal resolution, offer promising new monitoring opportunities. How fire is associated with land transformation in Madagascar has been discussed in the literature but only recently quantitatively assessed (43), demonstrating that tree loss anomalies are highest in environments where landscapes-scale fire (>21 ha) does not occur, and where the role of small-scale fires (<21 ha) requires close and urgent investigation. We show that trends in anthropogenic fire are variable, increasing in some areas of forest vegetation in the north, east, and west but decreasing in grassland-woodland mosaic vegetation across central Madagascar (Fig. 4A, B). Forest loss also reflects this pattern, primarily occurring in the humid forest biome in the east, but also in dry forest and spiny forest in the west (Fig. 4C, D). Deforestation and land use conversion

remain key challenges to conservation in Madagascar, and improved remote-sensing will accelerate monitoring and developing understanding on the effectiveness of PAs and other conservation measures.

#### Ex situ *conservation and restoration*

Living plant collections in botanic gardens and seed banks represent invaluable sources of taxonomic and genetic diversity for immediate conservation and research, and should continue to support restoration efforts. Globally, 29.6% of all known native Malagasy plant species (23.1% of endemic species and 23.1% of native threatened species) are held in botanic gardens, with 15.5% held in Madagascar (10), where their cultivation is sometimes linked to educational programs and community engagement, essential to raising awareness of biodiversity and conservation issues. The Millennium Seed Bank Partnership in Madagascar, initiated in 1996, hosts collections of an estimated 3,500 native Malagasy species, including members of four of the five endemic plant families and all seven of the iconic baobab species (*Adansonia* spp.). The single Malagasy plant species listed as Extinct in the Wild, *Aloe silicicola*, now only survives in one living collection outside Madagascar.

For native terrestrial and freshwater vertebrates, 9% of amphibians, 17% of mammals, 20% of reptiles, 21% of freshwater fishes, and 33% of birds are currently held in zoological collections (18% overall) (9, 80). Many are part of active breeding programs: a subset of these (3% of amphibians, 7% of reptiles, 11% of freshwater fishes, 13% of mammals, and 23% of birds) were successfully bred during 2020 (9). Unsurprisingly, the species held in captive breeding facilities are biased towards the more charismatic, well-known taxa (81). For example, among amphibians,

13 of the 34 species in zoos belong to the genus *Mantella*, a group of strikingly colored diurnal frogs, even though *Mantella* contains only 4% of Madagascar's amphibian fauna. Freshwater fishes, amphibians, and reptiles are highly suitable for targeted *ex situ* breeding and reintroduction programs (82-85). For species in these groups and others with high levels of micro-endemism, such conservation programs continue to represent a major safeguard against extinction (86). This complies with the One Plan Approach to species conservation proposed by the IUCN SSC Conservation Planning Specialist Group, which supports the development of conservation and management plans for all populations of a species, even outside of their natural range (87). It should be noted that the success of reintroduction relies also on the maintenance of natural habitat and functional diversity at potential reintroduction sites, along with minimizing risks associated with invasive species and infectious diseases. In addition, particularly for mammals, vulnerability of captive-bred populations to predation can also jeopardize the success of reintroductions (88).

*Progress towards international conservation commitments*

Madagascar continues to make progress towards Convention on Biological Diversity targets, but like most countries falls short of meeting them in full (89). Of particular relevance here is that Madagascar did not formally meet Aichi Target 11 to protect at least 17% of its total land area (Fig. 3) – as was the case for 48% of the parties reporting their progress (89). If areas designated as important for biodiversity but not currently under formal protection were also given protection, the total percentage of PA coverage would rise from the current 10.4% to 13% (Fig. 3B). However, given that even the existing network is widely considered to be chronically under-resourced, this action is not a priority for the near future (90, 91).

Target 4 of the Global Strategy for Plant Conservation seeks to protect 15% of each vegetation type. This has been achieved for mangrove (currently at 29.4%), spiny forest (21.5%), humid forest (18.5%), and tapia (17.9%), but not for dry forest (13.3%), subhumid forest (5.7%), and grassland-woodland mosaic (1.8%) (Table S6 (9)). However, expansion of the areas of those vegetation types under protection may not be feasible due to limited financial resources, the large degree of fragmentation and geographical spread of habitats, and the long administrative process involved in extending PAs or designating additional areas, as well as a lack of political will. It also may not be desirable until it can be demonstrated that the existing PAs are well-resourced, achieving conservation objectives and providing benefits to communities. Restoration within currently protected areas may provide a longer-term pathway to meeting this goal, particularly where there are rapidly realizable socio-economic benefits such as sustainable silk production from wild native silkworms (*Borocera cajani*) associated with tapia (*Uapaca bojeri*) in the Itremo Massif PA and Ambatofinandrahana KBA. Other targets are more difficult to assess due to lack of data. For example, there is very little evidence to assess success in the control of invasive alien species, with some exceptions such as the ongoing but promising house crow (*Corvus splendens*) eradication (92).


Although most of the Aichi and GSPC targets were either not achieved or cannot be assessed, a marked success is that Madagascar has comfortably achieved GSPC Target 7 (at least 75% of known threatened plant species conserved *in situ*), with our analyses indicating this percentage is currently at 80%.

*Realizing benefits of biodiversity for people*



492 The majority of Madagascar's over 28 million inhabitants live outside of, but often very close to,  
493 PAs (93) (Figs. 3A; S1). These communities face challenges connected to widespread poverty,  
494 which itself is related to degradation of natural capital in the landscape, limited access to formal  
495 education and health care, crime, corruption, weak governance, and regulatory issues including  
496 land tenure (15, 94, 95). For example, southern Madagascar is severely affected by food and water  
497 insecurity, which catalyzes political and social instability, exacerbates economic insecurity, and  
498 has led to large-scale migration within the country (96). This instability likewise hampers the  
499 operations of local, national, and international conservation organizations, which could be  
500 compounded further by adverse effects from climate change (59). As the human population in the  
501 country is expected to reach 42–105 million by the end of this century, of which half will be under  
502 15 years of age, and with the majority under the poverty threshold (97), the conservation success  
503 of PAs will be inextricably linked to the effective provision of livelihoods, food security, and  
504 natural capital – a situation echoed across all Malagasy ecosystems and the world over (98).

505



506

**Fig. 3. Madagascar’s terrestrial protected areas (PAs) in the context of human population**

507

**density and changes in coverage of vegetation type over time. (A) PAs with IUCN protected**

508

status (99), “orphan” status, or no formal protection status (e.g., unprotected Key Biodiversity

509

Areas [KBAs]), shown in the context of nearby marine PAs, surrounding bathymetry (100), coral

510

reefs (101), cities, roads, and population density (102). (B) The evolution of PA coverage over

511


time, showing the potential increase in area protected that could be gained if the designated areas

512

(those identified as important for biodiversity but not currently under formal protection, mostly

513

KBAs) were protected in the future (74, 75).



**Fig 4. Recent changes and patterns in burned area and tree cover in Madagascar.** (A) Average burned area in the period 2003–2019. (B) Statistically significant trends in burned area (MODIS) (103) from 2006–2016, not explained by precipitation change (TRMM) (104), dates chosen for comparison with Goodman et al. (72). Red indicates an increasing trend; blue indicates a decreasing trend. (C) Change in tree cover from 2000–2012 (105). (D) Vegetation map, inferred

and simplified from Moat & Smith (106). The legend indicates the percentage of each vegetation category currently covered by the protected area network.

### **Looking back, moving forward**

Despite decades of research and applied conservation programs supported through substantial financial investments (95, 107), Madagascar's remarkable biodiversity continues to face severe challenges (Figs. 1, 2). It is reasonable to ask whether more of the same – even if better resourced and underpinned with greater scientific understanding and technology – is likely to deliver a tangible reversal in Madagascar's trajectory of biodiversity loss, or whether new approaches are required to bring transformative change (108), including greater emphasis on monitoring interventions and addressing underlying drivers through key leverage points. The responsibility for averting humanitarian and biodiversity crises is a shared global challenge (36, 109), with solutions needed at all societal levels – including via local communities, engagement of the private sector, sound leadership and policy from regional and national government, steady international support for conservation, and increased recognition of how historic and ongoing global and national inequalities have contributed to the current situation. Scientific data and evidence will continue to make a vital contribution, but it is crucial that this is done in an interdisciplinary context, with open communication channels to relevant government departments and third sector organizations.

*Decades of progress in biodiversity science and conservation*

We now have a clearer and more detailed understanding than ever before of the past and present diversity and distribution of Madagascar's biodiversity, and the threats it faces (*1*) (Fig. 1). The underlying data are the product of decades of research – with an increasing number of Malagasy biologists involved. This body of research and the evidence we have collated and presented here makes a clear case for Madagascar as one of the world's foremost conservation priorities.

Despite multiple competing demands on land, the Malagasy government, in collaboration with a broad group of conservation organizations and donors, has succeeded in designating 10.4% of the country as terrestrial PAs in a network that is largely representative of Madagascar's diverse biomes (Fig. 3, 4). Most terrestrial and freshwater vertebrate species with known distributions have ranges that overlap with least one PA (94.7% of reptiles, 97.2% of amphibians, 98.1% of mammals, 98.9% of freshwater fishes, 100% of birds, and 97.1% for all groups combined), as do the majority of plants, but to a lesser extent (67.7%) (*9*). For threatened species with known distributions, the percentages are similar for vertebrates (94.3% of reptiles, 99.3% of amphibians, 97.7% of mammals, 100% of freshwater fishes, 100% of birds, and 97.7% for all groups combined) and markedly higher for plants (79.6%). Nonetheless, there are still many threatened species with ranges that do not overlap with existing PA network, including one amphibian, three mammals, seven reptiles, and 559 plants (*9*), and many more that have not yet been assessed but may be threatened. The ranges of all birds overlapped with at least one PA; this was also true when we filtered the analysis to only include resident and breeding areas (*9*).

Since the loss of Madagascar's terrestrial megafauna (here defined as vertebrates above 10 kg), there have been few documented modern extinctions, but many species have perilously reduced population sizes. The continued increase in new species descriptions suggests there may be undocumented extinctions, especially in poorly studied taxa (*1*). Despite this, with limited

resources and/or capacity, Madagascar has made important progress towards achieving international climate, biodiversity, and sustainable development goals, providing a foundation on which to build in the coming decades.

Success stories for individual species highlight how positive collaborative efforts can avert extinction. Examples include work on the Madagascar pochard (*Aythya innotata*) (110), which shows a 30% probability that extinction was prevented due to conservation action, the success story of the community-based protection of the tahina palm or dimaka (*Tahina spectabilis*) where local communities were involved in propagation and population reinforcement (111), and the work to prevent the extinction of the ploughshare tortoise (*Astrochelys yniphora*) through a captive breeding program (112).

Other notable successes have come from Madagascar's "biodiversity conservation boom", which started in the 1980s, including a growth in the number of students pursuing university-level education in environmental sciences, biodiversity conservation and management, and related fields, at both public and private universities. The result is an increasingly robust national capacity for the conservation and management of biodiversity that extends to international conservation organizations, which have been able to actively recruit Malagasy professionals to the highest administrative and executive positions. Going beyond this, the gap in scientific leadership that underpins conservation evidence is being incrementally filled by Malagasy biodiversity scientists. Researchers from outside Madagascar are increasingly collaborating with Malagasy researchers for mutual benefit. The requirement for international collaborators to provide financial and technical support for Malagasy researchers and their research infrastructure via collaboration protocols, set out in the national strategy for scientific research in Madagascar (113), reinforces the importance of this.

As in many low-income countries, insufficient public funding means that the number of Malagasy professionals is still insufficient to serve the country's needs, there are relatively few PhD positions available to students, and those that are trained at higher levels often move away from academia and into the private sector. Access to up-to-date biodiversity data has also been a limiting factor (15). A further challenge is how to successfully engage multiple parts of society in conservation. Efforts that are genuinely socially integrated have been shown to produce more effective and resilient practices, policies, and decision-making, especially in the face of unstable environmental, political, and health situations (114). The Madagascar Fauna and Flora Group, the Lemur Conservation Foundation, Durrell Wildlife Conservation Trust, The Peregrine Fund Madagascar, Madagascar Biodiversity Center, and Madagasikara Voakajy, as well as the work of the Royal Botanic Gardens, Kew, and Missouri Botanical Garden, are all examples of successful collaborations involving researchers, conservation partners and local communities to protect biodiversity and empower local people.

*The future of biodiversity in Madagascar*

Meeting the Convention on Biological Diversity's Post-2020 Global Biodiversity Framework 2030 targets and milestones and achieving the 2050 goals (115) will be challenging – in Madagascar and globally. Evaluating successes and failures over previous decades and learning from these to prioritize effective conservation investment will be particularly important. To embrace diverse views and promote inclusivity in the identification of future directions, we discussed our results and current literature among our co-authors and consulted with Malagasy and external researchers, conservation leaders, and politicians, to arrive at five main opportunities for the future, which we now present.

1) Investment in conservation and restoration must be based on evidence, effectiveness, and future challenges. Since the 1980s, billions of US dollars from international donors and conservation organizations, in cooperation with the Malagasy government, have been dedicated to protecting the country's biodiversity and creating today's network of PAs (107, 116). However, the effectiveness of many interventions is poorly understood because impact evaluations are absent or lacking rigor. Evaluating the effectiveness of conservation activities is challenging, but the subject of increasingly sophisticated research efforts (76, 78, 117). Nevertheless, it is imperative investments reinforce evidence-based and regularly evaluated interventions, requiring greater collaboration and co-design between local communities, regional and national authorities, researchers, the private sector, and other stakeholders. A particular opportunity is to frame these evaluations around community-based conservation interventions that address challenges faced by people and nature in unison. For example, nature-based solutions (118) for diversified, locally adapted and sustainable agriculture can help address livelihood needs, while more efficient stoves can substantially decrease the demand on charcoal from native forests for cooking and heating, and further may reduce the health hazards of smoke inhalation. Such initiatives increase food and energy security (119) while providing resilience to climate stochasticity (120). Similarly, coordinated, community-based fire management and awareness raising can be used to help mitigate risk to fire-sensitive forests. On-site management is especially important for fire mitigation, as a study during the COVID 19 pandemic has shown (121). Fire management also presents the opportunity to mitigate the impact of exotic species by targeting the removal of flammable invasives (e.g., *Pinus*), and guide appropriate tree-planting initiatives to avoid fire-prone plantations near areas of particular biological importance. Such measures can improve the quality of grazing land for livestock, while reducing carbon emissions from fire and helping to protect biodiverse habitats.



633

634 2) Expanded biodiversity monitoring is key to safeguarding Madagascar's most valuable natural  
635 assets. Existing biodiversity data are sufficient to characterize major conservation challenges and  
636 robustly support the orientation of conservation efforts in Madagascar. Calling for the collection  
637 of additional data risks delivering diminished returns on investment for conservation planning  
638 (122). Nevertheless, from collating the information for this review, we acknowledge a clear need  
639 to address gaps in understudied ecosystems, taxa, and genetically distinct populations, noting that  
640 many newly described species are already threatened (123) and in need of immediate protection.  
641 Monitoring is also crucial for the detection of new non-native and potentially invasive species, as  
642 well as providing important data for the management of those that have already taken hold.  
643 Increasing connections with international trading partners without concurrent improvements in  
644 capacity for biosecurity increases Madagascar's vulnerability to such species (124), and strategies  
645 to monitor and mitigate these risks while delivering near-term benefits are needed.

646 Although there are initiatives that provide broad overviews of conservation effectiveness (e.g.  
647 (117)), many conservation interventions lack impact evaluations, in part due to a lack of robust,  
648 long-term monitoring data for biodiversity and social outcomes. The major gap is a lack of capacity  
649 for robust biodiversity monitoring. An example of the increasing value of data and coherency in  
650 conservation efforts is the development of the Madagascar Protected Areas website (125), which  
651 consolidates much of the information about Madagascar's extensive network of PAs. But as with  
652 many initiatives, the key is in long-term financing and maintenance of these portals and ensuring  
653 that data flows freely and openly to similar, global initiatives like Protected Planet (73).

654 Biological monitoring needs to be based on consistent, repeatable methodologies, with shared data.  
655 This information provides the science-based evidence needed to leverage international funding

and government policy support. Monitoring is one area where new technologies will play a key role, such as through the increasing availability of near real-time satellite images and small and cost-effective unmanned aerial vehicles, which can increase visual access to remote areas (126). Similarly, DNA-based biodiversity surveys, including environmental sampling, can greatly improve the speed of site-inventories and identification of unknown and understudied taxa. Advances in monitoring must be delivered with improved and centralized management. This should include open-source and transdisciplinary data on biodiversity, social and conservation governance and performance. These data should be in formats that are accessible and useful to practitioners, to identify relevant baselines, and support evidence-based decisions for conservation and restoration.

3) Improving the effectiveness of existing PAs is more important than creating new ones. Madagascar has an extensive, evidence-based, and highly representative network of terrestrial PAs (Fig. 3, 4). Madagascar's existing PAs already include at least partial ranges of a substantial proportion of Malagasy taxa, including most Malagasy EDGE species. Focusing on improving their quality and effectiveness will likely lead to positive biodiversity outcomes (127), further increasing the already measurable impact that PAs have had on biodiversity. By strengthening PAs, biodiversity can be conserved across ecosystem, species, and genetic levels, all of which are integral in long-term conservation, as discussed above. Investment in restoration of degraded areas within and beyond the existing network (see Opportunity 4 below) will provide multiple benefits for biodiversity and people. This could help increase the resilience of habitats to future drivers of biodiversity loss including climate change, while increasing potential ranges of many species in parallel. Demonstrating the benefits of strengthened PAs to people is a likely prerequisite for societal support to maintain and improve upon the existing network, while mitigating risk of future

downgrading, downsizing, or degazettement (legal removal of conservation status) (128). Financial benefits that come with strengthened PAs must be distributed appropriately and equitably within the country's political and social contexts, with the full inclusion of local communities at all stages (127, 129).

4) Conservation and restoration should not focus solely on the PA network. Madagascar's PAs are islands of natural capital in a landscape of degraded natural resources (130) and therefore provide vital resources for communities living adjacent to them. Traditional "fortress conservation" – seeking to protect areas by limiting access – is therefore both undesirable and unlikely to be effective. To further reduce the detrimental human impacts that exist in all PAs (107) (Table S8 (9)), we argue for strategies to enhance the natural capital of the surrounding landscapes, to reduce pressure on PAs as providers of basic resources, and to increase buffer zones for the species that live in and around them. This could include increasing ecosystem provision, such as productive soils, food, fibers, and other materials and services such as water flow regulation and carbon capture. Such measures would serve to address some of the largest threats to species, including the expansion of agriculture and overexploitation (Fig. 2).

In particular, ecological restoration could benefit people and biodiversity, particularly when targeted to the 89.6% of the country that is not protected. It offers potential to provide new livelihood opportunities that are far from, and independent of, the resources within PAs, further reducing pressure on the system (131). Importantly, restoration should not only target those ecosystems that traditionally receive the most conservation attention because they hold the greatest biodiversity, for example forests. Other vegetation types such as grasslands, where most agriculture takes place, are equally vital. Restoration should be carried out following best practice

and in places where people will benefit most, not necessarily only adjacent to PAs. Further, restoration should include maximizing biodiversity recovery to meet multiple goals, using resilient species, and working together with local communities (49, 132).

For the species and their inherent genetic diversity not covered by the PA network, particularly those that are challenging to conserve, such as freshwater fishes and palms, *ex situ* conservation in zoological and botanical gardens is a vital tool to support conservation and restoration. For plants, efforts should especially focus on the 32.3% of plant species that fall outside of the PA network, and the species that have cultural or economic value for people (e.g., crop wild relatives). Promoting biobanking for animals and intensifying it for seeds, spores, and fungi will not only support conservation but also contribute material and knowledge to restoration and research (88).

5) Conservation actions must address the root causes of biodiversity loss. Our analysis showed that the most frequently listed threats to Madagascar's biodiversity come from overexploitation and agriculture, predominantly a result of forest loss and potentially tied to increases in small-scale anthropogenic fire in forests (Fig. 4A, B; see also (43)), significantly affecting humid forest areas in the east and dry forest and spiny forest in the west (Fig. 4C, D). This trend is likely to continue unless the root causes of this forest loss are addressed. Conservationists and their funders must recognize that food, social security, health, and well-being are the utmost priorities for rural communities, and that PAs will always be vulnerable when surrounded by impoverished people living in landscapes with eroded natural capital (133). Politicians and economists must recognize that sustainable and equitable development in Madagascar is inextricably linked to, and dependent on, the maintenance of ecosystem function and the goods and services they provide. Initiatives that address these issues by working with local communities to identify tailored solutions in health,

education, and green entrepreneurship are increasingly successful and should be expanded, but generally lack data and evidence from monitoring (see Opportunity 2). Promising approaches include voluntary savings and loans; inclusive, sustainable agricultural development schemes that promote stable land ownership and build – rather than destroy – natural capital and the ecosystem services it provides; implementation of conservation interventions, including research and monitoring; and PA management that maximizes local employment (107, 132). Such efforts will facilitate improved livelihoods for many, while reducing pressure on the PAs themselves, bringing tangible benefits to communities, and contributing to sustainable management (107, 134).

## Conclusions

The alarming status of Madagascar's biodiversity is the result of multifaceted, unsustainable practices including historic and contemporary exploitation. In the eyes of much of the world, Madagascar's biodiversity is a unique global asset that needs "saving"; in the daily lives of many of the Malagasy people, it is a rapidly diminishing source of the most basic needs for subsistence. Achieving a sustainable future that benefits people and biodiversity is possible by building on, and expanding, integrated, inclusive conservation efforts. Biodiversity is the greatest opportunity and most valuable asset for Madagascar's future development.

## References and notes

1. A. Antonelli, R. J. Smith, A. L. Perrigo, A. Crottini, J. Hackel, W. Testo, H. Farooq, M. F. T. Jiménez, N. Andela, T. Andermann, A. M. Andriamanohera, S. Andriambololonera, S. P. Bachman, C. D. Bacon, W. J. Baker, F. Belluardo, C. Birkinshaw, J. S. Borrell, S. Cable, N. A. Canales, J. D. Carrillo, R. Clegg, C. Clubbe, R. S. C. Cooke, G. Damasco, S. Dhanda, D. Edler, S. Faurby, P. d. L. Ferreira, B. L. Fisher, F. Forest, L. M. Gardiner,

- 749 S. M. Goodman, O. M. Grace, T. B. Guedes, M. C. Henniges, R. Hill, C. E. R. Lehmann,  
750 P. P. L. II, L. Marline, P. Matos-Maraví, J. Moat, B. Neves, M. G. C. Nogueira, R. E.  
751 Onstein, A. S. T. Papadopoulos, O. A. Perez-Escobar, L. N. Phelps, P. B. Phillipson, S.  
752 Pironon, N. A. S. Przelomska, M. Rabarimanarivo, D. Rabehevitra, J.  
753 Raharimampionona, M. T. Rajaonah, F. Rajaonary, L. R. Rajaovelona, M. Rakotoarinivo,  
754 A. A. Rakotoarisoa, S. E. Rakotoarisoa, H. N. Rakotomalala, F. Rakotonasolo, B. A.  
755 Ralaiveloarisoa, M. Ramirez-Herranz, J. E. N. Randriamamonjy, T. Randriamboavonjy,  
756 V. Randrianasolo, A. Rasolohery, A. N. Ratsifandrihamanana, N. Ravololomanana, V.  
757 Razafiniary, H. Razanajatovo, E. Razanatsoa, M. Rivers, F. Sayol, D. Silvestro, M. S.  
758 Vorontsova, K. Walker, B. E. Walker, P. Wilkin, J. Williams, T. Ziegler, A. Zizka, H.  
759 Ralimanana, Madagascar's extraordinary biodiversity: Evolution, distribution, and use.  
760 [first manuscript in our pair of reviews - Manuscript number: abf0869] (2022).
- 761 2. T. Andermann, S. Faurby, S. T. Turvey, A. Antonelli, D. Silvestro, The past and future  
762 human impact on mammalian diversity. *Sci. Adv.* **6**, eabb2313 (2020).
- 763 3. IUCN. The IUCN Red List of Threatened Species. Version 2021-1. (2021);  
764 <https://www.iucnredlist.org>.
- 765 4. IUCN, *IUCN Red List categories and criteria, version 3.1, second edition*. (Gland,  
766 Switzerland; Cambridge, UK, 2012).
- 767 5. E. Beech, M. Rivers, Rabarimanarivo, M., Ravololomanana, N., Manjato, N.,  
768 Lantoarisoa, F., Andriambololoner, S., Ramandimbisoa, B., Ralimanana, H.,  
769 Rakotoarisoa, S., Razanajatovo, H., Razafiniary, V., Andriamanohera, A.,  
770 Randrianasolo, V., Rakotonasolo, F., Rakotoarisoa, A., Randriamamonjy, N.,  
771 Rajaovelona, L., Rakotomalala, N., Randriamboavonjy, T., Rajaonah, M., Rabehevitra,  
772 D., Ramarosandrata, A.V., Rakotoarinivo, M., B.H. Ravaomanalina and Jeannoda, V.,  
773 *The Red List of Trees of Madagascar*. (Botanic Gardens Conservation International,  
774 Richmond, UK, 2021).
- 775 6. S. M. Goodman, W. L. Jungers, *Les Animaux et Ecosystemes de l'Holocene Disparus de*  
776 *Madagascar*. (Association Vahatra, Antananarivo, 2013).
- 777 7. F. Sayol, M. J. Steinbauer, T. M. Blackburn, A. Antonelli, S. Faurby, Anthropogenic  
778 extinctions conceal widespread evolution of flightlessness in birds. *Sci. Adv.* **6**, eabb6095  
779 (2020).
- 780 8. Y. Malhi, C. E. Doughty, M. Galetti, F. A. Smith, J.-C. Svenning, J. W. Terborgh,  
781 Megafauna and ecosystem function from the Pleistocene to the Anthropocene. *Proc. Natl.*  
782 *Acad. Sci. USA* **113**, 838-846 (2016).
- 783 9. Materials, methods, and supplementary text are available as supplementary materials.
- 784 10. Botanic Gardens Conservation International. PlantSearch online database. (2021);  
785 [https://tools.bgci.org/plant\\_search.php](https://tools.bgci.org/plant_search.php).
- 786 11. Zoological Society of London. EDGE of Existence. [www.edgeofexistence.org](http://www.edgeofexistence.org).
- 787 12. N. J. B. Isaac, S. T. Turvey, B. Collen, C. Waterman, J. E. M. Baillie, Mammals on the  
788 EDGE: Conservation Priorities Based on Threat and Phylogeny. *PLOS ONE* **2**, e296  
789 (2007).
- 790 13. Zoological Society of London. EDGE of Existence. (2021);  
791 <https://www.edgeofexistence.org/explore-edge-species-country/>.
- 792 14. A. Rakotoarison, M. D. Scherz, F. Glaw, J. Köhler, F. Andreone, M. Franzen, J. H. Glos,  
793 O. T. Jono, A. Mori, S. H. Ndriantsoa, N. R. Raminosoa, J. C. Riemann, M.-O. Rödel, G.  
794 M. Rosa, D. R. Vieites, A. Crottini, M. Vences, Describing the smaller majority:

- Integrative taxonomy reveals twenty-six new species of tiny microhylid frogs (genus *Stumpffia*) from Madagascar. *Vertebrate Zoology* **67**, 271–398 (2017).
15. M. S. Vorontsova, P. P. Lowry II, S. R. Andriambololona, L. Wilmé, A. Rasolohery, R. Govaerts, S. Z. Ficinski, A. M. Humphreys, Inequality in plant diversity knowledge and unrecorded plant extinctions: An example from the grasses of Madagascar. *PLANTS, PEOPLE, PLANET* **3**, 45-60 (2021).
16. J. P. Benstead, P. H. De Rham, J.-L. Gattolliat, F.-M. Gibon, P. V. Loiselle, M. Sartori, J. S. Sparks, M. L. J. Stiassny, Conserving Madagascar's Freshwater Biodiversity. *Bioscience* **53**, 1101-1111 (2003).
17. R. D. E. MacPhee, D. A. Burney, N. A. Wells, Early Holocene Chronology and environment of Ampasambazimb, a Malagasy subfossil lemur site. *Int. J. Primatol.* **6**, 463-489 (1985).
18. S. Faurby, W. L. Eiserhardt, J.-C. Svenning, Strong effects of variation in taxonomic opinion on diversification analyses. *Methods Ecol. Evol.* **7**, 4-13 (2016).
19. IUCN. The IUCN Red List of Threatened Species. Version 2020-3. (2020); <https://www.iucnredlist.org>.
20. A. Zizka, T. Andermann, D. Silvestro, IUCNN - deep learning approaches to approximate species' extinction risk. *Divers. Distrib.*, DOI: <https://doi.org/10.1111/ddi.13450> (2021).
21. P. de Villemereuil, A. Rutschmann, K. D. Lee, J. G. Ewen, P. Brekke, A. W. Santure, Little adaptive potential in a threatened passerine bird. *Curr. Biol.* **29**, 889-894.e883 (2019).
22. R. Lande, Genetics and demography in biological conservation. *Science* **241**, 1455-1460 (1988).
23. M. C. Rivers, N. A. Brummitt, E. Nic Lughadha, T. R. Meagher, Do species conservation assessments capture genetic diversity? *Glob. Ecol. Conserv.* **2**, 81-87 (2014).
24. B. Charlesworth, D. Charlesworth, The genetic basis of inbreeding depression. *Genet. Res.* **74**, 329-340 (1999).
25. A. J. Helmstetter, S. Cable, F. Rakotonasolo, R. Rabarijaona, M. Rakotoarinivo, W. L. Eiserhardt, W. J. Baker, A. S. T. Papadopoulos, The demographic history of Madagascan micro-endemics: Have rare species always been rare? *Proc. R. Soc. Lond. B: Biol. Sci.*, 20210957 (2021).
26. E. Quéméré, X. Amelot, J. Pierson, B. Crouau-Roy, L. Chikhi, Genetic data suggest a natural prehuman origin of open habitats in northern Madagascar and question the deforestation narrative in this region. *Proc. Natl. Acad. Sci. USA* **109**, 13028-13033 (2012).
27. L. M. Gardiner, M. Rakotoarinivo, L. R. Rajaovelona, C. Clubbe, Population genetics data help to guide the conservation of palm species with small population sizes and fragmented habitats in Madagascar. *PeerJ* **5**, e3248 (2017).
28. G. L. Olivieri, V. Sousa, L. Chikhi, U. Radespiel, From genetic diversity and structure to conservation: Genetic signature of recent population declines in three mouse lemur species (*Microcebus* spp.). *Biol. Conserv.* **141**, 1257-1271 (2008).
29. R. C. Williams, M. B. Blanco, J. W. Poelstra, K. E. Hunnicutt, A. A. Comeault, A. D. Yoder, Conservation genomic analysis reveals ancient introgression and declining levels of genetic diversity in Madagascar's hibernating dwarf lemurs. *Heredity* **124**, 236-251 (2020).

30. A. Shapcott, M. Rakotoarinivo, R. J. Smith, G. Lysakova, M. F. Fay, J. Dransfield, Can we bring Madagascar's critically endangered palms back from the brink? Genetics, ecology and conservation of the critically endangered palm *Beccariophoenix madagascariensis*. *Bot. J. Linn. Soc.* **154**, 589-608 (2007).
31. P. A. Hagl, R. Gargiulo, M. F. Fay, C. Solofondranohatra, J. Salmona, U. Suescun, N. Rakotomalala, C. E. R. Lehmann, G. Besnard, A. S. T. Papadopoulos, M. S. Vorontsova, Geographical structure of genetic diversity in *Loudetia simplex* (Poaceae) in Madagascar and South Africa. *Bot. J. Linn. Soc.* **196**, 81-99 (2020).
32. T. van der Valk, D. Díez-del-Molino, T. Marques-Bonet, K. Guschanski, L. Dalén, Historical genomes reveal the genomic consequences of recent population decline in eastern gorillas. *Curr. Biol.* **29**, 165-170.e166 (2019).
33. C. R. Peart, S. Tusso, S. D. Pophaly, F. Botero-Castro, C.-C. Wu, D. Auriolles-Gamboa, A. B. Baird, J. W. Bickham, J. Forcada, F. Galimberti, N. J. Gemmell, J. I. Hoffman, K. M. Kovacs, M. Kunasranta, C. Lydersen, T. Nyman, L. R. de Oliveira, A. J. Orr, S. Sanvito, M. Valtonen, A. B. A. Shafer, J. B. W. Wolf, Determinants of genetic variation across eco-evolutionary scales in pinnipeds. *Nat. Ecol. Evol.* **4**, 1095-1104 (2020).
34. A. Crottini, P. Orozco-terWengel, F. C. E. Rabemananjara, J. S. Hauswaldt, M. Vences, Mitochondrial introgression, color pattern variation, and severe demographic bottlenecks in three species of malagasy poison frogs, genus *Mantella*. *Genes* **10**, 317 (2019).
35. M. Craul, L. Chikhi, V. Sousa, G. L. Olivieri, A. Rabesandratana, E. Zimmermann, U. Radespiel, Influence of forest fragmentation on an endangered large-bodied lemur in northwestern Madagascar. *Biol. Conserv.* **142**, 2862-2871 (2009).
36. IPBES, "Summary for policymakers of the global assessment report on biodiversity and ecosystem services of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services," (Bonn, Germany, 2019).
37. IUCN. Threats Classification Scheme (Version 3.2). (2022); <https://www.iucnredlist.org/resources/threat-classification-scheme>.
38. G. Vieilledent, C. Grinand, F. A. Rakotomalala, R. Ranaivosoa, J.-R. Rakotoarijaona, T. F. Allnutt, F. Achard, Combining global tree cover loss data with historical national forest cover maps to look at six decades of deforestation and forest fragmentation in Madagascar. *Biol. Conserv.* **222**, 189-197 (2018).
39. GFW. Global Forest Watch - Madagascar. (2021); <https://gfw.global/3uplmN5>.
40. S. L. Maxwell, R. A. Fuller, T. M. Brooks, J. E. M. Watson, Biodiversity: The ravages of guns, nets and bulldozers. *Nature* **536**, 143-145 (2016).
41. G. Vieilledent, M. Nourtier, C. Grinand, M. Pedrono, A. Clausen, T. Rabetrano, J.-R. Rakotoarijaona, B. Rakotoarivelo, F. A. Rakotomalala, L. Rakotomalala, A. Razafimpahanana, J. M. Ralison, F. Achard, It's not just poverty: unregulated global market and bad governance explain unceasing deforestation in Western Madagascar. *bioRxiv*, 2020.2007.2030.229104 (2020).
42. D. A. Martin, F. Andrianisaina, T. R. Fulgence, K. Osen, A. A. N. A. Rakotomalala, E. Raveloaritiana, M. R. Soazafy, A. Wurz, R. Andriafanomezantsoa, H. Andriamaniraka, A. Andrianarimisa, J. Barkmann, S. Dröge, I. Grass, N. Guerrero-Ramirez, H. Hänke, D. Hölscher, B. Rakouth, H. L. T. Ranarijaona, R. Randriamanantena, F. M. Ratsoavina, L. H. R. Ravaomanarivo, D. Schwab, T. Tschardtke, D. C. Zemp, H. Kreft, Land-use trajectories for sustainable land system transformations: Identifying leverage points in a global biodiversity hotspot. *Proc. Natl. Acad. Sci. USA* **119**, e2107747119 (2022).



- 887 43. L. N. Phelps, N. Andela, M. Gravey, D. S. Davis, C. A. Kull, K. Douglass, C. E. R.  
888 Lehmann, Madagascar's fire regimes challenge global assumptions about landscape  
889 degradation. *Global Change Biol.* **00**, 1-17 (2022).
- 890 44. C. A. Kull, J. Tassin, S. Moreau, H. Rakoto Ramiarantsoa, C. Blanc-Pamard, S. M.  
891 Carrière, The introduced flora of Madagascar. *Biol. Invasions* **14**, 875-888 (2012).
- 892 45. R. Baohanta, J. Thioulouse, H. Ramanankierana, Y. Prin, R. Rasolomampianina, E.  
893 Baudoin, N. Rakotoarimanga, A. Galiana, H. Randriambanona, M. Lebrun, R.  
894 Duponnois, Restoring native forest ecosystems after exotic tree plantation in Madagascar:  
895 combination of the local ectotrophic species *Leptolena bojeriana* and *Uapaca bojeri*  
896 mitigates the negative influence of the exotic species *Eucalyptus camaldulensis* and  
897 *Pinus patula*. *Biol. Invasions* **14**, 2407-2421 (2012).
- 898 46. A. Lehavana, Distribution, ecological and economic impacts and competition of the  
899 invasive alien aquatic weeds (*Pontederia crassipes* Mart., *Pistia stratiotes* L., *Salvinia*  
900 *molesta* D.S. Mitch. and *Azolla filiculoides* Lam.) in Madagascar, Doctoral thesis,  
901 Rhodes University, South Africa (2020).
- 902 47. A. Ghulam, I. Porton, K. Freeman, Detecting subcanopy invasive plant species in tropical  
903 rainforest by integrating optical and microwave (InSAR/PolInSAR) remote sensing data,  
904 and a decision tree algorithm. *ISPRS Journal of Photogrammetry and Remote Sensing* **88**,  
905 174-192 (2014).
- 906 48. A. Gérard, J. U. Ganzhorn, C. A. Kull, S. M. Carrière, Possible roles of introduced plants  
907 for native vertebrate conservation: the case of Madagascar. *Restor. Ecol.* **23**, 768-775  
908 (2015).
- 909 49. A. Di Sacco, K. A. Hardwick, D. Blakesley, P. H. S. Brancalion, E. Breman, L. Cecilio  
910 Rebola, S. Chomba, K. Dixon, S. Elliott, G. Ruyonga, K. Shaw, P. Smith, R. J. Smith, A.  
911 Antonelli, Ten golden rules for reforestation to optimize carbon sequestration,  
912 biodiversity recovery and livelihood benefits. *Global Change Biol.* **27**, 1328-1348 (2021).
- 913 50. C. M. M. DeSisto, D. S. Park, C. C. Davis, V. Ramananjato, J. L. Tonos, O. H.  
914 Razafindratsima, An invasive species spread by threatened diurnal lemurs impacts  
915 rainforest structure in Madagascar. *Biol. Invasions* **22**, 2845-2858 (2020).
- 916 51. S. M. Goodman, Soarimalala, Introduction to mammals, in *New Natural History of*  
917 *Madagascar*, S. M. Goodman, Ed. (Princeton University Press, Princeton, 2022), pp.  
918 1737-1769.
- 919 52. J. P. Benstead, M. L. J. Stiassny, P. N. Loiselle, K. J. Riseng, N. Raminosoa, River  
920 conservation in Madagascar, in *Global Perspectives on River Conservation: Science,*  
921 *Policy, and Practice*, P. J. Boon, B. R. Davies, G. E. Petts, Eds. (Wiley, Chichester,  
922 2000), pp. 205-231.
- 923 53. P. N. Reinthal, M. L. J. Stiassny, The freshwater fishes of Madagascar: A study of an  
924 endangered fauna with recommendations for a conservation strategy. *Conserv. Biol.* **5**,  
925 231-243 (1991).
- 926 54. B. M. Marshall, N. R. Casewell, M. Vences, F. Glaw, F. Andreone, A. Rakotoarison, G.  
927 Zancolli, F. Woog, W. Wüster, Widespread vulnerability of Malagasy predators to the  
928 toxins of an introduced toad. *Curr. Biol.* **28**, R654-R655 (2018).
- 929 55. M. C. Bletz, G. M. Rosa, F. Andreone, E. A. Courtois, D. S. Schmeller, N. H. C.  
930 Rabibisoa, F. C. E. Rabemananjara, L. Raharivololoniaina, M. Vences, C. Weldon, D.  
931 Edmonds, C. J. Raxworthy, R. N. Harris, M. C. Fisher, A. Crottini, Widespread presence  
932 of the pathogenic fungus *Batrachochytrium dendrobatidis* in wild amphibian  
933 communities in Madagascar. *Sci. Rep.* **5**, 8633 (2015).

56. J. L. Brown, A. D. Yoder, Shifting ranges and conservation challenges for lemurs in the face of climate change. *Ecol. Evol.* **5**, 1131-1142 (2015).
57. T. Spencer, A. S. Laughton, N. C. Flemming, J. C. Ingram, T. P. Dawson, Climate change impacts and vegetation response on the island of Madagascar. *Philos. Trans. Roy. Soc. A: Math. Phys. Eng. Sci.* **363**, 55-59 (2005).
58. J.-N. Wan, N. J. Mbari, S.-W. Wang, B. Liu, B. N. Mwangi, J. R. E. Rasoarahona, H.-P. Xin, Y.-D. Zhou, Q.-F. Wang, Modeling impacts of climate change on the potential distribution of six endemic baobab species in Madagascar. *Plant Divers.* **43**, 117-124 (2021).
59. S. R. Weiskopf, J. A. Cushing, T. L. Morelli, B. J. E. Myers, Climate change risks and adaptation options for Madagascar. *Ecol. Soc.* **26**, 36 (2021).
60. J. Busch, R. Dave, L. Hannah, A. Cameron, A. Rasolohery, P. Roehrdanz, G. Schatz, Climate change and the cost of conserving species in Madagascar. *Conserv. Biol.* **26**, 408-419 (2012).
61. G. Chomicki, M. Weber, A. Antonelli, J. Bascompte, E. T. Kiers, The impact of mutualisms on species richness. *Trends Ecol. Evol.* **34**, 698-711 (2019).
62. L. P. Koh, R. R. Dunn, N. S. Sodhi, R. K. Colwell, H. C. Proctor, V. S. Smith, Species coextinctions and the biodiversity crisis. *Science* **305**, 1632-1634 (2004).
63. M. L. Moir, P. A. Vesk, K. E. Brennan, D. A. Keith, L. Hughes, M. A. McCarthy, Current constraints and future directions in estimating coextinction. *Conserv. Biol.* **24**, 682-690 (2010).
64. R. K. Colwell, R. R. Dunn, N. C. Harris, Coextinction and persistence of dependent species in a changing world. *Annu. Rev. Ecol., Evol. Syst.* **43**, 183-203 (2012).
65. D. M. Hansen, M. Galetti, The forgotten megafauna. *Science* **324**, 42-43 (2009).
66. S. Faurby, M. Davis, R. Ø. Pedersen, S. D. Schowanek, A. Antonelli, J.-C. Svenning, PHYLACINE 1.2: The Phylogenetic Atlas of Mammal Macroecology. *Ecology* **99**, 2626-2626 (2018).
67. C. A. Kull, The roots, persistence, and character of Madagascar's conservation boom, in *Conservation and Environmental Management in Madagascar*, I. R. Scales, Ed. (Routledge, 2014), pp. 146-171.
68. C. J. Gardner, M. E. Nicoll, C. Birkinshaw, A. Harris, R. E. Lewis, D. Rakotomalala, A. N. Ratsifandrihamanana, The rapid expansion of Madagascar's protected area system. *Biol. Conserv.* **220**, 29-36 (2018).
69. J. P. G. Jones, O. S. Rakotonarivo, J. H. Razafimanahaka, Forest conservation on Madagascar: past, present, and future, in *The New Natural History of Madagascar*, S. M. Goodman, Ed. (Princeton University Press, Princeton, 2022), pp. 2130-2140.
70. S. M. Goodman, H. M. Rakotondratsimba, J. C. Ranivo Rakotoson, Protected areas, in *The New Natural History of Madagascar*, S. M. Goodman, Ed. (Princeton University Press, Princeton, 2022), pp. 2091-2107.
71. M. Virah-Sawmy, Gardner, C.J. & Ratsifandrihamanana, A.N., The Durban Vision in practice: experiences in the participatory governance of Madagascar's new protected areas, in *Conservation and Environmental Management in Madagascar*, I. R. Scales, Ed. (Routledge, 2014).
72. S. M. Goodman, M. J. Raherilalao, S. Wohlhauser, *The Terrestrial Protected Areas of Madagascar: Their History, Description, and Biota*. (Association Vahatra, Antananarivo, 2018).

73. Protected Planet. The World Database on Protected Areas. (2020);  
<https://www.protectedplanet.net/en/thematic-areas/wdpa?tab=WDPA>.
74. Key Biodiversity Areas Partnership. World Database of Key Biodiversity Areas. (2021);  
<http://www.keybiodiversityareas.org/kba-data/request>.
75. P. Kullberg, E. Di Minin, A. Moilanen, Using key biodiversity areas to guide effective expansion of the global protected area network. *Glob. Ecol. Conserv.* **20**, e00768 (2019).
76. J. Geldmann, A. Manica, N. D. Burgess, L. Coad, A. Balmford, A global-level assessment of the effectiveness of protected areas at resisting anthropogenic pressures. *Proc. Natl. Acad. Sci. USA* **116**, 23209-23215 (2019).
77. J. Eklund, L. Coad, J. Geldmann, M. Cabeza, What constitutes a useful measure of protected area effectiveness? A case study of management inputs and protected area impacts in Madagascar. *Conservation Science and Practice* **1**, e107 (2019).
78. J. Eklund, F. G. Blanchet, J. Nyman, R. Rocha, T. Virtanen, M. Cabeza, Contrasting spatial and temporal trends of protected area effectiveness in mitigating deforestation in Madagascar. *Biol. Conserv.* **203**, 290-297 (2016).
79. Protected Planet. Management effectiveness (PAME). (2022);  
<https://www.protectedplanet.net/en/thematic-areas/protected-areas-management-effectiveness-pame?tab=METT>.
80. Species 360. Zoological Information Management Software. (2021);  
<https://zims.species360.org>.
81. A. Miralles, M. Raymond, G. Lecointre, Empathy and compassion toward other species decrease with evolutionary divergence time. *Sci. Rep.* **9**, 19555 (2019).
82. J. R. Ali, M. Huber, Mammalian biodiversity on Madagascar controlled by ocean currents. *Nature* **463**, 653-656 (2010).
83. R. A. Griffiths, L. Pavajeau, Captive breeding, reintroduction, and the conservation of amphibians. *Conserv. Biol.* **22**, 852-861 (2008).
84. T. Ziegler, Frank-Klein, N., Ommer, S., Hürche, R. Loiselle, P.V. & Vences, M, Keeping and breeding of threatened endemic Malagasy freshwater fishes at Cologne Zoo (Germany): a contribution towards the advancement of a conservation breeding network. *Der Zoologische Garten N.F.* **88**, 123-155 (2020).
85. L. Leiss, A. Rauhaus, A. Rakotoarison, C. Fusari, M. Vences, T. Ziegler, Review of threatened Malagasy freshwater fishes in zoos and aquaria: The necessity of an ex situ conservation network—A call for action. *Zoo Biol.*, 10.1002/zoo.21661 (2021).
86. P. V. Loiselle, Captive breeding for the freshwater fishes of Madagascar, in *The Natural History of Madagascar*, S. M. Goodman, Benstead, J. P., Ed. (University of Chicago Press, Chicago, 2003), pp. 1569-1574.
87. K. Traylor-Holzer, K. Leus, O. Byers, Integrating ex situ management options as part of a One Plan Approach to species conservation, in *The Ark and Beyond: The Evolution of Zoo and Aquarium Conservation.*, B. A. Minter, J. Maienschein, J. P. Collins, Eds. (University of Chicago Press, Chicago, 2018), pp. 129-141.
88. A. Britt, B. R. Iambana, C. R. Welch, A. S. Katz, Project Betampona: Re-stocking of *Varecia variegata variegata* into the Betampona Reserve, in *The Natural History of Madagascar*, S. M. Goodman, J. Benstead, Eds. (University of Chicago Press, Chicago, USA, 2001).
89. Secretariat of the Convention on Biological Diversity, "Global Biodiversity Outlook 5," (Montreal, 2020); [www.cbd.int/GB05](http://www.cbd.int/GB05).

- 1026 90. M. D. Barnes, L. Glew, C. Wyborn, I. D. Craigie, Prevent perverse outcomes from global  
1027 protected area policy. *Nat. Ecol. Evol.* **2**, 759-762 (2018).
- 1028 91. R. L. Pressey, P. Visconti, M. C. McKinnon, G. G. Gurney, M. D. Barnes, L. Glew, M.  
1029 Maron, The mismeasure of conservation. *Trends Ecol. Evol.* **36**, 808-821 (2021).
- 1030 92. Critical Ecosystem Partnership Fund. Indian House Crow Eradication and Invasive  
1031 Species Surveillance in Madagascar - Madagascar Fauna and Flora Group. (2021);  
1032 [https://www.cepf.net/grants/grantee-projects/indian-house-crow-eradication-and-](https://www.cepf.net/grants/grantee-projects/indian-house-crow-eradication-and-invasive-species-surveillance-madagascar)  
1033 [invasive-species-surveillance-madagascar](https://www.cepf.net/grants/grantee-projects/indian-house-crow-eradication-and-invasive-species-surveillance-madagascar).
- 1034 93. INSTAT. instat Madagascar. (2022); <https://www.instat.mg/>.
- 1035 94. J. P. G. Jones, J. Ratsimbazafy, A. N. Ratsifandrihamanana, J. E. M. Watson, H. T.  
1036 Andrianandrasana, M. Cabeza, J. E. Cinner, S. M. Goodman, F. Hawkins, R. A.  
1037 Mittermeier, A. L. Rabearisoa, O. S. Rakotonarivo, J. H. Razafimanahaka, A. R.  
1038 Razafimpahanana, L. Wilmé, P. C. Wright, Madagascar: Crime threatens biodiversity.  
1039 *Science* **363**, 825-825 (2019).
- 1040 95. J. P. G. Jones, J. Ratsimbazafy, A. N. Ratsifandrihamanana, J. E. M. Watson, H. T.  
1041 Andrianandrasana, M. Cabeza, J. E. Cinner, S. M. Goodman, F. Hawkins, R. A.  
1042 Mittermeier, A. L. Rabearisoa, O. S. Rakotonarivo, J. H. Razafimanahaka, A. R.  
1043 Razafimpahanana, L. Wilmé, P. C. Wright, Last chance for Madagascar's biodiversity.  
1044 *Nat. Sustain.* **2**, 350-352 (2019).
- 1045 96. United Nations World Food Programme. Southern Madagascar faces drought-driven  
1046 hunger, threatening millions. (30 November 2020,); [https://www.wfp.org/news/southern-](https://www.wfp.org/news/southern-madagascar-faces-drought-driven-hunger-threatening-millions)  
1047 [madagascar-faces-drought-driven-hunger-threatening-millions](https://www.wfp.org/news/southern-madagascar-faces-drought-driven-hunger-threatening-millions).
- 1048 97. S. E. Vollset, E. Goren, C.-W. Yuan, J. Cao, A. E. Smith, T. Hsiao, C. Bisignano, G. S.  
1049 Azhar, E. Castro, J. Chalek, A. J. Dolgert, T. Frank, K. Fukutaki, S. I. Hay, R. Lozano, A.  
1050 H. Mokdad, V. Nandakumar, M. Pierce, M. Pletcher, T. Robalik, K. M. Steuben, H. Y.  
1051 Wunrow, B. S. Zlavog, C. J. L. Murray, Fertility, mortality, migration, and population  
1052 scenarios for 195 countries and territories from 2017 to 2100: a forecasting analysis for  
1053 the Global Burden of Disease Study. *The Lancet* **396**, 1285-1306 (2020).
- 1054 98. T. T. Gatiso, L. Kulik, M. Bachmann, A. Bonn, L. Bösch, D. Eirdosh, A. Freytag, S.  
1055 Hanisch, M. Heurich, T. Sop, K. Wesche, M. Winter, H. S. Kühl, Effectiveness of  
1056 protected areas influenced by socio-economic context. *Nat. Sustain.*, (2022).
- 1057 99. N. Dudley, *Guidelines for applying protected area management categories including*  
1058 *IUCN WCPA best practice guidance on recognising protected areas and assigning*  
1059 *management categories and governance types.*, (IUCN, Gland, 2013), pp. 86.
- 1060 100. C. Amante, B. W. Eakins, "ETOPO1 1 Arc-minute global relief model: Procedures, data  
1061 sources and analysis," *NOAA Technical Memorandum NESDIS NGDC-24* (National  
1062 Oceanic and Atmospheric Administration, 2009);  
1063 <https://www.ngdc.noaa.gov/mgg/global/relief/ETOPO1/docs/ETOPO1.pdf>.
- 1064 101. UNEP-WCMC; WorldFish Centre; WRI; TNC. Global distribution of warm-water coral  
1065 reefs, compiled from multiple sources including the Millennium Coral Reef Mapping  
1066 Project. Version 4.1. Includes contributions from IMaRS-USF and IRD (2005), IMaRS-  
1067 USF (2005) and Spalding et al. (2001). (2021); <https://doi.org/10.34892/t2wk-5t34>.
- 1068 102. Center for International Earth Science Information Network - CIESIN - Columbia  
1069 University. Gridded Population of the World, Version 4 (GPWv4): Population Density,  
1070 Revision 11. (2018); <https://doi.org/10.7927/H49C6VHW>.

103. L. Giglio, C. Justice, L. Boschetti, D. Roy. MCD64A1 v006, MODIS/Terra+Aqua Burned Area Monthly L3 Global 500 m SIN Grid. (2015); <https://doi.org/10.5067/MODIS/MCD64A1.006>.
104. G. J. Huffman, D. T. Bolvin, E. J. Nelkin, D. B. Wolff, R. F. Adler, G. Gu, Y. Hong, K. P. Bowman, E. F. Stocker, The TRMM Multisatellite Precipitation Analysis (TMPA): Quasi-Global, Multiyear, Combined-Sensor Precipitation Estimates at Fine Scales. *Journal of Hydrometeorology* **8**, 38-55 (2007).
105. G. Vieilledent, C. Grinand, F. A. Rakotomalala, R. Ranaivosoa, J.-R. Rakotoarijaona, T. F. Allnutt, F. Achard. Output data from: Combining global tree cover loss data with historical national forest-cover maps to look at six decades of deforestation and forest fragmentation in Madagascar. (2018); <https://doi.org/10.18167/DVN1/AUBRRC>.
106. J. Moat, P. Smith, *Atlas of the Vegetation of Madagascar (Atlas de La Vegetation de Madagascar)*. (Royal Botanic Gardens, Kew, 2007).
107. World Bank, "Madagascar—Third Environment Program Support Project. Independent Evaluation Group, Project Performance Assessment Report 158221," (World Bank, Washington DC, USA, 2021).
108. I. R. Scales, The future of conservation and development in Madagascar: time for a new paradigm? *Madagascar Conservation & Development* **9**, 5-12 (2014).
109. S. Díaz, J. Settele, E. S. Brondízio, H. T. Ngo, J. Agard, A. Arneth, P. Balvanera, K. A. Brauman, S. H. M. Butchart, K. M. A. Chan, L. A. Garibaldi, K. Ichii, J. Liu, S. M. Subramanian, G. F. Midgley, P. Miloslavich, Z. Molnár, D. Obura, A. Pfaff, S. Polasky, A. Purvis, J. Razzaque, B. Reyers, R. R. Chowdhury, Y.-J. Shin, I. Visseren-Hamakers, K. J. Willis, C. N. Zayas, Pervasive human-driven decline of life on Earth points to the need for transformative change. *Science* **366**, eaax3100 (2019).
110. F. C. Bolam, L. Mair, M. Angelico, T. M. Brooks, M. Burgman, C. Hermes, M. Hoffmann, R. W. Martin, P. J. K. McGowan, A. S. L. Rodrigues, C. Rondinini, J. R. S. Westrip, H. Wheatley, Y. Bedolla-Guzmán, J. Calzada, M. F. Child, P. A. Cranswick, C. R. Dickman, B. Fessl, D. O. Fisher, S. T. Garnett, J. J. Groombridge, C. N. Johnson, R. J. Kennerley, S. R. B. King, J. F. Lamoreux, A. C. Lees, L. Lens, S. P. Mahood, D. P. Mallon, E. Meijaard, F. Méndez-Sánchez, A. R. Percequillo, T. J. Regan, L. M. Renjifo, M. C. Rivers, N. S. Roach, L. Roxburgh, R. J. Safford, P. Salaman, T. Squires, E. Vázquez-Domínguez, P. Visconti, J. C. Z. Woinarski, R. P. Young, S. H. M. Butchart, How many bird and mammal extinctions has recent conservation action prevented? *Conserv. Lett.* **14**, e12762 (2021).
111. L. M. Gardiner, D. Rabehevitra, R. Letsara, A. Shapcott, *Tahina spectabilis*: an exciting new discovery in Madagascar ten years on. *Palms* **61**, 69-82 (2017).
112. Durrell Wildlife Conservation Trust. Durrell Index: Ploughshare tortoise. (2022); <https://www.durrell.org/wildlife/species-index/ploughshare-tortoise/>.
113. Ministère de l'Enseignement Supérieur et de la Recherche Scientifique à Madagascar, "Stratégie Nationale de la Recherche Scientifique à Madagascar," (2013); [http://www.recherches.gov.mg/IMG/pdf/strategie\\_nationale\\_de\\_la\\_recherche.pdf](http://www.recherches.gov.mg/IMG/pdf/strategie_nationale_de_la_recherche.pdf).
114. E. Razanatsoa, S. Andriantsaralaza, S. M. Holmes, O. S. Rakotonarivo, A. N. Ratsifandrihamanana, L. Randriamiharisoa, M. Ravaloharimanitra, N. Ramahefamanana, D. Tahirinirainy, J. Raharimampionona, Fostering local involvement for biodiversity conservation in tropical regions: Lessons from Madagascar during the COVID-19 pandemic. *Biotropica* **53**, 994-1003 (2021).



- 1117 115. Convention on Biological Diversity and United Nations Environment Program, "First  
1118 draft of the post-2020 global biodiversity framework," (2020);  
1119 <https://www.cbd.int/doc/c/abb5/591f/2e46096d3f0330b08ce87a45/wg2020-03-03-en.pdf>.
- 1120 116. P. O. Waeber, L. Wilmé, J.-R. Mercier, C. Camara, P. P. Lowry, II, How effective have  
1121 thirty years of internationally driven conservation and development efforts been in  
1122 Madagascar? *PLOS ONE* **11**, e0161115 (2016).
- 1123 117. J. Börner, D. Schulz, S. Wunder, A. Pfaff, The effectiveness of forest conservation  
1124 policies and programs. *Annu. Rev. Resour.* **12**, 45-64 (2020).
- 1125 118. N. Seddon, A. Smith, P. Smith, I. Key, A. Chausson, C. Girardin, J. House, S. Srivastava,  
1126 B. Turner, Getting the message right on nature-based solutions to climate change. *Global*  
1127 *Change Biol.* **27**, 1518-1546 (2021).
- 1128 119. O. M. Grace, J. C. Lovett, C. J. N. Gore, J. Moat, I. Ondo, S. Pironon, M. K. Langat, O.  
1129 A. Pérez-Escobar, A. Ross, M. Suzan Abbo, K. K. Shrestha, B. Gowda, K. Farrar, J.  
1130 Adams, R. Cámara-Leret, M. Diazgranados, T. Ulian, S. Sagala, E. Rianawati, A. Hazra,  
1131 O. R. Masera, A. Antonelli, P. Wilkin, Plant Power: Opportunities and challenges for  
1132 meeting sustainable energy needs from the plant and fungal kingdoms. *PLANTS,*  
1133 *PEOPLE, PLANET* **2**, 446-462 (2020).
- 1134 120. C. Funk, M. D. Dettinger, J. C. Michaelsen, J. P. Verdin, M. E. Brown, M. Barlow, A.  
1135 Hoell, Warming of the Indian Ocean threatens eastern and southern African food security  
1136 but could be mitigated by agricultural development. *Proc. Natl. Acad. Sci. USA* **105**,  
1137 11081-11086 (2008).
- 1138 121. J. Eklund, J. P. G. Jones, M. Räsänen, J. Geldmann, A.-P. Jokinen, A. Pellegrini, D.  
1139 Rakotobe, O. S. Rakotonarivo, T. Toivonen, A. Balmford, Elevated fires during COVID-  
1140 19 lockdown and the vulnerability of protected areas. *Nat. Sustain.* **5**, 603-609 (2022).
- 1141 122. H. S. Grantham, A. Moilanen, K. A. Wilson, R. L. Pressey, T. G. Rebelo, H. P.  
1142 Possingham, Diminishing return on investment for biodiversity data in conservation  
1143 planning. *Conserv. Lett.* **1**, 190-198 (2008).
- 1144 123. J. Liu, F. Slik, S. Zheng, D. B. Lindenmayer, Undescribed species have higher extinction  
1145 risk than known species. *Conserv. Lett.*, e12876 (2022).
- 1146 124. IUCN. Motion 116 - Building Madagascar's capacity to counter the threat from invasive  
1147 species. Motion passed at IUCN World Conservation Congress, Marseille, 3-11  
1148 September, 2021. (2020); <https://www.iucncongress2020.org/motion/116>.
- 1149 125. Madagascar Protected Areas. Protected Areas of Madagascar. (2022);  
1150 <https://protectedareas.mg/>.
- 1151 126. L. P. Koh, S. A. Wich, Dawn of drone ecology: low-cost autonomous aerial vehicles for  
1152 conservation. *Tropical Conservation Science* **5**, 121-132 (2012).
- 1153 127. S. L. Maxwell, V. Cazalis, N. Dudley, M. Hoffmann, A. S. L. Rodrigues, S. Stolton, P.  
1154 Visconti, S. Woodley, N. Kingston, E. Lewis, M. Maron, B. B. N. Strassburg, A.  
1155 Wenger, H. D. Jonas, O. Venter, J. E. M. Watson, Area-based conservation in the twenty-  
1156 first century. *Nature* **586**, 217-227 (2020).
- 1157 128. R. E. Golden Kroner, S. Qin, C. N. Cook, R. Krithivasan, S. M. Pack, O. D. Bonilla, K.  
1158 A. Cort-Kansinally, B. Coutinho, M. Feng, M. I. Martínez Garcia, Y. He, C. J. Kennedy,  
1159 C. Lebreton, J. C. Ledezma, T. E. Lovejoy, D. A. Luther, Y. Parmanand, C. A. Ruíz-  
1160 Agudelo, E. Yereña, V. Morón Zambrano, M. B. Mascia, The uncertain future of  
1161 protected lands and waters. *Science* **364**, 881-886 (2019).
- 1162 129. I. J. Bateman, G. M. Mace, The natural capital framework for sustainably efficient and  
1163 equitable decision making. *Nat. Sustain.* **3**, 776-783 (2020).

- 1164 130. C. Birkinshaw, P. P. Lowry II, J. Raharimampionona, J. Aronson, Supporting Target 4 of  
1165 the Global Strategy for Plant Conservation by integrating ecological restoration into  
1166 Missouri Botanical Garden's conservation program in Madagascar. *Annals of the*  
1167 *Missouri Botanical Garden* **99**, 139-146 (2013).
- 1168 131. J. Aronson, A. F. Clewell, J. N. Blignaut, S. J. Milton, Ecological restoration: A new  
1169 frontier for nature conservation and economics. *J. Nat. Conserv.* **14**, 135-139 (2006).
- 1170 132. L. Robson, "The history of PHE in Madagascar: looking back over the last 25 years and  
1171 forward to the next chapter," (Blue Ventures for the Madagascar PHE Network, London,  
1172 2014); phemadagascar.org.
- 1173 133. S. Naeem, R. Chazdon, J. E. Duffy, C. Prager, B. Worm, Biodiversity and human well-  
1174 being: an essential link for sustainable development. *Proc. R. Soc. Lond. B: Biol. Sci.*  
1175 **283**, 20162091 (2016).
- 1176 134. M. Poudyal, J. P. G. Jones, O. S. Rakotonarivo, N. Hockley, J. M. Gibbons, R.  
1177 Mandimbiniaina, A. Rasoamanana, N. S. Andrianantenaina, B. S. Ramamonjisoa, Who  
1178 bears the cost of forest conservation? *PeerJ* **6**, e5106 (2018).
- 1179 135. H. Ralimanana, A. L. Perrigo, R. J. Smith, J. S. Borrell, A. Crottini, S. Faurby, J. Hackel,  
1180 M. T. Rajaonah, T. Randriamboavonjy, W. Testo, M. S. Vorontsova, N. Andela, T.  
1181 Andermann, A. M. Andriamanohera, S. Andriambololonera, S. P. Bachman, C. D.  
1182 Bacon, W. J. Baker, F. Belluardo, C. Birkinshaw, S. Cable, N. A. Canales, J. D. Carrillo,  
1183 R. Clegg, C. Clubbe, R. S. C. Cooke, G. Damasco, S. Dhanda, D. Edler, H. Farooq, P. d.  
1184 L. Ferreira, F. Forest, B. L. Fisher, L. M. Gardiner, S. M. Goodman, O. M. Grace, T. B.  
1185 Guedes, M. C. Henniges, R. Hill, C. E. R. Lehmann, P. P. L. II, L. Marline, P. Matos-  
1186 Maraví, J. Moat, B. Neves, M. G. C. Nogueira, R. E. Onstein, A. S. T. Papadopoulos, O.  
1187 A. Perez, L. N. Phelps, P. B. Phillipson, S. Pironon, N. A. S. Przelomska, M.  
1188 Rabarimanarivo, D. Rabehevitra, J. Raharimampionona, F. Rajaonary, L. R. Rajaovelona,  
1189 M. Rakotoarinivo, A. A. Rakotoarisoa, S. E. Rakotoarisoa, H. N. Rakotomalala, F.  
1190 Rakotonasolo, B. A. Ralaiveloarisoa, M. Ramirez-Herranz, J. E. N. Randriamamonjy, V.  
1191 Randrianasolo, A. Rasolohery, A. N. Ratsifandrihamanana, N. Ravololomanana, V.  
1192 Razafiniary, H. Razanajatovo, E. Razanatsoa, M. Rivers, F. Sayol, D. Silvestro, M. F. T.  
1193 Jiménez, K. Walker, B. E. Walker, P. Wilkin, J. Williams, T. Ziegler, A. Zizka, A.  
1194 Antonelli. Madagascar's extraordinary biodiversity: a data repository [Data set]. (2021);  
1195 <https://doi.org/10.5281/zenodo.6586742>.
- 1196 136. A. Zizka, D. Silvestro, P. Vitt, T. M. Knight, Automated conservation assessment of the  
1197 orchid family with deep learning. *Conserv. Biol.* **35**, 897-908 (2021).
- 1198 137. Madagascar Catalogue. Catalogue of the Vascular Plants of Madagascar. (2021);  
1199 <http://www.efloras.org/madagascar>. [Accessed March 2021].
- 1200 138. C. Maldonado, C. I. Molina, A. Zizka, C. Persson, C. M. Taylor, J. Albán, E. Chilquillo,  
1201 N. Rønsted, A. Antonelli, Estimating species diversity and distribution in the era of Big  
1202 Data: to what extent can we trust public databases? *Global Ecol. Biogeogr.* **24**, 973-984  
1203 (2015).
- 1204 139. A. Zizka, C. F. Antunes, A. Calvente, M. Rocio Baez-Lizarazo, A. Cabral, J. F. R.  
1205 Coelho, M. Colli-Silva, M. R. Fantinati, M. F. Fernandes, T. Ferreira-Araújo, F. Gondim  
1206 Lambert Moreira, N. M. C. Santos, T. A. B. Santos, R. C. dos Santos-Costa, F. C.  
1207 Serrano, A. P. Alves da Silva, A. de Souza Soares, P. G. Cavalcante de Souza, E. Calisto  
1208 Tomaz, V. F. Vale, T. L. Vieira, A. Antonelli, No one-size-fits-all solution to clean  
1209 GBIF. *PeerJ*, 8:e9916 (2020).

- 1210 140. GBIF.org. GBIF Occurrence Download. Tracheophyta, iNaturalist. (2020);  
1211 <https://doi.org/10.15468/dl.jbh5dk>.
- 1212 141. GBIF.org. GBIF Occurrence Download. Tracheophyta, Missouri. (2020);  
1213 <https://doi.org/10.15468/dl.7skt3j>.
- 1214 142. A. Zizka, D. Silvestro, T. Andermann, J. Azevedo, C. Duarte Ritter, D. Edler, H. Farooq,  
1215 A. Herdean, M. Ariza, R. Scharn, S. Svantesson, N. Wengström, V. Zizka, A. Antonelli,  
1216 CoordinateCleaner: Standardized cleaning of occurrence records from biological  
1217 collection databases. *Methods Ecol. Evol.* **10**, 744-751 (2019).
- 1218 143. B. H. Daru, D. S. Park, R. B. Primack, C. G. Willis, D. S. Barrington, T. J. S. Whitfield,  
1219 T. G. Seidler, P. W. Sweeney, D. R. Foster, A. M. Ellison, C. C. Davis, Widespread  
1220 sampling biases in herbaria revealed from large-scale digitization. *New Phytol.* **217**, 939-  
1221 955 (2018).
- 1222 144. C. Meyer, P. Weigelt, H. Kreft, Multidimensional biases, gaps and uncertainties in global  
1223 plant occurrence information. *Ecol. Lett.* **19**, 992-1006 (2016).
- 1224 145. A. Zizka, A. Antonelli, D. Silvestro, sampbias, a method for quantifying geographic  
1225 sampling biases in species distribution data. *Ecography* **44**, 25-32 (2021).
- 1226 146. S. P. Bachman, J. Moat, A. Hill, J. de la Torre, B. Scott, Supporting Red List threat  
1227 assessments with GeoCAT: geospatial conservation assessment tool. *ZooKeys* **150**,  
1228 (2011).
- 1229 147. J. Moat. rCAT: Conservation Assessment Tools. R package version 0.1.5. (2017);  
1230 <https://cran.r-project.org/package=rCAT>.
- 1231 148. S. E. Fick, R. J. Hijmans, WorldClim 2: new 1-km spatial resolution climate surfaces for  
1232 global land areas. *Int. J. Climatol.* **37**, 4302-4315 (2017).
- 1233 149. D. M. Olson, E. Dinerstein, E. D. Wikramanayake, N. D. Burgess, G. V. N. Powell, E. C.  
1234 Underwood, J. A. D'amico, I. Itoua, H. E. Strand, J. C. Morrison, C. J. Loucks, T. F.  
1235 Allnutt, T. H. Ricketts, Y. Kura, J. F. Lamoreux, W. W. Wettengel, P. Hedao, K. R.  
1236 Kassem, Terrestrial ecoregions of the world: a new map of life on Earth: A new global  
1237 map of terrestrial ecoregions provides an innovative tool for conserving biodiversity.  
1238 *Bioscience* **51**, 933-938 (2001).
- 1239 150. A. Antonelli, A. Zizka, F. A. Carvalho, R. Scharn, C. D. Bacon, D. Silvestro, F. L.  
1240 Condamine, Amazonia is the primary source of Neotropical biodiversity. *Proc. Natl.*  
1241 *Acad. Sci. USA* **115**, 6034-6039 (2018).
- 1242 151. O. Venter, E. W. Sanderson, A. Magrath, J. R. Allan, J. Beher, K. R. Jones, H. P.  
1243 Possingham, W. F. Laurance, P. Wood, B. M. Fekete, M. A. Levy, J. E. M. Watson,  
1244 Global terrestrial Human Footprint maps for 1993 and 2009. *Scientific Data* **3**, 160067  
1245 (2016).
- 1246 152. M. Diazgranados, Allkin, B., Black N., Cámara-Leret, R., Canteiro C., Carretero J.,  
1247 Eastwood R., Hargreaves S., Hudson A., Milliken W., Nesbitt, M., Ondo, I., Patmore, K.,  
1248 Pironon, S., Turner, R., Ulian, T. World Checklist of Useful Plant Species. (2020);  
1249 doi:10.5063/F1CV4G34.
- 1250 153. D. Silvestro, T. Andermann, Prior choice affects ability of Bayesian neural networks to  
1251 identify unknowns. *ArXiv:2005.04987 [Cs, Stat]*, (2020).
- 1252 154. K. He, X. Zhang, S. Ren, J. Sun, in *2015 IEEE International Conference on Computer*  
1253 *Vision (ICCV)*. (2015), pp. 1026-1034.
- 1254 155. R. Webportal. Rebioma. (2019); <http://data.rebioma.net/>.
- 1255 156. UNEP-WCMC. Alliance for Zero Extinction sites. (2020); [https://www.biodiversitya-](https://www.biodiversitya-z.org/content/alliance-for-zero-extinction-sites-aze)  
1256 [z.org/content/alliance-for-zero-extinction-sites-aze](https://www.biodiversitya-z.org/content/alliance-for-zero-extinction-sites-aze).



- 1257 157. H. Wickham. rvest. (2021); <https://rvest.tidyverse.org/>.
- 1258 158. S. M. Goodman, J. R. Rakotoson, P. M. Razafimahatratra, M. J. Raheirilalao, Introduction  
1259 to part I, in *The terrestrial protected areas of Madagascar: Their history, description,*  
1260 *and biota*, S. M. Goodman, M. J. Raheirilalao, S. Wohlhauser, Eds. (Association Vahatra,  
1261 Antananarivo, 2018), pp. 33–78.
- 1262 159. GADM. GADM database of Global Administrative Areas v 3.6. (2021);  
1263 <https://gadm.org/data.html>.
- 1264 160. Madagascar Catalogue. Catalogue of the Vascular Plants of Madagascar. (2022);  
1265 <http://www.efloras.org/madagascar>. [Accessed May 2022].
- 1266 161. L. Giglio, L. Boschetti, D. P. Roy, M. L. Humber, C. O. Justice, The Collection 6  
1267 MODIS burned area mapping algorithm and product. *Remote Sens. Environ.* **217**, 72-85  
1268 (2018).
- 1269 162. S. Archibald, C. E. R. Lehmann, C. M. Belcher, W. J. Bond, R. A. Bradstock, A. L.  
1270 Daniau, K. G. Dexter, E. J. Forrester, M. Greve, T. He, S. I. Higgins, W. A. Hoffmann, B.  
1271 B. Lamont, D. J. McGlenn, G. R. Moncrieff, C. P. Osborne, J. G. Pausas, O. Price, B. S.  
1272 Ripley, B. M. Rogers, D. W. Schwilk, M. F. Simon, M. R. Turetsky, G. R. Van der Werf,  
1273 A. E. Zanne, Biological and geophysical feedbacks with fire in the Earth system.  
1274 *Environmental Research Letters* **13**, 033003 (2018).
- 1275 163. S. Archibald, C. E. R. Lehmann, J. L. Gómez-Dans, R. A. Bradstock, Defining pyromes  
1276 and global syndromes of fire regimes. *Proc. Natl. Acad. Sci. USA* **110**, 6442-6447 (2013).
- 1277 164. S. T. Alvarado, N. Andela, T. S. F. Silva, S. Archibald, Thresholds of fire response to  
1278 moisture and fuel load differ between tropical savannas and grasslands across continents.  
1279 *Global Ecol. Biogeogr.* **29**, 331-344 (2020).
- 1280 165. R. A. Bradstock, A biogeographic model of fire regimes in Australia: current and future  
1281 implications. *Global Ecol. Biogeogr.* **19**, 145-158 (2010).
- 1282 166. N. Andela, D. C. Morton, L. Giglio, Y. Chen, G. R. van der Werf, P. S. Kasibhatla, R. S.  
1283 DeFries, G. J. Collatz, S. Hantson, S. Kloster, D. Bachelet, M. Forrest, G. Lasslop, F. Li,  
1284 S. Mangeon, J. R. Melton, C. Yue, J. T. Randerson, A human-driven decline in global  
1285 burned area. *Science* **356**, 1356-1362 (2017).
- 1286 167. M. J. Andrade-Núñez, T. M. Aide, Using nighttime lights to assess infrastructure  
1287 expansion within and around protected areas in South America. *Environmental Research*  
1288 *Communications* **2**, 021002 (2020).
- 1289 168. X. Li, Y. Zhou, M. Zhao, X. Zhao, A harmonized global nighttime light dataset 1992–  
1290 2018. *Scientific Data* **7**, 168 (2020).
- 1291 169. T. F. Allnutt, S. Ferrier, G. Manion, G. V. N. Powell, T. H. Ricketts, B. L. Fisher, G. J.  
1292 Harper, M. E. Irwin, C. Kremen, J.-N. Labat, D. C. Lees, T. A. Pearce, F.  
1293 Rakotondrainibe, A method for quantifying biodiversity loss and its application to a 50-  
1294 year record of deforestation across Madagascar. *Conserv. Lett.* **1**, 173-181 (2008).
- 1295 170. R Core Team, *R: A language and environment for statistical computing*. (R Foundation  
1296 for Statistical Computing, Vienna, Austria, 2020).
- 1297 171. H. Wickham, M. Averick, J. Bryan, W. Chang, L. D. A. McGowan, R. François, G.  
1298 Grolemond, A. Hayes, L. Henry, J. Hester, M. Kuhn, T. L. Pedersen, E. Miller, S. M.  
1299 Bache, K. Müller, J. Ooms, D. Robinson, D. P. Seidel, V. Spinu, K. Takahashi, D.  
1300 Vaughan, C. Wilke, K. Woo, H. Yutani, Welcome to the Tidyverse. *Journal of Open*  
1301 *Source Software* **4**, 1686 (2019).
- 1302 172. K. Müller, H. Wickham. tibble: Simple Data Frames. (2021); [https://CRAN.R-](https://CRAN.R-project.org/package=tibble)  
1303 [project.org/package=tibble](https://CRAN.R-project.org/package=tibble).

- 1304 173. H. Wickham, François, R., Henry, L., Müller, K. dplyr: A Grammar of Data  
1305 Manipulation. (2021); <https://CRAN.R-project.org/package=dplyr>.
- 1306 174. H. Wickham. tidyr: Tidy Messy Data. (2020); [https://CRAN.R-](https://CRAN.R-project.org/package=tidyr)  
1307 [project.org/package=tidyr](https://CRAN.R-project.org/package=tidyr).
- 1308 175. H. Wickham, J. Hester. readr: Read Rectangular Text Data. (2020); [https://CRAN.R-](https://CRAN.R-project.org/package=readr)  
1309 [project.org/package=readr](https://CRAN.R-project.org/package=readr).
- 1310 176. H. Wickham, Chang, W., Henry, L., Pedersen, T. L., Takahashi, K., Wilke, C., Woo, K.,  
1311 Yutani, H., Dunnington, D. ggplot2: Create Elegant Data Visualisations Using the  
1312 Grammar of Graphics. (2020); <https://CRAN.R-project.org/package=ggplot2>.
- 1313 177. D. Pierce. ncdf4: Interface to Unidata netCDF (Version 4 or Earlier) Format Data Files.  
1314 (2019); <http://cirrus.ucsd.edu/pierce/ncdf>.
- 1315 178. R. S. Bivand, E. J. Pebesma, V. Gomez-Rubio, *Applied spatial data analysis with R*,  
1316 *Second edition.*, (Springer, 2013).
- 1317 179. E. Pebesma. sf: Simple Features for R. (2021); <https://CRAN.R-project.org/package=sf>.
- 1318 180. R. J. Hijmans. raster: Geographic Data Analysis and Modeling. R package version 3.4-5.  
1319 (2020); <https://CRAN.R-project.org/package=raster>.
- 1320 181. A. South. rnaturalearth: World Map Data from Natural Earth. (2017);  
1321 <https://github.com/ropenscilabs/rnaturalearth>.
- 1322 182. A. South. rnaturalearthdata: World Vector Map Data from Natural Earth Used in  
1323 rnaturalearth. (2017); <https://github.com/ropenscilabs/rnaturalearthdata>.
- 1324 183. E. Pebesma. stars: Spatiotemporal Arrays, Raster and Vector Data Cubes. (2020); [https://](https://cloud.r-project.org/package=stars)  
1325 [cloud.r-project.org/package=stars](https://cloud.r-project.org/package=stars).
- 1326 184. S. Chamberlain, E. Szoecs, Z. Foster, Z. Arendsee. taxize: Taxonomic Information from  
1327 Around the Web. (2020); <https://CRAN.R-project.org/package=taxize>.
- 1328 185. M. Freiberg, M. Winter, A. Gentile, A. Zizka, A. N. Muellner-Riehl, A. Weigelt, C.  
1329 Wirth, LCVP, The Leipzig catalogue of vascular plants, a new taxonomic reference list  
1330 for all known vascular plants. *Scientific Data* **7**, 416 (2020).
- 1331 186. C. R. Harris, K. J. Millman, S. J. van der Walt, R. Gommers, P. Virtanen, D. Cournapeau,  
1332 E. Wieser, J. Taylor, S. Berg, N. J. Smith, R. Kern, M. Picus, S. Hoyer, M. H. Van  
1333 Kerkwijk, M. Brett, A. Haldane, J. Fernández del Río, M. Wiebe, P. Peterson, P. Gérard-  
1334 Marchant, K. Sheppard, T. Reddy, W. Weckesser, H. Abbasi, C. Gohlke, T. E. Oliphant,  
1335 Array programming with NumPy. *Nature* **585**, 357-362 (2020).
- 1336 187. J. Reback, jbrockmendel, W. McKinney, J. V. d. Bossche, T. Augspurger, P. Cloud, S.  
1337 Hawkins, g. Sinhrks, M. Roeschke, A. Klein, T. Petersen, J. Tratner, C. She, W. Ayd, P.  
1338 Hoefler, S. Naveh, M. Garcia, J. Schendel, A. Hayden, D. Saxton, R. Shadrach, M. E.  
1339 Gorelli, V. Jancauskas, F. Li, attack68, A. McMaster, P. Battiston, S. Seabold, K. Dong.  
1340 pandas-dev/pandas: Pandas 1.3.0. (2021);  
1341 <https://zenodo.org/record/5060318#.YRWfgIhKhPY>.
- 1342 188. K. Jordahl, J. V. d. Bossche, M. Fleischmann, J. Wasserman, J. McBride, J. Gerard, J.  
1343 Tratner, M. Perry, A. G. Badaracco, C. Farmer, G. A. Hjelle, A. D. Snow, M. Cochran, S.  
1344 Gillies, L. Culbertson, M. Bartos, N. Eubank, maxalbert, A. Bilogur, S. Rey, C. Ren, D.  
1345 Arribas-Bel, L. Wasser, L. J. Wolf, M. Journois, J. Wilson, A. Greenhall, C. H. Filipe, F.  
1346 Leblanc. geopandas/geopandas: v0.9.0. (2020);  
1347 <https://zenodo.org/record/4569086#.YRWgdYhKhPY>.
- 1348 189. S. Gillies. Rasterio: access to geospatial raster data. (2021);  
1349 <https://pypi.org/project/rasterio/1.2.6/>.

190. S. Hoyer, J. Hamman, M. Roos, keewis, D. Cherian, C. Fitzgerald, K. Fujii, F. Maussion, M. Hauser, crusaderky, S. Clark, A. Kleeman, T. Kluyver, J. Munroe, A. Amici, T. Nicholas, A. Barghini, A. Banihirwe, gimperiale, Z. Hatfield-Dodds, R. Abernathey, Illviljan, R. Bell, johnmotani, M. Roszko, P. J. Wolfram, J. Signell, K. Mühlbauer, Y. B. Sinai, B. Bovy. pydata/xarray: v0.18.2. (2021); <https://zenodo.org/record/4774304#.YRWi4YhKhPY>.
191. A. D. Snow, D. Brochart, T. Chegini, A. Amici, D. Hoese, J. Hamman, R. Bell, RichardScottOZ, S. Henderson, T. G. Badger, T. Augspurger, pmallas, remi-braun. corteva/rioxarray: 0.4.3 Release. (2021); <https://zenodo.org/record/5032168#.YRWj54hKhPY>.
192. GDAL/OGR contributors. GDAL/OGR Geospatial Data Abstraction software Library. (2021); <https://gdal.org>.
193. T. A. Caswell, M. Droettboom, A. Lee, E. S. d. Andrade, J. Hunter, T. Hoffmann, E. Firing, J. Klymak, D. Stansby, N. Varoquaux, J. H. Nielsen, B. Root, R. May, P. Elson, J. K. Seppänen, D. Dale, J.-J. Lee, D. McDougall, A. Straw, P. Hobson, C. Gohlke, hannah, T. S. Yu, E. Ma, A. F. Vincent, S. Silvester, C. Moad, N. Kniazev, E. Ernest, P. Ivanov. matplotlib/matplotlib: REL: v3.4.2. (2021); <https://zenodo.org/record/4743323#.YRWk64hKhPY>.
194. M. L. Waskom. seaborn: statistical data visualization. (2021); <https://doi.org/10.21105/joss.03021>.

## Acknowledgments:

Carly Cowell (RBG Kew) provided information on METT assessments and monitoring and Karen Freeman (Madagascar Fauna and Flora Group) provided information on invasive vertebrates. Anna Rauhaus (Cologne Zoo), Joel Kamphausen, and Laura Leiss (both of University of Cologne) supported the zoo database analyses. Inessa Voet illustrated the graphical abstract. Julia P.G. Jones (Bangor University) contributed to discussions that shaped and improved the analyses and text. We thank James Aronson (Missouri Botanical Garden), David Ashley (UK Ambassador to Madagascar and Comoros), Adolphe Lehavana (Missouri Botanical Garden), Jonah Ratsimbazafy (Houston Zoo and the Groupe d'Etude et de Recherche sur les Primates de Madagascar), Serge Ratsirahonana (Fondation pour les Aires Protégées et la Biodiversité de Madagascar), and George E. Schatz (Missouri Botanical Garden) for their insightful feedback on the first full draft of the manuscript, and three anonymous reviewers and the editors for valuable feedback. We stress that

1383 all views expressed in this article are only those of the authors. **Data and materials availability:**

1384 All data is available in the manuscript or the supplementary materials.

1385 **Funding:** AA acknowledges financial support from the Swedish Research Council (2019-05191),  
1386 the Swedish Foundation for Strategic Research (FFL15-0196) and a grant from the Kew  
1387 Foundation. TA was supported by the SciLifeLab & Wallenberg Data Driven Life Science  
1388 Program, grant no. KAW 2020.0239. CDB and MFTJ were supported by a grant from the Swedish  
1389 Research Council to CDB, grant no. 2017-04980. FB was supported by Fundação para a Ciência  
1390 e a Tecnologia, grant no. PD/BD/128493/2017. JB, JH, SP, and BEW were supported by a Future  
1391 Leader Fellowship from RBG Kew. NAC was financed by H2020 MSCA-ITN-ETN Plant.ID, a  
1392 European Union's Horizon 2020 research and innovation programme under grant agreement no.  
1393 765000. JDC was supported by the Swiss National Science Foundation, grant no. P400PB\_186733  
1394 and P4P4PB\_199187. AC was supported by Portuguese National Funds through the Foundation  
1395 for Science and Technology, FCT, grant no. 2020.00823.CEECIND and project PTDC/BIA-  
1396 EVL/31254/2017. GD was supported by Vinnova, grant no. 2019-02717. DE was supported by  
1397 the Swedish Research Council, grant no. 2016-00796. SF was supported by the Swedish Research  
1398 Council, grant no. 2017-03862. DE was supported by the Swedish Research Council, grant no.  
1399 2016-00796, and the Swedish Foundation for Strategic Research, grant no. FFL15-0196. BLF was  
1400 supported by National Science Foundation, grant no. DEB- 1655076. PLF was supported by the  
1401 MEMOVA project, EU Operational Programme Research, Development and Education, grant no.  
1402 CZ.02.2.69/0.0/0.0/18\_053/0016982. TBG is supported by a young researcher grant from São  
1403 Paulo Research Foundation (FAPESP, #2021/07161-6, #2022/09428-2), Brazil. MCH and RH are  
1404 Natural Environment Research Council funded Ph.D. students with the London NERC DTP  
1405 (NERC Ref: NE/L002485/1). PMM was funded by the Czech Science Foundation, grant no. GJ20-

1406 18566Y. BN was supported by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior,  
1407 Brasil (CAPES) – (PDSE 88881.132750/2016-01). REO acknowledges the support of iDiv funded  
1408 by the German Research Foundation (DFG–FZT 118, 202548816). ASTP was supported by the  
1409 Natural Environment Research Council (NERC), grant no. NE/R001081/1. LNP was supported by  
1410 a Swiss National Science Foundation Early Postdoc Mobility Grant, grant no. P2LAP2\_187745.  
1411 MRH was supported by a beca de doctorado del Instituto de Ecología y Biodiversidad, Chile;  
1412 Programa de Doctorado en Biología y Ecología Aplicada, Universidad Católica del Norte,  
1413 Universidad de La Serena, La Serena, Chile and a ANID National Scholarship (grant no.  
1414 21181931); Institute of Ecology and Biodiversity (Project ANID FB210006). ER was supported  
1415 by the NRF / SASSCAL (Southern African Science Service Centre), grant no. 118589, and the  
1416 NRF / African Origins Platform, grant no. 117666. FS was supported by European Union's  
1417 Horizon 2020 research and innovation program under the Marie Skłodowska-Curie grant  
1418 agreement, grant no. 838998. DS received funding from the Swiss National Science Foundation  
1419 (PCEFP3\_187012; FN-1749) and from the Swedish Research Council (VR: 2019-04739). MSV  
1420 was supported by a GBIF: Biodiversity Information for Development grant. KW was supported  
1421 by a Techne Arts and Humanities Research Council Doctoral Training Partnership. PW was  
1422 supported by the Darwin Initiative, grant nos 22-005 and EIDPO049.

1423 **Author contributions:** The paper was project-led, with overall writing responsibility, by AA,  
1424 ALP, HRal, and RJS. The content was divided into three sections, with corresponding working  
1425 groups, each making full use of the expertise and direct in-country experience of the 25 Malagasy  
1426 co-authors. The sections were organized, written, and led by ALP, HRal, JSB, MSV, MTR, RJS,  
1427 SF, and TR. Formal analyses were carried out by ARas, AZ, DE, DS, HF, JSB, LNP, MCH, MRi,  
1428 MSV, RSCC, SF, TA, and WT. Visualizations and figures were done by DE, FS, HF, HRaz, JSB,

1429 LNP, and RSCC. This paper and its sister manuscript on evolution, distribution and use (1) were  
1430 based on the outputs of a consortium focusing on Madagascar's biodiversity. The data curation,  
1431 investigation and resources therefore formed the foundation for the project as a whole and we  
1432 therefore list all co-authors involved in these activities across both papers. Data curation was  
1433 carried out by AC, AMA, ARak, ARas, AZ, BAR, BLF, BN, CDB, DE, GD, DR, ER, FRaj, FRak,  
1434 FS, HNR, HRal, HRaz, JDC, JENR, JH, JM, KW, LM, LNP, LRR, MFTJ, MGCN, MRab, MRH,  
1435 MRi, MSV, MTR, NA, NAC, NASP, NR, OAP, PLF, PMM, PBP, PW, RC, REO, RH, SA, SC,  
1436 SD, SER, SMG, SP, TBG, TR, TZ, VRan, VRaz, and WT. Investigation was carried out by AC,  
1437 AMA, ARak, ARas, ASTP, BAR, BN, FB, FRak, HNR, HRaz, MGCN, OAP, OMG, PLF, PMM,  
1438 PW, RA, RH, RSCC, SER, and TBG. Methodology was developed by AC, AZ, BEW, CERL, DS,  
1439 JH, JM, JSB, LNP, MSV, SPB, and TA. Resources were provided by AC, DR, ER, JR, LM, MRak,  
1440 MSV, MTR, NASP, and TR. Software was designed by AZ, BEW, DE, DS, and HF. Validations  
1441 were carried out by AC, DE, JM, and WJB. All co-authors were involved in the writing, revision  
1442 and editing of the text.

1443 **Competing interests:** Authors declare no competing interests.

1444 **Data and materials availability:** All data is available in the main text or the Supplementary  
1445 Materials available on the *Science* website.

1446

## 1447 **Supplementary Materials**

1448 Materials and Methods

1449 Figure S1

1450 Tables S1–S8

1451   References 135-194

1452

1453