A record-breaking extreme heat event caused unprecedented warming of lakes in China

Wang, Weijia; Shi, Kun; Wang, Xiwen; Wang, Siqi; Zhang, Dong; Peng, Yuanyuan; Li, Na; Zhang, Yunlin; Zhang, Yibo; Qin, Boqiang; lestyn Woolway, R.; Jeppesen, Erik

Science bulletin

DOI: 10.1016/j.scib.2023.03.001

Published: 30/03/2023

Peer reviewed version

Cyswllt i'r cyhoeddiad / Link to publication

Dyfyniad o'r fersiwn a gyhoeddwyd / Citation for published version (APA): Wang, W., Shi, K., Wang, X., Wang, S., Zhang, D., Peng, Y., Li, N., Zhang, Y., Zhang, Y., Qin, B., lestyn Woolway, R., & Jeppesen, E. (2023). A record-breaking extreme heat event caused unprecedented warming of lakes in China. *Science bulletin*, *68*(6), 578-582. https://doi.org/10.1016/j.scib.2023.03.001

Hawliau Cyffredinol / General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

· Users may download and print one copy of any publication from the public portal for the purpose of private study or research.

- You may not further distribute the material or use it for any profit-making activity or commercial gain
 You may freely distribute the URL identifying the publication in the public portal ?

Take down policy If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Science Bulletin

3 4	1	A record-breaking extreme heat event caused unprecedented warming of lakes in China
5		
6 7	2	Weijia Wang ^{a, b} , Kun Shi ^{a*} , Xiwen Wang ^{a, c} , Siqi Wang ^d , Dong Zhang ^{a, b} , Yuanyuan Peng ^{a, b} ,
8 9 10	3	Na Li ^{a, b} , Yunlin Zhang ^a , Yibo Zhang ^a , Boqiang Qin ^a , R. Iestyn Woolway ^e , Jeppesen Erik ^{f-i}
11 12 13	4	^a Taihu Laboratory for Lake Ecosystem Research, State Key Laboratory of Lake Science and
14 15	5	Environment, Nanjing Institute of Geography and Limnology, Chinese Academy of Sciences,
16 17 18	6	Nanjing 210008, China
19 20 21	7	^b University of Chinese Academy of Sciences, Beijing 100049, China
22 23	8	° Key Laboratory of Western China's Environmental Systems (Ministry of Education), College
24 25 26	9	of Earth and Environmental Sciences, Lanzhou University, Lanzhou, 730000, China
27 28	10	^d State Key Laboratory of Information Engineering in Surveying, Mapping, and Remote
29 30 31	11	Sensing (LIESMARS), Wuhan University, Wuhan, 430079, China
32 33 34	12	^e School of Ocean Sciences, Bangor University, Menai Bridge, Anglesey, UK
35 36	13	^f Department of Ecoscience, Aarhus University, Aarhus, 6000C, Denmark
37 38	14	^g Sino-Danish Centre for Education and Research, Beijing, China
39 40	15	^h Limnology Laboratory, Department of Biological Sciences and Centre for Ecosystem
41 42	16	Research and Implementation (EKOSAM), Middle East Technical University, Ankara, 06800,
43 44	17	Türkiye
45 46	18	ⁱ Institute of Marine Sciences, Middle East Technical University, Erdeneli-Mersin, 33731,
47 48 49	19	Türkiye
50 51 52	20	*Corresponding author: Kun Shi, Nanjing Institute of Geography and Limnology, Chinese
53 54	21	Academy of Sciences, 73 East Beijing Road, Nanjing 210008, P. R. China, Tel: + 86-25-
55 56	22	86882198, Email: <u>kshi@niglas.ac.cn</u>
57 58 59	23	

24	Strong evidence confirms that extreme climatic events, such as heat waves, storms and
25	droughts, are becoming more frequent due to anthropogenic climate change ^[1] . These extreme
26	events may have overwhelmingly negative impacts on lake ecosystems ^[2] . Specifically, by
27	warming lake surface waters, extreme heat events are expected to influence many within-lake
28	physical, chemical and biological processes ^[3] . Critically, extreme heat events can alter the lake
29	environment in only a fraction of the time that is needed for achieving the same effect of long-
30	term climate change ^[4] . In some cases, extreme heat events can exceed the resilience capacity
31	of a lake ecosystem and lead to a loss of functionality and can even trigger catastrophic regime
32	shifts ^[5] . From mid-June to late August 2022, a record-breaking extreme heat event, with the
33	longest duration and highest intensity since recording began in 1961, swept across China ^[6] . The
34	direct driver of this heat event was the Western Pacific Subtropical High, which normally sits
35	as a smaller pressure system to the east of Yangtze River Basin. However, in summer 2022, the
36	system was strong and hovered over Yangtze River Basin, thereby preventing cold air from the
37	north and moisture from the Indian Ocean from reaching the region ^[6] . Here, we used satellite
38	observations from 2000 to 2022 to demonstrate the effects of the unprecedented warming of
39	China's lakes caused by this extreme event.

We found that the average air temperature (AT) across China in summer (June to August)
2022 was 24.12 °C, which was 0.75 °C higher than the base-period average between 2000 and
2021 (Figure. 1, Table S1). Specifically, AT was above 30.00 °C and 35.00 °C (threshold of
high AT) for 80 and 42 days, respectively, which is 5 and 22 days more than the corresponding
total cumulative yearly days from 2000 to 2021. Yangtze River Basin, Huai River Basin,
Yellow River Basin and Southeast River Basin also experienced a significant increase in AT,

Page 5 of 47

Science Bulletin

46	with the summer mean temperature being 1.70 °C, 1.46 °C, 1.11 °C and 0.95 °C higher than
47	the corresponding temperature for the period 2000 - 2021, respectively. Inland River basin
48	experienced the warmest heat event in July, with AT reaching 40.00 °C. Moreover, there were
49	a total of 42 days in summer when ATs ≥ 35.00 °C and 80 days when ATs ≥ 30.00 °C, which
50	is 22 days and 5 days more than during 2000 – 2021, respectively. Temperature extremes were
51	also observed in Yangtze River Basin, with temperatures reaching 37.83 °C and a total of 24
52	and 71 days with temperatures exceeding 35.00 °C and 30.00 °C, respectively, which is 23 and
53	21 days more than the corresponding number of days observed from 2000 to 2021.
54	The summer every a local surface air temperature (SAT) shows the studied lakes in China
54	The summer average local surface an temperature (SAT) above the studied lakes in China
55	exhibited heterogeneity in 2022 compared to that between 2000 and 2021 (Figure 1, Table S2),
56	being extremely high in the east and relatively low over the Tibetan Plateau. Specifically, the
57	SAT in summer 2022 was highest over Lake Dongtinghu (SAT = 30.08 °C) in Yangtze River
58	Basin. The SAT anomalies in summer 2022 showed varying levels of increase in most lakes (<i>n</i>
59	= 105, 88.24%), with an average nationwide increase of 1.20 (min: -0.60; max: 2.95) °C
60	compared to those in 2000 - 2021. An exception to this was observed in several lakes
61	predominately located in the Songhua and Liao River Basins ($n = 14, 11.76\%$).

Lake surface water temperatures (LSWTs) across China during the summer of 2022
demonstrated a spatial pattern consistent with those observed in SAT (Figure. 2, Table S3).
When compared to mean temperatures between 2000 and 2021, the LSWT in 2022 showed a
significant increase nationwide, being 1.63 °C warmer on average. The increase in LSWT is
considerably higher than the 0.75 °C increase in AT (Figure. S2). Thirty-three lakes

2
כ ⊿
4
5
6
7
8
9
10
11
12
13
14
15
16
17
10
10
עו 20
20
21
22
23
24
25
26
27
28
29
30
31
32
32
27
54 25
35
36
37
38
39
40
41
42
43
44
45
46
47
48
10
50
50
ן כ בי
5Z
53
54
55
56
57
58
59
60

1

67	experienced LSWT anomalies above 2.00 °C during the summer of 2022, most notably in Huai
68	River Basin, Yangtze River Basin, Inland River Basin and Southwest River Basin. Lakes
69	situated in the Yangtze River Basin experienced the highest LSWT of 32.24 °C in August,
70	which was 2.20 °C higher than the average maximum LSWT during 2000 – 2021. This was
71	followed by Huaihe River basin, where the highest temperature reached 30.33°C. Huai River
72	Basin experienced the largest increase in LSWT (average = 2.01 °C; min: 1.89 °C; max:
73	2.21 °C), followed by Southwest River Basin (average = 1.81 °C; min: 0.55 °C; max: 2.44 °C),
74	and Songhua and Liao River Basins experienced the smallest increase in LSWT (average =
75	0.33 °C; min: 0.12 °C; max: 0.45 °C). Seventy-four lakes (62.18%) had significantly greater
76	LSWT increases than SAT. These lakes are mainly located in Yangtze River Basin, Huai River
77	Basin, Inland River Basin and Southwest River Basin (Figure. S3).
78	The increase in average LSWT in 2022 was greatest in August, peaking at 1.78 °C,

79 followed by July (1.38 °C), and was lowest in June (1.12 °C) (Figure. S4, Table S3). LSWT 80 anomalies in Songhua and Liao River Basin, Hai River Basin, Pearl River Basin, and Southeast 81 River Basin were negative in June, indicating that the summer season began later than usual in 82 northeastern and southeastern China. Except a few lakes in Inland River Basin, lakes across the 83 whole country were subjected to high temperatures in July, and the anomalies in Southwest 84 River Basin reached the highest level of 2.28 °C. In August, LSWT anomalies in Songhua and 85 Liao River Basin and Pearl River Basin started to decrease, while they were still high in other 86 basins, up to 2.71 °C in Huai River Basin and above 2.00 °C in Southeast River Basin, Yangtze 87 River Basin, and Southwest River Basin.

Page 7 of 47

Science Bulletin

88	The average LSWT increase of 1.63 °C in 2022 was much higher than the cumulative
89	increase of the past 20 years (Figure. 2, Table S4). The average rate of increase in LSWT in
90	China from 2000 to 2021 was 0.11 °C/decade, resulting in a twenty-year average increase of
91	0.22 °C. There were 17 lakes with a significant increase in LSWT. All basins except Southeast
92	River Basin exhibited a warming trend in LSWT from 2000 to 2021, with an average increase
93	of 0.15 °C/decade, being highest in Huai River Basin with 0.36 °C/decade. Incorporating 2022
94	into the time series (i.e., the period 2000 - 2022), an LSWT warming trend appeared in all
95	basins with an average of 0.24 °C/decade and up to 0.47°C/decade in Huai River Basin. The
96	extreme heat event in 2022 had the largest impact on the LSWT trend in Southwest River Basin;
97	thus, LSWT here changed from 0.09 to 0.24 °C/decade, followed by Southeast River Basin,
98	changing from -0.08 to 0.05 °C/decade.
99	Increases in LSWT may have a cascade of ecological and environmental consequences
100	such as alterations in lake stratification and mixing regimes, water level decline, a decrease in
101	dissolved oxygen concentrations, highly frequent cyanobacterial blooms, and loss of habitat for
102	cold-water species ^[5] . Specifically, in terms of the physical environment of lakes, surface water
103	temperature will be more responsive to extreme heat events compared to bottom waters leading

102 cold-water species^[5]. Specifically, in terms of the physical environment of lakes, surface water 103 temperature will be more responsive to extreme heat events compared to bottom waters, leading 104 to an increase in thermal stability. Subsequently, the timing of seasonal overturn in 105 autumn/winter will likely be delayed, leading to a prolonged summer stratification^[7]. In 106 addition, an increase in LSWT may also lead to higher evaporation rates, and thus reduce lake 107 volumes and the spatial extent of surface water in the absence of substantial water inflows^[8].

Chemical processes in lakes can be influenced, both directly and indirectly, by changes in

LSWT^[9]. For instance, dissolved oxygen concentrations in lakes decrease directly due to warmer water temperatures and indirectly due to enhanced thermal stratification (e.g., increase in surface water temperature with minimal change at deeper depth). Specifically, deoxygenation in lakes can arise due to a reduction in the solubility of dissolved oxygen (i.e., due to increasing water temperature) and warming-induced changes in the strength and duration of stratification. Notably, thermal stratification can reduce the supply of oxygen from the lake surface into deeper waters, where oxygen is consumed by respiration and remineralization processes ^[10]. Anoxic conditions at depth can subsequently lead to nutrient release from lake sediments, accelerating eutrophication and the production and emission of potent greenhouse gases such as methane^[11].

The catastrophic mortality of fish, benthic organisms and aquatic vegetation can be a direct result of LSWT increasing to extreme levels (i.e., values outside the norm)^[12]. The increase in LSWT may also lead to the advancement of phytoplankton phenology, which will increase the biomass of phytoplankton before the onset of the light-dependent growth of other aquatic vegetation, potentially enlarging the area and duration of algal blooms. This may lead to a shift from a clear state dominated by aquatic vegetation to a turbid state dominated by algae^[13]. In addition to these direct threats to lake ecosystems, subtle changes in physical or chemical processes within a lake as a result of LSWT modification can have significant and broader ecological impacts. The survival of aquatic life is threatened when the water level and the surface area of lakes are reduced, and oxygen is diminished^[14]. Following the extreme heat event, one of the worst droughts in around 60 years occurred in Yangtze River Basin. The drought was so intense that the water levels in China's largest freshwater lake, Lake Poyanghu,

Science Bulletin

dropped by 55.07% from 19.43 m in June to 8.73 m at the end of August (2022). Correspondingly, the water surface area plummeted from 4186 km² in August 2020 to 1117 km², leaving birds without habitats and aggravating fish die-off (Figure, S5)^[6]. Moreover, this extreme heat event directly affected the drinking water supply in Jiangxi and Hunan provinces and indirectly caused a declined freshwater flow into the ocean from the Yangtze River. This event triggered an extremely rare summer seawater intrusion, also known as a salty tide, which led to the close of water resource in Shanghai. If this positive feedback continues, the ecological balance will likely be destroyed, and the impact on aquatic life, as well as humans, could be far worse than anticipated^[15]. In summary, the extreme heat event in summer 2022 caused unprecedented warming of lakes in China, with an average LSWT increase of 1.63°C, which is considerably higher than the total increase in LSWT observed during the past 20 years. The effects were most apparent in Southwest and Yangtze River Basins where the warming was particularly pronounced. Increasing LSWT may lead to notable changes in the physical and chemical characteristics of lakes, such as an intensified thermal stratification, a decline in water level, a reduction of surface area, a decrease in dissolved oxygen, and an increase in greenhouse gas production and emission. Furthermore, these changes may have biological consequences such as phenological advancement and outbreaks of algal blooms and even catastrophic die-offs of lake biota. Given that extreme events induced by global climate change are projected to be more frequent in the future^[10], more attention should be paid to the response of lake systems to extreme events. In future research, high-frequency and real-time monitoring, experiments and modeling are needed to quantify the impacts of extreme events on lake ecosystems.

154 The authors declare that they have no conflicts of interest.

155 Author contributions

Weijia Wang: Conceptualization, Data curation, Methodology, Writing - original draft. Shi Kun: Supervision, Conceptualization, Methodology and Editing. Xiwen Wang: Data curation and Validation. Siqi Wang: Data curation, Methodology. Dong Zhang: Software, Validation. Yuanyuan Peng: Data curation, Software. Na Li: Data curation, Methodology, Validation. Yunlin Zhang: Methodology and Editing. Yibo Zhang: Methodology, Software, Validation. Boqiang Qin: Conceptualization, Validation and Editing. R. Iestyn Woolway: Supervision and Editing. Jeppesen Erik: Supervision and Editing. All authors reviewed the results and approved the final version of the manuscript.

164 Acknowledgments

This study was supported by the National Natural Science Foundation of China
(U22A20561 and 41922005), the Tibetan Plateau Scientific Expedition and Research Program,
(2019QZKK0202), the NIGLAS foundation (E1SL002), the Water Resource Science and
Technology Project in Jiangsu Province (2020057), the Key Research and Development
Program of Jiangsu Province (Social Development) (BE2022857), the TÜBITAK BIDEB2232
Program (118C250), and the UKRI Natural Environment Research Council (NERC)
Independent Research Fellowship (NE/T011246/1).

Science Bulletin

2		
3 4	173	References
5		
6 7 8	174	[1] Hegerl GC, Hanlon H, Beierkuhnlein C. Climate science elusive extremes. Nat
9 10 11	175	Geosci 2011; 4: 142-143.
12 13 14	176	[2] Williamson CE, Saros JE, Vincent WF, et al. Lakes and reservoirs as sentinels,
15 16 17	177	integrators, and regulators of climate change. Limnol Oceanogr 2009; 54: 2273-2282.
18 19 20	178	[3] Li X, Peng S, Deng X, et al. Attribution of lake warming in four shallow lakes in the
20 21 22	179	middle and lower yangtze river basin. Environ Sci Technol 2019; 53: 12548-12555.
23 24 25	180	[4] Anderson EJ, Stow CA, Gronewold AD, et al. Seasonal overturn and stratification
26 27 28	181	changes drive deep-water warming in one of earth's largest lakes. Nat Commun 2021; 12.
29 30 31	182	[5] Woolway RI, Dokulil MT, Marszelewski W, et al. Warming of central european lakes
32 33 34	183	and their response to the 1980s climate regime shift. Clim Change 2017; 142: 505-520.
35 36 37	184	[6] Mallapaty S. China's extreme weather challenges scientists trying to study it. Nature
38 39 40	185	2022; 609: 888-888.
41 42 43	186	[7] Boehrer B, Schultze M. Stratification of lakes. Rev Geophys 2008; 46.
44 45 46	187	[8] Zhan S, Song C, Wang J, et al. A global assessment of terrestrial evapotranspiration
40 47 48	188	increase due to surface water area change. Earths Future 2019; 7: 266-282.
49 50 51	189	[9] Liao F, Wang GC, Yang N, et al. Groundwater discharge tracing for a large ice-
52 53 54	190	covered lake in the tibetan plateau: Integrated satellite remote sensing data, chemical
55 56 57 58 59 60	191	components and isotopes. J Hydrol 2022; 609.

1 2

192 [10] Woolway RI, Kraemer BM, Lenters JD, et al. Global lake responses to climate change.

193 Nat Rev Earth Environ 2020; 1: 388-403.

194 [11] Walter KM, Zimov SA, Chanton JP, et al. Methane bubbling from siberian thaw lakes

as a positive feedback to climate warming. Nature 2006; 443: 71-75.

196 [12] Till A, Rypel AL, Bray A, et al. Fish die-offs are concurrent with thermal extremes

197 in north temperate lakes. Nat Clim Change 2019; 9: 637-641.

198 [13] Shi K, Zhang YL, Qin BQ, et al. Remote sensing of cyanobacterial blooms in inland

199 waters: Present knowledge and future challenges. Sci Bull 2019; 64: 1540-1556.

200 [14] Paerl HW, Huisman J. Climate - blooms like it hot. Science 2008; 320: 57-58.

201 [15] Qin BQ, Paerl HW, Brookes JD, et al. Why lake taihu continues to be plagued with

202 cyanobacterial blooms through 10 years (2007-2017) efforts. Sci Bull 2019; 64: 354-356.

Figure Captions Figure. 1. Spatial distribution of average summer air temperature between 2000 and 2021(a1)

217	and during the summer of 2022 (a2) across China. Spatial distribution of local surface air
218	temperature (SAT) during summer between 2000 and 2021(a3) and during the summer of 2022
219	(a4) across the country. (a5-a6) Anomalies of air temperature and SAT in summer 2022. (b1-
220	b10) The temporal variation in the average summer SAT for the nine basins and the whole
221	country.
222	Figure. 2. (a1) Spatial distribution of lake surface water temperature (LSWT) in summer 2022
223	and (a2) the anomalies of the LSWT in summer 2022 compared to those between 2000 and
224	2021. (b1-b10) The temporal variation in average summer LSWT for the nine basins and the
225	whole country. Histograms show the differences in summer mean LSWT (c1) and the decadal
226	change rate of LSWT (c2) between 2000 – 2021 and 2022 for the nine basins and the national
227	scale.
228	
229	



230

231



www.scibull.com



24 Supplementary materials

25 Materials and methods

In this study, we made lake surface water temperature (LSWT) observations of large lakes in China between 2000 and 2022 from the Terra Moderate Resolution Imaging Spectroradiometer (MODIS-terra) land surface temperature/emissivity 8-day product (MOD11A2 version 6.1) provided by NASA's Earth Observing System Data and Information System (EOSDIS, https://earthdata.nasa.gov). The daytime land surface temperature (LST) (approximately 10:30 local time), nighttime LST (approximately 22:30 local time), and corresponding quality control (QC) images were extracted from MODIS LST images after resampling to a 1 km pixel size. The average of the daytime LST and nighttime LST was taken as the daily LST for this study, before which pixels of poor quality were excluded using OC images^[1]. We discard these data and set them to null if less than half the total lake pixels remained after removing those deemed poor quality. The variation between MODIS temperature products and in situ measured water temperature has been verified to be less than 2 °C^[2, 3]. Meteorological data were obtained from the National Centers for Environmental Information (NCEI) (https://www.ncei.noaa.gov/) by inverse distance weight interpolation of the daily site data.

Polygons provided by HydroLAKES (https://www.hydrosheds.org/pages/hydrolakes)
were used to determine the lakes (≥ 100 km²) investigated in our study. The global surface
water occurrence (GSWO) dataset, which provides the probability of the presence of surface
water between 1984 and 2020 at a spatial resolution of 30 m, was utilized within the lake

Page 29 of 47

Science Bulletin

boundaries of HydroLAKES to determine the permanent water surface extent of the studied lakes^[4]. To eliminate boundary effects, i.e., pixels that may be a mixture of land and water, the boundary obtained by the above treatment was buffered inward by 1 km. Excluding lakes that dried up and bays connected to the ocean that were misidentified, 119 large lakes were confirmed for this study. Nine basin polygons in China (namely, Songhua and Liao Basin, Inland River Basin, Yellow River Basin, Hai River Basin, Huai River Basin, Yangtze River Basin, Southeast River Basin, Pearl River Basin, and Southwest River Basin) were obtained from the Resource and Environment Science and Data Center (https://www.resdc.cn/). Water levels in Lake Poyanghu was obtained from the Poyang Lake Wetland Observation and Research Station (http://pyl.cern.ac.cn/). The water surface extent data of Poyang Lake for the summer of 2000 – 2022 was obtained by modified normalized difference water index (MNDWI) and calculated using Landsat^[5]. The Ostu algorithm was used to determine the threshold^[6]. Due to the low temporal resolution and susceptibility to weather, we took the median value of all usable Landsat images from June to August of each year to calculate the water surface extent for Lake Poyanghu.

LSWT data from the MOD11A2 and GSWO datasets were obtained and processed through the Google Earth Engine (GEE, https://earthengine.google.com/). A lake specific LSWT in our study was calculated as the average temperature of all 1 km pixels within the lake. The original vacancies and the vacant data after quality control were filled using a random forest regression model with five input parameters: year, day of year, longitude, latitude, and elevation^[7, 8]. The result of the model was satisfactory, with the error between the estimated and observed values controlled at approximately 2 °C (Figure. S1). Air temperatures (ATs) were

67	acquired by inverse distance interpolation to 1 km spatial resolution images using station data
68	in China from the NCEI dataset. The average local surface air temperature (SAT) was
69	calculated similarly to LSWT, which was obtained by extracting all available pixels of each
70	lake from the AT dataset and calculating the mean value. The anomalies in AT, LSWT and
71	SAT were calculated as the difference between the 2022 values and the averages of 2000 -
72	2021. On the basis of the reconstructed LSWT, AT and SAT dataset, the long-term variation in
73	the study area was analyzed using the Mann-Kendall trend analysis approach ^[9, 10] .
74	
75	Deferences
75	References
76	[1] Guo L, Zheng H, Wu Y, et al. An integrated dataset of daily lake surface water temperature
77	over the Tibetan plateau. Earth Syst Sci Data 2022; 14: 3411-3422.
78	[2] Xie C, Zhang X, Zhuang L, et al. Analysis of surface temperature variation of lakes in
79	China using MODIS land surface temperature data. Sci Rep 2022; 12: 2415.
80	[3] Wang R, Yan X, Niu ZG, et al. Long-term changes in inland water surface temperature
81	across China based on remote sensing data. J Hydrometeorol 2021; 22: 523-532.
82	[4] Pekel JF, Cottam A, Gorelick N, et al. High-resolution mapping of global surface water
83	and its long-term changes. Nature 2016; 540: 418-422.
84	[5] Li YZ, Gong XQ, Guo Z, et al. An index and approach for water extraction using Landsat-
85	OLI data. Int J Remote Sens, 2016; 37: 3611-3635.

86 [6] Xu HQ. Modification of normalised difference water index (NDWI) to enhance open water

3	
4	
5	
0 7	
, 8	
9	
10	
11	
12 13	
14	
15	
16	
17	
18 19	
20	
21	
22	
23 24	
25	
26	
27	
28 29	
30	
31	
32	
33 34	
35	
36	
37	
39	
40	
41	
42 43	
44	
45	
46	
47 49	
40 49	
50	
51	
52 52	
⊃3 54	
55	
56	
57	
58	

59 60

- 87 features in remotely sensed imagery. Int J Remote Sens, 2006; 27: 3025-3033.
- [7] Breiman L. Random forests. Mach Learn 2001; 45: 5-32. 88
- 89 [8] Woolway RI, Jennings E, Shatwell T, et al. Lake heatwaves under climate change. Nature

90 2021; 589: 402-418.

91 [9] Kendall MG. Rank correlation methods, 2nd ed. 1955,

Int 92 [10] Mann HB. Nonparametric tests against trend. Econometrica 1945; 13: 245-259.
 Table S1. Statistics of AT comparison between summer 2022 and summer 2000 – 2021 by month.

		2000 – 2021			2022				
		Mean AT	lean AT Max AT		Number of	Mean AT	Max AT	Number of	Number of
		(°C)	(°C)	days above	days above	(°C)	(°C)	days above	days above
				30 °C (d)	35 °C (d)			30 °C (d)	35°C (d)
	Southeast River Basin	26.03	32.27	10.82	0.00	25.89	33.33	7.00	0.00
	Hai River Basin	23.34	32.75	5.77	0.09	24.17	35.89	11.00	1.00
June	Huai River Basin	24.79	32.60	6.00	0.09	26.99	35.67	16.00	1.00
	Yellow River Basin	19.29	32.28	5.05	0.05	20.95	34.44	14.00	0.00
	Inland River Basin	21.01	36.80	23.64	4.45	22.29	39.89	28.00	15.00
	Songhua and Liao River Basin	20.28	30.64	1.45	0.00	19.74	31.61	1.00	0.00
	Southwest River Basin	18.10	33.52	15.14	0.50	17.79	32.11	5.00	0.00

 Science Bulletin

	Yangtze River Basin	22.20	33.03	10.41	0.00	22.97	32.83	16.00	0.00
	Pearl River Basin	27.10	32.65	14.95	0.00	26.89	31.72	13.00	0.00
	China	22.46	36.91	24.50	4.68	23.08	39.89	28.00	15.00
	Southeast River Basin	28.60	33.80	21.41	0.09	29.89	35.72	25.00	3.00
	Hai River Basin	25.15	33.20	8.64	0.09	25.18	32.78	7.00	0.00
	Huai River Basin	27.00	33.54	10.95	0.05	27.67	33.67	14.00	0.00
	Yellow River Basin	21.16	33.14	8.23	0.14	21.32	33.94	7.00	0.00
Inly	Inland River Basin	23.07	38.21	27.77	10.68	23.69	40.00	30.00	16.00
July	Songhua and Liao River Basin	22.91	31.23	2.14	0.00	23.40	29.11	0.00	0.00
	Southwest River Basin	18.45	32.98	10.36	0.09	19.36	33.44	14.00	0.00
	Yangtze River Basin	24.48	34.62	22.32	0.50	25.78	36.39	25.00	5.00
	Pearl River Basin	27.90	33.35	17.50	0.00	28.59	34.78	22.00	0.00

	China	24.30	38.25	28.50	10.73	24.99	40.00	30.00	16.00
	Southeast River Basin	28.17	33.11	16.68	0.09	29.86	35.56	26.00	5.00
	Hai River Basin	23.65	31.18	3.09	0.00	23.93	33.22	12.00	0.00
	Huai River Basin	26.31	32.55	7.09	0.05	27.83	35.06	21.00	1.00
	Yellow River Basin	19.59	31.77	4.27	0.00	21.09	34.44	14.00	0.00
	Inland River Basin	21.49	36.70	23.27	4.91	21.59	39.11	22.00	11.00
August	Songhua and Liao River	21.27	29.94	0.86	0.00	20.31	30.61	2.00	0.00
	Basin								
	Southwest River Basin	18.21	31.98	7.18	0.00	19.42	32.00	9.00	0.00
	Yangtze River Basin	23.90	34.53	17.32	0.86	26.93	37.83	30.00	19.00
	Pearl River Basin	27.54	32.92	16.05	0.00	27.85	34.78	15.00	0.00
	China	23.35	36.80	24.00	5.05	24.31	39.11	30.00	19.00
Summe	Southeast River Basin	27.60	34.01	48.91	0.18	28.55	35.72	58.00	8.00

 Science Bulletin

r	Hai River Basin	24.05	33.48	17.50	0.18	24.43	35.89	30.00	1.0
	Huai River Basin	26.03	34.00	24.05	0.18	27.50	35.67	51.00	2.0
	Yellow River Basin	20.01	33.62	17.55	0.18	21.12	34.44	35.00	0.0
	Inland River Basin	21.86	38.45	74.68	20.05	22.52	40.00	80.00	42.
	Songhua and Liao River Basin	21.49	32.03	4.45	0.00	21.15	31.61	3.00	0.0
	Southwest River Basin	18.25	33.92	32.68	0.59	18.85	33.44	28.00	0.0
	Yangtze River Basin	23.53	35.05	50.05	1.36	25.23	37.83	71.00	24.
	Pearl River Basin	27.51	33.50	48.50	0.00	27.78	34.78	50.00	0.0
	China	23.37	38.45	74.68	20.05	24.12	40.00	80.00	42.

www.scibull.com

96 Table S2. Statistics of average SAT comparison between summer 2022 and summer 2000 –

97 2021 by month.

		200	0 - 2021				2022	
Basin	June	July	August	Summer	June	July	August	Summer
Southeast River	24 50	28.12	27.68	26 77	25 32	30.70	31 32	29.11
Basin	24.30	20.12	27.00	20.77	23.32	50.70	51.52	29.11
Hai River Basin	23.42	25.51	24.22	24.38	23.66	25.84	24.32	24.61
Huai River Basin	25.06	27.52	26.74	26.44	27.36	28.39	28.59	28.11
Yellow River Basin	17.02	19.10	17.94	18.02	18.44	19.35	19.95	19.25
Inland River Basin	14.45	16.20	15.36	15.33	15.55	17.11	16.86	16.51
Songhua and Liao	20.60	23 12	21.32	21.68	20.11	23 59	20.23	21 31
River Basin	20.00	23.12	21.52	21.00	20.11	23.37	20.25	21.51
Southwest River	14 17	15 19	14 57	14 64	14 75	16 23	16.06	15 68
Basin					5			
Yangtze River	22.91	25 79	25 10	24 60	24.01	27.05	27.88	26 32
Basin	, .	_0.17	_0.10			_,	27.00	_0.02
Pearl River Basin	24.80	25.42	25.08	25.10	24.21	26.50	25.74	25.48
China	17.54	19.55	18.70	18.59	18.55	20.49	20.36	19.80

4

Science Bulletin

5	
6	
7	
8	
9	
10	
11	
12	
13	
14	
14	
15	
16	
17	
18	
19	
20	
21	
22	
23	
24	
25	
26	
27	
27	
20	
29	
30	
31	
32	
33	
34	
35	
36	
37	
38	
30	
40	
40	
41 42	
42	
43	
44	
45	
46	
47	
48	
49	
50	
51	
52	
52	
55	
54	
22	
56	
57	
58	
59	
60	

Table S3. Statistics of LSWT comparison between summer 2022 and summer 2000 – 2021 by

101 month.

		2000 -	- 2021	202	22
		Mean	Max	Mean	Max
		LSWT	LSWT	LSWT	LSWT
	Southeast River Basin	23.37	25.14	22.97	25.52
	Hai River Basin	23.19	25.41	22.46	25.36
	Huai River Basin	24.09	26.48	26.17	28.10
	Yellow River Basin	17.26	27.16	18.29	28.32
Iuno	Inland River Basin	10.12	25.71	11.54	28.74
June	Songhua and Liao River Basin	20.07	23.71	20.06	24.38
	Southwest River Basin	9.65	20.49	10.61	18.78
	Yangtze River Basin	21.88	27.60	22.74	28.96
	Pearl River Basin	21.87	26.18	21.44	26.05
	China	19.05	27.60	19.59	28.96
	Southeast River Basin	26.65	28.16	28.80	29.33
	Hai River Basin	25.26	27.12	26.04	27.05
Inly	Huai River Basin	26.07	28.44	27.31	28.87
July	Yellow River Basin	19.82	28.14	20.07	27.78
	Inland River Basin	12.85	27.50	14.24	29.28
	Songhua and Liao River Basin	22.74	25.33	23.56	25.75

	Southwest River Basin	10.23	20.77	12.51	2
	Yangtze River Basin	24.36	29.67	25.85	3
	Pearl River Basin	22.83	27.52	24.37	2
	China	21.20	29.67	22.53	3
	Southeast River Basin	26.65	27.88	29.00	2
	Hai River Basin	24.37	26.31	25.57	2
	Huai River Basin	25.49	27.56	28.20	3
	Yellow River Basin	19.02	26.69	20.86	2
A	Inland River Basin	12.88	26.25	14.49	2
August	Songhua and Liao River Basin	21.29	24.20	21.47	2
	Southwest River Basin	10.96	20.98	13.15	2
	Yangtze River Basin	24.00	29.36	26.30	3
	Pearl River Basin	23.02	27.21	23.89	2
	China	20.85	29.36	22.55	3
	Southeast River Basin	25.56	28.33	26.92	2
	Hai River Basin	24.27	27.44	24.69	2
Summe	Huai River Basin	25.22	28.65	27.23	3
r	Yellow River Basin	18.70	28.49	19.74	2
-	Inland River Basin	11.95	27.62	13.42	29
	Songhua and Liao River Basin	21.37	25.60	21.69	2:
	Southwest River Basin	10.28	21.45	12.09	2

1 2					
3 <u> </u>	Yangtze River Basin	23.41	30.04	24.96	32.24
6 7	Pearl River Basin	22.57	27.72	23.23	29.73
8 9 10	China	20.37	30.04	21.55	32.24
11 12 102 13					
14 15 103					
16 17 18					
19 20 21					
22 23					
24 25 26					
27 28 20					
30 31					
32 33 34					
35 36					
37 38 39					
40 41					
42 43 44					
45 46					
47 48 49					
50 51 52					
52 53 54					
55 56					
57 58 59					
60		13			

Table S4. Comparison of rate of change in SAT and LSWT in 2000 – 2021 with 2000-2022

105	for the nine basins and China.	

	Rate of SA	T change	Rate of LS	WT change
	(°C/de	cade)	(°C/de	ecade)
Basin	2000 - 2021	2000 - 2022	2000 - 2021	2000 - 2022
Southeast River Basin	0.23	0.30	-0.08	0.05
Hai River Basin	0.63	0.75	0.15	0.21
Huai River Basin	0.11	0.12	0.36	0.47
Yellow River Basin	0.18	0.31	0.05	0.15
Inland River Basin	0.07	0.20	0.04	0.17
Songhua and Liao River Basin	0.29	0.36	0.18	0.18
Southwest River Basin	-0.16	-0.21	0.09	0.24
Yangtze River Basin	0.43	0.47	0.20	0.30
Pearl River Basin	0.13	0.24	0.36	0.39
China	0.32	0.32	0.11	0.22

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
10
10
20
∠∪ ว1
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
40
47
48
49 50
50
51
52
53
54
55
56
57
58

Table S5. Slope and <i>p</i> -value of the linear relationship of the change in the summer average

109 LSWT and SAT during 2000 – 2022.

	S	AT	LSWT	
Basin	Slope	<i>p</i> -value	Slope	<i>p</i> -value
Southeast River Basin	0.02	<i>P</i> < 0.01	0.08	<i>P</i> < 0.01
Hai River Basin	0.02	<i>P</i> < 0.01	0.01	<i>P</i> < 0.01
Huai River Basin	0.05	<i>P</i> < 0.01	0.03	<i>P</i> < 0.01
Yellow River Basin	0.01	<i>P</i> < 0.01	0.02	<i>P</i> < 0.01
Inland River Basin	0.02	<i>P</i> < 0.01	0.04	<i>P</i> < 0.01
Songhua and Liao River Basin	0.02	<i>P</i> < 0.01	-0.01	<i>P</i> < 0.01
Southwest River Basin	0.02	<i>P</i> < 0.01	0.05	<i>P</i> < 0.01
Yangtze River Basin	0.04	P < 0.01	0.03	<i>P</i> < 0.01
Pearl River Basin	0.04	P < 0.01	0.03	<i>P</i> < 0.01
China	0.02	P < 0.01	0.03	<i>P</i> < 0.01

110

111



Figure S1. Performance of the random forest regression model for LSWT supplemented with

114 missing data (daytime (a-c), nighttime (d-f)). The green circle represents the training dataset,

and the red circle represents the test dataset.



(b)

Nº0t

Nº08







Figure S5. Variation of water level and water surface area of Lake Poyanghu in summer from
2000 to 2022 (a). Landsat images of Lake Poyanghu, the largest freshwater lake in China, on
August 27, 2020 (b) and August 27, 2022 (c). Migratory birds gathering in a small puddle in
the drought-stricken lake in August 2022 (d). Dead fish in drought-stricken lakebed (e). Photos

1		
2		
4	136	d and e are from reports on the China News Network (https://www.chinanews.com.cn/).
5		
6	137	
/ 8		
9		
10		
11		
12		
14		
15		
16		
1/ 18		
19		
20		
21 22		
22 23		
24		
25		
26 27		
27		
29		
30		
31		
33		
34		
35		
30 37		
38		
39		
40 41		
41		
43		
44		
45 46		
47		
48		
49 50		
50 51		
52		
53		
54 55		
ээ 56		
57		
58		
59 60		
00		21

