

#### **Potential sources and occurrence of macro-plastics and microplastics pollution in farmland soils: A typical case of China**

Ren, Siyang; Wang, Kai; Zhang, Jinrui; Li, Jingjing; Zhang, Hanyue; Qi, Ruimin; Xu, Wen; Yan, Changrong; Liu, Xuejun; Zhang, Fusuo; Jones, Davey L.; Chadwick, Dave R

#### **Critical Reviews in Environmental Science and Technology**

DOI:

[10.1080/10643389.2023.2259275](https://doi.org/10.1080/10643389.2023.2259275)

Published: 02/04/2024

Peer reviewed version

[Cyswllt i'r cyhoeddiad / Link to publication](https://research.bangor.ac.uk/portal/en/researchoutputs/potential-sources-and-occurrence-of-macroplastics-and-microplastics-pollution-in-farmland-soils-a-typical-case-of-china(7404b0e6-5ffd-49fa-98ee-09476ef501c7).html)

Dyfyniad o'r fersiwn a gyhoeddwyd / Citation for published version (APA): [Ren, S.](https://research.bangor.ac.uk/portal/en/researchers/siyang-ren(a4e1aa4f-3428-4287-bf74-0e1229ce914b).html), Wang, K., Zhang, J., Li, J., Zhang, H., Qi, R., Xu, W., Yan, C., Liu, X., Zhang, F[., Jones,](https://research.bangor.ac.uk/portal/en/researchers/davey-jones(247f4973-4af0-4656-9ec0-e008f86111cb).html) [D. L.](https://research.bangor.ac.uk/portal/en/researchers/davey-jones(247f4973-4af0-4656-9ec0-e008f86111cb).html)[, & Chadwick, D. R.](https://research.bangor.ac.uk/portal/en/researchers/dave-chadwick(1b57f584-5a03-45dd-b9af-3c4b0d1b4a7e).html) (2024). [Potential sources and occurrence of macro-plastics and](https://research.bangor.ac.uk/portal/en/researchoutputs/potential-sources-and-occurrence-of-macroplastics-and-microplastics-pollution-in-farmland-soils-a-typical-case-of-china(7404b0e6-5ffd-49fa-98ee-09476ef501c7).html) [microplastics pollution in farmland soils: A typical case of China](https://research.bangor.ac.uk/portal/en/researchoutputs/potential-sources-and-occurrence-of-macroplastics-and-microplastics-pollution-in-farmland-soils-a-typical-case-of-china(7404b0e6-5ffd-49fa-98ee-09476ef501c7).html). Critical Reviews in Environmental Science and Technology, 54(7), 533-556. <https://doi.org/10.1080/10643389.2023.2259275>

#### **Hawliau Cyffredinol / General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

 • Users may download and print one copy of any publication from the public portal for the purpose of private study or research.

- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

#### **Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



4 Siyang Ren<sup>a,b</sup>, Kai Wang<sup>a\*</sup>, Jinrui Zhang<sup>a,c</sup>, Jingjing Li<sup>a</sup>, Hanyue Zhang<sup>a,d</sup>, Ruimin Qi<sup>e</sup>, Wen

- 5 Xu<sup>a</sup>, Changrong Yan<sup>f,g</sup>, Xuejun Liu<sup>a,c\*</sup>, Fusuo Zhang<sup>a</sup>, Davey L. Jones<sup>b,h</sup>, David R. Chadwick<sup>b</sup>
- 
- *<sup>a</sup> College of Resources and Environmental Sciences; National Academy of Agriculture Green*
- *Development; Key Laboratory of Plant-Soil Interactions of Ministry of Education; National*
- *Observation and Research Station of Agriculture Green Development (Quzhou, Hebei), China*
- *Agricultural University, Beijing, 100193, P. R. China*
- *<sup>b</sup> School of Natural Sciences, Bangor University, Bangor, LL57 2UW, UK*
- *<sup>c</sup> College of Resources and Environmental Sciences; Beijing Key Laboratory of Farmland Soil*
- *Pollution Prevention and Remediation, China Agricultural University, Beijing, 100193, P. R.*
- *China*
- *<sup>d</sup> Soil Physics and Land Management Group, Wageningen University & Research, Wageningen,*
- *6700 AA, the Netherlands*
- *<sup>e</sup> School of Environmental Science and Engineering, Southern University of Science and*
- *Technology, Shenzhen 518055, P. R. China*
- *<sup>f</sup> Institute of Environment and Sustainable Development in Agriculture, Chinese Academy of*
- *Agricultural Sciences, Beijing 100081, P. R. China*
- *<sup>g</sup> Key Laboratory of Prevention and Control of Residual Pollution in Agricultural Film, Ministry*
- *of Agriculture and Rural Affairs, Beijing 100081, P. R. China*
- *hSoilsWest, Centre for Sustainable Farming Systems, Food Futures Institute, Murdoch*

*University, Murdoch, WA 6105, Australia*



#### **Potential sources and occurrence of macro-plastic and microplastics pollution in farmland**

## **soils: A typical case of China**

#### <span id="page-3-0"></span>**Abstract**

 Plastic debris (including macro-plastics, microplastics (MPs), and nanoplastics), defined as an emerging contaminant, has been proven to significantly affect soil ecosystem functioning. Accordingly, there is an urgent need to robustly quantify the pollution situation and potential sources of plastics in soils. China as the leading producer and user of agricultural plastics is analysed as a typical case study to highlight the current situation of farmland macro-plastics and MPs. Our study summarized information on the occurrence and abundance of macro-plastics and MPs in Chinese farmland soils for the first time based on 163 publications with 728 sample sites. The results showed that the average concentration of macro-plastics, and the abundance of MPs in Chinese farmlands 45 were 103 kg ha<sup>-1</sup> and 4537 items  $kg^{-1}$  (dry soil), respectively. In addition, this study synthesized the latest scientific evidence on sources of macro-plastics and MPs in farmland soils. Agricultural plastic films and organic wastes are the most reported sources, indicating that they contribute significantly to plastic debris in agricultural soils. Furthermore, the modelling methods for quantifying macro-plastics and MPs in soils and estimating the stock and flow of plastic materials within agricultural systems were also summarized.

 Keywords: macro-plastics, microplastics, abundance, source apportionment, quantitative method, farmland soils

# **Graphical abstract**



 





#### <span id="page-7-0"></span>**1. Introduction**

 Plastics are widely used across almost all sectors of society due to their versatility relatively cheap cost, light weight and durability (Jambeck et al., 2015; Plastics Europe-The Facts 2022). The production of plastics is increasing with global cumulative production predicted to reach up to 33 billion tons by 2050 (Sharma et al., 2020). As most plastics have a relatively short functional lifespan, the disposal of plastics represents a major global problem with a large proportion of plastics not being recycled (Luan et al., 2021; Wang et al., 2021). Currently, it is estimated that 79% of plastic waste enters either landfills or the natural environment where it represents a threat to terrestrial, freshwater, and marine ecosystems (Geyer et al., 2017). Although macro-plastics (particles size > 5 mm) represent the primary type of waste entering the environment, they gradually degrade into smaller fragments in response to ultraviolet (UV) irradiation, mechanical abrasion, and biodegradation (Barnes et al., 2009; Yang et al., 2022). Microplastics (MPs) are defined as particles < 5 mm and > 1 μm, including fragments, fibers, particles, foams, and films, 106 while plastic particles with the size between 1 nm and 1  $\mu$ m are defined as nanoplastics (NPs) (Frias & Nash, 2019; Thompson et al., 2004). Since MPs and NPs are small in size, and present in large quantities and degrade slowly, they are easily absorbed, inhaled, or ingested by organisms, leading to bioaccumulation (Barnes et al., 2009; Leslie et al., 2022; Wu et al., 2022). Studies have found the presence of MPs and NPs in plants, soil fauna, human feces, and blood (Leslie et al., 2022; Li et al., 2020; Lwanga et al., 2017; Zhang et al., 2021). Current evidence indicates that MPs can be transferred to the human body through the food chain as well as via inhalation and are likely to give rise to a range of cytotoxic effects that are now becoming evident, albeit still incomplete (Wu et al., 2022; Brachner et al., 2020; Hua et al., 2022).

 Previous reports have focused mainly on aquatic ecosystems with MPs as an emerging contaminant (Cozar et al., 2014; Rochman et al., 2016). Recently, researchers have expanded their focus to terrestrial environments (Kumar et al., 2020; Li et al., 2020). Our recent meta-analysis study quantified the effect of plastic residues and MPs on indicators of global soil ecosystem functioning (i.e. soil physicochemical properties, plant and soil animal health, abundance, and diversity of soil microorganisms) (Zhang et al., 2022). The results showed that plastic residues and MPs can alter plant growth and soil physicochemical properties. For example, plastic residues and MPs decreased root biomass, plant height, soil dissolved organic carbon, and soil total nitrogen (N) content by 14%, 13%, 9%, and 7%, respectively (Zhang et al., 2022). It should be noted, however, that significant bias may occur in meta-analyses as neutral/non-significant results are often not reported in the literature (Coursol & Wagner, 1986). Further, many studies investigating the effect of MPs on soil ecosystem responses have used high rates of MPs contamination in soil (> 1% w/w) (Meng et al., 2021; Ng et al., 2021), which may not reflect levels (much lower than 1% w/w) that occur in typical agronomic field conditions (Huang et al., 2020; Liu et al. 2019). In addition, plastic contaminants and their additives have been shown to inhibit the growth and development of soil animals. The decrease in growth rate, movement rate (e.g. frequency of body bending and head thrashing), feeding rate, and reproduction rate reveals the disturbed locomotor behaviors of animals caused by plastic residues and MPs (Zhang et al., 2022; Wang et al., 2021). Furthermore, several studies have reported that plastic 134 particles within the size of 0.08–2.00  $\mu$ m (i.e. NPs) can penetrate the stele of rice, cucumber, wheat, and lettuce, leading to efficient uptake of smaller plastic particles (Li et al., 2021; Liu et al., 2022). It indicates that MPs and NPs can be transferred to livestock and the human body

 through the terrestrial food chain, causing a potential threat to livestock and human health, and natural ecosystem food webs (Lwanga et al., 2017; Zhou et al., 2021).

139 MPs can also be vectors for the attachment and transmission of other contaminants (e.g. hydrophobic organic contaminants, heavy metals, harmful microorganisms), posing a potential threat to (human, livestock, soil fauna) organismal health and the wider environment (Brennecke et al., 2016; Wang et al., 2019; Zettler et al., 2013). For example, the heavy metals 143 cadmium (Cd) and lead (Pb) were detected in MPs samples  $(n = 924)$  from two beaches in Southwest England with maximum concentrations of 3.4 and 5.3 mg  $g^{-1}$ , respectively (Massos & Turner, 2017). In addition, they evaluated the maximum bio-accessible concentrations of Cd and Pb in the proventriculus-gizzard of seabirds (*Fulmarus glacialis*) and found concentrations 147 exceeded the safe dietary intake limit by a factor of about 50 and 4, respectively (Massos & Turner, 2017). Furthermore, high concentrations of zinc (Zn, 9407 mg kg<sup>-1</sup>) and polycyclic 149 aromatic hydrocarbons (PAH, 47 mg kg<sup>-1</sup>) have been detected in MPs samples recovered from earthworms and the surrounding soil, and MPs exposure resulted in the steep rise of the abundance of pathogenic microorganisms in the worm intestinal tract (Ding et al., 2020).

 Many studies have indicated that the additives contained in MPs may represent a greater threat to terrestrial ecosystems than the plastic polymer itself (Hahladakis et al., 2018; Halden, 2010). There have been very few studies on the effects of MPs on human health, however, some of the additives used in plastics manufacturing, e.g. plasticizers and heavy metals, have been shown to interfere with gene expression, cell metabolism (e.g. signal transduction, enzyme function), and animals and humans development as well as reproduction (e.g. endocrine disrupting properties) (Rist et al., 2018). For instance, the common plasticizer of bisphenol A  (BPA), as an endocrine disruptor, could disrupt the endocrine system and various functions of organisms, including the thyroid, reproductive system, and metabolism (Halden, 2010).

 As plastic waste (including macro-plastics and MPs) in terrestrial environments poses a potential threat to food security, human health, and the health of our natural environment, it is important to control plastic input and manage legacy plastic in soils. Greater effort is needed to quantify the sources of macro-plastics and MPs and the fate of different plastic fragments. A study by Jambeck et al. (2015) reported that 80% of marine plastic residuals arise from land, suggesting that soil is not only an important sink of MPs but also an important source (Rachman, 2018). The accumulation of macro-plastics and MPs waste in soils is the result of various human activities and environmental origins, such as discarded plastic litter (Rillig, 2012), plasticulture practices (e.g. plastic mulch films, greenhouse films, irrigation pipes, and associated infrastructure) (Bläsing & Amelung, 2018, Gündoğdu et al., 2022; Huang et al., 2020; Wang et al., 2022), sewage sludge application (Long et al., 2019), coated fertilizers (Katsumi et al., 2021), organic fertilizer and agricultural compost (Weithmann et al., 2018), atmospheric deposition (Allen et al., 2019), digested food waste (Porterfield et al., 2023), and rubber tire wear (Evangeliou et al., 2020). Of these, plastic films represents an important source of macro- plastics and MPs in agricultural soils and has attracted extensive research and discussion (Qi et al., 2020). Plastic films are used extensively throughout the world in both horticulture and arable cropping (e.g. mulch films and polytunnels) as well as within livestock production (e.g. silage wrapping). Typically, these plastics are not recycled efficiently for several reasons including (i) difficulty in removing them from the soil after use, (ii) contamination by soil and vegetation residues, (iii) loss from the field due to wind erosion, (iv) lack of recycling  infrastructure, and (v) poor financial incentives (Li et al., 2021; Mekonnen et al., 2016). These barriers to recycling have led to the accumulation of legacy plastic in soils.

 An increasing body of research has focused on the abundance and distribution, migration pathways, and ecological environmental impact of MPs. Several recent reviews have covered these topics, but there is little information about the sources of macro-plastics and MPs, generation rates of MPs, and movement of MPs in and through the soil environment, especially for farmland soils (Qi et al., 2018; Yang et al., 2021). At the current rate of increase in plastic production between 2020 and 2021 (about 4%) (Plastics Europe-The Facts 2022), understanding the sources and consequences of macro-plastics and MPs represents a priority in terms of understanding the potential risks as well as the design and implementation of effective mitigation 191 strategies. Therefore, the aims of this review are to provide the latest understanding of the sources, abundance, and distribution of both macro-plastics and MPs in agricultural farmlands with a focus on quantitative methods and knowledge gaps. Furthermore, China as an example, which has become the world's biggest consumer and disposer of plastic films to be mapped the different sources and abundance of macro-plastics and MPs in the Chinese farmland soils based on the data from published literature (Figure 1 and 2).

#### <span id="page-11-0"></span>**2. Distribution and potential source of macro-plastics and MPs in Chinese farmlands**

 China is suffering from serious plastic pollution, particularly within agricultural environments where recycling is problematic (Plastics Europe-The Facts 2022; Qi et al., 2020). In 2021, the use of plastic mulch film (PMFs) in China was 1.3 million tons, representing 75% of global plastic agri-use (NBSC, 2019; Yan, 2022). Further, the area covered by plastic mulch film in China in  2021 was 17.3 million hectares, equivalent to 70% of the area of the United Kingdom (NBSC, 2022). However, the recovery rate of plastic mulch film from the field at the end of cropping was < 60% (Zhao et al., 2017). Because of the low plastic film thickness of 6‒8 μm commonly used in China, it is difficult to completely retrieve it from soil (Yan et al., 2006), and what is retrieved is contaminated with soil, limiting opportunities for recycling. Based on previous studies, summary information on the occurrence and potential sources of macro-plastics and MPs is important to control them in agricultural soils.

#### <span id="page-12-0"></span>*2.1 Literature search and data collection*

 To understand the distribution of soil macro-plastics and MPs in China's agricultural soils, this review searched the literature from three scientific databases (i.e. Web of Science, EI Compendex, and China Knowledge Resource Integrated Database) for "search terms" including (plastic residue or plastic debris) and (macro-plastic or microplastic or nanoplastic) and (soil or terrestrial) (specific search strings in Table S1). The literature search was limited to papers published before 217 the  $31<sup>st</sup>$  of January 2022. In summary, these papers were chosen according to the following selection criteria: (a) the study must be practical measurement data, without extra addition of non- agricultural plastics and MPs; (b) these sampling data must have a specific location name (to cities or counties) or longitude and latitude; (c) the samples must be collected in the farmland soil of China; (d) must be collected in bare soil, rather than in the greenhouse. Finally, 163 articles (including 123 studies for macro-plastics and 40 studies for MPs) were selected from more than 4,800 publications using strict inclusion criteria (Table S2 and S3). Details of search strings and the number of publications were presented in Table S1 and S4 in Supporting Information. The

 database covered 30 provinces with 728 data points to show the situation of macro-plastics and MPs in Chinese agricultural soils (Figure 1 and 2). However, there is a limitation that due to the longitude and latitude of some points being located close together, they overlapped and appeared to only be one location presented in Figure 2. This situation was shown in Jiangsu (data num = 5), Guangdong (13), Jilin (7), Hebei (16), Heilongjiang (4) and Inner Mongolia (13) Province.

#### <span id="page-13-0"></span>*2.2 Distribution and occurrence of macro-plastics and MPs in Chinese farmlands soil*

<span id="page-13-1"></span>*2.2.1 The occurrence of macro-plastics in Chinese farmlands soil*

 The concentration of macro-plastics was investigated in 24 provinces of China collected from 123 articles (see the list of articles in Table S2) as shown in Figure 1. The concentration of 236 macro-plastics found in soil  $(0 - 80 \text{ cm depth})$  varied from 0.2 to 421.6 kg ha<sup>-1</sup> with an average 237 value of 103.3 kg ha<sup>-1</sup>, with the median of 54.7 kg ha<sup>-1</sup>. The province's highest concentration (421.6 kg ha<sup>-1</sup>) was found in Xinjiang province (Northwestern China). The highest average 239 concentration of macro-plastics  $(142.7 \text{ kg ha}^{-1})$  was found in Northwestern China (e.g. Xinjiang, 240 Ningxia, and Gansu province), followed by 37.4 kg ha<sup>-1</sup> in Northern China (e.g. Hebei, 241 Shandong, and Shanxi province), and  $30.7 \text{ kg}$  ha<sup>-1</sup> in the soils of Southern China (e.g. Hubei, Yunnan province and Shanghai) (Table S2). This result was consistent with previous results that investigated 384 soil samples collected from 19 provinces, and the macro-plastics concentrations 244 in the typical areas covered with mulch films soil samples ranged from 0.1 to 324.5 kg ha<sup>-1</sup> and the highest concentration was observed in Northwestern China (i.e. Xinjiang, Gansu province and 246 Inner Mongolia) (Huang et al., 2020). More than 40% of soil samples  $(n = 51)$  were from the



269 included 31 provinces and based on the samples collected in 2017.

 The chemical composition of most macro-plastics generated from plastic mulch film was suggested to be polyethylene (PE) since the majority of plastic mulch film was made of PE in China. In addition, chemical compositions of polyvinyl chloride (PVC) and ethylene vinyl acetate copolymer (EVA) were also reported in several studies as shown in Table S2.

#### <span id="page-15-0"></span>*2.2.2 The occurrence of MPs in Chinese farmlands soil*

 The abundance of MPs was investigated in 28 provinces of Chinese farmland through the data collected from 40 articles (see the list of articles in Table S3) shown in Figure 2. The abundance 278 of MPs varied from 1.6 to  $6.2 \times 10^5$  items kg<sup>-1</sup> (dry soil) with an average abundance of 4536.6 279 items kg<sup>-1</sup>, with the median of 1640.0 items kg<sup>-1</sup>. The highest average MPs abundance (4,817.9 280 items kg<sup>-1</sup>) was found in Southern China, followed by 4,156.1 items kg<sup>-1</sup> in Northern China, 281 3,602.7 items kg<sup>-1</sup> in Northwestern China, and 82.3 items kg<sup>-1</sup>in the Qinghai–Tibet Plateau (Table S3). Interestingly, Northwestern China with a higher concentration of macro-plastics was not the highest abundance of MPs. This might be due to the lower temperatures and plough frequency in Northwestern China, causing the lower generation rate from macro-plastics to MPs. The most commonly researched regions for MPs were Hubei province (abundance of MPs from 286 327.5 to  $6.2 \times 10^5$  items kg<sup>-1</sup>), and Shanghai city (abundance of MPs from 1.6 to 2153.0 items  $\text{kg}^{-1}$ ) (see more information in Table S3). The number of sampling points in these two regions was the highest, with 59 (Hubei province) and 123 (Shanghai city), indicating that more research of MPs has been carried out in Hubei province (Zhou et al., 2019; Wang et al., 2021) and Shanghai City (Lv et al., 2019; Zhou et al., 2020) compared to other Chinese regions.

 However, there were only a few studies of MPs in Northwestern China (such as Xinjiang and Gansu provinces), where the concentration of macro-plastics was relatively high. Therefore, further studies are recommended that focus on quantifying terrestrial MPs pollution in Northwestern China.

 As the types of MPs are important to their environmental fate and ecotoxicity (Zhang et al., 2022), the chemical composition of MPs in Chinese farmland soils investigated in the literatures are summarized in Table S3. The mainly components of MPs in Chinese farmland soils were PE, polypropylene (PP), polyester (PES), polystyrene (PS) and polyamide (PA). 299 There were  $2 - 27$  types of plastic materials in the Northern China and PP and PE were mostly 300 widely used, while more types  $(2 - 60)$  of plastic materials were investigated in the Southern China. This may be due to the well-watered condition and higher population density in the region, resulting in a complex source of polymers (Chen et al., 2022). 10 of 11 literatures (which studied the chemical composition of MPs in the Northwestern China) showed that the main plastic type was PE. This was probably attributed to the large consumption of plastic mulch films in this area, and the majority of this film was made of PE in China. Therefore, PE mulch films probable be an important source of soil MPs in Chinese farmland soils.

#### <span id="page-16-0"></span>*2.3 The potential sources of MPs in Chinese farmlands soil*

 Most of the macro-plastics in farmland soils come from the damage of agricultural plastic products. However, the sources of MPs in soils are much more various. In the last 10 years, the agricultural sector has become increasingly important as a source of MPs in soils. The most potential agricultural sources (i.e. agricultural mulch films, compost and sewage sludge, and  atmospheric deposition) have been focused and mapped the distribution and abundance of MPs from these sources in China (Figure 2 and Table S3). It should be noted that the data points of MPs caused by other sources, such as coated fertilizer and food waste (which are not well investigated in previous literature but may contribute to MPs in soils) are presented in yellow circles in the revised Figure 2.

 This research explored the distribution and abundance of MPs from the agricultural plastic films with 146 sampling points (Figure 2(green)). It highlighted that most of the research on agricultural plastic films has focused on the regions of Northern and Northwest China (Zhang et al., 2021; Wang et al., 2021). In contrast, most of the research on compost and sewage sludge was concentrated in Southern China as shown in Figure. 2(purple) with 57 soil samplings, where there are more wastewater treatment plants and larger quantities of sludge production (Yang et al., 2021). Only a few cities have measured MPs in the atmosphere, including Shanghai, Dongguan, Dalian, Tianjin, Wenzhou, and Yantai, with a total of 15 samples as shown in Figure. 2(blue). The average abundance of MPs from agricultural plastic films, composts and sludge, and atmospheric 327 deposition were 4,231.1 items kg<sup>-1</sup>, 1,002.3 items kg<sup>-1</sup>, and  $7.9 \times 10^4$  items m<sup>-2</sup> yr<sup>-1</sup>. Taking into account all the potential sources on MPs, plastic mulch films represent the most important source in Chinese farmland soils.

#### <span id="page-17-0"></span>**3. Contribution of different sources to soil macro-plastics and MPs wastes**

 The sources of macro-plastics mainly consist of input from improper disposal of agricultural plastic practices and solid waste (e.g., domestic waste) from the surroundings of farmland soil (Hurley & Nizzetto, 2018; Qi et al., 2020). In contrast, the sources of MPs are more complex.

 MPs are generally categorized into two types: primary and secondary MPs. Primary MPs mainly refer to micro-size plastic particles that are manufactured intentionally for commercial uses, and they often act as raw materials for industrial production (Bläsing & Amelung, 2018; Yang et al., 2021). The rapid growth in the fibers from man-made textiles, microbeads from personal care products, and fragments produced during the plastic manufacturing process mean that they are significant sources of primary MPs (Fu & Wang, 2019). Furthermore, increasing quantities of primary MPs are being introduced to agricultural soils via organic wastes and wastewater residue 342 input (Yang et al., 2021). However, the most common MPs in the environment are secondary MPs (Qi et al., 2020; Cole et al., 2011). Secondary MPs are generated from the degradation and decomposition of larger macro-plastics products and debris into micro-nanoplastics by abiotic (e.g. high temperature, wind-blown and ultraviolet radiation, (Horton et al., 2017; Rezaei et al., 2019) and biotic factors (e.g. microbial decomposition, Yuan et al., 2020). Compared to primary MPs, secondary MPs are more difficult to determine the source and rates of generation. The following sections of this study provide a summary of the potential sources and quantitative estimate modelling methods for macro-plastics and MPs.

#### <span id="page-18-0"></span>*3.1 Plasticulture practices*

#### <span id="page-18-1"></span>*3.1.1 Plastic mulch films*

 Plastic mulch films are used for several reasons that improve crop yields, e.g. for increasing the soil temperature and water use efficiency, promoting seed germination, inhibiting weed growth, and reducing soil erosion (Gao et al., 2019). However, with the increasing use of agricultural plastics film, especially in several developing countries (e.g. China, India, Egypt, and Vietnam),

 there is an increasing legacy of plastic mulch film residue accumulation, including macro-plastics and MPs (Plastics Europe-The Facts 2022; Maraveas, 2020).

 In Asia, the largest usage of agricultural filmsis China with consumption of 2.5 million tons, accounting for > 70% of Asia's, and almost 50% of the worldwide in 2018 (NBSC, 2019; Le Moine, 2018; FAOUN, 2021). The use of plastic mulch film in China has increased nearly three times from 375 thousand tons in 1993 to 1320 thousand tons in 2021, however, the recovery rate after crop harvest is under 60% (Zhao et al., 2017). This equates to around 1.18 million 364 tons of legacy plastic (equivalent to the area of  $0.16 - 0.22$  million km<sup>2</sup>) that has been left in the 365 soil over this period (Atlas of the Second National Agricultural Pollution Source Census, 2022). A study by Ren et al. (2021) reported that the amounts of MPs in Chinese surface farmland 367 soils (0 – 10 cm) ranged from  $4.9 \times 10^6$  to  $1.0 \times 10^7$  tons in 2018, and agricultural mulch films 368 contributed  $10 - 30\%$  of the total inventory of MPs in the Chinese farmland. Of this, it was 369 estimated that  $1.2 \times 10^5 - 2.2 \times 10^5$  and  $3.4 \times 10^4 - 6.6 \times 10^4$  tons of MPs from Chinese farmland soils entered the surface water and ocean each year, respectively. In addition to the plastic polymer, chemicals added to agricultural films during their production, e.g. phthalates (phthalic acid esters, PAEs) can be released into farmland soil by plastic debris (Wang et al., 2016). According to the study by Zhang et al. (2021), 91.5 tons of PAEs migrated into Chinese soils from agricultural films in 2017, with a risk of these being taken up by vegetables and entering the human body via the food chain.

 Biodegradable plastic mulches (BDMs) have been developed as substitutes to conventional PE mulch films and are formulated to reduce the persistence of residues in soil (Yang et al., 2022). Because of the growing awareness of the persistence of synthetic plastic mulch films in the

 environment, BDMs have gradually entered the mulch film market in China (Plastics Europe- The Facts 2022; He et al., 2018). However, studies have reported that MPs formation is more rapid from biodegradable mulch than from traditional non-degradable mulch films. For example, 982 plastic films were exposed to UV irradiation of 2.1 MJ m<sup>-2</sup> in a lab experiment by Yang et al. (2022). This level of UV exposure simulated the cumulative irradiance level of 70 days of natural summer solar light in Northern China. The average quantity of MPs released from 385 biodegradable, and non-degradable mulch films were 475, and 155 particles cm<sup>-2</sup>, respectively.

 In summary, the mulching duration, amounts of mulch films and plastic material are important factors that affect the plastic fragmentation. As an important source of macro-plastics and MPs in agricultural soil, the level of plastic accumulation by agricultural plastic mulch films, especially BDMs, is alarming but has received relatively little attention to date. A few studies have shown that BDMs can contribute more MPs to soil compared with conventional PE mulch films at the same time period (Yang et al., 2022; Zhou et al., 2023). Therefore, further research is needed to investigate the fate and generation process of MPs from mulch films, including BDMs.

#### <span id="page-20-0"></span>*3.1.2 Greenhouse films*

 Greenhouses represent the largest proportion of agricultural plastic films used in plant production worldwide (Le Moine, 2018). It was estimated that the global average quantities of greenhouse 398 films used 3500 kg ha<sup>-1</sup> in 2019, which represented 47% of agricultural film demand (Le Moine, 2018; FAOUN, 2021). Greenhouses are used to prolong the growing season in temperate regions of the world, and most plastic greenhouses are concentrated around Asia (FAOUN, 2021). Since

 there is no direct amount of greenhouse films used in China and agricultural films is mainly used as mulch films and greenhouse films in China, the amount of greenhouse films is represented by the difference between the amount of agricultural films and mulch film (Zhang et al., 2021). In 2021, China's use of greenhouse films reached 1.04 million tons, accounting for 44% of the total amount of agricultural film (NBSC. 2022).

 A study by Wang et al. (2022) investigated MPs contamination from three different types of greenhouses (abandoned greenhouse, normal greenhouse, and simple greenhouse). The degree of 408 MPs contamination was found to follow the order: abandoned greenhouse (2215.56 items  $kg^{-1}$ ) > 409 normal greenhouse (891.11 items  $kg^{-1}$ ) > simple greenhouse (632.50 items  $kg^{-1}$ ). The composition of MPs from these different greenhouses was also different. The most important components of the abandoned greenhouses were rayon (RY) (10.3%) and poly (ethylene 412 terephthalate) (PET) (7.7%). In the simple greenhouses, poly (1-tetradecene) (PTD) (14.2%) and RY (10.0%) were more common. The most abundant polymer type was PP, PE, and polypropylene polyethylene copolymer (PP: PE) in all the three greenhouses. These polymers accounted for > 50% of the total (Wang et al., 2022).

 The environmental concern about MPs from greenhouse plastic film covers is less compared to PMFs. There is a tradition of recycling greenhouse films in China, since the greenhouse films (thickness of 8‒50 µm) is durable and easier to be recycled than mulch films (thickness of 6‒8 µm). The recovery rate of plastic mulch films was less than 60% in China (Zhao et al., 2017), and the target recovery rate of agricultural films will be 85% by 2025 (Development and Reform Commission of the People's Republic of China. 2021). It can be inferred that the recovery rate of greenhouse films would be higher than 85% by 2025.

#### <span id="page-22-0"></span>*3.1.3 Irrigation pipes and associated infrastructure*

 The source of macro-plastics and MPs in farmland soil is not limited to the use of plastic films, also included abandoned irrigation pipes, mismanaged agrochemical containers, and disposable crop protection packaging (Gündoğdu et al., 2022).

 A recent study in Turkey showed the concentration of MPs from disposable greenhouse 429 plastic films and irrigation pipes in agricultural soil ranged from 0.3 to 32 particles  $kg^{-1}$  with an 430 average of 11.1 particles  $kg^{-1}$  (Gündoğdu et al., 2022). Furthermore, MPs with additives could act as vectors for pollutants, e.g. dibutyl phthalate which has been shown to be released from PVC pipe fragments in water, representing an added risk to both organisms and the environment (Ye et al., 2020). In addition, the pollution level is highly correlated with the amount of disposable drip irrigation pipes and greenhouses in the contaminated sites (Katsumi et al., 2021; Ye et al., 2020). Results showed that irrigation pipes and associated infrastructure could be potential sources of MPs in irrigated farmland soils, which should be taken into account when estimating the concentration of MPs in soil (Gündoğdu et al., 2022; Pérez-Reverón et al., 2022). Furthermore, the topics of lifespan and waste management of agricultural plastic infrastructure and source identification of MPs in soil based on their physical properties (e.g. size, shape, and the type) are also worthy of further study.

#### <span id="page-22-1"></span>*3.1.4 Coated fertilizer use*

 Polymer-coated fertilizer (PCF) comprises a nutrient core wrapped by a polymer coating and is designed to release nutrients to plants at a gradual and controlled rate (Du et al., 2006). PCF is

445 composed of microcapsules with a thickness of  $10 - 80 \mu m$  and a diameter of  $2 - 5 \mu m$ , which are not recovered after use. These microcapsules are primary MPs and can further degrade into NPs (Katsumi et al., 2021; Bian et al., 2022; Trenkel 2010). The use of PCF in China is increasing at a rate of 10% − 15% per year (Li et al., 2022). It is expected that the output of the Chinese PCF 449 will reach  $7.6 \times 10^6 - 11.3 \times 10^6$  tons by 2025, with the microcapsules input to soils potentially 450 amounting to  $0.4 \times 10^6 - 0.6 \times 10^6$  tons (Yang et al., 2009).

 Although PCF can reduce nutrient leaching loss and ammonia emissions, the fate and impact of the residual polymer coating are attracting the attention of fertilizer companies and environmental researchers (Lian et al., 2021). For example, Katsumi et al. (2021) investigated the accumulation of microcapsules derived from coated fertilizer in 19 paddy fields in Japan with 455 concentrations found to range from  $6 - 369$  mg kg<sup>-1</sup> (mean 144 mg kg<sup>-1</sup>). The result showed that legacy plastics from microcapsules will continue to accumulate in farmland soil as long as conventional PCF is used. The spatial distribution of MPs from PCF is also strongly affected by irrigation, and the soil around drainage outlets has been found to be a hot spot (Katsumi et al., 2021). Several studies have also measured the release of macro-plastics in the environment by PCF made from different co-polyesters (Lubkowski et al., 2016). For example, experiments have shown that the residual amount of PCF film shells left in the soil was 50 kg ha<sup>-1</sup> every year, which accounting for 50% of the average annual nutrient consumption input in the European Union (100 kg ha−1 ) (Lubkowski et al., 2016). Furthermore, it is estimated that the concentration of macro-464 plastics from PCF in the soil can reach 500 kg ha<sup>-1</sup> after continuous application of PCF for 10 years (Li et al., 2022). Whilst previous research has focused on the benefits of PCF on plant growth, soil properties, soil microbial communities, and reduced risk of nutrient losses to water  and air, further attention is needed to assess the contribution of this source to plastic pollution in China (Bian et al., 2022; Lian et al., 2021). The release of MPs may become a potential food safety problem for the long-term application of PCF in farmland.

## <span id="page-24-0"></span>*3.2 Organic wastes and wastewater residue input*

#### <span id="page-24-1"></span>*3.2.1 Sewage sludge*

 During wastewater treatment, most MPs are removed from the wastewater stream and become concentrated in the sludge (biosolids) fraction (Ziajahromi et al., 2016). In many countries, this nutrient-rich semi-solid waste product is applied to agricultural land as a soil improver and fertilizer (Hurley & Nizzetto, 2018; Corradini et al., 2019). Agricultural soils in Europe and North America may receive more than 63,000 and 44,000 tons of MPs per year through sludge applications, respectively (Nizzetto et al., 2016). However, very little is known about the fate and transport of MPs in sludge in the terrestrial environment (Ng et al., 2018; de Souza Machado et al., 2018).

 In Europe and North America, about 50% of sewage sludge is processed for agricultural use, and it is estimated that 125 – 850 tons of MPs per million peoples are added annually to European agricultural soils either through direct application of sewage sludge or as processed biosolid (Nizzetto et al., 2016). A recent study by Lofty et al. (2022) estimated a maximum application 485 rate of 4.8 g of MPs m<sup>-2</sup> yr<sup>-1</sup> or  $1.15 \times 10^4$  MPs particles m<sup>-2</sup> yr<sup>-1</sup> in Europe from sewage sludge applied to agricultural soil by measuring the MPs content of sewage sludge at wastewater treatment plants (WWTPs). These studies strongly suggest that the practice of spreading sludge on agricultural land could potentially make them one of the largest global reservoirs of primary

MPs pollution (Lofty et al., 2022).

 In China, the situation is similar to other regions of the world. It was estimated that more than seven million tons of dry sludge were generated from wastewater treatment in China in 2020 (MEPC, 2017). However, > 80% of this sludge is disposed of improperly (i.e. dumped) with only 2.4% of the sludge applied to land (Yang et al., 2015). Li et al. (2018) investigated the occurrence of MPs in sludge by analyzing 79 sewage sludge samples collected from 28 WWTPs in 11 Chinese provinces. The results showed that on average, the concentration of sludge-based MPs 496 entering the soil and the wider environment in China was estimated to be  $1.56 \times 10^{14}$  particles per year, which is the same order of magnitude of MPs released into European farmland soils (i.e. 8.6  $\times 10^{13} - 7.1 \times 10^{14}$  particles per year) (Li et al., 2018; Lofty et al., 2022). Yang et al. (2021) investigated the contributions of three types of sludge (i.e. fresh municipal sludge, mainly industrial sludge, and dry heat-treated municipal sludge), which were repeatedly applied to farmland soil for nine years in Jiangsu province, Southwest China. The results showed that the 502 input of sludge led to an accumulation of MPs in the soil, as high as 149.2 particles  $kg^{-1}$  (compared with the control treatment). These findings confirm that sewage sludge recycling to land represents an important source of plastic pollution in the environment.

 At present, the reported treatment methods of MPs in sewage sludge are generally divided into two types: i) physical and chemical methods, and ii) anaerobic digestion methods (Wu et al., 2022). Of these, physical and chemical methods often cause MPs to break into smaller plastic fragments. Whilst there is evidence that some MPs (such as PET and polyurethane reactive, PUR) can be partially degraded under anaerobic digestion (Mahon et al., 2017), most MPs are not degraded, mainly depending on their chemical structure, molecular weight as well

 as the type of plastic additives in MPs (Moharir & Kumar, 2019). In the future, the technique of efficiently removing MPs from sewage sludge should be developed to reduce the pollution caused by MPs.

 The Chinese government has proposed that the daily capacity for harmless treatment of 515 sludge (with moisture content >80%) should be no less than  $2.0 \times 10^4$  tons by 2025 (Development and Reform Commission of the People's Republic of China, 2022). The harmless treatment rate of urban sludge is expected to reach above 90%. These policies would affect the sources and occurrence of MPs, which can be further investigated in the future study.

#### <span id="page-26-0"></span>*3.2.2 Compost*

 Organic resources such as compost are rich in plant nutrients and organic carbon and are hence widely used as soil amendments to improve soil properties and soil nutrient content (Cherif et al., 2009). However, there is increasing evidence that soils receive plastic input through the application of compost (Bläsing & Amelung, 2018). Because of improper disposal and insufficient waste separation of plastic from organic matter, macro-plastics in compost can accumulate in the soil and risk entering the food chain via crop plants. In China, Zhang et al. (2022) investigated the abundance, shape, composition, and size of MPs from organic fertilizers using attenuated total Fourier transformed infrared spectroscopy. The results showed that mature 529 compost application to agricultural fields goes along with MPs load of  $3.5 \times 10^{12} - 6.6 \times 10^{12}$  items per year. Another study in Germany showed that compost application led to an annual input of > 1mm plastic plastics to arable fields that reached up to 35 billion – 2.2 trillion (Weithmann et al., 2018). In recent years, several countries have strongly encouraged farmers to use organic

 fertilizers and successively formulated subsidy policies for composts. For instance, the Chinese 534 Ministry of Agriculture provides a subsidy for households of 1,500 RMB ha<sup>-1</sup> (equal to 215 US 535 dollars ha<sup>-1</sup>) to use >3,750 kg ha<sup>-1</sup> commercial composts in some pilot areas (Ministry of Agriculture and Rural Affairs of the People 's Republic of China. 2018). However, more attention needs to be paid to the potential contribution of compost to macro-plastics and MPs in agricultural soils.

 The analysis result shows that plastic mulch films are the most studied potential sources of macro-plastics and MPs in Chinese farmland soils, while compost and sewage sludge may be important sources in Europe, since this type of fertilization is commonly used in European counties.

#### <span id="page-27-0"></span>*3.2.3 Food waste digestate*

 Diverting food waste from landfills to anaerobic digestion can facilitate the conversion of energy into usable forms and produce nutrient-rich soil improvers (Cheong et al., 2020; Xu et al., 2018). However, concerns arise due to the presence of plastic packaging in many food waste streams, which may inadvertently introduce macro- and micro-plastics into agricultural soils (Porterfield et al., 2023). For example, the abundance of MPs was  $3.0 \times 10^5$  pieces kg<sup>-1</sup> in food waste collected from grocery stores in the USA (Golwala et al., 2021), and it was  $4.1 \times 10^3$  particles kg<sup>-1</sup> in the food compost sample in Lithuania (Sholokhova et al., 2021). In addition, some biodegradable plastic packages (e.g. Polylactic acid, PLA) are widely used in food packaging and disposable tableware and the usage of biodegradable plastic packages are increasing (Lu et al., 2022). However, the aging and fragmentation process of PLA also could be enhanced within

 thermophilic anaerobic digestion with kitchen waste, generating large amounts of macro- plastics and MPs. Research on the occurrence and relative importance of MPs from food waste is in an early stage and this potential pathway of macro-plastics and MPs to agricultural soils needs further clarification (Porterfield et al., 2023).

## <span id="page-28-0"></span>*3.3 Other sources of plastic contamination*

#### <span id="page-28-1"></span>*3.3.1 Atmospheric deposition*

 Atmospheric transport and deposition of MPs is one of the major pathways for plastic fragments entering the soil environment (Allen et al., 2019; Brahney et al., 2020). It is estimated that 564 atmospheric deposition rates of MPs range from  $1.1 \times 10^4$  to  $4.1 \times 10^5$  items m<sup>-2</sup> yr<sup>-1</sup> globally (Allen et al., 2019; Brahney et al., 2020; Bergmann et al., 2019). These particles typically enter the atmosphere through mechanical processes, such as dust entrainment during strong wind events or wave breaking of sea surface spray (Seinfeld & Pandis, 2008). Brahney et al. (2021) created a model to calculate the atmospheric component of the plastic cycle, estimating the current average daily total atmospheric burden (content) of MPs over the land regions of the western United States to be ca. 100 tons. The largest contributor to modeled plastic deposition (84%) in the western United States is road dust. In comparison, agriculturally derived plastics in dust entrained into the atmosphere from agricultural fields are thought to contribute 5% to annual total deposition in the same region (Brahney et al., 2021). In China, Liu et al. (2019) measured indoor and outdoor dust samples collected from 39 major cities of China. The mass concentrations of PET and polycarbonate (PC) MPs were determined, and the concentrations of PET and PC MPs 576 in dust were  $1.6 \times 10^3 - 1.2 \times 10^5$  mg kg<sup>-1</sup> and 4.6 mg kg<sup>-1</sup> (indoors), 212 – 9,020 mg kg<sup>-1</sup> and

2.0 mg kg−1 (outdoors), respectively (Liu et al., 2019). Although it is difficult for MPs > 50 μm to enter the respiratory tract, these particles can enter the gastrointestinal tract where adsorbed contaminants may be released, posing a potential threat to human health (Brahney et al., 2020; Bergmann et al., 2019; Liu et al., 2019).

<span id="page-29-0"></span>*3.3.2 Rubber tire wear*

 Rubber is also considered a class of plastic, and physical abrasion of tires significantly contributes to the release of MPs into the environment (Lassen et al., 2015). In addition, tire residues are also present in sewage sludge where road run-off enters the wastewater network (Essel et al., 2015). Rubber MPs size and generation rate from tires depends on their composition (Kole et al., 2017). Evangeliou et al. (2020) found high transport efficiencies of rubber MPs to remote regions worldwide. Their results showed that about 34% of the emitted coarse tire wear particles (TWPs) 589 and 30% of the emitted coarse brake wear particles (BWPs) (100 kt yr<sup>-1</sup> and 40 kt yr<sup>-1</sup>, respectively) were subsequently deposited in the world's oceans. However, knowledge about the fate of tire-derived MPs entering Chinese farmland, especially those from farm machinery, is currently lacking (Evangeliou et al., 2020).

#### <span id="page-29-1"></span>*3.3.3 Water-flow and irrigation*

 Many studies have indicated that large quantities of MPs are present in irrigation source (Chen et al., 2022; Pérez-Reverón et al., 2022). Research showed that the MPs concentration in irrigation water was significantly correlated with the abundance of MPs in agricultural soil, and the MPs concentration of soils in direct contact with irrigation water was significantly higher  than that in deeper soils (Katsumi et al., 2021). A study in Spain showed that the shape, color, size and type of MPs in soil samples collected from cropland were similar to those in the irrigation water used on the crops (Pérez-Reverón et al., 2022). This evidence indicates that irrigation water is an important source of MPs in farmland soils. Moreover, comparing MPs abundance in the different source of irrigation water, the concentration of MPs in recycled 604 wastewater (159 items  $kg^{-1}$ ) was around three times higher than that in the desalinated brackish 605 water (46 items kg<sup>-1</sup>) (Pérez-Reverón et al., 2022). In addition, the MPs abundance in agricultural soil irrigated by underground water and rainwater is significantly lower than irrigated with surface water (Chen et al., 2022). Meanwhile, the soil is also a potential MPs sink and MPs in soils could be transported by water off into surface water and ocean (Ren et al., 2021). It is important to explore the natural and anthropogenic processes affecting the fate of MPs in irrigation water.

#### <span id="page-30-0"></span>**4. Modelling method for quantifying the source of macro-plastics and MPs**

#### <span id="page-30-1"></span>*4.1 Material Flow Analysis (MFA)*

 MFA is a useful tool applied to better understand pathways of substance. It is an analytical method to quantify flows and stocks of materials or substances in a well-defined system (Bornhöft et al., 2016). Several studies have quantified the possible flows of plastic into the soil from different sources by using this method (Kawecki & Nowack, 2020; Liu et al., 2020; Sieber et al., 2020). MFA has been employed to analyze the flows of plastics in China (Luan et al., 2021; Liu et al., 2020; Luan et al., 2022). However, few studies have focused on plastic emissions and flows in an agricultural context. Of the relevant studies, Zhou et al. (2013) analyzed the emission and  accumulation of PVC waste in the environment, and Bai et al. (2018) estimated and predicted the annual input of plastic waste from China into the ocean. Nevertheless, most of these studies have used emission factors to estimate the losses of plastics directly or estimated the amounts of plastic waste through municipal solid waste indirectly, rarely covering all life cycle processes and not distinguishing between plastic types.

 Luan et al. (2021) conducted a study to assess MPs and macro-plastics losses throughout the plastic life cycle by using a dynamic MFA approach. The losses were analyzed based on different polymers (including PE, PP, PS), PVC, acrylonitrile-butadiene-styrene (ABS), and PET, sources (including personal care products, laundry process, indoor dust, fishery waste), environmental media (ocean and soil) and lifecycle processes (i.e. production, use, recycling, and end-of-life treatment). Based on field research and published literature, localized emission factors were obtained to systematically and comprehensively estimate the plastic losses to the environment in China. The results showed that MPs and macro-plastics losses entering the environment were 634 352.1 kt y<sup>-1</sup> and 12.7 Mt yr<sup>-1</sup>. Of these losses, PET accounted for the highest proportion (29.1%) and 32.2%), and the net loss to the ocean and soil were 4.0 Mt and 173.7 Mt, respectively in 2020 (Luan et al., 2022).

 Based on the research mentioned above, our study summarized the necessary steps of this evaluation method (i.e. MFA) to calculate the main stocks and flows of soil macro-plastics and MPs derived from different sources of agricultural activities. There is an essential process in the MFA where the input of secondary MPs is calculated, i.e. generation rate. According to this rate, the concentration of MPs converted from plastic products can be calculated. As mechanical abrasion (MA) and ultraviolet (UV) irradiation are key factors controlling plastic degradation rates, they are discussed in more details below.

#### <span id="page-32-0"></span>*4.1.1. Generation rate.*

 Generation rate is an essential part of MFA, which refers to the mass ratio of MPs generated from macro-plastics. This is a necessary index in the calculation of the flows and stocks of materials in different environments. The methods for calculating generation rates of MPs from macro-plastics vary (Ren et al., 2020; Sieber et al., 2020). Plastic fragmentation may be caused by solar ultraviolet (UV) irradiation, physical abrasion (abiotic), or biological attack (Cole et al., 2011; Kershaw, 2015). Therefore, it is important to provide an improved estimation of MPs generation in farmlands according to the generation rate of MPs.

 *Mechanical abrasion (MA)*. MA of plastics is likely to be common in many cropping environments. For example, plastic packaging of seeds, crops, and fertilizers is abraded by external forces during transportation and use, resulting in plastic debris left in the environment. Further, as mentioned previously, a large amount of plastic debris is left in the soil after the crop harvest due to the fragile nature of the mulch films (Zhao et al., 2017; Rezaei et al., 2019). Farmland may be the most favorable environment for plastic weathering and fragmentation because of photodegradation and MA of plastics by soil (Ren et al., 2020). Studies to measure MPs generation via MA are limited.

 Some studies have demonstrated the impact of polymer type on MP generation rate, especially following UV exposure. Song et al. (2017) calculated MPs fragmentation rates via the combined effects of MA (using a mechanical roller) and UV exposure for LDPE, PP, and  expanded polystyrene (EPS). Their results showed that there was a minimal effect of MA (crushing) on MPs generation from LDPE and PP when the plastic was not exposed to UV (fragmented particle generation was 8.7 and 10.7 particles, respectively), but PP generated more MPs following UV radiation. However, MPs generation from EPS was mainly via MA, which resulted in 4220 MPs particles after only 2 months of mechanical friction by a roller without any UV irradiation (Song et al., 2017). Ren et al. (2020) utilized a range of sizes of sandpaper to abrade different components of plastic film to simulate MA and demonstrated a positive relationship between fragmentation rates of different plastic mulch films and relative light transmittance (RLT, %). These studies provide new insight for future calculation of MA rates and MPs production.

 *Ultraviolet (UV) irradiation*. One of the most important degradation factors of plastic polymers fracture is UV radiation, which induces oxidation and molecular chain scission for plastics breakdown in the environment (Laycock et al., 2017; Uheida et al., 2021). The random chain scission and cross-linking of plastic polymers results in the progressive formation of micro-, and nano-size plastic particles from macro-plastics fragments (Qi et al., 2020). Yang et al. (2022) found that when different types of macro-plastics were incubated in soil exposed to the same 682 cumulative UV irradiation (2.1 MJ m<sup>-2</sup>), the average quantity of MPs generated from biodegradable and oxo-degradable plastic mulch films was greater than from conventional non- degradable mulch films. These results help us to understand the kinetics and mechanisms of different types of mulch films (Yang et al., 2022). It is worth noting that the generation of MPs from biodegradable plastics in the terrestrial environment is more significant (Bao et al., 2022; Qin et al., 2021).

#### <span id="page-34-0"></span>*4.2 The other quantitation methods of macro-plastics and MPs sources*

 Several studies have listed the possible flows of plastic into soils based on an analysis of plastic use in agriculture (Brandes et al., 2021; Ren et al., 2021; Wang et al., 2021). The empirical formulas for estimating MPS generation from agricultural plastics are summarized in Table 1. Brandes et al. (2021) used data-driven models alongside data on MPs composition from the literature in combination with national statistics on sewage sludge, compost, and plastic waste production, as well as specialty cropping areas, to estimate the spatial distributions of cumulative MPs mass inputs into agricultural soils in Germany. Based on the Nomenclature of territorial units 697 for statistics (NUTS3) scale, the results showed that MPs input range for soils was  $0 - 15.7$  kg  $h^2$ ,  $h^2$ , 0 – 3.79 kg ha<sup>-1</sup>, and 0 – 5.18 kg ha<sup>-1</sup> from sludge, compost, and plasticulture used in agriculture, respectively. Of these, the contribution of MPs followed the series sewage sludge > compost > plasticulture in Germany (Brandes et al., 2021). Ren et al. (2021) found PMFs contributed 10 – 30% of the total MPs from all sources in farmland soils according to a Monte Carlo simulation. In addition, Wang et al. (2021) developed a novelty model to calculate the distribution of MPs in the soil environment based on the aging process of different types of MPs (fibers, films, fragments and granules). Based on this model, a distinct downsizing phenomenon from fibers, films, and fragments to granules was observed, and human interference accelerated the fragmentation of MPs. However, the quantitative contribution of different plastic sources to Chinese farmland soil MPs, specifically the rate and kinetics of MPs formation, is still in the primary stage.

#### <span id="page-35-0"></span>*4.3 Uncertainty statistics*

711 Quantifying MPs in soil mainly relies on the use of multiple statistical sources of data, which may lead to uncertainty of model outputs. Other sources of uncertainty include soil analytical data, as different studies have used different MPs extraction and detection methods (Bläsing & Amelung, 2018; Li et al., 2020). Accordingly, systematic uncertainty analyses are necessary to confirm the reliability of the method. A unified approach developed by Laner et al. (2016) could be used for characterizing the uncertainty of data based on this method, where the quantitative calculation of the data uncertainties used coefficients of variation (CV) (Laner et al., 2016).

 Monte Carlo (MC) simulation can be used to investigate the effects of parameters and data changes, combined with CV (Wang & Ma, 2018). The MC method employs observed data to simulate the distribution, and then the uncertainty range is calculated. This is common in MFA research when there are sufficient data (Luan et al., 2022; Tsai & Krogmann, 2013). It is noteworthy that the MC can be best used if the dataset contains more than 30 records (Montangero & Belevi, 2008).

#### <span id="page-35-1"></span>**5. Gaps in knowledge and priorities for future research**

 Taking China as an example, this research reviewed the source, location, and occurrence of macro-plastics and MPs pollution in soils. It needs to be noted that a limitation of this research is the uncertainty introduced by the plastic data from various studies. In these studies, different MPs extraction and detection methods are employed, which makes the data less comparable. Therefore, we recommend that future studies could adopt a standardized extraction and  detection protocols to reduce this uncertainty. This research also concludes that there are many ways that plastic fragments can enter the soil, including plasticulture practices, irrigation pipes and associated infrastructure, organic wastes and wastewater residues, atmospheric deposition, rubber tire wear and irrigation water. Of these, agricultural plastic mulch films are the most important source of macro-plastics and MPs in Chinese farmland soils, while wastewater-derived sludge may be more important in European countries. Furthermore, a MFA modeling approach can be used to quantify the flows and stocks of macro-plastics and MPs. In terrestrial systems, research on MPs is growing rapidly. However, more real-world studies are needed to narrow the knowledge gaps in the following aspects:

 1. Tracking the transport and fate of MPs from different sources in soil. At present, most of the quantitative studies are estimated by models, with few actual field measurements. The use of isotopically labeled plastics would be useful to track their fate, stocks, and their biodegradation rates. In addition, current research is also restricted by the limited availability of published data. Consequently, more samples to be required from soil, irrigation water, compost and sludge, and other potential sources in the typical regions using plastic films. This will help better parameterize models and reflect regional conditions.

 2. There is still a lack of information on the impacts of MPs on soil health, especially regarding non-point sources. Furthermore, to compare soil MPs concentrations and forms of MPs, sampling, extraction, and detection methods should be standardized. In addition, most soil MPs contents are expressed as the number of MPs per unit weight of soil (e.g. items  $kg^{-1}$ ), rather than the mass of MPs per unit weight of soil (e.g. mg kg<sup>-1</sup>). Future research should consider the use of mass units more often, to allow a more accurate description of the mass

 transfer of MPs through the environment and organisms (i.e. humans, livestock and soil fauna).

 3. Establishing spatially explicit models to predict the transfer and fate of MPs entering soils, linking specific source inputs to movement within the landscape and soil profile. These models need to account for different soils, climates, and management practices and describe MPs migration, generation rates, and the ultimate fate. Such models could then be used to underpin practical guidance to farmers and regulators to promote more sustainable use of plastics, and their alternatives, in agriculture.

 4. Strengthening research on the influence of residual plastic film or MPs on crop growth, yield and crop quality. The mechanistic basis of how residual plastic film or MPs affect nutrient cycling in the soil also needs to be explored, to determine the optimal use of plastic mulch films for different crops. Further efforts are also needed to improve the efficacy of the retrieval of used plastic mulch films after crop harvest.

 5. Strengthen the research on the impact of agricultural plastic use and recycle policies. The formulation of policies can greatly improve the standard of farmers' use of agricultural plastics. In China, for example, the governments have introduced various regulations and incentive mechanisms following research on the hazards of plastic film residues in recent years. In February 2022, the National Development and Reform Commission and the Ministry of Ecology and Environment issued the "*Key Points for the Treatment of Plastic Pollution*", which called for standardize use and recycling of plastic film, focusing on the promotion and application of biodegradable plastic film and thickened high-strength plastic film in key film areas. Through improvements to the recycling network system and  effectively controlling agricultural film pollution in farmland, the recovery rate of agricultural film should be stabilized at more than 80%. In addition, the "*Opinions on Further Strengthening the Treatment of Plastic Pollution*" issued by the Central Office of the Reform and Reform prohibited the production, sale and use of several specific plastic products (e.g. plastic shopping bags with a thickness of less than 25 μm, PE agricultural mulch film with a thickness of less than 10 μm) in 2019, and the "*Notice on Solidly Promoting the Treatment of Plastic Pollution*" issued in 2020 by the National Development and Reform Commission, have clearly required the strengthening of the plastic prohibition and restriction policies in the fields of agriculture, retail and catering. These show that China is making efforts to reduce the environmental pollution caused by plastic waste from the aspect of policy control. 6. Discussion the economic evidence of plastic pollution. Based on the survey results of 1067 cotton farmers in Xinjiang Province, China (Liu et al., 2020), most farmers are willing to recycling plastic film rather than incineration or landfill with the supports, such as reducing input cost of agricultural plastic film recycling by government subsidies, improving the mechanization of agricultural film recycling and setting up lectures and training about use and recycling of agricultural plastic film. In addition, there is already evidence suggesting extensive negative impacts of plastic waste on ecosystem services (Beaumont et al., 2019). 792 Researchers have estimated the loss of  $1 - 5\%$  in marine ecosystem services due to plastic pollution. This decrease equates to about \$0.5 to \$2.5 trillion per year. In other words, each 794 metric ton of plastic waste costs about  $\$3.3 \times 10^4$  (Beaumont et al., 2019). This illustrates that plastic pollution is not just damaging the environment, but that the economic losses caused by plastic pollution can also be substantial.

#### <span id="page-39-0"></span>**6. Conclusions**

 This review summarized the potential sources, current concentration and abundance, and modelling methods of quantification and flows of macro-plastics and MPs pollution in soils. Plastic pollution appears to be ubiquitous in soil and evidence suggests that it may represent an environmental risk to soil quality and soil functioning. Furthermore, as China is the biggest agricultural plastic film manufacturer and user it will face more serious pollution risks from legacy plastic in the farmland soil. Therefore, there is an urgent need to quantify the emission of plastic debris from different agricultural activities to the environment and to control the input of plastic from different sources into farmland soil. In addition, tracking the fate of MPs and NPs in soils, analyzing the impact of plastic waste on soil ecological health based on realistic concentration of MPs and NPs in the field, and studying the influence of policy and economic evidence on plastic pollution also deserve of future research.

#### <span id="page-39-1"></span>**Acknowledgments**

 This research was supported by the National Natural Science Foundation of China under Grant [number 42277097]; the UKRI Global Challenges Research Fund (GCRF) and the Natural Environment Research Council project, "Do agricultural microplastics undermine food security and sustainable development in less economically developed countries?" under Grant 816 [NE/V005871/1]; the International Cooperation and Exchange of the National Natural Science Foundation of China under Grant [NSFC-UNEP: 32261143459], and the High-level Team Project of China Agricultural University.

# <span id="page-40-0"></span>**Disclosure statement**

The authors report there are no competing interests to declare.

#### <span id="page-41-0"></span>**References**

- 826 1. Allen, S., Allen, D., Phoenix, V.R., Le Roux, G., Jimenez, P.D., Simonneau, A., Binet, S. and Galop, D. (2019). Atmospheric transport and deposition of microplastics in a remote mountain catchment. *Nature Geoscience* 12, 339. doi: 10.1038/s41561-019-0335-5
- 829 2. Atlas of the Second National Agricultural Pollution Source Census. (2022). Beijing: China **Agriculture Press.**
- 3. Bai, M., Zhu, L., An, L., Peng, G. and Li, D. (2018). Estimation and prediction of plastic waste annual input into the sea from China. *Acta Oceanologica Sinica* 37, 26-39. doi: 10.1007/s13131-018-1279-0
- 4. Bao, R., Pu, J., Xie, C., Mehmood, T., Chen, W., Gao, L., Lin, W., Su, Y., Lin, X., and Peng, L. (2022). Aging of biodegradable blended plastic generates microplastics and attached bacterial communities in air and aqueous environments. *Journal of Hazardous Materials,* 434, 128891. doi: 10.1016/j.jhazmat.2022.128891
- 5. Barnes, D.K., Galgani, F., Thompson, R.C. and Barlaz, M. (2009). Accumulation and fragmentation of plastic debris in global environments. *Philosophical transactions of the royal society B: biological sciences* 364, 1985-1998. doi: 10.1098/rstb.2008.0205
- 6. Beaumont, N. J., Aanesen, M., Austen, M. C., Börger, T., Clark, J. R., Cole, M., Hooper, T., Lindeque, P. K., Pascoe, C., and Wyles, K. J. (2019). Global ecological, social and economic impacts of marine plastic. *Marine pollution bulletin* 142, 189-195. doi: 10.1016/j.marpolbul.2019.03.022
- 7. Bergmann, M., Mützel, S., Primpke, S., Tekman, M.B., Trachsel, J. and Gerdts, G. (2019). White and wonderful? Microplastics prevail in snow from the Alps to the Arctic. *Science Advances* 5, eaax1157. doi: 10.1126/sciadv.aax1157
- 848 8. Beriot, N., Peek, J., Zornoza, R., Geissen, V., and Lwanga, E. H. (2021). Low density- microplastics detected in sheep faeces and soil: A case study from the intensive vegetable farming in Southeast Spain. *Science of the Total Environment* 755, 142653. doi: 10.1016/j.scitotenv.2020.142653
- 9. Bian, W., An, L., Zhang, S., Feng, J., Sun, D., Yao, Y., Shen, T., Yang, Y. and Zhang, M. (2022). The long-term effects of microplastics on soil organomineral complexes and bacterial communities from controlled-release fertilizer residual coating. *Journal of Environmental Management* 304, 114193. doi: 10.1016/j.jenvman.2021.114193
- 10. Bläsing, M. and Amelung, W. (2018). Plastics in soil: Analytical methods and possible sources. *Science of the Total Environment* 612, 422-435. doi: 10.1016/j.scitotenv.2017.08.086
- 11. Bornhöft, N.A., Sun, T.Y., Hilty, L.M. and Nowack, B. (2016). A dynamic probabilistic material flow modeling method. *Environmental Modelling Software* 76, 69-80. doi: 10.1016/j.envsoft.2015.11.012
- 12. Boughattas, I., Hattab, S., Zitouni, N., Mkhinini, M., Missawi, O., Bousserrhine, N., and Banni, M. (2021). Assessing the presence of microplastic particles in Tunisian agriculture soils and their potential toxicity effects using Eisenia andrei as bioindicator. *Science of The Total Environment* 796, 148959. doi: 10.1016/j.scitotenv.2021.148959
- 13. Brahney, J., Mahowald, N., Prank, M., Cornwell, G., Klimont, Z., Matsui, H. and Prather, K.A. (2021). Constraining the atmospheric limb of the plastic cycle. *Proceedings of the*
- *National Academy of Sciences* 118. doi: 10.1073/pnas.2020719118
- 14. Brahney, J., Wetherbee, G., Sexstone, G.A., Youngbull, C., Strong, P. and Heindel, R.C. (2020). A new sampler for the collection and retrieval of dry dust deposition. *Aeolian Research* 45, 100600. doi: 10.1016/j.aeolia.2020.100600
- 15. Brandes, E., Henseler, M. and Kreins, P. (2021). Identifying hot-spots for microplastic contamination in agricultural soils—a spatial modelling approach for Germany. *Environmental Research Letters* 16, 104041. doi: 10.1073/pnas.2020719118
- 16. Chen, L., Yu, L., Li, Y., Han, B., Zhang, J., Tao, S., & Liu, W. (2022). Spatial Distributions, Compositional Profiles, Potential Sources, and Intfluencing Factors of Microplastics in Soils from Different Agricultural Farmlands in China: A National Perspective. Environmental Science & Technology, 56(23), 16964-16974. doi: 10.1021/acs.est.2c07621
- 17. Cheong, J.C., Lee, J.T., Lim, J.W., Song, S., Tan, J.K., Chiam, Z.Y., Yap, K.Y., Lim, E.Y., Zhang, J. and Tan, H.T. (2020). Closing the food waste loop: Food waste anaerobic digestate as fertilizer for the cultivation of the leafy vegetable, xiao bai cai (Brassica rapa). *Science of the Total Environment* 715, 136789. doi: 10.1016/j.scitotenv.2020.136789
- 18. Cherif, H., Ayari, F., Ouzari, H., Marzorati, M., Brusetti, L., Jedidi, N., Hassen, A. and Daffonchio, D. (2009). Effects of municipal solid waste compost, farmyard manure and chemical fertilizers on wheat growth, soil composition and soil bacterial characteristics under Tunisian arid climate. *European Journal of soil biology* 45, 138-145. doi: 10.1016/j.ejsobi.2008.11.003
- 19. Cole, M., Lindeque, P., Halsband, C. and Galloway, T. (2011). Microplastics as contaminants in the marine environment: a review. *Marine Pollution Bulletin* 62, 2588- 2597. doi: 10.1016/j.marpolbul.2011.09.025
- 20. Corradini, F., Meza, P., Eguiluz, R., Casado, F., Huerta-Lwanga, E. and Geissen, V. (2019). Evidence of microplastic accumulation in agricultural soils from sewage sludge disposal. *Science of the Total Environment* 671, 411-420. doi: 10.1016/j.scitotenv.2019.03.368
- 21. Coursol, A. and Wagner, E.E. (1986). Effect of positive findings on submission and acceptance rates: a note on meta-analysis bias. doi: 10.1037/0735-7028.17.2.136
- 22. Cozar, A., Echevarria, F., Gonzalez-Gordillo, J.I., Irigoien, X., Ubeda, B., Hernandez-Leon, S., Palma, A.T., Navarro, S., Garcia-de-Lomas, J., Ruiz, A., Fernandez-de-Puelles, M.L. and Duarte, C.M. (2014). Plastic debris in the open ocean. *Proceedings of the National Academy of Sciences of the United States of America* 111, 10239-10244. doi: 10.1073/pnas.1314705111
- 23. de Souza Machado, A.A., Lau, C.W., Till, J., Kloas, W., Lehmann, A., Becker, R., Rillig, M.C. (2018). Impacts of microplastics on the soil biophysical environment. *Environmental Science Technology* 52, 9656-9665. doi: 10.1021/acs.est.8b02212
- 24. Development and Reform Commission of the People 's Republic of China. Notice on Issuing the Action Plan for Plastic Pollution Control during the 14th Five Year Plan Period. [2021-09-08] [https://www.gov.cn/zhengce/zhengceku/2021-09/16/content\\_5637606.htm](https://www.gov.cn/zhengce/zhengceku/2021-09/16/content_5637606.htm)
- 25. Development and Reform Commission of the People 's Republic of China. The implementation plan of harmless treatment and resource utilization of sludge. [2022-09- 22]https://www.gov.cn/zhengce/zhengceku/2022-09/28/content\_5713319.htm
- 26. Ding, J., Zhu, D., Wang, H.-T., Lassen, S.B., Chen, Q.-L., Li, G., Lv, M. and Zhu, Y.-G.
- particles. *Environmental Science Technology* 54, 7450-7460. doi: 10.1021/acs.est.0c00917
- 27. Dong, H.D., Liu, T., Han, Z.Q., Sun, Q.M. and Li, R. (2015). Determining time limits of continuous film mulching and examining residual effects on cotton yield and soil properties. *Journal of Environmental Biology* 36, 677-684.
- 28. Du, C.-w., Zhou, J.-m. and Shaviv, A. (2006). Release characteristics of nutrients from polymer-coated compound controlled release fertilizers. *Journal of Polymers the Environment* 14, 223-230. doi: 10.1007/s10924-006-0025-4
- 29. Essel, R., Engel, L., Carus, M. and Ahrens, R.H. (2015). Sources of microplastics relevant to marine protection in Germany. *Texte* 64, 656-660. doi: https://www.umweltbundesamt.de/sites/default/files/medien/378/publikationen/texte\_64\_ 922 2015 sources of microplastics relevant to marine protection 1.pdf
- 30. Evangeliou, N., Grythe, H., Klimont, Z., Heyes, C., Eckhardt, S., Lopez-Aparicio, S. and Stohl, A. (2020). Atmospheric transport is a major pathway of microplastics to remote regions. *Nature Communications* 11, 3381. doi: 10.1038/s41467-020-17201-9
- 31. FAOUN. (2021). Assessment of Agricultural Plastics and their Sustainability: A Call for Action. Rome. Rome. doi: 10.4060/cb7856en
- 32. Frias, J. and Nash, R. (2019). Microplastics: Finding a consensus on the definition. *Marine Pollution Bulletin* 138, 145-147. doi: 10.1016/j.marpolbul.2018.11.022
- 33. Fu, Z. and Wang, J. (2019). Current practices and future perspectives of microplastic pollution in freshwater ecosystems in China. *Science of the Total Environment* 691, 697- 712. doi: 10.1016/j.scitotenv.2019.07.167
- 34. Gao, H.H., Yan, C.R., Liu, Q., Ding, W.L., Chen, B.Q. and Li, Z. (2019). Effects of plasticmulching and plastic residue on agricultural production: A meta-analysis. *Science of the Total Environment* 651, 484-492. doi: 10.1016/j.scitotenv.2018.09.105
- 35. Geyer, R., Jambeck, J.R. and Law, K.L. (2017). Production, use, and fate of all plastics ever made. *Science Advances* 3, e1700782. doi: 10.1126/sciadv.1700782
- 36. Golwala, H., Zhang, X., Iskander, S. M., and Smith, A. L. (2021). Solid waste: An overlooked source of microplastics to the environment. *Science of the Total Environment*  769, 144581. doi: 10.1016/j.scitotenv.2020.144581
- 37. Gündoğdu, R., Önder, D., Gündoğdu, S. and Gwinnett, C. (2022). Microplastics Derived From Disposable Greenhouse Plastic Films and Irrigation Pipes: A Case Study From Turkey. *Environmental Science and Pollution Research*. doi: doi.org/10.21203/rs.3.rs-1282764/v1
- 38. Hahladakis, J.N., Velis, C.A., Weber, R., Iacovidou, E. and Purnell, P. (2018). An overview of chemical additives present in plastics: Migration, release, fate and environmental impact during their use, disposal and recycling. *Journal of Hazardous Materials* 344, 179-199. doi: 10.1016/j.jhazmat.2017.10.014
- 39. Halden, R.U. (2010). Plastics and health risks. *Annual review of public health* 31, 179-194. doi: 10.1146/annurev.publhealth.012809.103714
- 40. He, W., Li, Z., Yan, C., He, X., Li, Q. and Liu, Q. (2018). Development and prospect of biodegradable mulch films in China. *XXI International Congress on Plastics in Agriculture: Agriculture, Plastics and Environment* 1252, 47-50. doi: 10.17660/ActaHortic.2019.1252.6
- 41. Horton, A. A., Walton, A., Spurgeon, D. J., Lahive, E., & Svendsen, C. (2017).
- understanding to identify the knowledge gaps and future research priorities. *Science of the total environment* 586, 127-141. doi: 10.1016/j.scitotenv.2017.01.190
- 42. Hua, T., Kiran, S., Li, Y. and Sang, Q.-X.A. (2022). Microplastics exposure affects neural development of human pluripotent stem cell-derived cortical spheroids. *Journal of Hazardous Materials* 435, 128884. doi: 10.1016/j.jhazmat.2022.128884
- 43. Huang, Y., Liu, Q., Jia, W., Yan, C. and Wang, J. (2020). Agricultural plastic mulching as a source of microplastics in the terrestrial environment. *Environmental Pollution* 260, 114096. doi: 10.1016/j.envpol.2020.114096
- 44. Hurley, R.R. and Nizzetto, L. (2018). Fate and occurrence of micro(nano)plastics in soils: Knowledge gaps and possible risks. *Current Opinion in Environmental Science & Health* 1, 6-11. doi: 10.1016/j.coesh.2017.10.006
- 45. Jambeck, J.R., Geyer, R., Wilcox, C., Siegler, T.R., Perryman, M., Andrady, A., Narayan, R. and Law, K.L. (2015). Plastic waste inputs from land into the ocean. *Science* 347, 768- 771. doi: 10.1126/science.1260352
- 46. Katsumi, N., Kusube, T., Nagao, S. and Okochi, H. (2021). Accumulation of microcapsules derived from coated fertilizer in paddy fields. *Chemosphere* 267, 129185. doi: 10.1016/j.chemosphere.2020.129185
- 47. Kawecki, D. and Nowack, B. (2020). A proxy-based approach to predict spatially resolved emissions of macro- and microplastic to the environment. *Science of the Total Environment*  748, 141137. doi: 10.1016/j.scitotenv.2020.141137
- 48. Kershaw, P. (2015). Sources, fate and effects of microplastics in the marine environment: a global assessment. Report No. 1020-4873.
- 49. Kim, S. W., Waldman, W. R., Kim, T. Y., & Rillig, M. C. (2020). Effects of different microplastics on nematodes in the soil environment: tracking the extractable additives using an ecotoxicological approach. *Environmental Science & Technology* 54(21), 13868- 13878. https://doi.org/10.1021/acs.est.0c04641
- 50. Kole, P.J., Lohr, A.J., Van Belleghem, F. and Ragas, A.M.J. (2017). Wear and Tear of Tyres: A Stealthy Source of Microplastics in the Environment. *International Journal of Environmental Research and Public Health* 14, 1265. doi: 10.3390/ijerph14101265
- 51. Kumar, M., Xiong, X.N., He, M.J., Tsang, D.C.W., Gupta, J., Khan, E., Harrad, S., Hou, D.Y., Ok, Y.S. and Bolan, N.S. (2020). Microplastics as pollutants in agricultural soils. *Environmental Pollution* 265, 11. doi: 10.1016/j.envpol.2020.114980
- 52. Laner, D., Feketitsch, J., Rechberger, H. and Fellner, J. (2016). A novel approach to characterize data uncertainty in material flow analysis and its application to plastics flows in Austria. *Journal of industrial ecology* 20, 1050-1063. doi: 10.1111/jiec.12326
- 53. Lassen, C., Hansen, S.F., Magnusson, K., Norén, F., Hartmann, N.I.B., Jensen, P.R., Nielsen, T.G. and Brinch, A. (2015). Microplastics-Occurrence, effects and sources of releases to the environment in Danmark. *Danish Environmental Protection Agency: Copenhagen, Denmark* 2, 2.
- 54. Laycock, B., Nikolić, M., Colwell, J.M., Gauthier, E., Halley, P., Bottle, S. and George, G. (2017). Lifetime prediction of biodegradable polymers. *Progress in Polymer Science* 71, 144-189. doi: 10.1016/j.progpolymsci.2017.02.004
- 55. Le Moine, B. (2018). Worlwide Plasticulture a focus on films. Paper presented at 21st CIPA Congress. Arcachon, France. https://cipa-plasticulture.com/wp-
- content/uploads/2018/06/Worldwide-Plasticulture\_Le-Moine\_CIPA.pptx
- 56. Leslie, H.A., Van Velzen, M.J., Brandsma, S.H., Vethaak, A.D., Garcia-Vallejo, J.J. and Lamoree, M.H. (2022). Discovery and quantification of plastic particle pollution in human blood. *Environment International* 163, 107199. doi: 10.1016/j.envint.2022.107199
- 57. Li, C., Sun, M., Xu, X., Zhang, L., Guo, J. and Ye, Y. (2021). Environmental village regulations matter: Mulch film recycling in rural China. *Journal of cleaner production* 299, 126796. doi: 10.1016/j.jclepro.2021.126796
- 58. Li, J., Song, Y. and Cai, Y. (2020). Focus topics on microplastics in soil: Analytical methods, occurrence, transport, and ecological risks. *Environmental Pollution* 257. doi: 10.1016/j.envpol.2019.113570
- 59. Li, J., Wang, Y., Chen, Y., Gu, D., Lin, R. and Yang, X. (2022). Research advances on the accumulation and degradation of microcapsules derived from polymer-coated controlled- release fertilizers and their effects on soil quality. *Journal of Plant Nutrition and Fertilizers* 28, 1113-1121. doi: 10.11674/zwyf.2021591
- 60. Li, L.Z., Luo, Y.M., Li, R.J., Zhou, Q., Peijnenburg, W., Yin, N., Yang, J., Tu, C. and Zhang, Y.C. (2020). Effective uptake of submicrometre plastics by crop plants via a crack-entry mode. *Nature Sustainability* 3, 929-937. doi: 10.1038/s41893-020-0567-9
- 61. Li, S., Ding, F., Flury, M., Wang, Z., Xu, L., Li, S., Jones, D.L. and Wang, J. (2022). Macro- and microplastic accumulation in soil after 32 years of plastic film mulching. *Environmental Pollution* 300, 118945. doi: 10.1016/j.envpol.2022.118945
- 62. Li, X., Chen, L., Mei, Q., Dong, B., Dai, X., Ding, G. and Zeng, E.Y. (2018). Microplastics in sewage sludge from the wastewater treatment plants in China. *Water Research* 142, 75- 85. doi: 10.1016/j.watres.2018.05.034
- 63. Li, Z., Li, Q., Li, R., Zhou, J. and Wang, G. (2021). The distribution and impact of polystyrene nanoplastics on cucumber plants. *Environmental Science Pollution Research* 28, 16042-16053. doi: 10.1007/s11356-020-11702-2
- 64. Lian, J., Liu, W., Meng, L., Wu, J., Zeb, A., Cheng, L., Lian, Y. and Sun, H. (2021). Effects of microplastics derived from polymer-coated fertilizer on maize growth, rhizosphere, and soil properties. *Journal of cleaner production* 318, 128571. doi: 10.1016/j.jclepro.2021.128571
- 65. Liu, C., Li, J., Zhang, Y., Wang, L., Deng, J., Gao, Y., Yu, L., Zhang, J. and Sun, H. (2019). Widespread distribution of PET and PC microplastics in dust in urban China and their estimated human exposure. *Environment International* 128, 116-124. doi: 10.1016/j.envint.2019.04.024
- 66. Liu, K., Wang, X., Fang, T., Xu, P., Zhu, L. and Li, D. (2019). Source and potential risk assessment of suspended atmospheric microplastics in Shanghai. *Science of the Total Environment* 675, 462-471. doi: 10.1016/j.scitotenv.2019.04.110
- 67. Liu, Y., Zhai, X., and Zhou, M. (2020) Analysis of farmers'cognition on the value of agricultural membrane recycling and influencing factors. *Journal of Arid Land Resources and Environment* 34(02):80-87.doi:10.13448/j.cnki.jalre.2020.39
- 68. Liu, Y., Guo, R., Zhang, S., Sun, Y. and Wang, F. (2022). Uptake and translocation of nano/microplastics by rice seedlings: Evidence from a hydroponic experiment. *Journal of Hazardous Materials* 421, 126700. doi: 10.1016/j.jhazmat.2021.126700
- 69. Liu, Y., Zhou, C., Li, F., Liu, H. and Yang, J. (2020). Stocks and flows of polyvinyl chloride
- (PVC) in China: 1980-2050. *Resources, Conservation Recycling* 154, 104584. doi: 10.1016/j.resconrec.2019.104584
- 70. Lofty, J., Muhawenimana, V., Wilson, C. and Ouro, P. (2022). Microplastics removal from a primary settler tank in a wastewater treatment plant and estimations of contamination onto European agricultural land via sewage sludge recycling. *Environmental Pollution* 304, 119198. doi: 10.1016/j.envpol.2022.119198
- 71. Lu, B., Jiang, C., Chen, Z., Li, A., Wang, W., Zhang, S., and Luo, G. (2022). Fate of polylactic acid microplastics during anaerobic digestion of kitchen waste: Insights on property changes, released dissolved organic matters, and biofilm formation. *Science of The Total Environment* 834, 155108. 10.1016/j.scitotenv.2022.155108
- 72. Luan, X., Cui, X., Zhang, L., Chen, X., Li, X., Feng, X., Chen, L., Liu, W. and Cui, Z. (2021). Dynamic material flow analysis of plastics in China from 1950 to 2050. *Journal of cleaner production* 327, 129492. doi: 10.1016/j.jclepro.2021.129492
- 73. Luan, X., Kou, X., Zhang, L., Chen, L., Liu, W. and Cui, Z. (2022). Estimation and prediction of plastic losses to the environment in China from 1950 to 2050. *Resources, Conservation Recycling* 184, 106386. doi: 10.1016/j.resconrec.2022.106386
- 74. Lubkowski, K., Smorowska, A., Markowska-Szczupak, A. and Ukielski, R. (2016). Copolyester-coated mineral fertilizers–preparation, characterization, and nutrient release. *Toxicological Environmental Chemistry* 98, 1163-1172. doi: 10.1080/02772248.2015.1130225
- 75. Lv, W., Zhou, W., Lu, S., Huang, W., Yuan, Q., Tian, M., Lv, W. and He, D. (2019). Microplastic pollution in rice-fish co-culture system: A report of three farmland stations in Shanghai, China. *Science of the Total Environment* 652, 1209-1218. doi: 10.1016/j.scitotenv.2018.10.321
- 76. Lwanga, E.H., Vega, J.M., Quej, V.K., Chi, J.D., del Cid, L.S., Chi, C., Segura, G.E., Gertsen, H., Salanki, T., van der Ploeg, M., Koelmans, A.A. and Geissen, V. (2017). Field evidence for transfer of plastic debris along a terrestrial food chain. *Scientific Reports* 7, 14071. doi: 10.1038/s41598-017-14588-2
- 77. Mahon, A. M., O'Connell, B., Healy, M. G., O'Connor, I., Officer, R., Nash, R., and Morrison, L. (2017). Microplastics in sewage sludge: effects of treatment. *Environmental Science & Technology* 51(2), 810-818.doi:10.1021/acs.est.6b04048
- 78. Maraveas, C. (2020). Environmental sustainability of plastic in agriculture. *Agriculture* 10, 310. doi: 10.3390/agriculture10080310
- **79.** Massos, A. and Turner, A. (2017). Cadmium, lead and bromine in beached microplastics. *Environmental Pollution* 227, 139-145. doi: 10.1016/j.envpol.2017.04.034
- 80. Mekonnen, T., Misra, M. and Mohanty, A.K. (2016). Fermented soymeals and their reactive blends with poly (butylene adipate-co-terephthalate) in engineering biodegradable cast films for sustainable packaging. *ACS Sustainable Chemistry Engineering* 4, 782-793. doi: 10.1021/acssuschemeng.5b00782
- 81. Meng, F., Yang, X., Riksen, M., Xu, M., and Geissen, V. (2021). Response of common bean (Phaseolus vulgaris L.) growth to soil contaminated with microplastics. *Science of the Total Environment,* 755, 142516. doi: 10.1016/j.scitotenv.2020.142516
- 82. MEPC. (2017). China Environmental Statistical Yearbook (in Chinese). Beijing: China Statistics Press.
- 83. Moharir, R. V., and Kumar, S. (2019). Challenges associated with plastic waste disposal and allied microbial routes for its effective degradation: A comprehensive review. *Journal of Cleaner Production* 208, 65-76. doi: 10.1016/j.jclepro.2018.10.059
- 84. Montangero, A. and Belevi, H. (2008). An approach to optimise nutrient management in environmental sanitation systems despite limited data. *Journal of Environmental Management* 88, 1538-1551. doi: 10.1016/j.jenvman.2007.07.033
- 85. Ministry of Agriculture and Rural Affairs of the People 's Republic of China. Reply to the No.2357 suggestion of the First Session of the Thirteenth National People 's Congress.[2018-07-10].http://www.moa.gov.cn/gk/jyta/201807/t20180713\_6154037.htm
- 86. NBSC. (2011−2022).China Rural Statistical Yearbook (in Chinese). Beijing: China Statistics Press.
- 87. Ng, E. L., Lin, S. Y., Dungan, A. M., Colwell, J. M., Ede, S., Lwanga, E. H., Meng, K., Geissen, V., Blackall, L. L., and Chen, D. (2021). Microplastic pollution alters forest soil microbiome. *Journal of hazardous materials*, 409, 124606. doi: 10.1016/j.jhazmat.2020.124606
- 88. Ng, E.L., Lwanga, E.H., Eldridge, S.M., Johnston, P., Hu, H.W., Geissen, V. and Chen, D.L. (2018). An overview of microplastic and nanoplastic pollution in agroecosystems. *Science of the Total Environment* 627, 1377-1388. doi: 10.1016/j.scitotenv.2018.01.341
- 89. Nizzetto, L., Futter, M. and Langaas, S. (2016). Are Agricultural Soils Dumps for Microplastics of Urban Origin? *Environmental Science & Technology* 50, 10777-10779. doi: 10.1021/acs.est.6b04140
- 90. Pérez-Reverón, R., González-Sálamo, J., Hernández-Sánchez, C., González-Pleiter, M., Hernández-Borges, J., and Díaz-Peña, F. J. (2022). Recycled wastewater as a potential source of microplastics in irrigated soils from an arid-insular territory (Fuerteventura, Spain). *Science of The Total Environment* 817, 152830. doi: 1113 10.1016/j.scitotenv.2021.152830
- 91. Plastics Europe-The Facts (2022). https://plasticseurope.org/knowledge-hub/plastics-the-facts-2022/
- 92. Porterfield, K. K., Hobson, S. A., Neher, D. A., Niles, M. T., and Roy, E. D. (2023). Microplastics in composts, digestates, and food wastes: a review. Journal of Environmental Quality 52(2), 225-240. doi: 10.1002/jeq2.20450
- 93. Qi, R.M., Jones, D.L., Li, Z., Liu, Q. and Yan, C.R. (2020). Behavior of microplastics and plastic film residues in the soil environment: A critical review. *Science of the Total Environment* 703, 12. doi: 10.1016/j.scitotenv.2019.134722
- 94. Qi, Y., Yang, X., Mejia Pelaez, A., Huerta Lwanga, E., Beriot, N., Gertsen, H., Garbeva, P. and Geissen, V. (2018). Macro- and micro- plastics in soil-plant system: Effects of plastic mulch film residues on wheat (Triticum aestivum) growth. *Science of the Total Environment* 645, 1048-1056. doi: 10.1016/j.scitotenv.2018.07.229
- 95. Qin, M., Chen, C., Song, B., Shen, M., Cao, W., Yang, H., Zeng, G., and Gong, J. (2021). A review of biodegradable plastics to biodegradable microplastics: Another ecological threat to soil environments?. *Journal of Cleaner Production* 312, 127816. doi: 10.1016/j.jclepro.2021.127816
- 96. Rachman, C.M. (2018). Microplastics research from sink to source. *Science* 360, 28-29.
- doi: 10.1126/science.aar7734
- 97. Ren, S.-Y., Kong, S.-F. and Ni, H.-G. (2021). Contribution of mulch film to microplastics in agricultural soil and surface water in China. *Environmental Pollution* 291, 118227. doi: 10.1016/j.envpol.2021.118227
- 98. Ren, S.-Y., Sun, Q., Ni, H.-G. and Wang, J. (2020). A minimalist approach to quantify emission factor of microplastic by mechanical abrasion. *Chemosphere* 245, 125630. doi: 10.1016/j.chemosphere.2019.125630
- 99. Rezaei, M., Riksen, M.J.P.M., Sirjani, E., Sameni, A. and Geissen, V. (2019). Wind erosion as a driver for transport of light density microplastics. *Science of the Total Environment* 669, 273-281. doi: 10.1016/j.scitotenv.2019.02.382
- 100.Rillig, M. C. (2012). Microplastic in terrestrial ecosystems and the soil? *Environmental Science & Technology* 46(12):6453-4. doi: 10.1021/es302011r
- 101.Rist, S., Almroth, B.C., Hartmann, N.B. and Karlsson, T.M. (2018). A critical perspective on early communications concerning human health aspects of microplastics. *Science of the Total Environment* 626, 720-726. doi: 10.1016/j.scitotenv.2018.01.092
- 102.Rochman, C.M., Browne, M.A., Underwood, A.J., van Franeker, J.A., Hompson, R.C.T. and Amaral-Zettler, L.A. (2016). The ecological impacts of marine debris: unraveling the demonstrated evidence from what is perceived. *Ecology* 97, 302-312. doi: 10.1890/14- 2070.1
- 103.Sharma, M.D., Elanjickal, A.I., Mankar, J.S. and Krupadam, R.J. (2020). Assessment of cancer risk of microplastics enriched with polycyclic aromatic hydrocarbons. *Journal of Hazardous Materials* 398, 10. doi: 10.1016/j.jhazmat.2020.122994
- 104.Sholokhova, A., Ceponkus, J., Sablinskas, V., and Denafas, G. (2021). Abundance and characteristics of microplastics in treated organic wastes of Kaunas and Alytus regional waste management centres, Lithuania. *Environmental Science and Pollution Research* 1- 10. doi: 10.1007/s11356-021-17378-6
- 105.Sieber, R., Kawecki, D. and Nowack, B. (2020). Dynamic probabilistic material flow analysis of rubber release from tires into the environment. *Environmental Pollution* 258, 113573. doi: 10.1016/j.envpol.2019.113573
- 106.Song, Y.K., Hong, S.H., Jang, M., Han, G.M., Jung, S.W. and Shim, W.J. (2017). Combined effects of UV exposure duration and mechanical abrasion on microplastic fragmentation by polymer type. *Environmental Science Technology* 51(8), 4368-4376. doi: 10.1021/acs.est.6b06155
- 107.Thompson, R.C., Olsen, Y., Mitchell, R.P., Davis, A., Rowland, S.J., John, A.W.G., McGonigle, D. and Russell, A.E. (2004). Lost at sea: Where is all the plastic? *Science* 304, 838-838. doi: 10.1126/science.1094559
- 108.Trenkel, M.E. (2010). *Slow-and controlled-release and stabilized fertilizers: an option for enhancing nutrient use efficiency in agriculture*: IFA, International fertilizer industry association.
- 109.Tsai, C.L. and Krogmann, U. (2013). Material flows and energy analysis of glass containers discarded in New Jersey, USA. *Journal of industrial ecology* 17, 129-142. doi: 1172 10.1111/j.1530-9290.2012.00509.x
- 110.Uheida, A., Mejía, H.G., Abdel-Rehim, M., Hamd, W. and Dutta, J. (2021). Visible light photocatalytic degradation of polypropylene microplastics in a continuous water flow system. *Journal of Hazardous Materials* 406, 124299. doi: 10.1016/j.jhazmat.2020.124299
- 111.Wang, C., Liu, Y., Chen, W.Q., Zhu, B., Qu, S. and Xu, M. (2021). Critical review of global plastics stock and flow data. *Journal of industrial ecology* 25, 1300-1317. doi: doi.org/10.1111/jiec.13125
- 112.Wang, J., Coffin, S., Sun, C.L., Schlenk, D. and Gan, J. (2019). Negligible effects of microplastics on animal fitness and HOC bioaccumulation in earthworm Eisenia fetida in soil. *Environmental Pollution* 249, 776-784. doi: 10.1016/j.envpol.2019.03.102
- 113.Wang, J., Li, J., Liu, S., Li, H., Chen, X., Peng, C., Zhang, P. and Liu, X. (2021). Distinct microplastic distributions in soils of different land-use types: A case study of Chinese farmlands. *Environmental Pollution* 269, 116199. doi: 10.1016/j.envpol.2020.116199
- 114.Wang, J., Lv, S.H., Zhang, M.Y., Chen, G.C., Zhu, T.B., Zhang, S., Teng, Y., Christie, P. and Luo, Y.M. (2016). Effects of plastic film residues on occurrence of phthalates and microbial activity in soils. *Chemosphere* 151, 171-177. doi: 10.1016/j.chemosphere.2016.02.076
- 115.Wang, K., Chen, W., Tian, J., Niu, F., Xing, Y., Wu, Y., Zhang, R., Zheng, J. and Xu, L. (2022). Accumulation of microplastics in greenhouse soil after long-term plastic film mulching in Beijing, China. *Science of the Total Environment* 828, 154544. doi: 10.1016/j.scitotenv.2022.154544
- 116.Wang, K., Li, J., Zhao, L., Mu, X., Wang, C., Wang, M., Xue, X., Qi, S. and Wu, L. (2021). Gut microbiota protects honey bees (Apis mellifera L.) against polystyrene microplastics exposure risks. *Journal of Hazardous Materials* 402, 123828. doi: 10.1016/j.jhazmat.2020.123828
- 117.Wang, L., Li, P., Zhang, Q., Wu, W. M., Luo, J., and Hou, D. (2021). Modeling the conditional fragmentation-induced microplastic distribution. *Environmental Science & Technology* 55(9), 6012-6021. doi: 10.1021/acs.est.1c01042
- 118.Wang, Y. and Ma, H.-w. (2018). Analysis of uncertainty in material flow analysis. *Journal of cleaner production* 170, 1017-1028. doi: 10.1016/j.jclepro.2017.09.202
- 119.Weithmann, N., Moller, J.N., Loder, M.G.J., Piehl, S., Laforsch, C. and Freitag, R. (2018). Organic fertilizer as a vehicle for the entry of microplastic into the environment. *Science Advances* 4, 7. doi: 10.1126/sciadv.aap8060
- 120.Wu, P., Lin, S., Cao, G., Wu, J., Jin, H., Wang, C., Wong, M.H., Yang, Z. and Cai, Z. (2022). Absorption, distribution, metabolism, excretion and toxicity of microplastics in the human body and health implications. *Journal of Hazardous Materials*, 129361. doi: 10.1016/j.jhazmat.2022.129361
- 121.Wu, S.L., Xu, A.M., Zhou, J., Xin, F. X., Yu, Z. Y., Dong, W. L., and Jiang, M. (2022). Microplastics in wastewater treatment: current status and future trends. *Journal of Biotechnology* 38(7): 2410-2422. doi: 10.13345/j.cjb.210918
- 122.Xu, F., Li, Y., Ge, X., Yang, L. and Li, Y. (2018). Anaerobic digestion of food waste– Challenges and opportunities. *Bioresource technology* 247, 1047-1058. doi: 10.1016/j.biortech.2017.09.020
- 123.Yan, C., Mei, X., He, W. and Zheng, S. (2006). Present situation of residue pollution of mulching plastic film and controlling measures. *Transactions of the Chinese Society of Agricultural Engineering* 22, 269-272.
- 124.Yan, C. (2022). The basic situation of agricultural plastic film application in China and the effect of plastic film on soil and agricultural production.
- https://www.thepaper.cn/newsDetail\_forward\_18374301
- 125.Yang, G., Zhang, G. and Wang, H. (2015). Current state of sludge production, management, treatment and disposal in China. *Water Research* 78, 60-73. doi: 10.1016/j.watres.2015.04.002
- 126.Yang, J., Li, L., Zhou, Q., Li, R., Tu, C. and Luo, Y. (2021). Microplastics Contamination of Soil Environment: Sources, Processes and Risks. *Acta Pedologica Sinica* 58, 281-298. doi: 10.11766/trxb202006090286
- 127.Yang, J., Li, L., Li, R., Xu, L., Shen, Y., Li, S., Tu, C., Wu, L., Christie, P. and Luo, Y. (2021). Microplastics in an agricultural soil following repeated application of three types of sewage sludge: A field study. *Environmental Pollution* 289, 117943. doi: 10.1016/j.envpol.2021.117943
- 128.Yang, X., Cao, Y., Jiang, R. and Zhang, F. (2009). Effect of components of polyethylene solution on release characteristics of controlled-release fertilizer. *Chemical Engineering (China)* 37, 44-47.
- 129.Yang, Y., Li, Z., Yan, C., Chadwick, D., Jones, D.L., Liu, E., Liu, Q., Bai, R. and He, W. (2022). Kinetics of microplastic generation from different types of mulch films in agricultural soil. *Science of the Total Environment* 814, 152572. doi: 10.1016/j.scitotenv.2021.152572
- 130.Ye, X., Wang, P., Wu, Y., Zhou, Y., Sheng, Y. and Lao, K. (2020). Microplastic acts as a vector for contaminants: the release behavior of dibutyl phthalate from polyvinyl chloride pipe fragments in water phase. *Environmental Science Pollution Research* 27, 42082- 42091. doi: 10.1007/s11356-020-10136-0
- 131.Yuan, J., Ma, J., Sun, Y., Zhou, T., Zhao, Y., and Yu, F. (2020). Microbial degradation and other environmental aspects of microplastics/plastics. *Science of the Total Environment* 715, 136968. doi: 10.1016/j.scitotenv.2020.136968
- 132.Zettler, E.R., Mincer, T.J. and Amaral-Zettler, L.A. (2013). Life in the "Plastisphere": Microbial Communities on Plastic Marine Debris. *Environmental Science & Technology* 47, 7137-7146. doi: 10.1021/es401288x
- 133.Zhan, Y. R., Chen, Y., Xu, W. T., and Qian, Q. (2017). The current status and recycling countermeasures of the wasted pesticide packaging materials in China. *Recyclable Resources and Circular Economy* 10(7): 17-21
- 134.Zhang, J., Ren, S., Xu, W., Liang, C., Li, J., Zhang, H., Li, Y., Liu, X., Jones, D.L. and Chadwick, D. (2022). Effects of plastic residues and microplastics on soil ecosystems: A global meta-analysis. *Journal of Hazardous Materials*, 129065. doi: 10.1016/j.jhazmat.2022.129065
- 135.Zhang, S., Li, Y., Chen, X., Jiang, X., Li, J., Yang, L., Yin, X., and Zhang, X. (2022). Occurrence and distribution of microplastics in organic fertilizers in China. *Science of the Total Environment,* 844, 157061. doi: 10.1016/j.scitotenv.2022.157061
- 136.Zhang, J.J., Wang, L., Trasande, L. and Kannan, K. (2021). Occurrence of Polyethylene Terephthalate and Polycarbonate Microplastics in Infant and Adult Feces. *Environmental Science & Technology Letters* 8, 989-994. doi: 10.1021/acs.estlett.1c00559
- 137.Zhang, Q.-Q., Ma, Z.-R., Cai, Y.-Y., Li, H.-R. and Ying, G.-G. (2021). Agricultural plastic pollution in China: generation of plastic debris and emission of phthalic acid esters from agricultural films. *Environmental Science Technology* 55, 12459-12470. doi:
- 10.1021/acs.est.1c04369
- 138.Zhao, Y., Chen, X., Wen, H., Zheng, X., Niu, Q. and Kang, J. (2017). Research Status and Prospect of Control Technology for Residual Plastic Film Pollution in Farmland. *Transactions of the Chinese Society for Agricultural Machinery* 48, 1-14. doi: 10.6041/j.issn.1000-1298.2017.06.001
- 139.Zhou, B., Wang, J., Zhang, H., Shi, H., Fei, Y., Huang, S., Tong, Y., Wen, D., Luo, Y., and Barceló, D. (2020). Microplastics in agricultural soils on the coastal plain of Hangzhou Bay, east China: Multiple sources other than plastic mulching film. *Journal of Hazardous Materials*, *388*, 121814. doi: 10.1016/j.jhazmat.2019.121814
- 140.Zhou, C.-Q., Lu, C.-H., Mai, L., Bao, L.-J., Liu, L.-Y. and Zeng, E.Y. (2021). Response of rice (Oryza sativa L.) roots to nanoplastic treatment at seedling stage. *Journal of Hazardous Materials* 401, 123412. doi: 10.1016/j.jhazmat.2020.123412
- 141.Zhou, J., Jia, R., Brown, R. W., Yang, Y., Zeng, Z., Jones, D. L., & Zang, H. (2023). The long-term uncertainty of biodegradable mulch film residues and associated microplastics pollution on plant-soil health. *Journal of Hazardous Materials*, 442, 130055. doi: 10.1016/j.jhazmat.2022.130055
- 142.Zhou, Y., Liu, X. and Wang, J. (2019). Characterization of microplastics and the association of heavy metals with microplastics in suburban soil of central China. *Science of the Total Environment* 694. doi: 10.1016/j.scitotenv.2019.133798
- 143.Zhou, Y., Yang, N., Hu, S. (2013). Industrial metabolism of PVC in China: A dynamic material flow analysis. *Resources, Conservation Recycling* 73, 33-40. doi: 10.1016/j.resconrec.2012.12.016
- 144.Ziajahromi, S., Neale, P.A. and Leusch, F.D. (2016). Wastewater treatment plant effluent as a source of microplastics: review of the fate, chemical interactions and potential risks to aquatic organisms. *Water science technology* 74, 2253-2269. doi: 10.2166/wst.2016.414