

Needles in haystacks: monitoring the potential escape of bioaerosolised antimicrobial resistance genes from wastewater treatment plants with air and phyllosphere sampling

George, Paul; Hillary, Luke; Leclerc, Samantha; Cooledge, Emily; Lemieux, Joanie; Duchaine, Caroline; Jones, Davey L.

Canadian Journal of Microbiology

DOI:
[10.1139/cjm-2023-0226](https://doi.org/10.1139/cjm-2023-0226)

Published: 12/04/2024

Peer reviewed version

[Cyswllt i'r cyhoeddiad / Link to publication](#)

Dyfyniad o'r fersiwn a gyhoeddwyd / Citation for published version (APA):
George, P., Hillary, L., Leclerc, S., Cooledge, E., Lemieux, J., Duchaine, C., & Jones, D. L. (2024). Needles in haystacks: monitoring the potential escape of bioaerosolised antimicrobial resistance genes from wastewater treatment plants with air and phyllosphere sampling. *Canadian Journal of Microbiology*. <https://doi.org/10.1139/cjm-2023-0226>

Hawliau Cyffredinol / General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

1 **Needles in haystacks: monitoring the potential escape of bioaerosolised antimicrobial**
2 **resistance genes from wastewater treatment plants with air and phyllosphere sampling**

3
4 Paul B. L. George^{a,b*}, Luke Hillary^c, Samantha Leclerc^{a,b}, Emily C. Cooledge^d, Joanie Lemieux^{a,b},
5 Caroline Duchaine^{a,b}, and Davey L. Jones^{d,e}

6
7 ^aDépartement de biochimie, de microbiologie et de bio-informatique, Université Laval, Quebec
8 City, QC, G1V 0A6, Canada

9 ^bCentre de recherche de l'Institut universitaire de cardiologie et de pneumologie, Quebec City,
10 QC, G1V 4G5, Canada

11 ^cDepartment of Plant Pathology, University of California, Davis, CA 95616, USA

12 ^dSchool of Environmental and Natural Sciences, Bangor University, Bangor, Gwynedd, LL57
13 2UW, UK

14 ^eFood Futures Institute, Murdoch University, Murdoch, WA 6150, Australia

15
16 *Corresponding author: Paul B.L. George (email: paul.george@bcm.ulaval.ca)

17 **Abstract**

18 Wastewater treatment plants are well known point sources of emissions of antimicrobial resistance
19 genes (ARGs) into the environment. Although most work to date has focused on ARG dispersal
20 via effluent, aerial dispersal in bioaerosols is a poorly understood, but likely important vector for
21 ARG dispersal. Recent evidence suggests that ARG profiles of the conifer needle phyllosphere
22 could be used to measure bioaerosol dispersal from anthropogenic sources. Here we assessed
23 airborne dispersal of ARGs from wastewater treatment plants in Wales, UK and Quebec, Canada,

24 using conifer needles as passive bioaerosol monitors. ARG profiles of wastewater were compared
25 to those of conifer phyllosphere using high-throughput qPCR. ARG richness was significantly
26 lower in conifer phyllosphere samples than wastewater samples, though no differences were
27 observed across the dispersal gradients. Mean copy number of ARGs followed a similar trend.
28 ARG profiles showed limited, but consistent patterns with increasing distance from wastewater
29 treatment plants, but these did not align with those of wastewater samples. For example,
30 proportional abundance of aminoglycosides decreased over the dispersal gradient in Wales,
31 whereas mobile genetic elements showed the inverse relationship. In summary, while distinct ARG
32 profiles exist along dispersal gradients, links to those of wastewater were not apparent.

33
34 Les usines de traitement des eaux usées sont de potentielles sources d'émission de gènes de
35 résistance aux antibiotiques (ARGs) dans l'environnement. La majorité des travaux publiés à ce
36 jour se concentrent principalement sur la dispersion des ARGs dans les effluents, mais la
37 dispersion des ARGs peut également se produire dans l'air sous forme de bioaérosols. L'air comme
38 mode de transport des ARGs est encore relativement peu étudié et mal compris. De récentes études
39 ont démontré que la phyllosphère des aiguilles de conifères pouvait être utilisée pour détecter et
40 identifier le profil d'ARGs retrouvé dans l'air à proximité de sources anthropiques. La présente
41 étude se concentrait sur l'évaluation de la dispersion des ARGs près d'usines de traitement des
42 eaux usées situées au Pays de Galles (Royaume-Uni) et au Québec (Canada), en utilisant des
43 aiguilles de conifères comme échantillonneurs passifs de bioaérosols. La qPCR à haut débit a été
44 utilisée pour comparer le profil d'ARGs des eaux usées avec celui de la phyllosphère des aiguilles
45 de conifères. La richesse en ARGs était significativement plus faible dans les échantillons de
46 phyllosphère des aiguilles que dans les échantillons d'eaux usées, bien qu'aucune différence n'ait

47 été observée parmi les différents gradients de dispersion. Le nombre moyen de copies d'ARGs
48 était relativement similaire. Les profils d'ARGs sur les aiguilles de conifères tendent à changer
49 plus la distance de la source augmente. Par exemple, la proportion de gènes de résistance aux
50 aminoglycosides tend à diminuer en s'éloignant de l'usine du Pays de Galles tandis que les
51 éléments génétiques mobiles (MGEs) suivaient une tendance contraire. En somme, bien que des
52 profils d'ARGs spécifiques peuvent être observés selon les gradients de dispersion, aucun lien clair
53 ne peut être établi entre les profils des aiguilles et ceux des eaux usées. D'autres recherches devront
54 être menées pour évaluer à plus long terme les changements dans les profils d'ARGs et la
55 biodiversité bactérienne pour déterminer de potentiels contaminants aéroportés.

56
57 **Keywords:** Conifers; Biomonitoring; Environmental AMR; High-throughput qPCR; Bioaerosols

59 **Introduction**

60 In 2019, an estimated 4.95 million deaths were caused by infections by antimicrobial
61 resistant (AMR) bacteria globally (Murray et al., 2022). Unfortunately, the consequences of
62 increasing AMR extend further than simple mortality. As of 2019, AMR is estimated to cost the
63 European Union €1.5 billion annually in lost economic productivity and healthcare spending
64 (Anderson et al., 2019). In Canada, the present costs of AMR are similar (1.4 billion CAD), but
65 are expected to increase to 8 billion CAD annually by 2050 (Council of Canadian Academics,
66 2019). Yet despite increasing efforts to monitor AMR bacterial infections in human and animal
67 health, there is a clear lack of monitoring capacity to track the spread of antibacterial resistance
68 genes (ARGs) into the environment (Pruden et al., 2021). Improving environmental monitoring
69 capacity for AMR would adhere to the One Health ethos, which calls for multidisciplinary and

70 intersectoral approaches to protect human, animal, and environmental health (Pokharel et al., 2020;
71 Robinson et al., 2016). Monitoring wastewater effluent using molecular methods can effectively
72 quantify ARG loads and track their dispersal through the environment (Pruden et al., 2021). In
73 turn, this could provide early detection of risks associated with AMR bacteria helping to prevent
74 outbreaks or promote prescribing-based interventions.

75 Indeed, the dispersal of AMR in the wider environment from anthropogenic sources has
76 received much attention, particularly the contamination of soils and waterways (Singer et al., 2016;
77 Wu et al., 2023). Wastewater treatment plants (WWTPs) in particular, are well-known point-
78 sources for the discharge of ARM bacteria and associated mobile genetic elements (MGEs)
79 carrying ARGs (Singer et al., 2016; Shi et al., 2023; de Nies et al., 2022). Several studies have
80 demonstrated that sites near WWTPs harbour greater quantities of ARGs, thus posing a risk for
81 primary human exposure and secondary exposure through contamination of the food chain
82 (Czekalski et al., 2014; Marti et al., 2013; Gaviria-Figueroa et al., 2019). Further, the totality of
83 ARGs and MGEs of wastewater effluent is often enhanced during the treatment process (Ju et al.,
84 2019; Corno et al., 2019; Raza et al., 2022; Shi et al., 2023; Thornton, 2020). As such, resilient
85 AMR bacteria can be recovered in the downstream water column (Amos et al., 2014; Munir et al.,
86 2011) and can propagate ARGs in the wider environment through horizontal gene transfer
87 (Jacquiod et al., 2017). Given the fact that the release of AMR bacteria from WWTPs can exceed
88 10^{10} CFU day⁻¹ (Manaia et al., 2010; Novo and Manaia, 2010), it is vital that these point sources
89 of AMR pollutants be monitored.

90 There is growing evidence that aerial dispersal of AMR bacteria may represent an
91 important vector for ARGs to enter the environment (Bai et al., 2022; Gaviria-Figueroa et al.,
92 2019). During the wastewater treatment process, microorganisms become aerosolised through

93 aeration, mechanical mixing, and bubble bursting (Kataki et al., 2022), with the airborne bacterial
94 biomass ranging between 10^6 - 10^8 genomic units (based on 16S rRNA copy numbers) per m^3 of air
95 (Bélanger Cayouette et al., 2022; Blais-Lecours et al., 2014; Mbareche et al., 2022). Many families
96 of ARGs have been detected in WWTP bioaerosols using metagenomics (Gaviria-Figueroa et al.,
97 2019; Yang et al., 2018) and quantitative PCR (qPCR) approaches (e.g. Bélanger Cayouette et al.,
98 2022; Ouradou et al., 2023; Wang et al., 2019). Though information on long-distance bioaerosol
99 dispersal is limited, Gaviria-Figueroa et al. (2019) used a modelling approach to estimate that over
100 a 24 h period more than 220,000 individual ARGs could be transported across a 10 km radius
101 surrounding an American WWTP capable of processing 15 million L of waste per day, with a
102 further 10,000 ARG copies transported up to 120 km away. This suggests that WWTP bioaerosols
103 have the potential impact microbial communities over an incredibly large area. More work is
104 therefore needed to monitor the long-distance dispersal of WWTPs bioaerosols and their potential
105 roles as environmental contaminants.

106 In comparison to waterborne vectors, the extent to which bioaerosols contribute to the
107 propagation of AMR in the wider environment remains unclear. Further, new modelling
108 approaches are needed to quantify the risk posed by airborne ARGs. Recent work has shown that
109 vegetation can be used as passive collectors when sampling bioaerosols (Theofel et al., 2020;
110 George et al., 2022). This mirrors the use of conifer trees as barriers around point sources of
111 pollution to limit emissions of fine particulate matter, ammonia, and odours (Adrizal et al., 2008;
112 Lin et al., 2006). For example, Galès et al. (2014) delineated the dispersal patterns of
113 *Saccharopolyspora rectivirgula* emissions from a composting plant along a gradient of pine trees.
114 Assessments of the conifer phyllosphere have also revealed different ARG profiles in agricultural
115 and forest regions of Quebec, Canada (George et al., 2022a). Harnessing the conifer phyllosphere

116 as a monitoring tool could allow for long-term monitoring of ARGs in bioaerosols, contributing to
117 the assessment of anthropogenic influences on AMR bacteria and ARG dispersal (Manibusan and
118 Mainelis, 2022). Indeed, this method draws inspiration from the use of biofilms as environmental
119 markers of ARG dispersal from wastewater effluent (Engemann et al., 2008; Marti et al., 2013). It
120 is assumed that ARGs will be transferred via colonisation and plasmid exchange on conifer needles
121 in the same way as ARGs are accumulated in biofilms as AMR bacteria settle out of the
122 contaminated water column (Engemann et al., 2008).

123 Here, the ARGs profiles of wastewater, air, and the conifer phyllosphere were assessed
124 along a dispersal gradient at two contrasting WWTPs in Quebec, Canada and Wales, UK. Our two
125 main objectives were to: *i*) assess whether the conifer phyllosphere could delineate ARG dispersal
126 from WWTPs; *ii*) assess differences in ARG profiles could be detected using a common panel of
127 ARGs. We hypothesised that the ARG profile of conifer needles would resemble those of WWTPs
128 but decrease with distance away from the WWTP. Further, it was hypothesised that our common
129 ARG panel was diverse enough to be applicable to both sampling regions.

130

131 **Materials and methods**

132 *Study sites*

133 The study focused on two exemplar sites, one located in Quebec, Canada (Site 1) and one
134 located in Bangor, North Wales, UK (Site 2). Both WWTPs are open air treatment systems with
135 primary settling and secondary aeration ponds where bioaerosols can directly enter the
136 environment.

137 Site 1: The WWTP in Quebec was located at Sainte-Brigitte-de-Laval (N47°00'00",
138 W71°12'00"). The site serves a population of 8,500 and has a daily flow of 1041.2 l day⁻¹. In

139 September 2022, wastewater samples were collected from one of the settling lagoons over the
140 course of 3 h. Air samples were collected from 3 points across the WWTP simultaneously with
141 wastewater samples in a 200 m diagonal transect within the WWTP. Conifer needles were
142 collected from trees surrounding the WWTP in a ~150 m radius. Due to the geography of the area
143 surrounding Sainte-Brigitte-de-Laval WWTP, it was not possible to follow the downwind
144 dispersal of bioaerosols in this region. Air and conifer needles were also collected from an upwind
145 control site (Lac-Beauport, QC, Canada) located ~10.5 km from the Sainte-Brigitte-de-Laval
146 WWTP, to serve as reference samples for the environment surrounding the WWTP (Fig. 1).

147 Site 2: The WWTP in North Wales was located at Treborth, Gwynedd, (N53°12'34",
148 W04°10'59"). The WWTP serves a population of 40,000 and has a daily average flow of 285 l s⁻¹.
149 In July 2022, air and conifer needle samples were collected from sites across a ~16 km transect
150 in North Wales, UK. Conifer needle samples were classified as upwind (Y Felinheli) (within a 3.8
151 km radius), in proximity to the WWTP (within a < 1 km radius), in Bangor city centre (~4.2 km
152 from WWTP), or distant (Beumaris and Abergwyngregyn sites, ~7.5 km and ~11 km from
153 WWTP site, respectively; Fig. 1). A control sample of conifer needles was taken 65 km away from
154 the WWTP at Lake Vyrnwy, Powys, in Mid Wales (Fig. 1).

155 *Wastewater sampling*

156 At both WWTPs, three wastewater samples were collected in polypropylene bottles over the
157 course of 1 h, with 100 ml of wastewater collected every 10 mins to produce one composite 700
158 ml sample. Samples were stored at 4 °C until analysis. Prior to DNA extraction, 100 ml of
159 wastewater per composite sample was centrifuged for 10 min at 10,000 x g at a temperature of
160 4°C. Supernatants were discarded and the pellet resuspended in PowerSoil Pro lysis buffer C1.
161 DNA extraction was performed immediately following processing, as described below.

162 Briefly, wastewater pH and electrical conductivity were determined using standard
163 electrodes. Ammonium (NH_4^+) and orthophosphate (PO_4^{3-}) were determined colorimetrically as
164 described in Mulvaney (1996) and Murphy and Riley (1962), respectively. Turbidity was
165 measured with an Orion™ AQ4500 Turbidimeter (ThermoFisher Scientific, UK). Chemical
166 properties of wastewater samples are presented in Table 1.

167 *Air sampling*

168 Simultaneous to wastewater sampling, air samples were collected over the course of 1 h
169 using a SASS 3100 Dry Air Sampler (Research International, Monroe, WA, USA) with charged
170 electret filters. At the WWTPs, each sample was collected from a different location across a 50 m
171 transect, with data from all 3 grouped together to reduce within-site variation. Reference air
172 samples were collected from upwind control sites (Y Felinheli and Lac-Beauport) and at
173 downwind sites in Wales, again each over the course of 1 h. Field blanks were collected at upwind
174 sites and WWTPs by exposing a clean filter to the environment but not running the sampling
175 apparatus. Filters were stored in sealed bags at room temperature until DNA extraction.

176 To facilitate DNA extraction, a protocol developed by the *Institut de recherche Robert-*
177 *Sauvé en santé et en sécurité du travail* was followed. Briefly, filters were immersed in 10 ml of
178 SASS extraction buffer (0.1 M sodium phosphate, pH 7.4) and vortexed for 10 min at 2,500 rev
179 min^{-1} . This solution was then centrifuged at 17,500 rev min^{-1} for 10 min to create a pellet for DNA
180 extraction.

181 *Conifer needle sampling*

182 Conifer needles were collected following the methods of George et al. (2022) and Galès et
183 al. (2014). Briefly, 2 branch ends from each tree ~20 cm in length were collected from 1.5–2.5 m
184 height and stored in sterile plastic bags and kept at 4 °C until processing. A total of 15 g of conifer

185 needles were removed and homogenised in a 50 ml tube of 0.05 % Tween20 saline buffer with a
 186 paddle mixer for 5 min. Debris were removed from this solution by differential centrifugation at
 187 250 rev min⁻¹ for 3 min. The remaining supernatant was centrifuged at 17,500 rev min⁻¹ for 30 min
 188 to create pellets from which DNA was extracted. In Wales, 38 needle samples were collected, of
 189 which 10 came from proximity of the Treborth WWTP, 11 came from Bangor city centre, 10 came
 190 from distant sites, and 7 came from the Lake Vyrnwy control site. In Quebec, 12 needles samples
 191 were collected from the perimeter of the WWTP, and 5 samples were collected at the Lac-Beauport
 192 control site. Due to geographic and access limitations, we could not replicate the dispersal transect
 193 as in Wales.

194 *DNA extraction & qPCR analyses*

195 DNA extraction was performed for all sample types using a DNeasy PowerSoil Pro Kits
 196 (Qiagen, Montreal, Canada and Hilden, Germany) following manufacturer's instructions. Total
 197 bacterial biomass was estimated using qPCR analyses of the 16S rRNA gene and FAM
 198 fluorescence probes following Bach et al. (2002), on a Bio-Rad CFX-384 Touch™ Real-Time
 199 PCR Detection System (Bio-Rad, Montreal, CA). The thermoprotocol was 95 °C for 3 min; then
 200 95 °C for 20 s; and 62 °C for 1 min for 40 cycles. Results were deemed valid if accompanying
 201 standard curves had efficiency values of 90-110%. This qPCR data was used to generate 16S rRNA
 202 copy numbers per g of conifer needles or per ml of air or water using the following formulae:

$$203 \quad 16S \text{ rRNA copy number per g needles} = \frac{\left(\frac{(Sq * 50)}{7.5}\right) * 50}{W}$$

$$204 \quad 16S \text{ rRNA copy number per ml(air or water)} = \frac{(Sq * 50)}{V}$$

205 where S_q is the starting quantity estimate from qPCR analyses from 2 μ l of DNA of a 100 μ l
206 extraction, 7.5 (ml) is the volume of supernatant used to make aliquots, 50 (ml) is the total volume
207 of starting solution, W is the weight of needles (g), and V is the volume of air or water (ml).

208 The presence of crAssphage, a viral marker of human faecal contamination, was also
209 assessed using qPCR to assess the spread of faecal matter in the environments surrounding
210 WWTPs (Farkas et al., 2019). This was accomplished using the CPQ_056 primer pairs (Table S1)
211 and a thermocycler protocol of 98 °C for 5 min, then 95 °C for 15 s, and finally 60 °C for 1 min
212 for 40 cycles (Stachler et al., 2017).

213 A Takara SmartChip high-throughput qPCR (HT-qPCR) system (TakaraBio USA, San
214 Jose, CA, USA) was used to assess the abundance and diversity of ARGs and MGEs. A total of
215 39 genes were assessed including aminoglycosides (3), beta-lactams (10), colistin (1), macrolides
216 (4), MGEs (3), quinolones (2), sulfonamides (2), tetracyclines (10), and vancomycins (4) as
217 explained in George et al. (2022). These genes were selected based on their frequent use in the
218 literature (Stedfelt et al., 2018) and at the suggestion of colleagues, as in the case of *mcr-1* and
219 certain carbapenemase genes, as part of an effort to monitor AMR across sectors in Canada
220 (George et al., 2022). Information on mode of action was retrieved from the Comprehensive
221 Antibiotic Resistance Database (<https://card.mcmaster.ca>). A full list of genes and primer
222 sequences can be found in Table S1. These primers used SYBR Green dye fluorescence except for
223 those of the *mcr1* and *blaCTX-M-1* genes, which used a FAM probe for fluorescence. Analyses on
224 the Takara SmartChip HT-qPCR system used a shared a thermocycler protocol of an initial step
225 of 95 °C for 3 min; then 95 °C for 10 s; 60 °C for 30 s; and 55 °C for 31 s; with a melt curve of 55
226 °C for 5 s + 0.5 °C/cycle for a total of 45 cycles. Copy numbers of ARGs/MGEs were calculated
227 following the comparative CT method (Schmittgen and Livak, 2008):

$$228 \quad \textit{Relative gene copy number} = 2^{-(Ct(ARG) - Ct(16S))}$$

229 Relative abundance of ARGs/MGEs was normalised by multiplying the ARG/MGE copy
230 number by 16S rRNA gene copy number.

231

232 *Statistical analyses*

233 Relative copy numbers were log₁₀-transformed for all subsequent statistical analyses. The
234 presence/absence and proportional abundance of target genes is presented in Table S2. Total 16S
235 biomass, ARG/MGE richness linear models with either sample type or location as categorical
236 factors. Differences in the composition of ARGs/MGEs across samples were assessed by non-
237 metric dimensional scaling (NMDS) with Bray-Curtis distances using the vegan package (Oksanen
238 et al., 2022). Differences between groups were assessed by PERMANOVA (adonis2). All
239 statistical analyses were performed using R v. 4.2.2 (R Core Team, 2022).

240

241 **Results**

242 Quantifiable levels of bacterial DNA were recovered from all samples, except for the Y
243 Felinheli upwind control sample. Total bacterial biomass based on 16S rRNA gene copies
244 followed identical trends in both areas (Fig. 2). Biomass was lower in air samples than in
245 wastewater and needle samples in both Wales ($F_{2,47} = 108.4; p < 0.001$) and Quebec ($F_{2,20} = 91.19;$
246 $p < 0.001$).

247 Richness of ARGs was significantly different between sample types in both sample
248 locations (Wales: $F_{2,46} = 35.36; p < 0.001$; Quebec: $F_{2,20} = 26.62; p < 0.001$). Richness was highest
249 in wastewater samples and lowest in air samples (Fig. 3A). Although no significant differences
250 were observed along distance gradients radiating from WWTPs, ARG richness was higher in

251 wastewater samples than in phyllosphere samples in both Wales and Quebec ($F_{7,54} = 18.37$; $p <$
252 0.001). Richness of ARGs was however higher in wastewater than needles (Fig. 3B). CrAssphage
253 was only quantifiable in a small number of samples besides those of wastewater, namely the
254 Treborth WWTP air and one needles sample from Bangor itself (Fig. S1).

255 Quinolone genes were overwhelmingly abundant in Quebec samples (Fig. S2) and they
256 overshadow the subtleties. When they were removed from the data file, more detailed trends
257 became evident. Wastewater profiles in both sites were dominated by MGEs. Aminoglycosides
258 and sulfonamides were more prevalent in Quebec whereas tetracycline and macrolide resistance
259 genes were more present in Welsh wastewater (Fig. 4A). This translated into a greater proportion
260 of transposases/integrases in Welsh wastewater and a greater proportion of antibiotic inactivation
261 genes in Quebec wastewater (Fig. 4B). Quebec phyllosphere samples appear to have a more even
262 distribution of ARGs when quinolones were removed compared to Welsh needle samples, which
263 had high proportions of aminoglycosides (Fig. 4A). These genes were primarily associated with
264 antibiotic inactivation (Fig. 4B). Air samples from Wales had a high proportion of tetracycline
265 resistance genes. In Quebec, aminoglycoside and vancomycin resistance genes were important
266 contributors (Fig. 4A).

267 There were significant differences ($F_{7,54} = 4.56$; $p < 0.001$) in ARG copy numbers along
268 dispersal gradients from WWTPs based on PERMANOVA results. When $\log_{10} + 1$ transformed
269 abundances were visualised via NMDS (stress = 0.15) wastewater samples were clustered together.
270 There was also evidence that Quebec phyllosphere samples clustered. However, phyllosphere
271 samples collected along the dispersal gradient in Wales were so widely distributed that they
272 overshadowed other trends (Fig. 5A). Copy numbers of ARGs decreased over distance from the
273 WWTPs. Yet there was a high abundance of ARGs from the Lake Vyrnwy control site (Fig. 5B).

274 This was likely influenced by the proportion of number of beta-lactamase genes in these samples
275 (Fig. 5C).

276 Proportional abundances of ARGs in needle samples were vastly different from
277 wastewater. Welsh wastewater samples were dominated by MGEs. Oddly, their abundance was
278 lowest in proximity to the WWTP and grew with distance. They had an inverse relationship with
279 aminoglycoside resistance genes. Beta-lactam resistance genes were especially prevalent in the
280 Lake Vyrnwy control site (Fig. 5C). In Quebec, there was a decline in aminoglycoside resistance
281 genes from the wastewater to the control site. Needles collected from around the WWTP had a
282 much different ARG profile to the wastewater. Macrolides were strongly present. The control area
283 at Lac-Beauport had a more even distribution of ARGs (Fig. 5C).

284 **Discussion**

285 Monitoring ARG emissions via bioaerosols is an increasingly important aspect of addressing the
286 AMR crisis (George et al., 2022b). Yet tracking emissions across landscapes remains challenging.
287 Assessing changes in ARG profiles of phyllosphere communities represents a potential low-cost
288 tool to monitor ARG pollution in bioaerosol emissions. Previous work has shown that AMR
289 profiles within the phyllosphere can be differentiated with distance from anthropogenic ARG
290 sources (George et al., 2022a; Huang et al., 2023; Xiang et al., 2020). In past studies, forest
291 phyllospheres commonly exhibited fewer ARGs when compared to urban or agricultural areas
292 (George et al., 2022a; Xiang et al., 2020). However, this phenomenon was not exhibited in the two
293 forested sites used in the present study. Rather, ARG richness was not significantly different across
294 phyllosphere samples.

295 High-throughput qPCR revealed distinct trends in the proportional abundances of ARG
296 across the dispersal gradient. The similarity between wastewater ARG profiles between British

297 and Canadian sites reflects the clustering of Western nations observed in recent global analyses
298 (Prieto Riquelme et al., 2022). Yet these trends do not extend to the ARG profiles of phyllosphere
299 samples. In Wales, the increasing prevalence of aminoglycoside resistance genes across the
300 dispersal gradient may reflect the impacts of other anthropogenic activities. For example, high-
301 throughput qPCR data has shown aminoglycoside resistance genes are important components of
302 animal husbandry facilities (Yang et al., 2018). Indeed, aminoglycosides are the third most
303 commonly used antibiotics in British sheep farms (Davies et al., 2017). This could indicate
304 bioaerosols from agricultural sources are more important contaminants in this area. Given the
305 prevalence of sheep raising in the surrounding areas, this may indicate that animal waste and
306 management are more important sources of bioaerosols than the WWTP.

307 This work was predicated on the hypothesis that WWTP would be the most prominent
308 ARG emission source. However, the dispersal patterns of bioaerosols from point sources like
309 WWTPs are difficult to determine. Although there are some estimates for ARG dispersal over long
310 distances (Bai et al., 2022; Gaviria-Figueroa et al., 2019), so far no study has conclusively tracked
311 such emissions. Such local dispersal dynamics are likely important in Wales, where precipitation
312 was higher during the sampling period. Realised dispersal ranges are likely to be much smaller
313 than mathematical predictions due to variation in particle composition or changes in air pressure
314 or windspeed, for instance (Brunet et al., 2018). Recent work has demonstrated that ARGs are
315 present in clouds, however, how their deposition in precipitation may influence terrestrial
316 communities is unknown (Rossi et al., 2023). Indeed, soils are important reservoirs of both
317 anthropogenic and naturally occurring ARGs (Cytryn, 2013; Singer et al., 2016). Comparisons
318 with the ARG profiles of soil were not made in the present study. Shared ARG profiles between
319 the phyllosphere and soil may indicate deposition of local bioaerosols, such as those generated by

320 rainfall (Joung et al., 2017). It is likely that agricultural bioaerosols and run-off are important
321 vectors for ARG transport in soils along the dispersal gradient (Singer et al., 2016).

322 To our knowledge the present work represents the first time phyllosphere ARG profiles
323 have been directly compared to source material. However, more intensive studies are needed to
324 validate the potential for monitoring airborne ARGs with conifer needles sampling. This must
325 include longer sampling campaigns with multiple sampling events, which are beyond the scope of
326 this study. Further, greater efforts must be made to account for confounding aerosol sources.
327 Although complementary air samples were taken, the equipment needed is expensive and
328 logistically difficult to run in long-term field experiments. Passive air samplers could be installed
329 in forested areas so that the aerial component of phyllosphere samples over a set time period can
330 be observed in isolation. This could be done using the Rutgers Electrostatic Passive Sampler
331 (REPS) system, for example. The REPS is compact, affordable, and validated for collecting viable
332 bacteria outdoors (Grogan and Mainelis, 2022). Further, soil samples should be periodically
333 collected from below selected conifer trees so that confounding effects of local bioaerosol
334 emissions can be eliminated. Incorporating these additional measures will better validate future
335 tests of this monitoring method.

336 **Conclusion**

337 Mitigating ARG emissions via bioaerosols will be a vital aspect of future efforts to combat the
338 AMR crisis. To achieve this, effective monitoring programmes must be developed. Other
339 monitoring programmes suggest that the conifer needle phyllosphere may be an effective
340 monitoring tool. However, our results indicate that further refinement is needed for this to be
341 effective. Although distinct trends in ARG profiles were observed across a dispersal gradient from
342 WWTPs, these findings were incongruous with those of wastewater. Future work should include

343 repeat sampling at multiple time points to account for seasonal variation in emissions and
344 phyllosphere community composition. Further, molecular approaches such as amplicon or shotgun
345 metagenomic sequencing should be employed to better identify bioindicators of contamination.
346 These lessons will inform future attempts to create ARG monitoring networks.

347 **Acknowledgements**

348 We are grateful to the city of Sainte-Brigitte-de-Laval and Dŵr Cymru Welsh Water for providing
349 access to WWTPs. Funding for this project was provided by a joint Quebec-Wales cooperation
350 programme through Welsh Government and Quebec's *Ministère des Relations internationales et*
351 *de la Francophonie*. Additional funds were provided by an NSERC Frontiers Grant (Grant #:
352 539024-2019) held by Pr Caroline Duchaine. Pr Duchaine holds the Tier 1 Canada Research Chair
353 on Bioaerosols. We thank Dr Robert W. Brown and Dr Nathalie Turgeon for their help collecting
354 conifer needle samples in Wales and Quebec, respectively. We thank Dr Peter A. Vanrollenghem
355 and the modelEAU group at Université Laval for chemical analyses of Quebec wastewater.

356 **Data availability**

357 All data are presented in the manuscript. Files can be provided upon reasonable request via the
358 corresponding author.

359 **Conflict of interest**

360 The authors have no financial or personal conflicts of interest in the work presented.

361 **Author contributions**

362 PBLG, DLJ, and CD conceived the project which was further refined by ECC and LH. ECC, LH,
363 PBLG, SL, and JL collected and processed samples in both Quebec and Wales. PBLG led
364 statistical analyses with help from LH. LH and PBLG led the writing of the first draft with
365 contributions from ECC, SL, and JL. All authors edited and revised subsequent drafts.

366

367 **References**

- 368 Amos, G.C.A., Hawkey, P.M., Gaze, W.H., and Wellington, E.M. 2014. Waste water effluent
369 contributes to the dissemination of CTX-M-15 in the natural environment. *J. Antimicrob.*
370 *Chemother.* **69(7)**: 1785–1791. doi:10.1093/jac/dku079
- 371 Anderson, M., Clift, C., Schulze, K., Sagan, A., Nahrgang, S., Ait Ouakrim, D., and Mossialos,
372 E. 2019. Averting the AMR crisis: What are the avenues for policy action for countries in
373 Europe? European Observatory on Health Systems and Policies, Copenhagen.
- 374 Bach, H.-J., Tomanova, J., Schloter, M., Munch, J.C. 2002. Enumeration of total bacteria and
375 bacteria with genes for proteolytic activity in pure cultures and in environmental samples
376 by quantitative PCR mediated amplification. *J. Microbiol. Methods.* **49(3)**: 235–245.
377 doi:10.1016/S0167-7012(01)00370-0
- 378 Bai, H., He, L.-Y., Wu, D.-L., Gao, F.-Z., Zhang, M., Zou, H.-Y., Yao, M.-S., et al. 2022. Spread
379 of airborne antibiotic resistance from animal farms to the environment: Dispersal pattern
380 and exposure risk. *Environ. Int.* **158**, 106927. doi:10.1016/j.envint.2021.106927
- 381 Bélanger Cayouette, A., Ouradou, A., Veillette, M., Turgeon, N., George, P.B.L., Corbin, S.,
382 Boulanger, C., et al. 2022. Effect of odor-treatment biofilter on bioaerosols from an
383 indoor wastewater treatment plant. *Atmosphere.* **13(10)**: 1723.
384 doi:10.3390/atmos13101723
- 385 Blais-Lecours, P., Perrott, P., Duchaine, C. 2015. Non-culturable bioaerosols in indoor settings:
386 Impact on health and molecular approaches for detection. *Atmos. Environ.* **110**: 45–53.
387 doi:10.1016/j.atmosenv.2015.03.039
- 388 Brunet, Y., Wéry, N., and Galès, A. 2018. "Short-Scale Transport of Bioaerosols" in Delort, A.-
389 M., and Amato, P (ed.), *Microbiology of Aerosols* John Wiley & Sons Inc., Hoboken,
390 pp.137-153.
- 391 Corno, G., Yang, Y., Eckert, E.M., Fontaneto, D., Fiorentino, A., Galafassi, S., Zhang, T., et al.
392 2019. Effluents of wastewater treatment plants promote the rapid stabilization of the
393 antibiotic resistome in receiving freshwater bodies. *Water Res.* **158**: 72–81.
394 doi:10.1016/j.watres.2019.04.031
- 395 Council of Canadian Academics. 2019. When Antibiotics Fail: The Expert Panel on the Potential
396 Socio-Economic Impacts of Antimicrobial Resistance in Canada. Council of Canadian
397 Academic, Ottawa, Canada
- 398 Cytryn, E. 2013. The soil resistome: The anthropogenic, the native, and the unknown. *Soil Biol.*
399 *Biochem.* **63**: 18–23. doi:10.1016/j.soilbio.2013.03.017
- 400 Czekalski, N., Gascón Díez, E., and Bürgmann, H. 2014. Wastewater as a point source of
401 antibiotic-resistance genes in the sediment of a freshwater lake. *ISME J.* **8(7)**: 1381–
402 1390. doi:10.1038/ismej.2014.8
- 403 Davies, P., Remnant, J.G., Green, M.J., Gascoigne, E., Gibbon, N., Hyde, R., Porteous, J.R., et
404 al. 2017. Quantitative analysis of antibiotic usage in British sheep flocks. *Vet Rec.*
405 **181(19)**: 511. doi: 10.1136/vr.104501
- 406 Engemann, C.A., Keen, P.L., Knapp, C.W., Hall, K.J., and Graham, D.W. 2008. Fate of
407 tetracycline resistance genes in aquatic systems: Migration from the water column to
408 peripheral biofilms. *Environ. Sci. Technol.* **42**: 5131–5136. doi:10.1021/es800238e

- 409 Farkas, K., Adriaenssens, E.M., Walker, D.I., McDonald, J.E., Malham, S.K., and Jones, D.L.
410 2019. Critical evaluation of crAssphage as a molecular marker for human-derived
411 wastewater contamination in the aquatic environment. *Food Environ. Virol.* **11**: 113–119.
412 doi:10.1007/s12560-019-09369-1
- 413 Galès, A., Latrille, E., Wéry, N., Steyer, J.-P., and Godon, J.-J. 2014. Needles of *Pinus*
414 *halepensis* as biomonitors of bioaerosol emissions. *PLoS ONE* **9**: e112182.
415 doi:10.1371/journal.pone.0112182
- 416 Gavia-Figueroa, A., Preisner, E.C., Hoque, S., Feigley, C.E., and Norman, R.S. 2019. Emission
417 and dispersal of antibiotic resistance genes through bioaerosols generated during the
418 treatment of municipal sewage. *Sci. Total Environ.* **686**: 402–412.
419 doi:10.1016/j.scitotenv.2019.05.454
- 420 George, P.B.L., Leclerc, S., Turgeon, N., Veillette, M., and Duchaine, C. 2022a. Conifer needle
421 phyllosphere as a potential passive monitor of bioaerosolised antibiotic resistance genes.
422 *Antibiotics* **11**: 907. doi:10.3390/antibiotics11070907
- 423 George, P.B.L., Rossi, F., St-Germain, M.-W., Amato, P., Badard, T., Bergeron, M.G.,
424 Boissinot, M., et al. 2022b. Antimicrobial resistance in the environment: Towards
425 elucidating the roles of bioaerosols in transmission and detection of antibacterial
426 resistance genes. *Antibiotics* **11**: 974. doi:10.3390/antibiotics11070974
- 427 Grogan, S.N.C.M., and Mainelis, G. 2022. Effect of sampling duration on culturable and viable
428 bioaerosol determination when using Rutgers Electrostatic Passive Sampler (REPS). *J.*
429 *Aerosol Sci.* **166**: 106066. doi:10.1016/j.jaerosci.2022.106066
- 430 Huang, X.-R., Neilson, R., Yang, L.-Y., Deng, J.-J., Zhou, S.-Y.-D., Li, H., Zhu, Y.-G., et al.
431 2023. Urban greenspace types influence the microbial community assembly and
432 antibiotic resistome more in the phyllosphere than in the soil. *Chemosphere* **338**: 139533.
433 doi:10.1016/j.chemosphere.2023.139533
- 434 Jacquiod, S., Brejnrod, A., Morberg, S.M., Abu Al-Soud, W., Sørensen, S.J., and Riber, L. 2017.
435 Deciphering conjugative plasmid permissiveness in wastewater microbiomes. *Mol. Ecol.*
436 **26**: 3556–3571. doi:10.1111/mec.14138
- 437 Joung, Y.S., Ge, Z., and Buie, C.R. 2017. Bioaerosol generation by raindrops on soil. *Nat.*
438 *Commun.* **8**: 14668. doi:10.1038/ncomms14668
- 439 Katak, S., Patowary, R., Chatterjee, S., Vairale, M.G., Sharma, S., Dwivedi, S.K., and Kamboj,
440 D.V. 2022. Bioaerosolization and pathogen transmission in wastewater treatment plants:
441 Microbial composition, emission rate, factors affecting and control measures.
442 *Chemosphere* **287**: 132180. doi:10.1016/j.chemosphere.2021.132180
- 443 Lin, X.-J., Barrington, S., Nicell, J., Choinière, D., and Vézina, A. 2006. Influence of windbreaks
444 on livestock odour dispersion plume in the field. *Agric. Ecosyst. Environ.* **116**: 263–272.
445 doi:10.1016/j.agee.2006.02.014
- 446 Manaia, C.M., Novo, A., Coelho, B., and Nunes, O.C. 2010. Ciprofloxacin resistance in
447 domestic wastewater treatment plants. *Water Air Soil Pollut.* **208**: 335–343.
448 doi:10.1007/s11270-009-0171-0
- 449 Manibusan, S., and Mainelis, G., 2022. Passive bioaerosol samplers: A complementary tool for
450 bioaerosol research. A review. *J. Aerosol Sci.* **163**: 105992.
451 doi:10.1016/j.jaerosci.2022.105992
- 452 Marti, E., Jofre, J., and Balcazar, J.L. 2013. Prevalence of antibiotic resistance genes and
453 bacterial community composition in a river influenced by a wastewater treatment plant.
454 *PLoS ONE* **8**: e78906. doi:10.1371/journal.pone.0078906

- 455 Mbareche, H., Dion-Dupont, V., Veillette, M., Brisebois, E., Lavoie, J., and Duchaine, C. 2022.
 456 Influence of seasons and sites on bioaerosols in indoor wastewater treatment plants and
 457 proposal for air quality indicators. *J. Air Waste Manag. Assoc.* **72**: 1000–1011.
 458 doi:10.1080/10962247.2022.2066735
- 459 Munir, M., Wong, K., and Xagorarakis, I. 2011. Release of antibiotic resistant bacteria and genes
 460 in the effluent and biosolids of five wastewater utilities in Michigan. *Water Res.* **45**: 681–
 461 693. doi:10.1016/j.watres.2010.08.033
- 462 Murray, C.J., Ikuta, K.S., Sharara, F., Swetschinski, L., Robles Aguilar, G., Gray, A., Han, C., et
 463 al. 2022. Global burden of bacterial antimicrobial resistance in 2019: a systematic
 464 analysis. *Lancet* **399**: 629–655. doi:10.1016/S0140-6736(21)02724-0
- 465 de Nies, L., Bhanu Busi, S., Kunath, B.J., May, P., Wilmes, P. 2022. Mobilome-driven
 466 segregation of the resistome in biological wastewater treatment. *eLife* **11**: e81196
 467 doi:10.7554/eLife.81196
- 468 Novo, A., Manaia, and C.M. 2010. Factors influencing antibiotic resistance burden in municipal
 469 wastewater treatment plants. *Appl. Microbiol. Biotechnol.* **87**: 1157–1166.
 470 doi:10.1007/s00253-010-2583-6
- 471 O'Neill, J. 2016. Tackling drug-resistant infections globally: final report and recommendations.
 472 The review on antimicrobial resistance.
- 473 Ouradou, A., Veillette, M., Bélanger Cayouette, A., Corbin, S., Boulanger, C., Dorner, S.,
 474 Duchaine, C., et al. 2023. Effect of odor treatment systems on bioaerosol microbial
 475 concentration and diversity from wastewater treatment plants. *Sci. Total Environ.* **874**:
 476 162419. doi:10.1016/j.scitotenv.2023.162419
- 477 Pokharel, S., Shrestha, P., and Adhikari, B. 2020. Antimicrobial use in food animals and human
 478 health: time to implement 'One Health' approach. *Antimicrob. Resist Infect. Control* **9**:
 479 181. doi:10.1186/s13756-020-00847-x
- 480 Prieto Riquelme, M.V., Garner, E., Gupta, S., Metch, J., Zhu, N., Blair, M.F., Arango-Argoty,
 481 G., et al. 2022. Demonstrating a comprehensive wastewater-based surveillance approach
 482 that differentiates globally sourced resistomes. *Environ. Sci. Technol.* **56**: 14982–14993.
 483 doi:10.1021/acs.est.1c08673
- 484 Pruden, A., Vikesland P.J., Davis, B.C., de Roda Husman, A.M. 2021. Seizing the moment:
 485 now is the time for integrated global surveillance of antimicrobial resistance in
 486 wastewater environments. *Curr. Opin. Microbiol.* **64**: 91-99. doi:
 487 10.1016/j.mib.2021.09.013
- 488 Raza, S., Shin, H., Hur, H.-G., and Unno, T. 2022. Higher abundance of core antimicrobial
 489 resistant genes in effluent from wastewater treatment plants. *Water Res.* **208**: 117882.
 490 doi:10.1016/j.watres.2021.117882
- 491 Robinson, T.P., Bu, D.P., Carrique-Mas, J., Fèvre, E.M., Gilbert, M., Grace, D., Hay, S.I., et al.
 492 2016. Antibiotic resistance is the quintessential One Health issue. *Trans. R. Soc. Trop.*
 493 *Med. Hyg.* **110**: 377–380. doi:10.1093/trstmh/trw048
- 494 Rossi, F., Péguilhan, R., Turgeon, N., Veillette, M., Baray, J.-L., Deguillaume, L., Amato, P., et
 495 al. 2023. Quantification of antibiotic resistance genes (ARGs) in clouds at a mountain site
 496 (puy de Dôme, central France). *Sci. Total Environ.* **865**: 161264.
 497 doi:10.1016/j.scitotenv.2022.161264
- 498 Schmittgen, T.D., and Livak, K.J. 2008. Analyzing real-time PCR data by the comparative CT
 499 method. *Nat. Protoc.* **3**: 1101–1108. doi:10.1038/nprot.2008.73

- 500 Shi, B., Zhao, R., Su, G., Liu, B., Liu, W., Xu, J., Li, Q., et al. 2023. Metagenomic surveillance
 501 of antibiotic resistome in influent and effluent of wastewater treatment plants located on
 502 the Qinghai-Tibetan Plateau. *Sci. Total Environ.* **870**: 162031.
 503 doi:10.1016/j.scitotenv.2023.162031
- 504 Singer, A.C., Shaw, H., Rhodes, V., and Hart, A. 2016. Review of antimicrobial resistance in the
 505 environment and its relevance to environmental regulators. *Front. Microbiol.* **7**: 1728 doi:
 506 10.3389/fmicb.2016.01728
- 507 Stachler, E., Kelty, C., Sivaganesan, M., Li, X., Bibby, K., Shanks, O.C. 2017. Quantitative
 508 CrAssphage assays for human fecal pollution measurement. *Environ. Sci. Technol.* **51**:
 509 9146-9154 doi: 10.1021/acs.est.7bo2703
- 510 Stedfelt, R.D., Guo, X., Stedtfeld, T.M., Sheng, H., Williams, M.R., Hauschild, K., et al. 2018.
 511 Primer set 2.0 for highly parallel qPCR array targeting antibiotic resistance genes and
 512 mobile genetic elements. *FEMS Microbiol. Ecol.* **94**(9): fiy130.
 513 doi:10.1093/femsec/fiy130.
- 514 Wang, Y., Wang, C., and Song, L. 2019. Distribution of antibiotic resistance genes and bacteria
 515 from six atmospheric environments: Exposure risk to human. *Sci. Total Environ.* **694**:
 516 133750. doi:10.1016/j.scitotenv.2019.133750
- 517 Wu, J., Wang, J., Li, Z., Guo, S., Li, K., Xu, P., Ok, Y.S., et al. 2023. Antibiotics and antibiotic
 518 resistance genes in agricultural soils: A systematic analysis. *Crit. Rev. Environ. Sci.*
 519 *Technol.* **53**: 847–864. doi:10.1080/10643389.2022.2094693
- 520 Xiang, Q., Zhu, D., Giles, M., Neilson, R., Yang, X.-R., Qiao, M., and Chen, Q.-L. 2020.
 521 Agricultural activities affect the pattern of the resistome within the phyllosphere
 522 microbiome in peri-urban environments. *J. Hazard. Mat.* **382**: 121068.
 523 doi:10.1016/j.jhazmat.2019.121068
- 524 Yang, Y., Zhou, R., Chen, B., Zhang, T., Hu, L., and Zou, S. 2018. Characterization of airborne
 525 antibiotic resistance genes from typical bioaerosol emission sources in the urban
 526 environment using metagenomic approach. *Chemosphere* **213**: 463–471. doi:
 527 10.1016/j.chemosphere.2018.09.066

528 Figure captions

529
 530 **Fig. 1.** Sampling locations visited in Quebec and Wales. Points represent the centre of
 531 cities/villages sampled. Molecular analyses were conducted at Université Laval in Quebec City
 532 (marked in blue).
 533

534 **Fig. 2.** Box plots of log₁₀-transformed bacterial biomass across sample types (air, phyllosphere,
 535 wastewater) in Wales and Quebec. Boxes denote the first and third quartiles. Horizontal lines
 536 mark medians. Whiskers represent 1.5X the interquartile range. Black dots represent outliers.
 537 Individual data points are displayed next to boxes.
 538

539 **Fig. 3.** Box plot of ARG richness across **A**) sample types and **B**) from conifer phyllosphere samples
 540 along distance gradients from WWTPs. Boxes denote the first and third quartiles. Horizontal lines
 541 mark medians. Whiskers represent 1.5X the interquartile range. Black dots represent outliers.

542 Individual data points are displayed next to boxes. SBL stands for Saint-Brigitte-de-Laval. Control
543 sites are labelled in grey; all other sites are labelled in black.

544

545 **Fig 4.** Proportionate abundances of **A)** ARG type and **B)** method of action for sample types (air,
546 phyllosphere, and wastewater) at two WWTPs located in Wales and Quebec. Note that quinolones
547 have been removed from this figure demonstrate trends in other genes. SBL stands for Saint-
548 Brigitte-de-Laval.





549


550 **Fig. 5.** ARGs profiles of wastewater and conifer phyllosphere samples. Shown are **A)** NMDS
551 ordination of phyllosphere and wastewater samples; **B)** the mean and standard error of ARG copy
552 numbers. Control sites are labelled in grey; all other sites are labelled in black; and **C)** proportional
553 abundances of ARGs. SBL stands for Saint-Brigitte-de-Laval.

554

Table 1. Chemical properties of the three wastewater samples collected at both the Treborth and Sainte-Brigitte-de-Laval WWTPs at the time of aerosol sampling. Values represent means (\pm SE).


Property	Treborth	Sainte-Brigitte-de-Laval
pH	7.60 \pm 0.02	6.98 \pm 0.03
Electrical conductivity (μ S/cm)	1018 \pm 17	652 \pm 2
Turbidity (NTU)	210.3 \pm 12.8	145.2 \pm 10.7
NH ₄ ⁺ (mg N/l)	30.3 \pm 0.6	20.0 \pm 0.3
PO ₄ ³⁻ (mg P/l)	8.1 \pm 0.62	19.4 \pm 0.71






A Location

■ Bangor
■ SBL



Sample Type

B



Phyllosphere Location

ARG type

- Aminoglycoside
- Beta-lactem
- Collistin
- Macrolide
- MGE
- Sulfonamide
- Tetracycline
- Vancomycin

ARG action

- Antibiotic inactivation
- Antibiotic target protection
- Efflux pump
- Target modification
- Transposase/integrase

