Acute isometric and dynamic exercise do not alter cerebral sympathetic nerve activity in healthy humans

Tymko, Michael M; Drapeau, Audrey; Vieira-Coelho, Maria Augusta; Labrecque, Lawrence; Imhoff, Sarah; Coombs, Geoff B; Langevin, Stephan; Fortin, Marc; Châteauvert, Nathalie; Ainslie, Philip N; Brassard, Patrice

Journal of Cerebral Blood Flow and Metabolism

DOI:

10.1177/0271678X241248228

E-pub ahead of print: 13/04/2024

Publisher's PDF, also known as Version of record

Cyswllt i'r cyhoeddiad / Link to publication

Dyfyniad o'r fersiwn a gyhoeddwyd / Citation for published version (APA): Tymko, M. M., Drapeau, A., Vieira-Coelho, M. A., Labrecque, L., Imhoff, S., Coombs, G. B., Langevin, S., Fortin, M., Châteauvert, N., Ainslie, P. N., & Brassard, P. (2024). Acute isometric and dynamic exercise do not alter cerebral sympathetic nerve activity in healthy humans. Journal of Cerebral Blood Flow and Metabolism. Advance online publication. https://doi.org/10.1177/0271678X241248228

Hawliau Cyffredinol / General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

• Users may download and print one copy of any publication from the public portal for the purpose of private study or research.

- You may not further distribute the material or use it for any profit-making activity or commercial gain
 You may freque distribute the LIPL identifying the publication in the public portal?
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Acute isometric and dynamic exercise do not alter cerebral sympathetic nerve activity in healthy humans

JCBFM

Journal of Cerebral Blood Flow & Metabolism 0(0) 1–12 © The Author(s) 2024 © W Article reuse guidelines:

sagepub.com/journals-permissions DOI: 10.1177/0271678X241248228 journals.sagepub.com/home/jcbfm



Michael M Tymko^{1,2}, Audrey Drapeau^{3,4}, Maria Augusta Vieira-Coelho^{5,6}, Lawrence Labrecque^{3,4}, Sarah Imhoff^{3,4}, Geoff B Coombs⁷, Stephan Langevin⁴, Marc Fortin⁴, Nathalie Châteauvert⁴, Philip N Ainslie⁸ and Patrice Brassard^{3,4}

Abstract

The impact of physiological stressors on cerebral sympathetic nervous activity (SNA) remains controversial. We hypothesized that cerebral noradrenaline (NA) spillover, an index of cerebral SNA, would not change during both submaximal isometric handgrip (HG) exercise followed by a post-exercise circulatory occlusion (PECO), and supine dynamic cycling exercise. Twelve healthy participants (5 females) underwent simultaneous blood sampling from the right radial artery and right internal jugular vein. Right internal jugular vein blood flow was measured using Duplex ultrasound, and tritiated NA was infused through the participants' right superficial forearm vein. Heart rate was recorded via electrocardiogram and blood pressure was monitored using the right radial artery. Total NA spillover increased during HG (P = 0.049), PECO (P = 0.006), and moderate cycling exercise (P = 0.03) compared to rest. Cerebral NA spillover remained unchanged during isometric HG exercise (P = 0.36), PECO after the isometric HG exercise (P = 0.45), and during moderate cycling exercise involving both small and large muscle mass do not engage cerebral SNA in healthy humans. Our findings suggest that cerebral SNA may be non-obligatory for exercise-related cerebrovascular adjustments.

Keywords

Cerebral blood flow, sympathetic nervous activity, noradrenaline spillover

Received 27 August 2023; Revised 30 January 2024; Accepted 26 March 2024

Introduction

Several physiological factors, such as aerobic/resistance exercise and rapid eye movement sleep, as well as pathological clinical conditions including autonomic dysreflexia, baroreflex failure, or uncontrolled arterial hypertension, require efficient counter-regulation against transient arterial blood pressure fluctuations. Without the neuroprotective influences of sympathetic activation or cerebral autoregulation (which refers to the relationship between cerebral perfusion pressure

Corresponding author:

¹Integrative Cerebrovascular and Environmental Physiology SB Laboratory, Department of Human Health and Nutritional Sciences, College of Biological Science, University of Guelph, Guelph, Canada ²Department of Medicine, Faculty of Medicine, University of British Columbia, Vancouver, Canada

³Department of Kinesiology, Faculty of Medicine, Université Laval, Québec, Canada

⁴Institut universitaire de cardiologie et de pneumologie de Québec-Université Laval, Québec, Canada

⁵Department of Biomedicine, Pharmacology and Therapeutics Unit, Faculty of Medicine, University of Porto, Portugal

⁶Department of Psychiatry and Mental Health, University Hospital Center of São João, Porto, Portugal

⁷Department of Kinesiology, Faculty of Science, McMaster University, Hamilton, Canada

⁸Centre for Heart, Lung and Vascular Health, School of Health and Exercise Sciences, University of British Columbia – Okanagan, Kelowna, Canada

Patrice Brassard, Department of Kinesiology, Faculty of Medicine, PEPS – Université Laval, 2300 rue de la Terrasse, Québec, QC GIV OA6, Canada. Email: patrice.brassard@kin.ulaval.ca

and blood flow during changes in arterial blood pressure), acute surges in arterial blood pressure could increase the risk of hyper-perfusion injury, predisposing to stroke or blood-brain barrier breakdown.¹ As such, tight regulation of cerebral blood flow (CBF) during acute changes in arterial blood pressure helps to maintain normal cerebral function.^{2–4}

Cerebral sympathetic nerve activity (SNA) increases with acute increases in arterial blood pressure induced pharmacologically or mechanically in anesthetized lambs, but does not change with acute reductions in arterial blood pressure.⁵ This finding suggests that elevations in cerebral SNA could serve as the effective dynamic regulator of CBF during transient increases in arterial blood pressure, at least in this animal model. However, much less work has been conducted in clinical and healthy human models. In humans with essential arterial hypertension, cerebral noradrenaline (NA) release is elevated,⁶ which in support of the existing animal literature, could be interpreted as a cerebral protective mechanism for overperfusion, or alternatively, elevated cerebral NA release itself could contribute to chronic arterial hypertension. A study in healthy humans demonstrated that moderate intensity cycling exercise, which is known to acutely increase arterial blood pressure, CBF, and peripheral SNA,7 resulted in no change in cerebral NA spillover, an index of cerebral specific SNA.^{8,9} The latter finding in healthy humans indicates that cerebral SNA does not contribute to CBF regulation during moderate intensity aerobic exercise; however, these data have yet to be replicated, and the effects of isometric exercise [e.g., handgrip (HG)] and post-exercise circulatory occlusion (PECO), which increases sympathetic outflow independent from exercise, has not been explored. A better understanding of how acute physiological stresses that are known to result in increased peripheral SNA and arterial blood pressure, such as dynamic cycling exercise, HG exercise, and PECO, may provide the basis for developing new approaches to the management of pathological conditions that require attenuation of CBF in arterial hypertension and/or loss of cerebrovascular regulation in humans. Each of these modalities to increase peripheral SNA and arterial blood pressure are distinct from one another (e.g., dynamic vs isometric exercise, large vs small muscle activation, active vs inactive tissue), potentially giving insight into whether different types of SNA activation can influence cerebral SNA.

The purpose of the current study was to measure the effects of an acute increase in arterial blood pressure using isometric HG exercise and PECO and dynamic cycling exercise on cerebral SNA using the NA spill-over technique. Based on previous work in humans,⁷ we hypothesized that cerebral NA spillover would not

change during both isometric and dynamic exercise, highlighting that cerebral SNA is not a regulatory mechanism during moderate intensity exercise in healthy humans.

Methods

Ethical approval and informed consent

The present study was approved by the *Comité* d'éthique de la recherche de l'Institut universitaire de cardiologie et de pneumologie de Québec (CER: 21557). All participants provided their written informed consent prior to participating. The study conformed to the standards set by the Tri-Council Policy Statement: Ethical Conduct for Research Involving Humans (TCPS 2) and the Declaration of Helsinki, except for registration in a database. The data that support the findings of this study are available from the corresponding author upon reasonable request.

Participants

Twelve young and healthy participants (five females) were recruited to participate in this study. A sample size calculation revealed that a minimum of 5 participants would be necessary to yield a statistically significant increase in arterial blood pressure between baseline and steady-state isometric HG exercise with a power of 0.80, a p value <0.05 and a Cohen's d effect size of 1.86.¹⁰ Twelve participants were recruited to consider potential participant drop out. Pilot testing revealed that a similar increase ($\sim 20\%$) in mean arterial blood pressure compared to isometric HG exercise was achievable during our dynamic supine cycling exercise protocol. All participants were recruited at the Université Laval campus (Québec City, Canada). Participants were free of known cardiovascular, respiratory and neurological disease, and non-smokers. Participants were not on any current medication except for oral contraceptives (n = 2). For unforeseen methodological issues (e.g., catheter obstruction [n = 1], and equipment failure [n = 2]), some participants did not complete the cycling portion of the study.

Experimental design

On an initial visit, participants came to the laboratory to have all procedures explained to them in person, sign informed consent to the study, complete three lefthanded maximal HG exercise (for the evaluation of their maximal voluntary contraction), and have their internal jugular veins (IJV) scanned via Duplex ultrasound to ensure adequate image quality for this vessel. On the second experimental visit, participants arrived at the laboratory having abstained from alcohol, caffeine and exercise for ≥ 12 hrs, and were fasted for a minimum of 2 hrs, but allowed to drink water ad libitum. All testing procedures were completed in ~5 hrs. Seven participants were tested in the morning (7 am), and 5 participants were tested in the afternoon (12 pm). Data collection took place over a period of two weeks.

Under sterilized conditions, participants had their right radial artery, right superficial forearm vein, and right IJV catheterized (see additional details below). After catheterization, the participants were transferred to the laboratory, placed supine, and rested for 1 hr to offset the nociceptive stimuli associated with catheterization. Participants were then instrumented for data collection (see details below). A continuous tracer infusion of ³H-labelled L-NA with high specific activity $(10 \text{ Ci mmol}^{-1})$, in order to avoid an elevation in arterial blood pressure and reflex lowering NA release.¹¹ was administered intravenously at $0.8 \,\mu\text{Ci min}^{-1}$ via a superficial vein of the right forearm to steady-state (45 min). The exercise data presented in this study were part of a larger series of separate experiments including lower body negative pressure without and with hypercapnia. After a 5-min resting period, the participants first completed a two-min isometric HG test to 30% of their maximal voluntary contraction of their left hand (previously measured). Participants completed the HG test in a supine position with a hand dynamometer connected to an acquisition system (Biopac System, Santa Barbara, California, USA) while seeing a screen on which the individual target workload (i.e. 30% of their maximal voluntary contraction) was marked by a horizontal red line. To ensure this target workload was maintained over the 2 min, an experimenter was placed closed to the screen and provided constant encouragement for participants to stay as close as possible to the target workload. Participants were instructed to avoid doing a Valsalva maneuver during the HG protocol. Immediately after the isometric HG test, a PECO test was completed by inflating a pressure cuff around the forearm to \sim 250 mmHg for three min. The PECO was conducted in order to assess the influence of increased global SNA via local metabolic afferent activity (i.e., muscle metaboreflex).¹² After the HG and PECO tests, the participants were then instructed to rest quietly for 10 min to recover. The participants feet were then placed into the pedals of a supine cycle ergometer attached to the hospital bed (LODE Lode BV, The Netherlands). After a 5-min resting period, the participants cycled at three exercise workloads that represented mild-to-moderate exercise for young and healthy males and females (75, 100, and 125 Watts for males; 50, 75, and 100 Watts for females), with each intensity lasting three min (Figure 1). Higher intensities of exercise were avoided due to concerns of obtaining reliable ultrasound measures. These exercise workloads are referred to as EX1, EX2, and EX3.



Figure 1. Schematic of the experimental protocol. Participants were instrumented with IJV, arterial (i.e., ART), and superficial vein of the forearm (i.e., ANTE) catheters upon arrival to the laboratory. The experimental protocol began with a 45-min infusion of tritiated NA (to steady-state) that continued throughout the entire protocol. The participants were then asked to perform a 2-min isometric handgrip test at 30% of their maximal voluntary contraction, which was followed by a 3-min post-exercise circulatory occlusion test. After a 10-min recovery, participants then performed a submaximal supine cycling exercise test that consisted of three stages for both male (75 W, 100 W, 125 W) and female (50 W, 75 W, 100 W) participants.

Experimental measurements

Cardiovascular measurements. All continuously recorded cardiovascular measurements were acquired at 1000 Hz using an analog-to-digital converter (Powerlab/16SP ML 880; ADInstruments, Colorado Springs, CO, USA) interfaced with a personal computer. Commercially available software was used to analyse cardiovascular variables (LabChart V7.1; ADInstruments). Electrocardiogram electrodes were placed in lead II configuration (Bioamp, ML132; ADInstruments) to measure heart rate. The radial artery catheter was connected to a commercially available arterial blood sampling kit integrated with a pressure transducer (VAMP Adult, Edwards Lifescience, Irvine, CA), allowing for beatby-beat measurements of arterial blood pressure. The radial arterial blood pressure waveform was corrected to brachial artery blood pressure using a manual sphygmomanometer. A one-min average was taken for these variables at the end of the rest period, during the isometric HG protocol (HG and PECO), and each of the cycling exercise stages (EX1, EX2, EX3).

Respiratory measurements and dynamic end-tidal forcing. Participants breathed through a mouthpiece while wearing a nose clip. Both partial pressure of end-tidal carbon dioxide ($P_{ET}CO_2$) and oxygen ($P_{ET}O_2$) were sampled at the mouth and recorded by a calibrated gas analyzer (model ML206, ADInstruments); breathing frequency, tidal volume, and minute ventilation were measured by a pneumotachograph (model HR 800 L, Hans Rudolph) connected in series to a bacteriological filter. PETCO2 and PETO2 were controlled by a portable dynamic end-tidal forcing system during the HG and PECO protocol. This system was not used to control end-tidal gases during cycling exercise, since the available system could not accommodate the high respiratory flow rates associated with cycling exercise. Details on this device are described in detail elsewhere.^{13,14} This system has been validated (against arterial blood gases) and used to control end-tidal gases during several different physiological stressors. A one-min average was taken for these variables at the end of the rest period, during the isometric HG protocol (HG and PECO), and each of the cycling exercise stages (EX1, EX2, EX3).

Extracranial blood flow. Simultaneous B-mode and Doppler ultrasonography were performed to assess continuous diameter and blood velocity recordings of the right IJV using a 10 MHz linear array probe attached to a high-resolution ultrasound machine (Terason t3200; Teratech, Burlington, MA, USA). The same experienced sonographer insonated the right IJV for all participants. The right IJV was

preferred since blood flow is usually greater in the right IJV compared to the left IJV. Of importance, this asymmetry in IJV drainage has no effect on brain NA spillover.¹⁵ The insonation angle remained unchanged at 60 degrees throughout the entire protocol. Ultrasound recordings were screen captured and saved for offline analysis. Blood flow analyses was performed using edge-detection software, which allows for the integration of synchronous diameter and velocity measurements to determine the mean beat-to-beat flow at 30 Hz independent of investigator bias.¹⁶ Mean blood flow was determined as half of the timeaveraged maximal velocity multiplied by the crosssectional luminal area for a minimum of 12 cardiac cycles.¹⁷ A one-min average, acquired from the final minute of each stage, was taken for these variables at the end of the rest periods, during the isometric HG protocol (HG and PECO), and each of the cycling exercise stages (EX1, EX2, EX3). IJV conductance was calculated using the following equation:

IJV conductance

$$=$$
 IJV blood flow/mean arterial pressure (1)

Arterial and venous catheterization. Three catheters were placed under sterile conditions prior to the experimental protocol: 1) an 18-gauge venous catheter into a superficial vein of the right forearm; 2) a 20-gauge arterial catheter (Arrow, Markham, ON, Canada) into the right radial artery, and 3) a 16-gauge central venous catheter (Arrow, Markham, ON, Canada) into the right IJV, which was advanced cephalad to the jugular bulb. Placement of the IJV was confirmed with an IJV oxygen saturation (S_JO₂) of <75%.

Arterial and venous blood analysis. Before sampling was started, the dead space volume was withdrawn and then an arterial sample (3 ml) was collected in preheparinized syringes (safePICO syringes, Radiometer, Copenhagen, Denmark). Blood samples were taken at the end of the rest periods, during the HG protocol (HG and PECO), and each of the cycling exercise stages (EX1, EX2, EX3). Air bubbles were immediately evacuated from the syringe, and blood gas analysis was performed within 5 min of sampling with a gas analyzer (ABL80 FLEX, Radiometer). The blood gas analyzer underwent a one-point calibration every 4 hrs, 2-point calibration every 8 hrs, and an additional automated 1-point calibration every 15 samples using manufacturer's standard quality checks, and ampule-based quality checks. All blood data were analyzed assuming a core body temperature of 37.0°C. Reported variables that were calibrated and analyzed included hemoglobin

concentration ([Hb]), (arterial) hematocrit (Hct), oxygen saturation (SO₂), partial pressure of oxygen (PO₂), partial pressure of carbon dioxide (PCO₂), pH, glucose, and lactate. Furthermore, arterial (CaO₂) and venous (CvO₂) oxygen content were calculated using the following equation:

Oxygen content (mL/dL) = $(1.34 \times [Hb] \times (SO_2/100)) + (0.0031 \times PO_2)$ (2)

Neurochemical assays. The assay of NA in plasma samples by means of high-performance liquid chromatography with electrochemical detection (HPLC-EC) was performed adapted from previously described methods.^{8,18} In brief, aliquots of 1000 µl of plasma were placed in 5-ml conical-based glass vials with 50 mg alumina, and the pH of the samples was adjusted to 8.6 by the addition of Tris buffer. Dihydroxybenzylamine hydrobromide was used as internal standard. The absorbed catecholamines were then eluted from the alumina with 300 µl of 0.2 M perchloric acid on Costar Spin-X microfilter tubes. From this eluate, 50 µl was injected into a high-pressure liquid chromatograph (Gilson Medical Electronics, Villiers le Bel, France) and 150 μ l were used for [³H]NE radioactivity measure.

 $[{}^{3}H]$ NA-radioactivity measure. Radioactivity was measured by liquid scintillation counting (liquid scintillation counter 1209 Rackbeta, LKB Wallac, Turku, Finland) in 150 µl aliquots from alumina eluate after addition of 10 ml of scintillation mixture (OptiPhase "HiSafe" 3, LKB, Loughborough, England). Final concentrations were corrected for loss during extraction using recoveries of internal standards.

Noradrenaline Spillover. At steady-state during the infusion of tritiated NA, the rate of spillover of NA into plasma and total NA clearance were calculated using the following equations:¹¹

NA spillover =
$$[{}^{3}H]NA$$
 infusion rate (dpm/min)
/plasma NA specific activity (dpm/pg) (4)

The rate of cerebral NA spillover was calculated using the Fick principle, which was corrected for the fractional extraction of [³H]NA:¹⁹

Cerebral NA spillover

$$= [(IJV [NA]-Arterial [NA]) + (Arterial [NA]
\times fractional extraction of cerebral [3H] NA)]
\times IJV plasma flow (5)$$

where IJV plasma flow represents IJV blood flow multiplied by 1 – Hct. Arterial Hct was substituted for IJV Hct in the plasma flow equation, which is further discussed in the "Methodological considerations" section of the manuscript. In the case of a measured negative fractional extraction of cerebral [³H]NA, we considered it as zero extraction on the ground of lack of plausibility.

Statistical analysis

Statistical analyses were performed using GraphPad Prism V8.0 (San Diego, CA) and reported as means \pm SD. Statistical significance was set at P < 0.05. Repeated measures one-way analysis of variance mixed effects models were used to detect statistical differences between rest, HG, and PECO (Tables 1, 2, and Figure 2), and rest, EX1, EX2, and EX3 (Tables 1, 2, and Figure 3). This statistical model accounts for the missing data, and when significant F-ratios were detected, post hoc comparisons were made using Tukey's test. The effect size (i.e., Cohen's d) values associated with our primary outcome (cerebral NA spillover) was computed using the estimated mean differences divided by the pooled standard deviation of the comparison.²⁰

Results

Participants

The participants included in this study data analysis (n = 12; 5 females) had a mean \pm SD age of 25.7 ± 2.8 years, height of $172.2 \pm 8.4 \text{ cm}$, and body mass of $71.7 \pm 13.8 \text{ kg}$. Our primary outcome measure, cerebral NA spillover, was obtained in only a portion of participants during the HG + PECO protocol (n = 9), and cycling exercise EX1 (n = 7), EX2 (n = 5), and EX3 (n = 5). Participant dropout was primarily due to inadequate image quality during exercise. Individual data for NA spillover and clearance are illustrated in Figures 2 and 3.

The effects of exercise on cardiovascular and respiratory variables

Table 1 illustrates cardiovascular and respiratory data during exercise. During isometric HG and PECO, IJV

IJV Diameter (mm) 9.1 ±		HG EX	PECO	Rest	Cyc EXI	Cyc EX2	Cyc EX3
	± 2.0	8.9 ± 2.1	8.6 ±2.0	7.5 ± 1.8	7.2 ± 1.9	6.8 ± 1.8	8.6±1.6
IJV IMean velocity (cm/s) 3.1 ±	± 3.4	$12.4\pm4.5^{*}$	11.7 ± 4.1	10.5 ± 3.3	10.0 ± 3.0	$\textbf{9.2}\pm\textbf{1.9}$	7.8 ± 3.8
IJV Mean blood flow (ml/min) 338.5 \pm	± 140.9	$421.7 \pm 108.8^{*}$	$379.2 \pm 127.3^{*}$	284.7 ± 153.1	276.1 ± 194.7	209.5 ± 114.0	255.2 ± 105.1
IJV Con (ml/min/mmHg) 3.4 ±	± 1.2	3.7 ± 1.0	3.4±1.1	3.0 ± 1.5	2.7 ± 1.7	$2.0\pm1.1*$	2.3 ± 0.8
Systolic BP (mmHg) I 25.4 \pm	± 12.1	$140.9 \pm 15.6^{*}$	$135.6 \pm 12.1^{*}$	123.9 ± 10.5	$169.3 \pm 15.1^{*}$	$177.4 \pm 20.4^{*}$	$179.7 \pm 18.2^{*}$
Diastolic BP (mmHg) 84.6 \pm	± 8.2	$97.4\pm12.8^{*\ddagger}$	$\textbf{92.2}\pm\textbf{8.8*}$	78.2 ± 5.7	83.7±9.1*	$84.3 \pm 7.7*$	$84.4\pm5.8^{*}$
Mean BP (mmHg) 98.9 ±	± 9.8	$117.0 \pm 13.7^{*\pm}$	$110.0 \pm 11.5^{*}$	93.5 ± 8.1	$105.1 \pm 12.0^{*}$	$109.2 \pm 11.3^{*}$	$112.9 \pm 11.5^{*\ddagger}$
Heart rate (bpm) $66.6 \pm$	± 11.6	$\textbf{82.6} \pm \textbf{11.3}^{\texttt{*}\ddagger}$	71.2 ± 11.4	62.5 ± 10.1	$105.2 \pm 8.4^{*}$	$127.3\pm13.4^{*\ddagger}$	$I50.8\pmI6.2^{*\ddagger}$
Ventilation (I/min) 22.7 ±	± 4.5	$33.8 \pm 10.8^{*}$	27.1 ± 8.4				
Tidal volume (I)	± 0.2	$1.5\pm0.5^{*}$	1.5 ± 0.2				
Breathing rate (breaths/min) 19.6 \pm	± 1.9	$22.7\pm4.6^{\ddagger}$	18.3 ± 4.4				
P _{ET} CO ₂ (mmHg) 43.7 ±	± 2.0	44.3 ± 1.9	$\textbf{44.3} \pm \textbf{1.8}$				
$P_{\text{ET}}O_2 \text{ (mmHg)} \qquad \qquad 86.6 \pm$	± 5.1	$\textbf{87.8}\pm\textbf{4.2}$	88.2 ± 4.7				

minute; mm: millimeter; mmHg: millimeters of mercury; PECO: post-exercise circulatory occlusion; P_{ET}CO₂: Partial pressure of end-tidal carbon dioxide; P_{ET}O₂: Partial pressure of end-tidal oxygen. *P < 0.05 vs rest. ⁺P < 0.05 vs Cyc EX2. ⁺P < 0.05: HG EX vs PECO, and vs Cyc EX1.

diameter remained unchanged (P=0.748), but IJV blood velocity was elevated during HG (P=0.0229); and IJV blood flow was elevated during both isometric HG (P=0.0035) and PECO (P=0.0359) compared to rest. During isometric HG and PECO, systolic (P<0.01), diastolic (P<0.001), and mean arterial blood pressure (P<0.001) all increased compared to rest; however, IJV conductance was unchanged (P=0.251). Heart rate was only elevated during isometric HG (P<0.001), but not during PECO (P=0.0658). Minute ventilation and tidal volume

were elevated during isometric HG (P < 0.001 and P = 0.0318, respectively), but not during PECO (P = 0.13 and P = 0.06, respectively). Breathing frequency did not change between rest and isometric HG (P = 0.09) or PECO (P = 0.59), but it was elevated during isometric HG compared to PECO (P = 0.0146). Both $P_{ET}CO_2$ and $P_{ET}O_2$ were controlled to resting values during isometric HG and PECO (P = 0.122

and P = 0.136, respectively). During supine cycling exercise, IJV diameter, blood velocity, and blood flow did not change compared to rest (P = 0.291, P = 0.126, and P = 0.141, respectively). Systolic, diastolic, and mean arterial blood pressure were all elevated compared to rest (all P < 0.001), and IJV conductance was reduced during EX2 compared to rest (P = 0.0459). Heart rate was elevated at each exercise intensity compared to rest (P < 0.001).

The effects of exercise on arterial and venous blood

Table 2 illustrates arterial and venous blood data. Arterial NA was elevated during isometric HG (P=0.0237), but not PECO (P=0.0848), and there were no changes in IJV NA during isometric HG and PECO (main effect: P=0.547) compared to rest. During isometric HG and PECO, there were no differences found for arterial or venous [Hb], oxygen content, oxygen saturation, PO₂, PCO₂, pH, glucose, lactate and arterial Hct (all P > 0.05).

Arterial NA and IJV NA were elevated at EX3 compared to both rest and EX1 (all P < 0.01). Both arterial [Hb] and Hct were elevated during supine cycling exercise compared to rest (main effects: P < 0.04). Internal jugular C_VO_2 , SO_2 , PO_2 , and PCO_2 all declined with supine cycling exercise (main effects: P < 0.05), while arterial CaO₂, and PO₂ increased compared to rest (P < 0.05). Arterial SO₂ remained unchanged (main effect: P = 0.643), and arterial PCO₂ decreased (P < 0.001). Internal jugular pH decreased during supine cycling exercise (main effect: P = 0.05). Internal jugular and arterial glucose did not change with supine cycling exercise (main effects: P = 0.907

Ć

i i i

	Rest	HG EX	PECO	Rest	Cyc EXI	Cyc EX2	Cyc EX3
IJV NA (pg/ml)	$\textbf{468.6} \pm \textbf{441.0}$	$\textbf{576.0} \pm \textbf{457.1}$	$\textbf{585.9} \pm \textbf{466.8}$	$\textbf{467.2} \pm \textbf{442.6}$	$\textbf{465.8} \pm \textbf{323.1}$	997.3 \pm 813.4	1275.3±831.3* [‡]
ART NA (pg/ml)	$\textbf{404.5} \pm \textbf{412.4}$	$\textbf{593.1} \pm \textbf{409.1}^{\texttt{*}}$	$\textbf{638.0} \pm \textbf{475.3}$	$\textbf{419.9} \pm \textbf{300.5}$	$\textbf{480.4} \pm \textbf{309.1}$	$\textbf{913.6} \pm \textbf{691.9}$	1259.0 \pm 838.0* $^{\!$
A-V NA (pg/ml)	-64.1 ± 115.6	$\textbf{17.1} \pm \textbf{90.7}$	$\textbf{52.1} \pm \textbf{146.6}$	-47.3 ± 184.3	-13.4 ± 109.3	-25.0 ± 177.7	-91.7 ± 324.5
ART [Hb] (g/dl)	14.9 ± 1.2	14.8 ± 1.1	$\textbf{14.6} \pm \textbf{1.3}$	14.7 ± 1.2	$14.8\pm1.3^{*}$	$15.2\pm1.4^{*\ddagger}$	$15.3\pm1.2^{*\ddagger}$
ART Hct (%)	$\textbf{45.6} \pm \textbf{3.7}$	$\textbf{45.3} \pm \textbf{3.5}$	$\textbf{44.9} \pm \textbf{4.0}$	$\textbf{45.1} \pm \textbf{3.6}$	$\textbf{45.5} \pm \textbf{3.9}^{\texttt{*}}$	$46.5\pm4.2^{*\pm}$	$\textbf{46.9} \pm \textbf{3.7}^{\texttt{*}\ddagger}$
CvO ₂ (ml/dl)	14.2 ± 1.8	14.6 ± 1.7	13.6 ± 2.5	$\textbf{13.1}\pm\textbf{1.9}$	$13.4\pm1.5^{*}$	13.3 ± 1.5	12.3 \pm 1.7 ^{†‡}
CaO ₂ (ml/dl)	19.7 ± 1.6	19.7 ± 1.5	19.5 ± 1.7	19.6 ± 1.6	$\textbf{19.8} \pm \textbf{1.7}$	$20.2\pm1.8^{*\ddagger}$	$20.4\pm1.6^{*\ddagger}$
A-V O ₂ content (ml/dl)	5.5 ± 1.3	5.1 ± 1.2	$\textbf{5.9} \pm \textbf{1.6}$	6.5 ± 1.7	$\textbf{6.4} \pm \textbf{0.9}$	$\textbf{6.9} \pm \textbf{1.0}$	8.1 \pm 1.1* ^{†‡}
IJV SO ₂ (%)	$\textbf{70.9} \pm \textbf{4.2}$	$\textbf{72.7} \pm \textbf{5.6}$	$\textbf{67.3} \pm \textbf{9.8}$	64.6 ± 7.3	65.4 ± 3.7	$\textbf{62.6} \pm \textbf{6.9}$	58.3 \pm 5.9 $^{*^{\dagger \ddagger}}$
ART SO ₂ (%)	$\textbf{97.3} \pm \textbf{0.6}$	$\textbf{97.8} \pm \textbf{0.6}$	$\textbf{97.6} \pm \textbf{0.7}$	$\textbf{97.8} \pm \textbf{0.4}$	$\textbf{97.8} \pm \textbf{0.4}$	$\textbf{97.8} \pm \textbf{0.7}$	$\textbf{98.0} \pm \textbf{0.5}$
A-V SO ₂ (%)	$\textbf{26.3} \pm \textbf{4.7}$	$\textbf{25.1} \pm \textbf{5.8}$	$\textbf{30.3} \pm \textbf{10.2}$	$\textbf{33.2} \pm \textbf{7.2}$	$\textbf{32.4} \pm \textbf{3.9}$	$\textbf{35.2} \pm \textbf{7.3}$	$39.7 \pm 6.1^{*1\pm}$
IJV PO ₂ (mmHg)	$\textbf{43.1} \pm \textbf{4.3}$	44.3 ± 4.8	40.3 ± 7.0	$\textbf{38.2} \pm \textbf{4.4}$	$\textbf{38.6} \pm \textbf{2.8}$	$\textbf{37.4} \pm \textbf{2.7}$	$35.7 \pm 2.8^{\ddagger}$
ART PO ₂ (mmHg)	$\textbf{95.7} \pm \textbf{5.2}$	102.1 ± 8.4	$\textbf{102.8} \pm \textbf{13.7}$	102.1 ± 7.1	102.8 ± 8.1	100.1 ± 6.1	$106.5\pm7.7^{*\dagger}$
A-V PO ₂ (mmHg)	$\textbf{52.6} \pm \textbf{7.6}$	$\textbf{57.8} \pm \textbf{10.7}$	$\textbf{62.6} \pm \textbf{18.5}$	$\textbf{63.9} \pm \textbf{5.9}$	64.2 ± 7.8	62.7 ± 6.3	$\textbf{70.8} \pm \textbf{9.2}^{\texttt{*}^{\ddagger\ddagger}}$
IJV PCO ₂ (mmHg)	52.1 ± 4.4	$\textbf{52.5} \pm \textbf{4.0}$	51.1 ± 4.1	51.1 ± 3.1	$\textbf{50.8} \pm \textbf{3.4}$	$\textbf{49.6} \pm \textbf{4.5}^{*}$	$47.5 \pm 5.0^{*^{\dagger \ddagger}}$
ART PCO ₂ (mmHg)	44.6 ± 2.5	$\textbf{43.4} \pm \textbf{3.3}$	$\textbf{42.0} \pm \textbf{6.1}$	40.5 ± 2.5	40.2 ± 3.3	$\textbf{38.3} \pm \textbf{4.2}^{\texttt{*}\ddagger}$	$\textbf{34.9} \pm \textbf{3.8}^{\texttt{*}^{\ddagger \ddagger}}$
A-V PCO ₂ (mmHg)	-7.5 ± 2.9	-9.1 ± 3.2	-9.1 ± 4.1	-9.7 ± 3.8	-10.6 ± 1.2	-11.2 ± 1.2	$-12.6 \pm 2.1^{*\pm}$
IJV pH	7.34 ± 0.02	$\textbf{7.35} \pm \textbf{0.02}$	7.35 ± 0.02	7.35 ± 0.01	7.34 ± 0.02	$\textbf{7.34} \pm \textbf{0.02}^{\texttt{*}\ddagger}$	$\textbf{7.33} \pm \textbf{0.02}^{\texttt{*}^\ddagger}$
ART pH	$\textbf{7.38} \pm \textbf{0.02}$	$\textbf{7.39} \pm \textbf{0.02}$	$\textbf{7.39} \pm \textbf{0.05}$	$\textbf{7.40} \pm \textbf{0.02}$	$\textbf{7.39} \pm \textbf{0.02}$	$\textbf{7.39} \pm \textbf{0.02}$	$\textbf{7.39} \pm \textbf{0.02}$
A-V pH	0.04 ± 0.01	0.05 ± 0.01	0.04 ± 0.03	0.05 ± 0.01	0.05 ± 0.01	0.05 ± 0.01	0.06 ± 0.01
IJV Glucose (mmol/l)	$\textbf{5.4} \pm \textbf{0.8}$	$\textbf{5.6} \pm \textbf{1.0}$	5.5 ± 1.1	5.1 ± 0.4	5.1 ± 0.2	5.1 ± 0.3	$\textbf{5.0} \pm \textbf{0.6}$
ART Glucose (mmol/l)	6.0 ± 1.2	$\textbf{6.0} \pm \textbf{1.0}$	$\textbf{6.1} \pm \textbf{1.0}$	5.6 ± 0.3	5.6 ± 0.3	5.6 ± 0.3	5.7 ± 0.6
A-V Glucose (mmol/l)	$\textbf{0.6}\pm\textbf{0.6}$	$\textbf{0.4}\pm\textbf{0.1}$	$\textbf{0.6}\pm\textbf{0.2}$	$\textbf{0.5}\pm\textbf{0.2}$	$\textbf{0.6}\pm\textbf{0.2}$	0.5 ± 0.1	$\textbf{0.7}\pm\textbf{0.2}^{*\dagger\ddagger}$
IJV Lactate (mmol/l)	0.7 ± 0.1	$\textbf{0.8}\pm\textbf{0.2}$	$\textbf{0.8}\pm\textbf{0.4}$	1.0 ± 0.6	1.9 ± 0.9	$3.3\pm1.9^{*\pm}$	$4.8\pm1.8^{*\dagger\ddagger}$
ART Lactate (mmol/l)	0.6 ± 0.1	$\textbf{0.8}\pm\textbf{0.2}$	$\textbf{0.8}\pm\textbf{0.5}$	$\textbf{0.9}\pm\textbf{0.7}$	1.9 ± 1.0	$3.4\pm2.1^{*\pm}$	$5.2 \pm 2.2^{*^{\dagger \ddagger}}$
A-V Lactate (mmol/l)	-0.04 ± 0.09	-0.02 ± 0.08	0.02 ± 0.10	-0.08 ± 0.15	-0.00 ± 0.03	0.1 ± 0.4	$\textbf{0.4}\pm\textbf{0.7*}$

Table 2. Arterial and venous blood data during isometric handgrip exercise and post-exercise circulatory occlusion and supine cycling exercise.

ART: arterial; CaO₂: arterial oxygen content; CvO₂: venous oxygen content; g/dl: grams per deciliter; [Hb]: hemoglobin concentration; Hct: hematocrit; HG EX: handgrip exercise; IJV: internal jugular vein; ml/dl: milliliters per deciliter; mmHg: millimeters of mercury; mmol/l: millimole per liter; NA: noradrenaline; pg/ml: picogram per millimeter; PCO₂: partial pressure of carbon dioxide; PECO: post-exercise circulatory occlusion; PO₂: partial pressure of oxygen; SO₂: oxygen saturation. *P < 0.05 vs rest. [†]P < 0.05 vs Cyc EX2. [‡]P < 0.05: HG EX vs PECO, and vs Cyc EX1.

and P = 0.818, respectively), but IJV and arterial lactate increased (main effects: P < 0.001).

The effects of exercise on noradrenaline spillover and clearance

Figure 2 illustrates NA data during isometric HG exercise. Total NA spillover was elevated during isometric HG (P=0.049) and persisted during the PECO (P=0.006); however, NA clearance remained unchanged compared to rest (P=0.130). Cerebral NA spillover remained unchanged during isometric HG exercise (P=0.36; Cohen's d=0.46) and during the PECO (P=0.45; Cohen's d=0.40), compared to rest.

Figure 3 illustrates NA data during supine cycling exercise. Total NA spillover during supine cycling exercise was higher during EX2 and EX3 compared to rest (P = 0.046, and P = 0.025, respectively). NA clearance was reduced at EX2 and EX3 compared to EX1 (P = 0.026 and P = 0.003, respectively). Despite changes in total NA spillover and clearance, cerebral NA spillover remained unchanged during supine cycling exercise (main effect: P = 0.94; Cohen's d = 0.10).

Discussion

Our study yielded several key findings: 1) in support to our initial hypothesis, cerebral NA spillover remained unchanged during isometric HG exercise and PECO, and supine cycling exercise; and 2) total NA spillover increased during isometric exercise and PECO, as well as during dynamic supine cycling exercise, compared to rest. Taken together, these results indicate that transient increases in arterial blood pressure induced by acute moderate exercise does not engage cerebral SNA in healthy humans. We discuss the implications of these results in detail in the sections below.

Sympathetic nerve activity during transient increases in arterial blood pressure

It is well-established that both isometric and dynamic exercise lead to an increase in heart rate, arterial blood pressure, and peripheral SNA. Traditionally, SNA signaling to the peripheral vasculature results in increased smooth muscle tone (i.e., vasoconstriction). The mechanisms behind this involve a complex interplay between central command,²¹ afferent signaling from mechanical and metabolic sensory receptors in skeletal muscles,²²



Figure 2. Isometric handgrip exercise on noradrenaline spillover and clearance. Mean \pm SD and individual data for total noradrenaline spillover (Panel a; n = 11), total noradrenaline clearance (Panel b; n = 12), and brain noradrenaline spillover (Panel c; n = 10 [Rest], 9 [HG], 9 [PECO]). Statistical comparisons that yielded P < 0.05 are illustrated on the Figure.



Figure 3. Supine cycling exercise on noradrenaline spillover and clearance. Mean \pm SD and individual data for total noradrenaline spillover (Panel a; n = 11 [Rest], 10 [EX1], 9 [EX2], 8 [EX3]), total noradrenaline clearance (Panel b; n = 11 [Rest], 9 [EX1], 9 [EX2], 8 [EX3]), and brain noradrenaline spillover (Panel c; n = 9 [Rest], 7 [EX1], 5 [EX2], 5 [EX3]). Statistical comparisons that yielded P < 0.05 are illustrated on the Figure.

feedback from stretch receptors in the carotid arteries and aortic arch,²³ low pressure stretch receptors in the heart, great veins, and pulmonary vessels,²⁴ arterial chemoreceptors,²⁵ and phrenic activity.²⁶ Microneurography and plasma NA are commonly used experimental techniques to measure SNA during both isometric (e.g., HG) and dynamic (e.g., cycling) exercise. Previous research has demonstrated that peripheral SNA robustly increases and even persists during PECO.^{27–31} Our study's findings are consistent with the current body of literature, as we observed an increase in total NA spillover during both isometric (Figure 2) and dynamic exercise (Figure 3).

The question of whether peripheral SNA accurately reflects cerebral SNA in humans at rest, or during physiological stress (e.g., exercise) has been a topic of controversy. This is because different vascular beds in the human body have varying distributions of sympathetic neurotransmitter receptor densities,³² and

possibly different sensitivities. Thus, it is plausible and perhaps even likely - that the cerebrovasculature responds differently compared to vascular beds located in the periphery given its unique features and functions, such as the circle of Willis and the brain confinement to the human skull. From a teleological standpoint, it would be physiologically beneficial for cerebral SNA to increase similarly to peripheral SNA in response to transiently elevated arterial blood pressure during exercise as a protective mechanism against overperfusion. Indeed, this hypothesis is supported by animal models that have demonstrated increases in SNA during acute increases in arterial blood pressure.^{5,33} However, there is evidence of interspecies differences in CBF regulation,^{33–35} which underscores the importance of conducting cerebral NA spillover studies in humans. The limited data collected in healthy humans suggests that peripheral and cerebral SNA are similar, at least under resting conditions.³⁶ The data from the current study

highlights that in contrast to available animal data, and in support of the limited data collected in healthy humans,⁷ transient increases in arterial blood pressure during moderate intensity isometric and dynamic exercise had no impact on cerebral SNA.

The influence of sympathetic nerve activity on the cerebrovasculature

The extent to which SNA regulates the cerebrovasculature remains under scrutiny,^{37,38} despite the rich innervation of the cerebral circulation by both sympathetic and parasympathetic nerve fibers. Various pharmacological blockades and ganglion excision studies conducted in humans over the past few decades have shown that altering sympathetic signaling to the cerebrovasculature can affect resting CBF and reactivity to physiological stressors such as carbon dioxide and transient decreases and increases in arterial blood pressure.^{37,38} However, the interpretation of these studies is challenging due to methodological limitations associated with the use of transcranial Doppler ultrasound and the simultaneous effects of drug interactions on physiological parameters that can directly influence cerebrovascular tone, such as arterial blood pressure and cardiac output.

Cerebral sympathetic nerve activity during moderate intensity isometric handgrip exercise and post-circulatory occlusion. During aerobic exercise, cardiovascular adjustments are necessary to ensure sufficient oxygen and nutrient

are necessary to ensure sufficient oxygen and nutrient supply to active skeletal muscle tissue and active brain regions, resulting in a 10-20% increase in cerebral perfusion.³⁹ However, it remains unclear whether SNAmediated protective mechanisms for overperfusion are present at these modest levels of transient increases in arterial blood pressure during isometric and dynamic exercise. Surprisingly, brain NA spillover remained unchanged during isometric exercise, indicating that SNA is not a regulatory mechanism for the cerebrovasculature during isometric exercise as used in the current study. Only one study has investigated the sympathetic control of CBF in healthy humans using pharmacological intervention (i.e., prazosin) during isometric HG exercise,¹⁰ which found that regional sympathetic neural restraint may be present. The discrepancy between our findings and previous research may be due to the NA spillover technique providing a whole brain measurement of SNA, rather than measuring a regional response. One interesting finding was that brain NA spillover also remained unchanged during PECO, after the removal of central command (i.e., voluntary muscle contraction). Future work should focus on the effects of non-exercise induced sympathetic stress on cerebral NA spillover.

Cerebral sympathetic nerve activity during mild-to-moderate intensity subine cycling exercise. Similar to previous work,⁷ we found that brain NA spillover remained unchanged during mild-to-moderate intensity supine cycling exercise. Other studies have investigated the effects of SNA on CBF regulation during dynamic exercise using pharmacological interventions. Two studies used β -adrenergic blockades (metoprolol and propranolol) and found that they had no effect on CBF regulation at rest or during cycling exercise.^{40,41} Another study measured the impact of ganglion blockade, which has demonstrated to reduce CBF at rest.⁴² during cycling exercise with simultaneous administration of a β -adrenergic blockade (e.g., metoprolol) or placebo. The data from the latter study indicate that under cycling exercise conditions where elevations in cardiac output are limited (e.g., after β -adrenergic blockade), global SNA signalling increases $\sim 2-3x$ (measured via arterial NA), and is associated with a reduction in the CBF response to exercise.⁴³ The complex pharmacological interventions and extreme conditions of limited cardiac output during dynamic exercise make these studies difficult to compare to the current study, but taken together, submaximal dynamic exercise in itself may not be a strong enough physiological stressor to elicit profound increases in cerebral SNA.

Methodological considerations

Although these are the only available data on cerebral NA spillover during both isometric and dynamic exercise, there are several methodological limitations that require further discussion. First, this invasive study includes a small sample size of healthy males and females and some data were lost during the protocol, primarily due to inadequate image quality during exercise. This work cannot be generalized to older participants or clinical populations (e.g. patients with arterial hypertension). Second, the current findings do not exclude the notion cerebral SNA acts as a protective mechanism for the cerebrovasculature in response to increases in arterial blood pressure in other physiological/clinical conditions. Indeed, it was not possible to measure cerebral SNA during transient surges in arterial blood pressure, but only following relatively slow steady-state increases (i.e. over 2 (HG exercise) to 3 (supine cycling exercise) minutes). A strength to the study is that HG and cycling exercise yielded similar increases in mean arterial blood pressure (18.3 and 20.7%, respectively). However, the duration and nature of the stimulation may influence the cerebrovascular response and SNA control of the cerebral circulation may be more efficient under dynamic than steady-state conditions.^{37,44} Third, the participants underwent various stressful procedures and interventions, including catheter placements, exercise, multiple ultrasound methods, and had to breathe on a custom end-tidal forcing system, resulting in resting arterial and venous NA concentrations that were higher than anticipated. However, the individual trends in NA concentrations (and calculated whole body NA spillover) in response to exercise were consistent with previous research. Fourth, IJV Hct, a parameter included in the calculation of IJV plasma flow, was not included in our panel of blood parameters analyzed, although arterial Hct was. We have substituted IJV Hct for arterial Hct in the plasma flow equation. Considering venous Hct is $\sim 3\%$ higher than arterial Hct,⁴⁵ our cerebral NA spillover values will be slightly overestimated. However, this should not have impacted within-participant comparisons. Fifth, we attempted to take measurements of CBF (e.g., internal carotid artery blood flow) during the exercise protocols, but due to the difficulty of obtaining simultaneous CBF and IJV blood flow, we decided to solely focus on the IJV measurements for the calculation of cerebral NA spillover. Future studies using cerebral NA spillover should aim to collect these complementary data to gain a full scope of the cerebrovascular response to physiological stressors. Sixth, due to the nature of the experimental measurements taken and the possibility of IJV collapse in the upright position, submaximal exercise was completed in the supine position, which could impact the performance and the physiological response to cycling exercise. Seventh, there is evidence that regional NA spillover underestimates true NA release, therefore, small changes in SNA via NA spillover may be difficult to detect.⁴⁶ Eighth, testing participants in the supine position loads cardiopulmonary baroreceptors, which may influence the sympathetic nerve response to exercise. Lastly, the cerebrovasculature is richly innervated by parasympathetic nerve fibers, yet our data do not account for any contribution of parasympathetic nerve activity on the cerebrovascular function. This is a crucial area of research that should be addressed in future studies.

Perspectives

These results represent a key milestone towards the examination of the impact of exercise on cerebral SNA in healthy humans. Further research aimed at improving our understanding of the role of SNA on cerebrovascular regulation in several clinical conditions associated with central sympathetic overactivity such as ischemic heart disease, chronic heart failure, autonomic disorders, and arterial hypertension are needed. Furthermore, the exercise stressors conducted in this study yield moderate elevations of peripheral SNA, and more powerful sympathetic stimuli should be explored in future studies in healthy humans. A better understanding of the physiological role of SNA on the cerebrovasculature during different physiological and clinical conditions in humans may also provide the basis for: 1) developing new approaches to pharmacologically manage pathological conditions characterized by cerebral sympathetic dysregulation and impaired CBF control and, 2) new therapeutic targets for early detection and prevention of cerebral hemorrhage/ischemic injuries and better management of CBF during anesthesia or in intensive care.

Conclusion

This is the first investigation to measure brain specific sympathetic nerve activity using the noradrenaline spillover technique during isometric and dynamic exercise in humans. The primary finding was that cerebral noradrenaline spillover remained unchanged during isometric (i.e., handgrip), post exercise cuff occlusion, and dynamic (i.e., cycling) exercises. The data from the current study opposes the available animal literature but are largely in support of the available healthy human data. Future research should focus on the potential effects of other mild-to-strong sympathetic stressors on cerebral sympathetic nerve activity in both healthy and disease populations to gain a better understanding of its role within the cerebrovasculature.

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: M.M.T. was funded by a NSERC CGS PhD award. G.C. was funded by a NSERC PGS PhD award. P.N.A. was funded by a Canada Research Chair award. P.B. was funded by a NSERC Discovery grant and the Foundation of the Institut universitaire de cardiologie et de pneumologie de Quebec.

Acknowledgements

We would like to thank the study participants for volunteering their time for this study.

Declaration of conflicting interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Authors' contributions

The experiments were performed in the laboratory of P.B. M.M.T., P.N.A., and P.B., contributed to conception or design of the work.

M.M.T., A.D., M.A.V.C., L.L., S.I., G.B.C., S.L., M.F., N.C., P.N.A., and P.B., contributed to acquisition, analysis or interpretation of data for the work.

M.M.T., A.D., M.A.V.C., L.L., S.I., G.B.C., S.L., M.F., N.C., P.N.A., and P.B., contributed to drafting the work or revising it critically for important intellectual content.

All authors approved the final version of the manuscript; agree to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved; and all persons designated as authors qualify for authorship, and all those who qualify for authorship are listed.

ORCID iD

Patrice Brassard D https://orcid.org/0000-0002-6254-5044

References

- Bailey DM, Evans KA, McEneny J, et al. Exerciseinduced oxidative-nitrosative stress is associated with impaired dynamic cerebral autoregulation and bloodbrain barrier leakage. *Exp Physiol* 2011; 96: 1196–1207.
- Willie CK, Tzeng YC, Fisher JA, et al. Integrative regulation of human brain blood flow. *J Physiol* 2014; 592: 841–859.
- Brassard P, Labrecque L, Smirl JD, et al. Losing the dogmatic view of cerebral autoregulation. *Physiol Rep* 2021; 9: e14982.
- Claassen J, Thijssen DHJ, Panerai RB, et al. Regulation of cerebral blood flow in humans: physiology and clinical implications of autoregulation. *Physiol Rev* 2021; 101: 1487–1559.
- Cassaglia PA, Griffiths RI and Walker AM. Sympathetic nerve activity in the superior cervical ganglia increases in response to imposed increases in arterial pressure. *Am J Physiol Regul Integr Comp Physiol* 2008; 294: R1255–1261.
- Ferrier C, Esler MD, Eisenhofer G, et al. Increased norepinephrine spillover into the jugular veins in essential hypertension. *Hypertension* 1992; 19: 62–69.
- Ide K, Horn A and Secher NH. Cerebral metabolic response to submaximal exercise. J Appl Physiol (1985) 1999; 87: 1604–1608.
- Esler M, Jennings G, Korner P, et al. Assessment of human sympathetic nervous system activity from measurements of norepinephrine turnover. *Hypertension* 1988; 11: 3–20.
- Mitchell DA, Lambert G, Secher NH, et al. Jugular venous overflow of noradrenaline from the brain: a neurochemical indicator of cerebrovascular sympathetic nerve activity in humans. *J Physiol* 2009; 587: 2589–2597.
- Fernandes IA, Mattos JD, Campos MO, et al. Selective alpha1-adrenergic blockade disturbs the regional distribution of cerebral blood flow during static handgrip exercise. *Am J Physiol Heart Circ Physiol* 2016; 310: H1541–1548.
- 11. Esler M, Jackman G, Bobik A, et al. Determination of norepinephrine apparent release rate and clearance in humans. *Life Sci* 1979; 25: 1461–1470.
- Alam M and Smirk FH. Observations in man upon a blood pressure raising reflex arising from the voluntary muscles. *J Physiol* 1937; 89: 372–383.

- Tymko MM, Ainslie PN, MacLeod DB, et al. End tidalto-arterial CO₂ and O₂ gas gradients at low- and highaltitude during dynamic end-tidal forcing. *Am J Physiol Regul Integr Comp Physiol* 2015; 308: R895–906.
- Tymko MM, Hoiland RL, Kuca T, et al. Measuring the human ventilatory and cerebral blood flow response to CO2: a technical consideration for the end-tidalto-arterial gas gradient. *J Appl Physiol (1985)* 2016; 120: 282–296.
- Ferrier C, Jennings GL, Eisenhofer G, et al. Evidence for increased noradrenaline release from subcortical brain regions in essential hypertension. *J Hypertens* 1993; 11: 1217–1227.
- Woodman RJ, Playford DA, Watts GF, et al. Improved analysis of brachial artery ultrasound using a novel edgedetection software system. *J Appl Physiol (1985)* 2001; 91: 929–937.
- Thomas KN, Lewis NC, Hill BG, et al. Technical recommendations for the use of carotid duplex ultrasound for the assessment of extracranial blood flow. *Am J Physiol Regul Integr Comp Physiol* 2015; 309: R707–720.
- Eisenhofer G, Goldstein DS, Ropchak TG, et al. Source and physiological significance of plasma 3,4-dihydroxyphenylglycol and 3-methoxy-4-hydroxyphenylglycol. *J Auton Nerv Syst* 1988; 24: 1–14.
- Esler M, Willett I, Leonard P, et al. Plasma noradrenaline kinetics in humans. J Auton Nerv Syst 1984; 11: 125–144.
- Maher JM, Markey JC and Ebert-May D. The other half of the story: effect size analysis in quantitative research. *CBE Life Sci Educ* 2013; 12: 345–351.
- Waldrop TG, Eldridge FL, Iwamoto GA, et al. Central neural control of respiration and circulation during exercise. *Comprehensive Physiology*: 333–380.
- 22. Kaufman MP and Hayes SG. The exercise pressor reflex. *Clin Auton Res* 2002; 12: 429–439.
- Raven PB, Potts JT and Shi X. Baroreflex regulation of blood pressure during dynamic exercise. *Exerc Sport Sci Rev* 1997; 25: 365–389.
- Fadel PJ and Raven PB. Human investigations into the arterial and cardiopulmonary baroreflexes during exercise. *Exp Physiol* 2012; 97: 39–50.
- Marshall JM. Peripheral chemoreceptors and cardiovascular regulation. *Physiol Rev* 1994; 74: 543–594.
- Rodman JR, Henderson KS, Smith CA, et al. Cardiovascular effects of the respiratory muscle metaboreflexes in dogs: rest and exercise. *J Appl Physiol (1985)* 2003; 95: 1159–1169.
- Mark AL, Victor RG, Nerhed C, et al. Microneurographic studies of the mechanisms of sympathetic nerve responses to static exercise in humans. *Circ Res* 1985; 57: 461–469.
- Victor RG, Seals DR and Mark AL. Differential control of heart rate and sympathetic nerve activity during dynamic exercise. Insight from intraneural recordings in humans. J Clin Invest 1987; 79: 508–516.
- Victor RG and Seals DR. Reflex stimulation of sympathetic outflow during rhythmic exercise in humans. *Am J Physiol* 1989; 257: H2017–2024.

- Seals DR, Victor RG and Mark AL. Plasma norepinephrine and muscle sympathetic discharge during rhythmic exercise in humans. J Appl Physiol (1985) 1988; 65: 940–944.
- Pott F, Ray CA, Olesen HL, et al. Middle cerebral artery blood velocity, arterial diameter and muscle sympathetic nerve activity during post-exercise muscle ischaemia. *Acta Physiol Scand* 1997; 160: 43–47.
- Gezalian MM, Mangiacotti L, Rajput P, et al. Cerebrovascular and neurological perspectives on adrenoceptor and calcium channel modulating pharmacotherapies. J Cereb Blood Flow Metab 2021; 41: 693–706.
- Heistad DD, Marcus ML and Gross PM. Effects of sympathetic nerves on cerebral vessels in dog, cat, and monkey. *Am J Physiol* 1978; 235: H544–552.
- 34. Sándor P. Nervous control of the cerebrovascular system: doubts and facts. *Neurochem Int* 1999; 35: 237–259.
- Edvinsson L, MacKenzie ET and McCulloch J. Cerebral blood flow and metabolism. Newark: Raven Press (ID), 1993.
- Lambert GW, Thompson JM, Turner AG, et al. Cerebral noradrenaline spillover and its relation to muscle sympathetic nervous activity in healthy human subjects. *J Auton Nerv Syst* 1997; 64: 57–64.
- Brassard P, Tymko MM and Ainslie PN. Sympathetic control of the brain circulation: appreciating the complexities to better understand the controversy. *Auton Neurosci* 2017; 207: 37–47.
- Koep JL, Taylor CE, Coombes JS, et al. Autonomic control of cerebral blood flow: fundamental comparisons between peripheral and cerebrovascular circulations in humans. J Physiol 2022; 600: 15–39.

- Tymko MM, Ainslie PN and Smith KJ. Evaluating the methods used for measuring cerebral blood flow at rest and during exercise in humans. *Eur J Appl Physiol* 2018; 118: 1527–1538.
- 40. Ogoh S, Dalsgaard MK, Secher NH, et al. Dynamic blood pressure control and middle cerebral artery mean blood velocity variability at rest and during exercise in humans. *Acta Physiol (Oxf)* 2007; 191: 3–14.
- Seifert T, Rasmussen P, Secher NH, et al. Cerebral oxygenation decreases during exercise in humans with beta-adrenergic blockade. *Acta Physiol (Oxf)* 2009; 196: 295–302.
- Umeyama T, Kugimiya T, Ogawa T, et al. Changes in cerebral blood flow estimated after stellate ganglion block by single photon emission computed tomography. *J Auton Nerv Syst* 1995; 50: 339–346.
- Ide K, Boushel R, Sørensen HM, et al. Middle cerebral artery blood velocity during exercise with beta-1 adrenergic and unilateral stellate ganglion blockade in humans. *Acta Physiol Scand* 2000; 170: 33–38.
- Ainslie PN and Brassard P. Why is the neural control of cerebral autoregulation so controversial? *F1000Prime Rep* 2014; 6: 14.
- Yang ZW, Yang SH, Chen L, et al. Comparison of blood counts in venous, fingertip and arterial blood and their measurement variation. *Clin Lab Haematol* 2001; 23: 155–159.
- Kopin IJ, Rundqvist B, Friberg P, et al. Different relationships of spillover to release of norepinephrine in human heart, kidneys, and forearm. *Am J Physiol* 1998; 275: R165–R173.