

## The amphibians and reptiles of Cusuco National Park, Northwest Honduras: updates from a long-term conservation programme

Laking, Alexandra E.; Solís, José M.; Brown, Tom; Maddock, Simon T.; Burdekin, Oliver; Taylor, Peter; Lonsdale, George; Green, Stephen E. W.; Martin, Thomas E.; Galdamez, Josue R.; Kolby, Jonathan E.; Erens, Jesse; Jocque, Merlijn

### Neotropical Biology and Conservation

DOI:  
[10.3897/neotropical.19.e120902](https://doi.org/10.3897/neotropical.19.e120902)

Published: 28/06/2024

Peer reviewed version

[Cyswllt i'r cyhoeddiad / Link to publication](#)

*Dyfyniad o'r fersiwn a gyhoeddwyd / Citation for published version (APA):*  
Laking, A. E., Solís, J. M., Brown, T., Maddock, S. T., Burdekin, O., Taylor, P., Lonsdale, G., Green, S. E. W., Martin, T. E., Galdamez, J. R., Kolby, J. E., Erens, J., & Jocque, M. (2024). The amphibians and reptiles of Cusuco National Park, Northwest Honduras: updates from a long-term conservation programme. *Neotropical Biology and Conservation*, 19(2), 37-62.  
<https://doi.org/10.3897/neotropical.19.e120902>

#### Hawliau Cyffredinol / General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

#### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

1 **Updates from a long-term conservation programme: The amphibians and reptiles of**  
2 **Cusuco National Park, Northwest Honduras**

3  
4 Alexandra E. Laking<sup>1</sup>, José M. Solís<sup>1,2</sup>, Tom Brown<sup>1,3</sup>, Simon T. Maddock<sup>1,4</sup>, Oliver  
5 Burdekin<sup>1,5</sup>, Peter Taylor<sup>1</sup>, George Lonsdale<sup>1</sup>, Stephen E.W. Green<sup>6</sup>, Thomas E. Martin<sup>1,7</sup>, Josue  
6 R. Galdamez<sup>1,3</sup>, Jonathan E. Kolby<sup>1,8,9</sup>, Jesse Erens<sup>1</sup>, Merlijn Jocque<sup>1,10</sup>

7  
8 **1** *Operation Wallacea, Wallace House, Old Bolingbroke, Lincolnshire, UK*

9 **2** *Grupo de Investigación de Reptiles y Anfibios de Honduras (GIRAH), Tegucigalpa,*  
10 *Honduras*

11 **3** *Kanahau Utila Research and Conservation Facility, Utila Island, Honduras*

12 **4** *School of Natural and Environmental Sciences, Newcastle University, Newcastle Upon*  
13 *Tyne, UK*

14 **5** *Oxford University Department for Continuing Education, University of Oxford, UK*

15 **6** *Newquay University Centre, Cornwall College, Wildflower Lane, Trenance Gardens,*  
16 *Newquay, Cornwall, TR7 2LZ, UK*

17 **7** *School of Natural Sciences, College of Environmental Sciences and Engineering, Bangor*  
18 *University, UK*

19 **8** *One Health Research Group, James Cook University, Australia*

20 **9** *Honduras Amphibian Rescue & Conservation Center, Lancetilla Botanical Garden, Tela,*  
21 *Honduras*

22 **10** *Aquatic and Terrestrial Ecology (ATECO), Royal Belgian Institute of Natural Sciences*  
23 *(RBINS), Vautierstraat 29, 1000 Brussels, Belgium*

24  
25  
26 **Abstract**

27 Mesoamerican cloud forests support a rich and unique biodiversity but face severe threats from  
28 increasing habitat degradation and climate change. Here, we present an updated overview of  
29 the amphibians and reptiles of Cusuco National Park (CNP), an isolated cloud forest in the  
30 Sierra de Omoa, Northwest Honduras. Based on surveys conducted over a 17-year period, we  
31 report the presence of 105 confirmed species of amphibians (30) and reptiles (75) within the  
32 reserve. This includes numerous threatened and regionally endemic amphibian species as well  
33 as several reptile species previously unrecorded within the park. Given it harbours  
34 approximately 26% of all recorded Honduran herpetofauna, our study highlights CNP as the  
35 most diverse forest region in Honduras with respect to its reptile and amphibian diversity  
36 documented to date. Our findings reinforce the plea to actively protect CNP as a globally  
37 valuable biodiversity hotspot and a centre of herpetofauna endemism. Furthermore, in the face  
38 of rapid deforestation across Mesoamerica, our findings highlight the need for expanded  
39 biodiversity studies across extant forest regions in Honduras to refine species distribution  
40 ranges and to facilitate timely and effective conservation measures.

41  
42 **Resumen**

43  
44 Los bosques nublados de Mesoamérica soportan una diversidad rica y única, pero por otro lado  
45 sufre de severas amenazas debido a la degradación del hábitat y el cambio climático. En este  
46 manuscrito presentamos un listado general de los anfibios y reptiles del parque Nacional  
47 Cusuco (CNP), un bosque nublado en la sierra de Omoa, noroccidente de Honduras. Basados  
48 en muestreos durante un periodo de 16 años reportamos la presencia de al menos 105 especies  
49 de anfibios (30) y reptiles (75) en la reserva. Dicha herpetofauna incluye numerosas especies  
50 endémicas y amenazadas de anfibios, así como algunos reptiles no registrados previamente en

51 el área. Esto alberga el 24% de toda la herpetofauna conocida para Honduras, nuestro estudio  
52 remarca que CNP es la región forestal en Honduras con mayor diversidad de anfibios y reptiles  
53 con respecto a la diversidad documentada hasta la fecha. Nuestros encuentros refuerzan el  
54 hecho que se debe proteger activamente el CNP como un centro de alto valor global de  
55 biodiversidad y como un núcleo de endemidad de herpetofauna. Además, en vista de la  
56 acelerada deforestación a través de los ecosistemas remanentes en Mesoamérica, nuestros datos  
57 son un llamado a realizar estudios a través de las regiones forestales existentes en Honduras,  
58 para refinar los rangos de distribución de las especies que permitan tomar las medidas efectivas  
59 de conservación.

60

61

62 **Keywords**

63 Biodiversity hotspot, Cloud Forest, Herpetofauna, IUCN status, Mesoamerica, Nuclear  
64 Central America, Population monitoring, Species list


65 **Introduction**

66

67 Positioned centrally in Mesoamerica, Honduras is characterised by an extensive interior  
68 highland area (the Chortis Block) that extends from western Guatemala to Nicaragua (Alvarado  
69 et al. 2007; Townsend 2014). Biodiversity in this region is shaped by a multitude of  
70 environmental gradients and includes a range of isolated mountains topped by evergreen cloud  
71 forests, which can be broadly defined as “tropical forests frequently covered in a cloud of mist”  
72 (Stadtmuller 1987). These high elevation forests exhibit highly specific bioclimatic conditions,  
73 constituting rare and unique ecosystems. Owing to their geographic isolation, Mesoamerican  
74 cloud forests form an array of ‘sky islands’ that provide habitat to a diverse and highly endemic  
75 fauna and flora. The diversity of the region is threatened by the rapid habitat degradation  
76 occurring throughout Mesoamerica’s forests (Brooks et al. 2002; Jung et al. 2022), with high-  
77 elevation ecosystems disproportionately impacted (Bubb et al. 2004). Despite their biological  
78 importance and elevated threat status, most remaining cloud forest habitats in Mesoamerica  
79 remain scientifically under-explored, hampering insights in the distribution of biodiversity  
80 across extended geographic areas (the Wallacean shortfall; Lomolino 2004). A better  
81 understanding of species diversity, abundance, and distributions is imperative to allow timely  
82 and effective implementation of conservation measures.

83

84 Cusuco National Park (CNP), in northwestern Honduras, is an isolated, biologically diverse,  
85 yet threatened cloud forest ecosystem. CNP supports a rich herpetofauna (reptile and  
86 amphibian) community, including many threatened and nationally or regionally endemic  
87 species. The park harbours four micro-endemic amphibian and four micro-endemic reptile  
88 species that are only known to occur at this single locality (Fig. 1). The forest ecosystem of  
89 CNP first gained protection for its value in protecting the watershed of the nearby city of San  
90 Pedro Sula. It is situated within the Mesoamerican biodiversity hotspot (Myers et al. 2000) and  
91 is globally recognised by the Alliance for Zero Extinction (2018) for the critical habitat it  
92 provides to its endemic amphibian fauna. Likewise, CNP was included in a global list of the  
93 most irreplaceable protected areas on the basis of its amphibian, bird and mammal diversity (Le  
94 Saout et al. 2013) and is considered a Key Biodiversity Area by the IUCN. This illustrates the  
95 vital role of extant cloud forest systems in providing ecosystem services to nearby human  
96 communities (Bubb et al. 2004; Bruijnzeel et al. 2010).



97 **Figure 1.** The four amphibian and four reptile micro-endemic species currently known only to  
 98 occur in Cusuco National Park. **A** *Bolitoglossa diaphora* **B** *Oedipina tomasi* **C** *Plectrohyla*  
 99 *dasypus* **D** *Plectrohyla exquisita* **E** *Anolis amplisquamosus* **F** *Geophis nephodrymus* **G**  
 100 *Omoadiphas aurula* **H** *Rhadinella pegosalyta* (Photographs provided by: Tom Brown A,B,E,  
 101 F,G,H, Achyuthan Srikanthan C,D)

102 More than 400 species of reptiles and amphibians have been recorded in Honduras to date, of  
103 which around 27% are endemic to the country (Solís et al. 2014; McCranie 2015). The first  
104 overview of the herpetofauna of CNP was presented by Wilson and McCranie (2004), which  
105 provided a list of 30 identified species based on surveys conducted from the late 1970s to the  
106 early 2000s. These findings prompted the start of a long-term monitoring program to assess the  
107 herpetological diversity of the national park. As a result, the number of recorded species grew  
108 over subsequent years and an updated inventory was published by Townsend et al. (2006),  
109 which increased the number of species identified in CNP to 50. This updated inventory was  
110 comprised of five salamanders, 12 frogs, 12 lizards, and 21 snakes, which were detailed in a  
111 subsequent field guide (Townsend and Wilson 2008). Whereas in the more recent works of  
112 Solís et al. (2017) and Martin et al. (2021) we included tentative overviews of the park's reptiles  
113 and amphibians, these were different in scope and comprise incomplete species accounts.  
114 Hence, we here provide detailed results of the herpetofauna surveys carried out in CNP across  
115 the subsequent 16 years spanning the period 2007–2023 and confirm the presence of 105  
116 species of amphibians and reptiles within the park's boundaries. In addition to providing an  
117 overview of all recorded herpetofauna, we discuss the global conservation significance of the  
118 reserve in view of continuing environmental change.

119

## 120 **Methods**


121

### 122 *Study area - Cusuco National Park and delineation*

123

124 CNP is located in the Sierra de Omoa, the northernmost extension of the Sierra del Merendón  
125 in the region of Cortés, northwestern Honduras (Fig. 2A). Ranging up to 2,243m in elevation,  
126 the park is located among the highest regions of the Sierra de Omoa, while its lower ranges  
127 extend to around 500m above sea level (m/asl) (Martin & Blackburn 2009). Consequently, its  
128 elevational gradient supports a large range of vegetation types. Tropical montane cloud forests  
129 are prevalent around 1,200–2,000m/asl, and are characterised both by broadleaved and pine  
130 vegetation, while elevations above 2,000m/asl are characterised by *bosque enano*, or elfin  
131 'dwarf' forests. Towards lower elevations between 500–1,200m/asl, broadleaved forests are  
132 dominant, with increasing fragmented forest sections and agricultural clearances further down  
133 the mountain slopes (Hamilton, Juvik & Scatena 1995). CNP and its direct environs are covered  
134 by a variety of protected areas (*cf.* Martin et al. 2021). On the basis of the Cloud Forest Act  
135 (1987), the original delineation of CNP comprised a core zone that includes all terrain above  
136 1,800m/asl and a buffer zone that expands 2km outwards from the 1,800m/asl boundary (Fig.  
137 2B) (Bonta 2005). This initial definition of CNP provided the focus of most early herpetofauna  
138 inventories in the reserve (see Wilson and McCranie 2004). Following this original demarcation  
139 however, the Corporación Hondureña de Desarrollo Forestal (COHDEFOR) published an  
140 updated management plan for CNP that substantially expanded the core zone and buffer zone  
141 (Fig. 2B). This latter delineation presents the study area in which all research efforts in the  
142 period 2007–2019 were concentrated, and hence the zonation that is used throughout this report  
143 although this interpretation of the Park's borders is not universally accepted).

144



145  
 146 **Figure 2.** The location of Cusuco National Park (CNP) in northwestern Honduras, the different  
 147 interpretations of its borders, and an overview of study camps. **A** CNP is situated in the northern  
 148 ranges of the Sierra del Merendón, highlighting the isolated position of the cloud forest reserve  
 149 in relation to other high-altitude regions. **B** Our study area as based on the management plan of  
 150 the Corporación Hondureña de Desarrollo Forestal (in blue), shown in reference to the original  
 151 delineation of CNP as based on the 87-1987 Cloud Forest Act (in green). **C** An overview of  
 152 field sites surveyed during the study period 2007-2023. A dashed line indicates the delineation  
 153 of the reserve's core zone. Digital elevation model from Jarvis et al. (2008); basemap from ESRI  
 154 (2017).

155  
 156  
 157  
 158

159 *Data collection*

160

161 Annual surveys were conducted from early-June to early-August each year between 2007 and  
162 2023 (with the exception of 2020–2021 due to the Covid-19 pandemic), as part of an ongoing  
163 long-term biodiversity monitoring programme run by Operation Wallacea, a non-governmental  
164 conservation and research organisation. These annual surveys are carried out by teams of  
165 students and volunteers under the supervision of a rotating team of experienced herpetologists.  
166 All survey activities were concentrated around seven field camps situated in the Park's core-  
167 and buffer zones (Fig. 2C). In each of these seven camps, three to four standardised transects  
168 were monitored across successive years to assess herpetofauna diversity throughout CNP.  
169 Across the research period, transects were studied by means of diurnal visual encounter surveys  
170 in the morning until midday in which the time, distance and number of participants was  
171 recorded to quantify search effort. This involved visually searching for amphibians and reptiles  
172 along the defined transects and when an individual was encountered recording the species, the  
173 distance along the transect and the perpendicular distance from the transect. Encountered  
174 species were caught when possible, and the sex, weight, snout-to-vent length (SVL), and  
175 photographs (dorsal, lateral, ventral and close up of the head) were obtained. Nocturnal  
176 opportunistic searches along portions of these transects or near water bodies were also  
177 conducted, the same information being recorded for these surveys. Within some study seasons,  
178 a single drift fence array with three pitfall traps was opportunistically set up in suitable habitat  
179 at each camp and checked daily to investigate the potential presence of small (semi-)fossorial  
180 species likely to remain undetected by active searches. As this research was conducted by  
181 people of varying levels of experience, only species occurrences with photographic evidence  
182 and or verification by an expert were included. Photographic vouchers of all amphibian and  
183 reptile species recorded in CNP are provided in Suppl. material 1: Fig. S1 and Suppl. material  
184 2: Fig. S2.

185

186 The conservation status of all included species was assessed based on the IUCN Red List  
187 (2022), and the Environmental Vulnerability Score (EVS) following Johnson et al. (2015).  
188 While the former was used as a central representation of species conservation status, the latter  
189 provides an integrative conservation index for all Mesoamerican reptile and amphibian species,  
190 based on their extant geographic distribution, habitat occupation and reproductive mode (for  
191 amphibians) or human persecution (for reptiles). As such, the EVS also provides a valuable  
192 indication of conservation status for species that remain unassessed by the IUCN Red List.

193


194 **Results**

195

196 Following recurring survey efforts in the period 2007–2023, a total of 105 amphibian and reptile  
197 species have been confirmed in CNP: 30 amphibians (Table 1) and 75 reptiles (Table 2). Fig. 3  
198 provides an overview of the cumulative number of species recorded over consecutive field  
199 seasons.

200





201  
 202 **Figure 3.** Species accumulation curves and logarithmic approximation of the reptile and  
 203 amphibian species recorded in Cusuco National Park in the study period 2007–2023 following  
 204 the earlier works of McCranie et al. (2004) and Townsend et al. (2006).  
 205


206  
 207 *Amphibians*

208  
 209 Since the inventory of 2006 (Townsend et al. 2006), an additional 13 amphibian species were  
 210 recorded within the 17-year study period, and of which a provisional overview was provided in  
 211 Solís et al. (2017). Among some of the most notable records, the secretive species *Nototriton*  
 212 *brodiei* Campbell & Smith, 1998 and *Ecnomiohyla salvaje* (Wilson, McCranie & Williams,  
 213 1985) were both found in the high-elevation habitats of the CNP core zone. *N. brodiei* is a  
 214 cryptic salamander that is typically found in dense leaf litter layers and in moss mats of intact  
 215 montane forest (Kolby et al. 2009). The canopy dwelling frog species *E. salvaje*, which is  
 216 endemic to the region, is a cryptic species that is likely facing significant threats. With only  
 217 seven total reported adult individuals in Honduras, five of which were observed in the core zone  
 218 of CNP from 2009–2018 (Solís et al. 2017; Thorp et al. 2021). The park represents a crucial

219 stronghold for this little-known species as forests in Quebrada Grande, Copán, have  
220 disappeared, making the forests of CNP be its main refuge.

221  
222 The other newly recorded amphibians were mostly observed within lower elevation habitats,  
223 largely within the buffer zone of CNP. Additions to the park's salamander diversity stemming  
224 from these lower areas include *Bolitoglossa dofleini* (Werner, 1903) and *Bolitoglossa nympha*  
225 Campbell, Smith, Streicher, Acevedo & Brodie, 2010, which are both endemic to nuclear  
226 Central America. The more widely distributed *Bolitoglossa mexicana* Duméril, Bibron &  
227 Duméril, 1854 was the final salamander species to be registered for CNP, with the first  
228 observation in 2013. Like the latter two salamanders, it seems to thrive in the areas of cattle  
229 pasture and agricultural plantations around Santo Tomas. Important anuran records include the  
230 Critically Endangered Honduran endemic *Craugastor coffeus* (McCranie & Köhler, 1999),  
231 observed around the field site of Santo Tomas (also see Kolby 2009). We furthermore highlight  
232 the presence of the endangered *Craugastor laevisimus* (Werner, 1896) (Suppl. material 1: Fig.  
233 S1, plate 3) within CNP, following evaluation of photographic and bioacoustic observations  
234 made around the field site of Santo Tomas. The toad species *Incilius campbelli* (Mendelson,  
235 1994), endemic to nuclear Central America, was the final anuran reported for CNP, with a single  
236 individual observed close to Santo Tomas in 2013. No additional amphibian species were  
237 recorded in CNP during the subsequent decade (Fig. 3).

238  
239 In addition to the four amphibian species that are endemic to CNP, four species in the park are  
240 endemic to Honduras, and 17 are endemic to nuclear Central America (Table 1). Almost half  
241 of the recorded amphibian species are presently listed as either Critically Endangered (five) or  
242 Endangered (eight), and four species are listed as Vulnerable (IUCN 2022). The threatened  
243 status of the amphibian diversity in CNP is reflected by their respective Environmental  
244 Vulnerability Scores (EVS), with a total of 18 amphibian species present in the park being  
245 attributed a high EVS (14–20), while just five species show a medium EVS (10–13), and six  
246 relatively widespread species show a relatively low EVS (3–9). The threatened status of the  
247 amphibian is also reflected in the location in which they species has been found within the park  
248 (core, buffer, or both) (Fig. 4).



250 **Figure 4.** The IUCN status of amphibian and reptile species recorded in the core and/or buffer  
 251 zone of Cusuco National Park in the period 2007-2023. The delimitation of the core and buffer  
 252 zone is based on the management plan of the Corporación Hondureña de Desarrollo Forestal  
 253 (see Figure 1). IUCN status: Critically Endangered (CR), Endangered (EN), Vulnerable (VU),  
 254 Near-Threatened (NT), Least Concern (LC), and Not Evaluated (NE).

255 **Table 1.** The amphibian fauna of Cusuco National Park (CNP). Species conservation status is based on the assessment criteria of the IUCN Red  
 256 List (IUCN, 2022) and environmental vulnerability scores (EVS) (Johnson et al. 2015). Species are listed alphabetically by family. Geographic  
 257 Distribution is characterised as either widespread (found outside of nuclear Central America), NCA (restricted to localities in nuclear Central  
 258 America), or endemic (restricted to Honduras), while endemic taxa in bold are only known from CNP. \* Indicates species which were formerly  
 259 thought to be endemic to CNP. Conservation Status follows the IUCN Red List (2022): CR, Critically Endangered; EN, endangered; NT, Near  
 260 Threatened; LC, Least Concern; NE, Not Evaluated. Environmental Vulnerability Score (EVS) indicate low (3–9), medium (10–13), or high (14–  
 261 20) vulnerability to environmental degradation. Presence of the species reported within previous studies are indicated by an x.  
 262

Nr.	Taxon	Geographic Distribution	Wilson & McCranie (2004)	Townsend et al. (2006)	CNP zonation	Conservation Status	EVS Score
<b>Order Caudata (Salamanders)</b>							
<b>Family Plethodontidae</b>							
1	<i>Bolitoglossa conanti</i>	NCA	x	x	Core, Buffer	VU	16
2	<i>Bolitoglossa diaphora</i>	<b>Endemic</b>	x	x	Core	EN	18
3	<i>Bolitoglossa dofleini</i>	NCA			Core, Buffer	NT	15
4	<i>Bolitoglossa dunni</i>	NCA	x	x	Core	EN	16
5	<i>Bolitoglossa mexicana</i>	NCA			Buffer	LC	8
6	<i>Bolitoglossa nympa</i>	NCA			Buffer	LC	16
7	<i>Cryptotriton nasalis</i>	NCA *	x	x	Core	EN	18
8	<i>Nototriton brodiei</i>	NCA*			Core	EN	17
9	<i>Oedipina tomasi</i>	<b>Endemic</b>		x	Core	CR	18
<b>Order Anura (Frogs)</b>							
<b>Family Bufonidae</b>							
10	<i>Incilius campbelli</i>	NCA			Buffer	LC	12
11	<i>Incilius valliceps</i>	Widespread	x		Buffer	LC	6
12	<i>Rhinella horribilis</i>	Widespread			Buffer	LC	6
<b>Family Centrolenidae</b>							
13	<i>Hyalinobatrachium fleischmanni</i>	Widespread			Buffer	LC	8
<b>Family Craugastoridae</b>							
14	<i>Craugastor cf. chac</i>	Widespread			Core, Buffer	LC	16
15	<i>Craugastor charadra</i>	NCA		x	Core, Buffer	VU	15
16	<i>Craugastor coffeus</i>	Endemic			Buffer	CR	18

17	<i>Craugastor sp. aff. nefrens</i>				Buffer	NE	
18	<i>Craugastor laevisimus</i>	NCA			Buffer	EN	12
19	<i>Craugastor laticeps</i>	NCA			Core, Buffer	LC	12
20	<i>Craugastor milesi</i>	Endemic	x	x	Core	CR	16
21	<i>Craugastor rostralis</i>	NCA	x	x	Core, Buffer	VU	16
<b>Family Hylidae</b>							
22	<i>Bromeliohyla bromeliacia</i>	NCA	x	x	Core, Buffer	LC	17
23	<i>Bromeliohyla melacaena</i>	Endemic		x	Core, Buffer	EN	20
24	<i>Duellmanohyla soralia</i>	NCA	x	x	Core, Buffer	EN	12
25	<i>Ecnomiohyla salvaje</i>	NCA			Core	EN	19
26	<i>Plectrohyla dasypus</i>	<b>Endemic</b>	x	x	Core, Buffer	CR	14
27	<i>Plectrohyla exquisita</i>	<b>Endemic</b>	x	x	Core, Buffer	CR	15
28	<i>Ptychohyla hypomykter</i>	NCA	x	x	Core, Buffer	VU	10
29	<i>Smilisca baudinii</i>	Widespread		x	Core, Buffer	LC	3
<b>Family Ranidae</b>							
30	<i>Rana maculata</i>	NCA	x		Core, Buffer	LC	5

264 *Reptiles*

265

266 Since the inventory of Townsend et al. in 2006, an additional 46 reptile species were recorded  
267 across the 17-year study period: 18 lizard species and 28 snake species. These records double  
268 the reptile diversity documented in CNP up until 2007 and constitute around 26% of the 264  
269 species known from Honduras (McCranie 2015). Novel records from within the reserve's high-  
270 elevation core zone include the snake species *Leptophis modestus* (Günther, 1872) (Suppl.  
271 material 2: Fig. S2, plate 7), a cloud forest specialist that is endemic to nuclear Central America.  
272 The majority of the newly added reptile species were recorded at relatively low altitudes, with  
273 observations centred around the field sites of Buenos Aires and Santo Tomas. These include  
274 the presence of *Amerotyphlops stadelmani* (Schmidt, 1936), a Honduran endemic blind snake  
275 that is known from just a few individual records and localities. A single individual was found  
276 dead around forest clearings near the intersection of the core- and buffer zone in the proximity  
277 of Santo Tomas. The occurrence of the Honduran endemic lizard, *Laemanctus julioi* McCranie,  
278 2018, was confirmed from a single individual captured near Buenos Aires in 2023. Considering  
279 a recent range extension (Antúnez Fonseca et al. 2021), CNP may be the most north-westerly  
280 record for *L. julioi* in the Atlantic versant of Honduras; an important record for a species that  
281 was previously considered exclusive to the south-central Pacific versant (McCranie 2018).  
282 Among the herpetofauna recorded in CNP, the gecko species *Hemidactylus frenatus* Duméril  
283 & Bibron, 1836 thus far presents the only confirmed alien introduced species.

284

285 Besides the four reptile micro-endemic species only found in CNP, six taxa are Honduran  
286 endemics, and 19 species are restricted to nuclear Central America (Table 2). Of all presently  
287 recorded reptile species, two are listed as Critically Endangered on the IUCN Red List, four are  
288 listed as Endangered, and seven species are listed as Vulnerable (IUCN 2022). However, the  
289 conservation status of four reptile species, mostly lizards, remain unassessed. Of the species  
290 that have been indicated with a threatened category in the IUCN Red List, seven species  
291 similarly show a high EVS (14–20). Among more widespread taxa, 34 species show a medium  
292 EVS (10–13), while 28 species were attributed a relatively low EVS (3–9). The threatened  
293 status of reptiles in CNP is reflected in the location in which the species has been found within  
294 the park (core, buffer, or both) (Fig. 4).

295 **Table 2.** The reptile fauna of Cusuco National Park. Species conservation status is based on the assessment criteria of the IUCN Red List (IUCN,  
 296 2022) and environmental vulnerability scores (EVS) (Johnson et al. 2015). Species are listed alphabetically by family. Geographic Distribution is  
 297 characterised as either widespread (found outside of nuclear Central America), NCA (restricted to localities in nuclear Central America), or endemic  
 298 (restricted to Honduras), while endemic taxa in bold are only known from Cusuco National Park. Conservation Status follows the IUCN Red List  
 299 (2022): CR, Critically Endangered; EN, endangered; NT, Near Threatened; LC, Least Concern; NE, Not Evaluated. Environmental Vulnerability  
 300 Score (EVS) indicate low (3–9), medium (10–13), or high (14–20) vulnerability to environmental degradation. Presence of the species reported  
 301 within previous studies are indicated by an x.  
 302

Nr.	Taxon	Geographic Distribution	Wilson & McCranie (2004)	Townsend et al. (2006)	CNP zonation	Conservation Status	EVS Score
<b>Order Squamata (Lizards)</b>							
<b>Family Anguidae</b>							
1	<i>Abronia moreletii</i>	NCA	x	x	Core, Buffer	LC	13
<b>Family Corytophanidae</b>							
2	<i>Basiliscus vittatus</i>	NCA			Buffer	LC	7
3	<i>Corytophanes cristatus</i>	Widespread			Buffer	LC	11
4	<i>Corytophanes hernandesii</i>	Widespread			Buffer	LC	13
5	<i>Laemantus julioi</i>	Endemic			Buffer	NE	NA
6	<i>Laemantus longipes</i>	NCA			Buffer	LC	10
<b>Family Dactyloidae</b>							
7	<i>Anolis amplisquamosus</i>	<b>Endemic</b>	x	x	Core, Buffer	CR	14
8	<i>Anolis biporcatus</i>	Widespread			Buffer	LC	8
9	<i>Anolis capito</i>	Widespread		x	Core, Buffer	LC	9
10	<i>Anolis cusuco</i>	Endemic	x	x	Core, Buffer	CR	12
11	<i>Anolis johnmeyeri</i>	Endemic	x	x	Core, Buffer	EN	12
12	<i>Anolis lemurinus</i>	Widespread			Buffer	LC	7
13	<i>Anolis mccraniei</i>	NCA			Core, Buffer	NE	NA
14	<i>Anolis ocelloscapularis</i>	Endemic		x	Core, Buffer	VU	11
15	<i>Anolis petersii</i>	NCA		x	Core, Buffer	NT	9
16	<i>Anolis rodriguezii</i>	NCA			Buffer	LC	8
17	<i>Anolis uniformis</i>	Widespread			Core, Buffer	LC	13
18	<i>Anolis unilobatus</i>	Widespread			Buffer	LC	NA

19	<i>Anolis yoroensis</i>	Endemic			Core	EN	11
<b>Family Dipoglossidae</b>							
20	<i>Siderolamprus montanus</i>	Endemic		x	Core, Buffer	EN	13
<b>Family Gekkonidae</b>							
21	<i>Coleonyx mitratus</i>	Widespread			Buffer	LC	14
22	<i>Hemidactylus frenatus</i>	Introduced			Core, Buffer	LC	NA
<b>Family Phrynosomatidae</b>							
23	<i>Sceloporus schmidti</i>	NCA	x	x	Core, Buffer	LC	11
24	<i>Sceloporus variabilis</i>	Widespread		x	Core, Buffer	LC	5
<b>Family Sphaenomorphidae</b>							
25	<i>Scincella cherriei</i>	Widespread	x		Core, Buffer	LC	4
26	<i>Scincella incerta</i>	NCA		x	Core, Buffer	LC	12
<b>Family Sphaerodactylidae</b>							
27	<i>Sphaerodactylus continentalis</i>	NCA			Buffer	NE	6
<b>Family Teiidae</b>							
28	<i>Holcosus festivus</i>	Widespread			Buffer	LC	10
<b>Family Xantusiidae</b>							
29	<i>Lepidophyma flavimaculatum</i>	NCA			Core, Buffer	LC	9
<b>Order Squamata (Snakes)</b>							
<b>Family Colubridae</b>							
30	<i>Dendrophidion rufiterminorum</i>	Widespread			Buffer	NE	12
31	<i>Drymarchon melanurus</i>	Widespread	x		Core, Buffer	LC	9
32	<i>Drymobius chloroticus</i>	Widespread	x	x	Core, Buffer	LC	11
33	<i>Drymobius margaritiferus</i>	Widespread			Core, Buffer	LC	9
34	<i>Lampropeltis abnorma</i>	Widespread		x	Core, Buffer	LC	9
35	<i>Leptophis modestus</i>	NCA			Core	VU	14
36	<i>Leptophis praestans</i>	Widespread	x	x	Core, Buffer	LC	9
37	<i>Mastigodryas dorsalis</i>	NCA	x	x	Core, Buffer	LC	12
38	<i>Mastigodryas melanolomus</i>	Widespread			Buffer	LC	9
39	<i>Oxybelis koheleri</i>	Widespread			Buffer	LC	9
40	<i>Phrynonax poecilonotus</i>	Widespread		x	Core, Buffer	LC	11



41	<i>Scolecophis atrocinctus</i>	Widespread			Core, Buffer	LC	12
42	<i>Senticolis triaspis</i>	Widespread			Buffer	LC	7
43	<i>Spilotes pullatus</i>	Widespread			Buffer	LC	9
44	<i>Stenorrhina degenhardtii</i>	Widespread	x		Core, Buffer	LC	9
45	<i>Stenorrhina freminvillei</i>	Widespread			Buffer	LC	NA
46	<i>Tantilla schistosa</i>	Widespread	x	x	Core	LC	9
47	<i>Tantillita lintoni</i>	NCA			Buffer	LC	12
<b>Family Dipsadidae</b>							
48	<i>Adelphicos quadrivirgatum</i>	Widespread		x	Core, Buffer	LC	6
49	<i>Amastridium sapperi</i>	Widespread			Buffer	LC	12
50	<i>Coniophanes imperialis</i>	Widespread			Core, Buffer	LC	10
51	<i>Geophis nephodrymus</i>	<b>Endemic</b>		x	Core	VU	15
52	<i>Geophis sartorii</i>	NCA			Core, Buffer	LC	12
53	<i>Hydromorphus concolor</i>	Widespread			Buffer	LC	12
54	<i>Imantodes cenchoa</i>	Widespread	x	x	Core, Buffer	LC	6
55	<i>Leptodeira septentrionalis</i>	Widespread			Core, Buffer	LC	9
56	<i>Ninia diademata</i>	Widespread			Core, Buffer	LC	7
57	<i>Ninia espinali</i>	NCA	x		Core	NT	11
58	<i>Ninia sebae</i>	Widespread			Core, Buffer	LC	5
59	<i>Omoadiphas aurula</i>	<b>Endemic</b>		x	Core, Buffer	VU	14
60	<i>Pliocercus elapoides</i>	Widespread			Core, Buffer	LC	9
61	<i>Rhadinella kinkelini</i>	NCA			Core, Buffer	LC	10
62	<i>Rhadinella montecristi</i>	NCA		x	Core	VU	12
63	<i>Rhadinella pegosalyta</i>	<b>Endemic</b>		x	Core	VU	15
64	<i>Sibon dimidiatus</i>	Widespread			Core	LC	10
65	<i>Sibon nebulatus</i>	Widespread			Core, Buffer	LC	5
66	<i>Xenodon rabdocephalus</i>	Widespread			Buffer	LC	11
<b>Family Elapidae</b>							
67	<i>Micrurus apiatus</i>	Widespread	x	x	Core, Buffer	LC	13
68	<i>Micrurus nigrocinctus</i>	Widespread			Core, Buffer	LC	9

<b>Family Sibynophiidae</b>							
69	<i>Scaphiodontophis annulatus</i>	Widespread		x	Core, Buffer	LC	10
<b>Family Typhlopidae</b>							
70	<i>Amerotyphlops stadelmani</i>	Endemic			Core	VU	11
<b>Family Viperidae</b>							
71	<i>Metlapilcoatlus mexicanus</i>	NCA		x	Core, Buffer	LC	11
72	<i>Bothriechis marchi</i>	Endemic	x	x	Core, Buffer	EN	15
73	<i>Bothriechis schlegelii</i>	Widespread			Buffer	LC	12
74	<i>Bothrops asper</i>	Widespread		x	Core, Buffer	LC	12
75	<i>Cerrophidion wilsoni</i>	NCA	x	x	Core, Buffer	LC	12

304 *Taxonomic changes, notes and decisions*

305

306 Since the last overview of amphibians and reptiles in CNP was published by Townsend et al.  
307 (2006), the Mesoamerican herpetofauna has been subject to numerous taxonomic discoveries  
308 and revisions. We provide here a brief outline of the taxonomic decisions made in our species  
309 list in view of recent changes in nomenclature and ongoing taxonomic debate, as well as recent  
310 molecular studies and range extensions.

311

312 For the amphibians, this includes several generic revisions, with the placement of the former  
313 *Isthmohyla melacaena* in the genus *Bromeliohyla* (Faivovich et al. 2018), as well as the revision  
314 of *Rana maculata* following paraphyletic relationships recovered in the genus *Lithobates* (Yuan  
315 et al. 2016). Among the amphibian species present in CNP, members of the genus *Craugastor*  
316 comprise significant cryptic diversity. A recent molecular assessment was therefore performed  
317 to characterise the species diversity within this genus, in which focus was put on a combined  
318 phylogenetic and morphological analyses of the *C. laticeps*-like species group (unpublished  
319 data). This indicated the presence of four disparate lineages, corresponding to earlier records of  
320 *C. rostralis*, *C. chac*, *C. laticeps* and *C. charadra*, in addition to the more readily distinguishable  
321 *C. milesi*, *C. laevissimus* and *C. coffeus*. However, as McCranie (2018) indicated, the presence  
322 of nominal *C. chac* in Northwest Honduras might be restricted to lower elevations, and hence  
323 we provisionally term this species lineage *C. cf. chac* warranting further studies on its status.  
324 An ongoing molecular assessment highlighted a disparate lineage from the *Craugastor*  
325 *campbelli* complex in the park, provisionally termed *C. aff. nefrens* and awaiting further studies  
326 on its status (M. Jocque, pers. comm.). Additionally, *Cryptotriton nasalis* was previously  
327 considered a CNP and Honduran endemic, but after having recently been discovered just across  
328 the border in Guatemala (McCranie and Rovito, 2014), is now designated as being endemic to  
329 nuclear Central America. However, similar to *Nototriton brodiei*, it likely resides in a highly  
330 confined distribution range and is therefore still of particular conservation concern and is still  
331 listed as Endangered by the IUCN.

332

333 For the reptiles, taxonomic changes include the recognition of the species *Anolis mccranie*,  
334 following its subdivision from the *Anolis tropidonotus* species complex (Köhler et al. 2016).  
335 We furthermore include *Diploglossus montanus* following paraphyletic relationships recovered  
336 with respect to the genus *Celestus* (Pyron et al. 2013) and recognize *Abronia moreletii* as  
337 synonymous with the genus *Mesaspis* (Gutiérrez-Rodríguez et al. 2020). *Sceloporus schmidtii*  
338 was included in our list as a valid species in northwest Honduras in place of *S. malachiticus*  
339 following McCranie et al. (2015), while the designation of *Scincella cherriei* and *S. incerta*  
340 follow generic revisions of the genus *Sphenomorphus* (see Linkem et al. 2011). *Lampropeltis*  
341 *abnorma* was included in our list after its split from *L. triangulum* (Ruane et al. 2014), while  
342 *Metlapilcoatlus mexicanus* was revised after polyphyletic relationships within the genus  
343 *Atropoides* (Campbell et al. 2019). *Geophis sartorii* is included following the revision of  
344 *Tropidodipsas* (Grünwald et al. 2021). *Siderolamprus montanus* is now included following  
345 generic revisions of *Diploglossus* (Schools and Hedges 2021). In addition, *Leptophis praestans*  
346 was included following taxonomic revisions in the *L. ahaetulla* complex (Albuquerque and  
347 Fernandez 2022), and *Micrurus apiatus* was included following revisions in the *M. diastema*  
348 complex (Reyes-Velasco et al. 2020). Comparable to the amphibian genus *Craugastor* as  
349 described above, members of the reptile genus *Anolis* include various cryptic taxa and species  
350 complexes, making field identification of anoles highly challenging. We therefore first  
351 referenced our observations to the comprehensive overview of known species localities  
352 provided by McCranie and Köhler (2015), which corroborated the presence of 12 species in the  
353 reserve's core and buffer zone. Initial genetic findings in combination with field identifications

354 also confirmed the presence of *A. uniformis* in the park (Suppl. material 2: Fig. S2, plate 4), of  
355 which the closest known localities were noted to occur around El Paraíso, northeast of CNP  
356 (McCranie and Köhler 2015). However, an ongoing DNA barcoding study has highlighted that  
357 five more species might be present in the park, including potential candidate species (O'Brien  
358 et al., unpublished). Hence, the current species list presents the most conservative estimate of  
359 anole diversity in the park. Note that here we adhere to the genus name *Anolis* as opposed to  
360 *Norops*, following recent controversy about this generic revision coined for the clade composed  
361 of “beta anoles” (see Poe 2013) and in which all species in CNP are classified.

362  
363

## 364 **Discussion**

365

366 More than 400 species of amphibians and reptiles are currently reported to occur in Honduras  
367 (Solís et al. 2014; McCranie 2015). Within the core- and buffer zones of CNP at least 105  
368 species have been recorded thus far, amounting to around 26% of recorded Honduran  
369 herpetofauna diversity. This highlights CNP as an exceptionally diverse national park in  
370 Honduras, and as a hotspot of Mesoamerican herpetofauna diversity and endemism. Its  
371 location at the northern edge of the biogeographically isolated Sierra del Merendón partially  
372 explains the presence of the relatively high number of micro-endemics (four amphibians and  
373 four reptiles). These observations are echoed by inventories of other species communities in  
374 CNP where, for instance, birds (Martin et al. 2016), bats (Medina-van Berkum et al. 2020) as  
375 well as non-volant mammals (Hoskins et al. 2018) were also found to exhibit remarkable  
376 species diversity across its extensive elevational range. However, it is important to note that the  
377 park has received an exceptional amount of research effort over more than 15 years compared  
378 to other Honduran cloud forests. There are several other protected zones in Honduras that  
379 exceed CNP in both land area and habitat diversity and show a similar elevational range (CNP  
380 reaching 2,243m/asl). For instance, Pico Bonito National Park covers an area of 565 km<sup>2</sup> and  
381 ranges up to an elevation of 2,480m/asl. Across a limited number of surveys, already 82 species  
382 of amphibians and reptiles were recorded in Pico Bonito National Park (McCranie and Solís  
383 2013). Another example is the protected yet highly imperilled area of Texiguat, where 39  
384 species of amphibians and reptiles have thus far been observed across a similar elevational  
385 gradient (Townsend et al. 2010). Hence, biodiversity assessments are urgently warranted in  
386 other extant cloud forest ecosystems in order to refine species distribution patterns across  
387 Honduras and the wider Mesoamerican biodiversity hotspot.

388

389 Early herpetofauna surveys in CNP were largely focused on the eastern sections of the present-  
390 day core zone (Wilson and McCranie 2004; Townsend et al. 2006). Since then, the monitoring  
391 focus was expanded to cover both the west side of the mountain as well as lower elevations,  
392 including transects situated in the warmer and drier habitats in the designated buffer zone of  
393 CNP (see Fig. 2C). Many additions to our updated species list, and in particular many reptile  
394 species, are a result of extended monitoring efforts in these previously underexplored habitats.  
395 This is reflected by a surge in new species records in the period 2007–2008 following the start  
396 of research activities in these field sites (Fig. 3). During the course of the study period, the rate  
397 of reptile and amphibian species detections slowed down considerably (Fig. 3), and we expect  
398 the large majority of the amphibian diversity inhabiting CNP to now be recorded. However,  
399 additional reptile species continued to be observed in recent field seasons in the period 2015–  
400 2023. In addition to observations of relatively thermophilic species at lower elevations, we  
401 expect that novel records in the reserve's core zone might still arise from lesser studied  
402 microhabitats such as the forest canopy and in the form of cryptic leaf-litter dwelling or (semi)  
403 fossorial species. Further molecular studies may furthermore highlight additional taxonomic

404 diversity among the various species groups, which is likely to increase the known species  
405 diversity within the park. Thus, although we included 105 species in our updated species list,  
406 this figure is likely a conservative estimate.

407  
408 While the herpetofauna diversity in CNP is unquestionably high, several species have been  
409 sighted only once during the study period, and records of several others are sporadically  
410 dispersed across many years. This pattern is particularly evident in snake species, which  
411 compose c. 43% of the known herpetofauna in CNP. The resulting variability highlights the  
412 challenge in accurately determining seasonal species occurrence, because detection can be  
413 subject to various environmental constraints. Consequently, assessing the true abundance, loss  
414 or replacement rate of species in CNP becomes a considerable conservation challenge.

415  
416 It is worth noting that while the accumulation of amphibian diversity plateaued in 2013, the  
417 known reptile diversity has continued to grow throughout the last decade. Furthermore, there  
418 remains potential additional records for several other snake species in the long-term database,  
419 such as *Oxyrhopus petolarius* (Linnaeus, 1758), *Ninia pavimentata* (Bocourt, 1883), and  
420 *Rhadinella anachoreta* (Smith & Campbell, 1994). However, for the present study, we were  
421 unable to confirm them with certainty due to lack of sufficient evidence, such as being identified  
422 by an expert, or unambiguous photographic evidence. Additionally, *Xenodon rabdocephalus*  
423 (Wied-Neuwied, 1824) is likely present in CNP, as indicated by photographs of two specimens  
424 (one roadkill) taken near the core and buffer zone in 2023 (Brown TW pers. obs), albeit captured  
425 by field guide outside the annual survey season.

426  
427 As a more comprehensive picture of the amphibian and reptile diversity in CNP is starting to  
428 take shape, continued monitoring efforts are becoming essential to assess the response of its  
429 herpetofauna community to the combined effects of ongoing environmental change. With the  
430 solidification of a long-term dataset, future analyses can start shifting their focus to changes in  
431 relative species abundance and species distributions within the reserve. This is especially  
432 prudent given the notion that cloud forests are highly vulnerable to climatic change, with  
433 shifting temperature and precipitation regimes causing upslope elevational shifts in vegetation  
434 patterns and animal distribution ranges (Foster 2001; Laurance et al. 2011). As many  
435 specialised cloud forests species inhabit narrow microclimatic niches and, thus, highly specific  
436 elevational ranges, these changes have the potential to cause near-term extinctions (the  
437 “escalator to extinction”, e.g., Freeman et al. 2018) and a shift in community composition. Such  
438 patterns have already started to form in CNP’s bird community (Neate-Clegg et al. 2018). As  
439 ectothermic vertebrate communities include relatively more high-elevation specialists  
440 (Laurance et al. 2011), these effects might be even more pronounced among herpetofauna.

441  
442 Habitat destruction has been accelerating at an alarming pace and threatens all wildlife and  
443 habitats in CNP. Being a relatively small reserve with a core zone of 7,690 ha which is largely  
444 isolated from other high-elevation habitats, this region is highly sensitive to disturbances.  
445 Despite its protected status as a national park, CNP has no permanent forest guards, and a  
446 growing population around the mountain together with a challenging economic situation over  
447 the past decade increases pressure on the remaining ecosystem. Deforestation for coffee  
448 plantations and livestock led to 7% of the park’s forested area being lost between 2000 and  
449 2017 (Hoskins 2019). This deforestation is increasingly encroaching within the reserve’s core  
450 zone, posing an immediate threat to this vulnerable cloud forest ecosystem and its endemic  
451 species. As such, the combined effects of habitat degradation and climate change are likely to  
452 strongly exacerbate biodiversity loss within a short timeframe (Ponce-Reyes et al. 2012). The  
453 reserve’s buffer zone accommodates limited settlement and licensed farming practices and has

454 little primary habitat left. Nevertheless, our findings across transects in the buffer zone indicate  
455 that the existing mosaic of disturbed forest fragments and anthropogenic landscape features still  
456 supports a wide variety of reptiles and amphibians. Besides the paramount importance of  
457 safeguarding the remaining pristine cloud forest in the core zone, sustained efforts are therefore  
458 also needed to preserve extant forest patches in the buffer zone. The protection of cloud forests  
459 such as CNP thus becomes increasingly dependent on successful community-based  
460 conservation schemes that incorporate sustainable socio-economic benefits to nearby  
461 livelihoods (Hostettler 2002; Bubb et al. 2004). For instance, payment for ecosystem services  
462 (PES) programs might aid in the participatory protection of cloud forests. Nevertheless, this  
463 only becomes possible once the environmental value of cloud forests becomes adequately  
464 recognised and exceed the short-term gains of habitat conversion (Martínez et al. 2009).

465  
466 The persistence of many amphibian species in CNP is particularly jeopardised. Of the 13  
467 species in the reserve that are listed as Critically Endangered or Endangered by the IUCN Red  
468 List, 11 are stated to be in continued decline, and in the other two the population trend is  
469 unknown. In addition to other environmental stressors, the amphibian diversity in CNP is  
470 imperilled by the spread of emerging infectious disease, most notably by the amphibian chytrid  
471 fungus *Batrachochytrium dendrobatidis* Longcore, Pessier & Nichols, 1999 also known as *Bd*.  
472 This pathogen causes the lethal disease chytridiomycosis, which has been linked to global  
473 amphibian population declines and extinctions and has had especially devastating effects on  
474 Neotropical amphibian communities (Scheele et al. 2019). Reports of declines of Critically  
475 Endangered amphibian species in CNP prompted an investigation in 2007, revealing  
476 widespread *Bd* presence in the park with a high prevalence of infection in Endangered and  
477 Critically Endangered species, such as the endemic *Plectrohyla dasypus* McCranie & Wilson,  
478 1981 and *Plectrohyla exquisita* McCranie & Wilson, 1998 (Kolby et al. 2010). An array of *Bd*  
479 dispersal mechanisms were identified in CNP, including detection not only in stream water and  
480 bromeliad reservoirs, but also potential aerial dispersion by rainwater, waterfall spray, and in  
481 residues on leaves resulting from contact with amphibian skin (Kolby et al. 2015 a,b). Stream-  
482 associated amphibians in CNP were found to be five times more susceptible to *Bd* infection  
483 than bromeliad-dependent species (Bloom et al. 2018). Yet, even strictly canopy dwelling  
484 species can be infected; for example, *Bd* was detected in 100% (4) of the sampled *Ecnomiohyla*  
485 *salvaje* (Wilson, McCranie, and Williams, 1985) during 2017–2018 (Thorp et al. 2021). Further  
486 investigation and continued disease monitoring in CNP is therefore crucial to detect long-term  
487 changes in host-pathogen dynamics. The recently established Honduras Amphibian Rescue and  
488 Conservation Centre (HARCC) is actively researching methods to protect vulnerable species  
489 within habitats where this pathogen is present. On the upside, *Craugastor milesi* (Schmidt,  
490 1933), a species previously listed as Extinct and now listed as Critically Endangered, was  
491 rediscovered in CNP in 2008 after last being seen in the 1980s when it was still locally abundant  
492 (see Kolby and McCranie 2009), thus providing prospects for its persistence in the park. A  
493 second observation of *C. milesi* (Kolby, Brown, Solís pers.obs) was registered in 2013, however  
494 subsequent monitoring efforts have failed to detect the species again, and it is possible the  
495 species might have disappeared from CNP in recent years (Solís et al. 2017). Its continued  
496 presence in the park thus remains to be reaffirmed. If the species has indeed disappeared from  
497 CNP, this would constitute the first local extinction among all previously documented  
498 herpetofauna diversity (Wilson and McCranie 2004; Townsend et al. 2006; the present study).

499  
500 Less information is available on the conservation status of reptile species present in CNP, with  
501 several remaining to be evaluated by the IUCN. However, an extended conservation assessment  
502 by García-Padilla et al. (2020) recently highlighted 11 amphibian and 12 reptile species in CNP  
503 as being among the most threatened ‘priority level’ species in Mesoamerica based on their

504 combined endemic distribution range and high EVS scores. Their present IUCN status is  
505 therefore not an indication of lower conservation status than the amphibians, but rather indicates  
506 the absence of data sufficient to designate a particular level of concern, and thus many species  
507 urgently warrant further research and conservation attention. Furthermore, it can be expected  
508 that amphibian declines have cascading effects on reptile diversity, with reptile declines  
509 following the decline and extirpation of their amphibian prey (Zipkin et al. 2020). All the CNP  
510 endemic reptiles have been assessed by the IUCN Red List, including *Anolis amplisquamosus*  
511 (McCranie, Wilson & Williams, 1993), which is listed as Critically Endangered and the three  
512 endemic snake species *Geophis nephodrymus* Townsend & Wilson, 2006, *Omoadiphas aurula*  
513 Köhler, McCranie & Wilson, 2001, and *Rhadinella pegosalyta* (McCranie, 2006), are presently  
514 considered Vulnerable.

515  
516 Much of the herpetofauna diversity in Mesoamerica likely remains undescribed, with high rates  
517 of species discoveries taking place in concert with high rates of habitat destruction (Johnson et  
518 al. 2015; García-Padilla et al. 2020). Besides the Wallacean shortfall, uncertainty about the  
519 actual species diversity existing in areas (the Linnean shortfall) is, therefore, another factor  
520 hindering conservation efforts in biodiversity hotspots (Whittaker et al. 2005). Additional field  
521 surveys are therefore not only necessary to fill crucial knowledge gaps in species occurrences,  
522 but also to gather the molecular, morphological and ecological data necessary to disentangle  
523 undescribed diversity. DNA barcoding studies based on a standardized set of loci could in this  
524 way provide an increasingly affordable and scalable means to facilitate species delimitation in  
525 understudied tropical assemblages (Vences et al. 2005; Menegon et al. 2017). In CNP, an  
526 assessment of the cryptic species diversity within *Anolis* is presently underway, and further  
527 studies are assessing the status of several candidate species. Moreover, by gathering  
528 comparative genetic data across time, it could be evaluated to what extent populations retain  
529 their genetic diversity and connectivity within and between increasingly fragmented and  
530 imperilled forest regions (e.g. Dixo et al. 2009). While the present paper is directed at providing  
531 a renewed overview of herpetofauna species richness in CNP, such complementary studies in  
532 the park, the broader Sierra de Omoa and other forest regions across Honduras would therefore  
533 strongly benefit our understanding of regional biodiversity patterns and dynamics.

534  
535

## 536 **Conclusion**

537

538 CNP is an irreplaceable hotspot of Mesoamerican biodiversity, providing habitat to a  
539 remarkable 105 species of reptiles and amphibians, including a high number of local, national  
540 and regional endemics. Ongoing threats to CNP, in particular its high deforestation rates, place  
541 strong conservation urgency on this unique ecosystem and its biodiversity. A detailed  
542 understanding of the importance of CNP's herpetofauna community has only been possible due  
543 to our long-running monitoring program. Increased surveys in other extant forest regions are  
544 thus necessary to provide essential baseline biodiversity data and inform the timely and targeted  
545 conservation efforts necessary to safeguard the future of Mesoamerica's irreplaceable  
546 herpetofauna diversity.

547

548

## 549 **Acknowledgements**

550

551 This work is dedicated to Don Macrín Ramírez, a guardian of CNP for more than half a century  
552 and a personal inspiration to many who had the honour of meeting him whilst visiting his  
553 extraordinary forest home. During the course of his life, he was fortunate to marvel at much of

554 the forest's biodiversity in person. We hope and aim that his legacy be kept alive. We would  
555 furthermore like to express our sincere gratitude to all staff members, research associates,  
556 students and park guides that contributed to the collection of amphibian and reptile observations  
557 in CNP over the years. We also thank Instituto Nacional de Conservación Forestal (ICF) for  
558 providing annual research permits for biodiversity monitoring in CNP, and Operation Wallacea  
559 and Expediciones y Servicios Ambientales Cusuco for providing logistical support for all  
560 survey work completed in the park.

561

562

## 563 **References**

564

565 Alliance for Zero Extinction (2018) Global AZE map: <https://zeroextinction.org/>

566 Albuquerque NR De, Fernandes DS (2022) Taxonomic revision of the parrot snake *Leptophis*  
567 *ahaetulla* (Serpentes, Colubridae). *Zootaxa* 5153: 001–069.

568 Alvarado GE, Dengo C, Martens U, Bundschuh J, Aguilar T, Bonis B (2007) Stratigraphy and  
569 geologic history. In: J. Bundschuh, G. Avarado (Eds) *Central America*. Taylor and Francis,  
570 Leiden, 345-394.

571 Antúnez Fonseca C, Zúniga L, Padilla Raudales, D, Vega, H, Funes, C (2021) Extended range  
572 and observations on the natural history of the casquehead lizard *Laemanctus julioi* from  
573 Honduras. *Herpetological Bulletin* 156, 18–22.

574 Blooi M, Laking AE, Martel A, Haesebrouck F, Jocque M, Brown T, Green S, Vences M, Bletz  
575 MC, Pasmans F (2017) Host niche may determine disease-driven extinction risk. *PLOS ONE*  
576 12(7): e0181051 <https://doi.org/10.1371/journal.pone.0181051>

577 Bonta M (2005) Becoming Forest, becoming local: transformations of a protected area in  
578 Honduras. *Geoforum* 36, 95–112. <https://doi.org/10.1016/j.geoforum.2004.03.011>

579 Brooks TM, Mittermeier RA, Mittermeier CG, da Fonseca GAB, Rylands AB, Konstant WR,  
580 Flick P, Pilgrim J, Oldfield S, Magin G, Hilton-Taylor C (2002) Habitat loss and extinction  
581 in the hotspots of biodiversity. *Conservation Biology* 16: 909-923.  
582 <https://doi.org/10.1046/j.1523-1739.2002.00530.x>

583 Bubb P, May I, Miles L, Sayer J (2004) *Cloud Forest Agenda*, UNEP-World Conservation  
584 Monitoring Centre: Cambridge, UK.

585 Bruijnzeel LA, Kappelle M, Muligan M, Scatena FN (2010) Tropical montane cloud forests:  
586 state of knowledge and sustainability perspectives in a changing world. In: Bruijnzeel LA,  
587 Scatena FN, Hamilton LS (Eds) *Tropical montane cloud forests: science for conservation*  
588 and management. Cambridge University Press, Cambridge,

589 Campbell JA, Frost DR, Castoe TA (2019) New generic name for jumping pitvipers (Serpentes:  
590 viperidae). *Revista Latinoamericana de Herpetología* 2: 52-53.  
591 <https://doi.org/10.22201/fc.25942158e.2019.2.73>

592 Dixo M, Metzger JP, Morgante JS, Zamudio KR (2009) Habitat fragmentation reduces genetic  
593 diversity and connectivity among toad populations in the Brazilian Atlantic Coastal Forest.  
594 *Biological Conservation* 142: 1560-1569. <https://doi.org/10.1016/j.biocon.2008.11.016>

595 Esri. "Light Gray Canvas Base" [basemap]. Scale Not Given. (2017).  
596 [https://basemaps.arcgis.com/arcgis/rest/services/World\\_Basemap\\_v2/VectorTileServer](https://basemaps.arcgis.com/arcgis/rest/services/World_Basemap_v2/VectorTileServer).  
597 Accessed: 2023-12-19.

598 Faivovich J, Pereyra MO, Celeste Luna M, Hertz A, Blotto BL, Vásquez-Almazán CR,  
599 McCranie JR, Sánchez DA, Baêta D, Araujo- Vieira K, Köhler G, Kubicki B, Campbell JA,  
600 Frost DR, Wheeler WC, Haddad CFB (2018) On the Monophyly and Relationships of  
601 Several Genera of Hylini (Anura: Hylidae: Hylinae), with Comments on Recent Taxonomic  
602 Changes in Hylids. *South American Journal of Herpetology* 13: 1-32.  
603 <https://doi.org/10.2994/SAJH-D-17-00115.1>



604 Freeman BG, Scholer MN, Ruiz-Guitierrez V, Fitzpatrick JW (2018) Climate change causes  
605 upslope shifts and mountaintop extirpations in a tropical bird community. *Proceedings of the*  
606 *National Academy of Sciences* 115: 11982-11987.  
607 <https://doi.org/10.1073/pnas.1804224115>

608 García-Padilla E, DeSantis DL, Rocha A, Mata-Silva V, Johnson JD, Wilson LD (2020)  
609 Conserving the Mesoamerican herpetofauna: the most critical case of the priority level one  
610 endemic species. *Amphibian & Reptile Conservation* 14: 73-132.

611 Gilroy D, Jones S, Vulinec K, Thompson P, Hoskins H, Creedy T, Jocque M, Green S, Lonsdale  
612 G, Phipps C, Burdekin O, Brown T (2017) Operation Wallacea Cusuco National Park,  
613 Honduras 2016 & 2017: End of Season Report. Operation Wallacea: Old Bolingbroke, UK.

614 Grünwald CI, Toribio-Jiménez S, Montaña-Ruvalcaba C, Franz-Chávez H, Peñaloza-Montaña  
615 MA, Barrera-Nava EY, Jones JM, Rodriguez CM, Hughes IM, Strickland JL, Reyes-Velasco  
616 J (2021) Two new species of snail-eating snakes of the genus *Tropidodipsas* (Serpentes,  
617 *Dipsadidae*) from southern Mexico, with notes on related species. *Herpetozoa* 34: 233-257.  
618 <https://doi.org/10.3897/herpetozoa.34.e69176>

619 Gutiérrez-Rodríguez J, Zaldívar-Riverón A, Solano-Zavaleta I, Campbell JA, Meza-Lázaro  
620 RN, Flores-Villela O, Nieto-Montes de Oca A (2020). Phylogenomics of the Mesoamerican  
621 Alligator-Lizard Genera *Abronia* and *Mesaspis* (Anguillidae: Gerrhonotinae) Reveals Multiple  
622 Independent Clades of Arboreal and Terrestrial Species. *Molecular Phylogenetics and*  
623 *Evolution* 154: 106963. <https://doi.org/10.1016/j.ympev.2020.106963>

624 Hamilton LS, Juvik JO, Scatena FN (Eds.) (1995) *Tropical Montane Cloud Forests*. Springer,  
625 New York.

626 Hoskins HMJ, Burdekin OJ, Dicks K, Slater KY, McCann NP, Jocque M, Castañeda F, Reid N  
627 (2018) Non-volant mammal inventory of Cusuco National Park, north-west Honduras:  
628 reporting the presence of Jaguar, *Panthera onca* (Linnaeus, 1758), and demonstrating the  
629 effects of zonal protection on mammalian abundance. *Check List* 14: 877-891.  
630 <https://doi.org/10.15560/14.5.877>

631 Hostettler S (2002) *Tropical montane cloud forests: a challenge for conservation*. *Bois forêts*  
632 *Tropiques* 274:19–31.

633 Hoskins, H.M.J. (2019). *Assessing conservation status of remnant mammal populations in a*  
634 *Neotropical cloud forest*. PhD Dissertation: Queens University Belfast

635 IUCN (2022) *The IUCN Red List of Threatened Species*. <https://www.iucnredlist.org>

636 Foster P (2001) The potential negative impacts of global climate change on tropical montane  
637 cloud forests. *Earth-Science Reviews* 55: 73-106. [https://doi.org/10.1016/S0012-](https://doi.org/10.1016/S0012-8252(01)00056-3)  
638 [8252\(01\)00056-3](https://doi.org/10.1016/S0012-8252(01)00056-3)

639 Jadin RC, Townsend JH, Castoe TA, Campbell JA (2012) Cryptic diversity in disjunct  
640 populations of Middle American Montane Pitvipers: a systematic reassessment of  
641 *Cerrophidion godmani*. *Zoologica Scripta* 41: 455-470. [https://doi.org/10.1111/j.1463-](https://doi.org/10.1111/j.1463-6409.2012.00547.x)  
642 [6409.2012.00547.x](https://doi.org/10.1111/j.1463-6409.2012.00547.x)

643 Jadin RC, Blair C, Orlofske SA, Jowers MJ, Rivas GA, Vitt LJ, Ray JM, Smith EN, Murphy  
644 JC (2020) Not withering on the evolutionary vine: systematic revision of the Brown Vine  
645 Snake (Reptilia: Squamata: *Oxybelis*) from its northern distribution. *Organisms Diversity &*  
646 *Evolution* 20: 723–746. <https://doi.org/10.1007/s13127-020-00461-0>

647 Jarvis A, Reuter HI, Nelson A, Guevara E (2008) Hole-filled seamless SRTM data V4,  
648 International Centre for Tropical Agriculture (CIAT), available from  
649 <http://srtm.csi.cgiar.org>. Accessed: 2023-12-19.

650 Johnson JD, Mata-Silva V, Wilson LD (2017) A conservation reassessment of the Central  
651 American herpetofauna based on the EVS measure. *Amphibian & Reptile Conservation* 9:  
652 1-94.

653 Jung H, Kalter R, Untiedt, A, Villalobos-Heredia C (2022) Mesoamerica Ecological  
654 Forecasting: Assessing Land Cover Change to Inform Management Planning for the  
655 Mesoamerican Biological Corridor.

656 Kolby JE (2009) Geographic distribution: *Craugastor coffeus* (Coffee Rain Frog).  
657 Herpetological Review 40: 107.

658 Kolby JE, McCranie JR (2009) Discovery of a Surviving Population of the Montane Streamside  
659 Frog *Craugastor milesi* (Schmidt). Herpetological Review 40: 282-283.

660 Kolby JE, McCranie JR, Rovito SM (2009) Geographic distribution: *Nototriton brodiei*.  
661 Herpetological Review, 40: 444.

662 Kolby JE, Padgett-Flohr GE, Field R (2010) Amphibian chytrid fungus (*Batrachochytrium*  
663 *dendrobatidis*) in Cusuco National Park, Honduras. Diseases of Aquatic Organisms 92: 245-  
664 251. <https://doi.org/10.3354/dao02055>

665 Kolby JE, Ramirez SD, Berger, L, Griffin DW, Jocque M, Skerratt LF (2015a) Presence of  
666 amphibian chytrid fungus (*Batrachochytrium dendrobatidis*) in rainwater suggests aerial  
667 dispersal is possible. Aerobiologia 31, 411–419 (2015). <https://doi.org/10.1007/s10453-015-9374-6>

668

669 Kolby JE, Ramirez SD, Berger L, Richards-Hrdlicka KL, Jocque M, Skerratt LF (2015b)  
670 Terrestrial Dispersal and Potential Environmental Transmission of the Amphibian Chytrid  
671 Fungus (*Batrachochytrium dendrobatidis*). PLoS One. 2015 Apr 30;10(4):e0125386.  
672 <https://doi.org/10.1371/journal.pone.0125386>

673 Köhler G, Townsend JH, Peterson CBP (2016) A taxonomic revision of the *Norops*  
674 *tropidonotus* complex (Squamata, Dactyloidae), with the resurrection of *N. spilorhipis*  
675 (Álvarez del Toro and Smith, 1956) and the description of two new species. Mesoamerican  
676 Herpetology 3: 8-41.

677 Laurance WF, Useche DC, Shoo LP, Herzog SK, Kessler M, Escobar F, Brehm G, Axmacher  
678 JC, Chen IC, Gamez LA, Hietz P, Fiedler K, Pyrcz T, Wolf J, Merkord CL, Cardelus C,  
679 Marshall AR, Ah-Peng C, Aplet GH, Arizmendi MD, Baker WJ, Barone J, Bruhl CA,  
680 Bussmann RW, Cicuzza D, Eilu G, Favila ME, Hemp A, Hemp C, Homeier J, Hurtado J,  
681 Jankowski J, Kattan G, Kluge J, Kromer T, Lees DC, Lehnert M, Longino JT, Lovett J,  
682 Martin PH, Patterson BD, Pearson RG, Peh KSH, Richardson B, Richardson M, Samways  
683 MJ, Senbeta F, Smith TB, Utteridge TMA, Watkins JE, Wilson R, Williams SE, Thomas  
684 CD (2011) Global warming, elevational ranges and the vulnerability of tropical biota.  
685 Biological Conservation 144: 548–557. <https://doi.org/10.1016/j.biocon.2010.10.010>

686 Le Saout S, Hoffmann M, Shi Y, Hughes A, Bernard C, Brooks TM, Bertzky B, Butchart SHM,  
687 Stuart SN, Badman T, Rodrigues ASL (2013) Protected areas and effective biodiversity  
688 conservation. Science 342: 803-805. <https://doi.org/10.1126/science.1239268>

689 Linkem CW, Diesmos AC, Brown RM (2011) Molecular systematics of the Philippine forest  
690 skinks (Squamata: Scincidae: *Sphenomorphus*): testing morphological hypotheses of  
691 interspecific relationships. Zoological Journal of the Linnean Society 163: 1217-1243.  
692 <https://doi.org/10.1111/j.1096-3642.2011.00747.x>

693 Lomolino MV (2004) Conservation biogeography. In: Lomolino MV, Heaney LR (Eds)  
694 Frontiers of Biogeography: new directions in the geography of nature. Sinauer Associates,  
695 Sunderland, Massachusetts, 293–296.

696 Martin TE, Blackburn GA (2009) The effectiveness of a Mesoamerican ‘paper park’ in  
697 conserving cloud forest avifauna. Biodivers. Conserv., 18, 3841–3859.  
698 <https://doi.org/10.1007/s10531-009-9683-6>

699 Martin TE, Jones SEI, Creedy TJ, Hoskins HMJ, McCann NP, Batke SP, Kelly DL, Kolby JE,  
700 Downing R, Zelaya SMS, Green SEW, Lonsdale G, Brown T, Waters S, Rodríguez-Vásquez  
701 F, McCravy KW, D’Souza ML, Crace D, Nuñez-mino JM, Haelewaters D, Medina-van  
702 Berkum P, Phipps CD, Barker RJ, Castañeda F, Reid N, Jocque M (2021) A review of the

703 ecological value of Cusuco National Park: an urgent call for conservation action in a highly  
704 threatened Mesoamerican cloud forest. *Journal of Mesoamerican Biology* 1: 6-50.

705 Martin TE, Rodríguez F, Simcox W, Dickson I, Van Dort, J, Reyes, E, Jones, SEI (2016)  
706 Notable range and altitudinal records from Cusuco National Park and environs, north-  
707 western Honduras. *Cotinga* 38: 32-39.

708 Martínez ML, Perez-Maqueo O, Vazquez G, Castillo-Campos G, Garcia-Franco J, Mehlreter  
709 K, Equihua M, Landgrave R (2009) Effects of land use change on biodiversity and ecosystem  
710 services in tropical montane cloud forests of Mexico. *Forest Ecology and Management* 258:  
711 1856-1863. <https://doi.org/10.1016/j.foreco.2009.02.023>

712 McCranie JR (2015) A checklist of the amphibians and reptiles of Honduras, with additions,  
713 comments on taxonomy, some recent taxonomic decisions, and areas of further studies  
714 needed. *Zootaxa* 3931: 352–386. <https://doi.org/10.11646/zootaxa.3931.3.2>

715 McCranie JR (2018) A discussion of the phenetic-based *Craugastor laticeps* species group  
716 (Anura: Brachycephaloidea: Craugastoridae) from north-central Honduras, with the  
717 description of two new species. *Herpetologica* 74: 169-180.  
718 <https://doi.org/10.1655/Herpetologica-D-17-00023.1>

719 McCranie JR, Köhler G (2015) The anoles (Reptilia: Squamata: Dactyloidae: *Anolis: Norops*)  
720 of Honduras. Systematics, distribution, and conservation. *Bulletin of the Museum of*  
721 *Comparative Zoology* 161: 1-292. <https://doi.org/10.3099/0027-4100-14.1.1>

722 McCranie JR, Rovito SM (2014) New species of salamander (Caudata: Plethodontidae:  
723 *Cryptotriton*) from Quebrada Cataguana, Francisco Morazán, Honduras, with comments on  
724 the taxonomic status of *Cryptotriton wakei*. *Zootaxa* 3795: 61-70.  
725 <https://doi.org/10.11646/zootaxa.3795.1.6>

726 McCranie JR, Solís JM (2014) Additions to the amphibians and reptiles of Parque Nacional  
727 Pico Bonito, Honduras, with an updated nomenclatural list. *Herpetology Notes* 6: 239-243.

728 Medina-van Berkum P, Vulinec K, Crace D, Gallego ZL, Martin TE (2020) Community  
729 Composition of Bats in Cusuco National Park, Honduras, a Mesoamerican Cloud Forest,  
730 Including New Regional and Altitudinal Records. *Neotropical Naturalist* 3: 1-24.

731 Menegon M, Cantaloni C, Rodriguez-Prieto A, Centomo C, Abdelfattah A, Rossato M,  
732 Bernardini M, Xumerle L, Loader S, Delledonne M (2017) On site DNA barcoding by  
733 nanopore sequencing. *PLoS One* 12: e0184741.  
734 <https://doi.org/10.1371/journal.pone.0184741>

735 Myers N, Mittermeier RA, Mittermeier CG, da Fonseca GAB, Kent J (2000) Biodiversity  
736 hotspots for conservation priorities. *Nature* 403: 853–858. <https://doi.org/10.1038/35002501>

737 Neate-Clegg MHC, Jones SEI, Burdekin O, Jocque M, Şekercioğlu ÇH (2018) Elevational  
738 changes in the avian community of a Mesoamerican cloud forest park. *Biotropica* 50: 805–  
739 815. <https://doi.org/10.1111/btp.12596>

740 Poe S (2013) 1986 Redux: New genera of anoles (Squamata: Dactyloidae) are unwarranted.  
741 *Zootaxa* 3626: 295–299.

742 Ponce-Reyes R, Reynoso-Rosales V-H, Watson JEM, VanDerWal J, Fuller RA, Pressey RL,  
743 Possingham HP (2012) Vulnerability of cloud forest reserves in Mexico to climate change.  
744 *Nature Climate Change* 2: 448–452. <https://doi.org/10.1038/nclimate1453>

745 Pyron AR, Burbrink FT, Wiens, JJ (2013) A phylogeny and revised classification of Squamata,  
746 including 4161 species of lizards and snakes. *BMC evolutionary biology* 13: 1-54.  
747 <https://doi.org/10.1186/1471-2148-13-93>

748 Reyes-Velasco J, Adams RH, Boissinot S, Parkinson CL, Campbell JA, Castoe TA, Smith EN  
749 (2020) Genome-wide SNPs clarify lineage diversity confused by coloration in coral snakes  
750 of the *Micrurus diastema* species complex (Serpentes: Elapidae). *Molecular Phylogenetics*  
751 *and Evolution* 147: 106770. <https://doi.org/10.1016/j.ympev.2020.106770>

- 752 Ruane S, Bryson Jr RW, Pyron RA, Burbrink FT (2014) Coalescent Species Delimitation in  
753 Milksnakes (Genus *Lampropeltis*) and Impacts on Phylogenetic Comparative Analyses.  
754 *Systematic Biology* 63: 231-250. <https://doi.org/10.1093/sysbio/syt099>
- 755 Scheele BC, Pasmans F, Skerratt LF, Berger L, Martel A, Beukema W, Acevedo AA, Burrowes  
756 PA, Carvalho T, Catenazzi A, De la Riva I, Fisher MC, Flechas SV, Foster CN, Frías-  
757 Álvarez P, Garner TWJ, Gratwicke B, Guayasamin JM, Hirschfeld M, Kolby JE, Kosch TA,  
758 La Marca E, Lindenmayer DB, Lips KR, Longo AV, Maneyro R, McDonald CA, Mendelson  
759 J, Palacios-Rodriguez P, Parra-Olea G, Richards-Zawacki CL, Rödel M-O, Rovito SM,  
760 Soto-Azat C, Toledo LF, Voyles J, Weldon C, Whitfield SM, Wilkinson M, Zamudio KR,  
761 Canessa S (2019) Amphibian fungal panzootic causes catastrophic and ongoing loss of  
762 biodiversity. *Science* 363: 1459-1463. <https://doi.org/10.1126/science.aav0379>
- 763 Schools M, Hedges SB (2021). Phylogenetics, classification, and biogeography of the  
764 Neotropical Forest lizards (Squamata, Diploglossidae). *Zootaxa* 4974: 201–257.  
765 <https://doi.org/10.11646/ZOOTAXA.4974.2.1>
- 766 Solís JM, Wilson LD, Townsend JH (2014) An updated list of the amphibians and reptiles of  
767 Honduras, with comments on their nomenclature. *Mesoamerican Herpetology* 1: 123-144.
- 768 Solís JM, Taylor P, López-Paredes J (2017). Confirmation of the occurrence of a poorly known  
769 species, *Ecnomiohyla salvaje* (Anura: Hylidae) from El Cusuco National Park, Honduras  
770 with an updated list of its amphibian species. *Alytes* 34: 62-68.
- 771 Stadtmüller T (1987) Cloud Forests in the Humid Tropics. A Bibliographic Review. The United  
772 Nations University, Tokyo and Centro Agronomico Tropical de Investigacion y Ensenanza,  
773 Turrialba, Costa Rica, 81 pp.
- 774 Thorp C, Phipps CD, Lonsdale G, Arrivillaga C, Brown TW, Snyder A (2021) *Ecnomiohyla*  
775 *salvaje* Hylidae (Wilson, McCranie & Williams 1985) in Cusuco National Park, Honduras:  
776 morphological descriptions of adults and metamorphs with notes on ecology, natural history,  
777 and the conservation implications of amphibian chytrid fungus (Bd). *Journal of*  
778 *Mesoamerican Biology* 1: 70-88
- 779 Townsend JH (2014) Characterizing the Chortís block biogeographic province: geological,  
780 physiographic, and ecological associations and herpetofauna diversity. *Mesoamerican*  
781 *Herpetology* 1: 204–252.
- 782 Townsend JH, Butler JM, Wilson LD, Austin JD (2010) A distinctive new species of moss  
783 salamander (Caudata: Plethodontidae: *Nototriton*) from an imperiled Honduran endemism  
784 hotspot. *Zootaxa* 2434: 1-16.
- 785 Townsend JH, Wilson LD (2008) Guide to the Amphibians and Reptiles of Cusuco National  
786 Park, Honduras / Guía de los Anfíbios y Reptiles de Parque Nacional Cusuco, Honduras.  
787 Bibliomania, Salt Lake City, UT.
- 788 Townsend JH, Wilson LD, Talley BL, Fraser DC, Plenderleith TL, Hughes SM (2006)  
789 Additions to the herpetofauna of Parque Nacional El Cusuco, Honduras. *Herpetological*  
790 *Bulletin* 96: 29-39.
- 791 Vences M, Thomas M, Bonett RM, Vieites DR (2005) Deciphering amphibian diversity  
792 through DNA barcoding: chances and challenges. *Philosophical Transactions of the Royal*  
793 *Society B* 360: 1859–1868. <https://doi.org/10.1098/rstb.2005.1717>
- 794 Whittaker RJ, Araújo MB, Jepson P, Ladle RJ, Watson JEM, Willis KJ (2005) Conservation  
795 Biogeography: assessment and prospect. *Diversity and Distributions* 11: 3-23.  
796 <https://doi.org/10.1111/j.1366-9516.2005.00143.x>
- 797 Wilson LD, McCranie JR (2004) The herpetofauna of Parque Nacional El Cusuco, Honduras.  
798 *Herpetological Bulletin* 87: 13–24.
- 799 Yuan ZY, Zhou WW, Chen X, Poyarkov Jr NA, Chen HM, Jang-Liaw NH, Chou WH, Matzke  
800 NJ, Iizuka K, Min MS, Kuzmin SL, Zhang YP, Cannatella DC, Hillis DM, Che J (2016)  
801 Spatiotemporal Diversification of the True Frogs (Genus *Rana*): A Historical Framework

802 for a Widely Studied Group of Model Organisms. *Systematic Biology* 65: 824-842.  
803 <https://doi.org/10.1093/sysbio/syw055>  
804 Zipkin EF, DiRenzo GV, Ray JM, Rossman M, Lips KR (2020) Tropical snake diversity  
805 collapses after widespread amphibian loss. *Science* 367: 814-816.  
806 <https://doi.org/10.1126/science.aay5733>

807 **Supplementary material**

808

809 **Supplementary material 1: Fig. S1.** Photographic vouchers of amphibian species recorded in  
810 Cusuco National Park during the research period 2007-2023.

811

812 **Supplementary material 2: Fig. S2.** Photographic vouchers of reptile species recorded in  
813 Cusuco National Park during the research period 2007-2023.