Disordered eating in male athletes: A meta-analysis

Chapman, J.; Woodman, Tim

Journal of Sports Sciences

DOI:

[10.1080/02640414.2015.1040824](https://doi.org/10.1080/02640414.2015.1040824)

Published: 28/04/2015

Peer reviewed version

[Cyswllt i'r cyhoeddiad / Link to publication](#)

Dyfyniad o'r fersiwn a gyhoeddwyd / Citation for published version (APA):

Chapman, J., & Woodman, T. (2015). Disordered eating in male athletes: A meta-analysis. *Journal of Sports Sciences*. <https://doi.org/10.1080/02640414.2015.1040824>

Hawliau Cyffredinol / General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

This is an Accepted Manuscript of an article published by Taylor & Francis in *Journal of Sports Sciences* on 28/04/2015, available online: <http://www.tandfonline.com> DOI 10.1080/02640414.2015.1040824.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Running Head: DISORDERED EATING IN MALE ATHLETES

**This article was accepted in its current form in the Journal of Sports Sciences on 11th
April 2015. Doi: 10.1080/02640414.2015.1040824**

Disordered eating in male athletes: A meta-analysis

James Chapman, Sussex Coast College; University of Brighton

Tim Woodman, Bangor University

Key Words: Eating disorder, Disordered eating, Men

Date of Submission: 19th September 2014

Date of resubmission: 26th January 2015

Date of resubmission: 9th March 2015

Abstract

We examined the propensity for male athletes to exhibit symptoms of disordered eating. Using meta-analytic techniques, we examined overall effect size, individual effect sizes for specific sport types, standard of athletic competition, and diagnostic tools from 31 studies. When all studies were considered as a homogeneous group, male athletes did not have symptoms of disordered eating that were significantly different from non-athletic controls. However, significant moderator effects emerged for sport type and measurement: (a) wrestling reported a greater incidence of disordered eating; and (b) studies that reported data from the Eating Attitudes Test yielded a significantly greater incidence of disordered eating in male athletes compared to non-athletes. Although some sports seem to present a higher risk of disordered eating compared to others, the effects are weak and heterogeneous. We make suggestions for the development of the research area, which has been severely hampered by the diagnostic tools that have been available for the study of men.

Disordered eating in male athletes: A meta-analysis

It has been suggested that athletes, especially those involved in sports that emphasise lean body mass and shape, may be at higher risk of experiencing disordered eating than people from the general population (Sundgot-Borgen & Torstveit, 2004). Although some male athletes (e.g. from mass-dependent sports such as wrestling) are thought to be at some risk (e.g. Sundgot-Borgen & Torstveit, 2004), the vast majority of disordered eating research has been conducted with women (e.g. Anshel, 2004; Pollock et al. 2010; Smolak, Murnen, & Ruble, 2000), which has resulted in a paucity of male-specific literature. The aim of this paper is to synthesise the research in disordered eating in male athletes with a view to establishing some priorities for future research.

An eating disorder is defined by behavioural eating symptoms (e.g. binge eating, dieting), psychological symptoms (e.g. body image concerns), and physical symptoms (e.g. weight loss), which can be clinical (e.g. anorexia nervosa, bulimia nervosa) or subclinical (American Psychiatric Association, 2013). Symptoms associated with eating disorders, such as episodes of binge eating or extreme dieting, are considered subclinical when the behaviours are present but insufficiently frequent or severe to be deemed clinical (Anderson & Petrie, 2012; Petrie, Greenleaf, Carter & Reel, 2007). Individuals who display such symptomatic sub-clinical behaviours are said to exhibit *disordered eating* behaviours. Disordered eating behaviours are considered clear risk factors for developing a clinical eating disorder when the behaviour is continued (Kotler, Cohen, Davies, Pine & Walsh; 2001).

Although disordered eating has traditionally been associated with women (Striegel-Moore & Bulik, 2007), research attention has recently begun to focus on men. In their review of disordered eating in athletes, Byrne and McLean (2001) proposed two arguments for why male athletes might be more at risk than non-athletes. First, in addition to the general demands of Western society to be lean, athletes may be under sport-specific pressure to

1 conform to a certain body type. For example, for sports such as gymnastics, lean and
2 muscular bodies are typically seen as an advantage. Second, patients diagnosed with an
3 eating disorder often demonstrate similar characteristics to those that have been observed in
4 high-level athletes. Specifically, perfectionism and the need for high achievement are found
5 in both eating disorder patients (Bardone-Cone, Abramson, Vohs, Heatherton, & Joiner,
6 2006) and in athletes (Haase, 2011).

7 Research examining disordered eating in athletes has been equivocal (Hausenblas &
8 McNally, 2004), with considerable ambiguity regarding male athletes. The largest study to
9 date in an elite male sample revealed that Norwegian athletes ($n = 687$) reported more clinical
10 and subclinical eating disorders (8%) than a matched control group ($n = 629$; 0.5%). In the
11 same study, anti-gravitation (e.g. high jump, long jump, triple jump) and “weight” class sport
12 (e.g. combat sports) athletes were at the highest risk, with 22% and 18% reporting a clinical
13 disorder, respectively (Sundgot-Borgen & Torstveit, 2004). There are, however, a number of
14 studies that have reported no significant differences between male athletes and controls (e.g.
15 Martinsen, Bratland-Sanda, Eriksson & Sundgot-Borgen, 2010). Other research has even
16 revealed that male athletes display behaviours that reflect lower levels of disordered eating
17 than their non-athletic counterparts. For example, Croll et al. (2006) reported that male
18 adolescent power sport athletes had a more balanced and nutritious diet than non-athletes.
19 Equally, Fogelholm and Hiilloskorpi (1999) reported that male athletes did not want to lose
20 body mass, as a loss of mass was considered a loss of strength, which suggests that the
21 motivation and behaviours of male athletes might somewhat protect them from developing
22 disordered eating.

23 Due to the equivocal nature of disordered eating prevalence findings, Hausenblas and
24 McNally (2004) suggested a number of variables that might moderate the relationship
25 between athletes and disordered eating, three of which are directly relevant to the present

1 analysis. The first potential moderator is the competitive standard of the athletic group. In a
2 female sample it was shown that an increase in competitive standard leads athletes to
3 encounter more intense training and greater pressure to maintain a specific body mass, which
4 might lead to a greater likelihood of disordered eating behaviours in athletes who operate at
5 an elite standard (Sundgot-Borgen, 1994). Conversely, disordered eating may in fact be less
6 likely in elite samples. Indeed, Darcy, Hardy, Lock, Hill, and Peebles (2013) found that male
7 recreational collegiate athletes reported higher disordered eating than those competing at
8 higher competitive standards.

9 Second, sport type has frequently been shown to be the strongest moderator of the
10 relationship between male athletes and disordered eating. Indeed, many studies have reported
11 that athletes competing in sports associated with “leanness” (e.g. anti-gravitation sports such
12 as high-jumping) are most at risk (Hausenblas & Carron, 1999; Sundgot-Borgen & Torstveit,
13 2004). The final potential moderator relevant to the current paper is the measurement tool
14 that is used to assess eating behaviour (Hausenblas & McNally, 2004). The two measures that
15 are most commonly used are the Eating Attitudes Test (EAT; Garner et al., 1982) and the
16 Eating Disorder Inventory (EDI; Garner, 1991). Although both measures assess symptoms of
17 disordered eating, there are differences between the sub-scales of each measure. These
18 differences relate to how males are thought to experience disordered eating and could help to
19 explain the equivocal findings in the literature. For example, Stanford and Lemberg (2012a)
20 argued that men’s body-related motives are driven more by a preoccupation to increase
21 leanness and muscularity than a desire to lose body mass (which appears associated more
22 with the female motive), and are thus more likely to regulate shape (rather than mass) through
23 excessive exercise and diet. Neither excessive exercise nor body shape are considered in the
24 aforementioned diagnostic tools, which were developed using female samples. However,

given that *food preoccupation* and *dieting behaviour* are categories that are assessed within the EAT, the EAT may be a somewhat more sensitive measure of men's eating behaviours.

The purpose of this paper is to apply meta-analytic procedures to investigate whether male athletes are more at risk of disordered eating relative to male controls, and whether there are moderators that might explain some of the inconsistent findings in the literature. We seek to explore three potential moderators: sport type, competitive standard, and measurement tool. Due to the (theoretically and empirically) equivocal research to date, we hypothesise that male athletes – when considered as a single homogenous group – will report levels of disordered eating that are comparable to those of controls. Conversely, we hypothesise that athletes who participate in sports that promote lean body mass and shape will report higher levels of disordered eating. Given the equivocal research on elite athletes, we do not formulate a specific hypothesis regarding competitive standard. Finally, we hypothesise that the global score on the EAT will demonstrate a disordered eating discriminatory capacity that is greater than that of other diagnostic measures because it contains categories that are likely more specific to men's disordered eating (e.g. *food preoccupation*) than other diagnostic subscales (e.g. EDI: *drive for thinness*).

Method

Studies included in the meta-analysis

A study was included in the meta-analysis if: a) it had a group of male athletes and a control group of male non-athletes; and b) the results included means and standard deviations, *F* values, *t* values, *r* values, or percentages for the relevant variables. All studies published before 1st January 2014 were eligible for inclusion in the analysis.

Articles were located using an online search of journal data-bases including: PsycInfo, Psych Lit, Sport Discus, Science Direct, Web of Knowledge, Pub Med, Ingenta-Connect, First Search, and Google Scholar. The key terms used to search for articles included: eating

disorders, eating problems, disordered eating, anorexia, bulimia, anorexia athletica, male athletes, male, men, exercise, and sport. On completion of the data-base searches, further electronic searches of key authors were conducted. The indexes for the last 15 years of the *International Journal of Eating Disorders* were also searched electronically. Bibliographies and reference sections of review articles together with each article collected were individually checked. Finally, in cases where studies were unobtainable or data that were relevant to the meta-analysis were not reported, the corresponding author of the study concerned was contacted directly via email or telephone.

Thirty-six studies met the inclusion criteria. Eight of the studies did not report the data required for the analysis and so the corresponding authors were contacted. In the event that the authors did not respond to a voicemail or email, a follow-up request for the data was made two weeks later. As none of the eight authors replied, 28 of the studies used in the meta-analysis reported all the data that were required for analysis purposes. Three of the eight incomplete studies simply reported “non-significant” results. Consequently, p was assumed to be 0.50 and r was assumed to be 0 in these cases (cf. Rosenthal, 1991). Thirty-one studies were thus retained for analysis.

Coding the studies

Participant characteristics were coded according to standard of competition (elite, non-elite) and specific sport. In line with Sundgot-Borgen and Torstveit (2004), sports were categorized into six sport groups: endurance sports (running, swimming, rowing, and cycling), mass-dependent sports (wrestling, karate, and judo), aesthetic sports (figure skating, cheerleading, diving, gymnastics, and bodybuilding), technical sports (bowling, golf, high jump, equestrian, long jump, shooting, and sailing), power sports (discus, javelin, power lifting, shot put, and sprinting), and ball sports (badminton, basketball, football, soccer, table tennis, handball, tennis, and volleyball). Elite athletes were operationalised as those athletes

who competed in national (including Division 1 collegiate athletes) or international competitions. Study characteristics were coded according to total sample size, number of participants in each experimental group and in the control group, the t statistic (or equivalent) for each comparison in the analysis, and means and standard deviations for each group on the diagnostic measure used, and the dependent variable or subcategory of the measure used.

Statistical analysis

The statistical methods used were those described in Rosenthal (1991). We calculated effect sizes for the 28 studies that met the inclusion criteria and used Zr as the measure of effect size (see Rosenthal, 1991). To calculate the significance of each effect size the standard normal deviate Z was used. We further calculate the average effect size Zr by weighting each study by its individual sample size. The average effect size Zr was then converted to a Z score for statistical significance. We also calculated heterogeneity estimates, in the first instance as Cochran's Q statistic, which is distributed as the χ^2 statistic with $k - 1$ degrees of freedom (where k = number of effect sizes). In response to the suggestion that the Cochran Q statistic is a poor predictor of heterogeneity in analyses that contain low study numbers, and in line with recommendations from Higgins, Thompson, Deeks and Altman (2003), we also calculated the I^2 heterogeneity statistic. Finally, moderator variables were tested for significance using z score calculations (see Rosenthal, 1991).

In the event that a study had more than one experimental group and only one control group, independent t -tests were used for each comparison. A mean of these t scores was then calculated before transforming the t score to a correlation coefficient using the mean sample sizes to satisfy the assumption of independence. In order to include the most effect sizes in the moderator analyses the individual t scores for sport type and competitive level were also transformed into independent correlation coefficients.

Results

We present a summary of all the studies included in the analysis in Table 1, and we present a forest plot for the effect sizes that were included in the overall analysis in Figure 1.

Overall difference between athletes and non-athletes

The comparison between athletes and controls revealed no significant difference for disordered eating, as hypothesised, $r = .07$, $z = 1.30$, $p = 0.19$. This overall r is based on r values ranging between $-.47$ and $.52$ with significant heterogeneity, $\chi^2(31) = 129.26$, $p < .001$, $I^2 = 76.02\%$. The mean total sample of the studies included in this analysis was 345.45.

Sport type

It was hypothesised that athletes from sports emphasising lean body mass and shape would report higher levels of disordered eating than their non-athletic counterparts and that these differences would be more pronounced than in sports that do not emphasise such body types. Separate analyses were conducted for endurance sports (eight effect sizes), mass-dependent sports (nine effect sizes), and aesthetic sports (seven effect sizes). Other sport types were not analysed due to an insufficient number of effect sizes (cf. Rosenthal, 1991).

When endurance athletes were compared to controls no significant difference was revealed for disordered eating, $r = .02$, $z = 0.27$, $p = .79$, and it was marked with significant heterogeneity $\chi^2(8) = 34$, $p < .001$, $I^2 = 76.47\%$. This overall r is based on r values ranging between $-.13$ and $.55$ with a mean total sample size of 178.50. The moderator analysis was also found to be non-significant, $z = .08$.

The comparison between mass-dependent sport athletes and controls revealed no significant difference for disordered eating, $r = .11$, $z = 1.46$, $p = 0.14$. The overall r is based on r values ranging between $-.65$ and $.39$. The mean total sample was 175.67. As much of the research has suggested that male athletes competing in mass-dependent sports such as wrestling are at increased risk (Rosendahl et al., 2009; Sundgot-Borgen & Torstveit, 2004), whereas there is little evidence to suggest such risk in martial artists, the analysis was

repeated without the martial arts samples. This second comparison revealed a significant disordered eating effect for wrestlers compared to non-athlete controls, $r = .14$, $z = 2.22$, $p = .03$. The overall r was based on r values ranging from .03 to .39, the mean sample size was 251.50, and heterogeneity was not significant $\chi^2 (6) = 8.8$, $p = .18$, $I^2 = 31.82\%$. However, mass-dependent sports, either including or excluding martial arts, were not found to be significant moderators; $z = .10$ and $z = .53$, respectively.

Finally, the comparison between aesthetic sport athletes and controls revealed no significant effect for disordered eating, $r = .11$, $z = 1.62$, $p = .11$. This overall r is based on r values ranging from -.15 and .23. There was significant heterogeneity $\chi^2 (7) = 14.82$, $p = 0.02$, $I^2 = 52.77\%$. The mean total sample was 181.29.

Competitive Standard

The comparison between elite athletes and controls revealed no significant effect for disordered eating, $r = .07$, $z = 1.14$, $p = .11$. This overall r is based on r values ranging from -.47 and .42. There was significant heterogeneity, $\chi^2 (11) = 49.18$, $p < .001$, $I^2 = 77.63\%$. The mean total sample was 263.64.

Measurement

The restricted number of studies allowed for three analyses to be conducted with a view to assessing the effect sizes for three dependent variables: the drive for thinness and bulimia subcategories of the EDI, and the total score from the EAT. Of the 12 studies that were included in the analysis and that used the EDI, only the *drive for thinness* subscale data were reported in all cases with the bulimia subscale reported on ten occasions. Similarly, in the 13 studies in which the EAT was used, data from the subscales were reported on four or fewer occasions. For the drive for thinness subscale comparison, no significant difference between athletes and non-athletes was revealed, $r = .04$, $z = .71$, $p = .48$. This overall r is

based on r values ranging from $-.15$ and $.52$. There was significant heterogeneity $\chi^2 (12) = 34.77, p < .001, I^2 = 65.49\%$. The mean total sample was 316.50.

The Bulimia comparison was also non-significant, $r = .09, z = 1.25, p = .19$. This result was not marked with significant heterogeneity $\chi^2 (10) = 14.86, p = .06, I^2 = 32.71$.

The total EAT comparison revealed a significant disordered eating difference between athletes and non-athletes with athletes appearing to be more at risk, $r = .12, z = 2.07, p = .04$.

This result was again marked with significant heterogeneity, $\chi^2 (13) = 63.01, p < .001, I^2 = 79.37\%$. The mean sample size was 299.00. The EAT comparison was not significantly different to the effect size produced from the drive for thinness or bulimia analyses; $z = .02$ and $.07$, respectively.

File drawer analyses

For the probability of the follow-up wrestling comparison to become non-significant, 54 studies with a mean probability of $.50$ would need to have been stored away. As this number of additional null-finding studies required is higher than the number of studies that were included in the analysis, we can consider it reasonably robust to the file drawer threat. . To become non-significant the EAT analysis would require 159 studies with a mean probability of $.50$ to be stored away. Thus, the EAT finding is also robust against the file drawer threat.

Discussion

The purpose of this review was to establish whether male athletes were more susceptible to disordered eating compared to male controls and whether there were moderating variables of this susceptibility to disordered eating. To this end, effect sizes were calculated for sport type, competitive standard and diagnostic tools. No significant difference was revealed across standards of competition. There was, however, a significant difference between wrestlers and controls. Regarding diagnostic tools, the total score on the EAT

emerged as a significant discriminator between athletes and controls. Conversely, the drive for thinness and bulimia subscales of the EDI were not significant discriminators.

It was hypothesised that athletes participating in sports emphasizing lean body mass and shape would report higher levels of disordered eating than controls. Previous research has suggested that male athletes from mass-dependent sports (Sundgot-Borgen & Torstveit, 2004), aesthetic sports (Krentz & Warschburger, 2011), and endurance sports (Riebl et al., 2007) are more likely to display disordered eating. In the present paper the effect size was significant for wrestling only, with non-significant effect sizes emerging for endurance and aesthetic sports. The finding that wrestlers are more at risk than non-athletes most likely reflects these athletes' desire to achieve a high muscle mass and low body fat mass to compete in a specific "weight" class, which may be below their more natural body mass. Much of the previous literature has focused on mass-dependent sports as a homogenous group, in which wrestling is grouped with sports such as boxing and the martial arts. In this review, however, both studies involving judo athletes produced a negative effect (Filaire et al., 2007; Rouveix et al., 2007) and another involving martial arts produced non-significant results (Costarelli & Stamou, 2009). This difference between sports is informative, as it suggests that different body mass-dependent sports might harbour different degrees of risk regarding eating behaviours and should not be treated as a homogeneous group.

Endurance sports have commonly been cited as high-risk for female athletes (Pollock et al., 2010) but male athletes involved in endurance sports were not significantly more at risk than non-athletes. At first view, the non-significant finding reported here suggests that the risk associated with endurance sports may be unique to female endurance athletes. There are two main problems associated with such an interpretation.

First, sports such as running and swimming are typically operationalised within a single category of "endurance" sports despite the wide variety of activities involved in these

sports (e.g. 100-metre run or swim). A more fine-grained analysis of each sport, rather than sport categories, is likely to yield a more sharply focused picture of the relative risks associated with each specific sport activity. Unfortunately, given the lack of data, such an analysis was not possible and more research is clearly warranted here.

Second, the finding that endurance sports are no more at risk of disordered eating than their non-athletic counterparts is inconsistent with research that has considered disordered eating beyond its *prevalence*. Indeed, Atkinson's (2011) interviews with male endurance athletes (long distance runners and triathletes) revealed a number of disordered eating incidences within this athletic group. The lack of consistency or clear evidence from the current meta-analysis might consequently be masking significant instances of disordered eating within each or any of the endurance (or other) sport groups. It is further noteworthy that there is a reluctance for male athletes to disclose eating issues, which will necessarily limit the knowledge gains that might be brought about by questionnaire-based research. In other words, there is a need to consider more in-depth analyses and potentially "to reduce the prevalence of prevalence studies" (Papathomas & Lavalley, 2012, p. 389).

The current analysis revealed no significant difference between athletes of aesthetic sports and controls. In a Norwegian sample of male aesthetic athletes (Sundgot-Borgen & Torstveit, 2004), none of the athletes were diagnosed with a clinical eating disorder and the authors suggested that the male athletes had healthier methods of restricting their body mass for competition. Another possible explanation for what appears to be an exclusively female disordered eating relationship (see Anderson & Petrie, 2012) is that a large number of the aesthetic sport sample in this analysis comprised bodybuilders. Bodybuilders are likely to report low scores on diagnostic tools that operationalise a desire to lose body mass as a risk indicator, as these athletes are judged on their ability to maintain and *increase* body mass and muscularity. Such athletes are likely to report any shape or weight concern only if presented

1 with a scale that reflects a desire for *leanness* and *muscularity* rather than a reduction in body
2 *mass* or a desire for *thinness*.

3 As hypothesised the meta-analysis revealed a significant difference between athletes
4 and non-athletes for studies that used the total score on the EAT. Conversely, there was no
5 significant difference in studies that used the drive for thinness or bulimia scales of the EDI.
6 A parsimonious explanation is that this measurement-derived discrepancy is a reflection of
7 the EAT being a better predictor for male eating problems than are the other scales. When
8 considered against the *drive for thinness* subscale it should be noted that thinness is
9 associated with body *mass* whereas the drive for leanness in men infers a desire for less body
10 *fat* and yet greater *muscle* mass, or at least a more visible abdominal muscle mass (Leit et al,
11 2001; Oliviardia, Pope, Borowiecki, & Cohane, 2004). Further, it is worth noting that both
12 the EDI and the EAT were developed with female samples. Although risk factors of eating
13 disorders are largely shared by male and female patients, the EDI and EAT focus on body
14 “weight” and shape concerns that are perhaps more common to females (dieting, thinness)
15 than they are to males. Regardless of how one might interpret the greater discriminatory
16 power of the EAT over the analysed subscales of the EDI, there is a clear need to move
17 toward more male-specific ways of operationalising and measuring men’s and boys’
18 disordered eating in the specific context of sport research (see also Woodman & Hemmings,
19 2008; Woodman & Steer, 2011).

20 A significant measurement step forward is Stanford and Lemberg’s (2012b) Eating
21 Disorder Assessment for Men (EDAM). The EDAM is a preliminary assessment tool that
22 contains items that are more specific to men’s symptoms (e.g. binge eating, muscle
23 dysmorphia). In research that compared the EDAM directly to the EDI-3, Stanford and
24 Lemberg (2012a) concluded that the EDI-3 does not capture the construct of *drive for*
25 *thinness* in males and that males are at risk of achieving a low score on this factor despite

1 exhibiting eating disorder symptoms. As such, EDI drive for thinness items such as “I am
2 preoccupied by the desire to be thinner” should be replaced with “I am preoccupied by the
3 desire to be lean and muscular.” Stanford and Lemberg’s research suggests that researchers
4 would do well to embrace the challenge of using male-specific methods and male-specific
5 research questions for a greater understanding of the male-specific body and eating concerns,
6 which appear markedly different to those of women and girls.

7 Athletes involved in elite sport were no different to those taking part in sports at a
8 lower standard. Thus, there was no overall support for the position that elite athletes may be
9 more at risk because of the greater competitive pressure to succeed and the more intense
10 training programs (Sundgot-Borgen 1994); equally, there was no overall support for the
11 finding that elite athletes may be less at risk (Darcy et al., 2013). Of course, this null finding
12 clearly warrants further research in light of the aforementioned male-specific arguments (i.e.,
13 measurement and specific sports).

14 There are several key limitations revealed through this meta-analysis, which should be
15 addressed in future research. First, many of the studies did not report whether the research
16 was conducted in or out of season. It has been argued that collegiate wrestlers only develop
17 transient disordered eating, which fluctuates over the course of a season (Dale & Landers,
18 1999); consequently, the timing of data collection for this sport may have influenced the
19 findings in either direction. Second, despite previous calls for more informative research
20 involving male athletes (Bryne & McLean, 2001; Lock, 2009), there was insufficient data
21 available to investigate age or experience as potential moderators. Finally, as mentioned
22 previously, although the way in which sports were coded was consistent with previous
23 research it was likely too simplistic for meaningful future comparisons and meta-analyses.
24 Specifically, we urge researchers to consider the detail of their sample(s) rather than to

1 categorise individuals within global sports according to simplistic categorisation criteria (e.g.
2 “endurance” sport).

3 In summary, in the studies contained in the current review, wrestling as a group of
4 sportsmen reported a higher incidence of disordered eating compared to non-athletic controls.
5 Only the EAT differentiated male athletes from non-athletes. The reliance on measurement
6 tools that were developed with women’s and girls’ disordered eating patterns in mind has
7 severely hampered this research field and there is a need to move beyond simple prevalence
8 studies that use such measures. Specifically, we urge researchers to use disordered eating
9 methodologies that are more specifically suitable for men and boys (e.g. Atkinson, 2011;
10 Papathomas & Lavalley, 2012; Stanford & Lemberg, 2012a).

11

12

References

Studies included in the meta-analysis are marked with an asterisk.

American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders* (5th ed.). Washington DC: Author. doi: 10.1176/appi.books.9780890425596.744053

Anderson, C. & Petrie, T. (2012). Prevalence of disordered eating and pathogenic weight control behaviors among NCAA division 1 female collegiate gymnasts and swimmers. *Research Quarterly for Exercise and Sport*, 83, 120-124.

Anshel, M. H. (2004). Sources of disordered eating patterns between ballet dancers and non-dancers. *Journal of Sport Behavior*, 27, 115-133.

Atkinson, M. (2011). Male athletes and the cult(ure) of thinness in sport. *Deviant Behavior*, 32, 224-256. doi: 10.1080/01639621003748860

Bardone-Cone, A. M., Abramson, L. Y., Vohs, K. D., Heatherton, T. F., & Joiner, T. E., Jr. (2006). Predicting bulimic symptoms: An interactive model of self-efficacy, perfectionism, and perceived weight status. *Behaviour Research & Therapy*, 44, 27-42. doi: 10.1016/j.brat.2004.09.009

*Boroughts, M., & Thompson, J. K. (2002). Exercise status and sexual orientation as moderators of body image disturbance and eating disorders in males. *International Journal of Eating Disorders*, 31, 307-311. doi: 10.1002/eat.10031

*Brugh, D. D., Williford, J. H., Chibucos, T. R., & Shewokis, P. A., (2001). The frequency of eating disorder tendencies among male high school wrestlers, non wrestler athletes, and non-athletes. *Journal of the American Dietetic Association*, 101, (Supp A86). doi: 10.1016/S0002-8223(01)80294-7

- 1 Byrne, S., & McLean, N. (2001). Eating disorders in athletes: A review of the literature.
2 *Journal of Science and Medicine in Sport*, 4, 145-159. doi: 10.1016/S1440-
3 2440(01)80025-6
- 4 *Byrne, S., & McLean, N. (2002). Elite athletes: Effects of the pressure to be thin. *Journal of*
5 *Science and Medicine in Sport*, 5, 80-94. doi: 10.1016/S1440-2440(02)80029-9
- 6 *Callejas, J. D., & Levine, A. M. (2003). Disordered eating risk among seventh grade male
7 athletes and non-athletes. *Journal of the American Dietetic Association*, 103,
8 September sup p. 112. doi: 10.1016/S0002-8223(08)70180-9
- 9 *Costarelli, V. & Stamou, D. (2009). Emotional intelligence, body image, and disordered
10 eating attitudes in combat sport athletics. *Journal of Exercise Science and Fitness*, 7,
11 104-111. doi: 10.1016/S1728-869X(09)60013-7
- 12 Croll, J. K., Neumark-Sztainer, D., Story, M., Wall, M., Perry, C., & Harnack, L. (2006).
13 Adolescents Involved in weight related and power team sports have better eating
14 patterns and nutrient intakes than non-sport involved adolescents. *Journal of the*
15 *American Dietetic Association*, 106, 709-717. doi: 10.1016/j.jada.2006.02.010
- 16 *Dale, K. S., & Landers, D. M. (1999). Weight control in wrestling: Eating disorders or
17 disordered eating? *Medicine and Science in Sport*, 31, 1382-1389.
- 18 *Darcy, A. M., Hardy, K. K., Lock, J., Hill, K. B., & Peebles, R. (2013). The Eating Disorder
19 Examination Questionnaire (EDE-Q) among university men and women at different
20 levels of athleticism. *Eating Behaviors*, 14, 378-381.
21 <http://dx.doi.org/10.1016/j.eatbeh.2013.04.002>
- 22 *Filaire, E., Rouviex, M., Bouget, M., & Pannafieux, C. (2007). Prevalence of eating
23 disorders in athletes. *Science & Sports*, 22, 135-142. doi:
24 10.1016/j.scispo.2007.03.002

- 1 Fisher, R.A. (1928). *Statistical Methods for Research Workers (2nd Edition)*. London: Oliver
2 & Boyd.
- 3 *Fogelholm, M, & Hiilloskorpi, H. (1999). Weight and diet concerns in Finnish female and
4 male athletes. *Medicine and Science in Sports and Exercise*, 31, 229-235.
- 5 *Fulkerson, J. A., Keel, P. K., Leon, G. R., & Dorr, T. (1999). Eating disordered behaviors
6 and personality characteristics of high school athletes and nonathletes. *International*
7 *Journal of Eating Disorders*, 26, 73-79. doi: 10.1002/(SICI)1098-108X
- 8 Garner, D. M. (1991). *Eating Disorder Inventory-2. Professional manual*. Odessa, FL:
9 Psychological Assessment Resources.
- 10 Garner, D. M., Olmstead, M. P., Bohr, Y., & Garfinkel, P. E. (1982). The Eating Attitudes
11 Test: Psychometric features and clinical correlates. *Psychological Medicine*, 12, 871-
12 878. doi: 10.1017/S0033291700049163
- 13 Haase, A. (2011). Weight perception in female athletes: Associations with disordered eating
14 correlates and behaviour. *Eating Behaviors*, 12, 64-67.
15 doi:10.1016/j.eatbeh.2010.09.004
- 16 Hausenblas, H. A., & Carron, A. V. (1999). Eating disorder indices and athletes: An
17 integration. *Journal of Sport and Exercise Psychology*, 21, 230-258.
- 18 *Hausenblas, H. A., & McNally, K. D. (2004). Eating disorder prevalence and symptoms for
19 track and field athletes and non-athletes. *Journal of Applied Sport Psychology*, 16,
20 274-286. doi: 10.1080/10413200490485630
- 21 Higgins, J., Thompson, S., Deeks, J. and Altman, D. (2003) Measuring inconsistency in
22 meta-analysis. *British Medical Journal*, 327, 557-560. doi: 10.1136/bmj.327.7414.557
- 23 Kotler, L.A., Cohen, P., Davies, M., Pine, D.S., & Walsh, B.T. (2001). Longitudinal
24 relationships between childhood, adolescent, and adult eating disorders. *Journal of the*

1 *American Academy of Child and Adolescent Psychiatry*. 40, 1434-1440. doi:

2 10.1097/00004583-2001120000-00014

3 *Krentz, M. & Warschburger, P. (2011) Sports-related correlates of disordered eating in

4 aesthetic sports. *Psychology of Sport and Exercise*, 12, 375-382. doi:

5 10.1016/j.psychsport.2011.03.004

6 Leit, R. A., Pope, H. G., & Gray, J. J. (2001). Cultural expectations of muscularity in men:

7 The evolution of playgirl centerfolds. *International Journal of Eating Disorders*, 29,

8 90-93. doi: 10.1002/1098-108X

9 Lock, J. D. (2009). Trying to fit square pegs in round holes: Eating disorders in males.

10 *Journal of Adolescent Health*, 44, 99-100. doi: 10.1016/j.jadohealth.2008.11.001

11 *Long-Anderson, S., Zager, K., Hetzler, R. K., Nahikian-Nelms, M., & Syler, G. (1996).

12 Comparison of Eating Disorder Inventory (EDI-2) scores of male bodybuilders to the

13 male college student subgroup. *International Journal of Sports Nutrition*, 6, 255-262.

14 doi: 10.1016/S0002-8223(95)00467-X

15 *Martinsen, M., Bratland-Sanda, S., Eriksson, A. K. & Sundgot-Borgen, J. (2010) Dieting to

16 win or be thin? A study of dieting and disordered eating among adolescent elite

17 athletes and non-elite controls. *British Journal of Sports Medicine*, 44, 70-76. doi:

18 10.1136/bjism.2009.068668

19 *Martinsen, M. & Sundgot-Borgen, J. (2013). Higher prevalence of eating disorders among

20 adolescent elite athletes than controls. *Medicine and Science in Sports and Exercise*,

21 45, 1188-1197. doi: 10.1249/MSS.0b013e318281a939

22 *Moss, L., & Bessinger, R. C. (1999). Disordered eating and body image perception in male

23 high school athletes and non-athletes. *Journal of the American Dietetic Association*,

24 99, Sep sup A-103. doi: 10.1016/S0002-8223(99)00761-0

- 1 *Nattiv, A., Puffer, J. C., & Green, G. A. (1997). Lifestyles and health risks of collegiate
2 athletes: A multi-center study. *Clinical Journal of Sport Medicine*, 7, 262-272.
- 3 *Nudelman, S., Rosen, J. C., & Leitenberg, H. (1988). Dissimilarities in eating attitudes,
4 body image distortion, depression, and self-esteem between high intensity male
5 runners and women with bulimia nervosa. *International Journal of Eating Disorders*,
6 7, 625-634. doi: 10.1002/1098-108X
- 7 Oliviardia, R., Pope, H. G., Borowiecki, J. J., & Cohane, G. H. (2004). Biceps and body
8 image: The relationship between muscularity and self-esteem, depression, and eating
9 disorder symptoms. *Psychology of Men and Masculinity*, 5, 112-120. doi:
10 10.1037/1524-9220.5.2.112
- 11 Papathomas, A., & Lavallee, D. (2012). Eating disorders in sport: A call for methodological
12 diversity. *Revista de Psicología del Deporte*, 21, 387-392.
13 <http://www.redalyc.org/articulo.oa?id=235126897020>
- 14 *Passman, L., & Thompson, J. K. (1988). Body image and eating disturbance in obligatory
15 runners, obligatory weightlifters, and sedentary individuals. *International Journal of*
16 *Eating Disorders*, 7, 759-769. doi: 10.1002/1098-108X
- 17 *Petrie, T. A. (1996). Differences between male and female collegiate lean sport athletes,
18 nonlean sport athletes, and nonathletes on behavioral and psychological indices of
19 eating disorders. *Journal of Applied Sport Psychology*, 8, 218-230. doi:
20 10.1080/10413209608406478
- 21 Petrie, T.A., Greenleaf, C., Carter, J., & Reel, J. (2007). Psychosocial corrolates of disordered
22 eating among male collegiate athletes. *Journal of Clinical Sport Psychology*, 1, 340-
23 357.
- 24 Pollock, N., Grogan, C., Perry, M., Pedlar, C., Cooke, K., Morrissey, D., & Dimitriou, L.
25 (2010). Bone-mineral density and other features of the female athlete triad in elite

1 endurance runners: A longitudinal and cross-sectional observational study.

2 *International Journal of Sport Nutrition and Exercise Metabolism*, 20, 418-426.

3 *Ravaldi, C., Vannacci, A., Zucchi, T., Mannucci, E., Cabras, P.L., Boldrini, M., Murciano,
4 L., Rotella, C.M., & Ricca, V. (2003). Eating disorders and body image disturbances
5 among ballet dancers, gymnasium users, and bodybuilders. *Psychopathology*, 36, 247-
6 254. doi: 10.1159/000073450

7 *Riebl, S.K., Subudhi, A. W., Broker, J. P., Schenck, K., & Berning, J. R. (2007). The
8 prevalence of subclinical eating disorders among male cyclists. *Journal of the*
9 *American Dietetic Association*, 107, 1214-1217. doi: 10.1016/j.jada.2007.04.017

10 *Rosendahl, J., Bormann, B., Aschenbrenner, K., Aschenbrenner, F. & Strauss, B. (2009).
11 Dieting and disordered eating in German high school athletes and non-athletes.
12 *Scandinavian Journal of Medicine and Science in Sports*, 19, 731-739. doi:
13 10.1111/j.1600-0838.2008.00821.x

14 Rosenthal, R. (1991). *Meta-analytic procedures for social research*. Sage: London.

15 *Rosenvinge, J. H., & Vig, C. (1993). Eating disorders and associated symptoms among
16 adolescent swimmers. *Scandinavian Journal of Medicine and Science in Sport*, 3,
17 164-169. doi: 10.1111/j.1600-0838.1993.tb00381.x

18 *Rouveix, M., Bouget, M., Pannafieux, C., Champely, S., & Filaire, E. (2007). Eating
19 attitudes body esteem, perfectionism and anxiety of judo athletes and nonathletes.
20 *International Journal of Sports Medicine*, 28, 340-345. doi: 10.1055/s-2006-924334
21 Smolak, L., Murnen, S. K., & Ruble, A. E. (2000). Female athletes and eating problems: A
22 meta-analysis. *International Journal of Eating Disorders*. 27, 371-380. doi:
23 10.1002/(SICI)1098-108X

- 1 Stanford, S. C., & Lemberg, R. (2012a). Measuring eating disorders in men: Development of
2 the Eating Disorder Assessment for Men (EDAM). *Eating Disorders: The Journal of*
3 *Treatment & Prevention*, 20, 427-436. doi: 10.1080/10640266.2012.715522
- 4 Stanford, S. C., & Lemberg, R. (2012b). A clinical comparison of men and women on the
5 Eating Disorder Inventory-3 (EDI-3) and the Eating Disorder Assessment for Men
6 (EDAM). *Eating Disorders*, 20, 379-394. doi: 10.1080/10640266.2012.715516
- 7 *Stanford-Martens, T. C., Davidson, M. M., Yakushko, O. F., Martens, M., Hinton, P., &
8 Beck, N. (2005). Clinical and subclinical eating disorders: An examination of
9 collegiate athletes. *Journal of Applied Sport Psychology*, 17, 79-86. doi:
10 10.1080/10413200590907586
- 11 Striegel-Moore, R. H., & Bulik, C. M. (2007). Risk factors for eating disorders. *American*
12 *Psychologist*, 62, 181-198. doi: 10.1037/0003-066X.62.3.181
- 13 Sundgot-Borgen, J. (1994). Risk and trigger factors for the development of eating disorders in
14 female elite athletes. *Medicine and Science in Sports and Exercise*, 26, 414-419.
- 15 *Sundgot-Borgen, J., & Torstveit, M. K. (2004). Prevalence of eating disorders in elite
16 athletes is higher than in the general population. *Clinical Journal of Sport and*
17 *Medicine*, 14, 25-37.
- 18 *Theil, A., Gotffried, H., & Hesse, F. W. (1993). Subclinical eating disorders in male
19 athletes. *Acta Psychiatrica Scandinavica*, 88, 259-265. doi: 10.1111/j.1600-
20 0447.1993.tb03454.x
- 21 *Wheeler, G. D., Wall, S. R., Belcastro, A. N., Conger, P., & Cumming, D. C. (1986). Are
22 anorexic tendencies prevalent in the habitual runner? *British Journal of Sports*
23 *Medicine*, 20, 77-81. doi: 10.1136/bjism.20.2.77

- 1 *Wichstrom, L., Skogen, K., & Oia, T. (1994). Social and Cultural Factors Related to Eating
2 Problems Among Adolescents in Norway. *Journal of Adolescence*, 17, 471-482. doi:
3 10.1006/jado.1994.1040
- 4 *Wilkins, J. A., Boland, F. J., & Albinson, J. A. (1991). Comparison of male and female
5 university athletes and nonathletes on eating disorder indices: Are athletes protected?
6 *Journal of Sport Behavior*, 14, 129–143.
- 7 Woodman, T., & Hemmings, S. (2008). Body image self-discrepancies and affect: Exploring
8 the feared body self. *Self & Identity*, 7, 413-429.
- 9 Woodman, T., & Steer, R. (2011). Body self-discrepancies and women's social physique
10 anxiety: The moderating role of the feared body. *British Journal of Psychology*, 102,
11 147-160.

Table 1: *Characteristics of samples used in the meta-analysis*

Author (Year)	<i>r</i> Value	Sport	Elite	Athletic Sample	Control Sample	Dependent Variable
Boroughs & Thompson (2002)	.07	Running/Bodybuilding	N	72	62	Dthin
Brugh et al. (2001)	.38	Wrestling/Non-wrestling athletes	N	261	60	EATtot
Bryne & McLean (2002)	.30	Thin/normal build sports	Y	161	55	Dthin
Callejas & Levine (2003)	.16	Varied	N	67	37	EATtot
Costarelli & Stamou (2009)	.00*	Martial arts	Y	6	8	EATtot
Dale & Landers (1999)	.14	Wrestling	N	85	75	Dthin
Darcy et al. (2013)	-.01	Varied	N	597	64	EDE-Q
Filaire et al. (2007)	.22	Judo/Cycling	N	27	17	EATtot
Fogelholm & Hiilloskorpi (1999)	.08	Varied**	Y	190	61	Dthin
Fulkerson et al. (1999)	.13	Varied	N	174	195	Dthin
Hausenblas & McNally (2004)	.09	Athletics	N	76	98	Dthin
Krentz & Warschburger (2011)	.23	Aesthetic sports	Y	35	35	EATtot
Long-Anderson et al. (1996)	-.15	Bodybuilding	N	68	50	Dthin

Author and Year	<i>r</i> Value	Sport	Elite	Athletic Sample	Control Sample	Dependent Variable
Martinsen et al. (2010)	-.08	Varied	Y	389	197	Dthin
Martinsen et al. (2013)	-.01	Varied	Y	274	82	EDE-Q
Moss & Bessinger (1999)	.08	Varied	N	203	340	EATtot
Nattiv et al. (1997)	.05	Varied	N	1502	289	SC-Wc
Nudleman et al. (1988)	-.13	Running	N	20	20	Dthin
Passman & Thompson (1988)	.52	Running/Bodybuilding	N	30	15	Dthin
Petrie (1996)	-.07	Lean/Non-lean sports	N	187	230	Dthin
Ravaldi et al. (2003)	.10	Bodybuilding	N	44	30	EDE-Dr
Riebl et al. (2007)	.42	Cycling	Y	61	63	EATtot
Rosendahl et al. (2009)	.03	Varied	N	366	122	EATtot
Rosenvinge & Vig (1993)	.00*	Swimming	Y	19	20	EATtot
Rouveix et al. (2007)	-.47	Judo	Y	12	17	EATtot
Stanford-Martens et al. (2005)	-.06	Lean/Non-lean sports	N	164	55	Q-EDDtot
Sundgot-Borgen & Torstveit (2004)	.06	Varied**	Y	687	629	Dthin
Theil et al. (1993)	.09	Wrestling/Rowing	Y	84	104	Dthin
Wheeler et al. (1986)	.00*	Running	Y	49	18	EATtot

Author and Year	<i>r</i> Value	Sport	Elite	Athletic Sample	Control Sample	Dependent Variable
Wichstrom et al. (1994)	.12	Varied**	N	1802	107	EATfp
Wilkins et al. (1991)	-.20	Varied	N	99	39	Dthin

Note: *r* is Fisher's (1928) transformed *Zr*, where a positive score indicates that the athletic group scored higher than the control group on the diagnostic questionnaire; - a negative score indicates that the control group scored higher; for Elite, Y = yes and N = no; Dthin = the drive for thinness subscale of the Eating Disorder Inventory, EATtot = the total score on the Eating Attitudes Test, SC-Wc = the weight concern subscale of a diagnostic tool developed for the research, EDE-Dr = the dietary restraint subscale of the Eating Disorder Enquiry Questionnaire, Q-EDDt = total score for the Questionnaire for Eating Disorder Diagnosis, EATfp = the food preoccupation subcategory of the Eating Attitudes Test; * = non-significant results were reported and included in the analysis as *r* = 0 based on Rosenthal's (1991) recommendation. ** = The study contained more than one experimental group allowing for additional effect size *r* values to be calculated and used in the moderator analyses. All reported *p* levels associated with *z* scores are based on two-tailed tests.

Figure 1: Forest plot of the effect sizes that were in the overall analysis.