

### Sediment Composition Influences Spatial Variation in the Abundance of Human Pathogen Indicator Bacteria within an Estuarine Environment

Perkins, T.L.; Clements, K.; Baas, J.H.; Jago, C.F.; Jones, D.L.; Malham, S.K.; McDonald, J.E.

### PLoS ONE

DOI: 10.1371/journal.pone.0112951

Published: 14/11/2014

Publisher's PDF, also known as Version of record

Cyswllt i'r cyhoeddiad / Link to publication

Dyfyniad o'r fersiwn a gyhoeddwyd / Citation for published version (APA): Perkins, T. L., Clements, K., Baas, J. H., Jago, C. F., Jones, D. L., Malham, S. K., & McDonald, J. E. (2014). Sediment Composition Influences Spatial Variation in the Abundance of Human Pathogen Indicator Bacteria within an Estuarine Environment. *PLoS ONE*, 9(11). https://doi.org/10.1371/journal.pone.0112951

Hawliau Cyffredinol / General rights Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

• Users may download and print one copy of any publication from the public portal for the purpose of private study or research.

- You may not further distribute the material or use it for any profit-making activity or commercial gain
   You may freely distribute the URL identifying the publication in the public portal ?

Take down policy If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

## Sediment Composition Influences Spatial Variation in the Abundance of Human Pathogen Indicator Bacteria within an Estuarine Environment



# Tracy L. Perkins<sup>1</sup>, Katie Clements<sup>2</sup>, Jaco H. Baas<sup>2</sup>, Colin F. Jago<sup>2</sup>, Davey L. Jones<sup>3</sup>, Shelagh K. Malham<sup>2,4</sup>, James E. McDonald<sup>1</sup>\*

1 School of Biological Sciences, Bangor University, Bangor, United Kingdom, 2 School of Ocean Sciences, Bangor University, Bangor, United Kingdom, 3 School of Environment, Natural Resources and Geography, Bangor University, Bangor, United Kingdom, 4 Centre for Applied Marine Sciences, Bangor University, Bangor, United Kingdom

### Abstract

Faecal contamination of estuarine and coastal waters can pose a risk to human health, particularly in areas used for shellfish production or recreation. Routine microbiological water quality testing highlights areas of faecal indicator bacteria (FIB) contamination within the water column, but fails to consider the abundance of FIB in sediments, which under certain hydrodynamic conditions can become resuspended. Sediments can enhance the survival of FIB in estuarine environments, but the influence of sediment composition on the ecology and abundance of FIB is poorly understood. To determine the relationship between sediment composition (grain size and organic matter) and the abundance of pathogen indicator bacteria (PIB), sediments were collected from four transverse transects of the Conwy estuary, UK. The abundance of culturable Escherichia coli, total coliforms, enterococci, Campylobacter, Salmonella and Vibrio spp. in sediments was determined in relation to sediment grain size, organic matter content, salinity, depth and temperature. Sediments that contained higher proportions of silt and/or clay and associated organic matter content showed significant positive correlations with the abundance of PIB. Furthermore, the abundance of each bacterial group was positively correlated with the presence of all other groups enumerated. Campylobacter spp. were not isolated from estuarine sediments. Comparisons of the number of culturable E. coli, total coliforms and Vibrio spp. in sediments and the water column revealed that their abundance was 281, 433 and 58-fold greater in sediments (colony forming units (CFU)/100 g) when compared with the water column (CFU/100 ml), respectively. These data provide important insights into sediment compositions that promote the abundance of PIB in estuarine environments, with important implications for the modelling and prediction of public health risk based on sediment resuspension and transport.

Citation: Perkins TL, Clements K, Baas JH, Jago CF, Jones DL, et al. (2014) Sediment Composition Influences Spatial Variation in the Abundance of Human Pathogen Indicator Bacteria within an Estuarine Environment. PLoS ONE 9(11): e112951. doi:10.1371/journal.pone.0112951

Editor: Newton C M. Gomes, University of Aveiro, Portugal

Received July 17, 2014; Accepted October 17, 2014; Published November 14, 2014

**Copyright:** © 2014 Perkins et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability: The authors confirm that all data underlying the findings are fully available without restriction. All relevant data are within the paper.

**Funding:** This research was supported by a Knowledge Economy Skills Scholarship (KESS), part-funded by the European Social Fund (ESF) through the European Union's Convergence program administered by the Welsh Government in association with Dŵr Cymru Cyf/Welsh Water Ltd. (http://www.higherskillswales.co.uk/ kess/index.php.en?menu = 0& catid = 0), and a Natural Environment Research Council consortium grant under the Macronutrient Cycles Research Programme (NE/ J011908/1) (http://www.nerc.ac.uk/research/funded/programmes/macronutrient/#xcollapse3). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

\* Email: j.mcdonald@bangor.ac.uk

### Introduction

Estuarine environments represent some of the most biologically productive systems in the biosphere and consequently provide a wealth of economic, social and natural ecosystem services that include food, employment, recreation and habitat [1]. However, the sustainability of such systems can be severely compromised along developed and urbanised coastlines, and this is predominantly due to anthropogenic influences [2]. Almost half of the world's population are thought to live within a few hundred kilometers of the coast [3] and consequently, anthropogenic activities have a significant influence upon the health of estuarine and ocean ecosystems. Furthermore, as the global climate changes, meteorological events such as storms and floods present further impacts upon estuarine environments, and increased rainfall will significantly impact the flow and transportation of microbial pollution from the terrestrial environment into the coastal zone [4]. Human pathogenic microorganisms are introduced into estuarine ecosystems via the release of effluent from wastewater treatment plants, ineffective septic tank systems and storm water runoff [5]. Agricultural runoff from livestock farming can also represent a major source of microbial pollution, particularly when excreted waste from poultry and livestock is re-applied to land [6]. Wildlife, especially migratory wildfowl and other birds also represent an important source of zoonotic bacterial pathogens in natural environments [7].

Human pathogenic bacteria often occur at low levels in the environment [8] and their specific detection is a laborious and costly process [9]. Consequently, the detection and enumeration of faecal indicator bacteria that are present in the gastrointestinal tract of warm-blooded animals in much higher quantities represent a more effective 'indicator' of risk [10]. Current legislation on water quality testing relies on the enumeration of faecal coliforms (including the well-characterised E. coli) and enterococci to assess and classify water quality [11]. However, several other groups of potentially pathogenic bacteria have been detected in estuarine and marine environments. These include Salmonella spp. [12], Campylobacter spp. [13] and Vibrio spp. [14]. Salmonella spp. are well-known foodborne pathogens and recognised as one of the main causes of gastroenteritis in humans [15], resulting in around 1.3 billion annual cases of infection worldwide [16] but their presence in bathing waters is not routinely monitored [17]. Pathogenic strains of Campylobacter spp. are also thought to be responsible for a large proportion of enteric illnesses in humans living in developed and industrialized countries [18]. Contamination of surface and coastal waters by Salmonella and Campylobacter spp. is thought to occur predominantly via faecal contamination from wildfowl and other birds, although other animals including domestic livestock are also reservoirs [19]. Vibrio spp. are ubiquitous in aquatic environments and some strains are pathogens of humans; V. cholera has been responsible for several previous pandemics, resulting in high human mortality rates [20], some of which originated from the consumption of contaminated seafood [21]. Consequently, filter-feeding shellfish such as mussels, scallops and oysters are especially susceptible to contamination with bacterial pathogens if grown in contaminated waters [22].

Spatial and temporal variations in bacterial abundance in an environment are controlled by the interactions of complex physical, chemical and biological parameters, such as available nutrients [23], organic matter [24], sediment grain size [25], clay content [26], heavy metal content [27], predation by protozoa [28], competition [29], temperature [30], salinity [31], sunlight intensity [32] and seasonal variations [33]. Consequently, association with particulate material and sediments offers several advantages in terms of the survival and persistence of bacterial pathogens, such as protection from UV light [34,35], protection from phage attack in saline conditions [36], shelter from predation [37] and a greater organic matter content compared to the water column [38]. However, hydrodynamic processes (e.g. tides and storms), recreational activities and mechanical disturbances such as commercial dredging all have the potential to re-suspend sediment particles and their associated pathogens back into the water column, resulting in periodic elevated levels of pollution [39,40]. Currently, classification of both bathing and shellfish water quality relies solely upon the enumeration of FIB in water samples, despite the well-established paradigm that allochthonous bacterial pathogens in the water column become preferentially attached to particulate material, which promotes their downward flux to the bottom sediments where they are typically found in greater abundance [41,42].

The Conwy catchment, North Wales, UK, has a population of approximately 112,000, 80% of which live in coastal resorts that represent the main economic and tourist areas, with over 5.4 million visitors per annum. The Conwy estuary directly impacts commercially important shellfish beds, bathing waters and beaches; hence deterioration of the water quality could have major socioeconomic consequences. The catchment covers an area of  $\sim$ 300 km<sup>2</sup> which consists of large areas of land utilised for agriculture [43], some of the highest mountains in the UK [44], in addition to residential and commercial areas. Consequently, the Conwy river and it's tributaries (such as the river Gyffin and river Ganol) receive wastewater effluent from several waste water treatment plants in addition to septic tank discharges. Microbio-

logical inputs from wildlife, agriculture, wastewater effluent, septic tank discharges and run-off from storm events therefore represent potential point and diffuse sources of FIB in the Conwy estuary.

Previous analysis of FIB concentrations in the water column across four transverse transects of the Conwy estuary revealed significant spatial variations in waterborne E. coli numbers on contrasting sides of the estuary (45). These data suggested that significantly different levels of microbial pollution were present in the east and west sides of the river (45) and consequently, the choice of sampling points for routine water quality monitoring could have a significant impact upon the classification of microbiological water quality in this area. The Conwy estuary has a dynamic tidal cycle, with resuspension and deposition of finer sediment particles observed on the shallower banks of the river, resulting in the formation of mud flats. Consequently, the localised resuspension of sediment associated FIB into the water column was one proposed explanation for such contrasting FIB counts in the water column of the Conwy estuary (45), yet the spatial relationships between sediments composition (grain size and organic matter content) and FIB abundance in the Conwy and beyond has received little attention.

Whilst most studies focus on enumeration of E. coli and enterococci in water samples for the assessment of public health risk, the aim of our study was to address the paucity of information regarding the ecology of FIB along with other potentially humanpathogenic bacteria in estuarine sediments, and to identify sediment characteristics that promote pathogen abundance in estuarine environments. A second aim was to determine if the enumeration of E. coli, coliforms and enterococci, that represent the current standard 'indicator' organisms for faecal contamination and water quality monitoring, are a suitable indicator for the co-occurence of other potentially pathogenic bacterial groups. This was achieved by determining spatial variations in the abundance of culturable E. coli, total coliforms, enterococci, Salmonella, Campylobacter and Vibrio spp. (to be used as a proxy for potential human bacterial pathogens in the estuary and referred to as 'pathogen indicator bacteria' (PIB) herein) in estuarine sediments of the Conwy Estuary, UK, in relation to sediment grain size, organic content and other physico-chemical factors.

#### **Materials and Methods**

#### Study site

Sample locations were selected to correspond to those tested in a previous study that observed spatial variations of waterborne E. *coli* within the study area [45], in addition to possible point and diffuse sources of pollution that include mudflats used by wading birds and agricultural land (transect 1), input from the river Ganol (transect 2), the river Gyffin (transect 3) and Conwy marina (transect 4), (Figure 1).

#### Estuarine sampling

Estuarine sediment and water samples were collected from four transverse transects of the estuary over two consecutive sampling days (transects 1 and 2 collected on the 14/05/12, transects 3 and 4 on the 15/05/12). The Bangor University research vessel "Macoma", a 7.9 m Cheetah catamaran survey boat was utilised for sample collection, which was initiated at slack water. For each sampling point, three replicate sediment samples were collected using a manually operated Van Veen sediment grab, approximately 50 g of bottom sediment from the centre of each grab was transferred into sterile 50 ml polypropylene tubes (VWR International Ltd., Leicestershire, UK), water samples were collected in



Figure 1. A map of the study site; the Conwy Estuary, North Wales, UK. Water and sediment samples were collected in triplicate from four transverse transects of the Conwy estuary (twenty one sampling sites). doi:10.1371/journal.pone.0112951.g001

triplicate from 0.2 m below the water surface using a sterile 1 L polypropylene container, then transported to the laboratory where water samples were processed within 4 h and sediment samples within 6 h.

## Isolation and cultivation of target bacterial groups from sediment

One gram of each sediment sample was transferred to a 7 ml sterile bijou tube (Starlab UK Ltd., Milton Keynes, UK) and suspended in 5 ml Ringers solution (Oxoid Ltd., Basingstoke, UK) to obtain a 1:5 (w/v) dilution. Each sample was vortexed for 90 s to disassociate and resuspend bacteria from the sediment. Aliquots of the resulting supernatant for each sample were transferred

aseptically onto agar plates containing a selective medium for *E. coli*, total coliforms, enterococci, *Vibrio* spp., *Campylobacter* spp., *Salmonella* spp. and total heterotrophs. The optimum volume of supernatant used to inoculate each selective medium was determined in a previous study (data not shown). All selective medium plates were inverted and incubated according to manufacturer's recommendations. Resulting colony forming units (CFUs) provided enumeration of bacterial groups. Details describing the selective media used and incubation times are described in Table S1 in the supplemental material. The CFU data for sample point 13 represents the average of only two sediment samples due to one of the replicates being lost as a result

of the presence of gravel clasts. All other CFU data represent the average of three independent replicate samples.

# Isolation and cultivation of target bacterial groups from water

Water samples were processed within 4 h in accordance with the Revised Bathing Water Directive (2006/7/EC). Enumeration of bacteria in water samples was achieved by using vacuum– filtration as described in [45]. Briefly, water samples were homogenised by shaking and 50 ml of water filtered under vacuum through a 0.2  $\mu$ m cellulose acetate membrane (Sartorius Stedim Biotech., Gottingen, Germany). Subsequently, the membranes were aseptically transferred onto sterile agar plates containing selective medium for the enumeration of *E. coli*, total coliforms, *Vibrio* spp. and heterotrophic bacteria. Agar plates were inverted and incubated according to manufacturer's recommendations (Table S1). Resulting Colony Forming Units (CFUs) were enumerated 24 h post incubation.

#### Sediment particle size analysis

Sediment grain size was determined by laser diffraction after 1 min sonication to separate particles, using a Malvern Hydro 2000 MU particle size analyser in conjunction with the Mastersizer 2000 software. Three replicate sediment samples from each site were pooled and homogenised. Approximately 1 g of sediment was added to the particle size analyser and 3 independent size determinations were made. This was repeated 3 times using the same pooled sample to determine an overall average.

### Determination of sediment organic matter content

The loss on ignition method (LOI) was used to determine organic matter content of sediment samples. Three replicate sediment samples from each site were pooled and homogenised. Approximately 20 g of fresh sediment from each sample was placed in a pre-weighed crucible and dried at  $95^{\circ}$ C for 24 h. Approximately 4 g of the resultant dried sediment was pre-weighed, transferred to another crucible and placed into a muffle furnace at  $550^{\circ}$ C for 4 h. Organic matter content was calculated as the difference between the weight of the dry sediment and weight of the residue post-combustion. This was repeated 3 times using the same pooled sample to determine an overall average. Moisture content per g of fresh sediment was determined by calculating the percentage difference between wet weight and dry weight after 24 h at  $95^{\circ}$ C (Table S2).

# *In situ* physico-chemical measurements of estuarine transect sample sites

Conductivity (calculated to practical salinity units (PSU)), temperature and depth measurements were recorded *in situ* using a YSI 600LS CTD scanner attached to a YSI 650 MDS data logger. This was deployed at each sample site in parallel with collection of water and sediment samples. Salinity and temperature measurements were recorded from the water immediately above the sediment bed and 0.2 m below the water surface. (Conductivity readings were calculated to determine salinity) (Table S3).

### DNA extraction, 16S rRNA gene PCR and sequencing of bacterial isolates

To validate the identity of bacterial colonies on selective microbiological medium, DNA was extracted from a total of 30 isolates that matched or were similar to the expected colony morphology/phenotype for *E. coli*, enterococci, *Campylobacter* and *Vibrio spp*. for 16S rRNA gene PCR amplification and sequencing. For *E. coli* and enterococci, little variation in colony morphology and phenotype was observed across all of the colonies counted, and so only a small number of isolates were sequenced for confirmation.

DNA was extracted from 30 isolated bacterial colonies using the ISOLATE genomic DNA mini kit (Bioline Reagents Ltd., London, UK) following the manufacturer's protocol. Agarose gel electrophoresis (1%) was used to visualise the DNA extracted from each bacterial isolate. Subsequently, the 16S rRNA gene of each isolate was amplified via PCR using the primer pair pA (5'-AGAGTTTGATCCTGGCTCAG-3') and pH (5'-AAGGAG-GTGATCCAGCCGCA-3') (45). PCR reactions consisted of 10 pmol of each primer (pA and pH), 1x MyTaq red mix (Bioline), approximately 100 ng of template DNA and ddH<sub>2</sub>O to a total reaction volume of 50 µl. The PCR conditions were as follows: an initial denaturation step at 95°C for 1 min, followed by 35 cycles of 95°C for 15 s, 55°C for 15 s, 72°C for 10 s and a final hold at 4°C. PCR amplicons were visualised using 1% agarose gel. The expected 16S rRNA gene amplicon size was approximately 1500 bp and amplicons of the expected size were subsequently excised from the agarose gel and purified using the Isolate PCR and Gel Kit (Bioline Reagents Ltd., London, UK) following the manufacturer's instructions. Purified 16S rRNA gene amplification products for each bacterial isolate were sequenced in both the forward and reverse orientation by Macrogen Europe (Netherlands). The Geneious Pro software package, Geneious 6.1 version created by Biomatters, (Available from http://www.geneious. com/) was used to quality clip each sequence and assemble the forward and reverse reads of each strain into a contiguous sequence. The sequence identity of each contig was determined using NCBI BLASTn. Identification of sequenced strains is given in Table S4 of the supplemental material.

#### Statistical analysis

Using the Statistical Package for Social Sciences SPSS v20, (IBM Corp., Armonk. NY), basic correlations were performed using the average data calculated for each site to determine the relationships between cultured bacterial abundance in sediments and water with different tested parameters. The non-parametric Spearman Rank Correlation Coefficient ( $r_s$ ) was used due to the data being not normally distributed.

#### Results

### The relationship between PIB abundance, sediment grain size and organic content

There were marked spatial differences in the abundance of sediment-associated PIB across all of the 21 sample sites within the estuary as determined by culture counts on selective medium (Table 1). Mean densities of E. coli and total coliforms in sediments ranged from 0 to  $2.4 \times 10^4$  CFU/100 g and 0 to  $5.4 \times 10^5$  CFU/100 g wet weight, respectively (for all mean bacterial densities from all sampling sites see Table S5 in the supplemental material). Enterococci and Salmonella abundance varied from 0 to  $1 \times 10^4$  CFU/100 g and 0 to  $2.5 \times 10^4$  CFU/100 g wet weight, respectively. Vibrio spp. were detected at all 21 sample sites  $(6.7 \times 10^3 \text{ to } 1.2 \times 10^6 \text{ CFU}/100 \text{ g wet weight})$  and this reflects their status as indigenous members of marine and aquatic environments. In addition, direct colony counts on Campylobacter selective media indicated the presence of 0 to  $3.5 \times 10^4$  CFU/100 g wet weight. However, 16S rRNA gene sequencing of putative Campylobacter spp. isolated from a subsequent sampling survey on the same selective medium revealed that none of the sequenced isolates were Campylobacter spp. (Table S4). Culturable heterotrophic bacteria were enumerated as a proxy for the abundance of the indigenous estuarine microbial community and their abundance ranged from  $6.6 \times 10^4$  to  $8.7 \times 10^6$  CFU/100 g wet weight. The enumeration of culturable heterotrophic bacteria alongside our target PIB groups enabled analysis of the relationship between total culturable heterotrophic bacteria counts and PIB counts in sediments, these data demonstrate that the abundance of heterotrophic bacteria within the sediments showed significant positive correlations with the abundance of all PIB groups measured (Table S6). Sediment grain size composition within sediment samples ranged from clay (0 to 18%), silt (0 to 65%), very fine sand (0 to 15%), fine sand (3 to 67%) medium sand (0 to 59%), coarse sand (0 to 17%) and very coarse sand (0 to 16%). The organic matter content of the sediment samples varied from 0.3 to 6% across the sample sites (Figure 2).

The abundance of each cultured bacterial group within the sediments showed significant positive correlations with the abundance of all other measured bacterial groups (Table S6). Vibrio spp. were also more abundant in sediments that had higher densities of FIB. In addition, the abundance of all isolated PIB groups showed a significant positive correlation with both sediment clay content (grain size  $<4 \ \mu m$ ) (E. coli, enterococci, total coliforms, and *Vibrio* spp.  $r_s = 0.543$  (p<0.011)  $r_s = 0.664$  (p< 0.001),  $r_s = 0.495$  (p<0.023),  $r_s = 0.663$  (p<0.001) respectively) and silt content (grain size 4 µm-63 µm) (E. coli, enterococci, total coliforms and *Vibrio* spp.,  $r_s = 0.570$  (p<0.007)  $r_s = 0.687$  (p< 0.001),  $r_s = 0.547$  (p<0.010),  $r_s = 0.688$  (p<0.001) respectively). Significant negative correlations were observed between the abundance of all PIB groups with fine sand  $(125 \,\mu\text{m}-250 \,\mu\text{m})$ and medium sand (250 µm-500 µm) (Table S7). Sediments with high amounts of clay and silt along with very fine sand contained the greatest proportion of organic material (significant positive correlation; organic matter content and clay,  $r_s = 0.917$  (p<0.001). organic matter content and silt,  $r_s = 0.926$  (p<0.001), organic matter content and very fine sand,  $r_s = 0.810$  (p<0.001). Significant negative correlations were evident between organic matter content and fine sand and also organic matter content and medium sand (Table S8). Sediments with high organic matter therefore also showed significant positive correlations with the abundance of all isolated PIB (Table S9).

# Comparison of PIB abundance between sediment and water samples

The average abundance data for culturable PIB (*E. coli*, total coliforms and *Vibrio* spp.) in sediment and water samples across all 21 sample sites revealed that *E. coli*, coliforms and *Vibrio* spp. were 281, 433 and 58-fold more abundant in the sediment (CFU/100 g) than the water column (CFU/100 ml), respectively (Table 1).

## The effect of depth, salinity and temperature on PIB abundance

Collection of water samples was initiated at slack water, with a range of water column depths from 0.5 to 12.3 m across all sites. Salinity and temperature measurements taken from above the bottom sediment varied between 9.3 to 30.0 psu and 10.8 to 11.2°C, respectively. At 0.2 m below the water surface the temperature varied from 10.8 to 11.2°C and salinity ranged from 10.1 to 27.8 psu. Bacterial abundance within the sediments revealed no significant correlations with any of the physicochemical parameters measured (Table S10). *Vibrio* spp. enumerated from the water samples were the only cultured species to show significant correlations with depth, temperature or salinity. Temperature had a significant negative correlation with the abundance of *Vibrio* spp. ( $r_s = 0.614 \text{ p} < 0.003$ ), while salinity had a significant positive correlation with abundance ( $r_s = 0.544 \text{ p} < 0.011$ ) (Table S11).

### Taxonomic identification of bacterial isolates using 16S rRNA gene sequencing

None of the isolates derived from the selective medium for *Campylobacter* (n = 7) were positively identified as the target group. However, 91% of *E. coli* (n = 11), 100% of enterococci (n = 3) and 67% of *Vibrio* (n = 9) isolates were positively identified as the target group (Table S4). No sequence data were obtained for *Salmonella* isolates and they should therefore be considered as presumptive *Salmonella* spp.

### Discussion

Results from this study demonstrate that sediments represent a significant reservoir for PIB and that sediment characteristics (grain size and organic matter content) influence spatial variations in PIB abundance in estuarine environments. Here, clay and silt fractions comprised a higher organic matter content in comparison to medium and coarse sand and harboured significantly higher densities of PIB. Previous studies have also concluded that the presence of finer sediment particles can have a positive impact on FIB abundance [25,46] and that organic material is a contributing factor to the prolonged persistence and survival of FIB in sediments [24,47,48]. However, almost all of the previous studies focus specifically on 'classic' faecal indicators such as *E. coli*, coliforms and enterococci, and here we have expanded knowledge on comparative PIB–sediment interactions with sediments for a greater repertoire of bacterial groups.

Results revealed there was not only a large spatial variation in PIB abundance that correlated to sediment grain size and organic matter content, but PIB abundance was also found to be significantly higher in the sediments when compared to the overlying water column, with levels of *E. coli* reaching over 3 orders of magnitude higher in the sediment. It is well documented that bacterial association with sediments circumvents the negative

**Table 1.** Average abundance of culturable pathogen indicator bacteria in sediment (CFU/100 g wet weight) and water (CFU/ 100 ml) across 21 estuarine sample sites.

Bacterial group	Sediment	Water	Fold - difference
E. coli	5.9×10 <sup>3</sup>	2.1×10 <sup>1</sup>	281
Total coliforms	1.3 x10 <sup>5</sup>	3.0×10 <sup>2</sup>	433
Vibrio spp.	4.5×10 <sup>5</sup>	7.8×10 <sup>3</sup>	58

doi:10.1371/journal.pone.0112951.t001



**Figure 2.** Bacterial abundance (CFU/100 g wet weight) compared to sediment grain size, organic matter content in sediments and bacterial abundance in the water column (CFU/100 ml), across four transverse transects. (A) Transect 1, (B) Transect 2, (C) Transect 3, (D) Transect 4. The X-axis represents sample points (n = 3 replicate samples for A, B, C and D except for sediment samples site 13, n = 2), mean values are plotted and error bars represent the SEM). doi:10.1371/journal.pone.0112951.g002

effects of environmental stresses, such as providing protection from UV light (34), which in turn can augment the survival and even growth of FIB. For example, DNA fingerprinting analyses performed on E. coli populations in beach sand and sediment revealed the possibility that some strains may have become naturalized to this environment [52] and growing evidence also suggests that there may be free-living strains of FIB surviving and multiplying within the water column of environmental waters, independently of a host [53].

The abundance of E. coli and enterococci in sediments had a significant positive correlation with the abundance of other PIB. Consequently, the enumeration of FIB as the 'classic' bacterial indicators currently conducted as part of routine water quality monitoring represent suitable indicators for the presence of other PIB groups. E. coli and enterococci are the predominant indicator species for monitoring faecal pollution of aquatic environments within the European Union (EU) despite numerous studies highlighting the differential environmental survival of different FIB groups [48,51], in addition to different strains of the same group [54]. In this study, the presence and abundance of enterococci had a stronger significant positive correlation with silt and clay when compared with other FIB, which may suggest greater survival times of enterococci under favourable conditions that certain sediment types may provide. However, it is well established that the survival times of different species, and even strains of the same species, varies considerably in aquatic environments, both between and within species, and this must be taken into consideration. For example, in comparison to other FIB, enterococci survive for longer periods in sediments [48] and within estuarine environments [31], and it has been proposed that the increase in survival under harsh conditions may be due to their membrane composition [55,56]. Such evidence suggests that enterococci may be a more robust indicator of the persistence of faecal contamination rather than more recent contamination within coastal and estuarine environments.

The enumeration of total heterotrophs revealed significant positive correlations with all other cultured PIB groups and therefore the same trends were seen within the indigenous microbial community (Table S6). It should be noted that despite obtaining high colony counts on selective microbiological media for Campylobacter spp., 16S rRNA gene sequence analysis of some of these isolates suggested that none were *Campylobacter* spp. These data highlight the potential pitfalls of culture-based analyses of microbial taxa using a selective microbiological medium, and care must be taken when interpreting microbial culture counts. Despite this, the sequenced isolates of enterococci, Vibrio spp. and E. coli indicate that these media were selective for the desired bacterial target groups (Table S4). Little variation in temperature and salinity could be explained by the small geographical variation along each transect. There was no significant correlation between temperature and salinity and the abundance of PIB, with the exception of Vibrio spp. enumerated from the water column. Here, Vibrio spp. had a significant positive correlation with salinity and a significant negative correlation with temperature, supporting previous proposals that temperature and salinity have important implications for Vibrio spp. population dynamics [30]. The depth at which sediment samples were taken had no impact on the abundance of PIB enumerated from the sediments.

This study highlights the risk of periodic elevated FIB and other PIB in the water column due to the resuspension of microbial contaminated sediments. Furthermore, the risk of microbial pollution from the resuspension of sediments is not taken into consideration when assessing microbial pollution of recreational and shellfish harvesting waters. The time and place of sampling in addition to tidal and hydrodynamic conditions may impact upon FIB concentrations in the water column. Despite several reports on the importance of sediments as a reservoir for FIB [28,29], the enumeration of waterborne FIB has received much more attention. Quilliam *et al.*, (2011) revealed significant spatial variations in waterborne *E. coli* numbers on contrasting sides of the same four transverse transects of the Conwy estuary studied here, indicating that significantly different levels of microbial pollution were present in the east and west sides of the river, and the localised re-suspension of sediment associated FIB into the water column was one proposed explanation for such contrasting FIB counts in the water column [45]. Despite this, the dynamics of sediment transport and re-suspension in relation to PIB concentrations is poorly understood. The attachment of PIB to particulate matter in the water column provides a platform for transportation and downward flux to the bed sediment, cyclical changes in tidal flow and the salinity of the water will also impact the downward flow of particle-associated PIB to the bed sediments. Conversely, under certain hydrodynamic conditions (e.g. tidal cycles and storm flow events), bed stress and turbulence can also impart the resuspension of sediments and subsequent transportation and deposition of particle associated FIB to other areas of the estuary. Due to the hydrodynamics of an estuarine system, fine sediments are usually deposited around the banks of the basin [49] and our data support this trend, with finer particles detected in greater abundance in sediments on the east and west sides of the estuary and coarser sand deposited in the central channel. FIB levels in water can be affected by other factors such as temperature [33], exposure to sunlight and salinity [31], nutrient concentrations [5], predation by protozoan [50] and competition [51]. However, here we demonstrate that sediments also contribute to the distribution of FIB and other PIB in an estuarine environment.

Experimental results confirm that estuarine sediments harbour PIB, which may potentiate their prolonged persistence and survival in this environment. This study also identifies areas of high microbial contamination within the Conwy estuary, which highlights the risk of sediment and PIB resuspension back into the water column under turbulent hydrodynamic conditions that result in sediment bed stress and promote the erosion of sediments. To our knowledge, this is the first comprehensive study of the cooccurrence of E. coli, coliforms, enterococci, Salmonella and Vibrio spp. (PIB) in relation to sediment composition. These data show that all PIB groups studied are strongly correlated with the presence of clay, silt and organic matter content in sediments. In addition, the presence of E. coli and total coliforms strongly correlates with the abundance of the other PIB tested, both allochthonous and autochthonous, suggesting that culture-based determinations of E. coli and coliform abundance in sediments represent a useful surrogate for the presence of other PIB groups.

#### Conclusion

Faecal contamination in aquatic environments is currently assessed by measuring culturable E. coli and enterococci counts in water samples only. Here, we demonstrate that sediment composition, specifically clay, silt and organic matter content, are linked with greater PIB abundance in sediments. The enhanced abundance of viable PIB in sediments therefore has implications for water quality and public health when considering the potential for resuspension of bed sediments, particularly finer particles such as clay and silt that we have demonstrated to contain higher concentrations of PIB. It may therefore be necessary to incorporate PIB loadings in bottom sediments into routine monitoring protocols and hydrodynamic models to adequately assess their risk to human health. The detection of spatial variations of PIB within sediments also highlights the necessity for further research on the interactions of pathogens with sediments and their role in the survival, persistence and transportation of PIB within environmental waters.

 Table S1
 Selective media used to enumerate target bacterial groups.

(DOCX)

Table S2 Sediment dry weight determined from  $1 \text{ g}^{-1}$  wet weight. (DOCX)

Table S3 Details the depth at which the sediment samples were collected, salinity measurements taken for both water samples (0.2 m from the surface) and directly above the sediment samples, calculated as Practical Salinity Units (PSU). Temperature was recorded at a depth of 0.2 m from the surface and directly above the sediment. (DOCX)

Table S4Identification of sequenced isolates.(DOCX)

Table S5 Bacteria counts, sediment (CFU/100 g) versus water (CFU/100 ml). Data shown as mean (n = 3, sample point 13 n = 2 for sediment). (DOCX)

Table S6 Correlation coefficient  $(r_s)$  matrix demonstrating the relationship between the abundance of each cultured bacterial group within estuarine sediments (n=21). (DOCX)

Table S7 Correlation coefficient  $(r_s)$  matrix demonstrating the relationship between the abundance of each cultured bacterial group within estuarine sediments and sediment grain size (n = 21). (DOCX)

### References

- Costanza R, d'Arge R, de Groot R, Farber S, Grasso M, et al. (1997) The value of the world's ecosystem services and natural capital. Nature 387: 253–260.
- Vitousek PM, Mooney HA, Lubchenco J, Melillo JM (1997) Human domination of Earth's ecosystems. Science 277: 494–499.
- Shuval H (2003) Estimating the global burden of thalassogenic diseases: human infectious diseases caused by wastewater pollution of the marine environment. J Water Health 1: 53–64.
- 4. GESAMP (2001) The sea of troubles. GESAMP Rep. Stud. 70: 19-20.
- Hong H, Qiu J, Liang Y (2010) Environmental factors influencing the distribution of total and fecal coliform bacteria in six water storage reservoirs in the Pearl River Delta Region, China. J Environ Sci (China) 22: 663–668.
- Topp E, Scott A, Lapen DR, Lyautey E, Duriez P (2009) Livestock waste treatment systems for reducing environmental exposure to hazardous enteric pathogens: some considerations. Bioresour Technol 100: 5395–5398.
- Obiri-Danso K, Jones K (2000) Intertidal sediments as reservoirs for hippurate negative Campylobacters, Salmonellae and faecal indicators in three EU recognised bathing waters in North West England. Wat Res 34: 519–527.
- Straub TM, Chandler DP (2003) Towards a unified system for detecting waterborne pathogens. J Microbiol Meth 53: 185–197.
- Edge TA, Boehm AB (2011) Classical and molecular methods to mearusre fecal bacteria. In: Sadowsky MJ, Whitman RL, editors. The fecal bacteria. Washington DC: ASM Press: pp. 241–273.
- Carrero-Colon M, Wickman GS, Turco RF (2011) Taxonomy, phylogeny and physiology of fecal indicator bacteria. In: Sadowsky MJ, Whitman RL, editors. The fecal bacteria. Washington DC: ASM Press: pp. 23–38.
- 2006 EC European Parliament, Directive 2006/7/EC of the European Parliament, concerning the management of bathing water quality and repealing directive 76/160/EEC. Off J Eur Communities 64: 37–51.
- Berthe T, Touron A, Leloup J, Deloffre J, Petit F (2008) Faecal-indicator bacteria and sedimentary processes in estuarine mudflats (Seine, France). Mar Pollut Bull 57: 59–67.
- Alonso JL, Alonso MA (1993) Presence of *Campylobacter* in marine waters of Valencia, Spain. Wat Res 27: 1559–1562.

Table S8 Correlation coefficient  $(r_s)$  matrix demonstrating the relationship between estuarine sediment grain size (%) and organic matter content (%) (n=21). (DOCX)

Table S9 Correlation coefficient  $(r_s)$  matrix demonstrating the relationship between the abundance of each cultured bacterial group within estuarine sediments and sediment organic matter content (n = 21). (DOCX)

Table S10 Correlation coefficient  $(r_s)$  matrix demonstrating the relationship between the abundance of each cultured bacterial group within estuarine sediments and physico-chemical parameters measured directly above the bottom sediments (n=2). (DOCX)

Table S11 Correlation coefficient  $(r_s)$  matrix demonstrating the relationship between the abundance of each cultured bacterial group within estuarine water and physico-chemical parameters measured at 0.2 m depth (n=21).

(DOCX)

#### Acknowledgments

We are grateful to Ben Winterbourn, Gwynn Parry Jones, Peter Hughes and Ben Powell for their assistance with estuarine sampling.

### **Author Contributions**

Conceived and designed the experiments: TLP JHB CFJ DLJ SKM JEM. Performed the experiments: TLP KC JHB JEM. Analyzed the data: TLP KC JHB JEM. Contributed reagents/materials/analysis tools: JHB CFJ SKM DLJ JEM. Contributed to the writing of the manuscript: TLP KC JHB DLJ SKM JEM.

- Cox AM, Gomez-Chiarri M (2012) Vibrio parahaemolyticus in Rhode Island coastal ponds and the estuarine environment of Narragansett bay. Appl Environ Microb 78: 2996–2999.
- Polo F, Figueras MJ, Inza I, Sala J, Fleisher JM, et al. (1998) Relationship between presence of *Salmonella* and indicators of faecal pollution in aquatic habitats. Fems Microbiol Lett 160: 253–256.
- Coburn B, Grassl GA, Finlay BB (2007) Salmonella, the host and disease: a brief review. Immunol Cell Biol 85: 112–118.
- Figueras MJ, Polo F, Inza I, Guarro J (1997) Past, present and future perspectives of the EU bathing water directive. Mar Pollut Bull 34: 148–156.
- Levin RE (2007) Campylobacter jejuni: A review of its characteristics, pathogenicity, ecology, distribution, subspecies characterization and molecular methods of detection. Food Biotechnol 21: 271–374.
- 19. Thorns CJ (2000) Bacterial food-borne zoonoses. Rev Sci Tech 19: 226-239.
- Dziejman M, Balon E, Boyd D, Fraser CM, Heidelberg JF, et al. (2002) Comparative genomic analysis of *Vibrio cholerae*: genes that correlate with cholera endemic and pandemic disease. Proc Natl Acad Sci U S A 99: 1556– 1561.
- Cabral JP (2010) Water microbiology. Bacterial pathogens and water. Int J Environ Res Public Health 7: 3657–3703.
- Bakr WMK, Hazzah WA, Abaza AF (2011) Detection of Salmonella and Vibrio species in some seafood in Alexandria. J Amer Sci 7: 663–668.
- Pommepuy M, Guillaud JF, Dupray E, Derrien A, Le Guyader F, et al. (1992) Enteric bacteria survival factors Wat Sci Tech 25: 93–103.
- Pote J, Haller L, Kottelat R, Sastre V, Arpagaus P, et al. (2009) Persistence and growth of faecal culturable bacterial indicators in water column and sediments of Vidy Bay, Lake Geneva, Switzerland. J Environ Sci (China) 21: 62–69.
- Garzio-Hadzick A, Shelton DR, Hill RL, Pachepsky YA, Guber AK, et al. (2010) Survival of manure-borne *E. coli* in streambed sediment: effects of temperature and sediment properties. Water Res 44: 2753–2762.
- Burton GA, Jr., Gunnison D, Lanza GR (1987) Survival of pathogenic bacteria in various freshwater sediments. Appl Environ Microbiol 53: 633–638.
- Jones GE (1964) Effect of chelating agents on the growth of *Escherichia coli* in seawater. J Bacteriol 87: 483–499.

- Davies CM, Long JA, Donald M, Ashbolt NJ (1995) Survival of fecal microorganisms in marine and freshwater sediments. Appl Environ Microbiol 61: 1888–1896.
- Marino RP, Gannon JJ (1991) Survival of fecal-coliforms and fecal streptococci in storm-drain sediment. Water Res 25: 1089–1098.
- Singleton FL, Attwell R, Jangi S, Colwell RR (1982) Effects of temperature and salinity on *Vibrio-cholerae* growth. Appl Environ Microbiol 44: 1047–1058.
- Bordalo AA, Onrassami R, Dechsakulwatana C (2002) Survival of faecal indicator bacteria in tropical estuarine waters (Bangpakong River, Thailand). J Appl Microbiol 93: 864–871.
- Davies CM, Evison LM (1991) Sunlight and the survival of enteric bacteria in natural waters. J Appl Bacteriol 70: 265–274.
- Faust MA, Aotaky AE, Hargadon MT (1975) Effect of physical parameters on the in situ survival of *Escherichia coli* MC-6 in an estuarine environment. Appl Microbiol 30: 800–806.
- Davies-Colley RJ, Donnison AM, Speed DJ, Ross CM, Nagels JW (1999) Inactivation of faecal indicator microorganisms in waste stabilisation ponds: Interactions of environmental factors with sunlight. Water Res 33: 1220–1230.
- Sinton LW, Davies-colley RJ, Bell RG (1994) Inactivation of enterococci and fecal-coliforms from sewage and meatworks effluents in seawater chambers. Appl Environ Microbiol 60: 2040–2048.
- Roper MM, Marshall KC (1974) Modification of the interaction between Escherichia coli and bacteriophage in saline sediments. Microbial Ecol 1: 1–13.
- Davies CM, Bavor HJ (2000) The fate of stormwater-associated bacteria in constructed wetland and water pollution control pond systems. J Appl Microbiol 89: 349–360.
- Gerba CP, McLeod JS (1976) Effect of sediments on the survival of *Escherichia* coli in marine waters. Appl Environ Microbiol 32: 114–120.
- An YJ, Kampbell DH, Breidenbach GP (2002) Escherichia coli and total coliforms in water and sediments at lake marinas. Environ Pollut 120: 771–778.
- Grimes DJ (1975) Release of sediment-bound fecal coliforms by dredging. Appl Microbiol 29: 109–111.
- Danovaro R, Fonda Umani S, Pusceddu A (2009) Climate change and the potential spreading of marine mucilage and microbial pathogens in the Mediterranean sea. PLoS One 4: e7006: doi:10.1371/journal.pone.0007006.
- Droppo IG, Liss SN, Williams D, Nelson T, Jaskot C, et al. (2009) Dynamic existence of waterborne pathogens within river sediment compartments. Implications for water quality regulatory affairs. Environ Sci Technol 43: 1737–1743.

- Thorn CE, Quilliam RS, Williams AP, Malham SK, Cooper D, et al. (2011) Grazing intensity is a poor indicator of waterborne *Escherichia coli* O157
- activity. Anaerobe 17: 330–333.
  44. Oliver LR, Seed R, Reynolds B (2008) The effect of high flow events on mussels (*Mytilus edulis*) in the Conwy estuary, North Wales, UK. Hydrobiologia 606: 117–127.
- Quilliam RS, Clements K, Duce C, Cottrill SB, Malham SK, et al. (2011) Spatial variation of waterborne *Escherichia coli* - implications for routine water quality monitoring. J Water Health 9: 734–737.
- Howell JM, Coyne MS, Cornelius PL (1996) Effect of sediment particle size and temperature on fecal bacteria mortality rates and the fecal coliform/fecal streptococci ratio. J Environ Qual 25: 1216–1220.
- Craig DL, Fallowfield HJ, Cromar NJ (2004) Use of microcosms to determine persistence of *Escherichia coli* in recreational coastal water and sediment and validation with in situ measurements. J Appl Microbiol 96: 922–930.
- Haller L, Amedegnato E, Pote J, Wildi W (2009) Influence of freshwater sediment characteristics on persistence of fecal indicator bacteria. Water Air Soil Pollut 203: 217–227.
- 49. Malham SK, Rajko-Nenow P, Howlett E, Tuson KE, Perkins TL, et al. (2014) The interaction of human microbial pathogens, particulate material and nutrients in estuarine environments and their impacts on recreational and shellfish waters. Environ Sci Process Impact 16: 2145–2155.
- Barcina I, Lebaron P, Vives-Rego J (1997) Survival of allochthonous bacteria in aquatic systems: a biological approach. FEMS Microbiol Ecol 23: 1–9.
- Korhonen LK, Martikainen PJ (1991) Survival of *Escherichia coli* and *Campylobacter jejuni* in untreated and filtered lake water. J Appl Bacteriol 71: 379–382.
- Ishii S, Hansen DL, Hicks RE, Sadowsky MJ (2007) Beach sand and sediments are temporal sinks and sources of *Escherichia coli* in Lake Superior. Environ Sci Technol 41: 2203–2209.
- Power ML, Littlefield-Wyer J, Gordon DM, Veal DA, Slade MB (2005) Phenotypic and genotypic characterization of encapsulated *Escherichia coli* isolated from blooms in two Australian lakes. Environ Microbiol 7: 631–640.
- Anderson KL, Whitlock JE, Harwood VJ (2005) Persistence and differential survival of fecal indicator bacteria in subtropical waters and sediments. Appl Environ Microbiol 71: 3041–3048.
- Evison LM (1989) Comparative studies on the survival of indicator organisms and pathogens in fresh and seawater. Wat Res 20: 309–352.
- Okpookwasili GC, Akujobi TC (1996) Bacteriological indicators of tropical water quality. Environ Toxic Water 11: 77–81.