



Is This Elderly Patient Dehydrated? Diagnostic Accuracy of Hydration Assessment Using Physical Signs, Urine, and Saliva Markers

Fortes, M.B.; Owen, J.A.; Raymond-Barker, P.; Bishop, C.; Elghenzai, S.; Oliver, S.J.; Walsh, N.P.

Journal of the American Medical Directors Association

DOI:

[10.1016/j.jamda.2014.09.012](https://doi.org/10.1016/j.jamda.2014.09.012)

Published: 22/10/2014

Other version

[Cyswllt i'r cyhoeddiad / Link to publication](#)

Dyfyniad o'r fersiwn a gyhoeddwyd / Citation for published version (APA):

Fortes, M. B., Owen, J. A., Raymond-Barker, P., Bishop, C., Elghenzai, S., Oliver, S. J., & Walsh, N. P. (2014). Is This Elderly Patient Dehydrated? Diagnostic Accuracy of Hydration Assessment Using Physical Signs, Urine, and Saliva Markers. *Journal of the American Medical Directors Association*, 16(3), 221-228. <https://doi.org/10.1016/j.jamda.2014.09.012>

Hawliau Cyffredinol / General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Manuscript Number: JAMDA-D-14-00269R1

Title: Is this elderly patient dehydrated? Diagnostic accuracy of hydration assessment using physical signs, urine and saliva markers

Article Type: Original Study

Keywords: Dehydration; diagnosis; older; hypovolemia; osmolality; clinical

Corresponding Author: Prof. Neil Peter Walsh, PhD

Corresponding Author's Institution: Bangor University

First Author: Matthew B Fortes

Order of Authors: Matthew B Fortes; Julian A Owen; Philippa Raymond-Barker, MSc; Claire Bishop; Salah Elghenzai; Samuel J Oliver, PhD; Neil P Walsh, PhD

Abstract: Objectives: Dehydration in older adults contributes to increased morbidity and mortality during hospitalization. As such, early diagnosis of dehydration may improve patient outcome and reduce the burden on healthcare. This prospective study investigated the diagnostic accuracy of routinely used physical signs, and non-invasive markers of hydration in urine and saliva. Design: Prospective diagnostic accuracy study. Setting: Hospital acute medical care unit and emergency department. Participants: One hundred and thirty older adults (59 males, 71 females, mean (SD) age = 78 (9) y). Measurements: Participants with any primary diagnosis underwent a hydration assessment within 30min of admittance to hospital. Hydration assessment comprised seven physical signs of dehydration (tachycardia (>100bpm), low systolic blood pressure (<100mmHg), dry mucous membrane, dry axilla, poor skin turgor, sunken eyes, and long capillary refill time (>2s)), urine color, urine specific gravity (USG), saliva flow rate (SFR) and saliva osmolality. Plasma osmolality (Posm) and the blood urea nitrogen to creatinine ratio (BUN:Cr) were assessed as reference standards of hydration, with 21% of participants classified with water-loss dehydration (Posm >295mOsm/kg), 19% classified with water-and-solute-loss dehydration (BUN:Cr >20) and 60% classified as euhydrated. Results: All physical signs showed poor sensitivity (0-44%) for detecting either form of dehydration, with only low systolic blood pressure demonstrating potential utility for aiding the diagnosis of water-and-solute-loss dehydration (diagnostic OR = 14.7). Neither urine color, USG, nor SFR could discriminate hydration status (area under the receiver operating characteristic curve, AUCROC = 0.49-0.57, P>0.05). In contrast, saliva osmolality demonstrated moderate diagnostic accuracy (AUCROC = 0.76, P<0.001) to distinguish both dehydration types (70% sensitivity, 68% specificity, OR =5.0 (95%CI 1.7-15.1) for water-loss dehydration, and 78% sensitivity, 72% specificity, OR =8.9 (95%CI 2.5-30.7) for water-and-solute-loss dehydration). Conclusions: With the exception of low systolic blood pressure, which could aid in the specific diagnosis of water-and-solute-loss dehydration, physical signs and urine markers show little utility to determine if an elderly patient is dehydrated. Saliva osmolality demonstrated superior diagnostic accuracy compared with physical signs and urine markers, and may have utility for the assessment of both water-loss and water-and-solute-loss dehydration in older individuals. It is particularly noteworthy that saliva osmolality was able to detect water-and-solute-loss dehydration, for which a measurement of plasma osmolality would have no diagnostic utility.



Prof. Neil Walsh
College of Health and Behavioural Sciences
Bangor University,
George Building,
Bangor, Gwynedd,
LL57 2PZ United Kingdom

11th September 2014

Tel: +44 (0) 1248 383480
E-mail: n.walsh@bangor.ac.uk

To the editor in chief of *Journal of the American Medical Directors Association*,

Re: Manuscript # JAMDA-D-14-00269, "Is this elderly patient dehydrated? Diagnostic accuracy of hydration assessment using physical signs, urine and saliva markers"

Thankyou for allowing us to resubmit the above manuscript to your journal. We have responded to the reviewers comments (see below), with changes in the manuscript highlighted in red text. We hope you feel that these changes have improved the manuscript.

Please don't hesitate to contact me if you require further information. I look forward to hearing from you.

Kind regards,

Prof. Neil Walsh.

Response to reviewers:

Reviewer #1:

Many thanks for your very insightful and constructive comments. Below are our responses to your comments, and changes within the manuscript are **highlighted in red text**. We hope you agree that these small changes have improved the manuscript.

- 1) Only 42% of screened individuals entered the study, and 31% had sufficient parameters to be analyzed. Although you show numbers in Figure 1, please comment on the large number of excluded subjects. Doesn't this affect the usefulness of the screening in clinical practice?

Thanks. As you have noted, of those that were screened (n =420), a large number of participants were excluded (58%). However, it should be noted that these were excluded due to ethical reasons of conducting the research as stipulated to us by the ethics committee (e.g. patients unable to provide consent (incapacity) for the research study (n=98), or that the research should not interfere with routine care of the patient and in those who had already began treatment (n = 88)), or because participants declined to take part (n = 54). We have now included this information at the start of the results sections, along with percentages of those excluded (lines 197-199). In terms of the application of the usefulness in clinical practice, the reasons outlined above do not preclude the usefulness of the measures in the current study being used in clinical practice (i.e. in all patients admitted to hospital).

In light of this being a proof of concept study for saliva indices, we did in this instance exclude participants who had potential confounding effects on saliva (e.g. oral trauma, recent dental surgery, swallowing problems etc), although it should be noted that only 2 participants were excluded for this (both had swallowing problems), and in light of your excellent point, we have now added this information to the results (lines 198-199,) and Figure 1, and have also now acknowledged in the

discussion that future studies should investigate whether saliva indices have utility, in patients with oral related problems (please see lines 341-344). Thanks.

- 2) "and allowing for an approximate one-third exclusion rate from data analysis (due to missing reference tests, and co-morbidities that preclude the use of the reference standards), a total of 178 participants were recruited into the study." It appears that the exclusion rate was higher than anticipated? Please comment.

Please note that the allowance for the one third exclusion rate (for missing reference tests and co-morbidities that affected the reference standards), was for those who might be excluded from the data analysis after they were already recruited into the study (i.e. n = 178 recruited). The N for which we analyzed data was n = 130, with 48 excluded from the data analysis. The proportion excluded from analysis of those recruited (48/178, 27%) is therefore actually lower, not higher than the anticipated 1/3 exclusion rate. Thanks.

- 3) "participants with a history of renal disease (n = 24), or who were in cardiac failure (n = 1) were excluded from data analysis." Please specify the criteria for renal disease and cardiac failure. What level of renal disease (stage?) or creatinine or other. For CHF, only "history" or other criteria? As you point out, the presence of renal disease, starvation, malnutrition (among others) limit the usefulness of the BUN/Cr ratio. It would be useful to discuss the level of renal disease that you excluded.

Thanks. For this study, we excluded from data analysis, all participants who had any known previous history of renal disease (CKD stage 1-5) or if they were in cardiac failure as diagnosed by the clinician. We have now clarified this and added this information to the methods section (Line 165). In line with comment 1 above, we have also now added a sentence to the discussion where we discuss how future studies should investigate the utility of these indices in these relatively small populations (lines 318-320). Thanks.

- 4) Please discuss relationship between saliva and blood osmolality. If the values are generally highly correlated, is there any benefit in using saliva rather than blood. Is it quicker, cheaper, easier to use saliva. Given a paucity of saliva in 25% of subjects, should blood be favored?

This is a very helpful observation and the changes we have made (described below) in response have improved the take home message of the manuscript. Many thanks.

As we have addressed in the manuscript (lines 55-61, 169-171 and in Figure 1), plasma osmolality is elevated in, and will only detect water-loss dehydration. In water-and-solute loss dehydration, plasma osmolality is either normal or low, and thus has no diagnostic utility for this type of dehydration. Given the differential response of plasma osmolality to these two types of dehydration, we feel it would be inappropriate to report, or rely on the correlation between saliva and plasma osmolality to determine saliva's utility as a diagnostic method. In the current study, saliva osmolality was able to detect a proportion of patients with water-and-solute dehydration (sensitivity 78%), and is an easier to perform and non-invasive so has advantages over blood sampling.

With this in mind, based on your excellent point, as this limitation of plasma (blood) osmolality for detecting water-and-solute-loss only dehydration was not as prominent as it should be in the manuscript, we have now added a sentence to the end of the abstract (lines 29-31) and to the discussion and conclusion where we address this (lines 312-314, 351-352).

We were able to collect a quantity of saliva in 126/130 patients (97%)- reported on lines 210 and 328, although as we have stated (lines 213, 327) we only had adequate saliva (at least 25ul) to assess osmolality using our osmometer in 75% of samples. However, we have addressed this limitation in the discussion, (line 327-333) where we say that micro osmometers are in development that can assess

osmolality on nano-gram quantities. We hope you feel that this is adequately addressed in the manuscript. Thanks.

Reviewer #2:

Many thanks for reviewing our manuscript.

Is this elderly patient dehydrated? Diagnostic accuracy of hydration assessment using physical signs, urine and saliva markers

Matthew B. Fortes¹, Julian A. Owen¹, Philippa Raymond-Barker¹, Claire Bishop², Salah Elghenzai², Samuel J. Oliver¹, and Neil P. Walsh¹

¹College of Health and Behavioural Sciences, Bangor University, Bangor, UK.

²Geriatric Medicine, Gwynedd Hospital, Betsi Cadwaladr University Health Board, Bangor, UK

Corresponding author:

Prof Neil Walsh,
College of Health and Behavioural Sciences,
Bangor University,
Bangor,
LL57 2PZ, UK.
Telephone: +(00) 44 1248 383480
Email: n.walsh@bangor.ac.uk

Alternative corresponding author:

Dr Matthew B. Fortes
College of Health and Behavioural Sciences,
Bangor University,
Bangor,
LL57 2PZ, UK.
Tel: +(00) 44 1248 388309
Email: m.fortes@bangor.ac.uk

Keywords: Dehydration, diagnosis, older, hypovolemia, osmolality, clinical

Running header: Dehydration assessment of older adults.

1 **Is this elderly patient dehydrated? Diagnostic accuracy of hydration assessment using physical signs,**
2 **urine and saliva markers**

3
4 **ABSTRACT**

5 **Objectives:** Dehydration in older adults contributes to increased morbidity and mortality during
6 hospitalization. As such, early diagnosis of dehydration may improve patient outcome and reduce the burden
7 on healthcare. This prospective study investigated the diagnostic accuracy of routinely used physical signs,
8 and non-invasive markers of hydration in urine and saliva. **Design:** Prospective diagnostic accuracy study.
9 **Setting:** Hospital acute medical care unit and emergency department. **Participants:** One hundred and thirty
10 older adults (59 males, 71 females, mean (SD) age = 78 (9) y). **Measurements:** Participants with any
11 primary diagnosis underwent a hydration assessment within 30min of admittance to hospital. Hydration
12 assessment comprised seven physical signs of dehydration (tachycardia (>100bpm), low systolic blood
13 pressure (<100mmHg), dry mucous membrane, dry axilla, poor skin turgor, sunken eyes, and long capillary
14 refill time (>2s)), urine color, urine specific gravity (USG), saliva flow rate (SFR) and saliva osmolality.
15 Plasma osmolality (Posm) and the blood urea nitrogen to creatinine ratio (BUN:Cr) were assessed as
16 reference standards of hydration, with 21% of participants classified with water-loss dehydration (Posm
17 >295mOsm/kg), 19% classified with water-and-solute-loss dehydration (BUN:Cr >20) and 60% classified as
18 euhydrated. **Results:** All physical signs showed poor sensitivity (0-44%) for detecting either form of
19 dehydration, with only low systolic blood pressure demonstrating potential utility for aiding the diagnosis of
20 water-and-solute-loss dehydration (diagnostic OR = 14.7). Neither urine color, USG, nor SFR could
21 discriminate hydration status (area under the receiver operating characteristic curve, $AUC_{ROC} = 0.49-0.57$,
22 $P>0.05$). In contrast, saliva osmolality demonstrated moderate diagnostic accuracy ($AUC_{ROC} = 0.76$,
23 $P<0.001$) to distinguish both dehydration types (70% sensitivity, 68% specificity, OR =5.0 (95%CI 1.7-15.1)
24 for water-loss dehydration, and 78% sensitivity, 72% specificity, OR =8.9 (95%CI 2.5-30.7) for water-and-
25 solute-loss dehydration). **Conclusions:** With the exception of low systolic blood pressure, which could aid
26 in the specific diagnosis of water-and-solute-loss dehydration, physical signs and urine markers show little
27 utility to determine if an elderly patient is dehydrated. Saliva osmolality demonstrated superior diagnostic
28 accuracy compared with physical signs and urine markers, and may have utility for the assessment of both
29 water-loss and water-and-solute-loss dehydration in older individuals. **It is particularly noteworthy that saliva**

30 osmolality was able to detect water-and-solute-loss dehydration, for which a measurement of plasma
31 osmolality would have no diagnostic utility.

32

33 INTRODUCTION

34

35 Dehydration in older adults is a significant clinical problem. A diagnosis of dehydration is associated with
36 the presence of co-morbidities, longer hospital stay, additional future hospitalization and higher mortality
37 rates¹⁻⁵. The point-prevalence of dehydration in community-dwelling older adults in the USA was reported
38 as 17-28%^{6;7}. In many cases, simple and inexpensive oral rehydration is sufficient to treat dehydration and
39 halt the progress of more serious fluid-deficit related illnesses such as acute kidney injury. However, upon
40 hospitalization, many patients may be denied the correct course of treatment due to physician misdiagnosis
41 of dehydration⁷. Therefore, accurate and early identification of dehydration in older adults admitted to
42 hospital is vital to alleviate ill-health and the significant economic burden of treating dehydration on
43 healthcare^{1;2}.

44

45 No single 'gold-standard' marker of hydration status exists⁸, although blood biochemistry including plasma
46 osmolality, electrolytes and blood urea nitrogen to creatinine ratio (BUN:Cr) represent criterion methods of
47 identifying dehydration in a clinical setting⁹⁻¹². However, blood sample collection is invasive and laboratory
48 analysis is time-consuming, often delaying the course of treatment by hours. To aid an initial diagnosis of
49 dehydration before requesting blood biochemistry confirmation, clinicians may use a variety of simple
50 screening measures, albeit in a non-systematic way, that may include; presenting signs and symptoms of
51 dehydration^{11;13;14}, patient history¹³, orthostatic blood pressure change¹⁵, and/or urinary parameters¹⁶.
52 Nevertheless, these screening methods are often characterized by poor diagnostic performance^{11;17-21}. To
53 confound hydration assessment further, the term 'dehydration' is poorly defined and is used to characterize
54 many water and solute deficits relating to whole body fluid deficits⁷. In order to simplify clinical practice
55 researchers have suggested the classification of clinical dehydration into two distinct types. Firstly, water-
56 loss dehydration (also termed hypertonic hypovolemia, or intracellular dehydration), which is hypertonic in
57 nature and occurs when water loss proportionally exceeds solute loss. Water loss dehydration is typically
58 defined as a plasma osmolality ≥ 295 mOsm/kg^{12;22}. Secondly, water-and-solute-loss dehydration (also

59 termed intravascular volume depletion or extracellular dehydration), which may be isotonic or hypotonic due
60 to equal, or greater proportional loss of solutes than water ^{10;12;23}, and typically defined as a BUN:Cr ≥ 20 in
61 the absence of hypertonicity ²². To the best of our knowledge, there are few ^{18;19}, rigorous studies that have
62 investigated the diagnostic accuracy of clinical physical signs and/or urine indices to detect dehydration in
63 hospitalized older adults using a criterion reference method, and none which have simultaneously assessed
64 the utility of any hydration marker to assess both types of dehydration.

65

66 In a series of studies (in young healthy adults) we have shown that rapid measurements made from non-
67 invasive collection of saliva fluid can be used to identify water-loss dehydration ²⁴⁻²⁶. For example,
68 decreases in whole saliva flow rate and increases in whole saliva osmolality were shown to track progressive
69 modest dehydration (equivalent to 1-3% body mass loss). The utility of these novel saliva markers of
70 dehydration has not yet been examined in a clinical, older adult population, although encouragingly, the
71 presence of a dry tongue was identified as the clinical sign most strongly associated with dehydration in an
72 elderly cohort ¹⁴. To this end, the purpose of this prospective study was to determine, and compare, the
73 diagnostic accuracy of clinical physical signs routinely used in hospital settings ^{11;13;14}, along with saliva
74 (flow rate and osmolality) and urine indices (color and specific gravity) ²⁷, to detect static (one-point in time)
75 water-loss, and water-and-solute-loss dehydration in a hospitalized, older adult cohort using primary
76 reference standards; plasma osmolality and BUN:Cr ^{10;12;22;28}.

77

78 **METHODS**

79

80 **Experimental design and procedures**

81 The study was conducted as a prospective, hospital-based cross-sectional study. All measures of hydration
82 status were performed within 30 minutes of admission, with no disruption to routine care in the following
83 order; examination of physical signs of dehydration, collection of saliva, blood and urine. For the reference
84 standards of whole body hydration assessment, a blood sample was collected by the clinical research fellow
85 or a specialist phlebotomist and analyzed for plasma osmolality (within 15min) and BUN:Cr (within 2h).
86 For consistency, all physical examinations and assessment of confidential medical information was carried
87 out by the same clinical research fellow (a junior doctor with five years clinical experience), who was

88 blinded to the results of the reference standards and the saliva and urine index test results when conducting
89 the physical examination. Saliva and urine samples were collected, and analyzed by an independent research
90 assistant who had been trained in the handling and assessment of saliva and urine samples by a postdoctoral
91 researcher, and who was blinded to the physical examination results. All osmolality analyses were made by
92 a trained research assistant. Details of the patients' medical condition, history and medication were recorded
93 retrospectively after the reference and index test results had been established.

94

95 **Participants**

96 A convenience sample of adults over 60 years of age admitted consecutively to the acute medical care unit or
97 emergency department of Gwynedd Hospital, Bangor, UK, with any primary diagnosis and capacity to
98 consent were enrolled between May and November 2011 during the times the investigators were available
99 (09:00h – 17:00h, Monday-Friday). Participant exclusion criteria included: oral trauma or dental surgery
100 within 14 days, swallowing problems, salivary gland tumors, if they were deemed too unwell by the medical
101 staff to participate in the study, if they were assessed as not having capacity to consent, or if they had already
102 begun any form of medical treatment or rehydration therapy (oral or intravenous). Participant flow through
103 the study is depicted in **Figure 1**. All participants recruited provided fully informed written consent, and the
104 study adhered to the Declaration of Helsinki and was approved by the North West Wales Research Ethics
105 Committee (Ref: 11/WA/0023).

106

107 **Assessment of hydration status**

108

109 **Reference standards**

110 *Blood sample collection and analysis*

111 Blood samples were collected from an antecubital or dorsal metacarpal vein without venostasis into one
112 serum separation vacutainer, and one lithium heparin coated vacutainer (Becton Dickinson, Oxford, UK).
113 Serum blood urea nitrogen and serum creatinine were assessed at the hospital clinical biochemistry
114 department using an automated biochemistry analyzer (Olympus AU 2700 chemistry immuno analyzer,
115 Beckman Coulter, USA). The lithium heparin treated blood was centrifuged immediately upon collection at
116 1500 g for 10 minutes at 4 °C. The plasma was aspirated and triplicate measurements of osmolality were

117 made immediately using a freezing point depression osmometer (Model 330 MO, Advanced Instruments,
118 Massachusetts, USA). Standard control solutions (290 mOsm/kg) were run through the osmometer and
119 checked daily to ensure acceptable limits of precision (± 2 mOsm/kg). The analytical coefficient of variation
120 for repeated sample plasma osmolality measurements was 0.7% (1.9 mOsm/kg).

121

122 **Index tests**

123 *Clinical assessment of physical signs of dehydration*

124 The clinical assessment consisted of seven physical signs of dehydration that are routinely used in Gwynedd
125 Hospital; tachycardia (resting heart rate >100 beats per minute), low resting systolic blood pressure
126 (<100 mmHg), dry mucous membrane (inside of the cheek, dry vs. wet), axillary dryness (assessed by
127 palpating the armpit, dry vs. moist), poor skin turgor (measured by pinching the skin on the dorsum of the
128 hand and observing if the tissue fold returned to normal immediately), presence of sunken eyes as assessed
129 by the clinical research fellow, and long capillary refill time (> 2 s, assessed by holding the patients hand at
130 heart level and blanching the participant's right index finger using moderate pressure and assessing the length
131 of time for the return of normal color). Each physical sign was assessed with the participant rested and
132 seated upright and assessed dichotomously.

133

134 *Saliva sample collection and analysis*

135 Unstimulated whole saliva samples were collected using a pre-weighed Versi-sal® collection device (Oasis
136 Technology, USA) as previously described²⁹. Participants firstly swallowed in order to empty the mouth of
137 residual saliva, before saliva was collected by placing the Versi-sal® collection device under the tongue.
138 Saliva collection was performed with minimal orofacial movements and accurately timed. After 4 min, the
139 collection device was inspected for volume of saliva by weighing it immediately (to the nearest milligram)
140 and subtracting the pre-weight. If the volume was insufficient for osmolality analysis ($< 25\mu$ l), the swab was
141 replaced under the tongue for a further 4 min. By assuming the density of saliva to be 1.00g/ml, saliva flow
142 rate (SFR) was calculated by dividing the volume collected by the time of collection²⁴. Saliva was
143 recovered from the collection device by centrifugation at 1500 g for 10 min, and assessed immediately in
144 duplicate for saliva osmolality using a freezing point depression osmometer (Model 330 MO, Advanced

145 Instruments, Massachusetts, USA). The analytical coefficient of variation for repeated sample saliva
146 osmolality measurements was 0.8% (0.9 mOsm/kg).

147

148 *Urine sample collection and analysis*

149 A mid-flow urine sample was collected and immediately analyzed for urine color ²⁷ and urine specific
150 gravity (USG) using a handheld refractometer (Atago URC-Osmo refractometer, Japan).

151

152 **Sample size calculation and data analysis**

153 The desired sample size for dehydrated participants ($n = 20$ water-loss only) was calculated using the
154 following equation:

155

156

157

158

$$n \geq \frac{(1.96)^2 p(1-p)}{x^2}$$

159 Where p = desired sensitivity (70%) as a proportion, and x = desired confidence interval (20%) as a

160 proportion ³⁰. Assuming a prevalence of impending water-loss dehydration (plasma osmolality \geq

161 295mOsm/kg) of 17% ⁷, and allowing for an approximate one-third exclusion rate from data analysis (due to

162 missing reference tests, and co-morbidities that preclude the use of the reference standards), a total of 178

163 participants were recruited into the study. Medical records for participants were accessed after enrolment,

164 and due to potential influencing effects on the reference standards assessed in this study, participants with a

165 history of renal disease (CKD stage 1-5, $n = 24$), or who were in cardiac failure as diagnosed by a clinician

166 ($n = 1$) were excluded from data analysis. Participants were also excluded from data analysis if the reference

167 tests were not available ($n = 11$), if they had an abnormally low (<10) BUN:Cr which may be indicative of

168 renal disease or the syndrome of inappropriate antidiuretic hormone ($n = 8$), or if they were taking

169 glucocorticoid medication ($n = 4$) which affects the validity of the BUN:Cr ¹⁰. Based on the reference

170 standards, participants with a presenting plasma osmolality ≥ 295 mOsm/kg were classified as having

171 impending water-loss dehydration ^{12;22}. Of the remaining participants, those with a BUN:Cr ≥ 20 in the

172 absence of hypertonicity ²² were classified as having water-and-solute-loss dehydration, and the remaining

173 participants formed the euhydrated control group (normal plasma osmolality and BUN:Cr).

174

175 To assess the diagnostic accuracy of saliva and urine indices, and clinical physical signs for assessment of
176 hydration status, both water-loss, and water-and-solute-loss dehydration groups were separately compared
177 with the euhydrated control group. Both dehydration groups were also combined to form a generic
178 dehydration group for comparison with euhydration. For all dichotomized clinical physical sign data, the
179 following were calculated; area under the receiver operating characteristic curve (AUC_{ROC}) as a measure of
180 global diagnostic accuracy, sensitivity, specificity, positive and negative likelihood ratios (LR), and the
181 diagnostic odds ratio (OR) generated by logistical regression. For continuous variable data (urine color,
182 USG, SFR and saliva osmolality), the degree to which each variable could discriminate between dehydration
183 and euhydration was assessed using AUC_{ROC} . For variables that could distinguish hydration status, the
184 single cut-off value that provided the optimal discrimination was identified as the point on the curve with the
185 largest vertical displacement from the reference line, and sensitivity, specificity, overall diagnostic accuracy,
186 positive and negative LR, and the diagnostic OR were calculated. For all diagnostic analyses 95%
187 confidence intervals were constructed. To compare AUC_{ROC} , a method was adopted that accounts for the
188 correlation between samples from the same individual³¹. Group data were analyzed using one-way ANOVA.
189 Data were analyzed using Microsoft excel (Microsoft, USA), SigmaPlot version 12.0 (Systat Software, Inc.
190 USA) and SPSS version 20 (IBM Corporation, USA) software. Significance was accepted as $P < 0.05$ for all
191 ANOVA, logistic regression and AUC_{ROC} analyses.

192 *****Insert Figure 1 about here*****

193

194 **RESULTS**

195

196 **Participant characteristics**

197 **A total of 420 participants were screened for inclusion, with 242 excluded, largely due to ethical**
198 **considerations of conducting the research, or declining to take part ($n = 240, 57\%$), or due to swallowing**
199 **problems ($n = 2, 1\%$). Therefore, 178 participants were enrolled into the study ($n = 85$ males, $n = 93$**
200 **females) with mean age (SD) 78 (9) y. After further exclusions for data analysis, data were analyzed for $n =$**
201 **130 participants ($n = 59$ males, $n = 71$ females; mean age 78 (9), range 60-101y), of which $n = 27$ (21%)**
202 **were classified as water-loss dehydrated, $n = 25$ (19%) were classified as water-and-solute-loss dehydrated,**
203 **and $n = 78$ (60%) were classified as euhydrated. Of the 27 participants in the water-loss only dehydration**

204 group, 10 also had an elevated BUN:Cr (≥ 20). There were no differences between the groups for age (**Table**
205 **1**). By design, participants with water-loss dehydration had elevated plasma osmolality, and participants
206 with water-and-solute-loss dehydration had elevated BUN:Cr compared with euhydrated control (**Table 1**).

207 *****Insert Table 1 about here*****

208

209 **Feasibility of collecting index tests**

210 All clinical physical sign assessments were conducted in all 130 participants. Saliva was collected in all but
211 four participants (1 water-loss dehydrated, 2 water-and-solute-loss dehydrated, and 1 euhydrated control).

212 For these four participants SFR was recorded as zero, and SFR data was therefore analyzed for $n = 130$.

213 There was adequate saliva ($> 25\mu\text{l}$) to assess saliva osmolality in 98 participants (75%). In comparison urine
214 samples could not be collected in 45 participants, who were unable to urinate within 30 min of the blood
215 collection. One participant provided a urine sample containing blood, confounding interpretation. Urine
216 color and specific gravity were therefore analyzed in 84 participants (65%).

217

218 **Diagnostic accuracy of clinical physical signs**

219 Diagnostic data for all seven clinical physical signs for both types of dehydration are shown in **Table 2** and
220 **Figure 2**. No clinical physical sign in isolation could discriminate between euhydration and either form of
221 dehydration (AUC_{ROC} range 0.44-0.57). Individually, all clinical physical signs performed poorly in terms of
222 detecting dehydration with sensitivity ranging from 0–44%. They did however generally perform better at
223 detecting euhydration, with specificity ranging from 60-99%. For detecting water-and-solute-loss
224 dehydration, a low resting systolic blood pressure ($< 100\text{mmHg}$) demonstrated high diagnostic odds and
225 likelihood ratios (14.7 (95% CI 1.6-138.3) and 12.5 (95% CI 1.5-107 respectively)), suggesting potential
226 utility in aiding the diagnosis of this specific type of dehydration.

227 *****Insert Table 2 about here*****

228

229 **Diagnostic accuracy of urine and saliva indices**

230 There were no differences between any of the three groups for urine color, USG or SFR (**Table 1**).

231 Furthermore, when assessed using ROC analyses, neither urine color, USG or SFR were able to discriminate
232 between dehydration and euhydration (AUC_{ROC} range 0.49-0.57, all $P > 0.05$, **Table 3**). Saliva osmolality

233 was greater in participants with both forms of dehydration than euhydrated control ($P < 0.001$, **Table 1**), but
234 more importantly, was able to distinguish both types of dehydration separately from euhydration ($AUC_{ROC} =$
235 0.76 , $P < 0.01$ for both types of dehydration individually and combined, **Table 3**). Based on the ROC
236 analysis, the saliva osmolality cut-off that provided the optimum balance between sensitivity and specificity
237 was calculated as: 95, 97, and 94 mOsm/kg for water-loss only, water-and-solute-loss only, and both forms
238 of dehydration combined, respectively. The diagnostic accuracy of saliva osmolality to detect all
239 dehydration types is displayed in **Table 4**. Saliva osmolality was able to identify water-loss dehydration,
240 water-and-solute-loss dehydration, and both forms of dehydration combined with a sensitivity of 70, 78 and
241 76%, and specificity of 68, 72, and 68%, respectively. Importantly, when AUC_{ROC} curves were compared,
242 the ability of saliva osmolality to discriminate hydration status was superior ($P < 0.05$) to all clinical physical
243 signs and urine indices for both types of dehydration in older adults (**Figure 2**).

244 *****Insert Table 3 and Table 4 about here***** *****Insert Figure 2 about here*****

245

246 **DISCUSSION**

247 Dehydration in older adults is a leading cause of hospitalizations, contributing to increased morbidity and
248 mortality during clinical care, and poorer functional status of the individual^{1-5;32}. As such, early
249 identification of hydration status is paramount to prevent the development of further co-morbidities, and to
250 reduce the burden on healthcare^{1;2}. This prospective study sought to investigate the diagnostic accuracy of
251 routinely used clinical physical signs and urine indices, and novel, simple, non-invasive saliva indices. The
252 main finding was that currently used clinical physical signs were not able to discriminate between
253 dehydration and euhydration, and thus provide little help to the physician making an initial hydration
254 assessment. The exception was a low systolic blood pressure which could aid in the specific diagnosis of
255 water-and-solute-loss dehydration. Whilst showing promise in young healthy cohorts²⁷, urine analysis
256 demonstrated no utility in identifying dehydration in an older adult cohort admitted to hospital. However,
257 the novel finding from the study was that saliva osmolality could discriminate between dehydration and
258 euhydration, and importantly, was sensitive to both water-loss, and water-and-solute-loss forms of
259 dehydration, demonstrating superior diagnostic accuracy than urinary parameters and currently used clinical
260 physical signs. Saliva collection is non-invasive, and easy to collect, and therefore, may have practical
261 utility as an initial screening method for impending dehydration in older adults.

262

263 Despite a relative paucity of clear supporting evidence, clinicians may rely on an array of simple physical
264 screening tests to aid the hydration assessment of patients admitted to hospital. Whilst showing some
265 clinical promise in young children^{33;34}, clinical physical signs often demonstrate poor diagnostic
266 performance when applied to older adults, likely due to; a loss of skin elasticity with advancing age affecting
267 skin turgor, smoking and cold environmental temperatures causing peripheral vasoconstriction which may
268 result in false positives for capillary refill time, and anticholinergic medications and a reliance on mouth
269 breathing in the elderly which can result in a dry oral mucosa^{11;17;35}. Findings from previous studies
270 investigating the utility of clinical physical signs should also be viewed with caution where they have
271 adopted a non-criterion reference standard, e.g. difference in weight gain after rehydration¹³, urinary
272 measures¹⁶, or relied on a clinician's overall diagnosis¹⁴ as opposed to a more, objective biochemical
273 criterion measure such as plasma osmolality^{7;10-12;17;22;36}. Furthermore, previous studies have been limited by
274 failing to characterize the diagnostic accuracy of clinical physical signs in assessing both forms of
275 dehydration commonly encountered in a clinical setting, i.e. water-loss, and water-and-solute-loss
276 dehydration¹⁰.

277

278 A particular strength of the current study was that both forms of dehydration were characterized
279 simultaneously using valid biochemical assessments as reference standards, including the preferred direct
280 measurement of plasma osmolality as opposed to calculated osmolality^{12;37}. We observed that no clinical
281 physical sign could discriminate between either type of dehydration and euhydration when assessed using
282 AUC_{ROC} , and thus, should not be used in isolation to diagnose hydration status in older adults admitted to
283 hospital. However, although not sensitive (16%), a low (<100mmHg) sitting systolic blood pressure, may
284 aid the physician in making a diagnosis of water-and-solute-loss dehydration owing to its very high
285 specificity (i.e. low false positive rate), high diagnostic odds ratio (OR =14.7), and high positive likelihood
286 ratio (OR =12.5). This finding is in line with the well-known effects of a loss of extracellular fluid
287 (intravascular volume depletion) on blood pressure responses. Although researchers have previously focused
288 on orthostatic blood pressure responses to assess hydration^{13;15}, altering posture may be impractical in a
289 clinical setting, particularly in bed-ridden patients. Therefore, a sitting blood pressure assessment may have
290 practical value for the clinician.

291

292 Urinary markers have been reported as valid methods to assess acute changes in hydration status in young
293 healthy people²⁷. In the current study, neither USG nor urine color were able to discriminate between
294 dehydration and euhydration. This is likely due in part, to the decreased renal function that is characteristic
295 of older age, and to a potential confounding effect on urine of the many types of medications that an older
296 adult cohort are likely to be prescribed. In support, previous studies have also shown that urine indices are
297 poor markers of hydration status in elderly patients^{13;19}, in critically ill patients²⁰ and in young children with
298 gastroenteritis³⁸. Urine collection is not always possible when required, and was only able to be collected in
299 65% of participants in the current study, and in only 79% of elderly patients in a recent clinical study¹³.
300 Taken together, we do not recommend the use of USG or urine color as screening tools for dehydration in
301 older adults.

302

303 To the best of our knowledge, this is the first study that has investigated the diagnostic accuracy of saliva
304 indices to assess dehydration in older adults admitted to hospital. Saliva sample collection is simple and
305 non-invasive and has previously been shown to track modest water-loss dehydration in young healthy males
306²⁴⁻²⁶. Saliva flow rate was not associated with either form of dehydration, but the novel finding of the current
307 study was that saliva osmolality was able to detect both forms of dehydration with sensitivity >70% and
308 diagnostic OR >5. Although a sensitivity to detect dehydration of 72-78% may only be described as “fair to
309 moderate”³⁹, it is important to stress that any novel diagnostic marker should be compared against what is
310 currently used in clinical practice, and in the case of the present study, a high saliva osmolality
311 (>94mOsm/kg) was able to detect more cases of both types of dehydration than any single clinical physical
312 sign or urinary marker without compromising specificity (**Figure 2**). **It is also worth re-iterating that saliva**
313 **osmolality was able to detect water-and-solute-loss dehydration, for which a measurement of plasma**
314 **osmolality would have no diagnostic utility**. Furthermore, the cohort in the current study reflects a
315 representative, older adult clinical population, admitted with any primary diagnosis, and we did not remove
316 participants taking medications (except for 4 patients taking glucocorticoid medications). Thus, the fact that
317 a single marker is able to achieve a sensitivity > 70% for both types of dehydration at one-point in time
318 regardless of medication is promising. **It remains unknown whether saliva osmolality can also identify both**
319 **types of dehydration in the relatively small proportion of patients in this study taking glucocorticoid**

320 medication, in patients with heart failure, and in those with various stages of kidney disease. Finally, since
321 we set our reference standard cut-off at the lower end of the dehydration continuum to reflect impending, or
322 pre-clinical dehydration^{10;12;22}, the measurement of saliva osmolality may have practical utility in identifying
323 those individuals with modest dehydration, so that further biochemistry analysis can confirm the presence of,
324 and type of dehydration, in order that specific, tailored rehydration is commenced to prevent the patient
325 developing more severe dehydration along with its associated co-morbidities and poorer outcome.

326

327 There are a few limitations of saliva that we must acknowledge. Firstly, in the current study, the requirement
328 of 25 µl of saliva sample for analysis meant that only 75% of the samples could be analyzed (although a
329 measurable quantity of saliva was collected from 97% of participants compared with only 65% of
330 participants able to provide a urine sample). However, point of care devices that utilize nano-technology for
331 the assessment of saliva osmolality are under development^{40;41}. For example, the osmolality of tears can
332 now be assessed using the principle of impedance on as little as 50nl^{42;43}. Thus, this limitation should not be
333 seen to detract from the future application of saliva osmolality to assess hydration status in clinical care.

334 Secondly, with saliva sampling in a clinical population, there may be a potential confounding effect of
335 anything which can affect saliva flow rate, e.g. anticholinergic medications, or recent food/fluid consumption
336^{44;45}. This is potentially important since a decrease in saliva flow explained in part, the increase in saliva
337 osmolality observed during acute dehydration in young healthy males²⁴⁻²⁶. However, we observed only a
338 small association between saliva flow rate and osmolality ($r = -0.40$), suggesting that in the current study,
339 saliva osmolality was largely independent of saliva flow rate. The physiological mechanisms responsible for
340 an increase in saliva osmolality during dehydration are unclear, but may be due to an increase in water
341 absorption in the saliva gland and/or neural factors²⁴⁻²⁶. Finally, although we excluded only 2 participants
342 with swallowing problems, further research should investigate the diagnostic utility of saliva indices in
343 patients with this, and other oral-related problems (e.g. oral trauma, recent dental surgery, salivary gland
344 tumors etc).

345

346 CONCLUSIONS

347 In conclusion, with the exception of low systolic blood pressure, which could aid in the specific diagnosis of
348 water-and-solute-loss dehydration, physical signs and urine markers show little utility to determine if an

349 elderly patient is dehydrated. Saliva osmolality demonstrated superior diagnostic accuracy compared with
350 physical signs and urine markers for the assessment of both water-loss and water-and-solute-loss
351 dehydration. **It is particularly noteworthy that saliva osmolality was able to detect water-and-solute-loss**
352 **dehydration, for which a measurement of plasma osmolality would have no diagnostic utility.** The
353 measurement of saliva osmolality has potential utility as a screening method to aid the diagnosis of
354 impending dehydration in older adults.

355

356

357 **COMPETING INTERESTS DECLARATION:**

358 The study was funded by HydraDx Inc, who were interested in identifying if saliva indices other than those
359 presented herein had utility for hydration assessment. MBF and PRB were employed as research assistants
360 by HydraDx on this study.

361

362 **ACKNOWLEDGEMENTS**

363 The authors would like to thank Emma Tye for her assistance with data collection, Dr Frank Bellizzi, Dr
364 Andrew Goldstein, Dr Ken Strahs and Dr John Donovan for their valuable input into the research protocol,
365 and to Betsi Cadwaladr University Health Board, for facilitating this research. We would also like to thank
366 all the patients who agreed to participate in this study.

367

368

369 **REFERENCES**

- 370 (1) Xiao H, Barber J, Campbell ES. Economic burden of dehydration among hospitalized elderly
371 patients. *Am J Health Syst Pharm* 2004;61:2534-2540.
- 372 (2) Warren JL, Bacon WE, Harris T, McBean AM, Foley DJ, Phillips C. The burden and outcomes
373 associated with dehydration among US elderly, 1991. *Am J Public Health* 1994;84:1265-1269.
- 374 (3) Bhalla A, Sankaralingam S, Dundas R, Swaminathan R, Wolfe CD, Rudd AG. Influence of raised
375 plasma osmolality on clinical outcome after acute stroke. *Stroke* 2000;31:2043-2048.
- 376 (4) O'Neill PA, Faragher EB, Davies I, Wears R, McLean KA, Fairweather DS. Reduced survival with
377 increasing plasma osmolality in elderly continuing-care patients. *Age Ageing* 1990;19:68-71.
- 378 (5) Rowat A, Graham C, Dennis M. Dehydration in hospital-admitted stroke patients: detection,
379 frequency, and association. *Stroke* 2012;43:857-859.

- 380 (6) Stookey JD. High prevalence of plasma hypertonicity among community-dwelling older adults:
381 results from NHANES III. *J Am Diet Assoc* 2005;105:1231-1239.
- 382 (7) Thomas DR, Tariq SH, Makhdomm S, Haddad R, Moinuddin A. Physician misdiagnosis of
383 dehydration in older adults. *J Am Med Dir Assoc* 2004;5:S30-S34.
- 384 (8) Armstrong LE. Assessing hydration status: the elusive gold standard. *J Am Coll Nutr* 2007;26:575S-
385 584S.
- 386 (9) Mange K, Matsuura D, Cizman B et al. Language guiding therapy: the case of dehydration versus
387 volume depletion. *Ann Intern Med* 1997;127:848-853.
- 388 (10) Thomas DR, Cote TR, Lawhorne L et al. Understanding clinical dehydration and its treatment. *J Am*
389 *Med Dir Assoc* 2008;9:292-301.
- 390 (11) McGee S, Abernethy WB, III, Simel DL. The rational clinical examination. Is this patient
391 hypovolemic? *JAMA* 1999;281:1022-1029.
- 392 (12) Hooper L, Attreed NJ, Campbell WW et al. Clinical and physical signs for identification of
393 impending and current water-loss dehydration in older people (Protocol). *Cochrane Database of*
394 *Systematic Reviews* 2012;2.
- 395 (13) Vivanti A, Harvey K, Ash S, Battistutta D. Clinical assessment of dehydration in older people
396 admitted to hospital: what are the strongest indicators? *Arch Gerontol Geriatr* 2008;47:340-355.
- 397 (14) Vivanti A, Harvey K, Ash S. Developing a quick and practical screen to improve the identification
398 of poor hydration in geriatric and rehabilitative care. *Arch Gerontol Geriatr* 2010;50:156-164.
- 399 (15) Chassagne P, Druesne L, Capet C, Menard JF, Bercoff E. Clinical presentation of hypernatremia in
400 elderly patients: a case control study. *J Am Geriatr Soc* 2006;54:1225-1230.
- 401 (16) Wakefield B, Menten J, Diggelmann L, Culp K. Monitoring hydration status in elderly veterans.
402 *West J Nurs Res* 2002;24:132-142.
- 403 (17) Weinberg AD, Minaker KL. Dehydration. Evaluation and management in older adults. Council on
404 Scientific Affairs, American Medical Association. *JAMA* 1995;274:1552-1556.
- 405 (18) Eaton D, Bannister P, Mulley GP, Connolly MJ. Axillary sweating in clinical assessment of
406 dehydration in ill elderly patients. *BMJ* 1994;308:1271.
- 407 (19) Rowat A, Smith L, Graham C, Lyle D, Horsburgh D, Dennis M. A pilot study to assess if urine
408 specific gravity and urine colour charts are useful indicators of dehydration in acute stroke patients. *J*
409 *Adv Nurs* 2011;67:1976-1983.
- 410 (20) Fletcher SJ, Slaymaker AE, Bodenham AR, Vucevic M. Urine colour as an index of hydration in
411 critically ill patients. *Anaesthesia* 1999;54:189-192.
- 412 (21) Shimizu M, Kinoshita K, Hattori K et al. Physical signs of dehydration in the elderly. *Intern Med*
413 2012;51:1207-1210.
- 414 (22) Stookey JD, Pieper CF, Cohen HJ. Is the prevalence of dehydration among community-dwelling
415 older adults really low? Informing current debate over the fluid recommendation for adults aged
416 70+years. *Public Health Nutr* 2005;8:1275-1285.
- 417 (23) Hooper L, Bunn D, Jimoh FO, Fairweather-Tait SJ. Water-loss dehydration and aging. *Mech Ageing*
418 *Dev* 2013;10.

- 419 (24) Oliver SJ, Laing SJ, Wilson S, Bilzon JL, Walsh NP. Saliva indices track hypohydration during 48h
420 of fluid restriction or combined fluid and energy restriction. *Arch Oral Biol* 2008;53:975-980.
- 421 (25) Walsh NP, Laing SJ, Oliver SJ, Montague JC, Walters R, Bilzon JL. Saliva parameters as potential
422 indices of hydration status during acute dehydration. *Med Sci Sports Exerc* 2004;36:1535-1542.
- 423 (26) Walsh NP, Montague JC, Callow N, Rowlands AV. Saliva flow rate, total protein concentration and
424 osmolality as potential markers of whole body hydration status during progressive acute dehydration
425 in humans. *Arch Oral Biol* 2004;49:149-154.
- 426 (27) Armstrong LE, Maresh CM, Castellani JW et al. Urinary indices of hydration status. *Int J Sport Nutr*
427 1994;4:265-279.
- 428 (28) Chevront SN, Ely BR, Kenefick RW, Sawka MN. Biological variation and diagnostic accuracy of
429 dehydration assessment markers. *Am J Clin Nutr* 2010;92:565-573.
- 430 (29) Fortes MB, Diment BC, Di Felice U, Walsh NP. Dehydration decreases saliva antimicrobial proteins
431 important for mucosal immunity. *Appl Physiol Nutr Metab* 2012;37:850-859.
- 432 (30) Banoo S, Bell D, Bossuyt P et al. Evaluation of diagnostic tests for infectious diseases: general
433 principles. *Nat Rev Microbiol* 2008;6:S16-S26.
- 434 (31) Hanley JA, McNeil BJ. A method of comparing the areas under receiver operating characteristic
435 curves derived from the same cases. *Radiology* 1983;148:839-843.
- 436 (32) Stookey JD, Purser JL, Pieper CF, Cohen HJ. Plasma hypertonicity: another marker of frailty? *J Am*
437 *Geriatr Soc* 2004;52:1313-1320.
- 438 (33) Gorelick MH, Shaw KN, Murphy KO. Validity and reliability of clinical signs in the diagnosis of
439 dehydration in children. *Pediatrics* 1997;99:E6.
- 440 (34) Steiner MJ, DeWalt DA, Byerley JS. Is this child dehydrated? *JAMA* 2004;291:2746-2754.
- 441 (35) Sarhill N, Walsh D, Nelson K, Davis M. Evaluation and treatment of cancer-related fluid deficits:
442 volume depletion and dehydration. *Support Care Cancer* 2001;9:408-419.
- 443 (36) Chevront SN, Kenefick RW. Dehydration: Physiology, Assessment and Performance Effects.
444 *Comprehensive Physiology* 2014;4:257-285.
- 445 (37) Walsh NP, Fortes MB, Raymond-Barker P et al. Is whole-body hydration an important consideration
446 in dry eye? *Invest Ophthalmol Vis Sci* 2012;53:6622-6627.
- 447 (38) Steiner MJ, Nager AL, Wang VJ. Urine specific gravity and other urinary indices: inaccurate tests
448 for dehydration. *Pediatr Emerg Care* 2007;23:298-303.
- 449 (39) Obuchowski NA, Lieber ML, Wians FH, Jr. ROC curves in clinical chemistry: uses, misuses, and
450 possible solutions. *Clin Chem* 2004;50:1118-1125.
- 451 (40) Woods DL, Mentis JC. Spit: saliva in nursing research, uses and methodological considerations in
452 older adults. *Biol Res Nurs* 2011;13:320-327.
- 453 (41) Miller CS, Foley JD, Bailey AL et al. Current developments in salivary diagnostics. *Biomark Med*
454 2010;4:171-189.

- 455 (42) Benelli U, Nardi M, Posarelli C, Albert TG. Tear osmolarity measurement using the TearLab
456 Osmolarity System in the assessment of dry eye treatment effectiveness. *Cont Lens Anterior Eye*
457 2010;33:61-67.
- 458 (43) Fortes MB, Diment BC, Di Felice U et al. Tear fluid osmolarity as a potential marker of hydration
459 status. *Med Sci Sports Exerc* 2011;43:1590-1597.
- 460 (44) Ely BR, Chevront SN, Kenefick RW, Sawka MN. Limitations of salivary osmolality as a marker of
461 hydration status. *Med Sci Sports Exerc* 2011;43:1080-1084.
- 462 (45) Perrier E, Demazieres A, Girard N et al. Circadian variation and responsiveness of hydration
463 biomarkers to changes in daily water intake. *Eur J Appl Physiol* 2013;113:2143-2151.

464

465

466

467

468

469

470

471

472 **FIGURE LEGENDS:**

473

474 **Figure 1.** Participant flow through the study. BUN, blood urea nitrogen.

475

476

477 **Figure 2.** ROC curve comparison between clinical physical signs, saliva and urine indices for the
478 assessment of dehydration. Data are shown for both forms of dehydration combined (A), water-loss
479 dehydration only (B) and water-and-solute-loss dehydration (C). The cut-off that provides the optimum
480 discrimination between sensitivity (true positive rate) and specificity (false positive rate) is plotted. BP, low
481 systolic blood pressure; SE, sunken eyes; CR, capillary refill time; Tc, Tachycardia; AD, axillary dryness,
482 ST, skin turgor, DM, dry mucous membrane; Sosm, saliva osmolality; SF, saliva flow rate; UrC, urine color;
483 USG, urine specific gravity. Vertical error lines represent sensitivity 95% CI, horizontal error lines represent
484 specificity 95% CI.

485

Table 1

Table 1. Group data for age, blood reference tests and urine and saliva index tests of hydration.					
		Water-loss only dehydrated (<i>n</i> = 27)	Water-and-solute-loss only dehydrated (<i>n</i> = 25)	Euhydrated controls (<i>n</i> = 78)	<i>P</i> value (one-way ANOVA)
	Age (Yr)	78.3 (9.6)	80.1 (9.6)	76.3 (7.7)	0.14
Reference tests	Plasma osmolality (mOsm/kg)	299 (6)†	283 (6)	283 (9)	<0.001
	BUN:Cr	18.8 (5.5)	24.3 (4.7)‡	15.7 (2.6)	<0.001
Index tests	Urine specific gravity	1.017 (0.006)	1.016 (0.007)	1.016 (0.006)	0.77
	Urine color	4.1 (1.6)	3.9 (1.8)	3.9 (1.7)	0.87
	Saliva flow rate (µl/min)	56 (55)	86 (183)	77 (90)	0.57
	Saliva osmolality (mOsm/kg)	136 (58)*	140 (66)*	92 (45)	<0.001
Values represent mean (standard deviation). BUN:Cr; blood urea nitrogen to creatinine ratio.					
† Significantly greater than water-and-solute-loss only dehydrated and euhydrated control groups (<i>P</i> < 0.001).					
‡ Significantly greater than water-loss only dehydrated and euhydrated control groups (<i>P</i> < 0.001).					
* Significantly greater than euhydrated control group (<i>P</i> < 0.01).					

Table 2

Table 2. Diagnostic accuracy of clinical signs to determine both forms of dehydration in combination, and separately (water-loss only, and water-and-solute loss dehydration) in older adults >60yr. Values in parentheses represent 95% Confidence intervals.													
Clinical assessment	All dehydration				Water-loss only dehydration				Water-and-solute-loss only dehydration				
	AUC _{ROC}	Positive LR	Negative LR	Diagnostic OR	AUC _{ROC}	Positive LR	Negative LR	Diagnostic OR	AUC _{ROC}	Positive LR	Negative LR	Diagnostic OR	
Low systolic BP (< 100 mmHg)	0.53	6.0	0.9	6.4	0.49	N/A	1	N/A	0.57	12.5	0.9	14.7*	
	(0.43-0.64)	(0.7-54.2)	(0.8-1.0)	(0.7-59.1)	(0.37-0.62)	N/A	(1)	N/A	(0.44-0.71)	(1.5-107.6)	(0.7-1.0)	(1.6-138.3)	
Tachycardia (HR > 100 bpm)	0.50	1.0	1.0	1.0	0.44	0.5	1.2	0.4	0.56	1.6	0.8	1.9	
	(0.40-0.60)	(0.5-1.9)	(0.8-1.2)	(0.4-2.3)	(0.32-0.56)	(0.2-1.5)	(1.0-1.4)	(0.1-1.5)	(0.43-0.70)	(0.8-3.0)	(0.6-1.1)	(0.7-5.0)	
Dry mucous membrane	0.51	1.1	1.0	1.1	0.51	1.0	1.0	1.0	0.52	1.1	0.9	1.2	
	(0.41-0.62)	(0.7-1.6)	(0.7-1.3)	(0.5-2.3)	(0.38-0.63)	(0.6-1.7)	(0.7-1.4)	(0.4-2.5)	(0.39-0.65)	(0.7-1.9)	(0.6-1.4)	(0.5-3.0)	
Axillary dryness	0.54	1.3	0.9	1.4	0.53	1.2	0.9	1.3	0.56	1.4	0.8	1.7	
	(0.44-0.64)	(0.8-2.0)	(0.7-1.2)	(0.7-3.0)	(0.40-0.65)	(0.6-2.1)	(0.7-1.3)	(0.5-3.1)	(0.43-0.70)	(0.8-2.4)	(0.6-1.2)	(0.7-4.2)	
Poor skin turgor	0.55	1.3	0.9	1.5	0.53	1.2	0.9	1.3	0.57	1.4	0.8	1.7	
	(0.45-0.65)	(0.8-2.0)	(0.6-1.1)	(0.7-3.1)	(0.40-0.66)	(0.7-2.0)	(0.6-1.3)	(0.5-3.2)	(0.44-0.70)	(0.8-2.3)	(0.5-1.2)	(0.7-4.3)	
Sunken eyes	0.51	1.2	1.0	1.2	0.43	0.6	1.1	0.5	0.56	1.9	0.9	2.2	
	(0.41-0.62)	(0.5-2.8)	(0.8-1.1)	(0.5-3.4)	(0.35-0.60)	(0.1-2.5)	(0.9-1.2)	(0.1-2.7)	(0.42-0.69)	(0.8-4.6)	(0.7-1.1)	(0.7-6.7)	
Capillary refill > 2 S	0.50	1.0	1.0	1.0	0.52	1.2	1.0	1.2	0.48	0.8	1.0	0.8	
	(0.40-0.60)	(0.5-2.0)	(0.8-1.2)	(0.4-2.4)	(0.39-0.64)	(0.5-2.7)	(0.8-1.2)	(0.5-3.5)	(0.36-0.61)	(0.3-2.3)	(0.9-1.3)	(0.2-2.7)	

AUR_{ROC}, area under the receiver operating characteristic curve; LR, likelihood ratio; OR, odds ratio; BP, blood pressure; HR, heart rate; * $P < 0.05$ significantly associated with hydration status by logistic regression analysis. N/A, not assessed as sensitivity was 0%.

Table 3

Table 3. Receiver operating characteristic (ROC) area under the curve (AUC) analysis for urine and saliva indices for the detection of dehydration in older adults (>60yr).			
		ROC analysis	
		AUC	P value
All dehydration			
	USG	0.53	0.67
		(0.39-0.66)	
	Urine color	0.52	0.79
		(0.39-0.65)	
	Saliva flow rate	0.56	0.25
		(0.46-0.66)	
	Saliva osmolality	0.76	<0.001
		(0.66-0.86)	
Water loss only dehydration			
	USG	0.55	0.53
		(0.39-0.72)	
	Urine color	0.54	0.61
		(0.38-0.70)	
	Saliva flow rate	0.55	0.46
		(0.43-0.67)	
	Saliva osmolality	0.76	<0.001
		(0.66-0.87)	
Water and solute loss dehydration			
	USG	0.50	0.98
		(0.32-0.69)	
	Urine color	0.49	0.91
		(0.31-0.67)	
	Saliva flow rate	0.57	0.28
		(0.44-0.71)	
	Saliva osmolality	0.76	0.001
		(0.62-0.89)	

Values in parentheses represent 95% confidence intervals. USG, urine specific gravity.

Table 4

Table 4. Diagnostic accuracy of saliva osmolality to determine both forms of dehydration in combination, and separately (water-loss only, and water-and-solute loss dehydration) in older adults >60yr.				
	Diagnostic accuracy	Positive LR	Negative LR	Diagnostic OR
Both forms of dehydration combined	71%	2.4	0.4	6.9
	(63-80%)	(1.6-3.6)	(0.2-0.6)	(2.8-17.5)
Water-loss dehydration	69%	2.2	0.4	5.0
	(59-79%)	(1.4-3.5)	(0.2-0.9)	(1.7-15.1)
Water-and-solute-loss dehydration	73%	2.8	0.3	8.9
	(63-83%)	(1.7-4.4)	(0.1-0.8)	(2.5-30.7)
Values in parentheses represent 95% Confidence intervals. LR, likelihood ratio; OR, odds ratio.				

Figure 1







Figure 2



Manuscript # JAMDA-D-14-00269, "Is this elderly patient dehydrated? Diagnostic accuracy of hydration assessment using physical signs, urine and saliva markers" – Response to reviewers

Reviewer #1:

Many thanks for your very insightful and constructive comments. Below are our responses to your comments, and changes within the manuscript are **highlighted in red text**. We hope you agree that these small changes have improved the manuscript.

- 1) Only 42% of screened individuals entered the study, and 31% had sufficient parameters to be analyzed. Although you show numbers in Figure 1, please comment on the large number of excluded subjects. Doesn't this affect the usefulness of the screening in clinical practice?

Thanks. As you have noted, of those that were screened (n =420), a large number of participants were excluded (58%). However, it should be noted that these were excluded due to ethical reasons of conducting the research as stipulated to us by the ethics committee (e.g. patients unable to provide consent (incapacity) for the research study (n=98), or that the research should not interfere with routine care of the patient and in those who had already begun treatment (n = 88)), or because participants declined to take part (n = 54). We have now included this information at the start of the results sections, along with percentages of those excluded (lines 197-199). In terms of the application of the usefulness in clinical practice, the reasons outlined above do not preclude the usefulness of the measures in the current study being used in clinical practice (i.e. in all patients admitted to hospital).

In light of this being a proof of concept study for saliva indices, we did in this instance exclude participants who had potential confounding effects on saliva (e.g. oral trauma, recent dental surgery, swallowing problems etc), although it should be noted that only 2 participants were excluded for this (both had swallowing problems), and in light of your excellent point, we have now added this information to the results (lines 198-199,) and Figure 1, and have also now acknowledged in the discussion that future studies should investigate whether saliva indices have utility, in patients with oral related problems (please see lines 341-344). Thanks.

- 2) "and allowing for an approximate one-third exclusion rate from data analysis (due to missing reference tests, and co-morbidities that preclude the use of the reference standards), a total of 178 participants were recruited into the study." It appears that the exclusion rate was higher than anticipated? Please comment.

Please note that the allowance for the one third exclusion rate (for missing reference tests and co-morbidities that affected the reference standards), was for those who might be excluded from the data analysis after they were already recruited into the study (i.e. n = 178 recruited). The N for which we analyzed data was n = 130, with 48 excluded from the data analysis. The proportion excluded from analysis of those recruited (48/178, 27%) is therefore actually lower, not higher than the anticipated 1/3 exclusion rate. Thanks.

- 3) "participants with a history of renal disease (n = 24), or who were in cardiac failure (n = 1) were excluded from data analysis." Please specify the criteria for renal disease and cardiac failure. What level of renal disease (stage?) or creatinine or other. For CHF, only "history"

or other criteria? As you point out, the presence of renal disease, starvation, malnutrition (among others) limit the usefulness of the BUN/Cr ratio. It would be useful to discuss the level of renal disease that you excluded.

Thanks. For this study, we excluded from data analysis, all participants who had any known previous history of renal disease (CKD stage 1-5) or if they were in cardiac failure as diagnosed by the clinician. We have now clarified this and added this information to the methods section (Line 165). In line with comment 1 above, we have also now added a sentence to the discussion where we discuss how future studies should investigate the utility of these indices in these relatively small populations (lines 318-320). Thanks.

- 4) Please discuss relationship between saliva and blood osmolality. If the values are generally highly correlated, is there any benefit in using saliva rather than blood. Is it quicker, cheaper, easier to use saliva. Given a paucity of saliva in 25% of subjects, should blood be favored?

This is a very helpful observation and the changes we have made (described below) in response have improved the take home message of the manuscript. Many thanks.

As we have addressed in the manuscript (lines 55-61, 169-171 and in Figure 1), plasma osmolality is elevated in, and will only detect water-loss dehydration. In water-and-solute loss dehydration, plasma osmolality is either normal or low, and thus has no diagnostic utility for this type of dehydration. Given the differential response of plasma osmolality to these two types of dehydration, we feel it would be inappropriate to report, or rely on the correlation between saliva and plasma osmolality to determine saliva's utility as a diagnostic method. In the current study, saliva osmolality was able to detect a proportion of patients with water-and-solute dehydration (sensitivity 78%), and is an easier to perform and non-invasive so has advantages over blood sampling.

With this in mind, based on your excellent point, as this limitation of plasma (blood) osmolality for detecting water-and-solute-loss only dehydration was not as prominent as it should be in the manuscript, we have now added a sentence to the end of the abstract (lines 29-31) and to the discussion and conclusion where we address this (lines 312-314, 351-352).

We were able to collect a quantity of saliva in 126/130 patients (97%)- reported on lines 210 and 328, although as we have stated (lines 213, 327) we only had adequate saliva (at least 25ul) to assess osmolality using our osmometer in 75% of samples. However, we have addressed this limitation in the discussion, (line 327-333) where we say that micro osmometers are in development that can assess osmolality on nano-gram quantities. We hope you feel that this is adequately addressed in the manuscript. Thanks.

Reviewer #2:

Many thanks for reviewing our manuscript.