

**Bangor University** 

DOCTOR OF PHILOSOPHY

Investigation of the role of mesoderm inducing genes in a cell culture model for the progression of colorectal cancer

Sarkar, Debalina

Award date: 2010

Awarding institution: Bangor University

Link to publication

#### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

• Users may download and print one copy of any publication from the public portal for the purpose of private study or research.

You may not further distribute the material or use it for any profit-making activity or commercial gain
You may freely distribute the URL identifying the publication in the public portal ?

# Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

# Investigation of the role of mesoderm inducing genes in a cell culture model for the progression of colorectal cancer

Debalina Sarkar

Bangor University School of Biological Sciences

PhD Thesis Submitted: 31<sup>st</sup> May 2010

Investigation of the role of mesoderm inducing genes in a cell culture model for the progression of colorectal cancer

Debalina Sarkar



#### Abstract

Cancer stem cells (CSC) are initiating cells in many cancer types that have similar properties like normal embryonic stem cells. Colorectal tumours can undergo epithelial to mesenchymal transition (EMT)-like processes at the invasive front, rendering them capable of invasion and metastasis. Recently, a possible link between cells undergoing EMT and cells with stem cell like properties has been identified in mammary Cancer stem cells (CSC). It is of fundamental importance to understand molecular events leading to establishment and maintenance of cancer initiating cells and how these relate to cellular transitions during tumourigenesis. However it is difficult to observe these processes *in vivo*, not least due to the transient nature of events during tumourigenesis.

We used an *in vitro* system to recapitulate changes occuring in CRC cells at the invasive front (mesenchymal-like cells) and central mass (epithelial-like cells) of tumours. We show for the first time, that the mesoderm inducer *BRACHYURY* differentially influences expression of the pluripotency gene *NANOG*, through association of BRACHYURY with upstream regulatory elements in the *NANOG* promoter in CRC cells. Binding of BRACHYURY to the *NANOG* promoter directly influences expression of *NANOG* and is predominant in mesenchymal-like cancer cells. We also demonstrate that the oncogene,  $\beta$ -catenin is upstream of both *BRACHYURY* and *NANOG* and influences the expression of markers of CSCs in the mesenchymal-like cells, whose presence has previously been linked to poor patient prognosis. Finally, we show that a similar pathway exists in two other cell lines (NTERA-2 and T84).

Control Chapter Canta Section	Contents –	Chapters an	d Sections
-------------------------------	------------	-------------	------------

Page	Section
1	Chapter 1 – Introduction
2	1.1 The normal colon
2	1.1.1 Structure and function of the colon
3	1.1.2 Cell types in the small intestine
5	1.2.3 Localisation of APC and B-catenin in normal human colon
6	1.2 Wnt signalling pathway
6	1.2.1 APC
7	1.2.2 B-catenin
10	1.2.3 Wnt and Wnt target genes
11	1.2.4 Wnt pathway in development
11	1.2.5 Wnt signalling in normal colon
12	1.2.6 Wnt pathway in cancer and other diseases
13	134 Colorectal cancer, the disease
14	1.4.1 Sporadic colon cancer
14	1.4.2 Hereditary colon cancer
14	1.4.2A Familial adenomatous poliposis (FAP):
16	1.4.2B Hereditary non-polyposis coli cancer (HNPCC)
17	1.4.3 Genes of colon cancer
17	1.4.4 APC and $\beta$ -catenin mutation in other forms of cancer
18	1.5 Epithelial-mesenchymal transition (EMT)
19	1.5.1 Growth factors regulating EMT
20	1.5.2 EMT and development
21	1.5.3 EMT and cancer
24	1.6 Stem cells
24	1.6.1 Types of stem cells
26	1.6.2 Properties of stem cells:
26	1.6.3 Leukemia inhibitory factor (LIF) and stem cells
27	1.6.4 NANOG and maintenance of stem cells
28	1.6.5 Intestinal stem cells
29	1.6.6 A) Intestinal stem cell microenvironment
29	1.6.6 B) Two models for intestinal stem cells
32	1.6.7 Cancer stem cells
35	1.6.8 Regulatory pathways of stem cells and cancer cells
38	1.6.9 Colon cancer stem cell markers
39	1.7 SW480 colorectal cancer cells transitioning between low and high density,
	an <i>in vitro</i> model to study CRC
41	1.8 Aim of the project
43	Chapter 2 – Materials and Methods
44	2.1 Source of cell lines
44	2.2 Routine cell culture
44	2.3 Freezing cells
45	2.4 Phenol-Chloroform extraction of DNA

46	2.5 Ethanol precipitation of DNA
46	2.6 Cell counting by Haemocytometer
46	2.7 Total RNA isolation
47	2.8 cDNA synthesis by MLV method
47	2.9 cDNA synthesis by SperscriptIII
48	2.10 Real time PCR using Biomix <
49	2.11 Quantitative Real time RCR
52	2.12 Selection of control genes for qRT-PCR
52	2.13 An worked example for analyzing qRT-PCR data
54	2.14 Western blot /
57	2.15 Immunofluorescence
58	2.16 Immunoprecipitation
59	2.17 RNAi (BRACHYURY, β-catenin, E-CADHERIN, TBX-2, TBX-3, FGF8)
60	2.17A Mechanism of RNAi
61	2.18 ChIP (Chromatin Immunoprecipitation)
63	2.19 DNA cloning and Luciferase assay
66	2.20 P-value and t-test
67	2.21 Mitotic index count
68	2.22 Cell Viability Asssay (Promega CellTiter-Glo <sup>™</sup> )
68	2.23 Cytoselect cell migration assay
69	2.24 Conditioned media experiment
69	2.25 Protein purification
	1
71	Chapter 3-A) The mesoderm inducer <b>BRACHYURY</b> is differentially
	expressed in colorectal cancer cells
72	3.1 Introduction
75	3.2 Validation of SW480 cell line for use as a model to study EMT
77	3.3 Screen of the mesoderm inducing genes by RT-PCR
77	3.4 qRT-PCR screen
87	Chapter 3-B) Localization of BRACHYURY in SW480 cells
89	3.5 Discussion
92	Chapter 4 – Determining the role of <i>BRACHYURY</i> in SW480 cells
93	a) Does <i>BRACHYURY</i> play a role in inducing EMT in the colorectal cancer
	cell line SW480?
93	4.1.1 A role for <i>BRACHYURY</i> in inducing EMT?
(97)	4.1.2 qRT-PCR screen to study the possible effects of BRACHYURY on four
	different categories of genes
99	4.1.2A Exploring a role for <i>BRACHYURY</i> in influencing the expression of
	influencing the expression of epithelial cell polarity markers:
101	
	4.1.2B Effect of BRACHYURY on the expression of a marker of intestinal
	4.1.2B Effect of <i>BRACHYURY</i> on the expression of a marker of intestinal epithelium
102	<ul><li>4.1.2B Effect of <i>BRACHYURY</i> on the expression of a marker of intestinal epithelium</li><li>4.1.2C Role of <i>BRACHYURY</i> in regulating expression of known target genes</li></ul>
102 105	<ul> <li>4.1.2B Effect of <i>BRACHYURY</i> on the expression of a marker of intestinal epithelium</li> <li>4.1.2C Role of <i>BRACHYURY</i> in regulating expression of known target genes</li> <li>4.1.2D Effect of <i>BRACHYURY</i> RNAi on inducer of EMT, <i>E-cadherin</i>, <i>Snail</i></li> </ul>

b) Role of *BRACHYURY* in influencing the regulation of *NANOG* and

maintaining a putative cancer stem cell population

109 4.2.1 The expression of *NANOG* is altered in response to changes in *BRACHYURY* in low density SW480 cells

- 4.2.2 Does *NANOG* regulate specific gene expression patterns in SW480 cells
- 120 c) A role of *BRACHYURY* in regulating the NF $\kappa$ B pathway?
- 4.3 A role for *BRACHYURY* in regulating the NFκB pathway
- d) Role of *BRACHYURY* in cell proliferation
- 124 4.4.1 Cell proliferation and cancer
- 124 4.4.2 Cell proliferation by mitotic index count
- 128 4.4.3 Cell proliferation by CellTiter-Glo Luminescent Cell Viability Assay
- e) Role of *BRACHYURY* in cell migration
- 134 4.5 Role of *BRACHYURY* in cell migration
- 138 Discussion
- 140 Chapter 5 The binding of BRACHYURY to the *NANOG* promoter region is dependent on β-catenin in SW480 cells
- 141 5.1 Introduction
- 144 5.2 Identification of BRACHYURY binding sites in the human *NANOG* promoter region
- 144 5.3 A) BRACHYURY binds to the NANOG promoter in human SW480 cells
- 148 5.3B) NANOG promoter sequence for Brachyury binding (human)
- 149 5.4 BRACHYURY binding to the NANOG promoter depends on β-catenin
- 151 5.5 BRACHYURY binding sites act as BRACHYURY-dependent enhancers of expression from the NANOG promoter
- 155 5.6 Discussion

# 157 Chapter 6 – The role of *BRACHYURY* in NTERA-2, SW626 and T84 cells

- 158 6.1 Role of *BRACHYURY* in NTERA-2 cells
- 159 6.2 Role of *BRACHYURY* in T84 cells
- 161 6.3 Role of *BRACHYURY* in SW626 cells
- 165 6.4 Discussion
- 166 Chapter 7 Preliminary characterization of an extracellular factor with BRACHYURY regulatory activity
- 167 7.1 Introduction
- 167 7.2 Results
- 174 7.3 Discussion

#### 175 Chapter 8 – Summary

- 176 8.1 SW480 cell line is a validated in vitro model to study molecular events of EMT
- 177 8.2 Why was BRACHYURY studied?
- 178 8.3 Possible roles of *BRACHYURY* in SW480 cells
- 180 8.4 BRACHYURY regulates NANOG
- 181 8.5  $\beta$ -catenin is upstream of BRACHYURY
- 181 8.6 BRACHYURY binds to NANOG promoter region in SW480 cells

- 183
- 8.7 *BRACHYURY* expression in other tumour types8.8 Conditioned media 'factor' might be responsible for differential expression 184 of BRACHYURY in SW480 cells
- 8.9 A model pathway derived 185

188-	References
238	
239-	Appendix

428

List of figures and tables:

Page	Figures, Tables and Graphs
4	Figure 1.1
9	Figure 1.2
15	Figure 1.3
22	Figure 1.4
30	Figure 1.5
33	Figure 1.6
36	Figure 1.7
45	Table 2.1
48	Table 2.2
50	Table 2.3
53	Table 2.4
55	Table 2.5
56	Table 2.6
59	Table 2.7
60	Table 2.8
62	Table 2.9
63	Table 2.10
67	Table 2.11
73	Table 3.1
76	Figure 3.1
78	Table 3.2
80	Figure 3.2
81	Figure 3.3
82	Table 3.3
83	Figure 3.4
84	Figure 3.5
85	Figure 3.6
86	Figure 3.7
87	Figure 3.8

95	Figure 4.1
96	Figure 4.2
98	Figure 4.3
100	Figure 4.4
103	Figure 4.5
106	Figure 4.6
107	Figure 4.7
111	Figure 4.8
113	Figure 4.9
117	Figure 4.10
118	Figure 4.11
121	Figure 4.12
123	Figure 4.13
126	Figure 4.14
127	Figure 4.15
129	Figure 4.16
133	Figure 4.17
136	Figure 4.18
137	Figure 4.19
143	Figure 5.1
145	Table 5.1
147	Figure 5.2
150	Figure 5.3
152	Figure 5.4
154	Figure 5.5
160	Figure 6.1
162	Figure 6.2
164	Figure 6.3
169	Figure 7.1
171	Figure 7.2
173	Figure 7.3
187	Figure 8.1

#### Abbreviations

AEBSF - 4 (2-aminoethyl) benzenesulfonyl fluoride ALCAM - Activated leukocyte cell adhesion molecule APC - Adenomatous Polyposis Coli APCL - Adenomatous Polyposis Coli Like ACF - Aberrant Crypt Foci Ascl2 - Achaete scute-like 2 BCL - B-cell lymphoma BMP - Bone Morphogenetic Protein CAPS - 3-(cyclohexylamino)-1-propane sulfonic acid CBC - Crypt base columnar CKI - Casein Kinase I CKIB - Casein Kinase I beta CCK - Cholecystokinin ChIP- Chromatin Immunoprecipitation CRC - Colorectal Cancer CRM1 - Chromosome Maintenance Region 1 CSC - Cancer stem cells DEP - Dishevelled, Eg-10, plekstin DIC - Differential Interference Contrast DIX - Dishevelled and Axin DLG - Discs Large DMEM - Dulbecco's Modified Eagle's Medium DMSO - Dimethlysulphoxide DNA - Deoxyribonucleic Acid Dsh - Dishevelled DTT – Dithiothrietol Dvl-Dishevelled EC – Embryonal Carcinoma ECACC - European Collection of Cell Cultures EDTA - (Ethylenedinitrilo) tetra acetic acid EMT - Epithelial-mesenchymal transition **EPH** - Ephrin receptors FAP – Familial Adenomatous Polyposis FBS - Foetal Bovine Serum FOX - Forkhead box GBP - GSK Binding Protein GSK - Glycogen Synthase Kinase GSK3β - Glycogen Synthase Kinase 3 beta HCC - Hepatocellular carcinoma HGF - Hepatocyte growth factor HMBA - Hexamethylene Bisacetamide HNPCC - Hereditary Non-Polyposis Coli Cancer HSC - Haematopoietic stem cell Ihh - Indian hedgehog

IL1 $\beta$  - Interleukin 1 $\beta$ IPSC- Induced pluripotent stem cells ISC - Intestinal stem cells LEF – Lymphocyte Enhancer Factor Lgr5 - leucine-rich repeat-containing G protein coupled receptor 5 LIF - Leukemia inhibitory factor LMB - Leptomycin B LOH - Loss of heterozygosity LRP - LDL (Low Density Lipoprotein) Receptor Related Protein MAPK - Mitogen activated protein kinase MDCK - Madin-Darby Canine Kidney Cells MCR - Mutation Cluster Region MEK - Mitogen-activated protein kinase/extracellular signal-regulated kinase MEM – Minimum Eagle's Medium MET - Mesenchymal to epithelial transition MLV - Murine leukemia virus MMP - Matrix Metalloproteinase MMR - Mismatch Repair mRNA - Messanger Ribonucleic Acid NES - Nuclear Export Sequence NFκB - Nuclear factor κB NLS - Nuclear Localisation Sequence NOD/SCID - non-obese diabetes/severe combined immunodeficiency OSE - Ovarian surface epithelial PBS - Phosphate Buffered Saline PCR – Polymerase Chain Reaction PDGF - Platelet-derived growth factor PDZ-PSD-95/DLG/ZO-1 PI - Propidium iodide PI3K - Phosphoinositide 3-kinase PP2A – Protein Phosphatase 2A RNA - Ribonucleic Acid RT - Reverse Transcription SCF – SKP-1, Cdc53, F-box SDS-PAGE - Sodium Dodecyl Sulphate - Polyacrylamide Gel Electrophoresis Ser - Serine Shh - Sonic hedgehog SHIP - SH2 domain-containing inositol phosphatase SMAD - Mothers against decapentaplegic (MAD) + C. elegans protein SMA STAT3 - Signal transducer and activator of transcription 3 TA - Transit amplifying TAD - Transcriptional activation domain TBM - Tubular basement membrane TCF - T-Cell Factor TGFβ - Transforming growth factor beta TLE - Transducing-Like Enhancer of split

# Gene names and Accession numbers

Gene name	Abbreviation	NCBI accession number
BRACHYURY	Т	NM_003181
FIBROBLAST GROWTH FACTOR 4	FGF4	NM_002007
FIBROBLAST GROWTH FACTOR RECEPTOR 1	FGFR1	NM_023110
FIBROBLAST GROWTH FACTOR RECEPTOR 2	FGFR2	NM_000141
FIBROBLAST GROWTH FACTOR RECEPTOR 3	FGFR3	NM 000142
OCTAMER-4	Oct-4	NM 002701
NODAL	NODAL	NM 018055
FIBROBLAST GROWTH FACTOR 8	FGF8	NM 033165
GROWTH DIFFERENTIATION FACTOR-3	GDF3	NM 020634
ACTIVINA	ACTIVIN A	NG 008004
ACTICIN B	ACTICIN B	NM 002193
BONE MORPHOGENETIC PROTEIN 4	BMP4	NM 001202
T-BOX TRANSCRIPTION FACTOR 2	TBX2	NM 005994
T-BOX TRANSCRIPTION FACTOR 3	TBX3	NM 005996
FIBROBLAST GROWTH FACTOR 19	FGF19	NM 005117
NUCLEAR FACTOR KAPPA-LIGHT-CHAIN-ENHANCER	ΝΕκΒ	NM 003998
OF ACTIVATED B CELLS		
SNAIL.	SNA	NM 005985
E-CADHERIN	CDH1	NM 004360
CAUDAL TYPE HOMEOBOX TRANSCRIPTION FACTOR 2	CDX2	NM 001265
CRUMBS HOMOLOG 3	CRB3	NM 139161
PROTEIN ASSOCIATED WITH LIN SEVEN 1	PALSI	AF397170
ZINC FINGER F-ROX-RINDING HOMEOBOX 1	ZEB1	NM 001128128
ZINC FINGER E-BOX-BINDING HOMEOBOX 2	ZEB2	NM 014795
WINGLESS-TYPE MMTV INTEGRATION SITE FAMILY	WNT11	NM 004626
MEMBER 11	1121000	_
EPITHELIAL V-LIKE ANTIGEN 1	EVA1	AF275945
INTERLEUKIN-2	IL2	NM 000586
INTERLEUKIN-15	IL15	NM 172174
ORTHOPEDIA	OTP	NM 032109
STRATIFIN	14-3-3 sigma	NM 006142
NOTCH HOMOLOG 1	NOTCH1	NM 017617
NOTCH HOMOLOG 2	NOTCH2	NM 024408
NOTCH HOMOLOG 3	NOTCH3	NM 000435
NOTCH HOMOLOG 4	NOTCH4	NM_004557
B-catenin	CTNNB1	NM 001904
SIGNAL TRANSDUCER AND ACTIVATOR OF	STAT3	NM_139276
I KANSCRIPTION 5	11.10	NIM 000572
INTERLEUMIN-10		NM_002659
PLASMINOGEN ACTIVATOR UROKINASE RECEPTOR	DCL 2 LIVE 1	NIM 138578
BULZLI	SOV2	NM 003106
SKY (SEX DETERMINING REGION 1)-BOX 2	CATA2	NM 032638
GATA BINDING PROTEIN 2(GLOBIN TRANSCRIPTION FACTOR 2)	GATAZ	052036
GATA BINDING PROTEIN 2(GLOBIN TRANSCRIPTION FACTOR 2)	GATA4	NM_002052
GATA BINDING PROTEIN 6(GLOBIN TRANSCRIPTION FACTOR 6)	GATA6	NM_005257
LAMIN B	LMNB	NM 005573
LAMIN A/C	LMNA/C	NM 170707
FORKHEAD BOX D3	FOXD3	NM 012183
C-MYC	МҮС	NM 002467
VIMENTIN	VIMENTIN	NM 003380

#### Acknowledgements

Funding for this work was provided by Tenovus Cancer Research and Gwynedd Haematology and Cancer Research Fund. The cell line NTERA-2 (clone D1) was a gift from P.W. Andrews (University of Sheffield). Thanks to everyone in lab G30, especially Soshila Ramayah, for being helpful and supportive, especially Dr. Melanie Davies for her help with most of the laboratory techniques. Thanks to Dr. J.A. Wakeman for her kind support, being a wonderful supervisor and for proof reading this thesis. I would also like to thank Dr. Ramsay MacFarlane and Dr. Jürgen Müller for their valuable advice on the project. Last but not the least, thanks to my parents, my sister and my fiancé for their enormous support and patience all through, without which I could not have achieved this piece of work.

# Chapter 1 - Introduction

#### Colorectal cancer, an overview:

Colorectal cancer (CRC) is the cancerous growth of the colon, rectum and appendix. It is the second most prevalent cancer in the Western world causing over 655,000 deaths per year. The tumour suppressor protein adenomatous polyposis coli (APC) is mutated in most cases of sporadic colorectal cancer; specifically mutations in APC have been found in 85% of the cases of nonhereditary colon cancers (Tucker and Pignetelli, 2000). APC has been described as 'the gene for colon cancer', describing its importance in this disease. CRC can be both hereditary (genetic) and sporadic (Non-heriditary) and 75% of CRCs are sporadic. Familial adenomatous polyposis (FAP) and hereditary non-polyposis coli cancer (HNPCC) are two non-sporadic forms of CRC. 95% of colorectal cancers are carcinomas, of which 95% are adenocarcinomas. Adenocarcinomas are generally considered to arise from adenomas. Adenomatous polyps are benign tumours which become malignant following malignant transformation. (Fenoglio et al., 1974; Morson et al., 1974; Shinya et al., 1979). There are many risk factors associated with colorectal cancer including family history of colon cancer, high fat diet (Potter et al. 1986), sedentary lifestyle (Slattery et al. 1988; White et al., 1996; Friedenreich et al., 2001) and smoking (Chao et al., 2000). The major signalling pathway altered in the event of colon cancer is the canonical Wnt signalling pathway and  $\beta$ -catenin is the key oncogene in this pathway. The canonical Wnt signalling pathway is described in section 1.2 of this chapter.

# 1.1 The normal colon

#### 1.1.1 Structure and function of the colon

The colon, also known as the intestine is located in the lower abdomen in humans and is an essential part of the gastrointestinal tract. The function of the colon is predominantly to absorb food and nutrients, to reclaim water and to store waste products before excretion. Finger-like projections, known as villi, protrude out of the walls of the small intestine with additional extensions, called the microvilli, which serve to increase the surface area of the intestine and increase its absorptive activity (Figure 1.1A). In the intestinal epithelial lining, glands called the crypts of Lieberkühn (or intestinal glands or intestinal crypts) are found which secrete many enzymes.

#### 1.1.2 Cell types in the small intestine

The epithelial lining of the small intestine consists of many distinct cell types to maintain normal structure and function of the tissue. The relative number and position of the cells are precisely regulated via exchange of signals between cells within the tissue. The common cell types of the intestinal epithelium (Figure 1.1B) are as follows:

**Enterocytes or absorptive cells:** These tall, columnar cells with microvilli are situated at the basal nucleus and are involved in the transport of substances. These cells are attached to each other and other cell types by junctional complexes (zonula occludens or tight junction, zonula adherens, and macula adherens). These cells are also involved in the absorption of amino acids, monosaccharides, monoglycerides and fatty acids across the microvilli membranes. These cells are continuously generated in the crypts to replace the cells at the tip of the villi. The enterocytes have a lifespan of about 5-6 days.

**Goblet cells:** These cells are mucus-secreting and absorptive; and are the second most abundant epithelial cells, found interspersed among the other cell types. The base of these cells holds the nucleus and organelles. These cells have lifespan of 5-6 days.

**Paneth cells:** These cells are located at the base of the crypts of Lieberkuhn. They have an oval basal nucleus and large, acidophilic granules at their apical end. The granules contain the antibacterial enzyme lysozyme, glycoproteins and zinc which are required for correct functioning of many enzymes. Paneth cells regulate the intestinal flora and have a lifespan of approximately four weeks.

3



Image taken from Nature Reviews, Genetics, 2006, vol7, 5

http://www.nature.com/nrg/journal/v7/n5/images/nrg1840-f1.jpg

# Figure 1.1

Intestinal villus and its cell types. A) Structure of the intestine in the crypt villi axis responsible for the renewal of its epithelium. Arrows show the upwards flow of cells out of the crypts. Stem cells lie at the crypt base. It is still not confirmed whether they are mixed with, or just above, the Paneth cells. Above the stem cells are the dividing progenitor cells, the transit-amplifying cells. At the top, lie the post-mitotic differentiated cells (which are the absorptive cells, goblet cells and enteroendocrine cells). B) Four types of terminally differentiated cells. Absorptive cells have a brush border and are at the apical surface. The other three types are all secretory and their apical cytoplasm is generally mucus-filled secretory granules.

**Enteroendocrine cells:** These cells are mostly found in the lower part of the crypts but can occur at all levels of the epithelium. They produce cholecystokinin (CCK), which stimulates pancreatic enzyme secretion and gall bladder contraction. They also secrete secret to stimulate pancreatic bicarbonate secretion and gastric inhibitory peptide, which inhibits gastric acid secretion.

**M or microfold cells:** These epithelial cells are found over Peyer's patches and other large lymphatic aggregations. They are flat shaped and their surface has many folds. They transport antigens to the nearby lymphoid cells where immune response to foreign antigens is regulated.

**Undifferentiated stem cells:** The crypt of the colon consists of a monolayer of epithelial cells. At the crypt base the stem cells are located which give rise to the progenitor cells which occupy the lower third of the crypt. Epithelial cells at the surface are renewed frequently as the progenitor cells migrate towards the surface where they differentiate. Differentiation to a goblet cell or enterocyte takes about 2 additional divisions and these cells then migrate from the crypt to the villus. At the tip of the villus, these epithelial cells finally shed off into the lumen by apoptosis and are replaced by the differentiating cells moving up the crypt. The entire process of renewal of the epithelial cell takes 3-5 days (Potten & Loeffler, 1990).

In addition to these cell types, the middle part of each villus contains a dense population of mesenchymal cell types which comprise of the stroma with pericytes, fibroblasts, machrophages and endothelial cells.

# 1.2.3 Localisation of APC and $\beta$ -catenin in normal human colon

APC is localised to the cytoplasm in differentiated epithelial cells above the crypt, and is localised to the nuclei of cells towards the base of crypts (Rosin-Arbesfeld *et al.*, 2003). Nuclear APC at the crypt base is related to the activation of the Wnt pathway in the proliferative cells of the crypt (van de Wetering *et al.*, 2002). Adenocarcinomas with truncating mutation of *APC* and loss of heterozygosity show nuclear localisation of APC (Rosin-Arbesfeld *et al.*, 2003). Studies show APC in the intestinal epithelium to be localised to the apical membrane (Miyashiro *et al.*, 1995; Midgley *et al.*, 1997; Reinacher-Schick & Gumbiner, 2001; Anderson *et al.*, 2002).; although the specificity of the antibodies used in these studies may lead one to question this apical localisation. Later, in 2005, *Brocardo* and co-workers confirmed that the polyclonal antibody used in 2003 by Rosin-Arbesfeld and co-workers was not reliable.

β-catenin is mostly found in the nuclei of cells within the proliferating compartment at the bottom third of crypts, with nuclear staining being strongest at the crypt base where cells are active for Wnt signallig (van de Wetering *et al.*, 2002). TCF-4, a TCF family member, is widely expressed in the intestinal epithelium and is required for the crypt stem cells (Korinek *et al.*, 1997). The βcatenin/TCF-4 complex is the key switch that controls proliferation versus differentiation in normal and malignant intestinal epithelial cells (van de Wetering *et al.*, 2002).

# 1.2 Wnt signalling pathway

The Wnt signalling pathway is of fundamental importance in development, promotion and maintenance of stem cells and also in disease; specifically, components of the canonical Wnt signalling pathway are critically altered in CRCs (Willert *et al.*, 2002). APC is a tumour suppressor protein, central to this pathway and  $\beta$ -catenin is a key oncogene in this pathway.

# 1.2.1 APC

Functions of the *APC* gene and its protein product include controlling the Wnt pathway, apoptosis, cell migration and adhesion and chromosomal segregation during mitosis (Fodde and Clevers 2001). The *APC* gene, located on the long arm of chromosome 5, consists of 15 exons and encodes a 312 kDa protein consisting of 2843 amino acids (Groden *et al.*, 1991; Kinzler *et al.*, 1991, Nishisho *et al.*, 1991). The central part of APC contains three repeats of 15 amino acids, and seven distinct repeats of 20 amino acids, both of which can bind

independently to  $\beta$ -catenin (Rubinfeld *et al.*, 1993; Rubinfeld *et al.*, 1995; Su *et al.*, 1993) but only the 20 amino acid repeats are involved in phosphorylation of  $\beta$ -catenin (Munemitsu *et al.*, 1995). APC can be directly phosphorylated by GSK-3 $\beta$  (Rubinfeld *et al.*, 1996) by binding to axin (Ikeda *et al.*, 2000). Phosphorylation of APC is important for degradation of  $\beta$ -catenin (Rubinfeld *et al.*, 1997). APC is a key player controling the degradation, subcellular localisation and transcriptional activity of  $\beta$ -catenin (Rosin-Arbesfeld *et al.*, 2003). APC is localised to kinetochores in mitotic cells (Kaplan *et al.*, 2001; Fodde *et al.*, 2001) and also in centrosomes in cancer cells (Tighe *et al.*, 2001) where it has a role in chromosome segregation. Truncated *APC* leads to increased abnormal mitoses and chromosomal aberration in embryonic stem cells (Fodde *et al.*, 2001) and many cancer cells (reviewed in Toncheva and Nacheva, 1998).

In a cell, when a gene allele loses its normal function in which the other allele was already inactivated, is known as loss of heterozygosity (LOH). In oncogenesis, after one allele of a tumour suppressor gene is inactivated, the mutation is then passed on to the zygote resulting in progeny that are heterozygous for that allele. LOH on chromosome 5q occurs for APC (Rosin-Arbesfeld *et al.*, 2003; Levy *et al.*, 1994; Lamlum *et al.*, 1999). LOH on chromosome 17p (17p3.1) occurs for the tumour suppressor protein, T53 and is found in 75% of CRCs (Forslunder *et al.*, 2001). Loss of the Smad family member 4, SMAD4 on chromosome 18q (18q21) is associated with missignalling of growth-suppressing transforming growth factor  $\beta$  (TGF- $\beta$ ) pathway. Smad4 mutations are found in 50-60% of LOH in CRC (Woodford-Richens *et al.*, 2001).

#### 1.2.2 β-catenin

 $\beta$ -catenin has role in intercellular adhesion alongside its function of transducing the Wnt signal. Its structure comprises of a 130 amino acid N-terminal domain, an armadillo repeat region of 12 imperfect repeats of 42 amino acids and a 100 amino acid C-terminal domain. The N-terminal domain is

involved in regulating its stability through a number of phosphorylation sites (three serines sites at S33, 37, 45 and one threonine site, T41) within this region. Deletions and certain point mutations of this region that alter these phosphorylation sites lead to activation of  $\beta$ -catenin (Yost *et al.*, 1996; Munemitsu *et al.*, 1996; Barth *et al.*, 1997).

In the absence of a Wnt signal, free cytoplasmic  $\beta$ -catenin is phosphorylated by CK1 and GSK3B in a multi-protein complex known as the degradation complex, consisting of axin (or its homologue conductin), GSK-3β  $(\Box \Box Glycogen synthase kinase 3 beta)$  and APC (reviewed in Bienz, 2002) (Figure 1.2A). The complex of axin and casein kinase (CK1) induces phosphorylation of  $\beta$ -catenin at Ser45 which initiates or primes phosphorylation of  $\beta$ -catenin at the other Wnt-dependent phosphorylation sites (Amit *et al.*, 2002; Hagen & Vidal-Puig, 2002; Liu et al., 2002; Sakanaka, 2002) and GSK-3β phosphorylation sites at Ser33, Ser37 and Thr41 (van Noort et al., 2002; Sadot et al., 2002). Once phosphorylated,  $\beta$ -catenin then interacts with the F-box protein  $\Box$   $\beta$ -TrCP (Slimb in *Drosophila*) (Hart *et al.*, 1999; Winston *et al.*, 1999; Liu *et al.*, 1999). The WD40 repeat domain of  $\beta$   $\beta$ -TrCP interacts specifically with the phosphorylated  $\beta$ -catenin and not with the unphosphorylated  $\beta$ catenin which makes the phosphorylated  $\beta$ -catenin ubiquitinated by this complex and this targets proteosomal degradation of  $\beta$ -catenin by 26S proteasome (reviewed in Maniatis 1999). Due to this degradation, β-catenin is not available to enter the nucleus, instead, it is mainly membrane bound as part of the cell adhesion complex with E-cadherin. In the absence of a Wnt signal, TCF/Lefs act as transcriptional repressors (Brannon et al., 1997; Bienz, 1998; Riese et al., 1997).

In the presence of a Wnt signal,  $\beta$ -catenin is stabilized and the destruction complex mentioned above is not formed. Rather, a complex is formed between dishevelled, GBP/Frat1, axin and Zw3/GSK-3 (Li *et al.*, 1999; Salic *et al.*, 2000; Farr *et al.*, 2000).  $\beta$ -catenin is released from this complex escaping phosphorylation and is therefore not targeted for proteosomal degradation (Salic

8



# Figure 1.2

Simplified diagram of the Wnt signalling pathway. A) In the absence of a Wnt signal,  $\beta$ -catenin is phosphorylated by casein kinase I $\beta$ /(CKI $\beta$ ) and Glycogen synthase kinase-3 (GSK-3 $\beta$ ), which enables its ubiquitination and proteasomal degradation. When a Wnt signal is present, DISHEVELLED proteins inhibit the phosphorylation of  $\beta$ -catenin which leads to nuclear entry of dephosphorylated  $\beta$ -catenin and activation of Wnt pathway target genes. B) Oncogenes and tumor suppressors of the Wnt pathway. Arrows or bars indicate activating or inhibitory effects, respectively. Green and red indicate proto-oncogenic and tumor suppressive activity, respectively, in human cancer.

*et al.*, 2000). The Wnt signal is initiated when Wnt ligands bind to Frizzled receptors, resulting in inhibition of the degradation complex via dishevelled (Cadigan & Nusse, 1997; Dale, 1998). Dishevelled proteins can directly interact with Axin (Li *et al.*, 1999) and phosphorylate it. The phosphorylation of Axin is reduced when there is a Wnt signal and this leads to decrease in the binding affinity of  $\beta$ -catenin to Axin which releases  $\beta$ -catenin from the degradation complex and leading to the levels of free  $\beta$ -catenin becoming raised in the cytoplasm (Willert *et al.*, 1999; Jho *et al.*, 1999). Dishevelled can also interact with casein kinase I $\beta$  (Peters *et al.*, 1999; Sakanaka *et al.*, 1999), which then acts to inhibit phosphorylation of  $\beta$ -catenin with its partner proteins, TCF/Lefs, allows them to act as transcriptional activators (reviewed in Bienz, 1998; Roose & Clevers, 1999) and leads to activation of Wnt pathway target genes (Figure 1.2).

#### 1.2.3 Wnt and Wnt target genes

So far, there are 19 Wnt genes identified in humans, 19 genes in mouse, 17 in Xenopus, 7 in Drosophila and 15 Wnt genes in Zebrafish (data taken from Wnt homepage, http://www.stanford.edu/~rnusse/wntwindow.html) discovered. In a microarray study in 2002 (Willert *et al.*), more than 50 genes were shown to be upregulated in response to Wnt signal, in human teratocarcinoma cells. These genes include *Cyclin D1*, *Ubc4/5E2*, *CBP/P300*, *MYC*, *ID2*,  $\beta$ -*TRCP*, *TLE/Groucho*, *MSX1*, *MSX2*, *Nucleophosmin*, *Frizzled*, *Follistatin* and *REST/NRSF* (Willert *et al.* 2002). Almost all Wnt target genes have putative TCF binding sites in their promoter region. Many of these target genes also respond to Bone morphogenic pathway (BMP) signaling pathway components along with Wnt pathway components. Wnt signaling is critical for activation of genes that promote stem cell fate and inhibit cellular differentiation and regulates a remarkable number of genes involved in its own signaling system (Willert *et al.*, 2002).

#### 1.2.4 Wnt pathway in development

The Wnt family of proteins is a group of highly conserved secreted signalling molecules. The Wnt pathway plays a key role during development of the embryo where these proteins regulate cell-to-cell interactions and also inhibit cellular differentiation. Mutations in Wnt genes or Wnt pathway molecules can lead to many developmental defects. In mouse, mutation in *Wnt-1* leads to loss of midbrain and cerebellum, mutation in *Wnt-2* leads to defects in the placenta, mutation in *Wnt-3A* enables loss of tailbud, mutation in *Wnt-4* and mutation of *Wnt-7A* gives rise to ventralization of limbs. In *Drosophila*, mutation in *wingless*, affects limb development. In *C. elegans, lin-44* mutation leads to defects in endoderm induction and spindle orientation (reviewed in Andreas Wodarz and Roel Nusse, 1998). Thus, Wnt signaling is fundamental to embryogenesis. However, as mentioned earlier, this pathway is also involved in many forms of cancer, including colorectal cancer.

#### 1.2.5 Wnt signalling in normal colon

The cells of the intestinal epithelium continuously self-renew and thereby act as a distinct model for the study of cell proliferation, differentiation, cell migration and also of cancer progression. At the bottom of the intestinal crypts, several Wnt factors are produced (Gregorieff *et al.*, 2005) which generate a morphogen-like gradient of Wnt signals along the crypt-villus axis. The Wnt signalling pathway drives the transcription of genes related to proliferation in the intestine such as *Sox9*, an intestinal crypt transcription factor (Blache *et al.*, 2004) and critical in maintaining the epithelial stem cells and early progenitor cells. The Wnt pathway also acts with other pathways, such as the Notch signalling to maintain the gut homeostasis and progenitor cells (reviewed in Fre *et al.*, 2009). There is evidence that the proliferating transit amplifying (TA) cells in the crypts are dependent on Wnt pathway stimulation and the accumulation of nuclear β-catenin by the progenitor cells at the crypt bottom indicates this stimulation (Van de Wetering *et al.*, 2002). Furthermore, complete proliferation loss due to overexpression of SKK-1 (Wnt inhibitor) and removal of Tcf4 and βcatenin further serve as evidence for active Wnt pathway in the gut (Korinek *et al.*, 1998; Pinto *et al.*, 2003; Kuhnert *et al.*, 2004). Hyper proliferation of the epithelium occurs following *APC* mutation (*APC* is a negative regulator of Wnt) or oncogenic  $\beta$ -catenin expression (Harada *et al.*, 1999; Romagnolo *et al.*, 1999; Smits *et al.*, 1998; Sansom *et al.*, 2004). Wnt signalling is also required for the terminal differentiation of the Paneth cells (Andreu *et al.*, 2005). Also, it is required to drive the expression of EphB2 and EphB3 (cell-sorting receptors) where β-catenin and TCF inversely control the receptors and their ligand ephrin-B1 along the crypt-villus axis (Batlle *et al.*, 2002). Recently, it was shown that *Achaete scute-like 2 (Ascl2*), which is transcription factor and Wnt target gene can alter the gut stem cell program; specifically, deletion of *Ascl2* can completely remove the stemness of stem cells (Van der Flier *et al.*, 2009).

#### 1.2.6 Wnt pathway in cancer and other diseases

Cancers exhibit many independent genetic defects. Many of the components of the Wnt signaling pathway are shown to be associated with human cancers or experimental cancer models. For example:

- Wnt-1 is a proto-oncogene activated by Mammary Tumor Virus in murine breast cancer (Nusse and Varmus, 1982).
- APC acts as a tumor suppressor gene in human CRC (reviewed in Polakis 1997).
- In human, mutations in β-catenin are found in colon cancer and in melanomas (Morin *et al*, 1997).
- The Drosophila gene legless is a homolog of *Bcl-9* (part of Wnt pathway), a gene is involved in B cell malignancies (Kramps, 2002).
- In human, mutations in *AXIN1* are found in liver cancers (Satoh *et al*, 2000; reviewed by Clevers, 2000).

- In mouse, *Tcf*1 can act as a tumor suppressor gene (Roose, 1999), and its mutation may lead to adenomas in the gut and mammary glands.
- A Wnt gene, *Wnt5A*, is mutated in lymphoid malignancies of mouse (Liang *et al*, 2003).

Other than cancer, mutation in Wnt genes may also lead to other diseases. Eg: mutation in *LRP5* leads to bone density defects (Gong, 2001; Little, 2002; Boyden, 2002), mutation in *LRP6* leads to early coronary diseases (Mani et al., 2007) and late onset of Alzheimer (De Ferrari, 2007), Wnt 10B mutation relates to obesity (Christodoulides 2006), *Axin2* mutation is involved in tooth agenesis (Lammli *et al*, 2004), *WTX2* relates to Wilm's tumour (Major, 2007, Rivera, 2007), *VANGL1* mutation leads to neural tube defects (Kibar, 2007) and *Tcf4* mutation may lead to TypeII diabetes (Grant 2006, Florez 2006, O'Rahilly 2006). The oncogenes and tumour suppressors of the Wnt pathway are demonstrated in Figure 1.2B. Thus, we see that components of the Wnt pathway are not only related to normal development of an organism but also critically involved in human cancers and many other diseases.

#### 1.4 Colorectal cancer, the disease

CRC initiates through a number of genetic alterations giving rise to genetic instability and which eventually leads to series of histological changes. The earliest stage of colon cancer is observed as aberrant crypt foci (ACF): two mutations in APC are present at this stage (Powell *et al.*, 1992; Jen *et al.*, 1994; Smith *et al.*, 1994a). These ACF are seen microscopically and consist of abnormally formed crypts containing cells of normal or abnormal morphology. The ACF may then lead to the formation of a polyp which is a benign mass that protrudes into the lumen of the colon. The Adenomatous polyps are also called adenomas and have abnormal cellular organisation whereas, the hyperplastic polyps have normal architecture and morphology. As colorectal cancer progresses, mutations in oncogenes and tumour suppressor genes occur and the chromosomal instability increases (Fodde and Clevers, 2001) (Figure 1.3).

For colorectal adenomas to progress towards malignancies, loss of response to TGF- $\beta$  is a key event (Polyak, 1996). TGF- $\beta$  signaling can inhibit the growth rate of the epithelial cells, for example human bronchial epithelial cells (Masui *et al.*, 1986), and renal tubular epithelial cells (Fine *et al.*, 1985). TGF- $\beta$  ligands are mostly found in the villi of the intestines (Oshima *et al.*, 1997).

Colorectal cancer can be studied under two major categories as follows:

- Sporadic colon cancer
- Hereditary colon cancer

#### 1.4.1 Sporadic colon cancer

Most cases of colon cancer are sporadic and are related to many factors such as diet, ethnicity, hormonal factors and environmental exposures. Interaction of some or all these factors may lead to increased risk of sporadic colon cancer. This type of colon cancer does not have a hereditary component and appears by random occurrence.

#### 1.4.2 Hereditary colon cancer

Around 10% of colon cancers fall under the category of being hereditary. If members of a family over a few generations are affected with colon cancer, there is an increased likelihood that family has a genetic mutation that can lead to hereditary colon cancer. The two main types of hereditary colon cancer are as follows:

- Familial adenomatous polyposis (FAP)
- Hereditary non-polyposis colorectal cancer (HNPCC)

#### 1.4.2A Familial adenomatous polyposis (FAP):

FAP is an uncommon hereditary colon cancer syndrome and is associated



Image taken from Fodde and Clevers, 2001

# Figure 1.3

The progression of colorectal cancer. Mutation in APC is observed in the earliest stage of CRC. Some aberrant crypt foci may lead to the formation of polyps (a benign mass which protrudes into the lumen), adenomatous (dysplastic) polyps (adenoma) may then progress to carcinoma of the colon. As CRC progresses, further mutations of oncogenes and tumour suppressor genes, and an increase in chromosomal instability are observed.

with a very high lifetime risk of colon cancer. The incidence of FAP is approximately 1 in 8000, 25% of which are novel germline mutations (Bisgaard et al., 1994). This condition leads to the formation of hundreds of polyps in the colon in the second or third decade of life, which has the ability to become malignant. [Note: A colon polyp is an area of tissue overgrowth in the colon or rectum and can be benign or malignant]. FAP patients may also be exposed to an increased risk for other types of cancers such as cancers of the small intestine, skin, thyroid and bile ducts (summarised in the Stanford cancer centre webpage, http://cancer.stanford.edu/information/geneticsAndCancer/types/fap.html). Most FAP patients have germline mutations in APC between codons 200 and 1600 and leading to truncated APC (Cottrell et al., 1992; Nagase & Nakamura, 1993; Laken et al., 1999). Colorectal tumours from FAP patients have been found to contain either additional somatic mutations in the second APC allele, or loss of heterozygosity at the APC locus (Miyoshi et al., 1992a; Solomon et al., 1987; Ichii et al., 1993; Levy et al., 1994; Lamlum et al., 1999). The location of the germline mutation in APC determines the type of mutations which eventually affects the second allele (Lamlum et al., 1999).

#### 1.4.2B Hereditary non-polyposis coli cancer (HNPCC)

Non-polyposis means that individuals, who have this syndrome, develop only a few polyps, maybe one or two, though, these polyps can become malignant or of the type that have the potential to become malignant. This hereditary disease develops due to germline mutations in the DNA mismatch repair (MMR) complex (Kinzler & Vogelstein, 1996; Kolodner, 1996; Markowitz, 2000). More than 90% of the cases of HNPCC involve mutations in *hMSG2* and *hMLH1*, two components of the mismatch repair complex (Yan *et al.*, 2000). Patients with this disease have 80% lifetime risk of developing colon cancer and also a greater risk of developing gastric and endometrial and ovarian cancers (Vasen *et al.*, 1990). In HNPCC, *APC* is mutated in 21% of cases (Konishi *et al.*, 1996) and  $\beta$ *catenin* is mutated in 43% of the cases (Miyaki *et al.*, 1999). Onset of this disease starts in a person who is in his/her early forties (Hodgson and Maher 1993).

#### 1.4.3 Genes of colon cancer

There are a number of key genes involved in the initiation and maintenance of colon cancer. Since, *APC* is a major tumour suppressor gene, its mutation is common in most CRCs. Inactivation of *APC* leads to genetic instability leading to genetic alterations thereby initiating colon cancer and maintaining its progression. Other tumour suppressor genes, *SMAD4* and *TP53* and the oncogene *KRAS* are major targets of the genetic alteration. Mutations in other genes, such as *KRAS*, have been seen in 50% of colon cancer patients, out of which 5% have mutations in *BRAF* (gene responsible for cell growth and cell signalling). *GLUT1* is a gene found to be in high amounts in cells with mutations in *KRAS* and *BRAF* (Fodde and Clevers 2001). *GLUT1* gene is a potential marker of hypoxia in patient with CRC (Chung *et al.*, 2009). *GLUT1* glucose transporter is associated with neoplastic progression in colon and also is a marker of poor prognosis (Haber *et al.*, 1998).

# 1.4.4: APC and β-catenin mutation in other forms of cancer

As already mentioned in section 1.1.2, mutations in *APC* and  $\beta$ -catenin are also found in cancers other than CRC. High levels of free  $\beta$ -catenin were seen in 7 out of 26 human melanoma cell lines (Rubinfeld *et al.*, 1997), 6 of which had mutations of the  $\beta$ -catenin gene. Aggressive fibromatosis (desmoid tumours) have also been shown to involve mutations in APC and point mutations in  $\Box\beta$ -catenin at either codon 41 or 45 (Tejpar *et al.*, 1999). Hepatocellular carcinoma (HCC) also involves mutation of  $\beta$ -catenin in 20% of cases (Huang *et al.*, 1999).  $\beta$ -catenin mutation is also common in hepatoblastoma, an embryonic liver tumour which occurs mainly in children under 2 years of age (Wei *et al.*, 2000). Miscoding mutations of both *APC* and  $\beta$ *-catenin* mutations have been observed in sporadic medulloblastomas (Huang *et al.*, 2000).

Thus, from the above sections, we have seen that there are various genes mutated in CRC and pathways like the Wnt signalling pathway are highly altered during this disease. Another important phenomenon associated with CRC is metastasis, by which a primary tumour invades the surrounding tissue and migrates to a secondary site in the body leading to a secondary tumour. Well differentiated carcinomas often lose epithelial differentiation during invasion at the invasive edge of the tumour and gain a mesenchymal-like phenotype by a process known as epithelial-mesenchymal transition (EMT); and then re-gain an epithelial phenotype following invasion by mesenchymal-epithelial transition (MET). Thus, this shows that malignancy in CRC is not only dependent on genetic alterations, but also on the tumour microenvironment since the stromal signals act in various combinations that can activate the EMT program (Weinberg 2007).

# 1.5 Epithelial-mesenchymal transition (EMT)

EMT is a vital and fundamental process in the morphogenesis of organisms. This process is also activated in cancer progression and also in many other diseases; and it occurs in response to extracellular signals (Gibbons *et al.*, 2009). EMT leads to a change in the shape of a cell and also may increase the cell's mobility as a cell transitions from an epithelial to a mesenchymal phenonotype. This transcriptional reprogramming process is associated with changes in epithelial cells which start expressing proteins that are associated with normal mesenchymal cells (reviewed in Thiery and Sleeman, 2006). Some characteristics of Epithelial-mesenchymal transition are as follows:

- The epithelial polarity is lost
- Cell:cell contact is lost/ disruption of intercellular contacts
- Aberration of cell substrate adhesion
- Basal membrane is degraded

- Cell invasion, migration and proliferation is increased
- Increase in cell motility

EMT is predominantly induced by stromal signals in the microenvironment (Weinberg, 2007). The major transcription factors involved in inducing EMT are *Zeb1/TCF8* (reviewed in Aigner *et al.*, 2007), *Zeb2* (Vandewalle *et al.*, 2005), *Snail* (reviewed in Usami *et al.*, 2008), *Slug* (reviewed in Kang and Massague, 2004), *E12/E47 FOXC2* (Mani *et al.*, 2007), and *Twist* (reviewed in Kang and Massague, 2004).

#### 1.5.1 Growth factors regulating EMT

Several growth factors (eg TGF $\beta$ , EGF (epithelial growth factor), etc) are known to induce EMT in epithelial cell lines by disrupting the tubular basement membrane (TBM). These factors can recognize the tyrosine kinase surface receptors and activate the Ras ((Lowenstein et al., 1992) pathway (observations in both in vitro and in vivo studies). [Note: Ras pathway is shown to be important in EMT]. MEK and Rac are downstream components of Ras that are shown to induce EMT in bladder carcinoma lines (Edme et al., 2002). Both the mitogen activated protein kinase (MAPK) pathway and the PI3K pathway can induce tumourigenicity. EMT was shown to be dependent on the activation of the MAPK pathway, though the PI3K pathway was related to scatter and cell proliferation (reviewed in Theiry 2003). Activation of c-met (a proto-oncogene that encodes mesenchymal-epithelial transition factor) induces inositol 5'phosphatase SHIP-1 which is important in branching morphogenesis. Overexpression of SHIP induces EMT in MDCK cells (Mancini et al., 2002). Thus, SHIP-1 is needed for EMT even though MAPK was activated. Chicken tumour virus no.10 regulator of kinase (Crk) is an adaptor protein that can also induce EMT in MDCK cells by being overexpressed in the absence of hepatocyte growth factor (HGF) (Lamorte et al., 2002).

Transforming growth factor  $\beta$  (TGF $\beta$ ) is involved in EMT both during embryogenesis and tumourigenesis. TGF $\beta$  has three isoforms in humans (TGF $\beta$ 1, TGF $\beta$ 2, and TGF $\beta$ 3); all three can regulate EMT. TGF $\beta$  was first shown to induce EMT in mammary epithelial cells (NMuMG cells) (Miettinen et al., 1994) where the epithelial cells were shown to differentiate into fibroblastic phenotype by the addition of TGF<sup>β</sup>1. TGF<sup>β</sup> along with extracellular matrix protein Laminin-5 (Ln-5) can also induce EMT (Giannelli et al., 2005). TGFB is known to be an inducer of EMT in association with the Ras pathway. In mouse, if the Ras pathway is active, the mammary epithelial cells undergo complete EMT (Janda et al., 2002). In MDCK cells, EMT is induced by TGFβ and Ras and and also by Smad3, which is involved in growth arrest, which becomes downregulated in these cells (Nicolas et al., 2003). Re-expression of Smad3 cannot reverse EMT in MDCK cells (Nicolas et al., 2003). TGFB signaling is also related to the activation of the proto-oncogene Src (Kim and Joo, 2002). Src is also involved in formation of a mesenchymal phenotype in colon cancers, though it cannot actually cause carcinoma (Shimizu et al., 2002).

Cytokine Interleukin 1 $\beta$  (IL1 $\beta$ ) is another growth factor regulating EMT and this was first shown in mesothelial cells, LP-9 cells which undergo EMT in response to IL1 $\beta$  and this was marked by changes in expression of EMT markers (*Vimentin, Twist, Snail, N-cadherin* and *cytokeratin*) (Masoud *et al.*, 2009). Two key hallmarks of cells undergoing EMT are reduction in levels of Ecadherin and nuclear localization of the oncogene,  $\beta$ -catenin (reviewed in Thiery 2002).

#### 1.5.2 EMT and development

EMT is a fundamental process in normal human development. The earliest event of EMT in development is during grastulation, when the mesenchymal cells and mesoderm are formed. Formation of placenta, neural crest and heart valve arise through EMT (reviewed in Polyak and Weinberg , 2009). The changes associated with EMT enable the cells of a developing embryo to respond to changes in the environment. Developmental stages involving EMT can be described as follows:

- During EMT in development, upper epiblast epithelium forms the primary mesenchyme. This leads to three layered blastocyst with ectoderm, mesoderm (primary mesenchyme) and the endoderm (Figure 1.4A steps 2 and 3).
- The first somite is formed by MET. Lateral mesenchyme forms the mesodermal epithelia of the body wall (Figure 1.4A step 4).
- Another EMT event is formed which makes the neural crest cells which then forms the nervous system (Figure 1.4A step 5) (Huang and Saint-Jeannet, 2004; Newgreen and McKeown, 2005).
- More steps of EMT then forms dorsal parts of the somite to form muscles and skin (Figure 1.4A step 6), peripheral neurons and glia, and melanocytes (reviewed in Hugo *et al.*, 2007).

EMT also occurs in ovaries of women. The mammalian ovary is covered with ovarian surface epitheliums (OSE) from which ovarian carcinomas arise. Normal OSE in the ovary undergoes EMT to maintain ovarian homeostasis (Auersperg *et al.*, 2001).

#### 1.5.3 EMT and cancer

EMT is seen in many forms of cancers such as colon cancer, prostate cancer, breast cancer and lung cancer. Most solid tumours arise from the epithelia, and the loss of epithelial-cell markers (e.g. E-cadherin and gamma catenin) is related to disease progression and tumour metastasis (Figure 1.4 B). Cancer cells can de-differentiate through EMT and attain mesenchymal-cell characteristics (e.g. expression of *Vimentin*, *Fibronectin*). Mesenchymal-Epithelial Transition (MET) is a reversible process, whereby mesenchymal tumor cells attain epithelial-like phenotype. MET enhances cell proliferation and growth of epithelial tumour cells at a secondary tumour site. Metastatic and invasive carcinoma is associated with loss of epithelial cell polarity and differen-



Reference: Hugo et al., 2007, review

# Figure 1.4

EMT and MET events in development and cancer. A) EMT and MET in early development. B) EMT and MET progression during carcinogenesis. Cancer cells undergo EMT escaping the environment (1, 2), create new malignancy (4). Mesenchymal cells intravasate (5), remain in circulation (6), extravasate (7). Persist as micrometastasis in bone marrow (8) and revert to epithelial cells via MET (9).

tiation. Data suggests that EMT has a role in carcinoma in micro metastatic tumour cells in bone marrow and lymph nodes (Pierga *et al.*, 2003). Sarcomatoid carcinoma may also have a possible EMT and MET balance although this has to be further explored.

During tumour progression, non-invasive benign tumour cells can gain the potential to become invasive and infiltrate surrounding tissues and migrate to distant sites leading to secondary tumours. Many EMT regulators can enhance the formation of a tumour (Thiery, 2002), eg. Expression of Snail1 increases the virulence of breast tumors, and high Snail1 expression relates to poor survival rates in human breast cancer (Moody *et al.*, 2005). Three transcriptional factors TWIST, GOOSECOID (homeobox transcriptional repressor), and *FOXC2* (transcription factor of the FOX family of forkhead helix-turn-helix DNAbinding proteins), that regulate developmental EMT also have important functions in enhancing metastasis. Twist expression is shown to be sufficient to induce *in vitro* EMT in breast cells (Yang *et al.*, 2004). Twist and Goosecoid regulate *FOXC2* that regulates EMT and development of organs (Carlsson and Mahlapuu, 2002). *FOXC2* also up-regulates mesenchymal gene transcription, rather than initiating EMT through E-cadherin repression.

Thus, we have seen that EMT is a fundamental process not only associated with embryo development and tissue regeneration but also in other forms of disease including various forms of cancer where, the invasion and metastasis is driven by the transition from a stable epithelial phenotype to a migratory, mesenchymal phenotype (Thiery and Sleeman, 2006). Recently, a possible link between cells undergoing EMT and cells with properties of cancer stem cells (CSCs) has been described in breast epithelial cells where breast cells from normal mouse and human mammary tissue were cultured and were induced to undergo an EMT, the cells acquired characteristics of stem cells (Mani *et al.*, 2008). The fundamentals of stem cells are described in section 1.6
## 1.6 Stem cells:

Stem cells have distinct potential to give rise to the many different cell types in the body during both embryogenesis and tissue specific growth. In a blastocyst (3- to 5-day-old embryo), the stem cells in the inner walls give rise to the entire body of the organism along with other specialized cell types and organs. The special properties of stem cells are also required for tissue repair to enable renewal of cells during an individual's lifetime. For example, as discussed previously, in the gut, adult stem cells divide regularly to replace the worn out/ damaged cells. Here, we will discuss the different types of stem cells and the factors required for the promotion and maintenance of stem cells.

## 1.6.1 Types of stem cells

Predominantly, there are two categories of stem cells, embryonic stem cells and somatic or adult stem cells. A new type of stem cell, called induced pluripotent stem cells (iPSCs), has also been characterized recently.

- Embryonic stem cells are the stem cells derived from four or five day old embryos (blastocyst) that develop from a fertilized egg. [Note: the blastocyst has the trophoblast which is the layer of cells that surrounds the blastocoel, a hollow cavity inside the blastocyst and the inner cell mass which is a group of cells that develop into the embryo]. Embryonic stem cells were first derived from early mouse embryos in 1981 (Evans and Kaufman) and first the human embryonic stem cells were derived and grown in 1998 (Thomson *et al.* 1998).
- An adult stem cell is an undifferentiated cell among differentiated cells in a tissue or an organ. These cells have the ability to renew themselves and differentiate into the specialized cells of that organ or tissue. Thus, these adult stem cells maintain and repair the tissue they are associated with. The area where they are found is called a "stem cell niche" which is

important in providing environmental cues and signals required for the maintenance of stem cells (reviewed in Scadden 2006). Both embryonic and adult stem cells each have advantages and disadvantages for their use in cell-based regenerative therapies (Hmadcha et al., 2009). One major difference between these two types of stem cells is that, embryonic stem cells are pluripotent (ability to give rise to all cell types of the body) and can differentiate to form all cell types of the body; whereas, differentiation of adult stem cells is limited to the cell types of the same tissue or organ (NIH stem cell website). Also, embryonic stem cells are easily grown in culture compared to adult stem cells which are rare in mature tissues (NIH stem cell website). Some organs have both embryonic and adult stem cells. e.g. the bone marrow contains two kinds of stem cells, hematopoietic stem cells that form all the types of blood cells in the body; and bone marrow stromal stem cells (mesenchymal stem cells) that can give rise to bone, cartilage and fat cells that support the formation of blood, and fibrous connective tissue (Rubin and Strayer, 2007).

Induced pluripotent stem cells (iPSCs) are derived from somatic adult cells and are obtained by genetic reprogramming of a fibroblast cell to an embryonic stem cell–like cell. This is done by forced expression of four genes, *Oct-3/4, Sox2, c-myc*, and *Klf4*, all of which are factors important for maintaining embryonic stem cells (Takahashi &Yamanaka, 2006). Mouse iPSCs were first reported in 2006 (Takahashi and Yamanaka) as having characteristics of pluripotent stem cells and expressing stem cell markers. Human iPSCs were first reported in 2007 (Takahashi *et al.*, 2007) as cells which expressed stem cell markers and which could also generate cells characteristic of all three germ layers. iPSCs are already in use for drug development and disease treatment (Wu *et al.*, 2008; Shizuru *et al.*, 2005; Rubin 2008; Rossi *et al.*, 2008; Nelson *et al.*, 2009). Recently, viruses have been used to introduce reprogramming factors into adult cells; however, this use of virus in treatment might itself be a cause of cancers. Thus, non-viral delivery strategies are currently being worked

on e.g. recently, virus free ESCs are made for Parkinson patients (Soldner *et al.*, 2009). Mouse iPS cells were also generated without using viral vectors (Okita *et al.*, 2008).

## 1.6.2 Properties of stem cells:

Stem cells can divide to generate two daughter cells, one rapidly cycling more specialized differentiated cell type and one stem cell that replace the parent stem cells (reviewed in Barker *et al.*, 2008). The former undergoes limited cell division and then terminally differentiates. The two important characteristics of stem cells are as follows:

- Stem cells are not specialized cells and can renew themselves by cell division.
- Stem cells can differentiate and induce tissue- or organ-specific cells with specialized functions

## 1.6.3 Leukemia inhibitory factor (LIF) and stem cells

Not much is known about the signaling pathways that regulate the human embryonic stem cells. LIF is a molecule involved in the maintenance of selfrenewal and pluripotency of mouse ES cells (Smith *et al.*, 1988; Williams *et al.*, 1988). LIF was found to maintain mouse ES cells in the absence of a feeder layer. However, LIF was not the only factor to maintain stem cell identity in mouse as it was shown that null mutants deficient in LIF/ Stat3 signaling did not show any defect in establishing the stem cell niche (Smith *et al.*, 2001). Also, LIF was shown to be incapable of maintaining human ES cell identity (Thomson *et al.*, 1998; Sato *et al.*, 2001). It was therefore suggested that other signaling pathways might also be involved in stem cell maintenance, such as FGF4, TGF $\beta$ , and Wnt. Some of these pathways are discussed below in terms of stem cell maintenance.

## 1.6.4 NANOG and maintenance of stem cells

One of the fundamental downstream effectors, required for promotion of pluripotency in both mouse and human ES cells, is the homeodomain transcription factor NANOG (Chambers *et al*, 2003; Mitsui, Tokuzawa *et al*, 2003; Hart AH, Hartley *et al*, 2004, Pan and Thompson, 2007). *Nanog* is involved in self-renewal and maintaining pluripotency of embryonic stem cells (Mitsui *et al.*, 2003; Chambers *et al.*, 2003). NANOG has also been shown to be expressed in tissue specific stem cells, such as the pancreas (Koblas *et al*, 2008), germ cell tumours (reviewed in A.T. Clark, 2007) and some cancers, such as prostate (Kasper 2006), breast cancer (Ezeha *et al.* 2007) and Ewings sarcoma (Suva *et al.*, 2009).

In human, NANOG is a 305 amino acid protein and is primarily located in the nucleus of a cell. It works along with two other transcription factors OCT4 and SOX2 to maintain pluripotency. [Note: Key transcription factors for maintaining murine stemness are Nanog, Oct3/4 and Sox2 (Nichols *et al.*, 1998; Niwa *et al.*, 2000; Boiani *et al.*, 2005)]. Over- expression of *NANOG* results in increased cell growth and transformation of cells (Piestun *et al.*, 2006). NANOG was also shown to be over-expressed in cervical cancers thereby aiding their progression (Ye *et al.*, 2008) and also in oral cancer along with OCT4 (Chiou *et al.*, 2008). NANOG has also been suggested as a marker for testicular cancer (Hoei-Hansen *et al.*, 2005).

Deletion of *Nanog* leads to embryonic lethality in mice (Mitsui *et al.*, 2003) and its' expression is related to autonomous self renewal of embryonic stem cells (Chambers *et al.*, 2003). It is needed for germ cell formation and construction of the inner cell mass (Chambers *et al.*, 2007). Cells are prone to differentiation on *Nanog* downregulation but in the permanent absence of *Nanog*, ES cells can self-renew (Chambers *et al.*, 2007). *Nanog* expression is not continuous in the embryonic pluripotent cells since it is present in the blastocyst during formation of pluripotent cells, downregulated during implantation (Chambers *et al.*, 2003) and expressed again in the posterior region (Hart *et al.*,

2002). Thus, Nanog is primarily involved in the formation of the inner cell mass and germ cell states. Nanog also stabilizes embryonic stem cells in culture by resisting or altering gene expression states (Chambers *et al.*, 2007).

#### 1.6.5 Intestinal stem cells

The epithelial cell lining of the gut proliferates very rapidly, representing one of the most rapidly renewing tissues in the human body; and stem cells are responsible for this phenomenon. The differentiated cells are renewed due to continuous supply of multipotent progenitor cells that originate from stem cells of the intestinal crypts. Stem cells are responsible for the proliferation of crypt cells in the intestine (Bjerknes et al., 2006; Marshman et al., 2002). The epithelium in the mouse small intestine renews every five days (reviewed in Laurens, et al., 2009). The stem cells at the bottom of the crypt enhance vigorous proliferation. The TA (transit- amplifying) cells or the 'rapidly cycling daughter cells' are easily distinguishable at the crypt and divide every 12-15 hour and this leads to 300 cells each day, per crypt (Marshman et al., 2002). These TA cells differentiate in an irreversible manner at the crypt-villus junction, and can divide up to 6 rounds of differentiation while going up the villi (Marshman et al., 2002). 3 days after the terminal differentiation, the epithelial cells undergo apoptosis at the top of villi (reviewed in Laurens et al., 2008). It is now known that every crypt has 4-6 stem cells (Potten et al., 1990). Intestinal stem cells (ISC) play an important role in the rapid expansion of the gut during development and also during tissue regeneration on injury, surgery or cancer. The stem cells normally undergo apoptosis in response to toxic stimuli thereby reducing the chances of acquiring mutation that could result in a tumour. It has been demonstrated that a single mutated stem cell can give rise to colorectal tumours (Simon et al., 2000). Colonic stem cells are defined by a number of markers, of which LGR5 (leucinerich repeat-containing G protein coupled receptor 5)/ GPR49 is one. This marker is used to define the location of intestinal stem cells at the crypt base of crypts (Barker et al., 2007; Sato et al., 2009).

## 1.6.6 A) Intestinal stem cell microenvironment

The tightly junctioned intestinal fibroblasts are responsible for the formation of the intestinal epithelial niche (Mills *et al.*, 2001). Here, the cells secrete growth factors and cytokines to enable the cells to proliferate (Powell *et al.*, 1999). In the crypt, BMP acts as a negative regulator of crypts and is active in the villus mesenchyme (Haramis *et al.*, 2004; Hardwick *et al.*, 2004). Intestinal epithelial cells express Sonic hedgehog (Shh) and Indian hedgehog (Ihh) ligands, inhibition of which leads to reduction in villi size and highly proliferative epithelium (Ramalho-Santos *et al.*, 2000).

## 1.6.6 B) Two models for intestinal stem cells

The two models of the intestinal stem cells are as follows:

## i) +4 model (Classical model):

Understanding from previous literature suggests that the crypts in the small intestine are bound by Paneth cells from below and behave like a tube of proliferating cells. The +4 model was suggested in late 1950s (Quastler and Sherman, 1959); according to which the intestinal stem cells are based at the +4position relative to crypt bottom and the Paneth cells occupy the first 3 positions (Cairnie et al. 1965). Experimental support for this model was suggested in 1974 (Potten et al., 1974) where label-retaining cells were observed at the +4 position in the crypt. Recently, this model was modified when the stem cells found among the ring of cells immediately above the Paneth cells, at position 4, where they occupy a stem cell microenvironment (Potten et al., 2002) (Figure 1.5 top panel). Potten and Roberts (1974) observed that the +4 cells are radiation sensitive and this sensitivity protects the stem cells from genetic damage/ alteration so that they do not acquire longer term changes. In this model, the first two to three generations of TA cells replace the damaged stem cells, enabling the stem cells to regain their stem cell property and occupy the +4 position again. Due to lack of knowledge of the cellular nature of these +4 cells, this model had constraints.



Reference: Barker et al., 2008

## Figure 1.5

The two models of intestinal stem cells: +4 position model (top), stem cell zone (bottom). In +4 position model, it was assumed that that the crypt base consists mainly of terminally differentiated Paneth cells and the stem cells were located just above the Paneth cells at the +4 position. Stem cell zone is a more recent model stating that small, undifferentiated, cycling cells (crypt base columnar cells) with the Paneth cells are the intestinal stem cells.

Apart from this, small and undifferentiated cells were identified hidden between the Paneth cells which were referred to as the crypt base columnar (CBC) cells (Cheng *et al.*, 1974a, b). This gave rise to the second model of the intestinal stem cells.

## ii) Stem cell zone model:

This model is based on Crypt Base Columnar (CBC) cells which are small and undifferentiated cells found between the Paneth cells (Cheng and Leblond 1974a, b; Bjerknes and Cheng 1981a, b, 1999). In this model, the CBC cells are considered to be the true stem cells and the short lived multipotent "mix" cells represent the direct derivative of the CBC cells. The common 'origin of differentiation' sites directly above the Paneth cells are occupied by the Mix cells. The maturing Paneth cells migrate downwards where the oldest cells are at the crypt base. The cells other than the Paneth cells move up the crypt-villi axis (Figure 1.5 bottom panel). Due to lack of distinct stem cell marker at the time, identification of these cells was not possible and this model suffered a setback. An attempt to mark these stem cells was tried using Musashi-1 and βintegrin, but, since they lacked specificity, the identification of stem cells was not possible (Kayahara et al., 2003, Spangrude et al., 1998). Some specific markers of +4 cells are phosphor-PTEN and phosphor-AKT, Dcamkl1, Sox4 and SFRP5 (He et al., 2004; Giannakis et al., 2006). But, their positional information remains inadequate.

Recently, the CRC stem cell marker LGR5, shows the expression of stem cells is restricted to CBC cells (Sato *et al.*, 2009). Another stem cell marker, *BMI I*, that encodes a chromatin remodeling protein shows the expression of the intestinal stem cells at the +4 position (Sangiorgi and Capecchi 2008). Thus, it might be possible that there are different activation states of the stem cells in the intestinal tract. However, this debate came to an end in 2009, with the conclusion that the CBC cells within the stem cell zone at the crypt base is the actual site of the intestinal epithelial stem cells (van der Flier *et al.*, 2009) where induced

31

deletion of the transcription factor Achaete Scute-Like 2 (ASCL2) leads to removal of LGR5 stem cells because ASCL2 binds to the LGR5 promoter. <u>1.6.7 Cancer stem cells</u>

It is well known that both tumour cells and normal stem cells can proliferate extensively and can lead to abnormal tissues. Both have a combination of cells with multiple phenotypes, proliferative capacities and different degrees of differentiation (Fidler et al., 1997; Fidler et al., 1982; Heppner et al., 1984; Nowell et al., 1986). Most tumours contain a heterogenous population of cancer cells with both tumourigenic and non-tumourigenic cancer cells. Cancer stem cells (CSCs) are cancer cells found in tumours or hematological cancers, which have similar properties to normal stem cells, i.e ability to develop into all the cell types in a cancer sample and to self renew (Reya et al., 2001). CSCs are therefore tumourigenic (tumour-forming) and can produce tumours through self-renewal (Clarke et al., 2005, 2006) and differentiation into various cell types. Such cells reside in a tumour as a population which can cause metastasis to give rise to a new tumour at a secondary site in the body. Thus, the CSCs are the rare cells in a tumour that are 'immortal' and produce 'normal' cancer cells those can divide many times to form a tumour, and also can give rise to new cancer stem cells.

It is already known that cancer results from various genetic mutations in a single cell may be over a period of many years (Fearon *et al.*, 1990). The stem cells are the only the long-lived cells in many tissues. Thus, it is stem cells in which these early mutations might have been accumulated. Mutations in stem cells along with/ or mutation in the progenitor cells can lead to the formation of cancer stem cells (reviewed in Bjerkvig *et al.*, 2005) (Figure 1.6). It might be considered that a tumour can be like an abnormal organ whose activity is induced by a cancer cell that acquired the capacity to proliferate indefinitely through mutation (Morrison *et al.*, 1997; Kummermehr *et al.*, 1997).

According to the stochastic cancer stem cell model, all tumour cells can form the entire tumour and the growth of the tumour depends on the genetic events, proliferation and cell death. But, according to the new model for cancer



Clarke et al., 2005

## Figure 1.6

Factors giving rise to cancer stem cells. Environmental factors may give rise to the progenitor, differentiated or stem cell niche. These can further lead to the formation of cancer stem cells and hence initiate tumour progression.

stem cells, only a subset of cancer cells, which are long-lived cancer stem cell, can sustain the tumour growth *in vivo* and form the entire tumour; and the shortlived differentiated cells in a tumour cannot maintain this tumour growth. This tumour growth depends on the stem cell renewal mechanisms. The origin of cancer stem cells is still not clear. But, it is widely accepted that they might result from specific mutations in normal or early stem cell progenitors. Also, there is a possibility that they are obtained from differentiated cells (Bjerkvig *et al.*, 2005).

The first evidence for CSCs was in 1997 where a sub-population of leukaemic cells was isolated and enriched, which expressed a specific surface marker CD34, not CD38 marker. Thus, it was established that the CD34+/CD38subpopulation is capable of initiating tumors in NOD/SCID (nonobese diabetic/severe combined immunodeficient) mice (Bonnet and Dick, 1997). [Note: NOD/SCID is a mouse model, efficient for human cells to engraft and also to proliferate and differentiate (Ito et al., 2002)]. Furthermore, Spangrude and co-workers (1998) identified the hematopoetic stem cells from bone marrow (Spangrude et al., 1998). Since then there has been evidence of cancer stem cells in many tumours including breast (Al-Hajj et al. 2002), colon (O'Brien et al., 200)], brain [Singh et al., 2003], ovary (Zhang et al., 2008), pancreas (Li et al., 2007), prostate (Maitland et al., 2008, Lang et al., 2009). Identification of cancer stem cells in solid tumours was done by several other groups in later years (Singh et al., 2004; Dalerba et al., 2007a). One of the most fascinating works done in this field was by Visvader and co-workers when they grew an entire mammary gland from single isolated stem cells (Visvader et al., 2006). Prior to this, single stem

cells were shown to give rise to epidermis, hair follicles and sebaceous gland (Blanpain *et al.*, 2004; Claudinot *et al.*, 2005). Genetic marking of the stem cells to visualize the modified stem cells has also been explored. One such example is the expression of a progesterone-responsive Cre recombinase enzyme in cells of bulge hair follicle regions. This was done by a transgenic keratin-15 promoter (Morris *et al.*, 2004) in mouse.

## 1.6.8 Regulatory pathways of stem cells and cancer cells

Since, both stem cells and cancer stem cells can self renew, one might guess that there are similarities in mechanisms of self-renewal in both ''normal'' stem cells and ''cancer'' stem cells. Evidence shows that many of these pathways that are cancer related may also regulate normal stem cells. The transformation of normal stem cells to cancer stem cells are controlled by various signaling pathways which includes Bmi-1 (active in cancer stem cells in brain tumours) (Dontu *et al.*, 2004), Notch pathway (demonstrated in neural and mammary stem cells (Dontu *et al.*, 2004), Sonic hedgehog and the Wnt pathway (Figure 1.7). Cell death also has a role in regulating homeostasis in hematopoietic stem cells (HSCs) (Domen, *et al.*, 1998, 2000). Shh, Notch and Wnt are also associated with oncogenes and thus may regulate stem cell self-renewal in CSCs (Taipale *et al.*, 2001). The role of these pathways is outlined below:

A) Notch pathway in stem cell maintenance: Notch signaling is involved in both normal embryonic development (Gridley *et al.*, 2003) and also in the regulation of self-renewal of stem cells along with other pathways such as Wnt and hedgehog pathways (Molofsky *et al.*, 2004). The Notch ligand, Notch-1, has a critical role in regeneration of the HSCs from the hematogenic endothelial cells during the early embyogenesis (Dzierzak *et al.*, 2003; Kumano *et al.*, 2003; Hadland *et al.*, 2004., Robert-Moreno *et al.*, 2004). When the Notch pathway is activated in HSCs, it increases the progenitor activity *in vivo* and *in vitro* indicating its role in maintaining multipotency (Varnum-Finney *et al.*, 2000). In genetically modified mice, the osteo-blastic cells were shown to produce high levels of Jagged 1 (Notch ligand) and to increase the numbers of HSCs on activation of Notch-1, *in vivo* (Calvi *et al.*, 2003). Notch receptors are also



Reference: Reya et al., 2001

## Figure 1.7

Signalling pathways regulating self-renewal mechanisms during normal stem cell development and transformation. Shh and Notch pathways contribute to the self-renewal of stem cells and/or progenitors in organs such as haematopoietic and nervous systems. These pathways can also lead to oncogenesis when dysregulated and can be associated with a human tumours such as colon carcinoma and epidermal tumours (Wnt), medulloblastoma and basal cell carcinoma (Shh), and T-cell leukaemias (Notch).

the gut (Schroder *et al.*, 2002). Active Notch signalling can also be seen in the muscle stem cells where activation of the Notch pathway can reinstate the regenerative potential in the aged mouse muscles (Conboy *et al.*, 2002; Conboy *et al.*, 2005).

B) <u>Shh in stem cell regulation</u>: The Shh pathway increases mesenchymal cell proliferation and regulates the differentiation of smooth muscle progenitor cells in mouse kidney (Yu *et al.*, 2002). Populations highly enriched for human HSCs exhibit increased response of self-renewal following Shh stimulation *in vitro* (Bhardwaj *et al.*, 2001). In neural cells, Shh signaling can regulate the number of stem cells and can also control the proliferation of early neuron production (Palma and Ruiz, 200; Palma *et al.*, 2005). Shh also has an important role in regulating epidermal stem cells of the skin (Kameda *et al.*, 2001) by maintaining a stem cell population for the proliferation of human putative epidermal stem cells. Studies also show that Shh is involved in the proliferation and cell-fate specification of mesenchymal stem cells (Kondo *et al.*, 2005).

C) Wnt in regulating stem cells: When ES cells self renew and maintain their pluripotency to differentiate into other cell types, many signal transduction pathways are involved,  $\beta$ -catenin/ Wnt pathway being one such pathway. Unknown factors in conditioned medium or Wnt activation by 6-bromoindirubin-3'-oxime (BIO), a GSK3β inhibitor, can also maintain the undifferentiated state of hESCs (Sato et al., 2004). The role of  $\beta$ -catenin in stem cell maintenance and early differentiation of embryonic stem cells was studied and the loss of βcatenin in ES cells was shown to alter the expression of stem cell markers (Anton et al., 2007). It was also concluded that the effect of Wnt signalling on ES cells varies according to the time-point of treatment. Despite the above mentioned evidence suggesting a role for  $\beta$ -catenin in stem cell maintenance, the actual mechanism is still not clear. Wnt3a conditioned medium, which is an inhibitor of the GSK3 $\beta$ , is shown to maintain murine and human stem cells (Sato *et al.*, 2004). Wnt genes were shown to inhibit differentiation by upregulating Stat3 in mouse (Hao et al., 2006). However, a different study showed that Wnt3a conditioned medium had no effect on Stat3 phosphorylation (Ogawa et al.,

mechanism is still not clear. Wnt3a conditioned medium, which is an inhibitor of the GSK3β, is shown to maintain murine and human stem cells (Sato *et al.*, 2004). Wnt genes were shown to inhibit differentiation by upregulating Stat3 in mouse (Hao *et al.*, 2006). However, a different study showed that Wnt3a conditioned medium had no effect on Stat3 phosphorylation (Ogawa *et al.*, 2006). It was also suggested that there is no direct cross-reactivity between the Wnt and Lif pathways (Ogawa *et al.*, 2006). Wnt3a has been shown to maintain the self renewal of hematopoietic cells (Reya 2002), however, recombinant purified Wnt3a does not stop the differentiation of mouse ES cells (Ogawa *et al.*, 2006; Singla *et al.*, 2006).

D) <u>TGF $\beta$  in regulation of hESCs</u>: The activation of TGF $\beta$ / activin/ nodal through SMAD2/3 was shown to be required for the maintenance of the undifferentiated state of hESCs (Daylon *et al.*, 2005). It has been shown that in a mouse embryo that lacks *Smad* (*Smad* modulates the TGF $\beta$  ligands) expression, results in defect of epiblast proliferation and also delays the growth of the inner cell mass (Sirard *et al.*, 1998). Also, mouse which lack ALK-3BMP type IA receptor can decrease the cell proliferation in the epiblast confirming that BMP signaling pathway is also important in the maintenance of mouse ES cells (Mishina *et al.*, 2002). BMP-4 is also shown to be important for the maintenance of ES cell self-renewal (Qi *et al.*, 2004).

## 1.6.9 Colon cancer stem cell markers

**CD133/ Prominin 1:** CD133 is a 5-transmembrane domain molecule with a molecular weight of 120 kDa (Miraglia *et al.*, 1997) and has been shown to be located on the apical plasma membrane in embryonal epithelial cells (Weigmann *et al.*, 1997; Corbeil *et al.*, 1999) and the plasma membrane of CRC cells (Ricci-Vitiani, *et al.*, 2007; reviewed in Corbeil *et al.*, 2000). Colon cancer cells were earlier demonstrated to have a subpopulation of CD133<sup>+</sup> cells in tumours. These were shown to initiate the tumour growth and were called colon-cancer-initiating cells or colon cancer stem cells (CSCs) (O'Brien *et al.*, 2007, Ricci-Vitiani *et al.*, 2007). On the other hand, CD133<sup>-</sup> cells showed none of these characteristics (Ricci-Vitiani *et al.*, 2007, Matilde *et al.*, 2007). CD133 is presently considered to be a potential marker of colon cancer stem cells though its expression is much wider (reviewed in Shmelkov *et al.*, 2008) such as in differentiated kidney and breast (reviewed in Montgomery and Shivdasani, 2009). A sub-population of tumour cells without CK20 (a differentiation marker) expression was shown to have strong CD133 expression *in situ* and the expression of CD133 in colon cancer is associated with low patient survivals (Horst *et al.*, 2008). Other than this evidence, CD133 localization *in situ* is not well studied.

LAMIN A: LAMIN A is a type V filamentous protein mostly localised to the inner nuclear membrane (Hutchison 2002). In 2008, LAMIN A was shown to be a critical biomarker for colon cancer stem cells (Weichert *et al.*, 2004). About one third of the patients with aggressive bowel cancer express LAMIN A. LAMIN A expression is altered in many forms of cancer such as lung tumours (Rowlands *et al.*, 1994), testicular germ cell tumours (Machiels *et al.*, 1997) and Hodgkins disease (Jansen *et al.*, 1997).

**CD166:** Activated leukocyte cell adhesion molecule (ALCAM) alternatively known as CD166, was shown to be over expressed in colorectal cancers where 58.6% of the cases were shown to have cytoplasmic expression of CD166 and 30.6% of the cases showed its membranous expression (Weichert *et al.*, 2004). It was previously shown to be expressed in melanoma (Léon *et al.*, 2000) and also in prostate cancer (Kristiansen *et al.*, 2003). CD166 is differentially expressed in normal and diseased colonic epithelium which makes it a good marker for colon cancer stem cells (Dalerba *et al.*, 2007b).

1.7 SW480 colorectal cancer cells transitioning between low and high density, an *in vitro* model to study CRC

It is not easy to study the events surrounding EMT during progression of cancer, partly due to the very transient nature of this process in situ. Thus, we used CRC cells from the cell line SW480, grown to low and high density, as a model for cells transitioning between the mesenchymal-like and epithelial-like cell states respectively (Brabletz et al, 2001; Davies et al, 2004). SW480 cells are human colorectal cancer cells derived from human colorectal adenocarcinoma. It is derived from lymph node metastasis (Leibovitz et al., 1976) and has single base pair mutation at codon 1338 of APC and a complete loss of the second allele. This mutation results in a stop codon and truncation at 1337 amino acids (Nishisho et al., 1991). In 2001, Brabletz and co-workers grew SW480 cells at low density with fibroblastoid phenotype with protruding lamellipodia. These cells resembled the mesenchyme-like cells at the invasive fronts of a colon tumour. With increasing density, these cells changed their phenotype to epithelial-like cells similar to those found at the central mass of a tumour. This phenotypic change was associated with β-catenin translocation from the nucleus to the cytoplasm and membranous co-localization of E-cadherin. There are several genes overexpressed at the invasive front of CRC, one of which is nuclear factor kB (NF-kB) (Horst et al., 2009). This dimeric transcription factor has role in immune response, cell growth control and survival of the cell. In our laboratory too, this transition of SW480 cells has been shown (Davies et al., 2004) where high amount of nuclear  $\beta$ -catenin were shown in low density SW480 cells whereas in high density SW480 cells,  $\beta$ -catenin was predominantly located at the cell membrane. As the SW480 cells are grown from low to high cell density, levels of E-cadherin expression was shown to increase and that the total level of APC expressed was decreased. Interaction of β-catenin and Ecadherin expression is also increased at high cell density and that between βcatenin and APC decreased. Thus, there has been an established series of evidence for the use of the human CRC cell line, SW480 grown at low and high density, as an *in vitro* model to study the CRC disease progression. An *in vitro* system that would mimic the changes in colorectal cancer cells during EMT would be of great importance to us in studying these changes further. Using this system, we show that the founding member of the T-box transcription factor family of genes, BRACHYURY, is differentially expressed in low density

40

(mesenchymal-like) cells. BRACHYURY, which we show to be regulated by  $\beta$ catenin, acts directly to maintain levels of the pluripotency determinant, NANOG and to maintain a CSC phenotype.

## 1.8 Aims and overview

Cancer stem cells (CSC) are initiating cells in many types of cancer that have similar properties to normal embryonic stem cells. Colorectal tumours also have the ability to undergo an epithelial to mesenchymal transition (EMT)-like processes at the invasive front, rendering them capable of invasion and metastasis. Recently, a possible link between these two processes, that is, cells undergoing EMT and cells with stem cell like properties, has been identified in mammary cells induced to undergo EMT, where such cells acquired SC characteristics. It is of importance to understand the molecular components leading to promotion or maintenance of cells involved in these disease causing processes; however since it is difficult to observe EMT in cancer due to the transient nature of these transition events during tumourigenesis, we use the colorectal cancer cell line, SW480, at low and high cell density as a model for mesenchymal-like cells at the invasive front and epithelial-like cells in the central mass of a colorectal tumour, respectively.

The initial aim was to carry out a limited screen for a set of developmental genes which are differentially normally expressed early on in mesodermal development, to determine whether they are differentially expressed in low and high density SW480 cells. We considered those genes that were expressed in low and high density SW480 cells might be involved in mesenchymal transitions in these cells and as such might be important in the progression of colorectal tumours. Our aim was to identify such differentially expressed genes and then to determine their possible role in inducing transitions between mesenchymal and epithelial cell types in SW480 cells. One such gene that we isolated in this screen was *BRACHYURY*, a developmental mesoderm inducing gene that is highly expressed in low density SW480 cells in contrast to expression levels in high density SW480 cells which are reduced to below 50%. In the first instance, our aim was to investigate whether *BRACHYURY* is involved in EMT; but if not, to determine its' possible role in processes such as cell migration, cell proliferation and/or other signalling pathways that might be relevant to the progression of colorectal tumours. It was our aim to then extend our finding in SW480 cells to address whether *BRACHYURY* was expressed similarly in other tumour cell lines.

# Chapter 2 - Materials and Methods

## 2.1 Source of cell lines

Cell lines (SW480, HCT116, LoVo, HT29, SW626 and T84) were obtained from the European Collection of Cell Cultures (ECACC). NTERA-2 (clone D1) cells were a gift from P.W. Andrews (University of Sheffield).

#### 2.2 Routine cell culture

SW480 and NTERA-2 cell lines were cultured in Dulbecco's modified Eagle's medium 1X (DMEM+GlutaMAX<sup>TM</sup>-I) (Invitrogen, GIBCO 61965) supplemented with 10% (v/v) foetal bovine serum (FBS) (Invitrogen, GIBCO 10270-098). HCT116 and HT-29 were cultured in 1X McCoy's 5A medium+GlutaMAX<sup>TM</sup>-I, (Invitrogen, GIBCO 36600) supplemented with 10% (v/v) FBS. LoVo cell were cultured in Ham's F-12 medium (Sigma, N6658) with 10% (v/v) FBS and T84 cells needed both DMEM with 10% (v/v) FBS and Ham's F-12 medium (Invitrogen, GIBCO N6658) with 10% (v/v) FBS in 1:1 ratio to grow. SW626 was maintained in Leibovitz L15 medium (Invitrogen, GIBCO 5520) with 10% (v/v) FBS. L-Glutamine (Invitrogen, GIBCO 25030, 200mM stock) is added if medium does not contain Glutamine. All cells were maintained in a 37°C incubator 5% CO<sub>2</sub> (apart from NTERA-2 cells which were maintained in 37°C incubator 10% CO<sub>2</sub> and SW626 cells which do not require CO<sub>2</sub>). For low density cells, SW480 cells were grown for 24 hours and for high density, cells were grown for 24 hours after they are confluent. The description of the origin and mutations of the cell lines are mentioned in table 2.1.

## 2.3 Freezing cells

Cells were grown in T75 flasks, washed with PBS and trypsinized. Medium was added after the cells started to detach and after a centrifugation of 5 minutes at 2000 rpm, cells from each flask were resuspended in 1ml freezing medium (1%

DMSO (w/v), 90% (v/v) FBS). Tubes were then transferred to a tub containing isopropanaol and incubated at -80°C for overnight. Next day, cells were transferred and stored in liquid nitrogen with proper passage numbers.

1 able 2.1 Description of the origin and mutations of the cell lines	used
--	------

Cell line	Age of patient (years)	Sex of patient	Site	Differentiati on	APC mutation 1	APC mutation 2	β-catenin mutation
SW480	50	male	colon	moderate	1338		
HCT116	1	male	colon	1	-	-	codon 45
LoVo	56	male	colon, distant lymph node	1	1114	1430	. =
HT29	43	female	colon	moderately well	853	-	-
Т84	76	male	colon, lung metastasis	/	1488	-	-
NTERA- 2	22	male	Testis, lung metastasis	1	1	/	1
SW626	46	female	Ovarian, colon metastasis	/	frameshift at codon 2941	inframe terminatio n at codon 9	1

/ = data not available; - = no mutation

## 2.4 Phenol-Chloroform extraction of DNA

First of all, chloroform equilibrated with isoamyl alcohol (24:1) was made. To the experimental samples, equal volumes of phenol and chloroform equilibrated with isoamyl alcohol were added (25:24:1). The mixture was nicely mixed by hand, followed by a spin of 13000 rpm for 10-15 minutes. The supernatant was collected in a separate tube and a same volume of chloroform was added. The supernatant was again collected in another tube, followed by a 13000 rpm spin for 5 minutes. The DNA was now ethanol precipitated.

#### 2.5 Ethanol precipitation of DNA

To the pellet, 2.5 to 3 times 100% (v/v) ethanol were added and then 3M sodium acetate of  $1/10^{\text{th}}$  the final volume was also added. This was left at -80°C for about 1 hour followed by a 13000 rpm spin for 10 minutes. The pellets were washed with 70% (v/v) ethanol, spun for 5-10 minutes and air dried. Finally, the DNA was resuspended in an appropriate volume of TE.

## 2.6 Cell counting by Haemocytometer

Cells were grown as required, medium aspirated, washed with PBS and trypsinized. The trypsin (Sigma, T3924) was deactivated by adding some media to it. 10µl of this were added to the end of the coverslips on a Hemocytometer. Under a microscope, the number of cells was counted on both the grids of the Haemocytometer. For each grid, cells were counted on the four corner squares.

Then, the average was taken for both the grids, divided by 4 (since we counted four corner squares) and multiplied by 25 (since, there were 25 squares in total). This total value was multiplied by  $10^4$  to get the number of cells per ml. The number of cells per flask can now be counted depending on how many ml there was in the flask.

#### 2.7 Total RNA isolation

Cells were grown in T75 flasks or in 6-well plates. Aspirate medium and wash the cells with PBS. 7.5ml of Trizol Reagent (Invitrogen, Cat#15596-026) was added to each T75 flask or 1ml Trizol to each well of 6-well plate. After pipetting up and down a few times, cells were transferred to separate tubes and incubated at room temperature for 5 minutes. To every 1ml, 0.2ml chloroform was added and shaken for 15 seconds in hand followed by 2-3 minutes incubation at room temperature. Cells were then centrifuged at 12,000g for 15 minutes at 4°C. The colourless upper

aqueous phase was transferred to a separate tube. To precipitate the RNA, 0.5 ml of isopropanol was added to per 1ml of this upper aqueous phase, incubated for 10 minutes at room temperature and then centrifuged at 12000g for 10 minutes at 4°C. The pellet was the collected and the RNA was washed with 1ml of 75% (v/v) ethanol and centrifuged at 7500g for 5 minutes at 4°C. The pellet was then dried on bench after which it was resuspended in 500 $\mu$ l RNAse free water and then incubated at 55-60°C for 10 minutes. Finally, the absorbance was measured using 1 $\mu$ l of RNA on a nanodrop (ND\_1000). The RNA isolated was run on 1% agarose gel to check the quality of the RNA.

## 2.8 cDNA synthesis by MLV method

To a sterile PCR tube, 1µl of oligo d(T) (500µg/ml), 1µg of RNA and 1µl of dNTP mix (10mM) were added and made upto 12µl with distilled water. This was mixed gently by pipetting and then heated to 65°C for 5 minutes, followed by a brief chill on ice. To this, 4µl of 5X first strand buffer, 2µl 0.1M DTT and 1µl of RNaseOUT were added, mixed gently and incubated at 37°C for 2 minutes. 1µl of M-MLV Reverse Transcriptase was added to gently mixing and 1µl of sterile water was added to the –RT experimental control. After incubating at 37°C for 50 minutes, followed by a 70°C heating for 15 minutes to deactivate the reaction, the sample was diluted with 130µl sterile water. The cDNA can be stored at -80°C for long term or can be used immediately for PCR. The cDNA made can be tested on a 2-3% (w/v) agarose gel.

## 2.9 cDNA synthesis by SuperscriptIII

The Superscript III 1st Strand Synthesis Kit (Invitrogen, Cat #18080-051) was used. 1µg of total RNA per 20µl of cDNA was used. Initial primer/ RNA mix was made by adding 1µl oligo d(T) primers, 10mM dNTP mix, 1µg RNA and made upto 3µl with DEPC treated water. This was heated to 95°C for 5 minutes. The enzyme mix was made by adding 2µl 10XRT buffer, 4µl of 25mM MgCl2, 2µl of

0.1 M DTT, 1µl RNaseOUT and 1µl SuperscriptII in the mentioned order. This was incubated at 50°C for 50 minutes followed by 85°C for 5 minutes. 1µl of RNaseH was then added and incubated for 20 minutes at 37°C. The cDNA can now be stored at -20°C.

## 2.10 Real time PCR using Biomix

In a sterile PCR tube, 25µl of BioMix<sup>™</sup> Red (Bioline-Cat#BIO-25005) was added along with 1µl each of the primers, 2µl of CDNA and 21µl of distilled water. The samples were run on a PCR machine on a suitable program. The initial heating was at 95°C for 5 minutes, denaturation at 95°C for 30 seconds, annealing for 30 seconds at suitable temperature for the primer set and the extension at 72°C for 30 seconds. The number of cycles was usually 30-35. Final elongation was done at 72°C for 10 minutes. The samples were then stored at 4°C. Refer table 2.2 for RT-PCR primer details.

Primers	Sequence	Primer base region	Primer length (bp)	Tm (°C)
FGF4 F	CTG CGG CGG CTC TAC TGC	566	18	62.8
FGF4 R	GAGGAAGTGGGTGACCTTCAT GG	929	23	64.2
FGFR3 F	CTC ATC ACT CTG CGT GGC TGG	1305	21	63.7
FGFR3 R	GCC CGG TCC TTG TCA ATG C	1747	19	61.0
FGFR2 F	ATG ACA CCGATG GTG CGG	1051	18	58.2
FGFR2 R	CCA CGT GCTTGA TCCACTGG	1870	20	61.4
FGF8 F	GAG TCC GAG GAG CCG AGA C	430	19	63.1
FGF8 R	GCC AGG CAG CAC CTA TCG	914	18	60.5
GDF3 F	GGC CAG GCA GTC CAA TTT CA	91	20	59.4
GDF3 R	CAG CAC CTT GTG GCC ATG G	222	19	61.0
ActivinA F	GGA TGT ACC CAA CTC TCA G	93104	20	56.7
ActivinA R	CCT AAC GGA CAC TCG TCA	93328	18	56.0
ActivinB F	TCG GCT CGA CTC GGC TCG	133	18	62.8
ActivinB R CAGGTAAAG CCA CAG GCTGGCC		456	22	65.8
BMP4 F	CTG CAG CTT CCC TGAGCC TTT C	324	22	64.0
BMP4 R	GAGGAG ATCGCCTCGTTCTCA GG	884	23	66.0
NODAL F	849	23	64.2	

#### Table 2.2: RT-PCR Primers

NODAL R	CTT GGC CAG ACT CCA CTG AGC	1271	21	63.7
BRACHYURY F	GGA GCC ACC GGT CCT CAC C	1341	19	65.3
BRACHYURY R	CTG GGA CAG CAC CGC TAC TGC	1851	21	65.7
TBX3 F	CAA GGT GCA CCT GGA GGC	1273	18	60.5
TBX3 R	CT CTT CGG CCA TTT CCA GTG	1839	20	59.4
TBX2 F	CCA GTT CCACAA GCTAGGCAC G	629	22	64.0
TBX2 R	CAC GTA GGT GCG GAA GGT GC	1044	20	63.5
FGF19 F	AGC TGC TTC CTG CGC ATC C	435	19	61.0
FGF19 R	CCA TGC TGT CGG TCT CCA GG	1056	20	63.5
FGFR1 F	CCA ACC GTG TGA CCA AAG TGG	919	21	61.8
FGFR1 R	GGT GAC CCT CCT TCA GCA GC	1160	20	63.5
NFkB F	GCT TGC AAA GAG GCA TGC C	1991	19	58.8
NFkB R	CAG CAG CAG CAA ACA TGG C	2481	19	58.8
SNAIL F	GCG GGA AGG CCT TCT CTA GG	624	20	63.5
SNAIL R	CCT CTT CCA GAG CCC AGG C	1064	19	63.1
Oct F	CTG GTG CCG TGA AGC TGG AG	410	20	63.5
Oct R	CGAGAGGAT TTT GAG GCT GCTG	848	22	62.1
E-cad F	ATT GTG GGC ATG AGC TGC	3762	18	56.0
E-cad R	ATC CTC AGC ATC AGT TTG C	4193	19	54.5

Note: Tm = melting temperature.

This was calculated using the formula 2[A+T] + 4[G+C], where, A= Adenine, T=Thymine, G=Guanine, C= Cytosine. Here, Tm was calculated using automated web based program (Oligoperfect designer, Invitrogen) based on the formula mentioned above (<u>http://tools.invitrogen.com/content.cfm?pageid=9716</u>). The annealing temperature was given by the lowest Tm for the PCR program + 3°C. Please refer to 'Gene names and Accession number' on page XI for the NCBI accession numbers of the genes. The gene nomenclature is followed according to MGI guideline (Eppig and Levan, 2009).

#### 2.11 Quantitative Real time RCR

After the cDNA was made by the MLV method, 1.1X cDNA loading mix was made by adding 1.5µl cDNA and 2.5µl distilled water in a sterile PCR tube. 1.1X Sybr Master mix for each primer set was made by adding 10µl 2X Sybr green PCR mastermix (Applied Biosystem, Cat#4309155, Lot#0803295), 2µl of the forward primer, 2µl of the reverse primer and 2µl of distilled water in a sterile tube. 4µl of the cDNA loading mix was added to the wells of 96-well plates in triplicate followed by addition of 16µl Sybr Master Mix to the cDNA. The 96-well plates were sealed by optically clear sealing film. Plates were either left in 4°C for few hours or directly loaded to AB7900HT machine. The program used on this machine was SDS v2.3 (an Applied Biosystem software used to get the Ct values of the samples from the AB7900HT machine) and the Ct values were collected and transferred to MSexcel sheets. The primers for the quantitative PCR are in table 2.3.

Note: The Ct (cycle threshold) value is the number of cycles required for the fluorescent signal to cross the threshold (which is sets by the user).  $\delta$ Ct gives the difference of the experimental and control Cts. The levels of the Ct value are inversely proportional to the amount of target nucleic acid in the sample. This would suggest that the lower the Ct level the greater the amount of target nucleic acid in the sample (Wisconsin diagnostic laboratory webpage,

http://www.wvdl.wisc.edu//PDF/WVDL.Info.PCR\_Ct\_Values.pdf)

Cts < 29 = abundant target nucleic acid in the sample

Cts of 30-37 = positive reactions with moderate amounts of target nucleic acid

Cts of 38-40 = weak reactions with minimal or no amounts

Primers	Sequence	Base no.	Tm (°C)	Product length (bp)	Primer length (bp)
CRB3 F	tctcccttcattgctgtgtg	588	59.83	72	20
CRB3 R	taagcactgggtggatctga	660	59.24	15	20
PALS1 F	gacacgtcgcagaatgagaa	82	59.99	07	20
PALS1 R	gttgcttgatggactgctga	178	59.99	97	20
EVA1 F	gggacagatgctcggttaaa	242	60.07	142	20
EVA1 R	tgggttggaagggatctatg	383	59.74	142	20
ZEB1 F	gcacaaccaagtgcagaaga	1832	60.03	146	20
ZEB1 R	actttgcctggttcaggaga	1977	59.84	140	20
ZEB2 F	gacctggacgtgaaggaaaa	4743	60.09	107	20
ZEB2 R	ggcacttgcagaaacacaga	4849	60.03	107	20
NANOG F	ttccttcctccatggatctg	377	60.00	(0)	20
NANOG R	ctttgggactggtggaagaa	436	60.08	00	20
WNT 11F	catggagctctgcttgtgaa	1271	60.14	70	20
WNT 11R	gcttccaagtgaaggcaaag	1342	59.99	12	20
IL-2 F	gaatcccaaactcaccagga	211	59.90	65	20
IL-2 R	tcagttctgtggccttcttg	275	59.01	0.5	20
IL-15 F	gaagccaactgggtgaatgt	984	59.97	116	20
IL-15 R	actttgcaactggggtgaac	1099	60.01	110	20
Orthopedia F	ctctgtgcccttggttgttt	2273	60.15	147	20
Orthopedia R	gaccagcacatgcagaaaga	2419	59.99	147	20
CDX2 F	tcttggggctgatgaagaag	1884	60.33	70	20
CDX2 R	agaccaacaacccaaacagc	1953	60.01	70	20
NOTCH1 F	tttctggggaaagacactgc	8566	60.23	102	20

## Table2.3: qRT-PCR Primers

NOTCH2 F   cctggaagtcattgccaagt   367   60.11   127   20     NOTCH2 R   ctgctttcccaacagcagtt   493   59.99   127   20     NOTCH3 R   atgcaggatcgaagaggagga   5975   59.80   86   20     NOTCH4 R   cccacgcaacagcagtggatgga   6060   59.91   85   20     NOTCH4 F   cccaggaatcggatgatggaa   4932   59.91   85   20     NOTCH4 R   ccacagcaacagtgcgatcaat   4932   59.91   85   20     FGFR1 R   tgctggtgcagcacaggatgagaat   1932   60.02   125   20     FGFR3 F   acctgaagatgggagcctttt   2999   60.07   145   20     TBX2 F   cggtgtgcagacgtttggacctttt   2999   60.07   145   20     TBX2 F   cggtgtgcagaggggagctttg   343   60.11   145   20     TBX2 F   cggtgtgcagaggggaggggagggggggagggggggggg	NOTCH1 R	tgtgttgctggagcatcttc	8667	59.99		20
NOTCH12 R   ctgctittcccaaacagcaget   493   59.99   127   20     NOTCH3 R   aagtggtccaacagcaget   6060   59.80   86   20     NOTCH3 R   ccaaggaatcggagaggag   5975   59.80   86   20     NOTCH4 F   cccaaggaatcggaatggaa   4848   60.00   85   20     NOTCH4 R   cccaaggaactggaatggaatggaatggaatggaatgga	NOTCH2 F	CH2 F cctggaagtcattgccaact 367 (		60.11	107	20
NOTCH3 F   a tg cag ag ta g ca ag g g g a   5975   59.80   86   20     NOTCH3 F   a ag tg g t ca aca ag cag cag cut   6060   59.91   85   20     NOTCH4 F   c cc ag g ca act g c t g a at t g a a   4848   60.00   85   20     NOTCH4 R   c ca ag ca act g c t g a at t g a at t g a at t g a at t g a g a	NOTCH2 R	ctgctttcccaaacagcttc	493	59.99	127	20
NOTCH3 R   a agiggiccacacagcagcit   6060   59.91   80   20     NOTCH4 F   cccaggaatclgagatggaa   4848   60.00   85   20     FGFR1 F   cgatgigcagagcatcact   1119   60.02   137   20     FGFR1 F   tggttaggaggcatcagacatag   1255   60.04   137   20     FGFR2 F   atgctgtaggaggagcatt   3028   60.02   125   20     FGFR3 R   tccgtgtaggaggagcatt   3028   60.07   145   20     FGFR3 R   tccgtgtagcaggaggagagcatt   3433   60.11   145   20     TBX2 F   cgtggtgtgctaggagcatt   1343   60.11   142   20     FGF8 R   ttggcctgagaactggaca   1150   60.44   20   20     FGF 8 R   ggacaccttggaca   800   59.84   127   20     BRACHYUNY F   ttgggctgagaactggaca   800   59.85   142   20     TBX3 F   ttttaggtcggcatagg   366   60.63   136   20   20	NOTCH3 F	atgcaggatagcaaggagga	5975	59.80		20
NOTCH4F   cccaggaatctgagatggaa   4848   60.00   85   20     NOTCH4R   ccacagcaactgctgacaat   4932   59.91   85   20     FGFRI F   cgatgigcaggcatcaact   1119   60.02   137   20     FGFR2R   atgctggttacgcagcatag   1255   60.04   137   20     FGFR2R   tccagttagacgtigggagcctit   2999   60.07   145   20     FGFR3R   acctgaagatgggagcctit   2999   60.07   145   20     TBX2R   cgatgtacatggttiggagcctit   2999   60.07   145   20     TBX2R   gtggatgtacatggttiggagcctit   253   59.88   85   20     14-3-3 oF   ttgtgctgaagcatgg   404   59.87   127   20     BACHYUNF f   tgggatgctggcagg   566   59.87   127   20     BACHYUNF f   ttgggccctglaaggcaagg   3734   60.02   136   20     SNAIL R   ccggacagggagattitga   1375   97   20   20   20	NOTCH3 R	aagtggtccaacagcagctt	6060	59.91	86	20
NOTCH4 R   ccacagcaaactgctgacat   4932   59.91   85   20     FGFR1 F   cgatgtgcagagcatcaact   1119   60.02   137   20     FGFR1 K   tgctgtcagcagcatag   3928   60.02   20   20     FGFR2 F   atgctgcgtcagagcatg   3928   60.02   20   20     FGFR3 R   atcctgaggatgggagcct1   2999   60.07   145   20     FGFR3 R   tccgttgtaccagctfttc   3143   60.11   145   20     TBX2 F   cgattgccgctatagttg   848   59.85   85   20     14-3.3 oF   ttgtggctgagactggag   140   20   20   20     FGF 8 F   ggacaccttggaagcggag   440   59.84   104   20     BRACHYURY F   ttgggctgaagacggag   430   59.85   142   20     BRACHYURY R   tcaggtctgcgttggaagacggag   3734   60.03   136   20     SNALL F   tttcagagccgtgtagg   3734   60.04   20   20     SNALL F <t< td=""><td>NOTCH4 F</td><td>cccaggaatctgagatggaa</td><td>4848</td><td>60.00</td><td>1</td><td>20</td></t<>	NOTCH4 F	cccaggaatctgagatggaa	4848	60.00	1	20
FGFRIF   c g at g t g c a g a c a t a a c t f GFRIR   1119   60.02   137   20     FGFR2R   a t g c t g c t g c a g c a a t g f g c t g a g a g a g a t g g g a g a c t t c c a g t a g g g g g g g g g g g g g g g g	NOTCH4 R	ccacagcaaactgctgacat	4932	59.91	85	20
FOFR R   tgctggttacgcaagcatag   1255   60.04   137   20     FGFR2 F   atgctgctgtagacgatag   3928   60.02   125   20     FGFR2 R   tccagttagacgitg gttg   4052   59.90   125   20     FGFR3 F   acctgaagtgggagcttt   2999   60.07   145   20     FGFR3 R   tccgttgtaccagcttttc   3143   60.11   145   20     TBX2 P   cgattgccgttaagtgtg   64   59.70   85   20     14.3-3 of R   tcaatctcggtagactggaca   1150   60.44   104   20     FGF 8 F   ggacaccttggaagcagag   440   59.87   127   20     BRACHYURY R   ttaggcgtagaactggaca   800   59.87   122   20     TBX3 F   ttitcagcccctgtgagg   3734   60.02   136   20     SNALL F   tccgacaagtgacgcatta   1487   59.72   93   20     E-cad R   ctggacagggagatttggag   1766   60.04   135   20     SNALL F	FGFR1 F	cgatgtgcagagcatcaact	1119	60.02		20
FGFR2 F   at got got got got got got got got got go	FGFR1 R	tgctggttacgcaagcatag	1255	60.04	137	20
FGFR2 R   tccagttagacgttgcgttg   4052   59.90   125   20     FGFR3 F   acctgaagatgggagccttt   2999   60.07   145   20     TBX2 F   cgattgcgctataagttc   764   59.70   85   20     TBX2 R   gtggatgtacatggtttg   848   59.85   20     14-3-3 oF   ttgtggtgagactggatt   848   59.84   104   20     FGF 8 F   ggacacttggagagggag   440   59.84   127   20     BRACHYURY F   ttagtggtgagaactggaca   800   59.85   142   20     TBX3 F   ttttcagagccgctgtaggt   3734   60.02   136   20     SNAIL F   tttcagcctcctfttggtg   1395   61.20   93   20     SNAIL F   ttcagcctcattgggagaaggaggatttga   1766   60.04   135   20     B-cade R   ggggtggaggatttga   1766   60.04   135   20     SNAIL F   tttcagcacgggaggatttga   1766   60.04   135   20     B-catenin F   acct	FGFR2 F	atgetgetgtcagacgattg	3928	60.02		20
FGFR3 F   a cctg aag at gg gag cctit   2099   60.07   145   20     FGFR3 R   tccg ttg accage cttttc   3143   60.11   145   20     TBX2 F   cg attg ccg cta taag ttc   3143   60.11   20   20     TBX2 R   gt gg at gt acat gcg ttt gg   848   59.85   85   20     14-3-3 oF   tt gt gc tg ag act gg aca   1150   60.44   104   20     PGF 8 F   gg acaccttt gg ag gc ag g   566   59.87   127   20     BRACHVURY F   ttg gc tg ag acat gg aca   941   60.18   142   20     TBX3 R   cact ga ttt g ag gc aca gg   3734   60.02   136   20     SNALL F   tttc ag ag ccg ctg ta gg t   374   60.02   136   20     SNALL F   ct ga a ag tg aca gc aca gg   3869   60.63   20   20     E-cad F   ct gg aca gg ga ga att ga act ga ac   1375   20   20   20     LAMIN A/C F   tt cca cat cat gg ga gg gt aca   1900   60.12	FGFR2 R	tccagttagacgttgcgttg	4052	59.90	125	20
FGFR3 R   tccgitgiaccagcetitte   3143   60.11   145   20     TBX2 F   cgaitgecgetataagtice   764   59.70   85   20     IH32 R   gtggatgiacatgegtitgg   848   59.85   85   20     I4-3-3 σ F   ttgggetgaagaeggag   1150   60.44   104   20     FGF 8 F   ggacaccttiggaagaggagg   566   59.87   127   20     BRACHYURY F   ttggggetgaagaggaggaggaggaggaggaggaggaggaggaggag	FGFR3 F	acctgaagatgggagccttt	2999	60.07		20
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	FGFR3 R	tccgttgtaccagccttttc	3143	60.11	145	20
TBX2 R   giggaigincaigogitigg   848   59.85   85   20     14-3-3 or F   tigtggcigagaaciggaca   1150   60.44   104   20     FGF 8 F   ggacaccttiggagacgaga   1253   59.98   104   20     FGF 8 F   ggacaccttiggagacgaga   566   59.87   127   20     BRACHYURY F   tigtgcigagaciggaca   800   59.85   142   20     TBX3 F   ticaatctcggtciggagi   3734   60.02   136   20     SNAIL F   tittcagagccgiggagagaggatigg   3784   60.02   136   20     SNAIL R   citcagacaggagaggagaggatiggag   1487   59.72   93   20     E-cad F   ciggacaggagagattigag   1395   61.20   93   20     B-catenin R   actcractgigagagagattigag   927   59.93   90   20     LAMINA/C F   ticcccatcataiggag   927   59.98   89   20     STAT3 F   cittggaacgaagggtaca   1974   59.96   146   20	TBX2 F	cgattgccgctataagttcc	764	59.70		20
14-3-3 o F tigiggcigagaactggaca 1150 60.44 104 20   14-3-3 o R tcaatctcggtcttgcactg 1253 59.98 127 20   FGF 8 F ggacacctttggaagacggag 440 59.84 127 20   BRACHYURY F ttgtggctgagaactggaca 800 59.85 142 20   BRACHYURY F ttgtggctgagactggaca 941 60.18 142 20   SNAIL F tttcagggccgttgggt 3734 60.02 136 20   SNAIL R cactgatttgaggccaaagg 3869 60.63 136 20   SNAIL R ctgacaagtgacagccatta 1447 72 93 20   E-cad F ctggacaggaggagtttga 1766 60.04 135 20   B-catenin F acttccactatcatgtgg 27 59.93 90 20   AdmiNA/C F ttctcctccttctttttce 2358 59.62 125 20   NF-KB F ctggaagcagaatgcaa 297 59.98 89 20   STAT3 F ctttggaacgaagggtaca 1974 59.96 125 20	TBX2 R	gtggatgtacatgcgtttgg	848	59.85	85	20
14-3-3 σR   tcaatcteggtottgeactg   123   59.98   104   20     FGF 8F   ggacacctttggaageagag   440   59.84   127   20     BRACHYURY F   ttgtggctgagaactggaca   800   59.85   142   20     BRACHYURY F   ttgtggctgagaactggaca   800   59.85   142   20     TBX3 F   ttttcagagcgctgtaggt   3734   60.02   136   20     SNAIL F   tttcagagcggtgaggtgsgagatttgg   1395   61.20   93   20     E-cad F   ctggacaggaggagatttgg   900   60.12   135   20     β-atenin F   acctttccatatcatacatacag   1900   60.12   135   20     LAMINA/C F   ttctcccatcatggag   927   59.93   90   20     LAMINA/C F   ttctctccccttcttttcc   238   59.62   125   20     LAMINA/C R   gggcagaaagcagaagcta   2482   59.73   125   20     STAT3 F   cctttggaacgaagggtaca   1974   59.96   146   20	14-3-3 σ F	ttgtggctgagaactggaca	1150	60.44		20
FGF 8 F   ggacacctitggaagcagag   440   59.84   127   20     BRACHYURY F   tigtggctgagaccgagt   566   59.87   127   20     BRACHYURY F   tigtggctgagaccgagt   566   59.87   142   20     BRACHYURY R   tcaatctcggtcttgcactg   941   60.18   142   20     TBX3 F   ttitcagcccctgttgggt   3734   60.02   136   20     SNAIL F   ttitcagcccctgttiggtg   1395   61.20   93   20     E-cad R   ggcgtgtacaggagttitga   1766   60.04   135   20     B-catenin F   acctitccatcatcggag   900   60.12   135   20     G-catenin R   aatccactggtgaaccaagg   1066   59.97   90   20     LAMINA/C F   ttcctcccttcttitcc   2358   59.62   125   20     LAMINA/C R   gggcagaaggagtaca   1015   59.98   89   20     STAT3 F   cctitggaacgagggagata   217   59.98   89   20     STAT3	14-3-3 σ R	tcaatctcggtcttgcactg	1253	59.98	104	20
FGF 8 R   aatctccgtgaagacgcagt   566   59.87   127   20     BRACHYURY F   ttgtggctgagaactggaca   800   59.85   142   20     TBX3 F   ttttcagagccgctgtaggt   3734   60.02   136   20     TBX3 R   cactgatttgaggcaaagg   3869   60.63   136   20     SNAIL F   tttcagcctcctgttggtg   1395   61.20   93   20     E-cad F   ctggacagtggaagactggag   1766   60.04   135   20     B-catenin F   acctttcccatcatcggag   927   59.93   90   20     G-catenin R   atccactcggtgaacagag   927   59.93   90   20     LAMINA/C F   ttctccctccttcctttcc   2358   59.62   125   20     NF-KB F   ctggaacgaagagtaga   219   60.08   146   20     STAT3 F   cctttggaacgaagggtaca   1974   59.96   146   20     STAT3 F   cctttggaacgaagggtaca   1974   59.96   146   20     IL-10 F </td <td>FGF 8 F</td> <td>ggacacctttggaagcagag</td> <td>440</td> <td>59.84</td> <td></td> <td>20</td>	FGF 8 F	ggacacctttggaagcagag	440	59.84		20
BRACHYURY F   trigtgecigagaactggaca   500   59.85   142   20     BRACHYURY R   tcaatclcggtcttgcactg   941   60.18   142   20     TBX3 F   ttttcagagccgctgtaggt   3734   60.02   136   20     SNAIL F   tttcaggcccctgttggtg   1395   61.20   93   20     SNAIL F   ttcagacccctgtttggtg   1395   61.20   93   20     E-cad F   ctggacagggggagacggggggggggggggggggggggg	FGF 8 R	aateteegtgaagacgeagt	566	59.87	127	20
BRACHYURY R   tragatceggtottgeactg   941   60.18   142   20     TBX3 F   tttteagaccgcgtaggt   3734   60.02   136   20     TBX3 R   cactgatttgagggcaaagg   3869   60.63   20     SNAIL F   tttteagacccectgtttggtg   1395   61.20   20     SNAIL R   ccgacaagtgacagcatta   1487   59.72   93   20     E-cad F   ctggacagggagattttga   1766   60.04   135   20     B-catenin F   actttcccatcatcgtgag   927   59.93   90   20     LAMINA/C F   ttcctccctctcttttcc   2358   59.62   125   20     NF-KB F   ctggaagacgaaggatca   2482   59.73   125   20     NF-KB R   cctttctgcttgcaaataggc   1015   59.98   89   20     STAT3 F   ccttteggaacgaagggtaca   1974   59.96   146   20     IL-10 F   cctttggaacgaagggtaca   1158   59.90   146   20     VIMENTIN F   gagaacttgccgtgaagc <td>BRACHYURY F</td> <td>ttotooctoagaactogaca</td> <td>800</td> <td>59.85</td> <td></td> <td>20</td>	BRACHYURY F	ttotooctoagaactogaca	800	59.85		20
TBX3 F   titte agage cgt gt agg   3734   60.02   136   20     TBX3 R   cact gatt tg agg caa agg   3869   60.63   136   20     SNAIL F   ttt cag cac gg agg cat tg gt   1395   61.20   93   20     E-cad F   ct g g ac agg gg ag gt tt ga   1766   60.04   135   20     E-cad R   gg cgt gt gt catt cacat cag   1900   60.12   135   20     B-catenin R   a at ccact gg tg ac cag g   27   59.93   90   20     LAMIN A/C F   tt cc ct ccat cat cg tg ag   27   59.93   90   20     LAMIN A/C R   gg g cag aa aa ag cag aa g ct a   2482   59.73   125   20     LAMIN A/C R   ct g g a c g a ag g gt a ca   1974   59.98   89   20     STAT3 F   ctt tg g a a c g a ag g gt a ca   1974   59.96   146   20     STAT3 F   ctt tg g a c g a ag g gt a ca   1158   59.90   90   20     IL-10 R   ct cca a agt g c g g g g a ca   1158 <t< td=""><td>BRACHYURY R</td><td>tcaatctcggtcttgcactg</td><td>941</td><td>60.18</td><td>142</td><td>20</td></t<>	BRACHYURY R	tcaatctcggtcttgcactg	941	60.18	142	20
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	TBX3 F	ttttcagagccgctgtaggt	3734	60.02		20
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	TBX3 R	cactgatttgagggcaaagg	3869	60.63	136	20
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	SNAIL F	tttcagcctcctgtttggtg	1395	61.20	· · · · · · · · · · · · · · · · · · ·	20
$\begin{array}{c c c c c c c c c c c c c c c c c c c $	SNAIL R	ccgacaagtgacagccatta	1487	59.72	93	20
$\begin{array}{c c c c c c c c c c c c c c c c c c c $	E-cad F	ctopacapppappattttpa	1766	60.04		20
$\begin{array}{c c c c c c c c c c c c c c c c c c c $	E-cad R	ggcgttgtcattcacatcag	1900	60.12	135	20
$\begin{array}{c c} \begin{array}{c c} \hline P-catenin R & a at cc a ct gg tg a a cc a ag c & 1066 & 59.97 & 90 & 20 \\ \hline LAMINA/C F & tt ct c ct cc ct t ct tt tc c & 2358 & 59.62 & 125 & 20 \\ \hline LAMIN A/C R & gg g cag a a a ag cag a ag ct a & 2482 & 59.73 & 125 & 20 \\ \hline NF-KB F & ct gg a a g ca cg a at g a c c a a \\ r & ct t ct g ct t g c a a g a g g g t a c a & 1015 & 59.98 & 89 & 20 \\ \hline STAT3 F & cctt t g g a a c g a g g g t a c a & 1974 & 59.96 & 146 & 20 \\ \hline STAT3 R & cct t ct g c a g g g g g a t a & 1118 & 59.90 & 146 & 20 \\ \hline IL-10 F & cct tt g g a a c g a g g g t a c a & 1158 & 59.90 & 90 & 20 \\ \hline IL-10 R & ct cc c a a a g t g g g g a t a & 11247 & 60.07 & 90 & 20 \\ \hline VIMENTIN F & g a g a a ct tt g c c g tt g a a g g c a a d t & 1247 & 60.07 & 90 & 20 \\ \hline VIMENTIN R & ct ca at g t c a g g g g a c a & 1247 & 60.07 & 90 & 20 \\ \hline VIMENTIN R & ct g g a c g g t g a a a d t & 1247 & 60.07 & 90 & 20 \\ \hline PLAUR F & cct g g a g ct tg a a a a t c t g & 389 & 59.96 & 105 & 20 \\ \hline PLAUR R & cg g c a g t c a at g a g g a a a d t & 1207 & 59.96 & 79 & 20 \\ \hline SOX2 R & ct g t a g g c c a a g g g a a c t & 1285 & 60.26 & 79 & 20 \\ \hline SOX2 R & ct g t a g c c c a a a t c g g a a a t c t g g a a c t t t g g d c c a a g g g a a c t & 1285 & 60.26 & 79 & 20 \\ \hline OCT-4 R & a g c g a t c a a g c g a a a t g & 3125 & 59.99 & 61 & 20 \\ \hline OCT-4 F & t a g c c t g g a g c g c g c a a t g g g a a c t & 911 & 60.15 & 118 & 20 \\ OCT-4 F & t a g c c t g g a g c g c g c a a c a g g & 1271 & 59.97 & 81 & 20 \\ \hline GATA2 R & a c c g g a a g t g c c a a g g a & 1271 & 59.97 & 81 & 20 \\ \hline GATA2 R & a c c g g a a g t g c c a a g g g a & 1271 & 59.97 & 81 & 20 \\ \hline GATA4 R & a c g g g t a c c a t t c g g c a c & 1351 & 59.99 & 81 & 20 \\ \hline GATA4 R & a c g g g t c a c t t t g g c t g g d g d & 3440 & 59.84 & 149 & 20 \\ \hline \end{array}$	β-catenin F	acctttcccatcatcgtgag	927	59.93	72:22	20
LAMINA/C F   ttctcctcctttcc   2358   59.62   125   20     LAMIN A/C R   gggcagaaaagcagaagcta   2482   59.73   125   20     NF-KB F   ctggaagcacgaatgacaga   927   59.98   89   20     STAT3 F   cctttggaacgaagggtaca   1974   59.96   146   20     IL-10 F   cctttggaacgaagggtaca   1158   59.90   90   20     VIMENTIN F   gagacacttgccgtggatta   1247   60.07   90   20     VIMENTIN F   gagacgtcaatgggccatct   6398   60.07   144   20     VIMENTIN R   ctcaatgtcaagggccatct   6398   60.07   144   20     PLAUR F   cctggagcttgaaagtggaaagt   943   60.26   105   20     BCL2L1 F   tggtcccttgcagtaggaaagt   943   60.26   79   20     SOX2 R   ctgtagcctggctacttgggaact   1285   59.99   61   20     SOX2 F   gttgcctggcttccttttg   8325   59.99   61   20 <td< td=""><td>β-catenin R</td><td>aatccactggtgaaccaagc</td><td>1066</td><td>59.97</td><td>90</td><td>20</td></td<>	β-catenin R	aatccactggtgaaccaagc	1066	59.97	90	20
$\begin{array}{c c c c c c c c c c c c c c c c c c c $	LAMINA/C F	ttctcctcccttccttttcc	2358	59.62		20
NF-KB F   ctggaagcacgaatgacaga   927   59.98   89   20     STAT3 F   ccttctgcttgcaaataggc   1015   59.98   89   20     STAT3 F   ccttctgcaacagagggtaca   1974   59.96   146   20     STAT3 R   ccttctccacccaagtgaa   2119   60.08   146   20     IL-10 F   ccttctggaacgacgggtaca   1158   59.90   90   20     VIMENTIN F   gagaactttgccgttgaagc   6255   60.00   144   20     VIMENTIN R   ctcggaggctgaaaatctgc   839   59.96   105   20     PLAUR F   cctggagctgaacaggggaaagt   943   60.26   105   20     BCL2L1 F   tggtcccttgcagctagttt   1207   59.96   79   20     SOX2 R   ctgtagccccaaatcggata   8265   59.92   61   20     SOX2 R   gttgcctggcttctcttttg   8325   59.99   61   20     OCT-4 R   agcgatcaggagggcacta   911   60.15   118   20     OCT-4 R </td <td>LAMIN A/C R</td> <td>gggcagaaaagcagaagcta</td> <td>2482</td> <td>59.73</td> <td>125</td> <td>20</td>	LAMIN A/C R	gggcagaaaagcagaagcta	2482	59.73	125	20
$\begin{array}{c c c c c c c c c c c c c c c c c c c $	NF-KB F	ctggaagcacgaatgacaga	927	59.98		20
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	NF-KB R	ccttctgcttgcaaataggc	1015	59.98	89	20
STAT3R   ccttctccacccaagtgaa   2119   60.08   146   20     IL-10 F   cctttggaacgaagggtaca   1158   59.90   90   20     IL-10 R   ctcccaaagtgctgggatta   1247   60.07   90   20     VIMENTIN F   gagaactttgccgttgaagc   6255   60.00   144   20     VIMENTIN R   ctccaatgtcaagggccatct   6398   60.07   144   20     PLAUR F   cctggagcttgaaaatctgc   839   59.96   105   20     BCL2L1 F   tggtcccttgcagctagttt   1207   59.96   79   20     SOX2 R   ctgtagccccaaagggaact   1285   60.26   79   20     SOX2 F   gttgcctggcttctctttg   8325   59.99   61   20     OCT-4 R   agcgatcaagcagcgactaat   911   60.15   118   20     GATA2 R   accggaagatgtccaacaag   1271   59.97   81   20     GATA4 R   acgggtcactatctggcaa   780   59.17   52   20     GATA4 F	STAT3 F	cctttggaacgaagggtaca	1974	59.96		20
IL-10 Fcctttggaacgaagggtaca115859.909020IL-10 Rctcccaaagtgctgggatta124760.079020VIMENTIN Fgagaactttgccgttgaagc625560.0014420VIMENTIN Rctcaatgtcaagggccatct639860.0714420PLAUR Fcctggagcttgaaaatctgc83959.9610520PLAUR Rcggcagtcaatgaggaaagt94360.2610520BCL2L1 Ftggtcccttgcagctagttt120759.967920BCL2L1Rattctgaggccaagggaact128560.267920SOX2 Rctgtagccccaaatcggata826559.926120SOX2 Fgttgcctggcttctctttg832559.996120OCT-4 Ragcgatcaagcggactat91160.1511820OCT-4 Ftagcctgggtaccaaagg127159.978120GATA2 Raccggaagatgtccaacaag127159.998120GATA4 Racgggtcactatctgtgcaa78059.175220GATA4 Ftgccttgggacttggag83159.7220GATA6 Rttggcttgggacttgagagt344059.8414920	STAT3R	ccttctccacccaagtgaa	2119	60.08	146	20
IL-10 Rctcccaaagtgctgggatta124760.079020VIMENTIN Fgagaactttgccgttgaagc625560.0014420VIMENTIN Rctcaatgtcaagggccatct639860.0714420PLAUR Fcctggagcttgaaaatctgc83959.9610520PLAUR Rcggcagtcaatgaggaaagt94360.2610520BCL2L1 Ftggtcccttgcagctagtt120759.967920BCL2L1 Rattctgaggccaagggaact128560.267920SOX2 Rctgtagccccaaatcggata826559.926120SOX2 Fgttgcctggcttctcttttg832559.996120OCT-4 Ragcgatcaagcagcgactat91160.1511820OCT-4 Ftagcctgggtaccaacag127159.978120GATA2 Raccggaagatgtccaacaag127159.998120GATA4 Racgggtcactatctggcaa78059.175220GATA4 Ftgccgttcatcttgggag83159.7220GATA6 Rttggcttgggacttgaggt344059.8414920	IL-10 F	cctttggaacgaagggtaca	1158	59.90	00	20
$\begin{array}{c c c c c c c c c c c c c c c c c c c $	IL-10 R	ctcccaaagtgctgggatta	1247	60.07	90	20
VIMENTIN Rctcaatgtcaagggccatct $6398$ $60.07$ $144$ $20$ PLAUR Fcctggagcttgaaaatctgc $839$ $59.96$ $105$ $20$ PLAUR Rcggcagtcaatgaggaaagt $943$ $60.26$ $105$ $20$ BCL2L1 Ftggtcccttgcagctagttt $1207$ $59.96$ $79$ $20$ BCL2L1Rattctgaggccaagggaact $1285$ $60.26$ $79$ $20$ SOX2 Rctgtagccccaaatcggata $8265$ $59.92$ $61$ $20$ SOX2 Fgttgcctggcttctcttttg $8325$ $59.99$ $61$ $20$ OCT-4 Ragcgatcaagcagcgactat $911$ $60.15$ $118$ $20$ OCT-4 Ftagcctgggtaccaaaagg $1271$ $59.97$ $81$ $20$ GATA2 Raccggaagatgtccaacaag $1271$ $59.99$ $81$ $20$ GATA4 Racgggtcactatctgtgcaa $780$ $59.17$ $52$ $20$ GATA4 Ftgccgttcatcttgtggtag $831$ $59.72$ $20$	VIMENTIN F	gagaactttgccgttgaagc	6255	60.00	1.4.4	20
PLAUR F PLAUR Rcctggagcttgaaaatctgc cggcagtcaatgaggaaagt $839$ 943 $59.96$ $60.26$ $105$ $20$ $20$ BCL2L1 F BCL2L1Rtggtcccttgcagctagttt attctgaggccaagggaact $1207$ $1285$ $59.96$ $60.26$ $79$ $20$ 	VIMENTIN R	ctcaatgtcaagggccatct	6398	60.07	144	20
PLAUR Rcggcagtcaatgaggaaagt94360.2610520BCL2L1 Ftggtcccttgcagctagttt120759.967920BCL2L1Rattctgaggccaagggaact128560.267920SOX2 Rctgtagccccaaatcggata826559.926120SOX2 Fgttgcctggcttctcttttg832559.996120OCT-4 Ragcgatcaagcagcgactat91160.1511820OCT-4 Ftagcctgggtaccaaaatg102859.8211820GATA2 Raccggaagatgtccaacaag127159.978120GATA4 Racgggtcactatctgtgcaa78059.175220GATA6 Rttggcttgggacttgagagt344059.8414920	PLAUR F	cctggagcttgaaaatctgc	839	59.96	105	20
$\begin{array}{c c c c c c c c c c c c c c c c c c c $	PLAUR R	cggcagtcaatgaggaaagt	943	60.26	105	20
BCL2L1Rattctgaggccaagggaact128560.267920SOX2 Rctgtagccccaaatcggata826559.926120SOX2 Fgttgcctggcttctcttttg832559.996120OCT-4 Ragcgatcaagcagcgactat91160.1511820OCT-4 Ftagcctgggtaccaaatg102859.8211820GATA2 Raccggaagatgtccaacaag127159.978120GATA2 Ftctcctgcatgcactttgac135159.998120GATA4 Racgggtcactatctgtgcaa78059.175220GATA6 Rttggcttgggacttgagagt344059.8414920	BCL2L1 F	teetcccttecaectaettt	1207	59.96	1.01000-03145	20
$\begin{array}{c c c c c c c c c c c c c c c c c c c $	BCL2L1R	attetgaggecaagggaact	1285	60.26	79	20
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	SOX2 R	ctata accesa a tega a ta	8265	50.02		20
SOR2 Igrtgeoriggettetetting $3323$ $33,37$ 20OCT-4 Ragcgatcaagcagcgactat911 $60.15$ $118$ $20$ OCT-4 Ftagcctgggtaccaaaatg $1028$ $59.82$ $118$ $20$ GATA2 Raccggaagatgtccaacaag $1271$ $59.97$ $81$ $20$ GATA2 Ftctcctgcatgcactttgac $1351$ $59.99$ $81$ $20$ GATA4 Racgggtcactatctgtgcaa $780$ $59.17$ $52$ $20$ GATA4 Ftgccgttcatcttgtggtag $831$ $59.72$ $20$ GATA6 Rttggcttgggacttgagagt $3440$ $59.84$ $149$ $20$	SOX2 F	attacctaacttctctttta	8205	59.92	61	20
OCT-4 Ftagcctgggtaccaageggegactat911 $60.13$ 11820OCT-4 Ftagcctgggtaccaagatg102859.8211820GATA2 Raccggaagatgtccaacaag127159.978120GATA2 Ftctcctgcatgcactttgac135159.998120GATA4 Racgggtcactatctgtgcaa78059.175220GATA6 Rttggcttgggacttgagagt344059.8414920	OCT-4 R	accatcaaccaccactat	011	60.15	-	20
$ \begin{array}{c c c c c c c c c c c c c c c c c c c $	OCT-4 F	tageetageagegaetat	1028	50.82	118	20
GATA2 Ftctcctgcatgcactttgac127159.978120GATA2 Ftctcctgcatgcactttgac135159.998120GATA4 Racgggtcactatctgtgcaa78059.175220GATA4 Ftgccgttcatcttgtggtag83159.725220GATA6 Rttggcttgggacttgagagt344059.8414920	GATA2 R	accoggaggtatocalatig	1271	59.02		20
GATA4 Racgggtcactatctgtgcaa78059.1720GATA4 Ftgccgttcatcttgtggtag83159.7220GATA6 Rttggcttgggacttgagagt344059.84149	GATA2 F	tetectocatocactitoac	1351	59.90	81	20
GATA4 Ftgccgttcatcttgtggtag83159.725220GATA6 Rttggcttgggacttgagagt344059.8414920	GATA4 R	acgogtcactatetotocaa	780	59.17		20
GATA6 R ttggcttgggacttgagagt 3440 59.84 149 20	GATA4 F	taccattcatcttataataa	831	59.72	52	20
140 20	GATA6 R	ttggcttggggcttggggtt	3440	59.84	2 10-21	20
GATA6 F   gaacteecegactgacaaaa   3588   60.09   177   20	GATA6 F	gaactccccgactgacaaaa	3588	60.09	149	20

FOXD3 F	actctgcctctccccaattt	1551	60.07	05	20
FOXD3 R	tttgctcgaccagcttaggt	1645	60.02	95	20
VIMENTIN F	gagaactttgccgttgaag	6255	60.00	144	20
VIMENTIN R	ctcaatgtcaagggccatct	6398	60.07	144	20
MYC F	agggatcgcgctgagtataa	118	59.83	74	20
MYC R	tgcctctcgctggaattact	191	59.98	74	20

Note: Tm = melting temperature.

Tm was calculated using the formula 2[A+T] + 4[G+C], where, A= Adenine, T=Thymine, G=Guanine, C= Cytosine. Here, Tm was calculated using automated web based program (Oligoperfect designer, Invitrogen) based on the formula mentioned above (<u>http://tools.invitrogen.com/content.cfm?pageid=9716</u>).

The annealing temperature was given by the lowest Tm for the PCR program + 3°C. Please refer 'Gene names and Accession number' on page XI for the NCBI accession numbers of the genes. The gene nomenclature is followed according to MGI guideline (Eppig and Levan, 2009).

#### 2.12 Selection of control genes for qRT-PCR:

LAMIN C and  $\beta$ -catenin were used as control genes for the qRT-PCR experiments. The levels of expression of these two genes (LAMIN C and  $\beta$ -catenin) were very similar in low and high density cells in all the cell lines used (Figure 3.2, Chapter 3) and also throughout the different cell lines (Figure 3.3, Chapter 3). [Note: Housekeeping genes such as  $\beta$ -actin varies in low and high density SW480 cells. So, it was not used as a control for our experiments].

Also, *LAMIN C* and  $\beta$ -catenin expression levels remain similar even after knocking down *BRACHYURY* in low density SW480 cells (Figure 3.2 and 3.3, Chapter 3). Therefore, since these genes do not change their expression levels on *BRACHYURY* RNAi, we used *LAMIN C* and  $\beta$ -catenin for RNAi experiments too.

## 2.13 A worked example for analyzing qRT-PCR data using $2^{-\delta\delta C}$ method

Reference: Livak and Schmittgen, 2001; Dharmaraj 2006; Hunt 2009; Michael, 2001 Note: This is an example. Refer to table 2.4

Table 2.4 qRT-PCR data analysis

	β- catenin			SNAIL						2^-δδCt
Sample	Ct	Mean Ct	S.D.	Ct	Mean Ct	S.D.	δCt	δδCt	S.D.	Fold increase (relative expression)
SW626	21.26	21.66	0.92	27.72	27.72	0.322	6.063	0	0.98	1
sub	22.71			27.35						
	21.00			27.94			1			
SW626	22.37	22.150	0.561	27.69	27.78	0.07	5.631	-	0.565	1.348
super	21.51			27.82	1			0.43	0.000	
	22.56			27.82						

Abbreviations: sub = sub-confluent cells; super = super-confluent cells; Ct = cycle threshold; S.D. = standard deviation.

 $\delta Ct$  = The difference between the mean Ct of the experimental and reference gene  $\delta\delta Ct$  = The difference between the  $\delta Ct$  of the condition we are comparing and the condition to which we are comparing.

Fold increase is calculated using the formula  $2^{-}\delta\delta$ Ct. This formula  $(2^{-}\delta\delta$ Ct) was derived by Applied Biosystems in 1997 by ABI prism 7700 sequence detection system)

Here, expression of *SNAIL* is analysed in sub- and super-confluent SW626 cells. The data was compared to  $\beta$ -catenin and the sub-confluent data was set to 1 to which super-confluent data was compared. Each sample has a triplicate value. The Mean Ct will give the average of those three Ct value. So, we have Mean Ct for the control gene (here,  $\beta$ -catenin Mean Ct) and Mean Ct of the experimental gene (here, *SNAIL* Mean Ct).  $\delta$ Ct will give the difference of the Mean Cts of the experimental and control gene.  $\delta$ SCt will give the difference between the  $\delta$ Ct of the condition we are comparing (here,  $\delta$ Ct of *SNAIL* in super-confluent SW626 cells) and the condition to which we are comparing (here,  $\delta$ Ct of *SNAIL* in sub-confluent SW626 cells). Here,  $\delta$ SCt of the *SNAIL* in sub-confluent cells is made to 0 by subtracting its own value. This gave the value for  $2^{-\kappa}\delta\delta^{Ct}$  for *SNAIL* expression in sub-confluent SW626 cells was compared to 1 and was calculated using  $2^{-\kappa}\delta\delta^{Ct}$ . This gave the expression to be 1.3 times stronger than *SNAIL* expression in sub-confluent SW626 cells (which is set as 1). The final standard deviation is given by standard deviation if the  $\beta$ -catenin and

standard deviation of *SNAIL* Ct value. A graph can now be plotted for *SNAIL* expression in sub-confluent SW626 cells as 1 and its expression in super-confluent SW626 cells as 1.3 and the final standard deviations will give the error bars. Note: When we have done multiple repeats of the experiment, we calculate the fold difference  $(2^{-\delta\delta Ct})$  for each experiment, and the take an average of all the  $2^{-\delta\delta Ct}$  values from all repeats to plot the graph. The final standard deviation (SD) is given by the calculating the SD of all repeats, which gives the error bars.

## 2.14 Western blot

Cells were grown as required, washed twice with 1X PBS and scraped to detach from the flask surface/ 6-well plate surfaces using a scraper. Then the cells were centrifuged at 2000g for 5 minutes and resuspended in lysis buffer (50mM Tris-HCl pH7.4, 200mM sodium chloride, 0.5% (v/v) Triton X-100, 1mM AEBSF (4-(2-aminoethyl)-benzenesulfonyl fluoride) and one complete, mini, EDTA-free protease inhibitor cocktail tablet (Roche- cat#11836170001) per 10ml of lysis buffer. Each tablet contains 0.02 mg/ml Pancreas-extract, 0.0005 mg/ml Thermolysin, 0.002 mg/ml Chymotrypsin, 0.02 mg/ml Trypsin, 0.33 mg/ml Papain. An equal volume of 2X (loading) Laemmli buffer (20% (v/v) glycerol, 0.2% (w/v) bromophenol blue, 4% (w/v) SDS (sodium dodecyl sulphate), 200mM DTT (Dithiothreitol), 100mM Tris-HCl pH6.8) was added and the lysates were boiled for 5-10 minutes. The lysates were then loaded on 1mm thick 4-12% (w/v) Tris glycine precast gels (Invitrogen- cat#EC60352BOX) or 0.75mm thick 7.5% (w/v) SDS-PAGE (sodium dodecyl sulphate - polyacrylamide gel electrophoresis). Precision Plus Protein Dual colour standards (BioRad- cat#161-0374) were also run as marker to detect the protein sizes. Gels were run at 125 volts for 2-3 hours. Proteins were transferred to methanol wet Immobilon-P PVDF membrane (Milliporecat#IPVH00010) at 500mA for 15-18 hours in 2x Towbin buffer (380mM Glycine, 50mM Tris) with(only in case of APC transfers) or without 0.02% (w/v) SDS. After transfer, the membrane was blocked in 10% (w/v) milk overnight at 4°C or 1hour at

room temperature. Immunodetection was performed in 10% (w/v) milk in PBS with 0.1% (v/v) Tween 20 (Sigma- cat#D-1379) for monoclonal primary antibodies and 0.5% (v/v) Tween 20 for polyclonal primary antibodies. Incubation with primary antibody solutions was at room temperature for 1 hour or 4°C overnight, after which a 15 minutes wash was done in milk solution at room temperature. Incubation with secondary antibodies was again at room temperature for 1 hour or 4°C overnight. Antibody details can be seen in tables 2.5 and 2.6. Following incubation with secondary antibody membranes were washed for 10 minutes in milk solution, followed by two 10 minutes washes in PBS/0.1% (v/v) Tween 20 (for monoclonal primaries) or PBS/0.5% (v/v) Tween 20 (for polyclonal primaries). Bands were gained on Kodak X-OMAT AR film with Super signal west pico chemiluminesent ECL substrate (Pierce- cat#34080).

Antibody	Clone no. / Cat no.	Lot no.	Source	Host	Clonalit y	Stock conc.	IF	Weste- rn	IP
Rel A	Ab1604	LV1447 918	Chemicon	rabbit	polyclon al	1mg/ ml		1/2000	
CD133	Ab19898	447711	Abcam	rabbit	polyclon al	1mg/ ml	1/510	1/500	1877
VIMENTI N	V9/ Ab8069-1	05430	Abcam	mouse	monoclo nal			1/1000	
BRACHY URY	AF2085	KQP010 7011	R&D systems	goat	polyclon al	0.2m g/ml	-	1/1000	-
NANOG	Ab21624	478537	Abcam	rabbit	polyclon al	0.2m g/ml	1/150	1/300	Ħ
LAMIN B	M-20/ sc- 6217	E0208	Santa-Cruz	goat	polyclon al	200µ g/ml	Ē	1/1000	Ħ

## Table 2.5 Primary Antibodies

55

NANOG	NNG-811/ Ab62734	505711	Abcam	mouse	monoclo nal	0.5m g/ml	1/520	<b></b>	E
BRACHY URY	Ab57480	476734	Abcam	mouse	monoclo nal	1mg/ ml	1/250		-
BRACHY URY	C-19/ sc- 717745	K2106	Santa Cruz	Goat	polyclon al	200µ g/ml			
BRACHY URY	N-19/ sc- 17743	G1006	Santa Cruz	mouse	monoclo nal	200µ g/ml			
14-3-3σ	C-18/ sc- 7683	112061	Santa Cruz	Goat	polyclon al	200µ g/ml			
β–catenin	H-102/ sc- 7199	12105	Santa-Cruz	rabbit	polyclon al	200µ g/ml	1/75	1/1000	
E- CADHERI N	36		Transducti on labs.	mouse IgG <sub>2a</sub>		250μ g/ml	1/250	1/2500	-
Ki-67	PP-67		Sigma	mouse IgM		-	1/500	-	-
LAMIN A+C	636/ sc- 7292	H2007	Santa-Cruz	mouse	monoclo nal	200µ g/ml	=	1/750	-
CD166/ Alacam	105902/ MAB6561	DXM01	R & D	Mouse	monoclo nal	500 ug/ ml	-	1/250	-
NANOG	AF1997	KKJ03	R & D	Goat	polyclon al	0.1 mg/ ml	-	1/ 100	-
Oct-4	Ab19857	641990	abcam	rabbit	polyclon al	0.1 mg/ ml	5ug/ ml	lug/ ml	-

## Table 2.6 Secondary Antibodies

Antibody	Conjugate	Stock conc.	Use	Source	
Goat anti-mouse	AlexaFlour 488	2mg/ml	IF (ML) 1/400	Molecular Probes	
Goat anti-mouse	AlexaFluor 568	2mg/ml	IF (ML) 1/400	Molecular Probes	
Goat anti-rabbit	AlexaFluor 488	2mg/ml	IF (ML) 1/200	Molecular Probes	

Goat anti-rabbit	AlexaFlour 568	2mg/ml	IF (ML) 1/200	Molecular Probes
Goat anti-mouse	HRP	400µg/ml	W 1/1000	Santa-Cruz
Goat anti-rabbit	HRP	400µg/ml	W 1/2000	Santa-Cruz
Bovine anti-goat	HRP	200µg/ml		Santa-Cruz
Donkey anti-rabbit	HRP		W 1/25000	Jackson Immunoresearch
Donkey anti-mouse	HRP		W 1/30000	Jackson Immunoresearch

Abbreviations for tables 2.5 and 2.6: "IP": Immunoprecipitation "W": Western "IF": Immunofluorescence "-": not known/not used

"ML": Suitable for multiple labelling (Highly cross-adsorbed) "HRP": Horseradish peroxidase.

#### 2.15 Immunofluorescence

Cells were washed in 1X phosphate buffered saline (PBS) then fixed with 4% (w/v) paraformaldehyde for 20 minutes at room temperature. Cells were then washed in 1X PBS and then permeabilised in 0.2% (v/v) Triton-X-100 for 15 minutes at room temperature. Cells were again washed with 1X PBS followed by blocking in PBS-5% (v/v) FBS for 1 hour at room temperature/ 4°C overnight. Cells were then incubated with primary antibodies (see table 2.5) diluted in 5% (v/v) FBS in PBS for overnight at 4°C. Cells were then rinsed in PBS five times for 5 minutes each and then incubated for 30 minutes at 37°C in secondary antibodies (see table 2.6) diluted in 5% (v/v) FBS in PBS. Following secondary antibody incubation, cells were again washed three times in PBS and stored in PBS until they were imaged using a Zeiss Axioplan 2 confocal microscope. For propidium iodide staining, 1µg/ml of propidium iodide was added to the PBS in the second wash step after secondary antibody incubation.

#### 2.16 Immunoprecipitation

Cells were grown as required, washed with PBS and detached from the flask surface using a cell scraper and collected in 10ml PBS. Cells were then centrifuged at 2500g for 3 minutes and resuspended in 200-500µl lysis buffer (50mM Tris-HCl pH7.4, 200mM sodium chloride, 0.5% (v/v) Triton X-100, 1mM AEBSF (4-(2aminoethyl)-benzenesulfonyl fluoride) and one complete, mini, EDTA-free protease inhibitor cocktail tablet (Roche- cat#11836170001) per 10ml of lysis buffer. Each tablet contained 0.02mg/ml Pancreas-extract, 0.0005mg/ml Thermolysin, 0.002mg/ml Chymotrypsin, 0.02mg/ml Trypsin, 0.33mg/ml Papain). Cells to be lysed with beadbeater were mixed with 200 µl of glass beads and lysed in a beadbeater (FastPrep FP120, Bio101) for 30 seconds on speed 5000 rpm. Whereas, the cells lysed without beadbeater were kept on ice for 30 minutes. All cells were then centrifuged at 15,000g at 4°C for 20 minutes. Supernatants were collected to fresh tubes. Around 15µl of the supernatant were kept separately and 15µl of 2X Laemmli buffer was added to load directly on the gel. Rest of the supernatant was divided into IP (to add the antibody) and IgG controls (negative controls to which normal IgG from the relevant species was added in place of the antibody, details in table 2.7). Both IP and IgG were precleared by adding 25µl raw Protein G linked to agarose beads (Upstate, Cat#16-266, and Lot #0610042813) and incubated for 1hr at 4°C with shaking, after which they were centrifuged for 2 minutes at 13000rpm. The supernatant was collected and the respective antibody (20µg/ml) and IgG were added. The lysates were then incubated at 4°C for 2-3 hours. Again 25µL Protein G linked to agarose beads was then added to the lysates, which were then incubated at 4°C with shaking for 1 hour. Following incubation with the lysate/antibody mix the protein G beads were washed three times in the lysis buffer. The pellets containing the Protein G beads were finally resuspended in 20µl Laemmli buffer. Following boiling for 5-10 minutes at 100°C, samples were loaded onto SDS-PAGE gels.

For western blotting, protocol is described in section 2.14. Bands of interest were cut from the gel using a scalpel and transferred in 1.5 ml eppendorf tubes which were then sent for mass spectrometry.

Tabl	e	2.7	1:	N	ori	nal	Ι	g	Gs
				_				-	_

Normal IgG	ID no.	Lo no.	Source	Stock conc.		
Normal mouse IgG	Sc-2025	B0606	Santa Cruz	200µg/ 0.5ml		
Normal rabbit IgG	Sc-2027	11306	Santa Cruz	200µg/ 0.5ml		
Normal goat IgG	Sc-2028	L0905	Santa Cruz	200µg/ 0.5ml		

## 2.17 RNAi (BRACHYURY, β-catenin, E-CADHERIN, TBX-2, TBX-3, FGF8)

Cells were grown in T75 flasks till confluent. Cells were then split 1in 4 dilution and 0.6ml of the cells were added to 12ml of fresh medium (with 10% (v/v)) FBS). 2ml of this cell and media mixture was added to each well of 6 well plates and/or 0.5ml to each well of 24-well plates. The plates were incubated in 37°C incubator with 5% CO<sub>2</sub>. Meanwhile, the RNAi mixture was prepared by adding 6µl of siRNA which was diluted 1/10 in serum free medium, 94 µl of serum free medium and 6µl of Hiperfect Transfection Reagent (Qiagen, Cat#301705, Lot#130172730) for each sample for the 6-well plates. The RNAi mixture for the 24-well plates was prepared by adding 1.5µl of siRNA which is diluted 1 in 10 in serum free medium, 98.5µl of serum free medium and 3µl of Hiperfect Reagent. Control RNAi mixture was prepared with negative control siRNA. These mixtures were then vortexed, centrifuged and incubated at room temperature for 20-25 minutes, after which the mixtures were slowly added to the plates incubated in the 37°C incubator by mild rotating and mixing. The cells were grown at 37°C for 1 day (subconfluent), 2 days (mid-point between subconfluent and confluent), 3 days (confluent) and 4 days (superconfluent cells). Cells were then fixed, permeabilised
and blocked for Immunofluorescence and/or used for western blots. Details of the siRNA can be found in table 2.8.

#### 2.17A: Mechanism of RNAi:

RNAi (RNA interference) works by introducing a gene- specific dsRNA into a cell that can degrade the homologous mRNA (Cottrell and Doering 2003). It is a post- transcriptional gene silencing mechanisms (Ullu and Tschudi 2000). For our experiments, RNA is made in a test tube by adding phage RNA polymerases that bind to phage promoters that are within expression vectors. The cDNA is transcribed through PCR and the sense and antisense strands are annealed to give double stranded RNA, which is then cut into 21-25 by small interfering RNAs (siRNA) by an enzyme, RNase II. RNase II also has a helicase section, which helps in unwinding of the dsRNA necessary for the binding the RNA to the target RNA that we want to inhibit the translation of (Shuey, McCallus, and Giordano 2002).

siRNA	Name	Cat#	siRNA type	Stock conc.
BRACHYURY	Hs_T_1	SI00738255	Predesigned	5nMol
TBX2	Hs_TBX2_2	SI00740782	Predesigned	5nMol
TBX3	Hs_TBX3_3	SI00083496	Predesigned	5nMol
FGF8	Hs_FGF8_1	SI00145593	Predesigned	5nMol
$\beta$ -catenin	Hs_CTNNB1_5	SI02662478	Predesigned	5nMol
E-CADHERIN	Hs_CDH1_13	SI02654029	Functionally validated	5nMol
Negative control	Alexa Fluor 488	1022563	Control	5nMol

Table 2.8: siRNAs

Note: All siRNA constructs were purchased from Qiagen.

#### 2.18 ChIP (Chromatin Immunoprecipitation)

Approximately  $1 \times 10^6$  cells were grown. The medium was aspirated and washed with PBS. Histones were cross-linked to DNA by adding formalydehyde to a final concentration of 1% (v/v) and incubated at room temperature for 20 minutes. To quench the reaction, Glycine was added directly to the medium to a final concentration of 125mM, shaken a bit and incubated at room temperature for 5 minutes. The medium was then aspirated and the cells were washed 2X with PBS. The cells were then scrapped and collected in a tube and centrifuged for 2000rpm for 4 minutes. The cells were then resuspended in 500µl SDS lysis buffer with protease inhibitors and kept on ice for 10 minutes. Chromatin was sheered by sonicating the cells on ice for 5 pulses of 10 secconds each at 8µm amplitude. To reverse the crosslinking, 8µl of 5M NaCl was added and incubated at 65°C for 4-5 hours. The DNA was recovered by phenol-chloroform extraction and ethanol precipitation. The samples were then run on 1% (w/v) agarose gel to optimize the sonication conditions.

Once the sonication condition has been optimized, the above protocol was repeated till the sonication step and then centrifuged for 10 minutes, 13000 rpm at 4°C. The supernatant was diluted 10 fold in Chip dilution buffer with protease inhibitors. Around 20  $\mu$ l of supernatant was kept separate as the 'Input' for the PCR later. Rest of the supernatant was divided into IP and IgG and was precleared with protein G for 1 hour at 4°C with agitation, followed by a brief centrifugation. Respective antibodies and IgGs were added to the supernatant and incubated rotating for overnight at 4°C.

Next day, protein G agarose was again added to the samples and kept rotating at 4°C for 1 hour. After a gentle centrifugation (700-1000 rpm, 4°C) for about 1 minute, the pellets were washed once with low salt immune complex buffer, once with high salt immune complex buffer, once with LiCl immune complex buffer and twice with TE buffer. The samples were now protein G/ Histone/ DNA complex. For Immunoprecipitaion/ Immunoblot experiment, they were loaded on a gel added with Laemmli buffer and heated for 5-10 minutes at 100°C. For PCR, to amplify the DNA bound to immunoprecipitated histones, the protein G/ Histone/ DNA complex was eluted by adding 250µl NaHCO<sub>3</sub>+SDS elution buffer, vortexed briefly and incubated at room temperature for 15 minutes with rotation. Agarose was the centrifuged and the supernatant was transferred to a separate tube and the elution step was repeated. After combining the eluates of both the elution steps, 5M NaCl was added and heated to 65°C for 4 hour to reverse the crosslinking. The details of the buffers are described in table 2.9.

Buffers	Contents	Conc.	pH
	SDS	1% (w/v)	
SDS lysis buffer	EDTA	10 mM	8.1
~	Tris.	50 mM	
	SDS	0.01% (w/v)	
	Triton X-100	1.1% (v/v)	
ChIP dilution Buffer	EDTA	1.2 mM	8.1
	Tris-HCl	16.7 mM	
	NaCl	167 mM	
	SDS	0.1% (w/v)	
T	Triton X-100	1% (v/v)	
Low sait immune	EDTA	2 mM	8.1
complex wash buller	Tris-HCl	20 mM	
	NaCl	150 mM	
	SDS	0.1% (w/v)	
	Triton X-100	1% (v/v)	
Fign sait immune	EDTA	2 mM	8.1
complex wash buffer	Tris-HCl	20 mM	
	NaCl	500 mM	
	LiCl	0.25 M	
T * 1 *	IGEPAL-CA630	1% (v/v)	
Lici immune complex	Deoxycholic acid	1% (w/v)	8.1
wash buffer	EDTA	1 mM	
	Tris.	10 mM	
TP I. Co	EDTA	1 mM	0.0
I E buffer	Tris-HCl	10 mM	8.0
NaHCO <sub>3</sub> +SDS elution	SDS	1% (w/v)	
buffer	NaHCO <sub>3</sub>	0.1M	

#### Table 2.9 ChIP assay buffers:

#### Table 2.10 Primer sequences for ChIP assay

Primer	Sequence	Length	GC content	Tm	
CH1 F	gtgtgggagcttttgagcatc	21mer	52.38%	66.2°C	
CH2 R	ccatcacgcctggctaattttg	22mer	50.00%	69.4°C	
CH3 F	cctcaagtgatccacgtgcctc	22mer	59.09%	70.4°C	
CH4 R	tgcacgtctgtggtcccagc	20mer	65.00%	72.5°C	

Note: Primer pair 1 was used for site 1 and gave a product length of 193bp and primer pair 2 was used for site 2 and gave a product length of 249bp. APC primers used were APC15a (ttagttttacaccgggggatgata) and APC15 (gcagaagacgacgcagatg) with  $Tm = 50^{\circ}C$ .

#### 2.19 DNA cloning and Luciferase assay

SW480 cells were grown to low density and the genomic DNA was extracted using Wizard SV Genomic DNA Purification System (Cat. Number A2360, Promega). The genomic DNA was stored at -20°C for future use. Primers were designed for the Brachyury binding sites (refer chapter 5 for the sites) in the Nanog promoter region. Details of the primers can be found in section 2.19. PCR was carried out using 2µl of 1/50 dilution of the genomic DNA prepared, 25µl BioMix™ Red (Bioline-Cat#BIO-25005), 1µl of each primer ser diluted 1/10 and 21µl distilled water in a 25µl PCR reaction. The annealing temperature was maintained at 56°C and the extension time was set accordingly (30 seconds per kilo base) and the PCR reaction was set for 40 cycles. PCR products were run on a 1% (w/v) agarose gel and the product bands were cut and purified using Geneclean kit (Cat. Number 1001-200, Q-biogene) and the DNA was eluted in TE buffer. DNA digestion reaction was set up at 37°C for 2 hours using 10µl DNA, 0.5µl KpnI restriction enzyme (Cat. Number R0142S, NEB), SacI restriction enzyme (Cat.Number R0156S, NEB), 2µl NEbuffer1 from 10X stock (Cat. Number B7001S, NEB), 0.2µl BSA from 100X stock (Cat.Number B9001S, NEB) and 6.8µl distilled water in a 20µl DNA digestion reaction. A vector digest was set up alongside 1µl PGL3 basic

vector of 20µg stock of concentration 1µg/µl (E175A, 19205606, Promega), thus using  $0.05\mu g$  vector in a 20 $\mu$ l reaction. Buffer1, BSA and the restriction enzymes were used with same concentration mentioned above and 15.8ul of distilled water is used to make up the volume of the reaction to 20µl. The digestion reactions (both experimental and vector digests) were purified using High Pure PCR Purification kit (Cat. Number 11732676, Roche). [Note: Heat inactivation of the restriction enzymes was not possible because KpnI is not feasible for heat inactivation, although SacI can be heat inactivated]. A small amount of the digested products were run on 1% (w/v) agarose gel to get an estimate of how much vetor: digest ratio is needed for ligation. Ligation mix was made up using 1µl of digested PGL3 basic vector with 1µl T4 DNA ligase (Cat. Number M0202S, NEB), 1.5µl T4 DNA ligase buffer 10X stock (Cat. Number B0202S, NEB). The amount of DNA digest used varied accordingly based on the fact that we used 1:3 ratio of vector: DNA digest, respectively. The reaction was made up to  $15\mu$ l with ultrapure water. The ligation reactions were incubated at 4°C for overnight. Next day, 5µl ligation mix was then transformed to LB-Ampicillin plates with 50µl competent cells (One shot top 10, Invitrogen). [For transformation, the ligation mix and competent cells were incubated on ice for 5-30 minutes, heated at 42°C for 30 seconds to 1 minute, recovered with 250µl LB, incubated at 37°C for 1 hour, shaking and the transformed to LB-Amp plates]. The plates were incubated for overnight at 37°C. Next day, colonies were picked up and were grown in 5ml LB for overnight at 37°C. Miniprep was made from 1ml of the overnight culture using miniprep kit (Cat. Number 27104, Qiagen). Rest of the overnight culture can be used for making glycerol stock (i.e 80% (v/v) sterile glycerol is added to the overnight culture in the ration 1:1 and stored at -80°C). The miniprep was digested again for 1 hour at 37°C using 5µl miniprep DNA with 0.1µl BSA, 1µl NEbuffer1, 0.5µl KpnI, 0.5µl SacI and the reaction was made up to 10µl with ultrapure water. The digested products were then run on 1% (w/v) agarose gel to check whether the insert and vector bands are of the expected size.

Luciferase assay:  $2x10^5$  SW480 cells were seeded and grown at a low density in 96 well plates  $(0.33 \text{ cm}^2 \text{ area})$ . When the cells were ready, the transfection complex was prepared using 5ug of DNA (clone) by diluting it in serum free DMEM to a total volume of 30µl and 0.1µg of Renilla vector, PGL4.7 [hRluc/SV40] (E6911, Promega) as a control vector. [Note: at this step, both experimental and control vectors should be added to the cells. Since, the insert already has PGL3 basic vector, thus only Renilla control vector is added here]. 2.5µl of Superfect Transfection Reagent (Cat. Number 301305, Qiagen) was added to transfect the DNA to the SW480 cells. This complex with DNA, experimental and control vectors and the transfection reagent was incubated at room temperature for 10-15 minutes. Meanwhile, the medium from the 96 well plates were aspirated and washed with 1X PBS. After 10-15 minutes incubation, 1ml of DMEM with serum was added to the transfection complex and this mix was added to the cells and incubated at 37°C incubator for 2-3 hours. After 2-3 hours, the medium was removed from the plates, washed with 1X PBS and then the fresh medium with serum was added to the cells. Dual-Glo Luciferase assay system (E2920, Promega) was used to measure the luciferase activity of the DNA. This was done by adding a volume of Luciferase Reagent equal to the culture medium volume to each well of the 96 well plate. The plates were incubated for 10 minutes in room temperature and then the luminescence was measured for the experimental samples using luminometer (Wallace Victor, 1420 multilabel counter). This gave the result for the luminescence of the clones (experimental). Next, Dual Glo stop and glo reagent was added directly to the plate equal to the original culture volume and incubated at room temperature for 10 minutes and measured the luminescence again. This gave the luminescence for the control (Renilla). This experiment was performed in triplicates for each sample. The data was then normalized using the formula:

Change in fold activity = [Average (F/R) from one sample]/ [Average (F/R) from another sample], where F= Experimental luminescence obtained; R= Control luminescence obtained.

Since, we analysed our values statistically, we plotted our F/R values on a scale rather than plotting the fold increase. The error bars were obtained using the standard deviations and the cut-off for significance was given by the p-values. P-values <0.05 were not considered statistically significant.

[Note: *Renilla* luciferase is a 36 kDa protein obtained from sea pansy (*Renilla reniformis*) which requires coelenterazine and oxygen to produces a blue light of 480nm. This enzyme is used in dual-reporter assays due to their differences in substrate requirements and light output from the experimental vector].

#### 2.20 P-value and t-test:

P-value is statistical hypothesis testing which gives the probability, with a value ranging from zero to one. It tells that what would be the probability that the random sampling will show a difference between sample means as large as observed. P-values can be two-tailed (i.e. what is the probability that the randomly selected samples will give means as far apart as observed with either group having the larger mean); or one-tailed ( i.e. what is the probability that the randomly selected samples will give means as far apart as observed with either group having the larger mean). We used 2-tailed t-test since we were looking for a difference in gene expression. We did not use 1-tailed t-test since it is for a difference only in one direction.

P-value is obtained from Student's t-test which is statistical test to find if the difference between two sample means is statistically significant. To calculate the p-value, from a t-test, MS-Excel was used, the syntax for which is t-test (array\_1, array\_2, tails, type). Array\_1 and \_2 are the data sets; tail value is 1 for one-tailed distribution and 2 for two-tailed distribution. The type will give the type of t-test to be performed.ie. Type of 1 for a paired t-test, type of 2 for a two-sample equal (homoscedastic) variance and type of 3 is for a two-sample unequal (heteroscedastic) variance (Microsoft office website, TTEST). As mentioned before,

P-values ≤0.05 is considered significant. We used type 3 test since, which is independent TTEST where we compared three repeats of the controls genes on a plate to three repeats of experimental gene on the same plate then it is 3 repeats of each and we cannot say that 1 well of one lot is equivalent to 1 well of another. So, we used independent TTEST. [Note: We have used paired t-test where there is definite relationship between the two data sets.]

Table 2.11: Clone details for the luciferase assay

Clone number	Primer combination	BRACHYURY binding sites	Expected product size
33	L2a + LR1	Sites 1 and 2	2271 bp
52	L2C + LR1	Site 1	2312 bp
53	L0F + LR1	No sites	634 bp

Note: The clone numbers are the clone reference numbers (arbitrary). The primers combinations are the name of the primers used (arbitrary) to get the respective clones. The BRACHYURY binding sites are the sites included in the respective clones. The expected product size is the size of the product obtained by using the respective primer pairs.

The primers are as follows:

LR1: attcca gagctc ctggtggtaggaagagtaaagg L0F: attcca ggtac gagacagtctggttctgttgtcacgc L2C: attcca ggtac gtcaggagt tcgagaccag cctc L2a: attcca ggtac gtgtgggagcttttgagcatc

attcca = random base pairs added on each site

#### 2.21 Mitotic index count

Cells were grown in 24 well plates with coverslips. Immunoflouresence protocol was followed (section 2.14) and the coverslips were mounted on a slide and imaged using a Zeiss Axioplan 2 confocal microscope. The number of cells in any stage of mitosis was counted against cells not in mitosis and thus percentage of cells undergoing mitosis was calculated for each sample.

#### 2.22 Cell Viability Asssay (Promega CellTiter-Glo<sup>™</sup>)

SW480 cells were grown in 24 well plates (1.65 cm<sup>2</sup>) for 1,2,3,4 and 5 days (1 being subconfluent and 4 being superconfluent). For each day, the medium was taken out from the plates, washed with PBS and 250µl trypsin was added to detach the cells. 750  $\mu$ l medium was added to this and number of cells per ml was counted by haemocytometer. Cells were now transferred to another 24 well plate with a few different dilutions which were needed for drawing a standard curve later. The plates were incubated at 37°C for 3-4 hrs, giving enough time for cells to start adhering to the base. Later, the medium was taken out from these plates and equal volume of medium and CellTiter-Glo<sup>™</sup> Reagent (Promega, Cat#G7571, Lot#258909) was added to the cells and was incubated at room temperature for 2 minutes. Equal volume of this mixture was transferred to opaque 96 well plates (0.37 cm<sup>2</sup>) in triplicates, mixed on shaker for 2 minutes and incubated at room temperature for 10 minutes. Reading was taken by a luminometer (Wallace Victor, 1420 multilabel counter) which gave luminescence against number of viable cells. [Note: Viable cells are checked by adding Trypan blue (Sigma) stain to the cells. The cells that are viable do not pick up the stain, since they have intact cell membrane].

#### 2.23 Cytoselect cell migration assay

This assay was run using Cytoselect 24-Well cell migration assay (Cell Biolabls, Cat. number CBA-101). Pore size used was  $8\mu m$  (suitable for SW480 cells). Assay plates stored at 4°C was warmed to room temperature for 10 minutes. Cell suspension was prepared in serum free medium (1.2 X 10<sup>6</sup> cells/ ml). 500µl of medium with serum was added to the lower well of the migration plate. 300µl of cell

suspension was added to each insert in the assay wells. 2.5  $\mu$ g/ ml of mitomycin C was added to the cells at this atge. Cells were grown in 37°C incubator. One time point analysed was 12 hours and the 2<sup>nd</sup> time point was 24 hours. After the required time, the plates were taken out of the incubator and medium was taken out from each insert. The inserts were then transferred to clean wells containing 225 $\mu$ l Cell Detachment Solution and incubated for 30 minutes at 37°C. The cells were completely dislodged from the underside of the membrane by tilting the inserts gently. Meanwhile, 4X Lysis Buffer/ CyQuant GR dye solution was prepared by diluting the dye 1:75 in the lysis buffer. 75 $\mu$ l of this buffer/ dye solution was added to each well with 225 $\mu$ l cell detachment solution and incubated for 20 minutes at room temperature. 200  $\mu$ l of this mixture was then transferred to a 96-well plate and the fluorescence was measured at 480 nm/ 520 nm.

#### 2.24 Conditioned media experiment

Cells were grown to a high density (4 days for SW480) at  $37^{\circ}$ C at 5% CO<sub>2</sub>. Medium from the high density cells were collected and then centrifuged to remove the debris. This medium is either snap frozen and stored at -80°C for long term use or is used immediately for growing fresh cells.

#### 2.25 Protein purification

A) Amicon Ultra-15 spin column (Chemicon/ Millipore, Cat. Number UFC9 050
96): This was centrifuged for 30 minutes at 4000g at room temperature. This contains a 10kDa filter that separates components with more than 10kDa size in the upper column and less than 10kDa in the lower column.

B) HiTrap Blue affinity column (GE healthcare, Cat. Number 17-0413-01): The dye ligand in this column is Cibacron Blue F3G-A which is covalently attached to the matrix by the triazine part of the dye molecule. Few proteins may interact biospecifically with the dye due to its structural similarity with nucleotide cofactors.

In this column, albumin and interferon can bind in a less specific manner by electrostatic and/or hydrophobic interactions with the aromatic anionic ligand. C) MonoS cationic exchange column (GE healthcare): This column was used at pH 6 and is negatively charged. Thus, it would bind the positively charged components. D) MonoQ anionic exchange column (GE healthcare): This column was used at pH 8 and is positively charged. Thus, it would bind the negatively charged components. Note: At the end of final run using any of the above mentioned columns, the columns are washed with elution buffer, binding buffer, water and 20% (v/v) ethanol of 20-25 ml each, in the order mentioned. [The details of binding and elution buffers are mentioned in chapter 7].

# Chapter 3 –a) The mesoderm inducer BRACHYURY is differentially expressed in colorectal cancer cells

#### 3.1 Introduction:

Cancers may arise from disruption of normal cell proliferation, differentiation and survival. The initiation and maintenance of cancer often involves the aberrant expression of genes that are associated with normal development. As mentioned previously (section 1.5), the process of EMT (or similar), a fundamental feature of normal development is also inappropriately involved in the progression of CRC. As such, the invasive fronts of colorectal tumours are characterized by epithelial de-differentiation, loss of intercellular adhesion and an increase in cell migration. This is associated with the activation of transcription factors related to epithelial-mesenchymal transition (EMT) (Brabletz *et al.*, 2005).

It is not easy to study the events involved in EMT during the progression of cancer and an *in vitro* system that re-capitulates this process would be invaluable to the study of these transition events. We used CRC cells from the cell line SW480, grown to low and high density, as a model for cells reversibly transitioning between the mesenchymal-like and epithelial-like cell states, respectively (Brabletz *et al.*, 2001; Davies *et al.*, 2004)

Our initial aim was to carry out a directed RT-PCR screen on a range of mesoderm inducing genes that may be candidates for inducing a transition from epithelial to mesenchymal cell types in high and low density SW480 cells. Specifically, we wanted to look for any differential expression of these genes in low density cells (mesenchymal-like) compared to high density cells (epithelial-like) SW480 cells. The following candidate genes were studied to determine if they are differentially expressed in low and high cell density SW480 cells. *E-CADHERIN, ACTIVIN A, ACTIVIN B, FGFR1, FGFR2, FGFR3, SNAIL, NFkB, BRACHYURY, NOTCH 1, NOTCH 2, NOTCH 3, NOTCH 4, TBX2, TBX3, FGF-4, FGF-19, BMP4, FGF-8, GDF3, NODAL, Oct-4* (refer to table 3.1).Alongside this study in SW480 cells (which have single copy of *APC* with a mutation at codon 1338 resulting in a stop codon and therefore a truncated protein of 1337 amino acids, Nishisho *et al.,* 1991), we also determined the expression of some of these mesoderm inducers in other human colorectal cancer cell lines including

Table 3.1:Summary of function	of the	developmental	genes	in early	development	we
used in our screen.						

Gene	Role	Reference
<i>E</i> -	E-cadherin is a tumor suppressor gene involved	Semb et. al., 1998;
CADHERIN	in calcium dependent cell-cell adhesion. It is	Wong et al. 2003
	downregulated in cells undergoing EMT.	
Activin	Activin is involved in early mesoderm	Dyson et al., 1997
	formation in Xenopus.	Chen et al., 2006;
	Involved in cell proliferation, cellular	Sulyok et al., 2004
	differentiation, apoptosis, metabolism,	
	homeostasis, immune response, wound repair	
	and endocrine function.	
FGFR1	Expressed during vey early neural development.	Walshe and Mason
(Fibroblast	Involved in limb induction, mammalian liver	2000;Jung et al. 1999;
growth	development, inner ear development in mouse.	Pirvola <i>et al.</i> 2002
receptor 1)		
FGFR2	Expressed in early neural development.	Walshe and Mason 2000
(Fibroblast	Involved in retinal development.	Moore et al. 2004
growth		
receptor 2)		
FGFR3	Expressed in early neural development.	Walshe and Mason 2000
(Fibroblast	Inhibits cell proliferation by interfering with	Riley et al., 2007; Antoine
growth	ribosomal biogenesis. Roles in nonsyndromic	<i>et al.</i> , 2005
factor	cleft lip and palate	
receptor 3)		
SNAIL	Induces EMT during development of mouse.	Ethan et al., 2001
(SNAII)	Downregulates the ectodermal gene expression	Twigg et al., 1999
	in the mesoderm	
ΝΓκΒ	Involved in B cell development during	Shaffer et al. 1997
	organogenesis.	Gilmore et al., 2006;
	Role in regulating the immune response to	Albensi et al., 2000
	infection.	
	Involved in synaptic plasticity and memory.	
BRACHYU	It is a transcription factor. In mouse, it is needed	Chesley 1935;
RY	for mesoderm formation and anteroposterior	Gluecksohn-
	axis extension	Schoenheimer 1944;
	in the mammalian embryo.	Spiegelman 1976
	Involved in formation and differentiation of	Naiche et al., 2005
	posterior mesoderm and axial development in	

	vertebrates.	
NOTCH	Notch family includes 4 receptors (Notch 1-4).	Tanigaki <i>et al.</i> 2001;
family	Notch is important for cell-cell interaction	Gaiano et al., 2002;
	involved in gene regulation which controls	Uyttendaele et al., 2001
	cellular differentiation during embryogenesis.	
TBX2,	In development of the atrioventricular canal.	Ribeiro et al., 2007
TBX3	Involved in cell proliferation during heart	
	development.	
FGF-4	Function in bone morphogenesis and limb	Powers et al., 2000
	development and secreted during early	
	mesoderm induction	
FGF-19	Initiates inner ear during early development and	Nishimura et al., 1999
	involved in embryonic development, cell	
	growth and morphogenesis.	
FGF-8	Secreted in midbrain and limb development,	Powers et al., 2000
	organogenesis, left-right axis determination,	
	embryo gastrulation	
NODAL	Secreted during mesoderm formation and axis	Gebbia et al., 1997
	specification during development.	
BMP4	Mediates dorsal/ ventral patterning during	Friedle et al., 1998
(Bone	Xenopus early embryogenesis, ventralises	Vainio et al., 1993
morphogen	mesoderm, involved in EMT during tooth	Hee et al., 2009
etic protein	development.	
4)		
GDF3	Regulates cell fate of the embryo during early	Ariel et al., 2006
(Growth	development and is present throughout early	
differentiati	embryogenesis in mouse.	Chen et al., 2006;
on factor-3)	Active during during embryonic development	Hexige <i>et al.</i> , 2005
	and also in adults during the development of	
	the brain, thymus, spleen, bone marrow and	
	adipose tissue	
Oct-4	Essential for early embryo development and is	Rosner et al., 1990.
	present during the the preimplantation period.	
	Critical for self-renewal of undifferentiated	Niwa <i>et al.</i> , 2000
	embryonic stem cells	

**Table 3.1:** This table lists the names and known developmental functions of the genes we screened to check for their differential expression in low and high density SW480 cells.

HCT116 cells (which has functional *APC*, but a mutation leading to constitutively active  $\beta$ -catenin, Kitaeya et al., 1997), LoVo (double *APC* mutation at 1,114 amino acid and 1,430 amino acid, Rowan et al., 2000), HT29 (double *APC* mutation at 853 amino acid and 1555 amino acid positions, Brocardo et al., 2005), T84 (APC mutation at 1,488 position, Rowan et al., 2000) and the germ cell tumour EC cell line NTERA-2.

#### 3.2 Validation of SW480 cell line for use as a model to study EMT:

SW480 cells grown to low and high density were previously shown to exhibit features of cells undergoing EMT (Brabletz *et al.*, 2001; Davies *et al.*, 2004). E-CADHERIN (both expression levels and co-localisation) and  $\beta$ -catenin (localisation) are key hallmarks (reviewed in Thiery 2002) of epithelial-tomesenchymal transition and have been shown to change in SW480 cells with cell density. In summary, in low density SW480 cells (mesenchymal-like), the cells have low levels of E-CADHERIN located in cytoplasm and nuclear  $\beta$ -catenin. In high density SW480 cells (epithelial-like), the cells exhibit higher levels of Ecadherin, mainly membrane associated, and predominantly membrane bound  $\beta$ catenin.

In accordance with previous literature, we were able to provide further evidence that SW480 cells grown at different densities might indeed be a good *in vitro* model to study the molecular events involved in EMT. Nuclear NF $\kappa$ B has previously been shown to be associated with induction and maintenance of EMT (Huber *et al.*, 2004). Figure 3.1 shows that one of the NF $\kappa$ B family members, REL A (Nuclear factor, NF $\kappa$ B p65 subunit), which is the most abundant form of NF $\kappa$ B, changes localisation in SW480 indicative of cells undergoing EMT. As the SW480 cells progress from low to high density, REL A translocates from the nucleus to the cytoplasm, respectively. The change in localization of REL A is in accordance with the changes in localization of  $\beta$ -catenin and E-CADHERIN and further validates the use of SW480 cells grown to different densities as a system to study EMT.



Confocal immunofluoresence microscopy showing localization of REL A in SW480 sub and superconfluent cells. A) Low density SW480 cells were stained with REL A primary antibody (green) which localizes to the nucleus of the low density SW480 cells. Nuclei are marked by staining with Propidium iodide (Pi) (red). B) High density cells where REL A (green) is located to the cytoplasm and the Pi (red) to the nucleus. The DIC image (in both A and B) shows the cells without any stain and the merged image shows the REL A and Pi staining together, giving a relative co-localisation. This translocation is indicative of EMT. [Note: The images were taken under 63X objective.]

#### 3.3. Screen of the mesoderm inducing genes by RT-PCR:

Based on the observations above, we set up a small scale and directed screen to determine if a selected group of predominantly mesoderm inducing genes, that are key inducers during development might be differentially expressed in low and high density SW480 cells. Those genes identified from this screen would then be studied to determine if they play a role in inducing EMT in SW480 cells. This screen was also extended to other cell lines to see if they behaved similarly.

The cells from the different cell lines were grown to low and high densities (as described in materials and methods section 2.2) and the cDNA synthesised (as described in section 2.9) Primers were designed for all the genes (refer to table 2.1) and PCR was carried out (as described in section 2.10). Table 3.2 summarizes the RT-PCR results illustrating the expression levels of various genes in the different CRC cell lines. *Brachyury* was found to be differentially expressed in SW480 cells, but was not expressed in other CRC cell lines studied. As expected, *E-CADHERIN* levels increase in SW480 high density cells. Other candidates were also differentially expressed in SW480 cells (*TBX2*, *TBX3*, *FGF-8*, *FGFR1*, 2 and 3), but the expression patterns were not consistent between cell lines.

# 3.4 qRT-PCR quantification:

In order to gain more detailed quantitative expression for these genes we also carried out qRT-PCR based on our RT-PCR screen results. For this, we followed the method described in materials and methods section 2.11 using Sybr green and analysed the data using  $2^{-\delta\delta C}_{T}$  method, as mentioned in section 2.13 of materials and methods. A worked example of how the raw data is converted into relative quantities is shown in section 2.13 of materials and methods. Section 2.20 of materials and methods describes how the p-values were calculated to check the significance of the data obtained.

Gene	SW480 sub	SW480 super	HCT 116 Sub	HCT 116 super	LoVo sub	LoVo super	HT29 sub	HT29 super	T84 Sub	T84 super
	1	1								
BRACHYURY	+++	+	-	-	-	8	-	-	-	-
E- CADHERIN	++	+++	+++	+++	++	++	+++	+++	+++	+++
ACTIVIN A	-	-	-	-	-	-	-	-	1	1
ACTIVIN B	-	-	2=	-	₩P.			-	1	1
FGFR1	++	+++	+++	+++	+	++	-	-	1	1
FGFR2	+	H)	++	+++	+	-	+	10 <b>4</b>	1	/
FGFR3	++	+	+	+	-	-		19	1	1
SNAIL	+	+	+++	+++	÷	-	-	-	1	1
NFĸB /	+++	+++	+++	++	-	-	=	-	1	1
NOTCH 1			+++	+++	-	+	+++	4	1	/
NOTCH 2	-	-		-	- #	-		-	1	1
<i>NOTCH 3</i>	8	=	-	-	-			-	1	7
NOTCH 4	-	-	-	-		-	- <b>-</b> -	-	1	1
TBX2	+++	++	-	+	-	-	<b>.</b>		1	1
TBX3	+++	+++	+++	++	3 <b>-</b>	-	+	-	1	1
FGF-4	-	2 <b>4</b> -	-	->	8 <b>9</b>		-		1	1
FGF19	+++	+++	+++	+++	<u>.</u>	×	-	-	1	1
BMP4	+++	+++	+++	+++			÷		1	/
FGF8	++	+	+++	+++	1572	-	-	( <b>.</b> .	1	1
GDF3	+	-	-	-	-	-	-		1	1
NODAL	+	-	+++	++	-	-	8		1	/
Oct4	+	-	1	Z	1	1	/	1	1	/
$\beta$ -catenin	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++

Abbreviations:

/ = not checked, - = no expression, += low expression, ++ = moderate expression, +++

= high expression, Sub = subconfluent cells, Super = superconfluent cells.

### **Table 3.2:**

Summary of the expression of genes in low and high density CRC cell lines as assessed by RT-PCR. The experiment was performed 5 times (N=5).  $\beta$ -catenin was used as a control since the level of  $\beta$ -catenin is high and similar in all the cell lines used. This table demonstrates the presence and absence of the genes found by running them on 1% agarose gel.

In general, the data for SW480 cells show gene expression patterns that resemble cells undergoing EMT, for example, high density SW480 cells have high E-CADHERIN levels, low SNAIL and ZEB1 expression relative to low density cells (Table 3.2). Table 3.4 is obtained by comparing low-density data to '1' and comparing high density data to '1'. None of the other cell lines follow such a distinct pattern nor do they display changes in localisation of  $\beta$ -catenin and E-CADHERIN (Brabletz *et al.*, 2001; Davies *et al.*, 2004). These observations further agree with SW480 cells being a good model in which to study EMT, when grown to different densities.

Note: Data for qRT-PCR was compared to either to *LAMINC* or  $\beta$ -catenin. Section 2.12 of material and methods and Figure 3.2, 3.3 shows that both *LAMINC* and  $\beta$ -catenin are proper controls for analysing the qRT-PCR data since their expression levels do not change with cell densities.

From table 3.2, we observe that BRACHYURY is differentially expressed in low and high density SW480 cells (low in high density cells and high in low density cells). Brachyury is the founding member of the T-box transcription factor family of genes and is involved in early mesoderm development (Wilson and Conlon, 1990). It has a conserved DNA-binding domain (Papaioannou et al., 1998). Apart from its role in cellular differentiation and mesoderm formation (Willison, 1990), Brachyury has recently been identified as a possible target for T-cell mediated cancer immunotherapy and was associated with a number of tumour cell types, including colorectal tumour cell lines (Palena et al., 2007). This complies with our finding of Brachyury being expressed in SW480 cells. Brachyury has also been shown to be expressed in chordomas and in the embryonal carcinoma (EC) cell line NTERA2, where it was found to be expressed in the absence of mesodermal differentiation (Vujovic et al., 2006; Gokhale et al., 2000). However, to date no role of Brachyury in cancer has been described. Results from RT-PCR and qRT-PCR analysis in SW480 cells, showed that BRACHYURY was differentially expressed in SW480 cells. The gRT-PCR data we obtained for SW480 cells indicates that BRACHYURY expression is decreased by 56% as the cells progress from low to high cell density (Figure 3.4). As mentioned earlier, we also analysed the expression of the panel of genes by qRT-PCR in other colorectal cancer cell lines; the results are summarised in Figures 3.5 (HCT116), Figure 3.6 (LoVo) and Figure 3.7 (HT29).



Figure 3.2: *LAMIN C* and  $\beta$ -catenin expression levels in SW480, HCT116, LoVo, T84, HT29 and SW626 cell lines. A)  $\beta$ -catenin expression levels are similar in all the cell lines mentioned and also in low and high density of each cell line mentioned above. The graph is obtained from Table 1A of Appendix. B) Expression levels of *LAMIN C* are also similar in all cell lines used and also in low and high density cells in each line. The graph is obtained from Table 1B of Appendix.

The error bars are obtained from the standard deviations. Each experiment was done in triplicate. The X-axis plots the different cell lines used and the Y-axis plots the average Ct values obtained from Tables 1A and B of Appendix.



Figure 3.3: *LAMIN C* and  $\beta$ -catenin expression levels on *BRACHYURY* RNAi in SW480 low density cells. A) *LAMIN C* expression levels are similar in SW480 low density cells with no RNAi treatment, cells treated with negative siRNA and on knocking down *BRACHYURY*. The graph is obtained from Tables 3.3A. B) Expression levels of  $\beta$ -catenin are also similar in SW480 low density cells with no RNAi treatment, cells treated with negative siRNA and on knocking down *BRACHYURY*. The graph is obtained from Tables 3.3A. B) Expression levels of  $\beta$ -catenin are also similar in SW480 low density cells with no RNAi treatment, cells treated with negative siRNA and on knocking down *BRACHYURY*. The graph is obtained from Tables 3.3 B. The error bars are obtained from the standard deviations. Each experiment was done in triplicate. The X-axis plots the different RNAi treatment and the Y-axis plots the average Ct values obtained from Tables 3.3 A and B. The error bars were obtained from the final standard deviations. [Note: Final SD is obtained by the formula [{(X-Y)^2+(X-Z)^2)/2}^0.5], where X is the average of two standard deviations; Y and Z are the two standards deviations]. Each experiment was done in triplicate.

Table 3.3 A

	LAMIN C	Mean Ct	S.D.
Untreated	24.437		
	24.395	24.5323	0.20
	24.765		
Non-interfering	24.53		
	24.74	24.5683	0.15
	24.435		
BRACHYURY RNAi	24.37		
	24.475	24.465	0.09
	24.55		

#### Table: 3.3 B

	β- catenin	Mean Ct	S.D.
Untreated	21.356		
	21.2861	21.5293	0.36
	21.946		
Non-interfering	21.267		
	21.7304	21.5108	0.23
	21.535		
BRACHYURY RNAi	21.3435		
	21.768	21.5168	0.22
	21.439		

#### Table 3.3:

**qRT-PCR data to show that both** *LAMIN C* and  $\beta$ -catenin were used as good controls. A) The table indicates that expression levels of *LAMIN C* does not change in SW480 cells treated with *BRACHYURY* siRNA compared to untreated cells, therefore a good control gene. B) The table indicates that expression levels of *LAMIN C* does not change in SW480 cells treated with  $\beta$ -catenin siRNA compared to untreated cells, therefore a good control gene.



qRT-PCR data obtained for genes in SW480 cells. A) Table illustrating the qRT-PCR data for selected mesoderm inducing genes in SW480 cells grown to low and high densities. Data is represented in terms of fold increase or fold reduction compared to sub-confluent cells (taken as 1). The data was compared to LAMINC and experiment was repeated 3 times (N=3), each time in triplicate. The table shows that *TBX2*, *TBX3*, *BRACHYURY*, *E-CADHERIN*, *FGF8*, *NANOG*, *ZEB1*, *SNAIL* and *CDX2* are differentially expressed in low and high density SW480 cells. B) Graphical representation of fold changes in qRT-PCR values in high density SW480 cells compared to low density cells. The error bars are obtained from the standard deviation from Figure 3.2A. The raw data with along with the data working is shown in appendix Table 2. Refer appendix Table 26 for the statistical significance data of the fold differences obtained for each gene in low and high density cells.



qRT-PCR data obtained for genes in HCT116 cells. A) Table illustrating the qRT-PCR data for selected mesoderm inducing genes in HCT116 cells grown to low and high densities. Data is represented in terms of fold increase or fold reduction compared to sub-confluent cells (taken as 1). The data was compared to LAMINC and experiment was repeated 3 times (N=3), each time in triplicate. *BRCAHYURY* was not examined in this screen since it was found to be absent in HCT116 cells. Genes such as *FGFR1*, *SNAIL*, *FGF8*, *E-CADHERIN*, *ZEB1*, *TBX2* and *TBX3* are found to be differentially expressed in low and high density HCT116 cells. B) Graphical representation of fold changes in qRT-PCR values in high density HCT116 cells compared to low density cells. The error bars are obtained from the standard deviation from Figure 3.3A. The raw data with along with the data working is shown in appendix table 3. Refer appendix Table 26 for the statistical significance data of the fold differences obtained for each gene in low and high density cells.



qRT-PCR data obtained for genes in LoVo cells. A) Table illustrating the qRT-PCR data for selected mesoderm inducing genes in LoVo cells grown to low and high densities. Data is represented in terms of fold increase or fold reduction compared to sub-confluent cells (taken as 1). The data was compared to LAMINC and experiment was repeated 3 times (N=3), each time in triplicate. Genes such as *FGFR1*, *FGFR3*, *E-CADHERIN* and *TBX2* are found to be differentially expressed in low and high density LoVo cells. B) Graphical representation of fold changes in qRT-PCR values in high density LoVo cells compared to low density cells. The error bars are obtained from the standard deviation from Figure 3.4A. The raw data with along with the data working is shown in appendix table 4. Refer appendix Table 26 for the statistical significance data of the fold differences obtained for each gene in low and high density cells.



qRT-PCR data obtained for genes in HT29 cells. A) Table illustrating the qRT-PCR data for selected mesoderm inducing genes in HT29 cells grown to low and high densities. Data is represented in terms of fold increase or fold reduction compared to sub-confluent cells (taken as 1). The data was compared to LAMINC and experiment was repeated 3 times (N=3), each time in triplicate. Genes such as *FGFR1*, *FGFR2*, *E-CADHERIN*, *ZEB1* and *TBX2* are found to be differentially expressed in low and high density HT29 cells. B) Graphical representation of fold changes in qRT-PCR values in high density HT29 cells compared to low density cells. The error bars are obtained from the standard deviation from Figure 3.5A. The raw data with along with the data working is shown in appendix table 5. Refer appendix Table 26 for the statistical significance data of the fold differences obtained for each gene in low and high density cells.

# b) Localization of BRACHYURY in SW480 cells

We focused our further studies on *BBRACHYURY* and determining its possible role in EMT in CRC cells. This was based on the following: 1) *Brachyury* had not previously been shown to be expressed in CRCs, its expression had been shown in chordomas and germ cell tumours. However, during this study, another group, Palena and co-workers (2007) showed that *Brachyury* was expressed in some of the CRC cell lines. The study by Palena and co-workers therefore further validated our screen. 2) *Brachyury* is a key mesoderm inducing gene during development and might have a possible role in inducing EMT during cancer. 3) From our RT-PCR data, we observed that SW480 cells were the only CRC cell line that expressed *Brachyury*. Also, SW480 cells were the only cell line we studied that shows features of cells undergoing EMT with respect to its cell density variation.

Knowledge of the localization of proteins in a cell is often key to our understanding of its functions. Futhermore, co-localization patterns allow us to determine the likelihood of one protein interacting with another. To understand the role of *Brachyury* in inducing EMT, we used indirect immunofluorescence (IF) in SW480 cells grown to different densities (see section 2.15 for IF protocol). Figure 3.8 shows the localization of BRACHYURY in these cells as determined by confocal microscopy. Since, BRACHYURY is a transcription factor, it is expected to be localised to the nucleus of the cells. Figure 3.6 shows that BRACHYURY is located in the nucleus of low density SW480 cells. However, in high density SW480 cells, very few cells are positive for BRACHYURY and those cells that are positive also have nuclear localization. These immunofluorescence results correlate with the qRT-PCR data showing high expression in low density cells but low expression of *BRACHYURY* in high density SW480 cells.



Confocal microscopy image of BRACHYURY localization in low and high density SW480 cells. A) BRACHYURY is located to the nucleus (green) of almost all cells in low density SW480 cells, confirming its nuclear localization and therefore, a transcription factor. B) In high density SW480 cells, very few cells positive for BRACHYURY (green). The DIC image shows the cells without any staining. This shows the differential expression of BRACHYURY in low and high density SW480 cells.

#### 3.5 Discussion:

A common theme has emerged over the last few years concerning genes that give rise to cancer, that is, many of these genes involved in initiation or maintenance of cancer are actually developmental genes that become misexpressed. One of key developmental processes involved in normal embryogenesis is EMT. But, it has also been shown that EMT is also active with the similar phenomenon in many forms of cancer, colorectal cancer being one of them. Our study and this chapter in particular, aimed to identify genes, from a limited group of genes that are involved in embryogenesis that might be involved in regulating the process of EMT in CRC. Changes in the cell density of the CRC cell line SW480 resemble the cells transitioning between mesenchymal-like and epithelial-like states. Our data show that mesoderm inducer, BRACHYURY, is differentially expressed in low and high density SW480 cells. BRACHYURY is highly expressed in the low density SW480 cells which are mesenchymal-like and representative of cells that are found at the invasive fronts of a tumour. Conversely, expression of BRACHYURY is very low in the high density SW480 cells which are epithelial-like and representative of cells at the central mass of a tumour. We found a number of other developmental genes which were also differentially expressed in SW480 and other CRC cell lines. We focused our studies on BRACHYURY because it plays a critical role in embryogenesis and it is not been extensively studied in cancer. Recent studies also showed (Palena, 2007) that *Brachyury* is present in SW480 cells, further validating our results. Human BRACHYURY (T) is located on human chromosome 6q27. This also makes it important to study because chromosome 6 is associated with colorectal cancers (Ezaki 2003). Presumably this should be mutation of or deletion of chromosome 6. Also, since chromosome 6 represents 5-6% of the DNA of cells, it is a possible site for many significant genetic alterations.

IF studies showed that BRACHYURY is present in the nucleus of all low density SW480 cells, but is present at low frequency in high density SW480 cells. We tested 4 CRC cell lines in total, but only one, SW480, expressed *BRACHYURY* leading us to suggest that expression of BRACHYURY is

89

associated with the ability of the cells to undergo EMT. Furthermore, SW480 cells showed a gene expression profile that was representative of cells undergoing EMT (in contrast to the other cell lines studied). The following chapters will focus on the possible role of *BRACHYURY* in colorectal cancer.

Further studies related to this screen, might involve work on the candidates *TBX2* and *TBX3* genes. The expression of transcription factor *TBX2* is amplified in pancreatic (Chen *et al.*, 2008) and breast cancer (Sinclair *et al.*, 2002). Its related factor TBX3 is also over-expressed in breast cancer. TBX2 is known to maintain proliferation in melanomas (Vance *et al.*, 2005). TBX3 is found to be a downstream target of the Wnt pathway and also shown to be expressed in the human CRC cell lines SW480 and HCT116 (Renard *et al.*, 2007). It is also involved in maintaining proliferation in liver cancers (Suzuki *et al.*, 2008).

*CDX2* (caudal related homeobox-2) could be another interesting member to be studied in SW480 cells. *CDX2* is often mutated in rare cases of CRC and it is also a marker of CRC. It is critical for maintenance of intestinal epithelial cells (Duluc *et al.*, 1997; Lorentz *et al.*, 1997). It also has tumourigenic potential in CRC cell lines SW480 and LoVo (Dang *et al.*, 2006).

As mentioned earlier, we were searching for developmental genes those are differentially expressed in low and high densities of SW480 and other colon cancer cell lines. Therefore, differential expression of such genes (mentioned in this chapter) would be interesting to study further which might be involved in EMT and thus in progression of cancer.

# Chapter 4 - Determining the role of *BRACHYURY* in SW480 cells

- a) Does *BRACHYURY* play a role in inducing EMT in the colorectal cancer cell line SW480?
- b) Role of *BRACHYURY* in regulating *NANOG* and maintaining a putative cancer stem cell population
- c) A role for *BRACHYURY* in regulating the NF $\kappa$ B  $\swarrow$  pathway?
- d) A role for *BRACHYURY* in cell proliferation?
- e) A role for BRACHYURY in cell migration?

# a) Does *BRACHYURY* play a role in inducing EMT in the colorectal cancer cell line SW480?

In Chapter 3, we have identified the differential expression of a key mesoderm inducing gene, *BRACHYURY*, from a screen of expressed developmental genes in the CRC cell line SW480. Specifically *BRACHYURY* was found to be relatively highly expressed in the low density mesenchymal-like cells (resembling the invasive fronts of a tumour) whereas its expression was very low in high density epithelial-like cells (resembling the central mass of a tumour). In this chapter, we further studied *BRACHYURY* to elucidate its function in SW480 cells.

#### 4.1.1 A role for BRACHYURY in inducing EMT?

EMT is an important process for growth of a tumour and metastasis (Boyer *et al.*, 2000). During this phenomenon, the cells undergo a transition from the epithelial phenotype to a highly motile mesenchymal phenotype (Huber *et al.*, 2005). E-CADHERIN is a member of the classic cadherin family and is a key component of the cell-junction adhesion complex which is critically altered during EMT (Tomita *et al.*, 2000, Wang *et al.*, 2006). Most inducers of EMT found to date are transcription factors that repress the cell-junction adhesion complex, E-CADHERIN, for example, *Snail* (Bolos *et al.*, 2003), *Slug* (Barrallo *et al.*, 2003) and *Twist* (Karreth *et al.*, 2004). A reduced level of E-CADHERIN also has been seen in various cancer metastases (Huber *et al.*, 2005).

Loss of E-CADHERIN expression is a hallmark of EMT along with the translocation and accumulation of  $\beta$ -catenin in the nucleus of cells (Hirohashi *et al.*, 2003).  $\beta$ -catenin is a central component of the Wnt signalling pathway and is also a subunit of the cadherin–adhesion complex.  $\beta$ -catenin has shown to be involved in EMT in normal development (such as in development of the heart)

(Liebner *et al.*, 2004) and also in the development of many cancers such as prostate cancer (Jiang *et al.*, 2007).

Based on our qRT-PCR data (Chapter 3), we predicted that *BRACHYURY* might play a crucial role in the process of EMT in SW480 cells. Therefore, we investigated whether *BRACHYURY* might be a direct inducer of EMT. For this, we investigated the following features of SW480 cells, following RNAi induced knock-down of *BRACHYURY*.

- i) Localisation of  $\beta$ -catenin and E-CADHERIN
- ii) mRNA levels of E-CADHERIN

# i) Localisation of β-catenin and E-CADHERIN in response to *BRACHYURY* knock-down

We performed a transient knock-down of *BRACHYURY* in SW480 cells (protocol in materials and methods section 2.17), followed by immunofluorescence (according to the protocol in Materials and Methods section 2.15) and the results were viewed by confocal microscopy. RNAi sequence for *BRACHYURY* knock-down is described in Table 2.10 of materials and methods.

If BRACHYURY is a direct inducer of EMT we might expect that knocking down *BRACHYURY* expression would affect the two key indicators of EMT, that is  $\beta$ -catenin localisation (a nuclear to cytoplasmic shift would be expected following RNAi in low density SW480 cells), and localisation of E-CADHERIN would be expected to change (E-CADHERIN re-localisation to membrane following *BRACHYURY* RNAi). The levels of *BRACHYURY* mRNA and protein were low following *BRACHYURY* RNAi (62% knock-down) as determined by qRT-PCR and Western blot analysis, respectively (Figure 4.1), confirming the success of the knock-down. [**Note**: Figure 4.1A is derived from Table 6 of appendix]. We found no change in  $\beta$ -catenin localisation from cytoplasm to the nucleus of the SW480 cells, nor in the localization and levels of *E*-CADHERIN following *BRACHYURY* knock-down; suggesting that *BRACHYURY* might not be a direct inducer of EMT in SW480 cells (Figure 4.2).



### Figure 4.1:

*BRACHYURY* RNAi in SW480 cells. A) Graph showing knock-down of *BRACHYURY* by qRT-PCR. Knock-down level was 62% (calculated by qRT-PCR). Raw data is detailed in appendix table 6. The X-axis gives the different RNAi conditions and the Y-axis gives the relative expression levels. B) Western blot analysis showing the reduction of BRACHYURY protein confirming the knock-down. U= cells not treated with siRNA; N=cells treated with non-interfering siRNA; B=cells treated with BRACHYURY RNAi.



# Figure 4.2:

Confocal microscopy showing  $\beta$ -catenin and E-CADHERIN localisation following *BRACHYURY* knock-down. A) Localistation of  $\beta$ -catenin (green) is mostly at the cell membranes following *BRACHYURY* knock-down in superconfluent SW480 cells. B) Membrane localisation of  $\beta$ -catenin in control SW480 cells that have not been treated with RNAi BRACHYURY. C) In low density SW480 cells, E-CADHERIN levels (green) are low following *BRACHYURY* knock-down and is localised to the cell junctions. (D) E-CADHERIN localised to the cell junctions in low levels in sub-confluent SW480 cells not treated with *BRACHYURY* siRNA. Therefore, this figure shows no change in sub-cellular localisation of E-CADHERIN and  $\beta$ -catenin in response to *BRACHYURY* RNAi, confirming that *BRACHYURY* is not directly involved in EMT. Note: Levels of *BRACHYURY* knock-down was determined by qRT-PCR to be 69%. The DIC images show the cells without any staining.
#### ii) Effect of *BRACHYURY* knock-down on expression of *E-CADHERIN* by <u>qRT-PCR in SW480 sub-confluent cells</u>

In the section above, we showed that BRACHYURY does not have an effect on two important hallmarks of EMT (B-catenin and E-CADHERIN localisation). We next determined a further indicator of EMT, the expression levels of E-CADHERIN following BRACHYURY RNAi, as assessed by qRT-PCR in SW480 sub-confluent cells. We show that BRACHYURY knock-down does not alter the expression of E-CADHERIN when compared to control cells treated with non-interfering siRNA and also untreated cells (Figure 4.3). LAMIN C was used as a control to compare the data (refer section 2.13 of materials and methods for its validation). Final data for Figure 4.3A is obtained in a similar manner as mentioned in material and methods section 2.13. Thus, we concluded that the levels of E-CADHERIN gene expression do not change effectively following BRACHYURY knock-down. [Note: BRACHYURY knock-down level is 64%]. Together, these results suggest BRACHYURY does not have a direct effect on inducing EMT in SW480 CRC cells. We, therefore explored the possible effects of BRACHYURY on four different categories of genes in low density SW480 cells, as described in the following sections.

## 4.1.2: The qRT-PCR screen to study the possible effects of *BRACHYURY* on four different categories of genes:

Data so far suggests that *BRACHYURY* may not be a direct inducer of EMT since knock-down of *BRACHYURY* does not have a direct effect on either E-CADHERIN or  $\beta$ -catenin expression and localisation. Therefore, as a next step to elucidate the possible function of *BRACHYURY* in SW480 cells, we studied the effect of *BRACHYURY* on the expression of following four categories of genes:

- Markers of epithelial cell polarity
- *CDX2* (marker of intestinal epithelium)
- Known gene targets of *BRACHYURY*
- Developmental genes which are inducers of EMT

RNAi conditions	E-CADHERIN (Relative expression)	BRACHYURY (Relative expression)	
Non-interfering	1 <u>+</u> 0.06	1 <u>+0.06</u>	
Untreated	1.05 <u>+</u> 0.12	1.06+0.12	
<b>Brachyury RNAi</b>	0.98+0.30	0.36+0.15	



*E-CADHERIN* and *BRACHYURY* expression following *BRACHYURY* knockdown in SW480 sub-confluent cells by qRT-PCR. A) Table showing expression of *E-CADHERIN* and *BRACHYURY* following knock- down of *BRACHYURY* as assessed by qRT-PCR and compared to LAMIN C. *BRACHYURY* RNAi does not show any effective change on levels of E-CADHERIN. *BRACHYURY* knockdown level obtained is 64%. B) Graphical representation of Figure 4.3A showing that *BRACHYURY* RNAi has no effect on expression of *E-CADHERIN* (2% reduction). The Y-axis gives the relative expression levels and the X-axis mentions the genes analysed under different RNAi conditions. Raw data and the calculations are shown in Table 7 of appendix. Experiment was repeated 3 times (N=3), each experiment was done in triplicate. After  $\pm$ , the final standard deviation is mentioned that gives the error bars in the graph.

## 4.1.2A: Exploring a role for *BRACHYURY* in influencing the expression of epithelial cell polarity markers:

As previously mentioned, during the process of EMT, epithelial cells lose their cell polarity and cell junction molecules and obtain a fibroblastic phenotype.

Zeb1 is a key transcription factor that is up-regulated during EMT (Liu et al., 2008), that disrupts the cell polarity by suppressing the expression of tight junction molecules such as Lgl2, Crumb3 (Crb3), HUGL2 and PATJ (Pals1-associated tight junction protein), (Aigner et al., 2007; Spaderna et al., 2008). Snail, a repressor of E-CADHERIN, also inhibits the expression Crumb3 and disrupts apical cell polarity complex (Whiteman et al., 2008). Eva1 (Epithelial V-like antigen. Eva1 is an epithelial cell polarity marker and is a transmembrane glycoprotein. Crb3 are transmembrane proteins (Bachmann et al., 2001) which are linked to Pals1, a membrane associated guanylate kinase protein (Kamberov et al., 2000). Zeb2 (Zinc finger E-box binding homeobox 2) (also known as SIP1), is another marker that represses various epithelial cell–cell junction genes thereby inducing EMT (Vandewalle et al., 2005).

We looked at the expression of some of these genes, as markers for cells undergoing EMT following *BRACHYURY* RNAi. We observed no change in the expression levels of these marker genes of cell polarity following knock-down of *BRACHYURY* in SW480 sub-confluent cells (Figure 4.4). **Note:** Raw data for the Figure 4.4 is presented in Table 8 of the appendix.

Data from Figure 4.4 shows that none of the cell polarity markers mentioned changes their expression levels in response to *BRACHYURY* knockdown. This suggests that *BRACHYURY* does not regulate genes involved in cell polarity and EMT and confirms our observations in the previous section. This data is graphically represented in Figure 4.4B. We also carried out similar *BRACHYURY* knock-down experiments in high density SW480 cells. In high density SW480 cells, knock-down of *BRACHYURY* resulted in a slight increase



qRT-PCR showing expression of EMT markers in low density SW480 cells following knock down of *BRACHYURY*. A) Table demonstrating the expression of these markers do not alter following knock-down of *BRACHYURY*. B) Graphical representation of Table A, showing no effect of the cell polarity marker genes in response to *BRACHYURY* RNAi in low density SW480 cells. The X-axis mentions the genes analysed by qRT-PCR under different RNAi conditions and the Y-axis gives the relative gene expression levels. Experiment was repeated 3 times (N=3), each done in triplicate. All data were compared to  $\beta$ -catenin. The raw data is shown in appendix Table 8. in expression of ZEB1 increased by 1.2 times. But, since the p-value for this reduction was >0.05, the increase in ZEB1 expression was not considered significant. (Refer to appendix Table 25 for the p-values). PALS1 expression increased by 1.02 times (p-value>0.05; no significant reduction); ZEB2 expression was increased by 1.11 times, EVA1 by 1.16 times and CRB3 expression increased by 1.19 times. But the p-values obtained for these expression levels were >0.05, thus were not considered statistically significant difference in gene expression. The biological significance of such changes in high density cells is likely to be complex and was not studied further since the focus of our experiments was on low density cells where BRACHYURY expression was high. [Note: Data for BRACHYURY RNAi on high density SW480 cells is not presented because, level of BRACHYURY in high density SW480 cells is very little; knocking down which will not give any effective data].

## 4.1.2B: Effect of *BRACHYURY* on the expression of a marker of intestinal epithelium

We next explored the expression of the key marker of intestinal epithelium (*CDX2*, Caudal type homeobox transcription factor 2) in response to *BRACHYURY* RNAi by qRT-PCR. *CDX2* is a developmental gene which is involved in the formation of the placenta in mice (Chawengsaksophak *et al.*, 2004) and is also a biomarker of gastrointestinal differentiation, especially colorectal cancer (Liu *et al.*, 2007). In 2004, De Lott and co-workers studied tissue microarrays of 71 colorectal adenocarcinomas, 71.8% of which were *CDX2* positive. 74.5% of these positive tumours were differentiated tumours and 65% were high grade tumours. In contrast, when other carcinomas were studied 1 out of 47 were positive for *CDX2* staining (lung adenocarcinoma) and 2 out of 43 were positive for *CDX2* (pancreatic carcinoma). Therefore, it was concluded that *CDX2* is a critical intestinal marker though it is not that sensitive for those colorectal carcinomas which are not well-differentiated. Figure 4.5 shows the effect of *BRACHYURY* knock-down by RNAi on expression of *CDX2*, as assessed by qRT-PCR. We show that the expression of the intestinal epithelial marker, *CDX2*, is reduced by 74% in response to *BRACHYURY* RNAi in low density but not in high density SW480 cells.

[Note: Figure 4.5 is obtained from appendix table 9]. This may suggest that *BRACHYURY* may play a role in regulating *CDX2* at the invasive fronts but not at the central mass of the tumour.

Invasive fronts of CRC undergo a loss of basement membrane and activation of transcription factors related to EMT. This is the main reason for distant metastasis (Conacci-Sorrell *et al.*, 2003; Franci *et al.*, 2006; Spaderna *et al.* 2006). In contrast to what we might predict from our data, *CDX2* expression was shown to be decreased in invasive cells at the stromal front of differentiated CRC (Brabletz *et al.*, 2004). In contrast to the studies of Brabletz and co-worker, and in support of our *in vitro* studies, Gross and co-workers (2008) showed that expression of *CDX2* is high at the invasive front and its expression opposes cell migration and dissemination in colorectal tumours. We therefore thought it is relevant to investigate a possible role of *BRACHYURY* in influencing colon cancer cell migration. This is further described in section 4.5 of this chapter.

## 4.1.2C: Role of BRACHYURY in regulating expression of known target genes

In this section, selected known target genes of BRACHYURY were studied in response to *BRACHYURY knock-down* by RNAi. *Orthopedia (otp)* in *Drosophila* is regulated by its *BRACHYURY* homologue Brachyenteron (Byn) (Simeone *et al.*, 1994; Singer *et al*, 1996). In *Drosophila*, Byn is involved in development of the hindgut and also to a certain extent in midgut development (Kispert *et al.* 1994; Singer *et al.*, 1996; Kusch and Reuter 1999). Otp was shown to be a direct target of Byn (Kusch *et al.*, 2002). The *cis*-regulatory region of *otp* 



Expression of *CDX2* following *BRACHYURY* RNAi by qRT-PCR in low density SW480 cells. A) Table showing that *CDX2* expression is reduced by 74% following *BRACHYURY* RNAi in sub-confluent SW480 cells. The CT value for CDX2 was compared to CT value of *LAMIN C*. The p-value for this reduction was 0.00016, therefore may be considered a significant reduction. B) Graphical representation of Table 4.5A along with error bars (obtained from SD). The X-axis mentions different RNAi conditions for *CDX2* and the Y-axis gives the relative *CDX2* expression levels.

Experiment was done 3 times (N=3); each experiment done in triplicate. Raw data is detailed in Table 9 of appendix. contains multiple binding sites of high (type A) and of low (type B) affinity to *Byn*.

In *Xenopus*, *BRACHYURY* (*Xbra*) is required for mesoderm formation and notochord development (Smith *et al.*, 1991). *Xwnt11* was shown to be a direct target of Xenopus BRACHYURY (Masazumi *et al.*, 2000). Wnt11 is a non-canonical Wnt (i.e. Wnt family member activating other pathways) has been shown to have increased expression in many prostate tumours and during prostate cancer progression (Zhu *et al.*, 2004). As summarised by Katoh and Katoh (2009), Wnt11 is also expressed and up-regulated in many other human cancers such as breast cancer, gastric cancer, esophageal cancer, primary colorectal cancer, neuroblastoma and Ewing sarcoma.

Expression of *IL-2* and *IL-15* were shown to be positively regulated by *BRACHYURY*, in mouse (Baldassarre *et al.*, 2001). *IL-2* and *IL-15* (structurally similar to *IL-2*) are two cytokine immune system signalling molecules and are widely studied in cancer cells. Both *IL-2* and *IL-15* can upregulate stimulatory molecules of chronic leukaemia cells (Spaner *et al.*, 2004; Brentjens *et al.*, 2003). In human, *IL-15* is expressed in metastatic colorectal cancer, activate NK cells and put forth an anti-tumour effect. It also causes growth of colon cancer cells by metastasis and tumour progression (Kuniyasu *et al.*, 2003).

*NANOG* is a key gene involved in the maintenance of pluripotency in stem cells. In mouse, NANOG has *BRACHYURY* and *Stat3* binding sites *in vivo* (Suzuki *et al.*, 2006) which regulates the expression of *NANOG*. *NANOG* has been shown to be expressed in a number of cancers such as primordial germ cell tumours (where it is involved in malignant cell transformation of gonads), osteosarcoma and lung cancer (Nirasawa *et al.*, 2009). *NANOG* was also shown to be expressed in human breast adenocarcinoma cell line, MCF-7 (Hart *et al.*, 2005; Ezeh *et al.*, 2005). *NANOG* has been shown to be a direct target of Tcf3 (Pereira, 2006) and Tcf3 interacts with β-catenin in the Wnt pathway (Korinek, 1998). In this section, we investigated the role of *BRACHYURY* in regulating selected target genes in SW480 sub-confluent cells. Knock-down of *BRACHYURY* in low density SW480 cells has the following effects (refer Figure 4.6): *WNT11* expression was reduced by 70%, *NANOG* expression reduced by 85%, *Orthopedia* by 75%, *IL2* by 74% and *IL15* by 64%. The p-value <0.05 for all these expressions; therefore these fold differences were considered statistically significant. [Note: p-values mentioned are significance of the gene knock-down compared to non-interfring siRNA treated cells. Refer appendix table 25 for the p-values obtained]. Based on these results, it seemed interesting to study some of these genes further. Specifically, we focussed our studied on the potential regulation of *NANOG* by *BRACHYURY* and this will be discussed in a subsequent section (section 4.2.1).

[Note: Figure 4.6 is obtained from appendix table 10)

## 4.1.2D Effect of *BRACHYURY* RNAi on inducer of EMT, *E-CADHERIN*, *SNAIL* and T-box genes, *TBX2* and *TBX3*

In this section, the expression levels of *E-CADHERIN*, *SNAIL* and the Tbox developmental genes (*TBX2* and *TBX3*) were studied in response to transient *BRACHYURY* knock-down in low density SW480 cells. We analysed expression of *TBX2* and *TBX3* in response to *BRACHYURY* RNAi because *BRACHYURY* is a T-box gene and we wanted to analyse the effect of BRACHYURY in other Tbox genes such as *TBX2* and *TBX3*. We analysed expression of *SNAIL* since *Snail* is a negative regulator of *E-CADHERIN*. The expression of T-box gene *TBX2* was shown to be reduced by 59% and *TBX3* expression was shown to be reduced by 57%, in response to *BRACHYURY* knock-down (Figure 4.7). P-value for these expressions were shown to be <0.05 and were considered statistically significant results.

Once again, *E-CADHERIN* expression was analysed along with the expression of *SNAIL*, which is a negative regulator of *E-CADHERIN*, both of which have shown to have no change in expression in response to *BRACHYURY* RNAi in low density SW480 cells (Figure 4.7). The p-values for these changes



Effect of *BRACHYURY* RNAi on target genes of *BRACHYURY* in SW480 subconfluent cells as assessed by qRT-PCR. A) Table demonstrating the relative gene expression in response to *BRACHYURY* knock-down on its target genes. B) Graphical representations for Figure 4.6A showing reduction in expression of these target genes on *BRACHYURY* RNAi. The X-axis mentions the genes analysed by qRT-PCR under different RNAi conditions and the Y-axis gives the relative gene expression levels.

All data are compared to *LAMIN C*. Experiments were done 3 times (N=3), each in triplicate. Raw data for this figure is detailed in appendix table 10. The final SD is mentioned after  $\pm$ 



#### Figure 4.7:

Expression of developmental genes in response to *BRACHYURY* RNAi in SW480 low density cells: A) Table demonstrating the relative expression level of the genes. B) Graphical representations of Table A showing reduction in expression of T-box genes on *BRACHYURY* knock-down in low density SW480 cells but not alteration in expression of *E-CADHERIN* and its negative regulator, *SNAIL*. The X-axis mentions the genes analysed by qRT-PCR under different RNAi conditions and the Y-axis gives the relative gene expression levels. Data was compared to *LAMIN C*. Appendix table 11 details the raw data for this figure. Experiment was done 3 times (N=3), each in triplicate. The final SD is mentioned after  $\pm$  were > 0.05, thus not showing any statistical significant data. [Note: p-values mentioned are significance of the gene knock-down compared to mock-treated SW480 cells]. These data were summarised in Figure 4.7. Figure 4.7 is obtained from appendix Table 11.

Changes in the expression of few of these genes appeared interesting, one of which was the reduction in expression of *NANOG*. Since, *NANOG* is involved in maintenance of pluripotency in stem cells; we focussed our studies on exploring a possible role for *BRACHYURY* in maintaining a cancer putative stem cell population. This is explained in part b of this chapter in full details.

## b) Role of *BRACHYURY* in influencing the regulation of *NANOG* and maintaining a putative cancer stem cell population

4.2.1 The expression of *NANOG* is altered in response to changes in *BRACHYURY* in low density SW480 cells:

One of the most significant changes in gene expression following *BRACHYURY* RNAi was reduction in *NANOG* expression (shown by qRT-PCR). *NANOG* is a key inducer of pluripotency involved in maintenance of stem cells and is expressed in tissue-specific stem cells, such as the pancreas (Koblas *et al.,* 2008). This led us to suggest that *NANOG* may be maintaining a population of so called cancer stem cells for the reasons outlined below.

CD166, CD133 and LAMIN A are three colorectal cancer stem cell markers we analysed in this study. LAMIN A was shown to be a novel biomarker for colorectal cancer stem cells (Naomi et al., 2008). It has recently been shown that expression of A-type lamins in CRCs is correlated with a poor prognosis due to increased tumour invasiveness (Willis et al., 2008). Furthermore, in addition to LAMIN A being expressed in differentiated epithelia, it has also been shown to be expressed in cells of the colonic crypt that reside in the stem cell niche. Due to this critical link between the presence of LAMIN A and cells having stem cell character, together with the link with poor prognosis for CRC patients, we investigated whether BRACHYURY might also regulate the levels of LAMIN A in CRC cells. Ricci-Vitiani and co-workers have demonstrated that CD133 can be used as a colorectal cancer stem cells marker (2007). However, not all tumours display this marker nor is every  $CD133^+$  cell a CSC (Dalerba et al., 2007: LaBarge and Bissel, 2008; Shmelkov et al., 2008)]. In mouse, CD133 expression was observed in differentiated cells such as proximal renal tubule cells, neurons in adult brain and pancreas (Zhu et al., 2009). This agrees with recent studies done by Shmelkov and co-workers (2009) in mouse tissues. In small intestine of mouse, Lgr5<sup>+</sup> stem cells were shown to express CD133 (Zhu et al., 2009). [Lgr5 marks intestinal stem cells (Barker et al., 2007).]

A more specific marker for CRC-SCs has been described as CD166 (Dalerba *et al.*, 2007). Almost 60% of CRCs express strong cytoplasmic staining of CD166, and 30% have membrane staining of CD166, and expression of CD166 in CRC is associated with a poor prognosis (Weichert *et al.*, 2004).

Firstly, we investigated the endogenous protein levels of BRACHYURY, NANOG and CRC stem cell markers (CD166, CD133, LAMINA) in low and high density SW480 cells grown for 1 day and 4 days, respectively (Figure 4.8a). Protein levels decrease to almost non-detectable when the cells transition from sub- to super-confluent stages.

We next investigated the protein levels of NANOG in response to BRACHYURY knock-down in low density SW480 cells. Our results show that NANOG levels are down-regulated in response to *BRACHYURY* knock-down in low density SW480 cells (Figure 4.8b). We also studied the expression of colorectal cancer stem cell markers in response to BRACHYURY RNAi. We show that levels of CRC stem cells markers (CD166, CD133, LAMIN A) are decreased following knock-down of *BRACHYURY* (Figure 4.8b).

#### β-catenin is upstream of BRACHYURY in SW480 sub-confluent cells:

β-catenin is a major oncogene which drives CRC and the Wnt signalling pathway (Wnt3) has been shown to regulate expression of *BRACHYURY* (Arnold *et al.*, 2000). We therefore knocked down β-catenin to determine whether it might be upstream of *BRACHYURY* in these cells. Knock-down of β-catenin by RNAi results in the down-regulation of BRACHYURY as shown by western blot analysis, in low density SW480 cells (Figure 4.8c). This down-regulated the levels of NANOG and CD133 in low density SW480 cells. LAMIN B was used as loading control in this experiment (Note: Figure 4.8 shows equal loading of LAMIN B, thus as a control).

These results led us to suggest that in low density mesenchymal-like cells, signalling through  $\beta$ -catenin induces BRACHYURY which in turn induces NANOG which acts to maintain a population of cells with stem cell-like

× a



Western blot analysis of levels of BRACHYURY, NANOG and CRC stem cell markers. a) Endogenous levels of BRACHYURY, NANOG, CD166 and CD133 at day 1 (sub-confluent) SW480 cells which decreased to almost non-detectable levels on day 4 (super-confluent) SW480 cells. b) Levels of NANOG and CRC stem cell markers (CD133, CD166, LAMN A) decrease in response to *BRACHYURY* RNAi in low density SW480 cells. LAMIN B was used as a loading control. The BRACHYURY lane confirms the knock-down. c) Western blot analysis showing  $\beta$ -catenin knock-down decreases the levels of BRACHYURY suggesting that  $\beta$ -catenin is upstream of BRACHYURY. This inturn, reduces the levels of NANOG and CRC stem cell marker. LAMIN B in each blot was used as loading control confirming equal amount of protein loaded in each lane. U = untreated cells, N=cells treated with non-interfering RNAi, B = BRACHYURY RNAi,  $\beta = \beta$ -catenin RNAi.

properties. (At high cell density in epithelial like cells, signalling through βcatenin might be low and subsequent expression of *BRACHYURY*, *NANOG* and cells with cancer stem cell phenotype also would be low). To confirm the relationship we observed by western blot analysis, between levels of BRACHYURY, NANOG and CRC stem cell markers, we studied the localisation of BRACHYURY, NANOG and CRC stem cell markers by immunoflouresence in low and high density SW480 cells. Specifically, we wanted to determine the levels and localisation of BRACHYURY, NANOG, CD133 and CD166 (CRC stem cell marker) in low and high density SW480 cells and to determine whether cells that express BRACHYURY also express NANOG, CD133 and CD166. [**Note**: β-catenin knock-down reduces BRACHYURY levels, thus β-catenin is upstream of BRACHYURY. But, *BRACHYURY* RNAi does not reduce the levels of β-catenin, confirming that βcatenin is upstream of BRACHYURY. Therefore, β-catenin can be used a control].

We show that all SW480 cells at low density express BRACHYURY and NANOG and also CD166 and CD133 (Figure 4.9). BRACHYURY and NANOG localisation was found to be nuclear (as expected as they are transcription factors), whereas, CD133 and CD166 confirms their surface antigen localisation. Thus, there is 100% co-localization of BRACHYURY, NANOG and the CRC stem cell markers in low density SW480 cells.

In contrast to low density cells, high density SW480 cells express BRACHYURY, NANOG, CD133 and CD166 very rarely. Moreover, the rare cells that express BRACHYURY also express NANOG. Furthermore, we showed that the cells that are positive for BRACHYURY and NANOG, are also positive for CD166 in high density SW480 cells (Figure 4.9).

In high density SW480 cells,  $\beta$ -catenin localisation mostly membrane bound with occasional nuclear localisation (personal communication Dr. Melanie Davies). We wanted to analyse whether the rare BRACHYURY positive cells in high density SW480 cells co-localise with occasional nuclear  $\beta$ -catenin (nuclear

	Brachyury	Nanog	Merged	DIC
Low density SW480	a		*	8
	Brachyury	Nanog	Merged	DIC
High density SW480	b e	•	•	
	Nanog	CD 166	Merged	DIC
Low density SW480	c	9	٠	¢
	Nanog	CD 166	Merged	DIC
High density SW480	d	6	4	
	CD 133	Brachvury	Merced	DIC
			mergeu	DIC
Low density SW480	e At	•	An Neigea	C:
Low density SW480	e • CD 133	Brachyury	Merged	DIC
Low density SW480 High density SW480	e CD 133 f	Brachyury	Merged	DIC
Low density SW480 High density SW480	e CD 133 f Nanog	Brachyury CD 133	Merged	DIC
Low density SW480 High density SW480 Low density SW480	e CD 133 f Nanog g	Brachyury CD 133	Merged Merged Merged	DIC
Low density SW480 High density SW480 Low density SW480	e CD 133 f Nanog g Nanog	Brachyury CD 133 CD 133	Merged Merged Merged Merged	DIC

**Figure 4.9** (a-h): Confocal IF images showing localisation of BRACHYURY, NANOG, CD133 and CD166 in low and high density SW480 cells.



#### Figure 4.9h

Figure 4.9: Indirect IF using respective antibodies and confocal microscopy shows that BRACHYURY and NANOG are localised to the nucleus, whereas CD166 and CD133 are surface antigen markers. a) BRACHYURY co-localises with NANOG in low density SW480 cells. b) BRACHYURY co-localises with NANOG in high density SW480 cells. c) NANOG co-localises with CD166 in all SW480 sub-confluent cells. d) NANOG co-localises with CD166 in all SW480 super-confluent cells. e) CD133 co-localises with BRACHYURY in all low density SW480 cells. (f) CD133 co-localises with BRACHYURY in rare high density SW480 cells. g) NANOG co-localises with CD133 in all SW480 sub-confluent cells. h) NANOG co-localises with CD133 in all SW480 sub-confluent cells. h) NANOG co-localises with CD133 in SW480 super-confluent cells. h) NANOG co-localises with CD133 in SW480 super-confluent cells. h) NANOG co-localises with CD133 in SW480 super-confluent cells. h) NANOG co-localises with CD133 in SW480 super-confluent cells. h) NANOG co-localises with CD133 in SW480 super-confluent cells. h) NANOG co-localises with CD133 in SW480 super-confluent cells. h) NANOG co-localises with CD133 in SW480 super-confluent cells. h) NANOG co-localises with CD133 in SW480 super-confluent cells. h) NANOG co-localises with CD133 in SW480 super-confluent cells. h) Rare BRACHYURY positive SW480 cells in high density are not the cells with nuclear  $\beta$ -catenin.

 $\beta$ -catenin means active Wnt signalling). Our data show that rare BRACHYURY positive cells in high density SW480 cells do not correspond to cells with nuclear  $\beta$ -catenin (Figure 4.9 i).

Thus, this section of the chapter has given us important information on localisation of BRACHYURY, NANOG and CRC stem cell markers in SW480 cells and their co-localisation with each other. We have studied further possible roles of BRACHYURY in later sections.

## 4.2.2 Does *NANOG* regulate specific gene expression patterns in SW480 cells?

The down-regulation of *NANOG* in human ES and EC (embryonal carcinoma) cells was shown to result in differentiation of the cells along an extraembryonic lineage, as determined by the expression of genes such as *GATA2*, *GATA4*, *GATA6* and *CDX2* (Hyslop *et al.*, 2005). Specifically, knockdown of *NANOG* increased the expression of the genes *GATA1*, *4*, *6*, (extraembryonic endoderm genes), *CDX2* and *GATA2* (tropho-ectoderm associated genes) showing that the down-regulation of *NANOG* induced differentiation in human ES and EC cells. *NANOG* knock-down (by RNAi) also decreased the expression of *Oct-4* but did not change the expression of *SOX2* indicating that this differentiation cannot be used to measure the pluripotency in these cells. We wanted to determine whether expression of these genes altered similarly, as a result of lower expression levels of *NANOG* in SW480 cells. Therefore, we looked at the expression of *GATA2*, *GATA4*, *GATA6*, *Oct-4* and *SOX2* in low density SW480 cells following RNAi induced knock-down of *BRACHYURY*.

The MYC family of transcription factors play an important role in cell proliferation, differentiation and in development of the embryo and in the progression of cancer (Henrikson and Lu<sup>°</sup>scher, 1996) and is also one of the key transcription factors used to induce pluripotency (Takahashi *et al.*, 2006). C-MYC has been shown to be a target of Wnt signalling pathway (He *et al.*, 1998). Since MYC is an important downstream target of  $\beta$ -catenin, we therefore, looked at expression of this gene following knock-down of *BRACHYURY*.

*FOXD3* is another embryonic stem cell transcription factor which is down-regulated during endoderm formation (Guo *et al.*, 2002). We therefore, also studied the expression of *FOXD3* in low and high density SW480 cells and also in response to *BRACHYURY* RNAi in low and high density SW480 cells.

Our results show that *GATA2*, *GATA6*, *MYC* and *FOXD3* do not alter in response to changes in SW480 cell density as compared to low density SW480 cells. Expression of *GATA4* was reduced by 57%, *Oct-4* by 50% and *SOX2* by 78% in SW480 high density cells compared to SW480 sub-confluent cells (Figure 4.10).

In response to *BRACHYURY* knock-down, expression of *SOX2* is reduced by 68% expression of *FOXD3* is reduced by 69% in SW480 sub-confluent cells, whereas expression of *MYC* increases by 9.8 times in low density SW480 cells (Figure 4.11).

To summarise, down-regulation of *NANOG* in SW480 cells does not induce differentiation and expression of genes associated with extra-embryonic lineages. Expression of the pluripotency genes *SOX2* and *FOXD3* is reduced, but not *Oct-4*. Furthermore, expression of *c-MYC* is increased by 9.8 fold suggesting that a complex regulatory mechanism is working in these cells.

Thus, we present data for the use of SW480 cells grown to low and high cell density as a model in which to study EMT/MET. We have shown that the expression of *BRACHYURY* is linked with the regulation of the pluripotency gene *NANOG*, with the maintenance of markers of CSCs and regulation of LAMIN A (a biomarker linked to poor prognosis) in low density SW480 cells.



qRT-PCR expression data for *GATA* genes, pluripotency marker genes, *MYC* and *FOXD3* in SW480 cells. A) This table summarises the relative expression of the above mentioned genes in low and high density SW480 cells. [Expressions in high density cells are in relative to that of sub-confluent cells]. B) Graphical representation of Table A showing the changes in gene expression of the genes in low and high density SW480 cells. The X-axis mentions the genes analysed by qRT-PCR in low and high density SW480 cells. The Y-axis gives the relative gene expression levels. [Experiments were done 3 times (N=3), each in triplicate. All data were compared to  $\beta$ -catenin. This figure is derived from Table of appendix 13. Refer appendix Table 26 for the statistical significance data of the fold differences obtained for each gene in low and high density cells.



#### Figure 4.11:

qRT-PCR of *GATA* genes, pluripotency marker genes, *MYC* and *FOXD3* in SW480 sub-confluent cells. A) Table demonstrating the expression of these genes in low density SW480 cells on BRACHYURY RNAi, as assessed by qRT-PCR. B) Graphical representation of Table A showing no change in *GATA* genes and *OCT4* but 9.8 fold increase in *c-MYC*. The X-axis mentions the genes analysed by qRT-PCR under different RNAi conditions and the Y-axis gives the relative gene expression levels.

[Experiments were done 3 times (N=3), each in triplicate. All data were compared to  $\beta$ -catenin. This figure is derived from Table of appendix 12. BRACHYURY knock-down ranged from 68-80%]. The Final SD is mentioned after  $\pm$ . We also show that the oncogene of the Wnt pathway,  $\beta$ -catenin, which is a modulator of 'stem' signaling pathways, affects the levels of BRACHYURY showing that *BRACHYURY* may be an important factor in transducing the  $\beta$ -catenin signaling pathway in the maintenance of cells with a CSC-like phenotype. We continue to explore the role of BRACHYURY in SW480 cells and the next section will describe the influence of BRACHYURY on NF $\kappa$ B signalling.

# c) A role of *BRACHYURY* in regulating the NFκB pathway?

#### 4.3 A role for BRACHYURY in regulating the NFKB pathway

NFκB (Nuclear factor kappa β) refers to a family of transcription factors and has crucial roles in the immune system (Ghosh *et al.*, 1998; Li and Verma 2002; Bonizzi and Karin 2004). The activation of the NFκB pathway is also involved in many cancers such as leukemia, lymphoma, colon cancer and ovarian cancer (Rayet and Gelinas 1999) and commonly occurs through de-regulation of upstream regulators. In mammals, the NFκB family is composed of the transcription factors, p50, p52, REL A (p65), c-REL And REL B (Moynagh 2005; Hoffmann *et al.*, 2006; Hayden *et al.*, 2004; Perkins *et al.*, 2007). REL A, c-REL and REL B contain C-terminal transcriptional activation domains (TADs), which are involved in activation of target gene expression. p50 and p52 do not contain C-terminal TADs and are derived from larger precursors, p105 and p100, respectively. Their homodimers repress transcription unless they are bound to a protein containing a TAD, such as REL A, c-REL or REL B or Bcl-3 (a related transcriptional co-activator).

Recently, it has been shown that *NANOG* inhibits the NF $\kappa$ B signalling pathway in mouse ES cells, thereby maintaining pluripotency (Watt, F. and Torres, J., 2008). As mentioned above, the NF $\kappa$ B family consists of 5 member proteins, out of these *REL A and REL B* were extensively expressed in ES cells, the other three members being present to be a lesser amount. *NANOG* was shown to bind to REL A in ES cells where it acts to inhibit NF $\kappa$ B regulated genes. Expression of NF $\kappa$ B regulated genes occurs on differentiation with dissociation of this complex. We wanted to determine whether *NANOG* was regulating targets of NF $\kappa$ B genes similarly in SW480 cells. We examined the expression of NF $\kappa$ B regulated genes in response to *BRACHYURY* knock-down in low density SW480 cells by qRT-PCR (Figure 4.12). These values are derived from the raw data presented in Table 14 of appendix.

The following NFkB regulated genes were studied:



Expression of NF $\kappa$ B regulated genes in response to *BRACHYURY* knock-down in SW480 sub-confluent cells. A) Table demonstrating the up-regulation of *VIMENTIN*, *PLAUR*, *BCL2L1* and *IL10* in response to *BRACHYURY* RNAi in low density SW480 cells obtained by qRT-PCR. This data was compared to cells with no treatment and was also controlled with mock treated SW480 low density cells. The table was derived from Appendix table 14. B) Graphical representation of the table 4.12A along with error bars obtained from their standard deviations. The X-axis mentions the genes analysed by qRT-PCR under different RNAi conditions and the Y-axis gives the relative gene expression levels. Experiments were repeated 3 times (N=3), each done in triplicate; and data was compared to *LAMIN C*. C) Western blot analysis demonstrating the increased levels of REL A, phosphorylated REL A and VIMENTIN in response to *BRACHYURY* RNAi in low density SW480 cells. LAMIN B was used as a loading control showing equal loading for all samples. The experiment was repeated 3 times (N=3). *VIMENTIN* is an intermediate filament protein and has metastatic and invasive roles in many cancers such as prostate cancer (Wei *et al.*, 2008) and colon cancer. *PLAUR* (plasminogen activator) has a role in metastasis in colon and liver cancers (Illemann *et al.*, 2009). *Il10* is an anti-inflammatory cytokine and also is critical for the development and progression of sporadic colon cancer (Tamara *et al.*, 2008). *BCL2L1* belongs to the Bcl-2 family which forms dimers and can behave as anti- and pro-apoptotic regulators in many cellular activities.

Figure 4.12 (A, B) demonstrates that the expression of NFkB regulated genes; PLAUR, BCL2L1 and VIMENTIN are upregulated following BRACHYURY knock-down in low density SW480 cells (by 2 fold increase) obtained by qRT-PCR, but there was no change in expression of IL10 (Figure 4.12 A, B). But, the p-value for the difference sin expression of *IL10* in response to the knock-down was shown to be <0.05 (appendix Table 25). Thus, this change in expression was considered significant along with the expression of PLAUR, BCL2L1 and VIMENTIN. We then carried out Western blot analysis to confirm the levels phosphorylated REL A (Ab1604, Chemicon), the p65 subunit of NFkB and also the expression of VIMENTIN, in response to BRACHYURY RNAi in low density SW480 cells (Figure 4.12 C). Our Western blot analysis show that both phosphorylated REL A (p-REL A) and VIMENTIN are upreguated on knocking down BRACHYURY showing possible active NFKB signalling in low density SW480 cells in response to knock-down of BRACHYURY, In summary, BRACHYURY may inhibit NFKB signalling in low density SW480 cells, and this may occur through activation of NANOG.

We also examined the expression of  $NF\kappa B$  and STAT3 in low density SW480 cells on *BRACHYURY* RNAi (Figure 4.13). Stat3 is a transcription factor shown to interact with NF $\kappa$ B1 (Yu *et al.*, 2002). Wnt pathway is shown to upregulate Stat3 and prevents differentiation of mouse embryonic stem cells (Hao *et al.*, 2003). So, we wanted to check the expression of *STAT3* in response to *BRACHYURY* RNAi in SW480 cells to know more about a possible role of *BRACHYURY* in NF $\kappa$ B signalling. Our results (Figure 4.13) show that knocking down *BRACHYURY* did not have any effect on the expression levels of *NF\kappaB* and *STAT3* in low density SW480 cells, showing no direct interaction of BRACHYURY with *NF\kappaB* and STAT3 in SW480 cells.



Expression of  $NF\kappa B$  and STAT3 in response to BRACHYURY RNAi in low density SW480 cells. A) Table demonstrating the no change in expression of these genes on knocking down BRACHYURY in sub-confluent SW480 cells. Raw data for table A is in Table 15 of appendix. Data on BRACHYURY RNAi was compared to cells treated with non-interfering siRNA. B) Graphical representation of the expression of the table A, showing reduction in  $NF\kappa B$ expression was 6% and STAT3 expression was increased by 1.16 times; p-values >0.05. Experiments were repeated 3 times (N=3), each done in triplicate; and data was compared to *LAMIN C*. The X-axis mentions the genes analysed by qRT-PCR under different RNAi conditions and the Y-axis gives the relative gene expression levels. For p-value, refer to appendix Table 25.

#### d) Role of BRACHYURY in cell proliferation

#### 4.4.1 Cell proliferation and cancer:

Cell proliferation is usually maintained by growth factors, receptor molecules in the cell, signalling pathways and transcription factors that bind to the DNA. Mutation in any of these genes can lead to changes in the cell proliferation, as frequently occurs in cancer. In cancer, genetically altered cells can proliferate abnormally to a secondary site (neoplasia) to continue forming tumour and thus spread the disease. Cell proliferation is often characterized by loss of normal tissue organisation and cell structure. In colon cancer, this type of proliferation is caused by chronic ulcerative colitis, infectious agents, and saturated fats (Preston-Martin, 1990).

Earlier in this chapter, we have seen that expression of *NANOG* is influenced by *BRACHYURY* in SW480 cells. *NANOG* is a transcription factor required to maintain the undifferentiated state of embryonic stem cells. Recently, exogenously transcribed *NANOG* was shown to enhance the proliferation of NIH3T3 cells by enabling the cells to enter S phase (Zhang *et al.*, 2005). [Note: NIH3T3 cells are mouse embryonic fibroblast cell line]. Thus, it was predicted that *NANOG* might have a similar function in ES cells. *NANOG*'s overexpression in many forms of cancer makes it an interesting candidate to study; furthermore, the oncogenic potential of NANOG was demonstrated when it was shown to transform NIH3T3 cells and increase the cell growth rate (Piestun *et al.*, 2006). We therefore, wanted to determine whether *BRACHYURY* has an effect on cell proliferation in SW480 cells via *NANOG*.

#### 4.4.2 Cell proliferation by mitotic index count

We have shown that the regulation of *NANOG* is influenced by *BRACHYURY* in SW480 cells (section 4.2); and since *NANOG* plays an important role in proliferation in NIH3T3 cells, we investigated the possible role of *BRACHYURY* in regulating cell proliferation in SW480 cells. We knocked down the expression of *BRACHYURY* in SW480 cells by RNAi and grew the 124 cells for 1, 2, 3, 4, 5 and 6 days (1 day = sub-confluent, 3 days = confluent, 4 days = super-confluent). We counted the number of Propidium iodide stained cells in all stages of mitosis to calculate the number of proliferating cells. [Mitotic index is a measure of the proliferating cells in a population, and is expressed as the ratio between the number of cells in mitosis and total number of cells]. This experiment was controlled by counting the cells in mitosis in cells without any treatment and also cells with non-interfering siRNA. Data in Figure 4.14 shows that the number of cells in mitosis decreases as SW480 cells grew from low to high density. On treating SW480 cells with BRACHYURY siRNA, the number of proliferating cells increase on day 4 (super-confluent) and day 5 (24 hours after super-confluency) as compared to cells with no treatment. (Raw data for these graphs are in Table 16 of appendix; and the data for each condition was normalised to medium without any cells). We applied the two tailed t- test to measure the statistical significance of this data (Table 17, appendix). Thus, we were able to show that the higher rate of proliferation of SW480 cells treated with BRACHYURY siRNA, 4 days after RNAi is statistically significant (Table 17, appendix). Furthermore, to determine whether the knock-down is active over the period of 6 days after BRACHYURY RNAi, we analysed the knock-down levels for each day of the experiment. We show that the levels of BRACHYURY knock-down remain high for days 1 (56%), 2 (72%), 3 (57%) and 4 (66%). On day 5 (38%) and 6 (38%) after BRACHYURY RNAi the knock-down efficiency starts to decrease (Figure 4.15).

In summary, there is no change in mitotic index in *BRACHYURY* knocked down cells for the first 3 days after RNAi; however after 4 days of *BRACHYURY* RNAi there is a small but statistically significant increase in number of cells in mitosis following *BRACHYURY* knock-down (Figure 4.14). The next day (day 5 after RNAi) showed a further increase in number of proliferating cells in response to *BRACHYURY* RNAi. But, Figure 4.15 shows that the knock-down levels of BRACHYURY for day 5 (38%) is not as high as the first 4 days after RNAi. Thus, the mitotic index result for day 5 is not considered. Day 6 after *BRACHYURY* RNAi shows no increase in number of proliferating cells on knocking down the expression of *BRACHYURY*.



A.

C.



Β.

Figure 4.14: Mean mitotic index count of SW480 cells. Graphical representation of mitotic index count of A) untreated, B) non-interfering siRNA, C) BRACHYURY knocked down SW480 cells over a period of 6 days after RNAi. D) This graph represents a comparison of the mitotic index count of cells treated with BRACHYURY siRNA (green) in comparison to mock-treated cells (red). X-axis = no. of days cells were grown after RNAi, Y-axis = Mitotic index count. Raw data for this figure is detailed in appendix Table 16. N=4 (for days 1 to 4) and N=3 (for days 5 and 6).

126



**Figure 4.15:** *BRACHYURY* knock-down levels after siRNA treatment. A) The relative gene knock-down levels of *BRACHYURY* compared to the SW480 cells treated with non-interfering siRNA. The raw data for this table is presented in **Table 18** of appendix. All data were compared to  $\beta$ -catenin. Experiment was done three times (N=3), each done in triplicate. B) Graphical representation table A showing high knock-down levels on days 1-4 of RNAi, but low RNAi levels on day 5 and 6 after RNAi. The X-axis mentions that BRACHYURY levels are analysed by qRT-PCR under different RNAi conditions on days 1 to 6 after RNAi. Y-axis gives the relative *BRACHYURY* expression levels.

#### 4.4.3 Cell proliferation by CellTiter-Glo Luminescent Cell Viability Assay:

To complement the data provided by the mitotic index count, we also performed a proliferation assay using CellTiter-Glo Luminescent Cell Viability Assay (Promega, Catalogue number G7571, see Materials and Methods section 2.22). This assay is based on quantifying the ATP present, thus can be used for measuring, cell viability and cell proliferation (Crouch *et al.*, 1993). We compared the proliferation of the SW480 cells treated with *BRACHYURY* siRNA to cells treated with non-interfering siRNA and untreated cells using this assay.

The number of proliferating SW480 cells on *BRACHYURY* knock-down was more compared to the untreated control conditions on days 1 to 4 after RNAi. We show that on days 1, 2 and 3 after RNAi, the number of proliferating SW480 cells on *BRACHYURY* knock-down was similar to that of untreated and non-interfering controls (Figure 4.16:A to F). But, 4 days after the RNAi treatment, when the SW480 cells are super-confluent, the number of proliferating SW480 cells on *BRACHYURY* knock-down was more compared to the untreated and non-interfering control conditions (Figure 4.16:G-H), thereby further validating our data obtained from the mitotic index count. [Raw data for Figure 4.16 is in Table 19 of appendix]

Figure 4.17 demonstrates the efficiency of *BRACHYURY* knock-down until 6 days after RNAi. We once again, show that the knock-down levels of *BRACHYURY* on day 5 (27%) and 6 (34%) is not as high compared to the first 4 days after the treatment (day 1=47%, day 2=69%, day 3=58% and day 4=57%). Thus, we obtained the cell proliferation data only for the first 4 days of the RNAi treatment, and not for day 5 and 6 of RNAi. Note: [Raw data for Figure 4.17 is in Table 20 of appendix].

We suggest that *BRACHYURY* might have a role in regulating cell proliferation in super-confluent SW480 cells although the mechanism through which this would work is unclear.



**Figure 4.16 A-B:** Graphs demonstrating the average luminescence obtained (Y-axis) against the number of viable cells (X-axis). This graph shows that the number of proliferating cells on knocking down *BRACHYURY*, on day 1 of treatment, is similar compared to that of untreated (A) and non-interfering siRNA (B) controls. For each data point, N=1, done in triplicate.



**Figure 4.16 C-D:** Graphs demonstrating the average luminescence obtained (Y-axis) against the number of viable cells (X-axis). This graph shows that the number of proliferating cells on knocking down *BRACHYURY*, on day 2 of treatment, is similar compared to that of untreated (A) and non-interfering siRNA (B) controls. For each data point, N=1, done in triplicate.



**Figure 4.16 E-F:** Graphs demonstrating the average luminescence obtained (Y-axis) against the number of viable cells (X-axis). This graph shows that the number of proliferating cells on knocking down *BRACHYURY*, on day 3 of treatment, is similar compared to that of untreated (A) and non-interfering siRNA (B) controls. For each data point, N=1, done in triplicate.



**Figure 4.16 G-H:** Graphs demonstrating the average luminescence obtained (Y-axis) against the number of viable cells (X-axis). This graph shows that the number of proliferating cells on knocking down *BRACHYURY*, on day 4 of treatment, is similar compared to that of untreated (A) and non-interfering siRNA (B) controls. For each data point, N=1, done in triplicate.

**Figure 4.16:** Cell proliferation assay by CellTiter-Glo Luminescent Cell Viability Assay kit to demonstrate the proliferation of SW480 cells on *BRACHYURY* RNAi compared to the untreated and non-interfering siRNA treated cells, as the SW480 cells reach from sub- to super-confluent stage.


#### Figure 4.17

Efficiency of *BRACHYURY* RNAi after RNAi in SW480 cells. A) Table showing the levels of BRACHYURY mRNA until 6 days after RNAi. This table shows that the BRACHYURY RNAi levels at day 5 and 6 after RNAi is less than that of days 1-4. B) Graphical representation table A showing high knock-down levels on days 1-4 of RNAi, but low RNAi levels on day 5 and 6 after RNAi. The X-axis mentions BRACHYURY levels are analysed for days 1-6 after RNAi by qRT-PCR under different RNAi conditions. Y-axis gives the relative *BRACHYURY* expression levels. Raw data is in appendix **Table 20**; N=3, in triplicate. All data were compared to  $\beta$ -catenin. p-values obtained for the fold reduction of BRACHYURY RNAi are 0.03 (Day 1), 0.00049 (Day 2), 0.0046 (Day 3), 0.05 (Day 4), 0.173 (Day 5), 0.088 (Day 6).

# e) Role of BRACHYURY in cell migration

#### 4.5: Role of BRACHYURY in cell migration:

Cancer cells become metastatic and move from primary to secondary tumour sites by the process of cell migration. Many signalling molecules are involved in this process. *CDX2* is a homeodomain transcription factor that regulates intestinal differentiation: it is a key intestinal gene down-regulated at the invasive cells of colon cancer (Brabletz *et al.*, 2004) and is also reduced during wound healing. *CDX2* has been shown to prevent cell migration and dissemination in colon cancer, both *in vitro* and *in vivo* (Gross *et al.*, 2008). In human COLO 205 cancer cells, *CDX2* expression is shown to induce E-CADHERIN activity and also increase the cell-cell adhesion (Keller *et al.*, 2004). Over-expression of *CDX2* along with *CDX1* in human colon cancer cell line HT29 (which is low in CDX expressions) showed a reduction of tumourigenic potential and migration; and an increase in sensitivity to apoptosis (Soubeyran *et al.*, 2000). This might show that the loss of expression of *CDX1* and *CDX2* might be related to the development and virulence of colorectal tumours (Soubeyran *et al.*, 2000).

Since, our data shows significant reduction of *CDX2* expression in response to *BRACHYURY* RNAi, we investigated a possible role of *BRACHYURY* in SW480 cell migration.

We used Cytoselect 24 well cell migration assay with 8µm pore size (Cell Biolabs, Cat. Number CBA-101) to analyse this experiment (Materials and Methods 2.23). According to the principle of this assay, the cells which have migrated will pass through a polycarbonate membrane and remain attached to the bottom side of the membrane until detached in cell detachment solution. After adding dye to the cell detachment solution containing the cells, the luminescence is then read at 420/ 530 nm wavelength.

We performed this assay on SW480 cells treated with MitoMYCin-C (to stop cell division and thus to synchronise the cell cycle). We analysed the cells

after *BRACHYURY* knock-down and compared it to the controls at time points 12 hours and 24 hours after RNAi. Our data show (Figure 4.18) that the number of migratory cells decrease over the time span, confirming our data obtained from the scratch wound assay. Also, there was a decrease in migration of SW480 cells treated with *BRACHYURY* RNAi after 12 hours of the treatment, compared to untreated and non-interfering controls. Also, after 24 hours, the number of migratory cells decreased in cells treated with *BRACHYURY* RNAi but this decrease was less compared to 12 hour time-point.

Only two repeats were done for this experiment and *BRACHYURY* knock-down levels were analysed for two sets of data (mentioned as trial 1 and trial 2 in Figure 4.18A). Trial 1 had *BRACHYURY* knock-down 43% for 12 hour time-point (Figure 4.19) and 47% for 24 hour time-point (Figure 4.19). Trial 2 had *BRACHYURY* knock-down to be 66% for 12 hour time point (Figure 4.19). and 62% for 24 hour time point (Figure 4.19) confirming the knock-down being active after 24 hours. Trial 1 (Figure 4.18) did not show any reduction in cell migration after on knocking down *BRACHYURY* at 12 hour time-point whereas trial 2 gave 44% reduction (Figure 4.18) of cell migration in SW480 cells treated with *BRACHYURY* siRNA, at a similar time-point (12 hour). At 24 hour time point, both trial 1 and 2 showed a small decrease in cell migration (~20%) on *BRACHYURY* knocked down SW480 cells.

An average of the two trials was calculated and Figure 4.18B was plotted based on the average values. At 12 hour time point (Figure 4.18B), since the error bar of the RNAi sample overlaps with that of the controls, statistically, this is not a significant difference although biologically, we did not explore its significance. This result was affected by the fact that trial 1 showed no decrease in migration of the cells on treating them with *BRACHYURY* siRNA. This might be due to the lower rate of *BRACHYURY* knock-down (43%) in trial 1.

At 24 hour time point (Figure 4.18B), since the error bar of the RNAi does not overlaps with that of the controls, statistically, this might be of significance, but we need more repeats of the experiment to conclude this result.

RFU (Arbitary Luminescence units)→ RNAI conditions	12 hours (trial 1 RFU)	12 hours (trial 2 RFU)	Average RFU (12 hours)	24 hours (trial 1 RFU)	24 hours (trial 2 RFU)	Average RFU (24 hours)
Untreated	10544	12100	11322	7508	7840	7674
Non-interfering	11442	11985	11713.5	7562	7825	7693.5
<i>BRACHYURY</i> RNAi	11733	7837	9785	6439	6849	6644





#### Figure 4.18

Cell migration assay using Cytoselect cell migration assay in SW480 cells. A) Table showing the cell migration rate decreases from 12 hours (blue) to 24 hours (pink) after plating the cells. This table gives the raw RFU values of the migration assay (after BRACHYURY RNAi) for trials 1 and 2, obtained from the luminometer. This table also gives the average of the RFU values obtained for trial 1 and trial 2 for each time point, which is then plotted on the graph B. On knocking down *BRACHYURY*, the rate of cell migration decreases, both 12 and 24 hours after RNAi. The X-axis mentions the time points and the different RNAi conditions, for which the migration assay was analysed. Y-axis gives the average RFU values (migration assay count). RFU = Relative Fluorescence Units. B) Graph depicting data from table A, showing lower cell migration rate on knocking down *BRACHYURY* in SW480 cells at time points 12h and 24h after RNAi. N=2, in triplicate.



**Figure 4.19:** BRACHYURY RNAi levels in SW480 cells for migration assay. A) This table shows that relative *BRACHYURY* expression on knocking down *BRACHYURY* at time points 12h and 24h. Graphs B-D gives the graphical representation of table A. X-axis=time points and RNAi conditions; Y axis=relative *BRACHYURY* expression. All data were compared to  $\beta$ -catenin. N=2, done in triplicate. Raw data for this figure can be obtained from Table 21 of appendix.

137

[Note: We did a cell count of the medium in which the inserts, containing the cells, were dipped in to confirm whether the migratory cells have crossed the membrane and moved to the solution instead of getting attached to the membrane. Since, we did not find any cells in the medium solution; we confirm that the migratory cells were only found at the membrane bottom and not in the medium].

Since, this data is preliminary; more repeats of this experiment should be performed to obtain p-value and the error bars confirming the significance of the data, which will depict a proper scenario of the role of *BRACHYURY* in cell migration.

#### **Discussion:**

*BRACHYURY* was originally shown to have a central role in mouse mesoderm development but its expression has also been shown in chordomas and in the embryonal carcinoma (EC) cell line NTERA2, where it is expressed in the absence of mesodermal differentiation. However, a role of *BRACHYURY* in cancer has not been determined.

In this chapter, we have explored possible roles of *BRACHYURY* in SW480 cells. We have shown that the expression of *BRACHYURY* influences expression of the pluripotency gene *NANOG*, and also the levels of markers of CSCs in low density SW480 cells. We also show that the oncogene  $\beta$ -catenin affects the levels of *BRACHYURY* expression, suggesting a possible role of BRACHYURY in the transducing the  $\beta$ -catenin signaling pathway and in the maintenance of cells with a CSC-like phenotype in SW480 cells.

 $\beta$ -catenin is a modulator of signaling pathways that regulate 'stemness' and also been shown to be required to maintain a population of CSCs in epidermal tumours. We show a possible mechanism whereby  $\beta$ -catenin signals

through *BRACHYURY* to maintain a population of cells potentially having stemlike character is determined by expression of *NANOG*.

NANOG is not absolutely required for stem cell self-renewal, but it increases the efficiency at which the process of self-renewal can be carried out. High levels of NANOG can translate to an efficient self-renewal and low levels of NANOG lead to an increased chance of differentiation. From our data, we show that SW480 cells at a high density, epithelial-like state, display BRACHYURY and NANOG positive cells very rarely. From the limited published studies to date, it is possible that not all tumours/cell lines express BRACHYURY. Those tumours that do not express BRACHYURY might have evolved an alternative route to activate NANOG which is then important in establishing cells bearing characteristics of CSCs. Also, BRACHYURY negative tumours might also be less aggressive. It would be interesting to determine the localisation pattern of NANOG and BRACHYURY within a larger panel of colorectal and other cancer tumours and to correlate such data with the presence of CSCs and patient outcomes. Chapter 5 - The binding of BRACHYURY to the *NANOG* promoter region is dependent on β-catenin in SW480 cells

#### **5.1 Introduction**

The T box gene family has been identified in Drosophila, mouse, human, Xenopus, sea urchin, chick, zebrafish C. elegans and the ascidian genomes (reviewed by Papaioannou et al., 1998). These genes have functional roles in early development during organogenesis (such as development of heart and limb, specification of mesoderm, formation of the vertebrae (Papaioannou and Silver, 1998; Smith, 1999; Kavka and Green, 1997) and are also involved in many developmental syndromes in humans (Li Q.Y. et al. 1997, Basson et al., 1997; Bamshad et al., 1997). Brachyury or T, is the founding member of the family of Tbox transcription factors and has been the most intensively studied. This gene encodes a sequence-specific DNA-binding protein which acts as a transcriptional activator (Conlon et al., 1996; Herrmann et al., 1990; Kispert and Herrmann, 1993; Kispert et al., 1995a). Brachyury is expressed throughout development of the mesoderm of mouse, chick, Xenopus and zebrafish embryo and then becomes restricted to the notochord development. (Kispert et al., 1995b; Schulte-Merker et al., 1992; Smith et al., 1991; Wilkinson et al., 1990). In the mouse, Brachyury has been shown to function in mesoderm development where genetic mutation leads to loss of posterior mesoderm formation and aberrant notochord differentiation (Chesley, 1935; Gluecksohn-Schoenheimer, 1938; Herrmann et al., 1990; Halpern et al., 1993; Schulte-Merker et al., 1994; Conlon et al., 1996). The structure, function and expression of Brachyury are highly conserved in mouse, zebrafish, Drosophila and human (reviewed in Papaioannou et al., 1998).

In 1993, Kispert and Herrmann isolated DNA fragments that bind Brachyury protein based on the binding site selection method. The core sequence required for Brachyury binding was found to be an 8 nucleotide sequence, TCACACCT. However, this can be extended either side to 10 nucleotides sequence by adding nucleotides to either sides of it. This can either form a palindrome or an inverted repeat (Kispert and Hermann, 1993; Papaioannau and Silver 1998; Conlon *et al.*, 2001).

The DNA binding ability of Brachyury is confined to the large N-terminal domain of 229 amino acid residues (Busch and Sassone-Corsi, 1990). This domain does not have similarity to any other DNA binding motifs (Harrison, 1991) and is highly conserved between the mouse T protein, and its *Xenopus* (Smith *et al.*, 1991) and zebrafish (Schulte-Merker *et al.*, 1991) homologues (Figure 5.1).

Putative Brachyury (GGGACACACCTAGGGTTCCC) and STAT-3 (TTCCTAGAA) binding sites were reported in the mouse *Nanog* promoter region, 4.91 kb upstream of the translation start site (Suzuki *et al.*, 2006). The Brachyury binding site was identified as a 20-bp sequence which formed an imperfect palindrome and has homology with the proposed Brachyury binding site for *Xenopus*. Further, a STAT3 binding site was also reported, 44 bp upstream of the T-binding site. Reporter assays suggested that the up-regulation of *Nanog* is governed by functional T and STAT3 sites upstream of the mouse *Nanog* gene which act as enhancer elements. When one or both STAT3 and T sites were mutated, the enhancer activity for *Nanog* transcription was lost. T was also found to associate with STAT3 only when the STAT3 translocation was stimulated by LIF (Kispert and Hermann, 1993; Papaioannau and Silver 1998; Conlon *et al.*, 2001).

As discussed above, *Nanog* has been shown to be a direct target of Brachyury binding in the mouse ES cells, and our data (shown in results chapter 4) suggests that *NANOG* might be a target of *BRACHYURY* in human CRC cells (SW480). To determine whether the regulation of *NANOG* by BRACHYURY might involve direct binding of BRACHYURY to the *NANOG* promoter, we carried out ChIP in the upstream regulatory region of the human *NANOG* gene in SW480 cells. In order to carry this out, it was necessary to search for the human *NANOG* upstream regulatory region (chromosome 12) in the *NANOG* contig region (gene sequence NM\_024865) on chromosome 12p13.31 for potential BRACHYURY binding sites. There are also 10-11 pseudogenes for *NANOG* in the human genome, some of which are also situated on chromosome 12. It was therefore important to carefully select the functional *NANOG* gene in human in place of any of the *NANOG* pseudogenes.



#### Figure 5.1

Brachyury (T) is conserved in human, mouse, *Xenopus* and zebrafish. This figure highlights the amino acids in the DNA binding region of Brachyury; *Xenopus Brachyury* (Xbra) amino acid sequence, aligned with the T box in human, mouse, *Xenopus* and zebrafish Brachyury. Yellow circles denote amino acids that contact DNA according to Muller and Herrmann, 1997. Lysine 149 is the amino acid which is associated with T box specificity. (Image taken from Conlon *et al.*, 2001).

#### **Results:**

Our experimental approach using ChIP to identify binding sites for Brachyury in the human *NANOG* promoter is described in the following sections.

# 5.2 Identification of BRACHYURY binding sites in the human *NANOG* promoter region:

BRACHYURY has been shown to bind to enhancer elements in the Nanog promoter region and to up-regulate the expression of Nanog in mouse ES cells in association with STAT3 (Suzuki et al., 2006). We wanted to investigate whether NANOG might similarly be regulated by BRACHYURY in SW480 cells. We therefore searched for putative T-binding sites in the sequences upstream of the human NANOG gene. The functional human NANOG gene is located on chromosome 12 (position 12p13.31) and contained in the contig NT 009714.17. We used this sequence to carry out our search for putative BRACHYURY binding sites. This search was based on the DNA consensus sequence of 8 nucleotides representing <sup>5</sup> TCACACCT<sup>3</sup> the core T-box consensus sequence; and its possible palindromes or inverted repeats (Kispert & Hermann, 1993). We found 8 possible sites for both <sup>5</sup>'AGGTGTGAA<sup>3</sup>' and <sup>5</sup>'TCACACCT<sup>3</sup>' sequences representing core binding sequences for BRACHYURY (refer to table 5.1 A and B). Out of these sites, we selected two putative T-box binding sequences, since these were arranged as inverted repeats; site 1, a 19 bp inverted repeat (-708 bp relative to the start codon of NANOG) and site 2, a 15 bp inverted repeat sequence (-2335 bp relative to the start codon of NANOG) (Figure 5.3A). These two sites were chosen based on the length of inverted repeat and proximity to NANOG translation start. [Note: The size of promoter region used was 2331 bp, starting from the BRACHYURY binding site upto the NANOG translation start site. This region of the NANOG promoter contained the two putative BRACHYURY binding sites. Refer section 5.3B].

## 5.3A: BRACHYURY binds to the NANOG promoter in human SW480 cells:

Following our identification of two putative BRACHYURY binding sites in

# A.

# BRACHYURY binding sequence 1: <sup>5'</sup><u>AGGTGTGAA</u><sup>3'</sup>

Positions/base site in the <i>NANOG</i> promoter region	BRACHYURY binding sequence	Inverted repeats	Palindrome sequence	No. of bases for palindrome
*705591	<u>AGGTGTGA</u> A	705604	CACACC	6
673181	AGGTGTGA	673151	CACCT	5
		673191	CACC	4
667152	AGGTGTGA	667162	CACCT	5
652812	AGGTGTGA	652782	CACCT	5
616613	AGGTGTGA	616579	CACCT	5
608761	AGGTGTGA	608731	CACCT	5
581831	AGGTGTGA	581846	CACCT	5
		581846	CACCT	5

#### B.

# BRACHYURY binding sequence 2: <sup>5'</sup><u>TCACACCT</u><sup>3</sup>

Positions/base site in the <i>NANOG</i> promoter region	Sequence	Inverted repeats	Palindrome sequence	No. of bases for palindrome
*703971	TCACACCT	703964	AGGTG	5
667771	TCACACCT			
662722	TCACACCT	662761	AGGTG	5
648293	TCACACCT			
598483	TCACACCT	598471	AGGTG	5
595553	TCACACCT			
575713	TCACACCT	575701	AGGTGTGA	8
572012	TCACACCT	575207	AGGTG	5

#### Table 5.1

Potential BRACHYURY binding sites in the promoter region of the human *NANOG* gene. Locations of possible BRACHYURY binding sites for the sequence A)  $5^{\circ}$ <u>AGGTGTGA</u>A<sup>3'</sup> and B)  $5^{\circ}$ <u>TCACACCT</u><sup>3</sup> identified in the upstream region of *NANOG* promoter region in the contig; and potential inverted repeat sequences in close proximity to these sites. Two sites at positions 705591 and 703971 were chosen due to their proximity to the *NANOG* gene and their having potential inverted repeat sequences. These two sites are marked with \*. Numbers refer to the positions on the *NANOG* contig region (Contig reference: NM\_024865, NCBI). described in section 5.3B.

relative close proximity to the *NANOG* promoter, we carried out ChIP (<u>Ch</u>romatin <u>I</u>mmunoprecipitation) using antibody to BRACHYURY (T), in low density and high density SW480 cells to determine whether these putative binding sites could act as binding sites for endogenous BRACHYURY. The ChIP protocol was followed as We optimized conditions for this as follows.

- Sonication conditions were optimized to 5 repeats of 10 seconds each. These conditions were chosen in order to obtain genomic DNA products ranging in size between 200-1000bp (Figure 5.2A).
- We also carried out IP for BRACHYURY with BRACHYURY antibody to optimize antibody concentration (IP protocol described in M & M section 2.16). This IP gave a very strong product for cell lysate derived from SW480 subconfluent cells, slightly weaker binding with SW480 confluent lysates and almost no binding with supeconfluent cells (Figure 5.2B) confirming our qRT-PCR data (chapter 3) that Brachyury is not abundant in superconfluent SW480 cells.

The ChIP assay was carried out on cell lysates from sub- to superconfluent SW480 cells. Specific binding was observed at both sites 1 and 2, although binding at site 1 may be slightly more efficient as indicated by slightly stronger PCR bands (Figure 5.3B); BRACHYURY binding at the two sites was observed only with lysates from day 1 (low density SW480 cells) and day 2, weaker binding (mid-density cells), and no binding was observed at day 4 (high density cells). The PCR products for the input DNA and ChIP, at sites 1 and 2, were confirmed as being the human *NANOG* promoter region by sequencing. We also included control primers in a ChIP reaction to a region in the *APC* gene (exon 15) [See M & M table 2.9 for APC primer sequence]. Sequencing confirmed this PCR product as *APC* and it did not yield any product following ChIP with BRACHYURY antibody, confirming that no non-specific product was pulled down in the BRACHYURY IP reaction.



#### Figure 5.2

Optimization of ChIP conditions. A) Optimization of sonication conditions with four different pulses and duration of sonication. The number of pulses and the time of sonication are mentioned below each lane in the image. B) The amount of BRACHYURY antibody was optimized using SW480 cell lysates from Day1 (sub-confluent), Day 3 (confluent) and Day 4 (super-confluent) confirming strongest binding on day 1 SW480 lysates and the weakest on day 4. The IgG was used as control without BRACHYURY antibody to confirm specific antibody binding.

#### 5.3B: NANOG promoter sequence for BRACHYURY binding (human)

caatacgtga aaattggtga atctccattc tgacctatta acacccagtg tgggagcttt tgagcatcag tttaaagcaa aaggtggtag cgccggcctg atgaggtggc tcacacctgt aatctcagta cttcgggagg ccgaggcagg cggatcacga ggtcaggagt tcgagaccag ceteaccaac attgagaaac cetgteteta etaaaaatae aaaattagee aggegtgatg gtgcctgcct gtaatcccag ctactcagga ggctgaggca ggagaatcgc ttgaacccag gagatggagg ttgcagtgag ccgagattgc accatcgcac ttcagcctgg gcaacaagag tgaaceteca tectaatttt aaaaaaaagg ggtagegeeg etcageaatg cattggeeae cattatagat ctctcccaac gcagtctatt atgtacaaaa tggagatact gataagactt cttggagtaa tactatataa agcttgccaa agtgccaggg ctgcttaaga aattgcttct tattacttag atctggggtt ctgggaatta tcaaagtact ttgaaaacaa tttttttaaa ggatatttta atatttgaaa aattttagac aaaagtgtcc ttttatttgt tcccaacagt ctctcctctt ccttcctcca tggatctgct tattcaggac agccctgatt cttccaccag tcccaaaggc aaacaaccca cttctgcaga gaagagtgtc gcaaaaaagg aagacaaggt cccggtcaag aaacagaaga ccagaactgt gttctcttcc acccagctgt gtgtactcaa tgatagattt cagagacaga aatacctcag cctccagcag atgcaagaac tctccaacat cctgaacctc agctacaaac aggtaggctt gttttgtcct tggaataagg tgaacaaaaa ttggactaat ttgcatggct aagacctctg tggatgcatg tagatgtgtg tactatgtgt ccgtacatcg cctcttgcaa ataatttatg aagatgaaat gcttttgtaa cttccttcac ctctttcttt ccttaaatat tctattatgt gaataattat gtcataattt aaccactttc ttgcacagac caatattgtg aaaatctttc caacgtttcc ttaataaaaa agaagtatct aactccactt accagggtag gagaaaccct aactcacact ggttctcatt aaaataaaaa gtgcattcaa taaggccaat cattgtttat tacttgttgg ccattttcct tcattaattg gettateete ttetaggett tattteette ttttettttg tttatttttg agaeggagte tcactctgtc tcccaggetg gagtgcagtg gcctcataat gagacatcat aatgacatta gcattacttg gtttaagtac aaatggatga aacctccacc tcctgggttc aagggattct cctgcctcag cctcccaagt agctgggatt acaggcgccc gccactacac ctggctaatt ttactgtatt ttttagtaga gatggggctt caccatgttg gccaggctgg ttttcaactc ctgacctcaa gtgatccacg tgcctctgcc tcctaaagtg ctaggattac aggtgtgaac taccacaccc ggcccttgcc tttattttct atccaaaatg ttcatttctc tctttttttc tgagacagtc tggttetgtt gtcacgcagc tggagtgcag tgatgcaatc ttggctcact gcagcettea etteccaggt gcaagegate etcecgtete ageeteetga gaagetggga ccacagacgt gcagcattac acceggetaa ttttttgtat ttttgtagac acggtgtttc ctcatgttgg ctagcctgta gcgaactcct gggctcaagc aatctgcccg ccttggcctt ccaaagtgtt gggattacag gcatgagata tggcacctgg ccaggaataa aagttagcaa tatactttga ttcaaagtac ctctgtatta tgaatatttt acaatttcta tcattttttc ctgcaggtga agacctggtt ccagaaccag agaatgaaat ctaagaggtg gcagaaaaac aactggccga agaatagcaa tggtgtgacg caggtaacag gaaacttcat tctgttcttt cetttcagtg atetttcaat ettgtccate eetgaaacae acaactecag tcacagacag ttctggttgt ccttgtaccc tttctgttaa tccctccttc tctttcagaa ggcctcagca cctacctacc ccagcettta etettectac caccagggat geetggtgaa ecegaetggg aaccttccaa tgtggagcaa ccagacctgg aacaattcaa cctggagcaa ccagacccag

Nanog translational site: atg

Two putative Brachyury binding sites: tcacacct and aggtgtgaa

L2a: attcca ggtac **gtgtgggagcttttgagcatc** L2C: attcca ggtac **gtgtgggagcttttgagcag** cctc L0F: attcca ggtac **gagacagtctggttctgttgtcacge** Reverse primer, LR1: attcca gagctc **ctggtggtaggaagagtaaagg** Construct 1(no sites) = L0F + LR1 = 635 bp Construct 2(1 sites) = L2C + LR1 = 2275 bp Construct 3(both sites) = L2a + LR1 = 2389 bp Size of the NANOG promoter = 2331 bp (from the BRACHYURY binding site, tcacacct to NANOG start site, atg)

#### 5.4 BRACHYURY binding to the NANOG promoter depends on β-catenin:

BRACHYURY has been shown to be a downstream target of Wnt 3a during mesoderm specification in the mouse embryo (Terry et al., 1999). Since, the Wnt signalling pathway is critically altered in CRC and  $\beta$ -catenin is the key oncogene of this pathway, we wanted to know if the binding of BRACHYURY to sequences in the NANOG promoter region was dependent on signaling through  $\beta$ -catenin. We therefore repeated the ChIP, but in the presence of interfering RNA to  $\beta$ -catenin; to knockdown  $\beta$ -catenin in the SW480 cells. Again, specific binding of BRACHYURY at both sites 1 and 2 was observed for day 1 and 2 in untreated and non-interfering control samples, but no binding was observed for samples treated with siRNA to knockdown  $\beta$ -catenin in day1 and 2 samples. None of the samples at day 4 (superconfluent SW480 cells) displayed binding of BRACHYURY to either binding site under any of the conditions used to knockdown  $\beta$ -catenin (control or experimental) (Figure 5.3C). These results are consistent with the expression patterns for BRACHYURY and NANOG observed previously, i.e. strong expression at day 1 (sub-confluent cells) and rare expression at day 4 (super-confluent cells); and demonstrate that binding sites for BRACHYURY exist within the NANOG promoter in human CRC cells and that BRACHYURY binding to these sites is dependent on β-catenin.

In summary, we demonstrated the presence of two putative T binding sites (19bp and 15bp long) at 708bp and 2335bp upstream of the *NANOG* promoter in human CRC cells. We were not able to find a putative STAT binding site near these



#### Figure 5.3

ChIP assay results for BRACHYURY. A) The putative two BRACHYURY binding sites in the upstream of human *NANOG* promoter with nucleotide base pair positions in relation to the promoter (+1). B) ChIP assay showing BRACHYURY binding to *NANOG* promoter in sub-confluent (Day 1), confluent (Day 2) and super-confluent (Day 4) SW480 cell lysates. C) ChIP with knocking down  $\beta$ -catenin indicating that  $\beta$ -catenin is essential for this binding of BRACHYURY to *NANOG* enhancer site in human. Site 1 and Site 2 are the two putative BRACHYURY binding sites in the *NANOG* promoter region. Abbreviations: U=untreated SW480 cells, N=non-interfering RNAi,  $\beta = \beta$ -catenin RNAi. Input DNA is the qRT-PCR product without IP. IgG=immunoglobulin G as a control for the IP.

sequences. The ChIP assay revealed that BRACHYURY can bind to these sites in the human *NANOG* enhancer region with one site apparently being slightly weaker for binding than the other. The binding was shown to be the strongest in subconfluent SW480 cell lysates at day 1 and this affinity becomes weaker and nondetectable as the cells reached higher density (input DNA was used as a control with no antisera directed to BRACHYURY). This correlates to our data presented in chapter 3 where the amount of BRACHYURY was shown to be high in SW480 subconfluent cells and lowest in the super-confluent cells. Our ChIP assay with  $\beta$ *catenin* knockdown confirmed that  $\beta$ -catenin is essential for this binding of BRACHYURY to the *NANOG* enhancer region in human and acts upstream of BRACHYURY.

# 5.5 BRACHYURY binding sites act as BRACHYURY -dependent enhancers of expression from the *NANOG* promoter:

Cloning, transfection and luciferase assay

So far, in this chapter, we have shown that BRACHYURY binds to sites within *NANOG* promoter and this is dependent on  $\beta$ -catenin. We next wanted to determine if these sites were functional i.e. whether binding of BRACHYURY to these sites could influence transcription. We made three constructs with and without these sites along with a construct with none of the sites for control. Our constructs (Figure 5.5) were as follows:

- Construct 1 = no BRACHYURY binding site
- Construct 2 = containing site 1
- Construct 3 = containing both sites 1 and 2

The PGL3-Basic vector used to make the constructs (1-3) is illustrated on Figure 5.4 with KpnI and SacI restriction sites in the multiple cloning site of the vector. The constructs were made by cloning the DNA into PGL3-Basic vector (4818bp) (Figure 5.4). Restriction enzymes KpnI and SacI were used for restriction digest since these two sites are present in the multiple cloning site of the PGL3 basic vector and do not within cut the *NANOG* regulatory sequence being studied. [Refer section



### Figure 5.4

PGL3 basic vector map used for cloning. The multiple cloning site contains the KpnI and SacI sites. The size of the vector is 4818bp.

2.19 of Materials and Methods for the cloning procedure]. The three constructs were co-transfected into SW480 cells in turn (transfection protocol- Qiagen, refer to section 2.19 of Materials and Methods) alongside a renilla contruct that contains renilla luciferin (pGL4.73 [hRluc/SV40]) under the control of the SV40 promoter. SW480 cells were grown for 24 hrs at 37°C. After 24 hrs, a luciferase assay (according to Promega protocol listed in M & M section 2.19) was performed and the intensities were recorded from the luminometer. The luminescence was then stopped by adding Stop & Glo reagent (E2920, Promega Dual-Glo Luciferase Assay kit) and a separate assay was carried out on the same cells to determine the luminescence intensity of the Renilla using the Promega Dual-Glo Luciferase Assay System protocol. The data was normalized to luminescence obtained from the Renilla vector to account for variation in transfection efficiencies. The experiment was controlled by including untreated cells, mock-treated cells (only transfection reagent and no DNA) and cells with transfection reagent and PGL3 basic vector only. Construct 1 contained 645 bp of sequence upstream of NANOG (relative to translation start site) with no BRACHYURY binding sites; construct 2 contained 2285 bp upstream of the start site and 1 BRACHYURY binding site (site 1); construct 3 contained 2399 bp of sequence upstream of the start site, and 2 BRACHYURY binding sites (site 1 and 2). Transient transfection of these constructs in low density SW480 cell showed an increase in the luciferase activity of construct 2 by 1.2 fold relative to control (promoterless luciferase construct, PGL3-Basic), and a 2.3 fold increase in luciferase activity (relative to a promoterless control, PGL3-Basic) for construct 3 that contained both the BRACHYURY binding sites (Figure 5.5A). The p-values showed that this increase in the luciferase activities of constructs 2 (p-value = 0.0008) and 3 (p-value = 0.0269) are statistically significant. The experiment was controlled by using cells with no transfection treated (untreated) and cells with only Superfect transfection reagent (Mock-treated) which gave similar luciferase activities to that of PGL3-Basic (which contains only PGL3-Basic vector and no DNA) (Figure 5.5A). The luminescence obtained from PGL3-Basic was adjusted to 1 and the luciferase activities of contructs (1-3) were compared to PGL3-



#### Figure 5.5

Luciferase assay. A) Image showing luciferase activity of construct 2 and 3 increased compared to cells with PGL3-Basic vector as a control. N=5 (for Pgl3-Basic and Construct 3); N= 4 (for constructs 1 and 2). P-values for final F/R value of Construct 1 compared to that of PGL3-Basic construct = 0.3394 (>0.05), p-value for Construct 2 compared to PGL3-Basic construct = 0.0008 (<0.05), p-value for Construct 3 compared to PGL3-Basic construct = 0.0269 (<0.05). B) Luciferase activity of constructs 2 and 3 in response to BRACHYURY RNAi decreases. The final F/R values for constructs 2 and 3 without RNAi were plotted on the graph against those of constructs 2 and 3 with RNAi . N=3. SDs gave the error bars. Knocking down the BRACHYURY reduces the luciferase activity of the Construct 2 by almost 19% (comparing 1.79 and 1.19) and that of the Construct 3 by 60% (comparing 0.961 and 0.77). The p-value obtained by comparing the final F/R values from the three trials of Construct 2 with RNAi to that of Construct 2 without RNAi was found to be 0.1362 (>0.05). The p-value obtained by comparing the final F/R values from the three trials of Construct 3 with RNAi to that of Construct 3 without RNAi was found to be 0.0012 (<0.05). '1' and '2' on the contructs mention the site 1 and 2 for BRACHYURY binding on Nanog promoter.

Basic luminescence. Further, we analyzed whether knocking down *BRACHYURY* in cells transfected with constructs 2 and 3 would show changes in the luciferase activities of the constructs (Figure 5.5B). This would tell us how efficient is the binding of BRACHYURY to these putative sites in *NANOG* promoter region.

Our knock-down data was plotted along with non-knockdown data to analyze the luciferase activities of the constructs 2 and 3. Our data show that knocking down the gene expression of *BRACHYURY* reduces the luciferase activity of the construct 2 by almost 19% and that of the construct 3 by 60%. The p-values for these data show that reduction in luciferase activity of construct 2 were statistically not significant (p-value = 0.1362), but that of construct 3 is significant (p-value = 0.0012). This shows that BRACHYURY can bind to the two putative BRACHYURY binding sites in the *NANOG* promoter, but binding to site 1 is less efficient than binding to site 2. [Raw data for Figure 5.5A in appendix Table 22A-D and for Figure 5.5B in Table 23A -D].

#### 5.6 Discussion:

The ChIP assay in this chapter shows a direct binding of BRACHYURY to putative specific regulatory elements in the *NANOG* promoter region. The ability of BRACHYURY to bind to these sites was removed on knocking down  $\beta$ -catenin demonstrating a dependence of binding upon  $\beta$ -catenin. In mouse ES cells, the binding of Brachyury to sites in the *Nanog* promoter was co-ordinate with the binding of STAT3 (Suzuki *et al.*, 2006). However, we did not find any STAT3 binding sites close to either of the T-binding sites that we located in the promoter region of the human *NANOG* gene. It has been shown that signaling through STAT3 does not promote self-renewal in human ES cells in culture (Daheron *et al.*, 2004).  $\beta$ -catenin signaling increases *Nanog* promoter activity in a Tcf3-independent manner in mouse ES cells suggesting that a complex control mechanism exists in mouse ES cells (Pereira *et al.*, 2006). Again, this is consistent with our data (chapter 4) where  $\beta$ -catenin acts to positively regulate the binding of BRACHYURY to the promoter region of *NANOG* in SW480 cells.

The luciferase assay data presented in this chapter indicates that expression levels from the luciferase construct are low when a 701bp *NANOG* upstream element is inserted (construct 2): these levels appear to be close to the control levels, suggesting that the promoter elements contained in the sequence are not efficient under the conditions used in this experiment. Construct 3 (with both BRACHYURY binding sites) appears to give an increase in the transcription activity that is 2.3 fold compared to controls, suggesting that enhancer activity may be contained in a 114bp sequence element (2399bp to 2285bp) that contains the second BRACHYURY binding site. Whilst ChIP data showed that BRACHYURY binds to both sites, it is likely that one site alone is insufficient to enhance transcription.

Removing the sites was a proper experiment preliminary. In future, replacing with random sequence and then mutating specific sequences with real sites will be performed. Future studies would involve more detailed analysis of these binding sites and their composition. Mutation analysis of the sites and further studies on the sequence elements upstream of the *NANOG* promoter will also be beneficial. Also, preparing constructs with site 2 only, and not site 1 will further confirm our results. Other constructs for this analysis would involve deletion or mutation of bases inbetween the two BRACHYURY binding sites.

# Chapter 6 - The role of *BRACHYURY* in NTERA-2, SW626 and T84 cells

In chapter 4, we have demonstrated a role for the mesoderm inducing gene, *BRACHYURY* in regulating the pluripotency gene *NANOG* and levels of the CSC markers CD166 and CD133 in the human colon cancer cell line, SW480. We next wanted to determine whether these results were specific to the CRC cell line SW480, or whether they could be applied more generally to other cell lines. From our results obtained, we hypothesized that this pathway may be initiated/ active in cell lines that were competent for differentiation along a mesodermal lineage. Therefore, we looked at this pathway in the germ cell tumour cell line, NTERA2.

#### 6.1 Role of BRACHYURY in NTERA-2 cells

NTERA-2 is an embryonal carcinoma (EC) cell line obtained from the lung metastasis of a testicular teratocarcinoma (Fogh & Trempe, 1975; Andrews *et al.*, 1984). It is a sub-clone obtained from TERA-2 which was derived by culturing embryonal carcinoma (EC) cells from the lung metastasis of testicular cancer (Andrews *et al.*, 1984).

The EC cells behave as pluripotent stem cells and give rise to all other cell types in the tumour. These cells may originate from a displaced embryonic stem (ES) cell which might have retained pluripotency into postnatal life. Alternatively, they may originate from germ cells which may proliferate without entering into meiosis.

We grew NTERA-2 cells to 70-80% confluency, collected cell lysates and assessed the levels of BRACHYURY by Western blot analysis. Once we confirmed the existence of BRACHYURY in these cells, we then knocked down *BRACHYURY* through RNAi in these cells (as described in materials and methods, section 2.17), collected the cell lysates and assessed the levels of NANOG, Oct-4 and CD166 by Western blot analysis. Our experiment shows that BRACHYURY is present in NTERA-2 cells, the knockdown of which reduces the levels of NANOG, BRACHYURY and Oct-4. We could not detect CD166 in these cells as expected since this is a specific marker for CRC-SC. Lamin B was used as a loading control confirming equal amount of protein loaded for experimental and control lanes (Figure 6.1B). Like NANOG, Oct-4 is another determinant of pluripotency (Pesce *et al.*, 2001; Nicholas *et al.*, 1998) and is known to be expressed in undifferentiated NTera2 cells (Rosner *et al.*, 1990). We were intrigued to see that Oct-4 was also regulated by BRACHYURY. Oct-4 has been shown to be expressed in some cancers such as bladder cancer (Xu *et al.*, 2007), human non-small cell lung cancer (Karnoubi *et al.*, 2009) and many other cancer (Tai *et al.*, 2005; Wang *et al.*, 2003; Monk *et al.*, 2001; Gidekel *et al.*, 2003; Looijenga *et al.*, 2003).

Thus, we have shown that *BRACHYURY* knockdown in NTERA2 cells decreased the levels of NANOG in similar manner to that observed in SW480 cells (Figure 6.1A). Based on our results from colon cancer cell line, SW480 and EC cell line NT2, we then checked the role of BRACHYURY in the colon cancer cell line T84 and the ovarian cancer cell line, SW626 which is a metastasis of a primary colon carcinoma. The cell lines SW626 and T84 were previously reported to express *BRACHYURY* (qRT-PCR) (Palena *et al.*, 2007).

#### 6.2 Role of BRACHYURY in T84 cells

T84 is a human colon cancer cell line derived from a lung metastasis of colon carcinoma in a 72 year old male. The tumour tissue was innoculated sub-cutaneously and transplanted into nude mice. T84 are epithelial cells which grow in monolayer and have receptors for many peptide hormones and neurotransmitters.

Earlier we have shown that the expression of *BRACHYURY* is very low or almost negligible in T84 cells (assessed by qRT-PCR). To check the protein levels,



#### Figure 6.1

Expression of BRACHYURY and other proteins in SW480, NT2 and T84 cells with respect to *BRACHYURY* RNAi. A) *BRACHYURY* knockdown in SW480 subconfluent cells regulates the expression of NANOG and colon cancer SC-markers CD133, CD166 and Lamin A. B) *BRACHYURY* RNAi in NTERA-2 cells also shows decrease in levels of NANOG, Oct-4 and colon cancer SC-marker CD166. C) Levels of BRACHYURY, NANOG and CD166 in sub-confluent T84 cells were shown here to be lower than that observed in SW480 sub-confluent cells. *BRACHYURY* knockdown by RNAi regulates the levels of NANOG and CD166 in sub-confluent T84 cells. The loading was controlled by LAMIN B whose levels do not change on knockdown of *BRACHYURY*. U=untreated cells, N= cells treated with non-interfering siRNA, B=*BRACHYURY* knockdown. we performed Western blot analysis to study the levels of BRACHYURY in low and high density T84 cells. We show that BRACHYURY is present inT84 cells but not differentially expressed in low and high density cells (data not shown). We also performed *BRACHYURY* RNAi experiment in low density T84 cells to investigate the effect of BRACHYURY on the levels of NANOG and CD166 (Figure 6.1C). We show that level of NANOG and the CRC stem cell marker CD166 are decreased by knocking down the expression of *BRACHYURY*. Therefore, this regulation of NANOG and the CRC stem cell marker CD166 on *BRACHYURY* knockdown is similar to that observed in SW480 cells (Figure 6.1A). This might suggest that a population of stem cells is maintained by BRACHYURY in the T84 sub-confluent cells by regulating NANOG and CD166 in these cells.

#### 6.3 Role of BRACHYURY in SW626 cells

SW626 cells are epithelial-like cells derived from a human ovarian tumour which was a metastasis of primary adenocarcinoma of the colon (Furlong *et al.*, 1999). This cell line was first initiated in January 1974 by A.Leibovitz at the Scotland White Clinic, Texas from a surgical specimen from a cystadenocarcinoma of the ovary of a 46 year old Caucasian female (Anti Cancer Research 1998; 18:1153).

In the first instance, we wanted to determine whether SW626 cells undergo an EMT-like transition similar to that observed in SW480 cells. Therefore, we grew SW626 cells to low and high cell density and performed immunofluorescence to study the localization of  $\beta$ -catenin in these cells. [Note:  $\beta$ -catenin localization shifts from membrane to nucleus in cells undergoing EMT.] But, here, we were able to show that there is no such transition in these cells and since the localization of  $\beta$ catenin is to the membrane in low density SW626 cells and to the cytoplasm in high density SW626 cells (Figure 6.2).



## Figure 6.2

Confocal image showing localization of  $\beta$ -catenin in low and high density SW480 cells using  $\beta$ -catenin antibody.  $\beta$ -catenin (green) is shown to be localized to the cell membrane in low density SW626 cells and to the cytoplasm in high density SW626 cells. Pi staining (red) stains the nucleus of the cells. DIC image shows the cells without any staining and merged image shows the localization of  $\beta$ -catenin in the cells with respect to the Pi staining that stains the nucleus of the cells.

We did Western blot analysis in low density SW626 cells to study the levels of BRACHYURY in these cells. BRACHYURY had previously been reported to be expressed in SW626 cells (Palena *et al.*, 2007) as detected by qRT-PCR. Our data show that BRACHYURY is not expressed in SW626 sub-confluent cells, but the cells did stain positive for NANOG and Oct-4 (Figure 6.3A). The CRC stem cell marker, CD166 could not be detected in these cells by Western blot (Figure 6.3A). Since, we could not detect BRACHYURY in SW626 sub-confluent cells by Western blot analysis; we examined its expression by qRT-PCR and we did not observe a great change in the expression of *BRACHYURY* in low density SW626 cells compared to high density SW626 cells (Figure 6.3B). But the absolute levels for the Ct (thermal cycle) of BRACHYURY was found to be 37 which was similar to water control used in this experiment, which confirms negligible expression of *BRACHYURY* in sub and super-confluent SW626 cells (Figure 6.3C).

This confirms our Western blot data that SW626 cells do not express BRACHYURY (Figure 6.3A). We also checked the expressions of *E-CADHERIN*, *SNAIL*, *NANOG*, *FGF8* and mesoderm forming gene *NODAL* in SW626 cells also showed no significant changes in low density SW626 cells compared to high density SW626 cells (Figure 6.3B). But, the cell polarity markers such as *ZEB1* and *ZEB2* showed a significant decrease in expression (by 50%) in high density SW626 cells compared to low density SW626 cells (Figure 6.3B). Intestinal cell-differentiation marker *CDX2* also showed decrease in expression in high density SW626 cells (by 78%) compared to low density SW626 cells. The absolute levels for each of these genes mentioned above are depicted in Figure 6.3C which shows that there is negligible expression of *FGF8* and *NODAL* in SW626 cells whereas, *SNAIL*, *CDX2*, *NANOG* and *E-CADHERIN* are highly expressed and *ZEB1* and *ZEB2* are moderately expressed in these cells. The absolute levels of these expressions were compared to water control which gave the CT value of 39.



Genes	SW626 Sub-confluent (Relative expression)	S.D.	SW626 Super-confluent (Relative expression)	S.D.
SNAIL	1	0.12	0.97	0.22
E-CADHERIN	1	0.09	1.02	0.20
BRACHYURY	1	0.24	0.99	0.38
ZEB1	1	0.22	0.54	0.21
ZEB2	1	0.08	0.55	0.20
NANOG	1	0.08	1.14	0.17
CDX2	1	0.16	0.50	0.17
FGF8	1	0.11	0.29	0.24
NODAL	1	0.10	0.97	0.14



**Figure 6.3:** Protein levels and gene expressions in SW626 cells. A) Western blot analysis depicting the levels of BRACHYURY, NANOG, Oct-4 and CD166 in low density SW626 cells. U=untreated cells, B=BRACHYURY RNAi, N=non-interfering cells. LAMIN B was used as loading control. B) Gene expressions in low and high density SW626 determined by qRT-PCR. Here, low density SW626 cells were adjusted to 1 and the high density expression was compared relative to normalized value of 1 in low density SW626 cells. Data was compared to  $\beta$ -catenin. N=3, each done in triplicate. C) Graph demonstrating the relative gene expression levels in low and high density SW626 cells.

[Note: Though *BRACHYURY*, *NODAL* and *FGF8* are plotted on the graph, their absolute Ct values (**appendix table 24**) show that there are negligible levels of these genes in SW626 cells. (Ct value for *BRACHYURY* was 39-40; *FGF8* = 34-36; *NODAL* =37). Water control (Ct value for water control=39). Part B of this figure is obtained from **appendix table 24**]. Refer appendix Table 26 for the statistical significance data of the fold differences obtained for each gene in low and high density cells.

#### 6.4 Discussion:

In this chapter, we studied the role of BRACHYURY in various carcinoma cells lines. In human embryonal carcinoma cell line, NTERA-2 and human colonic adenocarcinoma cell line T84, NANOG is shown to be regulated by BRACHYURY in a manner to that seen in SW480 cells. Thus, it shows that the mechanism of BRACHYURY regulating NANOG is not only shown in SW480 cells but also in other forms of cancer cells. It is possible that this regulatory pathway of BRACHYURY regulating NANOG is detected in cell lines that are competent for mesodermal differentiation, such as NTERA2, whilst T84 did not really undergo EMT-like transitions which were observed in SW480 cells. We note that the levels of BRACHYURY in these T84 cells are considerably reduced compared to SW480 cells. It will be beneficial and necessary to determine the status of other pathways in these cells to try elucidating the mechanism by which this pathway operates or is important in.

# Chapter 7 - Preliminary characterization of an extracellular factor with *BRACHYURY* regulatory activity

#### 7.1 Introduction:

In chapter 3, we showed that the expression of *BRACHYURY* in low density SW480 cells is twice as higher compared to high density SW480 cells. Based on this result, we proceeded to show that the density dependent expression of *BRACHYURY* is able to bind and regulate expression from the *NANOG* promoter control region, in low density SW480 cells (chapter 4). We were interested to explore the possibility that factors controlling the levels of BRACHYURY might be secreted into the medium and as such they might be isolated and identified. To approach this, we grew the SW480 cells to a high density, collected the medium from the cells (conditioned medium, materials and methods section 2.24) and removed dead cell debris with a centrifugation step. Conditioned medium is often used for studying growth factor and cytokines secreted into the medium. This conditioned medium was then used to plate out and grow low density SW480 cells, under various conditions as described below.

#### 7.2 Results:

SW480 cells were plated out at low density ( $2X10^6$  cells/ T75) under the following four conditions:

- 100% (v/v) conditioned medium collected from SW480 cells grown at a high density (grown for 24 hours after confluency) with no additional FBS. [Note: cell count for high density cells are 2X10<sup>9</sup> cells/ T75].
- 50% (v/v) conditioned medium collected from SW480 cells grown at a high density (grown for 24 hours after confluency) with 50% (v/v) fresh DMEM supplemented with 10% (v/v) FBS. Therefore, conditioned medium:fresh medium:FBS = 1:1:0.1
- 100% (v/v) conditioned medium with fresh 10% (v/v) FBS. Therefore, conditioned medium:fresh medium = 1:1

100% (v/v) fresh DMEM with 10% (v/v) FBS (control). Therefore, fresh medium:FBS = 1:0.1

Figure 7.1A demonstrates that the levels of BRACHYURY is reduced in subconfluent SW480 cells grown in 100% (v/v) conditioned medium irrespective of adding fresh FBS demonstrating that this effect is not brought about by depletion of components in FBS. The sub-confluent cells grown in fresh DMEM (control) and the sub-confluent cells grown in 50% (v/v) conditioned medium show levels of BRACHYURY similar to control levels. This suggests that there is a dose dependent effect and only 100% (v/v) conditioned medium decreases the level of BRACHYURY in low density SW480 cells. This result was indicative of a factor that may be present in the conditioned medium (secreted or otherwise) which is responsible for the reduction in levels of BRACHYURY.

To explore this further, we next grew sub-confluent SW480 cells in 100% (v/v) and 50% (v/v) conditioned medium which had been boiled at 100°C for 5 minutes (cooled to room temperature and centrifuged to remove debris), before adding to the SW480 cells. It was expected that this would denature the majority of proteins. This experiment was controlled by growing the cells in 100% (v/v) and 50% (v/v) conditioned medium which was not boiled. Figure 7.1B shows that the expression of *BRACHYURY* is reduced even after boiling the 100% (v/v) conditioned medium. The boiled and non-boiled 50% (v/v) conditioned medium still has no effect on this activity. This suggested that the 'factor' in the conditioned medium responsible for regulating BRACHYURY is highly stable and does not get deactivated on boiling.

We have shown (chapter 5) that BRACHYURY regulates the expression of *NANOG* in low density SW480 cells and also that  $\beta$ -catenin act upstream of BRACHYURY in these cells. Thus, we explored the expression of NANOG and  $\beta$ -catenin in the low density SW480 cells grown in boiled and non-boiled 100% (v/v) and 50% (v/v) conditioned medium. Figure 7.1 B shows that  $\beta$ -catenin levels remain high in the cells grown in 100% (v/v) conditioned medium irrespective of boiling the medium. NANOG levels have were still high on growing SW480 cells in 100% (v/v)


#### Figure 7.1

Western blot analysis showing the effects of growing sub-confluent SW480 cells in conditioned medium. A) Levels of BRACHYURY in low density SW480 cells grown under different conditions. Lane 1 = 100% (v/v) conditioned medium collected from SW480 cells grown at a high density with no additional FBS; Lane 2 = 50% (v/v) conditioned medium collected from SW480 cells grown at a high density with 50% (v/v) fresh DMEM with 10% (v/v) FBS; Lane 3 = 100% (v/v) conditioned medium with 10% (v/v) FBS; Lane 4 = 100% (v/v) fresh DMEM with 10% (v/v) fresh DMEM with 10% (v/v) fresh DMEM with 10% (v/v) FBS. B) Western blot analysis demonstrating the levels of NANOG and  $\beta$ -catenin in SW480 sub-confluent cells grown under various conditions. Lane 1=SW480 cells grown in fresh medium; Lane 2 = cells grown in 100% (v/v) conditioned medium; Lane 4 = cells grown in 100% (v/v) conditioned medium boiled for 5 minutes; Lane 5 = cells grown in 50% (v/v) conditioned medium boiled for 5 minutes.

conditioned medium but its levels went down on boiling the 100% (v/v) conditioned medium. Again, 50% (v/v) conditioned medium has no effect on these levels though NANOG levels are decreased in SW480 cells grown in conditioned medium boiled for 5 minutes. Thus, it suggests that mechanisms regulating BRACHYURY and NANOG levels in SW480 cells are complex and involve multiple components. Whilst most proteins would be denatured by boiling, peptides may still remain fully active. We therefore, next carried out a series of purification steps aimed at identifying the nature of the 'factor' in the conditioned medium. These purification steps will now be described.

We performed a range of fractionation procedures (materials and methods section 2.24) to help elucidate the conditions best suited to isolate the peptide/ factor of interest from the conditioned medium (Figure 7.2). The following columns were used:

- Amicon ultra-15 centrifugal units (Millipore) with membrane of 10 kDa: The conditioned medium was centrifuged in this unit at 4000 g for 30 minutes at room temperature, which gave an upper fraction with molecules bigger than 10kDa and a lower fraction of molecules smaller than 10kDa.
- HiTrap<sup>™</sup> Blue HP Column (GE healthcare): This affinity column has Blue Sepharose to purify albumin, enzymes (including NAD<sup>+</sup> and NADP<sup>+</sup>), coagulation factors, interferons, and related proteins. This purification yields a flow through fraction, not bound to the column, and elution fractions which have affinity for the column. The binding buffer used was 20 mM sodium phosphate buffer pH 7.0 and the elution buffer was sodium phosphate buffer pH 7.0 + 2 M NaCl. The pH of the sample (boiled and filtered conditioned medium) was adjusted to 7 using 1M HCl.
- MonoS column (GE healthcare): This is a cationic exchange column which binds anionic molecules to the column. We used this column to separate biomolecules are that negatively charged (elution fractions). We obtained a flow through fraction which contains molecules with a positive and neutral



**Figure 7.2**: Fractions obtained from the protein fractionation. A) Flow through (FT1, 2) and elution (E1) fractions obtained from HiTrap Blue HP. B) Flow through (FT3, 4) and elution (E2, 3) fractions obtained from cationic exchange column. C) Flow through (FT5, 6) and elution (E4, 5) fractions obtained from anionic exchange column. The X-axis gives the volume (ml) of sample flown through. The Y-axis gives the OD (blue) and the pressure (green). The flow through are the fractions that are not bound to the column and elution fractions are bound to the column. FT= flow through, E= elution. [Note: Total volume loaded for each of these columns was 25ul].

charge. The binding buffer for this column was 20 mM MES (2-(N-morpholino) ethane sulfonic acid) pH 6.0 and elution buffer was 20 mM MES pH 6.0 + 1 M NaCl. The pH of the sample (boiled and filtered conditioned medium) was adjusted to 6 using 1M HCl.

 MonoQ column (GE healthcare): This is an anionic exchange column to isolate the cationic components from the sample (elution fraction) whilst the negative and neutral charged molecules are released as flow through fractions. The binding buffer for this was 20 mM Tris-HCl pH 8.0 and the elution buffer was 20 mM Tris-HCl pH 8.0 + 1M NaCl. The sample was adjusted to pH 8.0 1M NaOH.

Separation of putative 'peptides' was carried out using column fractionation and each of these fractions was then examined for BRACHYURY levels. Figure 7.3A shows the details of initial volume of boiled and filtered conditioned medium run through each column and the final volume of each of the fractions. We grew SW480 sub-confluent cells in each of these fractions diluted (as appropriate, according to relative concentration following fractionation) in fresh medium, adjusted to a total volume of 2ml. Our results (Figure 7.3B) confirm that the levels of BARCHYURY remain high when SW480 cells are grown at low density in fresh medium with FBS, whereas BARCHYURY is not detected when cells are grown in conditioned medium and filtered conditioned medium. The level of BARCHYURY was found to be low in the anion exchange column but the elution fractions of the anionic exchange column shows high levels of BARCHYURY. In contrast, there is no BRACHYURY detected in SW480 cells grown in medium isolated from the 'eluted' fraction from the cationic exchange column. This suggests that the elution fractions from the cationic exchange column, containing molecules with an anionic charge have biological activity associated with reducing the levels of BARCHYURY.

From Figure 7.3B, we also confirm that level of BARCHYURY remains high in the upper fraction of the ultra 15 spin column and is reduced in the lower

Column	Initial volume run through the column (ml)	Fractions collected	Volume of fractions collected (ml)
Ultra-15 spin column	15 ml	Upper fraction Lower fraction	1 ml 14 ml
Hitrap-blue- affinity column	25 ml	Flow through FT1 Flow through FT2 Elution E1	6 ml 6 ml 2-3 ml
Mono S- cationic exchange column	25 ml	Flow through FT3 Flow through FT4 Elution E2 (500 mM) Elution E3 (1M)	5-6 ml 6 ml 4 ml .5 ml
Mono Q-anionic exchange column	25 ml	Flow through FT5 Flow through FT6 Elution E4 (500 mM) Elution E5 (1M)	6 ml 6 ml 1 ml 1 ml
Brachyury Lamin B	123456	7 8 9 10 11 1	12 13

#### Figure 7.3

Result of the peptide/ protein fractionation. A) Table demonstrating the details of fractions obtained from the protein fractionation. B) Western blot analysis of BRACHYURY levels in low density SW480 cells grown in different fractions of conditioned medium; Lane 1 = Conditioned medium, lane 2 = fresh medium, lane 3 = upper fraction of the ultra 15 spin column (>10 kD), lane 4 = lower fraction of the ultra 15 spin column (<10 kD), lane 5 = Flow through of affinity column, lane 6 = Flow through of exchange column, lane 7 = Flow through of anionic exchange column, lane 8 = Elution affinity column, lane 9 = Elution of cationic column (peak 1), lane 10 = elution of cationic column (peak 2), lane 11 = elution of anionic column (peak 1), lane 12 = elution of anionic column (fractionational filtered medium). [Note: All boiled, conditioned medium was filtered using 115-ml filter unit, (Cat. Number 121-0020, Nalgene) prior to loading on the columns. Lane 13 was included to confirm that the biological activity was not lost following filtration].

fraction of the same column. This suggests that the 'factor' is of less than a 10kDa size.

#### 7.3 Discussion:

We have shown that *BRACHYURY* is differentially expressed in SW480 cells, in a density dependent manner. Specifically, BRACHYURY is present in relatively high amounts and in relatively low levels in SW480 super-confluent cells. Our aim was to identify whether a possible 'factor' carrying biological activity to regulate levels of BRACHYURY could be secreted in the medium and therefore, isolated when SW480 cells are grown at high density. Thus, we collected this medium (conditioned medium) and carried out preliminary fractionation to try to identify factors present in the conditioned medium that might be responsible for reducing the levels of BRACHYURY. Our results from protein fractionation from the conditioned medium suggest that BRACHYURY is not detected in the elution fraction of a cationic exchange column. This elution contains anions since it is not bound to cationic column. This suggests that there might be a possible anionic factor responsible for the low level of BRACHYURY in super-confluent SW480 cells. It would be important to examine this anionic fraction in greater detail, isolate its components and determine whether it is a protein, peptide or any other biological compound. Also, our result confirms that the possible 'factor' is likely to be of less than 10kDa size which guides us to look for components less than 10kDa size.

Most proteins are not heat resistant, although few of them are. Our experiment shows that on boiling the 100% (v/v) conditioned medium, the BRACHYURY activity remain high suggesting that the 'factor' possibly is not a protein, unless it is heat resistant protein.

# Chapter 8 - Summary

# 8.1 SW480 cell line: validation as a model to study molecular events involved in cells undergoing EMT and MET

Changes in cell density of the CRC cell line, SW480 lead to changes in gene expression and protein localization, resulting in a phenotype that resembles cells transitioning between mesenchymal-like and epithelial-like states (Brabletz et al., 2001; Davies et al., 2004). In low density SW480 cells (mesenchymal-like), the cells have low levels of E-CADHERIN located in the cytoplasm and nuclear  $\beta$ -catenin. In high density SW480 cells (epithelial-like), the cells exhibit higher levels of E-CADHERIN, mainly membrane associated, and membrane bound  $\beta$ -catenin. Such changes in levels of expression of E-CADHERIN and localization of E-cad and Bcatenin are hallmarks of cells transitioning between epithelial and mesenchymal states. In accordance with previous literature, we provided further evidence that SW480 cells grown at different densities are a good *in vitro* model to study the molecular events involved in EMT. Nuclear NFkB has previously been shown to be associated with induction and maintenance of EMT (Huber et al., 2004). One of the NFκB family members, Rel A (Nuclear factor, NFκB p65 subunit), which is the most abundant forms of NFkB, translocates from the nucleus to the cytoplasm as the SW480 cells progress from low to high density. The change in localization of Rel A is in accordance with the changes in localization of β-catenin and E-CADHERIN and further validates the use of SW480 cells grown to different densities as a system to study EMT.

We have also shown that in low density, mesenchymal-like SW480 cells, *BRACHYURY* is expressed at levels that are double those observed in the epitheliallike high density SW480 cells. We also found a number of other developmental genes which were also differentially expressed in the SW480 cell line and that were indicative of cells undergoing EMT/MET. These genes include *TBX2*, *TBX3*, *CDX2*, *ZEB1*, *SNAIL*, *FGF8*, *BRCAHYURY*, *E-CADHERIN* and *NANOG*.

#### 8.2 Why was BRACHYURY studied?

Although we found a range of developmental genes differentially expressed in low and high density SW480 cells (such as *TBX2*, *TBX3*, *CDX2*, *ZEB1*, *SNAIL*, *FGF8*), we focused our studies on *BRACHYURY* because it plays a critical role in embryogenesis and at the time of staring this thesis, had not had a published association with colorectal cancer. Recently, Fernando and co-workers (2010) have published that *BRACHYURY* is expressed in many cancer cell lines including SW480 and also in few lung cancer cell line, such as H460. This now gives a broader chance of studying BRACHYURY in cancer.

*BRACHYURY* is the founding member of the T-box transcription factor family of genes and is involved in early mesoderm development (Wilson and Conlon, 1990; Vujovic *et al.*, 2006). *BRACHYURY* has recently been identified as a possible target for T-cell mediated cancer immunotherapy and was associated with many tumour cell types, including colorectal tumour cell lines (Palena *et al.*, 2007). This further validates our results. *BRACHYURY* is also expressed in chordomas and in the embryonal carcinoma (EC) cell line NTERA2, where it was found to be expressed in the absence of mesodermal differentiation (Vujovic *et al.*, 2006; Gokhale *et al.*, 2000). However, no role of *BRACHYURY* in cancer has been described as yet. Human *BRACHYURY (T)* is located on chromosome 6, which makes it important to study because chromosome 6 is associated with colorectal cancers (Ezaki 2003).

Another critical observation was that *BRACHYURY* was not expressed in other human colorectal cancer cell lines examined such HCT116, LoVo and HT29. This makes the study of BRACHYURY in SW480 even more interesting because SW480 is obtained from a very aggressive tumour. Therefore, we wanted to investigate the role of BRACHYURY specifically in SW480 cells.

#### 8.3 Possible roles of BRACHYURY in SW480 cells

We have investigated possible roles of BRACHYURY in SW480 cells using a range of techniques. We wanted to examine whether BRACHYURY is directly involved in EMT. To address this, we knocked down BRACHYURY in SW480 cells, and checked the translocation and accumulation of β-catenin and the loss of E-CADHERIN expression (the two key hallmarks of EMT). (Hirohashi et al., 2003). Since, we did not see any change in the localization and expression of  $\beta$ -catenin and E-CADHERIN in response to BRACHYURY RNAi in these cells; we concluded that BRACHYURY might not be sufficient on its own to induce EMT. Further, we concluded this observation by knocking down BRACHYURY in low density SW480 cells, and observing no change in the expression of cell polarity marker genes such as ZEB1, ZEB2, PALS1, EVA1 and CRB. Fernando and colleagues (2010) stated that BRACHYURY is involved in the induction of EMT. But, their experiment was done by over-expressing BRACHYURY in their cells which induced EMT; whereas, our experiments were done without over-expressing BRACHYURY. This explains the fact that we did not see any change in EMT markers on BRACHYURY RNAi. If we over-express a gene, there is always a possibility that we change balance of other factors in the cell. Therefore, in our system, BRACHYURY is released but other factors might inhibit its EMT inducing properties, which might be active on overexpressing BRACHYURY.

Further, we investigated the role of *BRACHYURY* in migration of SW480 sub-confluent cells. From our screen of differentially expressed developmental genes in low density SW480 cells, we found that *CDX2* (a marker of intestinal epithelium) expression was reduced by 78% in high density SW480 cells compared to the low-density cells. *CDX2* is a biomarker of gastrointestinal differentiation, especially colorectal cancer (Liu *et al.*, 2007). Therefore, we studied *CDX2* expression in low density SW480 cells in response to *BRAHCYURY* knockdown, which reduced the expression of *CDX2* by 74%. This may suggest that *BRACHYURY* might play a role in regulating *CDX2* at the invasive fronts but not at

the central mass of the tumour because we do not expect BRACHYURY to be present in the central mass of the tumour. Gross and co-workers (2008) showed that expression of *CDX2* is high at the invasive front and its expression opposes cell migration in colorectal tumours. Therefore, we investigated a possible role of *BRACHYURY* in influencing colon cancer cell migration. (This study is currently in progress). Fernando and co-workers (2010) found that *BRACHYURY* induces cell migration in the pancreatic cell line PANC-1. This validates our preliminary migration assay data whereby, migration of SW480 cells was reduced by knocking down *BRACHYURY* expression.

The role of BRACHYURY was also studied in NFKB pathway. The activation of the NFkB pathway is involved in many cancers including colon cancer (Rayet and Gelinas 1999). It was also shown that NANOG inhibits the NFkB signalling pathway in mouse ES cells, thereby maintaining pluripotency (Watt, F. and Torres, J., 2008). The NFkB family consists of 5 member proteins, out of which REL A and REL B were extensively expressed in ES cells. NANOG was shown to bind to REL A in ES cells where it acts to inhibit NFkB regulated genes (Watt, F. and Torres, J., 2008). Therefore, we wanted to determine whether NANOG was regulating targets of NFkB genes similarly in SW480 cells, for which, we examined the expression of NFkB regulated genes (such as VIEMNTIN, PLAUR, BCL2L1 and IL10) in response to BRACHYURY knockdown in low density SW480 cells by qRT-PCR. Our observations show that on BRACHYURY RNAi in low density SW480 cells, there is an upregulation of NFkB regulated genes. Also, our Western blot analysis shows that phosphorylated REL A (p-REL A) and VIMENTIN are up-regulated on knocking down BRACHYURY in low density SW480 cells. This suggests a possible activation of NFkB signalling in low density SW480 cells in response to knockdown of BRACHYURY. Therefore, BRACHYURY may be involved in inhibition of NFKB signalling in low density SW480 cells, which may occur through activation of NANOG. Since, NFkB signaling is activated in many form of tumours, this observation was interesting, which would suggest that at low density SW480, cells

are invasive, where BRACHYURY is in high amounts, and it can inhibit the NF $\kappa$ B signaling which can make the cells less aggressive.

#### 8.4 BRACHYURY regulates NANOG

*Nanog* is one of the genes which were found to be differentially expressed in low and high density SW480 cells. Its expression was reduced by 70% in high density SW480 cells compared to low density SW480 cells. *Nanog* is shown to be expressed in a number of cancers such as primordial germ cell tumours (where it is involved in malignant cell transformation of gonads), osteosarcoma and lung cancer (Nirasawa *et al.*, 2009). *Nanog* is also a key gene involved in the maintenance of pluripotency of stem cells.

In mouse, NANOG has *Brachyury* and *Stat3* binding sites *in vivo* (Suzuki *et al.*, 2006) which regulate the expression of *Nanog* in mouse ES cells. Knocking down expression of *BRACHYURY* in low density SW480 cells, results in 85% reduction in the expression of *NANOG* in these cells. Therefore, we present data that link the expression of *BRACHYURY* with the regulation of the pluripotency gene *NANOG* and the maintenance of markers of CSCs in low density SW480 cells. Therefore, we further studied the relation between *BRACHYURY* and *NANOG* in the low density SW480 cells. We also confirmed the influence of BRACHYURY on the regulation of Nanog for the germ cell tumour cell line, NTERA2, and the CRC cell line T84, similar to SW480 cells.

Therefore, it might be possible that in invasive cells (such as low density SW480), *BRACHYURY* expression is high which regulates the expression of pluripotency gene, NANOG thereby maintaining a population of cancer stem cells. In a tumour, a population of cancer cells can turn nearby cells cancerous. This will make a benign tumour malignant. Therefore, targeting a cancer cell population is very important for drug targets.

#### 8.5 β-catenin is upstream of BRACHYURY

β-catenin is a modulator of signaling pathways that regulate 'stemness' (Gokhale *et al.*, 2000) and is required to maintain a discrete population of CSCs in epidermal tumours (Malanchi *et al.*, 2008; Braun *et al.*, 2008). BRACHYURY is shown to be a target of β-catenin signaling (Arnold, et al., 2000) in mouse ES and NIH3T3 cells. In low density SW480 cells, β-catenin is localized to the nucleus of the cells confirming active Wnt signaling. In these cells, BRACHYURY is highly expressed. Therefore, to study whether β-catenin is upstream of BRACHYURY in low density SW480, we knocked down β-catenin in low density SW480 cells. Our western blot data shows that knock-down of β-catenin in low density SW480 cells reduced the BRACHYURY levels in these cells. This demonstrates a possible mechanism whereby β-catenin signals through BRACHYURY to maintain a population of cells potentially having stem-like character, as determined by expression of *NANOG*.

#### 8.6 BRACHYURY binds to NANOG promoter region in SW480 cells

In mouse ES cells, the binding of BRACHYURY to sites in the *Nanog* promoter was co-ordinated with the binding of STAT3 (Suzuki *et al.*, 2006). In addition to this, our ChIP assay showed direct binding of BRACHYURY to specific regulatory elements in the *Nanog* promoter region, and this binding was diminished following knockdown of  $\beta$ -catenin. However, we did not find any STAT3 binding sites close to either of the T-binding sites that we located in the promoter region of the human *Nanog* gene. This observation agrees with the fact that signalling through STAT3 does not promote self-renewal in human ES cells in culture (Le *et al.*, 2008).

In cancer cells, *NANOG* mRNA is derived mostly from the retrogene *NANOGP8* (*NANOG* pseudogene) (Jeter et al., 2008); and *NANOGP8* is expressed in many forms of cancer (Zhang et al., 2006) but not all. Therefore, we are currently doing ChIP assay to clarify whether we are detecting binding of BRACHYURY to *NANOGP8* promoter region. Before doing this, we will first make sure whether SW480 cells do express *NANOGP8* or not. If *NANOGP8* is not expressed in SW480 cells, then we will conclude that we are looking at the binding of BRACHYURY to *NANOG* promoter region. If we see any expression of *NANOGP8* in SW480 cells, we will do the ChIP assay for clarification.

Pereira and co-workers (2008) suggested that  $\beta$ -catenin signaling increase *Nanog* promoter activity in a Tcf3-independent manner in mouse ES cells. This suggests a complex control mechanism existing in these cells. This observation is consistent with our data where  $\beta$ -catenin acts to positively regulate the expression of *Nanog* in SW480 cells.

The luciferase assay data presented indicates that the expression levels from the luciferase construct are low when a 701bp *NANOG* upstream element is inserted (Construct 2): whose level is close to the control levels, suggesting that the promoter elements contained in the sequence are not efficient under the conditions used in this experiment. Construct 3 (with both BRACHYURY binding sites) gives an increase in the transcription activity that is 2.3 fold compared to controls, suggesting that enhancer activity may be contained in a 114bp sequence element (2399bp to 2285bp) that contains the second BRACHYURY binding site. Whilst ChIP data showed that BRACHYURY binds to both sites, it is likely that one site alone is insufficient to enhance transcription.

Further, it will be interesting to make a construct with site 2 and not site 1. This will tell us whether the binding affinity is dependent on site 2 or site 1 alone. Also, deletion or mutation of the bases in-between the two sites will also suggest whether the activity is due to the bases in-between the two sites and not site 1 or site 2 alone. Direct mutation of site 1 and 2 will also be done in the laboratory.

High levels of NANOG translate to efficient self-renewal and low levels of Nanog lead to an increased chance of differentiation (Lluis *et al.*, 2008). Although,

Nanog is not absolutely required for self-renewal of ESC, it is found to be essential for the development of primordial germ cells (PGCs), where it is estimated to have a role in re-setting the epigenetic state. It is possible that the presence of NANOG in low density SW480 cells serves to deter differentiation. Changes in the media or growth conditions over time might lead to changes in expression profiles that favour differentiation with a corresponding decrease in levels of NANOG.

Furthermore, it is also possible that NANOG plays a role in low density SW480 cells, in altering the epigenetic state of the cells to either protect an 'immature cell state' (through high expression) or alternatively to allow differentiation (through low expression).

The initial tumour-generating events in CRC are of defined nature, but the phenotype of the cells alters with progression of the tumour, irrespective of genetic mutations. This observation may imply that epigenetic changes are involved, or alternatively, that the cells respond to extrinsic cues (Brabletz *et al.*, 2005). In accordance with this, our work shows that the SW480 cancer cells can transition reversibly between two states (depending on cell density), one state bearing characteristics/markers of CSCs, and one state bearing epithelial-like character. The expression of *NANOG* in SW480 cells might therefore permit a cell to express stem cell markers (such as CD166) and also to have stem cell characteristics. We also detected expression of other genes associated with the undifferentiated state such as *OCT4*, *SOX2* and *FOXD3* in SW480 cells at low density.

#### 8.7 BRACHYURY expression in other tumour types

From the limited published studies, it is likely that not all tumours/cell lines express *BRACHYURY*. It may therefore be that those tumours that do not express *BRACHYURY* have evolved an alternative route to activate NANOG which might be important in establishing cells bearing characteristics of CSCs. Alternatively,

tumours not expressing *BRACHYURY* and *NANOG* may be less aggressive or may not have the ability to transition between different differentiation states. It would be interesting to determine the generality of the presence and localisation pattern of NANOG and BRACHYURY in a larger panel of colorectal tumours and tissue and to correlate such data with the presence of CSCs and patient outcomes.

We have also examined expression of BRACHYURY in the human ovarian tumour cell line SW626 cells by Western blot analysis and qRT-PCR. In these cells, BRACHYURY is not present but the expression of NANOG is very high. We suggest that in tumour types where BRACHYURY is not present, they develop an alternative route to activate NANOG which is centrally important for progression of the tumour by establishment of tumour initiating stem cells.

We have shown that in human embryonal carcinoma cell line, Ntera-2 and human colonic adenocarcinoma cell line T84, NANOG is shown to be regulated by BRACHYURY in a similar manner to that seen in SW480 sub-confluent cells. Therefore, it shows that the mechanism of BRACHYURY regulating NANOG is not restricted to SW480 cells. It is possible that this regulatory pathway of BRACHYURY regulating NANOG is detected in cell lines that are competent for mesodermal differentiation, such as NTERA2, whereas T84 did not really undergo EMT-like transitions which were observed in SW480 cells. The levels of BRACHYURY in these T84 cells are considerably reduced compared to SW480 cells. It will be necessary to determine the status of other pathways in these cells to try examining the mechanism by which this pathway operates.

### 8.8 Conditioned media 'factor' might be responsible for differential expression of BRACHYURY in SW480 cells

We have already mentioned that the expression of *BRACHYURY* in low density SW480 cells is twice as higher compared to high density SW480 cells. We

wanted to explore the possibility that factors controlling the levels of BRACHYURY might be secreted into the media and as such they might be isolated and identified.

The results we observed in the protein purification experiment, suggest that the possible 'factor' influencing *BRACHYURY* expression is an anionic factor. It would be important to examine this anionic fraction in greater detail, isolate its components and determine whether it is a protein, peptide or any other biological compound. Our result also suggests that the 'factor' is likely to be of less than 10kDa size and thus, we should look for components less than 10kDa size. Further study of this 'factor' will be beneficial in determining an element that is involved in regulation of BRACHYURY in SW480 colon cancer cell.

Also, when we repeated our experiment on boiled lysates, the 'factor' was still sustained. This might suggest that the 'factor' is unlikely to be a protein, since most proteins would degrade by heat. The possibility lies that the 'factor' could be a peptide or even peptidase.

#### 8.9 A model pathway derived:

Based on our study so far, we have derived a model pathway (Figure 8.1A) explaining the results we obtained so far. We have shown SW480 transitioning between low (mesenchymal-like) and high density (epithelial-like) and BRACHYURY expression and levels transitions between these two density types of SW480 cells. Since, we have shown that  $\beta$ -catenin signals upstream of BRACHYURY, we predict that  $\beta$ -catenin activates BRACHYURY in low density SW480 cells, which in turn activates the expression of NANOG to maintain a population of colorectal cancer stem cells defined by CSC markers (CD133 and CD166). A tumour needs cancer stem cell to make a benign one malignant. In high density SW480 cells,  $\beta$ -catenin signaling is low and therefore, BRACHYURY level

is also low. This gives very rare NANOG and CRC stem cell marker cells at high density SW480 cells. Based on these observations, we predict a model pathway (Figure 8.1B) whereby, Wnt pathway activates BRACHYURY in SW480 cells; which in turn activates NANOG and the CRC stem cell markers. Nevertheless, this pathway has to be further verified in other tumour cell types.



**Figure 8.1**: Model pathway derived. SW480 cells transition between low and high density. A) At low density,  $\beta$ -catenin, BRACHYURY, NANOG and CRC stem cell markers are highly expressed. At high density SW480 cells,  $\beta$ -catenin signaling is low and expression of BRACHYURY, NANOG and CRC stem cell markers are very rare. CRC = colorectal cancer. B) We predict that Wnt pathway activates BRACHYURY in SW480 cells which in turn activates the pluripotency gene, NANOG and the CRC stem cell markers.

### References

Aberle H., Bauer A., Stappert J., Kispert A., Kemler R. (1997)  $\beta$ -catenin is a target of the ubiquitin-proteasome pathway. EMBO J. 16(1):3797-3804

Aberle H., Butz S., Stappert J., Weissig H., Kemler R. & Hoschuetzky H. (1994) Assembly of the cadherin-catenin complex in vitro with recombinant proteins. J. Cell Sci. 107(1):3655-3663

Ahmed F, Wyckoff J, Lin EY, Wang W, Wang Y, Hennighausen L, Miyazaki J, Jones J, Pollard JW, Condeelis JS et al.(2002) GFP expression in the mammary gland for imaging of mammary tumor cells in transgenic mice. Cancer Res. 62(1):7166-7169

Aigner K, Dampier B, Descovich L, Mikula M, Sultan A, Schreiber M, Mikulits W et al. (2007) The transcription factor ZEB1 (deltaEF1) promotes tumour cell dedifferentiation by repressing master regulators of epithelial polarity. Oncogene 26(1): 6979-6988

Aigner K, Dampier B, Descovich L, Mikula M, Sultan A, Schreiber M, Mikulits W, Brabletz T, Strand D, Obrist P, Sommergruber W, Schweifer N, Wernitznig A, Beug H, Foisner R, Eger A (2007) The transcription factor ZEB1 (deltaEF1) promotes tumour cell dedifferentiation by repressing master regulators of epithelial polarity. Oncogene 26(49):6979-6988

Akashi, K. & Weissman, I. L. in Developmental Biology of Hematopoiesis (ed. Zon, L. I.) 15–34 (Oxford Univ. Press, New York, 2001)

Albensi BC, Mattson MP. (2000) Evidence for the involvement of TNF and NF-κB in hippocampal synaptic plasticity Synapse 35(2):151–159

Al-Hajj M, Wicha MS, Benito-Hernandez A, Morrison SJ, Clarke MF (2003) Prospective identification of tumorigenic breast cancer cells.Proceedings of the National Academy of Sciences of the United States of America 100(7):3983–3988

Allenspach EJ, Maillard I, Aster JC, Pear WS. (2002) Notch signaling in cancer. Cancer Biol Ther. 1(5):466-476

Anderson C.B., Neufeld K.L. & White R.L. (2002) Subcellular distribution of Wnt pathway proteins in normal and neoplastic colon. Proc. Natl. Acad. Sci. USA 99(1):8683-8688

Andreas Wodarz and Roel Nusse. (1998). Mechanisms of Wnt signaling in devleopment. Annu. Rev. Cell Dev. Biol. 14(1):59-88

Andreu, P., Colnot, S., Godard, C., Gad, S., Chafey, P., Niwa-Kawakita, M., Laurent-Puig, P., Kahn, A., Robine, S., Perret C., et al. (2005) Crypt-restricted

proliferation and commitment to the Paneth cell lineage following Apc loss in the mouse intestine. Development 132(1):1443–1451

Andrews P.W., Damjanov I., Simon D., Banting G., Carlin C., Dracopoli N.C. & Fogh J. (1984) Pluripotent embryonal carcinoma clones derived from the human teratocarcinoma cell line TERA-2: Differentiation in vivo and in vitro. Lab. Invest. 50(1):147-162

Andrews, P.W. (1984) Retinoic acid induced neuronal differentiation of a cloned human embryonal carcinoma cell line in vitro. Dev. Biol. 103(1):285-293

Antoine M, Reimers K, Wirz W, Gressner AM, Müller R, Kiefer P. (2005) Fibroblast growth factor 3, a protein with a dual subcellular fate, is interacting with human ribosomal protein S2. Biochem Biophys Res Commun. 338(2):1248-1255

Anton R, Hans A. Kestler, and Michael Kühla. (2007) β-Catenin signaling contributes to stemness and regulates early differentiation in murine embryonic stem cells. Federation of European Biochemical Societies. 581(27):5247-5254

Ariel J. Levine and Ali H. (2006) GDF3, a BMP inhibitor, regulates cell fate in stem cells and early embryos. Development. 133(1):209-216

Arnold J, Jörg Stappert, Andreas Bauer, Andreas Kispert, Bernhard G. Herrmann and Rolf Kemler. (2000) Brachyury is a target gene of the Wnt/ $\beta$ -catenin signaling pathway. Mechanisms of Development. 91(1-2):249-258

Arnold SJ, Stappert J, Bauer A, Kispert A, Herrmann BG, Kemler R. (2000) Brachyury is a target gene of the Wnt/beta-catenin signaling pathway. Mech Dev. 91(1):249-258

Asai, R., Okano, H., and Yasugi, S. (2005) Correlation between Musashi-1 and chairy-1 expression and cell proliferation activity in the developing intestine and stomach of both chicken and mouse. Dev. Growth Differ. 47(1):501–510

Auersperg N, Wong AST, Choi KC, Kang KS, and Leung PCK. (2001) Ovarian surface epithelium: biology, endocrinology, and pathology. Endocr Rev 22(1):255-288

Bachmann, A., Schneider, M., Theilenberg, E., Grawe, F. and Knust, E. (2001) Drosophila Stardust is a partner of Crumbs in the control of epithelial cell polarity. Nature 414(1):638 -643

Baldassarre G, Fedele M, Battista S, Vecchione A, Klein-Szanto AJ, Santoro M, Waldmann TA, Azimi N, Croce CM, Fusco A. (2001) Onset of natural killer cell lymphomas in transgenic mice carrying a truncated HMGI-C gene by the chronic

stimulation of the IL-2 and IL-15 pathway. Proc Natl Acad Sci U S A. 98(14):7970-7975

Bamshad, M., et al. (1997) Mutations in human TBX3 alter limb, apocrine and genital development in ulnar-mammary syndrome. Nat Genet 16(1):311–315

Barker N, van Es JH, Kuipers J, Kujala P, van den Born M, Cozijnsen M, Haegebarth A, Korving J, Begthel H, Peters PJ, Clevers H. (2007) Identification of stem cells in small intestine and colon by marker gene Lgr5. Nature. 449(7165):1003-1007

Barrallo-Gimeno A, Nieto MA. (2005) The Snail genes as inducers of cell movement and survival: implications in development and cancer. Development 132(1):3151–3161

Basson, C.T., et al. (1997). Mutations in human cause limb and cardiac malformations in Holt-Oram syndrome. Nat Genet 15(1):30–35

Batlle, E., Henderson, J.T., Beghtel, H., van den Born, M.M., Sancho, E., Huls, G., Meeldijk, J., Robertson, J., van de Wetering, M., Pawson, T., et al. (2002)  $\beta$ -Catenin and TCF mediatecell positioning in the intestinal epithelium by controlling the expression of EphB/ephrinB. Cell 111(1): 251–263

Behrens, J., Jerchow, B. A., Wurtele, M., Grimm, J., Asbrand, C., Wirtz, R., Kuhl, M., Wedlich, D. & Birchmeier, W. (1998) Caught up in a Wnt storm: Wnt signaling in cancer. Science. 280(1):596–599

Bhardwaj, G. et al. (2001) Sonic hedgehog induces the proliferation of primitive human hematopoietic cells via BMP regulation. Nature Immunol. 2(1):172–180

Bienz M. & Clevers H. (2000) Linking colon cancer to Wnt signalling. Cell 103(1):311-320

Bienz M. (1998) TCF: Transcriptional activator or repressor? Curr. Opin. Cell. Biol. 10(1):366-372

Bienz M. (2002) The subcellular destinations of APC proteins. Nat. Rev. Mol. Cell Biol. 3(1):328-338

Bisgaard M.L., Fenger K., Bulow S., Niebuhr E. & Mohr J. (1994) Familial adenomatous polyposis. Hum. Mutat. 3(1):121-125

Bjerknes M, Cheng H. (2005) Gastrointestinal stem cells. II. Intestinal stem cells. Am J Physiol Gastrointest Liver Physiol. 289(1):G381-G387

Bjerknes, M. and Cheng, H. (1999) Clonal analysis of mouse intestinal epithelial progenitors. Gastroenterology. 116(1):7-14

Bjerknes, M. and Cheng, H. (2006) Intestinal epithelial stem cells and progenitors. Methods Enzymol. 419(1):337-383

Bjerknes, M., and Cheng, H. (1981b) The stem-cell zone of the small intestinal epithelium. III. Evidence from columnar, enteroendocrine, and mucous cells in the adult mouse. Am. J. Anat. 160(1):77-91

Bjerkvig R, Tysnes BB, Aboody KS, Najbauer J, Terzis AJ. (2005) Opinion: the origin of the cancer stem cell: current controversies and new insights. Nat Rev Cancer. 5(11):899-904. Review

Blache P, van de Wetering M, Duluc I, Domon C, Berta P, Freund JN, Clevers H, Jay P. (2004) SOX9 is an intestine crypt transcription factor, is regulated by the Wnt pathway, and represses the CDX2 and MUC2 genes. J Cell Biol.166(1):37-47

Blanpain C, Lowry W.E, Geoghegan A, Polak L, and Fuchs E. (2004) Self-renewal, multipotency, and the existence of two cell populations within an epithelial stem cell niche. Cell. 118(5):530-532

Boiani M and H.R. Scholer. (2005) Regulatory networks in embryo-derived pluripotent stem cells, Nat. Rev. Mol. Cell Biol. 6(1):872–884

Bolos V, Peinao H, Perez-Moreno MA, Fraga MF, Estella M, Cano H. (2003) The transcription factor Slug represses E-cadherin expression and induces epithelial to mesenchymal transitions: a comparison with Snail and E47 repressors. J Cell Sci 116(1):499–511

Bonizzi G. Karin M. (2004) The two NFkB activation pathways and their role in innate and adaptive immunity. Trends Immunol. 25(1):280-288

Bonnet D, Dick JE. (1997) Human acute myeloid leukemia is organized as a hierarchy that originates from a primitive hematopoietic cell. Nat Med. 3(7):730-7

Booth C, Potten CS. (2000) Gut instincts: thoughts on intestinal epithelial stem cells. J Clin Invest.105(1):1493-1499

Boyden LM, Mao J, Belsky J, Mitzner L, Farhi A, Mitnick MA, Wu D, Insogna K, Lifton RP. (2002) High bone density due to a mutation in LDL-receptor-related protein 5. N Engl J Med. 16;(20):1513-1521

Boyer B, Valles AM, Edme N. (2000) Induction and regulation of epithelialmesenchymal transitions. Biochem Pharmacol 60(1):1091–1099 Brabletz T, Jung A and Kirchner T (2002) Beta-catenin and the morphogenesis of colorectal cancer. Virchows Arch 441(1):1-11

Brabletz T, Jung A Spaderna S, Hlubek F and Kirchner T (2005) Opinion: migrating cancer stem cells - an integrated concept of malignant tumour progression. Nat. Rev. Cancer. 5(1):744-749

Brabletz T, Jung A, Reu S, Porzner M, Hlubek F, Kunz-Schughart LA, Knuechel R, Kirchner T. (2001) Variable beta-catenin expression in colorectal cancers indicates tumor progression driven by the tumor environment. Proc Natl Acad Sci U S A. 98(1):10356-10361

Brabletz T, Spaderna S, Kolb J, Hlubek F, Faller G, Bruns CJ et al. (2004). Downregulation of the homeodomain factor cdx2 in colorectal cancer by collagen type I: an active role for the tumor environment in malignant tumor progression. Cancer Res 64(1): 6973–6977

Brabletz T., Jung A., Dag S., Hlubek F. & Kirchner T. (1999)  $\beta$ -catenin regulates the expression of the matrix metalloproteinase-7 in human colorectal cancer. Am. J. Pathol. 155(1):1033-1038

Brocardo, Nathke & Henderson. (2005) Redefining the subcellular location and transport of APC: new insights using a panel of antibodies. EMBO. 6(2):184-190

Braun K. (2008) Cutaneous Cancer Stem Cells: β-Catenin Strikes Again. Cell Stem Cell. 2(5):406-408

Brentjens RJ, Latouche JB, Santos E, Marti F, Gong MC, Lyddane C, King PD, Larson S, Weiss M, Rivière I, Sadelain M. (2003) Eradication of systemic B-cell tumors by genetically targeted human T lymphocytes co-stimulated by CD80 and interleukin-15. Nat Med. 9(3):279-286

Cadigan K.M. & Nusse R. (1997) Wnt signalling: a common theme in animal development. Genes. Dev. 11(1):3286-3305

Cai L, Ye Z, Zhou BY, Mali P, Zhou C, Cheng L. (2007) Promoting human embryonic stem cell renewal or differentiation by modulating Wnt signal and culture conditions. Cell Res. 17(1):62-72

Cairnie AB, Lamerton LF, and Steel GG. (1965) Cell proliferation studies in the intestinal epithelium of the rat. II. Theoretical aspects. Exp Cell Res. 39(1):539-553

Calvi LM, Adams GB, Weibrecht KW et al. (2003) Osteoblastic cells regulate the haematopoietic stem cell niche. Nature. 425(1):841–846

"Cancer". World Health Organization. February 2006. http://www.who.int/mediacentre/factsheets/fs297/en/. Retrieved on 2007-05-24

Cardiff RD, Chodosh LA. (2005) The transcriptional repressor Snail promotes mammary tumor recurrence. Cancer Cell 8(1):197-209

Carlsson, P., and M. Mahlapuu. (2002) Forkhead transcription factors: key players in development and metabolism. Dev. Biol. 250(1):1-23

Chakladar A, Dubeykovskiy A, Wojtukiewicz LJ, Pratap J, Lei S, Wang TC. (2005) Synergistic activation of the murine gastrin promoter by oncogenic Ras and betacatenin involves SMAD recruitment.Biochem Biophys Res Commun. 336(1):190-196

Chambers I, Colby D, Robertson M, Nichols J, Lee S, Tweedie S, Smith A. (2003) Functional expression cloning of Nanog, a pluripotency sustaining factor in embryonic stem cells. Cell 113(5):643–655

Chambers I, Silva J, Colby D, Nichols J, Nijmeijer B, Robertson M, Vrana J, Jones K, Grotewold L, Smith A. (2007) Nanog safeguards pluripotency and mediates germline development. Nature 450(1):1230-1234

Chao A, Thun MJ, Jacobs EJ. (2000) Cigarette smoking and colorectal cancer mortality in the cancer prevention study II. J Natl Cancer Inst 92 (23):1888-1896

Chawengsaksophak K, de Graaff W, Rossant J, Deschamps J, Beck F (May 2004) Cdx2 is essential for axial elongation in mouse development. Proc. Natl. Acad. Sci. U.S.A. 101 (20):7641-7645

Chen C, Ware S, Sato A, Houston-Hawkins D, Habas R, Matzuk M, Shen M, Brown C (2006) The Vg1-related protein Gdf3 acts in a Nodal signaling pathway in the pregastrulation mouse embryo. Development 133(2):319–329

Chen P, Tian D, Liu M. (2008) The role of Tbx2 in pancreatic cancers and its regulation by Wnt/ $\beta$ -catenin signaling. The Chinese-German Journal of Clinical Oncology. 7(7):404-409

Chen YG, Wang Q, Lin SL, Chang CD, Chuang J, Chung J, Ying SY (2006) Activin signaling and its role in regulation of cell proliferation, apoptosis, and carcinogenesis.Exp. Biol. Med. (Maywood) 231(5):534–544

Cheng H and Leblond CP. (1974) Origin, differentiation and renewal of the four main epithelial cell types in the mouse small intestine. V. Unitarian theory of the origin of the four epithelial cell types. Am J Anat. 141(1):537-562

Cheng, H. and Leblond, C.P. (1974a) Origin, differentiation and renewal of the four main epithelial cell types in the mouse small intestine. I. Columnar cell. Am. J. Anat. 141(1):461-479

Chesley, P. (1935) Development of the short-tailed mutant in the house mouse. J. Exp. Zool. 71(1):429-459

Chiou SH, Yu CC, Huang CY, Lin SC, Liu CJ, Tsai TH, Chou SH, Chien CS, Ku HH, Lo JF. (2008) Positive correlations of Oct-4 and Nanog in oral cancer stem-like cells and high-grade oral squamous cell carcinoma. Clin Cancer Res. 14(13):4085-4095

Christman JW, Sadikot RT, Blackwell TS (2000) The role of nuclear factor-kappa B in pulmonary diseases. Chest. 117(1):1482–1487

Christodoulides C, Scarda A, Granzotto M, Milan G, Dalla Nora E, Keogh J, De Pergola G, Stirling H, Pannacciulli N, Sethi JK, Federspil G, Vidal-Puig A, Farooqi IS, O'Rahilly S, Vettor R. (2006) WNT10B mutations in human obesity. Diabetologia. 49(4):678-684

Chung FY, Huang MY, Yeh CS, Chang HJ, Cheng TL, Yen LC, Wang JY, Lin SR. (2009) GLUT1 gene is a potential hypoxic marker in colorectal cancer patients. BMC Cancer. 9(1):241

Clark AT. (2007) The stem cell identity of testicular cancer. Stem Cell Rev. 3(1):49-59

Clarke MF, Fuller M. (2006) Stem cells and cancer: two faces of eve. Cell. 124(6):1111-1115

Clarke MF. (2005) A self-renewal assay for cancer stem cells. Cancer Chemother Pharmacol.56 Suppl 1(1):64-8. Review

Claudinot, S., Nicolas, M., Oshima, H., Rochat, A., and Barrandon, Y. (2005) Longterm renewal of hair follicles from clonogenic multipotent stem cells. Proc. Natl. Acad. Sci. 102(1):14677–14682

Clevers H. (2000) Axin and hepatocellular carcinomas. Nat Genet. 24(3):206-208

Conacci-Sorrell M, Simcha I, Ben Yedidia T, Blechman J, Savagner P, Ben Ze'ev A. (2003). Autoregulation of E-cadherin expression by cadherin-cadherin interactions: the roles of beta-catenin signaling, Slug, and MAPK. J Cell Biol 163(1):847–857

Conboy IM, Conboy MJ, Wagers AJ et al. (2005) Rejuvenation of aged progenitor cells by exposure to a young systemic environment. Nature. 433(1):760–764

Conboy IM, Rando TA. (2002) The regulation of Notch signaling controls satellite cell activation and cell fate determination in postnatal myogenesis. Dev Cell. 3(1):397-409

Conlon Frank L, Fairclough Lynne, Brenda, Price M. J, Elena S. Casey and Smith J. C. (2001). Determinants of T box protein specificity. Development. 128(1):3749-3758

Conlon, F. L., Sedgwick, S. G., Weston, K. M. and Smith, J. C. (1996) Inhibition of Xbra transcription activation causes defects in mesodermal patterning and reveals autoregulation of Xbra in dorsal mesoderm. Development 122(1):2427-2435

Corbeil D, Röper K, Hannah MJ, Hellwig A, Huttner WB. (1999) Selective localization of the polytopic membrane protein prominin in microvilli of epithelial cells - a combination of apical sorting and retention in plasma membrane protrusions. J Cell Sci. 112(7):1023-1033

Corbeil D, Röper K, Hellwig A, Tavian M, Miraglia S, Watt S, Simmons P, Huttner W. (2000) The Human AC133 Hematopoietic Stem Cell Antigen Is also Expressed in Epithelial Cells and Targeted to Plasma Membrane Protrusions. J Biol Chem. 275(8):5512-5520

Cottrell TR, Doering TL. 2003. Silence of the strands: RNA Interference in eukaryotic pathogens. Trends in Microbiology 11:37-43. <<u>http://reviews.bmn.com/rsearch/section/record?uid=TIM.bmn05510\_0966842x\_v0</u>011i01\_02000045&rendertype=text>. Accessed Feb 8 2010

Cottrell S., Bicknell D., Kaklamanis L. & Bodmer W.F. (1992) Molecular analysis of APC mutations in familial adenomatous polyposis and sporadic colon carcinomas. Lancet 340(1):626-630

Crosnier C, Stamataki D, Lewis J. (2006) Organizing cell renewal in the intestine: stem cells, signals and combinatorial control. Nat Rev Gene. 7(5):349-359

Crouch SP, Kozlowski R, Slater KJ, Fletcher J. (1993) The use of ATP bioluminescence as a measure of cell proliferation and cytotoxicity. J Immunol Methods. 160 (1):81-88

Dahéron L, Opitz SL, Zaehres H, Lensch MW, Andrews PW, Itskovitz-Eldor J, Daley GQ. (2004) LIF/STAT3 signaling fails to maintain self-renewal of human embryonic stem cells. Stem Cells 22(1):770-778

Dale T.C. (1998) Signal transduction by the Wnt family of ligands. Biochem. J. 329(1):209-223

Dalerba P, Cho RW, Clarke MF. (2007a) Cancer stem cells: models and concepts. Annu Rev Med. 58(1):267-284

Dalerba P, Dylla S.J, Park I.K, Liu R., Wang X, Cho R.W, Hoey T, Gurney A, Huang E.H, Simeone D.M, et al. (2007b) Phenotypic characterization of human colorectal cancer stem cells. Proc. Natl. Acad. Sci. 104(1):10158-10163

Dang, F Chen, C Ying, S Y Chun, S A Knock, H D Appelman, D T Dang. (2006) CDX2 has tumorigenic potential in the human colon cancer cell lines LOVO and SW48. Oncogene. 25(1):2264-2272

Davies ML, Roberts GT, Spiller DG, Wakeman JA. (2004) Density-dependent location and interactions of truncated APC and beta-catenin.Oncogene 23:1412-1419 Daylon J, Ariel JL, Besser D and Hemmati-Brivanlou A. (2005) TGF $\beta$ /activin/nodal signaling is necessary for the maintenance of pluripotency in human embryonic stem cells. Development. 132(1):1273-1282

De Ferrari GV, Papassotiropoulos A, Biechele T, Wavrant De-Vrieze F, Avila ME, Major MB, Myers A, Sáez K, Henríquez JP, Zhao A, Wollmer MA, Nitsch RM, Hock C, Morris CM, Hardy J, Moon RT. (2007) Common genetic variation within the low-density lipoprotein receptor-related protein 6 and late-onset Alzheimer's disease. Proc Natl Acad Sci U S A. 104(22):9434-9439

De Lott LB, Morrison C, Suster S, Cohn DE, Frankel WL. (2005). CDX2 is a useful marker of intestinal-type differentiation: a tissue microarray-based study of 629 tumors from various sites. Arch Pathol Lab Med. 129(9):1100-1105

Dharmaraj, S. (2006). RT-PCR: The Basics. Available: http://www.ambion.com/techlib/basics/rtpcr/index.html. Accessed 08 February 2010

Dobrovolskaia-Zavadskaia N, Sceances CR. (1927) Sur la mortification spontane' de la chez la souris nouveau-ne' et sur l'existence d'un caracte're (facteur) hereditaire. Sco Biol. 97(1):114–116

Domen, J. & Weissman, I. (2000) Hematopoietic stem cells need two signals to prevent apoptosis; BCL-2 can provide one of these, Kitl/c-Kit signaling the other. J. Exp. Med. 192:1707–1718

Domen, J., Gandy, K. L. & Weissman, I. L. (1998) Systemic overexpression of BCL-2 in the hematopoietic system protects transgenic mice from the consequences of lethal irradiation. Blood 91:2272–2282

Dongho Choi, Hyo Won Lee, Kyung Yul Hur, Jae Joon Kim, Gyeong-Sin Park, Si-Hyong Jang, Young Soo Song, Ki-Seok Jang, and Seung Sam Paik. (2009). Cancer stem cell markers CD133 and CD24 correlate with invasiveness and differentiation in colorectal adenocarcinoma. World J Gastroenterol.15(18):2258–2264

Dontu G, Jackson KW, McNicholas E, Kawamura MJ, Abdallah WM, Wicha MS / (2004) Role of Notch signaling in cell-fate determination of human mammary stem/progenitor cells. Breast cancer research: BCR. 6(6):R605–615.

Duluc I, Lorentz O, Fritsch C, Leberquier C, Kedinger M and Freund JN (1997) Changing intestinal connective tissue interactions alters homeobox gene expression in epithelial cells. J Cell Sci 110(1):1317–1324

Dyson S, Gurdon J.B. (1997) Activin signalling has a necessary function in Xenopus early development. Current Biology. 7(1):81-84

Dzierzak E. (2003) Ontogenic emergence of definitive hematopoietic stem cells. Curr Opin Hemato.10(1):229-234

Edlund S, Lee SY, Grimsby S, Zhang S, Aspenström P, Heldin CH, Landström M. (2005) Interaction between Smad7 and beta-catenin: importance for transforming growth factor beta-induced apoptosis. Mol Cell Biol 25(4):1475-88

Edme N, Downward J, Thiery JP, Boyer B. (2002) Involvement of MAPK and Rac in epithelial cell scattering, J Cell Sci.115(1)

Edwards YH, Putt W, Lekoape KM, Stott D, Fox M, Hopkinson DA, Sowden J. (1996) The human homolog T of the mouse T(Brachyury) gene; gene structure, cDNA sequence, and assignment to chromosome 6q27. Genome Res. 6(3):226-233

Ee HC, Erler T, Bhathal PS, Young GP, James RJ. (1995). Cdx-2 homeodomain protein expression in human and rat colorectal adenoma and carcinoma. Am J Pathol 147(1):586-592

Eppig JT, Levan G. (2009, Revised). Guidelines for Nomenclature of Genes, Genetic Markers, Alleles, and Mutations in Mouse and Rat. Available: http://www.informatics.jax.org/mgihome/nomen/gene.shtml. Accessed 19 February 2010

Essers MA, de Vries-Smits LM, Barker N, Polderman PE, Burgering BM, Korswagen HC. (2005) Functional interaction between beta-catenin and FOXO in oxidative stress signaling. Science. 308(5725):1181-1184 Ethan A. Carver, Rulang Jiang, Yu Lan, Kathleen F. Oram, and Thomas Gridley. (2001) The Mouse Snail Gene Encodes a Key Regulator of the Epithelial-Mesenchymal Transition. Molecular and Cellular Biology. 21(23):8184-8188

Ettensohn CA, Illies MR, Oliveri P, De Jong DL. (2003) Alx1, a member of the Cart1/Alx3/Alx4 subfamily of paired-class homeodomain proteins, is an essential component of the gene network controlling skeletogenic fate specification in the sea urchin embryo. Development. 130(1):2917-2928

Evans M, Kaufman M (1981) Establishment in culture of pluripotent cells from mouse embryos. Nature. 292(5819):154–156

Ezaki T, Watanabe M, Inoue N, Kanai T, Ogata H, Iwao Y, Ishii H, and Hibi T. (2003) A Specific Genetic Alteration on Chromosome 6 in Ulcerative Colitisassociated Colorectal Cancers. Cancer Research. 63(1):3747-3749

Ezeh UI, Turek PJ, Reijo RA, Clark AT. (2005) Human embryonic stem cell genes OCT4, NANOG, STELLAR, and GDF3 are expressed in both seminoma and breast carcinoma.Cancer. 104(10):2255-2265

Ezeha U, R.L. Baehnera, K.L. Chewa, J.J. Fridlyanda, F.M. Waldmana and M.I. Cedars. (2007) The prognostic role of the human embryonic stem cell protein, NANOG, in breast cancer . Fertility and Sterility. 88(1):399

Farr G.H. 3rd, Ferkey D.M., Yost C., Pierce S.B., Weaver C. & Kimelman D. (2000) Interaction among GSK-3, GBP, axin, and APC in Xenopus axis specification. J. Cell Biol. 148:691-702

Fearon ER, Vogelstein B. (1990) A genetic model for colorectal tumorigenesis. Cell. 61(5):759-767. Review

Feldmann M., Brennan F. M., Maini R. N. (1996) Role of cytokines in rheumatoid arthritis. Annu. Rev. Immunol. 14(1):397-440

Fenoglio CM, Lane N. (1974) The anatomical precursor of colorectal carcinoma. Cancer 34(3): suppl: 819-823

Fernando RI, Litzinger M, Trono P, Hamilton DH, Schlom J, Palena C (2010) The T-box transcription factor Brachyury promotes epithelial-mesenchymal transition in human tumor cells. 120(2):533–544

Fidler, I. J. & Hart, I. R. (1982) Biological diversity in metastatic neoplasms: origins and implications. Science 217(1):998–1003

Fidler, I. J. & Kripke, M. L. (1977) Metastasis results from preexisting variant cells within a malignant tumor. Science 197(1):893–895

Fine, L. G., Holley, R. W., Nasri, H., and Badie-Dezfooly, B. (1985) BSC-1 growth inhibitor transforms a mitogenic stimulus into a hypertrophie stimulus for renal proximal tubular cells: relationship to Na<sup>+</sup>/H<sup>+</sup> antiport activity. Proc.Nati. Acad. Sci. USA, 82(1):6163-6166

Florez JC, Jablonski KA, Bayley N, Pollin TI, de Bakker PI, Shuldiner AR, Knowler WC, Nathan DM, Altshuler D. (2006) TCF7L2 polymorphisms and progression to diabetes in the Diabetes Prevention Program. N Engl J Med. 20;355(3):241-250

Fodde R, Smits R, Clevers H. (2001) APC, signal transduction and genetic instability in colorectal cancer. Nat Rev Cancer 1(1):55-56

Fodde R. and Brabletz, T., (2007) Wnt/beta-catenin signaling in cancer stemness and malignant behavior. Curr Opin Cell Biol. 19(1):150-158

Fogh J. & Trempe G.: (1975) New human tumour cell lines. In: Fogh, J. (Ed): Human tumour cells in vitro. Plenum Press. NY. 115-159

Forslund A, Lönnroth C, Andersson M, Brevinge H, Lundholm K. (2001) Mutations and allelic loss of p53 in primary tumor DNA from potentially cured patients with colorectal carcinoma. J Clin Oncol. 19(11):2829-2836

Franci C, Takkunen M, Dave N, Alameda F, Gomez S, Rodriguez R et al. (2006). Expression of Snail protein in tumor-stroma interface. Oncogene 25(1):5134–5144

Frea S, Pallavib S, Huyghea M, Laec M, Janssend K, Robinea S, Artavanis-Tsakonasb S, Louvarda D. (2009) Notch and Wnt Signals cooperatively control cell proliferation and tumorigenesis in the intestine. Proc. Natl. Acad. Sci.106 (15):6309-6314

Friedenreich CM. (2001) Physical activity and cancer prevention: from observational to intervention research. Cancer Epidemiol Biomarkers Prev 10(4): 287-301

Friedle H, Rastegar S, Paul H, Kaufmann E, and Knöchel W. (1998) Xvent-1 mediates BMP-4-induced suppression of the dorsal-lip-specific early response gene XFD-1' in Xenopus embryos. EMBO J. 17(8):2298–2307

Fujimoto, K., Beauchamp, R.D., and Whitehead, R.H. (2002) Identification and isolation of candidate human colonic clonogenic cells based on cell surface integrin expression. Gastroenterology. 123(1):1941–1948

Furlong et al., (1999) Evidence for the colonic origin of ovarian cancer cell line SW626. J. Natl. Cancer Inst. 91(1):1327-1328

Furuuchi K.(2000) Development of a detection system (APC yeast color assay) of APC mutations by color change of yeast. Hokkaido Igaku Zasshi. 75(6):385-397

Gaiano N, Fishell G (2002) The role of notch in promoting glial and neural stem cell fates. Annu. Rev. Neurosci. 25:471–90

Gaiano N, Nye JS, Fishell G (2000) Radial glial identity is promoted by Notch1 signaling in the murine forebrain. Neuron 26(2):395–404

Gailani M. R., Bale A. E. (1997) Developmental genes and cancer : Role of patched in basal cell carcinoma of the skin. Journal of the National Cancer Institute. 89(15):1103-1109

Garnis C, Campbell J, Davies J, Macaulay C, Lam S, Lam WL. (2005) Involvement of multiple developmental genes on chromosome 1p in lung tumorigenesis. Hum Mol Genet.15(4):475-82

Gebbia M, Ferrero GB, Pilia G, et al. (1997) X-linked situs abnormalities result from mutations in ZIC3. Nat. Genet. 17(3):305-308

Ghosh S., May M. J., Kopp E. B. (1998) NFkB and Rel proteins: evoluntionary conserved mediators of immune responses. Annu. Rev. Immunol. 16(1):225-260

Giannakis, M., Stappenbeck, T.S., Mills, J.C., Leip, D.G., Lovett, M., Clifton, S.W., Ippolito, J.E., Glasscock, J.I., Arumugam, M., Brent, M.R (2006) Molecular properties of adult mouse gastric and intestinal epithelial progenitors in their niches. J. Biol. Chem. 281(1):11292–11300

Giannelli G, Bergamini C, Fransvea E, Sgarra C, Antonaci S. (2005) Laminin-5 with transforming growth factor-beta1 induces epithelial to mesenchymal transition in hepatocellular carcinoma. Gastroenterology. 129(5):1375-1383

Gibbons DL, Lin W, Creighton CJ, Rizvi ZH, Gregory PA, Goodall GJ, Thilaganathan N, Du L, Zhang Y, Pertsemlidis A, Kurie JM (2009) Contextual extracellular cues promote tumor cell EMT and metastasis by regulating miR-200 family expression. Genes Dev. 23(18):2140-2151

Gidekel S, Pizov G. (2003) Oct-3/4 is a dose-dependent oncogenic fate determinant, Cancer Cell 4(1):361–370 Giles, R. H., van Es, J. H., and Clevers, H. (2003) Biochim Biophys Acta. 1653(1):1-24

Gilmore TD (2006) Introduction to NF-kB: players, pathways, perspectives. Oncogene 25(51):6680–6684

Gluecksohn-Schoenheimer, S. (1944) The development of normal and homozygous Brachy (T/T) mouse embryos in the extraembryonic coelom of the chick. Proc. Natl. Acad. Sci. 30(1):134-140

Gokhale PJ, Giesberts AM, Andrews PW. (2000) Brachyury is expressed by human teratocarcinoma cells in the absence of mesodermal differentiation. Cell Growth Differ. 11(1):157-162

Gong Y, Slee RB, Fukai N, Rawadi G, Roman-Roman S, Reginato AM, Wang H, Cundy T, Glorieux FH, Lev D, Zacharin M, Oexle K, Marcelino J, Suwairi W, Heeger S, Sabatakos G, Apte S, Adkins WN, Allgrove J, Arslan-Kirchner M, Batch JA, Beighton P, Black GC, Boles RG, Boon LM, Borrone C, Brunner HG, Carle GF, Dallapiccola B, De Paepe A, Floege B, Halfhide ML, Hall B, Hennekam RC, Hirose T, Jans A, Jüppner H, Kim CA, Keppler-Noreuil K, Kohlschuetter A, LaCombe D, Lambert M, Lemyre E, Letteboer T, Peltonen L, Ramesar RS, Romanengo M, Somer H, Steichen-Gersdorf E, Steinmann B, Sullivan B, Superti-Furga A, Swoboda W, van den Boogaard MJ, Van Hul W, Vikkula M, Votruba M, Zabel B, Garcia T, Baron R, Olsen BR, Warman ML (2001) LDL receptor-related protein 5 (LRP5) affects bone accrual and eye development. Cell. 16;107(4):513-23

Gottardi C.J., Wong E. & Gumbiner B.M. (2001) E-cadherin suppresses cellular transformation by inhibiting beta-catenin signaling in an adhesion-independent manner. J. Cell Biol. 153(1):1049-1060

Grant SF, Thorleifsson G, Reynisdottir I, Benediktsson R, Manolescu A, Sainz J, Helgason A, Stefansson H, Emilsson V, Helgadottir A, Styrkarsdottir U, Magnusson KP, Walters GB, Palsdottir E, Jonsdottir T, Gudmundsdottir T, Gylfason A, Saemundsdottir J, Wilensky RL, Reilly MP, Rader DJ, Bagger Y, Christiansen C, Gudnason V, Sigurdsson G, Thorsteinsdottir U, Gulcher JR, Kong A, Stefansson K. (2006)Variant of transcription factor 7-like 2 (TCF7L2) gene confers risk of type 2 diabetes. Nat Genet. 38(3):320-323

Gregorieff, A., Pinto, D., Begthel, H., Destree, O., Kielman, M., and Clevers, H. (2005) Expression pattern of Wnt signalingcomponents in the adult intestine. Gastroenterology. 129(1):626–638

Gridley T. (2003) Notch signaling and inherited disease syndromes. Hum Mol Genet.12(1):R9–R13.

Groden J., Thliveris A., Samowitz W., Carlson M., Gelbert L., Albertsen H., Joslyn G., Stevens J., Spirio L., Robertson M., et al. (1991) Identification and characterization of the familial adenomatous polyposis coli gene. Cell. 66(1):589-600

Gross I, Duluc I, Benameur T, Calon A, Martin E, Brabletz T, Kedinger M, Domon-Dell C, Freund JN.(2008) The intestine-specific homeobox gene Cdx2 decreases mobility and antagonizes dissemination of colon cancer cells. Oncogene. 27(1):107-115

Gross, I Duluc, T Benameur, A Calon, E Martin, T Brabletz, M Kedinger, C Domon-Dell and J-N Freund (2008). The intestine-specific homeobox gene Cdx2 decreases mobility and antagonizes dissemination of colon cancer cells. Oncogene 27(1):107–115

Guo Y, Costa R, Ramsey H, Starnes T, Vance G, Robertson K, Kelley M, Reinbold R, Scholer H, Hromas R . (2002). The embryonic stem cell transcription factors Oct-4 and FoxD3 interact to regulate endodermal-specific promoter expression. PNAS. 99(6):3663-3667

Haber RS, Rathan A, Weiser KR, Pritsker A, Itzkowitz SH, Bodian C, Slater G, Weiss A, Burstein DE (1998) Glut1 glucose transporter expression in colorectal carcinoma, A marker for poor prognosis. Cancer. 83(1):34-40

Hadjihannas M, Martina, Boris J, Walter B, Wolfgang D (2005) Aberrant Wnt  $\beta$ catenin signalling can induce chromosomal instability in colon cancer. PNAS 8(1):1711–1717

Hadland BK, Huppert SS, Kanungo J et al. (2004) A requirement for Notch1 distinguishes 2 phases of definitive hematopoiesis during development. Blood. 2004(1):104:3097–3105

Han G, Li AG, Liang YY, Owens P, He W, Lu S, Yoshimatsu Y, Wang D, Ten Dijke P, Lin X, Wang XJ. (2006) Smad7-induced beta-catenin degradation alters epidermal appendage development. Dev Cell. 11(3):301-312

Hao J, Li TG, Qi X, Zhao DF, Zhao GQ. (2006) WNT/beta-catenin pathway upregulates Stat3 and converges on LIF to prevent differentiation of mouse embryonic stem cells.Dev Biol. 290(1):81-91

Harada, N., Tamai, Y., Ishikawa, T., Sauer, B., Takaku, K., Oshima, M., and Taketo, M.M. (1999) Intestinal polyposis in mice with a dominant stable mutation of the  $\beta$ -catenin gene. EMBO J. 18(1):5931-5942

Haramis AP, Begthel H, Van Den Born M, van Es J, Jonkheer S, et al. (2004) De novo crypt formation and juvenile polyposis on BMP inhibition in mouse intestine. Science. 303(1):1684-1686

Hardwick JC, Van Den Brink GR, Bleuming SA, Ballester I, Van Den Brande JM, et al. (2004) Bone morphogenetic protein 2 is expressed by, and acts upon, mature epithelial cells in the colon. Gastroenterology 126(1):111–121

Hart L. A., Krishnan V. L., Adcock I. M., Barnes P. J., Chung K. F. (1998) Activation and localization of transcription factor, nuclear factor-kappaB, in asthma. Am J Respir Crit Care Med.158(1):1585–1592

Hart M., Concordet J.P., Lassot I., Albert I., del los Santos R., Durand H., Perret C., Rubinfeld B., Margottin F., Benarous R. & Polakis P. (1999) The F-box protein  $\beta$ -TrCP associates with phosphorylated  $\beta$ -catenin and regulates its activity in the cell. Curr. Biol. 9(1):207-210

Hart, A. H., Hartley, L., Ibrahim, M. & Robb, L. (2004) Identification, cloning and expression analysis of the pluripotency promoting Nanog genes in mouse and human. Dev. Dyn. 230(1):187–198

Hart AH, Hartley L, Parker K, Ibrahim M, Looijenga LH, Pauchnik M, Chow CW, Robb L. (2005) The pluripotency homeobox gene NANOG is expressed in human germ cell tumors. Cancer 104(10):2092-2098

Hatano, S. Y. et al. (2005) Pluripotential competence of cells associated with Nanog activity. Mech. Dev. 122(1):67-79

Hayden, M. S. & Ghosh, S. (2004) Signaling to NFkB. Genes Dev. 18(1):2195-2224

He, T.C., Sparks, A.B., Rago, C., Hermeking, H., Zawel, L., da Costa, L.T., Morin, P.J., Vogelstein, B. and Kinzler, K.W. (1998) Identification of c-MYC as a target of the APC pathway. Science 281(1):1509–1512

He, X.C., Zhang, J., Tong, W.G., Tawfik, O., Ross, J., Scoville, D.H., Tian, Q., Zeng, X., He, X., Wiedemann, L.M., et al. (2004) BMP signaling inhibits intestinal stem cell self-renewal through suppression of Wnt signaling. Nat.Genet. 36:1117–1121

Heath, J.P. (1996) Epithelial cell migration in the intestine. CellBiol. Int. 20(1):139-146

Hecht A & Kemler R. (2000) Curbing the nuclear activities of  $\beta$ -catenin. Control over Wnt target gene expression. EMBOReports 1(1):24–28
Hee CK, Nicoll SB. (2009) Endogenous bone morphogenetic proteins mediate 1alpha, 25-dihydroxyvitamin D(3)-induced expression of osteoblast differentiation markers in human dermal fibroblasts. J Orthop Res. 27(2):162-168

Hemmati HD, Nakano I, Lazareff JA, Masterman-Smith M, Geschwind DH, Bronner-Fraser M, Kornblum HI (December 2003) Cancerous stem cells can arise from pediatric brain tumors. Proceedings of the National Academy of Sciences of the United States of America. 100(25):15178–1583

Henderson B.R. (2000) Nuclear-cytoplasmic shuttling of APC regulates  $\beta$ -catenin subcellular localization and turnover. Nat. Cell Biol. 2(1):653-660

Henderson BR, Galea M, Schuechner S. (2002) Lymphoid enhancer factor-1 blocks denomatous polyposis coli-mediated nuclear export and degradation of beta-catenin. Regulation by histone deacetylase 1. J Biol Chem 277(1):24258-24264

Heppner, G. H. (1984) Tumor heterogeneity. Cancer Res. 44(1):2259-2265

Herrmann B G; Labeit S; Poustka A; King T R; Lehrach H. (1990) Cloning of the T gene required in mesoderm formation in the mouse. Nature. 343(6259):617-622

Herrmann BG. (1991) Expression pattern of the Brachyury gene in wholemount Tw'l/T l mutant embryos. Development. 113(1):913-917

Hexige S, Guo J, Ma L, Sun Y, Liu X, Ma L, Yan X, Li Z, Yu L. (2005) Expression pattern of growth/differentiation factor 3 in human and murine cerebral cortex, hippocampus as well as cerebellum. Neurosci Lett 389(2):83-87

Hirohashi S, Kanai Y. (2003) Cell adhesion system and human cancer morphogenesis. Cancer Sci.94(1):575-581

Hmadcha A., Juan DB, Jane W, Mohamed A, Bernat S. (2009) The immune boundaries for stem cell based therapies: problems and prospective solutions. J Cell Mol Med. 8A(1):1464-1475

Hodgson, S. V. and Maher, E. R. (1993) A practical guide to human cancer genetics. Cambridge University Press, Cambridge

Hoei-Hansen CE, Almstrup K, Nielsen JE, Brask Sonne S, Graem N, Skakkebaek NE, Leffers H, Rajpert-De Meyts E. (2005) Stem cell pluripotency factor NANOG is expressed in human fetal gonocytes, testicular carcinoma in situ and germ cell tumours. Histopathology. 47(1):48-56

Hoffmann A., Natoli G., Ghosh G. (2006) Transcriptional regulation via the NFkB signaling module. Oncogene. 25(1)6706-6716

Horst D, Budczies J, Brabletz T, Kirchner T, Hlubek F. (2009) Invasion associated up-regulation of nuclear factor kappaB target genes in colorectal cancer. Cancer. [Epub ahead of print]

Horst D, L Kriegl, J Engel, T Kirchner and A Jung. (2008) CD133 expression is an independent prognostic marker for low survival in colorectal cancer. British Journal of Cancer. 99(1):1285-1289

Hottaa, Hiroki Takahashib, Tomomi Asakuraa, Banjo Saitoha, Naohito Takatoria, Yutaka Satoua and Nori Satoh. (2000) Characterization of Brachyury-Downstream Notochord Genes in the Ciona intestinalis Embryo. Developmental Biology. 224(1):69-80

http://www.stanford.edu/~rnusse/pathways/targets.html

Hu MC, Rosenblum ND. (2005) Smad1, beta-catenin and Tcf4 associate in a molecular complex with the Myc promoter in dysplastic renal tissue and cooperate to control Myc transcription. Development 132(1):215-225

Huang, X., and J.P. Saint-Jeannet. (2004) Induction of the neural crest and the opportunities of life on the edge. Dev. Biol. 275(1):1–11

Huang H., Fujii H., Sankila A., Mahler-Araujo B.M., Matsuda M., Cathomas G & Ohgaki H. (1999) □-catenin mutations are frequent in human hepatocellular carcinomas associated with hepatitic C virus infection. Am. J. Pathol. 155(1):1795-1801

Huang H., Mahler-Araujo B.M., Sankila A., Chimelli L., Yonekawa Y., Kleihues P. & Ohgaki H. (2000) APC mutations in sporadic medulloblastomas. Am. J. Pathol. 156(1):433-437

Huber MA, Azoitei N, Beug H, Wirth T. (2004) NF- $\kappa$ B is essential for epithelialmesenchymal transition and metastasis in a model of breast cancer progression. The Journal of clinical investigation. 114(4):569-581

Huber MA, Kraut N, Beug H. (2005) Molecular requirements for epithelialmesenchymal transition during tumor progression. Curr Opin Cell Biol 17(1):548-558

Hugo H, Ackland ML, Blick T, Lawrence MG, Clements JA, Williams ED, Thompson EW. (2007) Epithelial--mesenchymal and mesenchymal--epithelial transitions in carcinoma progression.J Cell Physiol. 213(2):374-383 Hunt, M. (2009). Real Time PCR. Available: http://pathmicro.med.sc.edu/pcr/realtime-home.htm. Accessed 08 February 2010

Hurlin P, Neal G.Copeland, Nancy A.Jenkins and Robert N.Eisenman. (1999). Mga, a dual-specificity transcription factor that interacts with Max and contains a T-domain DNAbinding motif. The EMBO Journal.18 (24):7019-7028

Hussein SM, Duff EK, Sirard C. (2003) Smad4 and beta-catenin co-activators functionally interact with lymphoid-enhancing factor to regulate graded expression of Msx2. J Biol Chem. 278(49):48805-14

Hutchison CJ (2002) Lamins: building blocks or regulators of gene expression? Nat Rev Mol Cell Biol 3:848–858

Hyslop L, Stojkovic M, Armstrong L, Walter T, Stojkovic P, Przyborski S, Herbert M, Murdoch A, Strachan T, Lako M. (2005). Downregulation of NANOG induces differentiation of human embryonic stem cells to extraembryonic lineages. Stem Cells. 23(8):1035-1043

Ichii S., Takeda S., Horii A., Nakatsuru S., Miyoshi Y., Emi M., Fujiwara Y., Koyama K., Furuyama J., Utsunomiya J., et al. (1993) Detailed analysis of genetic alterations in colorectal tumors from patients with and without familial adenomatous polyposis (FAP). Oncogene 8(1):2399-2405

Ilaria Malanchi, Hector Peinado, Deepika Kassen, Thomas Hussenet, Daniel Metzger, Pierre Chambon, Marcel Huber, Daniel Hohl, Amparo Cano, Walter Birchmeier & Joerg Huelsken. (2008) Cutaneous cancer stem cell maintenance is dependent on bold beta-catenin signalling. Nature. 452(1):650-653

Illemann M, Bird N, Majeed A, Laerum OD, Lund LR, Danø K, Nielsen BS. (2009) Two distinct expression patterns of urokinase, urokinase receptor and plasminogen activator inhibitor-1 in colon cancer liver metastases. Int J Cancer. 124(8):1860-1870

Inês Ribeiro, Yasuhiko Kawakami, Dirk Büscher, Ángel Raya, Joaquín Rodríguez-León, Masanobu Morita, Concepción Rodríguez Esteban, and Juan Carlos Izpisúa Belmonte. (2007) Tbx2 and Tbx3 Regulate the Dynamics of Cell Proliferation during Heart Remodeling. PLoS ONE. 2(4):398

Inomata M, Ochiai A, Akimoto S, Kitano S and Hirohashi S (1996) Alteration of bcatenin expression in colonic epithelial cells of familial adenomatous polyposis patients. Cancer Res 56(1):2213–2217

Ito M and et al. (2002) NOD/SCID/ $\gamma^{null}$  mouse: an excellent recipient mouse model for engraftment of human cells. Blood 100(1):3175–3182

J. Hao, T.G. Li, X. Qi, D.F. Zhao and G.Q. Zhao. (2006) WNT/beta-catenin pathway up-regulates Stat3 and converges on LIF to prevent differentiation of mouse embryonic stem cells, Dev. Biol. 290(1):81–91

Janda E, Lehmann K, Killisch I, Jechlinger M, Herzig M, Downward J, Beug H, Grunert S. (2002) Ras and TGFb cooperatively regulate epithelial cell plasticity and metastasis: dissection of Ras signaling pathways. J Cell Biol. 156(1):299-313

Jansen MPHM, Machiels BM, Hopman AHN, Broers JLV, Bot FJ, Arends JW, Ramaekers FCS and Schouten HC (1997) Comparison of A- and B-type lamin expression in nodular sclerosing Hodgkin's disease and reactive lymph nodes. Histopathology 4(1):304–312

Jean Paul Thiery. (2003) Epithelial-mesenchymal transitions in development and pathologies. Current Opinion in Cell Biology. 15(1):740-746

Jen J., Powell S.M., Papadopoulos N., Smith K.J., Hamilton S.R., Vogelstein B. & Kinzler K.W. (1994) Molecular determinants of dysplasia in colorectal lesions. Cancer Res. 54(1):5523-5526

Jeter CR, Badeaux M, Choy G, Chandra D, Patrawala L,Liu C, Calhoun-Davis T, Zaehres H, Q Daley, Tang DG. (2008) Functional Evidence that the Self-Renewal Gene NANOG Regulates Human Tumor Development. Stem cells 27(1):993–1005

Jiang YG, Luo Y, He DL, Li X, Zhang LL, Peng T, Li MC, Lin YH.(2007). Role of Wnt/beta-catenin signaling pathway in epithelial-mesenchymal transition of human prostate cancer induced by hypoxia-inducible factor-1alpha. Int J Urol.14(11):1034-1039

Jim Smith (1999) T-box genes: what they do and how they do it. Trends in Genetics 15(4):154-158

Jong P, Young KC, Choon HS, Chulso M. (2008) Epigenetic silencing of human T (brachyury homologue) gene in non-small-cell lung cancer. Biochemical and Biophysical Research Communications. 365(2):221-226

Jonkers J, Korswagen HC, Acton D, Breuer M, Berns A. (1997) Activation of a novel proto-oncogene, Frat1, contributes to progression of mouse T-cell lymphomas. EMBO J. 16(3):441-450

Jung, J, Zheng, M, Goldfarb M, Zaret K. S. (1999) Initiation of mammalian liver development from endoderm by fibroblast growth factors. Science 284(1):1998-2003

Kaidi A, Williams AC, Paraskeva C. (2007) Interaction between beta-catenin and HIF-1 promotes cellular adaptation to hypoxia. Nat Cell Biol. 9(2):210-217

Kaimaktchiev V, Terracciano L, Tornillo L, Spichtin H, Stoios D, Bundi M et al. (2004) The homeobox intestinal differentiation factor CDX2 is selectively expressed in gastrointestinal adenocarcinomas. Mod Pathol 17(1):1392–1399

Kamberov, E., Makarova, O., Roh, M., Liu, A., Karnak, D., Straight, S. and Margolis, B. (2000). Molecular cloning and characterization of Pals, proteins associated with mLin-7. J. Biol. Chem. 275(1):11425-11431

Kameda T, Hatakeyama S, Terada K and Sugiyama T (2001) Acceleration of the formation of cultured epithelium using the sonic hedgehog expressing feeder cells. Tissue Eng 7(1):545–555

Kang Y and Massague J (2004) Epithelial-Mesenchymal Transitions. Twist in development and metastasis. Cell 118(1):277-279

Karoubi G, Gugger M, Schmid R, Dutly A. (2009) OCT4 expression in human nonsmall cell lung cancer: implications for therapeutic intervention, Interact Cardiovasc Thorac Surg., 8(4):393-397

Karreth F, Tuveson DA. (2004) Twist induces an epithelial-mesenchymal transition to facilitate tumour metastasis. Cancer Biol Ther 3(1):1058–1059

Kasper S. (2008) Exploring the origins of the normal prostate and prostate cancer stem cell. Stem Cell Rev. 4(3):193-201

Katoh M, Katoh M. (2009) Integrative genomic analyses of WNT11: transcriptional mechanisms based on canonical WNT signals and GATA transcription factors signaling. Int J Mol Med. 24(2):247-251

Kaur, P. and Potten, C.S. (1986) Cell migration velocities in the crypts of the small intestine after cytotoxic insult are not dependent on mitotic activity. Cell Tissue Kinet.19(1):601–610

Kayahara, T., Sawada, M., Takaishi, S., Fukui, H., Seno, H., Fukuzawa, H., Suzuki, K., Hiai, H., Kageyama, R., Okano, H., et al. (2003) Candidate markers for stem and early progenitor cells, Musashi-1 and Hes1, are expressed in crypt base columnar cells of mouse small intestine. FEBS Lett.535(1):131–135

Keller M, Ezaki T, Guo R, and John P. Lynch. (2004). Cdx1 or Cdx2 expression activates E-cadherin-mediated cell-cell adhesion and compaction in human COLO 205 cells. Am J Physiol Gastrointest Liver Physiol. 287(1):G104-G114

Kibar Z, Torban E, McDearmid JR, Reynolds A, Berghout J, Mathieu M, Kirillova I, De Marco P, Merello E, Hayes JM, Wallingford JB, Drapeau P, Capra V, Gros P. (2007) Mutations in VANGL1 associated with neural-tube defects. N Engl J Med. 5(14):1432-1437

Kinzler K. & Vogelstein B. (1996) Lessons from hereditary colorectal cancer. Cell 87(1):159-170

Kim JT, Joo CK. (2002) Involvement of cell–cell interactions in the rapid stimulation of Cas tyrosine phosphorylation and Src kinase activity by transforming growth factor b1. J Biol Chem. 277(1):31938-31948

Kinzler, K. W. & Vogelstein, B. Lessons from hereditary colorectal cancer. (1996) Cell 87(1):159-170

Kirchner T, Brabletz T. (2000) Patterning and nuclear beta-catenin expression in the colonic adenoma-carcinoma sequence. Analogies with embryonic gastrulation. Am J Pathol. 157(1):1113-1121

Kispert A, Herrmann BG, Leptin M, Reuter R. (1994) Homologs of the mouse Brachyury gene are involved in the specification of posterior terminal structures in Drosophila, Tribolium, and Locusta. Genes & Dev.8(1):2137–2150

Kispert, A and Herrmann, B G. (1993) The Brachyury gene encodes a novel DNA binding protein. EMBO J. 12(8):3211–3220

Kispert, A., Koschorz, B. and Herrmann, B. G. (1995a) The T protein encoded by Brachyury is a tissue-specific transcription factor. EMBO J. 14(1):4763-4772

Kitaeva MN, Grogan L, Williams JP, Dimond E, Nakahara K, Hausner P, DeNobile JW, Soballe PW, Kirsch IR. (1997) Mutations in beta-catenin are uncommon in colorectal cancer occurring in occasional replication error-positive tumors. Cancer Res. 57(20):4478-4481

Koblas T, Pektorova L, Zacharovova K, Berkova Z, Girman P, Dovolilova E, Karasova L, Saudek F. (2008) Differentiation of CD133-positive pancreatic cells into insulin-producing islet-like cell clusters. Transplant Proc. 40(1):415-418

Kolodner R. (1996) Biochemistry and genetics of eukaryotic mismatch repair. Genes Dev. 10(1):1433-1442

Kondo T, Johnson SA, Yoder MC, Romand R and Hashino E (2005) Sonic hedgehog and retinoic acid synergistically promote sensory fate specification from bone marrow-derived pluripotent stem cells. Proc Natl Acad Sci.USA 102(1):4789-4794 Konishi M., Kikuchi-Yanoshita R., Tanaka K., Muraoka M., Onda A., Okumura Y., Kishi N., Iwama T., Mori T., Koike M., Ushio K., Chiba M., Nomizu S., Konishi F., Ustunomiya J. & Miyaki M. (1996) Molecular nature of colon tumours in hereditary nonpolyposis colon cancer, familial poyposis and sporadic colon cancer. Gastroenterology 111(1):307-317

Korinek V, Barker N, Morin PJ, van Wichen D, de Weger R, Kinzler KW, Vogelstein B and Clevers H (1997) Constitutive transcriptional activation by a bcatenin-Tcf complex in APC2/2 colon carcinoma. Science 275(1):1784–1787

Korinek, V., Barker, N., Moerer, P., van Donselaar, E., Huls, G., Peters, P.J., and Clevers, H. (1998) Depletion of epithelial stem-cell compartments in the small intestine of mice lacking Tcf-4. Nat. Genet. 19(1):379–383

Korinek V et al., (1998) Two members of the Tcf family implicated in Wnt/betacatenin signaling during embryogenesis in the mouse. Mol Cell Biol.18(3):1248-1256

Kosinski C, Li VS, Chan AS, Zhang J, Ho C, Tsui WY, Chan TL, Mifflin RC, Powell DW, Yuen ST, Leung SY, Chen X. (2007) Gene expression patterns of human colon tops and basal crypts and BMP antagonists as intestinal stem cell niche factors. Proc Natl Acad Sci U S A. 104(39):15418-15423

Kramps T, Peter O, Brunner E, Nellen D, Froesch B, Chatterjee S, Murone M, Züllig S, Basler K. (2002) Wnt/wingless signaling requires BCL9/legless-mediated recruitment of pygopus to the nuclear beta-catenin-TCF complex.Cell. 109(1):47-60

Kristiansen G, Pilarsky C, Wissmann C, Stephan C, Weissbach L, Rosenthal A. (2003) ALCAM/CD166 is up-regulated in low-grade prostate cancer and progressively lost in high-grade lesions. The Prostate. 54(1):34-43

Kuhnert, F., Davis, C.R., Wang, H.T., Chu, P., Lee, M., Yuan, J., Nusse, R., and Kuo, C.J. (2004) Essential requirement for Wnt signaling in proliferation of adult small intestine and colon revealed by adenoviral expression of Dickkopf-1. Proc. Natl.Acad. Sci. 101(1):266–271

Kumano K, Chiba S, Kunisato A et al. (2003) Notch1 but not Notch2 is essential for generating hematopoietic stem cells from endothelial cells. Immunity. 18(1):699-711

Kummermehr, J. & Trott, K.-R. in Stem Cells (ed. Potten, C. S.) (1997) 1(1):363-399 (Academic, New York)

Kuniyasu H, Ohmori H, Sasaki T, Sasahira T, Yoshida K, Kitadai Y, Fidler IJ. (2003) Production of interleukin 15 by human colon cancer cells is associated with

induction of mucosal hyperplasia, angiogenesis, and metastasis. Clin Cancer Res.9(13):4802-4810

Kusch T, Reuter R. (1999) Functions for Drosophila brachyenteron and forkhead in mesoderm specification and cell signalling. Development.126:3991–4003

Kusch, Tobias Storck, Uwe Walldorf and Rolf Reuter. (2002) Brachyury proteins regulate target genes through modular binding sites in a cooperative fashion. Genes Dev.16(4):518–529

LaBarge MA, Bissell MJ. (2008) Is CD133 a marker of metastatic colon cancer stem cells? J Clin Invest. 118(6):2021-2024

Laken S.J., Papadopoulos N., Petersen G.M., Gruber S.B., Hamilton S.R., Giardiello F.M., Brensinger J.D., Vogelstein B. & Kinzler K.W. (1999) Analysis of masked mutations in familial adenomatous polyposis. Proc. Natl. Acad. Sci USA 96(1):2322-2326

Lamlum H., Ilyas M., Rowan A., Clark S., Johnson V., Bell J., Frayling I., Efstathiou J., Pack K., Payne S., Roylance R., Gorman P., Sheer D., Neale K., Phillips R., Talbot I., Bodmer W. & Tomlinson I. (1999) The type of somatic mutation at APC in familial adenomatous polyposis is determined by the site of the germline mutation: a new facet to Knudson's 'two-hit' hypothesis. Nature Med. 5(1):1071-1075

Lammi L, Arte S, Somer M, Jarvinen H, Lahermo P, Thesleff I, Pirinen S, Nieminen P. (2004) Mutations in AXIN2 cause familial tooth agenesis and predispose to colorectal cancer. Am J Hum Genet. 74(5):1043-50

Lamorte L, Royal I, Naujokas M, Park M. (2002) Crk adapter proteins promote an epithelial–mesenchymal-like transition and are required for HGF-mediated cell spreading and breakdown of epithelial adherens junctions. Mol Biol Cell. 13(1):1449-1461

Lang Sh, Frame F, Collins A. (January 2009) Prostate cancer stem cells. J. Pathol. 217(2):299–306

Laurens G. van der Flier, Hans Clevers (2009) Stem Cells, Self-Renewal, and Differentiation in the Intestinal Epithelium. Annual Review of Physiology. Vol. 71(1):241-260

Le NH, Franken P, Fodde R. (2008) Tumour-stroma interactions in colorectal cancer: converging on beta-catenin activation and cancer stemness. Br J Cancer. 98(1):1886-1893. Review

Leibovitz A., Stinson J.C., McCombs W.B. 3<sup>rd</sup>, McCoy C.E., Mazur K.C. & Mabry N.D. (1976) Classification of human colorectal adenocarcinoma cell lines. Cancer Res. 36(1):4562-4569

Len H, FrankenN P, Fodde R. (2000) Tumour-stroma interactions in colorectal cancer : converging on  $\beta$ -catenin activation and cancer sternness. British Journal of Cancer. 98(12):1886-1893

Léon C. L. T. van Kempen, Joost J. van den Oord, Goos N. P. van Muijen, Ulrich H. Weidle, Henri P. J. Bloemers and Guido W. M. Swart. (2000) Activated Leukocyte Cell Adhesion Molecule/CD166, a Marker of Tumor Progression in Primary Malignant Melanoma of the Skin . American Journal of Pathology. 156(1):769-774

Levy D.B., Smith K.J., Beazer-Barclay Y., Hamilton S.R., Vogelstein B. & Kinzler K.W. (1994) Inactivation of both APC alleles in human and mouse tumors. Cancer Res. 54(1):5953-5958

Li C, Heidt DG, Dalerba P, Burant CF, Zhang L, Adsay V, Wicha M, Clarke MF, Simeone DM (February 2007) Identification of pancreatic cancer stem cells. Cancer research 67(3):1030-1037

Li L., Yuan H., Weaver C.D., Mao J., Farr G.H. III, Sussman D.J., Jonkers J., Kimelman D. & Wu D. (1999) Axin and Frat1 interact with Dvl and GSK, bridging Dvl to GSK in Wnt-mediated regulation of LEF-1. EMBO J. 18(1):4233-4240

Li Q., Verma I. M. (2002) NFkB regulation in the immune system. Nat. Rev. Immunol. 2(1):725-734

Li, Q.Y., et al. (1997). Holt-Oram syndrome is caused by mutations in TBX5, a member of the Brachyury (T) gene family. Nat Genet 15(1):21-29

Liang H, Chen Q, Coles AH, Anderson SJ, Pihan G, Bradley A, Gerstein R, Jurecic R, Jones SN. Cancer Cell. (2003) Wnt5a inhibits B cell proliferation and functions as a tumor suppressor in hematopoietic tissue. 4(5):349-360

Liang H, Chen Q, Coles AH, Anderson SJ, Pihan G, Bradley A, Gerstein R, Jurecic R, Jones SN. Liebner Stefan; Cattelino Anna; Gallini Radiosa; Rudini Noemi; Iurlaro Monica; Piccolo Stefano; Dejana Elisabetta. (2004)  $\beta$ -catenin is required for endothelial-mesenchymal transformation during heart cushion development in the mouse. The Journal of cell biology. 166(3):359-367

Liu C., Kato Y., Zhang Z., Do V.M., Yankner B.A. & He X. (1999)  $\beta$ -TrCP couples  $\beta$ -catenin phosphorylation-degradation and regulates Xenopus axis formation. Proc. Natl. Acad. Sci. USA. 96(1):6273-6278

Liu Q, Teh M, Ito K, Shah N, Ito Y, Yeoh KG (December 2007). CDX2 expression is progressively decreased in human gastric intestinal metaplasia, dysplasia and cancer. Mod. Pathol. 20(12):1286-1297

Liu Y, El-Naggar S, Darling DS, Higashi Y, Dean DC. (2008) Zeb1 links epithelialmesenchymal transition and cellular senescence. Development 135(1):579-588

Livak KJ and Schmittgen TD. (2001) Analysis of relative gene expression data using Real-Time quantitative PCR and the  $2^{-\Delta \Delta C}$ <sub>T</sub> Method

Lluis F, Pedone E, Pepe S, Cosma MP. (2008) Periodic activation of Wnt/betacatenin signaling enhances somatic cell reprogramming mediated by cell fusion. Cell Stem Cell. 3:493-507. Methods. 25(4):402-408

Looijenga L.H., Stoop H., de Leeuw H.P. (2003) POU5F1 (OCT3/4) identifies cells with pluripotent potential in human germ cell tumors. Cancer Res. 63(1):2244–2250

Lorentz O, Duluc I, Arcangelis AD, Simon-Assmann P, Kedinger M and Freund JN (1997) Key role of the CDX2 homeobox gene in extracellular matrix-mediated intestinal cell differentiation. J Cell Biol 139(6):1553-1565

Lowenstein, E. J., Daly, R. J., Batzer, A. G., Li, W., Margolis, B., Lammers, R., Ullrich, A., Skolnik, E. Y., Bar-Sagi, D., and Schlessinger, J. (1992) Cell 70(1):431-442

Lustig, B., Jerchow, B., Sachs, M., Weiler, S., Pietsch, T., Karsten, U., van de Wetering, M., Clevers, H., Schlag, P. M., Birchmeier, W. & Behrens, J. (2002) Negative feedback loop of Wnt signaling through upregulation of conductin/axin2 in colorectal and liver tumors. Mol. Cell. Biol. 22(1):1184–1193

Machiels BM, Ramaekers FCS, Helma JH, Kuijpers HJH, Groenewould JS, Oosterhuis JO and Looijenga LHJ (1997) Nuclear lamin expression in normal testis and testicular germ cell tumours of adolescents and adults. J Cell Biochem 62(1):275–289

Madison BB, Braunstein K, Kuizon E, Portman K, Qiao XT, Gumucio DL. (2005) Epithelial hedgehog signals pattern the intestinal crypt-villus axis. Development 132(1):279–289

Mai, M., Qian, C., Yokomizo, A., Smith, D. I. & Liu, W. (1999) Cloning of the human homolog of conductin (AXIN2), a gene mapping to chromosome 17q23-q24. Genomics 55(1):341–344

Maitland NJ, Collins AT (June 2008) Prostate cancer stem cells: a new target for therapy. J. Clin. Oncol. 26(17):2862–2870

Major MB, Camp ND, Berndt JD, Yi X, Goldenberg SJ, Hubbert C, Biechele TL, Gingras AC, Zheng N, Maccoss MJ, Angers S, Moon RT. (2007) Wilms tumor suppressor WTX negatively regulates WNT/beta-catenin signaling. Science. 18;316(5827):1043-1046

Mancini A, Koch A, Wilms R, Tamura T. (2002) The SH2-containing inositol-5phosphatase (SHIP)-1 is implicated in the control of cell–cell junction and induces dissociation and dispersion of MDCK cells. Oncogene. 21(1):1477-1484

Mani SA, Yang J, Guo W, Brooks M, Reinhard F, Zhang C, Shipitsin M, CampbellLL, Polyak K, Brisken C, Yang J, Weinberg RA (2007) Mesenchyme Forkhead 1 (FOXC2) plays a key role in metastasis and is associated with aggressive basal-like breast cancers. PNAS 104(24):10069-10074

Mani SA, Guo W, Liao MJ, Eaton E, Ayyanan A, Zhou A, Brooks M, Reinhard F, Zhang C, Shipitsin M, CampbellLL, Polyak K, Brisken C, Yang J, Weinberg RA (2008) The epithelial-mesenchymal transition generates cells with properties of stem cells. Cell. 133(1):704-715

Maniatis T. (1999) A ubiquitin ligase complex essential for the NF- $\kappa$ B, Wnt/Wingless, and Hedgehog signalling pathways. Genes Dev. 13(1):505-510

Marcial A, Carlson L, and Madara L. (1984) Partitioning of Paracellular Conductance Along the Ileal Crypt-Villus Axis: A Hypothesis Based on Structural Analysis with Detailed Consideration of Tight Junction Structure-Function Relationships. J. Membrane Biol.80(1):59-70

Markowitz S, Wang J, Myeroff L, Parsons R, Sun L, Lutterbaugh J, Fan RS, Zborowska E, Kinzler KW, Vogelstein B, et al. (1995) Inactivation of the type II TGF-b receptor in colon chim. Biophys. Acta 1242(1):185–199

Markowitz S. (2000) TGF- $\beta$  receptors and DNA repair genes, coupled targets in a pathway of human colon carcinogenesis. Biochim. Biophys. Acta. 1470(1):M13-M20

Marshman E, Booth C, Potten CS. (2002) The intestinal epithelial stem cell. Bioessays 24(1):91-98

Marshman, E., Booth, C., and Potten, C.S. (2002) The intestinal epithelial stem cell. Bioessays 24(1):91-98

Masazumi Tada and J. C. Smith. (2000) Xwnt11 is a target of Xenopus Brachyury: regulation of gastrulation movements via Dishevelled, but not through the canonical Wnt pathway. Development. 127(1):2227-2238

Masoud F, Mehrotra P, and Nowak R. (2009) The Cytokine Interleukin 1beta Induces Epithelial to Mesenchymal Transition (EMT) in Peritoneal Mesothelial Cells Through Direct and Indirect Effects Involving Basigin. Biology of reproduction. 81(1):606

Masui, T., Wakefield, L. M., Lechner, J. F., LaVeck, M. A., Sporn, M. B. and Harris, C. C. (1986) Type  $\beta$  transforming growth factor is the primary differentiation-inducing serum factor for normal human bronchial epithelial cells. Proc. Nati. Acad. Sci. USA, 83(1):2438-2442

Merrill BJ. (2008) Develop-WNTs in somatic cell reprogramming.Cell Stem Cell. 3(1):465-466

Michael, W. (2001). A new mathematical model for relative quantification in realtime RT-PCR. Available: http://www.gene-quantification.de/relative.html. Accessed 08 February 2010

Microsoft office website. TTEST. Available: http://office.microsoft.com/enus/excel/HP052093251033.aspx. Accessed 10 February 2010

Midgley C.A., White S., Howitt R., Save V., Dunlop M.G., Hall P.A., Lane D.P., Wyllie A.H. & Bubb V.J. (1997) APC expression in normal human tissues. J. Pathol. 181(1):426-433

Miettinen PJ, Ebner R, Lopez AR. (1994) TGF-beta induced transdifferentiation of mammary epithelial cells to mesenchymal cells: Involvement of type I receptors. J Cell Biol.127(1):2021–2036

Miller JR, Hocking AM, Brown JD & Moon RT. (1999) Mechanism and function of signal transduction by the Wnt/ $\beta$ -catenin and Wnt/Ca2+ pathways. Oncogene 18(1):7860–7872

Mills JC, Gordon JI. (2001) The intestinal stem cell niche: There grows the neighborhood. Proc. Natl. Acad. Sci. USA 98:12334–12336

Min C, Eddy SF, Sherr DH, Sonenshein GE. (2008) NF-kappaB and epithelial to mesenchymal transition of cancer. J Cell Biochem. 104(1):7337-7344

Miraglia S, Godfrey W, Yin AH, Atkins K, Warnke R, Holden JT, Bray RA, Waller EK, Buck DW. (1997) A novel five-transmembrane hematopoietic stem cell antigen: isolation, characterization, and molecular cloning. Blood. 90(12):5013-5021

Mishina Y, Hanks MC, Miura S, Tallquist MD, Behringer RR. (2002) Generation of Bmpr/Alk3 conditional knockout mice. Genesis. 32(1):69-72

Mitsui K, Tokuzawa Y, Itoh H, Segawa K, Murakami M, Takahashi K, Maruyama M, Maeda M, Yamanaka S (2003) The homeoprotein Nanog is required for maintenance of pluripotency in mouse epiblast and ES cells. Cell. 113(5):631-642

Mitsui, K. et al. (2003) The homeoprotein Nanog is required for maintenance of pluripotency in mouse epiblast and ES cells. Cell. 113(1):631–642

Miyaki M., Konishi M., Kikuchi-Yanoshita R., Enomoto M., Tanaka K., Muraoka M., Takahashi H., Amanda Y., Fukayama M, et al. (1994) Characteristics of somatic mutation of the adenomatous polyposis coli gene in colorectal tumours. Cancer Res. 54(1):3011-3020

Miyaki M., Iijima T., Kimura J., Yasuno M., Mori T., Hayashi Y., Koike M., Shitara N., Iwama T. & Kuroki T. (1999) Frequent mutation of  $\beta$ -catenin and APC genes in primary colorectal tumours from patients with hereditary nonpolyposis colorectal cancer. Cancer Res. 59(1):4506-4509

Miyashiro I., Senda T., Matsumine A., Baeg G.-H., Kuroda T., Shimano T., Miura S., Noda T., Kobayashi S., Monden M., Toyoshima K. & Akiyama T. (1995) Subcellar localization of the APC protein: Immunoelectron microscopic study of the association of the APC protein with catenin. Oncogene 11(1):89-95

Miyoshi Y., Ando H., Nagase H., Nishisho I., Horii A., Miki Y., Mori T., Utsunomiya J., Baba S., Petersen G., Hamilton S.R., Kinzler K.W., Vogelstein B. & Nakamura Y. (1992a) Germ-line mutations of the APC gene in 53 familial adenomatous polyposis patients. Proc. Natl. Acad. Sci. USA 89(1):4452-4456

Molofsky AV, Pardal R, Morrison SJ. (2004) Diverse mechanisms regulate stem cell self-renewal. Curr Opin Cell Biol.16(1):700–707

Monk M. and Holding C., (2001) Human embryonic genes re-expressed in cancer cells, Oncogene 20(1):8085–8091

Montgomery RK, Shivdasani RA. (200(Prominin1 (CD133) as an intestinal stem cell marker: promise and nuance. Gastroenterology. 136(7):2051-2054

Moody SE, Perez D, Pan TC, Sarkisian CJ, Portocarrero CP, Sterner CJ, Notorfrancesco KL, Moon, R. T., Brown, J. D., and Torres, M. (1997) WNTs modulate cell fate and behavior during vertebrate development. Trends Genet 13(4):157-162

Moore K. B, Mood K, Daar I. O, Moody, S. A. (2004) Morphogenetic movements underlying eye field formation require interactions between the FGF and ephrinB1 signaling pathways. Dev. Cell 6(1):55-67

Moore, S. (2009) Developmental genes and cancer in children. Pediatric Blood & Cancer. 52(7):755-760

Morin PJ, Sparks AB, Korinek V, Barker N, Clevers H, Vogelstein B, Kinzler KW. (1997) Activation of beta-catenin-Tcf signaling in colon cancer by mutations in beta-catenin or APC. Science. 21(5307):1787-1790

Morin, P. J. (1999) Beta-catenin signaling and cancer. Bioessays. 21:1021-1030 Morin, P.J., Sparks, A.B., Korinek, V., Barker, N., Clevers, H., Vogelstein, B., and Kinzler, K.W. (1997) Activation of \_-catenin-Tcf signaling in colon cancer by mutations in  $\beta$ catenin or APC. Science. 275(1):1787–1790

Morris, R.J., Liu, Y., Marles, L., Yang, Z., Trempus, C., Li, S., Lin, J.S., Sawicki, J.A., and Cotsarelis, G. (2004) Capturing and profiling adult hair follicle stem cells. Nat. Biotechnol. 22(1):411–417

Morrison, S. J., Shah, N. M. & Anderson, D. J. (1997) Regulatory mechanisms in stem cell biology. Cell 88(1):287–298

Morson B (2006) The polyp-cancer sequence in the large bowel. Gastroenterology. 130(1):287

Moynagh P. N. (2005) The NFkB pathway. J Cell Sci. 118(1):4389-4392

Munemitsu S., Albert I., Souza B., Rubinfeld B. & Polakis P. (1995) Regulation of intracellular  $\beta$ -catenin levels by the adenomatous polyposis coli (APC) tumor suppressor protein. Proc. Natl. Acad. Sci. USA 92(1):3046-3050

Nagase H. & Nakamura Y. (1993) Mutations of the APC (adenomatous polyposis coli) gene. Hum. Mutat. 2(1):425-434

Naiche LA, Harrelson Z, Kelly RG, Papaioannou VE. (2005) T-box genes in vertebrate development. Annu Rev Genet. 39 (Review) 219-239

Naomi D. Willis, Thomas R. Cox, Syed F. Rahman-Casañs, Kim Smits, Stefan A. Przyborski, Piet van den Brandt, Manon van Engeland, Matty Weijenberg, Robert. Wilson, Adriaan de Bruïne, and Ch. (2008) Lamin A/C Is a Risk Biomarker in Colorectal Cancer. PLoS ONE. 3(8):2988

Nateri AS, Spencer-Dene B, Behrens A. (2005) Interaction of phosphorylated c-Jun with TCF4 regulates intestinal cancer development. Nature. 437(7056):281-285

Nelson TJ, Behfar A, Yamada S, Martinez-Fernandez A, Terzic A.(2009) Stem Cell Platforms for Regenerative Medicine. Clin Transl Sci.1:2(3):222-227 Neurath M. F., Fuss I., Schurmann G., Pettersson S., Arnold K., Muller-Lobeck H., Strober W., Herfarth C., Buschenfelllde K. H. (1998) Cytokine gene transcrition by NF-kappa B family members in patients with inflammatory bowel disease. Ann. N. Y. Acad. Sci. 859(1):149-159

Newgreen, D.F., and S.J. McKeown. (2005) Neural crest cells migration. In Rise and Fall of Epithelial Phenotype: Concepts of Epithelial-Mesenchymal Transition. P. Savagner, editor. Landes Bioscience, Texas. 1(1):29-39

Nichols, J., Zevnik, B., Anastassiadis, K., Niwa, H., Klewe-Nebenius, D., Chambers, I., et al., (1998) Formation of pluripotent stem cells in the mammalian embryo depends on the POU transcription factor Oct4. Cell. 95(1):379-391

Nicolas FJ, Lehmann K, Warne PH, Hill CS, Downward J. (2003) Epithelial-tomesenchymal transition in Madin-Darby canine kidney cells is accompanied by down-regulation of Smad3 expression, leading to resistance to transforming growthfactor- b-induced growth arrest. J Biol Chem. 278(1):3251-3256

Nick Barker, MarNick Barker, Marc van de Wetering, and Hans Clevers. (2008) The intestinal stem cell. Genes Dev. 22(1):1856-1864

NIH Stem cell website. Stem cell Basics. Available: http://stemcells.nih.gov/info/basics/basics5.asp. Accessed 04 February 2010

Nirasawa S, Kobayashi D, Tsuji N, Kuribayashi K, Watanabe N. (2009) Diagnostic relevance of overexpressed Nanog gene in early lung cancers. Oncol Rep. 22(3):587-591

Nishimura T, Utsunomiya Y, Hoshikawa M, et al. (1999) Structure and expression of a novel human FGF, FGF-19, expressed in the fetal brain. Biochim. Biophys. Acta 1444(1):148–151

Nishisho I., Nakamura Y., Miyoshi Y., Miki Y., Ando H., Koyama K., Utsunomiya J., Baba S. & Hedge P. (1991) Mutations of chromosome 5q21 genes in FAP and colorectal cancer patients. Science 253:665-559

Nishisho I., Nakamura Y., Miyoshi Y., Miki Y., Ando H., Koyama K., Utsunomiya J., Baba S. & Hedge P. (1991) Mutations of chromosome 5q21 genes in FAP and colorectal cancer patients. Science 253(1):665-559

Nishita M, Hashimoto MK, Ogata S, Laurent MN, Ueno N, Shibuya H, Cho KW. (2000) Interaction between Wnt and TGF-beta signalling pathways during formation of Spemann's organizer. Nature. 403(6771):781-785

Niwa, H., Miyazaki, J., Smith, A.G., (2000) Quantitative expression of Oct-3/4 defines differentiation, dedifferentiation or self-renewal of ES cells.Nat. Genet. 24(1):372-376

Nowell, P. C. (1986) Mechanisms of tumor progression. Cancer Res. 46(1):2203-2207

Nusse R, Varmus H. (1982) Many tumors induced by the mouse mammary tumor virus contain a provirus integrated in the same region of the host genome. Cell. 31(1):99-109

Nusse, R. & Varmus, H. E. (1997) Many tumors induced by the mouse mammary tumor virus contain a provirus integrated in the same region of the host genome. Cell 31(1):99-109

O'Brien C.A, Pollett A, Gallinger S, and Dick J.E. (2007) A human colon cancer cell capable of initiating tumour growth in immunodeficient mice. Nature 445(1):106-110

O'Day Danton. (2009). Human development. Available: http://www.utm.utoronto.ca/~w3bio380/lecture10.htm. Accessed 15 June 2009

Okita K, Nakagawa M, Hyenjong H, Ichisaka T, Yamanaka S. (2008) Generation of mouse induced pluripotent stem cells without viral vectors. Science.7:322(5903):949-953

O'Rahilly S, Wareham NJ. (2006) Genetic variants and common diseases--better late than never. N Engl J Med. 20;355(3):306-308

O'Reilly, M.-A.J., Smith, J.C., and Cunliffe, V. (1995). Patterning of the mesoderm in Xenopus: Dose-dependent and synergistic effects of Brachyury and Pintallavis. Development 121(1):1351-1359

Ogawa K, R. Nishinakamura, Y. Iwamatsu, D. Shimosato and H. Niwa. (2006) Synergistic action of Wnt and LIF in maintaining pluripotency of mouse ES cells, Biochem. Biophys. Res. Commun. 343(1):159-166

Orford K., Crockett C., Jensen J.P., Weissman A.M. & Byers S.W. (1997) Serine phosphorylation-regulated ubiquitination and degradation of  $\beta$ -catenin. J. Biol. Chem. 272(1):24735-24738

Orsulic S., Huber O., Aberle H., Arnold S. & Kemler R. (1999) E-cadherin binding prevents  $\beta$ -catenin nuclear localization and  $\beta$ -catenin/LEF-1-mediated transactivation. J. Cell Sci. 112(1):1237-1245

Oshima, H., Oshima, M., Kobayashi, M., Tsutsumi, M., and Taketo, Liu, W., Dong, X., Mai, M., Seelan, R.S., Taniguchi, K., Krishnadath, M.M. (1997) Morphological and molecular processes of polyp formation in Apc (delta716) knockout mice. Cancer Research. 57(9):1644-1649

Ozawa M, Baribault H & Kemler R (1989) The cytoplasmic domain of the cell adhesion molecule uvomorulin associates with three independent proteins structurally related in different species.EMBO Journal 8(1):1711–1717

Palena C, Polev DE, Tsang KY, Fernando RI, Litzinger M, Krukovskaya LL, Baranova AV, Kozlov AP, Schlom J. (2007) The human T-box mesodermal transcription factor Brachyury is a candidate target for T-cell-mediated cancer immunotherapy. Clin Cancer Res. 13(1):2471-2478

Palma V and Ruiz i Altaba A (2004) Hedgehog-GLI signaling regulates the behavior of cells with stem cell properties in the developing neocortex. Development. 131(1)337-345

Palma V, Lim DA, Dahmane N, Sáchez P, Brionne TC, Herzberg CD, Gitton Y, Carleton A, Álvarez-Buylla A and Altaba AR (2005) Sonic hedgehog controls stem cells behavior in the postnatal and adult brain. Development. 132(1):335-344

Pan G, Thomson JA. (2007) Nanog and transcriptional networks in embryonic stem cell pluripotency.Cell Res. 17(1):42-49

Papaioannou VE, Silver LM. (1998) The T-box gene family. Bioessays 20(1):9-19

Paul J. Gokhale, Aukje M. Giesberts and Peter W. Andrews. (2000) Brachyury Is Expressed by Human Teratocarcinoma Cells in the Absence of Mesodermal Differentiation. Cell Growth & Differentiation. 11(1):157-162

Pesce M., Scholer H.R, (2001) Oct-4: gatekeeper in the beginnings of mammalian development, Stem Cells 19(1):271-278

Pereira, Fei Yi, and Bradley J. Merrill. (2006) Repression of Nanog Gene Transcription by Tcf3 Limits Embryonic Stem Cell Self-Renewal. Molecular and Cellular Biology. 26(20):7479-7491

Perkins N. D. (2007) Integrating cell-signalling pathways with NF $\kappa$ B and IKK function. Nature. 8(1):42-62

Peters J.M., McKay R.M., McKay J.P. & Graff J.M. (1999) Caesin kinase 1 transduces Wnt signals. Nature 401(1):345-350

Petersen OW, Nielsen HL, Gudjonsson T, Villadsen R, Rank F, Niebuhr E, Bissell MJ, Ronnov-Jessen L(2003) Epithelial-tomesenchymal transition in human breast cancer can provide a nonmalignant stroma. Am J Pathol. 162(1):391-402

Pierga JY, Bonneton C, Magdelenat H, Vincent-Salomon A, Nos C, Pouillart P, Thiery JP. (2003) Clinical significance of proliferative potential of occult metastatic cells in bone marrow of patients with breast cancer. Br J Cancer. 89:539-545

Piero Dalerba, Scott J. Dylla, In-Kyung Park, Rui Liu, Xinhao Wang, Robert W. Cho, Timothy Hoey, Austin Gurney, Emina H. Huang, Diane M. Simeone, Andrew A. Shelton. (2007) Phenotypic characterization of human colorectal cancer stem cells. PNAS. 104(24):10158–10163

Piestun D, Kochupurakkal BS, Jacob-Hirsch J, Zeligson S, Koudritsky M, Domany E, Amariglio N, Rechavi G, Givol D. (2006) Nanog transforms NIH3T3 cells and targets cell-type restricted genes. Biochem Biophys Res Commun. 343(1):279-285

Piestun D, Kochupurakkal BS, Jacob-Hirsch J, Zeligson S, Koudritsky M, Domany E, Amariglio N, Rechavi G, Givol D (2006) Nanog transforms NIH3T3 cells and targets cell-type restricted genes. Biochem. Biophys. Res. Commun. 343(1):279–285

Pinto, D., Gregorieff, A., Begthel, H., and Clevers, H. (2003) Canonical Wnt signals are essential for homeostasis of the intestinal epithelium. Genes & Dev. 17(1):1709-1713

Pirvola, U.; Ylikoski, J.; Trokovic, R.; Hebert, J. M.; McConnell, S. K.; Partanen, J. (2002) FGFR1 is required for the development of the auditory sensory epithelium. Neuron 35(1):671-680

Polakis P. (1997) The adenomatous polyposis coli (APC) tumor suppressor. Biochim. Biophys. Acta. 1332:F127-147

Polakis, P. (2000) Wnt signaling and cancer. Genes Dev. 14(15):1837-1851

Polyak K, Weinberg R (2009) Transitions between epithelial and mesenchymal states: acquisition of malignant and stem cell traits. Natture reviews, Cancer 9(1):265-273

Polyak, K. (1996) Negative regulation of cell growth by TGFb. Biochimica et biophysica acta. 1242(3):185-199

Potten CS, Kovacs L, Hamilton E. (1974) Continuous labelling studies on mouse skin and intestine. Cell. Tissue Kinet. 7(1):271–283

Potten CS, and Roberts S.A, (1994) Clonogen content of intestinal crypts:

its deduction using a microcolony assay on whole mount preparations and its dependence on radiation dose. Int. J. Rad. Biol. 65(1):477-481

Potten, C.S., Booth, C., Tudor, G.L., Booth, D., Brady, G., Hurley, P., Ashton, G., Clarke, R., Sakakibara, S., and Okano, H. (2003) Identification of a putative intestinal stem cell and early lineage marker; musashi-1. Differentiation. 71(1):28-41

Potten CS, Loeffler M. (1990) Stem cells: attributes, cycles, spirals, pitfalls and uncertainties. Lessons for and from the crypt. Development. 110(4):1001-1020

Potten CS, Owen G, and Booth D. (2002) Intestinal stem cells protect their genome by selective segregation of template DNA strands. J Cell Sci 115(1):2381–2388

Potten CS. (1977) Extreme sensitivity of some intestinal crypt cells to X and gamma irradiation. Nature 269(1):518-521

Potten, C.S., Kovacs, L., and Hamilton, E. (1974) Continuous labelling studies on mouse skin and intestine. Cell Tissue Kinet. 7(1):271–283

Potter J (1992) Reconciling the epidemiology, physiology, and molecular biology of colon cancer. JAMA 268(12):1573-1537

Potter JD, McMichael AJ (1986) Diet and cancer of the colon and rectum: a casecontrol study. J Natl Cancer Inst 76(4):557-569

Powell DW, Mifflin RC, Valentich JD, Crowe SE, Saada JI, West AB. (1999) Myofibroblasts. II. Intestinal subepithelial myofibroblasts. Am. J. Physiol. 277(1):C183–201

Powers CJ, McLeskey SW, Wellstein A (2000) Fibroblast growth factors, their receptors and signaling. Endocr. Relat. Cancer 7(3):165-197

Preston-Martin S, Malcolm C. Pike, Ronald K. Ross, Peter A. Jones, and Brian E. Henderson. (1990) Increased Cell Division as a Cause of Human Cancer. Cancer Research. 50(1):7415-7421

Prince M.E, Sivanandan R, Kaczorowski A, Wolf G.T, Kaplan M.J Dalerba, P, Weissman, I.L, Clarke M.F, and Ailles L.E. (2007) Identification of a subpopulation of cells with cancer stem cell properties in head and neck squamous cell carcinoma. Proc. Natl. Acad. Sci. 104(1):973–978

Qi X, Li TG, Hao J, et al. (2004) BMP4 supports self-renewal of embryonic stem cells by inhibiting mitogen-activated protein kinase pathways. Proc Natl Acad Sci USA. 101(1):6027-6032

Qiu,J.M., Roberts,S.A. and Potten,C.S. (1994) Cell migration in the small and large bowel shows a strong circadian rhythm. Epithelial Cell Biol. 3(1):137-148

Quastler H and Sherman FG. (1959) Cell population kinetics in the intestinal epithelium of the mouse. Exp Cell Res. 17(1):420–438

Ramalho-Santos M, Melton DA, McMahon AP. (2000) Development. Hedgehog signals regulate multiple aspects of gastrointestinal development.127(12):2763-2772

Raphael Rubin and David S. Strayer (2007). Rubin's Pathology: Clinicopathologic Foundations of Medicine. Lippincott Williams & Wilkins. pp. 90. <u>ISBN</u> 0781795168.

Rayet B., Gelinas C. (1999) Aberrant rel/nfkb genes and activity in human cancer. Oncogene. 18, 6938-6947

Reinacher-Schick A. & Gumbiner B.M. (2001) Apical membrane localization of the adenomatous polyposis coli tumor suppressor protein and subcellular distribution of the  $\beta$ -catenin destruction complex in polarized epithelial cells. J. Cell Biol. 152(1):491-502

Renard, Charlotte L, Armengol C. (2007) Tbx3 Is a Downstream Target of the Wnt/ $\beta$ -Catenin Pathway and a Critical Mediator of  $\beta$ -Catenin Survival Functions in Liver Cancer . Cancer Research. 67(1):901

Reubinoff, B. E., Pera, M., Fong, C., Trounson, A. and Bongso, A. (2000) Embryonic stem cell lines from human blastocysts: somatic differentiation in vitro. Nat. Biotechnol. 18(1):399-404

Reya T, Morrison S, Clarke M & Weissman I. (2001) Stem cells, cancer, and cancer stem cells. Nature. 414(1):105-111

Reya T. (2003) A role for Wnt signalling in self-renewal of haematopoietic stem cells. Nature 423(1):409-414

Reya T. and Clevers H. (2005) Wnt signalling in stem cells and cancer. Nature 14(1):843-850

Ricci-Vitiani L, Lombardi DG, Pilozzi E, Biffoni M, Todaro M, Peschle C, De Maria R. (2007) Identification and expansion of human colon-cancer-initiating cells. Nature.;445(1):111–115

Rijsewijk F, Schuermann M, Wagenaar E, Parren P, Weigel D, Nusse R (1987) The Drosophila homolog of the mouse mammary oncogene int-1 is identical to the segment polarity gene wingless. Cell 50(4):649-657

Riley BM, Mansilla MA, Ma J, Daack-Hirsch S, Maher BS, Raffensperger LM, Russo ET, Vieira AR, Dodé C, Mohammadi M, Marazita ML, Murray JC. (2007) Developmental genes and cancer : Role of patched in basal cell carcinoma of the skinImpaired FGF signaling contributes to cleft lip and palate. Proc Natl Acad Sci U S A. 104(11):4512-4517

Rivera MN, Kim WJ, Wells J, Driscoll DR, Brannigan BW, Han M, Kim JC, Feinberg AP, Gerald WL, Vargas SO, Chin L, Iafrate AJ, Bell DW, Haber DA. (2007) An X chromosome gene, WTX, is commonly inactivated in Wilms tumor. Science. 2;315(5812):642-645

Robert-Moreno A, Espinosa L, de la Pompa JL et al. (2005) RBPjkappa dependent Notch function regulates Gata2 and is essential for the formation of intra-embryonic hematopoietic cells. Development. 132(1):1117–1126

Roberts G.T., Davies M.L. & Wakeman J.A. (2003) Interaction between Ku80 protein and a widely used antibody to adenomatous polyposis coli. Br. J. Cancer 88(1):202-205

Rolf Bjerkvig, Berit B. Tysnes, Karen S. Aboody, Joseph Najbauer & A. J. A. Terzis. (2005) The origin of the cancer stem cell: current controversies and new insights. Nature Reviews Cancer. 5(1):899-904

Romagnolo, B., Berrebi, D., Saadi-Keddoucci, S., Porteu, A., Pichard, A.L., Peuchmaur, M., Vandewalle, A., Kahn, A., and Perret, C. (1999) Intestinal dysplasia and adenoma in transgenic mice after overexpression of an activated  $\beta$ -catenin. Cancer Res. 59(1):3875-3879

Roman-Blas J. A., Jimenez S. A. (2006) NFkappaB as a potential therapeutic target in osteoarthritis and rheumatoid arthritis. Osteoarthritis Cartilage. 14(1):839-848

Roose J, Huls G, van Beest M, Moerer P, van der Horn K, Goldschmeding R, Logtenberg T, Clevers H. (1999) Synergy between tumor suppressor APC and the beta-catenin-Tcf4 target Tcf1. Science. 17;285(5435):1923-1926

Roose J. & Clevers H. (1999) TCF transcription factors: molecular switches in caecinogenesis. Biochim. Biophys. Acta. 1424:M23-37

Rosin-Arbesfeld R., Cliffe A., Brabletz T. & Bienz M. (2003) Nuclear export of the APC tumour suppressor controls  $\beta$ -catenin function in transcription. EMBO J. 22(1):1101-1113

Rosin-Arbesfeld R., Townsley F. & Bienz M. (2000) The APC tumour suppressor has a nuclear export function. Nature 46(1):1009-1012

Rosner MH, Vigano MA, Ozato K, Timmons PM, Poirier F, Rigby PW, Staudt LM. (1990) A POU-domain transcription factor in early stem cells and germ cells of the mammalian embryo. Nature. 345(1):686–692

Rossi DJ, Jamieson CH, Weissman IL. (2008) Stems cells and the pathways to aging and cancer. Cell. Review. 132(4):681-696

Rowan H, Wheeler J. (2000) APC mutations in sporadic colorectal tumors: A mutational "hotspot" and interdependence of the "two hits". PNAS. 97(7):3352-3357

Rowlands DC, Bunce CM, Crocker J, Ayres JG, Johnson GD, Ling N and Brown G (1994) Expression of a NE protein recognised by the monoclonal antibody BU31 in lung tumours: relationship to Ki67 antigen expression. J Pathol 173(1): 89-96

Rubin LL. (2008) Stem cells and drug discovery: the beginning of a new era? Cell.22; 132(4):549-552

Rubinfeld B, Souza B, Albert I, Müller O, Chamberlain SH, Masiarz FR, Munemitsu S, Polakis P. (1993) Association of the APC gene product with beta-catenin. Science.10:262(5140):1731-1734

Rubinfeld B., Souza B., Albert I., Munemitsu S. & Polakis P. (1995) The APC protein and E-cadherin form similar but independent complexes with  $\beta$ -catenin,  $\beta$ -catenin and plakoglobin. J. Biol. Chem. 270(1):5549-5555

Rubinfeld B., Albert I., Porfiri E., Fiol C., Munemitsu S. & Polakis P. (1996) Binding of GSK3 $\beta$  to the APC- $\beta$ -catenin complex and regulation of complex assembly. Science 272(1):1023-1026

Rubinfeld B., Albert I., Porfiri E., Munemitsu S. & Polakis P. (1997) Loss of bcatenin regulation by the APC tumor suppressor protein correlates with loss of structure due to somatic mutation of the gene. Cancer Res. 57(1):4624-4630

Saika S, Miyamoto T, Tanaka S, Tanaka T, Ishida I, Ohnishi Y, Ooshima A, Ishiwata T, Asano G, Chikama T et al. (2003) Response of lens epithelial cells to injury: role of lumican in epithelial– mesenchymal transition. Invest Ophthalmol Vis Sci. 44(1):2094-2102

Sakanaka C., Leong P., Xu L., Harrison S.D. & Williams L.T. (1999) Caesin kinase 1 $\beta$  in the Wnt pathway: Regulation of  $\beta$ -catenin function. Proc. Natl. Acad. Sci. USA. 96(1):12548-12552

Salic A., Lee E., Mayer L. & Kirschner M.W. (2000) Control of  $\beta$ -catenin stability: reconstitution of the cytoplasmic steps of the wnt pathway in Xenopus egg extracts.Mol. Cell. 5(1):523-532

Sangiorgi and M. Capecchi. (2008) Bmi1 is expressed in vivo in intestinal stem cells. Nat Genet. 40(1):915–920

Sansom, O.J., Reed, K.R., Hayes, A.J., Ireland, H., Brinkmann, H., Newton, I.P., Batlle, E., Simon-Assmann, P., Clevers, H.,Nathke, I.S., et al. (2004) Loss of Apc in vivo immediately perturbs Wnt signaling, differentiation, and migration.Genes & Dev. 18(1):1385–1390

Sato N, Meijer L, Skaltsounis L, Greengard P. and Brivanlou A. (2004) Maintenance of pluripotency in human and mouse embryonic stem cells through activation of Wnt signaling by a pharmacological GSK-3- specific inhibitor. Nat. Med. 10(1):55-63

Sato, T. et al. (2009) Single Lgr5 stem cells build crypt-villus structures in vitro without a mesenchymal niche. Nature. 459(1):262-265

Satoh S, Daigo Y, Furukawa Y, Kato T, Miwa N, Nishiwaki T, Kawasoe T, Ishiguro H, Fujita M, Tokino T, Sasaki Y, Imaoka S, Murata M, Shimano T, Yamaoka Y, Nakamura Y. (2000) AXIN1 mutations in hepatocellular carcinomas, and growth suppression in cancer cells by virus-mediated transfer of AXIN1.Nat Genet. 24(3):245-250

Scadden D. (2006) The stem-cell niche as an entity of action, Nature. 441(7097): 1075-1079

Scholzen T, Gerdes J (2000) The Ki-67 protein: from the known and the unknown. J. Cell. Physiol. 182(3):311-322

Schonhoff SE, Giel-Moloney M, Leiter AB. Minireview: (2004) Development and differentiation of gut endocrine cells. Endocrinology. 145(1):2639-2644

Schreiber S., Nikolaus S., Hampe J. (1998) Activation of nuclear factor kappa B inflammatory bowel disease. Gut. 42(1):477-484

Schroder N, Gossler A. (2002) Expression of Notch pathway components in fetal and adult mouse small intestine. Gene Expr Patterns. 2(1):247–250

Schulte-Merker S, Ho RK, Herrmann BG, Nsslein-Volhard, C. (1992) The protein product of the zebrafish homologue of the mouse T gene is expressed in nuclei of the germ ring and the notochord of the early embryo. Development. 116(1):1021-1032

Sebastian J. Arnold, Jörg Stappert, Andreas Bauer, Andreas Kispert, Bernhard G. Herrmann and Rolf Kemler. (2000) Brachyury is a target gene of the Wnt/ $\beta$ -catenin signaling pathway. Mechanisms of Development. 91(1-2): 249-258

Semb H, Christofori G (1998) The tumor-suppressor function of E-cadherin. Am. J. Hum. Genet. 63(6):1588–1593

Shackleton M, Vaillant F, Simpson K.J, Stingl J, Smyth G.K, Asselin-Labat M.L, Wu L, Lindeman G.J, and Visvader J.E. (2006) Generation of a functional mammary gland from a single stem cell. Nature 439(1):84-88

Shaffer AL, Peng A, Schlissel MS. (1997) In vivo occupancy of the kappa light chain enhancers in primary pro- and pre-B cells: a model for kappa locus activation. Immunity. 6(1):131–143.

Sharma R P, Chopra V L (1976). "Effect of the Wingless (wg1) Mutation on Wing and Haltere Development in Drosophila melanogaster. Developmental Biology 48(1): 461-465

Shuey DJ, McCallus DE, Giordano T. 2002. RNAi: gene-silencing in theraputic intervention. Drug Discovery Today 7: 1040-1046. <<u>http://reviews.bmn.com/rsearch/section/record?uid=DDT.etd01541\_13596446\_v00</u> 07i20\_00002474&node=TOC%40%40DDT%40008%4005%40008\_05>. Accessed Feb 7 2010

Shimizu Y, Yamamichi N, Saitoh K, Watanabe A, Ito T, Yamamichi-Nishina M, Mizutani M, Yahagi N, Suzuki T, Sasakawa C et al. (2003) Kinetics of v-srcinduced epithelial–mesenchymal transition in developing glandular stomach. Oncogene. 22(1):884-893

Shinya H, Wolff W (1979) Morphology, anatomic distribution and cancer potential of colonic polyps. Ann Surg 190(6):679-683

Shizuru JA, Negrin RS, Weissman IL. (2005) Hematopoietic stem and progenitor cells: clinical and preclinical regeneration of the hematolymphoid system. Annu Rev Med. Reveiew. 56(1):509-538

Shmelkov SV, Butler JM, Hooper AT, Hormigo A, Kushner J, Milde T, St Clair R, Baljevic M, White I, Jin DK, Chadburn A, Murphy AJ, Valenzuela DM, Gale NW, Thurston G, Yancopoulos GD, D'Angelica M, Kemeny N, Lyden D, Rafii S. (2008) CD133 expression is not restricted to stem cells, and both CD133<sup>+</sup> and CD133<sup>-</sup> metastatic colon cancer cells initiate tumors. J Clin Invest. 118(1):2111-2120

Simeone A, D'Apice MR, Nigro V, Casanova J, Graziani F, Acampora D, Avantaggiato V. (1994) Orthopedia, a novel homeobox-containing gene expressed in the developing CNS of both mouse and Drosophila. Neuron.13(1):83–101

Simon P. Bach, Andrew G. Renehan and Christopher S. Potten. (2000) Stem cells: the intestinal stem cell as a paradigm . Carcinogenesis. 21(3):469-476

Sinclair C, Adem C, Naderi A, Soderberg C, Johnson, Wadum L, Couch V, Sellers T, Schaid D, Slezak J, Fredericksen Z, Ingle . (2002). TBX2 Is Preferentially Amplified in BRCA1- and BRCA2-related Breast Tumors. Cancer Research. 62(1):3587-3591

Singer JB, Harbecke R, Kusch T, Reuter R, Lengyel JA. (1996) Drosophila brachyenteron regulates gene activity and morphogenesis in the gut. Development.122(1):3707–3718

Singh SK, Clarke ID, Terasaki M, Bonn VE, Hawkins C, Squire J, Dirks PB (2003) Identification of a cancer stem cell in human brain tumors. Cancer research 63(18):5821–5828

Singh, S.K., Hawkins, C., Clarke, I.D., Squire, J.A., Bayani, J.Hide, T., Henkelman, R.M., Cusimano, M.D., and Dirks, P.B. (2004) Identification of human brain tumour initiating cells. Nature. 432(1):396–401

Singla D, D.J. Schneider, M.M. LeWinter and B.E. Sobel. (2006) Wnt3a but not wnt11 supports self-renewal of embryonic stem cells, Biochem. Biophys. Res. Commun. 345(1):789–795

Sirard C, de la Pompa JL, Elia A, et al. (1998) The tumor suppressor gene Smad4/Dpc4 is required for gastrulation and later for anterior development of the mouse embryo. 12(1):107-119

Skibber J, Minsky B and Hoff, (2001) Cancer of the colon. In: V. DeVita, S. Hellman and S. Rosenberg, Editors, Cancer: Principles and Practice of Oncology, Lippincott Williams and Wilkins, Philadelphia 1(1):1216–1271

Slattery ML, Schumacher MC, Smith KR. (1988) Physical activity, diet, and risk of colon cancer in Utah. Am J Epidemiol 128(5):989-999

Smith, A. G. (2001) Embryo-derived stem cells: of mice and men. Annu. Rev. Cell Dev. Biol. 17(1):435-462

Smith, A.G., Heath, J.K., Donaldson, D.D., Wong, G.G., Moreau, J., Stahl, M., Rogers, D., (1988) Inhibition of pluripotential embryonic stem cell differentiation by purified polypeptides. Nature. 336(1):688–690

Smith, J. C., Price, B. M. J., Green, J. B. A., Weigel, D. and Herrmann, B. G. (1991) Expression of a Xenopus homolog of Brachyury (T) is an immediate-early response to mesoderm induction. Cell 67(1):79-87

Smits, R., Kielman, M.F., Breukel, C., Zurcher, C., Neufeld, K., Jagmohan-Changur, S., Hofland, N., van Dijk, J., White, R., Edelmann, W., et al. (1999) Apc1638T: A mouse model delineatingcritical domains of the adenomatous polyposis coli protein involved in tumorigenesis and development. Genes & Dev. 13(1):1309-1321

Soldner F., Hockemeyer D., Beard C., Gao Q., Jaenisch R. (2009) Parkinson's disease patient-derived induced pluripotent stem cells free of viral reprogramming factors. Cell: 136(5):964-977

Solomon E., Voss R., Hall V., Bodmer W.F., Jass J.R., Jeffreys A.J., Lucibello F.C., Patel I. & Rider S.H. (1987) Chromosome 5 allele loss in human colorectal carcinomas. Nature 328(1):616-619

Soubeyran P, Mallo GV, Moucadel V, Dagorn JC, Iovanna JL (2000) Overexpression of Cdx1 and Cdx2 homeogenes enhances expression of the HLA-I in HT-29 cells. Mol Cell Biol Res Commun. 3(5):271-276

Spaderna S, Schmalhofer O, Hlubek F, Berx G, Eger A, Merkel S et al. (2006). A transient, EMT-linked loss of basement membranes indicates metastasis and poor survival in colorectal cancer. Gastroenterology 131(1):830–840

Spaderna S, Schmalhofer O, Wahlbuhl M, Dimmler A, Bauer K, Sultan A, Hlubek F et al. (2008) The transcriptional repressor ZEB1 promotes metastasis and loss of cell polarity in cancer. Cancer Res. 68(1):537-544

Spaner DE, Hammond C, Mena J, Shi Y. (2004) Effect of IL-2R beta-binding cytokines on costimulatory properties of chronic lymphocytic leukaemia cells: implications for immunotherapy. Br J Haematol. 127(5):531-542

Spangrude, G.J., Heimfeld, S., and Weissman, I.L. (1988) Purification and characterization of mouse hematopoietic stem cells. Science. 241(1):58–62.

Spiegelman, M. (1976) Electron microscopy of cell associations in T-locus mutants. Embryogenesis in mammals. Ciba Found. Syrup. 40(1):199-220

Staal F.J.T., van Noort M., Strous G.J. & Clevers H.C. (2002) Wnt signals are transmitted through N-terminally dephosphorylated  $\beta$ -catenin. EMBO Rep. 3(1):63-68

Stappenbeck, T.S., Mills, J.C., and Gordon, J.I. (2003) Molecular features of adult mouse small intestinal epithelial progenitors.Proc. Natl. Acad. Sci. 100(1):1004-1009

Stoker M, Perryman M. (1985) An epithelial scatter factor released by embryo fibroblasts. J Cell Science. 327(1):239-242

Su L.K., Burrell M., Hill D.E., Gyuris J., Brent R., Wiltshire R., Trent J., Vogelstein B. & Kinzler KW. (1995) APC binds to the novel protein EB1. Cancer Res. 55(1):2972-2977

Su L.K., Vogelstein B. & Kinzler K.W. (1993) Association between wild-type and mutant APC gene products. Cancer Res. 53(1):2728-2731

Sulyok S, Wankell M, Alzheimer C, Werner S (2004) Activin: an important regulator of wound repair, fibrosis, and neuroprotection. Mol. Cell. Endocrinology. 225(1-2):127-132

Suvà ML, Riggi N, Stehle JC, Baumer K, Tercier S, Joseph JM, Suvà D, Clément V, Provero P, Cironi L, Osterheld MC, Guillou L, Stamenkovic I. (2009) Identification of cancer stem cells in Ewing's sarcoma. Cancer Res. 69(1):1776-1781

Suzuki A, Raya A, Kawakami Y, et al. (2006) Nanog binds to Smad1 and blocks bone morphogenetic protein-induced differentiation of embryonic stem cells. Proc Natl Acad Sci USA 103(1):10294–10299

Suzuki A, Sayaka Sekiya, Dirk Büscher3, Juan Carlos Izpisúa Belmonte, Hideki Taniguchi. (2008) Tbx3 controls the fate of hepatic progenitor cells in liver development by suppressing p19ARF expression. Development. 135(1):1589-1595

Tada, M., Casey, E., Fairclough, L. and Smith, J. C. (1998) Bix1, a direct target of Xenopus T-box genes, causes formation of ventral mesoderm and endoderm. Development. 125:3997-4006

Taipale, J. & Beachy, P. A. (2001) The Hedgehog and Wnt signaling pathways in cancer. Nature 411(1):349–354

Tai M.H., Chang C.C, Kiupel M. (2005) Oct4 expression in adult human stem cells: evidence in support of the stem cell theory of carcinogenesis, Carcinogenesis 26(1):495–502

Takahashi K, Tanabe K, Ohnuki M, Narita M, Ichisaka T, Tomoda K, Yamanaka S. (2007) Induction of pluripotent stem cells from adult human fibroblasts by defined factors. Cell. 131(5):861-872

Takahashi, K. & Yamanaka, S. (2006) Induction of pluripotent stem cells from mouse embryonic and adult fibroblast cultures by defined factors. Cell.126 (4):663-676

Tamara C, Senka Rados, Simun Krizanac and Sanja Kapitanovic. (2008) Influence of interleukin-8 and interleukin-10 on sporadic colon cancer development and progression. Carcinogenesis. 29(8):1572-1580

Tanigaki K, Nogaki F, Takahashi J, Tashiro K, Kurooka H, Honjo T (2001) Notch1 and Notch3 instructively restrict bFGF-responsive multipotent neural progenitor cells to an astroglial fate. Neuron 29(1):45-55

Tejpar S., Nollet F., Li C., Wunder J.S., Michils G., dal Cin P., Van Custem E., Bapat B., van Roy F., Cassiman J.J. & Alman B.A. (1999) Predominance of  $\beta$ catenin mutations and  $\Box$ -catenin dysregulation in sporadic aggressive fibromatosis (desmoid tumour). Oncogene. 18(1):6615-6620

Terry P. Yamaguchi, Shinji Takada, Yoshiaki Yoshikawa, Nongying Wu, and Andrew P. McMahon. (1999) T (Brachyury) is a direct target of Wnt3a during paraxial mesoderm specification . Genes & Dev.13(1):3185-3190

Thiery JP, Sleeman JP. (2006) Complex networks orchestrate epithelialmesenchymal transitions. Nat Rev Mol Cell Biol. 7(1):131-142

Thiery JP. (2002) Epithelial-mesenchymal transitions in tumor progression. Nat Rev Cancer. 2(1):442-454

Thomson, J. A., Itskovitz-Eldor, J., Shapiro, S., Waknitz, M., Swiergiel, J., Marshall, V. and Jones, J. (1998) Embryonic stem cell lines derived from human blastocysts. Science 282(1):1145-1147

Todaro M, Perez Alea M, Di Stefano A, Medema JP. (2007) Colon Cancer Stem Cells Dictate Tumor Growth and Resist Cell Death by Production of Interleukin-4. Cell Stem Cell. 1(4):389-402

Tomita K, van Bokhoven A, van Leenders GJ, et al. (2000) Cadherin switching in human prostate cancer progression. Cancer Res 60(1):3650-3654

Tomlinson I, Ilyas M and Novelli M. (1997) Molecular genetics of colon cancer. Cancer Metastasis Rev 16(1):67-79

Toncheva D, Nacheva M. (1998) Chromosomal instability in cancer patients. Hum Genet. 103(1):367-369

Torres J, Watt FM. (2008) Nanog maintains pluripotency of mouse embryonic stem cells by inhibiting NFkappaB and cooperating with Stat3. Nat Cell Biol.10(2):194-201

Trachette L. Jackson. The Jackson cacer modelling group. Available: www.math.lsa.umich.edu/%7Etjacks/research.html. Accessed 31 March, 2009

Tucker E.L. & Pignatelli M. (2000) Catenins and their associated proteins in colorectal cancer. Histol. Histopathol. 15(1):251-260

Twigg SR, Wilkie AO. (1999) Characterisation of the human snail (SNAI1) gene and exclusion as a major disease gene in craniosynostosis. Hum. Genet.105(4):320-326

Ullu E, Tschudi C. 2000. RNAi- applications in parasitology. New technologies for life sciences: A Trends Guide 2000: 43-46. <<u>http://reviews.bmn.com/rsearch/section/record?uid=SUPP.etd00615\_14711931\_v0</u>001i01\_00000073&rendertype=text>. Accessed Feb 7 2010

Usami Y, Satake S, Nakayama F, Matsumoto M, Ohnuma K, Komori T, Semba S, Ito A, Yokozaki H (2008) Snail-associated epithelial-mesenchymal transition promotes oesophageal squamous cell carcinoma motility and progression. J Pathol. 215(3):330-339

Uyttendaele H, J. Ho, J. Rossant, and J. Kitajewski. (2001) Vascular patterning defects associated with expression of activated Notch4 in embryonic endothelium. Proc Natl Acad Sci U S A. 98(10):5643-5648

Vainio S, Karavanova I, Jowett A, Thesleff I.Cell. (1993) Identification of BMP-4 as a signal mediating secondary induction between epithelial and mesenchymal tissues during early tooth development. 8;75(1):45-58

Valizadeh A, Karayiannakis AJ, El-Hariry I, Kmiot W and Pignatelli M (1997) Expression of E-cadherin-associated molecules (a, b, and g-catenin and p120) in colorectal polyps. American Journal of Pathology. 150(1):1977-1984

Van de Wetering M, Sancho E, Verweij C, de Lau W, Oving I, Hurlstone A, van der Horn K, Batlle E, Coudreuse D, Haramis AP, Tjon-Pon-Fong M, Moerer P, van den Born M, Soete G, Pals S, Eilers M, Medema R, Clevers H. (2002) The betacatenin/TCF-4 complex imposes a crypt progenitor phenotype on colorectal cancer cells. Cell. 111(1):241-250

Van der Flier, L. G. et al. (2009) Transcription factor Achaete scute-like 2 controls intestinal stem cell fate. Cell. 136(1):903–912

Van der Flier, L.G., Sabates-Bellver, J., Oving, I., Haegebarth, A., De Palo, M., Anti, M., Van Gijn, M.E., Suijkerbuijk, S., Vande Wetering, M., Marra, G., et al. (2007) The Intestinal Wnt/TCF Signature. Gastroenterology. 132(1):628–632

Van Es JH, Jay P, Gregorieff A, van Gijn ME, Jonkheer S, Hatzis P, Thiele A, van den Born M, Begthel H, Brabletz T, Taketo MM, Clevers H. (2005) Wnt signalling induces maturation of Paneth cells in intestinal crypts. Nat Cell Biol. 7(4):381-386

Vance KW, Carreira S, Brosch G, Goding CR. (2005) Tbx2 is overexpressed and plays an important role in maintaining proliferation and suppression of senescence in melanomas. Cancer Res. 65(6):2260-2268

Vandewalle C, Joke Comijn, Bram De Craene, Petra Vermassen, Erik Bruyneel, Henriette Andersen, Eugene Tulchinsky, Frans Van Roy and Geert Berx. (2005). SIP1/ZEB2 induces EMT by repressing genes of different epithelial cell–cell junctions. Nucleic Acids Res. 33(20):6566–6578

Varnum-Finney, B. (2000) Pluripotent, cytokine-dependent, hematopoietic stem cells are immortalized by constitutive Notch1 signaling. Nature Med. 6(1):1278–1281

Vasen HF, Offerhaus GJ, den Hartog Jager FC, Menko FH, Nagengast FM, Griffioen G, van Hogezand RB, Heintz AP (1990) The tumour spectrum in hereditary non-polyposis colorectal cancer: a study of 24 kindreds in the Netherlands. Int J Cancer. 46(1):31-34

Vujovic, S. and Henderson, S. and Presneau, N. and Odell, E. and Jacques, T.S. and Tirabosco, R. and Boshoff, C. and Flanagan, A.M. (2006) Brachyury, a crucial regulator of notochordal development, is a novel biomarker for chordomas. The Journal of Pathology. 209(2):157-165

Walshe J, Mason I. (2000) Expression of FGFR1, FGFR2 and FGFR3 during early neural development in the chick embryo. Mech Dev. 90(1):103-310

Wang P., Branch D.R, Bali M, Schultz G.A. (2003) The POU homeodomain protein OCT3 as a potential transcriptional activator for fibroblast growth factor-4 (FGF-4) in human breast cancer cells, Biochem. J. 375(1):199–205

Wang HL, Wang J, Xiao SY. (2002) Elevated protein expression of cyclin D1 and Fra-1 but decreased expression of c-Myc in human colorectal adenocarcinomas overexpressing beta-catenin. Int J Cancer. 101(1):301-310

Wang B, Zhang Z, Ke Y. (2006) Conversion of cadherin isoforms in cultured human gastric carcinoma cells. World J Gastroenterol 12(1):966–970

Watt, F and Torres, J. (2008) Nanog maintains pluripotency of mouse embryonic stem cells by inhibiting NF $\kappa$ B and cooperating with Stat3. Nature Cell Biology. 10(1):194-201

Wei Y., Fabre M., Branchereau S., Gauthier F., Perilongo G. & Buenida M.A. (2000) Activation of  $\beta$ -catenin in epithelial and mesenchymal hepatoblastomas. Oncogene. 19:498-504

Wei J, Xu G, Wu M, Zhang Y, Li Q, Liu P, Zhu T, Song A, Zhao L, Han Z, Chen G, Wang S, Meng L, Zhou J, Lu Y, Wang S, Ma D. (2008) Overexpression of vimentin contributes to prostate cancer invasion and metastasis via src regulation. Anticancer Res. 28(1A):327-334

Weichert W, T Knösel, J Bellach, M Dietel and G Kristiansen. (2004) ALCAM/CD166 is overexpressed in colorectal carcinoma and correlates with shortened patient survival. Journal of Clinical Pathology. 57(1):1160-1164

Weigmann A, Corbeil D, Hellwig A, Huttner WB. (1997) Prominin, a novel microvilli-specific polytopic membrane protein of the apical surface of epithelial cells, is targeted to plasmalemmal protrusions of non-epithelial cells. Proc Natl Acad Sci U S A. 94(23):12425-12430

Weinberg R (2007) The Biology of Cancer. Garland Science. ISBN 0-8153-4076-1; ISBN 0-8153-4078-8

White E, Jacobs EJ, Daling JR (1996) Physical activity in relation to colon cancer in middle-aged men and women. Am J Epidemiol 144(1):42-50

Whiteman EL, Liu CJ, Fearon ER, Margolis B. (2008) The transcription factor snail represses Crumbs3 expression and disrupts apico-basal polarity complexes. Oncogene 27(1):3875-3879

Wicha MS, Liu S, Dontu G. (2006) Cancer stem cells: an old idea-a paradigm shift. Cancer Res. 66(1):1883-1890

Willert J, Epping M, Pollack J, Brown P, Nusse R. (2002) A transcriptional response to Wnt protein in human embryonic carcinoma cells. BMC developmental biology 2(1):8

Willert K & Nusse R (1998) β-Catenin: a key mediator of Wnt signalling. Current Opinion in Genetics and Development 8(1):95-102

William H. Fishman. (1976) Activation of Developmental Genes in Neoplastic Transformation. Cancer Research. 36(1):3423-3428

Williams, R.L., Hilton, D.J., Pease, S., Willson, T.A., Stewart, C.L., Gearing, D.P., et al., (1988) Myeloid leukaemia inhibitory factor maintains the developmental potential of embryonic stem cells. Nature. 336(1):684-687

Willis ND, Cox TR, Rahman-Casañs SF, Smits K, Przyborski SA, van den Brandt P, van Engeland M, Weijenberg M, Wilson RG, de Bruïne A, Hutchison CJ. (2008) Lamin A/C is a risk biomarker in colorectal cancer. PLoS ONE. 3(8):e2988

Willison K. (1990) The mouse Brachyury gene and mesoderm formation. Trends Genet. 6(1):104-105

Wilson, Val and Conlon, F. (2002) The T-box family. Genome Biol. 3(6), review Winston J.T., Strack P., Beer-Romero P., Chu C.Y., Elledge S.J. & Harper J.W. (1999) The SCF $\beta$ -TrCP-ubiquitin ligase complex associates specifically with phosphorylated destruction motifs in I $\beta$ B $\beta$  ubiquitinatin in vitro. Genes Dev. 13(1):270-283

Winton, D.J. and Ponder, B.A. (1990) Stem-cell organization in mouse small intestine. Proc. Biol. Sci. 241(1):13–18

Wong AS, Gumbiner BM (2003) Adhesion-independent mechanism for suppression of tumor cell invasion by E-cadherin. J. Cell Biol. 161(6):1191–1203

Woodford-Richens, K.L., Rowan, A.J., Gorman, P., Halford, S., Bicknell, D.C., Wasan, H.S., Roylance, R.R., Bodmer, W.F., Tomlinson. (2001) SMAD4 mutations in colorectal cancer probably occur before chromosomal instability, but after divergence of the microsatellite instability pathway. I.P. Proc. Natl. Acad. Sci. U.S.A. 98(17):9719-9723

Wu DC, Boyd AS, Wood KJ. (2008) Embryonic stem cells and their differentiated derivatives have a fragile immune privilege but still represent novel targets of immune attack. Stem Cells.26(8):1939-1950

X.E. He et al., (2007) PTEN-deficient intestinal stem cells initiate intestinal polyposis. Nat Genet, 39:189–198

Xu K, Zhu Z, Zeng F. (2007) Expression and significance of Oct4 in bladder cancer. J Huazhong Univ Sci Technolog Med Sci.,27(6):675-677

Xue C, Plieth D, Venkov C, Xu C, Neilson EG. (2003) The gatekeeper effect of epithelial–mesenchymal transition regulates the frequency of breast cancer metastasis. Cancer Res. 63(1):3386-3394

Yamaguchi TP, Takada S, Yoshikawa Y, Wu N, McMahon AP. (1999) T (Brachyury) is a direct target of Wnt3a during paraxial mesoderm specification. Genes Dev.

## 13:3185-3190

Yamamoto Y., Gaynor R. B. (2001) Therapeutic potential of inhibition of the NFkappaB pathway in the treatment of inflammation and cancer. J Clin Invest.107(1):135-142

Yamanaka S, et al. (2007) Generation of germline-competent induced pluripotent stem cells. Nature. 448(1):313-317

Yan H., Papadopoulos N., Marra G., Perrarra C., Jiricny J., Boland C.R., de la Chapelle A., Berg K., Eshleman J., Yuan W., Markowitz S., Laken S.J., Lengauer C., Kinzler K.W. & Vogelstein B. (2000) Conversion of diploidy to haploidy. Nature 403(1):723-724

Yanez-Mo M, Lara-Pezzi E, Selgas R, Ramirez-Huesca M, Dominguez-Jimenez C, Jimenez-Heffernan JA, Aguilera A, Sanchez-Tomero JA, Bajo MA, Alvarez V et al. (2003) Peritoneal dialysis and epithelial-to-mesenchymal transition of mesothelial cells. N Engl J Med. 348(1):403-413

Yang J, Wen Z, Paul M, Xi C, Xi He and Chunming L (2006) APC differentially regulates  $\beta$ -catenin phosphorylation and ubiquitination in colon cancer cells. JBC 281(26):17751-17757

Yang, J., S.A. Mani, J.L. Donaher, S. Ramaswamy, R.A. Itzykson, C. Come, . Savagner, I. Gitelman, A. Richardson, and R.A. Weinberg. (2004) Twist, a master regulator of morphogenesis, plays an essential role in tumor metastasis. Cell. 117(1):927–939

Ye F, Zhou C, Cheng Q, Shen J, Chen H. (2008) Stem-cell-abundant proteins Nanog, Nucleostemin and Musashi1 are highly expressed in malignant cervical epithelial cells. BMC Cancer. 8(1):108

Ying, Q. L., Nichols, J., Chambers, I. & Smith, A. (2003) BMP induction of Id proteins suppresses differentiation and sustains embryonic stem cell self-renewal in collaboration with STAT3. Cell. 115(1):281–292

Yu J, Carroll TJ, McMahon AP. (2002) Sonic hedgehog regulates proliferation and differentiation of mesenchymal cells in the mouse metanephric kidney. Development. 129(22):5301-5312

Yu, Zhiyuan; Zhang Wenzheng, Kone Bruce C (2002). Signal transducers and activators of transcription 3 (STAT3) inhibits transcription of the inducible nitric oxide synthase gene by interacting with nuclear factor kappaB. Biochem. J. 367(1): 97-105

Zhang J, Wang X, Chen B, Suo G, Zhao Y, Duan Z, Dai J. (2005) Expression of Nanog gene promotes NIH3T3 cell proliferation. Biochem Biophys Res Commun. 338(2):1098-1102

Zhang J, Wang X, Li M, Chen B, Wang B, Dai J. (2006) NANOGP8 is a retrogene expressed in cancers. FEBS journal 273(1):1723-1730

Zhang S, Balch C, Chan MW, Lai HC, Matei D, Schilder JM, Yan PS, Huang TH, Nephew KP. (2008) Identification and characterization of ovarian cancer-initiating cells from primary human tumors. Cancer research 68(11):4311–4320

Zhu H, Mazor M, Kawano Y, Walker MM, Leung HY, Armstrong K, Waxman J, Kypta RM. (2004) Analysis of Wnt gene expression in prostate cancer: mutual inhibition by WNT11 and the androgen receptor. Cancer Res. 64(21):7918-7926

Zhu L, Gibson P, Currle DS, Tong Y, Richardson RJ, Bayazitov IT, Poppleton H, Zakharenko S, Ellison DW, Gilbertson RJ. (2009) Prominin 1 marks intestinal stem cells that are susceptible to neoplastic transformation. Nature 457(7229):603-607