

New locality records and notes on the scalation of the Alpine Punjab skink, *Eurylepis taeniolata* Blyth 1854, from Himachal Pradesh, India

Owens, John Benjamin; Kuttalam, Sourish Rajagopalan; Santra, Vishal; Graham, Stuart; Wüster, Wolfgang; Mukherjee, Nilanjan ; Bharti, Omesh; Malhotra, Anita

Reptiles & Amphibians

DOI:

[10.17161/randa.v31i1.21541](https://doi.org/10.17161/randa.v31i1.21541)

Published: 24/12/2024

Publisher's PDF, also known as Version of record

[Cyswllt i'r cyhoeddiad / Link to publication](#)

Dyfyniad o'r fersiwn a gyhoeddwyd / Citation for published version (APA):

Owens, J. B., Kuttalam, S. R., Santra, V., Graham, S., Wüster, W., Mukherjee, N., Bharti, O., & Malhotra, A. (2024). New locality records and notes on the scalation of the Alpine Punjab skink, *Eurylepis taeniolata* Blyth 1854, from Himachal Pradesh, India. *Reptiles & Amphibians*, 31(1), 1-2. Article e21541. <https://doi.org/10.17161/randa.v31i1.21541>

Hawliau Cyffredinol / General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



New Locality Records and Notes on the Scalation of the Alpine Punjab Skink, *Eurylepis taeniolata* Blyth 1854, from Himachal Pradesh, India

John Benjamin Owens^{1,2}, Sourish Kuttalam^{1,2,3}, Vishal Santra^{2,3}, Stuart Graham¹, Wolfgang Wüster¹, Nilanjan Mukherjee³, Omesh Bharti⁴, and Anita Malhotra¹

¹Molecular Ecology and Evolution at Bangor (MEEB), School of Natural Sciences, Bangor University, Environment Centre Wales, Bangor LL57 2UW, Wales, UK (jhw19vln@bangor.ac.uk [corresponding author])

²Captive & Field Herpetology Ltd., 13 Hirfron, Llaingoch, Holyhead, Anglesey LL65 1YU, Wales, UK (captiveandfieldherpetology@gmail.com)

³Society for Nature Conservation, Research and Community Engagement (CONCERN), Nalikul, Hooghly, West Bengal 712407, India

⁴State Institute of Health and Family Welfare, Shimla 171009, Himachal Pradesh, India

The Alpine Punjab Skink, *Eurylepis taeniolata* Blyth 1854, ranges from Jordan to northwestern India (Feizi et al. 2016). Three subspecies are currently recognized, *E. t. arabica* (Szczerbak 1990), *E. t. parthianica* (Szczerbak 1990), and *E. t. taeniolatus* Blyth 1854 (Feizi et al. 2016). *Eurylepis t. taeniolatus* occurs in Pakistan and northwestern India (type locality: Punjab, Pakistan) (Feizi et al. 2016). This species has been recorded from only a few locations in northwestern India and accurate herpetofaunal records from Himachal Pradesh are lacking (Griffith et al. 2000; Saikia et al. 2007, Datta-Roy et al. 2017). Herein we report two additional records of *E. t. taeniolatus* from Himachal Pradesh, India.

The first individual was under a large boulder in a patch of alpine forest near Banjar, Kullu District, Himachal Pradesh, India (31.645833, 77.379722; elev. 1,431 m asl), at 0951 h on 5 July 2018. This individual was captured (permit number FFE-FB-F (10)-3/2017), photographed (Fig. 1) to collect morphometric data and confirm the identity of the species, and released at the site of capture. A photographic voucher was deposited in the University of Kansas Digital Archive (KUDA 14507), with the identity of the species independently confirmed by James Hicks.

A second individual was observed while active on a boulder along a river bank at 1047 h on 21 July 2018 at the same location as the first individual and was not captured or photographed. Habitat consisted of alpine coniferous forest with large boulders and bordered a large river. The nearest previously documented locality is in Sirmaur District approximately 110 km SSE of our record (Fig. 2); however, an exact location was not provided for this record (Saikia et al. 2007).

The identity of the species was determined by comparing our photographs with those of *E. t. parthianica* from Iran (Feizi et al. 2016) and using the following characteristics (Datta-Roy et al. 2017): midbody scale rows 23, supralabial



Figure 1. An adult Alpine Punjab Skink (*Eurylepis t. taeniolatus*) in alpine forest in the Kullu District, Himachal Pradesh, India, where it was found under a large boulder. Photograph by John Benjamin Owens.




Figure 2. Map of Himachal Pradesh, India, showing the location where Alpine Punjab Skinks (*Eurylepis taeniolatus*) were observed in Kullu District (green dot) and the approximate location of the nearest previously recorded observation in Sirmour District (shaded blue).

scales 7–8, infralabial scales 7, loreal scales 2, enlarged vertebral dorsal scales 77 (Fig. 3), lamellae under third toe 9, lamellae under fourth toe 8, and third toe shorter than fourth. Our individual differed from descriptions in Datta-Roy et al. (2017) only in having 23 instead of 21 midbody scale rows.



Figure 3. An Alpine Punjab Skink (*Eurylepis t. taeniolatus*) from the Kullu District, Himachal Pradesh, India, showing diagnostic head scales (left) and enlarged vertebral dorsal scales (right). Photographs by John Benjamin Owens.

We returned to the site in July 2019 but failed to locate any additional individuals. The habitat had changed considerably since the previous year, with an increase in dense vegetation cover and signs of anthropogenic activities. Our observations demonstrate the need for future surveys to determine the distribution of this species in Himachal Pradesh.

Acknowledgements

We thank the Himachal Pradesh Forest Department for providing the permit (number FFE-FB-F (10)-3/2017) that allowed us to capture the individual. We also thank Dr. Anatoli Togruidou for assistance with fieldwork, Maya Master for help with the map, and Dr. James Hicks for confirming the identity of this lizard.

Literature Cited

- Datta-Roy, A., V. Deepak, R. Chaitanya, C. Murthy, H. Bhosale, A. Lajmi, P. Karanth, K. Kunte, and V. Giri. 2017. On the taxonomic status of *Eurylepis poonaensis* (Squamata: Scincidae): resolving a long-standing conundrum. *Salamandra* 53: 389–397.
- Feizi, H., N. Rastegar-Pouyani, E. Rastegar Pouyani, and N. Heidari. 2016. Systematics of the genera *Eumeces* Wiegmann, 1834 and *Eurylepis* Blyth 1854 (Sauria: Scincidae): A review. *Iranian Journal of Animal Biosystematics* 12: 225–237. <https://doi.org/10.22067/ijab.v12i2.47546>.
- Griffith, H., A. Ngo, and R.W. Murphy. 2000. A cladistic evaluation of the cosmopolitan genus *Eumeces* Wiegmann (Reptilia, Squamata, Scincidae). *Russian Journal of Herpetology* 7: 1–16. <https://doi.org/10.30906/1026-2296-2000-7-1-1-16>.
- Saikia, U., D.K. Sharma, and R.M. Sharma. 2007. Checklist of the reptilian fauna of Himachal Pradesh, India. *Reptile Rap* 8: 6–9.