



Age-dependent phenotypes of cognitive impairment as sequelae of SARS-CoV-2 infection

Alemán, Gabriela Gonzalez ; Vavougiou, George D; Tartaglia, Carmela; Uvais, Nalakath A; d'Avossa, Giovanni; Neidre, Daria; de Erausquin, Gabriel

Frontiers in Aging Neuroscience

Published: 07/01/2025

Publisher's PDF, also known as Version of record

[Cyswllt i'r cyhoeddiad / Link to publication](#)

Dyfyniad o'r fersiwn a gyhoeddwyd / Citation for published version (APA):

Alemán, G. G., Vavougiou, G. D., Tartaglia, C., Uvais, N. A., d'Avossa, G., Neidre, D., & de Erausquin, G. (2025). Age-dependent phenotypes of cognitive impairment as sequelae of SARS-CoV-2 infection. *Frontiers in Aging Neuroscience*, 16(1), 1-11.

Hawliau Cyffredinol / General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



OPEN ACCESS

EDITED BY

Fabrizia D'Antonio,
Sapienza University of Rome, Italy

REVIEWED BY

Christian Barbato,
National Research Council (CNR), Italy
Beatrice Paradiso,
University of Milan, Italy

*CORRESPONDENCE

Gabriel A. de Erausquin
✉ deerausquin@uthscsa.edu

†These authors have contributed equally to this work

RECEIVED 13 May 2024

ACCEPTED 28 October 2024

PUBLISHED 07 January 2025

CITATION

Gonzalez Aleman G, Vavougiou GD, Tartaglia C, Uvais NA, Guekht A, Hosseini AA, Lo Re V, Ferreccio C, D'Avossa G, Zamponi HP, Figueredo Aguiar M, Yecora A, Ul Haq Katshu MZ, Stavrou VT, Boutlas S, Gourgouljanis KI, Botero C, González Insúa F, Perez-Lloret S, Zinchuk M, Gersamija A, Popova S, Bryzgalova Y, Sviatskaya E, Russelli G, Avorio F, Wang S, Edison P, Niimi Y, Sohrabi HR, Mukaetova Ladinska EB, Neidre D and de Erausquin GA (2025) Age-dependent phenotypes of cognitive impairment as sequelae of SARS-CoV-2 infection. *Front. Aging Neurosci.* 16:1432357. doi: 10.3389/fnagi.2024.1432357

COPYRIGHT

© 2025 Gonzalez Aleman, Vavougiou, Tartaglia, Uvais, Guekht, Hosseini, Lo Re, Ferreccio, D'Avossa, Zamponi, Figueredo Aguiar, Yecora, Ul Haq Katshu, Stavrou, Boutlas, Gourgouljanis, Botero, González Insúa, Perez-Lloret, Zinchuk, Gersamija, Popova, Bryzgalova, Sviatskaya, Russelli, Avorio, Wang, Edison, Niimi, Sohrabi, Mukaetova Ladinska, Neidre and de Erausquin. This is an open-access article distributed under the terms of the [Creative Commons Attribution License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

Age-dependent phenotypes of cognitive impairment as sequelae of SARS-CoV-2 infection

Gabriela Gonzalez Aleman^{1†}, George D. Vavougiou^{2,3†}, Carmela Tartaglia^{4,5†}, Nalakath A. Uvais⁶, Alla Guekht^{7,8}, Akram A. Hosseini^{9,10}, Vincenzina Lo Re^{11,12}, Catterina Ferreccio^{13,14}, Giovanni D'Avossa¹⁵, Hernan P. Zamponi¹⁶, Mariana Figueredo Aguiar¹⁷, Agustin Yecora^{16,17}, Mohammad Zia Ul Haq Katshu¹⁸, Vasileios T. Stavrou^{2,3}, Stylianos Boutlas², Konstantinos I. Gourgouljanis², Camila Botero¹, Francisco González Insúa¹, Santiago Perez-Lloret¹⁹, Mikhail Zinchuk⁷, Anna Gersamija⁷, Sofya Popova⁷, Yulia Bryzgalova⁷, Ekaterina Sviatskaya⁷, Giovanna Russelli^{11,12}, Federica Avorio^{11,12}, Sophia Wang^{20,21}, Paul Edison^{22,23}, Yoshiki Niimi²⁴, Hamid R. Sohrabi²⁵, Elizabeta B. Mukaetova Ladinska^{26,27}, Daria Neidre²⁸ and Gabriel A. de Erausquin^{28,29,30*}

¹Department of Psychology, School of Psychology and Psychopedagogy, Universidad Catolica Argentina, Buenos Aires, Argentina, ²Department of Neurology, Medical School, University of Cyprus, Nicosia, Cyprus, ³Department of Respiratory Medicine, University of Thessaly, Larissa, Greece, ⁴Tanz Centre for Research in Neurodegenerative Diseases, University of Toronto, Toronto, ON, Canada, ⁵Memory Clinic, Department of Neurology, Toronto Western Hospital, Toronto, ON, Canada, ⁶Department of Psychiatry, Iqraa International Hospital and Research Centre, Calicut, India, ⁷Department of Neurology, Moscow Research and Clinical Centre for Neuropsychiatry, Moscow, Russia, ⁸Department of Neurology, Pirogov Russian National Research Medical University, Moscow, Russia, ⁹Nottingham University Hospitals NHS Trust, Nottingham, United Kingdom, ¹⁰Nottingham Sir Peter Mansfield Imaging Centre, University of Nottingham, Nottingham, United Kingdom, ¹¹Neurology Service, Department of Diagnostic and Therapeutic Services, IRCCS ISMETT, Palermo, Italy, ¹²Department of Experimental Medicine and Clinical Neuroscience, University of Pittsburgh Medical Center (UPMC), Palermo, Italy, ¹³Department of Public Health School of Medicine, Pontificia Universidad Catolica de Chile, Santiago, Chile, ¹⁴Advanced Center for Chronic Diseases, ACCDiS, Santiago, Chile, ¹⁵School of Psychology and Sports Sciences, Bangor University, Bangor, United Kingdom, ¹⁶Secretariat for Mental Health and Addictions, Ministry of Health, Government of Jujuy, San Salvador de Jujuy, Argentina, ¹⁷Instituto San Lazaro de Neurociencias, Fundacion de Lucha contra los Trastornos Neurologicos y Psiquiatricos en Minorias, FULTRA, San Salvador de Jujuy, Argentina, ¹⁸Institute of Mental Health, University of Nottingham, Nottinghamshire Healthcare NHS Foundation Trust, Nottingham, United Kingdom, ¹⁹Health Observatory, Vice Rectorate for Research, Universidad Catolica Argentina, Buenos Aires, Argentina, ²⁰Department of Psychiatry, Indiana University School of Medicine, Indianapolis, IN, United States, ²¹Indiana Alzheimer's Disease Research Center, Indianapolis, IN, United States, ²²Department of Brain Sciences, Faculty of Medicine, Imperial College London, London, United Kingdom, ²³Cardiff University, Cardiff, United Kingdom, ²⁴Faculty of Medicine, University of Tokyo, Tokyo, Japan, ²⁵Murdoch University Centre for Healthy Ageing, School of Psychology, Murdoch University, Murdoch, WA, Australia, ²⁶Department of Psychology and Visual Sciences, University of Leicester, Leicester, United Kingdom, ²⁷The Evington Centre, Leicester General Hospital, Leicester, United Kingdom, ²⁸Laboratory for Brain Development, Modulation and Repair, Glenn Biggs Institute for Alzheimer's and Neurodegenerative Diseases, University of Texas Health San Antonio, San Antonio, TX, United States, ²⁹Laboratory of Electrophysiology Imaging, Radiology Research Institute, University of Texas Health San Antonio, San Antonio, TX, United States, ³⁰Department of Neurology, Joe & Teresa Long School of Medicine, University of Texas Health San Antonio, San Antonio, TX, United States

Cognitive changes associated with PASC may not be uniform across populations. We conducted individual-level pooled analyses and meta-analyses of cognitive

assessments from eight prospective cohorts, comprising 2,105 patients and 1,432 controls from Argentina, Canada, Chile, Greece, India, Italy, Russia, and the UK. The meta-analysis found no differences by country of origin. The profile and severity of cognitive impairment varied by age, with mild attentional impairment observed in young and middle-aged adults, but memory, language, and executive function impairment in older adults. The risk of moderate to severe impairment doubled in older adults. Moderately severe or severe impairment was significantly associated with infection diagnoses (chi-square = 26.57, $p \leq 0.0001$) and the severity of anosmia (chi-square = 31.81, $p \leq 0.0001$). We found distinct age-related phenotypes of cognitive impairment in patients recovering from COVID-19. We identified the severity of acute illness and the presence of olfactory dysfunction as the primary predictors of dementia-like impairment in older adults.

KEYWORDS

COVID-19, cognitive decline, international cohort, age-dependent, long COVID

Introduction

COVID-19 is a disease caused by the coronavirus SARS-CoV-2. SARS-CoV-2 infection can cause severe respiratory disease, resulting in pneumonia and death. COVID-19 has now spread to 222 countries and territories worldwide, with more than 704 million cases and >7 million deaths [worldometers.info/coronavirus]. Since its emergence, our understanding of COVID-19 has evolved to recognize its systemic features and potentially long-lasting consequences. As the exposure of the population to SARS-CoV-2 accelerated, these symptoms became increasingly recognized as a syndrome known as post-acute sequelae of COVID-19 (PASC) or long COVID (Nalbandian et al., 2021; Proal and VanElzakker, 2021).

Epidemiological studies suggest that PASC continues to have a global impact. In 2021, a systematic review of studies involving >250,000 survivors of COVID-19 found that PASC was evident in more than half, and PASC symptoms that persisted for 6 months or longer after the acute phase of infection affected multiple organs, leading to respiratory, cardiovascular and neurological symptoms (Groff et al., 2021). In an analysis of nearly 1.5 million COVID-19 patients followed up for 2 years after the initial infection, the risk for cognitive impairment remained higher in the group exposed to COVID-19 (Taquet et al., 2022). In 2022, the Human Phenotype Ontology meta-analyzed findings from 81 PASC cohorts, indicating that cognitive impairment and other neuropsychiatric symptoms were consistent symptoms in PASC (Deer et al., 2021).

Neuropsychiatric PASC may be distinct from other multi-systemic manifestations. This may be explained by a biological interplay between SARS-CoV-2 infection and neurodegeneration (de Erausquin et al., 2021; Li et al., 2022). For instance, levels of neurodegenerative biomarkers such as total tau, p-tau181, glial fibrillary acidic protein (GFAP), and neurofilament light (NFL) were increased following COVID-19 infection in patients without a history of neurodegenerative disease (Frontera et al., 2022). Notably, plasma NFL, a marker of axonal damage, was high after SARS-CoV-2 infection in the absence of neurological manifestations, (Verde et al., 2022) but in association with increased levels of inflammatory cytokines (Hirzel et al., 2022).

An interpretation of these findings suggests that SARS-CoV-2 may damage the central nervous system either directly (Meinhardt et al., 2021), or mediated by an abnormal immune response (Jarius et al., 2022). Regardless of the underlying mechanism, data from the UK Biobank show brain volume reductions in limbic and olfactory pathways (Douaud et al., 2022), placing anosmia and cognitive deficits within the same syndromic spectrum (Deer et al., 2021; de Erausquin et al., 2021; Zamponi et al., 2021; Galderisi et al., 2024). Taken together, these associations between clinical phenotypes, imaging, and underlying biology indicate that neurocognitive disorders may be a distinct phenomenon among other PASC manifestations (de Erausquin et al., 2021; Frontera and Simon, 2022).

The impact of the pandemic has not been uniform across human groups or regions of the world. Notably, and beyond obvious comorbidities that increase the risk of various diseases, such as obesity or diabetes mellitus (Lee et al., 2024), susceptibility to the infection's deleterious effects has been higher across groups from specific ancestries. In the U.S.A., the highest morbidity and mortality rates were found in Amerindian minorities, followed by African Americans and Hispanics (Lundberg et al., 2023). Similarly, Amerindian (Maya) ancestry was associated with Southern Mexico's highest morbidity and mortality rates (Rangel-Méndez et al., 2023). In the U.K., the highest morbidity and mortality rates were reported for Bangladeshi and Pakistani minorities, followed by Black Africans and Black Caribbean [Office for National Statistics (ONS), 2023]. Notably, these differential effects have disappeared after the Omicron variant of SARS-CoV-2 became prevalent [Office for National Statistics (ONS), 2023]. Therefore, substantial human host factors influence SARS-CoV-2 infection, and several recent studies point to genetics as the primary contributor (Nakanishi et al., 2021; Garg et al., 2024). A genetic predisposition may be causally linked to some (but not all) PASC symptoms (Shenoy et al., 2023; Tang et al., 2023).

Genetically based protein alterations can lead to large differences in differential response to SARS-CoV-2 infection, and some manifestations of PASC may be influenced by genetic variants shared with other diseases. For cognitive decline, the APOE gene has long been implicated in the risk of Alzheimer's disease, and

TABLE 1 Description of cohorts participants.

	N	Controls	Patients	Age	Education (in years)
Argentina	866	137	729	67.13 ± 5.7	10.23 ± 4.72
Canada	97	0	97	52.23 ± 11.16	14.68 ± 3.62
Chile	467	361	106	57.1 ± 8.81	
Greece	252	47	205	54 ± 11.66	
India	340	169	171	37.29 ± 14.41	11.65 ± 3.61
Italy	28	6	22	57.23 ± 10.96	10.64 ± 4.07
Russia	1,416	689	727	47.63 ± 14.65	
UK	71	23	48	58.63 ± 11.49	
	3,537	1,432	2,105		

the e4/e4 genotype has been shown to increase susceptibility to COVID-19, excess neuroinflammation, and reduced antiviral response (Kuo et al., 2020; Farrer et al., 1997; Severe Covid-19 GWAS Group et al., 2020). Thus, APOE may play a causal role in the severity of neurological symptoms in infection. SARS-CoV-2 infection significantly alters molecular pathways implicated in brain inflammation, and certain viral entry factors are highly expressed in cells in the blood-brain barrier. Indeed, PASC cognitive sequelae may resemble early Alzheimer's disease, and olfactory dysfunction may be an accurate predictor of the severity of cognitive sequelae (Gonzalez-Aleman et al., 2022).

In sum, cognitive changes associated with PASC may not be uniform across populations, and refining clinical phenotypes is essential to pursuing biomarkers of genetic studies in the future. We carried out individual-level pooled data analyses from Argentina, Canada, Chile, Greece, India, Italy, Russia, and the United Kingdom (UK), harmonizing following published protocols (de Erausquin et al., 2022) to define cognitive impairment profiles.

Results

Description of the overall sample

The whole sample consisted of 3,537 (2,105 infected and 1,432 uninfected controls, Table 1). Participants ranged in age from 18 to 97 years. The mean duration of formal education was 10.66 ± 4.5 years, and the mean age was 52.81 ± 15.76 years; 36.5% were men and 64.5% were women.

Pooled analysis of individual-level data

The cognitive performance of pooled, individual-level data was compared first task-wise for the entire sample. The infection performed significantly worse than the controls on the Trail Making Test (Part B), Symbol Digit Coding, Memory Delayed Recall, Recognition, Verbal Semantic Fluency (animals), and Naming (Figure 1). By contrast, a meta-analysis to test for differences between patients and controls by country of origin found that all patients showed impairment in executive function

as indicated by performance only on the Trail Making Test-part B (Figure 2), suggesting loss of cognitive flexibility.



Based on visual inspection of frequency plots by age, a bimodal distribution of overall impairment was identified, and an empirical cut-off point was identified at 59 years of age (Figure 3). Specifically, the distribution of mild cognitive impairment (1 domain only) by age was examined, along with the distribution of moderate to severe cognitive impairment (2 or 3 domains) (Figure 3). We tested the hypothesis of independence of these distributions and found it to be highly significant ($t = -10.95$, $df = 642.86$, $p \leq 0.0001$). The median age in the mild cognitive decline group was 57.6, and in the moderate to severe cognitive impairment group, it was 65.8 years. Therefore, subjects 60 years of age or older (older adults) and subjects younger than 59 years of age (middle-aged and young adults) were treated as independent samples henceforth. The results are shown in Tables 2, 3.

Older adults were significantly impaired when compared to controls on Digit Span Forward performance ($t = -4.52$, $p \leq 0.001$); delayed recall ($t = 3.6$, $p < 0.001$); recognition tasks ($t = 6.94$, $p < 0.001$); and the semantic verbal fluency task ($t = 2.28$, $p = 0.02$). In young and middle-aged adults, we found milder deficits in recognition and naming only ($t = 2.75$ and $p = 0.006$ and $t = 3.3$ and $p = 0.001$, respectively). Findings suggest mild attentional focus in young and middle-aged adults, whereas older adults display impairment in memory, language, and executive function.

Severity of impairment: age-dependent phenotypes

To classify the severity of impairment, we used the deficit criterion as a difference of 1.5 standard deviations in performance on the tests administered. In young and middle-aged adults, we obtained four groups with distinct cognitive performance (Table 4):

- normal cognition (68.2%).
- impairment in one cognitive dimension (23.4%).
- impairment in two cognitive dimensions (6.5%).
- impairment in three cognitive dimensions impaired (1.8%).




By strong contrast, older adults showed a distribution skewed toward more severe impairment:

- normal cognition (68.2%).
- impairment in one cognitive dimension (30.1%).

- impairment in two cognitive dimensions (11.3%).
- impairment in three cognitive dimensions (3.7%).

The odds ratio (OR) of moderate to severe cognitive impairment (2 or 3 cognitive dimensions affected) was calculated



for the entire sample (OR = 1.75), for young and middle-aged adults (OR = 0.71), and older adults (OR = 1.49). Thus, the risk was doubled in older adults compared to middle-aged and young adults.

Predictors of risk of moderately severe or severe impairment

Moderately severe or severe impairment (2 or 3 dimensions) was significantly associated with infection diagnoses (chi-square = 26.57, $p \leq 0.0001$) and with the presence and severity of anosmia (chi-square = 31.81, $p \leq 0.0001$). No association was found between gender and cognitive dysfunction (chi-square = 0.74; $p = 0.38$) and between gender and anosmia (chi-square = 2.37; $p = 0.12$). The distribution of subjects by country is shown in Table 1, along with the average age and years of formal education; neither of these variables contributed to risk. On the other hand, the risk of moderate to severe cognitive impairment was nearly doubled in the presence of anosmia compared with its absence (OR = 1.89) (Table 5). Lastly, we also examined OR for cognitive impairment according to the severity of COVID-19 during the acute illness, considering the need for hospitalization, oxygen therapy, and admission to the intensive care unit. The risk of cognitive decline doubled when patients were hospitalized (OR = 2.06) or required intensive care (OR = 2.01). If they required oxygen therapy, the odds ratio was 1.47 (Table 6). Confidence intervals for each odds ratio are reported in Tables 5, 6, along with a normal deviation (z-statistic) calculated as $\ln(\text{OR})/\text{SE}\{\ln(\text{OR})\}$ and the corresponding p -value.

Discussion

We studied multiple samples of adults exposed to SARS-CoV-2 infection and collected them in a variety of settings (community as well as polyclinics or hospitals) in developed and underdeveloped countries, including individuals from multiple under-represented ancestries. Our analysis confirms that cognitive deficits are frequently present, but with different phenotypes determined by age at exposure and severity of anosmia. Older age and anosmia were found to be risk factors for severe phenotypes of cognitive impairment consistent with dementia-like syndrome. On the other hand, meta-analysis identified impaired verbal fluency in cases vs. controls independent of age. Notably, we did not detect inter-sample differences, which could be attributed to ancestry effects.

Attentional skills and executive functioning were significantly impaired in younger patients (59 and younger) across cohorts, but deficits were both infrequent and mild. A systematic literature review and retrospective meta-analysis on slightly <1,000 mostly European patients revealed that following COVID-19, patients suffer from impaired memory, attention, and executive function verbal and specifically verbal fluency, regardless of acute disease severity (Daroische et al., 2021), but nearly half of the assessments were carried out by phone, reducing accuracy. More consistent with our findings, a community sample of young and middle-aged (<54 years old) from Canada identified significant impairments in executive function (measured and reported) associated with acute infection severity (Hall et al., 2022), and identical results were reported in a smaller sample in Russia (Manukyan et al., 2022), and the USA (Apple et al., 2022). Of note, mild subjective impairment in younger adults is disproportionate to objective findings on formal

TABLE 2 Pooled data all cohorts.

	N	Mean \pm	SD	ΔX	SE	t	df	p-value
Trail-making test–part A				4.26	3.53	1.2	1225	0.2300
Patients	933	111.09 \pm	56.72					
Controls	294	106.83 \pm	52.81					
Trail-making test–part B				–1.82	4.14	–0.44	624.58	0.6600
Patients	915	111.48 \pm	56.71					
Controls	288	113.3 \pm	75.90					
Span forward				–6.39	1.41	–4.52	605.985	0.0010
Patients	915	99.73 \pm	24.46					
Controls	296	93.34 \pm	19.93					
Span backward				–1.12	1.75	–0.64	1302	0.5200
Patients	1004	93.43 \pm	26.51					
Controls	300	92.3 \pm	27.04					
Symbol digit coding				1.21	2.60	0.45	704.542	0.6400
Patients	895	91.12 \pm	50.74					
Controls	284	92.34 \pm	34.31					
Short-term memory				–0.65	1.80	–0.35	534.964	0.7200
Patients	901	96.82 \pm	30.94					
Controls	276	96.16 \pm	25.99					
Delayed recall memory				10.26	2.85	3.6	1,424	0.0010
Patients	1024	96.89 \pm	47.99					
Controls	402	107.15 \pm	48.75					
Recognition				10.78	1.68	6.94	298,960	0.0010
Patients	779	98.62 \pm	14.12					
Controls	249	109.4 \pm	25.28					
Verbal fluency-letter				3.9	3.17	1.23	571	0.2100
Patients	255	88.9 \pm	39.97					
Controls	318	92.81 \pm	35.82					
Verbal fluency-animals				3.94	1.72	2.28	773.266	0.0230
Patients	1058	93.04 \pm	28.05					
Controls	458	97 \pm	32.07					
Naming				3.65	1.84	1.97	620.871	0.0500
Patients	781	95.33 \pm	36.37					
Controls	264	98.98 \pm	23.42					

Cognitive performance in older adults (60 years old and above) vs. non-infected controls. Bold lettering indicates statistical significance.

testing and is associated with pre-existing psychiatric complaints (Apple et al., 2022).

Older adults were at over twice the risk of suffering memory, language, and executive function impairment that would be indistinguishable from early Alzheimer's disease with mild to moderate dementia. Once again, this finding is consistent with multiple previous reports in the literature (de Erausquin et al., 2021; Galderisi et al., 2024; Li et al., 2022; Tang et al., 2023; Gonzalez-Aleman et al., 2022; Miners et al., 2020) and many others. Similarly, as in most of the published literature, anosmia

was significantly associated with cognitive impairment (Gonzalez-Aleman et al., 2022; Global Burden of Disease Long COVID Collaborators et al., 2022; Llana et al., 2023, 2022; Sohrabi et al., 2012; Yahiaoui-Doktor et al., 2019) and specifically with memory loss in recovering COVID-19 patients (Gonzalez-Aleman et al., 2022; Llana et al., 2022; Sohrabi et al., 2012). The prevalence of dementia-like cognitive impairment in these cohorts strongly contrasts with the published prevalence of only 5% Alzheimer's type dementia in people aged 65 to 74 (Alzheimer's Association, 2024). Regardless of the presumed mechanism of PASC-related cognitive

TABLE 3 Pooled data all cohorts.

	N	Mean \pm	SD	ΔX	SE	t	df	p-value
Trail-Making Test- part A				-0.33	2.59	-0.13	1057	0.89
Patients	556	91.27 \pm	41.11					
Controls	503	90.49 \pm	43.42					
Trail-Making Test- part B				1.62	2.29	0.7	1155	0.89
Patients	640	89.56 \pm	37.48					
Controls	517	91.18 \pm	40.32					
Span forward				3.17	2.97	1.06	1022	0.28
Patients	518	102.11 \pm	19.56					
Controls	506	105.28 \pm	64.83					
Span backward				-0.19	1.43	-0.13	1214	0.89
Patients	667	102.35 \pm	24.91					
Controls	549	102.15 \pm	24.80					
Encryption				2.8	1.85	1.51	971.193	0.13
Patients	522	104.63 \pm	27.53					
Controls	486	107.43 \pm	31.01					
Short-term memory				0.92	1.06	0.86	1180	0.38
Patients	621	103.18 \pm	18.81					
Controls	561	104.1 \pm	17.57					
Delayed recall memory				-4.3	2.50	-1.6	1658	0.09
Patients	885	117.49 \pm	18.81					
Controls	852	113.19 \pm	46.05					
Recognition				5.4	2.01	2.75	609	0.006
Patients	269	109.69 \pm	21.42					
Controls	342	115.09 \pm	27.03					
Verbal fluency-letter				1.12	1.98	0.56	1328	0.57
Patients	628	105.06 \pm	36.76					
Controls	702	106.19 \pm	35.49					
Verbal fluency-animals				1.62	1.98	1.07	1814	0.28
Patients	908	103.08 \pm	31.15					
Controls	908	104.7 \pm	33.24					
Naming				3.46	1.00	3.3	501.359	0.001
Patients	299	99.65 \pm	15.52					
Controls	376	103.12 \pm	10.49					

Cognitive performance in young and middle-aged adults (60 years old and above) vs. non-infected controls. Bold lettering indicates statistical significance.

impairment, with current reported rates of infection worldwide, a sharp increase in dementia-like syndromes would place a massive strain on health systems already burdened by this problem in aging populations (Alzheimer's Association, 2024).

The relationship between neurodegeneration and olfactory dysfunction is well established. For instance, poor olfactory discrimination has been shown to predict cognitive decline in older community-dwelling adults (Sohrabi et al., 2012). In the population-based LIFE-Adult Study (n = 6,783), better olfactory function was associated with better performance on cognitive tests

even after adjustment for known confounders (Yahiaoui-Doktor et al., 2019).

Older age, anosmia, and cognitive decline are correlated with Alzheimer's disease and related dementia (ADRD). Entorhinal cortex thinning and anosmia are early hallmarks of memory decline (Murphy, 2019). Indeed, memory function and olfaction are intrinsically connected via neuroanatomical and functional pathways, leading to the hypothesis that exposure of the olfactory network to SARS-CoV-2 infection could account for tandem anosmia and cognitive impairment in at least a subset of COVID-19

and produce ADRD-like neuropathology (de Erausquin et al., 2021; Zamponi et al., 2021; Gonzalez-Aleman et al., 2022; de Erausquin et al., 2022). The current findings confirm the interconnection of anosmia and cognitive impairment following SARS-CoV-2 infection in a multinational sample.

Several independent studies have also provided support for the neuropathological and functional substrate of this hypothesis. In mammalian animal models, the olfactory networks have been

identified as a potential route of neuroinvasion for SARS-CoV-2, which was shown to infect both brain endothelial cells and neurons in rhesus monkey (Jiao et al., 2021) and nasal epithelial, olfactory sensory neurons, and hippocampal dendritic spines in rodents (Kishimoto-Urata et al., 2022). In humans, a post-mortem study of COVID-19 patients identified viral particles in the olfactory neurons and their projections (Thakur et al., 2021). These findings were associated with microgliosis and elevated neuroinflammatory markers in cerebrospinal fluid Thakur et al., 2021, and both findings have been independently confirmed (Boutajangout et al., 2021; Poloni et al., 2021). An extensive review of neuropathological findings in 184 patients who died from complications of acute COVID-19 found microglial activation in close to half and detectable virus in the human cerebrum, cerebellum, cranial nerves, olfactory bulb, as well as in the olfactory epithelium (Lou et al., 2021).

Whether viral particles can persist in surviving patients remains an open and much more complex question to address. As suggested above, inflammatory processes independent of neuroinvasion may mediate an alternate mechanism. Analysis of frontal lobe transcriptomes donated by COVID-19 patients revealed that exposure to SARS-CoV-2 was associated with molecular signatures of aging mediated by type I/III interferon (Mavrikaki et al., 2022). The same innate immune pathway is associated with Alzheimer's disease pathobiology and immune-mediated neuronal injury in rodents (Roy et al., 2022). More directly relevant to this discussion, post-mortem brain tissue donated by COVID-19 patients revealed abnormalities in shared pathways associated with Alzheimer's disease, including those involved innate immunity, tau hyperphosphorylation, as well as a site-specific calbindin hypoactivity in the hippocampus (Reiken et al., 2022). Lastly, interleukins IL-1 β , IL-6, and tumor necrosis factor plasma levels are elevated in PASC sufferers (Schulthei et al., 2022), and are also correlated to cognitive impairment in Alzheimer's disease (Silva et al., 2021), and with severity of hyposmia after COVID-19 (Liang et al., 2022).

Stimulation of the peripheral immune system by a disease-associated molecular pattern (DAMP) has been shown to result in an upregulation of complement-related genes in the hippocampus, even in the case of sterile DAMPs (Michalovicz et al., 2015). In the context of COVID-19, induction of type I interferon signaling and proinflammatory cytokines may affect brain function

TABLE 4 Frequency of cognitive impairment in all cohorts with (patients) and without infection (controls).

	Controls	Patients	Total
No impairment	1,032	1,381	2,413
Mild impairment (1 cognitive dimension affected)	306	523	829
Moderate impairment (2 cognitive dimensions affected)	75	156	231
Severe impairment (3 cognitive dimensions affected)	13	51	64

TABLE 5 Risk of severe cognitive impairment with moderate vs. severe anosmia.

	Anosmia	
	Severe	Moderate
No impairment	525	720
Mild impairment	145	244
Moderate impairment	38	117
Severe impairment	9	47
Chi-Square =	30,81 (df = 3; p-value = 0.000)	
Odds ratio for severe impairment =	1.89 (95% CI: 1.35-2.64)	
z statistic =	3.766; p-value = 000002	

Includes cohorts from Argentina, Greece, Italy, and Russia.

TABLE 6 Risk of severe cognitive impairment in young vs. older adults (top) and stratified by severity of acute COVID-19 illness.

	OR	95% CI	Z statistics	p-value
Cognitive impairment vs. age				
All sample	1.7578	1.35-2.28	4.218	0.0001
Older adults	1.4923	1.07-2.08	2.374	0.0176
Young and middle-aged adults	0.7191	0.42-1.24	1.192	0.2334
Cognitive impairment vs. severity of COVID-19				
Oxygen therapy (N = 154)	1.4696	0.97-2.22	1.832	0.07
Hospitalization (N = 81)	2.0653	1.44-2.96	3.952	0.0001
Intensive care unit (N = 9)	2.0065	0.91-4.39	1.738	0.08

via vascular pathways; alternatively, direct infection of the olfactory neuroepithelium may alter brain function via molecular mimicry or inflammatory events through its projections to areas in the extended olfactory network. These events and the molecules involved may trigger neurodegenerative diseases such as neuropathology (Vavougiou et al., 2022). Aside from molecules that can mediate these events directly and, in a dose-response manner (e.g., IL-6), adult neurogenesis sites such as the hippocampi and the olfactory network would be particularly vulnerable, accounting for the selective vulnerability of these networks and the resulting anosmia-cognitive impairment phenotype we report on. Of course, our results do not exclude other possible interpretations, such as vascular lesions or dysfunction of the olfactory pathway.

The results of our study should be interpreted in the context of its strengths and potential limitations. The main strength of our study is the use of prospective, harmonized data from a global collaboration of scientists working in different settings and with many underrepresented populations. The pooled analysis of individual-level data provides a unique insight into the phenotypes and clinical associations of cognitive impairment following COVID-19, which show several consistent aspects across centers. This was possible due to the inclusion of long-running cohorts functioning in the real-world setting initiated at the beginning of the pandemic. Another significant strength is that in most published studies, cognitive dysfunction may affect hospitalized patients in general, aside from diagnosis, whereas our cohorts include largely community-based samples with only an accurately proportional number of hospitalized cases.

Conversely, the most relevant limitation of our study is that this is a pooled analysis of heterogeneous studies rather than a multicentre study. Our approach's merit lies in assessing what amounts to real-world data and different approaches in diverse healthcare settings. For example, the population in Thessaly, Greece, represents an admixture of rural and urban areas and a small island complex. Similarly, data from the Canadian site is multi-ethnic as they serve a diverse community in an urban center, and the data from Argentina describe an Amerindian population rarely included in multinational studies. The heterogeneity of variables included in the analysis per study is another potential limitation. Studies included varying degrees of clinical, demographic, anthropometric, neuropsychological, fitness, and respiratory function assessments. Covariate analyses have accounted for these differences as far as reasonable.

We found distinct age-related phenotypes of cognitive impairment in patients recovering from COVID-19. Premorbid complaints emerged as the major predictor of mild impairment in young and middle-aged adults. In contrast, the severity of acute illness and the presence of olfactory dysfunction were the primary predictors of dementia-like impairment in older adults.

Data availability statement

The raw data supporting the conclusions of this article will be made available by the authors, without undue reservation.

Ethics statement

The studies involving humans were approved by multiple committees as samples from several countries are included. Approvals are available. The studies were conducted in accordance with the local legislation and institutional requirements. The participants provided their written informed consent to participate in this study.

Author contributions

GE: Conceptualization, Formal analysis, Funding acquisition, Investigation, Methodology, Project administration, Writing – original draft, Writing – review & editing. GG: Data curation, Formal analysis, Investigation, Methodology, Writing – original draft, Writing – review & editing. GV: Data curation, Formal analysis, Funding acquisition, Investigation, Methodology, Writing – original draft, Writing – review & editing. CT: Data curation, Formal analysis, Funding acquisition, Investigation, Methodology, Writing – review & editing. NU: Data curation, Funding acquisition, Investigation, Methodology, Writing – review & editing. AGu: Data curation, Formal analysis, Funding acquisition, Investigation, Methodology, Writing – review & editing. AH: Data curation, Formal analysis, Funding acquisition, Investigation, Methodology, Writing – review & editing. VL: Data curation, Formal analysis, Funding acquisition, Investigation, Methodology, Writing – review & editing. CF: Data curation, Formal analysis, Funding acquisition, Investigation, Methodology, Writing – review & editing. GD'A: Formal analysis, Methodology, Writing – review & editing. HZ: Data curation, Investigation, Writing – review & editing. MF: Data curation, Funding acquisition, Investigation, Methodology, Writing – review & editing. AY: Data curation, Funding acquisition, Investigation, Methodology, Writing – review & editing. MU: Data curation, Formal analysis, Funding acquisition, Investigation, Methodology, Writing – review & editing. VS: Data curation, Formal analysis, Funding acquisition, Investigation, Methodology, Writing – review & editing. SB: Data curation, Investigation, Methodology, Writing – review & editing. KG: Conceptualization, Funding acquisition, Investigation, Methodology, Writing – review & editing. CB: Data curation, Formal analysis, Writing – review & editing. FG: Data curation, Formal analysis, Methodology, Writing – review & editing. SP-L: Formal analysis, Methodology, Writing – review & editing. MZ: Data curation, Investigation, Methodology, Writing – review & editing. AGE: Data curation, Investigation, Methodology, Writing – review & editing. SP: Data curation, Investigation, Methodology, Writing – review & editing. YB: Data curation, Investigation, Methodology, Writing – review & editing. ES: Data curation, Investigation, Writing – review & editing. GR: Data curation, Investigation, Writing – review & editing. FA: Data curation, Investigation, Writing – review & editing. SW: Methodology, Writing – review & editing. PE: Methodology, Writing – review & editing. YN: Methodology, Writing – review & editing. HS: Methodology, Writing – review & editing. EBM-L: Conceptualization, Methodology, Writing – review & editing. DN: Investigation, Project administration, Writing – review & editing.

Funding

The author(s) declare financial support was received for the research, authorship, and/or publication of this article. The authors were funded completely or partially by: Fundacion FULTRA, Argentina (GG, HZ, AY, MF), National Institute of Aging Grants # U19AG076581, P30AG059305, and P30AG066546 (GE), ISIDORe (PID: 23245) (GV), COVALENT study, included herein received funding from EU-OSHA (SB, KG), University Hospital Foundation (CT), IQRAA International Hospital and Research Center, Calicut, Kerala, India (NU, MU), Italian Health Ministry, Ricerca Corrente 2023 (VL, GR, FA), ANID/CONICYT FONDECYT Regular 1212066; ANID/FONDAP/15130011 (CF), NIA K23AG06255, NIA R21AG074179, NIA P30AG072976 (SW), JPND/Italian Ministry/(European Union) Medical Research Council and National Institute for Health Research, UK (PE), Western Australia Department of Jobs, Tourism, Science and Innovation grant for the Effect of COVID-19 on mental health of the elderly (HS). In addition, GV, KG, and VS would like to thank the personnel of the COVID-19 outpatient clinic and as well as research collaborators on long COVID research: Nico Biagi, Dimitrios Mysiris, Kyriaki Astara, Eleana Pitaraki, Konstantinos

Tourlakopoulos, Eirini Papayianni, Glykeria Tsimirona, Ílias E. Dimeas, Pelagia Foka, Olga Papaggeli, Georgia Xiromerisiou, Sotirios Zarogiannis, Zoe Daniil 2, and Georgios Hadjigeorgiou.

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

The author(s) declared that they were an editorial board member of Frontiers, at the time of submission. This had no impact on the peer review process and the final decision.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

References

- Alzheimer's Association (2024). Alzheimer's Disease Facts and Figures. *Alzheimers Dement.* 2024:20. Available at: <https://www.alz.org/media/Documents/alzheimers-facts-and-figures.pdf>
- Apple, A. C., Oddi, A., Peluso, M. J., Asken, B. M., Henrich, T. J., Kelly, J. D., et al. (2022). Risk factors and abnormal cerebrospinal fluid associate with cognitive symptoms after mild COVID-19. *Ann. Clin. Transl. Neurol.* 9, 221–226. doi: 10.1002/acn3.51498
- Boutajangout, A., Frontera, J., Debure, L., Vedvyas, A., Faustina, A., Wisniewski, T., et al. (2021). Plasma biomarkers of neurodegeneration and neuroinflammation in hospitalized COVID-19 patients with and without new neurological symptoms. *Alzheimers Dement.* 17:57892. doi: 10.1002/alz.057892
- Daroische, R., Hemminghyth, M. S., Eilertsen, T. H., Breivite, M. H., and Chwiczczuk, L. J. (2021). Cognitive impairment after COVID-19—a review on objective test data. *Front. Neurol.* 12:699582. doi: 10.3389/fneur.2021.699582
- de Erausquin, G. A., Snyder, H., Brugha, T. S., Seshadri, S., Carrillo, M., Sagar, R., et al. (2022). Chronic neuropsychiatric sequelae of SARS-CoV-2: protocol and methods from the Alzheimer's Association Global Consortium. *Alzheimers Dement.* 8:e12348. doi: 10.1002/trc2.12348
- de Erausquin, G. A., Snyder, H., Carrillo, M., Hosseini, A. A., Brugha, T. S., Seshadri, S., et al. (2021). The chronic neuropsychiatric sequelae of COVID-19: the need for a prospective study of viral impact on brain functioning. *Alzheimer's and Dementia* 17, 1056–1065. doi: 10.1002/alz.12255
- Deer, R. R., Rock, M. A., Vasilevsky, N., Carmody, L., Rando, H., Anzalone, A. J., et al. (2021). Characterizing long COVID: deep phenotype of a complex condition. *EBioMedicine* 74:103722. doi: 10.1016/j.ebiom.2021.103722
- Douaud, G., Lee, S., Alfaro-Almagro, F., et al. (2022). SARS-CoV-2 is associated with changes in brain structure in UK Biobank. *Nature* 604, 697–707. doi: 10.1038/s41586-022-04569-5
- Farrer, L. A., Cupples, L. A., Haines, J. L., Hyman, B., Kukull, W. A., Mayeux, R., et al. (1997). Effects of age, sex, and ethnicity on the association between apolipoprotein E genotype and Alzheimer disease - a meta-analysis. *JAMA-J Am Med Assoc* 278, 1349–1356. doi: 10.1001/jama.1997.03550160069041
- Frontera, J. A., Boutajangout, A., Masurkar, A. V., Betensky, R. A., Ge, Y., Vedvyas, A., et al. (2022). Comparison of serum neurodegenerative biomarkers among hospitalized COVID-19 patients versus non-COVID subjects with normal cognition, mild cognitive impairment, or Alzheimer's dementia. *Alzheimers Dement.* 18, 899–910. doi: 10.1002/alz.12556
- Frontera, J. A., and Simon, N. M. (2022). Bridging knowledge gaps in the diagnosis and management of neuropsychiatric sequelae of COVID-19. *JAMA Psychiatry* 79, 811–817. doi: 10.1001/jamapsychiatry.2022.1616
- Galderisi, S., Perrotelli, A., Giuliani, L., Pisaturo, M. A., Monteleone, P., Pagliano, P., et al. (2024). Cognitive impairment after recovery from COVID-19: frequency, profile, and relationships with clinical and laboratory indices. *Eur. Neuropsychopharmacol.* 79, 22–31. doi: 10.1016/j.euroneuro.2023.11.001
- Garg, E., Arguello-Pascuali, P., Vishnyakova, O., Halevy, A. R., Yoo, S., Brooks, J. D., et al. (2024). Canadian COVID-19 host genetics cohort replicates known severity associations. *PLoS Genet.* 20:e1011192. doi: 10.1371/journal.pgen.1011192
- Global Burden of Disease Long COVID Collaborators, Hanson, S. W., Abbafati, C., Aerts, J. G., Al-Aly, Z., Ashbaugh, C., et al. (2022). Estimated global proportions of individuals with persistent fatigue, cognitive, and respiratory symptom clusters following symptomatic COVID-19 in 2020 and 2021. *JAMA* 328, 1604–1615. doi: 10.1001/jama.2022.18931
- Gonzalez-Aleman, G., Zamponi, H. P., Juarez-Aguayso, L., Bahl, A., Sanchez, D., Reyes-Dumeyer, D., et al. (2022). Olfactory dysfunction but not COVID-19 severity predicts severity of cognitive sequelae following SARS-CoV-2 infection in Amerindian older adults. *Alzheimer's Dementia* 18:e066868. doi: 10.1002/alz.066868
- Groff, D., Sun, A., Ssentongo, A. E., Ba, D. M., Parsons, N., Poudel, G. R., et al. (2021). Short-term and long-term rates of postacute sequelae of SARS-CoV-2 infection: a systematic review. *JAMA Network Open* 4:e2128568. doi: 10.1001/jamanetworkopen.2021.28568
- Hall, P. A., Meng, G., Hudson, A., Sakib, M. N., Hitchman, S. C., MacKillop, J., et al. (2022). Cognitive function following SARS-CoV-2 infection in a population-representative Canadian sample. *Brain Behav Immun Health* 21:100454. doi: 10.1016/j.bbih.2022.100454
- Hirzel, C., Grandgirard, D., Surial, B., Wider, M. F., Leppert, D., Kuhle, J., et al. (2022). Neuro-axonal injury in COVID-19: the role of systemic inflammation and SARS-CoV-2 specific immune response. *Ther. Adv. Neurol. Disord.* 15:17562864221080528. doi: 10.1177/17562864221080528
- Jarius, S., Pache, F., Körtvelyessy, P., Jelčić, I., Stettner, M., Franciotta, D., et al. (2022). Cerebrospinal fluid findings in COVID-19: a multicenter study of 150 lumbar punctures in 127 patients. *J. Neuroinflammation* 19:19. doi: 10.1186/s12974-021-02339-0
- Jiao, L., Yang, Y., Yu, W., Zhao, Y., Long, H., Gao, J., et al. (2021). The olfactory route is a potential way for SARS-CoV-2 to invade the central nervous system of rhesus monkeys. *Signal Transd. Target. Ther.* 6:169. doi: 10.1038/s41392-021-00591-7
- Kishimoto-Urata, M., Urata, S., Kagoya, R., Imamura, F., Nagayama, S., Reyna, R. A., et al. (2022). Prolonged and extended impacts of SARS-CoV-2 on the olfactory neurocircuit. *Sci. Rep.* 12:5728. doi: 10.1038/s41598-022-09731-7

- Kuo, C. L., Pilling, L. C., Atkins, J. L., Masoli, J. A. H., Delgado, J., Kuchel, G. A., et al. (2020). APOE e4 genotype predicts severe COVID-19 in the UK Biobank Community Cohort. *J. Gerontol. A Biol. Sci. Med. Sci.* 75, 2231–2232. doi: 10.1093/gerona/glaa131
- Lee, A., Seo, J., Park, S., Cho, Y., Kim, G., Li, J., et al. (2024). Type 2 diabetes and its genetic susceptibility are associated with increased severity and mortality of COVID-19 in UK Biobank. *Commun Biol.* 7:122. doi: 10.1038/s42003-024-05799-1
- Li, C., Liu, J., Lin, J., and Shang, H. (2022). COVID-19 and risk of neurodegenerative disorders: A Mendelian randomization study. *Transl. Psychiatry* 12:283. doi: 10.1038/s41398-022-02052-3
- Liang, Y., Mao, X., Kuang, M., Zhi, J., Zhang, Z., Bo, M., et al. (2022). Interleukin-6 affects the severity of olfactory disorder: a cross-sectional survey of 148 patients who recovered from Omicron infection using the Sniffin' Sticks test in Tianjin, China. *Int. J. Infect. Dis.* 123, 17–24. doi: 10.1016/j.ijid.2022.07.074
- Llana, T., Mendez, M., Garcés-Arilla, S., Hidalgo, V., Mendez-Lopez, M., Juan, M. C., et al. (2023). Association between olfactory dysfunction and mood disturbances with objective and subjective cognitive deficits in long-COVID. *Front. Psychol.* 14:1076743. doi: 10.3389/fpsyg.2023.1076743
- Llana, T., Mendez, M., Zorzo, C., Fidalgo, C., Juan, M.-., C., et al. (2022). Anosmia in COVID-19 could be associated with long-term deficits in the consolidation of procedural and verbal declarative memories. *Front. Neurosci.* 16:1082811. doi: 10.3389/fnins.2022.1082811
- Lou, J. J., Movassaghi, M., Gordy, D., Olson, M. G., Zhang, T., Khurana, M. S., et al. (2021). Neuropathology of COVID-19 (neuro-COVID): clinicopathological update. *Free Neuropathol.* 2:2–2. doi: 10.17879/freeneuropathology-2021-2993
- Lundberg, D. J., Wrigley-Field, E., Cho, A., Raquib, R., Nsoesie, E. O., Paglino, E., et al. (2023). COVID-19 mortality by race and ethnicity in US metropolitan and nonmetropolitan areas, March 2020 to February (2022). *JAMA Netw Open.* 6:e2311098. doi: 10.1001/jamanetworkopen.2023.11098
- Manukyan, P., Deviatierikova, A., Velichkovsky, B. B., and Kasatkin, V. (2022). The impact of mild COVID-19 on executive functioning and mental health outcomes in young adults. *Healthcare* 10:1891. doi: 10.3390/healthcare10101891
- Mavrikaki, M., Lee, J. D., Solomon, I. H., and Slack, F. J. (2022). Severe COVID-19 is associated with molecular signatures of aging in the human brain. *Nature Aging* 2, 1130–1137. doi: 10.1038/s43587-022-00321-w
- Meinhardt, J., Radke, J., Dittmayer, C., Franz, J., Thomas, C., Mothes, R., et al. (2021). Olfactory transmucosal SARS-CoV-2 invasion as a port of central nervous system entry in individuals with COVID-19. *Nat. Neurosci.* 24, 168–175. doi: 10.1038/s41593-020-00758-5
- Michalovicz, L. T., Lally, B., and Konat, G. W. (2015). Peripheral challenge with a viral mimic upregulates expression of the complement genes in the hippocampus. *J. Neuroimmunol.* 285, 137–142. doi: 10.1016/j.jneuroim.2015.06.003
- Miners, S., Kehoe, P. G., and Love, S. (2020). Cognitive impact of COVID-19: looking beyond the short term. *Alzheimers. Res. Ther.* 12:170. doi: 10.1186/s13195-020-00744-w
- Murphy, C. (2019). Olfactory and other sensory impairments in Alzheimer disease. *Nature Rev. Neurol.* 15, 11–24. doi: 10.1038/s41582-018-0097-5
- Nakanishi, T., Pigazzini, S., Degenhardt, F., Cordioli, M., Butler-Laporte, G., Maya-Miles, D., et al. (2021). Age-dependent impact of the major common genetic risk factor for COVID-19 on severity and mortality. *J. Clin. Invest.* 131:e152386. doi: 10.1172/JCI152386
- Nalbandian, A., Sehgal, K., Gupta, A., Madhavan, M. V., McGroder, C., Stevens, J. S., et al. (2021). Post-acute COVID-19 syndrome. *Nat. Med.* 27, 601–615. doi: 10.1038/s41591-021-01283-z
- Office for National Statistics (ONS) (2023). *Updating Ethnic and Religious Contrasts in Deaths Involving Coronavirus (COVID-19):24 January 2020 to 23 November 2022*. Washington, DC: U.S. Census Bureau.
- Poloni, T. E., Medici, V., Moretti, M., Visonà, S. D., Cirrincione, A., Carlos, A. F., et al. (2021). COVID-19-related neuropathology and microglial activation in elderly with and without dementia. *Brain Pathol.* 31:e12997. doi: 10.1111/bpa.12997
- Proal, A. D., and VanElzakker, M. B. (2021). Long COVID or post-acute sequelae of COVID-19 (PASC): an overview of biological factors that may contribute to persistent symptoms. *Front. Microbiol.* 12:698169. doi: 10.3389/fmicb.2021.698169
- Rangel-Méndez, J. A., Jaimes-Gurrusquieta, L., Sánchez-Cruz, J. F., Zarate-Murillo, M. I., Domínguez-Méndez, J., Laviada-Molina, H. A., et al. (2023). Characteristics of COVID-19 and mortality-associated factors during the first year of the pandemic in southeast Mexico. *Future Microbiol.* 18, 1147–1157. doi: 10.2217/fmb-2023-0052
- Reiken, S., Sittenfeld, L., Dridi, H., Liu, Y., Liu, X., Marks, A. R., et al. (2022). Alzheimer's-like signaling in brains of COVID-19 patients. *Alzheimer's Dement.* 18, 955–965. doi: 10.1002/alz.12558
- Roy, E. R., Chiu, G., Li, S., Propson, N. E., Kanchi, R., Wang, B., et al. (2022). Concerted type I interferon signaling in microglia and neural cells promotes memory impairment associated with amyloid β plaques. *Immunity* 55, 879–894. doi: 10.1016/j.immuni.2022.03.018
- Schultheiß, C., Willscher, E., Paschold, L., Gottschick, C., Klee, B., Henkes, S.-S., et al. (2022). The IL-1 β , IL-6, and TNF cytokine triad is associated with post-acute sequelae of COVID-19. *Cell Rep Med.* 3:100663. doi: 10.1016/j.xcrm.2022.100663
- Severe Covid-19 GWAS Group, Ellinghaus, D., Degenhardt, F., Bujanda, L., Buti, M., Albrillos, A., et al. (2020). Genomewide association study of severe Covid-19 with respiratory failure. *N. Engl. J. Med.* 383, 1522–1534. doi: 10.1056/NEJMoa2020283
- Shenoy, P. U., Udupa, H., Ks, J., Babu, S., Jain, N., Das, R., et al. (2023). The impact of COVID-19 on pulmonary, neurological, and cardiac outcomes: evidence from a Mendelian randomization study. *Front Public Health.* 11:1303183. doi: 10.3389/fpubh.2023.1303183
- Silva, N. M. L. E., Gonçalves, R. A., Pascoal, T. A., Lima-Filho, R. A. S., Resende, E., Vieira, E. L. M., et al. (2021). Pro-inflammatory interleukin-6 signaling links cognitive impairments and peripheral metabolic alterations in Alzheimer's disease. *Transl. Psychiatry* 11:251. doi: 10.1038/s41398-021-01349-z
- Sohrabi, H. R., Bates, K. A., Weinborn, M. G., Johnston, A. N., Bahramian, A., Taddei, K., et al. (2012). Olfactory discrimination predicts cognitive decline among community-dwelling older adults. *Transl. Psychiatry* 2, e118–e118. doi: 10.1038/tp.2012.43
- Tang, C. M., Li, G., and Cheung, C. L. (2023). COVID-19 and cognitive performance: a Mendelian randomization study. *Front Public Health.* 11:1185957. doi: 10.3389/fpubh.2023.1185957
- Taquet, M., Sillett, R., Zhu, L., Mendel, J., Camplisson, I., Dercon, Q., et al. (2022). Neurological and psychiatric risk trajectories after SARS-CoV-2 infection: an analysis of 2-year retrospective cohort studies including 1 284 437 patients. *Lancet Psychiatry* 9, 815–827. doi: 10.1016/S2215-0366(22)00260-7
- Thakur, K. T., Miller, E. H., Glendinning, M. D., Al-Dalahmah, O., Banu, M. A., Boehme, A. K., et al. (2021). COVID-19 neuropathology at Columbia University Irving Medical Center/New York Presbyterian Hospital. *Brain* 144, 2696–2708. doi: 10.1093/brain/awab148
- Vavougiou, G. D., de Erausquin, G. A., and Snyder, H. M. (2022). Type I interferon signaling in SARS-CoV-2 associated neurocognitive disorder (SAND): mapping host-virus interactions to an etiopathogenesis. *Front. Neurol.* 13:1063298. doi: 10.3389/fneur.2022.1063298
- Verde, F., Milone, I., Bulgarelli, I., Peverelli, S., Colombrita, C., Maranzano, A., et al. (2022). Serum neurofilament light chain levels in Covid-19 patients without major neurological manifestations. *J. Neurol.* 2022, 1–11. doi: 10.1007/s00415-022-11233-5
- Yahiaoui-Doktor, M., Luck, T., Riedel-Heller, S. G., Loeffler, M., Wirkner, K., Engel, C., et al. (2019). Olfactory function is associated with cognitive performance: results from the population-based LIFE-Adult Study. *Alzheimers. Res. Ther.* 11:43. doi: 10.1186/s13195-019-0494-z
- Zamponi, H. P., Juárez-Aguaysoy, L., Kukoc, G., Domínguez, M. E., Pini, B., Padilla, E. G., et al. (2021). Olfactory dysfunction and chronic cognitive impairment following SARS-CoV-2 infection in a sample of older adults from the Andes mountains of Argentina. *Alzheimers. Dement.* 17:e057897. doi: 10.1002/alz.057897