



## Factors Associated with Weight Change in Adults with Severe Mental Illness: Results from a Large Cross-Sectional Survey

Traviss-Turner, Gemma D; Lee, Ellen; Pratt, Peter; Hill, Andrew; Peckham, Emily

Published: 01/04/2025

Publisher's PDF, also known as Version of record

[Cyswllt i'r cyhoeddiad / Link to publication](#)

*Dyfyniad o'r fersiwn a gyhoeddwyd / Citation for published version (APA):*

Traviss-Turner, G. D., Lee, E., Pratt, P., Hill, A., & Peckham, E. (2025). *Factors Associated with Weight Change in Adults with Severe Mental Illness: Results from a Large Cross-Sectional Survey*. Preprints.

### Hawliau Cyffredinol / General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Article

Not peer-reviewed version

---

# Factors Associated with Weight Change in Adults with Severe Mental Illness: Results from a Large Cross-Sectional Survey

---

[Gemma D. Traviss-Turner](#)<sup>\*</sup>, [Ellen Lee](#), Peter Pratt, [Andrew J. Hill](#), [Emily Peckham](#)

Posted Date: 1 April 2025

doi: 10.20944/preprints202504.0025.v1

Keywords: weight gain; obesity; antipsychotic medication; severe mental illness



Preprints.org is a free multidisciplinary platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This open access article is published under a Creative Commons CC BY 4.0 license, which permit the free download, distribution, and reuse, provided that the author and preprint are cited in any reuse.

Article

# Factors Associated with Weight Change in Adults with Severe Mental Illness: Results from a Large Cross-Sectional Survey

Short title: Weight change in severe mental illness

Gemma D Traviss-Turner <sup>1,\*</sup>, Ellen Lee <sup>2</sup>, Peter Pratt <sup>3</sup>, Andrew J Hill <sup>1</sup> and Emily Peckham <sup>4</sup>

<sup>1</sup> Leeds Institute of Health Sciences, Clarendon Way, University of Leeds, Leeds, UK

<sup>2</sup> School of Medicine and Public Health, University of Sheffield, Regent Court, 30 Regent St, Sheffield, UK

<sup>3</sup> National Speciality Advisor Mental Health Pharmacy, NHS England, Wellington House, London, UK

<sup>4</sup> School of Health Sciences, Bangor University, Gwynedd, North Wales, UK

\* Correspondence: G.Traviss@leeds.ac.uk; <https://orcid.org/0000-0002-1770-6216>

**Abstract: Background/objectives:** Individuals with severe mental illness (SMI) have a reduced life expectancy of 15-25 years. This is due to a number of modifiable and non-modifiable risk factors. Levels of overweight and obesity in this group are 1.8 times higher than in the general population and diet quality is poor. Excess weight is linked to a range of serious long-term physical and mental health conditions. This paper presents the findings of a large cross-sectional survey of adults living with SMI. The survey aimed to understand the current weight, weight gain and diet quality of this group and secondly, to explore the association between weight change, diet quality, antipsychotic medication and weight management. **Methods:** Five hundred and twenty-nine participants (58% male, mean age 49.3) living with SMI completed the survey. **Results:** Results showed 42% of the sample were living with obesity and almost half reported having gained 6kg or more at least once in their adult life. Overall, 6% of the sample reported eating no fruit or vegetables and the same proportion had the highest consumption of carbonated drinks. There was no difference by weight category. Those taking antipsychotic medication and currently managing their weight were more likely to experience weight gain or fluctuation. **Conclusions:** These results suggest that excess weight and poor diet quality are a major problem in adults with SMI and that current weight management provision is ineffective in addressing the specific needs of people living with SMI.

**Keywords:** weight gain; obesity; antipsychotic medication; severe mental illness

## 1. Introduction

Individuals with severe mental illness (SMI) including schizophrenia and bipolar disorder experience poorer physical health than the general population. This group have a reduced life expectancy of 15-25 years [1–4]. Higher death rates are attributed to chronic physical health conditions including liver disease, respiratory disease, cardiovascular disease and cancer and also the behavioural consequences of living with SMI such as substance abuse and suicide [5].

This mortality gap is due to a number of modifiable and non-modifiable risk factors. Many are linked to the illness itself and associated medication and others are associated with the socio-economic environment in which these individuals live. People living in the most deprived areas are three times more likely to experience SMI than those living in the least deprived areas [6].

One area that requires attention in this group is the high levels of overweight and obesity. People with SMI are 1.8 times more likely to have obesity, that is a body mass index (BMI)  $\geq 30\text{kg/m}^2$  (PHE, 2018). Obesity is concerning as it is linked with other serious long term physical health complications such as diabetes, hypercholesterolaemia, metabolic syndrome, increased cardiovascular and

cerebrovascular morbidity and mortality. It is also associated with psychological co-morbidity including mood disorder, reduced quality of life and poor drug compliance [7–10].

The risk factors for higher rates of obesity in this group are poorly understood. People with SMI have higher rates of poverty and are more likely to live in obesogenic environments. These are characterised by limited access and resource for healthy food and where the built environment is not analogous with physical activity [5,11]. Diet quality is poorer in individuals with SMI and they are more likely to consume cheaper foods that are energy-dense, higher in salt and sugar and consume less fruit and vegetables [7]. In a large survey of adults with SMI, almost 8.5% reported eating no fruit or vegetables [12,13]. Lack of consumption of fruit and vegetables was more common in younger males, those who were unemployed, living in deprived neighbourhoods and those who did not rate healthy living as important [13]. Low food literacy and disordered eating patterns (binge eating, emotional eating and strict dietary restriction) are also common in people with SMI [7,14–16].

A unique risk factor for this group in terms of weight management is the routine use of antipsychotic medication. There is an established literature on the relationship between both typical antipsychotics (such as chlorpromazine) and newer atypical antipsychotic medications (clozapine, olanzapine risperidone) and significant and rapid weight gain [8]. However, there is variation in the propensity of these drugs to cause weight gain. The World Health Organisation guidelines recommend clinicians considers psychotropic medication with lower propensity to weight gain for people with SMI who are overweight, with obesity or at risk of developing obesity [17].

With this in mind, we conducted a large cross-sectional survey of adults living with SMI, which aimed to:

1. understand the characteristics of people with SMI in terms of current weight, weight change and diet quality of people with SMI
2. explore the association between participant weight change and;
  - a. diet quality and current weight category
  - b. antipsychotic medication
  - c. current weight management

Throughout this paper we use the umbrella term '*weight change*'. This term includes '*weight gain*' which we define as gaining 6kg or more and '*weight fluctuation*' which refers to gaining  $\geq 6$ kg more than once.

## Materials and Methods

### *Study Design*

This was a cross-sectional study in which a survey was completed online, by post, or with a researcher either over the telephone or in person. The study was conducted according to the guidelines laid down in the Declaration of Helsinki and all procedures involving human subjects/patients were approved by the West of Scotland Research Ethics Service in June 2018 (reference 18/WS/0107).

### *Participants*

To be eligible to complete the survey, participants had to be aged 18 years or over and have a documented diagnosis of schizophrenia, schizoaffective disorder or bipolar (DSM5 or ICD-10). The study aimed to recruit 500 participants based on expected response rate and experience from previous research.

### *Recruitment and Procedures*

There were two phases of recruitment for this study, prior to the Covid-19 pandemic (March 2020) and after the onset of the Covid-19 pandemic (from June 2021 until March 2022). The pre-pandemic recruitment method involved mass mailouts to people who had taken part in the Closing

the Gap: Health and Wellbeing Cohort (described in Peckham et al 2023 [12]) and had consented to recontact. Mailouts were sent from five sites, three secondary care NHS mental health trusts and two clinical commissioning groups. Those invited to take part were sent a postal invitation pack containing an invitation letter, participant information sheet the survey and a pre-paid envelope. Those who were interested in taking part were asked to complete and return the survey in the pre-paid envelope provided. After the commencement of the pandemic it became impractical to conduct mass mailouts due to limited access to office space, therefore a mainly remote method of recruitment was adopted when the study recommenced in June 2021. From June 2021 to March 2022 participants were recruited either from research databases held by NHS mental health trusts or through clinical contacts. Those who expressed an interest in taking part in the study were invited, in order of preference, to complete the survey over the phone with a researcher or when the pandemic restrictions were lifted complete the survey face to face with a researcher, complete the survey via an online link sent by SMS message or email, or via a hard copy sent in the post. Voluntary completion of the survey indicated consent to take part in the study. The survey took approximately 20 minutes to complete and participants maintained anonymity.

### *Measures*

The bespoke survey was developed by a multi-disciplinary research team of researchers and people with lived experience of SMI, a copy of which can be found on the Open Science Framework (<https://osf.io/rgc62/>). The survey had three sections: (A) physical activity and sedentary behaviour; (B) health; (C) socio-demographics. An overview of the measures included in this article are provided below. Analyses of physical activity and sedentary behaviour are reported elsewhere [18]. Minor amendments were made to the survey questions prior to the second wave of recruitment, PAQ1 refers to the initial version and PAQ2 the later version.

#### Health Measures

Participants were asked to rate their mental health in last 12 months as excellent, good, moderate, poor or very poor. Smoking status was recorded via the question 'do you smoke?' with responses; yes, no, I have never smoked, and no, but I used to smoke. Participants were then asked to rate their average level of fatigue in the last two weeks along with how much of their waking time they have spent feeling fatigued in the last two weeks, on a 5 point likert scale. The scores for each domain were multiplied to give a composite score of fatigue. Depression severity was measured using the Patient Health Questionnaire-8 (PHQ).

BMI was calculated via self-reported height and weight with a BMI of less than 18.5 being coded as underweight, 18.5-24.9 being healthy weight, 25-29.9 being overweight and 30 or having obesity. Self-reported lowest ever adult weight and highest ever adult weight and weight fluctuation were also recorded. To explore weight management participants were asked whether they were currently managing their weight and if so, i. whether it was to lose weight, maintain weight or gain weight and ii. how they were managing their weight (diet, exercise, other).

Diet was assessed through asking people how many portions of fruit and vegetables they eat per typical day, with options of 1, 2, 3, 4 or 5 or more. In addition, the number of carbonated beverages they consumed per day, with options of 1, 2, 3, 4, or 5 or more.

Participants were asked to list any medication they were taking with the option to leave that section blank if they didn't wish to say what medication they were taking. The complete list of medications reported was reviewed by the National Speciality Advisor Mental Health Pharmacy for NHS England (PP), who categorised them as being antipsychotic medication or non-antipsychotic medication. Antipsychotic medications were then categorised as low, medium or high propensity to weight gain based on findings of the Lancet review by Huhn et al of 402 studies and 53463 participants [19].

#### Sociodemographic Measures

Information was collected on age, gender and ethnicity.

### Data Analysis

All analyses were performed using Stata 17 software (StataCorp, 2021. Stata Statistical Software: Release 17. College Station, TX: StataCorp LLC.). Descriptive statistics (e.g., % [n], mean [SD], median [IQR]) were used to summarise respondent characteristics for the full cohort and by BMI classification.

Measures of weight change (weight gain, weight fluctuation, weight management) were compared between participants that reported using antipsychotic medication and those that did not via univariable and multivariable (adjusted for age and sex) logistic regression. Odds ratios and corresponding 95% confidence intervals were calculated. Participants who did not report any medication use were excluded from these analyses.

Weight gain and weight fluctuation were compared between participants who reported currently managing weight and those who reported not currently managing weight using logistic regression analogous to the above methods. Weight change (difference between lowest and highest adult weight) was compared between participants managing their weight and not managing their weight using univariable and multivariable (adjusted for age and sex) linear regression. Mean differences and corresponding 95% confidence intervals were calculated.

All analyses were performed using available data, characteristics were descriptively compared between participants that reported their medication use and those that did not (See <https://osf.io/rgc62/> for PAQ analysis plan)

## Results

### Participant Characteristics

Five hundred and twenty nine people completed the survey, 196 completed a postal survey, 106 people completed the survey online and 186 people completed the survey over the phone or in person (41). One hundred and fifty five completed PAQ1 and 374 people completed PAQ2 (the later version). Participant demographics and characteristics are detailed in Table 1.

**Table 1.** Demographics and characteristics of N=529 survey respondents.

|                            | Characteristic                     | n (%)                    |
|----------------------------|------------------------------------|--------------------------|
| <b>Age (years)</b>         | Number (%) of responses            | 519 (98%)                |
|                            | Mean (SD)                          | 49.3 (13.1)              |
|                            | Min., Max.                         | 20, 86                   |
| Gender                     | Female                             | 212 (40%)                |
|                            | Male                               | 308 (58%)                |
|                            | Prefer not to say                  | 4 (1%)                   |
|                            | Transgender                        | 3 (1%)                   |
|                            | Missing                            | 2 (0%)                   |
| Weight & Weight management | BMI (kg/m <sup>2</sup> ) Mean (SD) | 31.2 (7.8)               |
|                            | underweight (<18.5)                | 2 (0%)                   |
|                            | healthy weight (18.5-24.9)         | 93 (18%)                 |
|                            | overweight (25-29.9)               | 119 (22%)                |
|                            | obese (>=30)                       | 223 (42%)                |
|                            | Missing                            | 92 (17%)                 |
|                            |                                    | Lowest adult weight (kg) |
|                            | Number (%) of responses            | 421 (80%)                |
|                            | Mean (SD)                          | 65.5 (17.9)              |
|                            | Min., Max.                         | 31, 174                  |

|              |   |             |
|--------------|---|-------------|
|              | Highest adult weight (kg)                               |             |
|              | Number (%) of responses                                 | 427 (81%)   |
|              | Mean (SD)   | 99.1 (26.8) |
|              | Min., Max.  | 44, 250     |
|              | Difference between lowest and highest adult weight (kg) |             |
|              | Number (%) of responses                                 | 399 (75%)   |
|              | Mean (SD)   | 34.4 (21.7) |
|              | Min., Max.  | 0, 158      |
|              | Currently managing weight                               | 327 (62%)   |
|              | Managing weight via diet                                | 168 (32%)   |
|              | Managing weight via exercise                            | 139 (26%)   |
|              | Reason for managing weight                              |             |
|              | Gain weight   | 12 (2%)     |
|              | Lose weight   | 223 (42%)   |
|              | Stay the same weight                                    | 92 (17%)    |
|              | Missing   | 202 (38%)   |
|              | Gained weight (>6kg)                                    |             |
|              | No  | 267 (50%)   |
|              | Yes, once   | 160 (30%)   |
|              | Yes, more than once (fluctuation)                       | 80 (15%)    |
|              | Missing   | 22 (4%)     |
| Employment   | Professionally active                                   | 99 (19%)    |
|              | Not professionally active                               | 356 (67%)   |
|              | Other   | 21 (4%)     |
|              | Missing   | 53 (10%)    |
| Ethnicity    | UK/Irish/other white                                    | 416 (79%)   |
|              | Asian   | 21 (4%)     |
|              | Black/African/Caribbean                                 | 26 (5%)     |
|              | Mixed multiple ethnic groups                            | 14 (3%)     |
|              | Other   | 12 (2%)     |
|              | Missing   | 40 (8%)     |
| Diet quality | Consumption of fruit and vegetables                     |             |
|              | 0   | 32 (6%)     |
|              | 1   | 102 (19%)   |
|              | 2   | 119 (22%)   |
|              | 3   | 122 (23%)   |
|              | 4   | 58 (11%)    |
|              | 5 or more   | 91 (17%)    |
|              | Missing   | 5 (1%)      |
|              | Carbonated drinks consumption                           |             |
|              | I do not drink carbonated soft drinks                   | 194 (37%)   |
|              | Less than one time per day                              | 113 (21%)   |
|              | 1 time per day  | 71 (13%)    |
|              | 2 times per day   | 45 (9%)     |
|              | 3 times per day   | 40 (8%)     |
|              | 4 times per day   | 21 (4%)     |
|              | 5 times per day   | 34 (6%)     |

|            |                                 |           |
|------------|---------------------------------|-----------|
|            | Missing                         | 11 (2%)   |
| Medication | Reported any medication use     | 403 (76%) |
|            | Reported taking antipsychotics  | 318 (60%) |
|            | Medicine weight gain propensity |           |
|            | Low                             | 83 (16%)  |
|            | Medium                          | 59 (11%)  |
|            | High                            | 151 (29%) |
|            | Missing                         | 236 (45%) |
| Depression | PHQ-8 Depression severity       |           |
|            | 0-4 (minimal)                   | 159 (30%) |
|            | 5-9 (mild)                      | 120 (23%) |
|            | 10-14 (moderate)                | 85 (16%)  |
|            | 15-19 (moderately severe)       | 72 (14%)  |
|            | 20-24 (severe)                  | 47 (9%)   |
|            | Missing                         | 46 (9%)   |

The association between diet quality, weight management and mental wellbeing characteristics are presented by BMI classification in Table 2.

**Table 2.** Participant characteristics by BMI classification (n=435).

| Characteristic                        | Healthy weight<br>(n=93) | Overweight<br>(n=119) | Obese<br>(n=223) |
|---------------------------------------|--------------------------|-----------------------|------------------|
| Consumption of fruit and vegetables   |                          |                       |                  |
| four or less portions a day           | 81 (87%)                 | 97 (82%)              | 174 (78%)        |
| Five or more portions a day           | 11 (12%)                 | 21 (18%)              | 48 (22%)         |
| Carbonated drinks consumption         |                          |                       |                  |
| I do not drink carbonated soft drinks | 35 (38%)                 | 50 (42%)              | 84 (38%)         |
| less than one time per day            | 24 (26%)                 | 23 (19%)              | 46 (21%)         |
| 1 time per day                        | 14 (15%)                 | 18 (15%)              | 21 (9%)          |
| 2 times per day                       | 7 (8%)                   | 9 (8%)                | 21 (9%)          |
| 3 times per day                       | 7 (8%)                   | 5 (4%)                | 25 (11%)         |
| 4 times per day                       | 2 (8%)                   | 6 (5%)                | 7 (3%)           |
| 5 times per day                       | 2 (8%)                   | 6 (5%)                | 17 (8%)          |
| Medicine weight gain propensity       |                          |                       |                  |
| low                                   | 10 (11%)                 | 26 (22%)              | 41 (18%)         |
| medium                                | 17 (18%)                 | 9 (8%)                | 27 (12%)         |
| high                                  | 25 (27%)                 | 39 (33%)              | 69 (31%)         |
| Reason for managing weight            |                          |                       |                  |
| gain weight                           | 5 (5%)                   | 1 (1%)                | 1 (0%)           |
| lose weight                           | 23 (25%)                 | 48 (40%)              | 122 (55%)        |
| stay the same weight                  | 28 (30%)                 | 27 (23%)              | 21 (9%)          |
| Gained weight (>6kg)                  |                          |                       |                  |
| No                                    | 69 (74%)                 | 66 (56%)              | 79 (35%)         |
| Yes, once                             | 17 (18%)                 | 36 (30%)              | 83 (37%)         |
| Yes, more than once                   | 4 (4%)                   | 14 (12%)              | 55 (25%)         |
| Reported any medication use           | 80 (86%)                 | 98 (82%)              | 185 (83%)        |
| Reported taking antipsychotics        | 55 (59%)                 | 78 (66%)              | 153 (69%)        |
| Currently managing weight             |                          |                       |                  |
| Managing weight via diet              | 60 (65%)                 | 77 (65%)              | 140 (63%)        |
| Managing weight via exercise          | 26 (28%)                 | 38 (32%)              | 87 (39%)         |
|                                       | 22 (24%)                 | 35 (29%)              | 62 (28%)         |



| PHQ-8 depression severity            |          |          |          |
|--------------------------------------|----------|----------|----------|
| 0-4 (minimal depression)             | 34 (37%) | 37 (31%) | 63 (28%) |
| 5-9 (mild depression)                | 18 (19%) | 33 (28%) | 48 (22%) |
| 10-14 (moderate depression)          | 11 (12%) | 20 (17%) | 36 (16%) |
| 15-19 (moderately severe depression) | 9 (10%)  | 18 (15%) | 31 (14%) |
| 20-24 (severe depression)            | 13 (14%) | 3 (3%)   | 25 (11%) |

NB: data is not presented for two underweight participants and 92 participants whose BMI couldn't be calculated due to missing height or weight data.

#### Association Between Medication Use and Weight Change

Seventy six percent (n=403/529) of participants reported their medication use. The characteristics of the participants that reported medication use are compared to those that did not report medication use in Supplementary Material table. Participants who reported their medication use were more likely to be male and professionally active.

Error! Reference source not found. presents associations between antipsychotic medication use and weight change for the participants that reported any medication use (N=403). The odds of weight gain (gaining >6kg and not losing again) and weight fluctuation (weight gain more than once) were slightly increased in the participants who reported taking antipsychotics (weight gain was reported in 51% of antipsychotic users and 41% of non users, adjusted OR 1.31), but the confidence intervals for both unadjusted and adjusted analyses did not exclude a small decrease. Participants current weight management (yes/no) was not associated with antipsychotic medication use (adjusted OR for managing weight; 0.90, 95% CI 0.53 to 1.53).

**Table 3.** Association between current antipsychotic medication use and weight change for participants that reported medication use (N=403).

| Weight change      |     | Uses   | Does not use                                | Adjusted <sup>b</sup>         |              |      |              |
|--------------------|-----|--|---|-------------------------------|--------------|------|--------------|
|                    |     | antipsychotic medication<br>(N=318)<br>n (%) | antipsychotic medication<br>(N=85)<br>n (%) | Unadjusted <sup>a</sup><br>OR | 95% CI       | OR   | 95% CI       |
| Weight gain        | No  | 146 (46%)                                    | 48 (56%)                                    | 1.52                          | 0.93 to 2.48 | 1.31 | 0.78 to 2.20 |
|                    | Yes | 162 (51%)                                    | 35 (41%)                                    |                               |              |      |              |
| Weight fluctuation | No  | 250 (79%)                                    | 71 (84%)                                    | 1.37                          | 0.70 to 2.70 | 1.62 | 0.80 to 3.25 |
|                    | Yes | 58 (18%)                                     | 12 (14%)                                    |                               |              |      |              |
| Weight management  | No  | 118 (37%)                                    | 27 (32%)                                    | 0.84                          | 0.50 to 1.40 | 0.90 | 0.53 to 1.53 |
|                    | Yes | 198 (62%)                                    | 54 (64%)                                    |                               |              |      |              |

a) Unadjusted based on logistic regression of weight measure against antipsychotic medication use (dependent variable), b) adjusted based on logistic regression of weight measure against antipsychotic medication use (dependent variable), age and sex (simplified to be male vs female, prefer not to say and transgender included in male category due to small numbers). CI confidence interval. OR (Odds ratio); odds of weight measure for those using antipsychotic medication compared to those not using antipsychotic medication.

#### Association Between Attempts at Weight Management and Weight Change

**Table 4.** Association between current weight management and weight gain/fluctuation/change (N=520).

|  | Managing weight | Not managing weight | Unadjusted | Adjusted |
|--|-----------------|---------------------|------------|----------|
|  |                 |                     |            |          |
|  |                 |                     |            |          |

|   |           | (N=327)     | (N=193)     |                              |               |                              |               |
|---|-----------|-------------|-------------|------------------------------|---------------|------------------------------|---------------|
| Weight measure  |           | n (%)       | n (%)       | ORa                          | 95% CI        | ORb                          | 95% CI        |
| Weight gain   | No        | 162 (50%)   | 103 (53%)   | 1.19                         | 0.83 to 1.71  | 1.32                         | 0.91 to 1.93  |
|   | Yes       | 155 (47%)   | 83 (43%)    |                              |               |                              |               |
| Weight fluctuation  | No        | 260 (80%)   | 164 (85%)   | 1.63                         | 0.96 to 2.77  | 1.73                         | 1.00 to 2.96  |
|   | Yes       | 57 (17%)    | 22 (11%)    |                              |               |                              |               |
|   |           |             |             | Mean difference <sup>c</sup> | 95% CI        | Mean difference <sup>c</sup> | 95% CI        |
| Weight change (Difference between lowest and highest adult weight (kg)) | N (%)     | 254 (78%)   | 142 (74%)   | 1.86                         | -2.62 to 6.35 | 2.36                         | -2.14 to 6.86 |
|   | Mean (SD) | 35.1 (21.5) | 33.3 (22.2) |                              |               |                              |               |

Odds ratio (OR) odds of weight gain/weight fluctuation for participants that are currently managing their weight compared to those that aren't based on (a) (unadjusted) logistic regression, (b) adjusted OR based on logistic regression with covariates of age and sex. Mean difference in weight change between participants that are currently managing their weight compared to those that aren't, based on (a) unadjusted linear regression and (b) linear regression adjusted for age and sex.

presents associations between current weight management and adult weight gain. Weight gain (gaining >6kg and not losing it again) was reported by 47% of the participants currently managing their weight and 43% of the participants not currently managing their weight. The odds of having gained were higher in the participants currently managing their weight (adjusted OR 1.32) but the confidence interval was not conclusive. Weight fluctuation was reported in 17% of the participants currently managing their weight and 11 % of the participants not managing their weight. Weight management was associated with higher odds of weight fluctuation (adjusted OR 1.73, 95% CI 1.00 to 2.96).

Weight change (difference between lowest and highest adult weight) was similar in the group that were currently managing weight and not managing weight (35.1 kg compared to 33.3kg). After adjustment for age and sex, the mean difference between the two group was estimated to be 2.36 kg (i.e. weight change is 2.36kg larger in the managing weight group), the confidence interval for this estimate was large (95% CI -2.14 to 6.86) lifetime weight change was not associated with current weight management.

## Discussion

The current study presents the findings of a large cross-sectional survey of adults living with SMI. The survey aimed to understand the current weight, weight change and diet quality of this group and explore the factors associated with weight change.

Levels of obesity were notably higher in the current sample (42%) than in our earlier cohort of adults with SMI (37.5%) [13] and also compared to the general population (28%) [20]. A mean BMI of 31.2kg/m<sup>2</sup> puts the sample overall in the 'obese' range. Diet quality was also evidently poor. Only 17% of our sample met the recommended five portions of fruit and vegetables a day and 6% reported consuming no fruit or vegetables. This is lower than in the general population (where 25% report meeting the recommendations) [21].

Results showed there was no association between weight category and diet quality in this cohort. Meaning that those of higher weight did not consume less fruit and vegetables or drink more carbonated drinks as suggested in the literature. This suggests that excess weight may not be a result of poor diet alone, but may be associated with other factors related to living with SMI such as the side effects of antipsychotic medication, lack of physical activity and/or food poverty. In line with this, Tew et al 2023 showed that those of a healthy weight were more likely to be sufficiently active compared to those living with obesity. Similarly, our current findings indicated that individuals living with obesity were more likely to be those taking antipsychotic medication and to have experienced weight gain and weight fluctuation.

Almost two-thirds (62%) of our sample reported currently managing their weight and this did not differ across weight categories. This is substantially higher than observed in the general population. A large global review showed prevalence rates of 42% and this increased with the prevalence of overweight and obesity [22]. Similarly, recent trends in England show 30% of healthy weight adults are actively trying to lose weight and this increases with overweight (53%) and obesity (76%) [23]. The findings in our cohort may be due to a sense of hopelessness about weight. People with SMI have told us it feels pointless trying to manage weight while taking weight inducing medication. Furthermore, restricting our questions to 'current weight management' may have overlooked multiple past attempts. Future research may want to look at people's optimism regarding their future success in weight management.

In the current study, individuals currently managing their weight were those who were more likely to have experienced weight fluctuation. Weight cycling as seen in this cohort can be serious and has been associated with greater risk of mortality, chronic diseases and depression [24]. There is a body of evidence developing around disordered eating in patients with SMI. Binge eating disorder, emotional eating and reactive eating patterns are commonly observed in people with schizophrenia and bipolar [7,14–16] with less than half of people receiving support [25]. This said, PHQ scores analysed by weight category, did not indicate emotional eating therefore a specific measure of eating behaviour would have been beneficial.

Together, these findings suggest that current weight management options are inadequate in addressing the specific needs of people with SMI who require more regular contact, with tailored materials and support [26]. A recent systematic review highlighted both the lack of effective weight management interventions and understanding of the unique barriers for people with SMI accessing these [27]. Behavioural approaches in isolation have largely been unsuccessful and without consideration of the impact of antipsychotic medication, they only address part of the complex issue.

## Strengths and Limitations

The major strength of this study is the large sample, recruited from a range of localities and clinical settings across the UK, which increases the generalisability of our findings. However, the study was cross-sectional in nature so we were unable to infer causality or the temporal relationship between food consumption, medication use, weight management and weight change. Furthermore, the survey relied on self-report therefore, whilst data was reasonably complete, accurate recall of medication use and food consumption can be poor. Height and weight were the most commonly missing metrics, which meant we were unable to calculate BMI for 92 participants.

We used fruit and vegetable and carbonated drink consumption as a proxy for diet quality however, a more complete assessment of diet quality would be beneficial such as diary methods. In this large-scale survey, we tried to ensure that the questions relating to diet were simple for people to understand and complete in case they were completing the survey unaided. As a result of the autonomy needed to complete the survey, the sample did not include inpatients, so may not reflect the more severe presentations and those on higher doses of antipsychotic medication.

## Conclusions

High levels of obesity are observed in adults with SMI. While diet quality is poor, it is not linked to weight status, suggesting other factors such as antipsychotic medication may be associated with excess weight in this group. Two thirds of these individuals were actively managing their weight, which demonstrates that people with SMI are concerned about their weight and weight gain, however current weight management provision for adults with SMI is inadequate. Further work needs to be done to develop tailored interventions for this group that take into account the side effects of antipsychotic medication.

**Supplementary Materials:** The following supporting information can be downloaded at the website of this paper posted on Preprints.org.

**Author Contributions:** All authors contributed to the study conception and design. Material preparation and data collection were performed by Emily Peckham, Andrew Hill and Gemma Traviss-Turner. Analysis was conducted by Ellen Lee and Peter Pratt. The first draft of the manuscript was written by Gemma Traviss-Turner, Emily Peckham and Ellen Lee. All authors read and approved the final manuscript.

**Financial support:** This research was supported by the NIHR Programme Grant for Applied Research programme (reference 201618) and the NIHR Yorkshire and Humber Applied Research Collaboration.

**Conflicts of interest:** The author(s) declare none.

## Abbreviations

The following abbreviations are used in this manuscript:

*BMI* – Body mass index

*DSM5* – Diagnostic and Statistical Manual version 5

*ICD-10* – International Classification of Disease version 10

*Kg* - Kilogram

*NHS* – National Health Service

*PAQ* - Physical Activity Questionnaire

*PHQ* – Patient Health Questionnaire

*SMI* – Severe mental illness

## References

1. Chesney E, Goodwin GM & Fazel S Risks of all-cause and suicide mortality in mental disorders: a meta-review. .
2. Hayes JF, Marston L, Walters K, et al. (2017) Mortality gap for people with bipolar disorder and schizophrenia: UK-based cohort study 2000–2014. *British Journal of Psychiatry* 211, 175–181. Cambridge University Press.
3. Dregan A, McNeill A, Gaughran F, et al. (2020) Potential gains in life expectancy from reducing amenable mortality among people diagnosed with serious mental illness in the United Kingdom. *PLoS One* 15. Public Library of Science.
4. Fiorillo A & Sartorius N (2021) Mortality gap and physical comorbidity of people with severe mental disorders: the public health scandal. *Ann Gen Psychiatry* 20, 52.
5. Public Health England (2018) Severe mental illness (SMI) and physical health inequalities: briefing. London: .
6. Knifton L & Inglis G (2020) Poverty and mental health: policy, practice and research implications. *BJPsych Bull* 44, 193–196. Royal College of Psychiatrists.
7. Teasdale SB, Ward PB, Rosenbaum S, et al. (2017) Solving a weighty problem: Systematic review and meta-analysis of nutrition interventions in severe mental illness. *British Journal of Psychiatry*, 110–118. Royal College of Psychiatrists.

8. Dayabandara M, Hanwella R, Ratnatunga S, et al. (2017) Antipsychotic-associated weight gain: Management strategies and impact on treatment adherence. *Neuropsychiatr Dis Treat*, 2231–2241. Dove Medical Press Ltd.
9. Dickerson FB, Brown CH, Kreyenbuhl JA, et al. (2006) Obesity among individuals with serious mental illness. *Acta Psychiatr Scand* 113, 306–313.
10. Von Hausswolff-Juhlin Y, Bjartveit M, Lindström E, et al. (2009) Schizophrenia and physical health problems. *Acta Psychiatr Scand* 119, 15–21. John Wiley & Sons, Ltd.
11. Townshend T & Lake A (2017) Obesogenic environments: Current evidence of the built and food environments. *Perspect Public Health*, 38–44. SAGE Publications Ltd.
12. Peckham E, Lorimer B, Spanakis P, et al. (2023) Health-risk behaviours among people with severe mental ill health: Understanding modifiable risk in the Closing the Gap Health Study. *British Journal of Psychiatry* 222, 160–166. Cambridge University Press.
13. Lorimer B, Traviss-Turner GD, Hill A, et al. (2023) Factors associated with low fruit and vegetable consumption among people with severe mental ill health. *Soc Psychiatry Psychiatr Epidemiol*, 1–5.
14. Blouin M, Tremblay A, Jalbert ME, et al. (2008) Adiposity and eating behaviors in patients under second generation antipsychotics. *Obesity* 16, 1780–1787. Nature Publishing Group.
15. Yum SY, Caracci G & Hwang MY (2009) Schizophrenia and Eating Disorders. *Psychiatric Clinics of North America* 32, 809–819.
16. Kouidrat Y, Amad A, Stubbs B, et al. (2018) Disordered eating behaviors as a potential obesogenic factor in schizophrenia. *Psychiatry Res* 269, 450–454.
17. World Health Organisation (2018) Management of physical health conditions in adults with severe mental disorders WHO GUIDELINES. Geneva: .
18. Tew GA, Bailey L, Beeken RJ, et al. (2023) Physical Activity in Adults with Schizophrenia and Bipolar Disorder: A Large Cross-Sectional Survey Exploring Patterns, Preferences, Barriers, and Motivating Factors. *Int J Environ Res Public Health* 20.
19. Huhn M, Nikolakopoulou A, Schneider-Thoma J, et al. (2019) Comparative efficacy and tolerability of 32 oral antipsychotics for the acute treatment of adults with multi-episode schizophrenia: a systematic review and network meta-analysis. *The Lancet* 394, 939–951. Elsevier.
20. NHS Digital (2021) Statistics on Obesity, Physical Activity and Diet (replaced by Statistics on Public Health) (Link: [/data-and-information/publications/statistical/statistics-on-obesity-physical-activity-and-diet](#)) Statistics on Obesity, Physical Activity and Diet, England 2021. Leeds: .
21. NHS Digital (2019) Health Survey for England 2018: Adult health related behaviours. Leeds: .
22. Santos I, Sniechotta FF, Marques MM, et al. (2017) Prevalence of personal weight control attempts in adults: a systematic review and meta-analysis. *Obesity Reviews*, 32–50. Blackwell Publishing Ltd.
23. Piernas C, Aveyard P & Jebb SA (2016) Recent trends in weight loss attempts: Repeated cross-sectional analyses from the health survey for England. *Int J Obes* 40, 1754–1759. Nature Publishing Group.
24. Madigan CD, Pavey T, Daley AJ, et al. (2018) Is weight cycling associated with adverse health outcomes? A cohort study. *Prev Med (Baltim)* 108, 47–52. Academic Press Inc.
25. Abbott S, Dindol N, Tahrani AA, et al. (2018) Binge eating disorder and night eating syndrome in adults with type 2 diabetes: A systematic review. *J Eat Disord*. BioMed Central Ltd.
26. Lee C, Piernas C, Stewart C, et al. (2022) Identifying effective characteristics of behavioral weight management interventions for people with serious mental illness: A systematic review with a qualitative comparative analysis. *Obesity Reviews*. John Wiley and Sons Inc.
27. Stevens H, Smith J, Bussey L, et al. (2022) Weight management interventions for adults living with overweight or obesity and severe mental illness: A systematic review and meta-analysis. *British Journal of Nutrition*. Cambridge University Press.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.